

- [85762] mutated in adult-onset type II citrullinaemia encodes a putative mitochondrial
- [85763] carrier protein. *Nature Genet.* 22: 159–163, 1999.
- [85764] 15035.
- [85765] Saheki, T.; Kobayashi, K.: Mitochondrial aspartate glutamate carrier
- [85766] (citrin) deficiency as the cause of adult-onset type II citrullinemia
- [85767] (CTLN2) and idiopathic neonatal hepatitis (NICCD). *J. Hum. Genet.* 47:
- [85768] 333–341, 2002.
- [85769] 15036.
- [85770] Hara, Y.; Wakamori, M.; Ishii, M.; Maeno, E.; Nishida, M.; Yoshida,
- [85771] T.; Yamada, H.; Shimizu, S.; Mori, E.; Kudoh, J.; Shimizu, S.; Kurose,
- [85772] H.; Okada, Y.; Imoto, K.; Mori, Y.: LTRPC2
- Ca(2+)-permeable channel
- [85773] activated by changes in redox status confers susceptibility to cell
- [85774] death. *Molec. Cell* 9: 163–173, 2002.

[85775] 15037.

[85776] Harteneck, C.; Plant, T. D.; Schultz, G.: From worm to man: three

[85777] subfamilies of TRP channels. Trends Neurosci. 23: 159–166, 2000.

[85778] 15038.

[85779] Kudoh, J.; Nagamine, K.; Asakawa, S.; Abe, I.; Kawasaki, K.; Maeda,

[85780] H.; Tsujimoto, S.; Minoshima, S.; Ito, F.; Shimizu, N.: Localization

[85781] of 16 exons to a 450-kb region involved in the autoimmune polyglandular

[85782] disease type I (APECED) on human chromosome 21q22.3. DNA Res. 4:

[85783] 45–52, 1997.

[85784] 15039.

[85785] Nagamine, K.; Kudoh, J.; Minoshima, S.; Kawasaki, K.; Asakawa,

[85786] S.; Ito, F.; Shimizu, N.: Molecular cloning of a novel putative Ca(2+)

[85787] channel protein (TRPC7) highly expressed in brain. Genomics 54:

- [85788] 124–131, 1998.
- [85789] 15040.
- [85790] Perraud, A.–L.; Fleig, A.; Dunn, C. A.; Bagley, L. A.; Launay,
- [85791] P.; Schmitz, C.; Stokes, A. J.; Zhu, Q.; Bessman, M. J.; Penner, R.;
- [85792] Kinet, J.–P.; Scharenberg, A. M.: ADP–ribose gating of the calcium–permeable
- [85793] LTRPC2 channel revealed by Nudix motif homology. Nature 411: 595–599,
- [85794] 2001.
- [85795] 15041.
- [85796] Sano, Y.; Inamura, K.; Miyake, A.; Mochizuki, S.; Yokoi, H.; Matsushime,
- [85797] H.; Furuichi, K.: Immunocyte Ca(2+) influx system mediated by LTRPC2. Science 293:
- [85798] 1327–1330, 2001.
- [85799] 15042.
- [85800] Koyama, N.; Ishibashi, K.; Kuwahara, M.; Inase, N.; Ichioka, M.;
- [85801] Sasaki, S.; Marumo, F.: Cloning and functional expression of human

- [85802] aquaporin8 cDNA and analysis of its gene. Genomics 54: 169–172, 1998.
- [85803] 1998.
- [85804] 15043.
- [85805] Viggiano, L.; Rocchi, M.; Svelto, M.; Calamita, G.: Assignment of the aquaporin–8 water channel gene (AQP8) to human chromosome 16p12. Cytogenet. Cell Genet. 84: 208–210, 1999.
- [85806] 15044.
- [85807] Sperandio, M. P.; Borsani, G.; Incerti, B.; Zollo, M.; Rossi, E.;
- [85808] Zuffardi, O.; Castaldo, P.; Taglialatela, M.; Andria, G.; Sebastio, G.: The gene encoding a cationic amino acid transporter (SLC7A4) maps to the region deleted in the velocardiofacial syndrome. Genomics 49: 230–236, 1998.
- [85809] 15045.
- [85810] Koegl, M.; Hoppe, T.; Schlenker, S.; Ulrich, H. D.; Mayer, T.

U.;

[85816] Jentsch, S.: A novel ubiquitination factor, E4, is involved in multiubiquitin

[85817] chain assembly. Cell 96: 635–644, 1999.

[85818] 15046.

[85819] Nonaka, S.; Tanaka, Y.; Okada, Y.; Takeda, S.; Harada, A.; Kanai,

[85820] Y.; Kido, M.; Hirokawa, N.: Randomization of left–right asymmetry

[85821] due to loss of nodal cilia generating leftward flow of extraembryonic

[85822] fluid in mice lacking KIF3B motor protein. Cell 95: 829–837, 1998.

[85823] 15047.

[85824] Yamazaki, H.; Nakata, T.; Okada, Y.; Hirokawa, N.: KIF3A/B: a

[85825] heterodimeric kinesin superfamily protein that works as a microtubule

[85826] plus end–directed motor for membrane organelle transport. J. Cell

[85827] Biol. 130: 1387–1399, 1995.

[85828] 15048.

- [85829] Suzuki, N.; Zara, J.; Sato, T.; Ong, E.; Bakhiet, N.; Oshima, R.
- [85830] G.; Watson, K. L.; Fukuda, M. N.: A cytoplasmic protein, bystin,
- [85831] interacts with trophinin, tastin, and cytokeratin and may be involved
- [85832] in trophinin-mediated cell adhesion between trophoblast and endometrial
- [85833] epithelial cells. Proc. Nat. Acad. Sci. 95: 5027-5032, 1998.
- [85834] 15049.
- [85835] Hurskainen, T. L.; Hirohata, S.; Seldin, M. F.; Apte, S. S.: ADAM-TS5,
- [85836] ADAM-TS6, and ADAM-TS7, novel members of a new family of zinc metalloproteases:
- [85837] general features and genomic distribution of the ADAM-TS family. J.
- [85838] Biol. Chem. 274: 25555-25563, 1999.
- [85839] 15050.
- [85840] Tang, B. L.; Hong, W.: ADAMTS: a novel family of proteases with
- [85841] an ADAM protease domain and thrombospondin 1 re-

peats. FEBS Lett. 445:

[85842] 223–225, 1999.

[85843] 15051.

[85844] Tortorella, M. D.; Burn, T. C.; Pratta, M. A.; Abbaszade, I.;
Hollis,

[85845] J. M.; Liu, R.; Rosenfeld, S. A.; Copeland, R. A.; Decicco, C.
P.;

[85846] Wynn, R.; Rockwell, A.; Yang, F.; and 16 others: Purifica-
tion and

[85847] cloning of aggrecanase-1: a member of the ADAMTS fam-
ily of proteins. Science 284:

[85848] 1664–1666, 1999.

[85849] 15052.

[85850] Shiratsuchi, T.; Oda, K.; Nishimori, H.; Suzuki, M.; Taka-
hashi,

[85851] E.; Tokino, T.; Nakamura, Y.: Cloning and characterization
of BAP3

[85852] (BAI-associated protein 3), a C2 domain-containing pro-
tein that interacts

[85853] with BAI1. Biochem. Biophys. Res. Commun. 251:
158–165, 1998.

[85854] 15053.

- [85855] Almeida, R.; Amado, M.; David, L.; Levery, S. B.; Holmes, E. H.;
- [85856] Merckx, G.; Geurts van Kessel, A.; Rygaard, E.; Hassan, H.; Bennett,
- [85857] E.; Clausen, H.: A family of human beta-4-galactosyltransferases:
- [85858] cloning and expression of two novel UDP-galactose:beta-N-acetylglucosamine
- [85859] beta-1,4-galactosyltransferases, beta-4-Gal-T2 and beta-4-Gal-T3. J.
- [85860] Biol. Chem. 272: 31979-31991, 1997.
- [85861] 15054.
- [85862] Sato, T.; Furukawa, K.; Bakker, H.; Van den Eijnden, D. H.; Van
- [85863] Die, I.: Molecular cloning of a human cDNA encoding beta-1,4-galactosyltransferase
- [85864] with 37% identity to mammalian UDP-Gal:GlcNAc beta-1,4-galactosyltransferase. Proc.
- [85865] Nat. Acad. Sci. 95: 472-477, 1998.
- [85866] 15055.
- [85867] Shaper, N.: Personal Communication. Baltimore, Md. 11/2/1999.

[85868] 15056.

[85869] Gundelfinger, E.: Personal Communication. Madgeburg, FRG. 1/8/1999.

[85870] 15057.

[85871] Hashida, H.; Goto, J.; Zhao, N.; Takahashi, N.; Hirai, M.; Kanazawa,

[85872] I.; Sakaki, Y.: Cloning and mapping of ZNF231, a novel brain-specific

[85873] gene encoding neuronal double zinc finger protein whose expression

[85874] is enhanced in a neurodegenerative disorder, multiple system atrophy

[85875] (MSA). Genomics 54: 50-58, 1998.

[85876] 15058.

[85877] Sparks, A. B.; Peterson, S. N.; Bell, C.; Loftus, B. J.; Hocking,

[85878] L.; Cahill, D. P.; Frassica, F. J.; Streeten, E. A.; Levine, M. A.;

[85879] Fraser, C. M.; Adams, M. D.; Broder, S.; Venter, J. C.; Vogelstein,

[85880] B.; Ralston, S. H.: Mutation screening of the TNFRSF11A gene encoding

- [85881] receptor activator of NF- κ B (RANK) in familial and sporadic Paget's
- [85882] disease of bone and osteosarcoma. *Calcif. Tissue Int.* 68: 151–155,
- [85883] 2001.
- [85884] 15059.
- [85885] Santorelli, F. M.; Patrono, C.; Fortini, D.; Tessa, A.; Comanducci,
- [85886] G.; Bertini, E.; Pierallini, A.; Amabile, G. A.; Casali, C.: Intrafamilial
- [85887] variability in hereditary spastic paraplegia associated with an SPG4
- [85888] gene mutation. *Neurology* 55: 702–705, 2000.
- [85889] 15060.
- [85890] Sauter, S.; Mitterski, B.; Klimpe, S.; Bonsch, D.; Schols, L.;
- [85891] Visbeck, A.; Papke, T.; Hopf, H. C.; Engel, W.; Deufel, T.; Epplen,
- [85892] J. T.; Neesen, J.: Mutation analysis of the spastin gene (SPG4) in
- [85893] patients in Germany with autosomal dominant hereditary spastic paraplegia. *Hum.*

- [85894] *Mutat.* 20: 127–132, 2002.
- [85895] 15061.
- [85896] Svenson, I. K.; Ashley–Koch, A. E.; Pericak–Vance, M. A.;
Marchuk,
- [85897] D. A.: A second leaky splice–site mutation in the spastin
gene. (Letter) *Am.*
- [85898] *J. Hum. Genet.* 69: 1407–1409, 2001.
- [85899] 15062.
- [85900] Flannery, C. R.; Hughes, C. E.; Schumacher, B. L.; Tudor,
D.; Aydelotte,
- [85901] M. B.; Kuettner, K. E.; Caterson, B.: Articular cartilage su-
perficial
- [85902] zone protein (SZP) is homologous to megakaryocyte stim-
ulating factor
- [85903] precursor and is a multifunctional proteoglycan with po-
tential growth–promoting,
- [85904] cytoprotective, and lubricating properties in cartilage
metabolism. *Biochem.*
- [85905] *Biophys. Res. Commun.* 254: 535–541, 1999.
- [85906] 15063.
- [85907] Ikegawa, S.; Sano, M.; Koshizuka, Y.; Nakamura, Y.: Isola–

tion,

[85908] characterization and mapping of the mouse and human PRG4 (proteoglycan

[85909] 4) genes. *Cytogenet. Cell Genet.* 90: 291–297, 2000.

[85910] 15064.

[85911] Merberg, D. M.; Fitz, L. J.; Temple, P.; Giannotti, J.; Murtha,

[85912] P.; Fitzgerald, M.; Scaltreto, H.; Kelleher, K.; Preissner, K.; Kriz,

[85913] R.; Jacobs, K.; Turner, K.: In: Preissner, K. T.; Rosenblatt, S.;

[85914] Kost, C.; Wegerhoff, J.; Mosher, D. F. (eds.): *Biology of Vitronectins*

[85915] and Their Receptors. Elsevier Science, B.V. 1993. Pp. 45–53.

[85916] 15065.

[85917] Schumacher, B. L.; Block, J. A.; Schmid, T. M.; Aydelotte, M. B.;

[85918] Kuettner, K. E.: A novel proteoglycan synthesized and secreted by

[85919] chondrocytes of the superficial zone of articular cartilage. *Arch.*

[85920] *Biochem. Biophys.* 15: 144–152, 1994.

[85921] 15066.

[85922] Pfeiffer, R.; et al; et al: Amino acid transport of
y(0,+)-type

[85923] by heterodimers of 4F2hc/CD98 and members of the gly-
coprotein-associated

[85924] amino acid transporter family. EMBO J. 18: 49-57, 1999.

[85925] 15067.

[85926] Pineda, M.; et al.; et al: Identification of a membrane pro-
tein

[85927] (LAT-2) that co-expresses with 4F2hc an L type amino
acid transport

[85928] activity with broad specificity for small and large zwitteri-
onic amino

[85929] acids. J. Biol. Chem. 274: 19738-19744, 1999.

[85930] 15068.

[85931] Sato, H.; Tamba, M.; Ishii, T.; Bannai, S.: Cloning and ex-
pression

[85932] of a plasma membrane cystine/glutamate exchange
transporter composed

[85933] of two distinct proteins. J. Biol. Chem. 274: 11455-11458,
1999.

[85934] 15069.

[85935] Torrents, D.; et al.; et al: Identification and characterization

[85936] of a membrane protein (y+L amino acid transporter-1) that associates

[85937] with 4F2hc to encode the amino acid transport activity y+L: a candidate

[85938] gene for lysinuric protein intolerance. J. Biol. Chem. 273: 32437-32445,

[85939] 1998.

[85940] 15070.

[85941] Dominguez, A.; Ramos-Morales, F.; Romero F.; Rios R. M.; Dreyfus,

[85942] F.; Tortolero, M.; Pintor-Toro, J. A.: hpttg, a human homologue of

[85943] rat pttg, is overexpressed in hematopoietic neoplasms: evidence for

[85944] a transcriptional activation function of hPTTG. Oncogene 17: 2187-2193,

[85945] 1998.

[85946] 15071.

[85947] Ishikawa, H.; Heaney, A. P.; Yu, R.; Horwitz, G. A.; Melmed, S.

[85948] : Human pituitary tumor-transforming gene induces angiogenesis. J.

[85949] Clin. Endocr. Metab. 86: 867-874, 2001.

[85950] 15072.

[85951] Jallepalli, P. V.; Waizenegger, I. C.; Bunz, F.; Langer, S.; Speicher,

[85952] M. R.; Peters, J.-M.; Kinzler, K. W.; Vogelstein, B.; Lengauer, C.

[85953] : Securin is required for chromosomal stability in human cells. Cell 105:

[85954] 445-457, 2001.

[85955] 15073.

[85956] Kakar, S. S.: Assignment of the human tumor transforming gene

[85957] TUTR1 to chromosome band 5q35.1 by fluorescence in situ hybridization. Cytogenet.

[85958] Cell Genet. 83: 93-95, 1998.

[85959] 15074.

[85960] Pei, L.: Pituitary tumor-transforming gene protein asso-

ciates

[85961] with ribosomal protein S10 and a novel human homologue of DnaJ in

[85962] testicular cells. J. Biol. Chem. 274: 3151–3158, 1999.

[85963] 15075.

[85964] Wang, Z.; Yu, R.; Melmed, S.: Mice lacking pituitary tumor transforming

[85965] gene show testicular and splenic hypoplasia, thymic hyperplasia, thrombocytopenia,

[85966] aberrant cell cycle progression, and premature centromere division. Molec.

[85967] Endocr. 15: 1870–1879, 2001. 1. Dominguez, A.; Ramos-Morales, F.;

[85968] Romero, F.; Rios, R. M.; Dreyfus,

[85969] 15076.

[85970] Yu, R.; Ren, S.-G.; Horwitz, G. A.; Wang, Z.; Melmed, S.: Pituitary

[85971] tumor transforming gene (PTTG) regulates placental JEG-3 cell division

[85972] and survival: evidence from live cell imaging. Molec. Endocr. 14:

[85973] 1137–1146, 2000.

[85974] 15077.

[85975] Zhang, X.; Horwitz, G. A.; Heaney, A. P.; Nakashima, M.;
Prezant,

[85976] T. R.; Bronstein, M. D.; Melmed, S.: Pituitary tumor trans-
forming

[85977] gene (PTTG) expression in pituitary adenomas. J. Clin. En-
docr. Metab. 84:

[85978] 761–767, 1999.

[85979] 15078.

[85980] Zhang, X.; Horwitz, G. A.; Prezant, T. R.; Valentini, A.;
Nakashima,

[85981] M.; Bronstein, M. D.; Melmed, S.: Structure, expression,
and function

[85982] of human pituitary tumor-transforming gene (PTTG).
Molec. Endocr. 13:

[85983] 156–166, 1999.

[85984] 15079.

[85985] Pennica, D.; Swanson, T. A.; Welsh, J. W.; Roy, M. A.;
Lawrence,

[85986] D. A.; Lee, J.; Brush, J.; Taneyhill, L. A.; Deuel, B.; Lew, M.;
Watanabe,

- [85987] C.; Cohen, R. L.; Melhem, M. F.; Finley, G. G.; Quirke, P.;
Goddard,
[85988] A. D.; Hillan, K. J.; Gurney, A. L.; Botstein, D.; Levine, A. J.:
[85989] WISP genes are members of the connective tissue growth
factor family
[85990] that are up-regulated in Wnt-1-transformed cells and
aberrantly expressed
[85991] in human colon tumors. Proc. Nat. Acad. Sci. 95:
14717-14722, 1998.
[85992] 15080.
[85993] Tanaka, S.; Sugimachi, K.; Saeki, H.; Kinoshita, J.; Ohga,
T.;
[85994] Shimada, M.; Maehara, Y.; Sugimachi, K.: A novel variant
of WISP1
[85995] lacking a von Willebrand type C module overexpressed in
scirrhous
[85996] gastric carcinoma. Oncogene 20: 5525-5532, 2001.
[85997] 15081.
[85998] Feng, L.; Novak, E. K.; Hartnell, L. M.; Bonifacino, J. S.;
Collinson,
[85999] L. M.; Swank, R. T.: The Hermansky-Pudlak syndrome 1
(HPS1) and HPS2

- [86000] genes independently contribute to the production and function of platelet
- [86001] dense granules, melanosomes, and lysosomes. Blood 99: 1651–1658,
- [86002] 2002.
- [86003] 15082.
- [86004] Vanhooren, J. C. T.; Marynen, P.; Mannaerts, G. P.; Van Veldhoven,
- [86005] P. P.: Evidence for the existence of a pristanoyl-CoA oxidase gene
- [86006] in man. Biochem. J. 325: 593–599, 1997.
- [86007] 15083.
- [86008] Gee, S.; Krauss, S. W.; Miller, E.; Aoyagi, K.; Arenas, J.; Conboy,
- [86009] J. G.: Cloning of mDEAH9, a putative RNA helicase and mammalian homologue
- [86010] of *Saccharomyces cerevisiae* splicing factor Prp43. Proc. Nat. Acad.
- [86011] Sci. 94: 11803–11807, 1997.
- [86012] 15084.
- [86013] Imamura, O.; Sugawara, M.; Furuichi, Y.: Cloning and

characterization

[86014] of a putative human RNA helicase gene of the DEAH-box protein family. *Biochem.*

[86015] *Biophys. Res. Commun.* 240: 335–340, 1997.

[86016] 15085.

[86017] Del Mastro, R. G.; Wang, L.; Simmons, A. D.; Gallardo, T. D.; Clines,

[86018] G. A.; Ashley, J. A.; Hilliard, C. J.; Wasmuth, J. J.; McPherson,

[86019] J. D.; Lovett, M.: Human chromosome-specific cDNA libraries: new

[86020] tools for gene identification and genome annotation.

Genome Res. 5:

[86021] 185–194, 1995.

[86022] 15086.

[86023] Sano, K.; Tanihara, H.; Heimark, R. L.; Obata, S.; Davidson, M.;

[86024] St. John, T.; Taketani, S.; Suzuki, S.: Protocadherins: a large family

[86025] of cadherin-related molecules in central nervous system. *EMBO J.* 12:

[86026] 2249–2256, 1993.

- [86027] 15087.
- [86028] Frigerio, J.-M.; Berthezene, P.; Garrido, P.; Ortiz, E.;
Barthelley,
- [86029] S.; Vasseur, S.; Sastre, B.; Seleznieff, I.; Dagorn, J. C.; Iovanna,
vanna,
- [86030] J. L.: Analysis of 2166 clones from a human colorectal
cancer cDNA
- [86031] library by partial sequencing. Hum. Molec. Genet. 4:
37-43, 1995.
- [86032] 15088.
- [86033] Frigerio, J.-M.; Dagorn, J.-C.; Iovanna, J. L.: Cloning, se-
quencing
- [86034] and expression of the L5, L21, L27a, L28, S5, S9, S10 and
S29 human
- [86035] ribosomal protein mRNAs. Biochim. Biophys. Acta 1262:
64-68, 1995.
- [86036] 15089.
- [86037] Dominguez, O.; Ashhab, Y.; Sabater, L.; Belloso, E.; Caro,
P.;
- [86038] Pujol-Borrell, R.: Cloning of ARE-containing genes by AU-
motif-directed
- [86039] display. Genomics 54: 278-286, 1998.

[86040] 15090.

[86041] Kostrub, C. F.; Knudsen, K.; Subramani, S.; Enoch, T.:
Hus1p,

[86042] a conserved fission yeast checkpoint protein, interacts
with Rad1p

[86043] and is phosphorylated in response to DNA damage. EMBO
J. 17: 2055–2066,

[86044] 1998.

[86045] 15091.

[86046] Lieberman, H. B.; Hopkins, K. M.; Nass, M.; Demetrick, D.;
Davey,

[86047] S.: A human homolog of the Schizosaccharomyces pombe
rad9+ checkpoint

[86048] control gene. Proc. Nat. Acad. Sci. 93: 13890–13895,
1996.

[86049] 15092.

[86050] Stocker, M.; Kerschensteiner, D.: Cloning and tissue dis-
tribution

[86051] of two new potassium channel alpha-subunits from rat
brain. Biochem.

[86052] Biophys. Res. Commun. 248: 927–934, 1998.

[86053] 15093.

[86054] Seki, N.; Hattori, A.; Hayashi, A.; Kozuma, S.; Hori, T.;
Saito,

[86055] T.: The human regulator of G-protein signaling protein 6
gene (RGS6)

[86056] maps between markers WI-5202 and D14S277 on chro-
mosome 14q24.3. J.

[86057] Hum. Genet. 44: 138-140, 1999.

[86058] 15094.

[86059] Leegwater, P. A. J.; Konst, A. A. M.; Kuyt, B.; Sandkuijl, L.
A.;

[86060] Naidu, S.; Oudejans, C. B. M.; Schutgens, R. B. H.; Pronk, J.
C.;

[86061] van der Knaap, M. S.: The gene for leukoencephalopathy
with vanishing

[86062] white matter is located on chromosome 3q27. Am. J.
Hum. Genet. 65:

[86063] 728-734, 1999.

[86064] 15095.

[86065] tom Dieck, S.; Sanmarti-Vila, L.; Langnaese, K.; Richter, K.;
Kindler,

- [86066] S.; Soyke, A.; Wex, H.; Smalla, K.-H.; Kampf, U.; Franzer, J.-T.;
- [86067] Stumm, M.; Garner, C. C.; Gundelfinger, E. D.: Bassoon, a novel zinc-finger
- [86068] CAG/glutamine-repeat protein selectively localized at the active zone
- [86069] of presynaptic nerve terminals. J. Cell Biol. 142: 499-509, 1998.
- [86070] 15096.
- [86071] George, A. L., Jr.; Komisarof, J.; Kallen, R. G.; Barchi, R. L.
- [86072] : Primary structure of the adult human skeletal muscle voltage-dependent
- [86073] sodium channel. Ann. Neurol. 31: 131-137, 1992.
- [86074] 15097.
- [86075] Heine, R.; Pika, U.; Lehmann-Horn, F.: A novel SCN4A mutation
- [86076] causing myotonia aggravated by cold and potassium. Hum. Molec. Genet. 2:
- [86077] 1349-1353, 1993.
- [86078] 15098.
- [86079] Meyer-Kleine, C.; Otto, M.; Zoll, B.; Koch, M. C.: Molecular

- [86080] and genetic characterization of German families with paramyotonia
- [86081] congenita and demonstration of founder effect in the Ravensberg families. Hum.
- [86082] Genet. 93: 707–710, 1994.
- [86083] 15099.
- [86084] Orrell, R. W.; Jurkat-Rott, K.; Lehmann-Horn, F.; Lane, R. J. M.: Familial cramp due to potassium-aggravated myotonia. J. Neurol.
- [86086] Neurosurg. Psychiat. 65: 569–572, 1998.
- [86087] 15100.
- [86088] Rojas, C. V.: Personal Communication. Pittsburgh, Pa. 2/6/1992.
- [86089] 15101.
- [86090] Rojas, C. V.; Wang, J.; Schwartz, L. S.; Hoffman, E. P.; Powell, B. R.; Brown, R. H., Jr.: A met-to-val mutation in the skeletal muscle Na⁺ channel alpha-subunit in hyperkalaemic periodic paralysis. Nature 354: 387–389, 1991.

[86094] 15102.

[86095] Winter, c.; tom Dieck, S.; Boeckers, T. M.; Bockmann, J.;
Kampf,

[86096] U.; Sanmarti-Vila, L.; Langnaese, K.; Altrock, W.; Stumm,
M.; Soyke,

[86097] A.; Wieacker, P.; Garner, C. C.; Gundelfinger, E. D.: The
presynaptic

[86098] cytomatrix protein Bassoon: sequence and chromosomal
localization

[86099] of the human BSN gene. Genomics 57: 389–397, 1999.

[86100] 15103.

[86101] Bui, T. D.; Levy, E. R.; Subramaniam, V. N.; Lowe, S. L.;
Hong,

[86102] W.: cDNA characterization and chromosomal mapping of
human Golgi

[86103] SNARE GS27 and GS28 to chromosome 17. Genomics 57:
285–288, 1999.

[86104] 15104.

[86105] Lowe, S. L.; Peter, F.; Subramaniam, V. N.; Wong, S. H.;
Hong,

[86106] W.: A SNARE involved in protein transport through the
Golgi apparatus. Nature 389:

- [86107] 881–884, 1997.
- [86108] 15105.
- [86109] Nagahama, M.; Orci, L.; Ravazzola, M.; Amherdt, M.; La-comis, L.;
- [86110] Tempst, P.; Rothman, J. E.; Sollner, T. H.: A v-SNARE im-plicated
- [86111] in intra-Golgi transport. J. Cell Biol. 133: 507–516, 1996.
- [86112] 15106.
- [86113] Subramaniam, V. N.; Peter, F.; Philp, R.; Wong, S. H.; Hong, W.
- [86114] : GS28, a 28-kilodalton Golgi SNARE that participates in ER-Golgi
- [86115] transport. Science 272: 1161–1163, 1996.
- [86116] 15107.
- [86117] Hay, J. C.; Chao, D. S.; Kuo, C. S.; Scheller, R. H.: Protein
- [86118] interactions regulating vesicle transport between the en-doplasmic
- [86119] reticulum and Golgi apparatus in mammalian cells. Cell 89: 149–158,
- [86120] 1997.

- [86121] 15108.
- [86122] Li, J.; Ding, S.-F.; Habib, N. A.; Fermor, B. F.; Wood, C. B.;
- [86123] Gilmour, R. S.: Partial characterization of a cDNA for human stearyl-CoA
- [86124] desaturase and changes in its mRNA expression in some normal and malignant
- [86125] tissues. *Int. J. Cancer* 57: 348-352, 1994.
- [86126] 15109.
- [86127] Ntambi, J. M.; Miyazaki, M.; Stoeckl, J. P.; Lan, H.; Kendziora,
- [86128] C. M.; Yandell, B. S.; Song, Y.; Cohen, P.; Friedman, J. M.; Attie,
- [86129] A. D.: Loss of stearyl-CoA desaturase-1 function protects mice against
- [86130] adiposity. *Proc. Nat. Acad. Sci.* 99: 11482-11486, 2002.
- [86131] 15110.
- [86132] Thiede, M. A.; Ozols, J.; Strittmatter, P.: Construction and sequence
- [86133] of cDNA for rat liver stearyl coenzyme A desaturase. *J. Biol. Chem.* 261:
- [86134] 13230-13235, 1986.

- [86135] 15111.
- [86136] Zhang, L.; Ge, L.; Parimoo, S.; Stenn, K.; Prouty, S. M.: Human
man
- [86137] stearyl-CoA desaturase: alternative transcripts generated
from a
- [86138] single gene by usage of tandem polyadenylation sites.
Biochem. J. 340:
- [86139] 255–264, 1999.
- [86140] 15112.
- [86141] Zheng, Y.; Eilertsen, K. J.; Ge, L.; Zhang, L.; Sundberg, J. P.;
- [86142] Prouty, S. M.; Stenn, K. S.; Parimoo, S.: Scd1 is expressed
in sebaceous
- [86143] glands and is disrupted in the asebia mouse. (Letter) Na-
ture Genet. 23:
- [86144] 268–270, 1999.
- [86145] 15113.
- [86146] Harding, H. P.; Zeng, H.; Zhang, Y.; Jungries, R.; Chung, P.;
Plesken,
- [86147] H.; Sabatini, D. D.; Ron, D.: Diabetes mellitus and exocrine
pancreatic
- [86148] dysfunction in Perk $-/-$ mice reveals a role for transla-

tional control

[86149] in secretory cell survival. *Molec. Cell* 7: 1153–1163, 2001.

[86150] 15114.

[86151] Harding, H. P.; Zhang, Y.; Bertolotti, A.; Zeng, H.; Ron, D.:

[86152] Perk is essential for translational regulation and cell survival during

[86153] the unfolded protein response. *Molec. Cell* 5: 897–904, 2000.

[86154] 15115.

[86155] Duh, F.-M.; Latif, F.; Weng, Y.; Geil, L.; Modi, W.; Stackhouse,

[86156] T.; Matsumura, F.; Duan, D. R.; Linehan, W. M.; Lerman, M. I.; Gnarr,

[86157] J. R.: cDNA cloning and expression of the human homolog of the sea

[86158] urchin fascin and *Drosophila* singed genes which encodes an actin-bundling

[86159] protein. *DNA Cell Biol.* 13: 821–827, 1994.

[86160] 15116.

[86161] Mosialos, G.; Birkenbach, M.; Ayehunie, S.; Matsumura, F.; Pinkus,

- [86162] G. S.; Kieff, E.; Langhoff, E.: Circulating human dendritic cells
- [86163] differentially express high levels of a 55-kd actin-bundling protein. Am. J. Path. 148: 593-600, 1996.
- [86164] 15117.
- [86165] Ono, S.; Yamakita, Y.; Yamashiro, S.; Matsudaira, P. T.; Gnarra, J. R.; Obinata, T.; Matsumura, F.: Identification of an actin binding region and a protein kinase C phosphorylation site on human fascin. J. Biol. Chem. 272: 2527-2533, 1997.
- [86166] 15118.
- [86167] Pinkus, G. S.; Pinkus, J. L.; Langhoff, E.; Matsumura, F.; Yamashiro, S.; Mosialos, G.; Said, J. W.: Fascin, a sensitive new marker for Reed-Sternberg cells of Hodgkin's disease: evidence for a dendritic or B cell derivation? Am. J. Path. 150: 543-562, 1997.

- [86175] 15119.
- [86176] Sonderbye, L.; Magerstadt, R.; Blatman, R. N.; Preffer, F. I.;
- [86177] Langhoff, E.: Selective expression of human fascin (p55)
by dendritic
- [86178] leukocytes. Adv. Exp. Med. Biol. 471: 41–46, 1997.
- [86179] 15120.
- [86180] Yamakita, Y.; Ono, S.; Matsumura, F.; Yamashiro, S.: Phos-
phorylation
- [86181] of human fascin inhibits its actin binding and bundling
activities. J.
- [86182] Biol. Chem. 271: 12632–12638, 1996.
- [86183] 15121.
- [86184] Yamashiro–Matsumura, S.; Matsumura, F.: Intracellular lo-
calization
- [86185] of the 55–kD actin–bundling protein in cultured cells:
spatial relationships
- [86186] with actin, alpha–actinin, tropomyosin, and fimbrin. J. Cell
Biol. 103:
- [86187] 631–640, 1986.
- [86188] 15122.

- [86189] Yamashiro–Matsumura, S.; Matsumura, F.: Purification and characterization
of an F–actin–bundling 55–kilodalton protein from HeLa
cells. J.
Biol. Chem. 260: 5087–5097, 1985.
15123.
- [86193] Gibbs, P. E. M.; McGregor, W. G.; Maher, V. M.; Nisson, P.;
Lawrence,
C. W.: A human homolog of the *Saccharomyces cerevisiae*
REV3 gene,
which encodes the catalytic subunit of DNA polymerase
zeta. Proc.
Nat. Acad. Sci. 95: 6876–6880, 1998.
15124.
- [86198] Morelli, C.; Mungall, A. J.; Negrini, M.; Barbanti–Brodano,
G.;
Croce, C. M.: Alternative splicing, genomic structure, and
fine chromosome
localization of REV3L. Cytogenet. Cell Genet. 83: 18–20,
1998.
15125.

- [86202] Xiao, W.; Lechler, T.; Chow, B. L.; Fontanie, T.; Agustus, M.;
- [86203] Carter, K. C.; Wei, Y.-F.: Identification, chromosomal mapping and
- [86204] tissue-specific expression of hREV3 encoding a putative human DNA
- [86205] polymerase zeta. *Carcinogenesis* 19: 945-949, 1998.
- [86206] 15126.
- [86207] Kahn, M. L.; Zheng, Y.-W.; Huang, W.; Bigornia, V.; Zeng, D.; Moff,
- [86208] S.; Farese, R. V., Jr.; Tam, C.; Coughlin, S. R.: A dual thrombin
- [86209] receptor system for platelet activation. *Nature* 394: 690-694, 1998.
- [86210] 15127.
- [86211] Xu, W.-F.; Andersen, H.; Whitmore, T. E.; Presnell, S. R.; Yee,
- [86212] D. P.; Ching, A.; Gilbert, T.; Davie, E. W.; Foster, D. C.: Cloning
- [86213] and characterization of human protease-activated receptor 4. *Proc.*
- [86214] *Nat. Acad. Sci.* 95: 6642-6646, 1998.

[86215] 15128.

[86216] Santoro, B.; Grant, S. G. N.; Bartsch, D.; Kandel, E. R.: In-
teractive

[86217] cloning with the SH3 domain of N-src identifies a new
brain specific

[86218] ion channel protein, with homology to Eag and cyclic nu-
cleotide-gated

[86219] channels. Proc. Nat. Acad. Sci. 94: 14815-14820, 1997.

[86220] 15129.

[86221] Santoro, B.; Liu, D. T.; Yao, H.; Bartsch, D.; Kandel, E. R.;
Siegelbaum,

[86222] S. A.; Tibbs, G. R.: Identification of a gene encoding a hy-
perpolarization-activated

[86223] pacemaker channel of brain. Cell 93: 717-729, 1998.

[86224] 15130.

[86225] Wainger, B. J.; DeGennaro, M.; Santoro, B.; Siegelbaum, S.
A.;

[86226] Tibbs, G. R.: Molecular mechanism of cAMP modulation of
HCN pacemaker

[86227] channels. Nature 411: 805-810, 2001.

[86228] 15131.

- [86229] Ludwig, A.; Zong, X.; Stieber, J.; Hullin, R.; Hofmann, F.; Biel,
- [86230] M.: Two pacemaker channels from human heart with profoundly different
- [86231] activation kinetics. EMBO J. 18: 2323–2329, 1999.
- [86232] 15132.
- [86233] Casari, G.; De Fusco, M.; Ciarmatori, S.; Zeviani, M.; Mora, M.;
- [86234] Fernandez, P.; De Michele, G.; Filla, A.; Coccozza, S.; Marconi, R.;
- [86235] Durr, A.; Fontaine, B.; Ballabio, A.: Spastic paraplegia and OXPHOS
- [86236] impairment caused by mutations in paraplegin, a nuclear-encoded mitochondrial
- [86237] metalloprotease. Cell 93: 973–983, 1998.
- [86238] 15133.
- [86239] De Michele, G.; De Fusco, M.; Cavalcanti, F.; Filla, A.; Marconi,
- [86240] R.; Volpe, G.; Monticelli, A.; Ballabio, A.; Casari, G.; Coccozza,
- [86241] S.: A new locus for autosomal recessive hereditary spastic paraplegia

- [86242] maps to chromosome 16q24.3. Am. J. Hum. Genet. 63: 135–139, 1998.
- [86243] 15134.
- [86244] Settasatian, C.; Whitmore, S. A.; Crawford, J.; Bilton, R. L.;
- [86245] Cleton-Jansen, A.-M.; Sutherland, G. R.; Callen, D. F.: Ge-
nomic structure
- [86246] and expression analysis of the spastic paraplegia gene,
SPG7. Hum.
- [86247] Genet. 105: 139–144, 1999.
- [86248] 15135.
- [86249] Nykjaer, A.; Dragun, D.; Walther, D.; Vorum, H.; Jacobsen,
C.;
- [86250] Herz, J.; Melsen, F.; Christensen, E. I.; Willnow, T. E.: An
endocytic
- [86251] pathway essential for renal uptake and activation of the
steroid 25-(OH)
- [86252] vitamin D3. Cell 96: 507–515, 1999.
- [86253] 15136.
- [86254] Quelle, D. E.; Ashmun, R. A.; Hannon, G. J.; Rehberger, P.
A.;
- [86255] Trono, D.; Richter, K. H.; Walker, C.; Beach, D.; Sherr, C. J.;

Serrano,

[86256] M.: Cloning and characterization of murine p16(INK4a) and p15(INK4b)

[86257] genes. *Oncogene* 11: 635–645, 1995.

[86258] 15137.

[86259] Quelle, D. E.; Cheng, M.; Ashmun, R. A.; Sherr, C. J.: Cancer-associated

[86260] mutations at the INK4a locus cancel cell cycle arrest by p16(INK4a)

[86261] but not by the alternative reading frame protein p19(ARF). *Proc.*

[86262] *Nat. Acad. Sci.* 94: 669–673, 1997.

[86263] 15138.

[86264] Quelle, D. E.; Zindy, F.; Ashmun, R. A.; Sherr, C. J.: Alternative

[86265] reading frames of the INK4a tumor suppressor gene encode two unrelated

[86266] proteins capable of inducing cell cycle arrest. *Cell* 83: 993–1000,

[86267] 1995.

[86268] 15139.

- [86269] Ranade, K.; Hussussian, C. J.; Sikorski, R. S.; Varmus, H. E.;
- [86270] Goldstein, A. M.; Tucker, M. A.; Serrano, M.; Hannon, G. J.;
Beach,
- [86271] D.; Dracopoli, N. C.: Mutations associated with familial
melanoma
- [86272] impair p16(INK4) function. *Nature Genet.* 10: 114–116,
1995.
- [86273] 15140.
- [86274] Schmitt, C. A.; Fridman, J. S.; Yang, M.; Lee, S.; Baranov,
E.;
- [86275] Hoffman, R. M.; Lowe, S. W.: A senescence program con-
trolled by p53
- [86276] and p16–INK4a contributes to the outcome of cancer
therapy. *Cell* 109:
- [86277] 335–346, 2002.
- [86278] 15141.
- [86279] Serrano, J.; Goebel, S. U.; Peghini, P. L.; Lubensky, I. A.;
- Gibril,
- [86280] F.; Jensen, R. T.: Alterations in the p16INK4a/CDKN2A tu-
mor suppressor
- [86281] gene in gastrinomas. *J. Clin. Endocr. Metab.* 85:
4146–4156, 2000.

[86282] 15142.

[86283] Serrano, M.; Hannon, G. J.; Beach, D.: A new regulatory motif

[86284] in cell-cycle control causing specific inhibition of cyclin D/CDK4. Nature 366:

[86285] 704–707, 1993.

[86286] 15143.

[86287] Serrano, M.; Lee, H.-W.; Chin, L.; Cordon-Cardo, C.; Beach, D.;

[86288] DePinho, R. A.: Role of the INK4a locus in tumor suppression and

[86289] cell mortality. Cell 85: 27–37, 1996.

[86290] 15144.

[86291] Sharpless, N. E.; Bardeesy, N.; Lee, K.-H.; Carrasco, D.; Castrillon,

[86292] D. H.; Aguirre, A. J.; Wu, E. A.; Horner, J. W.; DePinho, R. A.:

[86293] Loss of p16(Ink4a) with retention of p19(Arf) predisposes mice to

[86294] tumorigenesis. Nature 413: 86–91, 2001.

[86295] 15145.

- [86296] Smith–Sorensen, B.; Hovig, E.: CDKN2A (p16–INK4A) so-
matic and
[86297] germline mutations. Hum. Mutat. 7: 294–303, 1996.
[86298] 15146.
- [86299] Stone, S.; Jiang, P.; Dayananth, P.; Tavtigian, S. W.;
Katcher,
[86300] H.; Parry, D.; Peters, G.; Kamb, A.: Complex structure and
regulation
[86301] of the p16(MTS1) locus. Cancer Res. 55: 2988–2994,
1995.
[86302] 15147.
- [86303] Stott, F. J.; Bates, S.; James, M. C.; McConnell, B. B.; Star-
borg,
[86304] M.; Brookes, S.; Palmero, I.; Ryan, K.; Hara, E.; Vousden, K.
H.;
- [86305] Peters, G.: The alternative product from the human
CDKN2A locus,
[86306] p14(ARF), participates in a regulatory feedback loop with
p53 and
[86307] MDM2. EMBO J. 17: 5001–5014, 1998.
[86308] 15148.

- [86309] Tsutsumi, M.; Tsai, Y. C.; Gonzalgo, M. L.; Nichols, P. W.; Jones,
- [86310] P. A.: Early acquisition of homozygous deletions of p16/p19 during
- [86311] squamous cell carcinogenesis and genetic mosaicism in bladder cancer. *Oncogene* 17:
- [86312] 3021–3027, 1998.
- [86313] 15149.
- [86314] van der Velden, P. A.; Sandkuijl, L. A.; Bergman, W.; Hille, E.
- [86315] T. M.; Frants, R. R.; Gruis, N. A.: A locus linked to p16 modifies
- [86316] melanoma risk in Dutch familial atypical multiple mole melanoma (FAMMM)
- [86317] syndrome families. *Genome Res.* 9: 575–580, 1999.
- [86318] 15150.
- [86319] Vasen, H. F. A.; Gruis, N. A.; Frants, R. R.; van der Velden,
- [86320] P. A.; Hille, E. T. M.; Bergman, W.: Risk of developing pancreatic
- [86321] cancer in families with familial atypical multiple mole melanoma associated

- [86322] with a specific 19 deletion of p16(p16–Leiden). *Int. J. Cancer* 87:
809–811, 2000.
- [86323] 15151.
- [86324] Wainwright, B.: Familial melanoma and p16—a hung jury.
Nature Genet. 8: 3–5, 1994.
- [86325] 15152.
- [86326] Walker, G. J.; Hussussian, C. J.; Flores, J. F.; Glendening, J.
M.; Haluska, F. G.; Dracopoli, N. C.; Hayward, N. K.; Fountain, J.
W.: Mutations of the CDKN2/p16(INK4) gene in Australian
melanoma kindreds. *Hum. Mol. Genet.* 4: 1845–1852, 1995.
- [86327] 15153.
- [86328] Wiest, J. S.; Franklin, W. A.; Otstot, J. T.; Forbey, K.;
Varella–Garcia,
M.; Rao, K.; Drabkin, H.; Gemmill, R.; Ahrent, S.; Sidransky, D.;
Saccomanno, G.; Fountain, J. W.; Anderson, M. W.: Identification

- [86336] of a novel region of homozygous deletion on chromosome 9p in squamous
- [86337] cell carcinoma of the lung: the location of a putative tumor suppressor
- [86338] gene. *Cancer Res.* 57: 1–6, 1997.
- [86339] 15154.
- [86340] Williamson, M. P.; Elder, P. A.; Shaw, M. E.; Devlin, J.; Knowles,
- [86341] M. A.: p16 (CDKN2) is a major deletion target at 9p21 in bladder
- [86342] cancer. *Hum. Molec. Genet.* 4: 1569–1577, 1995.
- [86343] 15155.
- [86344] Woloschak, M.; Yu, A.; Xiao, J.; Post, K. D.: Frequent loss of
- [86345] the P16(INK4a) gene product in human pituitary tumors. *Cancer Res.* 56:
- [86346] 2493–2496, 1996.
- [86347] 15156.
- [86348] Zhang, Y.; Xiong, Y.: Mutations in human ARF exon 2 disrupt its
- [86349] nucleolar localization and impair its ability to block nu-

clear export

[86350] of MDM2 and p53. *Molec. Cell* 3: 579–591, 1999.

[86351] 15157.

[86352] Zhang, Y.; Xiong, Y.; Yarbrough, W. G.: ARF promotes
MDM2 degradation

[86353] and stabilizes p53: ARF–INK4a locus deletion impairs both
the Rb and

[86354] p53 tumor suppression pathways. *Cell* 92: 725–734,
1998.

[86355] 15158.

[86356] Zhu, G.; Duffy, D. L.; Eldridge, A.; Grace, M.; Mayne, C.;
O'Gorman,

[86357] L.; Aitken, J. F.; Neale, M. C.; Hayward, N. K.; Green, A. C.;
Martin,

[86358] N. G.: A major quantitative–trait locus for mole density is
linked

[86359] to the familial melanoma gene CDKN2A: a maximum–like–
lihood combined

[86360] linkage and association analysis in twins and their sibs.
Am. J.

[86361] *Hum. Genet.* 65: 483–492, 1999.

[86362] 15159.

- [86363] Pennica, D.; Swanson, T. A.; Shaw, K. J.; Kuang, W.-J.; Gray, C.
- [86364] L.; Beatty, B. G.; Wood, W. I.: Human cardiotrophin-1: protein and
- [86365] gene structure, biological and binding activities, and chromosomal
- [86366] localization. *Cytokine* 8: 183-189, 1996.
- [86367] 15160.
- [86368] Bonnefont, J.-P.; Taroni, F.; Cavadini, P.; Cepanec, C.; Brivet,
- [86369] M.; Saudubray, J.-M.; Leroux, J.-P.; Demaugre, F.: Molecular analysis
- [86370] of carnitine palmitoyltransferase II deficiency with hepatocardiomyopathy
- [86371] expression. *Am. J. Hum. Genet.* 58: 971-978, 1996.
- [86372] 15161.
- [86373] Candia, A. F.; Hu, J.; Crosby, J.; Lalley, P. A.; Noden, D.; Nadeau,
- [86374] J. H.; Wright, C. V. E.: Mox-1 and Mox-2 define a novel homeobox
- [86375] gene subfamily and are differentially expressed during early mesodermal

- [86376] patterning in mouse embryos. Development 116: 1123–1136, 1992.
- [86377] 15162.
- [86378] Gorski, D. H.; LePage, D. F.; Patel, C. V.; Copeland, N. G.; Jenkins, N. A.; Walsh, K.: Molecular cloning of a diverged homeobox gene that is rapidly down-regulated during the G0/G1 transition in vascular smooth muscle cells. Molec. Cell Biol. 13: 3722–3733, 1993.
- [86382] 15163.
- [86383] Grigoriou, M.; Kastrinaki, M.–C.; Modi, W. S.; Theodorakis, K.;
- [86384] Mankoo, B.; Pachnis, V.; Karagogeos, D.: Isolation of the human MOX2 homeobox gene and localization to chromosome 7p22.1–p21.3. Genomics 26: 550–555, 1995.
- [86387] 15164.
- [86388] LePage, D. F.; Altomare, D. A.; Testa, J. R.; Walsh, K.: Molecular

[86389] cloning and localization of the human GAX gene to 7p21.

Genomics 24:

[86390] 535–540, 1994.

[86391] 15165.

[86392] Mankoo, B. S.; Collins, N. S.; Ashby, P.; Grigorieva, E.;

Pevny,

[86393] L. H.; Candia, A.; Wright, C. V. E.; Rigby, P. W. J.; Pachnis,

V.

[86394] : Mox2 is a component of the genetic hierarchy controlling
limb muscle

[86395] development. Nature 400: 69–73, 1999.

[86396] 15166.

[86397] Hayashi, Y. K.; Chou, F.-L.; Engvall, E.; Ogawa, M.; Mat-

suda, C.;

[86398] Hirabayashi, S.; Yokochi, K.; Ziober, B. L.; Kramer, R. H.;

Kaufman,

[86399] S. J.; Ozawa, E.; Goto, Y.; Nonaka, I.; Tsukahara, T.; Wang,

J.; Hoffman,

[86400] E. P.; Arahata, K.: Mutations in the integrin alpha-7 gene
cause

[86401] congenital myopathy. Nature Genet. 19: 94–97, 1998.

[86402] 15167.

[86403] Mayer, U.; Saher, G.; Fassler, R.; Bornemann, A.; Echter-
meyer,

[86404] F.; von der Mark, H.; Miosge, N.; Poschl, E.; von der Mark,
K.: Absence

[86405] of integrin alpha-7 causes a novel form of muscular dys-
trophy. Nature

[86406] Genet. 17: 318-323, 1997.

[86407] 15168.

[86408] Wang, W.; Wu, W.; Desai, T.; Ward, D. C.; Kaufman, S. J.:
Localization

[86409] of the alpha-7 integrin gene (ITGA7) on human chromo-
some 12q13: clustering

[86410] of integrin and Hox genes implies parallel evolution of
these gene

[86411] families. Genomics 26: 563-570, 1995.

[86412] 15169.

[86413] Lindor, N. M.; Furuichi, Y.; Kitao, S.; Shimamoto, A.; Arndt,
C.;

[86414] Jalal, S.: Rothmund-Thomson syndrome due to RECQ4 he-
licase mutations:

- [86415] report and clinical and molecular comparisons with Bloom syndrome
- [86416] and Werner syndrome. *Am. J. Med. Genet.* 90: 223–228, 2000.
- [86417] 15170.
- [86418] Puranam, K. L.; Blackshear, P. J.: Cloning and characterization
- [86419] of RECQL, a potential human homologue of the *Escherichia coli* DNA
- [86420] helicase RecQ. *J. Biol. Chem.* 269: 29838–29845, 1994.
- [86421] 15171.
- [86422] Puranam, K. L.; Kennington, E.; J.-Sait, S. N.; Shows, T. B.; Rochelle,
- [86423] J. M.; Seldin, M. F.; Blackshear, P. J.: Chromosomal localization
- [86424] of the gene encoding the human DNA helicase RECQL and its mouse homologue. *Genomics* 26:
- [86425] 595–598, 1995.
- [86426] 15172.
- [86427] Hagen, G.; Muller, S.; Beato, M.; Suske, G.: Cloning by recognition

- [86428] site screening of two novel GT box binding proteins: a family of Sp1
- [86429] related genes. *Nucleic Acids Res.* 20: 5519–5525, 1992.
- [86430] 15173.
- [86431] Lamour, V.; Lecluse, Y.; Desmaze, C.; Spector, M.; Bode-scot, M.;
- [86432] Aurias, A.; Osley, M. A.; Lipinski, M.: A human homolog of the S.
- [86433] cerevisiae HIR1 and HIR2 transcriptional repressors cloned from the
- [86434] DiGeorge syndrome critical region. *Hum. Molec. Genet.* 4: 791–799,
- [86435] 1995.
- [86436] 15174.
- [86437] Llevadot, R.; Scambler, P.; Estivill, X.; Pritchard, M.: Ge-nomic
- [86438] organization of TUPLE1/HIRA: a gene implicated in DiGe-orge syndrome. *Mammalian*
- [86439] *Genome* 7: 911–914, 1996.
- [86440] 15175.
- [86441] Lorain, S.; Quivy, J.-P.; Monier-Gavelle, F.; Scamps, C.;

Lecluse,

[86442] Y.; Almouzni, G.; Lipinski, M.: Core histones and HIRIP3, a novel

[86443] histone-binding protein, directly interact with WD repeat protein

[86444] HIRA. *Molec. Cell. Biol.* 18: 5546–5556, 1998.

[86445] 15176.

[86446] Magnaghi, P.; Roberts, C.; Lorain, S.; Lipinski, M.; Scambler,

[86447] P. J.: HIRA, a mammalian homologue of *Saccharomyces cerevisiae* transcriptional

[86448] co-repressors, interacts with Pax3. *Nature Genet.* 20: 74–77, 1998.

[86449] 15177.

[86450] Mattei, M.-G.; Halford, S.; Scambler, P. J.: Mapping of the TUPLE1

[86451] gene to mouse chromosome 16A-B1. *Genomics* 23: 717–718, 1994.

[86452] 15178.

[86453] Roberts, C.; Daw, S. C. M.; Halford, S.; Scambler, P. J.: Cloning

[86454] and developmental expression analysis of chick Hira

(Chira), a candidate

[86455] gene for DiGeorge syndrome. Hum. Molec. Genet. 6:
237-245, 1997.

[86456] 15179.

[86457] Wilming, L. G.; Snoeren, C. A. S.; van Rijswijk, A.;
Grosveld,

[86458] F.; Meijers, C.: The murine homologue of HIRA, a DiGe-
orge syndrome

[86459] candidate gene, is expressed in embryonic structures af-
fected in human

[86460] CATCH22 patients. Hum. Molec. Genet. 6: 247-258, 1997.

[86461] 15180.

[86462] Akao, Y.; Matsuda, Y.: Identification and chromosome
mapping of

[86463] the mouse homologue of the human gene (DDX6) that en-
codes a putative

[86464] RNA helicase of the DEAD box protein family. Cytogenet.
Cell Genet. 75:

[86465] 38-44, 1996.

[86466] 15181.

[86467] Akao, Y.; Seto, M.; Takahashi, T.; Kubonishi, I.; Miyoshi, I.;

[86468] Nakazawa, S.; Tsujimoto, Y.; Croce, C. M.; Ueda, R.: Molecular cloning
[86469] of the chromosomal breakpoint of a B-cell lymphoma with
the t(11;14)(q23;q32)
[86470] chromosome translocation. Cancer Res. 51: 1574–1576,
1991.
[86471] 15182.
[86472] Akao, Y.; Seto, M.; Yamamoto, K.; Iida, S.; Nakazawa, S.;
Inazawa,
[86473] J.; Abe, T.; Takahashi, T.; Ueda, R.: The RCK gene associated with
[86474] t(11;14) translocation is distinct from the MLL/ALL-1 gene
with t(4;11)
[86475] and t(11;19) translocations. Cancer Res. 52: 6083–6087,
1992.
[86476] 15183.
[86477] Akao, Y.; Tsujimoto, Y.; Finan, J.; Nowell, P. C.; Croce, C.
M.
[86478] : Molecular characterization of a t(11;14)(q23;q32) chromosome translocation
[86479] in a B-cell lymphoma. Cancer Res. 50: 4856–4859, 1990.
[86480] 15184.

- [86481] Lu, D.; Yunis, J. J.: Cloning, expression and localization of
- [86482] an RNA helicase gene from a human lymphoid cell line with chromosomal
- [86483] breakpoint 11q23.3. *Nucleic Acids Res.* 20: 1967–1972, 1992.
- [86484] 15185.
- [86485] Seto, M.; Yamamoto, K.; Takahashi, T.; Ueda, R.: Cloning and expression
- [86486] of a murine cDNA homologous to the human RCK/P54, a lymphoma-linked
- [86487] chromosomal translocation junction gene on 11q23. *Gene* 166: 293–296,
- [86488] 1995.
- [86489] 15186.
- [86490] Tunnacliffe, A.; Perry, H.; Radice, P.; Budarf, M. L.; Emanuel,
- [86491] B. S.: A panel of sequence tagged sites for chromosome band 11q23. *Genomics* 17:
- [86492] 744–747, 1993.
- [86493] 15187.
- [86494] Ricker, K.; Moxley, R. T.; Rohkamm, R.: Rippling muscle

disease. Arch.

[86495] Neurol. 46: 405–408, 1989.

[86496] 15188.

[86497] Dumon, K. R.; Ishii, H.; Fong, L. Y. Y.; Zanesi, N.; Fidanza, V.;

[86498] Mancini, R.; Vecchione, A.; Baffa, R.; Trapasso, F.; During, M. J.;

[86499] Huebner, K.; Croce, C. M.: FHIT gene therapy prevents tumor development

[86500] in Fhit-deficient mice. Proc. Nat. Acad. Sci. 98: 3346–3351, 2001.

[86501] 15189.

[86502] Ikeda, A.; Zheng, Q. Y.; Zuberi, A. R.; Johnson, K. R.; Nagert,

[86503] J. K.; Nishina, P. M.: Microtubule-associated protein 1A is a modifier

[86504] of tubby hearing (moth1). Nature Genet. 30: 401–405, 2002.

[86505] 15190.

[86506] Roy, S. K.; Hu, J.; Meng, Q.; Xia, Y.; Shapiro, P. S.; Reddy, S.

[86507] P. M.; Platanius, L. C.; Lindner, D. J.; Johnson, P. F.;

Pritchard,

[86508] C.; Pages, G.; Pouyssegur, J.; Kalvakolanu, D. V.: MEKK1 plays a

[86509] critical role in activating the transcription factor C/EBP-beta-dependent

[86510] gene expression in response to IFN-gamma. Proc. Nat. Acad. Sci. 99:

[86511] 7945-7950, 2002.

[86512] 15191.

[86513] Lindsten, T.; Ross, A. J.; King, A.; Zong, W.-X.; Rathmell, J.

[86514] C.; Shiels, H. A.; Ulrich, E.; Waymire, K. G.; Mahar, P.; Frauwirth,

[86515] K.; Chen, Y.; Wei, M.; and 9 others: The combined functions of

[86516] proapoptotic Bcl-2 family members Bak and Bax are essential for normal

[86517] development of multiple tissues. Molec. Cell 6: 1389-1399, 2000.

[86518] 15192.

[86519] Matikainen, T.; Perez, G. I.; Jurisicova, A.; Pru, J. K.; Schlezinger,

[86520] J. J.; Ryu, H.-Y.; Laine, J.; Sakai, T.; Korsmeyer, S. J.;

Casper,

[86521] R. F.; Sherr, D. H.; Tilly, J. L.: Aromatic hydrocarbon receptor-driven

[86522] Bax gene expression is required for premature ovarian failure caused

[86523] by biohazardous environmental chemicals. *Nature Genet.* 28: 355–360,

[86524] 2001.

[86525] 15193.

[86526] Wei, M. C.; Zong, W.-X.; Cheng, E. H.-Y.; Lindsten, T.; Panoutsakopoulou,

[86527] V.; Ross, A. J.; Roth, K. A.; MacGregor, G. R.; Thompson, C. B.; Korsmeyer,

[86528] S. J.: Proapoptotic BAX or BAK: a requisite gateway to mitochondrial

[86529] dysfunction and death. *Science* 292: 727–730, 2001.

[86530] 15194.

[86531] Dasari, V. R.; Sandhu, A. K.; Mills, D. C. B.; Athwal, R. S.; Kunapuli,

[86532] S. P.: Mapping of the P2U purinergic receptor gene to human chromosome

[86533] 11q13.5–14.1. *Somat. Cell Molec. Genet.* 22: 75–79,

1996.

[86534] 15195.

[86535] Katzur, A. C.; Koshimizu, T.-A.; Tomic, M.; Schultze-Mosgau, A.;

[86536] Ortmann, O.; Stojilkovic, S. S.: Expression and responsiveness of

[86537] P2Y2 receptors in human endometrial cancer cell lines. J. Clin. Endocr.

[86538] Metab. 84: 4085-4091, 1999.

[86539] 15196.

[86540] Parr, C. E.; Sullivan, D. M.; Paradiso, A. M.; Lazarowski, E. R.;

[86541] Burch, L. H.; Olsen, J. C.; Erb, L.; Weisman, G. A.; Boucher, R. C.;

[86542] Turner, J. T.: Cloning and expression of a human P(2U) nucleotide

[86543] receptor, a target for cystic fibrosis pharmacotherapy. Proc. Nat.

[86544] Acad. Sci. 91: 3275-3279, 1994.

[86545] 15197.

[86546] Somers, G. R.; Hammet, F.; Woollatt, E.; Richards, R. I.;

Southey,

[86547] M. C.; Venter, D. J.: Chromosomal localization of the human P2Y(6)

[86548] purinoceptor gene and phylogenetic analysis of the P2y purinoceptor

[86549] family. Genomics 44: 127–130, 1997.

[86550] 15198.

[86551] Tai, C.-J.; Kang, S. K.; Cheng, K. W.; Choi, K.-C.; Nathwani, P.

[86552] S.; Leung, P. C. K.: Expression and regulation of P2U–purinergic

[86553] receptor in human granulosa–luteal cells. J. Clin. Endocr. Metab. 85:

[86554] 1591–1597, 2000.

[86555] 15199.

[86556] Santamarina–Fojo, S.; Peterson, K.; Knapper, C.; Qiu, Y.; Freeman,

[86557] L.; Cheng, J.-F.; Osorio, J.; Remaley, A.; Yang, X.-P.; Hauschild,

[86558] C.; Prades, C.; Chimini, G.; Blackmon, E.; Francois, T.; Duverger,

[86559] N.; Rubin, E. M.; Rosier, M.; Deneffe, P.; Fredrickson, D. S.;

Brewer,

- [86560] H. B., Jr.: Complete genomic sequence of the human ABCA1 gene: analysis
[86561] of the human and mouse ATP-binding cassette A promoter. Proc. Nat.
[86562] Acad. Sci. 97: 7987–7992, 2000. Note: Erratum: Proc. Nat.
Acad. Sci.
[86563] 99: 1098 only, 2002.
[86564] 15200.
- [86565] Szakacs, G.; Langmann, T.; Ozvegy, C.; Orso, E.; Schmitz, G.;
[86566] Varadi, A.; Sarkadi, B.: Characterization of the ATPase cycle of
[86567] human ABCA1: implications for its function as a regulator rather than
[86568] an active transporter. Biochem. Biophys. Res. Commun. 288: 1258–1264,
[86569] 2001.
[86570] 15201.
- [86571] Utech, M.; Hobbel, G.; Rust, S.; Reinecke, H.; Assmann, G.; Walter,
[86572] M.: Accumulation of RhoA, RhoB, RhoG, and Rac1 in fi-

broblasts from

[86573] Tangier disease subjects suggests a regulatory role of Rho family

[86574] proteins in cholesterol efflux. Biochem. Biophys. Res. Commun. 280:

[86575] 229–236, 2001.

[86576] 15202.

[86577] Zhao, L.-X.; Zhou, C.-J.; Tanaka, A.; Nakata, M.; Hirabayashi,

[86578] T.; Amachi, T.; Shioda, S.; Ueda, K.; Inagaki, N.: Cloning, characterization

[86579] and tissue distribution of the rat ATP-binding cassette (ABC) transporter

[86580] ABC2/ABCA2. Biochem J. 350: 865–872, 2000.

[86581] 15203.

[86582] Zwarts, K. Y.; Clee, S. M.; Zwinderman, A. H.; Engert, J. C.;

[86583] Singaraja, R.; Loubser, O.; James, E.; Roomp, K.; Hudson, T. J.; Jukema,

[86584] J. W.; Kastelein, J. J. P.; Hayden, M. R.: ABCA1 regulatory variants

[86585] influence coronary artery disease independent of effects

on plasma

[86586] lipid levels. Clin. Genet. 61: 115–125, 2002.

[86587] 15204.

[86588] Ing, Y. L.; Leung, I. W. L.; Heng, H. H. Q.; Tsui, L.-C.; Las-sam,

[86589] N. J.: MLK-3: identification of a widely-expressed protein kinase

[86590] bearing an SH3 domain and a leucine zipper-basic region domain. Oncogene 9:

[86591] 1745–1750, 1994.

[86592] 15205.

[86593] Bernard, O. A.; Mauchauffe, M.; Mecucci, C.; Van Den Berghe, H.;

[86594] Berger, R.: A novel gene, AF-1p, fused to HRX in t(1;11)(p32;q23),

[86595] is not related to AF-4, AF-9 nor ENL. Oncogene 9: 1039–1045, 1994.

[86596] 15206.

[86597] Fazioli, F.; Minichiello, L.; Matoska, V.; Castagnino, P.; Miki,

[86598] T.; Wong, W. T.; Di Fiore, P. P.: Eps8, a substrate for the

epidermal

[86599] growth factor receptor kinase, enhances EGF-dependent mitogenic signals.

[86600] EMBO J. 12: 3799–3808, 1993.

[86601] 15207.

[86602] Fazioli, F.; Minichiello, L.; Matoskova, B.; Wong, W. T.; Di Fiore,

[86603] P. P.: eps 15, a novel tyrosine kinase substrate, exhibits transforming

[86604] activity. Molec. Cell. Biol. 13: 5814–5828, 1993.

[86605] 15208.

[86606] Fazioli, F.; Wong, W. T.; Ullrich, S. J.; Sakaguchi, K.; Appella,

[86607] E.; Di Fiore, P. P.: The ezrin-like family of tyrosine kinase substrates:

[86608] receptor-specific pattern of tyrosine phosphorylation and relationship

[86609] to malignant transformation. Oncogene 8: 1335–1345, 1993.

[86610] 15209.

[86611] Wong, W. T.; Kraus, M. H.; Carlomagno, F.; Zelano, A.;

Druck, T.;

[86612] Croce, C. M.; Huebner, K.; Di Fiore, P. P.: The human
eps15 gene,

[86613] encoding a tyrosine kinase substrate, is conserved in evolution and

[86614] maps to 1p31-p32. *Oncogene* 9: 1591-1597, 1994.

[86615] 15210.

[86616] Frydman, J.; Nimmesgern, E.; Erdjument-Bromage, H.;
Wall, J. S.;

[86617] Tempst, P.; Hartl, F.-U.: Function in protein folding of
TRiC, a

[86618] cytosolic ring complex containing TCP-1 and structurally
related subunits. *EMBO*

[86619] *J.* 11: 4767-4778, 1992.

[86620] 15211.

[86621] Joly, E. C.; Seigny, G.; Todorov, I. T.; Bibor-Hardy, V.:
cDNA

[86622] encoding a novel TCP1-related protein. *Biochim. Biophys.*
Acta 1217:

[86623] 224-226, 1994.

[86624] 15212.

- [86625] Sevigny, G.; Joly, E. C.; Bibor-Hardy, V.; Lemieux, N.: Assignment
- [86626] of the human homologue of the mTRiC-P5 gene (TRiC5) to band 1q23 by
- [86627] fluorescence in situ hybridization. Genomics 22: 634-636, 1994.
- [86628] 15213.
- [86629] Sevigny, G.; Lemieux, N.; Steyaert, A.; Bibor-Hardy, V.: Structure
- [86630] of the gene coding for the mouse TRiC-P5 subunit of the cytosolic

- [86631] chaperonin TRiC. Genomics 31: 107–110, 1996.
- [86632] 15214.
- [86633] Walkley, N. A.; Demaine, A. G.; Malik, A. N.: Cloning,
structure
- [86634] and mRNA expression of human Cctg, which encodes the
chaperonin subunit
- [86635] CCT-gamma. Biochem. J. 313: 381–389, 1996.
- [86636] 15215.
- [86637] Hoenicka, J.; Vidal, L.; Morales, B.; Ampuero, I.; Jimenez-
Jimenez,
- [86638] F. J.; Berciano, J.; del Ser, T.; Jimenez, A.; Ruiz, P. G.; de
Yebeles,
- [86639] J. G.: Molecular findings in familial Parkinson disease in
Spain. Arch.
- [86640] Neurol. 59: 966–970, 2002.
- [86641] 15216.
- [86642] Kitada, T.; Asakawa, S.; Hattori, N.; Matsumine, H.; Yama-
mura,
- [86643] Y.; Minoshima, S.; Yokochi, M.; Mizuno, Y.; Shimizu, N.:
Mutations
- [86644] in the parkin gene cause autosomal recessive juvenile

parkinsonism. Nature 392:

[86645] 605–608, 1998.

[86646] 15217.

[86647] Matsumine, H.; Saito, M.; Shimoda–Matsubayashi, S.;
Tanaka, H.;

[86648] Ishikawa, A.; Nakagawa–Hattori, Y.; Yokochi, M.;
Kobayashi, T.; Igarashi,

[86649] S.; Takano, H.; Sanpei, K.; Koike, R.; Mori, H.; Kondo, T.;
Mizutani,

[86650] Y.; Schaffer, A. A.; Yamamura, Y.; Nakamura, S.; Kuzuhura,
S.; Tsuji,

[86651] S.; Mizuno, Y.: Localization of a gene for an autosomal re-
cessive

[86652] form of juvenile parkinsonism to chromosome 6q25.2–27.
Am. J. Hum.

[86653] Genet. 60: 588–596, 1997.

[86654] 15218.

[86655] Tassin, J.; Durr, A.; de Broucker, T.; Abbas, N.; Bonifati, V.;

[86656] De Michele, G.; Bonnet, A.–M.; Broussolle, E.; Pollak, P.;
Vidailhet,

[86657] M.; De Mari, M.; Marconi, R.; Medjbeur, S.; Filla, A.; Meco,

G.; Agid,

[86658] Y.; Brice, A.; French Parkinson's Disease Genetics Study Group;

[86659] European Consortium on Genetic Susceptibility in Parkinson's Disease

[86660] : Chromosome 6-linked autosomal recessive early-onset parkinsonism:

[86661] linkage in European and Algerian families, extension of the clinical

[86662] spectrum, and evidence of a small homozygous deletion in one family. Am.

[86663] J. Hum. Genet. 63: 88-94, 1998.

[86664] 15219.

[86665] Kozu, T.; Henrich, B.; Schafer, K. P.: Structure and expression

[86666] of the gene (HNRPA2B1) encoding the human hnRNP protein A2/B1. Genomics 25:

[86667] 365-371, 1995.

[86668] 15220.

[86669] Kordeli, E.; Lambert, S.; Bennett, V.: Ankyrin-G: a new ankyrin

[86670] gene with neural-specific isoforms localized at the axonal

initial

[86671] segment and node of Ranvier. J. Biol. Chem. 270:
2352–2359, 1995.

[86672] 15221.

[86673] Hsu, L. C.; Chang, W.-C.; Yoshida, A.: Human aldehyde
dehydrogenase

[86674] genes, ALDH7 and ALDH8: genomic organization and
gene structure comparison. Gene 189:

[86675] 89–94, 1997.

[86676] 15222.

[86677] Hsu, L. C.; Chang, W.-C.; Yoshida, A.: Cloning of a cDNA
encoding

[86678] human ALDH7, a new member of the aldehyde dehydro-
genase family. Gene 151:

[86679] 285–289, 1994.

[86680] 15223.

[86681] Aruga, J.; Yokota, N.; Hashimoto, M.; Furuichi, T.; Fukuda,
M.;

[86682] Mikoshiba, K.: A novel zinc finger protein, Zic, is involved
in neurogenesis,

[86683] especially in the cell lineage of cerebellar granule cells. J.

Neurochem. 63:

[86684] 1880–1890, 1994.

[86685] 15224.

[86686] Salero, E.; Perez–Sen, R.; Aruga, J.; Gimenez, C.; Zafra, F.:

[86687] Transcription factors Zic1 and Zic2 bind and transactivate the apolipoprotein

[86688] E gene promoter. J. Biol. Chem. 276: 1881–1888, 2001.

[86689] 15225.

[86690] Yokota, N.; Aruga, J.; Takai, S.; Yamada, K.; Hamazaki, M.; Iwase,

[86691] T.; Sugimura, H.; Mikoshiba, K.: Predominant expression of human

[86692] Zic in cerebellar granule cell lineage and medulloblastoma. Cancer

[86693] Res. 56: 377–383, 1996.

[86694] 15226.

[86695] Bevins, C. L.; Jones, D. E.; Dutra, A.; Schaffzin, J.; Muenke,

[86696] M.: Human enteric defensin genes: chromosomal map position and a

[86697] model for possible evolutionary relationships. Genomics

31: 95–106,

[86698] 1996.

[86699] 15227.

[86700] Jones, D. E.; Bevins, C. L.: Defensin-6 mRNA in human
Paneth cells:

[86701] implications for antimicrobial peptides in host defense of
the human

[86702] bowel. FEBS Lett. 315: 187–192, 1993.

[86703] 15228.

[86704] Mallow, E. B.; Harris, A.; Salzman, N.; Russell, J. P.; DeBer-
ardinis,

[86705] R. J.; Ruchelli, E.; Bevins, C. L.: Human enteric defensins:
gene

[86706] structure and developmental expression. J. Biol. Chem.
271: 4038–4045,

[86707] 1996.

[86708] 15229.

[86709] Jones, D. E.; Bevins, C. L.: Paneth cells of the human small
intestine

[86710] express an antimicrobial peptide gene. J. Biol. Chem. 267:
23216–23225,

- [86711] 1992.
- [86712] 15230.
- [86713] Porter, E. M.; Liu, L.; Oren, A.; Anton, P. A.; Ganz, T.: Localization
- [86714] of human intestinal defensin 5 in Paneth cell granules. Infect. Immun. 65:
- [86715] 2389–2395, 1997.
- [86716] 15231.
- [86717] Quayle, A. J.; Porter, E. M.; Nussbaum, A. A.; Wang, Y. M.; Brabec,
- [86718] C.; Yip, K.-P.; Mok, S. C.: Gene expression, immunolocalization,
- [86719] and secretion of human defensin-5 in human female reproductive tract. Am.
- [86720] J. Path. 152: 1247–1258, 1998.
- [86721] 15232.
- [86722] Matsuzono, Y.; Kinoshita, N.; Tamura, S.; Shimozawa, N.; Hamasaki,
- [86723] M.; Ghaedi, K.; Wanders, R. J. A.; Suzuki, Y.; Kondo, N.; Fujiki,
- [86724] Y.: Human PEX19: cDNA cloning by functional comple-

mentation, mutation

[86725] analysis in a patient with Zellweger syndrome, and potential role

[86726] in peroxisomal membrane assembly. Proc. Nat. Acad. Sci. 96: 2116–2121,

[86727] 1999.

[86728] 15233.

[86729] Kinoshita, N.; Ghaedi, K.; Shimozawa, N.; Wanders, R. J. A.; Matsuzono,

[86730] Y.; Imanaka, T.; Okumoto, K.; Suzuki, Y.; Kondo, N.; Fujiki, Y.:

[86731] Newly identified Chinese hamster ovary cell mutants are defective

[86732] in biogenesis of peroxisomal membrane vesicles (peroxisomal ghosts),

[86733] representing a novel complementation group in mammals J. Biol. Chem. 273:

[86734] 24122–24130, 1998.

[86735] 15234.

[86736] Chen, K.-S.; Gunaratne, P. H.; Hoheisel, J. D.; Young, I. G.; Gabor

[86737] Miklos, G. L.; Greenberg, F.; Shaffer, L. G.; Campbell, H.

D.; Lupski,

[86738] J. R.: The human homologue of the *Drosophila melanogaster* flightless-I

[86739] gene (flil) maps within the Smith-Magenis microdeletion critical region

[86740] in 17p11.2. *Am. J. Hum. Genet.* 56: 175–182, 1995.

[86741] 15235.

[86742] Heisterkamp, N.; Kaartinen, V.; van Soest, S.; Bokoch, G. M.; Groffen,

[86743] J.: Human ABR encodes a protein with GAP-rac activity and homology

[86744] to the DBL nucleotide exchange factor domain. *J. Biol. Chem.* 268:

[86745] 16903–16906, 1993.

[86746] 15236.

[86747] Heisterkamp, N.; Morris, C.; Groffen, J.: ABR, an active BCR-related

[86748] gene. *Nucleic Acids Res.* 17: 8821–8831, 1989.

[86749] 15237.

[86750] McDonald, J. D.; Daneshvar, L.; Willert, J. R.; Matsumura, K.;

- [86751] Waldman, F.; Cogen, P. H.: Physical mapping of chromosome 17p13.3
- [86752] in the region of a putative tumor suppressor gene important in medulloblastoma. Genomics 23:
- [86753] 229–232, 1994.
- [86754] 15238.
- [86755] Shimomura, H.; Sanke, T.; Hanabusa, T.; Tsunoda, K.; Furuta, H.;
- [86756] Nanjo, K.: Nonsense mutation of islet-1 gene (Q310X) found in a type
- [86757] 2 diabetic patient with a strong family history. Diabetes 49: 1597–1600,
- [86758] 2000.
- [86759] 15239.
- [86760] Tanizawa, Y.; Riggs, A. C.; Dagogo-Jack, S.; Vaxillaire, M.; Froguel,
- [86761] P.; Liu, L.; Donis-Keller, H.; Permutt, M. A.: Isolation of the human
- [86762] LIM/homeodomain gene islet-1 and identification of a simple sequence
- [86763] repeat 1. Diabetes 43: 935–941, 1994.

- [86764] 15240.
- [86765] Corti, O.; Finocchiaro, G.; Rossi, E.; Zuffardi, O.; DiDonato,
[86766] S.: Molecular cloning of cDNAs encoding human carnitine
acetyltransferase
[86767] and mapping of the corresponding gene to chromosome
9q34.1. Genomics 23:
[86768] 94–99, 1994.
- [86769] 15241.
- [86770] Kalaria, R. N.; Harik, S. I.: Carnitine acetyltransferase ac-
tivity
[86771] in the human brain and its microvessels is decreased in
Alzheimer's
[86772] disease. Ann. Neurol. 32: 583–586, 1992.
- [86773] 15242.
- [86774] van der Leij, F. R.; Huijkman, N. C. A.; Boomsma, C.;
Kuipers,
[86775] J. R. G.; Bartelds, B.: Genomics of the human carnitine
acyltransferase
[86776] genes. Molec. Genet. Metab. 71: 139–153, 2000.
- [86777] 15243.

- [86778] Barkardottir, R. B.; Sarantaus, L.; Arason, A.; Vehmanen, P.; Bendahl,
- [86779] P.-O.; Kainu, T.; Syrjakoski, K.; Krahe, R.; Huusko, P.; Pyrhonen,
- [86780] S.; Holli, K.; Kallioniemi, O.-P.; Egilsson, V.; Kere, J.; Nevanlinna,
- [86781] H.: Haplotype analysis in Icelandic and Finnish BRCA2 999del5 breast
- [86782] cancer families. *Europ. J. Hum. Genet.* 9: 773–779, 2001.
- [86783] 15244.
- [86784] Breast Cancer Linkage Consortium: Cancer risks in BRCA2 mutation
- [86785] carriers. *J. Nat. Cancer Inst.* 91: 1310–1316, 1999.
- [86786] 15245.
- [86787] Connor, F.; Smith, A.; Wooster, R.; Stratton, M.; Dixon, A.; Campbell,
- [86788] E.; Tait, T.-M.; Freeman, T.; Ashworth, A.: Cloning, chromosomal
- [86789] mapping and expression pattern of the mouse Brca2 gene. *Hum. Molec.*
- [86790] *Genet.* 6: 291–300, 1997.

- [86791] 15246.
- [86792] Couch, F. J.; Rommens, J. M.; Neuhausen, S. L.; Belanger, C.; Dumont, M.; Abel, K.; Bell, R.; Berry, S.; Bogden, R.; Cannon-Albright, L.; Farid, L.; Frye, C.; and 30 others: Generation of an integrated transcription map of the BRCA2 region on chromosome 13q12-q13. *Genomics* 36: 86-99, 1996.
- [86797] 15247.
- [86798] Davies, A. A.; Masson, J.-Y.; McIlwraith, M. J.; Stasiak, A. Z.; Stasiak, A.; Venkitaraman, A. R.; West, S. C.: Role of BRCA2 in control of the RAD51 recombination and DNA repair protein. *Molec. Cell* 7: 273-282, 2001.
- [86802] 15248.
- [86803] Easton, D. F.; Ford, D.; Bishop, D. T.; Breast Cancer Linkage

- [86804] Consortium: Breast and ovarian cancer incidence in
BRCA1-mutation
- [86805] carriers. Am. J. Hum. Genet. 56: 265-271, 1995.
- [86806] 15249.
- [86807] Easton, D. F.; Steele, L.; Fields, P.; Ormiston, W.; Averill,
[86808] D.; Daly, P. A.; McManus, R.; Neuhausen, S. L.; Ford, D.;
Wooster,
[86809] R.; Cannon-Albright, L. A.; Stratton, M. R.; Goldgar, D. E.:
Cancer
[86810] risks in two large breast cancer families linked to BRCA2
on chromosome
[86811] 13q12-13. Am. J. Hum. Genet. 61: 120-128, 1997.
- [86812] 15250.
- [86813] Fackenthal, J. D.; Cartegni, L.; Krainer, A. R.; Olopade, O. I.
[86814] : BRCA2 T2722R is a deleterious allele that causes exon
skipping. Am.
[86815] J. Hum. Genet. 71: 625-631, 2002.
- [86816] 15251.
- [86817] Garcia-Marco, J. A.; Caldas, C.; Price, C. M.; Wiedemann, L.
M.;

- [86818] Ashworth, A.; Catovsky, D.: Frequent somatic deletion of the 13q12.3
- [86819] locus encompassing BRCA2 in chronic lymphocytic leukemia. Blood 88:
- [86820] 1568–1575, 1996.
- [86821] 15252.
- [86822] Gayther, S. A.; Mangion, J.; Russell, P.; Seal, S.; Barfoot, R.;
- [86823] Ponder, B. A. J.; Stratton, M. R.; Easton, D.: Variation in risks
- [86824] of breast and ovarian cancer associated with different germline mutations
- [86825] of the BRCA2 gene. Nature Genet. 15: 103–105, 1997.
- [86826] 15253.
- [86827] Gronberg, H.; Ahman, A.-K.; Emanuelsson, M.; Bergh, A.; Damber,
- [86828] J.-E.; Borg, A.: BRCA2 mutation in a family with hereditary prostate
- [86829] cancer. Genes Chromosomes Cancer 30: 299–301, 2001.
- [86830] 15254.
- [86831] Gudmundsson, J.; Johannesdottir, G.; Arason, A.; Bergthorsson,

- [86832] J. T.; Ingvarsson, S.; Egilsson, V.; Barkardottir, R. B.: Fre-
quent
- [86833] occurrence of BRCA2 linkage in Icelandic breast cancer
families and
- [86834] segregation of a common BRCA2 haplotype. Am. J. Hum.
Genet. 58:
- [86835] 749–756, 1996.
- [86836] 15255.
- [86837] Healey, C. S.; Dunning, A. M.; Teare, M. D.; Chase, D.;
Parker,
- [86838] L.; Burn, J.; Chang–Claude, J.; Mannermaa, A.; Kataja, V.;
Huntsman,
- [86839] D. G.; Pharoah, P. D. P.; Luben, R. N.; Easton, D. F.; Pon-
der, B.
- [86840] A. J.: A common variant in BRCA2 is associated with both
breast cancer
- [86841] risk and prenatal viability. Nature Genet. 26: 362–364,
2000.
- [86842] 15256.
- [86843] Gengyo–Ando, K.; Kitayama, H.; Mukaida, M.; Ikawa, Y.: A
murine
- [86844] neural–specific homolog corrects cholinergic defects in

Caenorhabditis

- [86845] *elegans* unc-18 mutants. J. Neurosci. 16: 6695–6702, 1996.
- [86846] 15257.
- [86847] Pevsner, J.; Hsu, S.-C.; Scheller, R. H.: n-Sec1: a neural-specific
- [86848] syntaxin-binding protein. Proc. Nat. Acad. Sci. 91: 1445–1449, 1994.
- [86849] 15258.
- [86850] Swanson, D. A.; Steel, J. M.; Valle, D.: Identification and characterization
- [86851] of the human ortholog of rat STXBP1, a protein implicated in vesicle
- [86852] trafficking and neurotransmitter release. Genomics 48: 373–376,
- [86853] 1998.
- [86854] 15259.
- [86855] Verhage, M.; Mala, A. S.; Plomp, J. J.; Brussaard, A. B.; Heeroma,
- [86856] J. H.; Vermeer, H.; Toonen, R. F.; Hammer, R. E.; van den Berg, T.
- [86857] K.; Missler, M.; Geuze, H. J.; Sudhof, T. C.: Synaptic as-

sembly of

[86858] the brain in the absence of neurotransmitter secretion.

Science 287:

[86859] 864–869, 2000.

[86860] 15260.

[86861] Monaco, C.; Negrini, M.; Sozzi, G.; Veronese, M. L.; Vorechovsky,

[86862] I.; Godwin, A. K.; Croce, C. M.: Molecular cloning and characterization

[86863] of LOH11CR2A, a new gene within a refined minimal region of LOH at

[86864] 11q23. Genomics 46: 217–222, 1997.

[86865] 15261.

[86866] Topper, J. N.; Cai, J.; Qiu, Y.; Anderson, K. R.; Xu, Y.–Y.; Deeds,

[86867] J. D.; Feeley, R.; Gimeno, C. J.; Woolf, E. A.; Tayber, O.; Mays,

[86868] G. G.; Sampson, B. A.; Schoen, F. J.; Gimbrone, M. A., Jr.; Falb,

[86869] D.: Vascular MADs: two novel MAD-related genes selectively inducible

[86870] by flow in human vascular endothelium. Proc. Nat. Acad.

Sci. 94:

[86871] 9314–9319, 1997.

[86872] 15262.

[86873] Festing, M. F. W. (ed.): Inbred strains in biomedical research. New

[86874] York: Oxford University Press , 1979. p. 255.

[86875] 15263.

[86876] Illa, I.; Serrano–Munuera, C.; Gallardo, E.; Lasa, A.; Rojas–Garcia,

[86877] R.; Palmer, J.; Gallano, P.; Baiget, M.; Matsuda, C.; Brown, R. H.

[86878] : Distal anterior compartment myopathy: a dysferlin mutation causing

[86879] a new muscular dystrophy phenotype. Ann. Neurol. 49: 130–134, 2001.

[86880] 15264.

[86881] Illarioshkin, S. N.; Ivanova–Smolenskaya, I. A.; Greenberg, C.

[86882] R.; Nylen, E.; Sukhorukov, V. S.; Poleshchuk, V. V.; Markova, E. D.;

[86883] Wrogemann, K.: Identical dysferlin mutation in limb–girdle

muscular

- [86884] dystrophy type 2B and distal myopathy. *Neurology* 55:
1931–1933,
[86885] 2000.
[86886] 15265.
- [86887] Liu, J.; Wu, C.; Bossie, K.; Bejaoui, K.; Hosler, B. A.; Gin-
grich,
[86888] J. C.; Ben Hamida, M.; Hentati, F.; Schurr, E.; de Jong, P. J.;
Brown,
[86889] R. H., Jr.: Generation of 3–Mb PAC contig spanning the
Miyoshi myopathy/limb
[86890] girdle muscular dystrophy (MM/LGMD2B) locus on chro-
mosome 2p13. *Genomics* 49:
[86891] 23–29, 1998.
[86892] 15266.
- [86893] Mahjneh, I.; Vannelli, G.; Bushby, K.; Marconi, G. P.: A
large
[86894] inbred Palestinian family with two forms of muscular dys-
trophy. *Neuromusc.*
[86895] *Disord.* 2: 277–283, 1992.
[86896] 15267.

- [86897] Weiler, T.; Bashir, R.; Anderson, L. V. B.; Davison, K.; Moss, J. A.; Britton, S.; Nylén, E.; Keers, S.; Vafiadaki, E.; Greenberg, C. R.; Bushby, K. M. D.; Wrogemann, K.: Identical mutation in patients with limb girdle muscular dystrophy type 2B or Miyoshi myopathy suggests a role for modifier gene(s). Hum. Molec. Genet. 8: 871–877, 1999.
- [86902] 15268.
- [86903] Hannon, G. J.; Beach, D.: p15(INK4B) is a potential effector of TGF- β -induced cell cycle arrest. Nature 371: 257–261, 1994.
- [86905] 15269.
- [86906] Stone, S.; Dayananth, P.; Jiang, P.; Weaver-Feldhaus, J. M.; Tavtigian, S. V.; Cannon-Albright, L.; Kamb, A.: Genomic structure, expression and mutational analysis of the P15 (MTS2) gene. Oncogene 11: 987–991, 1995.
- [86909] 1995.

- [86910] 15270.
- [86911] Pennica, D.; Arce, V.; Swanson, T. A.; Vejsada, R.; Pollock, R.
- [86912] A.; Armanini, M.; Dudley, K.; Phillips, H. S.; Rosenthal, A.; Kato, A. C.; Henderson, C. E.: Cardiotrophin-1, a cytokine present in embryonic muscle, supports long-term survival of spinal motoneurons. *Neuron* 17: 63-74, 1996.
- [86913] 15271.
- [86914] Pennica, D.; King, K. L.; Shaw, K. J.; Luis, E.; Rullamas, J.; Luoh, S.-M.; Darbonne, W. C.; Knutzon, D. S.; Yen, R.; Chien, K. R.; Baker, J. B.; Wood, W. I.: Expression cloning of cardiotrophin 1, a cytokine that induces cardiac myocyte hypertrophy. *Proc. Nat. Acad. Sci.* 92: 1142-1146, 1995.
- [86915] 15272.
- [86916] Ghosh, A.: Learning more about NMDA receptor regula-

tion. Science 295:

[86924] 449–451, 2002.

[86925] 15273.

[86926] Grunwald, I. C.; Korte, M.; Wolfer, D.; Wilkinson, G. A.; Unsicker,

[86927] K.; Lipp, H.-P.; Bonhoeffer, T.; Klein, R.: Kinase-independent requirement

[86928] of EphB2 receptors in hippocampal synaptic plasticity.

Neuron 32:

[86929] 1027–1040, 2001.

[86930] 15274.

[86931] Henderson, J. T.; Georgiou, J.; Jia, Z.; Robertson, J.; Elowe,

[86932] S.; Roder, J. C.; Pawson, T.: The receptor tyrosine kinase EphB2

[86933] regulates NMDA-dependent synaptic function. Neuron 32: 1041–1056,

[86934] 2001.

[86935] 15275.

[86936] Himanen, J.-P.; Rajashankar, K. R.; Lackmann, M.; Cowan, C. A.;

[86937] Henkemeyer, M.; Nikolov, D. B.: Crystal structure of an

Eph receptor-ephrin

[86938] complex. *Nature* 414: 933–938, 2001.

[86939] 15276.

[86940] Sakano, S.; Serizawa, R.; Inada, T.; Iwama, A.; Itoh, A.; Kato,

[86941] C.; Shimizu, Y.; Shinkai, F.; Shimizu, R.; Kondo, S.; Ohno, M.; Suda,

[86942] T.: Characterization of a ligand for receptor protein-tyrosine kinase

[86943] HTK expressed in immature hematopoietic cells. *Oncogene* 13: 813–822,

[86944] 1996.

[86945] 15277.

[86946] Takasu, M. A.; Dalva, M. B.; Zigmond, R. E.; Greenberg, M. E.

[86947] : Modulation of NMDA receptor-dependent calcium influx and gene expression

[86948] through EphB receptors. *Science* 295: 491–495, 2002.

[86949] 15278.

[86950] Wang, H. U.; Chen, Z.-F.; Anderson, D. J.: Molecular distinction

- [86951] and angiogenic interaction between embryonic arteries and veins revealed
- [86952] by ephrin-B2 and its receptor Eph-B4. *Cell* 93: 741-753, 1998.
- [86953] 15279.
- [86954] Slama, A.; Brivet, M.; Boutron, A.; Legrand, A.; Saudubray, J.-M.;
- [86955] Demaugre, F.: Complementation analysis of carnitine palmitoyltransferase
- [86956] I and II defects. *Pediat. Res.* 40: 542-546, 1996.
- [86957] 15280.
- [86958] Doudney, K.; Murdoch, J. N.; Paternotte, C.; Bentley, L.; Gregory,
- [86959] S.; Copp, A. J.; Stanier, P.: Comparative physical and transcript
- [86960] maps of approximately 1 Mb around loop-tail, a gene for severe neural
- [86961] tube defects on distal mouse chromosome 1 and human chromosome 1q22-q23. *Genomics* 72:
- [86962] 180-192, 2001.
- [86963] 15281.

- [86964] Kibar, Z.; Vogan, K. J.; Groulx, N.; Justice, M. J.; Underhill,
[86965] D. A.; Gros, P.: Ltap, a mammalian homolog of *Drosophila*
Strabismus/Van
[86966] Gogh, is altered in the mouse neural tube mutant loop-
tail. *Nature*
[86967] *Genet.* 28: 251–255, 2001.
[86968] 15282.
[86969] Mullick, A.; Trasler, D.; Gros, P.: High-resolution linkage
map
[86970] in the vicinity of the Lp locus. *Genomics* 26: 479–488,
1995.
[86971] 15283.
[86972] Murdoch, J. N.; Doudney, K.; Paternotte, C.; Copp, A. J.;
Stanier,
[86973] P.: Severe neural tube defects in the loop-tail mouse re-
sult from
[86974] mutation of *Lpp1*, a novel gene involved in floor plate
specification. *Hum.*
[86975] *Molec. Genet.* 10: 2593–2601, 2001.
[86976] 15284.
[86977] Stanier, P.; Henson, J. N.; Eddleston, J.; Moore, G. E.; Copp,

- [86978] A. J.: Genetic basis of neural tube defects: the mouse gene loop-tail
- [86979] maps to a region of chromosome 1 syntenic with human 1q21-q23. Genomics 26:
- [86980] 473-478, 1995.
- [86981] 15285.
- [86982] Yu, J. X.; Chao, L.; Chao, J.: Protasin is a novel human serine
- [86983] proteinase from seminal fluid: purification, tissue distribution,
- [86984] and localization in prostate gland. J. Biol. Chem. 269: 18843-18848,
- [86985] 1994.
- [86986] 15286.
- [86987] Yu, J. X.; Chao, L.; Chao, J.: Molecular cloning, tissue-specific
- [86988] expression, and cellular localization of human protasin mRNA. J.
- [86989] Biol. Chem. 270: 13483-13489, 1995.
- [86990] 15287.
- [86991] Yu, J. X.; Chao, L.; Ward, D. C.; Chao, J.: Structure and

chromosomal

[86992] localization of the human prostatic (PRSS8) gene. Genomics 32: 334–340,

[86993] 1996.

[86994] 15288.

[86995] Carlberg, C.; Hooft van Huijsduijnen, R.; Staple, J. K.; DeLamar, T.

[86996] J. F.; Becker-Andre, M.: RZR, a new family of retinoid-related orphan

[86997] receptors that function as both monomers and homodimers. Mol. Endocrinol. 8:

[86998] 757–770, 1994.

[86999] 15289.

[87000] Ueda, H. R.; Chen, W.; Adachi, A.; Wakamatsu, H.; Hayashi, S.;

[87001] Takasugi, T.; Nagano, M.; Nakahama, K.; Suzuki, Y.; Sugano, S.; Iino,

[87002] M.; Shigemoto, Y.; Hashimoto, S.: A transcription factor response

[87003] element for gene expression during circadian night. Nature 418:

[87004] 534–539, 2002.

[87005] 15290.

[87006] Masternak, K.; Barras, E.; Zufferey, M.; Conrad, B.;
Corthals,

[87007] G.; Aebersold, R.; Sanchez, J.-C.; Hochstrasser, D. F.;
Mach, B.;

[87008] Reith, W.: A gene encoding a novel RFX-associated trans-
activator

[87009] is mutated in the majority of MHC class II deficiency pa-
tients. Nature

[87010] Genet. 20: 273-277, 1998.

[87011] 15291.

[87012] Prange, C. K.; Pennacchio, L. A.; Lieuallen, K.; Fan, W.;
Lennon,

[87013] G. G.: Characterization of the human neurocan gene,
CSPG3. Gene 221:

[87014] 199-205, 1998.

[87015] 15292.

[87016] Rauch, U.; Grimpe, B.; Kulbe, G.; Arnold-Ammer, I.; Beier,
D. R.;

[87017] Fassler, R.: Structure and chromosomal localization of the
mouse

- [87018] neurocan gene. Genomics 28: 405–410, 1995.
- [87019] 15293.
- [87020] Rauch, U.; Karthikeyan, L.; Maurel, P.; Margolis, R. U.;
Margolis,
- [87021] R. K.: Cloning and primary structure of neurocan, a devel-
opmentally
- [87022] regulated, aggregating chondroitin sulfate proteoglycan of
brain. J.
- [87023] Biol. Chem. 267: 19536–19547, 1992.
- [87024] 15294.
- [87025] Drummond, J. T.; Genschel, J.; Wolf, E.; Modrich, P.: DHFR/
MSH3
- [87026] amplification in methotrexate-resistant cells alters the
hMutS-alpha/hMutS-beta
- [87027] ratio and reduces the efficiency of base-base mismatch
repair. Proc.
- [87028] Nat. Acad. Sci. 94: 10144–10149, 1997.
- [87029] 15295.
- [87030] Fujii, H.; Shimada, T.: Isolation and characterization of
cDNA
- [87031] clones derived from the divergently transcribed gene in

the region

[87032] upstream from the human dihydrofolate reductase gene. J.

Biol. Chem. 264:

[87033] 10057–10064, 1989.

[87034] 15296.

[87035] Inokuchi, K.; Ikejima, M.; Watanabe, A.; Nakajima, E.;

Orimo, H.;

[87036] Nomura, T.; Shimada, T.: Loss of expression of the human

MSH3 gene

[87037] in hematological malignancies. Biochem. Biophys. Res.

Commun. 214:

[87038] 171–179, 1995.

[87039] 15297.

[87040] Linton, J. P.; Yen, J. Y.-J.; Selby, E.; Chen, Z.; Chinsky, J.

[87041] M.; Liu, K.; Kellems, R. E.; Crouse, G. F.: Dual bidirectional

promoters

[87042] at the mouse dhfr locus: cloning and characterization of

two mRNA

[87043] classes of the divergently transcribed Rep-1 gene. Molec.

Cell. Biol. 9:

[87044] 3058–3072, 1989.

- [87045] 15298.
- [87046] Marra, G.; Iaccarino, I.; Lettieri, T.; Roscilli, G.; Delmastro,
[87047] P.; Jiricny, J.: Mismatch repair deficiency associated with
overexpression
[87048] of the MSH3 gene. Proc. Nat. Acad. Sci. 95: 8568–8573,
1998.
[87049] 15299.
- [87050] Nakajima, E.; Orimo, H.; Ikejima, M.; Shimada, T.: Nine-bp
repeat
[87051] polymorphism in exon 1 of the hMSH3 gene. Jpn. J. Hum.
Genet. 40:
[87052] 343–345, 1995.
[87053] 15300.
- [87054] Risinger, J. I.; Umar, A.; Boyd, J.; Berchuck, A.; Kunkel, T.
A.;
- [87055] Barrett, J. C.: Mutation of MSH3 in endometrial cancer and
evidence
[87056] for its functional role in heteroduplex repair. Nature
Genet. 14:
[87057] 102–109, 1996.
[87058] 15301.

- [87059] Smith, M. L.; Mitchell, P. J.; Crouse, G. F.: Analysis of the
[87060] mouse Dhfr/Rep-3 major promoter region by using
linker-scanning and
[87061] internal deletion mutations and DNase I footprinting.
Molec. Cell.
[87062] Biol. 10: 6003-6012, 1990.
[87063] 15302.
[87064] Watanabe, A.; Ikejima, M.; Suzuki, N.; Shimada, T.: Ge-
nomic organization
[87065] and expression of the human MSH3 gene. Genomics 31:
311-318, 1996.
[87066] 15303.
[87067] Yin, J.; Kong, D.; Wang, S.; Zou, T.-T. Souza, R. F.; Smolin-
ski,
[87068] K. N.; Lynch, P. M.; Hamilton, S. R.; Sugimura, H.; Powell,
S. M.;
[87069] Young, J.; Abraham, J. M.; Meltzer, S. J.: Mutation of
hMSH3 and
[87070] hMSH6 mismatch repair genes in genetically unstable hu-
man colorectal
[87071] and gastric carcinomas. Hum. Mutat. 10: 474-478, 1997.

- [87072] 15304.
- [87073] Burger, B.; Uhlhaas, S.; Mangold, E.; Propping, P.; Friedl, W.;
- [87074] Jenne, D.; Dockter, G.; Back, W.: Novel de novo mutation of MADH4/SMAD4
- [87075] in a patient with juvenile polyposis. (Letter) Am. J. Med. Genet. 110:
- [87076] 289–291, 2002.
- [87077] 15305.
- [87078] Sakamoto, A.; Ono, K.; Abe, M.; Jasmin, G.; Eki, T.; Murakami,
- [87079] Y.; Masaki, T.; Toyo-oka, T.; Hanaoka, F.: Both hypertrophic and
- [87080] dilated cardiomyopathies are caused by mutation of the same gene,
- [87081] delta-sarcoglycan, in hamster: an animal model of disrupted dystrophin-associated
- [87082] glycoprotein complex. Proc. Nat. Acad. Sci. 94: 13873–13878, 1997.
- [87083] 15306.
- [87084] Kurima, K.; Peters. L. M.; Yang, Y.; Riazuddin, S.; Ahmed, Z. M.;

- [87085] Naz, S.; Arnaud, D.; Drury, S.; Mo, J.; Makishima, T.; Ghosh, M.;
- [87086] Menon, P. S. N.; and 13 others: Dominant and recessive deafness
- [87087] caused by mutations of a novel gene, TMC1, required for cochlear hair-cell
- [87088] function. *Nature Genet.* 30: 277–284, 2002.
- [87089] 15307.
- [87090] Bryant, P. J.; Huettner, B.; Held, L. I., Jr.; Ryerse, J.; Szidonya,
- [87091] J.: Mutations at the fat locus interfere with cell proliferation
- [87092] control and epithelial morphogenesis in *Drosophila*. *Dev. Biol.* 129:
- [87093] 541–554, 1988.
- [87094] 15308.
- [87095] Dunne, J.; Hanby, A. M.; Poulsom, R.; Jones, T. A.; Sheer, D.;
- [87096] Chin, W. G.; Da, S. M.; Zhao, Q.; Beverley, P. C. L.; Owen, M. J.
- [87097] : Molecular cloning and tissue expression of FAT, the human homologue

[87098] of the *Drosophila* fat gene that is located on chromosome 4q34–q35

[87099] and encodes a putative adhesion molecule. *Genomics* 30: 207–223,

[87100] 1995.

[87101] 15309.

[87102] Hortsch, M.; Goodman, C. S.: Cell and substrate adhesion molecules

[87103] in *Drosophila*. *Ann. Rev. Cell. Biol.* 7: 505–557, 1991.

[87104] 15310.

[87105] Mahoney, P. A.; Weber, U.; Onofrechuk, P.; Biessmann, H.; Bryant,

[87106] P. J.; Goodman, C. S.: The fat tumor suppressor gene in *Drosophila*

[87107] encodes a novel member of the cadherin gene superfamily. *Cell* 67:

[87108] 853–868, 1991.

[87109] 15311.

[87110] Browning, J. L.; Ngam-ek, A.; Lawton, P.; DeMarinis, J.; Tizard,

[87111] R.; Chow, E. P.; Hession, C.; O'Brine-Greco, B.; Foley, S. F.;

Ware,

[87112] C. F.: Lymphotoxin beta, a novel member of the TNF family that forms

[87113] a heteromeric complex with lymphotoxin on the cell surface. Cell 72:

[87114] 847-856, 1993.

[87115] 15312.

[87116] Crowe, P. D.; VanArsdale, T. L.; Walter, B. N.; Ware, C. F.; Hession,

[87117] C.; Ehrenfels, B.; Browning, J. L.; Din, W. S.; Goodwin, R. G.; Smith,

[87118] C. A.: A lymphotoxin-beta-specific receptor. Science 264: 707-710,

[87119] 1994.

[87120] 15313.

[87121] Nakamura, T.; Tashiro, K.; Nazarea, M.; Nakano, T.; Sasayama, S.;

[87122] Honjo, T.: The murine lymphotoxin-beta receptor cDNA: isolation by

[87123] the signal sequence trap and chromosomal mapping. Genomics 30: 312-319,

[87124] 1995.

- [87125] 15314.
- [87126] Warzocha, K.; Renard, N.; Charlot, C.; Bienvenu, J.;
Coiffier,
- [87127] B.; Salles, G.: Identification of two lymphotoxin beta iso-
forms expressed
- [87128] in human lymphoid cell lines and non-Hodgkin's lym-
phomas. Biochem.
- [87129] Biophys. Res. Commun. 238: 273-276, 1997.
- [87130] 15315.
- [87131] Tashiro, K.; Tada, H.; Heilker, R.; Shirozu, M.; Nakano, T.;
Honjo,
- [87132] T.: Signal sequence trap: a cloning strategy for secreted
proteins
- [87133] and type I membrane proteins. Science 261: 600-603,
1993.
- [87134] 15316.
- [87135] George, A.; Sabsay, B.; Simonian, P. A.; Veis, A.: Charac-
terization
- [87136] of a novel dentin matrix acidic phosphoprotein: implica-
tions for induction
- [87137] of biomineralization . J. Biol. Chem. 268: 12624-12630,
1993.

[87138] 15317.

[87139] Hirst, K. L.; Simmons, D.; Feng, J.; Aplin, H.; Dixon, M. J.;
MacDougall,

[87140] M.: Elucidation of the sequence and the genomic organi-
zation of the

[87141] human dentin matrix acidic phosphoprotein 1 (DMP1)
gene: exclusion

[87142] of the locus from a causative role in the pathogenesis of
dentinogenesis

[87143] imperfecta type II. Genomics 42: 38-45, 1997.

[87144] 15318.

[87145] MacDougall, M.; DuPont, B. R.; Simmons, D.; Leach, R. J.:
Assignment

[87146] of DMP1 to human chromosome 4 band q21 by in situ hy-
bridization. Cytogenet.

[87147] Cell. Genet. 74: 189 only, 1996.

[87148] 15319.

[87149] Lu, Z.; Xu, S.; Joazeiro, C.; Cobb, M. H.; Hunter, T.: The
PHD

[87150] domain of MEKK1 acts as an E3 ubiquitin ligase and medi-
ates ubiquitination

- [87151] and degradation of ERK1/2. *Molec. Cell* 9: 945–956, 2002.
- [87152] 15320.
- [87153] Xu, G. M.; Sikaneta, T.; Sullivan, B. M.; Zhang, Q.; Andreucci, M.; Stehle, T.; Drummond, I.; Arnaout, M. A.: Polycystin-1 interacts with intermediate filaments. *J. Biol. Chem.* 276: 46544–46552, 2001.
- [87156] 15321.
- [87157] Zerres, K.; Rudnik-Schoneborn, S.; Deget, F.; Beetz, R.; Brodehl, J.; Kuwertz-Broking, E.; Muller-Wiefel, D.; Rascher, W.; Ruder, H.; Scharer, K.; Weber, H. P.: Childhood onset autosomal dominant polycystic kidney disease in sibs: clinical picture and recurrence risk. *J. Med. Genet.* 30: 583–588, 1993.
- [87161] 15322.
- [87162] 15322.
- [87163] Barnard, R. C.; Pascall, J. C.; Brown, K. D.; McKay, I. A.; Williams,

- [87164] N. S.; Bustin, S. A.: Coding sequence of ERF-1, the human homologue
[87165] of Tis11b/cMG1, members of the Tis11 family of early re-
[87166] sponse genes. Nucleic
[87167] Acids Res. 21: 3580 only, 1993.
[87167] 15323.
- [87168] Bustin, S. A.; Xiao-Feng, N.; Barnard, R. C.; Kumar, V.;
Pascall,
[87169] J. C.; Brown, K. D.; Leigh, I. M.; Williams, N. S.; McKay, I.
A.:
[87170] Cloning and characterisation of ERF1, a human member of
the Tis11
[87171] family of early-response genes. DNA Cell Biol. 13:
449-459, 1994.
[87172] 15324.
- [87173] Maclean, K. N.; See, C. G.; McKay, I. A.; Bustin, S. A.: The
human
[87174] immediate early gene BRF1 maps to chromosome
14q22-q24. Genomics 30:
[87175] 89-90, 1995.
[87176] 15325.

- [87177] Ning, Z.-Q.; Norton, J. D.; Li, J.; Murphy, J. J.: Distinct mechanisms
- [87178] for rescue from apoptosis in Ramos human B cells by signaling through
- [87179] CD40 and interleukin-4 receptor: a role for inhibition of an early
- [87180] response gene, Berg36. *Europ. J. Immun.* 26: 2356-2363, 1996.
- [87181] 15326.
- [87182] Maas, S.; Kim, Y.-G.; Rich, A.: Genomic clustering of tRNA-specific
- [87183] adenosine deaminase ADAT1 and two tRNA synthetases. *Mammalian Genome* 12:
- [87184] 387-393, 2001.
- [87185] 15327.
- [87186] Bolz, H.; von Brederlow, B.; Ramirez, A.; Bryda, E. C.; Kutsche,
- [87187] K.; Nothwang, H. G.; Seeliger, M.; Cabrera, M. C.-S.; Vila, M. C.;
- [87188] Molina, O. P.; Gal, A.; Kubisch, C.: Mutation of CDH23, encoding
- [87189] a new member of the cadherin gene family, causes Usher

syndrome type

[87190] 1D. Nature Genet. 27: 108–112, 2001.

[87191] 15328.

[87192] Bork, J. M.; Peters, L. M.; Riazuddin, S.; Bernstein, S. L.;
Ahmed,

[87193] Z. M.; Ness, S. L.; Polomeno, R.; Ramesh, A.; Schloss, M.;
Srisailpathy,

[87194] C. R. S.; Wayne, S.; Bellman, S.; and 16 others: Usher syn-
drome

[87195] 1D and nonsyndromic autosomal recessive deafness
DFNB12 are caused

[87196] by allelic mutations of the novel cadherin-like gene
CDH23. Am. J.

[87197] Hum. Genet. 68: 26–37, 2001.

[87198] 15329.

[87199] Derynck, R.; Gelbart, W. M.; Harland, R. M.; Heldin, C.–H.;
Kern,

[87200] S. E.; Massague, J.; Melton, D. A.; Mlodzik, M.; Padgett, R.
W.; Roberts,

[87201] A. B.; Smith, J.; Thomsen, G. H.; Vogelstein, B.; Wang,
X.–F.: Nomenclature:

[87202] vertebrate mediators of TGF-beta family signals. (Letter)

Cell 87:

[87203] 173 only, 1996.

[87204] 15330.

[87205] Friedl, W.; Kruse, R.; Uhlhaas, S.; Stolte, M.; Schartmann, B.;

[87206] Keller, K. M.; Jungck, M.; Stern, M.; Loff, S.; Back, W.;
Propping,

[87207] P.; Jenne, D. E.: Frequent 4-bp deletion in exon 9 of the
SMAD4/MADH4

[87208] gene in familial juvenile polyposis patients. Genes Chromosomes Cancer 25:

[87209] 403–406, 1999.

[87210] 15331.

[87211] Hahn, S. A.; Schutte, M.; Hoque, T. M. S.; Moskaluk, C. A.;
da

[87212] Costa, L. T.; Rozenblum, E.; Weinstein, C. L.; Fischer, A.;
Yeo, C.

[87213] J.; Hruban, R. H.; Kern, S. E.: DPC4, a candidate tumor
suppressor

[87214] gene at human chromosome 18q21.1. Science 271:
350–354, 1996.

[87215] 15332.

- [87216] Howe, J. R.; Shellnut, J.; Wagner, B.; Ringold, J. C.; Sayed,
[87217] M. G.; Ahmed, A. F.; Lynch, P. M.; Amos, C. I.; Sistonen, P.;
Aaltonen,
[87218] L. A.: Common deletion of SMAD4 in juvenile polyposis is
a mutational
[87219] hotspot. Am. J. Hum. Genet. 70: 1357–1362, 2002.
[87220] 15333.
[87221] Inman, G. J.; Nicolas, F. J.; Hill, C. S.: Nucleocytoplasmic
[87222] shuttling of Smads 2, 3, and 4 permits sensing of TGF-
beta receptor
[87223] activity. Molec. Cell 10: 283–294, 2002.
[87224] 15334.
[87225] Kim, S. K.; Fan, Y.; Papadimitrakopoulou, V.; Clayman, G.;
Hittelman,
[87226] W. N.; Hong, W. K.; Lotan, R.; Mao, L.: DPC4, a candidate
tumor suppressor
[87227] gene, is altered infrequently in head and neck squamous
cell carcinoma. Cancer
[87228] Res. 56: 2519–2521, 1996.
[87229] 15335.

- [87230] Kinzler, K. W.; Vogelstein, B.: Landscaping the cancer terrain. Science 280:
- [87231] 1036–1037, 1998.
- [87232] 15336.
- [87233] MacGrogan, D.; et al.; et al: Comparative mutational analysis
- [87234] of DPC4 (Smad4) in prostatic and colorectal carcinomas. Oncogene 15:
- [87235] 1111–1114, 1997.
- [87236] 15337.
- [87237] Roth, S.; Johansson, M.; Loukola, A.; Peltomaki, P.; Jarvinen,
- [87238] H.; Mecklin, J.–P.; Aaltonen, L. A.: Mutation analysis of SMAD2,
- [87239] SMAD3, and SMAD4 genes in hereditary non–polyposis colorectal cancer. J.
- [87240] Med. Genet. 37: 298–300, 2000.
- [87241] 15338.
- [87242] Schutte, M.; Hruban, R. H.; Hedrick, L.; Cho, K. R.; Nadasdy,
- [87243] G. M.; Weinstein, C. L.; Bova, G. S.; Isaacs, W. B.; Cairns, P.;

Nawroz,

[87244] H.; Sidransky, D.; Casero, R. A., Jr.; Meltzer, P. S.; Hahn, S. A.;

[87245] Kern, S. E.: DPC4 gene in various tumor types. Cancer Res. 56:

[87246] 2527–2530, 1996.

[87247] 15339.

[87248] Shioda, T.; Lechleider, R. J.; Dunwoodie, S. L.; Li, H.; Yahata,

[87249] T.; de Caestecker, M. P.; Fenner, M. H.; Roberts, A. B.; Is-selbacher,

[87250] K. J.: Transcriptional activating activity of Smad4: roles of SMAD

[87251] hetero-oligomerization and enhancement by an associating transactivator. Proc.

[87252] Nat. Acad. Sci. 95: 9785–9790, 1998.

[87253] 15340.

[87254] Sirard, C.; de la Pompa, J. L.; Elia, A.; Itie, A.; Mirtsos, C.;

[87255] Cheung, A.; Hahn, S.; Wakeham, A.; Schwartz, L.; Kern, S. E.; Rossant,

[87256] J.; Mak, T. W.: The tumor suppressor gene Smad4/Dpc4 is

required

[87257] for gastrulation and later for anterior development of the mouse embryo. Genes

[87258] Dev. 12: 107–119, 1998.

[87259] 15341.

[87260] Tagaki, Y.; et al.; et al: Somatic alterations of the DPC4 gene

[87261] in human colorectal cancers in vivo. Gastroenterology 111: 1369–1372,

[87262] 1996.

[87263] 15342.

[87264] Takaku, K.; Oshima, M.; Miyoshi, H.; Matsui, M.; Seldin, M. F.;

[87265] Taketo, M. M.: Intestinal tumorigenesis in compound mutant mice of

[87266] both Dpc4 (Smad4) and Apc genes. Cell 92: 645–656, 1998.

[87267] 15343.

[87268] Thiagalingam, S.; Lengauer, C.; Leach, F. S.; Schutte, M.; Hahn,

[87269] S. A.; Overhauser, J.; Willson, J. K. V.; Markowitz, S.;

Hamilton,

- [87270] S. R.; Kern, S. E.; Kinzler, K. W.; Vogelstein, B.: Evaluation of
- [87271] candidate tumour suppressor genes on chromosome 18 in colorectal cancers. Nature
- [87272] Genet. 13: 343–346, 1996.
- [87273] 15344.
- [87274] Zawel, L.; Dai, J. L.; Buckhaults, P.; Zhou, S.; Kinzler, K. W.;
- [87275] Vogelstein, B.; Kern, S. E.: Human Smad3 and Smad4 are sequence-specific
- [87276] transcription activators. Molec. Cell 1: 611–617, 1998.
- [87277] 15345.
- [87278] Zhou, S.; Buckhaults, P.; Zawel, L.; Bunz, F.; Riggins, G.; Le
- [87279] Dai, J.; Kern, S. E.; Kinzler, K. W.; Vogelstein, B.: Targeted deletion
- [87280] of Smad4 shows it is required for transforming growth factor beta
- [87281] and activin signaling in colorectal cancer cells. Proc. Nat. Acad.
- [87282] Sci. 95: 2412–2416, 1998.
- [87283] 15346.

- [87284] Abbs, S.; Roberts, R. G.; Mathew, C. G.; Bentley, D. R.; Bo-
brow,
- [87285] M.: Accurate assessment of intragenic recombination fre-
quency within
- [87286] the Duchenne muscular dystrophy gene. *Genomics* 7:
602-606, 1990.
- [87287] 15347.
- [87288] Ahn, A. H.; Kunkel, L. M.: The structural and functional di-
versity
- [87289] of dystrophin. *Nature Genet.* 3: 283-291, 1993.
- [87290] 15348.
- [87291] Alwine, J. C.; Kemp, D. J.; Stark, G. R.: Method for detec-
tion
- [87292] of specific RNAs in agarose gels by transfer to diazoben-
zyloxymethyl-paper
- [87293] and by hybridization with DNA probes. *Proc. Nat. Acad.*
Sci. 74:
- [87294] 5350-5354, 1977.
- [87295] 15349.
- [87296] Angelini, C.; Beggs, A. H.; Hoffman, E. P.; Fanin, M.;
Kunkel,

- [87297] L. M.: Enormous dystrophin in a patient with Becker muscular dystrophy. *Neurology* 40: 808–812, 1990.
- [87298] 15350.
- [87299] 15350.
- [87300] Badorff, C.; Berkely, N.; Mehrotra, S.; Talhouk, J. W.; Rhoads, R. E.; Knowlton, K. U.: Enteroviral protease 2A directly cleaves dystrophin and is inhibited by a dystrophin-based substrate analogue. *J. Biol. Chem.* 275: 11191–11197, 2000.
- [87301] 15351.
- [87302] Badorff, C.; Lee, G.-H.; Lamphear, B. J.; Martone, M. E.; Campbell, K. P.; Rhoads, R. E.; Knowlton, K. U.: Enteroviral protease 2A cleaves dystrophin: evidence of cytoskeletal disruption in an acquired cardiomyopathy. *Nature Med.* 5: 320–326, 1999.
- [87303] 15352.
- [87304] Bakker, E.; Pearson, P. L.: Mutation of the Duchenne mus-

cular

- [87311] dystrophy gene associated with meiotic recombination.
(Letter) Clin.
- [87312] Genet. 30: 347–349, 1986.
- [87313] 15353.
- [87314] Bakker, E.; Hofker, M. H.; Goor, N.; Mandel, J. L.; Wroge-
mann,
- [87315] K.; Davies, K. E.; Kunkel, L. M.; Willard, H. F.; Fenton, W.
A.; Sandkuyl,
- [87316] L.; Majoor-Krakauer, D.; van Essen, A. J.; Jahoda, M. G. J.;
Sachs,
- [87317] E. S.; van Ommen, G. J. B.; Pearson, P. L.: Prenatal diagno-
sis and
- [87318] carrier detection of Duchenne muscular dystrophy with
closely linked
- [87319] RFLPs. Lancet I: 655–658, 1985.
- [87320] 15354.
- [87321] Bakker, E.; Van Broeckhoven, C.; Bonten, E. J.; van de
Vooren,
- [87322] M. J.; Veenema, H.; Van Hul, W.; Van Ommen, G. J. B.;
Vandenberghe,
- [87323] A.; Pearson, P. L.: Germline mosaicism and Duchenne

muscular dystrophy

[87324] mutations. Nature 329: 554–556, 1987.

[87325] 15355.

[87326] Bar, S.; Barnea, E.; Levy, Z.; Neuman, S.; Yaffe, D.; Nudel, U.

[87327] : A novel product of the Duchenne muscular dystrophy gene which greatly

[87328] differs from the known isoforms in its structure and tissue distribution. Biochem.

[87329] J. 272: 557–560, 1990.

[87330] 15356.

[87331] Barbieri, A. M.; Soriani, N.; Tubiello, G. M.; Ferrari, M.; Carrera,

[87332] P.: A nonsense mutation (gln-673-term) in exon 17 of the human dystrophin

[87333] gene detected by heteroduplex analysis. Hum. Genet. 96: 343–344,

[87334] 1995.

[87335] 15357.

[87336] Bartlett, R. J.; Pericak-Vance, M. A.; Koh, J.; Yamaoka, L. H.;

- [87337] Chen, J. C.; Hung, W.-Y.; Speer, M. C.; Wapenaar, M. C.;
Van Ommen,
- [87338] G. J. B.; Bakker, E.; Pearson, P. L.; Kandt, R. S.; Siddique,
T.;
- [87339] Gilbert, J. R.; Lee, J. E.; Sirotkin-Roses, M. J.; Roses, A. D.:
- [87340] Duchenne muscular dystrophy: high frequency of dele-
tions. *Neurology* 38:
- [87341] 1-4, 1988.
- [87342] 15358.
- [87343] Barton-Davis, E. R.; Cordier, L.; Shoturma, D. I.; Leland, S.
- [87344] E.; Sweeney, H. L.: Aminoglycoside antibiotics restore dys-
trophin
- [87345] function to skeletal muscles of mdx mice. *J. Clin. Invest.*
104:
- [87346] 375-381, 1999.
- [87347] 15359.
- [87348] Bastianutto, C.; Bestard, J. A.; Lahnakoski, K.; Broere, D.;
- De
- [87349] Visser, M.; Zaccolo, M.; Pozzan, T.; Ferlini, A.; Muntoni, F.;
- Patarnello,
- [87350] T.; Klamut, H. J.: Dystrophin muscle enhancer 1 is impli-

cated in

[87351] the activation of non-muscle isoforms in the skeletal muscle of patients

[87352] with X-linked dilated cardiomyopathy. Hum. Molec. Genet. 10: 2627-2635,

[87353] 2001.

[87354] 15360.

[87355] Baumbach, L. L.; Chamberlain, J. S.; Ward, P. A.; Farwell, N.

[87356] J.; Caskey, C. T.: Molecular and clinical correlation of deletion

[87357] leading to Duchenne and Becker muscular dystrophies. Neurology 39:

[87358] 465-474, 1989.

[87359] 15361.

[87360] Baumbach, L. L.; Ward, P. A.; Fenwick, R.; Caskey, C. T.: Analysis

[87361] of mutations at the Duchenne muscular dystrophy locus provides no

[87362] evidence for illegitimate recombination in deletion formation. (Abstract) Am.

[87363] J. Hum. Genet. 45 (suppl.): A173, 1989.

[87364] 15362.

[87365] Beggs, A. H.; Koenig, M.; Boyce, F. M.; Kunkel, L. M.: De-
tection

[87366] of 98% of DMD/BMD gene deletions by polymerase chain
reaction. Hum.

[87367] Genet. 86: 45–48, 1990.

[87368] 15363.

[87369] Berko, B. A.; Swift, M.: X-linked dilated cardiomyopathy.
New

[87370] Eng. J. Med. 316: 1186–1191, 1987.

[87371] 15364.

[87372] Bettecken, T.; Muller, C. R.: Identification of a 220-kb in-
sertion

[87373] into the Duchenne gene in a family with an atypical course
of muscular

[87374] dystrophy. Genomics 4: 592–596, 1989.

[87375] 15365.

[87376] Bies, R. D.: X-linked dilated cardiomyopathy. (Letter) New
Eng.

[87377] J. Med. 330: 368–369, 1994.

- [87378] 15366.
- [87379] Bies, R. D.; Caskey, C. T.; Fenwick, R.: An intact cysteine-rich
domain is required for dystrophin function. J. Clin. Invest.
90:
666-672, 1992.
- [87382] 15367.
- [87383] Bittner, R. E.; Streubel, B.; Shorny, S.; Schaden, G.; Voit, T.;
Hoger, H.: Coisogenic all-plus-one immunization: a
model for identifying
missing proteins in null-mutant conditions. Antibodies to
dystrophin
in mdx mouse after transplantation of muscle from normal coisogenic
donor. Neuropediatrics 25: 176-182, 1994.
- [87388] 15368.
- [87389] Bleyl, S. B.; Mumford, B. R.; Brown-Harrison, M.-C.;
Pagotto, L.
- [87390] T.; Carey, J. C.; Pysher, T. J.; Ward, K.; Chin, T. K.:
Xq28-linked
noncompaction of the left ventricular myocardium: prena-

tal diagnosis

[87392] and pathologic analysis of affected individuals. Am. J. Med. Genet. 72:

[87393] 257–265, 1997.

[87394] 15369.

[87395] Bleyl, S. B.; Mumford, B. R.; Thompson, V.; Carey, J. C.; Pysker,

[87396] T. J.; Chin, T. K.; Ward, K.: Neonatal, lethal noncompaction of the

[87397] left ventricular myocardium is allelic with Barth syndrome. Am. J.

[87398] Hum. Genet. 61: 868–872, 1997.

[87399] 15370.

[87400] Mohapatra, B.; Verma, S.; Shankar, S.; Suri, A.: Molecular cloning

[87401] of human testis mRNA specifically expressed in haploid germ cells,

[87402] having structural homology with the A-kinase anchoring proteins. Biochem.

[87403] Biophys. Res. Commun. 244: 540–545, 1998.

[87404] 15371.

- [87405] Turner, R. M. O.; Johnson, L. R.; Haig-Ladewig, L.; Gerton, G.
- [87406] L.; Moss, S. B.: An X-linked gene encodes a major human sperm fibrous
- [87407] sheath protein, hAKAP82: genomic organization, protein kinase A-RII
- [87408] binding, and distribution of the precursor in the sperm tail. J.
- [87409] Biol. Chem. 273: 32135-32141, 1998.
- [87410] 15372.
- [87411] Dever, T. E.; Wei, C.-L.; Benkowski, L. A.; Browning, K.; Merrick,
- [87412] W. C.; Hershey, J. W. B.: Determination of the amino acid sequence
- [87413] of rabbit, human, and wheat germ protein synthesis factor eIF-4C by
- [87414] cloning and chemical sequencing. J. Biol. Chem. 269: 3212-3218,
- [87415] 1994.
- [87416] 15373.
- [87417] Knebelmann, B.; Breillat, C.; Forestier, L.; Arrondel, C.; Jaccassier,

- [87418] D.; Giatras, I.; Drouot, L; Deschenes, G.; Grunfeld, J.-P.; Broyer,
- [87419] M.; Gubler, M.-C.; Antignac, C.: Spectrum of mutations in the COL4A5
- [87420] collagen gene in X-linked Alport syndrome. Am. J. Hum. Genet. 59:
- [87421] 1221-1232, 1996.
- [87422] 15374.
- [87423] Perkoff, G. T.; Stephens, F. E.; Dolowitz, D. A.; Tyler, F. H.
- [87424] : A clinical study of hereditary interstitial pyelonephritis. Arch.
- [87425] Intern. Med. 88: 191-200, 1951.
- [87426] 15375.
- [87427] Lee, S.-H.; Kim, W.-H.; Kim, H.-K.; Woo, K.-M.; Nam, H.-S.; Kim,
- [87428] H.-S.; Kim, J.-G.; Cho, M.-H.: Altered expression of the fragile
- [87429] histidine triad gene in primary gastric adenocarcinomas. Biochem.
- [87430] Biophys. Res. Commun. 284: 850-855, 2001.
- [87431] 15376.

- [87432] Markkanen, A.; Heinonen, K.; Knuutila, S.; de la Chapelle, A.
- [87433] : Methotrexate-induced increase in gap formation in human chromosome
- [87434] band 3p14. Hereditas 96: 317–319, 1982.
- [87435] 15377.
- [87436] Markkanen, A.; Knuutila, S.; de la Chapelle, A.: Inducible fragile
- [87437] site on chromosome 3. (Letter) Hum. Genet. 65: 217 only, 1983.
- [87438] 15378.
- [87439] Bell, M. V.; Hirst, M. C.; Nakahori, Y.; MacKinnon, R. N.; Roche,
- [87440] A.; Flint, T. J.; Jacobs, P. A.; Tommerup, N.; Tranebjaerg, L.; Froster-Iskenius,
- [87441] U.; Kerr, B.; Turner, G.; Lindenbaum, R. H.; Winter, R.; Pembrey,
- [87442] M.; Thibodeau, S.; Davies, K. E.: Physical mapping across the fragile
- [87443] X: hypermethylation and clinical expression of the fragile X syndrome. Cell 64:
- [87444] 861–866, 1991.

- [87445] 15379.
- [87446] Matsumoto, T.; Imamura, O.; Yamabe, Y.; Kuromitsu, J.;
Tokutake,
- [87447] Y.; Shimamoto, A.; Suzuki, N.; Satoh, M.; Kitao, S.;
Ichikawa, K.;
- [87448] Kataoka, H.; Sugawara, K.; Thomas, W.; Mason, B.; Tsuchi-
hashi, Z.;
- [87449] Drayna, D.; Sugawara, M.; Sugimoto, M.; Furuichi, Y.;
Goto, M.: Mutation
- [87450] and haplotype analyses of the Werner's syndrome gene
based on its
- [87451] genomic structure: genetic epidemiology in the Japanese
population. Hum.
- [87452] Genet. 100: 123–130, 1997.
- [87453] 15380.
- [87454] Filmus, J.; Shi, W.; Wong, Z. M.; Wong, M. J.: Identification
- [87455] of a new membrane-bound heparan sulphate proteogly-
can. Biochem. J. 311:
- [87456] 561–565, 1995.
- [87457] 15381.
- [87458] Zhou, J.; Hertz, J. M.; Tryggvason, K.: Mutation in the al-

pha-5(IV)

[87459] collagen chain in juvenile-onset Alport syndrome without hearing loss

[87460] or ocular lesions: detection by denaturing gradient gel electrophoresis

[87461] of a PCR product. Am. J. Hum. Genet. 50: 1291-1300, 1992.

[87462] 15382.

[87463] Bohler, M. C.; Seger, R. A.; Mouy, R.; Vilmer, E.; Fischer, A.;

[87464] Griscelli, C.: A study of 25 patients with chronic granulomatous

[87465] disease: a new classification by correlating respiratory burst, cytochrome

[87466] b and flavoprotein. J. Clin. Immun. 6: 136-145, 1986.

[87467] 15383.

[87468] Basler, E.; Grompe, M.; Parenti, G.; Caskey, C. T.; Ballabio,

[87469] A.: Identification of point mutations in three patients with steroid

[87470] sulfatase deficiency. (Abstract) Am. J. Hum. Genet. 47 (suppl.):

[87471] A150, 1990.

[87472] 15384.

[87473] Arranz, V.; Harper, F.; Florentin, Y.; Puvion, E.; Kress, M.;
Ernoul-Lange,

[87474] M.: Human and mouse MOK2 proteins are associated with
nuclear ribonucleoprotein

[87475] components and bind specifically to RNA and DNA
through their zinc

[87476] finger domains. *Molec. Cell. Biol.* 17: 2116–2126, 1997.

[87477] 15385.

[87478] Arranz, V.; Kress, M.; Ernoul-Lange, M.: Localization of
zinc

[87479] finger Mok2 gene to mouse chromosome 6, a new region
of homology with

[87480] human chromosome 19. *Mammalian Genome* 7: 77–78,
1996.

[87481] 15386.

[87482] Ernoul-Lange, M.; Arranz, V.; Le Conait, M.; Berger, R.;
Kress,

[87483] M.: Human and mouse Kruppel-like (MOK2) orthologue
genes encode two

[87484] different zinc finger proteins. *J. Molec. Evol.* 41: 784–794,
1995.

[87485] 15387.

[87486] Yasunaga, S.; Grati, M.; Cohen-Salmon, M.; El-Amraoui, A.; Mustapha,

[87487] M.; Salem, N.; El-Zir, E.; Loiselet, J.; Petit, C.: A mutation in

[87488] OTOF, encoding otoferlin, a FER-1-like protein, causes DFNB9, a nonsyndromic

[87489] form of deafness. *Nature Genet.* 21: 363-369, 1999.

[87490] 15388.

[87491] Berry, A.; Scott, H. S.; Kudoh, J.; Talior, I.; Korostishevsky,

[87492] M.; Wattenhofer, M.; Guipponi, M.; Barras, C.; Rossier, C.; Shibuya,

[87493] K.; Wang, J.; Kawasaki, K.; Asakawa, S.; Minoshima, S.; Shimizu, N.;

[87494] Antonarakis, S.; Bonne-Tamir, B.: Refined localization of autosomal

[87495] recessive nonsyndromic deafness DFNB10 locus using 34 novel microsatellite

[87496] markers, genomic structure, and exclusion of six known genes in the

[87497] region. *Genomics* 68: 22-29, 2000.

[87498] 15389.

- [87499] Scott, H. S.; Kudoh, J.; Wattenhofer, M.; Shibuya, K.; Berry, A.;
- [87500] Chrast, R.; Guipponi, M.; Wang, J.; Kawasaki, K.; Asakawa, S.; Minoshima, S.; Younus, F.; and 10 others: Insertion of beta-satellite repeats
- [87502] identifies a transmembrane protease causing both congenital and childhood
- [87503] onset autosomal recessive deafness. *Nature Genet.* 27: 59–63, 2001.
- [87504] 15390.
- [87505] Good, P. J.; Chen, Q.; Warner, S. J.; Herring, D. C.: A family
- [87506] of human RNA-binding proteins related to the *Drosophila* Bruno transcriptional
- [87507] regulator. *J. Biol. Chem.* 275: 28583–28592, 2000.
- [87508] 15391.
- [87509] Roberts, R.; Timchenko, N. A.; Miller, J. W.; Reddy, S.; Caskey,
- [87510] C. T.; Swanson, M. S.; Timchenko, L. T.: Altered phosphorylation
- [87511] and intracellular distribution of a (CUG)_n triplet repeat RNA-binding

- [87512] protein in patients with myotonic dystrophy and in myotonic protein
- [87513] kinase knockout mice. Proc. Nat. Acad. Sci. 94: 13221–13226, 1997.
- [87514] 15392.
- [87515] Li, Z.; Jiang, H.; Xie, W.; Zhang, Z.; Smrcka, A. V.; Wu, D.:
- [87516] Roles of PLC- β 2 and - β 3 and PI3K- γ in chemoattractant-mediated
- [87517] signal transduction. Science 287: 1046–1049, 2000.
- [87518] 15393.
- [87519] Makita, N.; Sloan-Brown, K.; Weghuis, D. O.; Ropers, H. H.; George,
- [87520] A. L., Jr.: Genomic organization and chromosomal assignment of the
- [87521] human voltage-gated Na(+) channel β -1 subunit gene (SCN1B). Genomics 23:
- [87522] 628–634, 1994.
- [87523] 15394.
- [87524] Scheffer, I. E.; Berkovic, S. F.: Generalized epilepsy with febrile
- [87525] seizures plus: a genetic disorder with heterogeneous clin-

ical phenotypes. Brain 120:

[87526] 479–490, 1997.

[87527] 15395.

[87528] Wallace, R. H.; Wang, D. W.; Singh, R.; Scheffer, I. E.;
George,

[87529] A. L., Jr.; Phillips, H. A.; Saar, K.; Reis, A.; Johnson, E. W.;
Sutherland,

[87530] G. R.; Berkovic, S. F.; Mulley, J. C.: Febrile seizures and
generalized

[87531] epilepsy associated with a mutation in the Na(+)-channel
beta-1 subunit

[87532] gene SCN1B. Nature Genet. 19: 366–370, 1998.

[87533] 15396.

[87534] Gassmann, M.; Casagrande, F.; Orioli, D.; Simon, H.; Lai,
C.; Klein,

[87535] R.; Lemke, G.: Aberrant neural and cardiac development in
mice lacking

[87536] the ErbB4 neuregulin receptor. Nature 378: 390–394,
1995.

[87537] 15397.

[87538] Golding, J. P.; Trainor, P.; Krumlauf, R.; Gassmann, M.: De-

fects

[87539] in pathfinding by cranial neural crest cells in mice lacking the neuregulin

[87540] receptor ErbB4. *Nature Cell Biol.* 2: 103–109, 2000.

[87541] 15398.

[87542] Zimonjic, D. B.; Alimandi, M.; Miki, T.; Popescu, N. C.; Kraus,

[87543] M. H.: Localization of the human HER4/erbB-4 gene to chromosome 2. *Oncogene* 10:

[87544] 1235–1237, 1995.

[87545] 15399.

[87546] Adibi, S. A.: The oligopeptide transporter (Pept-1) in human intestine:

[87547] biology and function. *Gastroenterology* 113: 332–340, 1997.

[87548] 15400.

[87549] Fei, Y. -J.; Kanai, Y.; Nussberger, S.; Ganapathy, V.; Leibach,

[87550] F. H.; Romero, M. F.; Singh, S. K.; Boron, W. F.; Hediger, M. A.:

[87551] Expression cloning of a mammalian proton-coupled

oligopeptide transporter. Nature 368:

[87552] 563–566, 1994.

[87553] 15401.

[87554] Liang, R.; Fei, Y.-J.; Prasad, P. D.; Ramamoorthy, S.; Han, H.;

[87555] Yang–Feng, T. L.; Hediger, M. A.; Ganapathy, V.; Leibach, F. H.:

[87556] Human intestinal H(+)/peptide cotransporter: cloning, functional expression,

[87557] and chromosomal localization. J. Biol. Chem. 270: 6456–6463, 1995.

[87558] 15402.

[87559] Perry, S.; Harries, H.; Scholfield, C.; Lock, T.; King, L.; Gibson,

[87560] G.; Goldfarb, P.: Molecular cloning and expression of a cDNA for

[87561] human kidney cysteine conjugate beta-lyase. FEBS Lett. 360: 277–280,

[87562] 1995.

[87563] 15403.

[87564] Kaul, S. C.; Wadhwa, R.; Matsuda, Y.; Hensler, P. J.;

Pereira-Smith,

[87565] O. M.; Komatsu, Y.; Mitsui, Y.: Mouse and human chromosomal assignments

[87566] of mortalin, a novel member of the murine hsp70 family of proteins. FEBS

[87567] Lett. 361: 269-272, 1995.

[87568] 15404.

[87569] Heiber, M.; Docherty, J. M.; Shah, G.; Nguyen, T.; Cheng, R.; Heng,

[87570] H. H. Q.; Marchese, A.; Tsui, L.-C.; Shi, X.; George, S. R.; O'Dowd,

[87571] B. F.: Isolation of three novel human genes encoding G protein-coupled

[87572] receptors. DNA Cell Biol. 14: 25-35, 1995.

[87573] 15405.

[87574] Mahadevan, M. S.; Baird, S.; Bailly, J. E.; Shutler, G. G.; Sabourin,

[87575] L. A.; Tsilfidis, C.; Neville, C. E.; Narang, M.; Korneluk, R. G.

[87576] : Isolation of a novel G protein-coupled receptor (GPR4) localized

[87577] to chromosome 19q13.3. Genomics 30: 84-88, 1995.

- [87578] 15406.
- [87579] Ye, R. D.; Prossnitz, E. R.; Zou, A.; Cochrane, C. G.: Characterization
- [87580] of a human cDNA that encodes a functional receptor for platelet activating
- [87581] factor. *Biochem. Biophys. Res. Commun.* 180: 105–111, 1991.
- [87582] 15407.
- [87583] Hossain, A.; Saunders, G. F.: The human sex-determining gene
- [87584] SRY is a direct target of WT1. *J. Biol. Chem.* 276: 16817–16823,
- [87585] 2001.
- [87586] 15408.
- [87587] Iida, T.; Nakahori, Y.; Komaki, R.; Mori, E.; Hayashi, N.; Tsutsumi,
- [87588] O.; Taketani, Y.; Nakagome, Y.: A novel missense mutation in the
- [87589] HMG box of the SRY gene in a patient with XY sex reversal. *Hum. Molec.*
- [87590] *Genet.* 3: 1437–1438, 1994.

- [87591] 15409.
- [87592] Jager, R. J.; Harley, V. R.; Pfeiffer, R. A.; Goodfellow, P. N.;
- [87593] Scherer, G.: A familial mutation in the testis-determining
gene SRY
- [87594] shared by both sexes. Hum. Genet. 90: 350–355, 1992.
- [87595] 15410.
- [87596] Kawakura, K.; Miyake, Y.-I.; Murakami, R.-K.; Kondoh, S.;
- Hirata,
- [87597] T.-I.; Kaneda, Y.: Deletion of the SRY region on the Y
chromosome
- [87598] detected in bovine gonadal hypoplasia (XY female) by PCR.
Cytogenet.
- [87599] Cell Genet. 72: 183–184, 1996.
- [87600] 15411.
- [87601] Lahr, G.; Maxson, S. C.; Mayer, A.; Just, W.; Pilgrim, C.;
- Reisert,
- [87602] I.: Transcription of the Y chromosomal gene, Sry, in adult
mouse
- [87603] brain. Molec. Brain Res. 33: 179–182, 1995.
- [87604] 15412.

- [87605] Li, B.; Zhang, W.; Chan, G.; Jancso-Radek, A.; Liu, S.; Weiss,
Weiss,
- [87606] M. A.: Human sex reversal due to impaired nuclear localization of
- [87607] SRY: a clinical correlation. J. Biol. Chem. 276:
46480-46484, 2001.
- [87608] 15413.
- [87609] Lovell-Badge, R.; Robertson, E.: XY female mice resulting
from
- [87610] a heritable mutation in the primary testis-determining
gene, Tdy. Development 109:
- [87611] 635-646, 1990.
- [87612] 15414.
- [87613] Margarit, E.; Coll, M. D.; Oliva, R.; Gomez, D.; Soler, A.;
Ballesta,
- [87614] F.: SRY gene transferred to the long arm of the X chromosome in a
- [87615] Y-positive XX true hermaphrodite. Am. J. Med. Genet. 90:
25-28,
- [87616] 2000.
- [87617] 15415.

- [87618] Mittwoch, U.: Sex determination and sex reversal: genotype, phenotype,
[87619] dogma and semantics. Hum. Genet. 89: 467–479, 1992.
[87620] 15416.
- [87621] Muller, J.; Schwartz, M.; Skakkebaek, N. E.: Analysis of the
[87622] sex-determining region of the Y-chromosome (SRY) in
sex reversed patients:
[87623] point-mutation in SRY causing sex-reversion in a 46,XY
female. J.
[87624] Clin. Endocr. Metab. 75: 331–333, 1992.
[87625] 15417.
- [87626] Peters, R.; King, C.-Y.; Ukiyama, E.; Falsafi, S.; Donahoe, P.
[87627] K.; Weiss, M. A.: An SRY mutation causing human sex re-
versal resolves
[87628] a general mechanism of structure-specific DNA recogni-
tion: application
[87629] to the four-way DNA junction. Biochemistry 34:
4569–4576, 1995.
[87630] 15418.
- [87631] Schaffler, A.; Barth, N.; Winkler, K.; Zietz, B.; Rummele, P.;

- [87632] Knuchel, R.; Scholmerich, J.; Palitzsch, K.-D.: Identification of
- [87633] a new missense mutation (gly95glu) in a highly conserved codon within
- [87634] the high-mobility group box of the sex-determining region Y gene:
- [87635] report on a 46,XY female with gonadal dysgenesis and yolk-sac tumor. J.
- [87636] Clin. Endocr. Metab. 85: 2287-2292, 2000.
- [87637] 15419.
- [87638] Schmitt-Ney, M.; Thiele, H.; Kaltwasser, P.; Bardoni, B.; Cisternino,
- [87639] M.; Scherer, G.: Two novel SRY missense mutations reducing DNA binding
- [87640] identified in XY females and their mosaic fathers. Am. J. Hum. Genet. 56:
- [87641] 862-869, 1995.
- [87642] 15420.
- [87643] Sinclair, A. H.; Berta, P.; Palmer, M. S.; Hawkins, R.; Griffiths,
- [87644] B. L.; Smith, M.; Foster, J. W.; Frischauf, A.-M.; Lovell-Badge, R.;

- [87645] Goodfellow, P. N.: A gene from the human sex-determining region encodes
- [87646] a protein with homology to a conserved DNA-binding motif.. Nature 346:
- [87647] 240-245, 1990.
- [87648] 15421.
- [87649] Su, H.; Lau, Y.-F. C.: Identification of the transcriptional

- [87650] unit, structural organization, and promoter sequence of the human
- [87651] sex-determining region Y (SRY) gene, using a reverse genetic approach. Am.
- [87652] J. Hum. Genet. 52: 24–38, 1993.
- [87653] 15422.
- [87654] Tucker, P. K.; Lundrigan, B. L.: Rapid evolution of the sex determining
- [87655] locus in Old World mice and rats. Nature 364: 715–717, 1993.
- [87656] 15423.
- [87657] Uehara, S.; Funato, T.; Yaegashi, N.; Suzuki, H.; Sato, J.; Sasaki,
- [87658] T.; Yajima, A.: SRY mutation and tumor formation on the gonads of
- [87659] XY pure gonadal dysgenesis patients. Cancer Genet. Cytogenet. 113:
- [87660] 78–84, 1999.
- [87661] 15424.
- [87662] Uehara, S.; Hashiyada, M.; Sato, K.; Nata, M.; Funato, T.; Okamura,

- [87663] K.: Complete XY gonadal dysgenesis and aspects of the SRY genotype
- [87664] and gonadal tumor formation. J. Hum. Genet. 47: 279–284, 2002.
- [87665] 15425.
- [87666] Veitia, R.; Ion, A.; Barbaux, S.; Jobling, M. A.; Souleyreau,
- [87667] N.; Ennis, K.; Ostrer, H.; Tosi, M.; Meo, T.; Chibani, J.; Fellous,
- [87668] M.; McElreavey, K.: Mutations and sequence variants in the testis-determining
- [87669] region of the Y chromosome in individuals with a 46,XY female phenotype. Hum.
- [87670] Genet. 99: 648–652, 1997.
- [87671] 15426.
- [87672] Werner, M. H.; Huth, J. R.; Gronenborn, A. M.; Clore, G. M.:
- [87673] Molecular basis of human 46X,Y (sic) sex reversal revealed from the
- [87674] three-dimensional solution structure of the human SRY-DNA complex. Cell 81:
- [87675] 705–714, 1995.
- [87676] 15427.

- [87677] Whitfield, L. S.; Lovell-Badge, R.; Goodfellow, P. N.: Rapid
[87678] sequence evolution of the mammalian sex-determining
gene SRY. *Nature* 364:
[87679] 713–715, 1993.
[87680] 15428.
[87681] Zeng, Y.; Ren, Z.; Zhang, M.; Huang, Y.; Zeng, F.; Huang,
S.:
[87682] A new de novo mutation (A113T) in HMG box of the SRY
gene leads to
[87683] XY gonadal dysgenesis. *J. Med. Genet.* 30: 655–657,
1993.
[87684] 15429.
[87685] Zenteno, J. C.; Lopez, M.; Vera, C.; Mendez, J. P.; Kofman-
Alfaro,
[87686] S.: Two SRY-negative XX male brothers without genital
ambiguity. *Hum.*
[87687] *Genet.* 100: 606–610, 1997.
[87688] 15430.
[87689] Henkemeyer, M.; Orioli, D.; Henderson, J. T.; Saxton, T.
M.; Roder,
[87690] J.; Pawson, T.; Klein, R.: Nuk controls pathfinding of com-

missural

- [87691] axons in the mammalian central nervous system. Cell 86: 35–46, 1996.
- [87692] 15431.
- [87693] Ikegaki, N.; Tang, X. X.; Liu, X.-G.; Biegel, J. A.; Allen, C.;
- [87694] Yoshioka, A.; Sulman, E. P.; Brodeur, G. M.; Pleasure, D. E.: Molecular
- [87695] characterization and chromosomal localization of DRT (EPHT3): a developmentally
- [87696] regulated human protein-tyrosine kinase gene of the EPH family. Hum.
- [87697] Molec. Genet. 4: 2033–2045, 1995.
- [87698] 15432.
- [87699] Saito, T.; Seki, N.; Matsuda, Y.; Kitahara, M.; Murata, M.; Kanda,
- [87700] N.; Nomura, N.; Yamamoto, T.; Hori, T.: Identification of the human
- [87701] ERK gene as a putative receptor tyrosine kinase and its chromosomal
- [87702] localization to 1p36.1: a comparative mapping of human, mouse, and
- [87703] rat chromosomes. Genomics 26: 382–384, 1995.

- [87704] 15433.
- [87705] Wybenga-Groot, L. E.; Baskin, B.; Ong, S. H.; Tong, J.; Pawson,
- [87706] T.; Sicheri, F.: Structural basis for autoinhibition of the EphB2
- [87707] receptor tyrosine kinase by the unphosphorylated jux-
- tamembrane region. Cell 106:
- [87708] 745-757, 2001.
- [87709] 15434.
- [87710] Elpeleg, O. N.; Hammerman, C.; Saada, A.; Shaag, A.; Golzand, E.;
- [87711] Hochner-Celnikier, D.; Berger, I.; Nadjari, M.: Antenatal presentation
- [87712] of carnitine palmitoyltransferase II deficiency. Am. J. Med. Genet. 102:
- [87713] 183-187, 2001.
- [87714] 15435.
- [87715] Finocchiaro, G.; Taroni, F.; Rocchi, M.; Liras Martin, A.; Colombo,
- [87716] I.; Tarelli, G. T.; DiDonato, S.: cDNA cloning, sequence analysis,

- [87717] and chromosomal localization of the gene for human carnitine palmitoyltransferase. Proc.
- [87718] Nat. Acad. Sci. 88: 661–665, 1991.
- [87719] 15436.
- [87720] Haap, M.; Thamer, C.; Machann, J.; Tschritter, O.; Loblein, K.;
- [87721] Kellerer, M.; Schick, F.; Jacob, S.; Haring, H.–U.; Stumvoll, M.:
- [87722] Metabolic characterization of a woman homozygous for the ser113leu
- [87723] missense mutation in carnitine palmitoyl transferase II. J. Clin.
- [87724] Endocr. Metab. 87: 2139–2143, 2002.
- [87725] 15437.
- [87726] Handig, I.; Dams, E.; Taroni, F.; Van Laere, S.; de Barse, T.;
- [87727] Willems, P. J.: Inheritance of the S113L mutation within an inbred
- [87728] family with carnitine palmitoyltransferase enzyme deficiency. Hum.
- [87729] Genet. 97: 291–293, 1996.
- [87730] 15438.

- [87731] Martin, M. A.; Rubio, J. C.; De Bustos, F.; Del Hoyo, P.; Campos,
- [87732] Y.; Garcia, A.; Bornstein, B.; Cabello, A.; Arenas, J.: Molecular
- [87733] analysis in Spanish patients with muscle carnitine palmitoyltransferase
- [87734] deficiency. *Muscle Nerve* 22: 941–943, 1999.
- [87735] 15439.
- [87736] Minoletti, F.; Colombo, I.; Liras Martin, A.; DiDonato, S.; Taroni,
- [87737] F.; Finocchiaro, G.; Pandolfo, M.: Localization of the human gene
- [87738] for carnitine palmitoyltransferase to 1p13–p11 by nonradioactive in
- [87739] situ hybridization. *Genomics* 13: 1372–1374, 1992.
- [87740] 15440.
- [87741] Montermini, L.; Wang, H.; Verderio, E.; Taroni, F.; DiDonato,
- [87742] S.; Finocchiaro, G: Identification of 5–prime regulatory regions
- [87743] of the human carnitine palmitoyltransferase II gene. *Biochim. Biophys.*

- [87744] Acta 1219: 237–240, 1994.
- [87745] 15441.
- [87746] Taroni, F.; Verderio, E.; Fiorucci, S.; Cavadini, P.; Finoc-
chiaro,
- [87747] G.; Uziel, G.; Lamantea, E.; Gellera, C.; DiDonato, S.:
Molecular
- [87748] characterization of inherited carnitine palmitoyltrans-
ferase II deficiency. Proc.
- [87749] Nat. Acad. Sci. 89: 8429–8433, 1992.
- [87750] 15442.
- [87751] Verderio, E.; Cavadini, P.; Montermini, L.; Wang, H.;
Lamantea,
- [87752] E.; Finocchiaro, G.; DiDonato, S.; Gellera, C.; Taroni, F.:
Carnitine
- [87753] palmitoyltransferase II deficiency: structure of the gene
and characterization
- [87754] of two novel disease-causing mutations. Hum. Molec.
Genet. 4: 19–29,
- [87755] 1995.
- [87756] 15443.
- [87757] Vladutiu, G. D.; Bennett, M. J.; Smail, D.; Wong, L.-J.; Tag-

gart,

[87758] R. T.; Lindsley, H. B.: A variable myopathy associated with heterozygosity

[87759] for the R503C mutation in the carnitine palmitoyltransferase II gene. Molec.

[87760] Genet. Metab. 70: 134-141, 2000.

[87761] 15444.

[87762] Vladutiu, G. D.; Taggart, R. T.; Smail, D.; Lindsley, H. B.; Hogan,

[87763] K.: A carnitine palmitoyl transferase II (CPT2) arg503cys mutation

[87764] confers malignant hyperthermia and variable myopathy. (Abstract) Am.

[87765] J. Hum. Genet. 63 (suppl.): A5 only, 1998.

[87766] 15445.

[87767] Yamamoto, S.; Abe, H.; Kohgo, T.; Ogawa, A.; Ohtake, A.; Hayashibe,

[87768] H.; Sakuraba, H.; Suzuki, Y.; Aramaki, S.; Takayanagi, M.; Hasegawa,

[87769] S.; Niimi, H.: Two novel gene mutations (glu174-to-lys, phe383-to-tyr)

[87770] causing the 'hepatic' form of carnitine palmitoyltrans-

ferase II deficiency. Hum.

[87771] Genet. 98: 116–118, 1996.

[87772] 15446.

[87773] Johnson, D. F.; Hamon, M.; Fischel-Ghodsian, N.: Characterization

[87774] of the human mitochondrial ribosomal S12 gene. Genomics 52: 363–368,

[87775] 1998.

[87776] 15447.

[87777] Shah, Z. H.; Migliosi, V.; Miller, S. C. M.; Wang, A.; Friedman,

[87778] T. B.; Jacobs, H. T.: Chromosomal locations of three human nuclear

[87779] genes (RPSM12, TUFM, and AFG3L1) specifying putative components of

[87780] the mitochondrial gene expression apparatus. Genomics 48: 384–388,

[87781] 1998.

[87782] 15448.

[87783] Zhu, K.; Baudhuin, L. M.; Hong, G.; Williams, F. S.; Cristina,

[87784] K. L.; Kabarowski, J. H. S.; Witte, O. N.; Xu, Y.: Sphingo-

sylphosphorylcholine

[87785] and lysophosphatidylcholine are ligands for the G protein-coupled

[87786] receptor GPR4. J. Biol. Chem. 276: 41325–41335, 2001.

[87787] 15449.

[87788] Vinik, B. S.; Kay, E. S.; Fiedorek, F. T., Jr.: Mapping of the

[87789] MEK kinase gene (Mekk) to mouse chromosome 13 and human chromosome

[87790] 5. Mammalian Genome 6: 782–783, 1995.

[87791] 15450.

[87792] Yujiri, T.; Sather, S.; Fanger, G. R.; Johnson, G. L.: Role of

[87793] MEKK1 in cell survival and activation of JNK and ERK pathways defined

[87794] by targeted gene disruption. Science 282: 1911–1914, 1998.

[87795] 15451.

[87796] Arriza, J. L.; Weinberger, C.; Cerelli, G.; Glaser, T. M.; Handelin,

[87797] B. L.; Housman, D. E.; Evans, R. M.: Cloning of human mineralocorticoid

[87798] receptor complementary DNA: structural and functional

kinship with

[87799] the glucocorticoid receptor. *Science* 237: 268–275, 1987.

[87800] 15452.

[87801] Berger, S.; Bleich, M.; Schmid, W.; Cole, T. J.; Peters, J.;
Watanabe,

[87802] H.; Kriz, W.; Warth, R.; Greger, R.; Schutz, G.: Mineralocor-
ticoid

[87803] receptor knockout mice: pathophysiology of Na+
metabolism. *Proc.*

[87804] *Nat. Acad. Sci.* 95: 9424–9429, 1998.

[87805] 15453.

[87806] Fan, Y.-S.; Eddy, R. L.; Byers, M. G.; Haley, L. L.; Henry, W.

[87807] M.; Nowak, N. J.; Shows, T. B.: The human mineralocorti-
coid receptor

[87808] gene (MLR) is located on chromosome 4 at q31.2. *Cyto-*
genet. Cell Genet. 52:

[87809] 83–84, 1989.

[87810] 15454.

[87811] Geller, D. S.; Farhi, A.; Pinkerton, N.; Fradley, M.; Moritz,
M.;

[87812] Spitzer, A.; Meinke, G.; Tsai, F. T. F.; Sigler, P. B.; Lifton, R.

- [87813] P.: Activating mineralocorticoid receptor mutation in hypertension
- [87814] exacerbated by pregnancy. Science 289: 119–123, 2000.
- [87815] 15455.
- [87816] Geller, D. S.; Rodriguez–Soriano, J.; Vallo Boado, A.; Schifter,
- [87817] S.; Bayer, M.; Chang, S. S.; Lifton, R. P.: Mutations in the mineralocorticoid
- [87818] receptor gene cause autosomal dominant pseudohypoaldosteronism type
- [87819] I. Nature Genet. 19: 279–281, 1998.
- [87820] 15456.
- [87821] Hellal–Levy, C.; Fagart, J.; Souque, A.; Wurtz, J.–M.; Moras, D.;
- [87822] Rafestin–Oblin, M.–E.: Crucial role of the H11–H12 loop in stabilizing
- [87823] the active conformation of the human mineralocorticoid receptor. Molec.
- [87824] Endocr. 14: 1210–1221, 2000.
- [87825] 15457.
- [87826] Le Menuet, D.; Isnard, R.; Bichara, M.; Viengchareun, S.;

Muffat-Joly,

- [87827] M.; Walker, F.; Zennaro, M.-C.; Lombes, M.: Alteration of cardiac
- [87828] and renal functions in transgenic mice overexpressing human mineralocorticoid
- [87829] receptor. J. Biol. Chem. 276: 38911–38920, 2001.
- [87830] 15458.
- [87831] Morrison, N.; Harrap, S. B.; Arriza, J. L.; Boyd, E.; Connor, J.
- [87832] M.: Regional chromosomal assignment of the human mineralocorticoid
- [87833] receptor gene to 4q31.1. Hum. Genet. 85: 130–132, 1990.
- [87834] 15459.
- [87835] Morrison, N.; Harrap, S. B.; Arriza, J. L.; Boyd, E.; Connor,
- [87836] J. M.: Regional chromosomal assignment of the human mineralocorticoid
- [87837] receptor gene to 4q31.1. (Abstract) Cytogenet. Cell Genet. 51: 1048,
- [87838] 1989.
- [87839] 15460.
- [87840] Bristow, J.; Tee, M. K.; Gitelman, S. E.; Mellon, S. H.; Miller,

[87841] W. L.: Tenascin-X: a novel extracellular matrix protein encoded by
[87842] the human XB gene overlapping P450c21B. J. Cell Biol.
122: 265–278,
[87843] 1993.
[87844] 15461.
[87845] Jay, P.; Goze, C.; Marsollier, C.; Taviaux, S.; Hardelin, J.-P.;
[87846] Koopman, P.; Berta, P.: The human SOX11 gene: cloning,
chromosomal
[87847] assignment and tissue expression. Genomics 29:
541–545, 1995.
[87848] 15462.
[87849] Laudet, V.; Stehelin, D.; Clevers, H.: Ancestry and diversity
[87850] of the HMG box superfamily. Nucleic Acids Res. 21:
2493–2501, 1993.
[87851] 15463.
[87852] Weigt, C.; Gaertner, A.; Wegner, A.; Korte, H.; Meyer, H. E.:
[87853] Occurrence of an actin-inserting domain in tensin. J.
Molec. Biol. 227:
[87854] 593–595, 1992.

- [87855] 15464.
- [87856] Charest, A.; Wagner, J.; Muise, E. S.; Heng, H. H. Q.; Tremblay,
- [87857] M. L.: Structure of the murine MPTP-PEST gene: genomic organization
- [87858] and chromosomal mapping. *Genomics* 28: 501-507, 1995.
- [87859] 15465.
- [87860] Takekawa, M.; Itoh, F.; Hinoda, Y.; Adachi, M.; Ariyama, T.; Inazawa,
- [87861] J.; Imai, K.; Yachi, A.: Chromosomal localization of the protein
- [87862] tyrosine phosphatase G1 gene and characterization of the aberrant
- [87863] transcripts in human colon cancer cells. *FEBS Lett.* 339: 222-228,
- [87864] 1994.
- [87865] 15466.
- [87866] Takekawa, M.; Itoh, F.; Hinoda, Y.; Arimura, Y.; Toyota, M.; Sekiya,
- [87867] M.; Adachi, M.; Imai, K.; Yachi, A.: Cloning and characterization

- [87868] of a human cDNA encoding a novel putative cytoplasmic protein-tyrosine-phosphatase. *Biochem.*
- [87869] *Biophys. Res. Commun.* 189: 1223-1230, 1992.
- [87870] 15467.
- [87871] Yang, Q.; Co, D.; Sommercorn, J.; Tonks, N. K.: Cloning and expression
- [87872] of PTP-PEST: a novel, human, nontransmembrane protein tyrosine phosphatase. *J.*
- [87873] *Biol. Chem.* 268: 6622-6628, 1993.
- [87874] 15468.
- [87875] Coopman, P. J. P.; Do, M. T. H.; Barth, M.; Bowden, E. T.; Hayes,
- [87876] A. J.; Basyuk, E.; Blancato, J. K.; Vezza, P. R.; McLeskey, S. W.;
- [87877] Mangeat, P. H.; Mueller, S. C.: The Syk tyrosine kinase suppresses
- [87878] malignant growth of human breast cancer cells. *Nature* 406: 742-747,
- [87879] 2000.
- [87880] 15469.
- [87881] Inatome, R.; Yanagi, S.; Takano, T.; Yamamura, H.: A criti-

cal

[87882] role for Syk in endothelial cell proliferation and migration.

Biochem.

[87883] Biophys. Res. Commun. 286: 195–199, 2001.

[87884] 15470.

[87885] Mocsai, A.; Zhou, M.; Meng, F.; Tybulewicz, V. L.; Lowell,

C. A.

[87886] : Syk is required for integrin signaling in neutrophils. Im-

munity 16:

[87887] 547–558, 2002.

[87888] 15471.

[87889] Muller, B.; Cooper, L.; Terhorst, C.: Molecular cloning of

the

[87890] human homologue to the pig protein-tyrosine kinase syk.

Immunogenetics 39:

[87891] 359–362, 1994.

[87892] 15472.

[87893] Taniguchi, T.; Kobayashi, T.; Kondo, J.; Takahashi, K.;

Nakamura,

[87894] H.; Suzuki, J.; Nagai, K.; Yamada, T.; Nakamura, S.; Yama-

mura, H.

- [87895] : Molecular cloning of a porcine gene syk that encodes a 72-kDa protein-tyrosine
- [87896] kinase showing high susceptibility to proteolysis. J. Biol. Chem. 266:
- [87897] 15790–15796, 1991.
- [87898] 15473.
- [87899] Zioncheck, T. F.; Harrison, M. L.; Isaacson, C. C.; Geahlen, R.
- [87900] L.: Generation of an active protein-tyrosine kinase from lymphocytes
- [87901] by proteolysis. J. Biol. Chem. 263: 19195–19202, 1988.
- [87902] 15474.
- [87903] Satre, M. A.; Zgombic-Knight, M.; Duester, G.: The complete structure
- [87904] of human class IV alcohol dehydrogenase (retinol dehydrogenase) determined
- [87905] from the ADH7 gene. J. Biol. Chem. 269: 15606–15612, 1994.
- [87906] 15475.
- [87907] Yokoyama, H.; Baraona, E.; Lieber, C. S.: Molecular cloning and

- [87908] chromosomal localization of the ADH7 gene encoding human class IV
- [87909] (sigma) ADH. Genomics 31: 243–245, 1996.
- [87910] 15476.
- [87911] Zgombic-Knight, M.; Foglio, M. H.; Duester, G.: Genomic structure
- [87912] and expression of the ADH7 gene encoding human class IV alcohol dehydrogenase,
- [87913] the form most efficient for retinol metabolism in vitro. J. Biol. Chem. 270: 4305–4311, 1995.
- [87914] 15477.
- [87915] Bashir, M. M.; Abrams, W. R.; Tucker, T.; Sellinger, B.; Budarf, M.; Emanuel, B.; Rosenbloom, J.: Molecular cloning and characterization
- [87916] of the bovine and human tuftelin genes. Connect. Tissue Res. 39: 13–24, 1998.
- [87917] 15478.
- [87918] Deutsch, D.: Structure and function of enamel gene prod-

ucts. Anat.

[87922] Rec. 224: 189–210, 1989.

[87923] 15479.

[87924] Beguin, P.; Nagashima, K.; Gonoï, T.; Shibasaki, T.; Takahashi,

[87925] K.; Kashima, Y.; Ozaki, N.; Geering, K.; Iwanaga, T.; Seino, S.:

[87926] Regulation of Ca(2+) channel expression at the cell surface by the

[87927] small G-protein kir/Gem. Nature 411: 701–706, 2001.

[87928] 15480.

[87929] Maguire, J.; Santoro, T.; Jensen, P.; Siebenlist, U.; Yewdell,

[87930] J.; Kelly, K.: GEM: an induced, immediate early protein belonging

[87931] to the Ras family. Science 265: 241–244, 1994.

[87932] 15481.

[87933] Santoro, T.; Maguire, J.; McBride, O. W.; Avraham, K. B.; Copeland,

[87934] N. G.; Jenkins, N. A.; Kelly, K.: Chromosomal organization and transcriptional

[87935] regulation of human GEM and localization of the human

and mouse Gem

[87936] loci encoding an inducible Ras-like protein. Genomics 30:
558-564,

[87937] 1995.

[87938] 15482.

[87939] Inoue, I.; Taniuchi, I.; Kitamura, D.; Jenkins, N. A.; Gilbert,

[87940] D. J.; Copeland, N. G.; Watanabe, T.: Characteristics of the
mouse

[87941] genomic histamine H1 receptor gene. Genomics 36:
178-181, 1996.

[87942] 15483.

[87943] Le Coniat, M.; Traiffort, E.; Ruat, M.; Arrang, J.-M.; Berger,

[87944] R.: Chromosomal localization of the human histamine
H1-receptor gene. Hum.

[87945] Genet. 94: 186-188, 1994.

[87946] 15484.

[87947] Ma, R. Z.; Gao, J.; Meeker, N. D.; Fillmore, P. D.; Tung, K.
S.

[87948] K.; Watanabe, T.; Zachary, J. F.; Offner, H.; Blankenhorn, E.
P.;

[87949] Teuscher, C.: Identification of Bphs, an autoimmune dis-

ease locus,

[87950] as histamine receptor H-1. Science 297: 620-623, 2002.

[87951] 15485.

[87952] Yamashita, M.; Fukui, H.; Sugama, K.; Horio, Y.; Ito, S.; Mizuguchi,

[87953] H.; Wada, H.: Expression cloning of a cDNA encoding the bovine histamine

[87954] H1 receptor. Proc. Nat. Acad. Sci. 88: 11515-11519, 1991.

[87955] 15486.

[87956] Gaudino, G.; Follenzi, A.; Naldini, L.; Collesi, C.; Santoro, M.;

[87957] Gallo, K. A.; Godowski, P. J.; Comoglio, P. M.: RON is a heterodimeric

[87958] tyrosine kinase receptor activated by the HGF homologue MSP. EMBO

[87959] J. 13: 3524-3532, 1994.

[87960] 15487.

[87961] Muraoka, R. S.; Sun, W. Y.; Colbert, M. C.; Waltz, S. E.; Witte,

[87962] D. P.; Degen, J. L.; Degen, S. J. F.: The Ron/STK receptor

tyrosine

[87963] kinase is essential for peri-implantation development in the mouse. J.

[87964] Clin. Invest. 103: 1277-1285, 1999.

[87965] 15488.

[87966] Ronsin, C.; Muscatelli, F.; Mattei, M.-G.; Breathnach, R.: A novel

[87967] putative receptor protein tyrosine kinase of the met family. Oncogene 8:

[87968] 1195-1202, 1993.

[87969] 15489.

[87970] Santoro, M. M.; Collesi, C.; Grisendi, S.; Gaudino, G.; Comoglio,

[87971] P. M.: Constitutive activation of the RON gene promotes invasive

[87972] growth but not transformation. Molec. Cell. Biol. 16: 7072-7083,

[87973] 1996.

[87974] 15490.

[87975] Wang, M.-H.; Ronsin, C.; Gesnel, M.-C.; Coupey, L.; Skeel, A.;

- [87976] Leonard, E. J.; Breathnach, R.: Identification of the RON gene product
- [87977] as the receptor for the human macrophage stimulating protein. Science 266:
- [87978] 117–119, 1994.
- [87979] 15491.
- [87980] Bauer, S.; Groh, V.; Wu, J.; Steinle, A.; Phillips, J. H.; Lanier, L. L.; Spies, T.: Activation of NK cells and T cells by NKG2D, a
- [87982] receptor for stress-inducible MICA. Science 285: 727–729, 1999.
- [87983] 15492.
- [87984] Diefenbach, A.; Jensen, E. R.; Jamieson, A. M.; Raulet, D. H.:
- [87985] Rae1 and H60 ligands of the NKG2D receptor stimulate tumour immunity. Nature 413:
- [87986] 165–171, 2001.
- [87987] 15493.
- [87988] Girardi, M.; Oppenheim, D. E.; Steele, C. R.; Lewis, J. M.; Glusac,
- [87989] E.; Filler, R.; Hobby, P.; Sutton, B.; Tigelaar, R. E.; Hayday,

A.

[87990] C.: Regulation of cutaneous malignancy by gamma-delta T cells. Science 294:

[87991] 605–609, 2001.

[87992] 15494.

[87993] Groh, V.; Rhinehart, R.; Randolph-Habecker, J.; Topp, M. S.; Riddell,

[87994] S. R.; Spies, T.: Costimulation of CD8-alpha-beta T cells by NKG2D

[87995] via engagement by MIC induced on virus-infected cells. Nature Immun. 2:

[87996] 255–260, 2001.

[87997] 15495.

[87998] Groh, V.; Steinle, A.; Bauer, S.; Spies, T.: Recognition of stress-induced

[87999] MHC molecules by intestinal epithelial gamma-delta T cells. Science 279:

[88000] 1737–1740, 1998.

[88001] 15496.

[88002] Groh, V.; Wu, J.; Yee, C.; Spies, T.: Tumour-derived soluble MIC

[88003] ligands impair expression of NKG2D and T-cell activation.

Nature 419:

[88004] 734–738, 2002.

[88005] 15497.

[88006] Li, P.; Morris, D. L.; Willcox, B. E.; Steinle, A.; Spies, T.;

[88007] Strong, R. K.: Complex structure of the activating immunoreceptor

[88008] NKG2D and its MHC class I-like ligand MICA. Nature Immun. 2: 443–451,

[88009] 2001.

[88010] 15498.

[88011] Nakashima, T.; Sekiguchi, T.; Kuraoka, A.; Fukushima, K.; Shibata,

[88012] Y.; Komiyama, S.; Nishimoto, T.: Molecular cloning of a human cDNA

[88013] encoding a novel protein, DAD1, whose defect causes apoptotic cell

[88014] death in hamster BHK21 cells. Molec. Cell. Biol. 13: 6367–6374,

[88015] 1993.

[88016] 15499.

[88017] Ylug, I. G.; See, C. G.; Fisher, E. M. C.: The DAD1 protein,
whose

[88018] defect causes apoptotic cell death, maps to human chro-
mosome 14. Genomics 26:

[88019] 433–435, 1995. Note: Erratum: Genomics: 28: 129 only,
1995.

[88020] 15500.

[88021] Yulug, I. G.; See, C. G.; Fisher, E. M. C.: The DAD1 protein,

[88022] whose defect causes apoptotic cell death, maps to human
chromosome

[88023] 14. Genomics 26: 433–435, 1995.

[88024] 15501.

[88025] Jepsen, K. J.; Wu, F.; Peragallo, J. H.; Paul, J.; Roberts, L.;

[88026] Ezura, Y.; Oldberg, A.; Birk, D. E.; Chakravarti, S.: A syn-
drome

[88027] of joint laxity and impaired tendon integrity in lumican-
and fibromodulin-deficient

[88028] mice. J. Biol. Chem. 277: 35532–35540, 2002.

[88029] 15502.

[88030] Sztrolovics, R.; Chen, X.-N.; Grover, J.; Roughley, P. J.; Ko-
renberg,

- [88031] J. R.: Localization of the human fibromodulin gene (FMOD) to chromosome
- [88032] 1q32 and completion of the cDNA sequence. *Genomics* 23: 715–717,
- [88033] 1994.
- [88034] 15503.
- [88035] Dalton, S.; Treisman, R.: Characterization of SAP–1, a protein
- [88036] recruited by serum response factor to the c–fos serum response element. *Cell* 68:
- [88037] 597–612, 1992.
- [88038] 15504.
- [88039] Mo, Y.; Vaessen, B.; Johnston, K.; Marmorstein, R.: Structures
- [88040] of SAP–1 bound to DNA targets from the E74 and c–fos promoters: insights
- [88041] into DNA sequence discrimination by Ets proteins. *Molec. Cell* 2:
- [88042] 201–212, 1998.
- [88043] 15505.
- [88044] Shipley, J.; Sheer, D.; Dalton, S.; Treisman, R.; Patel, K.:

Mapping

- [88045] of the human SAP1 (SRF accessory protein 1) gene and SAP2, a gene
- [88046] encoding a related protein, to chromosomal bands 1q32 and 12q23, respectively. Genomics 23:
- [88047] 710–711, 1994.
- [88048] 15506.
- [88049] Ikeuchi, T.; Asaka, T.; Saito, M.; Tanaka, H.; Higuchi, S.; Tanaka,
- [88050] K.; Saida, K.; Uyama, E.; Mizusawa, H.; Fukuhara, N.; Nonaka, I.;
- [88051] Takamori, M.; Tsuji, S.: Gene locus for autosomal recessive distal
- [88052] myopathy with rimmed vacuoles maps to chromosome 9. Ann. Neurol. 41:
- [88053] 432–437, 1997.
- [88054] 15507.
- [88055] Erickson, J. D.; Varoqui, H.; Schafer, M. K.–H.; Modi, W.; Diebler,
- [88056] M.–F.; Weihe, E.; Rand, J.; Eiden, L. E.; Bonner, T. I.; Usdin, T.
- [88057] B.: Functional identification of a vesicular acetylcholine

transporter

[88058] and its expression from a 'cholinergic' gene locus. J. Biol. Chem. 269:

[88059] 21929–21932, 1994.

[88060] 15508.

[88061] Probst, M. R.; Beer, M.; Beer, D.; Jenö, P.; Meyer, U. A.; Gasser,

[88062] R.: Human liver arylacetamide deacetylase: molecular cloning of a

[88063] novel esterase involved in the metabolic activation of arylamine carcinogens

[88064] with high sequence similarity to hormone-sensitive lipase. J. Biol.

[88065] Chem. 269: 21650–21656, 1994.

[88066] 15509.

[88067] Yamazaki, K.; Kusano, K.; Tadano, K.; Tanaka, I.: Radiation hybrid

[88068] mapping of human arylacetamide deacetylase (AADAC) locus to chromosome 15510.

[88069] Worman, H. J.; Yuan, J.; Blobel, G.; Georgatos, S. D.: A lamin

[88070] B receptor in the nuclear envelope. Proc. Nat. Acad. Sci.

85: 8531–8534,

[88071] 1988.

[88072] 15511.

[88073] Tanahashi, H.; Tabira, T.: Alzheimer's disease-associated
presenilin

[88074] 2 interacts with DRAL, an LIM-domain protein. Hum.
Molec. Genet. 9:

[88075] 2281–2289, 2000.

[88076] 15512.

[88077] Erkinheimo, T.-L.; Saukkonen, K.; Narko, K.; Jalkanen, J.;
Ylikorkala,

[88078] O.; Ristimäki, A.: Expression of cyclooxygenase-2 and
prostanoid

[88079] receptors by human myometrium. J. Clin. Endocr. Metab.
85: 3468–3475,

[88080] 2000.

[88081] 15513.

[88082] Hla, T.; Neilson, K.: Human cyclooxygenase-2 cDNA. Proc.
Nat.

[88083] Acad. Sci. 89: 7384–7388, 1992.

[88084] 15514.

- [88085] Jones, D. A.; Carlton, D. P.; McIntyre, T. M.; Zimmerman, G. A.;
- [88086] Prescott, S. M.: Molecular cloning of human prostaglandin endoperoxide
- [88087] synthase type II and demonstration of expression in response to cytokines. J.
- [88088] Biol. Chem. 268: 9049–9054, 1993.
- [88089] 15515.
- [88090] Kraemer, S. A.; Meade, E. A.; DeWitt, D. L.: Prostaglandin endoperoxide
- [88091] synthase gene structure: identification of the transcriptional start
- [88092] site and 5-prime-flanking regulatory sequences. Arch. Biochem. Biophys. 293:
- [88093] 391–400, 1992.
- [88094] 15516.
- [88095] Lassus, P.; Wolff, H.; Andersson, S.: Cyclooxygenase-2 in human
- [88096] perinatal lung. Pediat. Res. 47: 602–605, 2000.
- [88097] 15517.
- [88098] Lim, H.; Paria, B. C.; Das, S. K.; Dinchuk, J. E.; Langenbach,

- [88099] R.; Trzaskos, J. M.; Dey, S. K.: Multiple female reproductive failures
- [88100] in cyclooxygenase 2-deficient mice. Cell 91: 197–208, 1997.
- [88101] 15518.
- [88102] Morrow, J. A.; Collie, I. T.; Dunbar, D. R.; Walker, G. B.; Shahid,
- [88103] M.; Hill, D. R.: Molecular cloning and functional expression of the
- [88104] human glycine transporter GlyT2 and chromosomal localisation of the
- [88105] gene in the human genome. FEBS Lett. 439: 334–340, 1998.
- [88106] 15519.
- [88107] Iacobazzi, V.; Palmieri, F.; Runswick, M. J.; Walker, J. E.: Sequences
- [88108] of the human and bovine genes for the mitochondrial 2-oxoglutarate
- [88109] carrier. DNA Seq. 3: 79–88, 1992.
- [88110] 15520.
- [88111] Piccininni, S.; Iacobazzi, V.; Lauria, G.; Rocchi, M.; Palmieri,

- [88112] F.: Assignment of the oxoglutarate carrier gene (SLC20A4) to human
- [88113] chromosome 17p13.3. Cytogenet. Cell Genet. 83: 256–257, 1998.
- [88114] 15521.
- [88115] Harvey, T. J.; Hooper, J. D.; Myers, S. A.; Stephenson, S.-A.;
- [88116] Ashworth, L. K.; Clements, J. A.: Tissue-specific expression patterns
- [88117] and fine mapping of the human kallikrein (KLK) locus on proximal 19q13.4. J.
- [88118] Biol. Chem. 275: 37397–37406, 2000.
- [88119] 15522.
- [88120] Chaudhry, F. A.; Reimer, R. J.; Krizaj, D.; Barber, D.; Storm-Mathisen,
- [88121] J.; Copenhagen, D. R.; Edwards, R. H.: Molecular analysis of system
- [88122] N suggests novel physiological roles in nitrogen metabolism and synaptic
- [88123] transmission. Cell 99: 769–780, 1999.
- [88124] 15523.

[88125] Yamada, K.; Nishida, K.; Hibi, M.; Hirano, T.; Matsuda, Y.:
Comparative

[88126] FISH mapping of Gab1 and Gab2 genes in human, mouse
and rat. Cytogenet.

[88127] Cell Genet. 94: 39–42, 2001.

[88128] 15524.

[88129] Afonina, E.; Stauber, R.; Pavlakis, G. N.: The human
poly(A)–binding

[88130] protein 1 shuttles between the nucleus and the cytoplasm.
J. Biol.

[88131] Chem. 273: 13015–13021, 1998.

[88132] 15525.

[88133] Deo, R. C.; Bonanno, J. B.; Sonenberg, N.; Burley, S. K.:
Recognition

[88134] of polyadenylate RNA by the poly(A)–binding protein. Cell
98: 835–845,

[88135] 1999.

[88136] 15526.

[88137] Gorlach, M.; Burd, C. G.; Dreyfuss, G.: The mRNA
poly(A)–binding

[88138] protein: localization, abundance, and RNA–binding speci–

ficity. Exp.

[88139] Cell Res. 211: 400–407, 1994.

[88140] 15527.

[88141] Grange, T.; Martins de Sa, C.; Oddos, J.; Pictet, R.: Human mRNA

[88142] polyadenylate binding protein: evolutionary conservation of a nucleic

[88143] acid binding motif. Nucleic Acids Res. 15: 4771–4787, 1987.

[88144] 15528.

[88145] Feral, C.; Guellaen, G.; Pawlak, A.: Human testis expresses a

[88146] specific poly(A)–binding protein. Nucleic Acids Res. 29: 1872–1883,

[88147] 2001.

[88148] 15529.

[88149] Danpure, C. J.; Purdue, P. E.; Fryer, P.; Griffiths, S.; Allsop,

[88150] J.; Lumb, M. J.; Guttridge, K. M.; Jennings, P. R.; Scheinman, J.

[88151] I.; Mauer, S. M.; Davidson, N. O.: Enzymological and mutational analysis

- [88152] of a complex primary hyperoxaluria type I phenotype involving alanine:glyoxylate
- [88153] aminotransferase peroxisome-to-mitochondrion mistargeting and intraperoxisomal
- [88154] aggregation. *Am. J. Hum. Genet.* 53: 417-432, 1993.
- [88155] 15530.
- [88156] Mori, M.; Oda, T.; Nishiyama, K.; Serikawa, T.; Yamada, J.; Ichiyama,
- [88157] A.: A single serine:pyruvate aminotransferase gene on rat chromosome
- [88158] 9q34-q36. *Genomics* 13: 686-689, 1992.
- [88159] 15531.
- [88160] Purdue, P. E.; Allsop, J.; Isaya, G.; Rosenberg, L. E.; Danpure,
- [88161] C. J.: Mistargeting of peroxisomal L-alanine:glyoxylate aminotransferase
- [88162] to mitochondria in primary hyperoxaluria patients depends upon activation
- [88163] of a cryptic mitochondrial targeting sequence by a point mutation. *Proc.*
- [88164] *Nat. Acad. Sci.* 88: 10900-10904, 1991.

- [88165] 15532.
- [88166] Purdue, P. E.; Lumb, M. J.; Allsop, J.; Minatogawa, Y.; Danpure,
- [88167] C. J.: A glycine-to-glutamate substitution abolishes alanine:glyoxylate
- [88168] aminotransferase catalytic activity in a subset of patients with primary
- [88169] hyperoxaluria type 1. *Genomics* 13: 215-218, 1992.
- [88170] 15533.
- [88171] Purdue, P. E.; Takada, Y.; Danpure, C. J.: Identification of
- [88172] mutations associated with peroxisome-to-mitochondrion mistargeting
- [88173] of alanine/glyoxylate aminotransferase in primary hyperoxaluria type
- [88174] 1. *J. Cell Biol.* 111: 2341-2351, 1990.
- [88175] 15534.
- [88176] Takada, Y.; Kaneko, N.; Esumi, H.; Purdue, P. E.; Danpure, C.
- [88177] J.: Human peroxisomal L-alanine:glyoxylate aminotransferase: evolutionary
- [88178] loss of a mitochondrial targeting signal by point mutation

of the

[88179] initiation codon. Biochem. J. 268: 517–520, 1990.

[88180] 15535.

[88181] von Schnakenburg, C.; Rumsby, G.: Primary hyperoxaluria
type

[88182] 1: a cluster of new mutations in exon 7 of the AGXT gene.
J. Med.

[88183] Genet. 34: 489–492, 1997.

[88184] 15536.

[88185] Kobayashi, H.; Hino, M.; Shimodahira, M.; Iwakura, T.;
Ishihara,

[88186] T.; Ikekubo, K.; Ogawa, Y.; Nakao, K.; Kurahachi, H.: Mis-
sense mutation

[88187] of TRPS1 in a family of tricho–rhino–phalangeal syndrome
type III. Am.

[88188] J. Med. Genet. 107: 26–29, 2002.

[88189] 15537.

[88190] Yoshida, A.; Minowa, M. T.; Takamatsu, S.; Hara, T.; Ike-
naga, H.;

[88191] Takeuchi, M.: A novel second isoenzyme of the human
UDP–N–acetylglucosamine:alpha–1,3–D–mannoside

- [88192] beta-1,4-N-acetylglucosaminyltransferase family: cDNA cloning, expression,
[88193] and chromosomal assignment. Glycoconj. J. 15:
1115-1123, 1998.
[88194] 15538.
- [88195] Jakobsson, P.-J.; Mancini, J. A.; Riendeau, D.; Ford-
Hutchinson,
[88196] A. W.: Identification and characterization of a novel micro-
somal
[88197] enzyme with glutathione-dependent transferase and per-
oxidase activities. J.
[88198] Biol. Chem. 272: 22934-22939, 1997.
[88199] 15539.
- [88200] Poliak, S.; Gollan, L.; Martinez, R.; Custer, A.; Einheber, S.;
[88201] Salzer, J. L.; Trimmer, J. S.; Shrager, P.; Peles, E.: Caspr2, a
[88202] new member of the neurexin superfamily, is localized at
the juxtaparanodes
[88203] of myelinated axons and associates with K⁺ channels.
Neuron 24:
[88204] 1037-1047, 1999.
[88205] 15540.

- [88206] Hata, M.; Ohtsuka, K.: Characterization of HSE sequences in human
- [88207] Hsp40 gene: structural and promoter analysis. *Biochim. Biophys. Acta* 1397:
- [88208] 43–55, 1998.
- [88209] 15541.
- [88210] Hata, M.; Okumura, K.; Seto, M.; Ohtsuka, K.: Genomic cloning
- [88211] of a human heat shock protein 40 (Hsp40) gene (HSPF1) and its chromosomal
- [88212] localization to 19p13.2. *Genomics* 38: 446–449, 1996.
- [88213] 15542.
- [88214] Ohtsuka, K.: Cloning of a cDNA for heat-shock protein hsp40, a
- [88215] human homologue of bacterial DnaJ. *Biochem. Biophys. Res. Commun.* 197:
- [88216] 235–240, 1993.
- [88217] 15543.
- [88218] Sultz, B. M.: Genetic control of leucocyte responses to endotoxin. *Nature* 219:
- [88219] 1253–1254, 1968.

- [88220] 15544.
- [88221] Donger, C.; Krejci, E.; Serradell, A. P.; Eymard, B.; Bon, S.;
- [88222] Nicole, S.; Chateau, D.; Gary, F.; Fardeau, M.; Massoulie, J.;
Guicheney,
- [88223] P.: Mutation in the human acetylcholinesterase-associated
collagen
- [88224] gene, COLQ, is responsible for congenital myasthenic
syndrome with
- [88225] end-plate acetylcholinesterase deficiency (type Ic). *Am. J.*
Hum.
- [88226] *Genet.* 63: 967–975, 1998.
- [88227] 15545.
- [88228] Krejci, E.; Coussen, F.; Duval, N.; Chatel, J.-M.; Legay, C.;
- Puype,
- [88229] M.; Vandekerckhove, J.; Cartaud, J.; Bon, S.; Massoulie, J.:
- Primary
- [88230] structure of a collagenic tail peptide of torpedo acetyl-
cholinesterase:
- [88231] co-expression with catalytic subunit induces the produc-
tion of collagen-tailed
- [88232] forms in transfected cells. *EMBO J.* 10: 1285–1293, 1991.

- [88233] 15546.
- [88234] Krejci, E.; Thomine, S.; Boschetti, N.; Legay, C.; Sketelj, J.;
- [88235] Massoulie, J.: The mammalian gene of acetyl-
cholinesterase-associated
- [88236] collagen. J. Biol. Chem. 272: 22840-22847, 1997.
- [88237] 15547.
- [88238] Ohno, K.; Brengman, J. M.; Felice, K. J.; Cornblath, D. R.;
Engel,
- [88239] A. G.: Congenital end-plate acetylcholinesterase defi-
ciency caused
- [88240] by a nonsense mutation and an A-to-G splice-donor-site
mutation at
- [88241] position +3 of the collagenlike-tail-subunit gene (COLQ):
how does
- [88242] G at position +3 result in aberrant splicing? Am. J. Hum.
Genet. 65:
- [88243] 635-644, 1999.
- [88244] 15548.
- [88245] Muravenko, O. V.; Gizatullin, R. Z.; Protopopov, A. I.;
Kashuba,
- [88246] V. I.; Zabarovsky, E. R.; Zelenin, A. V.: Assignment of

CDK5R2 coding

- [88247] for the cyclin-dependent kinase 5, regulatory subunit 2 (NCK5A1 protein)
- [88248] to human chromosome band 2q35 by fluorescent in situ hybridization. Cytogenet.
- [88249] Cell Genet. 89: 160-161, 2000.
- [88250] 15549.
- [88251] Nilden, F.; Backstrom, A.; Bark, C.: Molecular cloning and characterisation
- [88252] of a mouse gene encoding an isoform of the neuronal cyclin-dependent
- [88253] kinase 5 (CDK5) activator. Biochim. Biophys. Acta 1398: 371-376,
- [88254] 1998.
- [88255] 15550.
- [88256] Tang, D.; Yeung, J.; Lee, K-Y.; Matsushita, M.; Matsui, H.; Tomizawa,
- [88257] K.; Hatase, O.; Wang, J. H.: An isoform of the neuronal cyclin-dependent
- [88258] kinase 5 (Cdk5) activator. J. Biol. Chem. 270: 26897-26903, 1995.
- [88259] 15551.

- [88260] Fujioka, M.; Takahashi, N.; Odai, H.; Araki, S.; Ichikawa, K.;
- [88261] Feng, J.; Nakamura, M.; Kaibuchi, K.; Hartshorne, D. J.;
Nakano, T.;
- [88262] Ito, M.: A new isoform of human myosin phosphatase tar-
geting/regulatory
- [88263] subunit (MYPT2): cDNA cloning, tissue expression, and
chromosomal
- [88264] mapping. *Genomics* 49: 59–68, 1998.
- [88265] 15552.
- [88266] Hallas, C.; Pekarsky, Y.; Itoyama, T.; Varnum, J.; Bichi, R.;
Rothstein,
- [88267] J. L.; Croce, C. M.: Genomic analysis of human and mouse
TCL1 loci
- [88268] reveals a complex of tightly clustered genes. *Proc. Nat.*
Acad. Sci. 96:
- [88269] 14418–14423, 1999.
- [88270] 15553.
- [88271] Pekarsky, Y.; Hallas, C.; Isobe, M.; Russo, G.; Croce, C. M.:
- [88272] Abnormalities at 14q32.1 in T cell malignancies involve
two oncogenes. *Proc.*
- [88273] *Nat. Acad. Sci.* 96: 2949–2951, 1999.

- [88274] 15554.
- [88275] Koga, Y.; Fabrizi, G. M.; Mita, S.; Arnaudo, E.; Lomax, M. I.;
- [88276] Aqua, M. S.; Grossman, L. I.; Schon, E. A.: Sequence of a
cDNA specifying
- [88277] subunit VIIc of human cytochrome c oxidase. *Nucleic
Acids Res.* 18:
- [88278] 684 only, 1990.
- [88279] 15555.
- [88280] Harju, K.; Glumoff, V.; Hallman, M.: Ontogeny of Toll-like
receptors
- [88281] Tlr2 and Tlr4 in mice. *Pediat. Res.* 49: 81–83, 2001.
- [88282] 15556.
- [88283] Qu, L. H.; Nicoloso, M.; Michot, B.; Azum, M. C.; Caizer-
gues-Ferrer,
- [88284] M.; Renalier, M. H.; Bachellerie, J. P.: U21, a novel small
nucleolar
- [88285] RNA with a 13 nt. complementarity to 28S rRNA, is en-
coded in an intron
- [88286] of ribosomal protein L5 gene in chicken and mammals.
Nucleic Acids
- [88287] *Res.* 22: 4073–4081, 1994.

[88288] 15557.

[88289] Smith, M.; Wasmuth, J.; McPherson, J. D.; Wagner, C.;
Grandy, D.;

[88290] Civelli, O.; Potkin, S.; Litt, M.: Cosegregation of an
11q22.3-9p22

[88291] translocation with affective disorder: proximity of the
dopamine D2

[88292] receptor gene relative to the translocation breakpoint.
(Abstract) Am.

[88293] J. Hum. Genet. 45 (suppl.): A220 only, 1989.

[88294] 15558.

[88295] Abuladze, N.; Lee, I.; Newman, D.; Hwang, J.; Boorer, K.;
Pushkin,

[88296] A.; Kurtz, I.: Molecular cloning, chromosomal localization,
tissue

[88297] distribution, and functional expression of the human pan-
creatic sodium

[88298] bicarbonate cotransporter. J. Biol. Chem. 273:
17689-17695, 1998.

[88299] 15559.

[88300] Burnham, C. E.; Amlal, H.; Wang, Z.; Shull, G. E.; Soleimani,
M.

- [88301] : Cloning and functional expression of a human kidney Na⁺:HCO₃⁻ cotransporter. J.
- [88302] Biol. Chem. 272: 19111–19114, 1997.
- [88303] 15560.
- [88304] Choi, I.; Romero, M. F.; Khandoudi, N.; Bril, A.; Boron, W. F.
- [88305] : Cloning and characterization of a human electrogenic Na⁽⁺⁾-HCO₃⁻
- [88306] cotransporter isoform (hhNBC). Am. J. Physiol. 276: C576–C584, 1999.
- [88307] 15561.
- [88308] Igarashi, T.; Inatomi, J.; Sekine, T.; Cha, S. H.; Kanai, Y.; Kunimi,
- [88309] M.; Tsukamoto, K.; Satoh, H.; Shimadzu, M.; Tozawa, F.; Mori, T.;
- [88310] Shiobara, M.; Seki, G.; Endou, H.: Mutations in SLC4A4 cause permanent
- [88311] isolated proximal renal tubular acidosis with ocular abnormalities.
- [88312] (Letter) Nature Genet. 23: 264–265, 1999.
- [88313] 15562.

- [88314] Romero, M. F.; Boron, W. F.: Electrogenic Na(+)/HCO(3-) cotransporters:
[88315] cloning and physiology. *Annu. Rev. Physiol.* 61: 699-723, 1999.
[88316] 15563.
- [88317] Soleimani, M.; Burnham, C. E.: Physiologic and molecular aspects
[88318] of the Na(+):HCO(3-) cotransporter in health and disease processes. *Kidney Int.* 57: 371-384, 2000.
[88319] 15564.
- [88320] Usui, T.; et al.; et al. : *Pflugers Arch.* 438: 458-462, 1999.
[88321] 15565.
- [88322] Hayes, S. E.; Conner, L. J.; Stramm, L. E.; Shi, Y.: Assignment
[88323] of pancreatic eIF-2a kinase (EIF2AK3) to human chromosome band 2p12
[88324] by radiation hybrid mapping and in situ hybridization. *Cytogenet. Cell Genet.* 86: 327-328, 1999.
[88325] 15566.

- [88328] Avraham, K. B.; Cho, B. C.; Gilbert, D.; Fujii, H.; Okamoto, K.;
- [88329] Shimazaki, T.; Ito, T.; Shoji, H.; Wakamatsu, Y.; Kondoh, H.; Takahashi,
- [88330] N.; Muramatsu, M.; Hamada, H.; Copeland, N. G.; Jenkins, N. A.: Murine
- [88331] chromosomal location of four class III POU transcription factors. *Genomics* 18:
- [88332] 131–133, 1993.
- [88333] 15567.
- [88334] Egan, S.; Herbrick, J.-A.; Tsui, L.-C.; Cohen, B.; Flock, G.; Beatty,
- [88335] B.; Scherer, S. W.: Mapping of the human lunatic fringe (LFNG) gene
- [88336] to 7p22 and manic fringe (MFNG) to 22q12. *Genomics* 54: 576–577,
- [88337] 1998.
- [88338] 15568.
- [88339] Johnston, S. H.; Rauskolb, C.; Wilson, R.; Prabhakaran, B.; Irvine,
- [88340] K. D.; Vogt, T. F.: A family of mammalian Fringe genes implicated

[88341] in boundary determination and the Notch pathway. Development 124:

[88342] 2245–2254, 1997.

[88343] 15569.

[88344] Moran, J. L.; Johnston, S. H.; Rauskolb, C.; Bhalerao, J.; Bowcock,

[88345] A. M.; Vogt, T. F.: Genomic structure, mapping, and expression analysis

[88346] of the mammalian lunatic, manic, and radical fringe genes. Mammalian

[88347] Genome 10: 535–541, 1999.

[88348] 15570.

[88349] Van Tine, B. A.; Knops, J.; Shaw, G. M.; May, W. A.: Assignment

[88350] of human MFNG, manic fringe Drosophila homolog, to 22q13.1 using tyramide

[88351] fluorescence in situ hybridization (T-FISH). Cytogenet. Cell Genet. 87:

[88352] 132–133, 1999.

[88353] 15571.

[88354] Laufer, E.; Dahn, R.; Orozco, O. E.; Yeo, C.-Y.; Pisenti, J.;

Henrique,

[88355] D.; Abbott, U. K.; Fallon, J. F.; Tabin, C.: Expression of
Radical

[88356] fringe in limb-bud ectoderm regulates apical ectodermal
ridge formation. Nature 386:

[88357] 366–373, 1997.

[88358] 15572.

[88359] Rodriguez-Esteban, C.; Schwabe, J. W. R.; De La Pena, J.;
Foy,

[88360] B.; Eshelman, B.; Izpisua Belmonte, J. C.: Radical fringe
positions

[88361] the apical ectodermal ridge at the dorsoventral boundary
of the vertebrate

[88362] limb. Nature 386: 360–366, 1997.

[88363] 15573.

[88364] Wiley, S. R.; Cassiano, L.; Lofton, T.; Davis-Smith, T.; Win-
kles,

[88365] J. A.; Lindner, V.; Liu, H.; Daniel, T. O.; Smith, C. A.;
Fanslow,

[88366] W. C.: A novel TNF receptor family member binds TWEAK
and is implicated

[88367] in angiogenesis. Immunity 15: 837–846, 2001.

[88368] 15574.

[88369] Hoey, T.; Sun, Y.-L.; Williamson, K.; Xu, X.: Isolation of two

[88370] new members of the NF-AT gene family and functional
characterization

[88371] of the NF-AT proteins. *Immunity* 2: 461-472, 1995.

[88372] 15575.

[88373] Arany, Z.; Newsome, D.; Oldread, E.; Livingston, D. M.;
Eckner,

[88374] R.: A family of transcriptional adaptor proteins targeted by
the

[88375] E1A oncoprotein. *Nature* 374: 81-84, 1995.

[88376] 15576.

[88377] Dong, C.; Zhu, S.; Wang, T.; Yoon, W.; Li, Z.; Alvarez, R. J.;

[88378] ten Dijke, P.; White, B.; Wigley, F. M.; Goldschmidt-Cler-
mont, P.

[88379] J.: Deficient Smad7 expression: a putative molecular de-
fect in scleroderma. *Proc.*

[88380] *Nat. Acad. Sci.* 99: 3908-3913, 2002.

[88381] 15577.

[88382] Kavsak, P.; Rasmussen, R. K.; Causing, C. G.; Bonni, S.;

Zhu, H.;

[88383] Thomsen, G. H.; Wrana, J. L.: Smad7 binds to Smurf2 to form an E3

[88384] ubiquitin ligase that targets the TGF-beta receptor for degradation. Molec.

[88385] Cell 6: 1365-1375, 2000.

[88386] 15578.

[88387] Lallemand, F.; Mazars, A.; Prunier, C.; Bertrand, F.; Korn-prost,

[88388] M.; Gallea, S.; Roman-Roman, S.; Cherqui, G.; Atfi, A.: Smad7 inhibits

[88389] the survival nuclear factor kappa-B and potentiates apoptosis in epithelial

[88390] cells. Oncogene 20: 879-884, 2001.

[88391] 15579.

[88392] Roijer, E.; Moren, A.; ten Dijke, P.; Stenman, G.: Assignment

[88393] of the Smad7 gene (MADH7) to human chromosome 18q21.1 by fluorescence

[88394] in situ hybridization. Cytogenet. Cell Genet. 81: 189-190, 1998.

[88395] 15580.

- [88396] Cravatt, B. F.; Giang, D. K.; Mayfield, S. P.; Boger, D. L.; Lerner, R. A.; Gilula, N. B.: Molecular characterization of an enzyme that degrades neuromodulatory fatty-acid amides. *Nature* 384: 83–87, 1996.
- [88399] 15581.
- [88400] Giang, D. K.; Cravatt, B. F.: Molecular characterization of human and mouse fatty acid amide hydrolases. *Proc. Nat. Acad. Sci.* 94: 2238–2242, 1997.
- [88403] 15582.
- [88404] Sipe, J. C.; Chiang, K.; Gerber, A. L.; Beutler, E.; Cravatt, B. F.: A missense mutation in human fatty acid amide hydrolase associated with problem drug use. *Proc. Nat. Acad. Sci.* 99: 8394–8399, 2002.
- [88407] 15583.
- [88408] Wan, M.; Cravatt, B. F.; Ring, H. Z.; Zhang, X.; Francke, U.: Conserved chromosomal location and genomic structure

of human and

[88410] mouse fatty-acid amide hydrolase genes and evaluation of
clasper as

[88411] a candidate neurological mutation. Genomics 54:
408–414, 1998.

[88412] 15584.

[88413] Baier, M.; Bannert, N.; Werner, A.; Lang, K.; Kurth, R.:
Molecular

[88414] cloning, sequence, expression, and processing of the in-
terleukin 16

[88415] precursor. Proc. Nat. Acad. Sci. 94: 5273–5277, 1997.

[88416] 15585.

[88417] Roh, M. H.; Makarova, O.; Liu, C.-J.; Shin, K.; Lee, S.; Lau-
rinec,

[88418] S.; Goyal, M.; Wiggins, R.; Margolis, B.: The Maguk protein,
Pals1,

[88419] functions as an adapter, linking mammalian homologues
of Crumbs and

[88420] Discs Lost. J. Cell Biol. 157: 161–172, 2002.

[88421] 15586.

[88422] Bannert, N.; Baier, M.; Werner, A.; Kurth, R.: Interleukin-16

- [88423] or not? Nature 381: 30 only, 1996.
- [88424] 15587.
- [88425] Bannert, N.; Kurth, R.; Baier, M.: The gene encoding mouse interleukin-16
- [88426] consists of seven exons and maps to Chromosome 7 D2-D3. Immunogenetics 49:
- [88427] 704-706, 1999.
- [88428] 15588.
- [88429] Cruikshank, W.; Center, D. M.: Modulation of lymphocyte migration
- [88430] by human lymphokines. II. Purification of a lymphotactic factor (LCF). J.
- [88431] Immun. 128: 2569-2574, 1982.
- [88432] 15589.
- [88433] Cruikshank, W. W.; Center, D. M.; Nisar, N.; Wu, M.; Natke, B.;
- [88434] Theodore, A. C.; Kornfeld, H.: Molecular and functional analysis
- [88435] of a lymphocyte chemoattractant factor: association of biologic function
- [88436] with CD4 expression. Proc. Nat. Acad. Sci. 91:

5109–5113, 1994.

[88437] 15590.

[88438] Drwinga, H. L.; Toji, L. H.; Kim, C. H.; Greene, A. E.;
Mulivor,

[88439] R. A.: NIGMS human/rodent somatic cell hybrid mapping
panels 1 and

[88440] 2. Genomics 16: 311–314, 1993.

[88441] 15591.

[88442] Hudson, T. J.; Stein, L. D.; Gerety, S. S.; Ma, J.; Castle, A.

[88443] B.; Silva, J.; Slonim, D. K.; Baptista, R.; Kruglyak, L.; Xu, S.
H.;

[88444] Hu, X.; Colbert, A. M. E.; and 39 others: An STS-based
map of the

[88445] human genome. Science 270: 1945–1954, 1995.

[88446] 15592.

[88447] Keane, J.; Nicoll, J.; Kim, S.; Wu, D. M. H.; Cruikshank, W.
W.;

[88448] Brazer, W.; Natke, B.; Zhang, Y.; Center, D. M.; Kornfeld,
H.: Conservation

[88449] of structure and function between human and murine IL–
16. J. Immun. 160:

[88450] 5945–5954, 1998.

[88451] 15593.

[88452] Tarnasky, H.; Gill, D.; Murthy, S.; Shao, X.; Demetrick, D. J.;

[88453] van der Hoorn, F. A.: A novel testis-specific gene, SPAG4, whose

[88454] product interacts specifically with outer dense fiber protein ODF27,

[88455] maps to human chromosome 20q11.2. *Cytogenet. Cell Genet.* 81: 65–67,

[88456] 1998.

[88457] 15594.

[88458] Hurlin, P. J.; Queva, C.; Eisenman, R. N.: Mnt, a novel Max-interacting

[88459] protein is coexpressed with Myc in proliferating cells and mediates

[88460] repression at Myc binding sites. *Genes Dev.* 11: 44–58, 1997.

[88461] 15595.

[88462] Lo Nigro, C.; Venesio, T.; Reymond, A.; Meroni, G.; Alberici, P.;

- [88463] Cainarca, S.; Enrico, F.; Stack, M.; Ledbetter, D. H.; Liscia, D.
- [88464] S.; Ballabio, A.; Carrozzo, R.: The human ROX gene: genomic structure
- [88465] and mutation analysis in human breast tumors. *Genomics* 49: 275–282,
- [88466] 1998.
- [88467] 15596.
- [88468] Meroni, G.; Reymond, A.; Alcalay, M.; Borsani, G.; Tanigami, A.;
- [88469] Tonlorenzi, R.; Lo Nigro, C.; Messali, S.; Zollo, M.; Ledbetter, D.
- [88470] H.; Brent, R.; Ballabio, A.; Carrozzo, R.: Rox, a novel bHLHZip protein
- [88471] expressed in quiescent cells that heterodimerizes with Max, binds
- [88472] a non-canonical E box and acts as a transcriptional repressor. *EMBO*
- [88473] *J.* 16: 2892–2906, 1997.
- [88474] 15597.
- [88475] Lennon-Dumenil, A.-M.; Barbouche, M.-R.; Vedrenne, J.; Prod'Homme,

- [88476] T.; Bejaoui, M.; Ghariani, S.; Charron, D.; Fellous, M.; Del-
lagi,
- [88477] K.; Alcaide-Loridan, C.: Uncoordinated HLA-D gene ex-
pression in a
- [88478] RFXANK-defective patient with MHC class II deficiency. J.
Immun. 166:
- [88479] 5681-5687, 2001.
- [88480] 15598.
- [88481] Lisowska-Groszpiere, B.; Fondaneche, M. C.; Rols, M. P.;
Griscelli,
- [88482] C.; Fischer, A.: Two complementation groups account for
most cases
- [88483] of inherited MHC class II deficiency. Hum. Molec. Genet. 3:
953-958,
- [88484] 1994.
- [88485] 15599.
- [88486] Gerloff, T.; Stieger, B.; Hagenbuch, B.; Madon, J.; Land-
mann, L.;
- [88487] Roth, J.; Hofmann, A. F.; Meier, P. J.: The sister of P-
glycoprotein
- [88488] represents the canalicular bile salt export pump of mam-
malian liver. J.

- [88489] Biol. Chem. 273: 10046–10050, 1998.
- [88490] 15600.
- [88491] Wang, R.; Salem, M.; Yousef, I. M.; Tuchweber, B.; Lam, P.; Childs,
- [88492] S. J.; Helgason, C. D.; Ackerley, C.; Phillips, M. J.; Ling, V.:
- [88493] Targeted inactivation of sister of P-glycoprotein gene (spgp) in mice
- [88494] results in nonprogressive but persistent intrahepatic cholestasis. Proc.
- [88495] Nat. Acad. Sci. 98: 2011–2016, 2001.
- [88496] 15601.
- [88497] Wiszniewski, W.; Fondaneche, M.-C.; Lambert, N.; Master-nak, K.;
- [88498] Picard, C.; Notarangelo, L.; Schwartz, K.; Bal, J.; Reith, W.; Alcaide,
- [88499] C.; de Saint Basile, G.; Fischer, A.; Lisowska-Groszpiere, B.: Founder
- [88500] effect for a 26-bp deletion in the RFXANK gene in North African major
- [88501] histocompatibility complex class II-deficient patients belonging to

- [88502] complementation group B. Immunogenetics 51: 261–267, 2000.
- [88503] 15602.
- [88504] Stec, I.; Wright, T. J.; van Ommen, G.–J. B.; de Boer, P. A. J.;
- [88505] van Haeringen, A.; Moorman, A. F. M.; Altherr, M. R.; den Dunnen,
- [88506] J. T.: WHSC1, a 90 kb SET domain–containing gene, expressed in early
- [88507] development and homologous to a Drosophila dysmorphia gene maps in
- [88508] the Wolf–Hirschhorn syndrome critical region and is fused to IgH in
- [88509] t(4;14) multiple myeloma. Hum. Molec. Genet. 7: 1071–1082, 1998.
- [88510] 15603.
- [88511] Buanne, P.; Incerti, B.; Guardavaccaro, D.; Avvantaggiato, V.;
- [88512] Simeone, A.; Tirone, F.: Cloning of the human interferon–related
- [88513] developmental regulator (IFRD1) gene coding for the PC4 protein, a
- [88514] member of a novel family of developmentally regulated

genes. Genomics 51:

[88515] 233–242, 1998.

[88516] 15604.

[88517] Chen, D.; Lathrop, W.; Dong, Y.: Molecular cloning of mouse Lrp7(Lr3)

[88518] cDNA and chromosomal mapping of orthologous genes in mouse and human. Genomics 55:

[88519] 314–321, 1999.

[88520] 15605.

[88521] Dong, Y.; Lathrop, W.; Weaver, D.; Qiu, Q.; Cini, J.; Bertolini,

[88522] D.; Chen, D.: Molecular cloning and characterization of LR3, a novel

[88523] LDL receptor family protein with mitogenic activity. Biochem. Biophys.

[88524] Res. Commun. 251: 784–790, 1998.

[88525] 15606.

[88526] Mao, J.; Wang, J.; Liu, B.; Pan, W.; Farr, G. H., III; Flynn, C.;

[88527] Yuan, H.; Takada, S.; Kimelman, D.; Li, L.; Wu, D.: Low-density lipoprotein

[88528] receptor-related protein-5 binds to axin and regulates the

canonical

[88529] Wnt signaling pathway. *Molec. Cell* 7: 801–809, 2001.

[88530] 15607.

[88531] Mao, B.; Wu, W.; Li, Y.; Hoppe, D.; Stannek, P.; Glinka, A.; Niehrs,

[88532] C.: LDL-receptor-related protein 6 is a receptor for Dickkopf proteins. *Nature* 411:

[88533] 321–325, 2001.

[88534] 15608.

[88535] Vinkemeier, U.; Obermann, W.; Weber, K.; Furst, D. O.: The globular

[88536] head domain of titin extends into the center of the sarcomeric M band:

[88537] cDNA cloning, epitope mapping and immunoelectron microscopy of two

[88538] titin-associated proteins. *J. Cell Sci.* 106: 319–330, 1993.

[88539] 15609.

[88540] van der Ven, P. F. M.; Speel, E. J. M.; Albrechts, J. C. M.; Ramaekers,

[88541] F. C. S.; Hopman, A. H. N.; Furst, D. O.: Assignment of the human

- [88542] gene for endosarcomeric cytoskeletal M-protein (MYOM2) to 8p23.3. Genomics 55: 253–255, 1999.
- [88543] 15610.
- [88544] 15610.
- [88545] Murakami, M.; Mori, M.: Identification of immunogenic regions in human thyrotropin receptor for immunoglobulin G of patients with Graves' disease. Biochem. Biophys. Res. Commun. 171: 512–518, 1990.
- [88546] 15611.
- [88547] Nagayama, Y.; Kaufman, K. D.; Seto, P.; Rapoport, B.: Molecular cloning, sequence and functional expression of the cDNA for the human thyrotropin receptor. Biochem. Biophys. Res. Commun. 165: 1184–1190, 1989.
- [88548] 15612.
- [88549] Hirohata, S.; Seldin, M. F.; Apte, S. S.: Chromosomal assignment

- [88555] of two ADAM genes, TACE (ADAM17) and MLTNB (ADAM19), to human chromosomes
- [88556] 2 and 5, respectively, and of Mltnb to mouse chromosome 11. Genomics 54:
- [88557] 178–179, 1998.
- [88558] 15613.
- [88559] Inoue, D.; Reid, M.; Lum, L.; Kratzschmar, J.; Weskamp, G.; Myung,
- [88560] Y. M.; Baron, R.; Blobel, C. P.: Cloning and initial characterization
- [88561] of mouse meltrin beta and analysis of the expression of four metalloprotease–disintegrins
- [88562] in bone cells. J. Biol. Chem. 273: 4180–4187, 1998.
- [88563] 15614.
- [88564] Petruzzella, V.; Tiranti, V.; Fernandez, P.; Ianna, P.; Carrozzo,
- [88565] R.; Zeviani, M.: Identification and characterization of human cDNAs
- [88566] specific to BCS1, PET112, SCO1, COX15, and COX11, five genes involved
- [88567] in the formation and function of the mitochondrial respiratory chain. Genomics 54:

[88568] 494–504, 1998.

[88569] 15615.

[88570] Sakuma, H.; Ikeda, A.; Oka, S.; Kozutsumi, Y.; Zanetta, J.-P.;

[88571] Kawasaki, T.: Molecular cloning and functional expression of a cDNA

[88572] encoding a new member of mixed lineage protein kinase from human brain. J.

[88573] Biol. Chem. 272: 28622–28629, 1997.

[88574] 15616.

[88575] Iyer, N. V.; Leung, S. W.; Semenza, G. L.: The human hypoxia-inducible

[88576] factor 1- α gene: HIF1A structure and evolutionary conservation. Genomics 52:

[88577] 159–165, 1998.

[88578] 15617.

[88579] Kline, D. D.; Peng, Y.-J.; Manalo, D. J.; Semenza, G. L.; Prabhakar,

[88580] N. R.: Defective carotid body function and impaired ventilatory responses

[88581] to chronic hypoxia in mice partially deficient for hypoxia-

inducible

[88582] factor 1- α . Proc. Nat. Acad. Sci. 99: 821-826, 2002.

[88583] 15618.

[88584] Lando, D.; Peet, D. J.; Whelan, D. A.; Gorman, J. J.;
Whitelaw,

[88585] M. L.: Asparagine hydroxylation of the HIF transactivation
domain:

[88586] a hypoxic switch. Science 295: 858-861, 2002.

[88587] 15619.

[88588] Marti, H. H.; Katschinski, D. M.; Wagner, K. F.; Schaffer, L.;

[88589] Stier, B.; Wenger, R. H.: Isoform-specific expression of hy-
poxia-inducible

[88590] factor-1- α during the late stages of mouse spermi-
genesis. Molec.

[88591] Endocr. 16: 234-243, 2002.

[88592] 15620.

[88593] Maxwell, P. H.; Wiesener, M. S.; Chang, G.-W.; Clifford, S.
C.;

[88594] Vaux, E. C.; Cockman, M. E.; Wykoff, C. C.; Pugh, C. W.;
Maher, E.

[88595] R.; Ratcliffe, P. J.: The tumour suppressor protein VHL tar-

gets hypoxia-inducible

[88596] factors for oxygen-dependent proteolysis. *Nature* 399: 271–275, 1999.

[88597] 15621.

[88598] Min, J.-H.; Yang, H.; Ivan, M.; Gertler, F.; Kaelin, W. G., Jr.;

[88599] Pavletich, N. P.: Structure of an HIF-1- α -pVHL complex: hydroxyproline

[88600] recognition in signaling. *Science* 296: 1886–1889, 2002.

[88601] 15622.

[88602] Semenza, G. L.: HIF-1 and human disease: one highly involved

[88603] factor. *Genes Dev.* 14: 1983–1991, 2000.

[88604] 15623.

[88605] Semenza, G. L.; Rue, E. A.; Iyer, N. V.; Pang, M. G.; Kearns,

[88606] W. G.: Assignment of the hypoxia-inducible factor 1- α gene to

[88607] a region of conserved synteny on mouse chromosome 12 and human chromosome

[88608] 14q. *Genomics* 34: 437–439, 1996.

[88609] 15624.

- [88610] Sutter, C. H.; Laughner, E.; Semenza, G. L.: Hypoxia-inducible
- [88611] factor 1- α protein expression is controlled by oxygen-regulated
- [88612] ubiquitination that is disrupted by deletions and missense mutations. Proc.
- [88613] Nat. Acad. Sci. 97: 4748-4753, 2000.
- [88614] 15625.
- [88615] Wenger, R. H.; Rolfs, A.; Kvietikova, I.; Spielmann, P.; Zimmermann,
- [88616] D. R.; Gassmann, M.: The mouse gene for hypoxia-inducible factor-1- α --genomic
- [88617] organization, expression and characterization of an alternative first
- [88618] exon and 5-prime flanking sequence. Europ. J. Biochem. 246: 155-165,
- [88619] 1997.
- [88620] 15626.
- [88621] Wenger, R. H.; Rolfs, A.; Marti, H. H.; Guenet, J.-L.; Gassmann,
- [88622] M.: Nucleotide sequence, chromosomal assignment and mRNA expression

- [88623] of mouse hypoxia-inducible factor-1-alpha. Biochem. Biophys. Res.
- [88624] Commun. 223: 54-59, 1996.
- [88625] 15627.
- [88626] Wenger, R. H.; Rolfs, A.; Spielmann, P.; Zimmermann, D. R.; Gassmann,
- [88627] M.: Mouse hypoxia-inducible factor-1-alpha is encoded by two different
- [88628] mRNA isoforms: expression from a tissue-specific and a housekeeping-type
- [88629] promoter. Blood 91: 3471-3480, 1998.
- [88630] 15628.
- [88631] Hovanessian, A. G.; Laurent, A. G.; Chebath, J.; Galabru, J.; Robert,
- [88632] N.; Svab, J.: Identification of 69-kd and 100-kd forms of 2-5A synthetase
- [88633] in interferon-treated human cells by specific monoclonal antibodies. EMBO
- [88634] J. 6: 1273-1280, 1987.
- [88635] 15629.
- [88636] Marie, I.; Galabru, J.; Svab, J.; Hovanessian, A. G.: Prepara-

tion

[88637] and characterization of polyclonal antibodies specific for the 69

[88638] and 100 k-dalton forms of human 2-5A synthetase.

Biochem. Biophys.

[88639] Res. Commun. 160: 580-587, 1989.

[88640] 15630.

[88641] Marie, I.; Hovanessian, A. G.: The 69-kDa 2-5A synthetase is composed

[88642] of two homologous and adjacent functional domains. J. Biol. Chem. 267:

[88643] 9933-9939, 1992.

[88644] 15631.

[88645] Kools, P.; Van Imschoot, G.; van Roy, F.: Characterization of

[88646] three novel human cadherin genes (CDH7, CDH19, and CDH20) clustered

[88647] on chromosome 18q22-q23 and with high homology to chicken cadherin-7. Genomics 68:

[88648] 283-295, 2000.

[88649] 15632.

- [88650] Amado, M.; Almeida, R.; Carneiro, F.; Lavery, S. B.; Holmes, E.
- [88651] H.; Nomoto, M.; Hollingsworth, M. A.; Hassan, H.; Schwientek, T.;
- [88652] Nielsen, P. A.; Bennett, E. P.; Clausen, H.: A family of human beta-3-galactosyltransferases:
- [88653] characterization of four members of a UDP-galactose:beta-N-acetyl-glucosamine/beta-N-acetyl-galactosamine
- [88654] beta-1,3-galactosyltransferase family. J. Biol. Chem. 273: 12770-12778,
- [88655] 1998.
- [88656] 15633.
- [88657] Hennet, T.; Dinter, A.; Kuhnert, P.; Mattu, T. S.; Rudd, P. M.;
- [88658] Berger, E. G.: Genomic cloning and expression of three murine UDP-galactose:beta-N-acetylglucosamine
- [88659] beta-1,3-galactosyltransferase genes. J. Biol. Chem. 273: 58-65,
- [88660] 1998.
- [88661] 15634.
- [88662] Kolbinger, F.; Streiff, M. B.; Katopodis, A. G.: Cloning of a

[88663] human UDP-galactose:2-acetamido-2-deoxy-D-glucose
3-beta-galactosyltransferase

[88664] catalyzing the formation of type 1 chains. J. Biol. Chem.
273: 433-440,

[88665] 1998.

[88666] 15635.

[88667] Hsu, D. R.; Economides, A. N.; Wang, X.; Eimon, P. M.;
Harland,

[88668] R. M.: The Xenopus dorsalizing factor gremlin identifies a
novel

[88669] family of secreted proteins that antagonize BMP activities.

Molec.

[88670] Cell 1: 673–683, 1998.

[88671] 15636.

[88672] Topol, L. Z.; Modi, W. S.; Koochekpour, S.; Blair, D. G.:

DRM–Gremlin

[88673] (CKTSF1B1) maps to human chromosome 15 and is highly expressed in

[88674] adult and fetal brain. Cytogenet. Cell Genet. 89: 79–84, 2000.

[88675] 15637.

[88676] Beaconsfield, P.; Rainsbury, R.; Kalton, G.: Glucose–6-phosphate

[88677] dehydrogenase deficiency and the incidence of cancer. Oncologia 19:

[88678] 11–19, 1965.

[88679] 15638.

[88680] Ben–Bassat, J.; Ben–Ishay, D.: Hereditary hemolytic anemia associated

[88681] with glucose–6–phosphate dehydrogenase deficiency (Mediterranean type). Israel

[88682] J. Med. Sci. 5: 1053–1059, 1969.

[88683] 15639.

[88684] Benabadji, M.; Merad, F.; Benmoussa, M.; Trabuchet, G.;
Junien,

[88685] C.; Dreyfus, J. C.; Kaplan, J. C.: Heterogeneity of glucose–
6-phosphate

[88686] dehydrogenase deficiency in Algeria. Hum. Genet. 40:
177–184, 1978.

[88687] 15640.

[88688] Benohr, H. C.; Klumpp, F.; Waller, H. D.: Glucose–
6-phosphat-Dehydrogenase

[88689] Typ Schwaben. Dtsch. Med. Wschr. 96: 1029–1032, 1971.

[88690] 15641.

[88691] Benohr, H. C.; Waller, H. D.: Eigenschaften der Glucose–
6-p-dehydrogenase,

[88692] Typ Tubingen. Klin. Wschr. 48: 71–74, 1970.

[88693] 15642.

[88694] Benohr, H. C.; Waller, H. D.; Arnold, H.; Blume, K. G.; Lohr,

[88695] G. W.: Glucose–6-P-Dehydrogenase Typ Bodensee (eine
neue Enzymvariante). Klin.

[88696] Wschr. 49: 1058–1062, 1971.

[88697] 15643.

[88698] Beutler, E.: Glucose 6–phosphate dehydrogenase deficiency, a

[88699] new Indian variant, G6PD Jammu. In: Sen, N. N.; Basu, A. K.: Trends

[88700] in Haematology. Calcutta: Chatterjea Memorial Committee (pub.)

[88701] 1975. Pp. 279–283.

[88702] 15644.

[88703] Beutler, E.: Personal Communication. La Jolla, Calif. 11/12/1990.

[88704] 15645.

[88705] Beutler, E.: G6PD deficiency. Blood 84: 3613–3636, 1994.

[88706] 15646.

[88707] Beutler, E.: Glucose–6–phosphate dehydrogenase deficiency. New

[88708] Eng. J. Med. 324: 169–174, 1991.

[88709] 15647.

[88710] Beutler, E.: Selectivity of proteases as a basis for tissue

distribution

[88711] of enzymes in hereditary deficiencies. Proc. Nat. Acad. Sci.
80:

[88712] 3767–3768, 1983.

[88713] 15648.

[88714] Beutler, E.: Glucose–6–phosphate dehydrogenase deficiency. In:

[88715] Wintrobe, M. M.: Red Cell Metabolism in Hemolytic Anemia. New York:

[88716] Plenum Press (pub.) 1978.

[88717] 15649.

[88718] Beutler, E.: The hemolytic effect of primaquine and related compounds:

[88719] a review. Blood 14: 103–139, 1959.

[88720] 15650.

[88721] Beutler, E.; Grooms, A. M.; Morgan, S. K.; Trinidad, F.: Chronic

[88722] severe hemolytic anemia due to G–6–PD Charleston: a new deficiency

[88723] variant. J. Pediat. 80: 1005–1009, 1972.

[88724] 15651.

- [88725] Beutler, E.; Hartman, K.; Gelbart, T.; Forman, L.: G-6-PD
Walter
- [88726] Reed: possible insight into 'structural' NADP in G-6-PD.
Am. J. Hemat. 23:
- [88727] 25-30, 1986.
- [88728] 15652.
- [88729] Beutler, E.; Keller, J. W.; Matsumoto, F.: A new glucose
6-phosphate
- [88730] dehydrogenase (G6PD) variant associated with nonsphero-
cytic hemolytic
- [88731] anemia: G6PD Atlanta. I.R.C.S. 4: 479, 1976.
- [88732] 15653.
- [88733] Beutler, E.; Kuhl, W.: The NT 1311 polymorphism of G6PD:
G6PD
- [88734] Mediterranean mutation may have originated indepen-
dently in Europe
- [88735] and Asia. Am. J. Hum. Genet. 47: 1008-1012, 1990.
- [88736] 15654.
- [88737] Beutler, E.; Kuhl, W.: Linkage between a PvuII restriction
fragment
- [88738] length polymorphism and G6PD A- (202A/376G): evi-

dence for a single

[88739] origin of the common G6PD A- mutation. Hum. Genet. 85: 9-11, 1990.

[88740] 15655.

[88741] Beutler, E.; Kuhl, W.; Gelbart, T.; Forman, L.: DNA sequence

[88742] abnormalities of human glucose-6-phosphate dehydrogenase variants. J.

[88743] Biol. Chem. 266: 4145-4150, 1991.

[88744] 15656.

[88745] Beutler, E.; Kuhl, W.; Ramirez, E.; Lisker, R.: Some Mexican

[88746] glucose-6-phosphate dehydrogenase variants revisited. Hum. Genet. 86:

[88747] 371-374, 1991.

[88748] 15657.

[88749] Beutler, E.; Kuhl, W.; Saenz, G. F.; Rodriguez, W.: Mutation

[88750] analysis of glucose-6-phosphate dehydrogenase (G6PD) variants in Costa

[88751] Rica. Hum. Genet. 87: 462-464, 1991.

[88752] 15658.

- [88753] Beutler, E.; Mathai, C. K.; Smith, J. E.: Biochemical variants
[88754] of glucose-6-phosphate dehydrogenase giving rise to
congenital nonspherocytic
[88755] hemolytic disease. Blood 31: 131-150, 1968.
[88756] 15659.
- [88757] Beutler, E.; Matsumoto, F.: A new glucose 6-phosphate
dehydrogenase
[88758] variant: G6PD (-) Los Angeles. I.R.C.S. 5: 89, 1977.
[88759] 15660.
- [88760] Beutler, E.; Matsumoto, F.; Daiber, A.: Nonspherocytic
hemolytic
[88761] anemia due to G-6-PD Panama. I.R.C.S. 2: 1389, 1974.
[88762] 15661.
- [88763] Beutler, E.; Rosen, R.: Nonspherocytic congenital
hemolytic anemia
[88764] due to a new G-6-PD variant: G-6-PD Alhambra. Pediatrics 45: 230-235,
[88765] 1970.
[88766] 15662.
- [88767] Beutler, E.; Westwood, B.; Prchal, J. T.; Vaca, G.; Bartsocas,

- [88768] C. S.; Baronciani, L.: New glucose-6-phosphate dehydrogenase mutations
- [88769] from various ethnic groups. Blood 80: 255-256, 1992.
- [88770] 15663.
- [88771] Beutler, E.; Yoshida, A.: Human glucose-6-phosphate dehydrogenase
- [88772] variants: a supplementary tabulation. Ann. Hum. Genet. 37: 151-156,
- [88773] 1973.
- [88774] 15664.
- [88775] Boivin, P.; Galand, C.: Nouvelles variantes de la glucose-6-phosphate
- [88776] dehydrogenase erythrocytaire. Rev. Franc. Etud. Clin. Biol. 13:
- [88777] 30-39, 1968.
- [88778] 15665.
- [88779] Botha, M. C.; Dern, R. J.; Mitchell, M.; West, C.; Beutler, E.
- [88780] : G6PD Capetown, a variant of glucose-6-phosphate dehydrogenase. Am.
- [88781] J. Hum. Genet. 21: 547-551, 1969.

- [88782] 15666.
- [88783] Boyer, S. H.; Graham, J. B.: Linkage between the X chromosome
- [88784] loci for glucose-6-phosphate dehydrogenase electrophoretic variation
- [88785] and hemophilia A. Am. J. Hum. Genet. 17: 320-324, 1965.
- [88786] 15667.
- [88787] Boyer, S. H.; Porter, I. H.; Weilbaecher, R. G.: Electrophoretic
- [88788] heterogeneity of glucose-6-phosphate dehydrogenase and its relationship
- [88789] to enzyme deficiency in man. Proc. Nat. Acad. Sci. 48: 1868-1876,
- [88790] 1962.
- [88791] 15668.
- [88792] Busch, D.; Bote, K.: Glucose-6-phosphate-dehydrogenase-Defect
- [88793] in Deutschland. II. Eigenschaften des Enzyms (Typ Freiburg). Klin.
- [88794] Wschr. 48: 74-78, 1970.
- [88795] 15669.

- [88796] Calabro, V.; Giacobbe, A.; Vallone, D.; Montanaro, V.; Cascone,
- [88797] A.; Filosa, S.; Battistuzzi, G.: Genetic heterogeneity at the glucose-6-phosphate
- [88798] dehydrogenase locus in southern Italy: a study on a population from
- [88799] the Matera district. Hum. Genet. 86: 49-53, 1990.
- [88800] 15670.
- [88801] Cappadoro, M.; Giribaldi, G.; O'Brien, E.; Turrini, F.; Mannu,
- [88802] F.; Ulliers, D.; Simula, G.; Luzzatto, L.; Arese, P.: Early phagocytosis
- [88803] of glucose-6-phosphate dehydrogenase (G6PD)-deficient erythrocytes
- [88804] parasitized by Plasmodium falciparum may explain malaria protection
- [88805] in G6PD deficiency. Blood 92: 2527-2534, 1998.
- [88806] 15671.
- [88807] Cappellini, M. D.; Sampietro, M.; Toniolo, D.; Carandina, G.;
- [88808] Pittalis, S.; Martinez di Montemuros, F.; Tavazzi, D.; Fiorelli, G.

- [88809] : Biochemical and molecular characterization of a new sporadic glucose-6-phosphate
- [88810] dehydrogenase variant described in Italy: G6PD Modena. Brit. J. Haemat. 87:
- [88811] 209-211, 1994.
- [88812] 15672.
- [88813] Carandina, G.; Moretto, E.; Zecchi, G.; Conighi, C.: Glucose
- [88814] 6-phosphate dehydrogenase Ferrara. A new variant of G6PD identified
- [88815] in Northern Italy. Acta Haemat. 56: 116-122, 1976.
- [88816] 15673.
- [88817] Carson, P. E.; Flanagan, C. L.; Ickes, C. E.; Alving, A. S.:
- [88818] Enzymatic deficiency in primaquine-sensitive erythrocytes. Science 124:
- [88819] 484-485, 1956.
- [88820] 15674.
- [88821] Poy, F.; Yaffe, M. B.; Sayos, J.; Saxena, K.; Morra, M.; Sumegi,
- [88822] J.; Cantley, L. C.; Terhorst, C.; Eck, M. J.: Crystal structures
- [88823] of the XLP protein SAP reveal a class of SH2 domains with

extended,

[88824] phosphotyrosine-independent sequence recognition.

Molec. Cell 4:

[88825] 555-561, 1999.

[88826] 15675.

[88827] Provisor, A. J.; Iaccone, J. J.; Chilcote, R. R.; Neiburger, R.

[88828] G.; Crussi, F. G.; Baehner, R. L.: Acquired agammaglobulinemia after

[88829] a life-threatening illness with clinical and laboratory features of

[88830] infectious mononucleosis in three related male children.
New Eng.

[88831] J. Med. 293: 62-65, 1975.

[88832] 15676.

[88833] Purtilo, D. T.: Pathogenesis and phenotypes of an X-linked recessive

[88834] lymphoproliferative syndrome. Lancet II: 882-885, 1976.

[88835] 15677.

[88836] Purtilo, D. T.: X-linked lymphoproliferative syndrome: an immunodeficiency

[88837] disorder with acquired agammaglobulinemia, fatal infec-

tious mononucleosis,

[88838] or malignant lymphoma. Arch. Path. Lab. Med. 105:
119-121, 1981.

[88839] 15678.

[88840] Purtilo, D. T.; Bhawan, J.; Hutt, L. M.; De Nicola, L.; Szy-
manski,

[88841] I.; Yang, J. P. S.; Boto, W.; Naier, R.; Thorley-Lawson, D.:
Epstein-Barr

[88842] virus in the X-linked recessive lymphoproliferative syn-
drome. Lancet I:

[88843] 798-801, 1978.

[88844] 15679.

[88845] Purtilo, D. T.; Cassel, C. K.; Yang, J. P. S.: Fatal infectious

[88846] mononucleosis in familial lymphohistiocytosis. (Letter)
New Eng.

[88847] J. Med. 201: 736 only, 1974.

[88848] 15680.

[88849] Purtilo, D. T.; Cassel, C. K.; Yang, J. P. S.; Harper, R.;
Stephenson,

[88850] S. R.; Landing, B. H.; Vewter, G. F.: X-linked recessive pro-
gressive

[88851] combined variable immunodeficiency (Duncan's disease).

Lancet I:

[88852] 935–941, 1975.

[88853] 15681.

[88854] Purtilo, D. T.; DeFlorio, D., Jr.; Hutt, L. M.; Bhawan, J.;

Yang,

[88855] J. P. S.; Otto, R. L.; Edwards, W.: Variable phenotypic expression

[88856] of an X-linked recessive lymphoproliferative syndrome.

New Eng. J.

[88857] Med. 297: 1077–1081, 1977.

[88858] 15682.

[88859] Purtilo, D. T.; Grierson, H. L.: Methods of detection of new

[88860] families with X-linked lymphoproliferative disease. Cancer Genet.

[88861] Cytogenet. 51: 143–153, 1991.

[88862] 15683.

[88863] Purtilo, D. T.; Sakamoto, K.; Barnabei, V.; Seeley, J.; Bechtold,

[88864] T.; Rogers, G.; Yetz, J.; Harada, S.; the XLP collaborators: Epstein–Barr

- [88865] virus-induced diseases in boys with the X-linked lymphoproliferative
- [88866] syndrome (XLP): update on studies of the registry. *Am. J. Med.* 73:
- [88867] 49-56, 1982.
- [88868] 15684.
- [88869] Purtilo, D. T.; Yang, J. P. S.; Allegra, S.; DeFlorio, D.; Hutt,
- [88870] L. M.; Soltani, M.; Vawter, G. F.: Hematopathology and pathogenesis
- [88871] of the X-linked recessive lymphoproliferative syndrome. *Am. J. Med.* 62:
- [88872] 225-233, 1977.
- [88873] 15685.
- [88874] Sanger, W. G.; Grierson, H. L.; Skare, J.; Wyandt, H.; Pirruccello,
- [88875] S.; Fordyce, R.; Purtilo, D. T.: Partial Xq25 deletion in a family
- [88876] with the X-linked lymphoproliferative disease (XLP). *Cancer Genet.*
- [88877] *Cytogenet.* 47: 163-169, 1990.
- [88878] 15686.

- [88879] Sayos, J.; Wu, C.; Morra, M.; Wang, N.; Zhang, X.; Allen, D.;
- [88880] van Schaik, S.; Notarangelo, L.; Gehat, R.; Roncarolo, M. G.; Oettgen,
- [88881] H.; De Vries, J. E.; Aversall, G.; Terhorst, C.: The X-linked lymphoproliferative-disease
- [88882] gene product SAP regulates signals induced through the co-receptor
- [88883] SLAM. Nature 395: 462-469, 1998.
- [88884] 15687.
- [88885] Scher, I.: The CBA/N mouse strain: an experimental model illustrating
- [88886] the influence of the X-chromosome on immunity. Adv. Immun. 33: 1-71,
- [88887] 1982.
- [88888] 15688.
- [88889] Schuster, V.; Kreth, H. W.: X-linked lymphoproliferative disease.In:
- [88890] Ochs, H. D.; Smith, C. I. E.; Puck, J. M. (eds.): Primary Immunodeficiency
- [88891] Diseases: A Molecular and Genetic Approach. New York: Oxford University

[88892] Press 1999. Pp. 222–232.

[88893] 15689.

[88894] Seemayer, T. A.; Gross, T. G.; Egeler, R. M.; Pirruccello, S.

[88895] J.; Davis, D. J.; Kelly, C. M.; Okano, M.; Lanyi, A.; Sumegi, J.:

[88896] X-linked lymphoproliferative disease: twenty-five years after the

[88897] discovery. *Pediat. Res.* 38: 471–478, 1995.

[88898] 15690.

[88899] Skare, J.; Grierson, H.; Wyandt, H.; Sanger, W.; Milunsky, J.;

[88900] Purtilo, D.; Sullivan, J.; Milunsky, A.: Genetics of the X-linked

[88901] lymphoproliferative syndrome. (Abstract) *Am. J. Hum. Genet.* 45 (suppl.):

[88902] A161 only, 1989.

[88903] 15691.

[88904] Skare, J.; Madan, S.; Glaser, J.; Purtilo, D.; Nitowsky, H.; Pulijaal,

[88905] V.; Milunsky, A.: First prenatal diagnosis of X-linked lymphoproliferative

- [88906] disease. Am. J. Med. Genet. 44: 79–81, 1992.
- [88907] 15692.
- [88908] Skare, J.; Milunsky, A.; Byron, K.; Sullivan, J.: The mutation
- [88909] causing X-linked lymphoproliferative syndrome lies in Xq26. (Abstract) Am.
- [88910] J. Hum. Genet. 41: A185 only, 1987.
- [88911] 15693.
- [88912] Skare, J.; Wu, B.-L.; Madan, S.; Pulijaal, V.; Purtilo, D.;
- Haber,
- [88913] D.; Nelson, D.; Sylla, B.; Grierson, H.; Nitowsky, H.; Glaser, J.;
- [88914] Wissink, J.; White, B.; Holden, J.; Housman, D.; Lenoir, G.; Wyandt,
- [88915] H.; Milunsky, A.: Characterization of three overlapping deletions
- [88916] causing X-linked lymphoproliferative disease. Genomics 16: 254–255,
- [88917] 1993.
- [88918] 15694.
- [88919] Skare, J. C.; Grierson, H. L.; Sullivan, J. L.; Nussbaum, R. L.;

- [88920] Purtilo, D. T.; Sylla, B. S.; Lenoir, G. M.; Reilly, D. S.; White, B. N.; Milunsky, A.: Linkage analysis of seven kindreds with the
- [88922] X-linked lymphoproliferative syndrome (XLP) confirms that the XLP
- [88923] locus is near DXS42 and DXS37. Hum. Genet. 82: 354-358, 1989.
- [88924] 15695.
- [88925] Skare, J. C.; Milunsky, A.; Byron, K. S.; Sullivan, J. L.: Mapping
- [88926] the X-linked lymphoproliferative syndrome. Proc. Nat. Acad. Sci. 84: 2015-2018, 1987.
- [88927] 15696.
- [88928] 15696.
- [88929] Skare, J. C.; Sullivan, J. L.; Milunsky, A.: Mapping the mutation
- [88930] causing the X-linked lymphoproliferative syndrome in relation to restriction
- [88931] fragment length polymorphisms on Xq. Hum. Genet. 82: 349-353, 1989.
- [88932] 15697.

- [88933] Steinherz, R.; Levy, Y.; Litwin, A.; Nitzan, M.; Friedman, E.;
- [88934] Levin, S.: X-linked lymphoproliferative syndrome: a new
kindred with
- [88935] variable phenotypic expression. *Am. J. Dis. Child.* 139:
191–193,
- [88936] 1985.
- [88937] 15698.
- [88938] Nehlin, J. O.; Hara, E.; Kuo, W.-L.; Collins, C.; Campisi, J.:
- [88939] Genomic organization, sequence, and chromosomal local-
ization of the
- [88940] human helix-loop-helix Id1 gene. *Biochem. Biophys. Res.*
Commun. 231:
- [88941] 628–634, 1997.
- [88942] 15699.
- [88943] Inoue, S.; Orimo, A.; Matsuda, Y.; Inazawa, J.; Emi, M.;
Nakamura,
- [88944] Y.; Hori, T.; Muramatsu, M.: Chromosome mapping of hu-
man (ZNF147)
- [88945] and mouse genes for estrogen-responsive finger protein
(efp), a member
- [88946] of the RING finger family. *Genomics* 25: 581–583, 1995.

[88947] 15700.

[88948] Law, D. J.; Prasad, M. A.; King, S. E.; Spranger, K. D.; Lee, Y.

[88949] H.; Fox, R. E.; Collins, E. E.; Gebuhr, T. C.; Miller, D. E.; Petty,

[88950] E. M.: Localization of the human estrogen-responsive finger protein

[88951] (EFP) gene (ZNF147) within a YAC contig containing the myeloperoxidase

[88952] (MPO) gene. Genomics 28: 361–363, 1995.

[88953] 15701.

[88954] Fujii, H.; Chen, S.-H.; Akatsuka, J.; Miwa, S.; Yoshida, A.: Use

[88955] of cultured lymphoblastoid cells for the study of abnormal enzymes:

[88956] molecular abnormality of a phosphoglycerate kinase variant associated

[88957] with hemolytic anemia. Proc. Nat. Acad. Sci. 78: 2587–2590, 1981.

[88958] 15702.

[88959] Lindenbaum, R. H.; Clarke, G.; Patel, C.; Moncrieff, M.; Hughes,

[88960] J. T.: Muscular dystrophy in an X;1 translocation female suggests
[88961] that Duchenne locus is on X chromosome short arm. J.
Med. Genet. 16:
[88962] 389–392, 1979.
[88963] 15703.
[88964] Loftin, C. D.; Trivedi, D. B.; Tiano, H. F.; Clark, J. A.; Lee,
[88965] C. A.; Epstein, J. A.; Morham, S. G.; Breyer, M. D.; Nguyen,
M.; Hawkins,
[88966] B. M.; Goulet, J. L.; Smithies, O.; Koller, B. H.; Langenbach,
R.
[88967] : Failure of ductus arteriosus closure and remodeling in
neonatal
[88968] mice deficient in cyclooxygenase–1 and cyclooxygenase–2.
Proc. Nat.
[88969] Acad. Sci. 98: 1059–1064, 2001.
[88970] 15704.
[88971] Macchia, L.; Di Paola, R.; Guerrese, M.–C.; Chiechi, L. M.;
Tursi,
[88972] A.; Caiaffa, M. F.; Haeggstrom, J. Z.: Expression of
prostaglandin

- [88973] endoperoxide H synthase 1 and 2 in human placenta at term. Biochem.
- [88974] Biophys. Res. Commun. 233: 496–501, 1997.
- [88975] 15705.
- [88976] Morham, S. G.; Langenbach, R.; Loftin, C. D.; Tiano, H. F.; Vouloumanos,
- [88977] N.; Jennette, J. C.; Mahler, J. F.; Kluckman, K. D.; Ledford, A.;
- [88978] Lee, C. A.; Smithies, O.: Prostaglandin synthase 2 gene disruption
- [88979] causes severe renal pathology in the mouse. Cell 83: 473–482, 1995.
- [88980] 15706.
- [88981] Neufang, G.; Furstenberger, G.; Heidt, M.; Marks, F.; Muller-Decker,
- [88982] K.: Abnormal differentiation of epidermis in transgenic mice constitutively
- [88983] expressing cyclooxygenase–2 in skin. Proc. Nat. Acad. Sci. 98: 7629–7634,
- [88984] 2001.
- [88985] 15707.

[88986] O'Banion, M. K.; Sadowski, H. B.; Winn, V.; Young, D. A.: A serum-
[88987] and glucocorticoid-regulated 4-kilobase mRNA encodes a
cyclooxygenase-related
[88988] protein. J. Biol. Chem. 266: 23261-23267, 1991.
[88989] 15708.
[88990] O'Banion, M. K.; Winn, V. D.; Young, D. A.: cDNA cloning
and
[88991] functional activity of a glucocorticoid-regulated inflam-
matory cyclooxygenase. Proc.
[88992] Nat. Acad. Sci. 89: 4888-4892, 1992.
[88993] 15709.
[88994] Oshima, M.; Dinchuk, J. E.; Kargman, S. L.; Oshima, H.;
Hancock,
[88995] B.; Kwong, E.; Trzaskos, J. M.; Evans, J. F.; Taketo, M. M.:
Suppression
[88996] of intestinal polyposis in Apc(delta-716) knockout mice by
inhibition
[88997] of cyclooxygenase 2 (COX-2). Cell 87: 803-809, 1996.
[88998] 15710.
[88999] Salmenkivi, K.; Haglund, C.; Ristimaki, A.; Arola, J.;

Heikkila,

[89000] P.: Increased expression of cyclooxygenase-2 in malignant pheochromocytomas. J.

[89001] Clin. Endocr. Metab. 86: 5615-5619, 2001.

[89002] 15711.

[89003] Tay, A.; Squire, J. A.; Goldberg, H.; Skorecki, K.: Assignment

[89004] of the human prostaglandin-endoperoxide synthase 2 (PTGS2) gene to

[89005] 1q25 by fluorescence in situ hybridization. Genomics 23: 718-719,

[89006] 1994.

[89007] 15712.

[89008] Swan, S. K.; Rudy, D. W.; Lasseter, K. C.; Ryan, C. F.; Buechel,

[89009] K. L.; Lambrecht, L. J.; Pinto, M. B.; Dilzer, S. C.; Obrda, O.; Sundblad,

[89010] K. J.; Gumbs, C. P.; Ebel, D. L.; Quan, H.; Larson, P. J.; Schwartz,

[89011] J. I.; Musliner, T. A.; Gertz, B. J.; Brater, D. C.; Yao, S.-L.:

[89012] Effect of cyclooxygenase-2 inhibition on renal function in

elderly

[89013] persons receiving a low-salt diet: a randomized controlled trial. Ann.

[89014] Intern. Med. 133: 1-9, 2000.

[89015] 15713.

[89016] Tazawa, R.; Xu, X.-M.; Wu, K. K.; Wang, L.-H.: Characterization

[89017] of the genomic structure, chromosomal location and promoter of human

[89018] prostaglandin H synthase-2 gene. Biochem. Biophys. Res. Commun. 203:

[89019] 190-199, 1994.

[89020] 15714.

[89021] Tsujii, M.; DuBois, R. N.: Alterations in cellular adhesion and

[89022] apoptosis in epithelial cells overexpressing prostaglandin endoperoxide

[89023] synthase 2. Cell 83: 493-501, 1995.

[89024] 15715.

[89025] Zhou, X.-L.; Lei, Z. M.; Rao, C. V.: Treatment of human endometrial

- [89026] gland epithelial cells with chorionic gonadotropin/luteinizing hormone
- [89027] increases the expression of the cyclooxygenase-2 gene. J. Clin. Endocr.
- [89028] Metab. 84: 3364–3377, 1999.
- [89029] 15716.
- [89030] Inoue, S.; Orimo, A.; Hosoi, T.; Kondo, S.; Toyoshima, H.; Kondo,
- [89031] T.; Ikegami, A.; Ouchi, Y.; Orimo, H.; Muramatsu, M.: Genomic binding-site
- [89032] cloning reveals an estrogen-responsive gene that encodes a RING finger
- [89033] protein. Proc. Nat. Acad. Sci. 90: 11117–11121, 1993.
- [89034] 15717.
- [89035] Maho, A.; Bensimon, A.; Vassart, G.; Parmentier, M.: Mapping of
- [89036] the CCXCR1, CX3CR1, CCBP2 and CCR9 genes to the CCR cluster within
- [89037] the 3p21.3 region of the human genome. Cytogenet. Cell Genet. 87:
- [89038] 265–268, 1999.

[89039] 15718.

[89040] Anderson, D. M.; Johnson, L.; Glaccum, M. B.; Copeland, N. G.;

[89041] Gilbert, D. J.; Jenkins, N. A.; Valentine, V.; Kirstein, M. N.; Shapiro,

[89042] D. N.; Morris, S. W.; Grabstein, K.; Cosman, D.: Chromosomal assignment

[89043] and genomic structure of IL15. Genomics 25: 701-706, 1995.

[89044] 15719.

[89045] Grabstein, K. H.; Eisenman, J.; Shanebeck, K.; Rauch, C.; Srinivasan,

[89046] S.; Fung, V.; Beers, C.; Richardson, J.; Schoenborn, M. A.; Ahdieh,

[89047] M.; Johnson, L.; Alderson, M. R.; Watson, J. D.; Anderson, D. M.;

[89048] Giri, J. G.: Cloning of a T cell growth factor that interacts with

[89049] the beta chain of the interleukin-2 receptor. Science 264: 965-968,

[89050] 1994.

[89051] 15720.

- [89052] Roberts, A. I.; Lee, L.; Schwarz, E.; Groh, V.; Spies, T.; Ebert,
E. C.; Jabri, B.: Cutting edge: NKG2D receptors induced by IL-15
costimulate CD28-negative effector CTL in the tissue microenvironment. J.
Immun. 167: 5527-5530, 2001.
- [89056] 15721.
- [89057] Robinson, P.; Okhuysen, P. C.; Chappell, C. L.; Lewis, D. E.; Shahab,
I.; Lahoti, S.; White, A. C., Jr.: Expression of IL-15 and IL-4
in
IFN-gamma-independent control of experimental human
Cryptosporidium
parvum infection. Cytokine 15: 39-46, 2001.
- [89061] 15722.
- [89062] Chen, X.; Vinkemeier, U.; Zhao, Y.; Jeruzalmi, D.; Darnell,
J.
E., Jr.; Kuriyan, J.: Crystal structure of a tyrosine phosphorylated
STAT-1 dimer bound to DNA. Cell 93: 827-839, 1998.

- [89065] 15723.
- [89066] Copeland, N. G.; Gilbert, D. J.; Schindler, C.; Zhong, Z.;
Wen,
- [89067] Z.; Darnell, J. E., Jr.; Mui, A. L.-F.; Miyajima, A.; Quelle, F.
W.;
- [89068] Ihle, J. N.; Jenkins, N. A.: Distribution of the mammalian
Stat gene
- [89069] family in mouse chromosomes. Genomics 29: 225–228,
1995.
- [89070] 15724.
- [89071] Darnell, J. E., Jr.; Kerr, I. M.; Stark, G. M.: Jak–STAT path-
ways
- [89072] and transcriptional activation in response to IFNs and
other extracellular
- [89073] signaling proteins. Science 264: 1415–1421, 1994.
- [89074] 15725.
- [89075] Durbin, J. E.; Hackenmiller, R.; Simon, M. C.; Levy, D. E.:
Targeted
- [89076] disruption of the mouse Stat1 gene results in compro-
mised innate immunity
- [89077] to viral disease. Cell 84: 443–450, 1996.

- [89078] 15726.
- [89079] Haddad, B.; Pabon-Pena, C. R.; Young, H.; Sun, W. H.: Assignment
[89080] of STAT1 to human chromosome 2q32 by FISH and radiation hybrids. Cytogenet.
[89081] Cell Genet. 83: 58-59, 1998.
- [89082] 15727.
- [89083] Ihle, J. N.: STATs: signal transducers and activators of transcription. Cell 84:
[89084] 331-334, 1996.
- [89085] 15728.
- [89086] Meraz, M. A.; White, J. M.; Sheehan, K. C. F.; Bach, E. A.; Rodig,
[89087] S. J.; Dighe, A. S.; Kaplan, D. H.; Riley, J. K.; Greenlund, A. C.
- [89088] Campbell, D.; Carver-Moore, K.; DuBois, R. N.; Clark, R.; Aguet, M.;
- [89089] Schreiber, R. D.: Targeted disruption of the Stat1 gene in mice reveals
[89090] unexpected physiologic specificity in the JAK-STAT signaling pathway. Cell 84:

- [89091] 431–442, 1996.
- [89092] 15729.
- [89093] Mowen, K. A.; Tang, J.; Zhu, W.; Schurter, B. T.; Shuai, K.; Herschman,
- [89094] H. R.; David, M.: Arginine methylation of STAT1 modulates IFN- α /beta-induced
- [89095] transcription. *Cell* 104: 731–741, 2001.
- [89096] 15730.
- [89097] Ramana, C. V.; Chatterjee-Kishore, M.; Nguyen, H.; Stark, G. R.
- [89098] : Complex roles of Stat1 in regulating gene expression. *Oncogene* 19:
- [89099] 2619–2627, 2000.
- [89100] 15731.
- [89101] Yamamoto, K.; Kobayashi, H.; Arai, A.; Miura, O.; Hiro-sawa, S.;
- [89102] Miyasaka, N.: cDNA cloning, expression and chromosome mapping of
- [89103] the human STAT4 gene: both STAT4 and STAT1 genes are mapped to 2q32.2–q32.3. *Cytogenet.*
- [89104] *Cell Genet.* 77: 207–210, 1997.

- [89105] 15732.
- [89106] Fukuda, M.: Biogenesis of the lysosomal membrane. Sub-cell. Biochem. 22:
[89107] 199–230, 1994.
- [89108] 15733.
- [89109] Fukuda, M.; Viitala, J.; Matteson, J.; Carlsson, S. R.:
Cloning
[89110] of the cDNAs encoding human lysosomal membrane glycoproteins, h-lamp-1
[89111] and h-lamp-2: comparison of their deduced amino acid sequences. J.
[89112] Biol. Chem. 263: 18920–18928, 1988.
- [89113] 15734.
- [89114] Konecki, D. S.; Foetisch, K.; Schlotter, M.; Lichter-Konecki, U.
[89115] : Complete cDNA sequence of human lysosome-associated membrane protein-2. Biochem.
[89116] Biophys. Res. Commun. 205: 1–5, 1994.
- [89117] 15735.
- [89118] Lindsay, S.; Bird, A. P.: Use of restriction enzymes to detect

- [89119] potential gene sequences in mammalian DNA. *Nature* 327: 336–338,
- [89120] 1987.
- [89121] 15736.
- [89122] Manoni, M.; Tribioli, C.; Lazzari, B.; DeBellis, G.; Patrosso,
- [89123] C.; Pergolizzi, R.; Pellegrini, M.; Maestrini, E.; Rivella, S.; Vezzoni,
- [89124] P.; Toniolo, D.: The nucleotide sequence of a CpG island demonstrates
- [89125] the presence of the first exon of the gene encoding the human lysosomal
- [89126] membrane protein LAMP2 and assigns the gene to Xq24. *Genomics* 9: 551–554, 1991.
- [89127] 15737.
- [89128] 15737.
- [89129] Takahashi, M.; Yamamoto, A.; Takano, K.; Sudo, A.; Wada, T.; Goto,
- [89130] Y. I.; Nishino, I.; Saitoh, S.: Germline mosaicism of a novel mutation
- [89131] in lysosome-associated membrane protein-2 deficiency (Danon disease). *Ann.*

- [89132] Neurol. 52: 122–125, 2002.
- [89133] 15738.
- [89134] Tanaka, Y.; Guhe, G.; Suter, A.; Eskelinen, E.–L.; Hartmann, D.;
- [89135] Lullmann–Rauch, R.; Janssen, P. M. L.; Blanz, J.; von Figura, K.;
- [89136] Saftig, P.: Accumulation of autophagic vacuoles and cardiomyopathy
- [89137] in LAMP–2–deficient mice. Nature 406: 902–906, 2000.
- [89138] 15739.
- [89139] Falls, H. F.: The role of the sex chromosome in hereditary ocular
- [89140] pathology. Trans. Am. Ophthal. Soc. 50: 421–467, 1952.
- [89141] 15740.
- [89142] Rosse, W. F.: Paroxysmal nocturnal hemoglobinuria: the biochemical
- [89143] defects and the clinical syndrome. Blood Rev. 3: 192–200, 1989.
- [89144] 15741.
- [89145] Rotoli, B.; Luzzatto, L.: Paroxysmal nocturnal haemoglobinuria. Clin.

- [89146] Haemat. 2: 113–138, 1989.
- [89147] 15742.
- [89148] Sauer, B.; Henderson, N.: Site-specific DNA recombination
in
- [89149] mammalian cells by the Cre recombinase of bacteriophage
P1. Proc.
- [89150] Nat. Acad. Sci. 85: 5166–5170, 1988.
- [89151] 15743.
- [89152] Savoia, A.; Ianzano, L.; Lunardi, C.; De Sandre, G.;
Carotenuto,
- [89153] M.; Musto, P.; Zelante, L.: Identification of three novel mu-
tations
- [89154] in the PIG-A gene in paroxysmal nocturnal haemoglobin-
uria (PNH) patients. Hum.
- [89155] Genet. 97: 45–48, 1996.
- [89156] 15744.
- [89157] Socie, G.; Mary, J.-Y.; de Gramont, A.; Rio, B.; Leporrier,
M.;
- [89158] Rose, C.; Heudier, P.; Rochant, H.; Cahn, J.-Y.; Gluckman,
E.: Paroxysmal
- [89159] nocturnal haemoglobinuria: long-term follow-up and

prognostic factors. Lancet 348:

[89160] 573–577, 1996.

[89161] 15745.

[89162] Stevens, V. L.; Raetz, C. R. H.: Defective glycosyl phosphatidylinositol

[89163] biosynthesis in extracts of three thy-1 negative lymphoma cell mutants. J.

[89164] Biol. Chem. 266: 10039–10042, 1991.

[89165] 15746.

[89166] Sugiyama, E.; DeGasperi, R.; Urakaze, M.; Chang, H.-M.; Thomas,

[89167] L. J.; Hyman, R.; Warren, C. D.; Yeh, E. T. H.: Identification of

[89168] defects in glycosylphosphatidylinositol anchor biosynthesis in the

[89169] thy-1 expression mutants. J. Biol. Chem. 266: 12119–12122, 1991.

[89170] 15747.

[89171] Takahashi, M.; Takeda, J.; Hirose, S.; Hyman, R.; Inoue, N.; Miyata,

[89172] T.; Ueda, E.; Kitani, T.; Medof, M. E.; Kinoshita, T.: Defi-

cient

- [89173] biosynthesis of N-acetylglucosaminyl-phosphatidylinositol, the first intermediate of glycosyl phosphatidylinositol anchor biosynthesis,
- [89174] in cell lines established from patients with paroxysmal nocturnal hemoglobinuria. J. Exp. Med. 177: 517-521, 1993.
- [89175] 15748.
- [89176] Takeda, J.; Miyata, T.; Kawagoe, K.; Iida, Y.; Endo, Y.; Fujita, T.; Takahashi, M.; Kitani, T.; Kinoshita, T.: Deficiency of the GPI anchor caused by a somatic mutation of the PIG-A gene in paroxysmal nocturnal hemoglobinuria. Cell 73: 703-711, 1993.
- [89177] 15749.
- [89178] Ueda, E.; Nishimura, J.; Kitani, T.; Nasu, K.; Kageyama, T.; Kim, Y. U.; Takeda, J.; Kinoshita, T.: Deficient surface expression of glycosylphosphatidylinositol-anchored proteins in B cell

lines established

[89186] from patients with paroxysmal nocturnal hemoglobinuria.

Int. Immun. 4:

[89187] 1263–1271, 1992.

[89188] 15750.

[89189] Ware, R. E.; Howard, T. A.; Kamitani, T.; Chang, H.–M.;

Yeh, E.

[89190] T. H.; Seldin, M. F.: Chromosomal assignment of genes involved in

[89191] glycosylphosphatidylinositol anchor biosynthesis: implications for

[89192] the pathogenesis of paroxysmal nocturnal hemoglobinuria. Blood 83:

[89193] 3753–3757, 1994.

[89194] 15751.

[89195] Ware, R. E.; Rosse, W. F.; Howard, T. A.: Mutations within the

[89196] Piga gene in patients with paroxysmal nocturnal hemoglobinuria. Blood 83:

[89197] 2418–2422, 1994.

[89198] 15752.

- [89199] Watanabe, R.; Inoue, N.; Westfall, B.; Taron, C. H.; Orlean, P.;
- [89200] Takeda, J.; Kinoshita, T.: The first step of glycosylphosphatidylinositol
- [89201] biosynthesis is mediated by a complex of PIG-A, PIG-H, PIG-C and GPI1. EMBO
- [89202] J. 17: 877-885, 1998.
- [89203] 15753.
- [89204] Watanabe, R.; Kinoshita, T.; Masaki, R.; Yamamoto, A.; Takeda,
- [89205] J.; Inoue, N.: PIG-A and PIG-H, which participate in glycosylphosphatidylinositol
- [89206] anchor biosynthesis, form a protein complex in the endoplasmic reticulum. J.
- [89207] Biol. Chem. 271: 26868-26875, 1996.
- [89208] 15754.
- [89209] Yeh, E. T. H.; Rosse, W. F.: Paroxysmal nocturnal hemoglobinuria
- [89210] and the glycosylphosphatidylinositol anchor. J. Clin. Invest. 93:
- [89211] 2305-2310, 1994.

- [89212] 15755.
- [89213] Yu, J.; Nagarajan, S.; Ueda, E.; Knez, J. J.; Petersen, R. B.;
- [89214] Medof, M. E.: Characterization of alternatively spliced PIG-A transcripts
- [89215] in normal and paroxysmal nocturnal hemoglobinuria cells. Brazilian
- [89216] J. Med. Biol. Res. 27: 195–201, 1994.
- [89217] 15756.
- [89218] Chen, S.-H.; Malcolm, L. A.; Yoshida, A.; Giblett, E. R.: Phosphoglycerate
- [89219] kinase: an X-linked polymorphism in man. Am. J. Hum. Genet. 23:
- [89220] 87–91, 1971.
- [89221] 15757.
- [89222] Cooper, D. W.; Johnston, P. G.; Murtagh, C. E.; Sharman, G. B.;
- [89223] Vandeberg, J. L.; Poole, W. E.: Sex-linked isozymes and sex-chromosome
- [89224] evolution and inactivation in kangaroos. In: Markert, C. L.: Isozymes.
- [89225] Developmental Biology. New York: Academic Press (pub.)

III: 1975.

[89226] Pp. 559–573.

[89227] 15758.

[89228] Cooper, D. W.; Vandeberg, J. L.; Sharman, G. B.; Poole, W.
E.:

[89229] Phosphoglycerate kinase polymorphism in kangaroos provides further

[89230] evidence for paternal inactivation. Nature N.B. 230:
155–157, 1971.

[89231] 15759.

[89232] Deys, B. F.; Grzeschik, K.–H.; Grzeschik, A.; Jaffe, E. R.;
Siniscalco,

[89233] M.: Human phosphoglycerate kinase and inactivation of
the X chromosome. Science 175:

[89234] 1002–1003, 1972.

[89235] 15760.

[89236] DiMauro, S.; Dalakas, M.; Miranda, A. S.: Phosphoglycerate
kinase

[89237] deficiency: another cause of recurrent myoglobinuria.
Ann. Neurol. 13:

[89238] 11–19, 1983.

[89239] 15761.

[89240] Echard, G.; Gillois, M.: G6PD--alpha-GAL--PGK--HPRT
synteny in

[89241] the rabbit, *Oryctolagus cuniculus*. (Abstract) Cytogenet.
Cell Genet. 25:

[89242] 148-149, 1979.

[89243] 15762.

[89244] Patil, S. R.; Bartley, J. A.; Murray, J. C.; Ionasescu, V. V.;

[89245] Pearson, P. L.: X-linked glycerol kinase, adrenal hypopla-
sia and

[89246] myopathy maps at Xp21. (Abstract) Cytogenet. Cell Genet.
40: 720-721,

[89247] 1985.

[89248] 15763.

[89249] Kamal, A.; Stokin, G. B.; Yang, Z.; Xia, C.; Goldstein, L. S.:

[89250] Axonal transport of amyloid precursor protein is mediated
by direct

[89251] binding to the kinesin light chain subunit of kinesin-I.
Neuron 28:

[89252] 449-459, 2000.

[89253] 15764.

[89254] Ahn, A. H.; Yoshida, M.; Anderson, M. S.; Feener, C. A.;
Selig,

[89255] S.; Hagiwara, Y.; Ozawa, E.; Kunkel, L. M.: Cloning of hu-
man basic

[89256] A1, a distinct 59-kDa dystrophin-associated protein en-
coded on chromosome

[89257] 8q23-24. Proc. Nat. Acad. Sci. 91: 4446-4450, 1994.

[89258] 15765.

[89259] Yang, B.; Ibraghimov-Beskrovnaya, O.; Moomaw, C. R.;
Slaughter,

[89260] C. A.; Campbell, K. P.: Heterogeneity of the 59-kDa dys-
trophin-associated

[89261] protein revealed by cDNA cloning and expression. J. Biol.
Chem. 269:

[89262] 6040-6044, 1994.

[89263] 15766.

[89264] Merendino, J. J., Jr.; Spiegel, A. M.; Crawford, J. D.;
O'Carroll,

[89265] A.-M.; Brownstein, M. J.; Lolait, S. J.: A mutation in the va-
sopressin

- [89266] V2-receptor gene in a kindred with X-linked nephrogenic diabetes insipidus. New
- [89267] Eng. J. Med. 328: 1538-1541, 1993.
- [89268] 15767.
- [89269] Moses, A. M.; Sangani, G.; Miller, J. L.: Proposed cause of marked
- [89270] vasopressin resistance in a female with an X-linked recessive V2 receptor
- [89271] abnormality. J. Clin. Endocr. Metab. 80: 1184-1186, 1995.
- [89272] 15768.
- [89273] Nakano, K. K.: Familial nephrogenic diabetes insipidus. Hawaii
- [89274] Med. J. 28: 205-208, 1969.
- [89275] 15769.
- [89276] Nomura, Y.; Onigata, K.; Nagashima, T.; Yutani, S.; Mochizuki,
- [89277] H.; Nagashima, K.; Morikawa, A.: Detection of skewed X-inactivation
- [89278] in two female carriers of vasopressin type 2 receptor gene mutation. J.
- [89279] Clin. Endocr. Metab. 82: 3434-3437, 1997.

- [89280] 15770.
- [89281] Pan, Y.; Metzenberg, A.; Das, S.; Jing, B.; Gitschier, J.: Mutations
- [89282] in the V2 vasopressin receptor gene are associated with X-linked nephrogenic
- [89283] diabetes insipidus. Nature Genet. 2: 103–106, 1992.
- [89284] 15771.
- [89285] Pasel, K.; Schulz, A.; Timmermann, K.; Linnemann, K.; Hoeltzenbein,
- [89286] M.; Jaaskelainen, J.; Gruters, A.; Filler, G.; Schoneberg, T.: Functional
- [89287] characterization of the molecular defects causing nephrogenic diabetes
- [89288] insipidus in eight families. J. Clin. Endocr. Metab. 85: 1703–1710,
- [89289] 2000.
- [89290] 15772.
- [89291] Rosenthal, W.; Seibold, A.; Antaramian, A.; Lonergan, M.; Arthus,
- [89292] M.-F.; Hendy, G. N.; Birnbaumer, M.; Bichet, D. G.: Molecular identification

- [89293] of the gene responsible for congenital nephrogenic diabetes insipidus. *Nature* 359:
- [89294] 233–235, 1992.
- [89295] 15773.
- [89296] Sadeghi, H.; Robertson, G. L.; Bichet, D. G.; Innamorati, G.;
- [89297] Birnbaumer, M.: Biochemical basis of partial nephrogenic diabetes
- [89298] insipidus phenotypes. *Molec. Endocr.* 11: 1806–1813, 1997.
- [89299] 15774.
- [89300] Sadeghi, H. M.; Innamorati, G.; Birnbaumer, M.: An X-linked NDI
- [89301] mutation reveals a requirement for cell surface V2R expression. *Molec.*
- [89302] *Endocr.* 11: 706–713, 1997.
- [89303] 15775.
- [89304] Sawin, C. T.: Endocrine classic. *Endocrinologist* 8: 143–148,
- [89305] 1998.
- [89306] 15776.

- [89307] Schoneberg, T.; Sandig, V.; Wess, J.; Gudermann, T.; Schultz,
G.: Reconstitution of mutant V2 vasopressin receptors by
adenovirus-mediated
gene transfer: molecular basis and clinical implication. J.
Clin. Invest. 100: 1547-1556, 1997.
15777.
- [89312] Schoneberg, T.; Schulz, A.; Biebermann, H.; Gruters, A.;
Grimm,
T.; Hubschmann, K.; Filler, G.; Gudermann, T.; Schultz, G.:
V2 vasopressin
receptor dysfunction in nephrogenic diabetes insipidus
caused by different
molecular mechanisms. Hum. Mutat. 12: 196-205, 1998.
15778.
- [89317] Nathan, D. G.; Baehner, R. L.; Weaver, D. K.: Failure of ni-
tro
blue tetrazolium reduction in the phagocytic vacuoles of
leukocytes
in chronic granulomatous disease. J. Clin. Invest. 48:
1895-1904,

- [89320] 1969.
- [89321] 15779.
- [89322] Nelson, C. E.; Dahl, M. V.; Goltz, R. W.: Arcuate dermal erythema
- [89323] in a carrier of chronic granulomatous disease. Arch. Derm. 113:
- [89324] 789–800, 1977.
- [89325] 15780.
- [89326] Newburger, P. E.; Cohen, H. J.; Rothchild, S. B.; Hobbins, J.
- [89327] C.; Malawista, S. E.; Mahoney, M. J.: Prenatal diagnosis of chronic
- [89328] granulomatous disease. New Eng. J. Med. 300: 178–181, 1979.
- [89329] 15781.
- [89330] Noack, D.; Heyworth, P. G.; Kyono, W.; Cross, A. R.: A second
- [89331] case of somatic triple mosaicism in the CYBB gene causing chronic
- [89332] granulomatous disease. Hum. Genet. 109: 234–238, 2001.
- [89333] 15782.

- [89334] Okamura, N.; Malawista, S. E.; Roberts, R. L.; Rosen, H.; Ochs, H. D.; Babior, B. M.; Curnutte, J. T.: Phosphorylation of the oxidase-related 48K phosphoprotein family in the unusual cytochrome-negative and X-linked cytochrome-positive types of chronic granulomatous disease. Blood 72: 811-816, 1988.
- [89339] 15783.
- [89340] Orkin, S. H.: Stem cell alchemy. Nature Med. 6: 1212-1213, 2000.
- [89342] 15784.
- [89343] Orkin, S. H.: X-linked chronic granulomatous disease: from chromosomal position to the in vivo gene product. Trends Genet. 3: 149-151, 1987.
- [89346] 15785.
- [89347] Orkin, S. H.: Molecular genetics of chronic granulomatous

disease. Annu.

[89348] Rev. Immun. 7: 277–307, 1989.

[89349] 15786.

[89350] Patino, P. J.; Perez, J. E.; Lopez, J. A.; Condino-Neto, A.; Grumach,

[89351] A. S.; Botero, J. H.; Curnutte, J. T.; Garcia de Olarte, D.:
Molecular

[89352] analysis of chronic granulomatous disease caused by defects in gp91-phox. Hum.

[89353] Mutat. 13: 29–37, 1999.

[89354] 15787.

[89355] Pollock, J. D.; Williams, D. A.; Gifford, M. A. C.; Li, L. L.;

[89356] Du, X.; Fisherman, J.; Orkin, S. H.; Doerschuk, C. M.; Din-
auer, M.

[89357] C.: Mouse model of X-linked chronic granulomatous disease, an inherited

[89358] defect in phagocyte superoxide production. Nature Genet.
9: 202–209,

[89359] 1995.

[89360] 15788.

[89361] Porter, C. D.; Parkar, M. H.; Levinsky, R. J.; Collins, M. K.

[89362] L.; Kinnon, C.: X-linked chronic granulomatous disease: correction

[89363] of NADPH oxidase defect by retrovirus-mediated expression of gp91-phox. Blood 82:

[89364] 2196-2202, 1993.

[89365] 15789.

[89366] Quie, P. G.; White, J. G.; Holmes, B.; Good, R. A.: In vitro

[89367] bactericidal capacity of human polymorphonuclear leukocytes: diminished

[89368] activity in chronic granulomatous disease of childhood. J. Clin.

[89369] Invest. 46: 668-679, 1967.

[89370] 15790.

[89371] Rae, J.; Newburger, P. E.; Dinanuer, M. C.; Noack, D.; Hopkins,

[89372] P. J.; Kuruto, R.; Curnutte, J. T.: X-linked chronic granulomatous

[89373] disease: mutations in the CYBB gene encoding the gp91-phox component

[89374] of respiratory-burst oxidase. Am. J. Hum. Genet. 62: 1320-1331,

[89375] 1998.

[89376] 15791.

[89377] Roos, D.: The genetic basis of chronic granulomatous disease. Immun.

[89378] Rev. 138: 121–157, 1994.

[89379] 15792.

[89380] Roos, D.; de Boer, M.; Kuribayashi, F.; Meischl, C.; Weening,

[89381] R. S.; Segal, A. W.; Ahlin, A.; Nemet, K.; Hossle, J. P.; Bernatowska-Matuszkiewicz,

[89382] E.; Middleton-Price, H.: Mutations in the X-linked and autosomal

[89383] recessive forms of chronic granulomatous disease. Blood 87: 1663–1681,

[89384] 1996.

[89385] 15793.

[89386] Royer-Pokora, B.; Kunkel, L. M.; Monaco, A. P.; Goff, S. C.; Newburger,

[89387] P. E.; Baehner, R. L.; Cole, F. S.; Curnutte, J. T.; Orkin, S. H.

[89388] : Cloning the gene for an inherited human disorder—chronic granulomatous

- [89389] disease--on the basis of its chromosomal location. *Nature* 322: 32-38,
- [89390] 1986.
- [89391] 15794.
- [89392] Schmalzer, E. A.; Miller, D. R.: Chronic granulomatous disease. *Prog. Med. Genet.* 1: 145-184, 1976.
- [89393] 15795.
- [89394] Segal, A. W.: Cytochrome b(-245) and its involvement in the molecular
- [89395] pathology of chronic granulomatous disease. *Hemat. Oncol. Clin. North Am.* 2: 213-223, 1988.
- [89396]
- [89397] 100. Segal, A. W.; Cross, A. R.; Garcia, R. C.; Borregaard, N.; Valerius,
- [89398] N. H.; Soothill, J. F.; Jones, O. T. G.: Absence of cytochrome b(-245)
- [89399] in chronic granulomatous disease: a multicenter European evaluation
- [89400] of its incidence and relevance. *New Eng. J. Med.* 308:

245–251, 1983.

[89403]

[89404] 101. Segal, A. W.; Peters, T. J.: Characterisation of the enzyme

[89405] defect in chronic granulomatous disease. Lancet I: 1363–1365, 1976.

[89406]

[89407] 102. Segal, A. W.; Webster, D.; Jones, O. T. G.; Allison, A. C.:

[89408] Absence of a newly described cytochrome b from neutrophils of patients

[89409] with chronic granulomatous disease. Lancet II: 446–449, 1978.

[89410]

[89411] 103. Soothill, J. F.: Inheritance of chronic granulomatous disease.

[89412] (Letter) Lancet II: 543–544, 1969.

[89413]

[89414] 104. Southwick, F. S.; van der Meer, J. W. M.: Recurrent cystitis

[89415] and bladder mass in two adults with chronic granuloma-

tous disease. Ann.

[89416] Intern. Med. 109: 118–121, 1988.

[89417]

[89418] 105. Stasia, M. J.; Lardy, B.; Maturana, A.; Rousseau, P.;
Martel,

[89419] C.; Bordigoni, P.; Demaurex, N.; Morel, F.: Molecular and
functional

[89420] characterization of a new X-linked chronic granulomatous
disease variant

[89421] (X91+) case with a double missense mutation in the cy-
tosolic gp91phox

[89422] C-terminal tail. Biochim. Biophys. Acta 1586: 316–330,
2002.

[89423]

[89424] 106. Taswell, H. F.; Lewis, J. C.; Marsh, W. L.; Wimer, B. M.;
Pineda,

[89425] A. A.; Brzica, S. M., Jr.: Erythrocyte morphology in genetic
defects

[89426] of the Rh and Kell blood group systems. Mayo Clin. Proc.
52: 157–159,

[89427] 1977.

[89428]

- [89429] 107. Tauber, A. I.; Borregaard, N.; Simons, E.; Wright, J.:
Chronic
[89430] granulomatous disease: a syndrome of phagocyte oxidase
deficiencies. Medicine 62:
[89431] 286–309, 1983.
[89432]
[89433] 108. Teahan, C.; Rowe, P.; Parker, P.; Totty, N.; Segal, A.
W.: The
[89434] X-linked chronic granulomatous disease gene codes for
the beta-chain
[89435] of cytochrome b(–245). Nature 327: 720–721, 1987.
[89436]
[89437] 109. Thompson, E. N.; Chandra, R. K.; Cope, W. A.;
Soothill, J. F.
[89438] : Leukocyte abnormality in both parents of a patient with
chronic
[89439] granulomatous disease. Lancet I: 799–800, 1969.
[89440]
[89441] 110. Thompson, E. N.; Soothill, J. F.: Chronic granuloma-
tous disease:
[89442] quantitative clinicopathological relationships. Arch. Dis.

Child. 45:

[89443] 24–32, 1970.

[89444]

[89445] 111. Tsuda, M.; Kaneda, M.; Sakiyama, T.; Inana, I.;
Owada, M.; Kiryu,

[89446] C.; Shiraishi, T.; Kakinuma, K.: A novel mutation at a
probable heme-binding

[89447] ligand in neutrophil cytochrome b(558) in atypical X-
linked chronic

[89448] granulomatous disease. Hum. Genet. 103: 377–381,
1998.

[89449]

[89450] 112. Windhorst, D. B.: Inheritance of chronic granuloma-
tous disease.

[89451] (Letter) Lancet II: 543–544, 1969.

[89452]

[89453] 113. Windhorst, D. B.; Holmes, B.; Good, R. A.: A newly
defined X-linked

[89454] trait in man with demonstration of the Lyon effect in car-
rier females. Lancet I:

[89455] 737–739, 1967.

[89456]

[89457] 114. Wolff, G.; Muller, C. R.; Jobke, A.: Linkage of genes
for chronic

[89458] granulomatous disease and Xg. Hum. Genet. 54: 269–271,
1980.

[89459] 15796.

[89460] Yen, P. H.; Ellison, J.; Salido, E. C.; Mohandas, T.; Shapiro,

[89461] L.: Isolation of a new gene from the distal short arm of the
human

[89462] X chromosome that escapes X-inactivation. Hum. Molec.
Genet. 1:

[89463] 47–52, 1992.

[89464] 15797.

[89465] Abrams, E. S.; Murdaugh, S. E.; Lerman, L. S.: Comprehen-
sive detection

[89466] of single base changes in human genomic DNA using de-
naturing gradient

[89467] gel electrophoresis and a GC clamp. Genomics 7:
463–475, 1990.

[89468] 15798.

[89469] Acquila, M.; Caprino, D.; Pecorara, M.; Baudo, F.; Morfini,

M.;

[89470] Mori, P. G.: Two novel mutations at 373 codon of FVIII gene detected

[89471] by DGGE. Thromb. Haemost. 69: 392–393, 1993.

[89472] 15799.

[89473] Alexander, B.; Goldstein, R.: Dual hemostatic defect in pseudohemophilia.

[89474] (Abstract) J. Clin. Invest. 32: 551 only, 1953.

[89475] 15800.

[89476] Aly, A. M.; Arai, M.; Hoyer, L. W.: Cysteamine enhances the procoagulant

[89477] activity of factor VIII–East Hartford, a dysfunctional protein due

[89478] to a light chain thrombin cleavage site mutation (arginine–1689 to

[89479] cysteine). J. Clin. Invest. 89: 1375–1381, 1992.

[89480] 15801.

[89481] Muller, C. R.; Migl, B.; Ropers, H.–H.; Happle, R.: Heterozygote

[89482] detection in steroid sulphatase deficiency. (Letter) Lancet I: 546–547,

- [89483] 1980.
- [89484] 15802.
- [89485] Muller, C. R.; Wahlstrom, J.; Ropers, H.-H.: Further evidence
- [89486] for the assignment of the steroid sulfatase X-linked ichthyosis locus
- [89487] to the telomer of Xp. (Letter) Hum. Genet. 58: 446, 1981.
- [89488] 15803.
- [89489] Munke, M.; Kruse, K.; Goos, M.; Ropers, H. H.; Tolksdorf, M.:
- [89490] Genetic heterogeneity of the ichthyosis, hypogonadism, mental retardation,
- [89491] and epilepsy syndrome: clinical and biochemical investigations on
- [89492] two patients with Rud syndrome and review of the literature. Europ.
- [89493] J. Pediat. 141: 8-13, 1983.
- [89494] 15804.
- [89495] Nomura, K.; Nakano, H.; Umeki, K.; Harada, K.; Kon, A.; Tamai,
- [89496] K.; Sawamura, D.; Hashimoto, I.: A study of the steroid

sulfatase

[89497] gene in families with X-linked ichthyosis using polymerase chain reaction. *Acta*

[89498] *Derm. Venereol.* 75: 340-342, 1995.

[89499] 15805.

[89500] Okano, M.; Kitano, Y.; Nakamura, T.; Matsuzawa, Y.: Detection

[89501] of heterozygotes of X-linked ichthyosis by measuring steroid sulphatase

[89502] activity of lymphocytes: mode of inheritance in three families. *Brit.*

[89503] *J. Derm.* 113: 645-649, 1985.

[89504] 15806.

[89505] Orel, H.: Die Vererbung der Ichthyosis congenita und der Ichthyosis

[89506] vulgaris. *Z. Kinderheilk.* 47: 312-340, 1929.

[89507] 15807.

[89508] Passarge, E.; Post, B.; Schopf, E.: Possible genetic heterogeneity

[89509] of X-linked ichthyosis. *Birth Defects Orig. Art. Ser.* VII(8): 46-49,

[89510] 1971.

[89511] 15808.

[89512] Pike, M. G.; Hammerton, M.; Edge, J.; Atherton, D. J.;
Grant,

[89513] D. B.: A family with X-linked ichthyosis and hypogonadism. Europ.

[89514] J. Pediat. 148: 442-444, 1989.

[89515] 15809.

[89516] Ropers, H.-H.; Migl, B.; Zimmer, J.; Fraccaro, M.; Maraschio,

[89517] P. P.; Westerveld, A.: Activity of steroid sulfatase in fibroblasts

[89518] with numerical and structural X chromosome aberrations. Hum. Genet. 57:

[89519] 354-356, 1981.

[89520] 15810.

[89521] Ropers, H.-H.; Wiberg, U.: Evidence for X-linkage and non-inactivation

[89522] of steroid sulphatase locus in wood lemming. Nature 296: 766-767,

[89523] 1982.

[89524] 15811.

[89525] Ross, J. B.; Allderdice, P. W.; Shapiro, L. J.; Aveling, J.;
Eales,

[89526] B. A.; Simms, D., Jr.: Familial X-linked ichthyosis, steroid
sulfatase

[89527] deficiency, mental retardation, and nullisomy for
Xp223-pter. Arch.

[89528] Derm. 121: 1524-1528, 1985.

[89529] 15812.

[89530] Salido, E. C.; Li, X. M.; Yen, P. H.; Martin, N.; Mohandas, T.

[89531] K.; Shapiro, L. J.: Cloning and expression of the mouse
pseudoautosomal

[89532] steroid sulphatase gene (Sts). Nature Genet. 13: 83-86,
1996.

[89533] 15813.

[89534] Schlamadinger, J.; Meyer, J. C.; Vajda, I.; Szabo, G.: X-
linked

[89535] recessive ichthyosis: reinvestigation of a family first de-
scribed

[89536] in 1928. Dermatologica 175: 217-223, 1987.

[89537] 15814.

[89538] Schmickel, R. D.: Chromosomal deletions and enzyme deficiencies. J.

[89539] *Pediat.* 108: 244–246, 1986.

[89540]

[89541] 100. Schnyder, U. W.: Inherited ichthyoses. *Arch. Derm.* 102: 240–252,

[89542] 1970.

[89543]

[89544] 101. Sever, R. J.; Frost, P.; Weinstein, G.: Eye changes in ichthyosis. *J.A.M.A.* 206:

[89545] 2283–2286, 1968.

[89546]

[89547] 102. Shapiro, L. J.: Personal Communication. Los Angeles, Calif.

[89548] 3/1987.

[89549]

[89550] 103. Shapiro, L. J.: Personal Communication. San Francisco, Calif.

[89551] 5/20/1997.

[89552]

[89553] 104. Shapiro, L. J.: Personal Communication. Torrance, Calif.

[89554] 1977.

[89555]

[89556] 105. Shapiro, L. J.; Mohandas, T.; Weiss, R.; Romeo, G.: Non-activation

[89557] of a X-chromosome locus in man. Science 204: 1224-1226, 1979.

[89558]

[89559] 106. Shapiro, L. J.; Weiss, R.: Diminished cholesterol sulfatase

[89560] activity in fibroblasts of placental sulfatase deficiency patients.

[89561] (Abstract) Vth Int. Conf. on Birth Defects, Montreal , 8/1977.

[89562]

[89563] 107. Shapiro, L. J.; Weiss, R.; Buxman, M. M.; Vidgoff, J.; Dimond,

[89564] R. L.: Enzymatic basis of typical X-linked ichthyosis. Lancet II:

[89565] 756-757, 1978.

[89566]

[89567] 108. Shapiro, L. J.; Weiss, R.; Webster, D.; France, J. T.: X-linked

[89568] ichthyosis due to steroid-sulphatase deficiency. Lancet I: 70-72,

[89569] 1978.

[89570]

[89571] 109. Shapiro, L. J.; Yen, P.; Pomerantz, D.; Martin, E.; Rolewic,

[89572] L.; Mohandas, T.: Molecular studies of deletions at the human steroid

[89573] sulfatase locus. Proc. Nat. Acad. Sci. 86: 8477-8481, 1989.

[89574]

[89575] 110. Shapiro, L. J.; Yen, P. H.: Reply to letter of Andria et al.,

[89576] 1987. (Letter) Ann. Neurol. 22: 99, 1987.

[89577]

[89578] 111. Shapiro, L. J.; Yen, P. H.; Marsh, B.; Mohandas, T.: Frequent

[89579] deletions at the steroid sulfatase (STS) locus. (Abstract)

Am. J.

[89580] Hum. Genet. 41: A238, 1987.

[89581]

[89582] 112. Solomon, I. L.; Schoen, E. J.: Sex-linked ichthyosis in XO gonadal

[89583] dysgenesis. (Letter) Lancet I: 1304-1305, 1971.

[89584]

[89585] 113. Stein, C.; Hille, A.; Seidel, J.; Rijnbout, S.; Waheed, A.; Schmidt,

[89586] B.; Geuze, H.; von Figura, K.: Cloning and expression of human steroid-sulfatase:

[89587] membrane topology, glycosylation, and subcellular distribution in

[89588] BHK-21 cells. J. Biol. Chem. 264: 13865-13872, 1989.

[89589]

[89590] 114. Steinmann, B.; Mieth, D.; Gitzelmann, R.: A newly recognized

[89591] cause of low urinary estriol in pregnancy: multiple sulfatase deficiency

[89592] of the fetus. Gynec. Obstet. Invest. 12: 107-109, 1981.

[89593]

- [89594] 115. Stoll, C.; Grosshans, E.; Binder, P.; Roth, M.: Hyper-trophic
- [89595] pyloric stenosis associated with X-linked ichthyosis in two brothers. Clin.
- [89596] Exp. Derm. 8: 61–64, 1983.
- [89597]
- [89598] 116. Sunohara, N.; Sakuragawa, N.; Satoyoshi, E.; Tanae, A.; Shapiro,
- [89599] L. J.: A new syndrome of anosmia, ichthyosis, hypogonadism, and various
- [89600] neurological manifestations with deficiency of steroid sulphatase
- [89601] and arylsulphatase C. Ann. Neurol. 19: 174–181, 1986.
- [89602]
- [89603] 117. Tabei, F.; Heinrichs, W. L.: Diagnosis of placental sulphatase
- [89604] deficiency. Am. J. Obstet. Gynec. 124: 409–414, 1976.
- [89605]
- [89606] 118. Tiepolo, L.; Zuffardi, O.; Fraccaro, M.; di Natale, D.; Gargantini,
- [89607] L.; Muller, C. R.; Ropers, H.–H.: Assignment by deletion

mapping

[89608] of the steroid sulfatase X-linked ichthyosis locus to
Xp223. Hum.

[89609] Genet. 54: 205–206, 1980.

[89610]

[89611] 119. Toder, R.; Rappold, G. A.; Schiebel, K.; Schempp, W.:
ANT3 and

[89612] STS are autosomal in prosimian lemurs: implications for
the evolution

[89613] of the pseudoautosomal region. Hum. Genet. 95: 22–28,
1995.

[89614]

[89615] 120. Traupe, H.; Happle, R.: Clinical spectrum of steroid
sulfatase

[89616] deficiency: X-linked recessive ichthyosis, birth complica-
tions, and

[89617] cryptorchidism. Europ. J. Pediat. 140: 19–21, 1983.

[89618]

[89619] 121. Traupe, H.; Muller-Migl, C. R.; Kolde, G.; Happle, R.;
Kovary,

[89620] P. M.; Hameister, H.; Ropers, H. H.: Ichthyosis vulgaris

with hypogenitalism

[89621] and hypogonadism: evidence for different genotypes by lipoprotein

[89622] electrophoresis and steroid sulfatase testing. Clin. Genet. 25:

[89623] 42-51, 1984.

[89624]

[89625] 122. Traupe, H.; Ropers, H.-H.: Cryptorchidism and hypogenitalism

[89626] in X-linked recessive ichthyosis vulgaris. (Letter) Hum. Genet. 60:

[89627] 206, 1982.

[89628]

[89629] 123. Turpin, R.; Desvignes, P.; Demassieux, J.-L.: Sur une variete

[89630] d'ichthyose hereditaire avec alteration du fond d'oeil. Nouveau syndrome

[89631] ectoblastique?. Sem. Hop. Paris 21: 343, 1945.

[89632]

[89633] 124. Valdes-Flores, M.; Kofman-Alfaro, S. H.; Jimenez-Vaca, A. L.;

- [89634] Cuevas–Covarrubias, S. A.: Carrier identification by FISH analysis
- [89635] in isolated cases of X-linked ichthyosis. *Am. J. Med. Genet.* 102:
- [89636] 146–148, 2001.
- [89637]
- [89638] 125. Valdes–Flores, M.; Kofman–Alfaro, S. H.; Jimenez Vaca, A. L.;
- [89639] Cuevas–Covarrubias, S. A.: A novel partial deletion of exons 2–10
- [89640] of the STS gene in recessive X-linked ichthyosis. *J. Invest. Derm.* 114:
- [89641] 591–593, 2000.
- [89642]
- [89643] 126. Vogel, W.; Grompe, M.; Storz, R.; Pentz, S.: A comparative study
- [89644] on steroid sulfatase and arylsulfatase C in fibroblast clones from
- [89645] 45,X/47,XXX and 69,XXY. *Hum. Genet.* 66: 367–369, 1984.
- [89646]

[89647] 127. Wells, R. S.; Jennings, M. C.: X-linked ichthyosis and ichthyosis

[89648] vulgaris: clinical and genetic distinctions in a second series of

[89649] families. J.A.M.A. 202: 485–488, 1967.

[89650]

[89651] 128. Went, L. N.; DeGroot, W. P.; Sanger, R.; Tippet, P.; Gavin,

[89652] J.: X-linked ichthyosis: linkage relationship with the Xg blood groups

[89653] and other studies in a large Dutch kindred. Ann. Hum. Genet. 32:

[89654] 333–346, 1969.

[89655]

[89656] 129. Wieacker, P.; Davies, K. E.; Mevorah, B.; Ropers, H. H.: Linkage

[89657] studies in a family with X-linked recessive ichthyosis employing a

[89658] cloned DNA sequence from the distal short arm of the X chromosome. Hum.

[89659] Genet. 63: 113–116, 1983.

[89660]

[89661] 130. Willard, H. F.; Holmes, M. T.: A sensitive and dependable assay

[89662] for distinguishing hamster and human X-linked steroid sulfatase activity

[89663] in somatic cell hybrids. Hum. Genet. 66: 272-275, 1984.

[89664]

[89665] 131. Wirth, B.; Herrmann, F. H.; Neugebauer, M.; Gillard, E. F.; Wulff,

[89666] K.; Stein, C.; von Figura, K.; Ferguson-Smith, M. A.; Gal, A.: Linkage

[89667] analysis in X-linked ichthyosis (steroid sulfatase deficiency). Hum.

[89668] Genet. 80: 191-192, 1988.

[89669]

[89670] 132. Yen, P. H.; Allen, E.; Marsh, B.; Mohandas, T.; Wang, N.; Taggart,

[89671] R. T.; Shapiro, L. J.: Cloning and expression of steroid sulfatase

[89672] cDNA and the frequent occurrence of deletions in STS deficiency: implications

[89673] for X-Y interchange. Cell 49: 443-454, 1987.

[89674]

[89675] 133. Yen, P. H.; Marsh, B.; Allen, E.; Tsai, S. P.; Ellison, J.; Connolly,

[89676] L.; Neiswanger, K.; Shapiro, L. J.: The human X-linked steroid sulfatase

[89677] gene and a Y-encoded pseudogene: evidence for an inversion of the

[89678] Y chromosome during primate evolution. Cell 55: 1123-1135, 1988.

[89679]

[89680] 134. Zettersten, E.; Man, M.-Q.; Sato, J.; Denda, M.; Farrell, A.;

[89681] Ghadially, R.; Williams, M. L.; Feingold, K. R.; Elias, P. M. Recessive

[89682] X-linked ichthyosis: role of cholesterol-sulfate accumulation in the

[89683] barrier abnormality. J. Invest. Derm. 111: 784-790, 1998.

[89684] 15815.

[89685] Holinski-Feder, E.; Chahrokh-Zadeh, S.; Jedelev, K. B.; Meindl,

[89686] A.; Steinbach, P.; Wohrle, D.: FRAXE testing. (Letter) Am. J. Hum.

[89687] Genet. 59: 1168–1169, 1996.

- [89688] 15816.
- [89689] Knight, S. J. L.; Flannery, A. V.; Hirst, M. C.; Campbell, L.;
- [89690] Christodoulou, Z.; Phelps, S. R.; Middleton-Price, H. R.;
- Bamicoat,
- [89691] A.; Pembrey, M. E.; Holland, J.; Bobrow, M.; Oostra, B. A.;
- Davies,
- [89692] K. E.: FRAXE mental retardation associated with trinucleotide repeat
- [89693] expansion and hypermethylation of a CpG island in Xq28. (Abstract) Am.
- [89694] J. Hum. Genet. 53 (suppl.): A79, 1993.
- [89695] 15817.
- [89696] Knight, S. J. L.; Flannery, A. V.; Hirst, M. C.; Campbell, L.;
- [89697] Christodoulou, Z.; Phelps, S. R.; Pointon, J.; Middleton-Price, H.
- [89698] R.; Barnicoat, A.; Pembrey, M. E.; Holland, J.; Oostra, B. A.;
- Bobrow,
- [89699] M.; Davies, K. E.: Trinucleotide repeat amplification and hypermethylation
- [89700] of a CpG island in FRAXE mental retardation. Cell 74: 127-134, 1993.

- [89701] 15818.
- [89702] Knight, S. J. L.; Ritchie, R. J.; Chakrabarti, L.; Cross, G.;
- [89703] Taylor, G. R.; Mueller, R. F.; Hurst, J.; Paterson, J.; Yates, J.
- [89704] R. W.; Dow, D. J.; Davies, K. E.: A study of FRAXE in men-
tally retarded
- [89705] individuals referred for fragile X syndrome (FRAXA) testing
in the
- [89706] United Kingdom. Am. J. Hum. Genet. 58: 906–913, 1996.
- [89707] 15819.
- [89708] Knight, S. J. L.; Voelckel, M. A.; Hirst, M. C.; Flannery, A.
- [89709] V.; Moncla, A.; Davies, K. E.: Triplet repeat expansion at
the FRAXE
- [89710] locus and X-linked mild mental handicap. Am. J. Hum.
Genet. 55:
- [89711] 81–86, 1994.
- [89712] 15820.
- [89713] Mila, M.; Sanchez, A.; Badenas, C.; Brun, C.; Jimenez, D.;
- Villa,
- [89714] M. P.; Castellvi-Bel, S.; Estivill, X.: Screening for FMR1 and
FMR2

- [89715] mutations in 222 individuals from Spanish special schools: identification
- [89716] of a case of FRAXE-associated mental retardation. Hum. Genet. 100:
- [89717] 503–507, 1997.
- [89718] 15821.
- [89719] Mulley, J. C.; Yu, S.; Loesch, D. Z.; Hay, D. A.; Donnelly, A.;
- [89720] Gedeon, A. K.; Carbonell, P.; Lopez, I.; Glover, G.; Gabar-
ron, I.;
- [89721] Yu, P. W. L.; Baker, E.; Haan, E. A.; Hockey, A.; Knight, S. J.
L.;
- [89722] Davies, K. E.; Richards, R. I.; Sutherland, G. R.: FRAXE and
mental
- [89723] retardation. J. Med. Genet. 32: 162–169, 1995.
- [89724] 15822.
- [89725] Murray, A.; Webb, J.; Dennis, N.; Conway, G.; Morton, N.:
Microdeletions
- [89726] in FMR2 may be a significant cause of premature ovarian
failure. J.
- [89727] Med. Genet. 36: 767–770, 1999.
- [89728] 15823.

- [89729] Russo, S.; Selicorni, A.; Bedeschi, M. F.; Natacci, F.; Viziello, P.; Fortuna, R.; Pagani, G.; Dalpra, L.; Larizza, L.: Molecular characterization of FRAXE-positive subjects with mental impairment in two unrelated Italian families. Am. J. Med. Genet. 75: 304–308, 1998.
- [89732] 15824.
- [89733] Sutherland, G. R.: Personal Communication. Adelaide, Australia 11/17/1993.
- [89734] 15825.
- [89735] Sutherland, G. R.; Baker, E.: Characterisation of a new rare fragile site easily confused with the fragile X. Hum. Molec. Genet. 1: 111–113, 1992.
- [89736] 15826.
- [89737] Abitbol, M.; Menini, C.; Delezoide, A.–L.; Rhyner, T.; Veke-mans, M.; Mallet, J.: Nucleus basalis magnocellularis and hippocampus are

- [89743] the major sites of FMR-1 expression in the human fetal brain. *Nature*
- [89744] *Genet.* 4: 147-153, 1993.
- [89745] 15827.
- [89746] Abrams, M. T.; Kaufmann, W. E.; Rousseau, F.; Oostra, B. A.; Wolozin,
- [89747] B.; Taylor, C. V.; Lishaa, N.; Morel, M.-L.; Hoogeveen, A.; Reiss,
- [89748] A. L.: FMR1 gene expression in olfactory neuroblasts from two males
- [89749] with fragile X syndrome. *Am. J. Med. Genet.* 82: 25-30, 1999.
- [89750] 15828.
- [89751] Allingham-Hawkins, D. J.; Babul-Hirji, R.; Chitayat, D.; Holden,
- [89752] J. J. A.; Yang, K. T.; Lee, C.; Hudson, R.; Gorwill, H.; Nolin, S.
- [89753] L.; Glicksman, A.; Jenkins, E. C.; Brown, W. T.; and 27 others:
- [89754] Fragile X premutation is a significant risk factor for premature ovarian
- [89755] failure: the international collaborative POF in fragile X

study--preliminary

[89756] data. Am. J. Med. Genet. 83: 322-325, 1999.

[89757] 15829.

[89758] Ashley, C. T.; Sutcliffe, J. S.; Kunst, C. B.; Leiner, H. A.;
Eichler,

[89759] E. E.; Nelson, D. L.; Warren, S. T.: Human and murine FMR-
1: alternative

[89760] splicing and translational initiation downstream of the
CGG-repeat. Nature

[89761] Genet. 4: 244-251, 1993.

[89762] 15830.

[89763] Ashley, C. T., Jr.; Wilkinson, K. D.; Reines, D.; Warren, S. T.

[89764] : FMR1 protein: conserved RNP family domains and selec-
tive RNA binding. Science 262:

[89765] 563-566, 1993.

[89766] 15831.

[89767] Bachner, D.; Steinbach, P.; Wohrle, D.; Just, W.; Vogel, W.;
Hameister,

[89768] H.; Manca, A.; Poustka, A.: Enhanced Fmr-1 expression in
testis.

[89769] (Letter) Nature Genet. 4: 115-116, 1993.

[89770] 15832.

[89771] Backes, M.; Genc, B.; Schreck, J.; Doerfler, W.; Lehmkuhl, G.;

[89772] von Gontard, A.: Cognitive and behavioral profile of fragile X boys:

[89773] correlations to molecular data. Am. J. Med. Genet. 95: 150-156,

[89774] 2000.

[89775] 15833.

[89776] Maglott, D. R.; Durkin, A. S.; Lane, S. A.; Callen, D. F.; Feldblyum,

[89777] T. V.; Nierman, W. C.: The gene for membrane protein E16 (D16S469E)

[89778] maps to human chromosome 16q24.3 and is expressed in human brain,

[89779] thymus, and retina. Genomics 23: 303-304, 1994.

[89780] 15834.

[89781] Kastury, K.; Druck, T.; Huebner, K.; Barletta, C.; Acampora, D.;

[89782] Simeone, A.; Faiella, A.; Boncinelli, E.: Chromosome locations of

- [89783] human EMX and OTX genes. Genomics 22: 41–45, 1994.
- [89784] 15835.
- [89785] Simeone, A.; Acampora, D.; Gulisano, M.; Stornaiuolo, A.; Boncinelli, E.: Nested expression domains of four homeobox genes in developing rostral brain. Nature 358: 687–690, 1992.
- [89787] 15836.
- [89788] Bishop, K. M.; Goudreau, G.; O'Leary, D. D. M.: Regulation of area identity in the mammalian neocortex by Emx2 and Pax6. Science 288: 344–349, 2000.
- [89789] 15837.
- [89790] Boncinelli, E.; Gulisano, M.; Spada, F.; Broccoli, V.: Emx and Otx gene expression in the developing mouse brain. Ciba Found. Symp. 193: 100–116, 1995.
- [89791] 15838.

- [89797] Bosetti, A.; Faiella, A.; Boncinelli, E.; Consalez, G. G.: Linkage
- [89798] mapping of Emx2 to mouse chromosome 19. *Mammalian Genome* 8: 71-72,
- [89799] 1997.
- [89800] 15839.
- [89801] Noonan, F. C.; Mutch, D. G.; Mallon, M. A.; Goodfellow, P. J.:
- [89802] Characterization of the homeodomain gene EMX2: sequence conservation,
- [89803] expression analysis, and a search for mutations in endometrial cancers. *Genomics* 76:
- [89804] 37-44, 2001.
- [89805] 15840.
- [89806] Acampora, D.; Mazan, S.; Avantaggiato, V.; Barone, P.; Tuorto,
- [89807] F.; Lallemand, Y.; Brulet, P.; Simeone, A.: Epilepsy and brain abnormalities
- [89808] in mice lacking the Otx1 gene. *Nature Genet.* 14: 218-222, 1996.
- [89809] 15841.

- [89810] Boncinelli, E.; Gulisano, M.; Broccoli, V.: Emx and Otx homeobox
genes in the developing mouse brain. J. Neurobiol. 24:
1356–1366,
1993.
15842.
- [89814] Frantz, G. D.; Weimann, J. M.; Levin, M. E.; McConnell, S.
K.:
Otx1 and Otx2 define layers and regions in developing
cerebral cortex
and cerebellum. J. Neurosci. 14: 5725–5740, 1994.
15843.
- [89818] Avraham, S.; Jiang, S.; Ota, S.; Fu, Y.; Deng, B.; Dowler, L.
L.;
White, R. A.; Avraham, H.: Structural and functional studies of the
intracellular tyrosine kinase MATK gene and its translated
product. J.
Biol. Chem. 270: 1833–1842, 1995.
15844.
- [89823] Bennett, B. D.; Cowley, S.; Jiang, S.; London, R.; Deng, B.;

Grabarek,

[89824] J.; Groopman, J. E.; Goeddel, D. V.; Avraham, H.: Identification

[89825] and characterization of a novel tyrosine kinase from megakaryocytes. J.

[89826] Biol. Chem. 269: 1068–1074, 1994.

[89827] 15845.

[89828] Klages, S.; Adam, D.; Class, K.; Fargnoli, J.; Bolen, J. B.; Penhallow,

[89829] R. C.: Ctk: a protein–tyrosine kinase related to Csk that defines

[89830] an enzyme family. Proc. Nat. Acad. Sci. 91: 2597–2601, 1994.

[89831] 15846.

[89832] Sakano, S.; Iwama, A.; Inazawa, J.; Ariyama, T.; Ohno, M.; Suda,

[89833] T.: Molecular cloning of a novel non–receptor tyrosine kinase, HYL

[89834] (hematopoietic consensus tyrosine–lacking kinase). Oncogene 9: 1155–1161,

[89835] 1994.

[89836] 15847.

- [89837] Zrihan-Licht, S.; Lim, J.; Keydar, I.; Sliwkowski, M. X.; Groopman, J. E.; Avraham, H.: Association of Csk-homologous kinase (CHK) (formerly MATK) with HER-2/ErbB-2 in breast cancer cells. J. Biol. Chem. 272: 1856-1863, 1997.
- [89840] 15848.
- [89841] Shimizu, S.; Narita, M.; Tsujimoto, Y.: Bcl-2 family proteins regulate the release of apoptogenic cytochrome c by the mitochondrial channel VDAC. Nature 399: 483-487, 1999.
- [89844] 15849.
- [89845] Nichols, A. F.: Personal Communication. Berkeley, Calif. 10/4/1995.
- [89846] 15850.
- [89847] Campbell, H. D.; Schimansky, T.; Claudianos, C.; Ozsarac, N.; Kasprzak, A. B.; Cotsell, J. N.; Young, I. G.; de Couet, H. G.; Gabor Miklos,

- [89850] G. L.: The *Drosophila melanogaster* flightless-I gene involved in
- [89851] gastrulation and muscle degeneration encodes gelsolin-like and leucine-rich
- [89852] repeat domains and is conserved in *Caenorhabditis elegans* and humans. *Proc.*
- [89853] *Nat. Acad. Sci.* 90: 11386–11390, 1993.
- [89854] 15851.
- [89855] Harmon, D. L.; Gardner-Medwin, D.; Stirling, J. L.: Two new mutations
- [89856] in a late infantile Tay-Sachs patient are both in exon 1 of the beta-hexosaminidase
- [89857] alpha subunit gene. *J. Med. Genet.* 30: 123–128, 1993.
- [89858] 15852.
- [89859] Buckle, V.; Mondello, C.; Darling, S.; Craig, I. W.; Goodfellow,
- [89860] P. N.: Homologous expressed genes in the human sex chromosome pairing
- [89861] region. *Nature* 317: 739–741, 1985.
- [89862] 15853.
- [89863] Slaugenhaupt, S. A.; Roca, A. L.; Liebert, C. B.; Altherr, M.

R.;

[89864] Gusella, J. F.; Reppert, S. M.: Mapping of the gene for the Mel1a-melatonin

[89865] receptor to human chromosome 4 (MTNR1A) and mouse chromosome 8 (Mtnr1a). Genomics 27:

[89866] 355-357, 1995.

[89867] 15854.

[89868] Hulsebos, T. J. M.; Cerosaletti, K. M.; Fournier, R. E. K.; Sinke,

[89869] R. J.; Rocchi, M.; Marzella, R.; Jenkins, N. A.; Gilbert, D. J.; Copeland,

[89870] N. G.: Identification of the human beta-A2 crystallin gene (CRYBA2):

[89871] localization of the gene on human chromosome 2 and of the homologous

[89872] gene on mouse chromosome 1. Genomics 28: 543-548, 1995.

[89873] 15855.

[89874] Grimsby, J.; Toth, M.; Chen, K.; Kumazawa, T.; Klaidman, L.; Adams,

[89875] J. D.; Karoum, F.; Gal, J.; Shih, J. C.: Increased stress response

- [89876] and beta-phenylethylamine in MAOB-deficient mice. *Nature Genet.* 17:
[89877] 206-210, 1997.
- [89878] 15856.
- [89879] Lenders, J. W. M.; Eisenhofer, G.; Abeling, N. G. G. M.;
Berger,
[89880] W.; Murphy, D. L.; Konings, C. H.; Bleeker Wagemakers, L.
M.; Kopin,
[89881] I. J.; Karoum, F.; van Gennip, A. H.; Brunner, H. G.: Specific
genetic
[89882] deficiencies of the A and B isoenzymes of monoamine ox-
idase are characterized
[89883] by distinct neurochemical and clinical phenotypes. *J. Clin.*
Invest. 97:
[89884] 1010-1019, 1996.
- [89885] 15857.
- [89886] Rice, J.; McGuffin, P.; Goldin, L. R.; Shaskan, E. G.; Ger-
shon,
[89887] E. S.: Platelet monoamine oxidase (MAO) activity: evidence
for a
[89888] single major locus. *Am. J. Hum. Genet.* 36: 36-43, 1984.

[89889] 15858.

[89890] Weinshilboum, R. M.: Biochemical genetics of catecholamines in

[89891] humans. Mayo Clin. Proc. 58: 319–330, 1983.

[89892] 15859.

[89893] Wyatt, R. J.; Murphy, D. L.; Belmaker, R.; Cohen, S.; Donnelly,

[89894] C. H.; Pollin, W.: Reduced monoamine oxidase activity in platelets:

[89895] a possible genetic marker for vulnerability to schizophrenia. Science 179:

[89896] 916–918, 1973.

[89897] 15860.

[89898] Hattori, Y.; Yamashiro, Y.; Ohba, Y.; Miyaji, T.; Morishita, M.;

[89899] Yamamoto, K.; Yamamoto, K.; Narai, S.; Kimura, A.: A new beta-thalassemia

[89900] mutation (initiation codon ATG-to-GTG) found in the Japanese population. Hemoglobin 15:

[89901] 317–325, 1991.

[89902] 15861.

- [89903] Holmes, L. B.: Norrie's disease--an X-linked syndrome of retinal
- [89904] malformation, mental retardation and deafness. New Eng. J. Med. 284:
- [89905] 367-368, 1971.
- [89906] 15862.
- [89907] Isashiki, Y.; Ohba, N.; Yanagita, T.; Hokita, N.; Doi, N.; Nakagawa,
- [89908] M.; Ozawa, M.; Kuroda, N.: Novel mutation at the initiation codon
- [89909] in the Norrie disease gene in two Japanese families. Hum. Genet. 95:
- [89910] 105-108, 1995.
- [89911] 15863.
- [89912] Isashiki, Y.; Ohba, N.; Yanagita, T.; Hokita, N.; Hotta, Y.; Hayakawa,
- [89913] M.; Fujiki, K.; Tanabe, U.: Mutations in the Norrie disease gene:
- [89914] a new mutation in a Japanese family. (Letter) Brit. J. Ophthalmol. 79:
- [89915] 703-708, 1995.

- [89916] 15864.
- [89917] Johnson, K.; Mintz-Hittner, H. A.; Conley, Y. P.; Ferrell, R.
- [89918] E.: X-linked exudative vitreoretinopathy caused by an arginine to
- [89919] leucine substitution (R121L) in the Norrie disease protein. Clin.
- [89920] Genet. 50: 113-115, 1996.
- [89921] 15865.
- [89922] Johnston, S. S.; Hanna, J. E.; Nevin, N. C.; Bryars, J. H.: Norrie's
- [89923] disease. Birth Defects Orig. Art. Ser. 18(6): 729-738, 1982.
- [89924] 15866.
- [89925] Katayama, S.; Wohlferd, M.; Golbus, M. S.: First demonstration
- [89926] of recombination between the gene for Norrie disease and probe L1.28. Am.
- [89927] J. Med. Genet. 30: 967-970, 1988.
- [89928] 15867.
- [89929] Kivlin, J. D.; Sanborn, G. E.; Wright, E.; Cannon, L.; Carey,

- [89930] J.: Further linkage data on Norrie disease. *Am. J. Med. Genet.* 26:
- [89931] 733–736, 1987.
- [89932] 15868.
- [89933] Lindsay, S.; Thiselton, D. L.; Bateman, J. B.; Ngo, J. T.; Sparkes,
- [89934] R. S.; Coleman, M.; Davies, K. E.; Bhattacharya, S. S.: Localisation
- [89935] of the gene for Norrie disease to between DXS7 and DXS426 on Xp. *Hum. Genet.* 88: 349–350, 1992.
- [89936] 15869.
- [89937] Meindl, A.; Berger, W.; Meitinger, T.; van de Pol, D.; Achatz,
- [89938] H.; Dorner, C.; Haasemann, M.; Hellebrand, H.; Gal, A.; Cremers, F.;
- [89939] Ropers, H.–H.: Norrie disease is caused by mutations in an extracellular
- [89940] protein resembling C–terminal globular domain of mucins. *Nature Genet.* 2:
- [89941] 139–143, 1992.

[89943] 15870.

[89944] Meindl, A.; Lorenz, B.; Achatz, H.; Hellebrand, H.;
Schmitz-Valckenberg,

[89945] P.; Meitinger, T.: Missense mutations in the NDP gene in
patients

[89946] with a less severe course of Norrie disease. Hum. Molec.
Genet. 4:

[89947] 489–490, 1995.

[89948] 15871.

[89949] Meitinger, T.; Meindl, A.; Bork, P.; Rost, B.; Sander, C.;
Haasemann,

[89950] M.; Murken, J.: Molecular modeling of the Norrie disease
protein

[89951] predicts a cystine knot growth factor tertiary structure.
Nature

[89952] Genet. 5: 376–380, 1993.

[89953] 15872.

[89954] Moreira-Filho, C. A.; Neustein, I.: A presumptive new vari-
ant

[89955] of Norrie's disease. J. Med. Genet. 16: 125–128, 1979.

[89956] 15873.

- [89957] Nance, W. E.; Hara, S.; Hansen, A.; Elliott, J.; Lewis, M.; Chown,
[89958] B.: Genetic linkage studies in a Negro kindred with Norrie's disease. Am.
[89959] J. Hum. Genet. 21: 423-429, 1969.
[89960] 15874.
- [89961] Ngo, J.; Spence, M. A.; Cortessis, V.; Bateman, J. B.; Sparkes,
[89962] R. S.: Duplicate report crossing over in Norrie disease family. (Letter) Am.
[89963] J. Med. Genet. 33: 286, 1989.
[89964] 15875.
- [89965] Ngo, J. T.; Bateman, J. B.; Cortessis, V.; Sparkes, R. S.; Mohandas,
[89966] T.; Inana, G.; Spence, M. A.: Norrie disease: linkage analysis using
[89967] a 4.2-kb RFLP detected by a human ornithine aminotransferase cDNA
[89968] probe. Genomics 4: 539-545, 1989.
[89969] 15876.
- [89970] Ngo, J. T.; Spence, M. A.; Cortessis, V.; Sparkes, R. S.;

Bateman,

[89971] J. B.: Recombinational event between Norrie disease and DXS7 loci. Clin.

[89972] Genet. 34: 43–47, 1988.

[89973] 15877.

[89974] Norrie, G.: Nogle Blindhedsaarsager: en oversigt. Hospitalstidende 76:

[89975] 141–147, 1933.

[89976] 15878.

[89977] Ohba, N.; Yamashita, T.: Primary vitreoretinal dysplasia resembling

[89978] Norrie's disease in a female: association with X autosome chromosomal

[89979] translocation. Brit. J. Ophthal. 70: 64–71, 1986.

[89980] 15879.

[89981] Phillips, C. I.; Newton, M.; Duvall, J.; Holloway, S.; Levy, A.

[89982] M.: Probably Norrie's disease due to mutation: two sporadic sibships

[89983] of two males each, a necropsy of one case, and, given Norrie's disease,

[89984] a calculation of the gene mutation frequency. Brit. J. Oph–

thal. 70:

[89985] 305–313, 1986.

[89986] 15880.

[89987] Rehm, H. L.; Gutierrez–Espeleta, G. A.; Garcia, R.; Jimenez, G.;

[89988] Khetarpal, U.; Priest, J. M.; Sims, K. B.; Keats, B. J. B.; Morton,

[89989] C. C.: Norrie disease gene mutation in a large Costa Rican kindred

[89990] with a novel phenotype including venous insufficiency. Hum. Mutat.

[89991] 9: 402–408, 1997.

[89992] 15881.

[89993] Schuback, D. E.; Chen, Z. Y.; Craig, I. W.; Breakefield, X. O.;

[89994] Sims, K. B.: Mutations in the Norrie disease gene. Hum. Mutat. 5:

[89995] 285–292, 1995.

[89996] 15882.

[89997] Shastry, B. S.: Identification of a recurrent missense mutation

- [89998] in the Norrie disease gene associated with a simplex case of exudative
- [89999] vitreoretinopathy. *Biochem. Biophys. Res. Commun.* 246: 35–38, 1998.
- [90000] 15883.
- [90001] Shastry, B. S.; Hejtmancik, J. F.; Plager, D. A.; Hartzler, M.
- [90002] K.; Trese, M. T.: Linkage and candidate gene analysis in X-linked
- [90003] familial exudative vitreoretinopathy. *Genomics* 27: 341–344, 1995.
- [90004] 15884.
- [90005] Shastry, B. S.; Hejtmancik, J. F.; Trese, M. T.: Identification
- [90006] of novel missense mutations in the Norrie disease gene associated
- [90007] with one X-linked and four sporadic cases of familial exudative vitreoretinopathy. *Hum.*
- [90008] *Mutat.* 9: 396–401, 1997.
- [90009] 15885.
- [90010] Sims, K. B.; Lebo, R. V.; Benson, G.; Shalish, C.; Schuback, D.;
- [90011] Chen, Z. Y.; Bruns, G.; Craig, I. W.; Golbus, M. S.; Breake-

field,

[90012] X. O.: The Norrie disease gene maps to a 150 kb region on chromosome

[90013] Xp11.3. Hum. Molec. Genet. 1: 83-89, 1992.

[90014] 15886.

[90015] Sims, K. B.; Ozelius, L.; Corey, T.; Rinehart, W. B.; Libera-

[90016] R.; Haines, J.; Chen, W. J.; Norio, R.; Sankila, E.; de la Chapelle,

[90017] A.; Murphy, D. L.; Gusella, J.; Breakefield, X. O.: Norrie disease

[90018] gene is distinct from the monoamine oxidase genes. Am. J. Hum. Genet. 45:

[90019] 424-434, 1989.

[90020] 15887.

[90021] Taylor, P. J.; Coates, T.; Newhouse, M. L.: Episkopi blindness:

[90022] hereditary blindness in a Greek Cypriot family. Brit. J. Ophthalmol. 43:

[90023] 340-344, 1959.

[90024] 15888.

- [90025] Torrente, I.; Mangino, M.; Gennarelli, M.; Novelli, G.; Giannotti,
[90026] A.; Vadala, P.; Dallapiccola, B.: Two new missense mutations (A105T
[90027] and C110G) in the norrin gene in two Italian families with
Norrie
[90028] disease and familial exudative vitreoretinopathy. (Letter)
Am. J.
[90029] Med. Genet. 72: 242-244, 1997.
[90030] 15889.
[90031] Burgoyne, P. S.: Genetic homology and crossing over in
the X and
[90032] Y chromosomes of mammals. Hum. Genet. 61: 85-90,
1982.
[90033] 15890.
[90034] Cooke, H. J.; Brown, W. R. A.; Rappold, G. A.: Hypervariable telomeric
[90035] sequences from the human sex chromosomes are pseudoautosomal. Nature 317:
[90036] 687-692, 1985.
[90037] 15891.

- [90038] Darling, S. M.; Banting, G. S.; Pym, B.; Wolfe, J.; Goodfellow,
low,
- [90039] P. N.: Cloning an expressed gene shared by the human
sex chromosomes. *Proc.*
- [90040] *Nat. Acad. Sci.* 83: 135–139, 1986.
- [90041] 15892.
- [90042] Dracopoli, N. C.; Rettig, W. J.; Albino, A. P.; Esposito, D.;
Archidiacono,
- [90043] N.; Rocchi, M.; Siniscalco, M.; Old, L. J.: Genes controlling
gp25/30
- [90044] cell-surface molecules map to chromosomes X and Y and
escape X-inactivation. *Am.*
- [90045] *J. Hum. Genet.* 37: 199–207, 1985.
- [90046] 15893.
- [90047] Gelin, C.; Aubrit, F.; Phalipon, A.; Raynal, B.; Cole, S.; Kac-
zorek,
- [90048] M.; Bernard, A.: The E2 antigen, a 32 kD glycoprotein in-
volved in
- [90049] T-cell adhesion processes, is the MIC2 gene product.
EMBO J. 8:
- [90050] 3253–3259, 1989.

[90051] 15894.

[90052] Geller, R. L.; Shapiro, L. J.; Mohandas, T. K.: Fine mapping
of

[90053] the distal short arm of the human X chromosome using X/
Y translocations. Am.

[90054] J. Hum. Genet. 38: 884–890, 1986.

[90055] 15895.

[90056] Goodfellow, P.; Banting, G.; Sheer, D.; Ropers, H. H.;
Caine,

[90057] A.; Ferguson-Smith, M. A.; Povey, S.; Voss, R.: Genetic evi-
dence

[90058] that a Y-linked gene in man is homologous to a gene on
the X chromosome. Nature 302:

[90059] 346–349, 1983.

[90060] 15896.

[90061] Goodfellow, P.; Pym, B.; Mohandas, T.; Shapiro, L. J.: The
cell

[90062] surface antigen locus, MIC2X, escapes X-inactivation. Am.
J. Hum.

[90063] Genet. 36: 777–782, 1984.

[90064] 15897.

- [90065] Goodfellow, P. J.; Darling, S. M.; Thomas, N. S.; Goodfellow,
low,
- [90066] P. N.: A pseudoautosomal gene in man. Science 234:
740–743, 1986.
- [90067] 15898.
- [90068] Goodfellow, P. N.; Tippet, P.: A human quantitative polymorphism
morphism
- [90069] related to Xg blood groups. Nature 289: 404–405, 1981.
- [90070] 15899.
- [90071] Levy, R.; Dilley, J.; Fox, R. I.; Warnke, R.: A human thymus-leukemia
leukemia
- [90072] antigen defined by hybridoma monoclonal antibodies.
Proc. Nat. Acad.
- [90073] Sci. 76: 6552–6556, 1979.
- [90074] 15900.
- [90075] Pettersen, R. D.; Bernard, G.; Olafsen, M. K.; Pourtein, M.;
Lie,
- [90076] S. O.: CD99 signals caspase-independent T cell death. J.
Immun. 166:
- [90077] 4931–4942, 2001.
- [90078] 15901.

- [90079] Ropers, H. H.; Zimmer, J.; Strobl, G.; Goodfellow, P.: The MIC2X
- [90080] (12E7) locus maps distally from STS on Xp. (Abstract) Cytogenet.
- [90081] Cell Genet. 40: 736 only, 1985.
- [90082] 15902.
- [90083] Simmler, M.-C.; Rouyer, F.; Vergnaud, G.; Nystrom-Lahti, M.; Ngo,
- [90084] K. Y.; de la Chapelle, A.; Weissenbach, J.: Pseudoautosomal DNA sequences
- [90085] in the pairing region of the human sex chromosomes. Nature 317:
- [90086] 692-697, 1985.
- [90087] 15903.
- [90088] Tippett, P.; Shaw, M.-A.; Green, C. A.; Daniels, G. L.: The 12E7
- [90089] red cell quantitative polymorphism: control by the Y-borne locus,
- [90090] Yg. Ann. Hum. Genet. 50: 339-347, 1986.
- [90091] 15904.
- [90092] Janz, R.; Sudhof, T. C.; Hammer, R. E.; Unni, V.; Siegel-

baum, S.

- [90093] A.; Bolshakov, V. Y.: Essential roles in synaptic plasticity for
- [90094] synaptogyrin I and synaptophysin I. *Neuron* 24: 687–700, 1999.
- [90095] 15905.
- [90096] McMahon, H. T.; Bolshakov, V. Y.; Janz, R.; Hammer, R. E.; Siegelbaum,
- [90097] S. A.; Sudhof, T. C.: Synaptophysin, a major synaptic vesicle protein,
- [90098] is not essential for neurotransmitter release. *Proc. Nat. Acad. Sci.* 93:
- [90099] 4760–4764, 1996.
- [90100] 15906.
- [90101] Brown, C. J.; Sekiguchi, T.; Nishimoto, T.; Willard, H. F.: Regional
- [90102] localization of CCG1 gene which complements hamster cell cycle mutation
- [90103] BN462 to Xq11–Xq13. *Somat. Cell Molec. Genet.* 15: 93–96, 1989.
- [90104] 15907.
- [90105] Derry, J. M. J.; Barnard, P. J.: Localization of the Ccg1 gene

- [90106] on the mouse X chromosome. (Abstract) Cytogenet. Cell Genet. 51:
- [90107] 988, 1989.
- [90108] 15908.
- [90109] Dikstein, R.; Ruppert, S.; Tjian, R.: TAFII250 is a bipartite
- [90110] protein kinase that phosphorylates the base transcription factor RAP74. Cell 84:
- [90111] 781–790, 1996.
- [90112] 15909.
- [90113] Giles, R. E.; Ruddle, F. H.: X-linkage of a human genetic locus
- [90114] that corrects the DNA synthesis lesion in tsC1AGOH mouse cells. Genetics 93:
- [90115] 975–996, 1979.
- [90116] 15910.
- [90117] Hisatake, K.; Hasegawa, S.; Takada, R.; Nakatani, Y.; Horikoshi,
- [90118] M.; Roeder, R. G.: The p250 subunit of native TATA box-binding factor
- [90119] TFIID is the cell-cycle regulatory protein CCG1. Nature 362: 179–181,

[90120] 1993.

[90121] 15911.

[90122] Jacobson, R. H.; Ladurner, A. G.; King, D. S.; Tjian, R.:
Structure

[90123] and function of a human TAFII250 double bromodomain
module. Science 288:

[90124] 1422–1425, 2000.

[90125] 15912.

[90126] Jha, K. K.; Ozer, H. L.: Genetic studies with a mutant
mouse cell,

[90127] ts-2 Balb-3T3, with a temperature-sensitive defect in
DNA synthesis.

[90128] (Abstract) Genetics 86: s32–s33, 1977.

[90129] 15913.

[90130] Jha, K. K.; Siniscalco, M.; Ozer, H. L.: Temperature-sensi-
tive

[90131] mutants of Balb-3T3 cells. III. Hybrids between ts2 and
other mouse

[90132] mutant cells affected in DNA synthesis and correction of
ts2 defect

[90133] by human X chromosome. Somat. Cell Genet. 6: 603–614,

1980.

[90134] 15914.

[90135] Allen, F. H.; Krabbe, S. M. R.; Corcoran, P. A.: A new phenotype

[90136] (McLeod) in the Kell blood-group system. Vox Sang. 6: 555-560, 1961.

[90137] 15915.

[90138] Bertelson, C. J.; Pogo, A. O.; Chaudhuri, A.; Marsh, W. L.; Redman,

[90139] C. M.; Banerjee, D.; Symmans, W. A.; Simon, T.; Frey, D.; Kunkel,

[90140] L. M.: Localization of the McLeod locus (XK) within Xp21 by deletion

[90141] analysis. Am. J. Hum. Genet. 42: 703-711, 1988.

[90142] 15916.

[90143] Danek, A.; Rubio, J. P.; Rampoldi, L.; Ho, M.; Dobson-Stone, C.;

[90144] Tison, F.; Symmans, W. A.; Oechsner, M.; Kalckreuth, W.; Watt, J.

[90145] M.; Corbett, A. J.; Hamdalla, H. H. M.; Marshall, A. G.; Sutton, I.;

[90146] Dotti, M. T.; Malandrini, A.; Walker, R. H.; Daniels, G.;

Monaco,

[90147] A. P.: McLeod neuroacanthocytosis: genotype and phenotype. Ann.

[90148] Neurol. 50: 755–764, 2001.

[90149] 15917.

[90150] Danek, A.; Witt, T. N.; Stockmann, H. B. A. C.; Weiss, B. J.; Schotland,

[90151] D. L.; Fischbeck, K. H.: Normal dystrophin in McLeod myopathy. Ann.

[90152] Neurol. 28: 720–722, 1990.

[90153] 15918.

[90154] Densen, P.; Wilkinson-Kroovand, S.; Mandell, G. L.; Sullivan, G.;

[90155] Oyen, R.; Marsh, W. L.: Kx: its relationship to chronic granulomatous

[90156] disease and genetic linkage with Xg. Blood 58: 34–37, 1981.

[90157] 15919.

[90158] de Saint-Basile, G.; Bohler, M. C.; Fischer, A.; Cartron, J.; Dufier,

[90159] J. L.; Griscelli, C.; Orkin, S. H.: Xp21 DNA microdeletion in

a patient

[90160] with chronic granulomatous disease, retinitis pigmentosa,
and McLeod

[90161] phenotype. Hum. Genet. 80: 85-89, 1988.

[90162] 15920.

[90163] Faillace, R. T.; Kingston, W. J.; Nanda, N. C.; Griggs, R. C.:

[90164] Cardiomyopathy associated with the syndrome of amy-
otrophic chorea

[90165] and acanthocytosis. Ann. Intern. Med. 96: 616-617, 1982.

[90166] 15921.

[90167] Frey, D.; Machler, M.; Seger, R.; Schmid, W.; Orkin, S. H.:
Gene

[90168] deletion in a patient with chronic granulomatous disease
and McLeod

[90169] syndrome: fine mapping of the Xk gene locus. Blood 71:
252-255,

[90170] 1988.

[90171] 15922.

[90172] Giblett, E. R.; Klebanoff, S. J.; Pincus, S. H.; Swanson, J.;

[90173] Park, B. H.; McCullough, J.: Kell phenotypes in chronic

granulomatous

[90174] disease: a potential transfusion hazard. Lancet I:
1235–1236, 1971.

[90175] 15923.

[90176] Hanaoka, N.; Yoshida, K.; Nakamura, A.; Furihata, K.; Seo,
T.;

[90177] Tani, Y.; Takahashi, J.; Ikeda, S.; Hanyu, N.: A novel
frameshift

[90178] mutation in the McLeod syndrome gene in a Japanese
family. J. Neurol.

[90179] Sci. 165: 6–9, 1999.

[90180] 15924.

[90181] Ho, M.; Chelly, J.; Carter, N.; Danek, A.; Crocker, P.;
Monaco,

[90182] A. P.: Isolation of the gene for McLeod syndrome that en-
codes a novel

[90183] membrane transport protein. Cell 77: 869–880, 1994.

[90184] 15925.

[90185] Almind, K.; Ahlgren, M. G.; Hansen, T.; Urhammer, S. A.;
Clausen,

[90186] J. O.; Pedersen, O.: Discovery of a Met300Val variant in

Shc and

[90187] studies of its relationship to birth weight and length, impaired insulin

[90188] secretion, insulin resistance, and type 2 diabetes mellitus. J. Clin.

[90189] Endocr. Metab. 84: 2241–2244, 1999.

[90190] 15926.

[90191] Harun, R. B.; Smith, K. K.; Leek, J. P.; Markham, A. F.; Norris,

[90192] A.; Morrison, J. F. J.: Characterization of human SHC p66 cDNA and

[90193] its processed pseudogene mapping to Xq12–q13.1. Genomics 42: 349–352,

[90194] 1997.

[90195] 15927.

[90196] McGlade, J.; Cheng, A.; Pelicci, G.; Pelicci, P. G.; Pawson, T.

[90197] : Shc proteins are phosphorylated and regulated by the v-src and v-fps

[90198] protein-tyrosine-kinases. Proc. Nat. Acad. Sci. 89: 8869–8873, 1992.

[90199] 15928.

- [90200] Migliaccio, E.; Giorgio, M.; Mele, S.; Pelicci, G.; Reboldi, P.;
- [90201] Pandolfi, P. P.; Lanfranccone, L.; Pelicci, P. G.: The p66(shc)
adaptor
- [90202] protein controls oxidative stress response and life span in
mammals. Nature 402:
- [90203] 309–313, 1999.
- [90204] 15929.
- [90205] Nemoto, S.; Finkel, T.: Redox regulation of forkhead pro-
teins
- [90206] through a p66shc-dependent signaling pathway. Science
295: 2450–2452,
- [90207] 2002.
- [90208] 15930.
- [90209] Pelicci, G.; Lanfranccone, L.; Grignani, F.; McGlade, J.; Cav-
allo,
- [90210] F.; Forni, G.; Nicoletti, I.; Grignani, F.; Pawson, T.; Pelicci,
P.
- [90211] G.: A novel transforming protein (SHC) with an SH2 do-
main is implicated
- [90212] in mitogenic signal transduction. Cell 70: 93–104, 1992.
- [90213] 15931.

- [90214] Yulug, I. G.; Egan, S. E.; See, C. G.; Fisher, E. M. C.: A human
- [90215] SHC-related sequence maps to chromosome 17, the SHC gene maps to chromosome
- [90216] 1. Hum. Genet. 96: 245-248, 1995.
- [90217] 15932.
- [90218] Zhang, L.; Camerini, V.; Bender, T. P.; Ravichandran, K. S.: A
- [90219] nonredundant role for the adapter protein Shc in thymic T cell development. Nature
- [90220] Immun. 3: 749-755, 2002.
- [90221] 15933.
- [90222] Selig, S.; Bruno, S.; Scharf, J. M.; Wang, C. H.; Vitale, E.; Gilliam,
- [90223] T. C.; Kunkel, L. M.: Expressed cadherin pseudogenes are localized
- [90224] to the critical region of the spinal muscular atrophy gene. Proc.
- [90225] Nat. Acad. Sci. 92: 3702-3706, 1995.
- [90226] 15934.
- [90227] Thompson, T. G.; DiDonato, C. J.; Simard, L. R.; Ingraham,

S. E.;

[90228] Burghes, A. H. M.; Crawford, T. O.; Rochette, C.; Mendell, J. R.;

[90229] Wasmuth, J. J.: A novel cDNA detects homozygous microdeletions in

[90230] greater than 50% of type I spinal muscular atrophy patients.. Nature

[90231] Genet. 9: 56-62, 1995.

[90232] 15935.

[90233] Abramovitz, M.; Boie, Y.; Nguyen, T.; Rushmore, T. H.; Bayne, M.

[90234] A.; Metters, K. M.; Slipetz, D. M.; Grygorczyk, R.: Cloning and expression

[90235] of a cDNA for the human prostanoid FP receptor. J. Biol. Chem. 269:

[90236] 2632-2636, 1994.

[90237] 15936.

[90238] Betz, R.; Lagercrantz, J.; Kedra, D.; Dumanski, J. P.; Nordenskjold,

[90239] A.: Genomic structure, 5-prime flanking sequences, and precise localization

[90240] in 1p31.1 of the human prostaglandin F receptor gene.

Biochem. Biophys.

[90241] Res. Commun. 254: 413–416, 1999.

[90242] 15937.

[90243] Sugimoto, Y.; Yamasaki, A.; Segi, E.; Tsuboi, K.; Aze, Y.;
Nishimura,

[90244] T.; Oida, H.; Yoshida, N.; Tanaka, T.; Katsuyama, M.; Ha-
sumoto, K.;

[90245] Murata, T.; Hirata, M.; Ushikubi, F.; Negishi, M.; Ichikawa,
A.; Narumiya,

[90246] S.: Failure of parturition in mice lacking the prostaglandin
F receptor. Science 277:

[90247] 681–683, 1997.

[90248] 15938.

[90249] Jean, L.; Smih, F.; Olivier, E.; Soury, E.; Simon–Chazottes,
D.;

[90250] Guenet, J. L.; Mattei, M. G.; Salier, J. P.: Comparative as-
signments

[90251] of the genes of the inter- α -inhibitor family in human
and mouse:

[90252] ITIH4 is close to ITIH1 and ITIH3, on HSA 3 and MMU 14.
Genomics 41:

[90253] 139–140, 1997.

[90254] 15939.

[90255] Nishimura, H.; Kakizaki, I.; Muta, T.; Sasaki, N.; Pu, P. X.; Yamashita,

[90256] T.; Nagasawa, S.: cDNA and deduced amino acid sequence of human PK-120,

[90257] a plasma kallikrein-sensitive glycoprotein. FEBS Lett. 357: 207-211,

[90258] 1995.

[90259] 15940.

[90260] Saguchi, K.; Tobe, T.; Hashimoto, K.; Sano, Y.; Nakano, Y.; Miura,

[90261] N.-H.; Tomita, M.: Cloning and characterization of cDNA for inter-alpha-trypsin

[90262] inhibitor family heavy chain-related protein (IHRP), a novel human

[90263] plasma glycoprotein. J. Biochem. 117: 14-18, 1995.

[90264] 15941.

[90265] Tobe, T.; Saguchi, K.; Hashimoto, K.; Miura, N.-H.; Tomita, M.;

[90266] Li, F.; Wang, Y.; Minoshima, S.; Shimizu, N.: Mapping of human inter-alpha-trypsin

- [90267] inhibitor family heavy chain-related protein gene (ITIH1)
to human
- [90268] chromosome 3p21-p14. Cytogenet. Cell. Genet. 71:
296-298, 1995.
- [90269] 15942.
- [90270] Geppert, M.; Khvotchev, M.; Krasnoperov, V.; Goda, Y.;
Missler,
- [90271] M.; Hammer, R. E.; Ichtchenko, K.; Petrenko, A. G.; Sudhof,
T. C.
- [90272] : Neurexin I-alpha is a major alpha-latrotoxin receptor
that cooperates
- [90273] in alpha-latrotoxin action. J. Biol. Chem. 273: 1705-1710,
1998.
- [90274] 15943.
- [90275] Ichtchenko, K.; Hata, Y.; Nguyen, T.; Ullrich, B.; Missler,
M.;
- [90276] Moomaw, C.; Sudhof, T. C.: Neuroligin 1: a splice site-
specific ligand
- [90277] for beta-neurexins. Cell 81: 435-443, 1995.
- [90278] 15944.
- [90279] Kleiderlein, J. J.; Nisson, P. E.; Jessee, J.; Li, W.-B.; Becker,

- [90280] K. G.; Derby, M. L.; Ross, C. A.; Margolis, R. L.: CCG repeats in
- [90281] cDNAs from human brain. Hum. Genet. 103: 666–673, 1998. Note: Erratum:
- [90282] Hum. Genet. 104: 113 only, 1999.
- [90283] 15945.
- [90284] Schoneberg, T.; Yun, J.; Wenkert, D.; Wess, J.: Functional rescue
- [90285] of mutant V2 vasopressin receptors causing nephrogenic diabetes insipidus
- [90286] by a co-expressed receptor polypeptide. EMBO J. 15: 1283–1291, 1996.
- [90287] 15946.
- [90288] Aly, A. M.; Higuchi, M.; Kasper, C. K.; Kazazian, H. H., Jr.; Antonarakis,
- [90289] S. E.; Hoyer, L. W.: Hemophilia A due to mutations that create new
- [90290] N-glycosylation sites. Proc. Nat. Acad. Sci. 89: 4933–4937, 1992.
- [90291] 15947.
- [90292] Greenberg, J. H.; Turner, C. G., II; Zegura, S. L.: The settlement

[90293] of the Americas: a comparison of the linguistic, dental,
and genetic

[90294] evidence. *Curr. Anthropol.* 27: 477–497, 1986.

[90295] 15948.

[90296] Margolis, R.: Personal Communication. Baltimore, Md.
3/29/2000.

[90297] 15949.

[90298] Missler, M.; Sudhof, T. C.: Neurexins: three genes and
1001 products. *Trends*

[90299] *Genet.* 14: 20–26, 1998.

[90300] 15950.

[90301] Ikeshima, H.; Imai, S.; Shimoda, K.; Hata, J.; Takano, T.:
Expression

[90302] of a MADS box gene, MEF2D, in neurons of the mouse
central nervous

[90303] system: implication of its binary function in myogenic and
neurogenic

[90304] cell lineages. *Neurosci. Lett.* 200: 117–120, 1995.

[90305] 15951.

[90306] Delhase, M.; Hayakawa, M.; Chen, Y.; Karin, M.: Positive
and negative

- [90307] regulation of I-kappa-B kinase activity through IKK-beta subunit phosphorylation. Science 284:
- [90308] 309-312, 1999.
- [90309] 15952.
- [90310] DiDonato, J. A.; Hayakawa, M.; Rothwarf, D. M.; Zandi, E.; Karin,
- [90311] M.: A cytokine-responsive IkappaB kinase that activates the transcription
- [90312] factor NF-kappaB. Nature 388: 548-554, 1997.
- [90313] 15953.
- [90314] Mercurio, F.; Zhu, H.; Murray, B. W.; Shevchenko, A.; Ben-
- [90315] nett,
- [90315] B. L.; Li, J.; Young, D. B.; Barbosa, M.; Mann, M.; Manning, A.; Rao,
- [90316] A.: IKK-1 and IKK-2: cytokine-activated I-kappa-B kinases essential
- [90317] for NF-kappa-B activation. Science 278: 860-866, 1997.
- [90318] 15954.
- [90319] Matsushima, A.; Kaisho, T.; Rennert, P. D.; Nakano, H.; Kurosawa,
- [90320] K.; Uchida, D.; Takeda, K.; Akira, S.; Matsumoto, M.: Es-

sential role

[90321] of nuclear factor (NF)-kappa-B-inducing kinase and inhibitor of kappa-B

[90322] (I-kappa-B) kinase alpha in NF-kappa-B activation through lymphotoxin

[90323] beta receptor, but not through tumor necrosis factor receptor I. J.

[90324] Exp. Med. 193: 631-636, 2001.

[90325] 15955.

[90326] Brzezinski, A.: Melatonin in humans. New Eng. J. Med. 336: 186-195,

[90327] 1997.

[90328] 15956.

[90329] Liu, R.-Y.; Zhou, J.-N.; van Heerikhuize, J.; Hofman, M. A.; Swaab,

[90330] D. F.: Decreased melatonin levels in postmortem cerebrospinal fluid

[90331] in relation to aging, Alzheimer's disease, and apolipoprotein E-epsilon-4/4

[90332] genotype. J. Clin. Endocr. Metab. 84: 323-327, 1999.

[90333] 15957.

- [90334] Nelson, C. S.; Ikeda, M.; Gompf, H. S.; Robinson, M. L.; Fuchs,
- [90335] N. K.; Yoshioka, T.; Neve, K. A.; Allen, C. N.: Regulation of melatonin
- [90336] 1a receptor signaling and trafficking by asparagine-124. Molec. Endocr. 15:
- [90337] 1306-1317, 2001.
- [90338] 15958.
- [90339] Reppert, S. M.; Weaver, D. R.: Melatonin madness. Cell 83: 1059-1062,
- [90340] 1995.
- [90341] 15959.
- [90342] Reppert, S. M.; Weaver, D. R.; Ebisawa, T.: Cloning and characterization
- [90343] of a mammalian melatonin receptor that mediates reproductive and circadian
- [90344] responses. Neuron 13: 1177-1185, 1994.
- [90345] 15960.
- [90346] Deng, Z.; Centola, M.; Chen, X.; Sood, R.; Vedula, A.; Fischel-Ghodsian,
- [90347] N.; Kastner, D. L.: Identification of two Kruppel-related

zinc finger

- [90348] genes (ZNF200 and ZNF210) from human chromosome 16p13.3. Genomics 53: 97–103, 1998.
- [90349] 15961.
- [90350] 15961.
- [90351] Muller-Pillasch, F.; Zimmerhackl, F.; Lacher, U.; Schultz, N.;
- [90352] Hameister, H.; Varga, G.; Friess, H.; Buchler, M.; Adler, G.; Gress,
- [90353] T. M.: Cloning of novel transcripts of the human guanine-nucleotide-exchange
- [90354] factor Mss4: in situ chromosomal mapping and expression in pancreatic
- [90355] cancer. Genomics 46: 389–396, 1997.
- [90356] 15962.
- [90357] Yu, H.; Schreiber, S. L.: Cloning, Zn(2+) binding, and structural
- [90358] characterization of the guanine nucleotide exchange factor human Mss4. Biochemistry 34:
- [90359] 9103–9110, 1995.
- [90360] 15963.

- [90361] Kordes, E.; Savelyeva, L.; Schwab, M.; Rommelaere, J.; Jau-
niaux,
- [90362] J.-C.; Cziepluch, C.: Isolation and characterization of hu-
man SGT
- [90363] and identification of homologues in *Saccharomyces cere-*
visiae and *Caenorhabditis*
- [90364] *elegans*. *Genomics* 52: 90–94, 1998.
- [90365] 15964.
- [90366] Yabe, D.; Nakamura, T.; Kanazawa, N.; Tashiro, K.; Honjo,
T.:
- [90367] Calumenin, a Ca(2+)-binding protein retained in the en-
doplasmic reticulum
- [90368] with a novel carboxyl-terminal sequence, *HDEF. J. Biol.*
Chem. 272:
- [90369] 18232–18239, 1997.
- [90370] 15965.
- [90371] Yabe, D.; Taniwaki, M.; Nakamura, T.; Kanazawa, N.;
Tashiro, K.;
- [90372] Honjo, T.: Human calumenin gene (CALU): cDNA isolation
and chromosomal
- [90373] mapping to 7q32. *Genomics* 49: 331–333, 1998.

[90374] 15966.

[90375] Angelin-Duclos, C.; Cattoretti, G.; Lin, K.-I.; Calame, K.:
Commitment

[90376] of B lymphocytes to a plasma cell fate is associated with
Blimp-1

[90377] expression in vivo. J. Immun. 165: 5462-5471, 2000.

[90378] 15967.

[90379] Chang, D. H.; Angelin-Duclos, C.; Calame, K.: BLIMP-1:
trigger

[90380] for differentiation of myeloid lineage. Nature Immun. 1:
169-176,

[90381] 2000.

[90382] 15968.

[90383] Huang, S.: Blimp-1 is the murine homolog of the human
transcriptional

[90384] repressor PRDI-BF1. (Letter) Cell 78: 9 only, 1994.

[90385] 15969.

[90386] Keller, A. D.; Maniatis, T.: Identification and characteriza-
tion

[90387] of a novel repressor of beta-interferon gene expression.
Genes Dev. 5:

- [90388] 868–879, 1991.
- [90389] 15970.
- [90390] Mock, B. A.; Liu, L.; Le Paslier, D.; Huang, S.: The B-lymphocyte
- [90391] maturation promoting transcription factor BLIMP1/PRDI-BF1 maps to
- [90392] D6S447 on human chromosome 6q21–q22.1 and the syntenic region of mouse
- [90393] chromosome 10. Genomics 37: 24–28, 1996.
- [90394] 15971.
- [90395] Ito, T.; Yang, M.; May, W. S.: RAX, a cellular activator for double-stranded
- [90396] RNA-dependent protein kinase during stress signaling. J. Biol. Chem. 274:
- [90397] 15427–15432, 1999.
- [90398] 15972.
- [90399] Patel, R. C.; Sen, G. C.: PACT, a protein activator of the interferon-induced
- [90400] protein kinase, PKR. EMBO J. 17: 4379–4390, 1998.
- [90401] 15973.

- [90402] Scott, A. F.: Personal Communication. Baltimore, Md. 2001.
- [90403] 15974.
- [90404] Shearman, L. P.; Zylka, M. J.; Weaver, D. R.; Kolakowski, L. F.,
- [90405] Jr.; Reppert, S. M.: Two period homologs: circadian expression and
- [90406] photic regulation in the suprachiasmatic nuclei. Neuron 19: 1261–1269,
- [90407] 1997.
- [90408] 15975.
- [90409] Toh, K. L.; Jones, C. R.; He, Y.; Eide, E. J.; Hinz, W. A.; Virshup,
- [90410] D. M.; Ptacek, L. J.; Fu, Y.-H.: An hPer2 phosphorylation site mutation
- [90411] in familial advanced sleep phase syndrome. Science 291: 1040–1043,
- [90412] 2001.
- [90413] 15976.
- [90414] Engelender, S.; Wanner, T.; Kleiderlein, J. J.; Wakabayashi, K.;

- [90415] Tsuji, S.; Takahashi, H.; Ashworth, R.; Margolis, R. L.; Ross, C.
- [90416] A.: Organization of the human synphilin-1 gene, a candidate for Parkinson's
- [90417] disease. *Mammalian Genome* 11: 763-766, 2000.
- [90418] 15977.
- [90419] Kitao, S.; Ohsugi, I.; Ichikawa, K.; Goto, M.; Furuichi, Y.; Shimamoto,
- [90420] A.: Cloning of two new human helicase genes of the RecQ family: biological
- [90421] significance of multiple species in higher eukaryotes. *Genomics* 54:
- [90422] 443-452, 1998.
- [90423] 15978.
- [90424] Mohaghegh, P.; Hickson, I. D.: DNA helicase deficiencies associated
- [90425] with cancer predisposition and premature ageing disorders. *Hum. Molec.*
- [90426] *Genet.* 10: 741-746, 2001.
- [90427] 15979.
- [90428] Chien, W.; Pei, L.: A novel binding factor facilitates nuclear

- [90429] translocation and transcriptional activation function of the pituitary
- [90430] tumor-transforming gene product. J. Biol. Chem. 275: 19422–19427,
- [90431] 2000.
- [90432] 15980.
- [90433] Yaspo, M.–L.; Aaltonen, J.; Horelli-Kuitunen, N.; Peltonen, L.;
- [90434] Lehrach, H.: Cloning of a novel human putative type Ia integral membrane
- [90435] protein mapping to 21q22.3. Genomics 49: 133–136, 1998.
- [90436] 15981.
- [90437] Yaspo, M.–L.; Gellen, L.; Mott, R.; Korn, B.; Nizetic, D.; Poustka,
- [90438] A. M.; Lehrach, H.: Model for a transcript map of human chromosome
- [90439] 21: isolation of new coding sequences from exon and enriched cDNA
- [90440] libraries. Hum. Molec. Genet. 4: 1291–1304, 1995.
- [90441] 15982.

- [90442] Su, K.; Kyaw, H.; Fan, P.; Zeng, Z.; Shell, B. K.; Carter, K. C.;
- [90443] Li, Y.: Isolation, characterization, and mapping of two human potassium
- [90444] channels. *Biochem. Biophys. Res. Commun.* 241: 675–681, 1997.
- [90445] 15983.
- [90446] Ghiso, J. A.; Holton, J.; Miravalle, L.; Calero, M.; Lashley, T.;
- [90447] Vidal, R.; Houlden, H.; Wood, N.; Neubert, T. A.; Rostagno, A.; Plant,
- [90448] G.; Revesz, T.; Frangione, B.: Systemic amyloid deposits in familial
- [90449] British dementia. *J. Biol. Chem.* 276: 43909–43914, 2001.
- [90450] 15984.
- [90451] Stromgren, E.; Dalby, A.; Dalby, M.; Ranheim, B. :*Acta Neurol.*
- [90452] *Scand.* 46 (suppl. 43): 97–98, 1970.
- [90453] 15985.
- [90454] Gruber, A. D.; Schreur, K. D.; Ji, H.–L.; Fuller, C. M.; Pauli,
- [90455] B. U.: Molecular cloning and transmembrane structure of

hCLCA2 from

[90456] human lung, trachea, and mammary gland. *Am. J. Physiol.* 276: C1261–C1270,

[90457] 1999.

[90458] 15986.

[90459] Claustres, M.; Gerrard, B.; White, M. B.; Desgeorges, M.; Kjellberg,

[90460] P.; Rollin, B.; Dean, M.: A new mutation (1078delT) in exon 7 of

[90461] the CFTR gene in a southern French adult with cystic fibrosis. *Genomics* 13:

[90462] 907–908, 1992.

[90463] 15987.

[90464] Claustres, M.; Laussel, M.; Desgeorges, M.; Giansily, M.; Culard,

[90465] J.-F.; Razakatsara, G.; Demaille, J.: Analysis of the 27 exons and

[90466] flanking regions of the cystic fibrosis gene: 40 different mutations

[90467] account for 91.2% of the mutant alleles in Southern France. *Hum.*

[90468] *Molec. Genet.* 2: 1209–1213, 1993.

[90469] 15988.

[90470] Costes, B.; Girodon, E.; Ghanem, N.; Flori, E.; Jardin, A.; Soufir,

[90471] J. C.; Goossens, M.: Frequent occurrence of the CFTR intron 8 (TG)_n

[90472] 5T allele in men with congenital bilateral absence of the vas deferens. Europ.

[90473] J. Hum. Genet. 3: 285-293, 1995.

[90474] 15989.

[90475] Cromwell, O.; Walport, M. J.; Morris, H. R.; Taylor, G. W.; Hodson,

[90476] M. E.; Batten, J.; Kay, A. B.: Identification of leukotrienes D and

[90477] B in sputum from cystic fibrosis patients. Lancet II: 164-165, 1981.

[90478] 15990.

[90479] Cuppens, H.; Lin, W.; Jaspers, M.; Costes, B.; Teng, H.; Vankeerberghen,

[90480] A.; Jorissen, M.; Droogmans, G.; Reynaert, I.; Goossens, M.; Nilius,

[90481] B.; Cassiman, J.-J.: Polyvariant mutant cystic fibrosis transmembrane

- [90482] conductance regulator genes: the polymorphic (TG)_n locus explains
- [90483] the partial penetrance of the T5 polymorphism as a disease mutation. J.
- [90484] Clin. Invest. 101: 487–496, 1998.
- [90485] 15991.
- [90486] Cuppens, H.; Marynen, P.; De Boeck, C.; De Baets, F.; Eggermont,
- [90487] E.; Van den Berghe, H.; Cassiman, J. J.: A child, homozygous for
- [90488] a stop codon in exon 11, shows milder cystic fibrosis symptoms than
- [90489] her heterozygous nephew. J. Med. Genet. 27: 717–719, 1990.
- [90490] 15992.
- [90491] Cutting, G. R.; Kasch, L. M.; Rosenstein, B. J.; Tsui, L.-C.;
- [90492] Kazazian, H. H., Jr.; Antonarakis, S. E.: Two patients with cystic
- [90493] fibrosis, nonsense mutations in each cystic fibrosis gene, and mild
- [90494] pulmonary disease. New Eng. J. Med. 323: 1685–1689, 1990.

- [90495] 15993.
- [90496] Cutting, G. R.; Kasch, L. M.; Rosenstein, B. J.; Zielenski, J.;
- [90497] Tsui, L.-C.; Antonarakis, S. E.; Kazazian, H. H., Jr.: A cluster
ter
- [90498] of cystic fibrosis mutations in the first nucleotide-binding
fold
- [90499] of the cystic fibrosis conductance regulator protein. Nature 346:
ture 346:
- [90500] 366-369, 1990.
- [90501] 15994.
- [90502] Daigneault, J.; Aubin, G.; Simard, F.; De Braekeleer, M.:
Genetic
- [90503] epidemiology of cystic fibrosis in Saguenay-Lac-St-Jean
(Quebec, Canada). Clin.
- [90504] Genet. 40: 298-303, 1991.
- [90505] 15995.
- [90506] Dean, M.; White, M. B.; Amos, J.; Gerrard, B.; Stewart, C.;
Khaw,
- [90507] K.-T.; Leppert, M.: Multiple mutations in highly conserved
residues
- [90508] are found in mildly affected cystic fibrosis patients. Cell

61:

[90509] 863–870, 1990.

[90510] 15996.

[90511] Dean, M.; White, M. B.; Gerrard, B.; Amos, J.; Milunsky, A.:

[90512] A 22–bp deletion in the coding region of the cystic fibrosis gene. *Genomics* 13:

[90513] 235–236, 1992.

[90514] 15997.

[90515] Delaney, S. J.; Alton, E. W. F. W.; Smith, S. N.; Lunn, D. P.;

[90516] Farley, R.; Lovelock, P. K.; Thomson, S. A.; Hume, D. A.; Lamb, D.;

[90517] Porteous, D. J.; Dorin, J. R.; Wainwright, B. J.: Cystic fibrosis

[90518] mice carrying the missense mutation G551D replicate human genotype–phenotype

[90519] correlations. *EMBO J.* 15: 955–963, 1996.

[90520] 15998.

[90521] de Meeus, A.; Guittard, C.; Desgeorges, M.; Carles, S.; Demaille,

[90522] J.; Claustres, M.: Linkage disequilibrium between the

M470V variant

[90523] and the IVS8 polyT alleles of the CFTR gene in CBAVD. J.

Med. Genet. 35:

[90524] 594–596, 1998.

[90525] 15999.

[90526] Denning, G. M.; Anderson, M. P.; Amara, J. F.; Marshall, J.;

Smith,

[90527] A. E.; Welsh, M. J.: Processing of mutant cystic fibrosis

transmembrane

[90528] conductance regulator is temperature-sensitive. Nature

358: 761–764,

[90529] 1992.

[90530] 16000.

[90531] Devor, D. C.; Schultz, B. D.: Ibuprofen inhibits cystic fibro-

sis

[90532] transmembrane conductance regulator-mediated Cl(–) se-

cretion. J.

[90533] Clin. Invest. 102: 679–687, 1998.

[90534] 16001.

[90535] Dickinson, P.; Smith, S. N.; Webb, S.; Kilanowski, F. M.;

Campbell,

- [90536] I. J.; Taylor, M. S.; Porteous, D. J.; Willemsen, R.; de Jonge, H.
- [90537] R.; Farley, R.; Alton, E. W. F. W.; Dorin, J. R.: The severe G480C
- [90538] cystic fibrosis mutation, when replicated in the mouse, demonstrates
- [90539] mistrafficking, normal survival and organ-specific bio-electrics. Hum.
- [90540] Molec. Genet. 11: 243-251, 2002.
- [90541] 16002.
- [90542] Nicolino, P. M.; Dupin, H.; Macebeo, V.; Treppoz, S.; Chatelain,
- [90543] P. G.; Wolcott-Rallison syndrome (diabetes mellitus and spondyloepiphyseal
- [90544] dysplasia): a plausible existence of a gene(s) important for the
- [90545] mutation of neonatal pancreatic beta cell function. (Abstract) Hormone
- [90546] Res. 50 (suppl. 3): 77 only, 1998.
- [90547] 16003.
- [90548] Shi, Y.; An, J.; Liang, J.; Hayes, S. E.; Sandusky, G. E.; Stramm,

- [90549] L. E.; Yang, N. N.: Characterization of a mutant pancreatic eIF-2-alpha
- [90550] kinase, PEK, and co-localization with somatostatin in islet delta
- [90551] cells. J. Biol. Chem. 274: 5723-5730, 1999.
- [90552] 16004.
- [90553] Shi, Y.; Vатtem, K. M.; Sood, R.; An, J.; Liang, J.; Stramm, L.;
- [90554] Wek, R. C.: Identification and characterization of pancreatic eukaryotic
- [90555] initiation factor 2 alpha-subunit kinase, PEK, involved in translational
- [90556] control. Molec. Cell. Biol. 18: 7499-7509, 1998.
- [90557] 16005.
- [90558] Csoka, A. B.; Scherer, S. W.; Stern, R.: Expression analysis of
- [90559] six paralogous human hyaluronidase genes clustered on chromosomes
- [90560] 3p21 and 7q31. Genomics 60: 356-361, 1999.
- [90561] 16006.
- [90562] Dolganov, G. M.; Maser, R. S.; Novikov, A.; Tosto, L.;

Chong, S.;

[90563] Bressan, D. A.; Petrini, J. H. J.: Human Rad50 is physically associated

[90564] with human Mre11: identification of a conserved multi-protein complex

[90565] implicated in recombinational DNA repair. *Molec. Cell Biol.* 16:

[90566] 4832–4841, 1996.

[90567] 16007.

[90568] Hopfner, K.-P.; Craig, L.; Moncalian, G.; Zinkel, R. A.; Usui,

[90569] T.; Owen, B. A. L.; Karcher, A.; Henderson, B.; Bodmer, J.-L.; McMurray,

[90570] C. T.; Carney, J. P.; Petrini, J. H. J.; Tainer, J. A.: The Rad50

[90571] zinc-hook is a structure joining Mre11 complexes in DNA recombination

[90572] and repair. *Nature* 418: 562–566, 2002.

[90573] 16008.

[90574] Luo, G.; Yao, M. S.; Bender, C. F.; Mills, M.; Bladl, A. R.; Bradley,

[90575] A.; Petrini, J. H. J.: Disruption of mRad50 causes embryonic stem

- [90576] cell lethality, abnormal embryonic development, and sensitivity to
- [90577] ionizing radiation. *Proc. Nat. Acad. Sci.* 96: 7376–7381, 1999.
- [90578] 16009.
- [90579] Trujillo, K. M.; Yuan, S.-S. F.; Lee, E. Y.-H. P.; Sung, P.: Nuclease
- [90580] activities in a complex of human recombination and DNA repair factors
- [90581] Rad50, Mre11, and p95. *J. Biol. Chem.* 273: 21447–21450, 1998.
- [90582] 16010.
- [90583] Clark, J.; Lu, Y. J.; Sidhar, S. K.; Parker, C.; Gill, S.; Smedley,
- [90584] D.; Hamoudi, R.; Linehan, W.; Shipley, J.; Cooper, C.: Fusion of
- [90585] splicing factor genes PSF and NonO (p54nrb) to the TFE3 gene in papillary
- [90586] renal cell carcinoma. *Oncogene* 15: 2233–2239, 1997.
- [90587] 16011.
- [90588] Heimann, P.; El Housni, H.; Ogur, G.; Weterman, M. A. J.; Petty,

- [90589] E. M.; Vassart, G.: Fusion of a novel gene, RCC17, to the TFE3 gene
- [90590] in t(X;17)(p11.2;q25.3)-bearing papillary renal cell carcinomas. Cancer
- [90591] Res. 61: 4130-4135, 2001.
- [90592] 16012.
- [90593] Henthorn, P. S.; Stewart, C. C.; Kadesch, T.; Puck, J. M.: The
- [90594] gene encoding human TFE3, a transcription factor that binds the immunoglobulin
- [90595] heavy-chain enhancer, maps to Xp11.22. Genomics 11: 374-378, 1991.
- [90596] 16013.
- [90597] Joyama, S.; Ueda, T.; Shimizu, K.; Kudawara, I.; Mano, M.; Funai,
- [90598] H.; Takemura, K.; Yoshikawa, H.: Chromosome rearrangement at 17q25
- [90599] and Xp11.2 in alveolar soft-part sarcoma: a case report and review
- [90600] of the literature. Cancer 86: 1246-1250, 1999.
- [90601] 16014.

- [90602] Ladanyi, M.; Lui, M. Y.; Antonescu, C. R.; Krause-Boehm, A.; Meindl,
- [90603] A.; Argani, P.; Healey, J. H.; Ueda, T.; Yoshikawa, H.; Meloni-Ehrig,
- [90604] A.; Sorensen, P. H. B.; Mertens, F.; Mandahl, N.; van den Berghe,
- [90605] H.; Sciot, R.; Dal Cin, P.; Bridge, J.: The
der(17)t(X;17)(p11;q25)
- [90606] of human alveolar soft part sarcoma fuses the TFE3 transcription factor
- [90607] gene to ASPL, a novel gene at 17q25. *Oncogene* 20: 48–57, 2001.
- [90608] 16015.
- [90609] Macchi, P.; Notarangelo, L.; Giliani, S.; Strina, D.; Repetto,
- [90610] M.; Sacco, M. G.; Vezzoni, P.; Villa, A.: The genomic organization
- [90611] of the human transcription factor 3 (TFE3) gene. *Genomics* 28: 491–494,
- [90612] 1995.
- [90613] 16016.
- [90614] Shipley, J. M.; Birdsall, S.; Clark, J.; Crew, J.; Gill, S.; Linehan,

- [90615] M.; Gnarr, J.; Gisher, S.; Craig, I. W.; Cooper, C. S.: Mapping
[90616] the X chromosome breakpoint in two papillary renal cell
carcinoma
[90617] cell lines with a t(X;1)(p11.2;q21.2) and the first report of
a female
[90618] case. Cytogenet. Cell. Genet. 71: 280–284, 1995.
[90619] 16017.
[90620] Sidhar, S. K.; Clark, J.; Gill, S.; Hamoudi, R.; Crew, A. J.;
[90621] Gwilliam, R.; Ross, M.; Linehan, W. M.; Birdsall, S.; Shipley,
J.;
[90622] Cooper, C. S.: The t(X;1)(p11.2;q21.2) translocation in
papillary
[90623] renal cell carcinoma fuses a novel gene PRCC to the TFE3
transcription
[90624] factor gene. Hum. Molec. Genet. 5: 1333–1338, 1996.
[90625] 16018.
[90626] Larola, G.; Cuesta, R.; Brewer, G.; Schneider, R. J.: Control
[90627] of mRNA decay by heat shock–ubiquitin–proteasome
pathway. Science 284:
[90628] 499–502, 1999.

[90629] 16019.

[90630] Deutsch, D.; Palmon, A.; Dafni, L.; Fisher, L.; Termine, J. D.;

[90631] Young, M.: Cloning, sequencing, and characterization of tuftelin:

[90632] a novel acidic enamel protein. Connect. Tissue Res. 27: 121 only,

[90633] 1992.

[90634] 16020.

[90635] Genomics 44: 248-250, 1997.

[90636] 16021.

[90637] Deiss, L. P.; Feinstein, E.; Berissi, H.; Cohen, O.; Kimchi, A.

[90638] : Identification of a novel serine/threonine kinase and a novel 15-kD

[90639] protein as potential mediators of the gamma interferon-induced cell

[90640] death. Genes Dev. 9: 15-30, 1995.

[90641] 16022.

[90642] Feinstein, E.; Druck, T.; Kastury, K.; Berissi, H.; Goodart, S.

[90643] A.; Overhauser, J.; Kimchi, A.; Huebner, K.: Assignment of

DAP1 and

[90644] DAPK: genes that positively mediate programmed cell death triggered

[90645] by IFN- γ --to chromosome regions 5p12.2 (sic) and 9q34.1, respectively. Genomics 29:

[90646] 305-307, 1995.

[90647] 16023.

[90648] Raveh, T.; Droguett, G.; Horwitz, M. S.; DePinho, R. A.; Kimchi,

[90649] A.: DAP kinase activates a p19-ARF/p53-mediated apoptotic checkpoint

[90650] to suppress oncogenic transformation. Nature Cell Biol. 3: 1-7,

[90651] 2001.

[90652] 16024.

[90653] Simpson, D. J.; Clayton, R. N.; Farrell, W. E.: Preferential loss

[90654] of death associated protein kinase expression in invasive pituitary

[90655] tumours is associated with either CpG island methylation or homozygous

[90656] deletion. Oncogene 21: 1217-1224, 2002.

- [90657] 16025.
- [90658] Hughes, K. A.; Hurlstone, A. F. L.; Tobias, E. S.; McFarlane, R.;
- [90659] Black, D. M.: Absence of ST7 mutations in tumor-derived cell lines
- [90660] and tumors. *Nature Genet.* 29: 380–381, 2001.
- [90661] 16026.
- [90662] Thomas, N. A.; Choong, D. Y. H.; Jokubaitis, V. J.; Neville, P.
- [90663] J.; Campbell, I. G.: Mutation of the ST7 tumor suppressor gene on
- [90664] 7q31.1 is rare in breast, ovarian and colorectal cancers. *Nature*
- [90665] *Genet.* 29: 379–380, 2001.
- [90666] 16027.
- [90667] Zenklusen, J. C.; Conti, C. J.; Green, E. D.: Mutational and functional
- [90668] analyses reveal that ST7 is a highly conserved tumor-suppressor gene
- [90669] on human chromosome 7q31. *Nature Genet.* 27: 392–398, 2001.

[90670] 16028.

[90671] Zenklusen, J. C.; Rodriguez, L. V.; LaCava, M.; Wang, Z.;
Goldstein,

[90672] L. S.; Conti, C. J.: Novel susceptibility locus for mouse
hepatomas:

[90673] evidence for a conserved tumor suppressor gene. Genome
Res. 6: 1070–1076,

[90674] 1996.

[90675] 16029.

[90676] Zenklusen, J. C.; Weitzel, J. N.; Ball, H. G.; Conti, C. J.: Al-
lelic

[90677] loss at 7q31.1 in human primary ovarian carcinomas sug-
gests the existence

[90678] of a tumor suppressor gene. Oncogene 11: 359–363,
1995.

[90679] 16030.

[90680] Lee, P. L.; Gelbart, T.; West, C.; Adams, M.; Blackstone, R.;
Beutler,

[90681] E.: Three genes encoding zinc finger proteins on human
chromosome

[90682] 6p21.3: members of a new subclass of the Kruppel gene
family containing

- [90683] the conserved SCAN box domain. Genomics 43: 191–201, 1997.
- [90684] 16031.
- [90685] Deutsch, D.; Palmon, A.; Fisher, L. W.; Kolodny, N.; Ter-
mine, J.
- [90686] D.; Young, M. F.: Sequencing of bovine enamelin (tuftelin),
a novel
- [90687] acidic enamel protein. J. Biol. Chem. 266: 16021–16028,
1991.
- [90688] 16032.
- [90689] Deutsch, D.; Palmon, A.; Young, M. F.; Selig, S.; Kearns, W.
G.;
- [90690] Fisher, L. W.: Mapping of the human tuftelin (TUFT1) gene
to chromosome
- [90691] 1 by fluorescence in situ hybridization. (Abstract) Mam-
malian Genome 5:
- [90692] 461–462, 1994.
- [90693] 16033.
- [90694] MacDougall, M.; Simmons, D.; Dodds, A.; Knight, C.; Luan,
X.; Zeichner–David,
- [90695] M.; Zhang, C.; Ryu, O. H.; Qian, Q.; Simmer, J. P.; Hu,
C.–C.: Cloning,

[90696] characterization, and tissue expression pattern of mouse tuftelin

[90697] cDNA. J. Dent. Res. 77: 1970–1978, 1998.

[90698] 16034.

[90699] Fujii, H.; Kanno, H.; Hirono, A.; Shiomura, T.; Miwa, S.: A single

[90700] amino acid substitution (157gly-to-val) in a phosphoglycerate kinase

[90701] variant (PGK Shizuoka) associated with chronic hemolysis and myoglobinuria. Blood 79:

[90702] 1582–1585, 1992.

[90703] 16035.

[90704] Fujii, H.; Krietsch, W. K. G.; Yoshida, A.: A single amino acid

[90705] substitution (asp-to-asn) in a phosphoglycerate kinase variant (PGK

[90706] Munchen) associated with enzyme deficiency. J. Biol. Chem. 255:

- [90707] 6421–6423, 1980.
- [90708] 16036.
- [90709] Fujii, H.; Yoshida, A.: Molecular abnormality of phospho-glycerate
- [90710] kinase–Uppsala associated with chronic nonspherocytic hemolytic anemia. Proc.
- [90711] Nat. Acad. Sci. 77: 5461–5465, 1980.
- [90712] 16037.
- [90713] Gellin, J.; Benne, F.; Renard, C.; Vaiman, M.; Hors–Cayla, M.
- [90714] C.; Gillois, M.: Pig gene mapping: syntenry, attempt to assign the
- [90715] histocompatibility complex (SLA). (Abstract) Cytogenet. Cell Genet. 25:
- [90716] 159, 1979.
- [90717] 16038.
- [90718] Grzeschik, K.–H.; Allderdice, P. W.; Grzeschik, A.; Opitz, J.
- [90719] M.; Miller, O. J.; Siniscalco, M.: Cytological mapping of human X–linked
- [90720] genes by use of somatic cell hybrids involving an X–

autosome translocation. Proc.

[90721] Nat. Acad. Sci. 69: 69–73, 1972.

[90722] 16039.

[90723] Guis, M. S.; Karadsheh, N.; Mentzer, W. C.: Phosphoglycerate

[90724] kinase San Francisco: a new variant associated with hemolytic anemia

[90725] but not with neuromuscular manifestations. Am. J. Hemat. 25: 175–182,

[90726] 1987.

[90727] 16040.

[90728] Huijing, F.; Eicher, E. M.; Coleman, D. L.: Location of phosphorylase

[90729] kinase (Phk) in the mouse X-chromosome. Biochem. Genet. 9: 193–196,

[90730] 1973.

[90731] 16041.

[90732] Hutz, M. H.; Michelson, A. M.; Antonarakis, S. E.; Orkin, S. H.;

[90733] Kazazian, H. H., Jr.: Restriction site polymorphism in the phosphoglycerate

- [90734] kinase gene on the X chromosome. Hum. Genet. 66: 217–219, 1984.
- [90735] 16042.
- [90736] Konrad, P. N. J.; McCarthy, D. J.; Mauer, A. M.; Valentine, W.
- [90737] N.; Paglia, D. E.: Erythrocyte and leukocyte phosphoglycerate kinase
- [90738] deficiency with neurologic disease. J. Pediat. 82: 456–460, 1973.
- [90739] 16043.
- [90740] Kozak, L. P.; McLean, G. K.; Eicher, E. M.: X-linkage of phosphoglycerate
- [90741] kinase in the mouse. Biochem. Genet. 2: 41–47, 1974.
- [90742] 16044.
- [90743] Kraus, A. P.; Langston, M. F., Jr.; Lynch, B. L.: Red cell phosphoglycerate
- [90744] kinase deficiency: a new cause of non-spherocytic hemolytic anemia. Biochem.
- [90745] Biophys. Res. Commun. 30: 173–177, 1968.
- [90746] 16045.
- [90747] Krietsch, W. K.; Krietsch, H.; Kaiser, W.; Dunnwald, M.;

Kuntz,

[90748] G. W.; Duhn, J.; Bucher, T.: Hereditary deficiency of phosphoglycerate

[90749] kinase: a new variant in erythrocytes and leucocytes, not associated

[90750] with hemolytic anemia. *Europ. J. Clin. Invest.* 7: 427–435, 1977.

[90751] 16046.

[90752] Krietsch, W. K. G.; Eber, S. W.; Haas, B.; Rubbelt, W.; Kuntz,

[90753] G. W. K.: Characterization of a phosphoglycerate kinase deficiency

[90754] variant not associated with hemolytic anemia. *Am. J. Hum. Genet.* 32:

[90755] 364–373, 1980.

[90756] 16047.

[90757] Lay, A. J.; Jiang, X.-M.; Kisker, O.; Flynn, E.; Underwood, A.;

[90758] Condrón, R.; Hogg, P. J.: Phosphoglycerate kinase acts in tumour

[90759] angiogenesis as a disulphide reductase. *Nature* 408: 869–873, 2000.

- [90760] 16048.
- [90761] Maeda, M.; Bawle, E. V.; Kulkarni, R.; Beutler, E.; Yoshida, A.
- [90762] : Molecular abnormalities of a phosphoglycerate kinase variant generated
- [90763] by spontaneous mutation. Blood 79: 2759–2762, 1992.
- [90764] 16049.
- [90765] Maeda, M.; Yoshida, A.: Molecular defect of a phosphoglycerate
- [90766] kinase variant (PGK–Matsue) associated with hemolytic anemia: leu-to-pro
- [90767] substitution caused by T/A-to-C/G transition in exon 3. Blood 77:
- [90768] 1348–1352, 1991.
- [90769] 16050.
- [90770] Meera Khan, P.; Westerveld, A.; Grzeschik, K.–H.; Deys, B. F.;
- [90771] Garson, O. M.; Siniscalco, M.: X-linkage of human phosphoglycerate
- [90772] kinase confirmed in man–mouse and man–Chinese hamster somatic cell

- [90773] hybrids. *Am. J. Hum. Genet.* 23: 614–623, 1971.
- [90774] 16051.
- [90775] Michelson, A. M.; Blake, C. C. F.; Evans, S. T.; Orkin, S. H.
- [90776] : Structure of the human phosphoglycerate kinase gene
and the intron-mediated
- [90777] evolution and dispersal of the nucleotide-binding domain.
Proc. Nat.
- [90778] *Acad. Sci.* 82: 6965–6969, 1985.
- [90779] 16052.
- [90780] Michelson, A. M.; Markham, A. F.; Orkin, S. H.: Isolation
and
- [90781] DNA sequence of a full-length cDNA clone for human X
chromosome-encoded
- [90782] phosphoglycerate kinase. *Proc. Nat. Acad. Sci.* 80:
472–476, 1983.
- [90783] 16053.
- [90784] Miwa, S.; Nakawhima, K.; Oda, S.; Ogawa, H.; Nagafuji, H.;
Arima,
- [90785] M.; Okuna, T.; Nakashima, T.: Phosphoglycerate kinase
deficiency
- [90786] hereditary nonspherocytic hemolytic anemia: report of a

case found

[90787] in a Japanese family. *Acta Haemat. Jpn.* 35: 571–574, 1972.

[90788] 16054.

[90789] Ricciuti, F. C.; Ruddle, F. H.: Assignment of three gene loci
[90790] (PGK, HGPRT, and G6PD) to the long arm of the human X-
chromosome by

[90791] somatic cell genetics. *Genetics* 74: 661–678, 1973.

[90792] 16055.

[90793] Rosa, R.; George, C.; Fardeau, M.; Calvin, M. C.; Rapin, M.;
Rosa,

[90794] J.: A new case of phosphoglycerate kinase deficiency: PGK
Creteil

[90795] associated with rhabdomyolysis and lacking hemolytic
anemia. *Blood* 60:

[90796] 84–91, 1982.

[90797] 16056.

[90798] Saidi, N.; Hors-Cayla, M. C.; Van Cong, N.; Benne, F.:
Sheep

[90799] gene mapping by somatic cell hybridization. (Abstract)
Cytogenet.

- [90800] Cell Genet. 25: 200, 1979.
- [90801] 16057.
- [90802] Schwab, A. J.; Krietsch, W. K. G.: Linkage between phosphoglycerate
- [90803] kinase and Xg in a large German kindred. Hum. Genet. 38: 217-221,
- [90804] 1977.
- [90805] 16058.
- [90806] Shows, T. B.; Brown, J. A.: Human X-linked genes regionally mapped
- [90807] utilizing X-autosome translocations and somatic cell hybrids. Proc.
- [90808] Nat. Acad. Sci. 72: 2125-2129, 1975.
- [90809] 16059.
- [90810] Sugie, H.; Sugie, Y.; Ito, M.; Fukuda, T.: A novel missense mutation
- [90811] (837T-C) in the phosphoglycerate kinase gene of a patient with a myopathic
- [90812] form of phosphoglycerate kinase deficiency. J. Child Neurol. 13:
- [90813] 95-97, 1998.

- [90814] 16060.
- [90815] Buj-Bello, A.; Biancalana, V.; Moutou, C.; Laporte, J.; Mandel,
- [90816] J.-L.: Identification of novel mutations in the MTM1 gene causing
- [90817] severe and mild forms of X-linked myotubular myopathy. Hum. Mutat. 14:
- [90818] 320-325, 1999.
- [90819] 16061.
- [90820] Cui, X.; De Vivo, I.; Slany, R.; Miyamoto, A.; Firestein, R.;
- [90821] Cleary, M. L.: Association of SET domain and myotubularin-related
- [90822] proteins modulates growth control. Nature Genet. 18: 331-337, 1998.
- [90823] 16062.
- [90824] Dickeson, S. K.; Helmkamp, G. M., Jr.; Yarbrough, L. R.: Sequence
- [90825] of a human cDNA encoding phosphatidylinositol transfer protein and
- [90826] occurrence of a related sequence in widely divergent eukaryotes. Gene 142:

- [90827] 301–305, 1994.
- [90828] 16063.
- [90829] Dickeson, S. K.; Lim, C. N.; Schuyler, G. T.; Dalton, T. P.;
Helmkamp,
- [90830] G. M., Jr.; Yarbrough, L. R.: Isolation and sequence of
cDNA clones
- [90831] encoding rat phosphatidylinositol transfer protein. J. Biol.
Chem. 264:
- [90832] 16557–16564, 1989.
- [90833] 16064.
- [90834] Hay, J. C.; Martin, T. F. J.: Phosphatidylinositol transfer
protein
- [90835] required for ATP–dependent priming of $\text{Ca}(2+)$ –activated
secretion. Nature 366:
- [90836] 572–575, 1993.
- [90837] 16065.
- [90838] Vihtelic, T. S.; Goebel, M.; Milligan, S.; O'Tousa, J. E.; Hyde,
- [90839] D. R.: Localization of Drosophila retinal degeneration B, a
membrane–associated
- [90840] phosphatidylinositol transfer protein. J. Cell Biol. 122:
1013–1022,

- [90841] 1993.
- [90842] 16066.
- [90843] Wirtz, K. W. A.: Phospholipid transfer proteins. *Annu. Rev. Biochem.* 60:
- [90844] 73–99, 1991.
- [90845] 16067.
- [90846] Fitzgibbon, J.; Pilz, A.; Gayther, S.; Appukuttan, B.; Dulai, K.
- [90847] S.; Delhanty, J. D. A.; Helmkamp, G. M., Jr.; Yarbrough, L. R.; Hunt, D. M.: Localization of the gene encoding human phosphatidylinositol
- [90848] transfer protein (PITPN) to 17p13.3: a gene showing homology to the
- [90849] *Drosophila* retinal degeneration B gene (rdgB). *Cytogenet. Cell Genet.* 67:
- [90850] 205–207, 1994.
- [90851] 16068.
- [90852] Hamilton, B. A.; Smith, D. J.; Mueller, K. L.; Kerrebrock, A. W.;
- [90853] Bronson, R. T.; van Berkel, V.; Daly, M. J.; Kruglyak, L.;

Reeve,

[90855] M. P.; Nemhauser, J. L.; Hawkins, T. L.; Rubin, E. M.; Lander, E.

[90856] S.: The vibrator mutation causes neurodegeneration via reduced expression

[90857] of P1TP-alpha: positional complementation cloning and extragenic suppression. Neuron 18:

[90858] 711-722, 1997.

[90859] 16069.

[90860] Ishida, N.; Miura, N.; Yoshioka, S.; Kawakita, M.: Molecular cloning

[90861] and characterization of a novel isoform of the human UDP-galactose

[90862] transporter, and of related complementary DNAs belonging to the nucleotide-sugar

[90863] transporter gene family. J. Biochem. 120: 1074-1078, 1996.

[90864] 16070.

[90865] Sugie, H.; Sugie, Y.; Nishida, M.; Ito, M.; Tsurui, S.; Suzuki,

[90866] M.; Miyamoto, R.; Igarashi, Y.: Recurrent myoglobinuria in a child

[90867] with mental retardation: phosphoglycerate kinase defi-

ciency. J.

[90868] Child Neurol. 4: 95–99, 1989.

[90869] 16071.

[90870] Sugie, H.; Sugie, Y.; Tsurui, S.; Ito, M.: Phosphoglycerate kinase

[90871] deficiency. (Letter) Neurology 44: 1364–1365, 1994.

[90872] 16072.

[90873] Tani, K.; Takizawa, T.; Yoshida, A.: Normal mRNA content in a

[90874] phosphoglycerate kinase variant with severe enzyme deficiency. Am.

[90875] J. Hum. Genet. 37: 931–937, 1985.

[90876] 16073.

[90877] Tonin, P.; Shanske, S.; Miranda, A. F.; Brownell, A. K.; Wyse,

[90878] J. P.; Tsujino, S.; Di Mauro, S.: Phosphoglycerate kinase deficiency:

[90879] biochemical and molecular genetic studies in a new myopathic variant. Neurology 43:

[90880] 387–391, 1993.

[90881] 16074.

- [90882] Valentin, C.; Birgens, H.; Craescu, C. T.; Brodum-Nielsen, K.;
- [90883] Cohen-Solal, M.: A phosphoglycerate kinase mutant (PGK Herlev; D285V)
- [90884] in a Danish patient with isolated chronic hemolytic anemia: mechanism
- [90885] of mutation and structure-function relationships. Hum. Mutat. 12:
- [90886] 280-287, 1998.
- [90887] 16075.
- [90888] Valentine, W. N.; Hsieh, H.-S.; Paglia, D. E.; Anderson, H. M.;
- [90889] Baughan, M. A.; Jaffe, E. R.; Garson, O. M.: Hereditary hemolytic
- [90890] anemia associated with phosphoglycerate kinase deficiency in erythrocytes
- [90891] and leukocytes: a probable X-chromosome-linked syndrome. New Eng.
- [90892] J. Med. 280: 528-534, 1969.
- [90893] 16076.
- [90894] Yoshida, A.; Miwa, S.: Characterization of a phosphoglycerate

- [90895] kinase variant associated with hemolytic anemia. Am. J. Hum. Genet. 26: 378–384, 1974.
- [90896] 16077.
- [90897] Yoshida, A.; Twele, T. W.; Dave, V.; Beutler, E.: Molecular abnormality of a phosphoglycerate kinase variant (PGK–Alabama). Blood Cells Mol. Dis. 21: 179–181, 1995.
- [90900] 16078.
- [90901] Yoshida, A.; Watanabe, S.; Chen, S.–H.; Giblett, E. R.; Malcolm, L. A.: Human phosphoglycerate kinase II: structure of a variant enzyme. J. Biol. Chem. 247: 446–449, 1972.
- [90902] 16079.
- [90903] Becker, M. A.; Kostel, P. J.; Meyer, L. J.; Seegmiller, J. E.: Human phosphoribosylpyrophosphate synthetase: increased enzyme specific activity in a family with gout and excessive purine synthesis. Proc.

- [90909] Nat. Acad. Sci. 70: 2749–2752, 1973.
- [90910] 16080.
- [90911] Becker, M. A.; Losman, M. J.; Rosenberg, A. L.; Mehlman, I.; Levinson, D. J.; Holmes, E. W.: Phosphoribosylpyrophosphate synthetase superactivity: a study of five patients with catalytic defects in the enzyme. Arthritis Rheum. 29: 880–888, 1986.
- [90912] 16081.
- [90913] Becker, M. A.; Losman, M. J.; Wilson, J.; Simmonds, H. A.: Superactivity of human phosphoribosyl pyrophosphate synthetase due to altered regulation by nucleotide inhibitors and inorganic phosphate. Biochim. Biophys. Acta 882: 168–176, 1986.
- [90914] 16082.
- [90915] Becker, M. A.; Meyer, L. J.; Seegmiller, J. E.: Gout with purine overproduction due to increased phosphoribosylphos-

phate synthetase

[90923] activity. Am. J. Med. 55: 232-242, 1973.

[90924] 16083.

[90925] Becker, M. A.; Meyer, L. J.; Wood, A. W.; Seegmiller, J. E.:
Purine

[90926] overproduction in man associated with increased phosphoribosylpyrophosphate

[90927] synthetase activity. Science 179: 1123-1126, 1973.

[90928] 16084.

[90929] Becker, M. A.; Puig, J. G.; Mateos, F. A.; Jimenez, M. L.;
Kim,

[90930] M.; Simmonds, H. A.: Inherited superactivity of phosphoribosylpyrophosphate

[90931] synthetase: association of uric acid overproduction and
sensorineural

[90932] deafness. Am. J. Med. 85: 383-390, 1988.

[90933] 16085.

[90934] Becker, M. A.; Raivio, K. O.; Bakay, B.; Adams, W. B.; Nyhan, W.

[90935] L.: Variant human phosphoribosylpyrophosphate synthetase altered

[90936] in regulatory and catalytic functions. J. Clin. Invest. 65:
109–120,

[90937] 1980.

[90938] 16086.

[90939] Becker, M. A.; Smith, P. R.; Taylor, W.; Mustafi, R.; Switzer,

[90940] R. L.: The genetic and functional basis of purine nucleotide feedback-resistant

[90941] phosphoribosylpyrophosphate synthetase superactivity. J. Clin. Invest. 96:

[90942] 2133–2141, 1995.

[90943] 16087.

[90944] Becker, M. A.; Taylor, W.; Smith, P. R.; Ahmed, M.: Overexpression

[90945] of the normal phosphoribosylpyrophosphate synthetase 1 isoform underlies

[90946] catalytic superactivity of human phosphoribosylpyrophosphate synthetase. J.

[90947] Biol. Chem. 271: 19894–19899, 1996.

[90948] 16088.

[90949] Becker, M. A.; Yen, R. C. K.; Goss, S. J.; Seegmiller, J. E.;

- [90950] Itkin, P.; Lazar, C.; Adams, W. B.: Localization of the structural
- [90951] gene for human phosphoribosylpyrophosphate synthetase on the X-chromosome.
- [90952] (Abstract) Clin. Res. 26: 500A, 1978.
- [90953] 16089.
- [90954] Becker, M. A.; Yen, R. C. K.; Itkin, P.; Goss, S. J.; Seeg-
miller,
- [90955] J. E.; Bakay, B.: Regional localization of the gene for hu-
man phosphoribosylpyrophosphate
- [90956] synthetase on the X-chromosome. Science 203:
1016-1019, 1979.
- [90957] 16090.
- [90958] Christen, H.-J.; Hanefeld, F.; Duley, J. A.; Simmonds, H. A.:
- [90959] Distinct neurological syndrome in two brothers with hype-
ruricaemia.
- [90960] (Letter) Lancet 340: 1167-1168, 1992.
- [90961] 16091.
- [90962] De Vries, A.; Sperling, O.: Familial gouty malignant uric
acid
- [90963] lithiasis due to mutant phosphoribosylpyrophosphatase

synthetase. Der

[90964] Urologe 12: 153–157, 1973.

[90965] 16092.

[90966] Iinuma, K.; Wada, Y.; Onuma, A.; Tanabu, M.: Electroencephalographic

[90967] study of an infant with phosphoribosylpyrophosphate synthetase deficiency. Tohoku

[90968] J. Exp. Med. 116: 53–55, 1975.

[90969] 16093.

[90970] Lebo, R. V.; Martin, D. W., Jr.: Electrophoretic heterogeneity

[90971] of 5–phosphoribosyl–1–pyrophosphate synthetase within and among humans. Biochem.

[90972] Genet. 16: 905–916, 1978.

[90973] 16094.

[90974] Nyhan, W. L.; James, J. A.; Teberg, A. J.; Sweetman, L.; Nelson,

[90975] L. G.: A new disorder of purine metabolism with behavioral manifestations. J.

[90976] Pediat. 74: 20–27, 1969.

[90977] 16095.

- [90978] Riccardi, V. M.: Personal Communication. Denver, Colo. 1974.
- [90979] 16096.
- [90980] Roessler, B. J.; Bell, G.; Heidler, S.; Seino, S.; Becker, M.;
- [90981] Palella, T. D.: Cloning of two distinct copies of human phosphoribosylpyrophosphate
- [90982] synthetase cDNA. Nucleic Acids Res. 18: 193 only, 1990.
- [90983] 16097.
- [90984] Roessler, B. J.; Nosal, J. M.; Smith, P. R.; Heidler, S. A.; Palella,
- [90985] T. D.; Switzer, R. L.; Becker, M. A.: Human X-linked phosphoribosylpyrophosphate
- [90986] synthetase superactivity is associated with distinct point mutations
- [90987] in the PRPS1 gene. J. Biol. Chem. 268: 26476-26481, 1993.
- [90988] 16098.
- [90989] Lund, A.; Udd, B.; Juvonen, V.; Andersen, P. M.; Ced-erquist, K.;
- [90990] Davis, M.; Gellera, C.; Kolmel, C.; Ronnevi, L.-O.; Sperfeld, A.-D.;

- [90991] Sorensen, S.-A.; Tranebjaerg, L.; Van Maldergem, L.; Watanabe, M.;
- [90992] Weber, M.; Yeung, L.; Savontaus, M.-L.: Multiple founder effects
- [90993] in spinal and bulbar muscular atrophy (SBMA, Kennedy disease) around
- [90994] the world. *Europ. J. Hum. Genet.* 9: 431–436, 2001.
- [90995] 16099.
- [90996] Edelhoff, S.; Ayer, D. E.; Zervos, A. S.; Steingrimsson, E.; Jenkins,
- [90997] N. A.; Copeland, N. G.; Eisenman, R. N.; Brent, R.; Distech, C. M.
- [90998] : Mapping of two genes encoding members of a distinct subfamily of
- [90999] MAX interacting proteins: MAD to human chromosome 2 and mouse chromosome
- [91000] 6, and MXI1 to chromosome 10 and mouse chromosome 19. *Oncogene* 9:
- [91001] 665–668, 1994.
- [91002] 16100.
- [91003] Ferguson, G. D.; Anagnostaras, S. G.; Silva, A. J.; Herschman,

- [91004] H. R.: Deficits in memory and motor performance in synaptotagmin
- [91005] IV mutant mice. *Proc. Nat. Acad. Sci.* 97: 5598–5603, 2000.
- [91006] 16101.
- [91007] Ferguson, G. D.; Chen, X.-N.; Korenberg, J. R.; Herschman, H. R.
- [91008] : The human synaptotagmin IV gene defines an evolutionary break point
- [91009] between syntenic mouse and human chromosome regions but retains ligand
- [91010] inducibility and tissue specificity. *J. Biol. Chem.* 275: 36920–36926,
- [91011] 2000.
- [91012] 16102.
- [91013] Holterhus, P.-M.; Wiebel, J.; Sinnecker, G. H. G.; Bruggenwirth,
- [91014] H. T.; Sippell, W. G.; Brinkmann, A. O.; Kruse, K.; Hiort, O.: Clinical
- [91015] and molecular spectrum of somatic mosaicism in androgen insensitivity
- [91016] syndrome. *Pediat. Res.* 46: 684–690, 1999.

- [91017] 16103.
- [91018] Foresta, C.; Bertella, A.; Moro, E.; Roverato, A.; Merico, M.;
- [91019] Ferlin, A.: Sertoli cell function in infertile patients with and
- [91020] without microdeletions of the azoospermia factors on the
Y chromosome
- [91021] long arm. J. Clin. Endocr. Metab. 86: 2414–2419, 2001.
- [91022] 16104.
- [91023] Kobayashi, K.; Mizuno, K.; Hida, A.; Komaki, R.; Tomita,
K.; Matsushita,
- [91024] I.; Namiki, M.; Iwamoto, T.; Tamura, S.; Minowada, S.;
- Nakahori, Y.;
- [91025] Nakagome, Y.: PCR analysis of the Y chromosome long
arm in azoospermic
- [91026] patients: evidence for a second locus required for sper-
matogenesis. Hum.
- [91027] Molec. Genet. 3: 1965–1967, 1994.
- [91028] 16105.
- [91029] Agulnik, A. I.; Zharkikh, A.; Boettger–Tong, H.; Bourgeron,
T.;
- [91030] McElreavey, K.; Bishop, C. E.: Evolution of the DAZ gene
family suggests

- [91031] that Y-linked DAZ plays little, or a limited, role in spermatogenesis
- [91032] but underlines a recent African origin for human populations. Hum.
- [91033] Molec. Genet. 7: 1371–1377, 1998.
- [91034] 16106.
- [91035] Anonymous: A missing piece on the Y. (Editorial) Nature Genet. 10:
- [91036] 367–368, 1995.
- [91037] 16107.
- [91038] Cooke, H. J.; Lee, M.; Kerr, S.; Ruggiu, M.: A murine homologue
- [91039] of the human DAZ gene is autosomal and expressed only in male and
- [91040] female gonads. Hum. Molec. Genet. 5: 513–516, 1996.
- [91041] 16108.
- [91042] Foresta, C.; Moro, E.; Garolla, A.; Onisto, M.; Ferlin, A.: Y
- [91043] chromosome microdeletions in cryptorchidism and idiopathic infertility. J.
- [91044] Clin. Endocr. Metab. 84: 3660–3665, 1999.

- [91045] 16109.
- [91046] Makova, K. D.; Li, W.-H.: Strong male-driven evolution of DNA
- [91047] sequences in humans and apes. *Nature* 416: 624–626, 2002.
- [91048] 16110.
- [91049] Menke, D. B.; Mutter, G. L.; Page, D. C.: Expression of DAZ, an
- [91050] azoospermia factor candidate, in human spermatogonia. (Letter) *Am.*
- [91051] *J. Hum. Genet.* 60: 237–241, 1997.
- [91052] 16111.
- [91053] Moro, E.; Ferlin, A.; Yen, P. H.; Franchi, P. G.; Palka, G.; Foresta,
- [91054] C.: Male infertility caused by a de novo partial deletion of the
- [91055] DAZ cluster on the Y chromosome. *J. Clin. Endocr. Metab.* 85: 4069–4073,
- [91056] 2000.
- [91057] 16112.
- [91058] Reijo, R.; Lee, T.-Y.; Salo, P.; Alagappan, R.; Brown, L. G.;

- [91059] Rosenberg, M.; Rozen, S.; Jaffe, T.; Straus, D.; Hovatta, O.;
de la
- [91060] Chapelle, A.; Silber, S.; Page, D. C.: Diverse spermatogenic
defects
- [91061] in humans caused by Y chromosome deletions encom-
passing a novel RNA-binding
- [91062] protein gene. *Nature Genet.* 10: 383–393, 1995.
- [91063] 16113.
- [91064] Reijo, R.; Seligman, J.; Dinulos, M. B.; Jaffe, T.; Brown, L.
- [91065] G.; Disteche, C. M.; Page, D. C.: Mouse autosomal ho-
molog of DAZ,
- [91066] a candidate male sterility gene in humans, is expressed in
male germ
- [91067] cells before and after puberty. *Genomics* 35: 346–352,
1996.
- [91068] 16114.
- [91069] Saxena, R.; Brown, L. G.; Hawkins, T.; Alagappan, R. K.;
Skaletsky,
- [91070] H.; Reeve, M. P.; Reijo, R.; Rozen, S.; Dinulos, M. B.; Dis-
teche,
- [91071] C. M.; Page, D. C.: The DAZ gene cluster on the human Y
chromosome

- [91072] arose from an autosomal gene that was transposed, repeatedly amplified
- [91073] and pruned. *Nature Genet.* 14: 292–299, 1996.
- [91074] 16115.
- [91075] Najmabadi, H.; Huang, V.; Yen, P.; Subbarao, M. N.; Bhasin, D;
- [91076] Banaag, L.; Naseeruddin, S.; de Kretser, D. M.; Baker, H. W. G.; McLachlan,
- [91077] R. I.; Loveland, K. A.; Bhasin, S.: Substantial prevalence of microdeletions
- [91078] of the Y-chromosome in infertile men with idiopathic azoospermia and
- [91079] oligozoospermia detected using a sequence-tagged site-based mapping
- [91080] strategy. *J. Clin. Endocr. Metab.* 81: 1347–1352, 1996.
- [91081] 16116.
- [91082] Wang, G.; Sawai, N.; Kotliarova, S.; Kanazawa, I.; Nukina, N.:
- [91083] Ataxin-3, the MJD1 gene product, interacts with the two human homologs
- [91084] of yeast DNA repair protein RAD23, HHR23A and HHR23B. *Hum. Molec.*

- [91085] Genet. 9: 1795–1803, 2000.
- [91086] 16117.
- [91087] Ng, J. M. Y.; Vrieling, H.; Sugasawa, K.; Ooms, M. P.;
Grootegoed,
- [91088] J. A.; Vreeburg, J. T. M.; Visser, P.; Beems, R. B.; Gorgels,
T. G.
- [91089] M. F.; Hanaoka, F.; Hoeijmakers, J. H. J.; van der Horst, G.
T. J.
- [91090] : Developmental defects and male sterility in mice lacking
the ubiquitin-like
- [91091] DNA repair gene mHR23B. Molec. Cell. Biol. 22:
1233–1245, 2002.
- [91092] 16118.
- [91093] Volker, M.; Mone, M. J.; Karmakar, P.; van Hoffen, A.;
Schul, W.;
- [91094] Vermeulen, W.; Hoeijmakers, J. H. J.; van Driel, R.; van
Zeeland,
- [91095] A. A.; Mullenders, L. H. F.: Sequential assembly of the nu-
cleotide
- [91096] excision repair factors in vivo. Molec. Cell 8: 213–224,
2001.
- [91097] 16119.

[91098] Jernstrom, H.; Lerman, C.; Ghadirian, P.; Lynch, H. T.; Weber,

[91099] B.; Garber, J.; Daly, M.; Olopade, O. I.; Foulkes, W. D.; Warner,

[91100] E.; Brunet, J.-S.; Narod, S. A.: Pregnancy and risk of early breast

[91101] cancer in carriers of BRCA1 and BRCA2. Lancet 354: 1846–1850, 1999.

[91102] 16120.

[91103] Khittoo, G.; Manning, A.; Mustun, H.; Appadoo, J.; Venkatasamy,

[91104] S.; Fagoonee, I.; Ghadirian, P.; Tonin, P. N.: Mutation analysis

[91105] of a Mauritian hereditary breast cancer family reveals the BRCA2 6503delTT

[91106] mutation previously found to recur in different ethnic populations. Hum.

[91107] Hered. 52: 55–58, 2001.

[91108] 16121.

[91109] Kinzler, K. W.; Vogelstein, B.: Gatekeepers and caretakers. Nature 386:

[91110] 761–763, 1997.

- [91111] 16122.
- [91112] Carrie, A.; Piccolo, F.; Leturcq, F.; de Toma, C.; Azibi, K.; Beldjord, C.; Vallat, J.-M.; Merlini, L.; Voit, T.; Sewry, C.; Urtizberea, J.
- [91114] A.; Romero, N.; Tome, F. M. S.; Fardeau, M.; Sunada, Y.; Campbell, K. P.; Kaplan, J.-C.; Jeanpierre, M.: Mutational diversity and hot spots in the alpha-sarcoglycan gene in autosomal recessive muscular dystrophy (LGMD2D). J. Med. Genet. 34: 470-475, 1997.
- [91118] 16123.
- [91119] Coral-Vazquez, R.; Cohn, R. D.; Moore, S. A.; Hill, J. A.; Weiss, R. M.; Davisson, R. L.; Straub, V.; Barresi, R.; Bansal, D.; Hrstka, R. F.; Williamson, R.; Campbell, K. P.: Disruption of the sarcoglycan-sarcospan complex in vascular smooth muscle: a novel mechanism for cardiomyopathy and muscular dystrophy. Cell 98: 465-474, 1999.

- [91124] 16124.
- [91125] Duggan, D. J.; Gorospe, J. R.; Fanin, M.; Hoffman, E. P.; Angelini,
- [91126] C.: Mutations in the sarcoglycan genes in patients with myopathy. New
- [91127] Eng. J. Med. 336: 618–624, 1997.
- [91128] 16125.
- [91129] Fadic, R.; Sunada, Y.; Waclawik, A. J.; Buck, S.; Lewandoski, P.
- [91130] J.; Campbell, K. P.; Lotz, B. P.: Deficiency of a dystrophin-associated
- [91131] glycoprotein (adhalin) in a patient with muscular dystrophy and cardiomyopathy. New
- [91132] Eng. J. Med. 334: 362–366, 1996.
- [91133] 16126.
- [91134] Kawai, H.; Akaike, M.; Endo, T.; Adachi, K.; Inui, T.; Mitsui,
- [91135] T.; Kashiwagi, S.; Fujiwara, T.; Okuno, S.; Shin, S.; Miyoshi, K.;
- [91136] Campbell, K. P.; Yamada, H.; Shimizu, T.; Matsumura, K.; Saito, S.
- [91137] : Adhalin gene mutations in patients with autosomal re-

cessive childhood

[91138] onset muscular dystrophy with adhalin deficiency. J. Clin. Invest. 96:

[91139] 1202–1207, 1995.

[91140] 16127.

[91141] Ljunggren, A.; Duggan, D.; McNally, E.; Boylan, K. B.; Gama, C.

[91142] H.; Kunkel, L. M.; Hoffman, E. P.: Primary adhalin deficiency as

[91143] a cause of muscular dystrophy in patients with normal dystrophin. Ann.

[91144] Neurol. 38: 367–372, 1995.

[91145] 16128.

[91146] McNally, E. M.; Yoshida, M.; Mizuno, Y.; Ozawa, E.; Kunkel, L.

[91147] M.: Human adhalin is alternatively spliced and the gene is located

[91148] on chromosome 17q21. Proc. Nat. Acad. Sci. 91: 9690–9694, 1994.

[91149] 16129.

[91150] Passos–Bueno, M. R.; Oliveira, J. R.; Bakker, E.; Anderson,

R.

[91151] D.; Marie, S. K.; Vainzof, M.; Roberds, S. L.; Campbell, K. P.; Zatz,

[91152] M.: Genetic heterogeneity for Duchenne-like muscular dystrophy (DLMD)

[91153] based on linkage and 50 DAG analysis. Hum. Molec. Genet. 2: 1945–1947,

[91154] 1993.

[91155] 16130.

[91156] Passos Bueno, M. R.; Moreira, E. S.; Vainzof, M.; Chamberlain,

[91157] J.; Marie, S. K.; Pereira, L.; Akiyama, J.; Roberds, S. L.; Campbell,

[91158] K. P.; Zatz, M.: A common missense mutation in the adhalin gene in

[91159] three unrelated Brazilian families with a relatively mild form of

[91160] autosomal recessive limb-girdle muscular dystrophy. Hum. Molec. Genet. 4:

[91161] 1163–1167, 1995.

[91162] 16131.

[91163] Piccolo, F.; Roberds, S. L.; Jeanpierre, M.; Leturcq, F.; Az-

ibi,

[91164] K.; Beldjord, C.; Carrie, A.; Recan, D.; Chaouch, M.; Reghis, A.;

[91165] El Kerch, F.; Sefiani, A.; Voit, T.; Merlini, L.; Collin, H.; Ey-
mard,

[91166] B.; Beckmann, J. S.; Romero, N. B.; Tome, F. M. S.; Fardeau, M.; Campbell,

[91167] K. P.; Kaplan, J.-C.: Primary adhalinopathy: a common
cause of autosomal

[91168] recessive muscular dystrophy of variable severity. Nature
Genet. 10:

[91169] 243–245, 1995.

[91170] 16132.

[91171] Roberds, S. L.; Anderson, R. D.; Ibraghimov-Beskrovnaya, O.; Campbell,

[91172] K. P.: Primary structure and muscle-specific expression on
the 50-kDa

[91173] dystrophin-associated glycoprotein (adhalin). J. Biol.
Chem. 268:

[91174] 23739–23742, 1993.

[91175] 16133.

[91176] Roberds, S. L.; Leturcq, F.; Allamand, V.; Piccolo, F.; Jean-

pierre,

[91177] M.; Anderson, R. D.; Lim, L. E.; Lee, J. C.; Tome, F. M. S.;
Romero,

[91178] N. B.; Fardeau, M.; Beckmann, J. S.; Kaplan, J.-C.; Camp-
bell, K. P.

[91179] : Missense mutations in the adhalin gene linked to auto-
somal recessive

[91180] muscular dystrophy. Cell 78: 625-633, 1994.

[91181] 16134.

[91182] Romero, N. B.; Tome, F. M. S.; Leturcq, F.; El Kerch, F.; Az-
ibi,

[91183] K.; Bachner, L.; Anderson, R. D.; Roberds, S. L.; Campbell,
K. P.;

[91184] Fardeau, M.; Kaplan, J.-C.: Genetic heterogeneity of severe
childhood

[91185] autosomal recessive muscular dystrophy with adhalin (50
kDa dystrophin-associated

[91186] glycoprotein) deficiency. Comp. Rend. Acad. Sci. (Paris)
317: 70-76,

[91187] 1994.

[91188] 16135.

[91189] Schara, U.; Gencik, M.; Mortier, J.; Langen, M.; Gencikova,

A.;

[91190] Epplen, J. T.; Mortier, W.: Alpha-sarcoglycanopathy previously misdiagnosed

[91191] as Duchenne muscular dystrophy: implications for current diagnostics

[91192] and patient care. *Europ. J. Pediat.* 160: 452–453, 2001.

[91193] 16136.

[91194] Perelman, B.; Dafni, N.; Naiman, T.; Eli, D.; Yaakov, M.; Yang

[91195] Feng, T. L.; Sinha, S.; Weber, G.; Khodaei, S.; Sancar, A.; Dotan,

[91196] I.; Canaani, D.: Molecular cloning of a novel human gene encoding

[91197] a 63-kDa protein and its sublocalization within the 11q13 locus. *Genomics* 41:

[91198] 397–405, 1997.

[91199] 16137.

[91200] Teitz, T.; Penner, M.; Eli, D.; Stark, M.; Bakhanashvili, M.; Naiman,

[91201] T.; Canaani, D.: Isolation by polymerase chain reaction of a cDNA

[91202] whose product partially complements the ultraviolet sen–

sitivity of

[91203] xeroderma pigmentosum group C cells. Gene 87:
295–298, 1990.

[91204] 16138.

[91205] Kramps, T.; Peter, O.; Brunner, E.; Nellen, D.; Froesch, B.;
Chatterjee,

[91206] S.; Murone, M.; Zullig, S.; Basler, K.: Wnt/Wingless signal-
ing requires

[91207] BCL9/legless-mediated recruitment of pygopus to the nu-
clear beta-catenin-TCF

[91208] complex. Cell 109: 47–60, 2002.

[91209] 16139.

[91210] Willis, T. G.; Zalcborg, I. R.; Coignet, L. J. A.; Wlodarska,
M.;

[91211] Stul, D. M.; Jadayel, D. M.; Bastard, C.; Treleaven, J. G.;
Catovsky,

[91212] D.; Silva, M. L. M.; Dyer, M. J. S.: Molecular cloning of
translocation

[91213] t(1;14)(q21;q32) defines a novel gene (BCL9) at chromo-
some 1q21. Blood 91:

[91214] 1873–1881, 1998.

[91215] 16140.

- [91216] Davidson, R. G.; Nitowsky, H. M.; Childs, B.: Demonstration of
- [91217] two populations of cells in the human female heterozygous for glucose-6-phosphate
- [91218] dehydrogenase variants. *Proc. Nat. Acad. Sci.* 50: 481-485, 1963.
- [91219] 16141.
- [91220] De Flora, A.; Morelli, A.; Benatti, U.; Giuntini, P.; Ferraris, A. M.; Galiano, S.; Ravazzolo, R.; Gaetani, G. F.: G6PD
- [91221] Napoli and
- [91222] Ferrara II: two new glucose-6-phosphate dehydrogenase variants having
- [91223] similar characteristics but different intracellular lability and specific
- [91224] activity. *Brit. J. Haemat.* 48: 417-423, 1981.
- [91225] 16142.
- [91226] Dern, R. J.; McCurdy, P. R.; Yoshida, A.: A new structural variant
- [91227] of glucose-6-phosphate dehydrogenase with a high production rate (G6PD
- [91228] Hektoen). *J. Lab. Clin. Med.* 73: 283-290, 1969.

- [91229] 16143.
- [91230] De Vita, G.; Alcalay, M.; Sampietro, M.; Cappelini, M. D.;
Fiorelli,
- [91231] G.; Toniolo, D.: Two point mutations are responsible for
G6PD polymorphism
- [91232] in Sardinia. Am. J. Hum. Genet. 44: 233–240, 1989.
- [91233] 16144.
- [91234] Du, C.; Xu, Y.; Hua, X.; Wu, Q.; Liu, L.; Wu, M.: Studies on
- [91235] erythrocyte glucose–6–phosphate dehydrogenase variants
in Chinese.
- [91236] III: Gd(–) Miaozi–Baisha. Acta Genet. Sinica 11(2):
153–158, 1984.
- [91237] 16145.
- [91238] Du, C.–S.; Hua, X.–Y.; Wu, Q.–L.; Li, C.–Q.; Zheng, J.–F.; Li,
- [91239] H.–L.: Studies on erythrocyte glucose–6–phosphate dehy-
drogenase variants
- [91240] in Chinese. IV. Gd(–) Gaohe associated with paroxysmal
nocturnal hemoglobinuria. Chinese
- [91241] J. Pathophysiol. 1: 12–15, 1985.
- [91242] 16146.

- [91243] Du, C. S.; Xu, Y. K.; Hua, X. Y.; Wu, Q. L.; Liu, L. B.: Glucose-6-phosphate
- [91244] dehydrogenase variants and their frequency in Guangdong, China. Hum.
- [91245] Genet. 80: 385-388, 1988.
- [91246] 16147.
- [91247] Eber, S. W.; Gahr, M.; Schroter, W.: Glucose-6-phosphate dehydrogenase
- [91248] (G6PD) Iserlohn and G6PD Regensburg: two new severe enzyme defects
- [91249] in German families. Blut 51: 109-115, 1985.
- [91250] 16148.
- [91251] Echard, G.; Gillois, M.: G6PD--alpha-GAL-PGK--HPRT syntenry in
- [91252] the rabbit, *Oryctolagus cuniculus*. (Abstract) Cytogenet. Cell Genet. 25:
- [91253] 148-149, 1979.
- [91254] 16149.
- [91255] Elizondo, J.; Saenz, G. F.; Paez, C. A.; Ramon, M.; Garcia, M.;
- [91256] Gutierrez, A.; Estrada, M.: G6PD Puerto Limon: a new defi-

cient variant

[91257] of glucose-6-phosphate dehydrogenase associated with congenital nonspherocytic

[91258] hemolytic anemia. Hum. Genet. 62: 110-112, 1982.

[91259] 16150.

[91260] Engstrom, P. F.; Beutler, E.: G-6-PD Tripler: a unique variant

[91261] associated with chronic hemolytic disease. Blood 36: 10-13, 1970.

[91262] 16151.

[91263] Epstein, C. J.: Mammalian oocytes: X-chromosome activity. Science 163:

[91264] 1078-1079, 1969.

[91265] 16152.

[91266] Ermakov, N. V.; Chernyak, N. B.; Tokarev, Y. N.: Properties of

[91267] new variant of glucose-6-phosphate dehydrogenase (Regar variant):

[91268] glucose metabolism in erythrocytes containing abnormal enzyme. Biokhimiia 48:

[91269] 577-583, 1983.

- [91270] 16153.
- [91271] Estrada, M.; Garcia, M.; Gutierrez, A.; Quintero, I.; Gonzalez,
lez,
- [91272] R.: G6PD Varadero. Vox Sang. 43: 102–104, 1982.
- [91273] 16154.
- [91274] Fairbanks, V. F.; Nepo, A. G.; Beutler, E.; Dickson, E. R.;
Honig,
- [91275] G.: Glucose–6–phosphate dehydrogenase variants: reex-
amination of
- [91276] G6PD Chicago and Cornell and a new variant (G6PD Pea
Ridge) resembling
- [91277] G6PD Chicago. Blood 55: 216–220, 1980.
- [91278] 16155.
- [91279] Faust, C. J.; Levinson, B.; Gitschier, J.; Herman, G. E.: Ex-
tension
- [91280] of the physical map in the region of the mouse X chromo-
some homologous
- [91281] to human Xq28 and identification of an exception to con-
served linkage. Genomics 13:
- [91282] 1289–1295, 1992.
- [91283] 16156.

- [91284] Feldman, R.; Gromisch, D. S.; Luhby, A. L.; Beutler, E.:
Congenital
- [91285] nonspherocytic hemolytic anemia due to glucose-
6-phosphate dehydrogenase
- [91286] East Harlem: a new deficient variant. J. Pediat. 90: 89-91,
1977.
- [91287] 16157.
- [91288] Fernandez, M.; Fairbanks, V. F.: Glucose-6-phosphate de-
hydrogenase
- [91289] deficiency in the Philippines: report of a new variant-
-G6PD Panay. Mayo
- [91290] Clin. Proc. 43: 645-660, 1968.
- [91291] 16158.
- [91292] Hostikka, S. L.; Eddy, R. L.; Byers, M. G.; Hoyhtya, M.;
Shows,
- [91293] T. B.; Tryggvason, K.: Identification of a distinct type IV
collagen
- [91294] alpha chain with restricted kidney distribution and assign-
ment of
- [91295] its gene to the locus of X chromosome-linked Alport syn-
drome. Proc.
- [91296] Nat. Acad. Sci. 87: 1606-1610, 1990.

[91297] 16159.

[91298] Bland, J. H. (General Chairman): Proceedings of seminars
on the

[91299] Lesch-Nyhan syndrome. Fed. Proc. 27: 1017-1112, 1968.

[91300] 16160.

[91301] Ferraris, A. M.; Broccia, G.; Meloni, T.; Forteleoni, G.; Gaetan-
tani,

[91302] G. F.: Glucose-6-phosphate dehydrogenase deficiency
and incidence

[91303] of hematologic malignancy. Am. J. Hum. Genet. 42:
516-520, 1988.

[91304]

[91305] 100. Filosa, E.: Personal Communication. La Jolla, Calif.
1989.

[91306]

[91307] 101. Filosa, S.; Calabro, V.; Vallone, D.; Poggi, V.; Mason,
P.; Pagnini,

[91308] D.; Alfinito, F.; Rotoli, B.; Martini, G.; Luzzatto, L.; Battis-
tuzzi,

[91309] G.: Molecular basis of chronic non-spherocytic haemolytic
anaemia:

[91310] a new G6PD variant (393arg-to-his) with abnormal K(m)
G6P and marked

[91311] in vivo instability. *Brit. J. Haemat.* 80: 111–116, 1992.

[91312]

[91313] 102. Filosa, S.; Giacometti, N.; Wangwei, C.; De Mattia, D.;
Pagnini,

[91314] D.; Alfinito, F.; Schettini, F.; Luzzatto, L.; Martini, G.: So-
matic-cell

[91315] selection is a major determinant of the blood-cell pheno-
type in heterozygotes

[91316] for glucose-6-phosphate dehydrogenase mutations caus-
ing severe enzyme

[91317] deficiency. *Am. J. Hum. Genet.* 59: 887–895, 1996.

[91318]

[91319] 103. Fiorelli, G.; Manoussakis, C.; Sampietro, M.; Pittalis,
S.; Guglielmino,

[91320] C. R.; Cappellini, M. D.: Different polymorphic variants of
glucose-6-phosphate

[91321] dehydrogenase (G6PD) in Italy. *Ann. Hum. Genet.* 53:
229–236, 1989.

[91322]

[91323] 104. Fite, E.; Morell, F.; Zuazu, J.; Julia, A.; Morera, J.: Leucocyte

[91324] glucose-6-phosphate dehydrogenase deficiency and necrotizing pneumonia. Europ.

[91325] J. Resp. Dis. 64: 150-154, 1983.

[91326]

[91327] 105. Francke, U.; Bakay, B.; Connor, J. D.; Coldwell, J. G.; Nyhan,

[91328] W. L.: Linkage relationships of X-linked enzymes glucose-6-phosphate

[91329] dehydrogenase and hypoxanthine guanine phosphoribosyltransferase. Am.

[91330] J. Hum. Genet. 26: 512-522, 1974.

[91331]

[91332] 106. Friedman, M. J.; Trager, W.: The biochemistry of resistance

[91333] to malaria. Sci. Am. 244(3): 154-164, 1981.

[91334]

[91335] 107. Frigerio, R.; Sole, G.; Olla, N.; Lovicu, M.; Passiu, G.; Carcassi,

[91336] U.: Cagliari II: a new G-6-PD variant. Haematologica 72:

241–243,

[91337] 1987.

[91338]

[91339] 108. Fujii, H.; Miwa, S.; Takegawa, S.; Takahashi, K.; Hirono, A.;

[91340] Takizawa, T.; Morisaki, T.; Kanno, H.; Taguchi, T.; Okamura, J.:

[91341] Gd(–) Gifu and Gd(–) Fukuoka: two new variants of glucose-6-phosphate

[91342] dehydrogenase found in Japan. Hum. Genet. 66: 276–278, 1984.

[91343]

[91344] 109. Fujii, H.; Miwa, S.; Tani, K.; Takegawa, S.; Fujinami, N.; Takahashi,

[91345] K.; Nakayama, S.; Konno, M.; Sato, T.: Glucose 6-phosphate dehydrogenase

[91346] variants: a unique variant (G6PD Kobe) showed an extremely increased

[91347] affinity for galactose 6-phosphate and a new variant (G6PD Sapporo)

[91348] resembling G6PD Pea Ridge. Hum. Genet. 58: 405–407, 1981.

[91349]

[91350] 110. Gaetani, G. F.; Galiano, S.; Melani, C.; Miglino, M.;
Forni,

[91351] G. L.; Napoli, G.; Perrone, L.; Ferraris, A. M.: A new glu-
cose-6-phosphate

[91352] dehydrogenase variant with congenital nonspherocytic
hemolytic anemia

[91353] (G6PD Genova): biochemical characterization and mo-
saicism expression

[91354] in the heterozygote. Hum. Genet. 84: 337-340, 1990.

[91355]

[91356] 111. Gahr, M.; Bornhalm, D.; Schroeter, W.: Biochemische
Eigenschaften

[91357] einer neuen Variante des Glucose-
6-phosphadehydrogenase (G6PD) Mangels

[91358] mit Favismus: G6PD Bielefeld. Klin. Wschr. 55: 379-384,
1977.

[91359]

[91360] 112. Gahr, M.; Bornhalm, D.; Schroeter, W.: Haemolytic
anemia due

[91361] to glucose 6-phosphate dehydrogenase (G6PD) deficiency:
demonstration

[91362] of two new biochemical variants, G6PD Hamm and G6PD Tarsus. Brit.

[91363] J. Haemat. 33: 363–370, 1976.

[91364]

[91365] 113. Gahr, M.; Schroeter, W.: Glucose 6–phosphate dehydrogenase (G6PD)

[91366] Hamburg, a new variant with chronic nonspherocytic hemolytic anemia. Europ.

[91367] J. Clin. Invest. 4: 187–191, 1974.

[91368]

[91369] 114. Gahr, M.; Schroeter, W.; Sturzenegger, M.; Bornhalm, D.; Marti,

[91370] H. R.: Glucose 6–phosphate dehydrogenase (G6PD) deficiency in Switzerland. Helv.

[91371] Paediat. Acta 31: 159–166, 1976.

[91372]

[91373] 115. Geerdink, R. A.; Horst, R.; Staal, G. E.: An Iraqi Jewish family

[91374] with a new red cell glucose 6–phosphate dehydrogenase variant (Gd–Bagdad)

[91375] and kernicterus. Israel J. Med. Sci. 9: 1040–1043, 1973.

[91376]

[91377] 116. Gellin, J.; Benne, F.; Renard, C.; Vaiman, M.; Hors-Cayla, M.

[91378] C.; Gillois, M.: Pig gene mapping: synteny, attempt to assign the

[91379] histocompatibility complex (SLA). (Abstract) Cytogenet. Cell Genet. 25:

[91380] 159, 1979.

[91381]

[91382] 117. Gomez-Gallego, F.; Garrido-Pertierra, A.; Bautista, J. M.: Structural

[91383] defects underlying protein dysfunction in human glucose-6-phosphate

[91384] dehydrogenase A- deficiency. J. Biol. Chem. 275: 9256-9262, 2000.

[91385]

[91386] 118. Gonzalez, R.; Estrada, M.; Garcia, M.; Gutierrez, A.: G6PD Ciudad

[91387] de la Habana: a new slow variant with deficiency found in a Cuban

[91388] family. Hum. Genet. 55: 133-135, 1980.

[91389]

[91390] 119. Gonzalez, R.; Wade, M.; Estrada, M.; Svarch, E.;
Colombo, B.

[91391] : G6PD Pinar del Rio: a new variant discovered in a Cuban
family. Biochem.

[91392] Genet. 15: 909–913, 1977.

[91393]

[91394] 120. Goss, S. J.; Harris, H.: Gene transfer by means of cell
fusion.

[91395] I. Statistical mapping of the human X–chromosome by
analysis of radiation

[91396] induced gene segregation. J. Cell Sci. 25: 17–37, 1977.

[91397]

[91398] 121. Gourdin, D.; Vergnes, H.; Bouloux, C.; Ruffie, J.;
Gherardi,

[91399] M.: Polymorphism of erythrocyte G6PD in the baboon. Am.
J. Phys.

[91400] Anthropol. 37: 281–288, 1972.

[91401]

[91402] 122. Gray, G. R.; Stamatoyannopoulos, G.; Naiman, S. C.;
Kliman, M.

- [91403] R.; Klebanoff, S. J.; Austin, T.; Yoshida, A.; Robinson, G. C. G.
- [91404] : Neutrophil dysfunction, chronic granulomatous disease, and non-spherocytic
- [91405] haemolytic anaemia caused by complete deficiency of glucose-6-phosphate
- [91406] dehydrogenase. *Lancet* II: 530-534, 1973.
- [91407]
- [91408] 123. Grossman, A.; Ramanathan, K.; Justice, P.; Gordon, J.; Shahidi,
- [91409] N. T.; Hsia, D. Y. Y.: Congenital nonspherocytic hemolytic anemia
- [91410] associated with erythrocyte G-6-PD deficiency in a Negro family. *Pediatrics* 37:
- [91411] 624-629, 1966.
- [91412]
- [91413] 124. Gutierrez, A.; Garcia, M.; Estrada, M.; Quintero, I.; Gonzalez,
- [91414] R.: Glucose-6-phosphate dehydrogenase (G6PD) Guantano and G6PD
- [91415] Caujeri: two new glucose-6-phosphate dehydrogenase-deficient variants

[91416] found in Cuba. Biochem. Genet. 25: 231–238, 1987.

[91417]

[91418] 125. Hall, K.; Schreeder, M. T.; Prchal, J. T.: G6PD
Huntsville:

[91419] a new glucose–6–phosphate dehydrogenase associated
with chronic hemolytic

[91420] anemia. Hum. Genet. 79: 90–91, 1988.

[91421]

[91422] 126. Harkonen, M.; Vuopio, P.: Red cell glucose–
6–phosphate dehydrogenase

[91423] deficiency in Finland. Ann. Clin. Res. 6: 187–197, 1974.

[91424]

[91425] 127. Harley, J. D.; Agar, N. S.; Yoshida, A.: Glucose
6–phosphate

[91426] dehydrogenase variant Gd(+) Alexandra associated with
neonatal jaundice

[91427] and Gd(–) Camperdown in a young man with lamellar
cataracts. J. Lab.

[91428] Clin. Med. 91: 295–300, 1978.

[91429]

- [91430] 128. Helge, H.; Borner, K.: Kongenitale nichtsphaerozytare haemolytische
- [91431] Anaemie, Katarakt und Glucose-6-phosphat-dehydrogenase-mangel. Dtsch.
- [91432] Med. Wschr. 91: 1584-1589, 1966.
- [91433]
- [91434] 129. Hirono, A.; Beutler, E.: Alternative splicing of human glucose-6-phosphate
- [91435] dehydrogenase messenger RNA in different tissues. J. Clin. Invest. 83:
- [91436] 343-346, 1989.
- [91437]
- [91438] 130. Hirono, A.; Beutler, E.: Molecular cloning and nucleotide sequence
- [91439] of cDNA for human glucose-6-phosphate dehydrogenase variant A(-). Proc.
- [91440] Nat. Acad. Sci. 85: 3951-3954, 1988.
- [91441]
- [91442] 131. Hirono, A.; Fujii, H.; Shima, M.; Miwa, S.: G6PD Nara: a new
- [91443] class 1 glucose-6-phosphate dehydrogenase variant with

an eight amino

[91444] acid deletion. Blood 82: 3250–3252, 1998.

[91445]

[91446] 132. Hirono, A.; Kawate, K.; Honda, A.; Fujii, H.; Miwa, S.:
A single

[91447] mutation 202G→A in the human glucose-6-phosphate de-
hydrogenase gene

[91448] (G6PD) can cause acute hemolysis by itself. (Letter) Blood
99: 1498

[91449] only, 2002.

[91450]

[91451] 133. Hirono, A.; Kuhl, W.; Gelbart, T.; Forman, L.; Fair-
banks, V.

[91452] F.; Beutler, E.: Identification of the binding domain for
NADP(+)

[91453] of human glucose-6-phosphate dehydrogenase by se-
quence analysis of

[91454] mutants. Proc. Nat. Acad. Sci. 86: 10015–10017, 1989.

[91455]

[91456] 134. Hitzeroth, H. W.; Bender, K.: Age-dependency of so-
matic selection

[91457] in South African Negro G-6-PD heterozygotes. Hum.
Genet. 58: 338-343,

[91458] 1981.

[91459]

[91460] 135. Honig, G. R.; Habacon, E.; Vida, L. N.; Matsumoto, F.;
Beutler,

[91461] E.: Three new variants of glucose-6-phosphate dehydro-
genase associated

[91462] with chronic nonspherocytic hemolytic anemia: G6PD Lin-
coln Park, G6PD

[91463] Arlington Heights, and G6PD West Town. Am. J. Hemat. 6:
353-360,

[91464] 1979.

[91465]

[91466] 136. Hook, E. B.; Stamatoyannopoulos, G.; Yoshida, A.;
Motulsky, A.

[91467] G.: Glucose-6-phosphate dehydrogenase Madrona: a slow
electrophoretic

[91468] glucose-6-phosphate dehydrogenase variant with kinetic
characteristics

[91469] similar to those of normal type. J. Lab. Clin. Med. 72:
404-409,

[91470] 1968.

[91471]

[91472] 137. Hors-Cayla, M. C.; Heuertz, S.; Van Cong, N.; Benne, F.: Cattle

[91473] gene mapping by somatic cell hybridization. (Abstract)
Cytogenet.

[91474] Cell Genet. 25: 165-166, 1979.

[91475]

[91476] 138. Howell, E. B.; Nelson, A. J.; Jones, O. W.: A new G-6-PD variant

[91477] associated with chronic non-spherocytic haemolytic
anaemia in a Negro

[91478] family. J. Med. Genet. 9: 160-164, 1972.

[91479]

[91480] 139. Hutz, M. H.; Yoshida, A.; Salzano, F. M.: Three rare
G-6-PD

[91481] variants from Porto Alegre, Brazil. Hum. Genet. 39:
191-197, 1977.

[91482]

[91483] 140. Ishwad, C. S.; Naik, S. N.: A new glucose-6-phosphate dehydrogenase

[91484] variant (G-6-PD Kalyan) found in a Koli family. Hum.

Genet. 66:

[91485] 171-175, 1984.

[91486]

[91487] 141. Johnson, G. J.; Kaplan, M. E.; Beutler, E.: G6PD Long
Prairie:

[91488] a new mutant exhibiting normal sensitivity to inhibition by
NADPH

[91489] and accompanied by nonspherocytic hemolytic anemia.
Blood 49: 247-251,

[91490] 1977.

[91491]

[91492] 142. Johnston, P. G.; VandeBerg, J. L.; Sharman, G. B.: In-
heritance

[91493] of erythrocyte glucose 6-phosphate dehydrogenase in the
red-necked

[91494] wallaby, *Macropus rufogriseus* (Desmarest) consistent
with paternal

[91495] X inactivation. Biochem. Genet. 13: 235-242, 1975.

[91496]

[91497] 143. Junien, C.; Kaplan, J.-C.; Meienhofer, M. C.; Maigret,

P.; Sender,

[91498] A.: G6PD Baudelocque: a new unstable variant characterized in cultured

[91499] fibroblasts. *Enzyme* 18: 48–59, 1974.

[91500]

[91501] 144. Kaeda, J. S.; Chhotray, G. P.; Ranjit, M. R.; Bautista, J. M.;

[91502] Reddy, P. H.; Stevens, D.; Naidu, J. M.; Britt, R. P.; Vulliamy, T.

[91503] J.; Luzzatto, L.; Mason, P. J.: A new glucose-6-phosphate dehydrogenase

[91504] variant, G6PD Orissa (44 ala-gly), is the major polymorphic variant

[91505] in tribal populations in India. *Am. J. Hum. Genet.* 57: 1335–1341,

[91506] 1995.

[91507]

[91508] 145. Kageoka, T.; Satoh, C.; Goriki, K.; Fujita, M.; Neriishi, S.;

[91509] Yamamura, K.; Kaneko, J.; Masunari, N.: Electrophoretic variants

[91510] of blood proteins in Japanese. IV. Prevalence and enzyme-

logic characteristics

[91511] of G6PD variants in Hiroshima and Nagasaki. Hum. Genet.
70: 101–108,

[91512] 1985.

[91513]

[91514] 146. Kahn, A.; Bernard, J.-F.; Cottreau, D.; Mazie, J.;
Boivin, P.

[91515] : Gd(–) Abrami, a deficient G6PD variant with hemizygous
expression

[91516] in blood cells of a woman with primary myelofibrosis. Hu-
mangenetik 30:

[91517] 41–46, 1975.

[91518]

[91519] 147. Kahn, A.; Boivin, P.; Hakim, J.; Lagneau, J.: Hetero-
geneite

[91520] des glucose-6-phosphate dehydrogenase erythrocytaire
deficitaires

[91521] dans la race noire: etude cinetique et description de deux
nouvelles

[91522] variants Gd(–) Dakar et Gd(–) Mali. Nouv. Rev. Franc.
Hemat. 11:

[91523] 741–758, 1971.

[91524]

[91525] 148. Kahn, A.; Boivin, P.; Lagneau, J.: Phenotypes de la glucose-6-phosphate

[91526] dehydrogenase erythrocytaire dans la race noire. Human-genetik 18:

[91527] 261-270, 1973.

[91528]

[91529] 149. Kahn, A.; Boulard, M.; Hakim, J.; Schaison, G.; Boivin, P.; Bernard,

[91530] J.: Anemie hemolytique congenitale non spherocytaire par deficit

[91531] en glucose 6-phosphate dehydrogenase erythrocytaire: description de

[91532] deux nouvelles variants: Gd(-) Saint Louis (Paris) et Gd(-) Hayem. Nouv.

[91533] Rev. Franc. Hemat. 14: 587-600, 1974.

[91534]

[91535] 150. Kahn, A.; Dao, C.; Cottreau, D.; Bilski-Pasquier, G.: 'Gd(-)

[91536] Hotel Dieu': a new G-6PD variant with chronic hemolysis in a Negro

[91537] patient from Senegal. Hum. Genet. 39: 353–358, 1977.

[91538]

[91539] 151. Kahn, A.; Exters, A.; Habedank, M.: Gd(–) Aachen, a new variant

[91540] of deficient glucose–6–phosphate dehydrogenase. Humangenetik 32:

[91541] 171–180, 1976.

[91542]

[91543] 152. Kahn, A.; Hakim, J.; Cottreau, D.; Boivin, P.: Gd (–) Matam.

[91544] An African glucose 6–phosphate dehydrogenase variant with enzyme deficiency.

[91545] Biochemical and immunological properties in various hemopoietic tissues. Clin.

[91546] Chim. Acta 59: 183–190, 1975.

[91547]

[91548] 153. Kahn, A.; North, M. L.; Cottreau, D.; Giron, G.; Lang, J. M.

[91549] : G6PD Vientiane: a new glucose–6–phosphate dehydrogenase variant

[91550] with increased stability. Hum. Genet. 43: 85–89, 1978.

[91551]

[91552] 154. Kahn, A.; North, M. L.; Messer, J.; Boivin, P.: G-6PD
'Ankara':

[91553] a new G-6PD variant with deficiency found in a Turkish
family. Humangenetik 27:

[91554] 247-250, 1975.

[91555]

[91556] 155. Kanno, H.; Huang, I.-Y.; Kan, Y. W.; Yoshida, A.: Two
structural

[91557] genes on different chromosomes are required for encod-
ing the major

[91558] subunit of human red cell glucose-6-phosphate dehydro-
genase. Cell 58:

[91559] 595-606, 1989.

[91560]

[91561] 156. Kanno, H.; Takano, T.; Fujii, H.; Tani, K.; Morisaki, T.;
Hirono,

[91562] A.; Kumakawa, T.; Ogura, H.; Takahashi, K.; Tsutsumi, H.;
Miwa, S.

[91563] : A new glucose-6-phosphate variant (G6PD Iwate) associ-
ated with congenital

- [91564] non-spherocytic hemolytic anemia. Acta Haemat. Jpn. 51: 715-719, 1987.
- [91565]
- [91566]
- [91567] 157. Kaplan, J. C.; Hanzlickova-Leroux, A.; Nicholas, A. M.; Rosa, R.; Weiler, C.; Lepercq, G.: A new glucose-6-phosphate dehydrogenase variant (G6PD Port-Royal). Enzyme 12: 25-32, 1971.
- [91570]
- [91571] 158. Kaplan, J. C.; Rosa, R.; Seringe, P.; Hoeffel, J. C.: Le polymorphisme genetique de la glucose-6-phosphate deshydrogenase erythrocytaire chez l'homme. II. Etude d'une nouvelle variete a activite diminuee: le type 'Kabyle.'. Enzym. Biol. Clin. 8: 332-340, 1967.
- [91575]
- [91576] 159. Kaplan, M.; Hammerman, C.; Vreman, H. J.; Stevenson, D. K.; Beutler, E.: Acute hemolysis and severe neonatal hyperbilirubine-
- [91577]

mia in glucose-6-phosphate

[91578] dehydrogenase deficient heterozygotes. J. Pediat. 139:
137-140,

[91579] 2001.

[91580]

[91581] 160. Kaplan, M.; Renbaum, P.; Levy-Lahad, E.; Hammer-
man, C.; Lahad,

[91582] A. Beutler, E.: Gilbert syndrome and glucose-6-phosphate
dehydrogenase

[91583] deficiency: a dose-dependent genetic interaction crucial
to neonatal

[91584] hyperbilirubinemia. Proc. Nat. Acad. Sci. 94:
12128-12132, 1997.

[91585]

[91586] 161. Kappas, A.; Drummond, G. S.; Valaes, T.: A single
dose of Sn-mesoporphyrin

[91587] prevents development of severe hyperbilirubinemia in
glucose-6-phosphate

[91588] dehydrogenase-deficient newborns. Pediatrics 108:
25-30, 2001.

[91589]

[91590] 162. Karadsheh, N. S.; Awidi, A. S.; Tarawneh, M. S.: Two

new glucose-6

[91591] phosphate dehydrogenase (G6PD) variants associated with hemolytic

[91592] anemia: G6PD Amman-1 and G6PD Amman-2. Am. J. Hemat. 22: 185-192,

[91593] 1986.

[91594]

[91595] 163. Kay, A. C.; Kuhl, W.; Prchal, J.; Beutler, E.: The origin of

[91596] glucose-6-phosphate-dehydrogenase (G6PD) polymorphisms in African-Americans. Am.

[91597] J. Hum. Genet. 50: 394-398, 1992.

[91598]

[91599] 164. Kirkman, H. N.; Kidson, C.; Kennedy, M.: Variants of human glucose-6-phosphate

[91600] dehydrogenase. Studies of samples from New Guinea. In: Beutler, E.

[91601] : Hereditary Disorders of Erythrocyte Metabolism. New York: Grune

[91602] and Stratton (pub.) 1968. Pp. 126-145.

[91603]

- [91604] 165. Kirkman, H. N.; Lie-Injo, L. E.: Variants of glucose 6-phosphate
- [91605] dehydrogenase in Indonesia. *Nature* 221: 959, 1969.
- [91606]
- [91607] 166. Kirkman, H. N.; McCurdy, P. R.; Naiman, J. L.: Functionally
- [91608] abnormal glucose-6-phosphate dehydrogenases. *Cold Spring Harbor Symp.*
- [91609] *Quant. Biol.* 29: 391-398, 1964.
- [91610]
- [91611] 167. Kirkman, H. N.; Ramot, B.; Lee, J. T.: Altered aggregational
- [91612] properties in a genetic variant of human glucose-6-phosphate dehydrogenase. *Biochem.*
- [91613] *Genet.* 3: 137-150, 1969.
- [91614]
- [91615] 168. Kirkman, H. N.; Riley, H. D., Jr.: Congenital nonspherocytic
- [91616] hemolytic anemia. *Am. J. Dis. Child.* 102: 313-320, 1961.
- [91617]

[91618] 169. Kirkman, H. N.; Rosenthal, I. M.; Simon, E. B.; Carson, P. E.;

[91619] Brinson, A. G.: 'Chicago I' variant of glucose-6-phosphate dehydrogenase

[91620] in congenital hemolytic disease. J. Lab. Clin. Med. 63: 715-725,

[91621] 1964.

[91622]

[91623] 170. Kirkman, H. N.; Schettini, E.; Pickard, B. M.: Mediter-ranean

[91624] variant of glucose-6-phosphate dehydrogenase. J. Lab. Clin. Med. 63:

[91625] 726-735, 1964.

[91626]

[91627] 171. Kirkman, H. N.; Simon, E. R.; Pickard, B. M.: Seattle variant

[91628] of glucose-6-phosphate dehydrogenase. J. Lab. Clin. Med. 66: 834-840,

[91629] 1965.

[91630]

[91631] 172. Kissin, C.; Cotte, J.: Etude d'un variant de glucose-

6-phosphate

[91632] deshydrogenase: I B type Constantine. Enzyme 11: 277-284, 1970.

[91633]

[91634] 173. Kitao, T.; Ito, K.; Hattori, K.; Matsuki, T.; Yoneyama, Y.:

[91635] G6PD Kanazawa: a new variant of glucose-6-phosphate dehydrogenase

[91636] associated with congenital nonspherocytic hemolytic anemia. Acta

[91637] Haemat. 68: 131-135, 1982.

[91638]

[91639] 174. Kojima, H.: Congenital nonspherocytic hemolytic disease (CNHD)

[91640] due to a G-6-PD variant: G-6-PD Kyoto. Acta Haemat. Jpn. 35: 32-38,

[91641] 1972.

[91642]

[91643] 175. Koliakos, G.; Kalomenopoulou, M.; Grammatikos, P.; Dimitriadou,

[91644] A.; Kouzi-Koliakos, K.; Zacharaki, R.; Skaragas, G.; Kokka,

A.; Trakatellis,

[91645] A.: A new glucose 6-phosphate dehydrogenase variant
(G6PD Thessaloniki)

[91646] in a patient with idiopathic myelofibrosis. Hum. Hered. 39:
141–149,

[91647] 1989.

[91648]

[91649] 176. Krasnopol'skaya, K. D.; Bochkov, N. P.: Genetic het-
erogeneity

[91650] of hereditary enzymopathies. Vestn. Akad. Med. Nauk.
SSSR 9: 56–64,

[91651] 1982.

[91652]

[91653] 177. Krasnopol'skaya, K. D.; Shatskaya, T. L.; Filippov, I.
K.; Annenkov,

[91654] G. A.; Zakharova, T. V.; Mekhtiev, N. K.; Movsum-Zade, K.
M.: Genetic

[91655] heterogeneity of G6PD deficiency: study of mutant alleles
in Shekii

[91656] district of Azerbaijan. Genetika 13: 1455–1461, 1977.

[91657]

- [91658] 178. Kumakawa, T.; Suzuki, S.; Fujii, H.; Miwa, S.: Frequency of
[91659] glucose 6-phosphate dehydrogenase (G6PD) deficiency in
Tokyo and a
[91660] new variant: G6PD Musashino. *Acta Haemat. Jpn.* 50:
25–28, 1987.
[91661]
- [91662] 179. Kurdi-Haidar, B.; Mason, P. J.; Berrebi, A.; Ankra-
Badu, G.;
[91663] Al-Ali, A.; Oppenheim, A.; Luzzatto, L.: Origin and spread
of the
[91664] glucose-6-phosphate dehydrogenase variant
(G6PD-Mediterranean) in
[91665] the Middle East. *Am. J. Hum. Genet.* 47: 1013–1019,
1990.
[91666]
- [91667] 180. Kwiatkowska, J.; Kacprzak-Bergman, I.: New erythro-
cyte glucose
[91668] 6-phosphate dehydrogenase variant. *Acta Haemat.* 46:
188–192, 1971.
[91669]
- [91670] 181. Kwok, C. J.; Martin, A. C. R.; Au, S. W. N.; Lam, V. M.

S.:

- [91671] G6PDdb, an integrated database of glucose-6-phosphate dehydrogenase
- [91672] (G6PD) mutations. Hum. Mutat. 19: 217-224, 2002.
- [91673]
- [91674] 182. Lee, K. T.; Thomas, W. A.; Janakidevi, K.; Kroms, M.; Reiner, J. M.; Borg, K. Y.: Mosaicism in female hybrid hares heterozygous for glucose-6-phosphate dehydrogenase (G-6-PD). I. General properties
- [91677] of a hybrid hare model with special reference to atherogenesis. Exp.
- [91678] Molec. Path. 34: 191-201, 1981.
- [91679]
- [91680] 183. Lenzerini, L.; Meera Khan, P.; Filippi, G.; Rattazzi, M. C.;
- [91681] Rat, A. K.: Characterization of glucose-6-phosphate dehydrogenase
- [91682] variants. I. Occurrence of a G6PD Seattle-like variant in Sardinia
- [91683] and its interaction with G6PD Mediterranean variant. Am.

J. Hum.

[91684] Genet. 21: 142–153, 1969.

[91685]

[91686] 184. Lisker, R.; Linares, C.; Motulsky, A. G.: Glucose-6-phosphate

[91687] dehydrogenase Mexico, a new variant with enzyme deficiency, abnormal

[91688] mobility and absence of hemolysis. J. Lab. Clin. Med. 29: 788–793,

[91689] 1972.

[91690]

[91691] 185. Lisker, R.; Perez-Briceno, R.; Beutler, E.: A new glucose-6-phosphate

[91692] dehydrogenase variant, Gd(-) Tepic, characterized by moderate enzyme

[91693] deficiency and mild episodes of hemolytic anemia. Hum. Genet. 69:

[91694] 19–21, 1985.

[91695]

[91696] 186. Lisker, R.; Perez-Briceno, R.; Rave, V.; Yoshida, A.: Glucosa-6-fosfato

- [91697] deshidrogenasa Gd(-) Distrito Federal: nueva variante asociada a deficiencia
- [91698] enzimatica moderada y anemia hemolitica ocasional. Rev. Invest. Clin. 33:
- [91699] 209-211, 1981.
- [91700]
- [91701] 187. Lisker, R.; Perez Briceno, R.; Agrilar, L.; Yoshida, A.: A variant
- [91702] glucose-6-phosphate dehydrogenase Gd(-) Chiapas associated with moderate
- [91703] enzyme deficiency and occasional hemolytic anemia. Hum. Genet. 43:
- [91704] 81-84, 1978.
- [91705]
- [91706] 188. Lisker, R.; Perez Briceno, R.; Zavala, C.; Navarrette, J. I.;
- [91707] Wessels, M.; Yoshida, A.: A glucose 6-phosphate dehydrogenase Gd(-)
- [91708] Castilla variant characterized by mild deficiency associated with
- [91709] drug-induced hemolytic anemia. J. Lab. Clin. Med. 90: 754-759, 1977.

[91710]

[91711] 189. Liu, Y.; Phelan, J.; Go, R. C. P.; Prchal, J. F.; Prchal, J.

[91712] T.: Rapid determination of clonality by detection of two closely-linked

[91713] X chromosome exonic polymorphisms using allele-specific PCR. J. Clin.

[91714] Invest. 99: 1984-1990, 1997.

[91715]

[91716] 190. Long, W. K.; Kirkman, H. N.; Sutton, H. H.: Electrophoretically

[91717] slow variants of glucose-6-phosphate dehydrogenase from red cells

[91718] of Negroes. J. Lab. Clin. Med. 65: 81-87, 1965.

[91719]

[91720] 191. Longo, L.; Vanegas, O. C.; Patel, M.; Rosti, V.; Li, H.; Waka,

[91721] J.; Merghoub, T.; Pandolfi, P. P.; Notaro, R.; Manova, K.; Luzzatto,

[91722] L.: Maternally transmitted severe glucose 6-phosphate dehydrogenase

[91723] deficiency is an embryonic lethal. EMBO J. 21: 4229-4239,

2002.

[91724]

[91725] 192. Luzzatto, L.: Genetic heterogeneity and pathophysiology of G6PD

[91726] deficiency. Brit. J. Haemat. 28: 151–156, 1974.

[91727]

[91728] 193. Luzzatto, L.: Personal Communication. London, England 4/1990.

[91729]

[91730] 194. Luzzatto, L.; Afolayam, A.: Enzyme properties of different types

[91731] of human erythrocyte glucose–6–phosphate dehydrogenase with characterization

[91732] of two new genetic variants. J. Clin. Invest. 47: 1833–1842, 1968.

[91733]

[91734] 195. Luzzatto, L.; Martini, G.: X-Linked Wiskott–Aldrich syndrome

[91735] in a girl. (Letter) New Eng. J. Med. 338: 1850–1851, 1998.

[91736]

[91737] 196. Luzzatto, L.; Usanga, E. A.; Bienzle, U.; Esan, G. F. J.; Fusuan,

[91738] F. A.: Imbalance in X–chromosome expression: evidence for a human

[91739] X-linked gene affecting growth of hemopoietic cells. Sci–

ence 205:

[91740] 1418–1420, 1979.

[91741]

[91742] 197. Luzzatto, L.; Usanga, E. A.; Reddy, S.: Glucose-6-phosphate

[91743] dehydrogenase deficient red cells: resistance to infection by malarial

[91744] parasites. Science 164: 839–842, 1969.

[91745]

[91746] 198. MacDonald, D.; Town, M.; Mason, P.; Vulliamy, T.; Luzzatto, L.;

[91747] Goff, D. K.: Deficiency in red blood cells. (Letter) Nature 350:

[91748] 115, 1991.

[91749]

[91750] 199. Maeda, M.; Constantoulakis, P.; Chen, C.–S.; Stamatoyannopoulos,

[91751] G.; Yoshida, A.: Molecular abnormalities of a human glucose-6-phosphate

[91752] dehydrogenase variant associated with undetectable enzyme activity

[91753] and immunologically cross-reacting material. Am. J. Hum. Genet. 51:

[91754] 386–395, 1992.

[91755]

[91756] 200. Mallouh, A. A.; Abu-Osba, Y. K.: Bacterial infections in children

[91757] with glucose–6–phosphate dehydrogenase deficiency. J. Pediat. 111:

[91758] 850–852, 1987.

[91759]

[91760] 201. Mamlok, R. J.; Mamlok, V.; Mills, G. C.; Daeschner, C. W., III;

[91761] Schmalstieg, F. C.; Anderson, D. C.: Glucose–6–phosphate dehydrogenase

[91762] deficiency, neutrophil dysfunction and Chromobacterium violaceum sepsis. J.

[91763] Pediat. 111: 852–854, 1987.

[91764]

[91765] 202. Mamlok, R. J.; Mills, G. C.; Goldblum, R. M.; Daeschner, C. W.

[91766] : Glucose–6–phosphate dehydrogenase Beaumont: a new

variant with severe

[91767] enzyme deficiency and chronic nonspherocytic hemolytic anemia. Enzyme 34:

[91768] 15–21, 1985.

[91769]

[91770] 203. Mandelli, F.; Amadori, S.; DeLaurenzi, A.; Kahn, A.; Isacchi,

[91771] G.; Papa, G.: Glucose–6–phosphate dehydrogenase Vellettri: a new variant

[91772] with reduced activity in a patient with congenital non-spherocytic

[91773] haemolytic anemia. Acta Haemat. 57: 121–126, 1977.

[91774]

[91775] 204. Marks, P. A.; Banks, J.; Gross, R.: Genetic heterogeneity of

[91776] glucose–6–phosphate dehydrogenase deficiency. Nature 194: 454–456,

[91777] 1962.

[91778]

[91779] 205. Martin, S. K.; Miller, L. H.; Alling, D.; Okoye, V. C.; Esan,

- [91780] G. J. F.; Osunkoya, B. O.; Deane, M.: Severe malaria and glucose-6-phosphate-dehydrogenase
- [91781] deficiency: a reappraisal of the malaria-G6PD hypothesis. Lancet I:
- [91782] 524-526, 1979.
- [91783]
- [91784] 206. Martini, G.; Toniolo, D.; Vulliamy, T.; Luzzatto, L.;
Dono, R.;
- [91785] Viglietto, G.; Paonessa, G.; D'Urso, M.; Persico, M. G.:
Structural
- [91786] analysis of the X-linked gene encoding human glucose
6-phosphate dehydrogenase. EMBO
- [91787] J. 5: 1849-1855, 1986.
- [91788]
- [91789] 207. Mason, P. J.: New insights into G6PD deficiency.
(Annotation) Brit.
- [91790] J. Haemat. 94: 585-591, 1996.
- [91791]
- [91792] 208. Mason, P. J.; Stevens, D. J.; Luzzatto, L.; Brenner, S.;
Aparicio,
- [91793] S.: Genomic structure and sequence of the Fugu rubripes

glucose-6-phosphate

[91794] dehydrogenase gene (G6PD). Genomics 26: 587-591, 1995.

[91795]

[91796] 209. McCann, S. R.; Smithwick, A. M.; Temperley, I. J.; Tipton, K.

[91797] : G6PD (Dublin): chronic non-spherocytic haemolytic anaemia resulting

[91798] from glucose-6-phosphate dehydrogenase deficiency in an Irish kindred. J.

[91799] Med. Genet. 17: 191-193, 1980.

[91800]

[91801] 210. McCurdy, P. R.: Use of genetic linkage for the detection of

[91802] female carriers of hemophilia. New Eng. J. Med. 285: 218-219, 1971.

[91803]

[91804] 211. McCurdy, P. R.; Blackwell, R. Q.; Todd, D.; Tso, S. C.; Tuchinda,

[91805] S.: Further studies on glucose-6-phosphate dehydrogenase deficiency

[91806] in Chinese subjects. J. Lab. Clin. Med. 75: 788-797, 1970.

[91807]

[91808] 212. McCurdy, P. R.; Kamel, K.; Selim, O.: Heterogeneity of red cell

[91809] glucose 6-phosphate dehydrogenase (G6PD) deficiency in Egypt. J.

[91810] Lab. Clin. Med. 84: 673-680, 1974.

[91811]

[91812] 213. McCurdy, P. R.; Kirkman, H. N.; Naiman, J. L.; Jim, R. T. S.;

[91813] Pickard, B. M.: A Chinese variant of glucose-6-phosphate dehydrogenase. J.

[91814] Lab. Clin. Med. 67: 374-385, 1966.

[91815]

[91816] 214. McCurdy, P. R.; Mahmood, L.: Red cell glucose-6-phosphate dehydrogenase

[91817] deficiency in Pakistan. J. Lab. Clin. Med. 76: 943-948, 1970.

[91818]

[91819] 215. McCurdy, P. R.; Maldonado, N. I.; Dillon, D. E.: Variants of

[91820] glucose-6-phosphate dehydrogenase (G-6-PD) associated

with G-6-PD

[91821] deficiency in Puerto Ricans. J. Lab. Clin. Med. 82:
432-437, 1973.

[91822]

[91823] 216. McCurdy, P. R.; Maldonado, N. I.; Dillon, D. E.; Conrad, M. E.

[91824] : Variants of glucose-6-phosphate dehydrogenase
(G-6-PD) associated

[91825] with G-6-PD deficiency in Puerto Ricans. J. Lab. Clin. Med.
82:

[91826] 432-437, 1973.

[91827]

[91828] 217. Meloni, T.; Carta, F.; Forteleoni, G.; Carta, A.; Ena, F.;
Meloni,

[91829] G. F.: Glucose 6-phosphate dehydrogenase deficiency and
cataract

[91830] of patients in northern Sardinia. Am. J. Ophthal. 110:
661-664,

[91831] 1990.

[91832]

[91833] 218. Mentzer, W. C., Jr.; Warner, R.; Addiego, J.; Smith, B.;

Walter,

- [91834] T.: G6PD San Francisco: a new variant of glucose-6-phosphate dehydrogenase
- [91835] associated with congenital nonspherocytic hemolytic anemia. Blood 55:
- [91836] 195–198, 1980.
- [91837]
- [91838] 219. Mesbah–Namin, S. A.; Sanati, M. H.; Mowjoodi, A.; Mason, P. J.;
- [91839] Vulliamy, T. J.; Noori–Daloui, M. R.: Three major glucose-6-phosphate
- [91840] dehydrogenase–deficient polymorphic variants identified in Mazandaran
- [91841] state of Iran. Brit. J. Haemat. 117: 763–764, 2002.
- [91842]
- [91843] 220. Miller, D. R.; Wollman, M. R.: A new variant of glucose 6–phosphate
- [91844] dehydrogenase deficiency hereditary hemolytic anemia, G6PD Cornell:
- [91845] erythrocyte, leukocyte and platelet studies. Blood 44: 277–284,
- [91846] 1974.

[91847]

[91848] 221. Mills, G. C.; Alperin, J. B.; Trimmer, K. B.: Studies on variant

[91849] glucose-6-phosphate dehydrogenase: G6PD Fort Worth. Biochem. Med. 13:

[91850] 264-275, 1975.

[91851]

[91852] 222. Milner, G.; Delamore, I. W.; Yoshida, A.: G-6-PD Manchester:

[91853] a new variant associated with chronic nonspherocytic hemolytic anemia. Blood 43:

[91854] 271-276, 1974.

[91855]

[91856] 223. Miwa, S.; Fujii, H.: Molecular basis of erythroenzy-mopathies

[91857] associated with hereditary hemolytic anemia: tabulation of mutant

[91858] enzymes. Am. J. Hemat. 51: 122-132, 1996.

[91859]

[91860] 224. Miwa, S.; Fujii, H.; Nakashima, K.; Miura, Y.; Yamada, K.; Hagiwara,

- [91861] T.; Fukuda, M.: Three new electrophoretically normal glucose-6-phosphate
- [91862] dehydrogenase variants associated with congenital nonspherocytic hemolytic
- [91863] anemia found in Japan: G6PD Ogikubo, Yokohama, and Akita. Hum. Genet. 45:
- [91864] 11-17, 1978.
- [91865]
- [91866] 225. Miwa, S.; Fujii, H.; Nakatsuji, T.; Ishida, Y.; Oda, E.; Kaneto,
- [91867] A.; Motokawa, M.; Ariga, Y.; Fukuchi, S.; Sasai, S.; Hiraoka, K.;
- [91868] Kashii, H.; Kodama, T.: Four new electrophoretically slow-moving
- [91869] glucose-6-phosphate dehydrogenase variants associated with congenital
- [91870] nonspherocytic hemolytic anemia found in Japan: Gd(-) Kurume, Gd(-)
- [91871] Fukushima, Gd(-) Yamaguchi, and Gd(-) Wakayama. Am. J. Hemat. 5:
- [91872] 131-138, 1978.
- [91873]

- [91874] 226. Miwa, S.; Nakashima, K.; Ono, J.; Fujii, H.; Suzuki, E.:
Three
[91875] glucose 6-phosphate dehydrogenase variants found in
Japan. Hum. Genet. 36:
[91876] 327-334, 1977.
[91877]
- [91878] 227. Miwa, S.; Ono, J.; Nakashima, K.; Abe, S.; Kageoka,
T.; Shinohara,
[91879] K.; Isobe, J.; Yamaguchi, H.: Two new glucose
6-phosphate dehydrogenase
[91880] variants associated with congenital nonspherocytic
hemolytic anemia
[91881] found in Japan: Gd(-) Tokushima and Gd(-) Tokyo. Am. J.
Hemat. 1:
[91882] 433-442, 1976.
[91883]
- [91884] 228. Modiano, G.; Battistuzzi, G.; Esan, G. J. F.; Testa, U.;
Luzzatto,
[91885] L.: Genetic heterogeneity of 'normal' human erythrocyte
glucose-6-phosphate
[91886] dehydrogenase: an isoelectrophoretic polymorphism.
Proc. Nat. Acad.

[91887] Sci. 76: 852–856, 1979.

[91888]

[91889] 229. Mohrenweiser, H. W.; Neel, J. V.: Frequency of thermostability

[91890] variants: estimation of total 'rare' variant frequency in human populations. Proc.

[91891] Nat. Acad. Sci. 78: 5729–5733, 1981.

[91892]

[91893] 230. Morelli, A.; Benatti, U.; Guida, L.; De Flora, A.: G6PD Cagliari:

[91894] a new low activity glucose 6–phosphate dehydrogenase variant characterized

[91895] by enhanced intracellular lability. Hum. Genet. 66: 62–65, 1984.

[91896]

[91897] 231. Morisaki, T.; Fujii, H.; Takegawa, S.; Tani, K.; Hirono, A.;

[91898] Takizawa, T.; Takahashi, K.; Shinogi, M.; Teshirogi, T.; Miwa, S.

[91899] : G6PD Sendagi: a new glucose–6–phosphate dehydrogenase variant associated

- [91900] with congenital hemolytic anemia. Hum. Genet. 65: 214–215, 1983.
- [91901]
- [91902] 232. Nafa, K.; Reghis, A.; Osmani, N.; Baghli, L.; Ben-abadji, M.;
- [91903] Kaplan, J.-C.; Vulliamy, T. J.; Luzzatto, L.: G6PD Aures: a new mutation
- [91904] (48 ile-to-thr) causing mild G6PD deficiency is associated with favism. Hum.
- [91905] Molec. Genet. 2: 81–82, 1993.
- [91906]
- [91907] 233. Nagel, R. L.; Ranney, H. M.: Genetic epidemiology of structural
- [91908] mutations of the beta-globin gene. Semin. Hemat. 27: 342–359, 1990.
- [91909]
- [91910] 234. Nakai, T.; Yoshida, A.: G6PD Heian. A glucose-6-phosphate dehydrogenase
- [91911] variant associated with hemolytic anemia found in Japan. Clin. Chim.
- [91912] Acta 51: 199–203, 1974.

[91913]

[91914] 235. Nakashima, K.; Ono, J.; Abe, S.; Miwa, S.; Yoshida, A.:
G6PD

[91915] Ube: a glucose 6-phosphate dehydrogenase variant found
in four unrelated

[91916] Japanese families. Am. J. Hum. Genet. 29: 24-30, 1977.

[91917]

[91918] 236. Nakatsuji, T.; Miwa, S.: Incidence and characteristics
of glucose-6-phosphate

[91919] dehydrogenase variants in Japan. Hum. Genet. 51:
297-305, 1979.

[91920]

[91921] 237. Nance, W. E.: Turner's syndrome, twinning, and an
unusual variant

[91922] of glucose-6-phosphate dehydrogenase. Am. J. Hum.
Genet. 16: 380-392,

[91923] 1964.

[91924]

[91925] 238. Necheles, T. F.; Snyder, L. M.; Strauss, W.: Glucose-
6-phosphate

[91926] dehydrogenase Boston. A new variant associated with

congenital nonspherocytic

[91927] hemolytic disease. Humangenetik 13: 218–221, 1971.

[91928]

[91929] 239. Niazi, G.; Adeyokunu, A.; Westwood, B.; Beutler, E.:
G6PD Aures:

[91930] a rare mutant of G6PD in Saudi Arabia: molecular and
clinical presentations. Saudi

[91931] Med. J. 17: 311–314, 1996.

[91932]

[91933] 240. Ninfali, P.; Baronciani, L.; Bardoni, A.; Bresolin, N.:
Muscle

[91934] expression of glucose–6–phosphate dehydrogenase defi-
ciency in different

[91935] variants. Clin. Genet. 48: 232–237, 1995.

[91936]

[91937] 241. Notaro, R.; Afolayan, A.; Luzzatto, L.: Human muta-
tions in glucose

[91938] 6–phosphate dehydrogenase reflect evolutionary history.
FASEB J. 14:

[91939] 485–494, 2000.

[91940]

- [91941] 242. Nowicki, L.; Strobel, S.; Martin, H.; Koschwitz, U.: Ueber eine
ber eine
- [91942] neue erythrocytaere glucose 6-phosphatdehydrogenase
Variante, Typ
- [91943] Frankfurt. Klin. Wschr. 52: 478-484, 1974.
- [91944]
- [91945] 243. Nsouly, G. M.; Prchal, J. T.: Characterization of a new
G6PD
- [91946] variant and its associated oxidative damage. Clin. Res. 29:
829,
- [91947] 1981.
- [91948]
- [91949] 244. O'Brien, S. J.: The extent and character of biochemical genetic
cal genetic
- [91950] variation in the domestic cat. J. Hered. 71: 2-8, 1980.
- [91951]
- [91952] 245. Ogura, H.; Morisaki, T.; Tani, K.; Kanno, H.; Tsutsumi, H.; Takahashi,
sumi, H.; Takahashi,
- [91953] K.; Miyamori, T.; Fujii, H.; Miwa, S.: A new glucose-
6-phosphate
- [91954] dehydrogenase variant (G6PD Tsukui) associated with

congenital hemolytic

[91955] anemia. Hum. Genet. 78: 369–371, 1988.

[91956]

[91957] 246. Ohno, S.: Sex Chromosomes and Sex-linked Genes.
Berlin and

[91958] New York: Springer (pub.) 1967.

[91959]

[91960] 247. Orzalesi, N.; Sorcinelli, R.; Guiso, G.: Increased incidence

[91961] of cataracts in male subjects deficient in glucose-
6-phosphate dehydrogenase. Arch.

[91962] Ophthal. 99: 69–70, 1981.

[91963]

[91964] 248. Othieno-Obel, A.: East African variant of glucose-
6-phosphate

[91965] dehydrogenase. East Afr. Med. J. 49: 230–234, 1972.

[91966]

[91967] 249. Pai, G. S.; Sprenkle, J. A.; Do, T. T.; Maren, C. E.; Mi-
geon,

[91968] B. R.: Localization of loci for hypoxanthine phosphoribo-

sytransferase

[91969] **and glucose-6-phosphate dehydrogenase and biochemical evidence of**

[91970] **nonrandom X chromosome expression from studies of a human X-autosome**

[91971] **translocation. Proc. Nat. Acad. Sci. 77: 2810-2813, 1980.**

[91972]

[91973] **250. Panich, V.: G6PD variants in Laotians. Hum. Hered. 24: 285-290,**

[91974] **1974.**

[91975]

[91976] **251. Panich, V.: G6PD Intanon, a new glucose 6-phosphate dehydrogenase**

[91977] **variant. Humangenetik 21: 203-205, 1974.**

[91978]

[91979] **252. Panich, V.: G6PD characterization in Thailand. Genetics 74**

[91980] **(suppl.): s208 only, 1973.**

[91981]

[91982] **253. Panich, V.: Glucose-6-phosphate dehydrogenase in**

Thailand. Hum.

[91983] Genet. 53: 227–228, 1980.

[91984]

[91985] 254. Panich, V.; Bumrungtrakul, P.; Jitjai, C.; Kamolmatayakul, S.;

[91986] Khoprasert, B.; Klaisuvan, C.; Kongmuang, U.; Maneechai, P.; Pornpatkul,

[91987] M.; Ruengrairatanaroje, P.; Surapruk, P.; Viriyayudhakorn, S.: Glucose–6–phosphate

[91988] dehydrogenase deficiency in South Vietnamese. Hum. Hered. 30: 361–364,

[91989] 1980.

[91990]

[91991] 255. Panich, V.; Na–Nakorn, S.: G6PD variants in Thailand. J. Med.

[91992] Assoc. Thai. 63: 537–543, 1980.

[91993]

[91994] 256. Panich, V.; Sungnate, T.: Characterization of glucose 6–phosphate

[91995] dehydrogenase in Thailand: the occurrence of 6 variants among 50 G6PD

[91996] deficient Thai. Humangenetik 18: 39–46, 1973.

[91997]

[91998] 257. Panich, V.; Sungnate, T.; Na-Nakorn, S.: Acute intravascular

[91999] hemolysis and renal failure in a new glucose 6-phosphate dehydrogenase

[92000] variant: G6PD Siriraj. J. Med. Assoc. Thai. 55: 726–731, 1972.

[92001]

[92002] 258. Panich, V.; Sungnate, T.; Wasi, P.; Na-Nakorn, S.: G-6-PD Mahidol:

[92003] the most common glucose-6-phosphate dehydrogenase variant in Thailand. J.

[92004] Med. Assoc. Thai. 55: 576–585, 1972.

[92005]

[92006] 259. Pawlak, A. L.; Mazurkiewicz, C. A.; Ordynski, J.; Ruzynkowa,

[92007] D.; Horst, A.: G6PD Poznan, variant with severe enzyme deficiency. Humangenetik 28:

[92008] 163–165, 1975.

[92009]

[92010] 260. Pawlak, A. L.; Zagorski, Z.; Rozyrkowa, D.; Horst, A.:
Polish

[92011] variant of glucose-6-phosphate dehydrogenase (G-6-PD
Lublin). Humangenetik 10:

[92012] 340-343, 1970.

[92013]

[92014] 261. Pekrun, A.; Eber, S. W.; Schroter, W.: G6PD Avenches
and G6PD

[92015] Moosburg: biochemical and erythrocyte membrane char-
acterization. Blut 58:

[92016] 11-14, 1989.

[92017]

[92018] 262. Perroni, L.; Tassara, P.; Baldi, M.; Reali, R.;
Scartezzini,

[92019] P.: G6PD variants detected in Genoa area. In: Weatherall, D.
J.; Fiorelli,

[92020] G.; Gorini, S.: Advances in Red Blood Cell Biology. New
York: Raven

[92021] Press (pub.) 1982. Pp. 409-416.

[92022]

[92023] 263. Persico, M. G.; Toniolo, D.; Nobile, C.; D'Urso, M.;

Luzzatto,

[92024] L.: cDNA sequences of human glucose 6-phosphate dehydrogenase cloned

[92025] in pBR322. *Nature* 294: 778-780, 1981.

[92026]

[92027] 264. Persico, M. G.; Viglietto, G.; Martini, G.; Toniolo, D.; Paonessa,

[92028] G.; Moscatelli, C.; Dono, R.; Vulliamy, T.; Luzzatto, L.; D'Urso,

[92029] M.: Isolation of human glucose-6-phosphate dehydrogenase (G6PD) cDNA

[92030] clones: primary structure of the protein and unusual 5-prime non-coding

[92031] region. *Nucleic Acids Res.* 14: 2511-2522, 1986.

[92032]

[92033] 265. Picat, C.; Etienne, J.; Boivin, P.; Le Prise, P.-Y.: Gd(-)

[92034] Rennes, a new deficient variant of glucose-6-phosphate dehydrogenase

[92035] associated with congenital nonspherocytic hemolytic anemia found in

[92036] France. *Hum. Genet.* 55: 125-127, 1980.

[92037]

[92038] 266. Pinto, P. V. C.; Newton, W. A., Jr.; Richardson, K. E.:
Evidence

[92039] for four types of erythrocyte glucose-6-phosphate dehydrogenase from

[92040] G-6-PD deficient human subjects. J. Clin. Invest. 45:
823-831, 1966.

[92041]

[92042] 267. Poggi, V.: Personal Communication. London, England
1989.

[92043]

[92044] 268. Poon, M.-C.; Hall, K.; Scott, C. W.; Prchal, J. T.: G6PD
Viangchan:

[92045] a new glucose 6-phosphate dehydrogenase variant from
Laos. Hum. Genet. 78:

[92046] 98-99, 1988.

[92047]

[92048] 269. Porter, I. H.; Boyer, S. H.; Watson-Williams, E. J.;
Adam, A.;

[92049] Szeinberg, A.; Siniscalco, M.: Variation of glucose-
6-phosphate dehydrogenase

[92050] in different populations. Lancet I: 895–899, 1964.

[92051]

[92052] 270. Porter, I. H.; Schulze, J.; McKusick, V. A.: Genetical linkage

[92053] between the loci for glucose–6–phosphate dehydrogenase deficiency

[92054] and colour–blindness in American Negroes. Ann. Hum. Genet. 26: 107–122,

[92055] 1962.

[92056]

[92057] 271. Prchal, J.; Moreno, H.; Conrad, M.; Vitek, A.: G–6–PD Dothan:

[92058] a new variant associated with chronic hemolytic anemia. I.R.C.S. 7:

[92059] 348, 1979.

[92060]

[92061] 272. Prchal, J. T.: Personal Communication. Birmingham, Ala. 1985.

[92062]

[92063] 273. Prchal, J. T.; Crist, W. M.; Malluh, A.; Vitek, A.; Tauxe, W.

- [92064] N.; Carroll, A. J.: A new glucose-6-phosphate dehydrogenase deficient
- [92065] variant in a patient with Chediak-Higashi syndrome. Blood 56: 476-480,
- [92066] 1980.
- [92067]
- [92068] 274. Prchal, J. T.; Hall, K.; Csepreghy, M.; Lilly, M.; Berkow, R.;
- [92069] Scott, C. W.: Two apparent glucose-6-phosphate dehydrogenase variants
- [92070] in normal XY males: G6PD Alabama. Am. J. Med. 84: 517-523, 1988.
- [92071]
- [92072] 275. Pretsch, W.; Charles, D. J.; Merkle, S.: X-linked glucose-6-phosphate-dehydrogenase
- [92073] deficiency in *Mus musculus*. Biochem. Genet. 26: 89-103, 1988.
- [92074]
- [92075] 276. Puck, J. M.; Willard, H. F.: X inactivation in females with
- [92076] X-linked disease. New Eng. J. Med. 338: 325-327, 1998.

[92077]

[92078] 277. Ramot, B.; Ben-Bassat, I.; Shchory, M.: New glucose-6-phosphate

[92079] dehydrogenase variants observed in Israel and their association with

[92080] congenital nonspherocytic hemolytic disease. J. Lab. Clin. Med. 74:

[92081] 895-901, 1969.

[92082]

[92083] 278. Ramot, B.; Brok, F.: A new glucose-6-phosphate dehydrogenase

[92084] mutant (Tel-Hashomer mutant). Ann. Hum. Genet. 28: 167-172, 1964.

[92085]

[92086] 279. Rattazzi, M. C.; Corash, L. M.; Van Zzanen, G. E.; Jaffe, E.

[92087] R.; Piomelli, S.: G6PD deficiency and chronic hemolysis: four new

[92088] mutants--relationships between clinical syndrome and enzyme kinetics. Blood 38:

[92089] 205-218, 1971.

[92090]

[92091] 280. Rattazzi, M. C.; Lenzerini, L.; Meera Khan, P.; Luzzatto, L.

[92092] : Characterization of glucose-6-phosphate dehydrogenase variants.

[92093] II. G6PD Kephallonia, G6PD Attica, and G6PD 'Seattle-like' found in

[92094] Greece. Am. J. Hum. Genet. 21: 154-167, 1969.

[92095]

[92096] 281. Ravindranath, Y.; Beutler, E.: Two new variants of glucose-6-phosphate

[92097] dehydrogenase associated with hereditary non-spherocytic hemolytic

[92098] anemia: G6PD Wayne and G6PD Huron. Am. J. Hemat. 24: 357-363, 1987.

[92099]

[92100] 282. Reys, L.; Manso, C.; Stamatoyannopoulos, G.: Genetic studies

[92101] on Southeastern Bantu of Mozambique. I. Variants of glucose-6-phosphate

[92102] dehydrogenase. Am. J. Hum. Genet. 22: 203-215, 1970.

[92103]

[92104] 283. Rinaldi, A.; Filippi, G.; Siniscalco, M.: Variability of
red

[92105] cell phenotypes between and within individuals in an un-
biased sample

[92106] of 77 heterozygotes for G6PD deficiency in Sardinia. Am.
J. Hum.

[92107] Genet. 28: 496–505, 1976.

[92108]

[92109] 284. Rosenstrauss, M.; Chasin, L. A.: Isolation of mam-
malian cell

[92110] mutants deficient in glucose–6–phosphate dehydrogenase
activity: linkage

[92111] to hypoxanthine phosphoribosyl transferase. Proc. Nat.
Acad. Sci. 72:

[92112] 493–497, 1975.

[92113]

[92114] 285. Roth, E. F., Jr.; Raventos–Suarez, C.; Rinaldi, A.;
Nagel, R.

[92115] L.: Glucose–6–phosphate dehydrogenase deficiency in-
hibits in vitro

- [92116] growth of *Plasmodium falciparum*. Proc. Nat. Acad. Sci.
80: 298–299,
[92117] 1983.
- [92118]
- [92119] 286. Roychoudhury, A. K.; Nei, M.: Human Polymorphic
Genes: World
[92120] Distribution. New York: Oxford Univ. Press (pub.) 1988.
- [92121]
- [92122] 287. Ruwando, C.; Khea, S. C.; Snow, R. W.; Yates, S. N. R.;
Kwiatkoweld,
[92123] D.; Gupta, S.; Warn, P.; Alisopp, G. E. M.; Gilbert, S. C.;
Peschu,
[92124] N.; Newbold, C. I.; Greenwood, S. M.; Marsh, K.; Hill, A. V.
S.:
[92125] Natural selection of hemi- and heterozygotes for G6PD
deficiency in
[92126] Africa by resistance to severe malaria. Nature 376:
246–249, 1995.
- [92127]
- [92128] 288. Saenz, G. F.; Chaves, M.; Berrantes, A.; Elizondo, J.;
Montero,

[92129] A. G.; Yoshida, A.: A glucose-6-phosphate dehydrogenase variant,

[92130] Gd(-) Santamaria found in Costa Rica. *Acta Haemat.* 72: 37-40, 1984.

[92131]

[92132] 289. Saidi, N.; Hors-Cayla, M. C.; Van Cong, N.; Benne, F.: Sheep

[92133] gene mapping by somatic cell hybridization. (Abstract) *Cytogenet.*

[92134] *Cell Genet.* 25: 200, 1979.

[92135]

[92136] 290. Samuel, A. P. W.; Saha, N.; Omer, A.; Hoffbrand, A. V.: Quantitative

[92137] expression of G6PD activity of different phenotypes of G6PD and haemoglobin

[92138] in a Sudanese population. *Hum. Hered.* 31: 110-115, 1981.

[92139]

[92140] 291. Sansone, G.; Perroni, L.; Testa, U.; Mareni, C.; Luz-zatto, L.

[92141] : New genetic variants of glucose 6-phosphate dehydrogenase (G6PD)

[92142] in Italy. Ann. Hum. Genet. 45: 97–104, 1981.

[92143]

[92144] 292. Sansone, G.; Perroni, L.; Yoshida, A.: Glucose–6-phosphate dehydrogenase

[92145] variants from Italian subjects associated with severe neonatal jaundice. Brit.

[92146] J. Haemat. 31: 159–165, 1975.

[92147]

[92148] 293. Sansone, G.; Perroni, L.; Yoshida, A.; Dave, V.: A new glucose

[92149] 6-phosphate dehydrogenase variant (Gd Trinacria) in two unrelated

[92150] families of Sicilian ancestry. Ital. J. Biochem. 26: 44–50, 1977.

[92151]

[92152] 294. Shatskaya, T. L.; Krasnopolskaya, K. D.; Annenkov, G. A.: A

[92153] description of new mutant forms of erythrocyte glucose–6-phosphate

[92154] dehydrogenase isolated at the territory of the Soviet Union. Genetika 11:

[92155] 116–122, 1975.

[92156]

[92157] 295. Shatskaya, T. L.; Krasnopol'skaya, K. D.; Idelson, L. J.:
The

[92158] new form of glucose 6–phosphate dehydrogenase (G6PD
'Kaluga') from

[92159] erythrocytes of a patient with chronic non–spherocytic
hemolytic anemia. Vopr.

[92160] Med. Khim. 22: 764–768, 1976.

[92161]

[92162] 296. Shatskaya, T. L.; Krasnopol'skaya, K. D.; Idelson, L. J.:
Mutant

[92163] forms of erythrocyte glucose 6–phosphate dehydrogenase
in Ashkenazi:

[92164] description of two new variants, G6PD Kirovograd and
G6PD Zhitomir. Humangenetik 33:

[92165] 175–178, 1976.

[92166]

[92167] 297. Shatskaya, T. L.; Krasnopol'skaya, K. D.; Tzoneva, M.;
Mavrudieva,

[92168] M.; Toncheva, D.: Variants of erythrocyte glucose–

6-phosphate dehydrogenase

[92169] (G6PD) in Bulgarian populations. Hum. Genet. 54: 115–117, 1980.

[92170]

[92171] 298. Shatskaya, T. L.; Krasnopolskaya, K. D.; Zakharova, T. V.: Regularities

[92172] of distribution of Gd-alleles in Azerbaijan. II. Identification of

[92173] G6PD mutant forms. Genetika 16: 2217–2225, 1980.

[92174]

[92175] 299. Shows, T. B.; Brown, J. A.: Human X-linked genes regionally

[92176] mapped utilizing X-autosome translocations and somatic cell hybrids. Proc.

[92177] Nat. Acad. Sci. 72: 2125–2129, 1975.

[92178]

[92179] 300. Shows, T. B.; Brown, J. A.; Chapman, V. M.: Comparative gene

[92180] mapping of HPRT, G6PD and PGK in man, mouse, and Muntjac deer. Birth

[92181] Defects Orig. Art. Ser. XII(7): 436–439, 1976.

[92182]

[92183] 301. Shows, T. B.; Tashian, R. E.; Brewer, G. J.: Erythrocyte glucose-6-phosphate

[92184] dehydrogenase in Caucasians: new inherited variant. Science 145:

[92185] 1056-1057, 1964.

[92186]

[92187] 302. Sidi, Y.; Aderka, D.; Brok-Simoni, F.; Benjamin, D.; Ramot, B.;

[92188] Pinkhas, J.: Viral hepatitis with extreme hyperbilirubinaemia, massive

[92189] hemolysis and encephalopathy in a patient with a new G6PD variant. Israel

[92190] J. Med. Sci. 16: 130-133, 1980.

[92191]

[92192] 303. Siegel, N. H.; Beutler, E.: Hemolytic anemia caused by G-6-PD

[92193] Carswell, a new variant. Ann. Intern. Med. 75: 437-439, 1971.

[92194]

[92195] 304. Smith, J. E.; Ryer, K.; Wallace, L.: Glucose-

6-phosphate dehydrogenase

[92196] deficiency in a dog. Enzyme 21: 379-382, 1976.

[92197]

[92198] 305. Snyder, L. M.; Necheles, T. F.; Reddy, W. J.: G-6-PD
Worcester:

[92199] a new variant, associated with X-linked optic atrophy. Am.
J. Med. 49:

[92200] 125-132, 1970.

[92201]

[92202] 306. Stamatoyannopoulos, G.; Kotsakis, P.; Voigtlander,
V.; Akrivakis,

[92203] A.; Motulsky, A. G.: Electrophoretic diversity of glucose-
6-phosphate

[92204] dehydrogenase among Greeks. Am. J. Hum. Genet. 22:
587-596, 1970.

[92205]

[92206] 307. Stamatoyannopoulos, G.; Voigtlander, V.; Akrivakis,
A.: Thessaly

[92207] variant of glucose-6-phosphate dehydrogenase. Human-
genetik 9: 23-25,

[92208] 1970.

[92209]

[92210] 308. Stamatoyannopoulos, G.; Voightlander, V.; Kotsakis, P.; Akrivakis,

[92211] A.: Genetic diversity of the 'Mediterranean' glucose-6-phosphate

[92212] dehydrogenase deficiency phenotype. J. Clin. Invest. 50: 1253-1261,

[92213] 1971.

[92214]

[92215] 309. Stamatoyannopoulos, G.; Yoshida, A.; Bacopoulos, C.; Motulsky,

[92216] A. G.: Athens variant of glucose-6-phosphate dehydrogenase. Science 157:

[92217] 831-833, 1967.

[92218]

[92219] 310. Stevens, D. J.; Wanachiwanawin, W.; Mason, P. J.; Vulliamy, T.

[92220] J.; Luzzatto, L.: G6PD Canton a common deficient variant in South

[92221] East Asia caused by a 459 arg-to-leu mutation. Nucleic Acids Res. 18:

[92222] 7190, 1990.

[92223]

[92224] 311. Stocco dos Santos, R. C.; Barretto, O. C. O.;
Nonoyama, K.; Castro,

[92225] N. H. C.; Ferraz, O. P.; Walter-Moura, J.; Vescio, C. C. S.;
Becak,

[92226] W.: X-linked syndrome: mental retardation, hip luxation,
and G6PD

[92227] variant (Gd(+)) Butantan). Am. J. Med. Genet. 39: 133-136,
1991.

[92228]

[92229] 312. Stockham, S. L.; Harvey, J. W.; Kinden, D. A.: Equine
glucose-6-phosphate

[92230] dehydrogenase deficiency. Vet. Path. 31: 518-527, 1994.

[92231]

[92232] 313. Streiff, F.; Vigneron, C.: Anémie hémolytique
chronique par

[92233] déficit en glucose 6-phosphate deshydrogenase dans une
famille d'origine

[92234] Lorraine. Nouv. Rev. Franc. Hemat. 11: 279-290, 1971.

[92235]

[92236] 314. Takahashi, K.; Fujii, H.; Takegawa, S.; Tani, K.; Hirono, A.;

[92237] Takizawa, T.; Kawakatsu, T.; Miwa, S.: A new glucose-6-phosphate

[92238] dehydrogenase variant (G6PD Nagano) associated with congenital hemolytic

[92239] anemia. Hum. Genet. 62: 368-370, 1982.

[92240]

[92241] 315. Takizawa, T.; Fujii, H.; Takegawa, S.; Takahashi, K.; Hirono,

[92242] A.; Morisaki, T.; Kanno, H.; Oka, R.; Yoshioka, H.; Miwa, S.: A unique

[92243] electrophoretic slow-moving glucose 6-phosphate dehydrogenase variant

[92244] (G6PD Asahikawa) with a markedly acidic pH optimum. Hum. Genet. 68:

[92245] 70-72, 1984.

[92246]

[92247] 316. Takizawa, T.; Huang, I.-Y.; Ikuta, T.; Yoshida, A.: Human glucose-6-phosphate

[92248] dehydrogenase: primary structure and cDNA cloning. Proc. Nat. Acad.

[92249] Sci. 83: 4157-4161, 1986.

[92250]

[92251] 317. Takizawa, T.; Yoneyama, Y.; Miwa, S.; Yoshida, A.: A single

[92252] nucleotide base transition is the basis of the common human glucose-6-phosphate

[92253] dehydrogenase variant A(+). Genomics 1: 228-231, 1987.

[92254]

[92255] 318. Takizawa, T.; Yoshida, A.: Molecular abnormality of the common

[92256] glucose-6-phosphate dehydrogenase variant, G6PD A(+), and restriction-fragment-length

[92257] polymorphism. (Abstract) Am. J. Hum. Genet. 41: A241, 1987.

[92258]

[92259] 319. Talalak, P.; Beutler, E.: G-6-PD Bangkok: a new variant found

[92260] in congenital nonspherocytic hemolytic disease (CNHD). Blood 33:

[92261] 772-776, 1969.

[92262]

- [92263] 320. Tanaka, K. R.; Beutler, E.: Hereditary hemolytic anemia due
[92264] to glucose-6-phosphate dehydrogenase Torrance: a new
variant. J.
[92265] Lab. Clin. Med. 73: 657-667, 1969.
[92266]
[92267] 321. Tang, T. K.; Huang, C.-S.; Huang, M. J.; Tam, K.-B.;
Yeh, C.-H.;
[92268] Tang, C.-J. C.: Diverse point mutations result in glucose-
6-phosphate
[92269] dehydrogenase (G6PD) polymorphism in Taiwan. Blood
79: 2135-2140,
[92270] 1992.
[92271]
[92272] 322. Testa, U.; Meloni, T.; Lania, A.; Battistuzzi, G.; Cuttillo, S.;
[92273] Luzzatto, L.: Genetic heterogeneity of glucose-
6-phosphate dehydrogenase
[92274] deficiency in Sardinia. Hum. Genet. 56: 99-105, 1980.
[92275]
[92276] 323. Thigpen, J. T.; Steinberg, M. H.; Beutler, E.; Gillespie,

G.

[92277] T., Jr.; Dreiling, B. J.; Morrison, F. S.: Glucose-6-phosphate dehydrogenase

[92278] Jackson, a new variant associated with hemolytic anemia.
Acta Haemat. 51:

[92279] 310-314, 1974.

[92280]

[92281] 324. Tishkoff, S. A.; Varkonyi, R.; Cahinhinan, N.; Abbes, S.; Argyropoulos,

[92282] G.; Destro-Bisol, G.; Drousiotou, A.; Dangerfield, B.; Lefranc, G.;

[92283] Loiselet, J.; Piro, A.; Stoneking, M.; Tagarelli, A.; Tagarelli, G.;

[92284] Touma, E. H.; Williams, S. M.; Clark, A. G.: Haplotype diversity

[92285] and linkage disequilibrium at human G6PD: recent origin of alleles

[92286] that confer malarial resistance. Science 293: 455-462, 2001.

[92287]

[92288] 325. Tokarev, Y. N.; Chernyak, N. B.; Batischev, A. I.; Lanzina, N.

[92289] V.; Alexeyev, G. A.: Etude des proprietes electrophoretiques et cinetiques

[92290] de la glucose-6-phosphate deshydrogenase (Gd) d'erythrocytes dans

[92291] les deficits hereditaires de l'enzyme: description d'une nouvelle

[92292] variante de glucose-6-phosphate deshydrogenase: la Gd Kremenchug. Nouv.

[92293] Rev. Franc. Hemat. 20: 557-564, 1978.

[92294]

[92295] 326. Toncheva, D.: Variants of glucose-6-phosphate dehydrogenase

[92296] in a Vietnamese population. Hum. Hered. 36: 348-351, 1986.

[92297]

[92298] 327. Toncheva, D.; Tzoneva, M.: Genetic polymorphism of G6PD in a

[92299] Bulgarian population. Hum. Genet. 67: 340-342, 1984.

[92300]

[92301] 328. Toniolo, D.; Martini, G.; Migeon, B. R.; Dono, R.: Expression

[92302] of the G6PD locus on the human X chromosome is associated with demethylation

[92303] of three CpG islands within 100 kb of DNA. EMBO J. 7: 401–406, 1988.

[92304]

[92305] 329. Town, M.; Athanasiou–Metaxa, M.; Luzzatto, L.: In–tragenic interspecific

[92306] complementation of glucose 6–phosphate dehydrogenase in human–hamster

[92307] cell hybrids. Somat. Cell Molec. Genet. 16: 97–108, 1990.

[92308]

[92309] 330. Town, M.; Bautista, J. M.; Mason, P. J.; Luzzatto, L.: Both

[92310] mutations in G6PD A– are necessary to produce the G6PD deficient phenotype. Hum.

[92311] Molec. Genet. 1: 171–174, 1992.

[92312]

[92313] 331. Usanga, E. A.; Bienzle, U.; Cancedda, K.; Fasuan, F. A.; Ajayi,

[92314] O.; Luzzatto, L.: Genetic variants of human erythrocyte glucose 6–phosphate

[92315] dehydrogenase: new variants in West Africa characterized by column

[92316] chromatography. Ann. Hum. Genet. 40: 279–286, 1977.

[92317]

[92318] 332. Vaca, G.; Ibarra, B.; Garcia Cruz, D.; Medina, C.; Romero, F.;

[92319] Cantu, J. M.; Beutler, E.: G-6-PD Jalisco and G-6-PD Morelia: two

[92320] new Mexican variants. Hum. Genet. 71: 82–85, 1985.

[92321]

[92322] 333. Vaca, G.; Ibarra, B.; Romero, F.; Olivares, N.; Cantu, J. M.;

[92323] Beutler, E.: G-6-PD Guadalajara: a new mutant associated with chronic

[92324] nonspherocytic hemolytic anemia. Hum. Genet. 61: 175–176, 1982.

[92325]

[92326] 334. Ventura, A.; Panizon, F.; Soranzo, M. R.; Veneziano, G.; Sansone,

[92327] G.; Testa, U.; Luzzatto, L.: Congenital dyserythropoietic anaemia

[92328] type II associated with a new type of G6PD deficiency (G6PD Gabrovizza). Acta

[92329] Haemat. 71: 227-234, 1984.

[92330]

[92331] 335. Vergnes, H.; Gherardi, M.; Bouloux, C.: Erythrocyte glucose-6-phosphate

[92332] dehydrogenase in the Niokolonko (Malinke of the Niokolo) of the Eastern

[92333] Senegal: identification of a slow variant with normal activity (Tacoma-like). Hum.

[92334] Hered. 25: 80-87, 1975.

[92335]

[92336] 336. Vergnes, H.; Gherardi, M.; Quilici, J. C.; Yoshida, A.; Giacardy,

[92337] R.: G6PD Luz-Saint-Sauveur: a new variant with abnormal electrophoretic

[92338] mobility, mild enzyme deficiency and absence of hemolytic disorders. I.R.C.S. 7:

[92339] 14, 1973.

[92340]

[92341] 337. Vergnes, H.; Gherardi, M.; Yoshida, A.: G6PD Lozere

and Trinacria-like:

[92342] segregation of two non-hemolytic variants in a French family. Hum.

[92343] Genet. 34: 293-298, 1976.

[92344]

[92345] 338. Vergnes, H.; Ribet, A.; Bommelaer, G.; Amadiou, J.; Brun, H.

[92346] : GD(-) Muret and GD(-) Colomiers, two new variants of glucose-6-phosphate

[92347] dehydrogenase associated with favism. Hum. Genet. 57: 332-334, 1981.

[92348]

[92349] 339. Vergnes, H.; Yoshida, A.; Gourdin, D.; Gherardi, M.; Bierme,

[92350] R.; Ruffie, J.: Glucose 6-phosphate dehydrogenase Toulouse: a new

[92351] variant with marked instability and severe deficiency discovered in

[92352] a family of Mediterranean ancestry. Acta Haemat. 51: 240-249, 1974.

[92353]

[92354] 340. Viglietto, G.; Montanaro, V.; Calabro, V.; Vallone, D.;

D'Urso,

[92355] M.; Persico, M. G.; Battistuzzi, G.: Common glucose-6-phosphate dehydrogenase

[92356] (G6PD) variants from the Italian population: biochemical and molecular

[92357] characterization. Ann. Hum. Genet. 54: 1-15, 1990.

[92358]

[92359] 341. Vives-Corróns, J.-L.; Kuhl, W.; Pujades, M. A.; Beutler, E.:

[92360] Molecular genetics of the glucose-6-phosphate dehydrogenase (G6PD)

[92361] Mediterranean variant and description of a new G6PD mutant, G6PD Andalus(1361A). Am.

[92362] J. Hum. Genet. 47: 575-579, 1990.

[92363]

[92364] 342. Vives-Corróns, J. L.; Feliu, E.; Pujades, M. A.; Cardellach,

[92365] F.; Rozman, C.; Carreras, A.; Jou, J. M.; Vallespi, M. T.; Zuazu,

[92366] F. J.: Severe glucose-6-phosphate dehydrogenase (G6PD) deficiency

[92367] associated with chronic hemolytic anemia, granulocyte

dysfunction,

[92368] and increased susceptibility to infections: description of a new molecular

[92369] variant (G6PD Barcelona). Blood 59: 428-434, 1982.

[92370]

[92371] 343. Vives-Corrons, J. L.; Pujades, A.: Heterogeneity of 'Mediterranean

[92372] type' glucose-6-phosphate dehydrogenase (G6PD) deficiency in Spain

[92373] and description of two new variants associated with favism. Hum.

[92374] Genet. 60: 216-221, 1982.

[92375]

[92376] 344. Vives-Corrons, J. L.; Pujades, A.; Curia, M. D.: Caracterizacion

[92377] molecular de la glucosa-6-fosfato deshidrogenasa (G6PD) en 24 casos

[92378] de deficit enzimatico y descripcion de una nueva variante (G6PD-Betica). Sangre 25:

[92379] 1049-1064, 1980.

[92380]

[92381] 345. Vives-Corróns, J. L.; Pujades, M. A.; Petit, J.; Colomer, D.;

[92382] Corbella, M.; Aguilar i Bascompte, J. L.; Merino, A.:
Chronic nonspherocytic

[92383] hemolytic anemia (CNSHA) and glucose 6 phosphate de-
hydrogenase (G6PD)

[92384] deficiency in a patient with familial amyloidotic polyneu-
ropathy (FAP):

[92385] molecular study of a new variant (G6PD Clinic) with
markedly acidic

[92386] pH optimum. Hum. Genet. 81: 161-164, 1989.

[92387]

[92388] 346. Vulliamy, T.; Beutler, E.; Luzzatto, L.: Variants of glu-
cose-6-phosphate

[92389] dehydrogenase are due to missense mutations spread
throughout the

[92390] coding region of the gene. Hum. Mutat. 2: 159-167,
1993.

[92391]

[92392] 347. Vulliamy, T.; Mason, P.; Luzzatto, L.: The molecular
basis of

[92393] glucose-6-phosphate dehydrogenase deficiency. Trends

Genet. 8: 138–143,

[92394] 1992.

[92395]

[92396] 348. Vulliamy, T.; Rovira, A.; Yusoff, N.; Colomer, D.; Luzzatto,

[92397] L.; Vives-Corrons, J.-L.: Independent origin of single and double

[92398] mutations in the human glucose 6-phosphate dehydrogenase gene. Hum.

[92399] Mutat. 8: 311–318, 1996.

[92400]

[92401] 349. Vulliamy, T. J.: Personal Communication. London, England

[92402] 1989.

[92403]

[92404] 350. Vulliamy, T. J.; D'Urso, M.; Battistuzzi, G.; Estrada, M.; Foulkes,

[92405] N. S.; Martini, G.; Calabro, V.; Poggi, V.; Giordano, R.; Town, M.;

[92406] Luzzatto, L.; Persico, M. G.: Diverse point mutations in the human

- [92407] glucose-6-phosphate dehydrogenase gene cause enzyme deficiency and
- [92408] mild or severe hemolytic anemia. Proc. Nat. Acad. Sci. 85: 5171-5175,
- [92409] 1988.
- [92410]
- [92411] 351. Vulliamy, T. J.; Kaeda, J. S.; Ait-Chafa, D.; Mangerini, R.;
- [92412] Roper, D.; Barbot, J.; Mehta, A. B.; Athanassiou-Metaxa, M.; Luzzatto,
- [92413] L.; Mason, P. J.: Clinical and haematological consequences of recurrent
- [92414] G6PD mutations and a single new mutation causing chronic nonspherocytic
- [92415] haemolytic anaemia. Brit. J. Haemat. 101: 670-675, 1998.
- [92416]
- [92417] 352. Vulliamy, T. J.; Othman, A.; Town, M.; Nathwani, A.; Falusi,
- [92418] A. G.; Mason, P. J.; Luzzatto, L.: Polymorphic sites in the African
- [92419] population detected by sequence analysis of the glucose-6-phosphate

[92420] dehydrogenase gene outline the evolution of the variants A and A-. Proc.

[92421] Nat. Acad. Sci. 88: 8568-8571, 1991.

[92422]

[92423] 353. Vulliamy, T. J.; Wanachiwanawin, W.; Mason, P. J.; Luzzatto,

[92424] L.: G6PD Mahidol, a common deficient variant in South East Asia is

[92425] caused by a (163)glycine-to-serine mutation. Nucleic Acids Res. 17:

[92426] 5868, 1989.

[92427]

[92428] 354. Vuopio, P.; Harkonen, M.; Helske, T.; Naeveri, H.: Red cell

[92429] glucose-6-phosphate dehydrogenase deficiency in Finland: characterization

[92430] of a new variant with severe enzyme deficiency. Scand. J. Haemat. 15:

[92431] 145-152, 1975.

[92432]

[92433] 355. Vuopio, P.; Harkonen, M.; Johnsson, P.; Nuutinen, M.:

Red cell

[92434] glucose-phosphate dehydrogenase deficiency in Finland.
Ann. Clin.

[92435] Res. 5: 168-178, 1973.

[92436]

[92437] 356. Waitz, R.; Boivin, P.; Oberling, F.; Casenave, J. P.;
North,

[92438] M. L.; Mayer, S.: Variante Gd(-) Strasbourg de la glucose-
6-phosphate

[92439] dehydrogenase. Nouv. Rev. Franc. Hemat. 10: 312-314,
1970.

[92440]

[92441] 357. Wang, Y. M.; Patterson, J. H.; Van Eys, J.: The poten-
tial use

[92442] of xylitol in glucose-6-phosphate dehydrogenase defi-
ciency anemia. J.

[92443] Clin. Invest. 50: 1421-1428, 1971.

[92444]

[92445] 358. Weimer, T. A.; Salzano, F. M.; Hutz, M. H.: Erythro-
cyte isozymes

[92446] and hemoglobin types in a southern Brazilian population.

J. Hum.

[92447] *Evol.* 10: 319–328, 1981.

[92448]

[92449] 359. Weimer, T. A.; Schuler, L.; Beutler, E.; Salzano, F. M.:
Gd(+)

[92450] Laguna, a new rare glucose-6-phosphate dehydrogenase
variant from

[92451] Brazil. *Hum. Genet.* 65: 402–404, 1984.

[92452]

[92453] 360. Weinreich, J.; Busch, D.; Gottstein, U.; Schaefer, J.;
Rohr,

[92454] J.: Ueber zwei neue Faelle von hereditaerer nichtsphaero-
cytaerer

[92455] haemolytischer Anaemie bei glucose-
6-phosphat-dehydrogenase-Defekt

[92456] in einer Nord Deutschen Familie. *Klin. Wschr.* 46:
146–149, 1968.

[92457]

[92458] 361. Welch, S. G.; McGregor, I. A.; Williams, K.: A new
variant of

[92459] human erythrocyte G6PD occurring at a high frequency

amongst the population

[92460] of two villages in The Gambia, West Africa. Hum. Genet.
40: 305–309,

[92461] 1978.

[92462]

[92463] 362. Westring, D. W.; Pisciotto, A. V.: Anemia, cataracts,
and seizures

[92464] in patients with glucose–6–phosphate dehydrogenase de-
ficiency. Arch.

[92465] Intern. Med. 118: 385–390, 1966.

[92466]

[92467] 363. WHO: Nomenclature of glucose–6–phosphate dehy-
drogenase in

[92468] man. Bull. WHO 36: 319–322, 1967. Note: See Also:
Canad. Med. Assoc.

[92469] J. 97: 422–424, 1967.

[92470]

[92471] 364. WHO: Scientific group on the standardization of pro-
cedures

[92472] for the study of glucose–6–phosphate dehydrogenase.
WHO Techn. Rep.

[92473] (pub.) Ser. No. 366: 1967.

[92474]

[92475] 365. Wilson, W. W.: Congenital hemolytic anemia due to a deficiency

[92476] of glucose 6-phosphate dehydrogenase. Rocky Mt. Med. J. 73: 160-162,

[92477] 1976.

[92478]

[92479] 366. Witt, I.; Yoshioka, S.: Biochemical characterization of a glucose-6-phosphate

[92480] dehydrogenase variant with favism: G-6-PD Zaehringen. Klin. Wschr. 50:

[92481] 205-209, 1972.

[92482]

[92483] 367. Wong, P. W. K.; Shih, L.-Y.; Hsia, D. Y. Y.: Characterization

[92484] of glucose-6-phosphate dehydrogenase among Chinese. Nature 208:

[92485] 1323-1324, 1965.

[92486]

- [92487] 368. Yermakov, N.; Tokarev, J.; Chernjak, N.; Schoenian, G.; Grieger, M.; Guckler, G.; Jacobasch, G.; Mahmudova, M.; Bahramov, S.: New stable mutant Gd(-) variants: G6PD Tashkent and G6PD Nucus: molecular basis of hereditary enzyme deficiency. Acta Biol. Med. Ger. 40: 559–562, 1981.
- [92492]
- [92493] 369. Yoshida, A.: Personal Communication. Duarte, Calif. 2/26/1996.
- [92494]
- [92495] 370. Yoshida, A.: Amino acid substitution (histidine to tyrosine) in a glucose-6-phosphatase dehydrogenase variant (G6PD Hektoen) associated with over-production. J. Mol. Biol. 52: 483–490, 1970.
- [92498]
- [92499] 371. Yoshida, A.: A single amino acid substitution (asparagine to

- [92500] aspartic acid) between normal (B plus) and the common Negro variant
- [92501] (A plus) of human glucose-6-phosphate dehydrogenase. Proc. Nat. Acad.
- [92502] Sci. 57: 835-840, 1967.
- [92503]
- [92504] 372. Yoshida, A.: Human glucose-6-phosphate dehydrogenase: purification
- [92505] and characterization of Negro type variant (A+) and comparison with
- [92506] normal enzyme (B+). Biochem. Genet. 1: 81-99, 1967.
- [92507]
- [92508] 373. Yoshida, A.; Baur, E. W.; Motulsky, A. G.: A Philippino glucose-6-phosphate
- [92509] dehydrogenase variant (G6PD Union) with enzyme deficiency and altered
- [92510] substrate specificity. Blood 35: 506-513, 1970.
- [92511]
- [92512] 374. Yoshida, A.; Beutler, E.: Human glucose-6-phosphate dehydrogenase
- [92513] variants: a supplementary tabulation. Ann. Hum. Genet.

41: 347-355,

[92514] 1978.

[92515]

[92516] 375. Yoshida, A.; Beutler, E.; Motulsky, A. G.: Table of human glucose-6-phosphate

[92517] dehydrogenase variants. Bull. WHO 45: 243-253, 1971.

[92518]

[92519] 376. Yoshida, A.; Stamatoyannopoulos, G.; Motulsky, A. G.: Negro

[92520] variant of glucose-6-phosphate dehydrogenase deficiency (A-) in man. Science 155:

[92521] 97-99, 1967.

[92522]

[92523] 377. Yoshida, A.; Takizawa, T.: The same extra FokI cleavage site

[92524] exists in glucose-6-phosphate dehydrogenase variants A(+) and A(-). Am.

[92525] J. Hum. Genet. 43: 131-133, 1988.

[92526]

[92527] 378. Yoshida, A.; Takizawa, T.; Prchal, J. T.: RFLP of the X

chromosome-linked

[92528] glucose-6-phosphate dehydrogenase locus in blacks. Am. J. Hum. Genet. 42:

[92529] 872-876, 1988.

[92530]

[92531] 379. Zinkham, W. H.: A deficiency of glucose-6-phosphate dehydrogenase

[92532] activity in lens from individuals with primaquine-sensitive erythrocytes. Bull.

[92533] Johns Hopkins Hosp. 109: 206-216, 1961.

[92534]

[92535] 380. Zuo, L.; Chen, E.; Du, C. S.; Chang, C. N.; Chiu, D. T. Y.:

[92536] Genetic study of Chinese G6PD variants by direct PCR sequencing. (Abstract) Blood 76

[92537] (suppl. 1): 51A, 1990.

[92538] 16161.

[92539] Antignac, C.; Gros, F.; Geffriaud, C.; Kashtan, C.; Proesmans,

[92540] W.; Tryggvason, K.; Gubler, M. C.: Alport syndrome (AS) and diffuse

- [92541] esophageal leiomyomatosis (DL): deletions in the 5-prime end and upstream
- [92542] region of the COL4A5 collagen gene. (Abstract) Am. J. Hum. Genet. 51
- [92543] (suppl.): A7, 1992.
- [92544] 16162.
- [92545] Antignac, C.; Knebelmann, B.; Drouot, L.; Gros, F.; Deschenes,
- [92546] G.; Hors-Cayla, M.-C.; Zhou, J.; Tryggvason, K.; Grunfeld, J.-P.;
- [92547] Broyer, M.; Gubler, M.-C.: Deletions in the COL4A5 collagen gene
- [92548] in X-linked Alport syndrome: characterization of the pathological
- [92549] transcripts in nonrenal cells and correlation with disease expression. J.
- [92550] Clin. Invest. 93: 1195-1207, 1994.
- [92551] 16163.
- [92552] Antignac, C.; Zhou, J.; Sanak, M.; Cochat, P.; Roussel, B.; Deschenes,
- [92553] G.; Gros, F.; Knebelmann, B.; Hors-Cayla, M.-C.; Tryggvason, K.; Gubler,

[92554] M.-C.: Alport syndrome and diffuse leiomyomatosis: deletions in the

[92555] 5-prime end of the COL4A5 collagen gene. *Kidney Int.* 42: 1178-1183,

[92556] 1992.

[92557] 16164.

[92558] Barker, D. F.; Denison, J. C.; Atkin, C. L.; Gregory, M. C.: Common

[92559] ancestry of three Ashkenazi-American families with Alport syndrome

[92560] and COL4A5 R1677Q. *Hum. Genet.* 99: 681-684, 1997.

[92561] 16165.

[92562] Barker, D. F.; Pruchno, C. J.; Jiang, X.; Atkin, C. L.; Stone,

[92563] E. M.; Denison, J. C.; Fain, P. R.; Gregory, M. C.: A mutation causing

[92564] Alport syndrome with tardive hearing loss is common in the western

[92565] United States. *Am. J. Hum. Genet.* 58: 1157-1165, 1996.

[92566] 16166.

[92567] Boye, E.; Vetrie, D.; Flinter, F.; Buckle, B.; Pihlajaniemi, T.;

- [92568] Hamalainen, E.-R.; Myers, J. C.; Bobrow, M.; Harris, A.:
Major rearrangements
- [92569] in the alpha-5-(IV) collagen gene in three patients with
Alport syndrome. Genomics 11:
- [92570] 1125-1132, 1991.
- [92571] 16167.
- [92572] Guo, C.; Van Damme, B.; Vanrenterghem, Y.; Devriendt,
K.; Cassiman,
- [92573] J.-J.; Marynen, P.: Severe Alport phenotype in a woman
with two missense
- [92574] mutations in the same COL4A5 gene and preponderant
inactivation of
- [92575] the X chromosome carrying the normal allele. J. Clin. In-
vest. 95:
- [92576] 1832-1837, 1995.
- [92577] 16168.
- [92578] Campuzano, V.; Montermini, L.; Molto, M. D.; Pianese, L.;
Cossee,
- [92579] M.; Cavalcanti, F.; Monros, E.; Rodius, F.; Duclos, F.; Mon-
ticelli,
- [92580] A.; Zara, F.; Canizares, J.; Koutnikova, H.; Bidichandani, S.
I.;

- [92581] Gellera, C.; Brice, A.; Trouillas, P.; De Michele, G.; Filla, A.;
- [92582] De Frutos, R.; Palau, F.; Patel, P. I.; Di Donato, S.; Mandel, J.
- [92583] –L.; Coccozza, S.; Koenig, M.; Pandolfo, M.: Friedreich's ataxia:
- [92584] autosomal recessive disease caused by an intronic GAA triplet repeat
- [92585] expansion. *Science* 271: 1423–1427, 1996.
- [92586] 16169.
- [92587] Boyd, M.; Lanyon, W. G.; Connor, J. M.: Screening for molecular
- [92588] pathologies in Lesch–Nyhan syndrome. *Hum. Mutat.* 2: 127–130, 1993.
- [92589] 16170.
- [92590] Brennand, J.; Chinault, A. C.; Konecki, D. S.; Melton, D. W.; Caskey,
- [92591] C. T.: Cloned cDNA sequences of the hypoxanthine–guanine phosphoribosyltransferase
- [92592] gene from a mouse neuroblastoma cell line found to have amplified
- [92593] genomic sequences. *Proc. Nat. Acad. Sci.* 79: 1950–1954, 1982.

[92594] 16171.

[92595] Cariello, N. F.; Scott, J. K.; Kat, A. G.; Thilly, W. G.; Keo-
havong,

[92596] P.: Resolution of a missense mutant in human genomic
DNA by denaturing

[92597] gradient gel electrophoresis and direct sequencing using
in vitro

[92598] DNA amplification: HPRT(Munich). Am. J. Hum. Genet. 42:
726-734,

[92599] 1988.

[92600] 16172.

[92601] Caskey, C. T.; Kruh, G. D.: The HPRT locus: review. Cell
16:

[92602] 1-9, 1979.

[92603] 16173.

[92604] Chinault, A. C.; Caskey, C. T.: The hypoxanthine phospho-
ribosyltransferase

[92605] gene: a model for the study of mutation in mammalian
cells. Prog.

[92606] Nucleic Acids Res. Molec. Biol. 31: 317-465, 1984.

[92607] 16174.

- [92608] Cianfriglia, M.; Miggiano, V. C.; Meo, T.; Muller, H. J.;
Muller,
- [92609] E.; Battistuzzi, G.: Evidence for syntenic between the rabbit
gene
- [92610] loci coding for HPRT, PGK and G6PD in mouse-rabbit so-
matic cell hybrids.
- [92611] (Abstract) Cytogenet. Cell Genet. 25: 142 only, 1979.
- [92612] 16175.
- [92613] Colgin, L. M.; Hackmann, A. F. M.; Emond, M. J.; Monnat,
R. J.,
- [92614] Jr.: The unexpected landscape of in vivo somatic mutation
in a human
- [92615] epithelial cell lineage. Proc. Nat. Acad. Sci. 99:
1437-1442, 2002.
- [92616] 16176.
- [92617] Cox, R. P.; Krauss, M. R.; Balis, M. E.; Dancis, J.: Evidence
- [92618] for transfer of enzyme product as the basis of metabolic
cooperation
- [92619] between tissue culture fibroblasts of Lesch-Nyhan disease
and normal
- [92620] cells. Proc. Nat. Acad. Sci. 67: 1573-1579, 1970.

[92621] 16177.

[92622] Dancis, J.; Yip, L. C.; Cox, R. P.; Piomelli, S.; Balis, M. E.

[92623] : Disparate enzyme activity in erythrocytes and leukocytes:
a variant

[92624] of hypoxanthine phosphoribosyltransferase deficiency
with an unstable

[92625] enzyme. J. Clin. Invest. 52: 2068–2074, 1973.

[92626] 16178.

[92627] Davidson, B. L.: Personal Communication. Ann Arbor,
Mich.

[92628] 1990.

[92629] 16179.

[92630] Davidson, B. L.; Chin, S.-J.; Wilson, J. M.; Kelley, W. N.;
Palella,

[92631] T. D.: Hypoxanthine–guanine phosphoribosyltransferase:
genetic evidence

[92632] for identical mutations in two partially deficient subjects.
J. Clin.

[92633] Invest. 82: 2164–2167, 1988.

[92634] 16180.

[92635] Davidson, B. L.; Pashmforoush, M.; Kelley, W. N.; Palella, T.

[92636] D.: Genetic basis of hypoxanthine–guanine phosphoribosyltransferase

[92637] deficiency in a patient with the Lesch–Nyhan syndrome (HPRT–Flint). Gene 63:

[92638] 331–336, 1988.

[92639] 16181.

[92640] Davidson, B. L.; Pashmforoush, M.; Kelley, W. N.; Palella, T.

[92641] D.: Human hypoxanthine–guanine phosphoribosyltransferase deficiency:

[92642] the molecular defect in a patient with gout (HPRT–Ashville). J. Biol.

[92643] Chem. 264: 520–525, 1989.

[92644] 16182.

[92645] Davidson, B. L.; Tarle, S. A.; Palella, T. D.; Kelley, W. N.:

[92646] Molecular basis of hypoxanthine–guanine phosphoribosyltransferase

[92647] deficiency in 10 subjects determined by direct sequencing of amplified

[92648] transcripts. J. Clin. Invest. 84: 342–346, 1989.

- [92649] 16183.
- [92650] Davidson, B. L.; Tarle, S. A.; Van Antwerp, M.; Gibbs, D. A.;
- [92651] Watts, R. W. E.; Kelley, W. N.; Palella, T. D.: Identification
of
- [92652] 17 independent mutations responsible for human hypox-
anthine-guanine
- [92653] phosphoribosyltransferase (HPRT) deficiency. Am. J. Hum.
Genet. 48:
- [92654] 951-958, 1991.
- [92655] 16184.
- [92656] De Gregorio, L.; Nyhan, W. L.; Serafin, E.; Chamoles, N. A.:
- [92657] An unexpected affected female patient in a classical
Lesch-Nyhan family. Molec.
- [92658] Genet. Metab. 69: 263-268, 2000.
- [92659] 16185.
- [92660] Demars, R. I.; Sarto, G. E.; Felix, J. S.; Benke, P.: Lesch-
Nyhan
- [92661] mutation: prenatal detection with amniotic fluid cells. Sci-
ence 164:
- [92662] 1303-1305, 1969.

[92663] 16186.

[92664] Dempsey, J. L.; Morley, A. A.; Seshadri, R. S.; Emmerson, B. T.;

[92665] Gordon, R.; Bhagat, C. I.: Detection of the carrier state for an

[92666] X-linked disorder, the Lesch-Nyhan syndrome, by the use of lymphocyte

[92667] cloning. Hum. Genet. 64: 288-290, 1983.

[92668] 16187.

[92669] Dobrovic, A.; Gareau, P.; Seifert, A.-M.; Messing, K.; Bradley,

[92670] W. E. C.: A HindIII RFLP for the HPRT pseudogene on chromosome 3

[92671] (HPRT1). Nucleic Acids Res. 15: 1346, 1987.

[92672] 16188.

[92673] Doetschman, T.; Gregg, R. G.; Maeda, N.; Hooper, M. L.; Melton,

[92674] D. W.; Thompson, S.; Smithies, O.: Targetted correction of a mutant

[92675] HPRT gene in mouse embryonic stem cells. Nature 330: 576-578, 1987.

[92676] 16189.

[92677] Echard, G.; Gillois, M.: G6PD-PGK-GAL-HPRT syntenly in the rabbit,

[92678] *Oryctolagus cuniculus*. (Abstract) Cytogenet. Cell Genet. 25: 148-149,

[92679] 1979.

[92680] 16190.

[92681] Edwards, A.; Caskey, C. T.: Personal Communication. Houston,

[92682] Tex. 8/1990.

[92683] 16191.

[92684] Edwards, A.; Voss, H.; Rice, P.; Civitello, A.; Stegemann, J.;

[92685] Schwager, C.; Zimmermann, J.; Erfle, H.; Caskey, C. T.; An-sorge, W.

[92686] : Automated DNA sequencing of the human HPRT locus. Genomics 6:

[92687] 593-608, 1990.

[92688] 16192.

[92689] Emmerson, B. T.; Thompson, C. J.; Wallace, D. C.: Partial deficiency

- [92690] hypoxanthine-guanine phosphoribosyltransferase: intermediate enzyme
- [92691] deficiency in heterozygote red cells. *Ann. Intern. Med.* 76: 285-288,
- [92692] 1972.
- [92693] 16193.
- [92694] Engle, S. J.; Womer, D. E.; Davies, P. M.; Boivin, G.; Sahota, A.; Simmonds, H. A.; Stambrook, P. J.; Tischfield, J. A.: HPRT-APRT-deficient
- [92696] mice are not a model for Lesch-Nyhan syndrome. *Hum. Mol. Genet.* 5:
- [92697] 1607-1610, 1996.
- [92698] 16194.
- [92699] Epstein, C. J.: Expression of the mammalian X chromosome before
- [92700] and after fertilization. *Science* 175: 1467-1468, 1972.
- [92701] 16195.
- [92702] Ernst, M.; Zametkin, A. J.; Matochik, J. A.; Pascualvaca, D.;
- [92703] Jons, P. H.; Hardy, K.; Hankerson, J. G.; Doudet, D. J.; Cohen, R.

- [92704] M.: Presynaptic dopaminergic deficits in Lesch–Nyhan disease. New
- [92705] Eng. J. Med. 334: 1568–1572, 1996.
- [92706] 16196.
- [92707] Fenwick, R. G., Jr.: Reversion of a mutation affecting the molecular
- [92708] weight of HGPRT: intragenic suppression and localization of X-linked
- [92709] genes. Somat. Cell Genet. 6: 477–494, 1980.
- [92710] 16197.
- [92711] Fox, I. H.; Dwosh, I. L.; Marchant, P. J.; Lacroix, S.; Moore,
- [92712] M. R.; Omura, S.; Wyhofsky, V.: Hypoxanthine–guanine phosphoribosyltransferase:
- [92713] characterization of a mutant in a patient with gout. J. Clin. Invest. 56:
- [92714] 1239–1249, 1975.
- [92715] 16198.
- [92716] Francke, U.; Bakay, B.; Connor, J. D.; Coldwell, J. G.; Nyhan,
- [92717] W. L.: Linkage relationships of X-linked enzymes glucose–6-phosphate

- [92718] dehydrogenase and hypoxanthine guanine phosphoribosyltransferase. Am.
- [92719] J. Hum. Genet. 26: 512–522, 1974.
- [92720] 16199.
- [92721] Francke, U.; Felsenstein, J.; Gartler, S. M.; Migeon, B. R.; Dancis,
- [92722] J.; Seegmiller, J. E.; Bakay, B.; Nyhan, W. L.: The occurrence of
- [92723] new mutants in the X-linked recessive Lesch–Nyhan disease. Am. J.
- [92724] Hum. Genet. 28: 123–137, 1976.
- [92725] 16200.
- [92726] Francke, U.; Felsenstein, J.; Gartler, S. M.; Nyhan, W. L.; Seegmiller,
- [92727] J. E.: Answer to criticism of Morton and Lalouel. (Letter) Am. J.
- [92728] Hum. Genet. 29: 307–310, 1977.
- [92729] 16201.
- [92730] Gartler, S. M.; Francke, U.: Half–chromatid mutation: transmission
- [92731] in humans? Am. J. Hum. Genet. 27: 218–223, 1975.

- [92732] 16202.
- [92733] Gibbs, R. A.; McFadyen, I. R.; Crawford, M. d'A.; de Muinck Keizer, E. E.; Headhouse-Benson, C. M.; Wilson, T. M.; Farrant, P. H.: First-trimester diagnosis of Lesch-Nyhan syndrome. *Lancet* II: 1180-1183, 1984.
- [92736] 16203.
- [92737] Graham, G. W.; Aitken, D. A.; Connor, J. M.: Prenatal diagnosis by enzyme analysis in 15 pregnancies at risk for the Lesch-Nyhan syndrome. *Prenatal Diag.* 16: 647-651, 1996.
- [92740] 16204.
- [92741] Greene, M. L.; Nyhan, W. L.; Seegmiller, J. E.: Hypoxanthine-guanine phosphoribosyltransferase deficiency and Xg blood group. *Am. J. Hum. Genet.* 22: 50-54, 1970.
- [92744] 16205.

- [92745] Henderson, J. F.; Kelley, W. N.; Rosenbloom, F. M.; Seeg-
miller,
[92746] J. E.: Inheritance of purine phosphoribosyltransferases in
man. Am.
[92747] J. Hum. Genet. 21: 61-70, 1969.
[92748] 16206.
[92749] Hoefnagel, D.; Andrew, E. D.; Mireault, N. G.; Berndt, W.
O.:
[92750] Hereditary choreoathetosis, self-mutilation and hyper-
uricemia in young
[92751] males. New Eng. J. Med. 273: 130-135, 1965.
[92752] 16207.
[92753] Lloyd, K. G.; Hornykiewicz, O.; Davidson, L.; Shannak, K.;
Farley,
[92754] I.; Goldstein, M.; Shibuya, M.; Kelley, W. N.; Fox, I. H.: Bio-
chemical
[92755] evidence of dysfunction of brain neurotransmitters in the
Lesch-Nyhan
[92756] syndrome. New Eng. J. Med. 305: 1106-1111, 1981.
[92757] 16208.
[92758] Migeon, B. R.; Der Kaloustian, V. M.; Nyhan, W. L.; Young,

W.

- [92759] J.; Childs, B.: X-linked hypoxanthine-guanine phosphoribosyl transferase deficiency: heterozygote has two clonal populations. *Science* 160: 425-427, 1968.
- [92761] 16209.
- [92762] 16209.
- [92763] Morton, N. E.; Lalouel, J. M.: Genetic epidemiology of Lesch-Nyhan disease. (Letter) *Am. J. Hum. Genet.* 29: 304-307, 1977.
- [92764] 16210.
- [92765] 16210.
- [92766] Nabholz, M.; Miggiano, V.; Bodmer, W.: Genetic analysis with human-mouse somatic cell hybrids. *Nature* 223: 358-363, 1969.
- [92767] 16211.
- [92768] 16211.
- [92769] Nyhan, W. L.; Bakay, B.; Connor, J. D.; Marks, J. F.; Keele, D. K.: Hemizygous expression of glucose-6-phosphate dehydrogenase in erythrocytes of heterozygotes for the Lesch-Nyhan syn-

drome. Proc.

[92772] Nat. Acad. Sci. 65: 214–218, 1970.

[92773] 16212.

[92774] Nyhan, W. L.; Olivier, W. J.; Lesch, M.: A familial disorder

[92775] of uric acid metabolism and central nervous system function. J. Pediat. 67:

[92776] 257–263, 1965.

[92777] 16213.

[92778] Nyhan, W. L.; Wong, D. F.: New approaches to understanding Lesch–Nyhan

[92779] disease. (Editorial) New Eng. J. Med. 334: 1602–1604, 1996.

[92780] 16214.

[92781] Burwinkel, B.; Amat, L.; Gray, R. G. F.; Matsuo, N.; Muroya, K.;

[92782] Narisawa, K.; Sokol, R. J.; Vilaseca, M. A.; Kilimann, M. W.: Variability

[92783] of biochemical and clinical phenotype in X-linked liver glycogenosis

[92784] with mutations in the phosphorylase kinase PHKA2 gene. Hum. Genet. 102:

[92785] 423–429, 1998.

[92786] 16215.

[92787] Burwinkel, B.; Shin, Y. S.; Bakker, H. D.; Deutsch, J.;
Lozano,

[92788] M. J.; Maire, I.; Kilimann, M. W.: Mutation hotspots in the
PHKA2

[92789] gene in X-linked liver glycogenosis due to phosphorylase
kinase deficiency

[92790] with atypical activity in blood cells (XLG2). Hum. Molec.
Genet. 5:

[92791] 653–658, 1996.

[92792] 16216.

[92793] Davidson, J. J.; Ozcelik, T.; Hamacher, C.; Willems, P. J.;
Francke,

[92794] U.; Kilimann, M. W.: cDNA cloning of a liver isoform of the
phosphorylase

[92795] kinase alpha subunit and mapping of the gene to
Xp22.2–p22.1, the

[92796] region of human X-linked liver glycogenosis. Proc. Nat.
Acad. Sci. 89:

[92797] 2096–2100, 1992.

- [92798] 16217.
- [92799] Davisson, M. T.: X-linked genetic homologies between mouse and
- [92800] man. *Genomics* 1: 213–227, 1987.
- [92801] 16218.
- [92802] Garibaldi, L. R.; Borrone, C.; De Martini, I.; Battistini, E.:
- [92803] Dextrothyroxine treatment of phosphorylase-kinase deficiency glycogenosis
- [92804] in four boys. *Helv. Paediat. Acta* 33: 435–444, 1978.
- [92805] 16219.
- [92806] Goji, K.; Morishita, Y.; Kodama, S.; Takahashi, T.; Matsuo, T.
- [92807] : Lymphocyte phosphorylase kinase activities in the sex-linked form
- [92808] of liver phosphorylase kinase deficiency. *Europ. J. Pediat.* 143:
- [92809] 179–182, 1985.
- [92810] 16220.
- [92811] Hendrickx, J.; Bosshard, N. U.; Willems, P.; Gitzelmann, R.: Clinical,

- [92812] biochemical and molecular findings in a patient with X-linked liver
- [92813] glycogenosis followed for 40 years. *Europ. J. Pediat.* 157: 919–923,
- [92814] 1998.
- [92815] 16221.
- [92816] Hendrickx, J.; Coucke, P.; Bossuyt, P.; Wauters, J.; Raeymaekers,
- [92817] P.; Marchau, F.; Smit, G. P. A.; Stolte, I.; Sardharwalla, I. B.;
- [92818] Berthelot, J.; Van den Bergh, I.; Berger, R.; Van Broeckhoven, C.;
- [92819] Baussan, C.; Wapenaar, M.; Fernandes, J.; Willems, P. J.: X-linked
- [92820] liver glycogenosis: localization and isolation of a candidate gene. *Hum.*
- [92821] *Molec. Genet.* 2: 583–589, 1993.
- [92822] 16222.
- [92823] Hendrickx, J.; Coucke, P.; Dams, E.; Lee, P.; Odievre, M.; Corbeel,
- [92824] L.; Fernandes, J. F.; Willems, P. J.: Mutations in the phosphorylase

- [92825] kinase gene PHKA2 are responsible for X-linked liver glycogen storage
- [92826] disease. Hum. Molec. Genet. 4: 77-83, 1995.
- [92827] 16223.
- [92828] Hendrickx, J.; Coucke, P.; Hors-Cayla, M.-C.; Smit, G. P. A.;
- [92829] Shin, Y. S.; Deutsch, J.; Smeitink, J.; Berger, R.; Lee, P.; Fernandes,
- [92830] J.; Willems, P. J.: Localization of a new type of X-linked liver
- [92831] glycogenosis to the chromosomal region Xp22 containing the liver alpha-subunit
- [92832] of phosphorylase kinase (PHKA2). Genomics 21: 620-625, 1994.
- [92833] 16224.
- [92834] Hendrickx, J.; Coucke, P.; Raeymaekers, P.; Willems, P. J.: X-linked
- [92835] liver glycogenosis: localization and isolation of a strong candidate
- [92836] gene. (Abstract) Am. J. Hum. Genet. 51 (suppl.): A190 only, 1992.
- [92837] 16225.

- [92838] Hendrickx, J.; Dams, E.; Coucke, P.; Lee, P.; Fernandes, J.; Willems,
- [92839] P. J.: X-linked liver glycogenosis type II (XLG II) is caused by
- [92840] mutations in PHKA2, the gene encoding the liver alpha subunit of phosphorylase
- [92841] kinase. Hum. Molec. Genet. 5: 649-652, 1996.
- [92842] 16226.
- [92843] Hendrickx, J.; Lee, P.; Keating, J. P.; Carton, D.; Sardharwalla,
- [92844] I. B.; Tuchman, M.; Baussan, C.; Willems, P. J.: Complete genomic
- [92845] structure and mutational spectrum of PHKA2 in patients with X-linked
- [92846] liver glycogenosis type I and II. Am. J. Hum. Genet. 64: 1541-1549,
- [92847] 1999.
- [92848] 16227.
- [92849] Hers, H.-G.; Van Hoof, F.; de Barsey, T.: Glycogen storage diseases. In:
- [92850] Scriver, C. R.; Beaudet, A. L.; Sly, W. S.; Valle, D.: The Metabolic

[92851] Basis of Inherited Disease. New York: McGraw-Hill (pub.)
(6th

[92852] ed.) I: 1989. Pp. 425-452.

[92853] 16228.

[92854] Hug, G.: Personal Communication. Cincinnati, Ohio 1974.

[92855] 16229.

[92856] Hug, G.; Schubert, W. K.; Chuck, G.: Deficient activity of
dephosphophosphorylase

[92857] kinase and accumulation of glycogen in the liver. J. Clin.
Invest. 48:

[92858] 704-715, 1969.

[92859] 16230.

[92860] Huijing, F.: Phosphorylase kinase deficiency. Biochem.
Genet. 4:

[92861] 187-194, 1970.

[92862] 16231.

[92863] Huijing, F.: Phosphorylase kinase in leucocytes of normal
subjects

[92864] and of patients with glycogen-storage disease. Biochim.
Biophys.

- [92865] Acta 148: 601–603, 1967.
- [92866] 16232.
- [92867] Huijing, F.: Glycogen–storage disease type VIa: low phosphorylase
- [92868] kinase activity caused by a low enzyme–substrate affinity. Biochim.
- [92869] Biophys. Acta 206: 199–201, 1970.
- [92870] 16233.
- [92871] Huijing, F.; Fernandez, J.: Liver glycogenosis and phosphorylase
- [92872] kinase deficiency. (Letter) Am. J. Hum. Genet. 22: 484–485, 1970.
- [92873] 16234.
- [92874] Heidet, L.; Dahan, K.; Zhou, J.; Xu, Z.; Cochat, P.; Gould, J.
- [92875] D. M.; Leppig, K. A.; Proesmans, W.; Guyot, C.; Guillot, M.; Roussel,
- [92876] B.; Tryggvason, K.; Grunfeld, J.–P.; Gubler, M.–C.; Antignac, C.:
- [92877] Deletions of both alpha–5(IV) and alpha–6(IV) collagen genes in Alport
- [92878] syndrome and in Alport syndrome associated with smooth

muscle tumours. Hum.

[92879] Molec. Genet. 4: 99–108, 1995.

[92880] 16235.

[92881] Morton, C. C.; Nussenzweig, M. C.; Sousa, R.; Sorenson, G. D.;

[92882] Pettengill, O. S.; Shows, T. B.: Mapping and characterization of

[92883] an X-linked processed gene related to MYCL1. Genomics 4: 367–375,

[92884] 1989.

[92885] 16236.

[92886] Ambler, M. W.; Neave, C.; Singer, D. B.: X-linked recessive myotubular

[92887] myopathy. II. Muscle morphology and human myogenesis. Hum. Path. 15:

[92888] 1107–1120, 1984.

[92889] 16237.

[92890] Ambler, M. W.; Neave, C.; Tutschka, B. G.; Pueschel, S. M.; Orson,

[92891] J. M.; Singer, D. B.: X-linked recessive myotubular myopathy. I.

- [92892] Clinical and pathologic findings in a family. Hum. Path.
15: 566–574,
[92893] 1984.
- [92894] 16238.
- [92895] Askanas, V.; Engel, W. K.; Reddy, N. B.; Barth, P. G.; Beth-
lem,
[92896] J.; Krauss, D. R.; Hibberd, M. E.; Lawrence, J. V.; Carter, L.
S.
- [92897] : X-linked recessive congenital muscle fiber hypotrophy
with central
[92898] nuclei: abnormalities of growth and adenylate cyclase in
muscle tissue
[92899] cultures. Arch. Neurol. 36: 604–609, 1979.
- [92900] 16239.
- [92901] Barth, P. G.; Dubowitz, V.: X-linked myotubular myopa-
thy--a long-term
[92902] follow-up study. Europ. J. Paediat. Neurol. 2: 49–56,
1998.
- [92903] 16240.
- [92904] Ilyin, G. P.; Rialland, M.; Pigeon, C.; Guguen-Guillouzo, C.:
[92905] cDNA cloning and expression analysis of new members of

the mammalian

[92906] F-box protein family. Genomics 67: 40–47, 2000.

[92907] 16241.

[92908] Chan, A. M.–L.; Miki, T.; Meyers, K. A.; Aaronson, S. A.: A human

[92909] oncogene of the RAS superfamily unmasked by expression cDNA cloning. Proc.

[92910] Nat. Acad. Sci. 91: 7558–7562, 1994.

[92911] 16242.

[92912] Drivas, G. T.; Shih, A.; Coutavas, E.; Rush, M. G.; D'Eustachio,

[92913] P.: Characterization of four novel ras-like genes expressed in a

[92914] human teratocarcinoma cell line. Molec. Cell. Biol. 10: 1793–1798,

[92915] 1990.

[92916] 16243.

[92917] Brugnera, E.; Georgiev, O.; Radtke, F.; Heuchel, R.; Baker, E.;

[92918] Sutherland, G. R.; Schaffner, W.: Cloning, chromosomal mapping and

- [92919] characterization of the human metal-regulatory transcription factor
- [92920] MTF-1. Nucleic Acids Res. 22: 3167-3173, 1994.
- [92921] 16244.
- [92922] Gunes, C.; Heuchel, R.; Georgiev, O.; Muller, K.-H.; Lichtlen,
- [92923] P.; Bluthmann, H.; Marino, S.; Aguzzi, A.; Schaffner, W.: Embryonic
- [92924] lethality and liver degeneration in mice lacking the metal-responsive
- [92925] transcriptional activator MTF-1. EMBO J. 17: 2846-2854, 1998.
- [92926] 16245.
- [92927] Radtke, F.; Heuchel, R.; Georgiev, O.; Hergersberg, M.; Gariglio,
- [92928] M.; Dembic, Z.; Schaffner, W.: Cloned transcription factor MTF-1
- [92929] activates the mouse metallothionein I promoter. EMBO J. 12: 1355-1362,
- [92930] 1993.
- [92931] 16246.

- [92932] Westin, W.; Schaffner, W.: A zinc-responsive factor interacts
- [92933] with a metal-regulated enhancer element (MRE) of the mouse metallothionein-I
- [92934] gene. EMBO J. 7: 3763-3770, 1988.
- [92935] 16247.
- [92936] Andersson, P.; McGuire, J.; Rubio, C.; Gradin, K.; Whitelaw, M.
- [92937] L.; Pettersson, S.; Hanberg, A.; Poellinger, L.: A constitutively
- [92938] active dioxin/aryl hydrocarbon receptor induces stomach tumors. Proc.
- [92939] Nat. Acad. Sci. 99: 9990-9995, 2002.
- [92940] 16248.
- [92941] Ema, M.; Matsushita, N.; Sogawa, K.; Ariyama, T.; Inazawa, J.;
- [92942] Nemoto, T.; Ota, M.; Oshimura, M.; Fujii-Kuriyama, Y.: Human arylhydrocarbon
- [92943] receptor: functional expression and chromosomal assignment to 7p21. J.
- [92944] Biochem. 116: 845-851, 1994.

[92945] 16249.

[92946] Le Beau, M. M.; Carver, L. A.; Espinosa, R., III; Schmidt, J. V.;

[92947] Bradfield, C. A.: Chromosomal localization of the human AHR locus

[92948] encoding the structural gene for the Ah receptor to 7p21-p15. Cytogenet.

[92949] Cell Genet. 66: 172-176, 1994.

[92950] 16250.

[92951] Micka, J.; Milatovich, A.; Menon, A.; Grabowski, G. A.; Puga, A.;

[92952] Nebert, D. W.: Human Ah receptor (AHR) gene: localization to 7p15

[92953] and suggestive correlation of polymorphism with CYP1A1 inducibility. Pharmacogenetics 7:

[92954] 95-101, 1997.

[92955] 16251.

[92956] Shimizu, Y.; Nakatsuru, Y.; Ichinose, M.; Takahashi, Y.; Kume,

[92957] H.; Mimura, J.; Fujii-Kuriyama, Y.; Ishikawa, T.: Benzo[a]pyrene

- [92958] carcinogenicity is lost in mice lacking the aryl hydrocarbon receptor. Proc.
- [92959] Nat. Acad. Sci. 97: 779–782, 2000.
- [92960] 16252.
- [92961] Chen, P.; Hao, W.; Rife, L.; Wang, X. P.; Shen, D.; Chen, J.; Ogden,
- [92962] T.; Van Boemel, G. B.; Wu, L.; Yang, M.; Fong, H. K. W.: A photic
- [92963] visual cycle of rhodopsin regeneration is dependent on Rgr. Nature
- [92964] Genet. 28: 256–260, 2001.
- [92965] 16253.
- [92966] Chen, X.-N.; Korenberg, J. R.; Jiang, M.; Shen, D.; Fong, H. K.
- [92967] W.: Localization of the human RGR opsin gene to chromosome 10q23. Hum.
- [92968] Genet. 97: 720–722, 1996.
- [92969] 16254.
- [92970] Jiang, M.; Pandey, S.; Fong, H. K. W.: An opsin homologue in the
- [92971] retina and pigment epithelium. Invest. Ophthalm. Visual Sci.

34:

[92972] 3669–3678, 1993.

[92973] 16255.

[92974] Morimura, H.; Saindelle–Ribeaudau, F.; Berson, E. L.;
Dryja, T.

[92975] P.: Mutations in RGR, encoding a light-sensitive opsin ho-
mologue,

[92976] in patients with retinitis pigmentosa. (Letter) Nature
Genet. 23:

[92977] 393–394, 1999.

[92978] 16256.

[92979] Shen, D.; Jiang, M.; Hao, W.; Tao, L.; Salazar, M.; Fong, H.
K.

[92980] W.: A human opsin-related gene that encodes a retinalde-
hyde-binding

[92981] protein. Biochemistry 33: 13117–13125, 1994.

[92982] 16257.

[92983] Sasada, R.; Ono, Y.; Taniyama, Y.; Shing, Y.; Folkman, J.;
Igarashi,

[92984] K.: Cloning and expression of cDNA encoding human be-
tacellulin, a

- [92985] new member of the EGF family. *Biochem. Biophys. Res. Commun.* 190: 1173–1179, 1993.
- [92986] 16258.
- [92987] Watanabe, T.; Shintani, A.; Nakata, M.; Shing, Y.; Folkman, J.;
- [92988] Igarashi, K.; Sasada, R.: Recombinant human betacellulin: molecular structure, biological activities, and receptor interaction. *J. Biol. Chem.* 269: 9966–9973, 1994.
- [92989] 16259.
- [92990] Asano, H.; Ishida, A.; Hasegawa, M.; Ono, T.; Yoshida, M. C.; Taniguchi, M.; Kanno, M.: The mouse Mel-18 'RING-finger' gene: genomic organization, promoter analysis and chromosomal assignment. *DNA Seq.* 3: 369–377, 1993.
- [92991] 16260.
- [92992] Ishida, A.; Asano, H.; Hasegawa, M.; Koseki, H.; Ono, T.;

Yoshida,

[92999] M. C.; Taniguchi, M.; Kanno, M.: Cloning and chromosome mapping of

[93000] the human Mel-18 gene which encodes a DNA-binding protein with a new

[93001] 'RING-finger' motif. Gene 129: 249-255, 1993.

[93002] 16261.

[93003] Tagawa, M.; Sakamoto, T.; Shigemoto, K.; Matsubara, H.; Tamura,

[93004] Y.; Ito, T.; Nakamura, I.; Okitsu, A.; Imai, K.; Taniguchi, M.: Expression

[93005] of novel DNA-binding protein with zinc finger structure in various

[93006] tumor cells. J. Biol. Chem. 265: 20021-20026, 1990.

[93007] 16262.

[93008] Hara, E.; Yamaguchi, T.; Nojima, H.; Ide, T.; Campisi, J.; Okayama,

[93009] H.; Oda, K.: Id-related genes encoding helix-loop-helix proteins

[93010] are required for G1 progression and are repressed in senescent human

[93011] fibroblasts. J. Biol. Chem. 269: 2139-2145, 1994.

- [93012] 16263.
- [93013] Mathew, S.; Chen, W.; Murty, V. V. V. S.; Benezra, R.; Chaganti,
- [93014] R. S. K.: Chromosomal assignment of human ID1 and ID2 genes. *Genomics* 30:
- [93015] 385–387, 1995.
- [93016] 16264.
- [93017] Chen, M. X.; McPartlin, A. E.; Brown, L.; Chen, Y. H.; Barker,
- [93018] H. M.; Cohen, P. T. W.: A novel human protein serine/threonine phosphatase,
- [93019] which possesses four tetratricopeptide repeat motifs and localizes
- [93020] to the nucleus. *EMBO J.* 13: 4278–4290, 1994.
- [93021] 16265.
- [93022] Xu, X.; Lagercrantz, J.; Zickert, P.; Bajalica-Lagercrantz, S.;
- [93023] Zetterberg, A.: Chromosomal localization and 5-prime sequence of
- [93024] the human protein serine/threonine phosphatase 5-prime gene. *Biochem.*
- [93025] *Biophys. Res. Commun.* 218: 514–517, 1996.

- [93026] 16266.
- [93027] Yong, W. H.; Ueki, K.; Chou, D.; Reeves, S. A.; von Deimling, A.;
- [93028] Gusella, J. F.; Mohrenweiser, H. W.; Buckler, A. J.; Louis, D. N.
- [93029] : Cloning of a highly conserved human protein serine-threonine phosphatase
- [93030] gene from the glioma candidate region on chromosome 19q13.3. Genomics 29:
- [93031] 533-536, 1995.
- [93032] 16267.
- [93033] Breitbart, R. E.; Liang, C.; Smoot, L. B.; Laheru, D. A.; Mahdavi,
- [93034] V.; Nadal-Ginard, B.: A fourth human MEF2 transcription factor, hMEF2D,
- [93035] is an early marker of the myogenic lineage. Development 118: 1095-1106,
- [93036] 1993.
- [93037] 16268.
- [93038] Hobson, G. M.; Krahe, R.; Garcia, E.; Siciliano, M. J.; Fumanage,

- [93039] V. L.: Regional chromosomal assignments for four members of the MADS
- [93040] domain transcription enhancer factor 2 (MEF2) gene family to human
- [93041] chromosomes 15q26, 19p12, 5q14, and 1q12-q23. Genomics 29: 704-711,
- [93042] 1995.
- [93043] 16269.
- [93044] Mao, Z.; Bonni, A.; Xia, F.; Nadal-Vicans, M.; Greenberg, M. E.
- [93045] : Neuronal activity-dependent cell survival mediated by transcription
- [93046] factor MEF2. Science 286: 785-790, 1999.
- [93047] 16270.
- [93048] Martin, J. F.; Miano, J. M.; Hustad, C. M.; Copeland, N. G.; Jenkins,
- [93049] N. A.; Olson, E. N.: A Mef2 gene that generates a muscle-specific
- [93050] isoform via alternative mRNA splicing. Molec. Cell. Biol. 14: 1647-1656,
- [93051] 1994.

- [93052] 16271.
- [93053] Molkenitin, J. D.; Black, B. L.; Martin, J. F.; Olson, E. N.: Co-operative
activation of muscle gene expression by MEF2 and myogenic bHLH proteins. Cell 83:
1125–1136, 1995.
- [93055] 16272.
- [93056] Naya, F. J.; Black, B. L.; Wu, H.; Bassel-Duby, R.; Richardson,
J. A.; Hill, J. A.; Olson, E. N.: Mitochondrial deficiency and cardiac
sudden death in mice lacking the MEF2A transcription factor. Nature
Med. 15Oct, 2002. Note: Advance Electronic Publication.
- [93060] 16273.
- [93061] Pollock, R.; Treisman, R.: Human SRF-related proteins: DNA-binding
properties and potential regulatory targets. Genes Dev. 5:
2327–2341,
1991.
- [93064] 16274.

- [93066] Suzuki, E.; Lowry, J.; Sonoda, G.; Testa, J. R.; Walsh, K.:
Structures
- [93067] and chromosome locations of the human MEF2A gene and
a pseudogene
- [93068] MEF2AP. Cytogenet. Cell Genet. 73: 244–249, 1996.
- [93069] 16275.
- [93070] Yu, Y.-T.; Breitbart, R. E.; Smoot, L. B.; Lee, Y.; Mahdavi,
V.;
- [93071] Nadal-Ginard, B.: Human myocyte-specific enhancer fac-
tor 2 comprises
- [93072] a group of tissue-restricted MADS box transcription fac-
tors. Genes
- [93073] Dev. 6: 1783–1798, 1992.
- [93074] 16276.
- [93075] Youn, H.-D.; Sun, L.; Prywes, R.; Liu, J. O.: Apoptosis of T
cells
- [93076] mediated by Ca(2+)-induced release of the transcription
factor MEF2. Science 286:
- [93077] 790–793, 1999.
- [93078] 16277.
- [93079] Krainc, D.; Haas, M.; Ward, D. C.; Lipton, S. A.; Bruns, G.;

Leifer,

[93080] D.: Assignment of human myocyte-specific enhancer binding factor

[93081] 2C (hMEF2C) to human chromosome 5q14 and evidence that MEF2C is evolutionarily

[93082] conserved. Genomics 29: 809–811, 1995.

[93083] 16278.

[93084] Leifer, D.; Krainc, D.; Yu, Y.-T.; McDermott, J.; Breitbart, R.

[93085] E.; Heng, J.; Neve, R. L.; Kosofsky, B.; Nadal-Ginard, B.; Lipton,

[93086] S. A.: MEF2C, a MADS/MEF2-family transcription factor expressed in

[93087] a laminar distribution in cerebral cortex. Proc. Nat. Acad. Sci. 90:

[93088] 1546–1550, 1993.

[93089] 16279.

[93090] Holmbeck, K.; Bianco, P.; Caterina, J.; Yamada, S.; Kromer, M.;

[93091] Kuznetsov, S. A.; Mankani, M.; Robey, P. G.; Poole, A. R.; Pidoux,

[93092] I.; Ward, J. M.; Birkedal-Hansen, H.: MT1-MMP-deficient

mice develop

[93093] dwarfism, osteopenia, arthritis, and connective tissue disease due

[93094] to inadequate collagen turnover. Cell 99: 81–92, 1999.

[93095] 16280.

[93096] Sato, H.; Takino, T.; Okada, Y.; Cao, J.; Shinagawa, A.; Yamamoto,

[93097] E.; Seiki, M.: A matrix metalloproteinase expressed on the surface

[93098] of invasive tumor cells. Nature 370: 61–65, 1994.

[93099] 16281.

[93100] Takino, T.; Sato, H.; Yamamoto, E.; Seiki, M.: Cloning of a human

[93101] gene potentially encoding a novel matrix metalloproteinase having

[93102] a C-terminal transmembrane domain. Gene 155: 293–298, 1995.

[93103] 16282.

[93104] McCright, B.; Rivers, A. M.; Audlin, S.; Virshup, D. M.: The B56

[93105] family of protein phosphatase 2A (PP2A) regulatory sub-

units encodes

[93106] differentiation-induced phosphoproteins that target PP2A to both nucleus

[93107] and cytoplasm. J. Biol. Chem. 271: 22081-22089, 1996.

[93108] 16283.

[93109] Van Hoof, C.; Aly, M. S.; Garcia, A.; Cayla, X.; Cassiman, J. J.;

[93110] Merlevede, W.; Goris, J.: Structure and chromosomal localization

[93111] of the human gene of the phosphotyrosyl phosphatase activator (PTPA)

[93112] of protein phosphatase 2A. Genomics 28: 261-272, 1995.

[93113] 16284.

[93114] Lory, P.; Ophoff, R. A.; Nahmias, J.: Towards a unified nomenclature

[93115] describing voltage-gated calcium channel genes. Hum. Genet. 100:

[93116] 149-150, 1997.

[93117] 16285.

[93118] Rosenfeld, M. R.; Wong, E.; Dalmau, J.; Manley, G.; Posner, J.

- [93119] B.; Sher, E.; Furneaux, H. M.: Cloning and characterization of a
- [93120] Lambert-Eaton myasthenic syndrome antigen. *Ann. Neurol.* 33: 113-120,
- [93121] 1993.
- [93122] 16286.
- [93123] Taviaux, S.; Williams, M. E.; Harpold, M. M.; Nargeot, J.; Lory,
- [93124] P.: Assignment of human genes for beta-2 and beta-4 subunits of voltage-dependent
- [93125] Ca(2+) channels to chromosomes 10p12 and 2q22-q23. *Hum. Genet.* 100:
- [93126] 151-154, 1997.
- [93127] 16287.
- [93128] Bontron, S.; Steimle, V.; Ucla, C.; Eibl, M. M.; Mach, B.: Two
- [93129] novel mutations in the MHC class II transactivator CIITA in a second
- [93130] patient from MHC class II deficiency complementation group A. *Hum.*
- [93131] *Genet.* 99: 541-546, 1997.
- [93132] 16288.

- [93133] Dziembowska, M.; Fondaneche, M.-C.; Vedrenne, J.; Barbieri, G.;
- [93134] Wiszniewski, W.; Picard, C.; Cant, A. J.; Steimle, V.; Charon, D.;
- [93135] Alca-Loridan, C.; Fischer, A.; Lisowska-Groszpiere, B.:
Three novel
mutations of the CIITA gene in MHC class II-deficient patients with
a severe immunodeficiency. Immunogenetics 53:
821-829, 2002.
- [93138] 16289.
- [93139] Harton, J. A.; Cressman, D. E.; Chin, K.-C.; Der, C. J.; Ting,
J. P.-Y.: GTP binding by class II transactivator: role in nuclear
import. Science 285: 1402-1405, 1999.
- [93142] 16290.
- [93143] Harton, J. A.; Ting, J. P.-Y.: Class II transactivator: mastering
the art of major histocompatibility complex expression.
Molec. Cell.
- [93145] Biol. 20: 6185-6194, 2000.

- [93146] 16291.
- [93147] Mach, B.; Steimle, V.; Martinez-Soria, E.; Reith, W.: Regulation
[93148] of MHC class II genes: lessons from a disease. *Annu. Rev. Immun.* 14:
[93149] 301–331, 1996.
- [93150] 16292.
- [93151] Mahanta, S. K.; Scholl, T.; Yang, F.-C.; Strominger, J. L.: Transactivation
[93152] by CIITA, the type II bare lymphocyte syndrome-associated factor,
[93153] requires participation of multiple regions of the TATA box binding
[93154] protein. *Proc. Nat. Acad. Sci.* 94: 6324–6329, 1997.
- [93155] 16293.
- [93156] Raval, A.; Howcroft, T. K.; Weissman, J. D.; Kirshner, S.; Zhu,
[93157] X.-S.; Yokoyama, K.; Ting, J.; Singer, D. S.: Transcriptional coactivator,
[93158] CIITA, is an acetyltransferase that bypasses a promoter requirement

[93159] for TAFII250. Molec. Cell 7: 105–115, 2001.

[93160] 16294.

[93161] Reith, W.: Personal Communication. Geneva, Switzerland
5/30/1997.

[93162] 16295.

[93163] Scholl, T.; Mahanta, S. K.; Strominger, J. L.: Specific complex

[93164] formation between the type II bare lymphocyte syndrome-associated

[93165] transactivators CIITA and RFX5. Proc. Nat. Acad. Sci. 94:
6330–6334,

[93166] 1997.

[93167] 16296.

[93168] Steimle, V.; Otten, L. A.; Zufferey, M.; Mach, B.: Complement-
mentation

[93169] cloning of an MHC class II transactivator mutated in
hereditary MHC

[93170] class II deficiency (or bare lymphocyte syndrome). Cell 75:
135–146,

[93171] 1993.

[93172] 16297.

- [93173] Zhu, X.-S.; Linhoff, M. W.; Li, G.; Chin, K.-C.; Maity, S. N.;
- [93174] Ting, J. P.-Y.: Transcriptional scaffold: CIITA interacts with NF- κ B,
- [93175] RFX, and CREB to cause stereospecific regulation of the class II major
- [93176] histocompatibility complex promoter. *Molec. Cell. Biol.* 20: 6051-6061,
- [93177] 2000.
- [93178] 16298.
- [93179] Emery, P.; Durand, B.; Mach, B.; Reith, W.: RFX proteins, a novel
- [93180] family of DNA binding proteins conserved in the eukaryotic kingdom. *Nucleic*
- [93181] *Acids Res.* 24: 803-807, 1996.
- [93182] 16299.
- [93183] Pugliati, L.; Reith, W.; Fey, S.; Mach, B.: Mapping the RF-X gene,
- [93184] encoding a DNA-binding protein controlling HLA class II gene expression,
- [93185] to 19p13.(Abstract) *Cytogenet. Cell Genet.* 51: 1061 only, 1989.

- [93186] 16300.
- [93187] Mizuta, M.; Inagaki, N.; Nemoto, Y.; Matsukura, S.; Takahashi,
- [93188] M.; Seino, S.: Synaptotagmin III is a novel isoform of rat synaptotagmin
- [93189] expressed in endocrine and neuronal cells. J. Biol. Chem. 269: 11675–11678,
- [93190] 1994.
- [93191] 16301.
- [93192] Li, X.-J.; Wang, D.-Y.; Zhu, Y.; Guo, R.-J.; Wang, X.-D.; Lubomir,
- [93193] K.; Mukai, K.; Sasaki, H.; Yoshida, H.; Oka, T.; Machinami, R.; Shinmura,
- [93194] K.; Tanaka, M.; Sugimura, H.: Mxi1 mutations in human neurofibrosarcomas. Jpn.
- [93195] J. Cancer Res. 90: 740–746, 1999.
- [93196] 16302.
- [93197] Prochownik, E. V.; Grove, L. E.; Deubler, D.; Zhu, X. L.; Stephenson,
- [93198] R. A.; Rohr, L. R.; Yin, X.; Brothman, A. R.: Commonly occurring

[93199] loss and mutation of the MXI1 gene in prostate cancer.

Genes Chromosomes

[93200] *Cancer* 22: 295–304, 1998.

[93201] 16303.

[93202] Schreiber–Agus, N.; Meng, Y.; Hoang, T.; Hou, H., Jr.;
Chen, K.;

[93203] Greenberg, R.; Cordon–Cardo, C.; Lee, H.–W.; DePinho, R.
A.: Role

[93204] of Mxi1 in ageing organ systems and the regulation of
normal and neoplastic

[93205] growth. *Nature* 393: 483–487, 1998.

[93206] 16304.

[93207] Shapiro, D. N.; Valentine, V.; Eagle, L.; Yin, X.; Morris, S.
W.;

[93208] Prochownik, E. V.: Assignment of the human MAD and
MXI1 genes to

[93209] chromosomes 2p12–p13 and 10q24–q25. *Genomics* 23:
282–285, 1994.

[93210] 16305.

[93211] Wechsler, D. S.; Hawkins, A. L.; Li, X.; Jabs, E. W.; Griffin,

[93212] C. A.; Dang, C. V.: Localization of the human Mxi1 tran–

scription

[93213] factor gene (MXI1) to chromosome 10q24-q25. Genomics 21: 669-672,

[93214] 1994.

[93215] 16306.

[93216] Wechsler, D. S.; Shelly, C. A.; Dang, C. V.: Genomic organization

[93217] of human MXI1, a putative tumor suppressor gene. Genomics 32: 466-470,

[93218] 1996.

[93219] 16307.

[93220] Zervos, A. S.; Gyuris, J.; Brent, R.: Mxi1, a protein that specifically

[93221] interacts with Max to bind Myc-Max recognition sites. Cell 72: 223-232,

[93222] 1993. Note: Erratum: Cell 79: 389 only, 1994.

[93223] 16308.

[93224] Boie, Y.; Rushmore, T. H.; Darmon-Goodwin, A.; Grygorczyk, R.;

[93225] Slipetz, D. M.; Metters, K. M.; Abramovitz, M.: Cloning and expression

- [93226] of a cDNA for the human prostanoid IP receptor. J. Biol. Chem. 269:
- [93227] 12173–12178, 1994.
- [93228] 16309.
- [93229] Ishikawa, T.; Tamai, Y.; Rochelle, J. M.; Hirata, M.; Namba, T.;
- [93230] Sugimoto, Y.; Ichikawa, A.; Narumiya, S.; Taketo, M. M.; Seldin, M.
- [93231] F.: Mapping of the genes encoding mouse prostaglandin D, E, and F
- [93232] and prostacyclin receptors. Genomics 32: 285–288, 1996.
- [93233] 16310.
- [93234] Murata, T.; Ushikubi, F.; Matsuoka, T.; Hirata, M.; Yamasaki, A.;
- [93235] Sugimoto, Y.; Ichikawa, A.; Aze, Y.; Tanaka, T.; Yoshida, N.; Ueno,
- [93236] A.; Ohishi, S.; Narumiya, S.: Altered pain perception and inflammatory
- [93237] response in mice lacking prostacyclin receptor. Nature 388: 678–682,
- [93238] 1997.

- [93239] 16311.
- [93240] Ogawa, Y.; Tanaka, I.; Inoue, M.; Yoshitake, Y.; Isse, N.; Nakagawa,
- [93241] O.; Usui, T.; Itoh, H.; Yoshimasa, T.; Narumiya, S.; Nakao, K.: Structural
- [93242] organization and chromosomal assignment of the human prostacyclin
- [93243] receptor gene. *Genomics* 27: 142–148, 1995.
- [93244] 16312.
- [93245] Hoffman, I.; Balling, R.: Chromosomal localization of the murine
- [93246] cadherin-11. *Mammalian Genome* 6: 304 only, 1995.
- [93247] 16313.
- [93248] Okazaki, M.; Takeshita, S.; Kawai, S.; Kikuno, R.; Tsujimura, A.;
- [93249] Kudo, A.; Amann, E.: Molecular cloning and characterization of OB-cadherin,
- [93250] a new member of cadherin family expressed in osteoblasts. *J. Biol.*
- [93251] *Chem.* 269: 12092–12098, 1994.
- [93252] 16314.

- [93253] Tanihara, H.; Sano, K.; Heimark, R. L.; St. John, T.; Suzuki, S.
- [93254] : Cloning of five human cadherins clarifies characteristic features
- [93255] of cadherin extracellular domain and provides further evidence for
- [93256] two structurally different types of cadherin. Cell Adhes. Commun. 2:
- [93257] 15–26, 1994.
- [93258] 16315.
- [93259] Courvalin, J.-C.; Lassoued, K.; Worman, H. J.; Blobel, G.: Identification
- [93260] and characterization of autoantibodies against the nuclear envelope
- [93261] lamin B receptor from patients with primary biliary cirrhosis. J.
- [93262] Exp. Med. 172: 961–967, 1990.
- [93263] 16316.
- [93264] Karl, H.: Die besondere Haufung von Merkmalsträgern der Pelger–Huet–Kernanomalie
- [93265] der Leukozyten in Gelenau im Erzgebirge. Thesis: Humboldt Univ. Berlin,

- [93266] Germany , 1967.
- [93267] 16317.
- [93268] Nachtsheim, H.: The Pelger-anomaly in man and rabbit. J. Hered. 41:
- [93269] 131-137, 1950.
- [93270] 16318.
- [93271] Schuler, E.; Lin, F.; Worman, H. J.: Characterization of the human
- [93272] gene encoding LBR, an integral protein of the nuclear envelope inner
- [93273] membrane. J. Biol. Chem. 269: 11312-11317, 1994.
- [93274] 16319.
- [93275] Christie, P. T.; Curley, A.; Nesbit, M. A.; Chapman, C.; Genet,
- [93276] S.; Harper, P. S.; Keeling, S. L.; Wilkie, A. O. M.; Winter, R. M.;
- [93277] Thakker, R. V.: Mutational analysis in X-linked spondyloepiphyseal
- [93278] dysplasia tarda. J. Clin. Endocr. Metab. 86: 3233-3236, 2001.
- [93279] 16320.

- [93280] Fiedler, J.; Bittner, M.; Puhl, W.; Brenner, R. E.: Mutations
[93281] in the X-linked spondyloepiphyseal dysplasia tarda (SEDL)
coding sequence
[93282] are not a common cause of early primary osteoarthritis in
men. (Letter) Clin.
[93283] Genet. 62: 94–95, 2002.
[93284] 16321.
[93285] Gecz, J.; Hillman, M. A.; Gedeon, A. K.; Cox, T. C.; Baker,
E.;
[93286] Mulley, J. C.: Gene structure and expression study of the
SEDL gene
[93287] for spondyloepiphyseal dysplasia tarda. Genomics 69:
242–251, 2000.
[93288] 16322.
[93289] Gedeon, A. K.; Colley, A.; Jamieson, R.; Thompson, E. M.;
Rogers,
[93290] J.; Sillence, D.; Tiller, G. E.; Mulley, J. C.; Gecz, J.: Identifi-
cation
[93291] of the gene (SEDL) causing X-linked spondyloepiphyseal
dysplasia tarda. Nature
[93292] Genet. 22: 400–404, 1999.

[93293] 16323.

[93294] Gedeon, A. K.; Tiller, G. E.; Le Merrer, M.; Heuertz, S.;
Tranebjaerg,

[93295] L.; Chitayat, D.; Robertson, S.; Glass, I. A.; Savarirayan, R.;
Cole,

[93296] W. G.; Rimoin, D. L.; Kousseff, B. G.; Ohashi, H.; Zabel, B.;
Munnich,

[93297] A.; Gecz, J.; Mulley, J. C.: The molecular basis of X-linked
spondyloepiphyseal

[93298] dysplasia tarda. Am. J. Hum. Genet. 68: 1386–1397,
2001.

[93299] 16324.

[93300] Grunebaum, E.; Arpaia, E.; MacKenzie, J. J.; Fitzpatrick, J.;
Ray,

[93301] P. N.; Roifman, C. M.: A missense mutation in the SEDL
gene results

[93302] in delayed onset of X linked spondyloepiphyseal dysplasia
in a large

[93303] pedigree. (Letter) J. Med. Genet. 38: 409–411, 2001.

[93304] 16325.

[93305] Mumm, S.; Christie, P. T.; Finnegan, P.; Jones, J.; Dixon, P.
H.;

- [93306] Pannett, A. A. J.; Harding, B.; Gottesman, G. S.; Thakker, R. V.;
- [93307] Whyte, M. P.: A five-base pair deletion in the sedlin gene causes
- [93308] spondyloepiphyseal dysplasia tarda in a six-generation Arkansas kindred. J.
- [93309] Clin. Endocr. Metab. 85: 3343–3347, 2000.
- [93310] 16326.
- [93311] Mumm, S.; Zhang, X.; Vacca, M.; D'Esposito, M.; Whyte, M. P.:
- [93312] The sedlin gene for spondyloepiphyseal dysplasia tarda escapes X-inactivation
- [93313] and contains a non-canonical splice site. Gene 273: 285–293, 2001.
- [93314] 16327.
- [93315] Takahashi, T.; Takahashi, I.; Tsuchida, S.; Oyama, K.; Ko-matsu,
- [93316] M.; Saito, H.; Takada, G.: An SEDL gene mutation in a Japanese kindred
- [93317] of X-linked spondyloepiphyseal dysplasia tarda. (Letter) Clin. Genet. 61:
- [93318] 319–320, 2002.

- [93319] 16328.
- [93320] Tiller, G. E.; Hannig, V. L.; Dozier, D.; Carrel, L.; Trevarthen,
- [93321] K. C.; Wilcox, W. R.; Mundlos, S.; Haines, J. L.; Gedeon, A. K.; Gecz,
- [93322] J.: A recurrent RNA-splicing mutation in the SEDL gene causes X-linked
- [93323] spondyloepiphyseal dysplasia tarda. *Am. J. Hum. Genet.* 68: 1398-1407,
- [93324] 2001.
- [93325] 16329.
- [93326] Whyte, M. P.; Gottesman, G. S.; Eddy, M. C.; McAlister, W. H.
- [93327] : X-linked recessive spondyloepiphyseal dysplasia tarda: clinical
- [93328] and radiographic evolution in a 6-generation kindred and review of
- [93329] the literature. *Medicine* 78: 9-25, 1999.
- [93330] 16330.
- [93331] Montini, E.; Andolfi, G.; Caruso, A.; Buchner, G.; Walpole, S.

- [93332] M.; Mariani, M.; Consalez, G.; Trump, D.; Ballabio, A.; Franco, B.
- [93333] : Identification and characterization of a novel serine-threonine
- [93334] kinase gene from the Xp22 region. *Genomics* 51: 427-433, 1998.
- [93335] 16331.
- [93336] Becker, M. A.; Heidler, S. A.; Bell, G. I.; Seino, S.; Le Beau, M. M.; Westbrook, C. A.; Neuman, W.; Shapiro, L. J.; Mohandas, T.
- [93338] K.; Roessler, B. J.; Palella, T. D.: Cloning of cDNAs for human phosphoribosylpyrophosphate
- [93339] synthetases 1 and 2 and X chromosome localization of PRPS1 and PRPS2
- [93340] genes. *Genomics* 8: 555-561, 1990.
- [93341] 16332.
- [93342] Braverman, N.; Lin, P.; Moebius, F. F.; Obie, C.; Moser, A.; Glossmann,
- [93343] H.; Wilcox, W. R.; Rimoin, D. L.; Smith, M.; Kratz, L.; Kelley, R.
- [93344] I.; Valle, D.: Mutations in the gene encoding 3-beta-hydroxysteroid-delta(8),delta(7)-isomerase

- [93345] cause X-linked dominant Conradi-Hunermann syndrome.
Nature Genet. 22:
[93346] 291-294, 1999.
- [93347] 16333.
- [93348] Cho, S. Y.; Kim, J. H.; Paik, Y. K.: Cholesterol biosynthesis
[93349] from lanosterol: differential inhibition of sterol delta
8-isomerase
[93350] and other lanosterol-converting enzymes by tamoxifen.
Molec. Cells 8:
[93351] 233-239, 1998.
- [93352] 16334.
- [93353] Clayton, P. T.; Kalter, D. C.; Atherton, D. J.; Besley, G. T.;
[93354] Broadhead, D. M.: Peroxisomal enzyme deficiency in X-
linked dominant
[93355] Conradi-Hunermann syndrome. J. Inherit. Metab. Dis. 12:
358-360,
[93356] 1989.
- [93357] 16335.
- [93358] Derry, J. M. J.; Gormally, E.; Means, G. D.; Zhao, W.;
Meindl,

- [93359] A.; Kelley, R. I.; Boyd, Y.; Herman, G. E.: Mutations in a delta(8)-delta(7)
- [93360] sterol isomerase in the tattered mouse and X-linked dominant chondrodysplasia
- [93361] punctata. Nature Genet. 22: 286-290, 1999.
- [93362] 16336.
- [93363] Grange, D. K.; Kratz, L. E.; Braverman, N. E.; Kelley, R. I.:
- [93364] CHILD syndrome caused by deficiency of 3-beta-hydroxysteroid-delta-8,delta-7-isomerase. Am.
- [93365] J. Med. Genet. 90: 328-335, 2000.
- [93366] 16337.
- [93367] Hanner, M.; Moebius, F. F.; Weber, F.; Grabner, M.; Striessnig,
- [93368] J.; Glossmann, H.: Phenylalkylamine Ca(2+) antagonist binding protein:
- [93369] molecular cloning, tissue distribution, and heterologous expression. J.
- [93370] Biol. Chem. 270: 7551-7557, 1995.
- [93371] 16338.
- [93372] Has, C.; Bruckner-Tuderman, L.; Muller, D.; Floeth, M.; Folkers,

- [93373] E.; Donnai, D.; Traupe, H.: The Conradi–Hunermann–Happle syndrome
- [93374] (CDPX2) and emopamil binding protein: novel mutations, and somatic
- [93375] and gonadal mosaicism. Hum. Molec. Genet. 9: 1951–1955, 2000.
- [93376] 16339.
- [93377] Holmes, R. D.; Wilson, G. N.; Hajra, A. K.: Peroxisomal enzyme
- [93378] deficiency in the Conradi–Hunerman (sic) form of chondrodysplasia
- [93379] punctata. New Eng. J. Med. (Letter) 316: 1608 only, 1987.
- [93380] 16340.
- [93381] Ikegawa, S.; Ohashi, H.; Ogata, T.; Honda, A.; Tsukahara, M.; Kubo,
- [93382] T.; Kimizuka, M.; Shimode, M.; Hasegawa, T.; Nishimura, G.; Nakamura,
- [93383] Y.: Novel and recurrent EBP mutations in X-linked dominant chondrodysplasia
- [93384] punctata. Am. J. Med. Genet. 94: 300–305, 2000.
- [93385] 16341.

- [93386] Kelley, R. I.; Wilcox, W. G.; Smith, M.; Kratz, L. E.; Moser, A.; Rimoin, D. S.: Abnormal sterol metabolism in patients with Conradi-Hunermann-Happle syndrome and sporadic lethal chondrodysplasia punctata. Am. J. Med. Genet. 83: 213-219, 1999.
- [93387] 16342.
- [93388] Rasmussen, U. B.; Wolf, C.; Mattei, M.-G.; Chenard, M.-P.; Bellocq, J.-P.; Chambon, P.; Rio, M.-C.; Basset, P.: Identification of a new interferon- α -inducible gene (p27) on human chromosome 14q32 and its expression in breast carcinoma. Cancer Res. 53: 4096-4101, 1993.
- [93389] 16343.
- [93390] Lan, M. S.; Li, Q.; Lu, J.; Modi, W. S.; Notkins, A. L.: Genomic organization, 5-prime-upstream sequence, and chromosomal localization of an insulinoma-associated intronless gene, IA-1. J. Biol. Chem. 269:

[93399] 14170–14174, 1994.

[93400] 16344.

[93401] Bennett, B. D.; Wang, Z.; Kuang, W.-J.; Wang, A.; Groopman, J.

[93402] E.; Goeddel, D. V.; Scadden, D. T.: Cloning and characterization

[93403] of HTK, a novel transmembrane tyrosine kinase of the EPH subfamily. J.

[93404] Biol. Chem. 269: 14211–14218, 1994.

[93405] 16345.

[93406] Berclaz, G.; Andres, A.-C.; Albrecht, D.; Dreher, E.; Ziemiecki,

[93407] A.; Gusterson, B. A.; Crompton, M. R.: Expression of the receptor

[93408] protein tyrosine kinase myk-1/htk in normal and malignant mammary

[93409] epithelium. Biochem. Biophys. Res. Commun. 226: 869–875, 1996.

[93410] 16346.

[93411] Gerety, S. S.; Wang, H. U.; Chen, Z.-F.; Anderson, D. J.: Symmetrical

- [93412] mutant phenotypes of the receptor EphB4 and its specific transmembrane
- [93413] ligand ephrin-B2 in cardiovascular development. *Molec. Cell* 4: 403-414,
- [93414] 1999.
- [93415] 16347.
- [93416] Moynihan, T. P.; Ardley, H. C.; Leek, J. P.; Thompson, J.; Brindle,
- [93417] N. S.; Markham, A. F.; Robinson, P. A.: Characterization of a human
- [93418] ubiquitin-conjugating enzyme gene UBE2L3. *Mammalian Genome* 7: 520-525,
- [93419] 1996.
- [93420] 16348.
- [93421] Flanagan, J. R.; Becker, K. G.; Ennist, D. L.; Gleason, S. L.;
- [93422] Driggers, P. H.; Levi, B.-Z.; Appella, E.; Ozato, K.: Cloning of
- [93423] a negative transcription factor that binds to the upstream conserved
- [93424] region of Moloney murine leukemia virus. *Molec. Cell. Biol.* 12:

- [93425] 38–44, 1992.
- [93426] 16349.
- [93427] Hariharan, N.; Kelley, D. E.; Perry, R. P.: Delta, a transcription
- [93428] factor that binds to downstream elements in several polymerase II
- [93429] promoters, is a functionally versatile zinc finger protein. Proc.
- [93430] Nat. Acad. Sci. 88: 9799–9803, 1991.
- [93431] 16350.
- [93432] Oei, S. L.; Shi, Y.: Transcription factor Yin Yang 1 stimulates
- [93433] poly(ADP-ribosyl)ation and DNA repair. Biochem. Biophys. Res. Commun. 284:
- [93434] 450–454, 2001.
- [93435] 16351.
- [93436] Park, K.; Atchison, M. L.: Isolation of a candidate repressor/activator,
- [93437] NF-E1 (YY-1, delta), that binds to the immunoglobulin kappa 3-prime
- [93438] enhancer and the immunoglobulin heavy-chain micro-E1

site.. Proc.

[93439] Nat. Acad. Sci. 88: 9804–9808, 1991.

[93440] 16352.

[93441] Shi, Y.; Seto, E.; Chang, L.-S.; Shenk, T.: Transcriptional repression

[93442] by YY1, a human GLI-Kruppel-related protein, and relief of repression

[93443] by adenovirus E1A protein. Cell 67: 377–388, 1991.

[93444] 16353.

[93445] Yao, Y.-L.; Dupont, B. R.; Ghosh, S.; Fang, Y.; Leach, R. J.; Seto,

[93446] E.: Cloning, chromosomal localization and promoter analysis of the

[93447] human transcription factor YY1. Nucleic Acids Res. 26: 3776–3783,

[93448] 1998.

[93449] 16354.

[93450] Zhu, W.; Lossie, A. C.; Camper, S. A.; Gumucio, D. L.: Chromosomal

[93451] localization of the transcription factor YY1 in the mouse and human. Mammalian

- [93452] Genome 5: 234–236, 1994.
- [93453] 16355.
- [93454] Hakimi, M.–A.; Bochar, D. A.; Schmiesing, J. A.; Dong, Y.; Barak,
- [93455] O. G.; Speicher, D. W.; Yokomori, K.; Shiekhattar, R.: A chromatin
- [93456] remodelling complex that loads cohesin onto human chromosomes. Nature 418:
- [93457] 994–998, 2002.
- [93458] 16356.
- [93459] Berglund, E. O.; Murai, K. K.; Fredette, B.; Sekerkova, G.; Marturano,
- [93460] B.; Weber, L.; Mugnaini, E.; Ranscht, B.: Ataxia and abnormal cerebellar
- [93461] microorganization in mice with ablated contactin gene expression. Neuron 24:
- [93462] 739–750, 1999.
- [93463] 16357.
- [93464] Nagasaki, K.; Manabe, T.; Hanzawa, H.; Maass, N.; Tsukada, T.;
- [93465] Yamaguchi, K.: Identification of a novel gene, LDOC1,

down-regulated

[93466] in cancer cell lines. Cancer Lett. 140: 227-234, 1999.

[93467] 16358.

[93468] Nomura, N.; Miyajima, N.; Sazuka, T.; Tanaka, A.;
Kawarabayasi,

[93469] Y.; Sato, S.; Nagase, T.; Seki, N.; Ishikawa, K.; Tabata, S.:
Prediction

[93470] of the coding sequences of unidentified human genes. I.
The coding

[93471] sequences of 40 new genes (KIAA0001-KIAA0040) de-
duced by analysis

[93472] of randomly samples cDNA clones from human immature
myeloid cell line

[93473] KG-1. DNA Res. 1: 27-35, 1994.

[93474] 16359.

[93475] Kikuno, R.; Nagase, T.; Ishikawa, K.; Hirose, M.; Miya-
jima, N.;

[93476] Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Prediction
of the

[93477] coding sequences of unidentified human genes. XIV. The
complete sequences

[93478] of 100 new cDNA clones from brain which code for large

proteins in

[93479] vitro. DNA Res. 6: 197–205, 1999.

[93480] 16360.

[93481] Polsky, D.; Young, A. Z.; Busam, K. J.; Alani, R. M.: The transcriptional

[93482] repressor of p16/Ink4a, Id1, is up-regulated in early melanomas. Cancer

[93483] Res. 61: 6008–6011, 2001.

[93484] 16361.

[93485] Dahl, N.; Hu, L. J.; Chery, M.; Fardeau, M.; Gilgenkrantz, S.;

[93486] Nivelon–Chevallier, A.; Sidaner–Noisette, I.; Mugneret, F.; Gouyon,

[93487] J. B.; Gal, A.; Kioschis, P.; d'Urso, M.; Mandel, J.–L.: Myotubular

[93488] myopathy in a girl with a deletion at Xq27–q28 and unbalanced X inactivation

[93489] assigns the MTM1 gene to a 600–kb region. Am. J. Hum. Genet. 56:

[93490] 1108–1115, 1995.

[93491] 16362.

[93492] Dahl, N.; Samson, F.; Thomas, N. S. T.; Hu, L. J.; Gong, W.;

Herman,

[93493] G.; Laporte, J.; Kioschis, P.; Poustka, A.; Mandel, J. L.: X linked

[93494] myotubular myopathy (MTM1) maps between DXS304 and DXS305, closely

[93495] linked to the DXS455 VNTR and a new, highly informative microsatellite

[93496] marker (DXS1684). J. Med. Genet. 31: 922-924, 1994.

[93497] 16363.

[93498] Darnfors, C.; Borje Larsson, H. E.; Oldfors, A.; Kyllerman, M.;

[93499] Gustavson, K.-H.; Bjursell, G.; Wahlstrom, J.: X-linked myotubular

[93500] myopathy: a linkage study. Clin. Genet. 37: 335-340, 1990.

[93501] 16364.

[93502] Darnfors, C.; Borje Larsson, H. E.; Oldfors, A.; Kyllerman, M.;

[93503] Gustavson, K.-H.; Bjursell, G.; Wahlstrom, J.: X-linked myotubular

[93504] myopathy: a linkage study.(Abstract) Cytogenet. Cell Genet. 51:

- [93505] 983, 1989.
- [93506] 16365.
- [93507] de Gouyon, B.; Chatterjee, A.; Monaco, A.; Quaderi, N.;
Brown,
- [93508] S. D. M.; Herman, G. E.: Comparative mapping on the
mouse X chromosome
- [93509] defines a myotubular myopathy equivalent region. Mam-
malian Genome 7:
- [93510] 575–579, 1996.
- [93511] 16366.
- [93512] de Gouyon, B. M.; Zhao, W.; Laporte, J.; Mandel, J.–L.; Met-
zenberg,
- [93513] A.; Herman, G. E.: Characterization of mutations in the
myotubularin
- [93514] gene in twenty six patients with X-linked myotubular my-
opathy. Hum.
- [93515] Molec. Genet. 6: 1499–1504, 1997.
- [93516] 16367.
- [93517] Engel, W. K.; Gold, G. N.; Karpatis, B.: Type I fiber hypotro-
phy
- [93518] and central nuclei. Arch. Neurol. 18: 435–444, 1968.

- [93519] 16368.
- [93520] Fidzianska, A.; Warlo, I.; Goebel, H. H.: Neonatal centronuclear
- [93521] myopathy with N-Cam decorated myotubes. Neuropediatrics 25: 158-161,
- [93522] 1994.
- [93523] 16369.
- [93524] Guiraud-Chaumeil, C.; Vincent, M. C.; Laporte, J.; Fardeau, M.;
- [93525] Samson, F.; Mandel, J.-L.: A mutation in the MTM1 gene invalidates
- [93526] a previous suggestion of nonallelic heterogeneity in X-linked myotubular
- [93527] myopathy. (Letter) Am. J. Hum. Genet. 60: 1542-1544, 1997.
- [93528] 16370.
- [93529] Hammans, S. R.; Robinson, D. O.; Moutou, C.; Kennedy, C. R.; Dennis,
- [93530] N. R.; Hughes, P. J.; Ellison, D. W.: A clinical and genetic study
- [93531] of a manifesting heterozygote with X-linked myotubular myopathy. Neuromusc.

[93532] Disord. 10: 133–137, 2000.

[93533] 16371.

[93534] Hane, B. G.; Rogers, R. C.; Schwartz, C. E.: Germline mosaicism

[93535] in X-linked myotubular myopathy. Clin. Genet. 56: 77–81, 1999.

[93536] 16372.

[93537] Heckmatt, J. Z.; Sewry, C. A.; Hodes, D.; Dubowitz, V.: Congenital

[93538] centronuclear (myotubular) myopathy: a clinical, pathological and

[93539] genetic study in eight children. Brain 108: 941–964, 1985.

[93540] 16373.

[93541] Herman, G. E.; Finegold, M.; de Gouyon, B.; Metzenberg, A.: Medical

[93542] complications in long-term survivors with X-linked myotubular myopathy. J.

[93543] Pediat. 134: 206–214, 1999.

[93544] 16374.

[93545] Herman, G. E.; Kopacz, K.; Zhao, W.; Mills, P. L.; Metzenberg,

- [93546] A.; Das, S.: Characterization of mutations in fifty North American
- [93547] patients with X-linked myotubular myopathy. Hum. Mutat. 19: 114–121,
- [93548] 2002.
- [93549] 16375.
- [93550] Janssen, E. A. M.; Hensels, G. W.; van Oost, B. A.; Hamel, B.
- [93551] C. J.; Kemp, S.; Baas, F.; Weber, J. W.; Barth, P. G.; Bolhuis, P.
- [93552] A.: The gene for X-linked myotubular myopathy is located in an 8
- [93553] Mb region at the border of Xq27.3 and Xq28. Neuromusc. Disord. 4:
- [93554] 455–461, 1994.
- [93555] 16376.
- [93556] Joseph, M.; Pai, G. S.; Holden, K. R.; Herman, G.: X-linked myotubular
- [93557] myopathy: clinical observations in ten additional cases. Am. J. Med.
- [93558] Genet. 59: 168–173, 1995.
- [93559] 16377.

- [93560] Keppen, L. D.; Husain, M. M.; Woody, R. C.: X-linked myotubular
- [93561] myopathy: intrafamilial variability and normal muscle biopsy in a
- [93562] heterozygous female. Clin. Genet. 32: 95-99, 1987.
- [93563] 16378.
- [93564] Kioschis, P.; Rogner, U. C.; Pick, E.; Klauck, S. M.; Heiss, N.;
- [93565] Siebenhaar, R.; Korn, B.; Coy, J. F.; Laporte, J.; Liechti-Gallati,
- [93566] S.; Poustka, A.: A 900-kb cosmid contig and 10 new transcripts within
- [93567] the candidate region for myotubular myopathy (MTM1). Genomics 33:
- [93568] 365-373, 1996.
- [93569] 16379.
- [93570] Laporte, J.; Biancalana, V.; Tanner, S. M.; Kress, W.; Schneider,
- [93571] V.; Wallgren-Pettersson, C.; Herger, F.; Buj-Bello, A.; Blondeau,
- [93572] F.; Liechti-Gallati, S.; Mandel, J.-L.: MTM1 mutations in X-linked

- [93573] myotubular myopathy. Hum. Mutat. 15: 393–409, 2000.
- [93574] 16380.
- [93575] Laporte, J.; Blondeau, F.; Buj-Bello, A.; Mandel, J.-L.: The
- [93576] myotubularin family: from genetic disease to phospho-
inositide metabolism. Trends
- [93577] Genet. 17: 221–228, 2001.
- [93578] 16381.
- [93579] Laporte, J.; Blondeau, F.; Buj-Bello, A.; Tentler, D.; Kretz,
- [93580] C.; Dahl, N.; Mandel, J.-L.: Characterization of the my-
otubularin
- [93581] dual specificity phosphatase gene family from yeast to
human. Hum.
- [93582] Molec. Genet. 7: 1703–1712, 1998.
- [93583] 16382.
- [93584] Laporte, J.; Guiraud-Chaumeil, C.; Tanner, S. M.; Blon-
deau, F.;
- [93585] Hu, L.-J.; Vicaire, S.; Liechti-Gallati, S.; Mandel, J.-L.: Ge-
nomic
- [93586] organization of the MTM1 gene implicated in X-linked
myotubular myopathy. Europ.

- [93587] J. Hum. Genet. 6: 325–330, 1998.
- [93588] 16383.
- [93589] Laporte, J.; Guiraud–Chaumeil, C.; Vincent, M.–C.; Mandel, J.–L.;
- [93590] Tanner, S. M.; Liechti–Gallati, S.; Wallgren–Pettersson, C.; Dahl,
- [93591] N.; Kress, W.; Bolhuis, P. A.; Fardeau, M.; Samson, F.; Bertini, E.;
- [93592] members of the ENMC International Consortium on Myotubular Myopathy
- [93593] : Mutations in the MTM1 gene implicated in X-linked myotubular myopathy. Hum.
- [93594] Molec. Genet. 6: 1505–1511, 1997.
- [93595] 16384.
- [93596] Lehesjoki, A.–E.; Sankila, E.–M.; Miao, J.; Somer, M.; Salonen,
- [93597] R.; Rapola, J.; de la Chapelle, A.: X linked neonatal myotubular
- [93598] myopathy: one recombination detected with four polymorphic DNA markers
- [93599] from Xq28. J. Med. Genet. 27: 288–291, 1990.

- [93600] 16385.
- [93601] Liechti-Gallati, S.; Muller, B.; Grimm, T.; Kress, W.; Muller, C.; Boltshauser, E.; Moser, H.; Braga, S.: X-linked centronuclear myopathy: mapping the gene to Xq28. Neuromusc. Disord. 1: 239-245, 1991.
- [93604] 16386.
- [93605] 16386.
- [93606] Meyers, K. R.; Golomb, H. M.; Hansen, J. L.; McKusick, V. A.: Familial neuromuscular disease with 'myotubes.'. Clin. Genet. 5: 327-337, 1974.
- [93608] 16387.
- [93610] Walker, J. L.; Dixon, J.; Fenton, C. R.; Hungerford, J.; Lynch, S. A.; Stenhouses, S. A. R.; Christian, A.; Craig, I. W.: Two new mutations in exon 3 of the NDP gene: S73X and S101F associated with severe and less severe ocular phenotype, respectively.

Hum. Mutat. 9:

[93614] 53–56, 1997.

[93615] 16388.

[93616] Lee, D. K.; Horikoshi, M.; Roeder, R. G.: Interaction of
TFIID

[93617] in the minor groove of the TATA element. Cell 67:
1241–1250, 1991.

[93618] 16389.

[93619] Ho, M. F.; Chalmers, R. M.; Davis, M. B.; Harding, A. E.;
Monaco,

[93620] A. P.: A novel point mutation in the McLeod syndrome
gene in neuroacanthocytosis. Ann.

[93621] Neurol. 39: 672–675, 1996.

[93622] 16390.

[93623] Wilson, R. B.; Kiledjian, M.; Shen, C.-P.; Benezra, R.;
Zwollo,

[93624] P.; Dymecki, S. M.; Desiderio, S. V.; Kadesch, T.: Repres-
sion of

[93625] immunoglobulin enhancers by the helix–loop–helix pro-
tein Id: implications

[93626] for B–lymphoid–cell development. Molec. Cell. Biol. 11:

6185-6191,

[93627] 1991.

[93628] 16391.

[93629] Warburg, M.: Norrie's disease: a new hereditary bilateral
pseudotumour

[93630] of the retina. Acta Ophthal. (Copenh) 39: 757-772, 1961.

[93631] 16392.

[93632] Warburg, M.: Norrie's disease (atrofia bulborum hereditaria). Acta

[93633] Ophthal. 41: 134-146, 1963.

[93634] 16393.

[93635] Warburg, M.: Norrie's disease, a congenital progressive
oculo-acoustico-cerebral

[93636] degeneration. Acta Ophthal. 89 (suppl.): 1-147, 1966.

[93637] 16394.

[93638] Warburg, M.; Hauge, M.; Sanger, R.: Norrie's disease and
the

[93639] Xg blood group system: linkage data. Acta Genet. Statist.
Med. 15:

[93640] 103-115, 1965.

- [93641] 16395.
- [93642] Whitnall, S. E.; Norman, R. M.: Microphthalmia and the visual
[93643] pathways: a case associated with blindness and imbecility,
and sex-linked. Brit.
[93644] J. Ophthal. 24: 229-244, 1940.
- [93645] 16396.
- [93646] Wilson, W. M. G.: Congenital blindness (pseudoglioma) oc-
curring
[93647] as a sex-linked developmental anomaly. Canad. Med. As-
soc. J. 60:
[93648] 580-584, 1949.
- [93649] 16397.
- [93650] Wolff, G.; Mayerova, A.; Wienker, T. F.; Atalianis, P.; Ioan-
nou,
[93651] P.; Warburg, M.: Clinical reinvestigation and linkage analy-
sis in
[93652] the family with Episkopi blindness (Norrie disease). J. Med.
Genet. 29:
[93653] 816-819, 1992.
- [93654] 16398.

- [93655] Wong, F.; Goldberg, M. F.; Hao, Y.: Identification of a non-sense
- [93656] mutation at codon 128 of the Norrie's disease gene in a male infant. Arch.
- [93657] Ophthal. 111: 1553-1557, 1993.
- [93658] 16399.
- [93659] Woodruff, G.; Newbury-Ecob, R.; Plaha, D. S.; Young, I. D.: Manifesting
- [93660] heterozygosity in Norrie's disease?. Brit. J. Ophthal. 77: 813-814,
- [93661] 1993.
- [93662] 16400.
- [93663] Zhu, D.; Antonarakis, S. E.; Schmeckpeper, B. J.; Diergaarde,
- [93664] P. J.; Greb, A. E.; Maumenee, I. H.: Microdeletion in the X-chromosome
- [93665] and prenatal diagnosis in a family with Norrie disease. Am. J. Med.
- [93666] Genet. 33: 485-488, 1989.
- [93667] 16401.
- [93668] Avner, P.; Bucan, M.; Arnaud, D.; Lehrach, H.; Rapp, U.: A-

raf

- [93669] oncogene localizes on mouse X chromosome to region
some 10–17 centimorgans
- [93670] proximal to hypoxanthine phosphoribosyltransferase
gene. Somat. Cell
- [93671] Molec. Genet. 13: 267–272, 1987.
- [93672] 16402.
- [93673] Beck, T. W.; Huleihel, M.; Gunnell, M.; Bonner, T. I.; Rapp,
U.
- [93674] R.: The complete coding sequence of the human A-raf-1
oncogene and
- [93675] transforming activity of a human A-raf carrying retrovirus.
Nucleic
- [93676] Acids Res. 15: 595–609, 1987.
- [93677] 16403.
- [93678] Huebner, K.; ar-Rushdi, A.; Griffin, C. A.; Isobe, M.; Kozak,
C.;
- [93679] Emanuel, B. S.; Nagarajan, L.; Cleveland, J. L.; Bonner, T. I.;
Goldsborough,
- [93680] M. D.; Croce, C. M.; Rapp, U.: Actively transcribed genes in
the
- [93681] raf oncogene group, located on the X chromosome in

mouse and human. Proc.

[93682] Nat. Acad. Sci. 83: 3934–3938, 1986.

[93683] 16404.

[93684] Lee, J.-E.; Beck, T. W.; Brennscheidt, U.; DeGennaro, L. J.;
Rapp,

[93685] U. R.: The complete sequence and promoter activity of the
human A-raf-1

[93686] gene (ARAF1). Genomics 20: 43–55, 1994.

[93687] 16405.

[93688] Mark, G. E.; Seeley, T. W.; Shows, T. B.; Mountz, J. D.: Pks,

[93689] a raf-related sequence in humans. Proc. Nat. Acad. Sci.
83: 6312–6316,

[93690] 1986.

[93691] 16406.

[93692] Popescu, N. C.; Mark, G. E.: Localization of the pKs gene, a
raf

[93693] related sequence on human chromosomes X and 7. Onco-
gene 4: 517–519,

[93694] 1989.

[93695] 16407.

- [93696] Dowdy, S. F.; Fasching, C. L.; Araujo, D.; Lai, K.-M.; Li-
vanos,
[93697] E.; Weissman, B. E.; Stanbridge, E. J.: Suppression of tu-
morigenicity
[93698] in Wilms' tumor by the p15.5-p14 region of chromosome
11. Science 254:
[93699] 293-295, 1991.
[93700] 16408.
[93701] Dowdy, S. F.; Lai, K.-M.; Weissman, B. E.; Matsui, Y.;
Hogan, B.
[93702] L. M.; Stanbridge, E. J.: The isolation and characterization
of a
[93703] novel cDNA demonstrating an altered mRNA level in non-
tumorigenic Wilms'
[93704] microcell hybrid cells. Nucleic Acids Res. 19: 5763-5769,
1991.
[93705] 16409.
[93706] Farmer, A. A.; Loftus, T. M.; Mills, A. A.; Sato, K. Y.; Neill,
[93707] J. D.; Tron, T.; Yang, M.; Trumpower, B. L.; Stanbridge, E.
J.: Extreme
[93708] evolutionary conservation of QM, a novel c-Jun associated
transcription

- [93709] factor. Hum. Molec. Genet. 3: 723–728, 1994.
- [93710] 16410.
- [93711] Kaneko, K.; Kobayashi, H.; Onodera, O.; Miyatake, T.;
Tsuji, S.
- [93712] : Genomic organization of a cDNA (QM) demonstrating an
altered mRNA
- [93713] level in nontumorigenic Wilms' microcell hybrid cells and
its localization
- [93714] to Xq28. Hum. Molec. Genet. 1: 529–533, 1992.
- [93715] 16411.
- [93716] Korn, B.; Sedlacek, Z.; Manca, A.; Kioschis, P.; Konecki, D.;
Lehrach,
- [93717] H.; Poustka, A.: A strategy for the selection of transcribed
sequences
- [93718] in the Xq28 region. Hum. Molec. Genet. 1: 235–242,
1992.
- [93719] 16412.
- [93720] van den Ouweland, A. M. W.; Verdijk, M.; Mannens, M. M.
A. M.;
- [93721] van Oost, B. A.: The QM gene is X-linked and therefore
not involved

- [93722] in suppression of tumorigenesis in Wilms' tumor. Hum. Molec. Genet. 90: 144-146, 1992.
- [93723] 16413.
- [93724] Weissman, B. E.; Saxon, P. J.; Pasquale, S. R.; Jones, G. R.; Geiser, A. G.; Stanbridge, E. J.: Introduction of a normal human chromosome 11 into a Wilms' tumor cell line controls its tumorigenic expression. Science 236: 175-180, 1987.
- [93725] 16414.
- [93726] Nishimoto, T.; Sekiguchi, T.; Kai, R.; Yamashita, K.; Takahashi, T.; Sekiguchi, M.: Large-scale selection and analysis of temperature-sensitive mutants for cell reproduction from BHK cells. Somat. Cell Genet. 8: 811-812, 1982.
- [93727] 16415.
- [93728] O'Brien, T.; Tjian, R.: Functional analysis of the human

TAFII250

[93736] N-terminal kinase domain. Molec. Cell 1: 905–911, 1998.

[93737] 16416.

[93738] Ruppert, S.; Wang, E. H.; Tjian, R.: Cloning and expression of

[93739] human TAF(II)250: a TBP-associated factor implicated in cell-cycle

[93740] regulation. Nature 362: 175–179, 1993.

[93741] 16417.

[93742] Schwartz, H. E.; Holmes, S.; Meiss, H. K.: Assignment of temperature-sensitive

[93743] mutations of BHK cells to the X-chromosome. (Abstract) J. Cell Biol. 75:

[93744] 393A, 1977.

[93745] 16418.

[93746] Schwartz, H. E.; Moser, G. C.; Holmes, S.; Meiss, H. K.: Assignment

[93747] of temperature-sensitive mutations of BHK cells to the X-chromosome. Somat.

[93748] Cell Genet. 5: 217–224, 1979.

[93749] 16419.

[93750] Sekiguchi, T.; Miyata, T.; Nishimoto, T.: Molecular cloning of
[93751] the cDNA of human X chromosomal gene (CCG1) which complements the
[93752] temperature-sensitive G(1) mutants, tsBN462 and ts13, of the BHK cell
[93753] line. EMBO J. 7: 1683–1687, 1988.
[93754] 16420.
[93755] Sekiguchi, T.; Nohiro, Y.; Nakamura, Y.; Hisamoto, N.; Nishimoto,
[93756] T.: The human CCG1 gene, essential for progression of the G1 phase,
[93757] encodes a 210-kilodalton nuclear DNA-binding protein. Molec. Cell.
[93758] Biol. 11: 3317–3325, 1991.
[93759] 16421.
[93760] Sekiguchi, T.; Yoshida, M. C.; Sekiguchi, M.; Nishimoto, T.:
[93761] Isolation of a human X chromosome-linked gene essential for progression
[93762] from G1 to S phase of the cell cycle. Exp. Cell Res. 169: 395–407,

[93763] 1987.

[93764] 16422.

[93765] Simchen, G.: Cell cycle mutants. Annu. Rev. Genet. 12:
161–191,

[93766] 1978.

[93767] 16423.

[93768] Slater, M. L.; Ozer, H. L.: Temperature-sensitive mutants
of

[93769] Balb–3T3 cells: description of a mutant affected in cellular
and polyoma

[93770] virus DNA synthesis. Cell 7: 289–295, 1976.

[93771] 16424.

[93772] Starr, D. B.; Hawley, D. K.: TFIID binds in the minor groove
[93773] of the TATA box. Cell 67: 1231–1240, 1991.

[93774] 16425.

[93775] Abel, A.; Walcott, J.; Woods, J.; Duda, J.; Merry, D. E.: Ex-
pression

[93776] of expanded repeat androgen receptor produces neuro-
logic disease in

[93777] transgenic mice. Hum. Molec. Genet. 10: 107–116, 2001.

[93778] 16426.

[93779] Bailey, C. K.; Andriola, I. F. M.; Kampinga, H. H.; Merry, D. E.

[93780] : Molecular chaperones enhance the degradation of expanded polyglutamine

[93781] repeat androgen receptor in a cellular model of spinal and bulbar

[93782] muscular atrophy. Hum. Molec. Genet. 11: 515–523, 2002.

[93783] 16427.

[93784] Batch, J. A.; Williams, D. M.; Davies, H. R.; Brown, B. D.; Evans,

[93785] B. A. J.; Hughes, I. A.; Patterson, M. N.: Androgen receptor gene

[93786] mutations identified by SSCP in fourteen subjects with androgen insensitivity

[93787] syndrome. Hum. Molec. Genet. 1: 497–503, 1992.

[93788] 16428.

[93789] Beitel, L. K.; Kazemi-Esfarjani, P.; Kaufman, M.; Lumbroso, R.;

[93790] DiGeorge, A. M.; Killinger, D. W.; Trifiro, M. A.; Pinsky, L.: Substitution

[93791] of arginine-839 by cysteine or histidine in the androgen

receptor

[93792] causes different receptor phenotypes in cultured cells and coordinate

[93793] degrees of clinical androgen resistance. J. Clin. Invest. 94: 546–554,

[93794] 1994.

[93795] 16429.

[93796] Belsham, D. D.; Pereira, F.; Greenberg, C. R.; Liao, S.; Wro-
gemann,

[93797] K.: Leu676-to-pro mutation of the androgen receptor
causes complete

[93798] androgen insensitivity syndrome in a large Hutterite kin-
dred. Hum.

[93799] Mutat. 5: 28–33, 1995.

[93800] 16430.

[93801] Bevan, C. L.; Brown, B. B.; Davies, H. R.; Evans, B. A. J.;
Hughes,

[93802] I. A.; Patterson, M. N.: Functional analysis of six androgen
receptor

[93803] mutations identified in patients with partial androgen in-
sensitivity

[93804] syndrome. Hum. Molec. Genet. 5: 265–273, 1996.

- [93805] 16431.
- [93806] Biancalana, V.; Serville, F.; Pommier, J.; Julien, J.; Hanauer, A.; Mandel, J. L.: Moderate instability of the trinucleotide repeat
- [93808] in spinobulbar muscular atrophy. Hum. Molec. Genet. 1: 255–258,
- [93809] 1992.
- [93810] 16432.
- [93811] Boehmer, A. L. M.; Brinkmann, A. O.; Niermeijer, M. F.; Bakker, L.; Halley, D. J. J.; Drop, S. L. S.: Germ–line and somatic mosaicism
- [93813] in the androgen insensitivity syndrome: implications for genetic counseling.
- [93814] (Letter) Am. J. Hum. Genet. 60: 1003–1006, 1997.
- [93815] 16433.
- [93816] Boehmer, A. L. M.; Brinkmann, A. O.; Nijman, R. M.; Verleun–Mooijman, M. C. T.; de Ruiter, P.; Niermeijer, M. F.; Drop, S. L. S.: Phenotypic
- [93818] variation in a family with partial androgen insensitivity

syndrome

[93819] explained by differences in 5- α dihydrotestosterone availability. J.

[93820] Clin. Endocr. Metab. 86: 1240–1246, 2001.

[93821] 16434.

[93822] Bouvattier, C.; Carel, J.-C.; Lecointre, C.; David, A.; Sultan,

[93823] C.; Bertrand, A.-M., Morel, Y.; Chaussain, J.-L.: Postnatal changes

[93824] of T, LH, and FSH in 46,XY infants with mutations in the AR gene. J.

[93825] Clin. Endocr. Metab. 87: 29–32, 2002.

[93826] 16435.

[93827] Brown, C. J.; Goss, S. J.; Lubahn, D. B.; Joseph, D. R.; Wilson,

[93828] E. M.; French, F. S.; Willard, H. F.: Androgen receptor locus on

[93829] the human X chromosome: regional localization to Xq11–12 and description

[93830] of a DNA polymorphism. Am. J. Hum. Genet. 44: 264–269, 1989.

[93831] 16436.

- [93832] Brown, T. R.; Lubahn, D. B.; Wilson, E. M.; Joseph, D. R.; French,
- [93833] F. S.; Migeon, C. J.: Deletion of the steroid-binding domain of the
- [93834] human androgen receptor gene in one family with complete androgen
- [93835] insensitivity syndrome: evidence for further genetic heterogeneity
- [93836] in this syndrome. *Proc. Nat. Acad. Sci.* 85: 8151-8155, 1988.
- [93837] 16437.
- [93838] Bruggenwirth, H. T.; Boehmer, A. L. M.; Ramnarain, S.; Verleun-Mooijman,
- [93839] M. C. T.; Satijn, D. P. E.; Trapman, J.; Grootegoed, J. A.; Brinkmann,
- [93840] A. O.: Molecular analysis of the androgen-receptor gene in a family
- [93841] with receptor-positive partial androgen insensitivity: an unusual
- [93842] type of intronic mutation. *Am. J. Hum. Genet.* 61: 1067-1077, 1997.
- [93843] 16438.

- [93844] Buchanan, G.; Yang, M.; Harris, J. M.; Nahm, H. S.; Han, G.; Moore,
- [93845] N.; Bentel, J. M.; Matusik, R. J.; Horsfall, D. J.; Marshall, V. R.;
- [93846] Greenberg, N. M.; Tilley, W. D.: Mutations at the boundary of the
- [93847] hinge and ligand binding domain of the androgen receptor confer increased
- [93848] transactivation function. *Molec. Endocr.* 15: 46–56, 2001.
- [93849] 16439.
- [93850] Butler, R.; Leigh, P. N.; McPhaul, M. J.; Gallo, J.–M.: Truncated
- [93851] forms of the androgen receptor are associated with polyglutamine expansion
- [93852] in X-linked spinal and bulbar muscular atrophy. *Hum. Molec. Genet.* 7:
- [93853] 121–127, 1998.
- [93854] 16440.
- [93855] Ho, M. F.; Monaco, A. P.; Blonden, L. A. J.; van Ommen, G. J.
- [93856] B.; Affara, N. A.; Ferguson-Smith, M. A.; Lehrach, H.: Fine mapping

- [93857] of the McLeod locus (XK) to a 150–380–kb region in Xp21.
Am. J. Hum.
- [93858] Genet. 50: 317–330, 1992.
- [93859] 16441.
- [93860] Jung, H. H.; Hergersberg, M.; Kneifel, S.; Alkadhi, H.;
Schiess,
- [93861] R.; Weigell–Weber, M.; Daniels, G.; Kollias, S.; Hess, K.:
McLeod
- [93862] syndrome: a novel mutation, predominant psychiatric
manifestations,
- [93863] and distinct striatal imaging findings. Ann. Neurol. 49:
384–392,
- [93864] 2001.
- [93865] 16442.
- [93866] Malandrini, A.; Fabrizi, G. M.; Truschi, F.; Di Pietro, G.;
Moschini,
- [93867] F.; Bartalucci, P.; Berti, G.; Salvadori, C.; Bucalossi, A.;
Guazzi,
- [93868] G.: Atypical McLeod syndrome manifested as X–linked
chorea–acanthocytosis,
- [93869] neuromyopathy and dilated cardiomyopathy: report of a
family. J.

[93870] Neurol. Sci. 124: 89–94, 1994.

[93871] 16443.

[93872] Marsh, W. L.: Personal Communication. New York, N. Y.
11/13/1979.

[93873] 16444.

[93874] Marsh, W. L.: The Kell blood groups and their relationship
to

[93875] chronic granulomatous disease. In: Antigens and Disease.
Am. Assoc.

[93876] Blood Banks Symposium. (pub.) 1977. Pp. 52–66.

[93877] 16445.

[93878] Marsh, W. L.: Linkage relationship of the Xg and Xk loci.
Cytogenet.

[93879] Cell Genet. 22: 531–533, 1978.

[93880] 16446.

[93881] Marsh, W. L.: Chronic granulomatous disease, the McLeod
syndrome,

[93882] and the Kell blood groups. Birth Defects Orig. Art. Ser.
XIV(6A):

[93883] 9–25, 1978.

- [93884] 16447.
- [93885] Marsh, W. L.: Chronic granulomatous disease, Kx antigen and the
- [93886] Kell blood groups. In: Brewer, G. J.: Progress in Clinical and Biological
- [93887] Research: The Red Cell. New York: Alan R. Liss (pub.) 1978. Pp.
- [93888] 493–507.
- [93889] 16448.
- [93890] Marsh, W. L.; Marsh, N. J.; Moore, A.; Symmans, W. A.; Johnson,
- [93891] C. L.; Redman, C. M.: Elevated serum creatine phosphokinase in subjects
- [93892] with McLeod syndrome. Vox Sang. 40: 403–411, 1981.
- [93893] 16449.
- [93894] Marsh, W. L.; Oyen, R.; Nichols, M. E.: Kx antigen, the McLeod
- [93895] phenotype, and chronic granulomatous disease: further studies. Vox
- [93896] Sang. 31: 356–362, 1976.
- [93897] 16450.

- [93898] Marsh, W. L.; Oyen, R.; Nichols, M. E.; Allen, F. H., Jr.:
Chronic
- [93899] granulomatous disease and the Kell blood groups. Brit. J.
Haemat. 29:
- [93900] 247–262, 1975.
- [93901] 16451.
- [93902] Schwartz, S. A.; Marsh, W. L.; Symmans, A.; et al: 'New'
clinical
- [93903] features of McLeod syndrome. (Abstract) Transfusion 22:
404 only,
- [93904] 1982.
- [93905] 16452.
- [93906] Stanfield, G. M.; Horvitz, H. R.: The ced-8 gene controls
the
- [93907] timing of programmed cell deaths in *C. elegans*. Molec.
Cell 5: 423–433,
- [93908] 2000.
- [93909] 16453.
- [93910] Supple, S. G.; Iland, H. J.; Barnett, M. H.; Pollard, J. D.:
- [93911] A spontaneous novel XK gene mutation in a patient with
McLeod syndrome. Brit.

- [93912] J. Haemat. 115: 369–372, 2001.
- [93913] 16454.
- [93914] Swash, M.; Schwartz, M. S.; Carter, N. D.; Heath, R.; Leak, M.;
- [93915] Rogers, K. L.: Benign X-linked myopathy with acanthocytes (McLeod
- [93916] syndrome): its relationship to X-linked muscular dystrophy. Brain 106:
- [93917] 717–733, 1983.
- [93918] 16455.
- [93919] Symmans, W. A.; Sheperd, C. S.; Marsh, W. L.; Oyen, R.; Shohet,
- [93920] S. B.; Linehan, B. J.: Hereditary acanthocytosis associated with
- [93921] the McLeod phenotype of the Kell blood group system. Brit. J. Haemat. 42:
- [93922] 575–583, 1979.
- [93923] 16456.
- [93924] Tippet, P.: Personal Communication. London, England 7/1981.
- [93925] 16457.

- [93926] Wimer, B. M.; Marsh, W. L.; Taswell, H. F.: Clinical characteristics
- [93927] of the McLeod blood group phenotype. (Abstract) Am. Soc. Hemat.,
- [93928] Boston , 12/1976.
- [93929] 16458.
- [93930] Wimer, B. M.; Marsh, W. L.; Taswell, H. F.; Galey, W. R.: Haematological
- [93931] changes associated with the McLeod phenotype of the Kell blood group
- [93932] system. Brit. J. Haemat. 36: 219–224, 1977.
- [93933] 16459.
- [93934] Adler, D. A.; Bressler, S. L.; Chapman, V. M.; Page, D. C.; Disteche,
- [93935] C. M.: Inactivation of the Zfx gene on the mouse X chromosome. Proc.
- [93936] Nat. Acad. Sci. 88: 4592–4595, 1991.
- [93937] 16460.
- [93938] Chong, S. S.; Kristjansson, K.; Cota, J.; Handyside, A. H.; Hughes,
- [93939] M. R.: Preimplantation prevention of X-linked disease: re-

liable and

[93940] rapid sex determination of single human cells by restriction analysis

[93941] of simultaneously amplified ZFX and ZFY sequences. Hum. Molec. Genet. 2:

[93942] 1187-1191, 1993.

[93943] 16461.

[93944] Luoh, S.-W.; Jegalian, K.; Lee, A.; Chen, E. Y.; Ridley, A.; Page,

[93945] D. C.: CpG islands in human ZFX and ZFY and mouse Zfx genes: sequence

[93946] similarities and methylation differences. Genomics 29: 353-363,

[93947] 1995.

[93948] 16462.

[93949] Muller, G.; Schempp, W.: Mapping the human ZFX locus to Xp21.3

[93950] by in situ hybridization. Hum. Genet. 82: 82-84, 1989.

[93951] 16463.

[93952] Page, D. C.; Disteche, C. M.; Simpson, E. M.; de la Chapelle, A.;

- [93953] Andersson, M.; Alitalo, T.; Brown, L. G.; Green, P.; Akots, G.: Chromosomal
- [93954] localization of ZFX--a human gene that escapes X inactivation--and
- [93955] its murine homologs. *Genomics* 7: 37-46, 1990.
- [93956] 16464.
- [93957] Palmer, M. S.; Berta, P.; Sinclair, A. H.; Pym, B.; Goodfellow,
- [93958] P. N.: Comparison of human ZFY and ZFX transcripts. *Proc. Nat.*
- [93959] *Acad. Sci.* 87: 1681-1685, 1990.
- [93960] 16465.
- [93961] Palmer, M. S.; Sinclair, A. H.; Berta, P.; Ellis, N. A.; Goodfellow,
- [93962] P. N.; Abbas, N. E.; Fellous, M.: Genetic evidence that ZFY is not
- [93963] the testis-determining factor. *Nature* 342: 937-939, 1989.
- [93964] 16466.
- [93965] Scherer, G.; Schempp, W.; Baccichetti, C.; Lenzini, E.; Dagna Bricarelli,

- [93966] F.; Carbone, L. D. L.; Wolf, U.: Duplication of an Xp segment that
- [93967] includes the ZFX locus causes sex inversion in man. Hum. Genet. 81:
- [93968] 291–294, 1989.
- [93969] 16467.
- [93970] Schneider–Gadicke, A.; Beer–Romero, P.; Brown, L. G.; Nussbaum,
- [93971] R.; Page, D. C.: ZFX has a gene structure similar to ZFY, the putative
- [93972] human sex determinant, and escapes X inactivation. Cell 57: 1247–1258,
- [93973] 1989.
- [93974] 16468.
- [93975] Seibold, A.; Brabet, P.; Rosenthal, W.; Birnbaumer, M.: Structure
- [93976] and chromosomal localization of the human antidiuretic hormone receptor
- [93977] gene. Am. J. Hum. Genet. 51: 1078–1083, 1992.
- [93978] 16469.
- [93979] Trousseau, A.: Polydipsia. Lecture LXV. In: Clinique Medi–

cale

[93980] de L'Hotel-Dieu de Paris. Vol. 2. Paris: J.-B. Balliere 1862.

[93981] P. 607.

[93982] 16470.

[93983] Tsukaguchi, H.; Matsubara, H.; Taketani, S.; Mori, Y.;
Seido,

[93984] T.; Inada, M.: Binding-, intracellular transport-, and
biosynthesis-defective

[93985] mutants of vasopressin type 2 receptor in patients with X-
linked nephrogenic

[93986] diabetes insipidus. J. Clin. Invest. 96: 2043-2050, 1995.

[93987] 16471.

[93988] Uttley, W. S.; Thistlethwaite, D.: Failure to detect the car-
rier

[93989] in congenital nephrogenic diabetes insipidus. Arch. Dis.
Child. 47:

[93990] 137-138, 1972.

[93991] 16472.

[93992] van den Ouweland, A.; Markslag, P.; Knoers, N.; Ropers, H.
H.;

[93993] Poustka, A.; Jans, D.; Fahrenholz, F.; van Oost, B.: Colocal-

ization

[93994] of the locus for human nephrogenic diabetes insipidus and the vasopressin

[93995] type-2 receptor gene in the Xq28 region. (Series) Miami Short

[93996] Reports. Advances in Gene Technology: The Molecular Biology of Human

[93997] Genetic Disease. New York: IRL Press (pub.) 1: 1991. Pp. 20

[93998] only.

[93999] 16473.

[94000] van den Ouweland, A. M. W.; Dreesen, J. C. F. M.; Verdijk, M.;

[94001] Knoers, N. V. A. M.; Monnens, L. A. H.; Rocchi, M.; van Oost, B. A.

[94002] : Mutations in the vasopressin type 2 receptor gene (AVPR2) associated

[94003] with nephrogenic diabetes insipidus. Nature Genet. 2: 99-102, 1992.

[94004] 16474.

[94005] van den Ouweland, A. M. W.; Knoop, M. T.; Knoers, V. V. A. M.;

- [94006] Markslag, P. W. B.; Rocchi, M.; Warren, S. T.; Ropers, H. H.;
Fahrenholz,
- [94007] F.; Monnens, L. A.; van Oost, B. A. H.: Colocalization of the
gene
- [94008] for nephrogenic diabetes insipidus (DIR) and the vaso-
pressin type
- [94009] 2 receptor gene (AVPR2) in the Xq28 region. Genomics 13:
1350–1352,
- [94010] 1992.
- [94011] 16475.
- [94012] van Lieburg, A. F.; Knoers, N. V. A. M.; Monnens, L. A. H.:
Clinical
- [94013] presentation and follow-up of 30 patients with congenital
nephrogenic
- [94014] diabetes insipidus. J. Am. Soc. Nephrol. 10: 1958–1964,
1999.
- [94015] 16476.
- [94016] van Lieburg, A. F.; Verdijk, M. A. J.; Schoute, F.; Ligten-
berg,
- [94017] M. J. L.; van Oost, B. A.; Waldhauser, F.; Dobner, M.; Mon-
nens, L.
- [94018] A. H.; Knoers, N. V. A. M.: Clinical phenotype of nephro-

genic diabetes

[94019] insipidus in females heterozygous for a vasopressin type
2 receptor

[94020] mutation. Hum. Genet. 96: 70–78, 1995.

[94021] 16477.

[94022] Wildin, R. S.; Antush, M. J.; Bennett, R. L.; Schoof, J. M.;
Scott,

[94023] C. R.: Heterogeneous AVPR2 gene mutations in congenital
nephrogenic

[94024] diabetes insipidus. Am. J. Hum. Genet. 55: 266–277,
1994.

[94025] 16478.

[94026] Hata, A.; Matsuura, T.; Setoyama, C.; Shimada, K.; Yokoi,
T.;

[94027] Akaboshi, I.; Matsuda, I.: A novel missense mutation in
exon 8 of

[94028] the ornithine transcarbamylase gene in two unrelated
male patients

[94029] with mild ornithine transcarbamylase deficiency. Hum.
Genet. 87:

[94030] 28–32, 1991.

[94031] 16479.

- [94032] Masutani, C.; Sugasawa, K.; Yanagisawa, J.; Sonoyama, T.;
Ui, M.;
- [94033] Enomoto, T.; Takio, K.; Tanaka, K.; van der Spek, P. J.;
Bootsma,
- [94034] D.; Hoeijmakers, J. H. J.; Hanaoka, F.: Purification and
cloning
- [94035] of a nucleotide excision repair complex involving the xe-
roderma pigmentosum
- [94036] group C protein and a human homologue of yeast RAD23.
EMBO J. 13:
- [94037] 1831-1843, 1994.
- [94038] 16480.
- [94039] Angelini, C.; Fanin, M.; Menegazzo, E.; Freda, M. P.; Dug-
gan, D.
- [94040] J.; Hoffman, E. P.: Homozygous alpha-sarcoglycan muta-
tion in two
- [94041] siblings: one asymptomatic and one steroid-responsive
mild limb-girdle
- [94042] muscular dystrophy patient. Muscle Nerve 21: 769-775,
1998.
- [94043] 16481.
- [94044] Derst, C.; Konrad, M.; Kockerling, A.; Karolyi, L.; Desch-

enes,

[94045] G.; Daut, J.; Karschin, A.; Seyberth, H. W.: Mutations in the ROMK

[94046] gene in antenatal Bartter syndrome are associated with impaired K(+)

[94047] channel function. Biochem. Biophys. Res. Commun. 203: 641–645, 1997.

[94048] 16482.

[94049] Derst, C.; Wischmeyer, E.; Preisig–Muller, R.; Spauschus, A.; Konrad,

[94050] M.; Hensen, P.; Jeck, N.; Seyberth, H. W.; Daut, J.; Karschin, A.

[94051] : A hyperprostaglandin E syndrome mutation in Kir1.1 (renal outer

[94052] medullary potassium) channels reveals a crucial residue for channel

[94053] function in Kir1.3 channels. J. Biol. Chem. 273: 23884–23891, 1998.

[94054] 16483.

[94055] Ho, K.; Nichols, C. G.; Lederer, W. J.; Lytton, J.; Vassilev, P.

[94056] M.; Kanazirska, M. V.; Hebert, S. C.: Cloning and expression of an

[94057] inwardly rectifying ATP-regulated potassium channel. Nature 362:

[94058] 31-38, 1993.

[94059] 16484.

[94060] International Collaborative Study Group for Bartter-like Syndromes

[94061] : Mutations in the gene encoding the inwardly-rectifying renal potassium

[94062] channel, ROMK, cause the antenatal variant of Bartter syndrome: evidence

[94063] for genetic heterogeneity. Hum. Molec. Genet. 6: 17-26, 1997.

[94064] 16485.

[94065] Krishnan, S. N.; Desai, T.; Ward, D. C.; Haddad, G. G.: Isolation

[94066] and chromosomal localization of a human ATP-regulated potassium channel. Hum.

[94067] Genet. 96: 155-160, 1995.

[94068] 16486.

[94069] Shuck, M. E.; Bock, J. H.; Benjamin, C. W.; Tsai, T.-D.; Lee, K.

- [94070] S.; Slightom, J. L.; Bienkowski, M. J.: Cloning and characterization
of multiple forms of the human kidney ROM-K potassium
channel. J.
- [94072] Biol. Chem. 269: 24261–24270, 1994.
- [94073] 16487.
- [94074] Simon, D. B.; Karet, F. E.; Rodriguez-Soriano, J.; Hamdan,
J. H.;
- [94075] DiPietro, A.; Trachtman, H.; Sanjad, S. A.; Lifton, R. P.: Ge-
netic
heterogeneity of Bartter's syndrome revealed by mutations
in the K(+)
channel, ROMK. Nature Genet. 14: 152–156, 1996.
- [94078] 16488.
- [94079] Yano, H.; Philipson, L. H.; Kugler, J. L.; Tokuyama, Y.;
Davis,
E. M.; Le Beau, M. M.; Nelson, D. J.; Bell, G. I.; Takeda, J.:
Alternative
splicing of human inwardly rectifying K(+) channel ROMK1
mRNA. Molec.
- [94082] Pharm. 45: 854–860, 1994.

- [94083] 16489.
- [94084] Campbell, H. D.; Fountain, S.; McLennan, I. S.; Berven, L. A.;
- [94085] Crouch, M. F.; Davy, D. A.; Hooper, J. A.; Waterford, K.; Chen, K.-S.;
- [94086] Lupski, J. R.; Ledermann, B.; Young, I. G.; Matthaei, K. I.: Fliih,
- [94087] a gelsolin-related cytoskeletal regulator essential for early mammalian
- [94088] embryonic development. *Molec. Cell. Biol.* 22: 3518–3526, 2002.
- [94089] 16490.
- [94090] Campbell, H. D.; Fountain, S.; Young, I. G.; Claudianos, C.; Hoheisel,
- [94091] J. D.; Chen, K.-S.; Lupski, J. R.: Genomic structure, evolution,
- [94092] and expression of human FLII, a gelsolin and leucine-rich-repeat family
- [94093] member: overlap with LLGL. *Genomics* 42: 46–54, 1997.
- [94094] 16491.
- [94095] Connelly, M. A.; Zhang, H.; Kieleczawa, J.; Anderson, C. W.: The

- [94096] promoters for human DNA-PK(cs) (PRKDC) and MCM4: divergently transcribed
- [94097] genes located at chromosome 8 band q11. Genomics 47: 71–83, 1998.
- [94098] 16492.
- [94099] Aly, A. M.; Hoyer, L. W.: Factor VIII–East Hartford (arginine1689
- [94100] to cysteine) has procoagulant activity when separated from von Willebrand
- [94101] factor. J. Clin. Invest. 89: 1382–1387, 1992.
- [94102] 16493.
- [94103] Antonarakis, S. E.; Copeland, K. L.; Carpenter, R. J., Jr.; Carta,
- [94104] C. A.; Hoyer, L. W.; Caskey, C. T.; Toole, J. J.; Kazazian, H. H.,
- [94105] Jr.: Prenatal diagnosis of haemophilia A by factor VIII gene analysis. Lancet I:
- [94106] 1407–1409, 1985.
- [94107] 16494.
- [94108] Antonarakis, S. E.; Kazazian, H. H.; Tuddenham, G. D.: Molecular

- [94109] etiology of factor VIII deficiency in hemophilia A. Hum. Mutat. 5: 1-22, 1995.
- [94110] 16495.
- [94111] Antonarakis, S. E.; Rossiter, J. P.; Young, M.; Horst, J.; de Moerloose, P.; Sommer, S. S.; Ketterling, R. P.; Kazazian, H. H., Jr.; Ne-grier, C.; Vinciguerra, C.; Gitschier, J.; Goossens, M.; and 54 others : Factor VIII inversions in severe hemophilia A: results from an international consortium. Blood 86: 2206-2212, 1995.
- [94112] 16496.
- [94113] Antonarakis, S. E.; Waber, P. G.; Kittur, S. D.; Patel, A. S.; Kazazian, H. H., Jr.; Mellis, M. A.; Counts, R. B.; Stamatoy-annopoulos, G.; Bowie, E. J. W.; Fass, D. N.; Pittman, D. D.; Wozney, J. M.; Toole, J. J.: Hemophilia A: detection of molecular defects and of carriers

- [94122] by DNA analysis. New Eng. J. Med. 313: 842–848, 1985.
- [94123] 16497.
- [94124] Antonarakis, S. E.; Youssoufian, H.; Kazazian, H. H.:
Molecular
- [94125] genetics of hemophilia–A in man (factor VIII deficiency).
Molec.
- [94126] Biol. Med. 4: 81–95, 1987.
- [94127] 16498.
- [94128] Arai, M.; Higuchi, M.; Antonarakis, S. E.; Kazazian, H. H.,
Jr.;
- [94129] Phillips, J. A., III; Janco, R. L.; Hoyer, L. W.: Characteriza-
tion
- [94130] of a thrombin cleavage site mutation (arg1689–to–cys) in
the factor
- [94131] VIII gene of two unrelated patients with cross–reacting
material–positive
- [94132] hemophilia A. Blood 75: 384–389, 1990.
- [94133] 16499.
- [94134] Arai, M.; Inaba, H.; Higuchi, M.; Antonarakis, S. E.;
Kazazian,
- [94135] H. H., Jr.; Fujimaki, M.; Hoyer, L. W.: Direct characteriza–

tion of

[94136] factor VIII in plasma: detection of a mutation altering a
thrombin

[94137] cleavage site (arginine372-to-histidine). Proc. Nat. Acad.
Sci. 86:

[94138] 4277-4281, 1989.

[94139] 16500.

[94140] Arrants, J. E.; Jordan, P. H., Jr.; Newcomb, T. F.: Von Wille-
brand's

[94141] disease: a cause for massive postoperative bleeding-
report of a case. Ann.

[94142] Surg. 156: 845-851, 1962.

[94143] 16501.

[94144] Arveiler, B.; Vincent, A.; Mandel, J.-L.: Toward a physical
map

[94145] of the Xq28 region in man: linking color vision, G6PD, and
coagulation

[94146] factor VIII genes to an X-Y homology region. Genomics 4:
460-471,

[94147] 1989.

[94148] 16502.

- [94149] Bardoni, B.; Sampietro, M.; Romano, M.; Crapanzano, M.; Mannucci,
[94150] P. M.; Camerino, G.: Characterization of a partial deletion
of the
[94151] factor VIII gene in a haemophiliac with inhibitor. Hum.
Genet. 79:
[94152] 86–88, 1988.
[94153] 16503.
[94154] Barker, D.; Schafer, M.; White, R.: Restriction sites con-
taining
[94155] CpG show a higher frequency of polymorphism in human
DNA. Cell 36:
[94156] 131–138, 1984.
[94157] 16504.
[94158] Barrai, I.; Cann, H. M.; Cavalli-Sforza, L. L.: Segregation
analysis
[94159] of hemophilia A and B. (Letter) Am. J. Hum. Genet. 31:
226–227,
[94160] 1979.
[94161] 16505.
[94162] Barrai, I.; Cann, H. M.; Cavalli-Sforza, L. L.; de Nicola, P.

- [94163] : The effect of parental age on rates of mutation for hemophilia and
- [94164] evidence for differing mutation rates for hemophilia A and B. Am.
- [94165] J. Hum. Genet. 20: 175–196, 1968.
- [94166] 16506.
- [94167] Barrow, E. M. S.; Graham, J. B.: Factor VIII. (Letter) Lancet I:
- [94168] 1312–1313, 1973.
- [94169] 16507.
- [94170] Baty, B. J.; Drayna, D.; Leonard, C. O.; White, R.: Prenatal
- [94171] diagnosis of factor VIII deficiency to help with the management of
- [94172] pregnancy and delivery. (Letter) Lancet I: 207 only, 1986.
- [94173] 16508.
- [94174] Becker, J.; Schwaab, R.; Moller-Taube, A.; Schwaab, U.; Schmidt,
- [94175] W.; Brackmann, H. H.; Grimm, T.; Olek, K.; Oldenburg, J.: Characterization
- [94176] of the factor VIII defect in 147 patients with sporadic hemophilia

- [94177] A: family studies indicate a mutation type-dependent sex ratio of
- [94178] mutation frequencies. *Am. J. Hum. Genet.* 58: 657-670, 1996.
- [94179] 16509.
- [94180] Bennett, B.; Ratnoff, O. D.: Deletion of the carrier state for
- [94181] classic hemophilia. *New Eng. J. Med.* 7: 342-345, 1974.
- [94182] 16510.
- [94183] Bennett, E.; Huehns, E. R.: Immunological differentiation of
- [94184] three types of hemophilia and identification of some female carriers. *Lancet* II:
- [94185] 956-958, 1970.
- [94186] 16511.
- [94187] Berg, L.-P.; Wieland, K.; Millar, D. S.; Schlosser, M.; Wagner,
- [94188] M.; Kakkar, V. V.; Reiss, J.; Cooper, D. N.: Detection of a novel
- [94189] point mutation causing haemophilia A by PCR/direct sequencing of ectopically-transcribed
- [94190] factor VIII mRNA. *Hum. Genet.* 85: 655-658, 1990.

- [94191] 16512.
- [94192] Bernardi, F.; Marchetti, G.; Bertagnolo, V.; Faggioli, L.;
Volinia,
- [94193] S.; Patracchini, P.; Bartolai, S.; Vannini, F.; Felloni, L.;
Rossi,
- [94194] L.; Panicucci, F.; Conconi, F.: RFLP analysis in families with
sporadic
- [94195] hemophilia A: estimate of the mutation ratio in male and
female gametes. Hum.
- [94196] Genet. 76: 253–256, 1987.
- [94197] 16513.
- [94198] Bernardi, F.; Volinia, S.; Patracchini, P.; Gemmati, D.;
Boninsegna,
- [94199] S.; Schwienbacher, C.; Marchetti, G.: A recurrent missense
mutation
- [94200] (arg-to-gln) and a partial deletion in factor VIII gene caus-
ing severe
- [94201] haemophilia A. Brit. J. Haemat. 71: 271–276, 1989.
- [94202] 16514.
- [94203] Bidichandani, S. I.; Shiach, C. R.; Lanyon, W. G.; Connor, J.
- [94204] M.: A novel splice donor mutation affecting position +3 in

intron

- [94205] 6 of the factor VIII gene. Hum. Molec. Genet. 3: 651–653, 1994.
- [94206] 16515.
- [94207] Biggs, R.; Rizza, C. R.: The sporadic case of haemophilia A. Lancet II:
- [94208] 431–433, 1976.
- [94209] 16516.
- [94210] Bird, A. P.: DNA methylation and the frequency of CpG in animal
- [94211] DNA. Nucleic Acids Res. 8: 1499–1504, 1980.
- [94212] 16517.
- [94213] Bloom, A. L.; Peake, I. R.: Molecular genetics of factor VIII
- [94214] and its disorders. Am. J. Path. 88: 319–340, 1977.
- [94215] 16518.
- [94216] Bond, T. P.; Levin, W. C.; Celandier, D. R.; Guest, M. M.: 'Mild
- [94217] hemophilia' affecting both males and females. New Eng. J. Med. 266:
- [94218] 220–223, 1962.

- [94219] 16519.
- [94220] Boyer, S. H.; Siggers, D. C.; Krueger, L. J.: A caveat to protein
- [94221] replacement therapy for genetic disease. Immunologic implications
- [94222] of accurate molecular diagnosis. Lancet II: 654-659, 1973.
- [94223] 16520.
- [94224] Briet, E.; Bakker, B.; Brocker-Vriends, A.: Somatic and germinal
- [94225] mosaicism for a deletion in the factor VIII gene. (Abstract) Brit.
- [94226] J. Haemat. 71 (suppl. 1): 1 only, 1989.
- [94227] 16521.
- [94228] Greenough, W. T.; Klintsova, A. Y.; Irwin, S. A.; Galvez, R.;
- [94229] Bates, K. E.; Weiler, I. J.: Synaptic regulation of protein synthesis
- [94230] and the fragile X protein. Proc. Nat. Acad. Sci. 98: 7101-7106,
- [94231] 2001.
- [94232] 16522.

- [94233] Griffiths, M. J.; Strachan, M. C.: A single lymphocyte culture
- [94234] for fragile X induction and prometaphase chromosome analysis. J.
- [94235] Med. Genet. 28: 837–839, 1991.
- [94236] 16523.
- [94237] Haataja, R.; Vaisanen, M.–L.; Li, M.; Ryyananen, M.; Leisti, J.
- [94238] : The fragile X syndrome in Finland: demonstration of a founder effect
- [94239] by analysis of microsatellite haplotypes. Hum. Genet. 94: 479–483,
- [94240] 1994.
- [94241] 16524.
- [94242] Hagerman, R. J.; Leehey, M.; Heinrichs, W.; Tassone, F.; Wilson,
- [94243] R.; Hills, J.; Grigsby, J.; Gage, B.; Hagerman, P. J.: Intention
- [94244] tremor, parkinsonism, and generalized brain atrophy in male carriers
- [94245] of fragile X. Neurology 57: 127–130, 2001.
- [94246] 16525.

- [94247] Hagerman, R. J.; Synhorst, D. P.: Mitral valve prolapse and aortic
- [94248] dilatation in the fragile X syndrome. Am. J. Med. Genet. 17: 123–131,
- [94249] 1984.
- [94250] 16526.
- [94251] Hagerman, R. J.; Van Housen, K.; Smith, A. C. M.; McGavran, L.
- [94252] : Consideration of connective tissue dysfunction in the fragile X
- [94253] syndrome. Am. J. Med. Genet. 17: 111–121, 1984.
- [94254] 16527.
- [94255] Hanzlik, A. J.; Osemlak-Hanzlik, M. M.; Hauser, M. A.; Kurnit,
- [94256] D. M.: A recombination-based assay demonstrates that the fragile
- [94257] X sequence is transcribed widely during development. Nature Genet. 3:
- [94258] 44–48, 1993.
- [94259] 16528.
- [94260] Harvey, J.; Judge, C.; Wiener, S.: Familial X-linked mental

retardation

[94261] with an X chromosome abnormality. J. Med. Genet. 14: 46-50, 1977.

[94262] 16529.

[94263] Hecht, F.; Cannizzaro, L. A.; Hecht, B. K.: Gene symbols for

[94264] fragile sites: a proposal. (Letter) Hum. Genet. 82: 394, 1989.

[94265] 16530.

[94266] Hecht, F.; Glover, T. W.: Antibiotics containing trimethoprim

[94267] and the fragile X chromosome. (Letter) New Eng. J. Med. 308: 285-286,

[94268] 1983.

[94269] 16531.

[94270] Hecht, F.; Jacky, P. B.; Sutherland, G. R.: The fragile X chromosome:

[94271] current methods. Am. J. Med. Genet. 11: 489-495, 1982.

[94272] 16532.

[94273] Hinds, H. L.; Ashley, C. T.; Sutcliffe, J. S.; Nelson, D. L.;

[94274] Warren, S. T.; Housman, D. E.; Schalling, M.: Tissue spe-

cific expression

[94275] of FMR-1 provides evidence for a functional role in fragile X syndrome. *Nature*

[94276] *Genet.* 3: 36-43, 1993.

[94277] 16533.

[94278] Hirst, M.; Grewal, P.; Flannery, A.; Slatter, R.; Maher, E.; Barton,

[94279] D.; Fryns, J.-P.; Davies, K.: Two new cases of FMR1 deletion associated

[94280] with mental impairment. *Am. J. Hum. Genet.* 56: 67-74, 1995.

[94281] 16534.

[94282] Hirst, M. C.; Knight, S. J. L.; Christodoulou, Z.; Grewal, P.

[94283] K.; Fryns, J. P.; Davies, K. E.: Origins of the fragile X syndrome

[94284] mutation. *J. Med. Genet.* 30: 647-650, 1993.

[94285] 16535.

[94286] Hofker, M. H.; Bergen, A. A. B.; Skraastad, M. I.; Carpenter,

[94287] N. J.; Veenema, H.; Connor, J. M.; Bakker, E.; van Ommen, G. J. B.;

[94288] Pearson, P. L.: Efficient isolation of X chromosome-spe-

cific single

[94289] copy probes from a cosmid library of a human X/hamster hybrid-cell

[94290] line: mapping of new probes close to the locus for X-linked mental

[94291] retardation. Am. J. Hum. Genet. 40: 312-328, 1987.

[94292] 16536.

[94293] Howard-Peebles, P. N.; Stoddard, G. R.: Familial X-linked mental

[94294] retardation with a marker X chromosome and its relationship to macroorchidism. Clin.

[94295] Genet. 17: 125-128, 1980.

[94296] 16537.

[94297] Hundscheid, R. D. L.; Sistermans, E. A.; Thomas, C. M. G.; Braat,

[94298] D. D. M.; Straatman, H.; Kiemeney, L. A. L. M.; Oostra, B. A.; Smits,

[94299] A. P. T.: Imprinting effect in premature ovarian failure confined

[94300] to paternally inherited fragile X premutations. Am. J. Hum. Genet. 66:

[94301] 413-418, 2000.

[94302] 16538.

[94303] Hundscheid, R. D. L.; Thomas, C. M. G.; Braat, D. D. M.;
Oostra,

[94304] B. A.; Smits, A. P. T.: Reply to the letters from Murray et al.
and

[94305] Vianna-Morgante and Costa. (Letter) Am. J. Hum. Genet.
67: 256-258,

[94306] 2000.

[94307] 16539.

[94308] ISCN: An International System for Human Cytogenetic
Nomenclature

[94309] (1978). Cytogenet. Cell Genet. 21: 309-404, 1978.

[94310] 16540.

[94311] Jacky, P. B.; Dill, F. J.: Expression in fibroblast culture of

[94312] the satellited-X chromosome associated with familial sex-
linked mental

[94313] retardation. Hum. Genet. 53: 267-269, 1980.

[94314] 16541.

[94315] Jacobs, P. A.: Personal Communication. Honolulu, Hawaii
1982.

- [94316] 16542.
- [94317] Jacobs, P. A.: The fragile X syndrome. (Editorial) J. Med. Genet. 28:
[94318] 809–810, 1991.
- [94319] 16543.
- [94320] Jacobs, P. A.; Glover, T. W.; Mayer, M.; Fox, P.; Gerrard, J.
[94321] W.; Dunn, H. G.; Herbst, D. S.: X-linked mental retardation: a study
[94322] of 7 families. Am. J. Med. Genet. 7: 471–479, 1980.
- [94323] 16544.
- [94324] Jacobs, P. A.; Hunt, P. A.; Mayer, M.; Wang, J.-C.; Boss, G.
R.;
- [94325] Erbe, R. W.: Expression of the marker(X)(q28) in lymphoblastoid cell
[94326] lines. Am. J. Hum. Genet. 34: 552–557, 1982.
- [94327] 16545.
- [94328] Jakala, P.; Hanninen, T.; Ryyanen, M.; Laakso, M.; Partanen,
[94329] K.; Mannermaa, A.; Soininen, H.: Fragile-X: neuropsychological test

[94330] performance, CGG triplet repeat lengths, and hippocampal volumes. J.

[94331] Clin. Invest. 100: 331–338, 1997.

[94332] 16546.

[94333] Jenkins, E. C.; Brown, W. T.; Brooks, J.; Duncan, C. J.; Rudelli,

[94334] R. D.; Wisniewski, H. M.: Experience with prenatal fragile X detection. Am.

[94335] J. Med. Genet. 17: 215–239, 1984.

[94336] 16547.

[94337] Jenkins, E. C.; Brown, W. T.; Duncan, C. J.; Brooks, J.: Demonstration

[94338] of the fragile X chromosome in amniotic fluid cells. (Abstract) Clin.

[94339] Res. 30: 292A, 1982.

[94340] 16548.

[94341] Jin, P.; Warren, S. T.: Understanding the molecular basis of

[94342] fragile X syndrome. Hum. Molec. Genet. 9: 901–908, 2000.

[94343]

[94344] 100. Johnson, V. P.; Carpenter, N. J.; Skorey, P. A.: Martin-Bell

[94345] syndrome segregating in a large kindred with normal transmitting males:

[94346] clinical, cytogenetic, and linkage study. Am. J. Med. Genet. 38:

[94347] 275-282, 1991.

[94348]

[94349] 101. Kaiser-McCaw, B.; Hecht, F.; Cadien, J. D.; Moore, B. C.: Fragile

[94350] X-linked mental retardation. Am. J. Hum. Genet. 7: 503-505, 1980.

[94351]

[94352] 102. Kenneson, A.; Zhang, F.; Hagedorn, C. H.; Warren, S. T.: Reduced

[94353] FMRP and increased FMR1 transcription is proportionally associated

[94354] with CGG repeat number in intermediate-length and pre-mutation carriers. Hum.

[94355] Molec. Genet. 10: 1449-1454, 2001.

[94356]

- [94357] 103. Khalifa, M. M.; Reiss, A. L.; Migeon, B. R.: Methylation status
[94358] of genes flanking the fragile site in males with the fragile-X syndrome:
[94359] a test of the imprinting hypothesis. Am. J. Hum. Genet. 46: 744-753,
[94360] 1990.
[94361]
[94362] 104. Khandjian, E. W.; Corbin, F.; Woerly, S.; Rousseau, F.: The
[94363] fragile X mental retardation protein is associated with ribosomes. Nature
[94364] Genet. 12: 91-93, 1996.
[94365]
[94366] 105. Kinnell, H. G.: Fragile-X disorder associated with antisocial
[94367] personality. (Letter) Lancet II: 1104, 1982.
[94368]
[94369] 106. Kirchgessner, C. U.; Warren, S. T.; Willard, H. F.: X in-activation
[94370] if the FMR1 fragile X mental retardation gene. J. Med.

Genet. 32:

[94371] 925–929, 1995.

[94372]

[94373] 107. Kirkilionis, A. J.; Chudley, A. E.; Greenberg, C. R.; Yan, D.

[94374] L.; McGillivray, B.; Hamerton, J. L.: Transmission of the fra(X)

[94375] haplotype from three nonpenetrant brothers to their affected grandsons. Am.

[94376] J. Med. Genet. 43: 588–591, 1992.

[94377]

[94378] 108. Klauck, S. M.; Munstermann, E.; Bieber–Martig, B.; Ruhl, D.;

[94379] Lisch, S.; Schmotzer, G.; Poustka, A.; Poustka, F.: Molecular genetic

[94380] analysis of the FRM–1 gene in a large collection of autistic patients. Hum.

[94381] Genet. 100: 224–229, 1997.

[94382]

[94383] 109. Knight, S. J. L.; Flannery, A. V.; Hirst, M. C.; Campbell, L.;

- [94384] Christodoulou, Z.; Phelps, S. R.; Pointon, J.; Middleton-Price, H.
- [94385] R.; Barnicoat, A.; Pembrey, M. E.; Holland, J.; Oostra, B. A.; Bobrow,
- [94386] M.; Davies, K. E.: Trinucleotide repeat amplification and hypermethylation
- [94387] of a CpG island in FRAXE mental retardation. Cell 74: 127-134, 1993.
- [94388]
- [94389] 110. Kooy, R. F.; Reyniers, E.; Verhoye, M.; Sijbers, J.; Bakker,
- [94390] C. E.; Oostra, B. A.; Willems, P. J.; Van Der Linden, A.: Neuroanatomy
- [94391] of the fragile X knockout mouse brain studied using in vivo high resolution
- [94392] magnetic resonance imaging. Europ. J. Hum. Genet. 7: 526-532, 1999.
- [94393]
- [94394] 111. Krawczun, M. S.; Jenkins, E. C.; Brown, W. T.: Analysis of the
- [94395] fragile-X chromosome: localization and detection of the fragile site

[94396] in high resolution preparations. Hum. Genet. 69:
209–211, 1985.

[94397]

[94398] 112. Kruyer, H.; Mila, M.; Glover, G.; Carbonell, P.;
Ballesta, F.;

[94399] Estivill, X.: Fragile X syndrome and the (CGG)_n mutation:
two families

[94400] with discordant MZ twins. Am. J. Hum. Genet. 54:
437–442, 1994.

[94401]

[94402] 113. Kunst, C. B.; Leeflang, E. P.; Iber, J. C.; Arnheim, N.;
Warren,

[94403] S. T.: The effect of FMR1 CGG repeat interruptions on mu-
tation frequency

[94404] as measured by sperm typing. J. Med. Genet. 34:
627–631, 1997.

[94405]

[94406] 114. Kunst, C. B.; Zerylnick, C.; Karickhoff, L.; Eichler, E.;
Bullard,

[94407] J.; Chalifoux, M.; Holden, J. J. A.; Torroni, A.; Nelson, D. L.;
Warren,

[94408] S. T.: FMR1 in global populations. Am. J. Hum. Genet. 58:

513–522,

[94409] 1996.

[94410]

[94411] 115. Lachiewicz, A. M.; Dawson, D. V.; Spiridigliozzi, G.
A.: Physical

[94412] characteristics of young boys with fragile X syndrome:
reasons for

[94413] difficulties in making a diagnosis in young males. Am. J.
Med. Genet. 92:

[94414] 229–236, 2000.

[94415]

[94416] 116. Lagerbauer, B.; Ostareck, D.; Keidel, E.–M.;
Ostareck–Lederer,

[94417] A.; Fischer, U.: Evidence that fragile X mental retardation
protein

[94418] is a negative regulator of translation. Hum. Molec. Genet.
10: 329–338,

[94419] 2001.

[94420]

[94421] 117. Laird, C. D.: Possible erasure of the imprint on a
fragile X

[94422] chromosome when transmitted by a male. *Am. J. Med. Genet.* 38: 391–395,

[94423] 1991.

[94424]

[94425] 118. Laird, C. D.: Proposed mechanism of inheritance and expression

[94426] of the human fragile X syndrome of mental retardation. *Genetics* 117:

[94427] 587–599, 1987.

[94428]

[94429] 119. Laird, C. D.; Jaffe, E.; Karpen, G.; Lamb, M.; Nelson, R.: Fragile

[94430] sites in human chromosomes as regions of late-replicating DNA. *Trends*

[94431] *Genet.* 3: 274–281, 1987.

[94432]

[94433] 120. Laird, C. D.; Lamb, M. M.; Thorne, J. L.: Two progenitor cells

[94434] for human oogonia inferred from pedigree data and the X-inactivation

[94435] imprinting model of the fragile-X syndrome. *Am. J. Hum.*

Genet. 46:

[94436] 696–719, 1990.

[94437]

[94438] 121. Langenbeck, U.; Schmidtke, J.; Bartels, I.; Hansmann, I.; Knuppel,

[94439] H.: Mean corpuscular hemoglobin is increased in Martin–Bell syndrome. Hum.

[94440] Genet. 66: 365–366, 1984.

[94441]

[94442] 122. Larsen, L. A.; Armstrong, J. S. M.; Gronskov, K.; Hjalgrim, H.;

[94443] Brondum–Nielsen, K.; Hasholt, L.; Norgaard–Pedersen, B.; Vuust, J.

[94444] : Analysis of FMR1 (CGG)*n* alleles and FRAXA microsatellite haplotypes

[94445] in the population of Greenland: implications for the population of

[94446] the New World from Asia. Europ. J. Hum. Genet. 7: 771–777, 1999.

[94447]

[94448] 123. Larsen, L. A.; Vuust, J.; Nystad, M.; Evseeva, I.; Van

Ghelue,

- [94449] M.; Tranebjaerg, L.: Analysis of FMR1 (CGG)_n alleles and DXS548-FRAXAC1
- [94450] haplotypes in three European circumpolar populations: traces of genetic
- [94451] relationship with Asia. *Europ. J. Hum. Genet.* 9: 724-727, 2001.
- [94452]
- [94453] 124. Laval, S. H.; Blair, H. J.; Hirst, M. C.; Davies, K. E.; Boyd,
- [94454] Y.: Mapping of FMR1, the gene implicated in fragile X-linked mental
- [94455] retardation, on the mouse X chromosome. *Genomics* 12: 818-821, 1992.
- [94456]
- [94457] 125. Lejeune, J.; Legrand, N.; Lafourcade, J.; Rethore, M.-O.; Raoul,
- [94458] O.; Maunoury, C.: Fragilite du chromosome X et effets de la trimethoprimine. *Ann.*
- [94459] *Genet.* 25: 149-151, 1982.
- [94460]
- [94461] 126. Lewis, H. A.; Musunuru, K.; Jensen, K. B.; Edo, C.;

Chen, H.;

[94462] Darnell, R. B.; Burley, S. K.: Sequence-specific RNA binding by a

[94463] Nova KH domain: implications for paraneoplastic disease and the fragile

[94464] X syndrome. Cell 100: 323-332, 2000.

[94465]

[94466] 127. Limprasert, P.; Jaruratanasirikul, S.; Vasiknanonte, P.: Unilateral

[94467] macroorchidism in fragile X syndrome. (Letter) Am. J. Med. Genet. 95:

[94468] 516-517, 2000.

[94469]

[94470] 128. Linden, M. G.; Tassone, F.; Gane, L. W.; Hills, J. L.; Hagerman,

[94471] R. J.; Taylor, A. K.: Compound heterozygous female with fragile X

[94472] syndrome. Am. J. Med. Genet. 83: 318-321, 1999.

[94473]

[94474] 129. Loesch, D. Z.; Hay, D. A.: Clinical features and reproductive

[94475] patterns in fragile X female heterozygotes. J. Med. Genet.
25: 407–414,

[94476] 1988.

[94477]

[94478] 130. Lubs, H.; Travers, H.; Lujan, E.; Carroll, A.: A large
kindred

[94479] with X-linked mental retardation, marker X and
macroorchidism. Am.

[94480] J. Med. Genet. 17: 145–157, 1984.

[94481]

[94482] 131. Lubs, H. A.; Watson, M.; Breg, R.; Lujan, E.: Restudy
of the

[94483] original marker X family. Am. J. Med. Genet. 17: 133–144,
1984.

[94484]

[94485] 132. Lubs, H. A., Jr.: A marker X chromosome. Am. J.
Hum. Genet. 21:

[94486] 231–244, 1969.

[94487]

[94488] 133. Lugenbeel, K. A.; Peier, A. M.; Carson, N. L.; Chudley,
A. E.;

- [94489] Nelson, D. L.: Intragenic loss of function mutations demonstrate
- [94490] the primary role of FMR1 in fragile X syndrome. *Nature Genet.* 10:
- [94491] 483–485, 1995.
- [94492]
- [94493] 134. Malter, H. E.; Iber, J. C.; Willemsen, R.; de Graaff, E.; Tarleton,
- [94494] J. C.; Leisti, J.; Warren, S. T.; Oostra, B. A.: Characterization
- [94495] of the full fragile X syndrome mutation in fetal gametes. *Nature*
- [94496] *Genet.* 15: 165–169, 1997.
- [94497]
- [94498] 135. Mandel, J.–L.; Hagerman, R.; Froster, U.; Brown, W. T.; Jenkins,
- [94499] E. C.; Jacobs, P.; Turner, G.; Lubs, H.; Neri, G.: Fifth International
- [94500] Workshop on the Fragile X and X-Linked Mental Retardation. *Am. J. Med. Genet.* 43: 5–27, 1992.
- [94501]
- [94502]

[94503] 136. Martin, J. P.; Bell, J.: A pedigree of mental defect showing

[94504] sex-linkage. J. Neurol. Psychiat. 6: 154–157, 1943.

[94505]

[94506] 137. Martin, R. H.; Lin, C. C.; Mathies, B. J.; Lowry, R. B.: X-linked

[94507] mental retardation with macro-orchidism and marker-X chromosomes. Am.

[94508] J. Med. Genet. 7: 433–441, 1980.

[94509]

[94510] 138. Mattei, J. F.; Mattei, M. G.; Aumeras, C.; Auger, M.; Giraud,

[94511] F.: X-linked mental retardation with the fragile X: a study of 15

[94512] families. Hum. Genet. 59: 281–289, 1981.

[94513]

[94514] 139. McCabe, E. R. B.; de la Cruz, F.; Clapp, K.: Workshop on Fragile

[94515] X: future research directions. Am. J. Med. Genet. 85: 317–322, 1999.

[94516]

- [94517] 140. McConkie–Rosell, A.; Lachiewicz, A. M.;
Spiridigliozzi, G. A.;
- [94518] Tarleton, J.; Schoenwald, S.; Phelan, M. C.; Goonewardena,
P.; Ding,
- [94519] X.; Brown, W. T.: Evidence that methylation of the FMR–1
locus is
- [94520] responsible for variable phenotypic expression of the
fragile X syndrome. Am.
- [94521] J. Hum. Genet. 53: 800–809, 1993.
- [94522]
- [94523] 141. Meijer, H.; de Graaff, E.; Merckx, D. M. L.; Jongbloed,
R. J.
- [94524] E.; de Die–Smulders, C. E. M.; Engelen, J. J. M.; Fryns, J.–P.;
- Curfs,
- [94525] P. M. G.; Oostra, B. A.: A deletion of 1.6 kb proximal to
the CGG
- [94526] repeat of the FMR1 gene causes the clinical phenotype of
the fragile
- [94527] X syndrome. Hum. Molec. Genet. 3: 615–620, 1994.
- [94528]
- [94529] 142. Meryash, D. L.; Cronk, C. E.; Sachs, B.; Gerald, P. S.:
An anthropometric

[94530] study of males with the fragile-X syndrome. Am. J. Med.
Genet. 17:

[94531] 159-174, 1984.

[94532]

[94533] 143. Mila, M.; Castellvi-Bel, S.; Gine, R.; Vazquez, C.;
Badenas,

[94534] C.; Sanchez, A.; Estivill, X.: A female compound heterozy-
gote (pre-

[94535] and full mutation) for the CGG FMR1 expansion. Hum.
Genet. 98: 419-421,

[94536] 1996.

[94537]

[94538] 144. Mila, M.; Sanchez, A.; Badenas, C.; Brun, C.; Jimenez,
D.; Villa,

[94539] M. P.; Castellvi-Bel, S.; Estivill, X.: Screening for FMR1 and
FMR2

[94540] mutations in 222 individuals from Spanish special
schools: identification

[94541] of a case of FRAXE-associated mental retardation. Hum.
Genet. 100:

[94542] 503-507, 1997.

[94543]

[94544] 145. Mingroni-Netto, R. C.; Angeli, C. B.; Auricchio, M. T. B. M.;

[94545] Leal-Mesquita, E. R.; Ribeiro-dos-Santos, A. K. C.; Ferrari, I.; Hutz,

[94546] M. H.; Salzano, F. M.; Hill, K.; Hurtado, A. M.; Vianna-Morgante,

[94547] A. M.: Distribution of CGG repeats and FRAXAC1/DXS548 alleles in

[94548] South American populations. Am. J. Med. Genet. 111: 243-252, 2002.

[94549]

[94550] 146. Morales, J.; Hiesinger, P. R.; Schroeder, A. J.; Kume, K.; Verstreken,

[94551] P.; Jackson, F. R.; Nelson, D. L.; Hassan, B. A.: Drosophila fragile

[94552] X protein, DFXR, regulates neuronal morphology and function in the

[94553] brain. Neuron 34: 961-972, 2002.

[94554]

[94555] 147. Morton, J. E.; Bunday, S.; Webb, T. P.; MacDonald, F.; Rindl,

[94556] P. M.; Bullock, S.: Fragile X syndrome is less common than previously

[94557] estimated. J. Med. Genet. 34: 1–5, 1997.

[94558]

[94559] 148. Morton, N. E.; Macpherson, J. N.: Population genetics of the

[94560] fragile–X syndrome: multiallelic model for the FMR1 locus. Proc.

[94561] Nat. Acad. Sci. 89: 4215–4217, 1992.

[94562]

[94563] 149. Mulley, J. C.; Yu, S.; Gedeon, A. K.; Donnelly, A.; Turner, G.;

[94564] Loesch, D.; Chapman, C. J.; Gardner, R. J. M.; Richards, R. I.; Sutherland,

[94565] G. R.: Experience with direct molecular diagnosis of fragile X. J.

[94566] Med. Genet. 29: 368–374, 1992.

[94567]

[94568] 150. Mulligan, L. M.; Phillips, M. A.; Forster–Gibson, C. J.; Beckett,

[94569] J.; Partington, M. W.; Simpson, N. E.; Holden, J. J. A.; White,

B.

[94570] N.: Genetic mapping of DNA segments relative to the locus for the

[94571] fragile-X syndrome at Xq27.3. *Am. J. Hum. Genet.* 37: 463-472, 1985.

[94572]

[94573] 151. Murray, A.; Ennis, S.; Morton, N.: No evidence for parent of

[94574] origin influencing premature ovarian failure in fragile X premutation

[94575] carriers. (Letter) *Am. J. Hum. Genet.* 67: 253-254, 2000.

[94576]

[94577] 152. Murray, A.; Webb, J.; Grimley, S.; Conway, G.; Jacobs, P.: Studies

[94578] of FRAXA and FRAXE in women with premature ovarian failure. *J. Med.*

[94579] *Genet.* 35: 637-640, 1998.

[94580]

[94581] 153. Nolin, S. L.; Lewis, F. A., III; Ye, L. L.; Houck, G. E., Jr.;

[94582] Glicksman, A. E.; Limprasert, P.; Li, S. Y.; Zhong, N.; Ashley, A.

[94583] E.; Feingold, E.; Sherman, S. L.; Brown, W. T.: Familial transmission

[94584] of the FMR1 GCC repeat. *Am. J. Hum. Genet.* 59: 1252–1261, 1996.

[94585]

[94586] 154. Nussbaum, R. L.; Ledbetter, D. H.: Fragile X syndrome: a unique

[94587] mutation in man. *Annu. Rev. Genet.* 20: 109–145, 1986.

[94588]

[94589] 155. Oberle, I.; Rousseau, F.; Heitz, D.; Kretz, C.; Devys, D.; Hanauer,

[94590] A.; Boue, J.; Bertheas, M. F.; Mandel, J. L.: Instability of a 550–base

[94591] pair DNA segment and abnormal methylation in fragile X syndrome. *Science* 252:

[94592] 1097–1102, 1991.

[94593]

[94594] 156. Oostra, B. A.; Chiurazzi, P.: The fragile X gene and its function. *Clin.*

[94595] *Genet.* 60: 399–408, 2001.

[94596]

- [94597] 157. Opitz, J. M.; Sutherland, G. R.: International work-shop on the
[94598] fragile X and X-linked mental retardation. Am. J. Med.
Genet. 17:
[94599] 5-94, 1984.
- [94600]
- [94601] 158. Opitz, J. M.; Westphal, J. M.; Daniel, A.: Discovery of a
connective
[94602] tissue dysplasia in the Martin-Bell syndrome. Am. J. Med.
Genet. 17:
[94603] 101-109, 1984.
- [94604]
- [94605] 159. Oudet, C.; Mornet, E.; Serre, J. L.; Thomas, F.;
Lentes-Zengerling,
[94606] S.; Kretz, C.; Deluchat, C.; Tejada, I.; Boue, J.; Boue, A.;
Mandel,
[94607] J. L.: Linkage disequilibrium between the fragile X muta-
tion and
[94608] two closely linked CA repeats suggests that fragile X chro-
mosomes
[94609] are derived from a small number of founder chromo-
somes. Am. J. Hum.

[94610] Genet. 52: 297–304, 1993.

[94611]

[94612] 160. Park, V.; Howard–Peebles, P.; Sherman, S.; Taylor, A.; Wulfsberg,

[94613] E.: Policy statement: American College of Medical Genetics. Fragile

[94614] X syndrome: diagnostic and carrier testing. Am. J. Med. Genet. 53:

[94615] 380–381, 1994.

[94616]

[94617] 161. Parrish, J. E.; Oostra, B. A.; Verkerk, A. J. M. H.; Richards,

[94618] C. S.; Reynolds, J.; Spikes, A. S.; Shaffer, L. G.; Nelson, D. L.

[94619] : Isolation of a GCC repeat showing expansion in FRAXF, a fragile

[94620] site distal to FRAXA and FRAXE. Nature Genet. 8: 229–235, 1994.

[94621]

[94622] 162. Peier, A. M.; McIlwain, K. L.; Kenneson, A.; Warren, S. T.; Paylor,

- [94623] R.; Nelson, D. L.: (Over)correction of FMR1 deficiency with YAC transgenics:
- [94624] behavioral and physical features. Hum. Molec. Genet. 9: 1145–1159,
- [94625] 2000.
- [94626]
- [94627] 163. Pellissier, M.–C.; Voelckel, M.–A.; Piquet, C.; Mattei, M.–G.;
- [94628] Mattei, J.–F.: Transmission of mental retardation with fragile X
- [94629] site by two normal transmitter brothers. Am. J. Med. Genet. 38:
- [94630] 367–369, 1991.
- [94631]
- [94632] 164. Pembrey, M. E.; Winter, R. M.; Davies, K. E.: A premutation
- [94633] that generates a defect at crossing over explains the inheritance
- [94634] of fragile X mental retardation. Am. J. Med. Genet. 21: 709–717,
- [94635] 1985.

[94636]

[94637] 165. Pieretti, M.; Zhang, F.; Fu, Y.-H.; Warren, S. T.; Oostra, B.

[94638] A.; Caskey, C. T.; Nelson, D. L.: Absence of expression of the FMR-1

[94639] gene in fragile X syndrome. Cell 66: 817-822, 1991.

[94640]

[94641] 166. Poustka, A.; Dietrich, A.; Langenstein, G.; Toniolo, D.; Warren,

[94642] S. T.; Lehrach, H.: Physical map of human Xq27-qter: localizing the

[94643] region of the fragile X mutation. Proc. Nat. Acad. Sci. 88: 8302-8306,

[94644] 1991.

[94645]

[94646] 167. Purrello, M.; Alhadeff, B.; Esposito, D.; Szabo, P.; Rocchi,

[94647] M.; Truett, M.; Masiarz, F.; Siniscalco, M.: The human genes for

[94648] hemophilia A and hemophilia B flank the X chromosome fragile site

[94649] at Xq27.3. EMBO J. 4: 725–729, 1985.

[94650]

[94651] 168. Pyeritz, R. E.; Stamberg, J.; Thomas, G. H.; Bell, B. B.; Zahka,

[94652] K. G.; Bernhardt, B. A.: The marker Xq28 syndrome (fragile X syndrome)

[94653] in a retarded man with mitral valve prolapse. Johns Hopkins Med.

[94654] J. 151: 231–237, 1982.

[94655]

[94656] 169. Quan, F.; Grompe, M.; Jakobs, P.; Popovich, B. W.: Spontaneous

[94657] deletion in the FMR1 gene in a patient with fragile X syndrome and

[94658] cherubism. Hum. Molec. Genet. 4: 1681–1684, 1995.

[94659]

[94660] 170. Quan, F.; Zonana, J.; Gunter, K.; Peterson, K. L.; Magenis, R.

[94661] E.; Popovich, B. W.: An atypical case of fragile X syndrome caused

[94662] by a deletion that includes the FMR1 gene. Am. J. Hum.

Genet. 56:

[94663] 1042-1051, 1995.

[94664]

[94665] 171. Ramos, F. J.; Emanuel, B. S.; Spinner, N. B.: Frequency of the

[94666] common fragile site at Xq27.2 under conditions of thymidylate stress:

[94667] implications for cytogenetic diagnosis of the fragile-X syndrome. Am.

[94668] J. Med. Genet. 42: 835-838, 1992.

[94669]

[94670] 172. Reiss, A. L.; Aylward, E.; Freund, L. S.; Joshi, P. K.; Bryan,

[94671] R. N.: Neuroanatomy of fragile X syndrome: the posterior fossa. Ann.

[94672] Neurol. 29: 26-32, 1991.

[94673]

[94674] 173. Reiss, A. L.; Freund, L.; Tseng, J. E.; Joshi, P. K.: Neuroanatomy

[94675] in fragile X females: the posterior fossa. Am. J. Hum. Genet. 49:

[94676] 279–288, 1991.

[94677]

[94678] 174. Reiss, A. L.; Lee, J.; Freund, L.: Neuroanatomy of fragile X

[94679] syndrome: the temporal lobe. *Neurology* 44: 1317–1324, 1994.

[94680]

[94681] 175. Reyniers, E.; Vits, L.; De Boulle, K.; Van Roy, B.; Van Velzen,

[94682] D.; de Graaff, E.; Verkerk, A. J. M. H.; Jorens, H. Z. J.; Darby,

[94683] J. K.; Oostra, B.; Willems, P. J.: The full mutation in the FMR-1

[94684] gene of male fragile X patients is absent in their sperm. *Nature*

[94685] *Genet.* 4: 143–148, 1993.

[94686]

[94687] 176. Richards, B. W.; Sylvester, P. E.; Brooker, C.: Fragile X-linked

[94688] mental retardation: the Martin–Bell syndrome. *J. Ment. Defic. Res.* 25:

[94689] 253–256, 1981.

[94690]

[94691] 177. Richards, R. I.; Holman, K.; Friend, K.; Kremer, E.; Hillen,

[94692] D.; Staples, A.; Brown, W. T.; Goonewardena, P.; Tarleton, J.; Schwartz,

[94693] C.; Sutherland, G. R.: Evidence of founder chromosomes in fragile

[94694] X syndrome. *Nature Genet.* 1: 257–260, 1992.

[94695]

[94696] 178. Richards, R. I.; Holman, K.; Kozman, H.; Kremer, E.; Lynch, M.;

[94697] Pritchard, M.; Yu, S.; Mulley, J.; Sutherland, G. R.: Fragile X syndrome:

[94698] genetic localisation by linkage mapping of two microsatellite repeats

[94699] FRAXAC1 and FRAXAC2 which immediately flank the fragile site. *J.*

[94700] *Med. Genet.* 28: 818–823, 1991.

[94701]

[94702] 179. Richards, R. I.; Holman, K.; Kozman, H.; Kremer, E.;

Lynch, M.;

[94703] Pritchard, M.; Yu, S.; Mulley, J.; Sutherland, G. R.: Fragile X syndrome:

[94704] genetic localisation by linkage mapping of two microsatellite repeats

[94705] FRAXAC1 and FRAXAC2 which immediately flank the fragile site. J.

[94706] Med. Genet. 28: 818–823, 1991.

[94707]

[94708] 180. Richards, R. I.; Sutherland, G. R.: Dynamic mutations: a new

[94709] class of mutations causing human disease. Cell 70: 709–712, 1992.

[94710]

[94711] 181. Rocchi, M.; Archidiacono, N.; Rinaldi, A.; Filippi, G.; Bartolucci,

[94712] G.; Fancello, G. S.; Siniscalco, M.: Mental retardation in heterozygotes

[94713] for the fragile-X mutation: evidence in favor of an X inactivation-dependent

[94714] effect. Am. J. Hum. Genet. 46: 738–743, 1990.

[94715]

[94716] 182. Rodewald, L.; Miller, D. C.; Sciorra, L.; Barabas, G.;
Lee, M.-L.

[94717] : Central nervous system neoplasm in a young man with
Martin-Bell

[94718] syndrome--fra(X)-XLMR. Am. J. Med. Genet. 26: 7-12,
1987.

[94719]

[94720] 183. Rousseau, F.; Heitz, D.; Biancalana, V.; Blumenfeld,
S.; Kretz,

[94721] C.; Boue, J.; Tommerup, N.; Van Der Hagen, C.; DeLozier-
Blanchet,

[94722] C.; Croquette, M.-F.; Gilgenkrantz, S.; Jalbert, P.; Voelckel,
M.-A.;

[94723] Oberle, I.; Mandel, J.-L.: Direct diagnosis by DNA analysis
of the

[94724] fragile X syndrome of mental retardation. New Eng. J. Med.
325:

[94725] 1673-1681, 1991.

[94726]

[94727] 184. Rousseau, F.; Heitz, D.; Mandel, J.-L.: The unstable
and methylatable

[94728] mutations causing the fragile X syndrome. Hum. Mutat. 1:

91–96,

[94729] 1992.

[94730]

[94731] 185. Rousseau, F.; Heitz, D.; Oberle, I.; Mandel, J.-L.: Se-
lection

[94732] in blood cells from female carriers of the fragile X syn-
drome: inverse

[94733] correlation between age and proportion of active X chro-
mosomes carrying

[94734] the full mutation. J. Med. Genet. 28: 830–836, 1991.

[94735]

[94736] 186. Rousseau, F.; Heitz, D.; Tarleton, J.; MacPherson, J.;
Malmgren,

[94737] H.; Dahl, N.; Barnicoat, A.; Mathew, C.; Mornet, E.; Tejada,
I.; Maddalena,

[94738] A.; Spiegel, R.; Schinzel, A.; Marcos, J. A. G.; Schorderet,
D. F.;

[94739] Schaap, T.; Maccioni, L.; Russo, S.; Jacobs, P. A.; Schwartz,
C.;

[94740] Mandel, J. L.: A multicenter study on genotype–phenotype
correlations

[94741] in the fragile X syndrome, using direct diagnosis with

probe StB12.3:

[94742] the first 2,253 cases. Am. J. Hum. Genet. 55: 225–237, 1994.

[94743]

[94744] 187. Russo, S.; Briscioli, V.; Cogliati, F.; Macchi, M.; Lalatta,

[94745] F.; Larizza, L.: An unusual fragile X sibship: female compound heterozygote

[94746] and male with a partially methylated full mutation. Clin. Genet. 54:

[94747] 309–314, 1998.

[94748]

[94749] 188. Ruvalcaba, R. H. A.; Myhre, S. A.; Roosen-Runge, E. C.; Beckwith,

[94750] J. B.: X-linked mental deficiency megalotestes syndrome. J.A.M.A. 238:

[94751] 1646–1650, 1977.

[94752]

[94753] 189. Schrandt-Stumpel, C.; Gerver, W.-J.; Meyer, H.; Engelen, J.;

[94754] Mulder, H.; Fryns, J.-P.: Prader-Willi-like phenotype in

fragile

[94755] X syndrome. Clin. Genet. 45: 175–180, 1994.

[94756]

[94757] 190. Sherman, S. L.: Premature ovarian failure among fragile X premutation

[94758] carriers: parent-of-origin effect? (Editorial) Am. J. Hum. Genet. 67:

[94759] 11–13, 2000.

[94760]

[94761] 191. Sherman, S. L.; Jacobs, P. A.; Morton, N. E.; Froster-Iskenius,

[94762] U.; Howard-Peebles, P. N.; Nielsen, K. B.; Partington, M. W.; Sutherland,

[94763] G. R.; Turner, G.; Watson, M.: Further segregation analysis of the

[94764] fragile X syndrome with special reference to transmitting males. Hum.

[94765] Genet. 69: 289–299, 1985.

[94766]

[94767] 192. Siomi, H.; Choi, M.; Siomi, M. C.; Nussbaum, R. L.; Dreyfuss,

- [94768] G.: Essential role for KH domains in RNA binding: impaired RNA binding
- [94769] by a mutation in the KH domain of FMR1 that causes fragile X syndrome. Cell 77:
- [94770] 33–39, 1994.
- [94771]
- [94772] 193. Siomi, H.; Siomi, M. C.; Nussbaum, R. L.; Dreyfuss, G.: The
- [94773] protein product of the fragile X gene, FMR1, has characteristics of
- [94774] an RNA-binding protein. Cell 74: 291–298, 1993.
- [94775]
- [94776] 194. Sittler, A.; Devys, D.; Weber, C.; Mandel, J.-L.: Alternative
- [94777] splicing of exon 14 determines nuclear or cytoplasmic localisation
- [94778] of fmr1 protein isoforms. Hum. Molec. Genet. 5: 95–102, 1996.
- [94779]
- [94780] 195. Smeets, H. J. M.; Smits, A. P. T.; Verheij, C. E.; Theelen, J.

[94781] P. G.; Willemsen, R.; van de Burgt, I.; Hoogeveen, A. T.;
Oosterwijk,

[94782] J. C.; Oostra, B. A.: Normal phenotype in two brothers with
a full

[94783] FMR1 mutation. Hum. Molec. Genet. 4: 2103–2108, 1995.

[94784]

[94785] 196. Smits, A. P. T.; Dreesen, J. C. F. M.; Post, J. G.;
Smeets, D.

[94786] F. C. M.; de Die-Smulders, C.; Spaans-van der Bijl, T.; Go-
vaerts,

[94787] L. C. P.; Warren, S. T.; Oostra, B. A.; van Oost, B. A.: The
fragile

[94788] X syndrome: no evidence for any recent mutations. J. Med.
Genet. 30:

[94789] 94–96, 1993.

[94790]

[94791] 197. Snow, K.; Doud, L. K.; Hagerman, R.; Pergolizzi, R. G.;
Erster,

[94792] S. H.; Thibodeau, S. N.: Analysis of a CGG sequence at the
FMR–1

[94793] locus in fragile X families and in the general population.
Am. J.

[94794] Hum. Genet. 53: 1217–1228, 1993.

[94795]

[94796] 198. Snyder, F. F.; Lin, C. C.; Harasym, C. A.; Jamro, H. K.;

Kushnig,

[94797] M. L.; Lowe, J. K.; O'Brien, S. I.: Evidence for close association

[94798] between the fragile X(q27-28) chromosome site and glucose-6-phosphate

[94799] dehydrogenase (G6PD) but not with hypoxanthine-guanine phosphoribosyltransferase

[94800] (HPRT). (Abstract) Cytogenet. Cell Genet. 37: 587, 1984.

[94801]

[94802] 199. Soysa, P.; Senanayahe, M.; Mikkelsen, M.; Poulsen, H.: Martin-Bell

[94803] syndrome fra(X)(q28) in a Sri Lankan family. J. Ment. Defic. Res. 26:

[94804] 251-257, 1982.

[94805]

[94806] 200. Steinbach, P.: Mental impairment in Martin-Bell syndrome is

[94807] probably determined by interaction of several genes: simple explanation

[94808] of phenotypic differences between unaffected and affected males with

[94809] the same X chromosome. Hum. Genet. 72: 248-252,

1986.

[94810]

[94811] 201. Stoll, C.: Problems in the diagnosis of fragile X syndrome in

[94812] young children are still present. Am. J. Med. Genet. 100: 110–115,

[94813] 2001.

[94814]

[94815] 202. Storm, K.; Handig, I.; Reyniers, E.; Oostra, B. A.; Kooy, R.

[94816] F.; Willems, P. J.: Incomplete EcoRI digestion may lead to false

[94817] diagnosis of fragile X syndrome. Hum. Genet. 102: 54–56, 1998.

[94818]

[94819] 203. Subramanian, P. S.; Nelson, D. L.; Chinault, A. C.: Large domains

[94820] of apparent delayed replication timing associated with triplet repeat

[94821] expansion at FRAXA and FRAXE. Am. J. Hum. Genet. 59: 407–416, 1996.

[94822]

- [94823] 204. Sun, Y. J.; Baumer, A.: Nonrandom X inactivation and selection
[94824] of fragile X full mutation in fetal fibroblasts. Am. J. Med. Genet. 86:
[94825] 162–164, 1999.
[94826]
- [94827] 205. Sutherland, G. R.: Personal Communication. Adelaide, Australia
[94828] 11/3/1993.
[94829]
- [94830] 206. Sutherland, G. R.: Fragile sites on human chromosomes: demonstration
[94831] of their dependence on the type of tissue culture medium. Science 197:
[94832] 265–266, 1977.
[94833]
- [94834] 207. Sutherland, G. R.: Heritable sites on human chromosomes. VII.
[94835] Children homozygous for the BrdU–requiring fra10q25 are phenotypically
[94836] normal. Am. J. Hum. Genet. 33: 946–949, 1981.

[94837]

[94838] 208. Sutherland, G. R.: Heritable fragile sites on human chromosomes.

[94839] II. Distribution, phenotypic effects, and cytogenetics. Am. J. Hum.

[94840] Genet. 31: 136–148, 1979.

[94841]

[94842] 209. Sutherland, G. R.: Personal Communication. Adelaide, Australia

[94843] 6/1989.

[94844]

[94845] 210. Sutherland, G. R.; Ashford, P. L. C.: X-linked mental retardation

[94846] with macro-orchidism and the fragile site at Xq27 or 28. Hum. Genet. 48:

[94847] 117–120, 1979.

[94848]

[94849] 211. Sutherland, G. R.; Baker, E.: Characterisation of a new rare

[94850] fragile site easily confused with the fragile X. Hum. Molec. Genet. 1:

[94851] 111–113, 1992.

[94852]

[94853] 212. Sutherland, G. R.; Gedeon, A.; Kornman, L.; Donnelly, A.; Byard,

[94854] R. W.; Mulley, J. C.; Kremer, E.; Lynch, M.; Pritchard, M.; Yu, S.;

[94855] Richards, R. I.: Prenatal diagnosis of fragile X syndrome by direct

[94856] detection of the unstable DNA sequence. New Eng. J. Med. 325: 1720–1722,

[94857] 1991.

[94858]

[94859] 213. Sved, J. A.; Laird, C. D.: Population genetic consequences of

[94860] the fragile X syndrome, based on the X-inactivation imprinting model. Am.

[94861] J. Hum. Genet. 46: 443–451, 1990.

[94862]

[94863] 214. Sved, J. A.; Laird, C. D.: The X-inactivation imprinting model

[94864] can explain the incidence of the fragile-X syndrome of

mental retardation

[94865] in mother-offspring pairs. Brain Dysfunction 1: 245-254, 1988.

[94866]

[94867] 215. Szabo, P.; Purrello, M.; Rocchi, M.; Archidiacono, N.; Alhadeff,

[94868] B.; Filippi, G.; Toniolo, D.; Martini, G.; Luzzatto, L.; Siniscalco,

[94869] M.: Cytological mapping of the human glucose-6-phosphate dehydrogenase

[94870] gene distal to the fragile-X site suggests a high rate of meiotic

[94871] recombination across this site. Proc. Nat. Acad. Sci. 81: 7855-7859,

[94872] 1984.

[94873]

[94874] 216. Tamanini, F.; Bontekoe, C.; Bakker, C. E.; van Unen, L.; Anar,

[94875] B.; Willemsen, R.; Yoshida, M.; Galjaard, H.; Oostra, B. A.; Hoogeveen,

[94876] A. T.: Different targets for the fragile X-related proteins revealed

[94877] by their distinct nuclear localizations. Hum. Molec. Genet.
8: 863–869,

[94878] 1999.

[94879]

[94880] 217. Tarleton, J.; Richie, R.; Schwartz, C.; Rao, K.;
Aylsworth, A.

[94881] S.; Lachiewicz, A.: An extensive de novo deletion removing
FMR1 in

[94882] a patient with mental retardation and the fragile X syn-
drome phenotype. Hum.

[94883] Molec. Genet. 2: 1973–1974, 1993.

[94884]

[94885] 218. Thibodeau, S. N.; Dorkins, H. R.; Faulk, K. R.; Berry,
R.; Smith,

[94886] A. C. M.; Hagerman, R.; King, A.; Davies, K. E.: Linkage
analysis

[94887] using multiple DNA polymorphic markers in normal fami-
lies and in families

[94888] with fragile X syndrome. Hum. Genet. 79: 219–227, 1988.

[94889]

[94890] 219. Thode, A.; Laing, S.; Partington, M. W.; Turner, G.: Is

there

[94891] a fragile(X) negative Martin-Bell syndrome?. Am. J. Med. Genet. 30:

[94892] 459-471, 1988.

[94893]

[94894] 220. Toledano-Alhadeff, H.; Basel-Vanagaite, L.; Magal, N.; Davidov,

[94895] B.; Ehrlich, S.; Drasinover, V.; Taub, E.; Halpern, G. J.; Ginott,

[94896] N.; Shohat, M.: Fragile X carrier screening and the prevalence of

[94897] premutation and full-mutation carriers in Israel. Am. J. Hum. Genet. 69:

[94898] 351-360, 2001.

[94899]

[94900] 221. Tommerup, N.; Poulsen, H.; Brondum-Nielsen, K.: 5-Fluoro-2-prime-deoxyuridine

[94901] induction of the fragile site on Xq28 associated with X-linked mental

[94902] retardation. J. Med. Genet. 33: 234-242, 1981.

[94903]

- [94904] 222. Turner, G.; Brookwell, R.; Daniel, A.; Selikowitz, M.; Zilibowitz,
- [94905] M.: Heterozygous expression of X-linked mental retardation and X-chromosome
- [94906] marker fra(X)(q27). New Eng. J. Med. 303: 662-664, 1980.
- [94907]
- [94908] 223. Turner, G.; Daniel, A.; Frost, M.: X-linked mental retardation,
- [94909] macro-orchidism, and the Xq27 fragile site. J. Pediat. 96: 837-841,
- [94910] 1980.
- [94911]
- [94912] 224. Turner, G.; Eastman, C.; Casey, J.; McLeay, A.; Procopis, P.;
- [94913] Turner, B.: X-linked mental retardation associated with macro-orchidism. J.
- [94914] Med. Genet. 12: 367-371, 1975.
- [94915]
- [94916] 225. Turner, G.; Till, R.; Daniel, A.: Marker X chromosomes, mental
- [94917] retardation and macro-orchidism. (Letter) New Eng. J.

Med. 299:

[94918] 1472, 1978.

[94919]

[94920] 226. Turner, G.; Webb, T.; Wake, S.; Robinson, H.: Prevalence of

[94921] fragile X syndrome. Am. J. Med. Genet. 64: 196–197, 1996.

[94922]

[94923] 227. Uchida, I. A.; Joyce, E. M.: Activity of the fragile X in heterozygous

[94924] carriers. Am. J. Hum. Genet. 34: 286–293, 1982.

[94925]

[94926] 228. van den Ouweland, A. M. W.; Deelen, W. H.; Kunst, C. B.; Uzielli,

[94927] M.–L. G.; Nelson, D. L.; Warren, S. T.; Oostra, B. A.; Halley, D.

[94928] J. J.: Loss of mutation at the FMR1 locus through multiple exchanges

[94929] between maternal X chromosomes. Hum. Molec. Genet. 3: 1823–1827,

[94930] 1994.

[94931]

[94932] 229. Van Roy, B. C.; De Smedt, M. C.; Raes, R. A.; Dumon, J. E.; Leroy,

[94933] J. G.: Fragile X trait in a large kindred: transmission also through

[94934] normal males. J. Med. Genet. 20: 286–289, 1983.

[94935]

[94936] 230. Verheij, C.; Bakker, C. E.; de Graaff, E.; Keulemans, J.; Willemsen,

[94937] R.; Verkerk, A. J. M. H.; Galjaard, H.; Reuser, A. J. J.; Hoogeveen,

[94938] A. T.; Oostra, B. A.: Characterization and localization of the FMR-1

[94939] gene product associated with fragile X syndrome. Nature 363: 722–724,

[94940] 1993.

[94941]

[94942] 231. Verheij, C.; de Graaff, E.; Bakker, C. E.; Willemsen, R.; Willems,

[94943] P. J.; Meijer, N.; Galjaard, H.; Reuser, A. J. J.; Oostra, B. A.;

[94944] Hoogeveen, A. T.: Characterization of FMR1 proteins iso-

lated from

[94945] different tissues. Hum. Molec. Genet. 4: 895–901, 1995.

[94946]

[94947] 232. Verkerk, A. J. M. H.; Pieretti, M.; Sutcliffe, J. S.; Fu, Y.-H.;

[94948] Kuhl, D. P. A.; Pizzuti, A.; Reiner, O.; Richards, S.; Victoria, M.

[94949] F.; Zhang, F.; Eussen, B. E.; van Ommen, G.-J. B.; Blonden, L. A.

[94950] J.; Riggins, G. J.; Chastain, J. L.; Kunst, C. B.; Galjaard, H.; Caskey,

[94951] C. T.; Nelson, D. L.; Oostra, B. A.; Warren, S. T.: Identification

[94952] of a gene (FMR-1) containing a CCG repeat coincident with a breakpoint

[94953] cluster region exhibiting length variation in fragile X syndrome. Cell 65:

[94954] 905–914, 1991.

[94955]

[94956] 233. Vianna-Morgante, A. M.; Costa, S. S.: Premature ovarian failure

[94957] is associated with maternally and paternally inherited pre-

mutation

- [94958] in Brazilian families with fragile X. (Letter) Am. J. Hum. Genet. 67: 254–255, 2000.
- [94959]
- [94960]
- [94961] 234. Vincent, J. B.; Gurling, H. M. D.: Point mutation in intron 10 of FMR1 is unlikely to be a cause of fragile X syndrome. (Letter) Hum. Mutat. 12: 431 only, 1998.
- [94964]
- [94965] 235. Voelckel, M. A.; Mattei, M. G.; N'Guyen, C.; Philip, N.; Birg, F.; Mattei, J. F.: Dissociation between mental retardation and fragile site expression in a family with fragile X-linked mental retardation. Hum. Genet. 80: 375–378, 1988.
- [94969]
- [94970] 236. Vogel, F.; Crusio, W. E.; Kovac, C.; Fryns, J.-P.; Freund, M.

[94971] : Selective advantage of fra (X) heterozygotes. Hum.

Genet. 86:

[94972] 25–32, 1990.

[94973]

[94974] 237. Wallace, D. C.; Torroni, A.: American Indian prehistory as written

[94975] in the mitochondrial DNA: a review. Hum. Biol. 64: 403–416, 1992.

[94976]

[94977] 238. Wang, Y.-C.; Li, S.-Y.: Response to Vincent and Gurling. (Letter) Hum.

[94978] Mutat. 12: 432 only, 1998.

[94979]

[94980] 239. Wang, Y.-C.; Lin, M.-L.; Lin, S. J.; Li, Y.-C.; Li, S.-Y.: Novel

[94981] point mutation within intron 10 of FMR-1 gene causing fragile X syndrome. Hum.

[94982] Mutat. 10: 393–399, 1997.

[94983]

[94984] 240. Warren, S. T.; Glover, T. W.; Davidson, R. L.; Jagadeeswaran,

[94985] P.: Linkage and recombination between fragile X-linked mental retardation

[94986] and the factor IX gene. (Letter) Hum. Genet. 69: 44–46, 1985.

[94987]

[94988] 241. Warren, S. T.; Knight, S. J. L.; Peters, J. F.; Stayton, C. L.;

[94989] Consalez, G. G.; Zhang, F. P.: Isolation of the human chromosomal

[94990] band Xq28 within somatic cell hybrids by fragile X site breakage. Proc.

[94991] Nat. Acad. Sci. 87: 3856–3860, 1990.

[94992]

[94993] 242. Warren, S. T.; Zhang, F.; Licameli, G. R.; Peters, J. F.: The

[94994] fragile X site in somatic cell hybrids: an approach for molecular

[94995] cloning of fragile sites. Science 237: 420–423, 1987.

[94996]

[94997] 243. Warren, S. T.; Zhang, F. P.; Sutcliffe, J. S.; Peters, J. F.

[94998] : Strategy for molecular cloning of the fragile X site DNA.

Am. J.

[94999] Med. Genet. 30: 613–623, 1988.

[95000]

[95001] 244. Weaver, D. D.; Sherman, S. L.: A counseling guide to the Martin–Bell

[95002] syndrome. (Letter) Am. J. Med. Genet. 26: 39–44, 1987.

[95003]

[95004] 245. Webb, T.; Jacobs, P. A.: Fragile Xq27.3 in female heterozygotes

[95005] for the Martin–Bell syndrome. J. Med. Genet. 27: 627–631, 1990.

[95006]

[95007] 246. Webb, T. P.; Bunday, S.; Thake, A.; Todd, J.: The frequency

[95008] of the fragile X chromosome among schoolchildren in Coventry. J.

[95009] Med. Genet. 23: 396–399, 1986.

[95010]

[95011] 247. Wheeler, R. F.; Roberts, S. H.: An improved lymphocyte culture

[95012] technique: deoxycytidine release of a thymidine block and

use of a

[95013] constant humidity chamber for slide making. J. Med.

Genet. 24: 113–115,

[95014] 1987.

[95015]

[95016] 248. Willemsen, R.; Anar, B.; De Diego Otero, Y.; de Vries,

B. B.

[95017] A.; Hilhorst–Hofstee, Y.; Smits, A.; van Looveren, E.;

Willems, P.

[95018] J.; Galjaard, H.; Oostra, B. A.: Noninvasive test for fragile X syndrome,

[95019] using hair root analysis. Am. J. Hum. Genet. 65: 98–103, 1999.

[95020]

[95021] 249. Willemsen, R.; Mohkamsing, S.; De Vries, B.; Devys, D.; van den

[95022] Ouweland, A.; Mandel, J. L.; Galjaard, H.; Oostra, B.: Rapid antibody

[95023] test for fragile X syndrome. Lancet 345: 1147–1148, 1995.

[95024]

[95025] 250. Willemsen, R.; Smits, A.; Mohkamsing, S.; van

Beerendonk, H.;

[95026] de Haan, A.; de Vries, B.; van den Ouweland, A.; Sistermans, A.; Galjaard,

[95027] H.; Oostra, B. A.: Rapid antibody test for diagnosing fragile X syndrome:

[95028] a validation of the technique. Hum. Genet. 99: 308–311, 1997.

[95029]

[95030] 251. Winter, R. M.; Pembrey, M. E.: Analysis of linkage relationships

[95031] between genetic markers around the fragile X locus with special reference

[95032] to the daughters of normal transmitting males. Hum. Genet. 74: 93–97,

[95033] 1986.

[95034]

[95035] 252. Wohrle, D.; Hennig, I.; Vogel, W.; Steinbach, P.: Mitotic stability

[95036] of fragile X mutations in differentiated cells indicates early post-conceptual

[95037] trinucleotide repeat expansion. Nature Genet. 4: 140–142, 1993.

[95038]

[95039] 253. Wohrle, D.; Kotzot, D.; Hirst, M. C.; Manca, A.; Korn, B.; Schmidt,

[95040] A.; Barbi, G.; Rott, H.-D.; Poustka, A.; Davies, K. E.; Steinbach,

[95041] P.: A microdeletion of less than 250 kb, including the proximal part

[95042] of the FMR-1 gene and the fragile-X site, in a male with the clinical

[95043] phenotype of fragile-X syndrome. Am. J. Hum. Genet. 51: 299-306,

[95044] 1992.

[95045]

[95046] 254. Wohrle, D.; Salat, U.; Hameister, H.; Vogel, W.; Steinbach, P.

[95047] : Demethylation, reactivation, and destabilization of human fragile

[95048] X full-mutation alleles in mouse embryocarcinoma cells. Am. J. Hum.

[95049] Genet. 69: 504-515, 2001.

[95050]

- [95051] 255. Yamauchi, M.; Nagata, S.; Seki, N.; Toyama, Y.; Harada, N.; Niikawa,
[95052] N.; Masuno, I.; Kajii, T.; Hori, T.: Prenatal diagnosis of
fragile
[95053] X syndrome by direct detection of the dynamic mutation
due to an unstable
[95054] DNA sequence. Clin. Genet. 44: 169–172, 1993.
[95055]
[95056] 256. Yeshaya, J.; Shalgi, R.; Shohat, M.; Avivi, L.: Replica-
tion
[95057] timing of the various FMR1 alleles detected by FISH: infer-
ences regarding
[95058] their transcriptional status. Hum. Genet. 102: 6–14, 1998.
[95059]
[95060] 257. Yu, S.; Mulley, J.; Loesch, D.; Turner, G.; Donnelly, A.;
Gedeon,
[95061] A.; Hillen, D.; Kremer, E.; Lynch, M.; Pritchard, M.; Suther-
land,
[95062] G. R.; Richards, R. I.: Fragile-X syndrome: unique genetics
of the
[95063] heritable unstable element. Am. J. Hum. Genet. 50:
968–980, 1992.

[95064]

[95065] 258. Yu, S.; Pritchard, M.; Kremer, E.; Lynch, M.; Nancarrow, J.;

[95066] Baker, E.; Holman, K.; Mulley, J. C.; Warren, S. T.; Schlessinger,

[95067] D.; Sutherland, G. R.; Richards, R. I.: Fragile X genotype characterized

[95068] by an unstable region of DNA. Science 252: 1179–1181, 1991.

[95069]

[95070] 259. Zhang, Y. Q.; Bailey, A. M.; Matthies, H. J. G.; Renden, R. B.;

[95071] Smith, M. A.; Speese, S. D.; Rubin, G. M.; Broadie, K.: Drosophila

[95072] fragile X-related gene regulates the MAP1B homolog Futsch to control

[95073] synaptic structure and function. Cell 107: 591–603, 2001.

[95074]

[95075] 260. Zhong, N.; Dobkin, C.; Brown, W. T.: A complex mutable polymorphism

[95076] located within the fragile X gene. Nature Genet. 5: 248–253, 1993.

[95077]

[95078] 261. Zoll, B.; Arnemann, J.; Krawczak, M.; Cooper, D. N.;
Pescia,

[95079] G.; Wahli, W.; Steinbach, P.; Schmidtke, J.: Evidence against
close

[95080] linkage of the loci for fraXq of Martin-Bell syndrome and
for factor

[95081] IX. Hum. Genet. 71: 122-126, 1985.

[95082] 16549.

[95083] Bartley, T. D.; Bogenberger, J.; Hunt, P.; Li, Y.-S.; Lu, H. S.;

[95084] Martin, F.; Chang, M.-S.; Samal, B.; Nichol, J. L.; Swift, S.;
Johnson,

[95085] M. J.; Hsu, R.-Y.; and 41 others: Identification and cloning
of

[95086] a megakaryocyte growth and development factor that is a
ligand for

[95087] the cytokine receptor Mpl. Cell 77: 1117-1124, 1994.

[95088] 16550.

[95089] Chang, M.; McNinch, J.; Basu, R.; Shutter, J.; Hsu, R.;
Perkins,

[95090] C.; Mar, V.; Suggs, S.; Welcher, A.; Li, L.; Lu, H.; Bartley, T.;

- [95091] Hunt, P.; Martin, F.; Samal, B.; Bogenberger, J.: Cloning and characterization
- [95092] of the human megakaryocyte growth and development factor (MGDF) gene. J.
- [95093] Biol. Chem. 270: 511–514, 1995.
- [95094] 16551.
- [95095] Chang, M.-S.; Hsu, R.-Y.; McNinch, J.; Copeland, N. G.; Jenkins,
- [95096] N. A.: The gene for murine megakaryocyte growth and development factor
- [95097] (thrombopoietin, Thpo) is located on mouse chromosome 16. Genomics 26:
- [95098] 636–637, 1995.
- [95099] 16552.
- [95100] de Sauvage, F. J.; Hass, P. E.; Spencer, S. D.; Malloy, B. E.;
- [95101] Gurney, A. L.; Spencer, S. A.; Darbonne, W. C.; Henzel, W. J.; Wong,
- [95102] S. C.; Kuang, W.-J.; Oles, K. J.; Hultgren, B.; Solberg, L. A., Jr.;
- [95103] Goeddel, D. V.; Eaton, D. L.: Stimulation of megakaryocytopoiesis

- [95104] and thrombopoiesis by the c-Mpl ligand. *Nature* 369: 533–538, 1994.
- [95105] 16553.
- [95106] Farese, A. M.; Hunt, P.; Grab, L. B.; MacVittie, T. J.: Combined
- [95107] administration of recombinant human megakaryocyte growth and development
- [95108] factor and granulocyte colony-stimulating factor enhances multilineage
- [95109] hematopoietic reconstitution in nonhuman primates after radiation-induced
- [95110] marrow aplasia. *J. Clin. Invest.* 97: 2145–2151, 1996.
- [95111] 16554.
- [95112] Foster, D. C.; Sprecher, C. A.; Grant, F. J.; Kramer, J. M.; Kuijper,
- [95113] J. L.; Holly, R. D.; Whitmore, T. E.; Heipel, M. D.; Bell, L. A.;
- [95114] Ching, A. F. T.; McGrane, V.; Hart, C.; O'Hara, P. J.; Lok, S.: Human
- [95115] thrombopoietin: gene structure, cDNA sequence, expression, and chromosomal
- [95116] localization. *Proc. Nat. Acad. Sci.* 91: 13023–13027, 1994.

- [95117] 16555.
- [95118] Ghilardi, N.; Wiestner, A.; Kikuchi, M.; Ohsaka, A.; Skoda, R.
- [95119] C.: Hereditary thrombocythaemia in a Japanese family is caused by
- [95120] a novel point mutation in the thrombopoietin gene. *Brit. J. Haemat.* 107:
- [95121] 310–316, 1999.
- [95122] 16556.
- [95123] Kaushansky, K.; Lok, S.; Holly, R. D.; Broudy, V. C.; Lin, N.;
- [95124] Bailey, M. C.; Forstrom, J. W.; Buddle, M. M.; Oort, P. J.; Hagen,
- [95125] F. S.; Roth, G. J.; Papayannopoulou, T.; Foster, D. C.: Promotion
- [95126] of megakaryocyte progenitor expansion and differentiation by the c-Mpl
- [95127] ligand thrombopoietin. *Nature* 369: 568–571, 1994.
- [95128] 16557.
- [95129] Kondo, T.; Okabe, M.; Sanada, M.; Kurosawa, M.; Suzuki, S.; Kobayashi,
- [95130] M.; Hosokawa, M.; Asaka, M.: Familial essential thrombo-

cythemia associated

- [95131] with one-base deletion in the 5-prime-untranslated region of the thrombopoietin
- [95132] gene. Blood 92: 1091-1096, 1998.
- [95133] 16558.
- [95134] Li, J.; Yang, C.; Xia, Y.; Bertino, A.; Glaspy, J.; Roberts, M.;
- [95135] Kuter, D. J.: Thrombocytopenia caused by the development of antibodies
- [95136] to thrombopoietin. Blood 98: 3241-3248, 2001.
- [95137] 16559.
- [95138] Lok, S.; Kaushansky, K.; Holly, R. D.; Kuijper, J. L.; Lofton-Day,
- [95139] C. E.; Oort, P. J.; Grant, F. J.; Heipel, M. D.; Burkhead, S. K.;
- [95140] Kramer, J. M.; Bell, L. A.; Sprecher, C. A.; Blumberg, H.; Johnson,
- [95141] R.; Prunkard, D.; Ching, A. F. T.; Mathewes, S. L.; Bailey, M. C.;
- [95142] Forstrom, J. W.; Buddle, M. M.; Osborn, S. G.; Evans, S. J.; Sheppard,
- [95143] P. O.; Presnell, S. R.; O'Hara, P. J.; Hagen, F. S.; Roth, G. J.;
- [95144] Foster, D. C.: Cloning and expression of murine throm-

bopoietin cDNA

[95145] and stimulation of platelet production in vivo. Nature 369:
565–568,

[95146] 1994.

[95147] 16560.

[95148] Metcalf, D.: Thrombopoietin: at last. Nature 369:
519–520,

[95149] 1994.

[95150] 16561.

[95151] Methia, N.; Louache, F.; Vainchenker, W.; Wendling, F.:
Oligodeoxynucleotides

[95152] antisense to the proto-oncogene c-mpl specifically inhibit
in vitro

[95153] megakaryocytopoiesis. Blood 82: 1395–1401, 1993.

[95154] 16562.

[95155] Ratajczak, M. Z.; Ratajczak, J.; Marlicz, W.; Pletcher, C. H.,

[95156] Jr.; Machalinski, B.; Moore, J.; Hung, H.; Gewirtz, A. M.:

Recombinant

[95157] human thrombopoietin (TPO) stimulates erythropoiesis by
inhibiting

[95158] erythroid progenitor cell apoptosis. Brit. J. Haemat. 98:

8-17,

[95159] 1997.

[95160] 16563.

[95161] Sohma, Y.; Akahori, H.; Seki, N.; Hori, T.; Ogami, K.; Kato, T.;

[95162] Shimada, Y.; Kawamura, K.; Miyazaki, H.: Molecular cloning and chromosomal

[95163] localization of the human thrombopoietin gene. FEBS Lett. 353: 57-61,

[95164] 1994.

[95165] 16564.

[95166] Wendling, F.; Maraskovsky, E.; Debili, N.; Florindo, C.; Teepe,

[95167] M.; Titeux, M.; Methia, N.; Breton-Gorius, J.; Cosman, D.; Vainchenker,

[95168] W.: c-Mpl ligand is a humoral regulator of megakaryocytopoiesis. Nature 369:

[95169] 571-574, 1994.

[95170] 16565.

[95171] Dualan, R.; Brody, T.; Keeney, S.; Nichols, A. F.; Admon, A.; Linn,

- [95172] S.: Chromosomal localization and cDNA cloning of the genes (DDB1
- [95173] and DDB2) for the p127 and p48 subunits of a human damage-specific
- [95174] DNA binding protein. *Genomics* 29: 62–69, 1995.
- [95175] 16566.
- [95176] Koken, M. H. M.; Reynolds, P.; Jaspers–Dekker, I.; Prakash, L.;
- [95177] Prakash, S.; Bootsma, D.; Hoeijmakers, J. H. J.: Structural and functional
- [95178] conservation of two human homologs of the yeast DNA repair gene RAD6. *Proc.*
- [95179] *Nat. Acad. Sci.* 88: 8865–8869, 1991.
- [95180] 16567.
- [95181] Knoll, A.; Schunkert, H.; Reichwald, K.; Danser, A. H. J.; Bauer,
- [95182] D.; Platzer, M.; Stein, G.; Rosenthal, A.: Human renin binding protein:
- [95183] complete genomic sequence and association of an intronic T/C polymorphism
- [95184] with the prorenin level in males. *Hum. Molec. Genet.* 6: 1527–1534,

[95185] 1997.

[95186] 16568.

[95187] Laan, M.; Paabo, S.: Mapping genes by drift-generated linkage

[95188] disequilibrium. (Letter) Am. J. Hum. Genet. 63: 654-656, 1998.

[95189] 16569.

[95190] Takahashi, S.; Inoue, H.; Miyake, Y.: The human gene for renin-binding

[95191] protein. J. Biol. Chem. 267: 13007-13013, 1992.

[95192] 16570.

[95193] Terwilliger, J. D.; Zollner, S.; Laan, M.; Paabo, S.: Mapping

[95194] genes through the use of linkage disequilibrium generated by genetic

[95195] drift: 'drift mapping' in small populations with no demographic expansion. Hum.

[95196] Hered. 48: 138-154, 1998.

[95197] 16571.

[95198] van den Ouweland, A. M. W.; Verdijk, M.; Kiochis, P.; Poustka,

- [95199] A.; van Oost, B. A.: The human renin-binding protein gene (RENBP)
- [95200] maps in Xq28. *Genomics* 21: 279–281, 1994.
- [95201] 16572.
- [95202] Warren, S. T.; Knight, S. J. L.; Peters, J. F.; Stayton, C. L.;
- [95203] Consalez, G. G.; Zhang, F.: Isolation of the human chromosomal band
- [95204] Xq28 within somatic cell hybrids by fragile X site breakage. *Proc.*
- [95205] *Nat. Acad. Sci.* 87: 3856–3860, 1990.
- [95206] 16573.
- [95207] Calvo, R. M.; Asuncion, M.; Sancho, J.; San Millan, J. L.; Escobar-Morreale,
- [95208] H. F.: The role of the CAG repeat polymorphism in the androgen receptor
- [95209] gene and of skewed X-chromosome inactivation, in the pathogenesis
- [95210] of hirsutism. *J. Clin. Endocr. Metab.* 85: 1735–1740, 2000.
- [95211] 16574.
- [95212] Sinclair, A. H.; Foster, J. W.; Spencer, J. A.; Page, D. C.; Palmer,

- [95213] M.; Goodfellow, P. N.; Marshall Graves, J. A.: Sequences homologous
- [95214] to ZFY, a candidate human sex-determining gene, are autosomal in marsupials.
- [95215] Nature 336: 780–783, 1988.
- [95216] 16575.
- [95217] Gao, X.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Gridley,
- [95218] T.: Assignment of the murine Notch2 and Notch3 genes to chromosomes
- [95219] 3 and 17. Genomics 49: 160–161, 1998.
- [95220] 16576.
- [95221] Larsson, C.; Lardelli, M.; White, I.; Lendahl, U.: The human NOTCH1,
- [95222] 2, and 3 genes are located at chromosome positions 9q34, 1p13–p11,
- [95223] and 19p13.2–p13.1 in regions of neoplasia-associated translocation. Genomics 24:
- [95224] 253–258, 1994.
- [95225] 16577.
- [95226] Dichgans, M.; Herzog, J.; Gasser, T.: NOTCH3 mutation in–

volving

[95227] three cysteine residues in a family with typical CADASIL.

Neurology 57:

[95228] 1714–1717, 2001.

[95229] 16578.

[95230] Joutel, A.; Corpechot, C.; Ducros, A.; Vahedi, K.; Chabriat, H.;

[95231] Mouton, P.; Alamowitch, S.; Domenga, V.; Cecillion, M.; Marechal,

[95232] E.; Maciazek, J.; Vayssiere, C.; Cruaud, C.; Cabanis, E.–A.; Ruchoux,

[95233] M. M.; Weissenbach, J.; Bach, J. F.; Bousser, M. G.; Tournier–Lasserre,

[95234] E.: Notch3 mutations in CADASIL, a hereditary adult-onset condition

[95235] causing stroke and dementia. Nature 383: 707–710, 1996.

[95236] 16579.

[95237] Joutel, A.; Dodick, D. D.; Parisi, J. E.; Cecillon, M.; Tournier–Lasserre,

[95238] E.; Bousser, M. G.: De novo mutation in the Notch3 gene causing CADASIL. Ann.

- [95239] **Neurol. 47: 388–391, 2000.**
- [95240] **16580.**
- [95241] **Joutel, A.; Vahedi, K.; Corpechot, C.; Troesch, A.; Chabriat, H.;**
- [95242] **Vayssiere, C.; Cruaud, C.; Maciazek, J.; Weissenbach, J.; Bousser,**
- [95243] **M.–G.; Bach, J.–F.; Tournier–Lasserre, E.: Strong clustering and**
- [95244] **stereotyped nature of Notch3 mutations in CADASIL patients. Lancet 350:**
- [95245] **1511–1515, 1997.**
- [95246] **16581.**
- [95247] **Rebay, I.; Fehon, R. G.; Artavanis–Tsakonas, S.: Specific truncations**
- [95248] **of Drosophila Notch define dominant activated and dominant negative**
- [95249] **forms of the receptor. Cell 74: 319–329, 1993.**
- [95250] **16582.**
- [95251] **Deed, R. W.; Hirose, T.; Mitchell, E. L. D.; Santibanez–Koref,**
- [95252] **M. F.; Norton, J. D.: Structural organisation and chromo–**

somal mapping

[95253] of the human Id-3 gene. Gene 151: 309-314, 1994.

[95254] 16583.

[95255] Ellmeier, W.; Aguzzi, A.; Kleiner, E.; Kurzbauer, R.; Weith, A.

[95256] : Mutually exclusive expression of a helix-loop-helix gene and N-myc

[95257] in human neuroblastomas and in normal development. EMBO J. 11: 2563-2571,

[95258] 1992.

[95259] 16584.

[95260] Kee, B. L.; Rivera, R. R.; Murre, C.: Id3 inhibits B lymphocyte

[95261] progenitor growth and survival in response to TGF-beta. Nature Immun. 2:

[95262] 242-247, 2001.

[95263] 16585.

[95264] Lyden, D.; Young, A. Z.; Zagzag, D.; Yan, W.; Gerald, W.; O'Reilly,

[95265] R.; Bader, B. L.; Hynes, R. O.; Zhuang, Y.; Manova, K.; Benezra, R.

- [95266] : Id1 and Id3 are required for neurogenesis, angiogenesis and vascularization
- [95267] of tumour xenografts. *Nature* 401: 670–677, 1999.
- [95268] 16586.
- [95269] Pan, L.; Sato, S.; Frederick, J. P.; Sun, X.–H.; Zhuang, Y.: Impaired
- [95270] immune responses and B–cell proliferation in mice lacking the Id3
- [95271] gene. *Molec. Cell. Biol.* 19: 5969–5980, 1999.
- [95272] 16587.
- [95273] Yeh, K.; Lim, R. W.: Genomic organization and promoter analysis
- [95274] of the murine Id3 gene. *Gene* 254: 163–171, 2000.
- [95275] 16588.
- [95276] Rubinfeld, B.; Munemitsu, S.; Clark, R.; Conroy, L.; Watt, K.;
- [95277] Crosier, W. J.; McCormick, F.; Polakis, P.: Molecular cloning of
- [95278] a GTPase activating protein specific for the Krev–1 protein p21–rap1. *Cell* 65:
- [95279] 1033–1042, 1991.

[95280] 16589.

[95281] Weiss, J.; Rubinfeld, B.; Polakis, P. G.; McCormick, F.;
Cavenee,

[95282] W. K.; Arden, K. C.: The RAP1GA1 locus for human
Rap1-GTPase activating

[95283] protein 1 maps to chromosome 1p36.1-p35. Cytogenet.
Cell Genet. 66:

[95284] 18-21, 1994.

[95285] 16590.

[95286] Braun, A.; Kammerer, S.; Weissenhorn, W.; Weiss, E. H.;
Cleve,

[95287] H.: Sequence of a putative human housekeeping gene
(HK33) localized

[95288] on chromosome 1. Gene 146: 291-295, 1994.

[95289] 16591.

[95290] Gotte, K.; Girzalsky, W.; Linkert, M.; Baumgart, E.; Kam-
merer,

[95291] S.; Kunau, W.-H.; Erdmann, R.: Pex19p, a farnesylated
protein essential

[95292] for peroxisome biogenesis. Molec. Cell. Biol. 18: 616-628,
1998.

- [95293] 16592.
- [95294] James, G. L.; Goldstein, J. L.; Pathak, R. K.; Anderson, R. G. W.; Brown, M. S.: PxF, a prenylated protein of peroxisomes J. Biol. Chem. 269: 14182–14190, 1994.
- [95297] 16593.
- [95298] Kammerer, S.; Arnold, N.; Gutensohn, W.; Mewes, H.–W.; Kunau, W.–H.;
- [95299] Hofler, G.; Roscher, A. A.; Braun, A.: Genomic organization and molecular
- [95300] characterization of a gene encoding HsPXF, a human peroxisomal farnesylated
- [95301] protein. Genomics 45: 200–210, 1997.
- [95302] 16594.
- [95303] Caskey, C. T.; Pizzuti, A.; Fu, Y.–H.; Fenwick, R. G., Jr.; Nelson,
- [95304] D. L.: Triplet repeat mutations in human disease. Science 256:
- [95305] 784–789, 1992.
- [95306] 16595.

- [95307] Chang, B.; Zheng, S. L.; Hawkins, G. A.; Isaacs, S. D.; Wiley, K. E.; Turner, A.; Carpten, J. D.; Bleecker, E. R.; Walsh, P. C.;
- [95309] Trent, J. M.; Meyers, D. A.; Isaacs, W. B.; Xu, J.: Polymorphic GGC repeats in the androgen receptor gene are associated with hereditary and sporadic prostate cancer risk. Hum. Genet. 110: 122–129, 2002.
- [95312] 16596.
- [95313] Chang, C.; Kokontis, J.; Liao, S.: Molecular cloning of human and rat complementary DNA encoding androgen receptors. Science 240: 324–326, 1988.
- [95316] 16597.
- [95317] Choong, C. S.; Kempainen, J. A.; Zhou, Z.-X.; Wilson, E. M.:
- [95318] Reduced androgen receptor gene expression with first exon CAG repeat expansion. Molec. Endocr. 10: 1527–1535, 1996.

- [95320] 16598.
- [95321] Choong, C. S.; Quigley, C. A.; French, F. S.; Wilson, E. M.:
- [95322] A novel missense mutation in the amino-terminal domain
of the human
- [95323] androgen receptor gene in a family with partial androgen
insensitivity
- [95324] syndrome causes reduced efficiency of protein translation.
J. Clin.
- [95325] Invest. 98: 1423-1431, 1996.
- [95326] 16599.
- [95327] Chu, J.; Zhang, R.; Zhao, Z.; Zou, W.; Han, Y.; Qi, Q.;
Zhang,
- [95328] H.; Wang, J.-C.; Tao, S.; Liu, X.; Luo, Z.: Male fertility is
compatible
- [95329] with an Arg840Cys substitution in the AR in a large Chi-
nese family
- [95330] with divergent phenotypes of AR insensitivity syndrome. J.
Clin.
- [95331] Endocr. Metab. 87: 347-351, 2002.
- [95332] 16600.
- [95333] Coetzee, G. A.; Ross, R. K.: Re: Prostate cancer and the

androgen

[95334] receptor. (Letter) J. Nat. Cancer Inst. 86: 872–873, 1994.

[95335] 16601.

[95336] Correa–Cerro, L.; Wöhr, G.; Haussler, J.; Berthon, P.;
Drelon,

[95337] E.; Mangin, P.; Fournier, G.; Cussenot, O.; Kraus, P.; Just,
W.; Paiss,

[95338] T.; Cantu, J. M.; Vogel, W.: (CAG)_nCAA and GGN repeats in
the human

[95339] androgen receptor gene are not associated with prostate
cancer in

[95340] a French–German population. Europ. J. Hum. Genet. 7:
347–362, 1999.

[95341] 16602.

[95342] Dowsing, A. T.; Yong, E. L.; Clark, M.; McLachlan, R. I.; de
Kretser,

[95343] D. M.; Trounson, A. O.: Linkage between male infertility
and trinucleotide

[95344] repeat expansion in the androgen–receptor gene. Lancet
354: 640–643,

[95345] 1999.

[95346] 16603.

- [95347] Elo, J. P.; Kvist, L.; Leinonen, K.; Isomaa, V.; Hentuu, P.;
Lukkarinen,
- [95348] O.; Vihko, P.: Mutated human androgen receptor gene de-
tected in a
- [95349] prostatic cancer patient is also activated by estradiol. J.
Clin.
- [95350] Endocr. Metab. 80: 3494–3500, 1995.
- [95351] 16604.
- [95352] Gaddipati, J. P.; McLeod, D. G.; Heidenberg, H. B.; Sester-
henn,
- [95353] I. A.; Finger, M. J.; Moul, J. W.; Srivastava, S.: Frequent de-
tection
- [95354] of codon 877 mutation in the androgen receptor gene in
advanced prostate
- [95355] cancers. Cancer Res. 54: 2861–2864, 1994.
- [95356] 16605.
- [95357] Gehring, U.; Tomkins, G. M.: Characterization of a hor-
mone receptor
- [95358] defect in the androgen–insensitivity mutant. Cell 3:
59–64, 1974.
- [95359] 16606.

- [95360] Gingrich, J. R.; Barrios, R. J.; Kattan, M. W.; Nahm, H. S.; Finegold,
[95361] M. J.; Greenberg, N. M.: Androgen-independent prostate
cancer progression
[95362] in the TRAMP model. *Cancer Res.* 57: 4687–4691, 1997.
[95363] 16607.
- [95364] Gingrich, J. R.; Greenberg, N. M.: A transgenic mouse
prostate
[95365] cancer model. *Toxicol. Path.* 24: 502–504, 1996.
[95366] 16608.
- [95367] Giovannucci, E.; Stampfer, M. J.; Krithivas, K.; Brown, M.; Brufsky,
[95368] A.; Talcott, J.; Hennekens, C. H.; Kantoff, P. W.: The CAG
repeat
[95369] within the androgen receptor gene and its relationship to
prostate
[95370] cancer. *Proc. Nat. Acad. Sci.* 94: 3320–3323, 1997.
[95371] 16609.
- [95372] Gottlieb, B.; Beitel, L. K.; Trifiro, M. A.: Variable expressiv-
ity
[95373] and mutation databases: the androgen receptor gene mu-

tations database. Hum.

[95374] **Mutat. 17: 382–388, 2001.**

[95375] **16610.**

[95376] **Gottlieb, B.; Trifiro, M.; Lumbroso, R.; Pinsky, L.: The an-
drogen**

[95377] **receptor gene mutations database. Nucleic Acids Res. 25:
158–162,**

[95378] **1997.**

[95379] **16611.**

[95380] **Gottlieb, B.; Trifiro, M.; Lumbroso, R.; Vasiliou, D. M.; Pin-
sky,**

[95381] **L.: The androgen receptor gene mutations database. Nu-
cleic Acids**

[95382] **Res. 24: 151–154, 1996.**

[95383] **16612.**

[95384] **Griffin, J. E.: Androgen resistance--the clinical and molec-
ular**

[95385] **spectrum. New Eng. J. Med. 326: 611–618, 1992.**

[95386] **16613.**

[95387] **Grino, P. B.; Isidro-Gutierrez, R. F.; Griffin, J. E.; Wilson,**

- [95388] J. D.: Androgen resistance associated with a qualitative abnormality
- [95389] of the androgen receptor and responsive to high dose androgen therapy. J.
- [95390] Clin. Endocr. Metab. 68: 578–584, 1989.
- [95391] 16614.
- [95392] Hardy, D. O.; Scher, H. I.; Bogenreider, T.; Sabbatini, P.; Zhang,
- [95393] Z.-F.; Nanus, D. M.; Catterall, J. F.: Androgen receptor CAG repeat
- [95394] lengths in prostate cancer: correlation with age of onset. J. Clin.
- [95395] Endocr. Metab. 81: 4400–4405, 1996.
- [95396] 16615.
- [95397] Hellwinkel, O. J.-C.; Holterhus, P.-M.; Struve, D.; Marschke,
- [95398] C.; Homburg, N.; Hiort, O.: A unique exonic splicing mutation in
- [95399] the human androgen receptor gene indicates a physiologic relevance
- [95400] of regular androgen receptor transcript variants. J. Clin. Endocr.

- [95401] Metab. 86: 2569–2575, 2001.
- [95402] 16616.
- [95403] Hickey, T.; Chandy, A.; Norman, R. J.: The androgen receptor
- [95404] CAG repeat polymorphism and X-chromosome inactivation in Australian
- [95405] Caucasian women with infertility related to polycystic ovary syndrome. J.
- [95406] Clin. Endocr. Metab. 87: 161–165, 2002.
- [95407] 16617.
- [95408] Hiort, O.; Sinnecker, G. H. G.; Holterhus, P.–M.; Nitsche, E.
- [95409] M.; Kruse, K.: Inherited and de novo androgen receptor gene mutations:
- [95410] investigation of single-case families. J. Pediat. 132: 939–943,
- [95411] 1998.
- [95412] 16618.
- [95413] Holterhus, P.–M.; Bruggenwirth, H. T.; Hiort, O.; Kleinkauf–Houcken,
- [95414] A.; Kruse, K.; Sinnecker, G. H. G.; Brinkmann, A. O.: Mo-
saicism due

- [95415] to a somatic mutation of the androgen receptor gene determines phenotype
- [95416] in androgen insensitivity syndrome. *J. Clin. Endocr. Metab.* 82:
- [95417] 3584–3589, 1997.
- [95418] 16619.
- [95419] Shibuya, H.; Yamaguchi, K.; Shirakabe, K.; Tonegawa, A.; Gotoh,
- [95420] Y.; Ueno, N.; Irie, K.; Nishida, E.; Matsumoto, K.: TAB1: an activator
- [95421] of the TAK1 MAPKKK in TGF- β signal transduction. *Science* 272:
- [95422] 1179–1182, 1996.
- [95423] 16620.
- [95424] Chadwick, B. P.; Obermayr, F.; Frischauf, A.-M.: FKHL15, a new
- [95425] human member of the forkhead gene family located on chromosome 9q22. *Genomics* 41:
- [95426] 390–396, 1997.
- [95427] 16621.
- [95428] De Felice, M.; Ovitt, C.; Biffali, E.; Rodriguez-Mallon, A.;

Arra,

[95429] C.; Anastassiadis, K.; Macchia, P. E.; Mattei, M.-G.; Mariano, A.;

[95430] Scholer, H.; Macchia, V.; Di Lauro, R.: A mouse model for hereditary

[95431] thyroid dysgenesis and cleft palate. *Nature Genet.* 19: 395–398,

[95432] 1998.

[95433] 16622.

[95434] Kaufmann, E.; Knochel, W.: Five years on the wings of fork head. *Mech.*

[95435] *Dev.* 57: 3–20, 1996.

[95436] 16623.

[95437] Manley, N. R.; Capecchi, M. R.: Hox group 3 paralogs regulate

[95438] the development and migration of the thymus, thyroid, and parathyroid

[95439] gland. *Dev. Biol.* 195: 1–15, 1998.

[95440] 16624.

[95441] Toubanc, J. E.: Comparison of epidemiological data on congenital

- [95442] hypothyroidism in Europe with those of other parts in the world. Horm.
- [95443] Res. 38: 230–235, 1992.
- [95444] 16625.
- [95445] Zannini, M.; Avantaggiato, V.; Biffali, E.; Arnone, M. I.; Sato,
- [95446] K.; Pischetola, M.; Taylor, B. A.; Phillips, S. J.; Simeone, A.; Di
- [95447] Lauro, R.: TTF–2, a new forkhead protein, shows a temporal expression
- [95448] in the developing thyroid which is consistent with a role in controlling
- [95449] the onset of differentiation. EMBO J. 16: 3185–3197, 1997.
- [95450] 16626.
- [95451] Boyd, J. M.; Subramanian, T.; Schaeper, U.; La Regina, M.; Bayley,
- [95452] S.; Chinnadurai, G.: A region in the C–terminus of adenovirus 2/5
- [95453] E1a protein is required for association with a cellular phosphoprotein
- [95454] and important for the negative modulation of T24–ras

mediated transformation,

[95455] tumorigenesis and metastasis. EMBO J. 469–478, 1993.

[95456] 16627.

[95457] Furusawa, T.; Moribe, H.; Kondoh, H.; Higashi, Y.: Identification

[95458] of CtBP1 and CtBP2 as corepressors of zinc finger-homeodomain factor

[95459] delta-EF1. Molec. Cell. Biol. 19: 8581–8590, 1999.

[95460] 16628.

[95461] Katsanis, N.; Fisher, E. M. C.: A novel C-terminal binding protein

[95462] (CTBP2) is closely related to CTBP1, an adenovirus E1A-binding protein,

[95463] and maps to human chromosome 21q21.3. Genomics 47: 294–299, 1998.

[95464] 16629.

[95465] Schaeper, U.; Boyd, J. M.; Verma, S.; Uhlmann, E.; Subramanian,

[95466] T.; Chinnadurai, G.: Molecular cloning and characterization of a

[95467] cellular phosphoprotein that interacts with a conserved C-

terminal

- [95468] domain of adenovirus E1A involved in negative modulation of oncogenic transformation. Proc. Nat. Acad. Sci. 92: 10467-10471, 1995.
- [95469] 16630.
- [95470] Sewalt, R. G. A. B.; Gunster, M. J.; van der Vlag, J.; Satijn, D. P. E.; Otte, A. P.: C-terminal binding protein is a transcriptional repressor that interacts with a specific class of vertebrate polycomb proteins. Molec. Cell. Biol. 19: 777-787, 1999.
- [95471] 16631.
- [95472] Zhang, Q.; Piston, D. W.; Goodman, R. H.: Regulation of corepressor function by nuclear NADH. Science 295: 1895-1897, 2002.
- [95473] 16632.
- [95474] Aya, K.; Tanaka, H.; Seino, Y.: Novel mutation in the nephrin gene of a Japanese patient with congenital nephrotic syn-

drome of the

[95481] Finnish type. *Kidney Int.* 57: 401–404, 2000.

[95482] 16633.

[95483] Beltcheva, O.; Martin, P.; Lenkkeri, U.; Tryggvason, K.: Mutation

[95484] spectrum in the nephrin gene (NPHS1) in congenital nephrotic syndrome. *Hum.*

[95485] *Mutat.* 17: 368–373, 2001.

[95486] 16634.

[95487] Guez, S.; Giani, M.; Melzi, M. L.; Antignac, C.; Assael, B. M.

[95488] : Adequate clinical control of congenital nephrotic syndrome by enalapril. *Pediat.*

[95489] *Nephrol.* 12: 130–132, 1998.

[95490] 16635.

[95491] Koziell, A.; Grech, V.; Hussain, S.; Lee, G.; Lenkkeri, U.; Tryggvason,

[95492] K.; Scambler, P.: Genotype/phenotype correlations of NPHS1 and NPHS2

[95493] mutations in nephrotic syndrome advocate a functional inter-relationship

[95494] in glomerular filtration. *Hum. Molec. Genet.* 11: 379–388,

2002.

[95495] 16636.

[95496] Lenkkeri, U.; Mannikko, M.; McCready, P.; Lamerdin, J.;
Gribouval,

[95497] O.; Niaudet, P.; Antignac, C.; Kashtan, C. E.; Holmberg, C.;
Olsen,

[95498] A.; Kestila, M.; Tryggvason, K.: Structure of the gene for
congenital

[95499] nephrotic syndrome of the Finnish type (NPHS1) and char-
acterization

[95500] of mutations. Am. J. Hum. Genet. 64: 51-61, 1999.

[95501] 16637.

[95502] Liu, L.; Done, S. C.; Khoshnoodi, J.; Bertorello, A.; Wartio-
vaara,

[95503] J.; Berggren, P.-O.; Tryggvason, K.: Defective nephrin traf-
ficking

[95504] caused by missense mutations in the NPHS1 gene: insight
into the mechanisms

[95505] of congenital nephrotic syndrome. Hum. Molec. Genet. 10:
2637-2644,

[95506] 2001.

[95507] 16638.

- [95508] Putaala, H.; Soininen, R.; Kilpelainen, P.; Wartiovaara, J.; Tryggvason, K.: The murine nephrin gene is specifically expressed in kidney, brain and pancreas: inactivation of the gene leads to massive proteinuria and neonatal death. Hum. Molec. Genet. 10: 1-8, 2001.
- [95512] 16639.
- [95513] Ruotsalainen, V.; Ljungberg, P.; Wartiovaara, J.; Lenkkeri, U.; Kestila, M.; Jalanko, H.; Holmberg, C.; Tryggvason, K.: Nephin is specifically located at the slit diaphragm of glomerular podocytes. Proc. Nat. Acad. Sci. 96: 7962-7967, 1999.
- [95517] 16640.
- [95518] Watanabe, T.; Inoue, S.; Hiroi, H.; Orimo, A.; Kawashima, H.; Muramatsu, M.: Isolation of estrogen-responsive genes with a CpG island library. Molec. Cell. Biol. 18: 442-449, 1998.

- [95521] 16641.
- [95522] Adra, C. N.; Kobayashi, H.; Rowley, J. D.; Lim, B.: Assign-
ment
- [95523] of the human GDID4 gene, a GDP/GTP-exchange regula-
tor, to chromosome
- [95524] 12p12.3. Genomics 24: 188-190, 1994.
- [95525] 16642.
- [95526] Leffers, H.; Nielsen, M. S.; Andersen, A. H.; Honore, B.;
Madsen,
- [95527] P.; Vandekerckhove, J.; Celis, J. E.: Identification of two
human
- [95528] Rho GDP dissociation inhibitor proteins whose overex-
pression leads
- [95529] to disruption of the actin cytoskeleton. Exp. Cell Res. 209:
165-174,
- [95530] 1993.
- [95531] 16643.
- [95532] Lelias, J.-M.; Adra, C. N.; Wulf, G. M.; Guillemot, J.-C.;
Khagad,
- [95533] M.; Caput, D.; Lim, B.: cDNA cloning of a human mRNA
preferentially

- [95534] expressed in hematopoietic cells and with homology to a GDP-dissociation
- [95535] inhibitor for the rho GTP-binding proteins. Proc. Nat. Acad. Sci. 90:
- [95536] 1479-1483, 1993.
- [95537] 16644.
- [95538] Scherle, P.; Behrens, T.; Staudt, L. M.: Ly-GDI, a GDP-dissociation
- [95539] inhibitor of the RhoA GTP-binding protein, is expressed preferentially
- [95540] in lymphocytes. Proc. Nat. Acad. Sci. 90: 7568-7572, 1993.
- [95541] 16645.
- [95542] Telford, E. A. R.; Wightman, P.; Leek, J.; Markham, A. F.; Lench,
- [95543] N. J.; Bonthron, D. T.: cDNA cloning, genomic organization, and chromosomal
- [95544] localization of a novel human gene that encodes a kinesin-related
- [95545] protein highly similar to mouse Kif3C. Biochem. Biophys. Res. Commun. 242:
- [95546] 407-412, 1998.

- [95547] 16646.
- [95548] Kalchman, M. A.; Graham, R. K.; Xia, G.; Koide, H. B.;
Hodgson,
- [95549] J. G.; Graham, K. C.; Goldberg, Y. P.; Gietz, R. D.; Pickart,
C. M.;
- [95550] Hayden, M. R.: Huntingtin is ubiquitinated and interacts
with a specific
- [95551] ubiquitin-conjugating enzyme. J. Biol. Chem. 271:
19385-19394, 1996.
- [95552] 16647.
- [95553] Chiariotti, L.; Benvenuto, G.; Fedele, M.; Santoro, M.;
Simeone,
- [95554] A.; Fusco, A.; Bruni, C. B.: Identification and characteriza-
tion
- [95555] of a novel RING-finger gene (RNF4) mapping at 4p16.3.
Genomics 47:
- [95556] 258-265, 1998.
- [95557] 16648.
- [95558] Ishikawa, K.; Nagase, T.; Suyama, M.; Miyajima, N.;
Tanaka, A.;
- [95559] Kotani, H.; Nomura, N.; Ohara, O.: Prediction of coding
sequences

- [95560] of unidentified human genes. X. The complete sequences of 100 new
- [95561] cDNA clones from brain which can code for large proteins in vitro. DNA
- [95562] Res. 5: 169–176, 1998.
- [95563] 16649.
- [95564] Waldegger, S.; Erdel, M.; Nagl, U. O.; Barth, P.; Raber, G.; Steuer,
- [95565] S.; Utermann, G.; Paulmichl, M.; Lang, F.: Genomic organization and
- [95566] chromosomal localization of the human SGK protein kinase gene. Genomics 51:
- [95567] 299–302, 1998.
- [95568] 16650.
- [95569] Webster, M. K.; Goya, L.; Ge, Y.; Maiyar, A. C.; Firestone, G.
- [95570] L.: Characterization of sgk, a novel member of the serine/threonine
- [95571] protein kinase gene family which is transcriptionally induced by glucocorticoids
- [95572] and serum. Molec. Cell. Biol. 13: 2031–2040, 1993.

- [95573] 16651.
- [95574] Aoki, K.; Ishida, R.; Kasai, M.: Isolation and characterization
of a cDNA encoding a translin-like protein, TRAX. FEBS
Lett. 401:
109–112, 1997.
- [95577] 16652.
- [95578] Meng, G.; Aoki, K.; Tokura, K.; Nakahara, K.; Inazawa, J.;
Kasai,
M.: Genomic structure and chromosomal localization of
the gene encoding
TRAX, a translin-associated factor X. J. Hum. Genet. 45:
305–308,
2000.
- [95582] 16653.
- [95583] Collins, C.; Rommens, J. M.; Kowbel, D.; Godfrey, T.; Tan-
ner, M.;
- [95584] Hwang, S.; Polikoff, D.; Nonet, G.; Cochran, J.; Myambo,
K.; Jay,
K. E.; Froula, J.; Cloutier, T.; Kuo, W.-L.; Yaswen, P.; Dair-
kee,

- [95586] S.; Giovanola, J.; Hutchinson, G. B.; Isola, J.; Kallioniemi, O.-P.;
- [95587] Palazzolo, M.; Martin, C.; Ericsson, C.; Pinkel, D.; Albertson, D.;
- [95588] Li, W.-B.; Gray, J. W.: Positional cloning of ZNF217 and NABC1: genes
- [95589] amplified at 20q13.2 and overexpressed in breast carcinoma. Proc.
- [95590] Nat. Acad. Sci. 95: 8703-8708, 1998.
- [95591] 16654.
- [95592] Eudy, J. D.; Yao, S.; Weston, M. D.; Ma-Edmonds, M.; Talmadge,
- [95593] C. B.; Cheng, J. J.; Kimberling, W. J.; Sumegi, J.: Isolation of
- [95594] a gene encoding a novel member of the nuclear receptor superfamily
- [95595] from the critical region of Usher syndrome type IIa at 1q41. Genomics 50:
- [95596] 382-384, 1998.
- [95597] 16655.
- [95598] Greschik, H.; Wurtz, J.-M.; Sanglier, S.; Bourguet, W.; van Dorsselaer,

- [95599] A.; Moras, D.; Renaud, J.-P.: Structural and functional evidence
- [95600] for ligand-independent transcriptional activation by the estrogen-related
- [95601] receptor 3. *Molec. Cell* 9: 303-313, 2002.
- [95602] 16656.
- [95603] Matsuyoshi, N.; Imamura, S.: Multiple cadherins are expressed
- [95604] in human fibroblasts. *Biochem. Biophys. Res. Commun.* 235: 355-358,
- [95605] 1997.
- [95606] 16657.
- [95607] Bertilsson, G.; Heidrich, J.; Svensson, K.; Asman, M.; Jendeberg,
- [95608] L.; Sydow-Backman, M.; Ohlsson, R.; Postlind, H.; Blomquist, P.; Berkenstam,
- [95609] A.: Identification of a human nuclear receptor defines a new signaling
- [95610] pathway for CYP3A induction. *Proc. Nat. Acad. Sci.* 95: 12208-12213,
- [95611] 1998.

- [95612] 16658.
- [95613] Blumberg, B.; Sabbagh, W., Jr.; Juguilon, H.; Bolado, J., Jr.;
- [95614] van Meter, C. M.; Ong, E. S.; Evans, R. M.: SXR, a novel
steroid
- [95615] and xenobiotic-sensing nuclear receptor. *Genes Dev.* 12:
3195-3205,
- [95616] 1998.
- [95617] 16659.
- [95618] Watkins, R. E.; Wisely, G. B.; Moore, L. B.; Collins, J. L.;
- Lambert,
- [95619] M. H.; Williams, S. P.; Willson, T. M.; Kliewer, S. A.; Red-
inbo, M.
- [95620] R.: The human nuclear xenobiotic receptor PXR: structural
determinants
- [95621] of directed promiscuity. *Science* 292: 2329-2333, 2001.
- [95622] 16660.
- [95623] Nemes, J. P.; Benzow, K. A.; Koob, M. D.: The SCA8 tran-
script
- [95624] is an antisense RNA to a brain-specific transcript encod-
ing a novel
- [95625] actin-binding protein (KLHL1). *Hum. Molec. Genet.* 9:

1543–1551,

[95626] 2000.

[95627] 16661.

[95628] Adato, A.; Raskin, L.; Petit, C.; Bonne–Tamir, B.: Deafness heterogeneity

[95629] in a Druze isolate from the Middle East: novel OTOF and PDS mutations,

[95630] low prevalence of GJB2 35delG mutation and indication for a new DFNB

[95631] locus. *Europ. J. Hum. Genet.* 8: 437–442, 2000.

[95632] 16662.

[95633] Yasunaga, S.; Grati, M.; Chardenoux, S.; Smith, T. N.; Friedman,

[95634] T. B.; Lalwani, A. K.; Wilcox, E. R.; Petit, C.: OTOF encodes multiple

[95635] long and short isoforms: genetic evidence that the long ones underlie

[95636] recessive deafness DFNB9. *Am. J. Hum. Genet.* 67: 591–600, 2000.

[95637] 16663.

[95638] Yasunaga, S.; Petit, C.: Physical map of the region sur–

rounding

[95639] the OTOFERLIN locus on chromosome 2p22–p23. Genomics 66: 110–112,

[95640] 2000.

[95641] 16664.

[95642] Boffa, M. B.; Reid, S.; Joo, E.; Nesheim, M. E.; Koschinsky, M.

[95643] L.: Characterization of the gene encoding human TAFI (thrombin–activable

[95644] fibrinolysis inhibitor; plasma procarboxypeptidase B). Biochemistry 38:

[95645] 6547–6558, 1999.

[95646] 16665.

[95647] Eaton, D. L.; Malloy, B. E.; Tsai, S. P.; Henzel, W.; Drayna, D.

[95648] : Isolation, molecular cloning, and partial characterization of a

[95649] novel carboxypeptidase B from human plasma. J. Biol. Chem. 266:

[95650] 21833–21838, 1991.

[95651] 16666.

- [95652] Koschinsky, M. L.; Boffa, M. B.; Nesheim, M. E.; Zinman, B.; Hanley,
- [95653] A. J. G.; Harris, S. B.; Cao, H.; Hegele, R. A.: Association of a
- [95654] single nucleotide polymorphism in CPB2 encoding the thrombin-activable
- [95655] fibrinolysis inhibitor (TAFI) with blood pressure. Clin. Genet. 60:
- [95656] 345-349, 2001.
- [95657] 16667.
- [95658] Tregouet, D.-A.; Aubert, H.; Henry, M.; Morange, P.; Visvikis,
- [95659] S.; Juhan-Vague, I.; Tired, L.: Combined segregation-linkage analysis
- [95660] of plasma thrombin-activatable fibrinolysis inhibitor (TAFI) antigen
- [95661] levels with TAFI gene polymorphisms. Hum. Genet. 109: 191-197, 2001.
- [95662] 16668.
- [95663] Tsai, S. P.; Drayna, D.: The gene encoding human plasma carboxypeptidase
- [95664] B (CPB2) resides on chromosome 13. Genomics 14:

549–550, 1992.

[95665] 16669.

[95666] Vanhoof, G.; Wauters, J.; Schatteman, K.; Hendriks, D.;
Goossens,

[95667] F.; Bossuyt, P.; Scharpe, S.: The gene for human car-
boxypeptidase

[95668] U (CPU)--a proposed novel regulator of plasminogen acti-
vation--maps

[95669] to 13q14.11. Genomics 38: 454–455, 1996.

[95670] 16670.

[95671] Zhao, L.; Morser, J.; Bajzar, L.; Nesheim, M.; Nagashima,
N.:

[95672] Identification and characterization of two thrombin-acti-
vatable fibrinolysis

[95673] inhibitor isoforms. Thromb. Haemost. 80: 949–955, 1998.

[95674] 16671.

[95675] McGwire, G. B.; Tan, F.; Michel, B.; Rehli, M.; Skidgel, R. A.

[95676] : Identification of a membrane-bound carboxypeptidase
as the mammalian

[95677] homolog of duck gp180, a hepatitis B virus-binding pro-
tein. Life

[95678] Sci. 60: 715–724, 1997.

[95679] 16672.

[95680] Riley, D. A.; Tan, F.; Miletich, D. J.; Skidgel, R. A.: Chromosomal

[95681] localization of the genes for human carboxypeptidase D (CPD) and the

[95682] active 50–kilodalton subunit of human carboxypeptidase N (CPN1). Genomics 50:

[95683] 105–108, 1998.

[95684] 16673.

[95685] Tan, F.; Rehli, M.; Krause, S. W.; Skidgel, R. A.: Sequence of

[95686] human carboxypeptidase D reveals it to be a member of the regulatory

[95687] carboxypeptidase family with three tandem active site domains. Biochem.

[95688] J. 327: 81–87, 1997.

[95689] 16674.

[95690] Rajadhyaksha, A. Riviere, M.; Van Vooren, P.; Szpirer, J.; Szpirer,

[95691] C.; Babin, J.; Bina, M.: Assignment of AR1, transcription

factor

[95692] 20 (TCF20), to human chromosome 22q13.3 with somatic cell hybrids

[95693] and in situ hybridization. Cytogenet. Cell Genet. 81: 176–177, 1998.

[95694] 16675.

[95695] Sanz, L.; Moscat, J.; Diaz-Meco, M. T.: Molecular characterization

[95696] of a novel transcription factor that controls stromelysin expression. Molec.

[95697] Cell. Biol. 15: 3164–3170, 1995.

[95698] 16676.

[95699] Ding, H.; Descheemaeker, K.; Marynen, P.; Nelles, L.; Carvalho,

[95700] T.; Carmo-Fonseca, M.; Collen, D.; Belayew, A.: Characterization

[95701] of a helicase-like transcription factor involved in the expression

[95702] of the human plasminogen activator inhibitor-1 gene. DNA Cell Biol. 15:

[95703] 429–442, 1996.

[95704] 16677.

- [95705] Lin, Y.; Sheridan, P. L.; Jones, K. A.; Evans, G. A.: The
HIP116
- [95706] SNF2/SWI2-related transcription factor gene (SNF2L3) is
located on
- [95707] human chromosome 3q25.1-q26.1 Genomics 27:
381-382, 1995.
- [95708] 16678.
- [95709] Moinova, H. R.; Chen, W.-D.; Shen, L.; Smiraglia, D.;
Olechnowicz,
- [95710] J.; Ravi, L.; Kasturi, L.; Myeroff, L.; Plass, C.; Parsons, R.;
Minna,
- [95711] J.; Willson, J. K. V.; Green, S. B.; Issa, J.-P.; Markowitz, S. D.
- [95712] : HLTF gene silencing in human colon cancer. Proc. Nat.
Acad. Sci. 99:
- [95713] 4562-4567, 2002.
- [95714] 16679.
- [95715] Sheridan, P. L.; Schorpp, M.; Voz, M. L.; Jones, K. A.:
Cloning
- [95716] of an SNF2/SWI2-related protein that binds specifically to
the SPH
- [95717] motifs of the SV40 enhancer and to the HIV-1 promoter. J.
Biol. Chem. 270:

- [95718] 4575–4587, 1995.
- [95719] 16680.
- [95720] Ambros, P. F.; Schmid, J.; Rumpler, S.; Binder, B. R.; de
Martin,
- [95721] R.: Localization of the human I-kappa-B kinase-beta
(IKBKB) to chromosome
- [95722] 8p11.2 by fluorescence in situ hybridization and radiation
hybrid
- [95723] mapping. Genomics 54: 575–576, 1998.
- [95724] 16681.
- [95725] Pasparakis, M.; Courtois, G.; Hafner, M.; Schmidt-Sup-
prian, M.;
- [95726] Nenci, A.; Toksoy, A.; Krampert, M.; Goebeler, M.;
Gillitzer, R.;
- [95727] Israel, A.; Krieg, T.; Rajewsky, K.; Haase, I.: TNF-mediated
inflammatory
- [95728] skin disease in mice with epidermis-specific deletion of
IKK2. Nature 417:
- [95729] 861–866, 2002.
- [95730] 16682.
- [95731] Rossi, A.; Kapahi, P.; Natoli, G.; Takahashi, T.; Chen, Y.;

Karin,

[95732] M.; Santoro, M. G.: Anti-inflammatory cyclopentenone
prostaglandins

[95733] are direct inhibitors of I-kappa-B kinase. Nature 403:
103-108,

[95734] 2000.

[95735] 16683.

[95736] Shindo, M.; Nakano, H.; Sakon, S.; Yagita, H.; Mihara, M.;
Okumura,

[95737] K.: Assignment of I-kappa-B kinase beta (IKBKB) to human
chromosome

[95738] band 8p12-p11 by in situ hybridization. Cytogenet. Cell
Genet. 82:

[95739] 32-33, 1998.

[95740] 16684.

[95741] Tang, G.; Yang, J.; Minemoto, Y.; Lin, A.: Blocking cas-
pase-3-mediated

[95742] proteolysis of IKK-beta suppresses TNF-alpha-induced
apoptosis. Molec.

[95743] Cell 8: 1005-1016, 2001.

[95744] 16685.

[95745] Woronicz, J. D.; Gao, X.; Cao, Z.; Rothe, M.; Goeddel, D. V.:

[95746] IkappaB kinase-beta: NF-kappa-B activation and complex formation with

[95747] IkappaB kinase-alpha and NIK. Science 278: 866-869, 1997.

[95748] 16686.

[95749] Yin, M.-J.; Yamamoto, Y.; Gaynor, R. B.: The anti-inflammatory

[95750] agents aspirin and salicylate inhibit the activity of I-kappa-B kinase-beta. Nature 396:

[95751] 77-80, 1998.

[95752] 16687.

[95753] Zandi, E.; Rothwarf, D. M.; Delhase, M.; Hayakawa, M.; Karin,

[95754] M.: The IkappaB kinase complex (IKK) contains two kinase subunits,

[95755] IKKalpha and IKKbeta, necessary for IkappaB phosphorylation and NF-kappa-B

[95756] activation. Cell 91: 243-252, 1997.

[95757] 16688.

[95758] Aoki, M.; Hamada, F.; Sugimoto, T.; Sumida, S.; Akiyama,

T.; Toyoshima,

[95759] K.: The human cot proto-oncogene encodes two protein serine/threonine

[95760] kinases with different transforming activities by alternative initiation

[95761] of translation. J. Biol. Chem. 268: 22723–22732, 1993.

[95762] 16689.

[95763] Bataille, V.; Hiles, R.; Bishop, J. A. N.: Retinoblastoma, melanoma

[95764] and the atypical mole syndrome. Brit. J. Derm. 132: 134–138, 1995.

[95765] 16690.

[95766] Chilosi, M.; Doglioni, C.; Yan, Z.; Lestani, M.; Menestrina, F.;

[95767] Sorio, C.; Benedetti, A.; Vinante, F.; Pizzolo, G.; Inghirami, G.

[95768] : Differential expression of cyclin-dependent kinase 6 in cortical

[95769] thymocytes and T-cell lymphoblastic lymphoma/leukemia. Am. J. Pathol. 152:

[95770] 209–217, 1998.

[95771] 16691.

- [95772] Costello, J. F.; Plass, C.; Arap, W.; Chapman, V. M.; Held, W.
- [95773] A.; Berger, M. S.; Su Huang, H. J.; Cavenee, W. K.: Cyclin-dependent
- [95774] kinase 6 (cdk6) amplification in human gliomas identified using two-dimensional
- [95775] separation of genomic DNA. Cancer Res. 57: 1250–1254, 1997.
- [95776] 16692.
- [95777] Guan, K.-L.; Jenkins, C. W.; Li, Y.; Nichols, M. A.; Wu, X.; O'Keefe,
- [95778] C. L.; Matera, A. G.; Xiong, Y.: Growth suppression by p18, a p16(INK4/MTS1)–
- [95779] and p14(INK4B/MTS2)–related CDK6 inhibitor, correlates with wild-type
- [95780] pRb function. Genes Dev. 8: 2939–2952, 1994.
- [95781] 16693.
- [95782] Lien, H.-C.; Lin, C.-W.; Huang, P.-H.; Chang, M.-L.; Hsu, S.-M.
- [95783] : Expression of cyclin-dependent kinase 6 (cdk6) and frequent loss
- [95784] of CD44 in nasal–nasopharyngeal NK/T-cell lymphomas:

comparison with

[95785] CD56-negative peripheral T-cell lymphomas. Lab. Invest.
80: 893-900,

[95786] 2000.

[95787] 16694.

[95788] Meyerson, M.; Harlow, E.: Identification of G1 kinase activ-
ity

[95789] for cdk6, a novel cyclin D partner. Molec. Cell. Biol. 14:
2077-2086,

[95790] 1994.

[95791] 16695.

[95792] Shennan, M. G.; Badin, A.-C.; Walsh, S.; Summers, A.;
From, L.;

[95793] McKenzie, M.; Goldstein, A. M.; Tucker, M. A.; Hogg, D.;
Lassam, N.

[95794] : Lack of germline CDK6 mutations in familial melanoma.
Oncogene 19:

[95795] 1849-1852, 2000.

[95796] 16696.

[95797] Traboulsi, E. I.; Zimmerman, L. E.; Manz, H. J.: Cutaneous
malignant

[95798] melanoma in survivors of heritable retinoblastoma. Arch.
Ophthalmol. 106:

[95799] 1059-1061, 1988.

[95800] 16697.

[95801] Blais, A.; Labrie, Y.; Pouliot, F.; Lachance, Y.; Labrie, C.:

- [95802] Structure of the gene encoding the human cyclin-dependent kinase inhibitor
- [95803] p18 and mutational analysis in breast cancer. Biochem. Biophys. Res.
- [95804] Commun. 247: 146–153, 1998.
- [95805] 16698.
- [95806] Lapointe, J.; Lachance, Y.; Labrie, Y.; Labrie, C.: A p18 mutant
- [95807] defective in CDK6 binding in human breast cancer cells. Cancer Res. 56:
- [95808] 4586–4589, 1996.
- [95809] 16699.
- [95810] Bontemps, Y.; Maquart, F.-X.; , Wegrowski, Y.: Human UDP-glucose
- [95811] dehydrogenase gene: complete cloning and transcription start mapping. Biochem.
- [95812] Biophys. Res. Commun. 275: 981–985, 2000.
- [95813] 16700.
- [95814] Hempel, J.; Perozich, J.; Romovacek, H.; Hinich, A.; Kuo, I.; Feingold,
- [95815] D. S.: UDP-glucose dehydrogenase from bovine liver: pri-

mary structure

[95816] and relationship to other dehydrogenases. Protein Sci. 3: 1074–1080,

[95817] 1994.

[95818] 16701.

[95819] Marcu, O.; Stathakis, D. G.; Marsh, J. L.: Assignment of the UGDH

[95820] locus encoding UDP–glucose dehydrogenase to human chromosome band

[95821] 4p15.1 by radiation hybrid mapping. Cytogenet. Cell Genet. 86: 244–245,

[95822] 1999.

[95823] 16702.

[95824] Spicer, A. P.; Kaback, L. A.; Smith, T. J.; Seldin, M. F.: Molecular

[95825] cloning and characterization of the human and mouse UDP–glucose dehydrogenase

[95826] genes. J. Biol. Chem. 273: 25117–25124, 1998.

[95827] 16703.

[95828] Walsh, E. C.; Stainier, D. Y. R.: UDP–glucose dehydrogenase required

- [95829] for cardiac valve formation in zebrafish. Science 293:
1670–1674,
- [95830] 2001.
- [95831] 16704.
- [95832] Akamizu, T.; Ikuyama, S.; Saji, M.; Kosugi, S.; Kozak, C.;
McBride,
- [95833] O. W.; Kohn, L. D.: Cloning, chromosomal assignment, and
regulation
- [95834] of the rat thyrotropin receptor: expression of the gene is
regulated
- [95835] by thyrotropin, agents that increase cAMP levels, and thy-
roid autoantibodies. Proc.
- [95836] Nat. Acad. Sci. 87: 5677–5681, 1990.
- [95837] 16705.
- [95838] Reyes, R.; Duprat, F.; Lesage, F.; Fink, M.; Salinas, M.; Far-
man,
- [95839] N.; Lazdunski, M.: Cloning and expression of a novel pH-
sensitive
- [95840] two pore domain K⁺ channel from human kidney. J. Biol.
Chem. 273:
- [95841] 30863–30869, 1998.

- [95842] 16706.
- [95843] Becker, W.; Weber, Y.; Wetzel, K.; Eirmbter, K.; Tejedor, F. J.;
- [95844] Joost, H.-G.: Sequence characteristics, subcellular localization,
- [95845] and substrate specificity of DYRK-related kinases, a novel family
- [95846] of dual specificity protein kinases. J. Biol. Chem. 273: 25893-25902,
- [95847] 1998.
- [95848] 16707.
- [95849] Good, D.; Busfield, F.; Duffy, D.; Lovelock, P. K.; Kesting, J.
- [95850] B.; Cameron, D. P.; Shaw, J. T. E.: Familial Paget's disease of bone:
- [95851] nonlinkage to the PDB1 and PDB2 loci on chromosomes 6p and 18q in
- [95852] a large pedigree. J. Bone Miner. Res. 16: 33-38, 2001.
- [95853] 16708.
- [95854] Hocking, L.; Slee, F.; Haslam, S. I.; Cundy, T.; Nicholson, G.;
- [95855] van Hul, W.; Ralston, S. H.: Familial Paget's disease of

bone: patterns

[95856] of inheritance and frequency of linkage to chromosome
18q. Bone 26:

[95857] 577–580, 2000.

[95858] 16709.

[95859] Marco-Mingot, M.; San-Millan, J. L.; Wuyts, W.; Bachiller-
Corral,

[95860] J.; Van Hul, W.; Morales-Piga, A. A.: Lack of mutations in
the RANK

[95861] gene in Spanish patients with Paget disease of bone.
(Letter) Clin.

[95862] Genet. 60: 86–88, 2001.

[95863] 16710.

[95864] Li, J.; Sarosi, I.; Yan, X.-Q.; Morony, S.; Capparelli, C.; Tan,

[95865] H.-L.; McCabe, S.; Elliott, R.; Scully, S.; Van, G.; Kaufman,
S.;

[95866] Juan, S.-C.; Sun, Y.; Tarpley, J.; Martin, L.; Christensen, K.;
McCabe,

[95867] J.; Kostenuik, P.; Hsu, H.; Fletcher, F.; Dunstan, C. R.;
Lacey, D.

[95868] L.; Boyle, W. J.: RANK is the intrinsic hematopoietic cell

surface

[95869] receptor that controls osteoclastogenesis and regulation of bone mass

[95870] and calcium metabolism. Proc. Nat. Acad. Sci. 97: 1566–1571, 2000.

[95871] 16711.

[95872] Nakagawa, N.; Kinosaki, M.; Yamaguchi, K.; Shima, N.; Yasuda, H.;

[95873] Yano, K.; Morinaga, T.; Higashio, K.: RANK is the essential signaling

[95874] receptor for osteoclast differentiation factor in osteoclastogenesis. Biochem.

[95875] Biophys. Res. Commun. 253: 395–400, 1998.

[95876] 16712.

[95877] Wuyts, W.; Van Wesenbeeck, L.; Morales-Piga, A.; Ralston, S.;

[95878] Hocking, L.; Vanhoenacker, F.; Westhovens, R.; Verbruggen, L.; Anderson,

[95879] D.; Hughes, A.; Van Hul, W.: Evaluation of the role of RANK and OPG

[95880] genes in Paget's disease of bone. Bone 28: 104–107, 2001.

- [95881] 16713.
- [95882] Devilder, M.-C.; Cadoret, E.; Cherel, M.; Moreau, I.; Ron-
deau,
- [95883] G.; Bezieau, S.; Moisan, J.-P.: cDNA cloning, gene charac-
terization
- [95884] and 13q14.3 chromosomal assignment of CHC1-L, a
chromosome condensation
- [95885] regulator-like guanine nucleotide exchange factor. Ge-
nomics 54:
- [95886] 99-106, 1998.
- [95887] 16714.
- [95888] Phillips, N. J.; Zeigler, M. R.; Deaven, L. L.: A cDNA from
the
- [95889] ovarian cancer critical region of deletion on chromosome
17p13.3. Cancer
- [95890] Lett. 102: 85-90, 1996.
- [95891] 16715.
- [95892] Schultz, D. C.; Vanderveer, L.; Berman, D. B.; Hamilton, T.
C.;
- [95893] Wong, A. J.; Godwin, A. K.: Identification of two candidate
tumor

- [95894] suppressor genes on chromosome 17p13.3. Cancer Res. 56: 1997–2002,
- [95895] 1996.
- [95896] 16716.
- [95897] Peyrard, M.; Parveneh, S.; Lagercrantz, S.; Ekman, M.; Fransson,
- [95898] I.; Sahlen, S.; Dumanski, J. P.: Cloning, expression pattern, and
- [95899] chromosomal assignment to 16q23 of the human gamma-adaptin gene (ADTG). Genomics 50:
- [95900] 275–280, 1998.
- [95901] 16717.
- [95902] Adamou, J. E.; Aiyar, N.; Van Horn, S.; Elshourbagy, N. A.: Cloning
- [95903] and functional characterization of the human vasoactive intestinal
- [95904] peptide (VIP)–2 receptor. Biochem. Biophys. Res. Commun. 209: 385–392,
- [95905] 1995.
- [95906] 16718.
- [95907] Harmar, A. J.; Marston, H. M.; Shen, S.; Spratt, C.; West, K.

M.;

[95908] Sheward, W. J.; Morrison, C. F.; Dorin, J. R.; Piggins, H. D.;
Reubi,

[95909] J.-C.; Kelly, J. S.; Maywood, E. S.; Hastings, M. H.: The
VPAC2 receptor

[95910] is essential for circadian function in the mouse suprachiasmatic nuclei. Cell 109:

[95911] 497–508, 2002.

[95912] 16719.

[95913] Mackay, M.; Fantes, J.; Scherer, S.; Boyle, S.; West, K.; Tsui,

[95914] L.-C.; Belloni, E.; Lutz, E.; Van Heyningen, V.; Harmar, A.
J.: Chromosomal

[95915] localization in mouse and human of the vasoactive intestinal peptide

[95916] receptor type 2 gene: a possible contributor to the holoprosencephaly

[95917] 3 phenotype. Genomics 37: 345–353, 1996.

[95918] 16720.

[95919] Svoboda, M.; Tastenoy, M.; Van Rampelbergh, J.;
Goossens, J.-F.;

[95920] De Neef, P.; Waelbroeck, M.; Robberecht, P.: Molecular

cloning and

[95921] functional characterization of a human VIP receptor from SUP-T1 lymphoblasts. *Biochem.*

[95922] *Biophys. Res. Commun.* 205: 1617-1624, 1994.

[95923] 16721.

[95924] Andre, E.; Conquet, F.; Steinmayr, M.; Stratton, S. C.; Porciatti,

[95925] V.; Becker-Andre, M.: Disruption of retinoid-related orphan receptor

[95926] beta changes circadian behavior, causes retinal degeneration and leads

[95927] to vacillans phenotype in mice. *EMBO J.* 17: 3867-3877, 1998.

[95928] 16722.

[95929] Paravicini, G.; Steinmayr, M.; Andre, E.; Becker-Andre, M.: The

[95930] metastasis suppressor candidate nucleotide diphosphate kinase NM23

[95931] specifically interacts with members of the ROR/RZR nuclear orphan

[95932] receptor subfamily. *Biochem. Biophys. Res. Commun.* 227: 82-87, 1996.

[95933] 16723.

[95934] Sirlin, J. L.: Vacillans, a neurological mutant in the house mouse

[95935] linked with brown. J. Genet. 54: 42–48, 1956.

[95936] 16724.

[95937] Bourdi, M.; Demady, D.; Martin, J. L.; Jabbour, S. K.; Martin,

[95938] B. M.; George, J. W.; Pohl, L. R.: cDNA cloning and baculovirus expression

[95939] of the human liver endoplasmic reticulum P58: characterization as

[95940] a protein disulfide isomerase isoform, but not as a protease or a

[95941] carnitine acyltransferase. Arch. Biochem. Biophys. 323: 397–403,

[95942] 1995.

[95943] 16725.

[95944] Briquet-Laugier, V.; Xia, Y.-R.; Rooke, K.; Mehrabian, M.; Lusi,

[95945] A. J.; Doolittle, M. H.: Mapping of three members of the mouse protein

- [95946] disulfide isomerase family. Mammalian Genome 9: 176–177, 1998.
- [95947] 16726.
- [95948] Hirano, N.; Shibasaki, F.; Sakai, R.; Tanaka, T.; Nishida, J.;
- [95949] Yazaki, Y.; Takenawa, T.; Hirai, H.: Molecular cloning of the human
- [95950] glucose-regulated protein ERp57/GRP58, a thiol-dependent reductase:
- [95951] identification of its secretory form and inducible expression by the
- [95952] oncogenic transformation. Europ. J. Biochem. 234: 336–342, 1995.
- [95953] 16727.
- [95954] Koivunen, P.; Helaakoski, T.; Annunen, P.; Veijola, J.; Raisanen,
- [95955] S.; Pihlajaniemi, T.; Kivirikko, K. I.: ERp60 does not substitute
- [95956] for protein disulphide isomerase as the beta-subunit of prolyl 4-hydroxylase. Biochem.
- [95957] J. 316: 599–605, 1996.
- [95958] 16728.

- [95959] Koivunen, P.; Horelli-Kuitunen, N.; Helaakoski, T.; Karvonen, P.;
- [95960] Jaakkola, M.; Palotie, A.; Kivirikko, K. I.: Structures of the human
- [95961] gene for the protein disulfide isomerase-related polypeptide ERp60
- [95962] and a processed gene and assignment of these genes to 15q15 and 1q21. Genomics 42:
- [95963] 397-404, 1997.
- [95964] 16729.
- [95965] Oliver, J. D.; van der Wal, F. J.; Bulleid, N. J.; High, S.: Interaction
- [95966] of the thiol-dependent reductase ERp57 with nascent glycoproteins. Science 275:
- [95967] 86-88, 1997.
- [95968] 16730.
- [95969] Berthelsen, J.; Viggiano, L.; Schulz, H.; Ferretti, E.; Con-
salez,
- [95970] G. G.; Rocchi, M.; Blasi, F.: PKNOX1, a gene encoding PREP1, a new
- [95971] regulator of Pbx activity, maps on human chromosome 21q22.3 and murine

- [95972] chromosome 17B/C. *Genomics* 47: 323–324, 1998.
- [95973] 16731.
- [95974] Chen, H.; Rossier, C.; Nakamura, Y.; Lynn, A.; Chakravarti, A.;
- [95975] Antonarakis, S. E.: Cloning of a novel homeobox–containing gene,
- [95976] PKNOX1, and mapping to human chromosome 21q22.3. *Genomics* 41: 193–200,
- [95977] 1997.
- [95978] 16732.
- [95979] Morita, K.; Furuse, M.; Fujimoto, K.; Tsukita, S.: Claudin multigene
- [95980] family encoding four–transmembrane domain protein components of tight
- [95981] junction strands. *Proc. Nat. Acad. Sci.* 96: 511–516, 1999.
- [95982] 16733.
- [95983] Chiang, P.–W.; Fogel, E.; Jackson, C. L.; Lieuallen, K.; Lennon,
- [95984] G.; Qu, X.; Wang, S.–Q.; Kurnit, D. M.: Isolation, sequencing, and
- [95985] mapping of the human homologue of the yeast transcrip–

tion factor,

[95986] SPT5. Genomics 38: 421–424, 1996.

[95987] 16734.

[95988] Chiang, P.-W.; Stubbs, L.; Zhang, L.; Kurnit, D. M.: Isolation

[95989] of murine SPT5 homologue: completion of the isolation and characterization

[95990] of human and murine homologues of yeast chromatin structural protein

[95991] complex SPT4, SPT5, and SPT6. Genomics 47: 426–428, 1998.

[95992] 16735.

[95993] Iwaki, A.; Nagano, T.; Nakagawa, M.; Iwaki, T.; Fukumaki, Y.:

[95994] Identification and characterization of the gene encoding a new member

[95995] of the alpha-crystallin/Small hsp family, closely linked to the alpha-B-crystallin

[95996] gene in a head-to-head manner. Genomics 45: 386–394, 1997.

[95997] 16736.

[95998] de Coo, R. F. M.; Buddiger, P.; Smeets, H. J. M.; van Oost,

B.

- [95999] A.: Molecular cloning and characterization of the human mitochondrial
- [96000] NADH:oxidoreductase 10-kDa gene (NDUFV3). Genomics 45: 434–437,
- [96001] 1997.
- [96002] 16737.
- [96003] Canu, N.; Possenti, R.; Ricco, A. S.; Rocchi, M.; Levi, A.: Cloning,
- [96004] structural organization analysis, and chromosomal assignment of the
- [96005] human gene for the neurosecretory protein VGF. Genomics 45: 443–446,
- [96006] 1997.
- [96007] 16738.
- [96008] Ferri, G.-L.; Possenti, R.: vgf: a neurotrophin-inducible gene
- [96009] expressed in neuroendocrine tissues. Trends Endocr. Metab. 7: 233–239,
- [96010] 1996.
- [96011] 16739.

[96012] Levi, A.; Eldridge, J. D.; Paterson, B. M.: Molecular cloning
[96013] of a gene sequence regulated by nerve growth factor. Sci-
ence 229:
[96014] 393–395, 1985.
[96015] 16740.
[96016] Salton, S. R. J.; Fischberg, D. J.; Dong, K. W.: Structure of
[96017] the gene encoding VGF, a nervous system-specific mRNA
that is rapidly
[96018] and selectively induced by nerve growth factor in PC12
cells. Molec.
[96019] Cell. Biol. 11: 2335–2349, 1991.
[96020] 16741.
[96021] Gautvik, K. M.; de Lecea, L.; Gautvik, V. T.; Danielson, P.
E.;
[96022] Tranque, P.; Dopazo, A.; Bloom, F. E.; Sutcliffe, J. G.:
Overview
[96023] of the most prevalent hypothalamus-specific mRNAs as
identified by
[96024] directional tag PCR subtraction. Proc. Nat. Acad. Sci. 93:
8733–8738,
[96025] 1996.

[96026] 16742.

[96027] Green, E. D.; Olson, M. V.: Chromosomal region of the
cystic

[96028] fibrosis gene in yeast artificial chromosomes: a model for
human genome

[96029] mapping. Science 250: 94–98, 1990.

[96030] 16743.

[96031] Fritzler, M. J.; Lung, C.-C.; Hamel, J. C.; Griffith, K. J.;
Chan,

[96032] E. K. L.: Molecular characterization of golgin-245, a novel
Golgi

[96033] complex protein containing a granin signature. J. Biol.
Chem. 270:

[96034] 31262–31268, 1995.

[96035] 16744.

[96036] Gencik, M.; Dahmen, N.; Wieczorek, S.; Kasten, M.; Bier-
brauer,

[96037] J.; Anghelescu, I.; Szegedi, A.; Menezes Saecker, A. M.;
Epplen, J.

[96038] T.: A prepro-orexin gene polymorphism is associated with
narcolepsy. Neurology 56:

- [96039] 115–117, 2001.
- [96040] 16745.
- [96041] Hagan, J. J.; Leslie, R. A.; Patel, S.; Evans, M. L.; Wattam, T.
- [96042] A.; Holmes, S.; Benham, C. D.; Taylor, S. G.; Routledge, C.; Hemmati,
- [96043] P.; Munton, R. P.; Ashmeade, T. E.; Shah, A. S.; Hatcher, J. P.; Hatcher,
- [96044] P. D.; Jones, D. N. C.; Smith, M. I.; Piper, D. C.; Hunter, A. J.;
- [96045] Porter, R. A.; Upton, N.: Orexin A activates locus coeruleus cell
- [96046] firing and increases arousal in the rat. Proc. Nat. Acad. Sci. 96:
- [96047] 10911–10916, 1999.
- [96048] 16746.
- [96049] Hara, J.; Beuckmann, C. T.; Nambu, T.; Willie, J. T.; Chemelli,
- [96050] R. M.; Sinton, C. M.; Sugiyama, F.; Yagami, K.; Goto, K.; Yanagisawa,
- [96051] M.; Sakurai, T.: Genetic ablation of orexin neurons in mice results

- [96052] in narcolepsy, hypophagia, and obesity. *Neuron* 30: 345–354, 2001.
- [96053] 16747.
- [96054] Hungs, M.; Lin, L.; Okun, M.; Mignot, E.: Polymorphisms in the
- [96055] vicinity of the hypocretin/orexin are not associated with human narcolepsy. *Neurology* 57:
- [96056] 1893–1895, 2001.
- [96057] 16748.
- [96058] Nishino, S.; Ripley, B.; Overeem, S.; Lammers, G. J.; Mignot, E.
- [96059] : Hypocretin (orexin) deficiency in human narcolepsy. *Lancet* 355:
- [96060] 39–40, 2000.
- [96061] 16749.
- [96062] Chen, L.-S. K.; Lo, C. F.; Numann, R.; Cuddy, M.: Characterization
- [96063] of the human and rat phospholemmann (PLM) cDNAs and localization of
- [96064] the human PLM gene to chromosome 19q13.1. *Genomics* 41: 435–443,

- [96065] 1997.
- [96066] 16750.
- [96067] Sakurai, T.; Amemiya, A.; Ishii, M.; Matsuzaki, I.; Chemelli,
- [96068] R. M.; Tanaka, H.; Williams, S. C.; Richardson, J. A.; Ko-
złowski,
- [96069] G. P.; Wilson, S.; Arch, J. R. S.; Buckingham, R. E.; Haynes,
A. C.;
- [96070] Carr, S. A.; Annan, R. S.; McNulty, D. E.; Liu, W.-S.; Terrett,
J.
- [96071] A.; Elshourbagy, N. A.; Bergsma, D. J.; Yanagisawa, M.:
Orexins and
- [96072] orexin receptors: a family of hypothalamic neuropeptides
and G protein-coupled
- [96073] receptors that regulate feeding behavior. Cell 92:
573-585, 1998.
- [96074] 16751.
- [96075] Heard, D. J.; Norby, P. L.; Holloway, J.; Vissing, H.: Human
ERR-gamma,
- [96076] a third member of the estrogen receptor-related receptor
(ERR) subfamily
- [96077] of orphan nuclear receptors: tissue-specific isoforms are
expressed

- [96078] during development in the adult. *Molec. Endocr.* 14: 382–392, 2000.
- [96079] 16752.
- [96080] Hong, H.; Yang, L.; Stallcup, M. R.: Hormone-independent transcriptional
- [96081] activation and coactivator binding by novel orphan nuclear receptor
- [96082] ERR3. *J. Biol. Chem.* 274: 22618–22626, 1999.
- [96083] 16753.
- [96084] Billin, A. N.; Eilers, A. L.; Queva, C.; Ayer, D. E.: Mlx, a novel
- [96085] Max-like BHLHZip protein that interacts with the Max network of transcription
- [96086] factors. *J. Biol. Chem.* 274: 36344–36350, 1999.
- [96087] 16754.
- [96088] Bjerknes, M.; Cheng, H.: TCFL4: a gene at 17q21.1 encoding a putative
- [96089] basic helix-loop-helix leucine-zipper transcription factor. *Gene* 181:
- [96090] 7–11, 1996.
- [96091] 16755.

- [96092] Cairo, S.; Merla, G.; Urbinati, F.; Ballabio, A.; Reymond, A.:
[96093] WBSCR14, a gene mapping to the Williams–Beuren syn-
drome deleted region,
[96094] is a new member of the Mlx transcription factor network.
Hum. Molec.
[96095] Genet. 10: 617–627, 2001.
[96096] 16756.
[96097] Fukuoka, S.–I.; Freedman, S. D.; Scheele, G. A.: A single
gene
[96098] encodes membrane–bound and free forms of GP–2, the
major glycoprotein
[96099] in pancreatic secretory (zymogen) granule membranes.
Proc. Nat. Acad.
[96100] Sci. 88: 2898–2902, 1991.
[96101] 16757.
[96102] Fukuoka, S.–I.; Freedman, S. D.; Yu, H.; Sukhatme, V. P.;
Scheele,
[96103] G. A.: GP–2/THP gene family encodes self–binding glyco-
sylphosphatidylinositol–anchored
[96104] proteins in apical secretory compartments of pancreas
and kidney. Proc.

- [96105] Nat. Acad. Sci. 89: 1189–1193, 1992.
- [96106] 16758.
- [96107] Fukuoka, S.-I.; Suzuki, M.; Okabayashi, K.; Takahashi, E.:
Assignment
- [96108] of pancreatic zymogen granule membrane protein GP2
(GP2) to human
- [96109] chromosome band 9q21.11 to q21.2 by in situ hybridiza-
tion. Cytogenet.
- [96110] Cell Genet. 79: 231–232, 1997.
- [96111] 16759.
- [96112] Gregory, R. J.; Rich, D. P.; Cheng, S. H.; Souza, D. W.; Paul,
- [96113] S.; Manavalan, P.; Anderson, M. P.; Welsh, M. J.; Smith, A.
E.: Maturation
- [96114] and function of cystic fibrosis transmembrane conduc-
tance regulator
- [96115] variants bearing mutations in putative nucleotide-binding
domains
- [96116] 1 and 2. Molec. Cell. Biol. 11: 3886–3893, 1991.
- [96117] 16760.
- [96118] Greil, I.; Wagner, K.; Rosenkranz, W.: A new missense mu-
tation

- [96119] G1249E in exon 20 of the cystic fibrosis transmembrane conductance
- [96120] regulator (CFTR) gene. Hum. Hered. 44: 238–240, 1994.
- [96121] 16761.
- [96122] Guillermit, H.; Fanen, P.; Ferec, C.: A 3–prime splice site consensus
- [96123] sequence mutation in the cystic fibrosis gene. Hum. Genet. 85: 450–453,
- [96124] 1990.
- [96125] 16762.
- [96126] Guillermit, H.; Jehanne, M.; Quere, I.; Audrezet, M. P.; Mercier,
- [96127] B.; Ferec, C.: A novel mutation in exon 3 of the CFTR gene. Hum.
- [96128] Genet. 91: 233–235, 1993.
- [96129] 16763.
- [96130] Haardt, M.; Benharouga, M.; Lechardeur, D.; Kartner, N.; Lukacs,
- [96131] G. L.: C–terminal truncations destabilize the cystic fibrosis transmembrane
- [96132] conductance regulator without impairing its biogenesis: a

novel class

[96133] of mutation. J. Biol. Chem. 274: 21873–21877, 1999.

[96134] 16764.

[96135] Hamosh, A.; King, T. M.; Rosenstein, B. J.; Corey, M.; Levi-son,

[96136] H.; Durie, P.; Tsui, L.-C.; McIntosh, I.; Keston, M.; Brock, D. J.

[96137] H.; Macek, M., Jr.; Zemkova, D.; Krasnicanova, H.; Vavrova, V.; Macek,

[96138] M., Sr.; Golder, N.; Schwarz, M. J.; Super, M.; Watson, E. K.; Williams,

[96139] C.; Bush, A.; O'Mahoney, S. M.; Humphries, P.; DeArce, M. A.; Reis,

[96140] A.; Burger, J.; Stuhmann, M.; Schmidtke, J.; Wulbrand, U.; Dork,

[96141] T.; Tummeler, B.; Cutting, G. R.: Cystic fibrosis patients bearing

[96142] both the common missense mutation gly-to-aspartate at codon 551 and the

[96143] delta-F508 mutation are clinically indistinguishable from delta-F508

[96144] homozygotes, except for decreased risk of meconium

ileus. Am. J.

[96145] Hum. Genet. 51: 245–250, 1992.

[96146] 16765.

[96147] Hamosh, A.; Trapnell, B. C.; Zeitlin, P. L.; Montrose-Rafizadeh,

[96148] C.; Rosenstein, B. J.; Crystal, R. G.; Cutting, G. R.: Severe deficiency

[96149] of cystic fibrosis transmembrane conductance regulator messenger RNA

[96150] carrying nonsense mutations R553X and W1316X in respiratory epithelial

[96151] cells of patients with cystic fibrosis. J. Clin. Invest. 88: 1880–1885,

[96152] 1991.

[96153] 16766.

[96154] Hefferon, T. W.; Broackes-Carter, F. C.; Harris, A.; Cutting,

[96155] G. R.: Atypical 5-prime splice sites cause CFTR exon 9 to be vulnerable

[96156] to skipping. Am. J. Hum. Genet. 71: 294–303, 2002.

[96157] 16767.

[96158] Highsmith, W. E.; Burch, L. H.; Boat, T. F.; Boucher, R. C.;

Silverman,

- [96159] L. M.; Knowles, M. R.: Identification of a homozygous point mutation
- [96160] in intron 19 in an inbred CF patient with mild disease and normal
- [96161] sweat chloride: creation of an alternative splice site resulting in
- [96162] base-sequence insertion in CFTR coding region between exons 19 and
- [96163] 20. (Abstract) *Pediat. Pulmonol. Suppl.* 6: 22 only, 1991.
- [96164] 16768.
- [96165] Highsmith, W. E., Jr.: Personal Communication. Chapel Hill,
- [96166] N.C. 1991.
- [96167] 16769.
- [96168] Iannuzzi, M. C.; Stern, R. C.; Collins, F. S.; Tom Hon, C.; Hidaka,
- [96169] N.; Strong, T.; Becker, L.; Drumm, M. L.; White, M. B.; Gerard, B.;
- [96170] Dean, M.: Two frameshift mutations in the cystic fibrosis gene. *Am.*
- [96171] *J. Hum. Genet.* 48: 227-231, 1991.

- [96172] 16770.
- [96173] Ivaschenko, T. E.; White, M. B.; Dean, M.; Baranov, V. S.: A
- [96174] deletion of two nucleotides in exon 10 of the CFTR gene
in a Soviet
- [96175] family with cystic fibrosis causing early infant death. Ge-
nomics 10:
- [96176] 298–299, 1991.
- [96177] 16771.
- [96178] Jensen, T. J.; Loo, M. A.; Pind, S.; Williams, D. B.; Goldberg,
- [96179] A. L.; Riordan, J. R.: Multiple proteolytic systems, including
the
- [96180] proteasome, contribute to CFTR processing. Cell 83:
129–135, 1995.
- [96181] 16772.
- [96182] Jiang, Q.; Engelhardt, J. F.: Cellular heterogeneity of CFTR
- [96183] expression and function in the lung: implications for gene
therapy
- [96184] of cystic fibrosis. Europ. J. Hum. Genet. 6: 12–31, 1998.
- [96185] 16773.
- [96186] Jones, C. T.; McIntosh, I.; Keston, M.; Ferguson, A.; Brock,

D.

[96187] J. H.: Three novel mutations in the cystic fibrosis gene detected

[96188] by chemical cleavage: analysis of variant splicing and a nonsense

[96189] mutation. Hum. Molec. Genet. 1: 11–17, 1992.

[96190]

[96191] 100. Kabra, M; Kabra, S. K.; Ghosh, M.; Khanna, A.; Arora, S.; Menon,

[96192] P. S. N.; Verma, I. C.: Is the spectrum of mutations in Indian patients

[96193] with cystic fibrosis different? (Letter) Am. J. Med. Genet. 93:

[96194] 161–163, 2000.

[96195]

[96196] 101. Kalin, N.; Claass, A.; Sommer, M.; Puchelle, E.; Tummeler, B.

[96197] : Delta-F508 CFTR protein expression in tissues from patients with

[96198] cystic fibrosis. J. Clin. Invest. 103: 1379–1389, 1999.

[96199]

[96200] 102. Kelley, K. A.; Stamm, S.; Kozak, C. A.: Expression and chromosome

[96201] localization of the murine cystic fibrosis transmembrane conductance

[96202] regulator. *Genomics* 13: 381–388, 1992.

[96203]

[96204] 103. Kerem, B.; Buchanan, J. A.; Durie, P.; Corey, M. L.; Levison,

[96205] H.; Rommens, J. M.; Buchwald, M.; Tsui, L.–C.: DNA marker haplotype

[96206] association with pancreatic sufficiency in cystic fibrosis. *Am. J.*

[96207] *Hum. Genet.* 44: 827–834, 1989.

[96208]

[96209] 104. Kerem, B.; Rommens, J. M.; Buchanan, J. A.; Markiewicz, D.; Cox,

[96210] T. K.; Chakravarti, A.; Buchwald, M.; Tsui, L.–C.: Identification

[96211] of the cystic fibrosis gene: genetic analysis. *Science* 245: 1073–1080,

[96212] 1989.

[96213]

[96214] 105. Kerem, B.; Zielenski, J.; Markiewicz, D.; Bozon, D.;
Gazit, E.;

[96215] Yahav, J.; Kennedy, D.; Riordan, J. R.; Collins, F. S.; Rom-
mens, J.

[96216] M.; Tsui, L.-C.: Identification of mutations in regions cor-
responding

[96217] to the 2 putative nucleotide (ATP)-binding folds of the
cystic fibrosis

[96218] gene. Proc. Nat. Acad. Sci. 87: 8447-8451, 1990.

[96219]

[96220] 106. Kerem, E.; Corey, M.; Kerem, B.; Rommens, J.;
Markiewicz, D.;

[96221] Levison, H.; Tsui, L.-C.; Durie, P.: The relation between
genotype

[96222] and phenotype in cystic fibrosis--analysis of the most
common mutation

[96223] (delta-F508). New Eng. J. Med. 323: 1517-1522, 1990.

[96224]

[96225] 107. Kerem, E.; Kalman, Y. M.; Yahav, Y.; Shoshani, T.;
Abeliovich,

- [96226] D.; Szeinberg, A.; Rivlin, J.; Blau, H.; Tal, A.; Ben-Tur, L.; Springer,
- [96227] C.; Augarten, A.; Godfrey, S.; Lerer, I.; Branski, D.; Friedman, M.;
- [96228] Kerem, B.: Highly variable incidence of cystic fibrosis and different
- [96229] mutation distribution among different Jewish ethnic groups in Israel. Hum.
- [96230] Genet. 96: 193–197, 1995.
- [96231]
- [96232] 108. Kiesewetter, S.; Macek, M., Jr.; Davis, C.; Curristin, S. M.;
- [96233] Chu, C.-S.; Graham, C.; Shrimpton, A. E.; Cashman, S. M.; Tsui, L.-C.;
- [96234] Mickle, J.; Amos, J.; Highsmith, W. E.; Shuber, A.; Witt, D. R.; Crystal,
- [96235] R. G.; Cutting, G. R.: A mutation in CFTR produces different phenotypes
- [96236] depending on chromosomal background. Nature Genet. 5: 274–278, 1993.
- [96237]
- [96238] 109. Klinger, K.; Horn, G. T.; Stanislovitis, P.; Schwartz, R.

H.;

[96239] Fujiwara, T. M.; Morgan, K.: Cystic fibrosis mutations in the Hutterite

[96240] brethren. Am. J. Hum. Genet. 46: 983–987, 1990.

[96241]

[96242] 110. Kobayashi, K.; Knowles, M. R.; Boucher, R. C.; O'Brien, W. E.;

[96243] Beaudet, A. L.: Benign missense variations in the cystic fibrosis

[96244] gene. Am. J. Hum. Genet. 47: 611–615, 1990.

[96245]

[96246] 111. Kocher, O.; Comella, N.; Tognazzi, K.; Brown, L. F.: Identification

[96247] and partial characterization of PDZK1: a novel protein containing

[96248] PDZ interaction domains. Lab. Invest. 78: 117–125, 1998.

[96249]

[96250] 112. Konstan, M. W.; Byard, P. J.; Hoppel, C. L.; Davis, P. B.: Effect

[96251] of high-dose ibuprofen in patients with cystic fibrosis. New Eng.

[96252] J. Med. 332: 848–854, 1995.

[96253]

[96254] 113. Laroche, D.; Travert, G.: Abnormal frequency of
delta-F(508)

[96255] mutation in neonatal transitory hypertrypsinaemia. (Letter)
Lancet 337:

[96256] 55 only, 1991.

[96257]

[96258] 114. Latham, T.; Grabowski, G. A.; Theophilus, B. D. M.;
Smith, F.

[96259] I.: Complex alleles of the acid beta-glucosidase gene in
Gaucher

[96260] disease. Am. J. Hum. Genet. 47: 79–86, 1990.

[96261]

[96262] 115. Leoni, G.; Pitzalis, S.; Podda, R.; Zanda, M.; Silvetti,
M.;

[96263] Caocci, L.; Cao, A.; Rosatelli, M. C.: A specific cystic fibro-
sis

[96264] mutation (T338I) associated with the phenotype of iso-
lated hypotonic

[96265] dehydration. J. Pediat. 127: 281–283, 1995.

[96266]

[96267] 116. Lerer, I.; Sagi, M.; Cutting, G. R.; Abeliovich, D.: Cystic

[96268] fibrosis mutations delta-F508 and G542X in Jewish patients. J. Med.

[96269] Genet. 29: 131–133, 1992.

[96270]

[96271] 117. Logan, J.; Hiestand, D.; Daram, P.; Huang, Z.; Muccio, D. D.;

[96272] Hartman, J.; Haley, B.; Cook, W. J.; Sorscher, E. J.: Cystic fibrosis

[96273] transmembrane conductance regulator mutations that disrupt nucleotide

[96274] binding. J. Clin. Invest. 94: 228–236, 1994.

[96275]

[96276] 118. Loirat, F.; Hazout, S.; Lucotte, G.: G542X as a probable Phoenician

[96277] cystic fibrosis mutation. Hum. Biol. 69: 419–425, 1997.

[96278]

[96279] 119. Marino, C. R.; Matovcik, L. M.; Gorelick, F. S.; Cohn, J. A.

- [96280] : Localization of the cystic fibrosis transmembrane conductance regulator
- [96281] in pancreas. *J. Clin. Invest.* 88: 712–716, 1991.
- [96282]
- [96283] 120. Marshal, R. D.; Koontz, J.; Sklar, J.: Detection of mutations
- [96284] by cleavage of DNA heteroduplexes with bacteriophage resolvases. *Nature*
- [96285] *Genet.* 9: 177–183, 1995.
- [96286]
- [96287] 121. McCombie, W. R.; Adams, M. D.; Kelley, J. M.; FitzGerald, M.
- [96288] G.; Utterback, T. R.; Khan, M.; Dubnick, M.; Kerlavage, A. R.; Venter,
- [96289] J. C.; Fields, C.: *Caenorhabditis elegans* expressed sequence tags
- [96290] identify gene families and potential disease gene homologues. *Nature*
- [96291] *Genet.* 1: 124–131, 1992.
- [96292]
- [96293] 122. Mickle, J. E.; Macek, M., Jr.; Fulmer-Smentek, S. B.;

Egan, M.

[96294] M.; Schwiebert, E.; Guggino, W.; Moss, R.; Cutting, G. R.: A mutation

[96295] in the cystic fibrosis transmembrane conductance regulator gene associated

[96296] with elevated sweat chloride concentrations in the absence of cystic

[96297] fibrosis. Hum. Molec. Genet. 7: 729–735, 1998.

[96298]

[96299] 123. Mickle, J. E.; Milewski, M. I.; Macek, M., Jr.; Cutting, G. R.

[96300] : Effects of cystic fibrosis and congenital bilateral absence of the

[96301] vas deferens-associated mutations on cystic fibrosis transmembrane

[96302] conductance regulator-mediated regulation of separate channels. Am.

[96303] J. Hum. Genet. 66: 1485–1495, 2000.

[96304]

[96305] 124. Morral, N.; Nunes, V.; Casals, T.; Estivill, X.: CA/GT microsatellite

[96306] alleles within the cystic fibrosis transmembrane conduc-

tance regulator

[96307] (CFTR) gene are not generated by unequal crossing-over.

Genomics 10:

[96308] 692–698, 1991.

[96309]

[96310] 125. Moyer, B. D.; Duhaime, M.; Shaw, C.; Denton, J.;
Reynolds, D.;

[96311] Karlson, K. H.; Pfeiffer, J.; Wang, S.; Mickle, J. E.; Milewski,
M.;

[96312] Cutting, G. R.; Guggino, W. B.; Li, M.; Stanton, B. A.: The
PDZ-interacting

[96313] domain of cystic fibrosis transmembrane conductance
regulator is required

[96314] for functional expression in the apical plasma membrane.
J. Biol.

[96315] Chem. 275: 27069–27074, 2000.

[96316]

[96317] 126. Nelson, P. V.; Carey, W. F.; Morris, C. P.: Identifica-
tion of

[96318] a cystic fibrosis mutation: deletion of isoleucine-506.
Hum. Genet. 86:

[96319] 391–393, 1991.

[96320]

[96321] 127. Nunes, V.; Bonizzato, A.; Gaona, A.; Dognini, M.;
Chillon, M.;

[96322] Casals, T.; Pignatti, P. F.; Novelli, G.; Estivill, X.; Gasparini,
[96323] P.: A frameshift mutation (2869insG) in the second trans-
membrane

[96324] domain of the CFTR gene: identification, regional distribu-
tion, and

[96325] clinical presentation. (Letter) Am. J. Hum. Genet. 50:
1140–1142,

[96326] 1992.

[96327]

[96328] 128. Nunes, V.; Casals, T.; Gaona, A.; Antinolo, G.; Ferrer-
Calvete,

[96329] J.; Perez-Frias, J.; Tardio, E.; Molano, J.; Estivill, X.: Cystic

[96330] fibrosis patients with mutation 1949del84 in exon 13 of
the CFTR gene

[96331] have a similar clinical severity as delF508 homozygotes.
Hum. Mutat. 1:

[96332] 375–379, 1992.

[96333]

- [96334] 129. Nunes, V.; Chillon, M.; Dork, T.; Tummler, B.; Casals, T.; Estivill, X.: A new missense mutation (E92K) in the first trans-membrane domain of the CFTR gene causes a benign cystic fibrosis phenotype. Hum. Molec. Genet. 2: 79–80, 1993.
- [96338]
- [96339] 130. Nunes, V.; Gasparini, P.; Novelli, G.; Gaona, A.; Bonizzato, A.; Sangiuolo, F.; Balassopoulou, A.; Gimenez, F. J.; Dognini, M.; Ravnik-Glavac, M.; Cikuli, M.; Mokini, V.; Komel, R.; Dal-lapiccola, B.; Pignatti, P. F.; Loukopoulos, D.; Casals, T.; Estivill, X.: Analysis of 14 cystic fibrosis mutations in five south European populations. Hum. Genet. 87: 737–738, 1991.
- [96345]
- [96346] 131. Orita, M.; Iwahana, H.; Kanazawa, H.; Hayashi, K.; Sekiya, T.

- [96347] : Detection of polymorphisms of human DNA by gel electrophoresis as
- [96348] single-strand conformation polymorphisms. Proc. Nat. Acad. Sci. 86:
- [96349] 2766–2770, 1989.
- [96350]
- [96351] 132. Orita, M.; Suzuki, Y.; Sekiya, T.; Hayashi, K.: Rapid and sensitive
- [96352] detection of point mutations and DNA polymorphisms using the polymerase
- [96353] chain reaction. Genomics 5: 874–879, 1989.
- [96354]
- [96355] 133. Orozco, L.; Friedman, K.; Chavez, M.; Lezana, J. L.; Villarreal,
- [96356] M. T.; Carnevale, A.: Identification of the I507 deletion by site-directed
- [96357] mutagenesis. Am. J. Med. Genet. 51: 137–139, 1994.
- [96358]
- [96359] 134. Osborne, L.; Knight, R.; Santis, G.; Hodson, M.: A mutation
- [96360] in the second nucleotide binding fold of the cystic fibrosis

gene. Am.

[96361] J. Hum. Genet. 48: 608–612, 1991.

[96362]

[96363] 135. Osborne, L.; Santis, G.; Schwarz, M.; Klinger, K.;
Dork, T.;

[96364] McIntosh, I.; Schwartz, M.; Nunes, V.; Macek, M., Jr.; Reiss,
J.;

[96365] 46 others: Incidence and expression of the N1303K muta-
tion of the

[96366] cystic fibrosis (CFTR) gene. Hum. Genet. 89: 653–658,
1992.

[96367]

[96368] 136. Pier, G. B.; Grout, M.; Zaidi, T.; Meluleni, G.;
Mueschenborn,

[96369] S. S.; Banting, G.; Ratcliff, R.; Evans, M. J.; Colledge, W. H.:

[96370] Salmonella typhi uses CFTR to enter intestinal epithelial
cells. Nature 393:

[96371] 79–82, 1998.

[96372]

[96373] 137. Pier, G. B.; Grout, M.; Zaldi, T. S.; Olsen, J. C.; John-
son,

- [96374] L. G.; Yankaskas, J. R.; Goldberg, J. B.: Role of mutant CFTR in
- [96375] hypersusceptibility of cystic fibrosis patients to lung infections. Science 271:
- [96376] 63–67, 1996.
- [96377]
- [96378] 138. Pignatti, P. F.; Bombieri, C.; Benetazzo, M.; Casartelli, A.;
- [96379] Trabetti, E.; Gile, L. S.; Martinati, L. C.; Boner, A. L.; Luisetti,
- [96380] M.: CFTR gene variant IVS8–5T in disseminated bronchiectasis. (Letter) Am.
- [96381] J. Hum. Genet. 58: 889–892, 1996.
- [96382]
- [96383] 139. Pind, S.; Riordan, J. R.; Williams, D. B.: Participation of
- [96384] the endoplasmic reticulum chaperone calnexin (p88,IP90) in the biogenesis
- [96385] of the cystic fibrosis transmembrane conductance regulator. J. Biol.
- [96386] Chem. 269: 12784–12788, 1994.

[96387]

[96388] 140. Ramjeesingh, M.; Li, C.; Garami, E.; Huan, L.-J.; Galley, K.;

[96389] Wang, Y.; Bear, C. E.: Walker mutations reveal loose relationship

[96390] between catalytic and channel-gating activities of purified CFTR (cystic

[96391] fibrosis transmembrane conductance regulator). *Biochemistry* 38:

[96392] 1463-1468, 1999.

[96393]

[96394] 141. Reddy, M. M.; Light, M. J.; Quinton, P. M.: Activation of the

[96395] epithelial Na(+) channel (ENaC) requires CFTR Cl(-) channel function. *Nature* 402:

[96396] 301-304, 1999.

[96397]

[96398] 142. Rich, D. P.; Anderson, M. P.; Gregory, R. J.; Cheng, S. H.; Paul,

[96399] S.; Jefferson, D. M.; McCann, J. D.; Klinger, K. W.; Smith, A. E.;

- [96400] Welsh, M. J.: Expression of cystic fibrosis transmembrane conductance
- [96401] regulator corrects defective chloride channel regulation in cystic
- [96402] fibrosis airway epithelial cells. Nature 347: 358–363, 1990.
- [96403]
- [96404] 143. Riordan, J. R.; Rommens, J. M.; Kerem, B.; Alon, N.; Rozmahel,
- [96405] R.; Grzelczak, Z.; Zielenski, J.; Lok, S.; Plavsic, N.; Chou, J. L.;
- [96406] Drumm, M. L.; Iannuzzi, M. C.; Collins, F. S.; Tsui, L.-C.: Identification
- [96407] of the cystic fibrosis gene: cloning and characterization of complementary
- [96408] DNA. Science 245: 1066–1073, 1989.
- [96409]
- [96410] 144. Romey, M.-C.; Aguilar-Martinez, P.; Demaille, J.; Claustres,
- [96411] M.: Rapid detection of single nucleotide deletions: application to
- [96412] the beta-6 (-A) mutation of the beta-globin gene and to

cystic fibrosis. Hum.

[96413] Genet. 92: 627–628, 1993.

[96414]

[96415] 145. Romey, M.–C.; Desgeorges, M.; Laussel, M.; Durand, M.–F.; Demaille,

[96416] J.; Claustres, M.: Two novel rare frameshift mutations (2423 del

[96417] G in exon 13 and 1215 del G in exon 7) and one novel rare sequence

[96418] variation (3271 + 18 C or T) identified in a patient with cystic fibrosis. Hum.

[96419] Molec. Genet. 3: 1003–1004, 1994.

[96420]

[96421] 146. Romey, M.–C.; Guittard, C.; Carles, S.; Demaille, J.; Claustres,

[96422] M.; Ramsay, M.: First putative sequence alterations in the minimal

[96423] CFTR promoter region. (Letter) J. Med. Genet. 36: 263–264, 1999.

[96424]

[96425] 147. Romey, M.–C.; Guittard, C.; Chazalotte, J.–P.;

Frossard, P.;

[96426] Dawson, K. P.; Patton, M. A.; Casals, T.; Bazarbachi, T.;
Girodon,

[96427] E.; Rault, G.; Bozon, D.; Seguret, F.; Demaille, J.; Claustres,
M.

[96428] : Complex allele (-102T-to-A+S549R(T-to-G) is associ-
ated with milder

[96429] forms of cystic fibrosis than allele S549R(T-to-G) alone.
Hum. Genet. 105:

[96430] 145-150, 1999.

[96431]

[96432] 148. Romey, M.-C.; Pallares-Ruiz, N.; Mange, A.; Mettling,
C.; Peytavi,

[96433] R.; Demaille, J.; Claustres, M.: A naturally occurring se-
quence variation

[96434] that creates a YY1 element is associated with increased
cystic fibrosis

[96435] transmembrane conductance regulator gene expression. J.
Biol. Chem. 275:

[96436] 3561-3567, 2000.

[96437]

[96438] 149. Rommens, J. M.; Iannuzzi, M. C.; Kerem, B.; Drumm,

M. L.; Melmer,

[96439] G.; Dean, M.; Rozmahel, R.; Cole, J. L.; Kennedy, D.; Hidakaka, N.;

[96440] Zsiga, M.; Buchwald, M.; Riordan, J. R.; Tsui, L.-C.; Collins, F.

[96441] S.: Identification of the cystic fibrosis gene: chromosome walking

[96442] and jumping. Science 245: 1059–1065, 1989.

[96443]

[96444] 150. Ronchetto, P.; Telleria Orriols, J. J.; Fanen, P.; Cremonesi,

[96445] L.; Ferrari, M.; Magnani, C.; Seia, M.; Goossens, M.; Romeo, G.; Devoto,

[96446] M.: A nonsense mutation (R1158X) and a splicing mutation (3849 +

[96447] 4A-to-G) in exon 19 of the cystic fibrosis transmembrane conductance

[96448] regulator gene. Genomics 12: 417–418, 1992.

[96449]

[96450] 151. Rosenfeld, M. A.; Yoshimura, K.; Trapnell, B. C.; Yoneyama, K.;

[96451] Rosenthal, E. R.; Dalemans, W.; Fukayama, M.; Bargon, J.;

Stier, L.

[96452] E.; Stratford-Perricaudet, L.; Perricaudet, M.; Guggino, W. B.; Pavirani,

[96453] A.; Lecocq, J.-P.; Crystal, R. G.: In vivo transfer of the human

[96454] cystic fibrosis transmembrane conductance regulator gene to the airway

[96455] epithelium. Cell 68: 143-155, 1992.

[96456]

[96457] 152. Rowntree, R. K.; Vassaux, G.; McDowell, T. L.; Howe, S.; McGuigan,

[96458] A.; Phylactides, M.; Huxley, C.; Harris, A.: An element in intron

[96459] 1 of the CFTR gene augments intestinal expression in vivo. Hum. Molec.

[96460] Genet. 10: 1455-1464, 2001.

[96461]

[96462] 153. Rozen, R.; De Braekeleer, M.; Daigneault, J.; Ferreira-Rajabi,

[96463] L.; Gerdes, M.; Lamoureux, L.; Aubin, G.; Simard, F.; Fujiwara, T.

[96464] M.; Morgan, K.: Cystic fibrosis mutations in French Cana-

dians: three

[96465] CFTR mutations are relatively frequent in a Quebec population with

[96466] an elevated incidence of cystic fibrosis. Am. J. Med. Genet. 42:

[96467] 360–364, 1992.

[96468]

[96469] 154. Rozen, R.; Ferreira–Rajabi, L.; Robb, L.; Colman, N.: L206W

[96470] mutation of the cystic fibrosis gene, relatively frequent in French

[96471] Canadians, is associated with atypical presentations of cystic fibrosis. Am.

[96472] J. Med. Genet. 57: 437–439, 1995.

[96473]

[96474] 155. Rozen, R.; Schwartz, R. H.; Hilman, B. C.; Stanislovis, P.;

[96475] Horn, G. T.; Klinger, K.; Daigneault, J.; De Braekeleer, M.; Kerem,

[96476] B.; Tsui, L.–C.; Fujiwara, T. M.; Morgan, K.: Cystic fibrosis mutations

[96477] in North American populations of French ancestry: analy–

sis of Quebec

[96478] French-Canadian and Louisiana Acadian families. Am. J. Hum. Genet. 47:

[96479] 606-610, 1990.

[96480]

[96481] 156. Russo, M. P.; Romeo, G.; Devoto, M.; Barbujani, G.; Cabrini,

[96482] G.; Giunta, A.; D'Alcamo, E.; Leoni, G.; Sangiuolo, F.; Magnani, C.;

[96483] Cremonesi, L.; Ferrari, M.: Analysis of linkage disequilibrium between

[96484] different cystic fibrosis mutations and three intragenic microsatellites

[96485] in the Italian population. Hum. Mutat. 5: 23-27, 1995.

[96486]

[96487] 157. Sangiuolo, F.; Cicero, S. L.; Maceratesi, P.; Quattrucci, S.;

[96488] Novelli, G.; Dallapiccola, B.: Molecular characterization of a frameshift

[96489] mutation in exon 19 of the CFTR gene. Hum. Mutat. 2: 422-424, 1993.

[96490]

[96491] 158. Sangiuolo, F.; Novelli, G.; Murru, S.; Dallapiccola, B.:
A serine-to-arginine

[96492] (AGT-to-CGT) mutation in codon 549 of the CFTR gene in
an Italian

[96493] patient with severe cystic fibrosis. Genomics 9: 788–789,
1991.

[96494]

[96495] 159. Sato, S.; Ward, C. L.; Kopito, R. R.: Cotranslational
ubiquitination

[96496] of cystic fibrosis transmembrane conductance regulator in
vitro. J.

[96497] Biol. Chem. 273: 7189–7192, 1998.

[96498]

[96499] 160. Sato, S.; Ward, C. L.; Krouse, M. E.; Wine, J. J.; Kopito,
R.

[96500] R.: Glycerol reverses the misfolding phenotype of the
most common

[96501] cystic fibrosis mutation. J. Biol. Chem. 271: 635–638,
1996.

[96502]

[96503] 161. Schaedel, C.; Kristoffersson, A.-C.; Kornfalt, R.;
Holmberg,

- [96504] L.: A novel cystic fibrosis mutation, Y109C, in the first transmembrane
- [96505] domain of CFTR. Hum. Molec. Genet. 3: 1001–1002, 1994.
- [96506]
- [96507] 162. Schwartz, M.; Anvret, M.; Claustres, M.; Geir Eiken, H.; Eiklid,
- [96508] K.; Schaedel, C.; Stolpe, L.; Tranebjaerg, L.: 394delTT: a Nordic
- [96509] cystic fibrosis mutation. Hum. Genet. 93: 157–161, 1994.
- [96510]
- [96511] 163. Schwiebert, E. M.; Egan, M. E.; Hwang, T.–H.; Fulmer, S. B.;
- [96512] Allen, S. S.; Cutting, G. R.; Guggino, W. B.: CFTR regulates outwardly
- [96513] rectifying chloride channels through an autocrine mechanism involving
- [96514] ATP. Cell 81: 1063–1073, 1995.
- [96515]
- [96516] 164. Scotet, V.; De Braekeleer, M.; Audrezet, M.–P.; Lode, L.; Verlingue,

- [96517] C.; Quere, I.; Mercier, B.; Dugueperoux, I.; Codet, J.-P.;
Moineau,
[96518] M.-P.; Parent, P.; Ferec, C.: Prevalence of CFTR mutations
in hypertrypsinaemia
[96519] detected through neonatal screening for cystic fibrosis.
Clin. Genet. 59:
[96520] 42–47, 2001.
[96521]
[96522] 165. Sharer, N.; Schwarz, M.; Malone, G.; Howarth, A.;
Painter, J.;
[96523] Super, M.; Braganza, J.: Mutations of the cystic fibrosis
gene in
[96524] patients with chronic pancreatitis. New Eng. J. Med. 339:
645–652,
[96525] 1998.
[96526]
[96527] 166. Sheppard, D. N.; Rich, D. P.; Ostedgaard, L. S.; Gre-
gory, R.
[96528] J.; Smith, A. E.; Welsh, M. J.: Mutations in CFTR associated
with
[96529] mild–disease form CI– channels with altered pore proper-
ties. Nature 362:

[96530] 160–164, 1993.

[96531]

[96532] 167. Shoshani, T.; Augarten, A.; Gazit, E.; Bashan, N.; Yahav, Y.;

[96533] Rivlin, Y.; Tal, A.; Seret, H.; Yaar, L.; Kerem, E.; Kerem, B.:
Association

[96534] of a nonsense mutation (W1282X), the most common mutation in the Ashkenazi

[96535] Jewish cystic fibrosis patients in Israel, with presentation of severe

[96536] disease. Am. J. Hum. Genet. 50: 222–228, 1992.

[96537]

[96538] 168. Shoshani, T.; Augarten, A.; Yahav, J.; Gazit, E.; Kerem, B.:

[96539] Two novel mutations in the CFTR gene: W1089X in exon 17B and 4010delTATT

[96540] in exon 21. Hum. Molec. Genet. 3: 657–658, 1994.

[96541]

[96542] 169. Shoshani, T.; Berkun, Y.; Yahav, Y.; Augarten, A.; Bashan, N.;

[96543] Rivlin, Y.; Gazit, E.; Sereth, H.; Kerem, E.; Kerem, B.: A new

mutation

[96544] in the CFTR gene, composed of two adjacent DNA alterations, is a common

[96545] cause of cystic fibrosis among Georgian Jews. Genomics 15: 236–237,

[96546] 1993.

[96547]

[96548] 170. Shoshani, T.; Kerem, E.; Szeinberg, A.; Augarten, A.; Yahav,

[96549] Y.; Cohen, D.; Rivlin, J.; Tal, A.; Kerem, B. S.: Similar levels

[96550] of mRNA from the W1282X and the delta-F508 cystic fibrosis alleles,

[96551] in nasal epithelial cells. J. Clin. Invest. 93: 1502–1507, 1994.

[96552]

[96553] 171. Siegel, D.; Irving, N. G.; Friedman, J. M.; Wainwright, B. J.

[96554] : Localization of the cystic fibrosis transmembrane conductance regulator

[96555] (Cftr) to mouse chromosome 6. Cytogenet. Cell Genet. 61: 184–185,

[96556] 1992.

[96557]

[96558] 172. Smit, L. S.; Nasr, S. Z.; Iannuzzi, M. C.; Collins, F. S.:
An

[96559] African-American cystic fibrosis patient homozygous for a
novel frameshift

[96560] mutation associated with reduced CFTR mRNA levels.
Hum. Mutat. 2:

[96561] 148–151, 1993.

[96562]

[96563] 173. Smit, L. S.; Strong, T. V.; Wilkinson, D. J.; Macek, M.,
Jr.;

[96564] Mansoura, M. K.; Wood, D. L.; Cole, J. L.; Cutting, G. R.;
Cohn, J.

[96565] A.; Dawson, D. C.; Collins, F. S.: Missense mutation
(G480C) in the

[96566] CFTR gene associated with protein mislocalization but
normal chloride

[96567] channel activity. Hum. Molec. Genet. 4: 269–273, 1995.

[96568]

[96569] 174. Snouwaert, J. N.; Brigman, K. K.; Latour, A. M.; Mal-
ouf, N. N.;

[96570] Boucher, R. C.; Smithies, O.; Koller, B. H.: An animal model
for

[96571] cystic fibrosis made by gene targeting. Science 257:
1083–1088,

[96572] 1992.

[96573]

[96574] 175. Stern, R. C.: The diagnosis of cystic fibrosis. New
Eng. J.

[96575] Med. 336: 487–491, 1997.

[96576]

[96577] 176. Strong, T. V.; Smit, L. S.; Nasr, S.; Wood, D. L.; Cole,
J. L.;

[96578] Iannuzzi, M. C.; Stern, R. C.; Collins, F. S.: Characterization
of

[96579] an intron 12 splice donor mutation in the cystic fibrosis
transmembrane

[96580] conductance regulator (CFTR) gene. Hum. Mutat. 1:
380–387, 1992.

[96581]

[96582] 177. Strong, T. V.; Smit, L. S.; Turpin, S. V.; Cole, J. L.;
Tom Hon,

[96583] C.; Markiewicz, D.; Petty, T. L.; Craig, M. W.; Rosenow, E. C., III;

[96584] Tsui, L.-C.; Iannuzzi, M. C.; Knowles, M. R.; Collins, F. S.:
Cystic

[96585] fibrosis gene mutation in two sisters with mild disease
and normal

[96586] sweat electrolyte levels. New Eng. J. Med. 325:
1630-1634, 1991.

[96587]

[96588] 178. Tata, F.; Stanier, P.; Wicking, C.; Halford, S.; Kruyer, H.;

[96589] Lench, N. J.; Scambler, P. J.; Hansen, C.; Braman, J. C.;
Williamson,

[96590] R.; Wainwright, B. J.: Cloning the mouse homolog of the
human cystic

[96591] fibrosis transmembrane conductance regulator gene. Ge-
nomics 10:

[96592] 301-307, 1991.

[96593]

[96594] 179. Teng, H.; Jorissen, M.; Van Poppel, H.; Legius, E.;
Cassiman,

[96595] J.-J.; Cuppens, H.: Increased proportion of exon 9 alterna-

tively

[96596] spliced CFTR transcripts in vas deferens compared with nasal epithelial

[96597] cells. Hum. Molec. Genet. 6: 85–90, 1997.

[96598]

[96599] 180. The Cystic Fibrosis Genotype–Phenotype Consortium: Correlation

[96600] between genotype and phenotype in patients with cystic fibrosis. New

[96601] Eng. J. Med. 329: 1308–1313, 1993.

[96602]

[96603] 181. Trapnell, B. C.; Chu, C.-S.; Paakko, P. K.; Banks, T. C.; Yoshimura,

[96604] K.; Ferrans, V. J.; Chernick, M. S.; Crystal, R. G.: Expression of

[96605] the cystic fibrosis transmembrane conductance regulator gene in the

[96606] respiratory tract of normal individuals and individuals with cystic

[96607] fibrosis. Proc. Nat. Acad. Sci. 88: 6565–6569, 1991.

[96608]

- [96609] 182. Trezise, A. E. O.; Szpirer, C.; Buchwald, M.: Localization of
- [96610] the gene encoding the cystic fibrosis transmembrane conductance regulator
- [96611] (CFTR) in the rat to chromosome 4 and implications for the evolution
- [96612] of mammalian chromosomes. *Genomics* 14: 869–874, 1992.
- [96613]
- [96614] 183. Van Doorninck, J. H.; French, P. J.; Verbeek, E.; Peters, R.
- [96615] H. P. C.; Morreau, H.; Bijman, J.; Scholte, B. J.: A mouse model
- [96616] for the cystic fibrosis delta-F508 mutation. *EMBO J.* 14: 4403–4411,
- [96617] 1995.
- [96618]
- [96619] 184. Varon, R.; Magdorf, K.; Staab, D.; Wahn, H.–U.; Krawczak, M.;
- [96620] Sperling, K.; Reis, A.: Recurrent nasal polyps as a monosymptomatic
- [96621] form of cystic fibrosis associated with a novel in-frame

deletion

[96622] (591del18) in the CFTR gene. Hum. Molec. Genet. 4: 1463–1464, 1995.

[96623]

[96624] 185. Varon, R.; Stuhmann, M.; Macek, M., Jr.; Kufardjieva, A.; Angelicheva,

[96625] D.; Magdorf, K.; Jordanova, A.; Savov, A.; Wahn, U.; Macek, M.; Lalov,

[96626] V.; Ivanova, T.; Ellemunter, H.; Vavrova, V.; Ferak, V.; Kayserova,

[96627] H.; Reis, A.; Kalaydjieva, L.: Pancreatic insufficiency and pulmonary

[96628] disease in German and Slavic cystic fibrosis patients with the R347P

[96629] mutation. Hum. Mutat. 6: 219–225, 1995.

[96630]

[96631] 186. Verlingue, C.; Mercier, B.; Lecoq, I.; Audrezet, M. P.; Laroche,

[96632] D.; Travert, G.; Ferec, C.: Retrospective study of the cystic fibrosis

[96633] transmembrane conductance regulator (CFTR) gene mutations in Guthrie

- [96634] cards from a large cohort of neonatal screening for cystic fibrosis. Hum.
- [96635] Genet. 93: 429–434, 1994.
- [96636]
- [96637] 187. Vidaud, M.; Fanen, P.; Martin, J.; Ghanem, N.; Nicolas, S.; Goossens, M.: Three point mutations in the CFTR gene in French cystic fibrosis patients: identification by denaturing gradient gel electrophoresis. Hum.
- [96639] Genet. 85: 446–449, 1990.
- [96640]
- [96641]
- [96642] 188. Vidaud, M.; Gattoni, R.; Stevenin, J.; Vidaud, D.; Amselem, S.; Chibani, J.; Rosa, J.; Goossens, M.: A 5-prime splice-region G-to-C mutation in exon 1 of the human beta-globin gene inhibits pre-mRNA splicing: a mechanism for beta(+)-thalassemia. Proc. Nat. Acad. Sci. 86: 1041–1045, 1989.
- [96646]

[96647]

[96648] 189. Wang, J.; Bowman, M. C.; Hsu, E.; Wertz, K.; Wong, L.-J. C.:

[96649] A novel mutation in the CFTR gene correlates with severe clinical

[96650] phenotype in seven Hispanic patients. J. Med. Genet. 37: 215-218,

[96651] 2000.

[96652]

[96653] 190. Wang, S.; Yue, H.; Derin, R. B.; Guggino, W. B.; Li, M.: Accessory

[96654] protein facilitated CFTR-CFTR interaction, a molecular mechanism to

[96655] potentiate the chloride channel activity. Cell 103: 169-179, 2000.

[96656]

[96657] 191. Waterston, R.; Martin, C.; Craxton, M.; Huynh, C.; Coulson, A.;

[96658] Hillier, L.; Durbin, R.; Green, P.; Shownkeen, R.; Halloran, N.; Metzstein,

[96659] M.; Hawkins, T.; Wilson, R.; Berks, M.; Du, Z.; Thomas, K.; Thierry-Mieg,

[96660] J.; Sulston, J.: A survey of expressed genes in *Caenorhabditis elegans*. *Nature*

[96661] *Genet.* 1: 114–123, 1992.

[96662]

[96663] 192. Wauters, J. G.; Hendrickx, J.; Coucke, P.; Vits, L.; Stuer, K.;

[96664] van Schil, L.; van der Auwera, B. J.; Van Elsen, A.; Dumon, J.; Willems,

[96665] P. J.: Frequency of the phenylalanine deletion (delta-F508) in the

[96666] CF gene of Belgian cystic fibrosis patients. *Clin. Genet.* 39: 89–92,

[96667] 1991.

[96668]

[96669] 193. Wei, L.; Vankeerberghen, A.; Cuppens, H.; Droogmans, G.; Cassiman,

[96670] J.-J.; Nilius, B.: Phosphorylation site independent single R-domain

[96671] mutations affect CFTR channel activity. *FEBS Lett.* 439: 121–126,

[96672] 1998.

[96673]

[96674] 194. Weixel, K. M.; Bradbury, N. A.: The carboxyl terminus of the

[96675] cystic fibrosis transmembrane conductance regulator binds to AP-2

[96676] clathrin adaptors. J. Biol. Chem. 275: 3655-3660, 2000.

[96677]

[96678] 195. White, M. B.; Amos, J.; Hsu, J. M. C.; Gerrard, B.; Finn, P.;

[96679] Dean, M.: A frame shift mutation in the cystic fibrosis gene. Nature 344:

[96680] 655-667, 1990.

[96681]

[96682] 196. White, M. B.; Leppert, M.; Nielsen, D.; Zielenski, J.; Gerrard,

[96683] B.; Stewart, C.; Dean, M.: A de novo cystic fibrosis mutation: CGA

[96684] (arg) to TGA (stop) at codon 851 of the CFTR gene. Genomics 11:

[96685] 778-779, 1991.

[96686]

- [96687] 197. White, S. M.; Lucassen, A.; Norbury, G.: Cystic fibrosis: a
[96688] further case of an asymptomatic compound heterozygote.
(Letter) Am.
[96689] J. Med. Genet. 103: 342–343, 2001.
[96690]
[96691] 198. Will, K.; Dork, T.; Stuhrmann, M.; Meitinger, T.;
Bertele-Harms,
[96692] R.; Tummler, B.; Schmidtke, J.: A novel exon in the cystic
fibrosis
[96693] transmembrane conductance regulator gene activated by
the nonsense
[96694] mutation E92X in airway epithelial cells of patients with
cystic fibrosis. J.
[96695] Clin. Invest. 93: 1852–1859, 1994.
[96696]
[96697] 199. Yang, Y.; Devor, D. C.; Engelhardt, J. F.; Ernst, S. A.;
Strong,
[96698] T. V.; Collins, F. S.; Cohn, J. A.; Frizzell, R. A.; Wilson, J. M.
[96699] : Molecular basis of defective anion transport in L cells
expressing

- [96700] recombinant forms of CFTR. Hum. Molec. Genet. 2: 1253–1261, 1993.
- [96701]
- [96702] 200. Yang, Y.; Janich, S.; Cohn, J. A.; Wilson, J. M.: The common
- [96703] variant of cystic fibrosis transmembrane conductance regulator is
- [96704] recognized by hsp70 and degraded in a pre-Golgi nonlysosomal compartment. Proc.
- [96705] Nat. Acad. Sci. 90: 9480–9484, 1993.
- [96706]
- [96707] 201. Yang, Y.; Raper, S. E.; Cohn, J. A.; Engelhardt, J. F.; Wilson,
- [96708] J. M.: An approach for treating the hepatobiliary disease of cystic
- [96709] fibrosis by somatic gene transfer. Proc. Nat. Acad. Sci. 90: 4601–4605,
- [96710] 1993.
- [96711]
- [96712] 202. Youil, R.; Kemper, B. W.; Cotton, R. G. H.: Screening for mutations

- [96713] by enzyme mismatch cleavage with T4 endonuclease VII.
Proc. Nat.
- [96714] Acad. Sci. 92: 87–91, 1995.
- [96715]
- [96716] 203. Zeiher, B. G.; Eichwald, E.; Zabner, J.; Smith, J. J.;
Puga,
- [96717] A. P.; McCray, P. B., Jr.; Capecchi, M. R.; Welsh, M. J.;
Thomas,
- [96718] K. R.: A mouse model for the delta-F508 allele of cystic fi-
brosis. J.
- [96719] Clin. Invest. 96: 2051–2064, 1995.
- [96720]
- [96721] 204. Zeitlin, P. L.; Crawford, I.; Lu, L.; Woel, S.; Cohen, M.
E.;
- [96722] Donowitz, M.; Montrose, M. H.; Hamosh, A.; Cutting, G. R.;
Gruenert,
- [96723] D.; Haganir, R.; Maloney, P.; Guggino, W. B.: CFTR protein
expression
- [96724] in primary and cultured epithelia. Proc. Nat. Acad. Sci. 89:
344–347,
- [96725] 1992.

[96726]

[96727] 205. Zhang, F.; Kartner, N.; Lukacs, G. L.: Limited proteolysis as

[96728] a probe for arrested conformational maturation of deltaF508 CFTR. Nature

[96729] Struct. Biol. 5: 180–183, 1998.

[96730]

[96731] 206. Zielenski, J.; Bozon, D.; Kerem, B.; Markiewicz, D.; Durie, P.;

[96732] Rommens, J. M.; Tsui, L.-C.: Identification of mutations in exons

[96733] 1 through 8 of the cystic fibrosis transmembrane conductance regulator

[96734] (CFTR) gene. Genomics 10: 229–235, 1991.

[96735]

[96736] 207. Zielenski, J.; Markiewicz, D.; Rininsland, F.; Rommens, J.; Tsui,

[96737] L.-C.: A cluster of highly polymorphic dinucleotide repeats in intron

[96738] 17b of the cystic fibrosis transmembrane conductance regulator (CFTR)

[96739] gene. Am. J. Hum. Genet. 49: 1256–1262, 1991.

[96740]

[96741] 208. Zielenski, J.; Patrizio, P.; Corey, M.; Handelin, B.;
Markiewicz,

[96742] D.; Asch, R.; Tsui, L.-C.: CFTR gene variant for patients
with congenital

[96743] absence of vas deferens. (Letter) Am. J. Hum. Genet. 57:
958–960,

[96744] 1995.

[96745] 16774.

[96746] Martin, F.; Schaller, A.; Eglite, S.; Schumperli, D.; Muller, B.

[96747] : The gene for histone RNA hairpin binding protein is lo-
cated on human

[96748] chromosome 4 and encodes a novel type of RNA binding
protein. EMBO

[96749] J. 16: 769–778, 1997.

[96750] 16775.

[96751] Wang, Z.-F.; Whitfield, M. L.; Ingledue, T. C., III; Dominski,

[96752] Z.; Marzluff, W. F.: The protein that binds the 3–prime end
of histone

- [96753] mRNA: a novel RNA-binding protein required for histone pre-mRNA processing. *Genes Dev.* 10: 3028-3040, 1996.
- [96754] 16776.
- [96755] Kooy, J.; Toh, B.-H.; Pettitt, J. M.; Erlich, R.; Gleeson, P. A.
- [96757] : Human autoantibodies as reagents to conserved Golgi components:
- [96758] characterization of a peripheral, 230-kDa compartment-specific Golgi
- [96759] protein. *J. Biol. Chem.* 267: 20255-20263, 1992.
- [96760] 16777.
- [96761] Marneros, A. G.; Mehenni, H.; Reichenberger, E.; Antonarakis, S.
- [96762] E.; Krieg, T.; Olsen, B. R.: Gene for the human transmembrane-type
- [96763] protein tyrosine phosphatase H (PTPRH): genomic structure, fine-mapping
- [96764] and its exclusion as a candidate for Peutz-Jeghers syndrome. *Cytogenet.*
- [96765] *Cell Genet.* 92: 213-216, 2001.
- [96766] 16778.

- [96767] Matozaki, T.; Suzuki, T.; Uchida, T.; Inazawa, J.; Ariyama, T.;
- [96768] Matsuda, K.; Horita, K.; Noguchi, H.; Mizuno, H.; Sakamoto, C.; Kasuga,
- [96769] M.: Molecular cloning of a human transmembrane-type protein tyrosine
- [96770] phosphatase and its expression in gastrointestinal cancers. J. Biol.
- [96771] Chem. 269: 2075–2081, 1994.
- [96772] 16779.
- [96773] Seo, Y.; Matozaki, T.; Tsuda, M.; Hayashi, Y.; Itoh, H.; Kasuga,
- [96774] M.: Overexpression of SAP-1, a transmembrane-type protein tyrosine
- [96775] phosphatase, in human colorectal cancers. Biochem. Biophys. Res.
- [96776] Commun. 231: 705–711, 1997.
- [96777] 16780.
- [96778] Snow, B. E.; Antonio, L.; Suggs, S.; Gutstein, H. B.; Siderovski,
- [96779] D. P.: Molecular cloning and expression analysis of rat Rgs12 and

- [96780] Rgs14. Biochem. Biophys. Res. Commun. 233: 770–777, 1997.
- [96781] 16781.
- [96782] Kimple, R. J.; Kimple, M. E.; Betts, L.; Sondek, J.; Siderovski,
- [96783] D. P.: Structural determinants for GoLoco-induced inhibition of nucleotide
- [96784] release by G- α subunits. Nature 416: 878–881, 2002.
- [96785] 16782.
- [96786] Takesono, A.; et al; et al: Receptor-independent activators of
- [96787] heterotrimeric G-protein signalling pathways. J. Biol. Chem. 274:
- [96788] 33202–33205, 1999.
- [96789] 16783.
- [96790] Traver, S.; Bidot, C.; Spassky, N.; Baltauss, T.; de Tand, M.-F.;
- [96791] Thomas, J.-L.; Zalc, B.; Janoueix-Lerosey, I.; de Gunzburg, J.: RGS14
- [96792] is a novel Rap effector that preferentially regulates the GTPase activity

[96793] of G- α -0. Biochem. J. 350: 19-29, 2000.

[96794] 16784.

[96795] Zheng, B.; Chen, D.; Farquhar, M. G.: MIR16, a putative membrane

[96796] glycerophosphodiester phosphodiesterase, interacts with RGS16. Proc.

[96797] Nat. Acad. Sci. 97: 3999-4004, 2000.

[96798] 16785.

[96799] Schmitz, F.; Konigstorfer, A.; Sudhof, T. C.: RIBEYE, a component

[96800] of synaptic ribbons: a protein's journey through evolution provides

[96801] insight into synaptic ribbon function. Neuron 28: 857-872, 2000.

[96802] 16786.

[96803] Dale, M.; Hammond, D. W.; Cox, A.; Nicklin, M. J. H.: The human

[96804] gene encoding the interleukin-1 receptor accessory protein (IL1RAP)

[96805] maps to chromosome 3q28 by fluorescence in situ hybridization and

- [96806] radiation hybrid mapping. *Genomics* 47: 325–326, 1998.
- [96807] 16787.
- [96808] Huang, J.; Gao, X.; Li, S.; Cao, Z.: Recruitment of IRAK to the
- [96809] interleukin 1 receptor complex requires interleukin 1 receptor accessory
- [96810] protein. *Proc. Nat. Acad. Sci.* 94: 12829–12832, 1997.
- [96811] 16788.
- [96812] Wesche, H.; Korherr, C.; Kracht, M.; Falk, W.; Resch, K.; Martin,
- [96813] M. U.: The interleukin–1 receptor accessory protein (IL–1RAcP) is
- [96814] essential for IL–1–induced activation of interleukin–1 receptor–associated
- [96815] kinase (IRAK) and stress–activated protein kinases (SAP kinases). *J.*
- [96816] *Biol. Chem.* 272: 7727–7731, 1997.
- [96817] 16789.
- [96818] Saha, P.; Chen, J.; Thome, K. C.; Lawlis, S. J.; Hou, Z.–H.; Hendricks,
- [96819] M.; Parvin, J. D.; Dutta, A.: Human CDC6/Cdc18 associates

with Orc1

[96820] and cyclin-cdk and is selectively eliminated from the nucleus at the

- [96821] onset of S phase. *Molec. Cell. Biol.* 18: 2758–2767, 1998.
- [96822] 16790.
- [96823] Williams, R. S.; Shohet, R. V.; Stillman, B.: A human protein
- [96824] related to yeast Cdc6p. *Proc. Nat. Acad. Sci.* 94: 142–147, 1997.
- [96825] 16791.
- [96826] Yan, Z.; DeGregori, J.; Shohet, R.; Leone, G.; Stillman, B.; Nevins,
- [96827] J. R.; Williams, R. S.: Cdc6 is regulated by E2F and is essential
- [96828] for DNA replication in mammalian cells. *Proc. Nat. Acad. Sci.* 95:
- [96829] 3603–3608, 1998.
- [96830] 16792.
- [96831] Brais, B.; Bouchard, J.-P.; Xie, Y.-G.; Rochefort, D. L.; Chretien,
- [96832] N.; Tome, F. M. S.; Lafreniere, R. G.; Rommens, J. M.; Uyama, E.;
- [96833] Nohira, O.; Blumen, S.; Korczyn, A. D.; Heutink, P.; Mathieu, J.;
- [96834] Duranceau, A.; Codere, F.; Fardeau, M.; Rouleau, G. A.:

Short GCG

- [96835] expansions in the PABP2 gene cause oculopharyngeal muscular dystrophy. Nature
- [96836] Genet. 18: 164–167, 1998. Note: Erratum: Nature Genet. 19: 404 only,
- [96837] 1998.
- [96838] 16793.
- [96839] Calado, A.; Tome, F. M. S.; Brais, B.; Rouleau, G. A.; Kuhn, U.;
- [96840] Wahle, E.; Carmo-Fonseca, M.: Nuclear inclusions in oculopharyngeal
- [96841] muscular dystrophy consist of poly(A) binding protein 2 aggregates
- [96842] which sequester poly(A) RNA. Hum. Molec. Genet. 9: 2321–2328, 2000.
- [96843] 16794.
- [96844] Fan, X.; Dion, P.; Laganier, J.; Brais, B.; Rouleau, G. A.: Oligomerization
- [96845] of polyalanine expanded PABPN1 facilitates nuclear protein aggregation
- [96846] that is associated with cell death. Hum. Molec. Genet. 10: 2341–2351,

- [96847] 2001.
- [96848] 16795.
- [96849] Ikeda, S.; Shiva, N.; Ikeda, A.; Smith, R. S.; Nusinowitz, S.;
- [96850] Yan, G.; Lin, T. R.; Chu, S.; Heckenlively, J. R.; North, M. A.;
Naggert,
- [96851] J. K.; Nishina, P. M.; Duyao, M. P.: Retinal degeneration but
not
- [96852] obesity is observed in null mutants of the tubby-like pro-
tein 1 gene. Hum.
- [96853] Molec. Genet. 9: 155–163, 2000.
- [96854] 16796.
- [96855] North, M. A.; Naggert, J. K.; Yan, Y.; Noben-Trauth, K.;
- Nishina,
- [96856] P. M.: Molecular characterization of TUB, TULP1, and
TULP2, members
- [96857] of the novel tubby gene family and their possible relation
to ocular
- [96858] diseases. Proc. Nat. Acad. Sci. 94: 3128–3133, 1997.
- [96859] 16797.
- [96860] Paloma, E.; Hjelmqvist, L.; Bayes, M.; Garcia-Sandoval, B.;
- Ayuso,

- [96861] C.; Balcells, S.; Gonzalez-Duarte, R.: Novel mutations in the TULP1
- [96862] gene causing autosomal recessive retinitis pigmentosa. Invest. Ophthalm.
- [96863] Vis. Sci. 41: 656-659, 2000.
- [96864] 16798.
- [96865] Aoki, N.; Ishii, T.; Ohira, S.; Yamaguchi, Y.; Negi, M.; Adachi,
- [96866] T.; Nakamura, R.; Matsuda, T.: Stage specific expression of milk
- [96867] fat globule membrane glycoproteins in mouse mammary gland: comparison
- [96868] of MFG-E8, butyrophilin, and CD36 with a major milk protein, beta-casein. Biochim.
- [96869] Biophys. Acta 1334: 182-190, 1997.
- [96870] 16799.
- [96871] Collins, C.; Nehlin, J. O.; Stubbs, J. D.; Kowbel, D.; Kuo, W.-L.;
- [96872] Parry, G.: Mapping of a newly discovered human gene homologous to
- [96873] the apoptosis associated-murine mammary protein, MFG-E8, to chromosome

- [96874] 15q25. Genomics 39: 117–118, 1997.
- [96875] 16800.
- [96876] Haggqvist, B.; Naslund, J.; Sletten, K.; Westermark, G. T.;
Mucchiano,
- [96877] G.; Tjernberg, L. O.; Nordstedt, C.; Engstrom, U.; Wester-
mark, P.
- [96878] : Medin: an integral fragment of aortic smooth muscle
cell-produced
- [96879] lactadherin forms the most common human amyloid.
Proc. Nat. Acad.
- [96880] Sci. 96: 8669–8674, 1999.
- [96881] 16801.
- [96882] Hanayama, R.; Tanaka, M.; Miwa, K.; Shinohara, A.; Iwa-
matsu, A.;
- [96883] Nagata, S.: Identification of a factor that links apoptotic
cells
- [96884] to phagocytes. Nature 417: 182–187, 2002.
- [96885] 16802.
- [96886] Larocca, D.; Peterson, J. A.; Urrea, R.; Kuniyoshi, J.; Bis-
train,
- [96887] A. M.; Ceriani, R. L.: A M(r) 46,000 human milk fat globule

protein

[96888] that is highly expressed in human breast tumors contains factor VIII-like

[96889] domains. *Cancer Res.* 51: 4994–4998, 1991.

[96890] 16803.

[96891] Stubbs, J. D.; Lekutis, C.; Singer, K. L.; Bui, A.; Yuzuki, D.;

[96892] Srinivasan, U.; Parry, G.: cDNA cloning of a mouse mammary epithelial

[96893] cell surface protein reveals the existence of epidermal growth factor-like

[96894] domains linked to factor VIII-like sequences. *Proc. Nat. Acad. Sci.* 87:

[96895] 8417–8421, 1990.

[96896] 16804.

[96897] Zhang, J.; Kuehl, P.; Green, E. D.; Touchman, J. W.; Watkins, P.

[96898] B.; Daly, A.; Hall, S. D.; Maurel, P.; Relling, M.; Brimer, C.; Yasuda,

[96899] K.; Wrighton, S. A.; Hancock, M.; Kim, R. B.; Strom, S.; Thummel,

[96900] K.; Russell, C. G.; Hudson, J. R., Jr.; Schuetz, E. G.; Bo-

guski, M.

- [96901] S.: The human pregnane X receptor: genomic structure and identification
- [96902] and functional characterization of natural allelic variants. Pharmacogenetics 11:
- [96903] 555–572, 2001.
- [96904] 16805.
- [96905] Berrueta, L.; Kraeft, S.-K.; Tirnauer, J. S.; Schuyler, S. C.;
- [96906] Chen, L. B.; Hill, D. E.; Pellman, D.; Bierer, B. E.: The adenomatous
- [96907] polyposis coli-binding protein EB1 is associated with cytoplasmic
- [96908] and spindle microtubules. Proc. Nat. Acad. Sci. 95: 10596–10601,
- [96909] 1998.
- [96910] 16806.
- [96911] Berrueta, L.; Tirnauer, J. S.; Schuyler, S. C.; Pellman, D.; Bierer,
- [96912] B. E.: The APC-associated protein EB1 associates with components
- [96913] of the dynactin complex and cytoplasmic dynein interme-

diate chain. Curr.

[96914] Biol. 9: 425–428, 1999.

[96915] 16807.

[96916] Su, L.-K.; Burrell, M.; Hill, D. E.; Gyuris, J.; Brent, R.; Wiltshire,

[96917] R.; Trent, J.; Vogelstein, B.; Kinzler, K. W.: APC binds to the novel

[96918] protein EB1. Cancer Res. 55: 2972–2977, 1995.

[96919] 16808.

[96920] Su, L.-K.; Qi, Y.: Characterization of human MAPRE genes and their

[96921] proteins. Genomics 71: 143–149, 2001.

[96922] 16809.

[96923] Tirnauer, J. S.; Bierer, B. E.: EB1 proteins regulate microtubule

[96924] dynamics, cell polarity, and chromosome stability. J. Cell Biol. 149:

[96925] 761–766, 2000.

[96926] 16810.

[96927] Arai, T.; Akiyama, Y.; Okabe, S.; Ando, M.; Endo, M.;

Yuasa, Y.

[96928] : Genomic structure of the human Smad3 gene and its infrequent alterations

[96929] in colorectal cancers. Cancer Lett. 122: 157–163, 1998.

[96930] 16811.

[96931] Zhang, Y.; Feng, X.-H.; Wu, R.-Y.; Derynck, R.: Receptor-associated

[96932] Mad homologues synergize as effectors of the TGF- β response. Nature 383:

[96933] 168–172, 1996.

[96934] 16812.

[96935] Zhu, Y.; Richardson, J. A.; Parada, L. F.; Graff, J. M.: Smad3

[96936] mutant mice develop metastatic colorectal cancer. Cell 94: 703–714,

[96937] 1998.

[96938] 16813.

[96939] Wicki, R.; Marenholz, I.; Mischke, D.; Schafer, B. W.; Heizmann,

[96940] C. W.: Characterization of the human S100A12 (calgranulin C, p6,

[96941] CAAF1, CGRP) gene, a new member of the S100 gene

cluster on chromosome

[96942] 1q21. Cell Calcium 20: 459–464, 1996.

[96943] 16814.

[96944] Tan, K. B.; Harrop, J.; Reddy, M.; Young, P.; Terrett, J.;
Emery,

[96945] J.; Moore, G.; Truneh, A.: Characterization of a novel TNF-
like ligand

[96946] and recently described TNF ligand and TNF receptor su-
perfamily genes

[96947] and their constitutive and inducible expression in
hematopoietic and

[96948] non-hematopoietic cells. Gene 204: 35–46, 1997.

[96949] 16815.

[96950] Gruter, P.; Tabernero, C.; von Kobbe, C.; Schmitt, C.;
Saavedra,

[96951] C.; Bachi, A.; Wilm, M.; Felber, B. K.; Izaurralde, E.: TAP,
the

[96952] human homolog of Mex67p, mediates CTE-dependent
RNA export from the

[96953] nucleus. Molec. Cell 1: 649–659, 1998.

[96954] 16816.

- [96955] Segref, A.; Sharma, K.; Doye, V.; Hellwig, A.; Huber, J.;
Luhrmann,
[96956] R.; Hurt, E.: Mex67p, a novel factor for nuclear mRNA ex-
port, binds
[96957] to both poly(A)+ RNA and nuclear pores. EMBO J. 16:
3256–3271, 1997.
[96958] 16817.
- [96959] Yoon, D.-W.; Lee, H.; Seol, W.; DeMaria, M.; Rosenzweig,
M.; Jung,
[96960] J. U.: Tap: a novel cellular protein that interacts with Tip of
herpesvirus
[96961] saimiri and induces lymphocyte aggregation. Immunity 6:
571–582,
[96962] 1997.
[96963] 16818.
- [96964] Bonini, J. A.; Martin, S. K.; Dralyuk, F.; Roe, M. W.; Philip-
son,
[96965] L. H.; Steiner, D. F.: Cloning, expression, and chromoso-
mal mapping
[96966] of a novel human CC-chemokine receptor (CCR10) that
displays high-affinity
[96967] binding for MCP-1 and MCP-3. DNA Cell Biol. 16:

1249–1256, 1997.

[96968] 16819.

[96969] Nibbs, R. J. B.; Wylie, S. M.; Yang, J.; Landau, N. R.; Graham,

[96970] G. J.: Cloning and characterization of a novel promiscuous human

[96971] beta-chemokine receptor D6. *J. Biol. Chem.* 272: 32078–32083, 1997.

[96972] 16820.

[96973] Karet, F. E.; Finberg, K. E.; Nayir, A.; Bakkaloglu, A.; Ozen,

[96974] S.; Hulton, S. A.; Sanjad, S. A.; Al-Sabban, E. A.; Medina, J. F.;

[96975] Lifton, R. P.: Localization of a gene for autosomal recessive distal

[96976] renal tubular acidosis with normal hearing (rdRTA2) to 7q33–34. *Am.*

[96977] *J. Hum. Genet.* 65: 1656–1665, 1999.

[96978] 16821.

[96979] Smith, A. N.; Skaug, J.; Choate, K. A.; Nayir, A.; Bakkaloglu,

[96980] A.; Ozen, S.; Hulton, S. A.; Sanjad, S. A.; Al-Sabban, E. A.;

Lifton,

- [96981] R. P.; Scherer, S. W.; Karet, F. E.: Mutations in ATP6N1B, encoding
[96982] a new kidney vacuolar proton pump 116-kD subunit, cause recessive
[96983] distal renal tubular acidosis with preserved hearing. Nature Genet. 26:
[96984] 71–75, 2000.
[96985] 16822.
- [96986] McKie, J. M.; Sutherland, H. F.; Harvey, E.; Kim, U.-J.; Scambler,
[96987] P. J.: A human gene similar to Drosophila melanogaster peanut maps
[96988] to the DiGeorge syndrome region of 22q11. Hum. Genet. 101: 6–12,
[96989] 1997.
[96990] 16823.
- [96991] Yagi, M.; Zieger, B.; Roth, G. J.; Ware, J.: Structure and expression
[96992] of the human septin gene HCDCREL-1. Gene 212: 229–236, 1998.
[96993] 16824.

- [96994] Zieger, B.; Tran, H.; Hainmann, I.; Wunderle, D.; Zgaga-Griesz,
[96995] A.; Blaser, S.; Ware, J.: Characterization and expression
analysis
[96996] of two human septin genes, PNUTL1 and PNUTL2. Gene
261: 197–203,
[96997] 2000.
[96998] 16825.
- [96999] Latif, F.; Duh, F.–M.; Bader, S.; Sekido, Y.; Li, H.; Geil, L.;
[97000] Zbar, B.; Minna, J. D.; Lerman, M. I.: The human homolog
of the rodent
[97001] immediate early response genes, PC4 and TIS7, resides in
the lung
[97002] cancer tumor suppressor gene region on chromosome
3p21. Hum. Genet. 99:
[97003] 334–341, 1997.
[97004] 16826.
- [97005] Brandt, S.; Jentsch, T. J.: CIC–6 and CIC–7 are two novel
broadly
[97006] expressed members of the CLC chloride channel family.
FEBS Lett. 377:

- [97007] 15–20, 1995.
- [97008] 16827.
- [97009] Eggermont, J.; Buyse, G.; Voets, T.; Tytgat, J.; De Smedt, H.;
- [97010] Droogmans, G.: Alternative splicing of ClC-6 (a member of the ClC
- [97011] chloride-channel family) transcripts generates three truncated isoforms
- [97012] one of which, ClC-6c, is kidney-specific. *Biochem. J.* 325: 269–276,
- [97013] 1997.
- [97014] 16828.
- [97015] Cleiren, E.; Benichou, O.; Van Hul, E.; Gram, J.; Bollerslav, J.;
- [97016] Singer, F. R.; Beaverson, K.; Aledo, A.; Whyte, M. P.; Yoneyama, T.;
- [97017] deVernejou, M.-C.; Van Hul, W.: Albers-Schonberg disease (autosomal
- [97018] dominant osteopetrosis, type II) results from mutations in the CLCN7
- [97019] chloride channel gene. *Hum. Molec. Genet.* 10: 2861–2867, 2001.

- [97020] 16829.
- [97021] Hedblom, E.; Kirkness, E. F.: A novel class of GABA-A receptor
subunit in tissues of the reproductive system. J. Biol.
Chem. 272:
15346–15350, 1997.
- [97023] 16830.
- [97024] Caloca, M. J.; Garcia-Bermejo, M. L.; Blumberg, P. M.;
Lewin, N.
- [97026] E.; Kremmer, E.; Mischak, H.; Wang, S.; Nacro, K.; Bienfait,
B.; Marquez,
V. E.; Kazanietz, M. G.: Beta-2-chimaerin is a novel target
for diacylglycerol:
binding properties and changes in subcellular localization
mediated
by ligand binding to its C1 domain. Proc. Nat. Acad. Sci.
96: 11854–11859,
1999.
- [97030] 16831.
- [97031] Leung, T.; How, B.-E.; Manser, E.; Lim, L.: Cerebellar beta-
2-chimaerin,

- [97033] a GTPase-activating protein for p21 Ras-related Rac is specifically
- [97034] expressed in granule cells and has a unique N-terminal SH2 domain J.
- [97035] Biol. Chem. 269: 12888-12892, 1994.
- [97036] 16832.
- [97037] Leung, T.; How, B.-E.; Manser, E.; Lim, L.: Germ cell beta-chimaerin,
- [97038] a new GTPase-activating protein for p21rac, is specifically expressed
- [97039] during the acrosomal assembly stage in rat testis. J. Biol. Chem. 268:
- [97040] 3813-3816, 1993.
- [97041] 16833.
- [97042] Yuan, S.; Miller, D. W.; Barnett, G. H.; Hahn, J. F.; Williams, B. R.: Identification and characterization of human beta 2-chimaerin:
- [97044] association with malignant transformation in astrocytoma. Cancer
- [97045] Res. 55: 3456-3461, 1995.
- [97046] 16834.

- [97047] Holland, P. C.; Dillon, M. J.; Pincott, J.; Simmonds, H. A.; Barratt,
T. M.: Hypoxanthine guanine phosphoribosyl transferase deficiency
presenting with gout and renal failure in infancy. Arch. Dis. Child. 58:
831–833, 1983.
- [97051] 16835.
- [97052] Jolly, D. J.; Esty, A. C.; Bernard, H. U.; Friedmann, T.: Isolation
of a genomic clone partially encoding human hypoxanthine phosphoribosyltransferase. Proc. Nat. Acad. Sci. 79: 5038–5041, 1982.
- [97055] 16836.
- [97056] Jolly, D. J.; Okayama, H.; Berg, P.; Esty, A. C.; Filpula, D.; Bohlen, P.; Johnson, G. G.; Shively, J. E.; Hunkapiller, T.; Friedmann, T.: Isolation and characterization of a full-length expressible cDNA
for human hypoxanthine phosphoribosyltransferase. Proc. Nat. Acad.

- [97060] Sci. 80: 477–481, 1983.
- [97061] 16837.
- [97062] Chung, E.; Hanukoglu, A.; Rees, M.; Thompson, R.; Dillon, M.; Hanukoglu, I.; Bistritzer, T.; Kuhnle, U.; Seckl, J.; Gardiner, R. M.: Exclusion of the locus for autosomal recessive pseudohypoaldosteronism type 1 from the mineralocorticoid receptor gene region on human chromosome 4q by linkage analysis. J. Clin. Endocr. Metab. 80: 3341–3345, 1995.
- [97063] 16838.
- [97064] Schwienbacher, C.; Sabbioni, S.; Campi, M.; Veronese, A.; Bernardi, G.; Menegatti, A.; Hatada, I.; Mukai, T.; Ohashi, H.; Barbanti-Brodano, G.; Croce, C. M.; Negrini, M.: Transcriptional map of 170-kb region at chromosome 11p15.5: identification and mutational analysis of the BWR1A gene reveals the presence of mutations in tumor

samples. Proc.

[97073] Nat. Acad. Sci. 95: 3873–3878, 1998.

[97074] 16839.

[97075] Bartsch, O.; Kress, W.; Wagner, A.; Seemanova, E.: The novel contiguous

[97076] gene syndrome of myotubular myopathy (MTM1), male hypogenitalism and

[97077] deletion in Xq28: report of the first familial case. Cytogenet. Cell

[97078] Genet. 85: 310–314, 1999.

[97079] 16840.

[97080] Nagase, T.; Ishikawa, K.; Suyama, M.; Kikuno, R.; Hiro-sawa, M.;

[97081] Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Prediction

[97082] of the coding sequences of unidentified human genes. XIII. The complete

[97083] sequences of 100 new cDNA clones from brain which code for large proteins

[97084] in vitro. DNA Res. 6: 63–70, 1999.

[97085] 16841.

- [97086] Philibert, R. A.; Winfield, S. L.; Sandhu, H. K.; Martin, B. M.;
- [97087] Ginns, E. I.: The structure and expression of the human
neurologin-3
- [97088] gene. *Gene* 246: 303-310, 2000.
- [97089] 16842.
- [97090] Zonana, J.; Clarke, A.; Sarfarazi, M.; Thomas, N. S. T.;
- Roberts,
- [97091] K.; Marymee, K.; Harper, P. S.: X-linked hypohidrotic ecto-
dermal
- [97092] dysplasia: localization within the region Xq11-21.1 by
linkage analysis
- [97093] and implications for carrier detection and prenatal diag-
nosis. *Am.*
- [97094] *J. Hum. Genet.* 43: 75-85, 1988.
- [97095] 16843.
- [97096] Arico, M.; Imashuku, S.; Clementi, R.; Hibi, S.; Teramura,
T.;
- [97097] Danesino, C.; Haber, D. A.; Nichols, K. E.: Hemophagocytic
lymphohistiocytosis
- [97098] due to germline mutations in SH2D1A, the X-linked lym-
phoproliferative

- [97099] disease gene. *Blood* 97: 1131–1133, 2001.
- [97100] 16844.
- [97101] Arkwright, P. D.; Makin, G.; Will, A. M.; Ayres, M.; Gokhale, D.
- [97102] A.; Fergusson, W. D.; Taylor, G. M.: X linked lymphoproliferative
- [97103] disease in a United Kingdom family. *Arch. Dis. Child.* 79: 52–55,
- [97104] 1998.
- [97105] 16845.
- [97106] Bar, R. S.; DeLor, C. J.; Clausen, K. P.; Hurtubise, P.; Henle, W.; Hewetson, J. F.: Fatal infectious mononucleosis in a family. *New*
- [97107] *Eng. J. Med.* 290: 363–367, 1974.
- [97108] 16846.
- [97109] Benoit, L.; Wang, X.; Pabst, H. F.; Dutz, J.; Tan, R.: Cutting
- [97110] edge: defective NK cell activation in X-linked lymphoproliferative
- [97111] disease. *J. Immun.* 165: 3549–3553, 2000.
- [97112] 16847.
- [97113]

- [97114] Brandau, O.; Schuster, V.; Weiss, M.; Hellebrand, H.; Fink, F.
- [97115] M.; Kreczy, A.; Friedrich, W.; Strahm, B.; Niemeyer, C.; Be-lohradsky,
- [97116] B. H.; Meindl, A.: Epstein-Barr virus-negative boys with non-Hodgkin
- [97117] lymphoma are mutated in the SH2D1A gene, as are patients with X-linked
- [97118] lymphoproliferative disease (XLP). Hum. Molec. Genet. 8: 2407-2413,
- [97119] 1999.
- [97120] 16848.
- [97121] Coffey, A. J.; Brooksbank, R. A.; Brandau, O.; Oohashi, T.; Howell,
- [97122] G. R.; Bye, J. M.; Cahn, A. P.; Durham, J.; Heath, P.; Wray, P.; Pavitt,
- [97123] R.; Wilkinson, J.; and 31 others: Host response to EBV infection
- [97124] in X-linked lymphoproliferative disease results from mutations in
- [97125] an SH2-domain encoding gene. Nature Genet. 20: 129-135, 1998.

- [97126] 16849.
- [97127] Czar, M. J.; Kersh, E. N.; Mijares, L. A.; Lanier, G.; Lewis, J.;
- [97128] Yap, G.; Chen, A.; Sher, A.; Duckett, C. S.; Ahmed, R.;
Schwartzberg,
- [97129] P. L.: Altered lymphocyte responses and cytokine produc-
tion in mice
- [97130] deficient in the X-linked lymphoproliferative disease gene
SH2D1A/DSHP/SAP. Proc.
- [97131] Nat. Acad. Sci. 98: 7449–7454, 2001.
- [97132] 16850.
- [97133] Dutz, J. P.; Benoit, L.; Wang, X.; Demetrick, D. J.; Junker,
A.;
- [97134] de Sa, D.; Tan, R.: Lymphocytic vasculitis in X-linked lym-
phoproliferative
- [97135] disease. Blood 97: 95–100, 2001.
- [97136] 16851.
- [97137] Grierson, H. L.; Skare, J.; Church, J.; Silberman, T.; Davis, J.
- [97138] R.; Kibrinsky, N.; McGregor, R.; Israels, S.; McCarty, J.;
Andrews,
- [97139] L. G.; Blecha, T.; Erdman, S.; Obringer, A.; Scharnhorst, D.;
Purtilo,

- [97140] D. T.: Evaluation of families wherein a single male manifests a phenotype
- [97141] of X-linked lymphoproliferative disease (XLP). Am. J. Med. Genet. 47:
- [97142] 458–463, 1993.
- [97143] 16852.
- [97144] Hambleton, G.; Cottom, D. G.: Familial lymphoma. Proc. Roy.
- [97145] Soc. Med. 62: 1095 only, 1969.
- [97146] 16853.
- [97147] Hamilton, J. K.; Paquin, L. A.; Sullivan, J. L.; Maurer, H. S.;
- [97148] Cruzei, F. G.; Provisor, A. J.; Steuber, C. P.; Hawkins, E.; Yawn,
- [97149] D.; Cornet, J.; Clausen, K.; Finkelstein, G. Z.; Landing, B.; Grunnet,
- [97150] M.; Purtilo, D. T.: X-linked lymphoproliferative syndrome registry
- [97151] report. J. Pediat. 96: 669–673, 1980.
- [97152] 16854.
- [97153] Harris, A.; Docherty, Z.: X-linked lymphoproliferative disease:

[97154] a karyotype analysis. Cytogenet. Cell Genet. 47: 92–94, 1988.

[97155] 16855.

[97156] Harris, A.; Lenoir, G. M.; Lankester, S. A.: X-linked lymphoproliferative

[97157] disease: linkage studies using DNA probes. Clin. Genet. 33: 162–168,

[97158] 1988.

[97159] 16856.

[97160] Hayoz, D.; Lenoir, G. M.; Nicole, A.; Pugin, P.; Regamey, C.:

[97161] X-linked lymphoproliferative syndrome: identification of a large family

[97162] in Switzerland. Am. J. Med. 84: 529–534, 1988.

[97163] 16857.

[97164] Klein, G.; Klein, E.: Sinking surveillance's flagship. Nature 395:

[97165] 441–445, 1998.

[97166] 16858.

[97167] Levine, P. H.; Kamaraju, L. S.; Connelly, R. R.; Berard, C. W.;

- [97168] Dorfman, R. F.; Magrath, I.; Easton, J. M.: The American Burkitt's
- [97169] Lymphoma Registry: eight years' experience. Cancer 49: 1016–1022,
- [97170] 1982.
- [97171] 16859.
- [97172] Loeffel, S.; Chang, C.-H.; Heyn, R.; Harada, S.; Lipscomb, H.;
- [97173] Sinangil, F.; Volsky, D. J.; McClain, K.; Ochs, H.; Purtilo, D. T.
- [97174] : Necrotizing lymphoid vasculitis in X-linked lymphoproliferative
- [97175] syndrome. Arch. Path. Lab. Med. 109: 546–550, 1985.
- [97176] 16860.
- [97177] Lyon, M. F.; Loutit, J. F.: X-linked factor in acquired immunodeficiency
- [97178] syndrome?. (Letter) Lancet I: 768 only, 1983.
- [97179] 16861.
- [97180] Mulley, J. C.; Turner, A. M.; Gedeon, A. K.; Berdoukas, V. A.;
- [97181] Huang, T. H. M.; Ledbetter, D. H.; Grierson, H.; Purtilo, D.

T.:

[97182] X-linked lymphoproliferative disease: prenatal detection
of an unaffected

[97183] histocompatible male. Clin. Genet. 42: 76-79, 1992.

[97184] 16862.

[97185] Eva, A.; Aaronson, S. A.: Isolation of a new human onco-
gene from

[97186] a diffuse B-cell lymphoma. Nature 316: 273-275, 1985.

[97187] 16863.

[97188] Anderson, S. R.; Warburg, M.: Norrie's disease. Arch. Oph-
thal. 66:

[97189] 614-618, 1961.

[97190] 16864.

[97191] Bergen, A. A. B.; Wapenaar, M. C.; Schuurman, E. J. M.;
Diergaarde,

[97192] P. J.; Lerach, H.; Monaco, A. P.; Bakker, E.; Bleeker-
Wagemakers,

[97193] E. M.; van Ommen, G. J. B.: Detection of a new submicro-
scopic Norrie

[97194] disease deletion interval with a novel DNA probe isolated
by differential

- [97195] Alu PCR fingerprint cloning. *Cytogenet. Cell Genet.* 62: 231–235,
- [97196] 1993.
- [97197] 16865.
- [97198] Berger, W.; Meindl, A.; van de Pol, T. J. R.; Cremers, F. P. M.;
- [97199] Ropers, H. H.; Doerner, C.; Monaco, A.; Bergen, A. A. B.; Lebo, R.;
- [97200] Warburg, M.; Zergollern, L.; Lorenz, B.; Gal, A.; Bleeker-Wagemakers,
- [97201] E. M.; Meitinger, T.: Isolation of a candidate gene for Norrie disease
- [97202] by positional cloning. *Nature Genet.* 1: 199–203, 1992.
- [97203] 16866.
- [97204] Berger, W.; van de Pol, D.; Bachner, D.; Oerlemans, F.; Winkens,
- [97205] H.; Hameister, H.; Wieringa, B.; Hendriks, W.; Ropers, H.–H.: An
- [97206] animal model for Norrie disease (ND): gene targeting of the mouse
- [97207] ND gene. *Hum. Molec. Genet.* 5: 51–59, 1996.

- [97208] 16867.
- [97209] Eva, A.; Vecchio, G.; Rao, C. D.; Tronick, S. R.; Aaronson, S.
- [97210] A.: The predicted DBL oncogene product defines a distinct class of
- [97211] transforming proteins. Proc. Nat. Acad. Sci. 85: 2061–2065, 1988.
- [97212] 16868.
- [97213] Galland, F.; Stefanova, M.; Lafage, M.; Birnbaum, D.: Localization
- [97214] of the 5-prime end of the MCF2 oncogene to human chromosome 15q15–q23.
- [97215] Cytogenet. Cell Genet. 60: 114–116, 1992.
- [97216] 16869.
- [97217] Grant, S. G.; Mattei, M.-G.; Galland, F.; Stephenson, D. A.; Keitz,
- [97218] B. T.; Birnbaum, D.; Chapman, V. M.: Localization of the mouse Mcf-2
- [97219] (Dbl) protooncogene within a conserved linkage group on the mouse
- [97220] X chromosome. Cytogenet. Cell Genet. 54: 175–181, 1990.

- [97221] 16870.
- [97222] Nguyen, C.; Pontarotti, P.; Birnbaum, D.; Chimini, G.; Rey, J.
- [97223] A.; Mattei, J.-F.; Jordan, B. R.: Large scale physical mapping in
- [97224] the q27 region of the human X chromosome: the coagulation factor IX
- [97225] gene and the mcf.2 transforming sequence are separated by at most
- [97226] 270 kilobase pairs and are surrounded by several 'HTF islands.' EMBO
- [97227] J. 6: 3285-3289, 1987.
- [97228] 16871.
- [97229] Nguyen, C.; Poustka, A.-M.; Djabali, M.; Roux, D.; Mattei, J.-F.;
- [97230] Lehrach, H.; Jordan, B. R.: Large-scale mapping and chromosome jumping
- [97231] in the q27 region of the human X chromosome. Genomics 5: 298-303,
- [97232] 1989.
- [97233] 16872.

- [97234] Noguchi, T.; Mattei, M.-G.; Oberle, I.; Planche, J.; Imbert, J.;
- [97235] Pelassy, C.; Birg, F.; Birnbaum, D.: Localization of the mcf.2 transforming
- [97236] sequence to the X chromosome. EMBO J. 6: 1301-1307, 1987.
- [97237] 16873.
- [97238] Ron, D.; Tronick, S. R.; Aaronson, S. A.; Eva, A.: Molecular cloning
- [97239] and characterization of the human DBL proto-oncogene: evidence that
- [97240] its overexpression is sufficient to transform NIH/3T3 cells. EMBO
- [97241] J. 7: 2465-2473, 1988.
- [97242] 16874.
- [97243] Srivastava, S. K.; Wheelock, R. H. P.; Aaronson, S. A.; Eva, A.
- [97244] : Identification of the protein encoded by the human dif-fuse B-cell
- [97245] lymphoma (dbl) oncogene. Proc. Nat. Acad. Sci. 83: 8868-8872, 1986.
- [97246] 16875.

- [97247] Tronick, S. R.; McBride, O. W.; Popescu, N. C.; Eva, A.:
Chromosomal
- [97248] localization of DBL oncogene sequences. Genomics 5:
546–553, 1989.
- [97249] 16876.
- [97250] Giovane, A.; Sobieszczuk, P.; Mignon, C.; Mattei, M.–G.;
Wasylyk,
- [97251] B.: Locations of the ets subfamily members net, elk1, and
sap1 (ELK3,
- [97252] ELK1, and ELK4) on three homologous regions of the
mouse and human
- [97253] genomes. Genomics 29: 769–772, 1995.
- [97254] 16877.
- [97255] Acland, G. M.; Blanton, S. H.; Hershfield, B.; Aguirre, G. D.:
- [97256] XLPR: a canine retinal degeneration inherited as an X–
linked trait. Am.
- [97257] J. Med. Genet. 52: 27–33, 1994.
- [97258] 16878.
- [97259] Aldred, M. A.; Teague, P. W.; Jay, M.; Bunday, S.; Redmond,
R.
- [97260] M.; Jay, B.; Bird, A. C.; Bhattacharya, S. S.; Wright, A. F.:

Retinitis

- [97261] pigmentosa families showing apparent X linked inheritance but unlinked
- [97262] to the RP2 or RP3 loci. J. Med. Genet. 31: 848-852, 1994.
- [97263] 16879.
- [97264] Allan, W.: Eugenic significance of retinitis pigmentosa. Arch.
- [97265] Ophthal. 18: 938-947, 1937.
- [97266] 16880.
- [97267] Bhattacharya, S. S.; Clayton, J. F.; Harper, P. S.; Hoare, G. W.;
- [97268] Jay, M. R.; Lyness, A. L.; Wright, A. F.: A genetic linkage study
- [97269] of a kindred with X-linked retinitis pigmentosa. Brit. J. Ophthal. 69:
- [97270] 340-347, 1985.
- [97271] 16881.
- [97272] Bhattacharya, S. S.; Wright, A. F.; Clayton, J. F.; Price, W. H.;
- [97273] Phillips, C. I.; McKeown, C. M. E.; Jay, M.; Bird, A. C.; Pearson,

- [97274] P. L.; Southern, E. M.; Evans, H. J.: Close genetic linkage between
- [97275] X-linked retinitis pigmentosa and a restriction fragment length polymorphism
- [97276] identified by recombinant DNA probe L1.28. Nature 309: 253–255,
- [97277] 1984.
- [97278] 16882.
- [97279] Bird, A. C.: X-linked retinitis pigmentosa. Brit. J. Ophthal. 59:
- [97280] 177–199, 1975.
- [97281] 16883.
- [97282] Bunday, S.; Crews, S. J.: A study of retinitis pigmentosa in the
- [97283] City of Birmingham. (Letter) J. Med. Genet. 23: 188–191, 1986.
- [97284] 16884.
- [97285] Chapple, J. P.; Hardcastle, A. J.; Grayson, C.; Spackman, L. A.;
- [97286] Willison, K. R.; Cheetham, M. E.: Mutations in the N-terminus of

- [97287] the X-linked retinitis pigmentosa protein RP2 interfere with the normal
- [97288] targeting of the protein to the plasma membrane. Hum. Molec. Genet. 9:
- [97289] 1919–1926, 2000.
- [97290] 16885.
- [97291] Chen, J.-D.; Halliday, F.; Keith, G.; Sheffield, L.; Dickinson, P.; Gray, R.; Constable, I.; Denton, M.: Linkage heterogeneity between
- [97292] X-linked retinitis pigmentosa and a map of 10 RFLP loci. Am. J. Hum.
- [97293] Genet. 45: 401–411, 1989.
- [97294] 16886.
- [97295] Chen, J. D.; Denton, M. J.; Donald, J.; Serravalle, S.; Colley, P.; Lam Po Tang, P. R. L.; Halliday, F. B.: Analysis of linkage relationships
- [97296] of X-linked retinitis pigmentosa with the following Xp loci: DXS7,
- [97297] OTC, DXS84, XJ-1, DXS164, and DXS28. (Abstract) Cytogenet. Cell Genet. 46:
- [97300] 592 only, 1987.

- [97301] 16887.
- [97302] Chen, J. D.; Halliday, F.; Keith, G.; Sheffield, L.; Fraser, N.;
- [97303] Dickinson, P.; Gray, R.; Constable, I.; Denton, M. J.: Two
loci for
- [97304] X-linked retinitis pigmentosa suggested by multipoint
linkage analysis
- [97305] in nine families. (Abstract) Cytogenet. Cell Genet. 51: 976
only,
- [97306] 1989.
- [97307] 16888.
- [97308] Clayton, J. F.; Wright, A. F.; Jay, M.; McKeown, C. M. E.;
Dempster,
- [97309] M.; Jay, B. S.; Bird, A. C.; Bhattacharya, S. S.: Genetic link-
age
- [97310] between X-linked retinitis pigmentosa and DNA probe
DXS7 [L1.28]:
- [97311] further linkage data, heterogeneity testing, and risk esti-
mation. Hum.
- [97312] Genet. 74: 168-171, 1986.
- [97313] 16889.
- [97314] Coleman, M.; Bhattacharya, S.; Lindsay, S.; Wright, A.; Jay,

M.;

- [97315] Litt, M.; Craig, I.; Davies, K.: Localization of the mi-
crosatellite
- [97316] probe DXS426 between DXS7 and DXS255 on Xp and link-
age to X-linked
- [97317] retinitis pigmentosa. Am. J. Hum. Genet. 47: 935-940,
1990.
- [97318] 16890.
- [97319] Ernst, W.; Clover, G.; Faulkner, D. J.: X-linked retinitis pig-
mentosa:
- [97320] reduced rod flicker sensitivity in heterozygous females.
Invest.
- [97321] Ophthal. Vis. Sci. 6: 812-816, 1981.
- [97322] 16891.
- [97323] Farrar, G. J.; Geraghty, M. T.; Moloney, J. M. B.; McConnell,
- [97324] D. J.; Humphries, P.: Linkage analysis of X linked retinitis
pigmentosa
- [97325] in the Irish population. J. Med. Genet. 25: 222-226, 1988.
- [97326] 16892.
- [97327] Fishman, G. A.; Farber, M. D.; Derlacki, D. J.: X-linked re-
tinitis

- [97328] pigmentosa: profile of clinical findings. Arch. Ophthalm.
106: 369–375,
[97329] 1988.
[97330] 16893.
[97331] Friedrich, U.; Warburg, M.; Jorgensen, A. L.: X-inactivation
[97332] pattern in carriers of X-linked retinitis pigmentosa: a
valuable means
[97333] of prognostic evaluation?. Hum. Genet. 92: 359–363,
1993.
[97334] 16894.
[97335] Friedrich, U.; Warburg, M.; Kruse, T. A.; Andreasson, S.: X-
linked
[97336] retinitis pigmentosa: new map studies of XLRP2, and a
possible human
[97337] centromere effect. Hum. Genet. 88: 683–687, 1992.
[97338] 16895.
[97339] Friedrich, U.; Warburg, M.; Wieacker, P.; Wienker, T. F.;
Gal,
[97340] A.; Ropers, H.–H.: X-linked retinitis pigmentosa: linkage
with the
[97341] centromere and a cloned DNA sequence from the proxi-

mal short arm of

[97342] the X chromosome. Hum. Genet. 71: 93–99, 1985.

[97343] 16896.

[97344] Gieser, D. K.; Fishman, G. A.; Cunha-Vaz, J.: X-linked recessive

[97345] retinitis pigmentosa and vitreous fluorophotometry: a study of female

[97346] heterozygotes. Arch. Ophthal. 98: 307–310, 1980.

[97347] 16897.

[97348] Grutzner, P.; Sanger, R.; Spivey, B. E.: Linkage studies in X-linked

[97349] retinitis pigmentosa. Humangenetik 14: 155–158, 1972.

[97350] 16898.

[97351] Heck, A. F.: Presumptive X-linked intermediate transmission of

[97352] retinal degenerations: variations and coincidental occurrence with

[97353] ataxia in a large family. Arch. Ophthal. 70: 143–149, 1963.

[97354] 16899.

[97355] Hoare, G. W.: Choroido-retinal dystrophy. Brit. J. Ophthal.

49:

[97356] 449–459, 1965.

[97357] 16900.

[97358] Bermingham, J. R., Jr.; Scherer, S. S.; O'Connell, S.; Arroyo,

[97359] E.; Kalla, K. A.; Powell, F. L.; Rosenfeld, M. G.: Tst–

1/Oct–6/SCIP

[97360] regulates a unique step in peripheral myelination and is
required

[97361] for normal respiration. *Genes Dev.* 10: 1751–1762, 1996.

[97362] 16901.

[97363] Jaegle, M.; Mandemakers, W.; Broos, L.; Zwart, R.; Karis,
A.; Visser,

[97364] P.; Grosveld, F.; Meijer, D.: The POU factor Oct–6 and
Schwann cell

[97365] differentiation. *Science* 273: 507–510, 1996.

[97366] 16902.

[97367] Sumiyama, K.; Washio–Watanabe, K.; Ono, T.; Yoshida, M.
C.; Hayakawa,

[97368] T.; Ueda, S.: Human class III POU genes, POU3F1 and
POU3F3, map to

[97369] chromosomes 1p34.1 and 3p14.1. *Mammalian Genome* 9:

180-181, 1998.

[97370] 16903.

[97371] Costache, M.; Apoil, P.-A.; Cailleau, A.; Elmgren, A.; Larson,

[97372] G.; Henry, S.; Blancher, A.; Iordachescu, D.; Oriol, R.; Mollicone,

[97373] R.: Evolution of fucosyltransferase genes in vertebrates. J. Biol.

[97374] Chem. 272: 29721-29728, 1997.

[97375] 16904.

[97376] Yanagidani, S.; Uozumi, N.; Ihara, Y.; Miyoshi, E.; Yamaguchi,

[97377] N.; Taniguchi, N.: Purification and cDNA cloning of GDP-L-Fuc:N-acetyl-beta-D-glucosaminide:alpha-1-6

[97378] fucosyltransferase (alpha-1-6 FucT) from human gastric cancer MKN45

[97379] cells. J. Biochem. 121: 626-632, 1997.

[97380] 16905.

[97381] Bekri, S.; Adelaide, J.; Merscher, S.; Grosgeorge, J.; Caroli-Bosc,

[97382] F.; Perucca-Lostanlen, D.; Kelley, P. M.; Pebusque, M.-J.;

Theillet,

- [97383] C.; Birnbaum, D.; Gaudray, P.: Detailed map of a region commonly
[97384] amplified at 11q13-q14 in human breast carcinoma. Cy-
togenet. Cell
[97385] Genet. 79: 125-131, 1997.
[97386] 16906.
- [97387] Goodwin, R. G.; Alderson, M. R.; Smith, C. A.; Armitage, R.
J.;
- [97388] VandenBos, T.; Jerzy, R.; Tough, T. W.; Schoenborn, M. A.;
Davis-Smith,
[97389] T.; Hennen, K.; Falk, B.; Cosman, D.; Baker, E.; Sutherland,
G. R.;
- [97390] Grabstein, K. H.; Farrah, T.; Giri, J. G.; Beckmann, M. P.:
Molecular
[97391] and biological characterization of a ligand for CD27 de-
fines a new
[97392] family of cytokines with homology to tumor necrosis fac-
tor. Cell 73:
[97393] 447-456, 1993.
[97394] 16907.
- [97395] Kawabata, M.; Saeki, K.: Multiple alternative transcripts of

the

[97396] human homologue of the mouse TRAD/R51H3/RAD51D gene, a member of the

[97397] rec A/RAD51 gene family. Biochem. Biophys. Res. Commun. 257: 156–162,

[97398] 1999.

[97399] 16908.

[97400] Pittman, D. L.; Weinberg, L. R.; Schimenti, J. C.: Identification,

[97401] characterization, and genetic mapping of Rad51d, a new mouse and human

[97402] RAD51/RecA-related gene. Genomics 49: 103–111, 1998.

[97403] 16909.

[97404] Bell, B.; Scheer, E.; Tora, L.: Identification of hTAFII80-delta

[97405] links apoptotic signaling pathways to transcription factor TFIID function. Molec.

[97406] Cell 8: 591–600, 2001.

[97407] 16910.

[97408] Hisatake, K.; Ohta, T.; Takada, R.; Guermah, M.; Horikoshi, M.;

- [97409] Nakatani, Y.; Roeder, R. G.: Evolutionary conservation of human TATA-binding-polypeptide-associated
- [97410] factors TAFII31 and TAFII80 and interactions of TAFII80 with other
- [97411] TAFs and with general transcription factors. Proc. Nat. Acad. Sci. 92:
- [97412] 8195–8199, 1995.
- [97413] 16911.
- [97414] Weinzierl, R. O. J.; Ruppert, S.; Dynlacht, B. D.; Tanese, N.;
- [97415] Tjian, R.: Cloning and expression of Drosophila TAFII60 and human
- [97416] TAFII70 reveal conserved interactions with other subunits of TFIID. EMBO
- [97417] J. 12: 5303–5309, 1993.
- [97418] 16912.
- [97419] de Winter, J. P.; Waisfisz, Q.; Rooimans, M. A.; van Berkel, C.
- [97420] G. M.; Bosnoyan–Collins, L.; Alon, N.; Carreau, M.; Bender, O.; Demuth,
- [97421] I.; Schindler, D.; Pronk, J. C.; Arwert, F.; Hoehn, H.; Digweed, M.;

- [97422] Buchwald, M.; Joenje, H.: The Fanconi anaemia group G gene FANCG
- [97423] is identical with XRCC9. Nature Genet. 20: 281–283, 1998.
- [97424] 16913.
- [97425] Demuth, I.; Wlodarski, M.; Tipping, A. J.; Morgan, N. V.; de Winter,
- [97426] J. P.; Thiel, M.; Grasl, S.; Schindler, D.; D'Andrea, A. D.; Altay,
- [97427] C.; Kayserili, H.; Zatterdale, A.; Kunze, J.; Ebell, W.; Mathew, C.
- [97428] G.; Joenje, H.; Sperling, K.; Digweed, M.: Spectrum of mutations
- [97429] in the Fanconi anaemia group G gene, FANCG/XRCC9. Europ. J. Hum.
- [97430] Genet. 8: 861–868, 2000.
- [97431] 16914.
- [97432] Futaki, M.; Watanabe, S.; Kajigaya, S.; Liu, J. M.: Fanconi anemia
- [97433] protein, FANCG, is a phosphoprotein and is upregulated with FANCA
- [97434] after TNF- α treatment. Biochem. Biophys. Res. Com-

mun. 281: 347–351,

[97435] 2001.

[97436] 16915.

[97437] Koomen, M.; Cheng, N. C.; van de Vrugt, H. J.; Godthelp, B. C.;

[97438] van der Valk, M. A.; Oostra, A. B.; Zdzienicka, M. Z.; Joenje, H.;

[97439] Arwert, F.: Reduced fertility and hypersensitivity to mitomycin C

[97440] characterize Fancg/Xrcc9 null mice. Hum. Molec. Genet. 11: 273–281,

[97441] 2002.

[97442] 16916.

[97443] Kuang, Y.; Garcia-Higuera, I.; Moran, A.; Mondoux, M.; Digweed,

[97444] M.; D'Andrea, A. D.: Carboxy terminal region of the Fanconi anemia

[97445] protein, FANCG/XRCC9, is required for functional activity. Blood 96:

[97446] 1625–1632, 2000.

[97447] 16917.

- [97448] Liu, N.; Lamerdin, J. E.; Tucker, J. D.; Zhou, Z.-Q.; Walter, C.
- [97449] A.; Albala, J. S.; Busch, D. B.; Thompson, L. H.: The human XRCC9
- [97450] gene corrects chromosomal instability and mutagen sensitivities in
- [97451] CHO UV40 cells. *Proc. Nat. Acad. Sci.* 94: 9232–9237, 1997.
- [97452] 16918.
- [97453] Saar, K.; Schindler, D.; Wegner, R. D.; Reis, A.; Wienker, T. F.;
- [97454] Hoehn, H.; Joenje, H.; Sperling, K.; Digweed, M.: Localisation of
- [97455] a Fanconi anaemia gene to chromosome 9p. *Europ. J. Hum. Genet.* 6:
- [97456] 501–508, 1998.
- [97457] 16919.
- [97458] Yamada, T.; Tachibana, A.; Shimizu, T.; Mugishima, H.; Okubo, M.;
- [97459] Sasaki, M. S.: Novel mutations of the FANCG gene causing alternative
- [97460] splicing in Japanese Fanconi anemia. *J. Hum. Genet.* 45:

159–166,

[97461] 2000.

[97462] 16920.

[97463] Yang, Y.; Kuang, Y.; Montes De Oca, R.; Hays, T.; Moreau, L.;

[97464] Lu, N.; Seed, B.; D'Andrea, A. D.: Targeted disruption of the murine

[97465] Fanconi anemia gene, Fancg/Xrcc9. Blood 98: 3435–3440, 2001.

[97466] 16921.

[97467] Tsai, K. J.; Chen, S. K.; Ma, Y. L.; Hsu, W. L.; Lee, E. H. Y.

[97468] : sgk, a primary glucocorticoid-induced gene, facilitates memory consolidation

[97469] of spatial learning in rats. Proc. Nat. Acad. Sci. 99: 3990–3995,

[97470] 2002.

[97471] 16922.

[97472] Waldegger, S.; Barth, P.; Raber, G.; Lang, F.: Cloning and characterization

[97473] of a putative human serine/threonine protein kinase transcriptionally

- [97474] modified during anisotonic and isotonic alterations of cell volume. Proc.
- [97475] Nat. Acad. Sci. 94: 4440–4445, 1997.
- [97476] 16923.
- [97477] Lynch, R. A.; Piper, M.; Bankier, A.; Bhugra, B.; Surti, U.; Liu,
- [97478] J.; Buckler, A.; Dear, P. H.; Menon, A. G.: Genomic and functional
- [97479] map of the chromosome 14 t(12;14) breakpoint cluster region in uterine
- [97480] leiomyoma. Genomics 52: 17–26, 1998.
- [97481] 16924.
- [97482] Adam–Klages, S.; Adam, D.; Wiegmann, K.; Struve, S.; Kolanus, W.;
- [97483] Schneider–Mergener, J.; Kronke, M.: FAN, a novel WD–repeat protein,
- [97484] couples the p55 TNF–receptor to neutral sphingomyelinase. Cell 86:
- [97485] 937–947, 1996.
- [97486] 16925.
- [97487] Zhang, Y.; Kreder, D.; Schwandner, R.; Krut, O.; Scherer,

- G.; Adam-Klages,
[97488] S.; Siebert, R.; Kronke, M.; Schlegelberger, B.: Assignment
of the
[97489] human FAN protein gene (NSMAF) to human chromosome
region 8q12-q13
[97490] by in situ hybridization. Cytogenet. Cell Genet. 87:
115-116, 1999.
[97491] 16926.
- [97492] Ide, H.; Katoh, M.; Sasaki, H.; Yoshida, T.; Aoki, K.; Nawa,
Y.;
- [97493] Osada, Y.; Sugimura, T.; Terada, M.: Cloning of human
bone morphogenetic
[97494] protein type 1B receptor (BMPR-1B) and its expression in
prostate
[97495] cancer in comparison with other BMPRs. Oncogene 14:
1377-1382, 1997.
[97496] 16927.
- [97497] ten Dijke, P.; Yamashita, H.; Ichijo, H.; Franzen, P.; Laiho,
M.;
- [97498] Miyazono, K.; Heldin, C.-H.: Characterization of type I re-
ceptors
[97499] for transforming growth factor-beta and activin. Science

264: 101–104,

[97500] 1994.

[97501] 16928.

[97502] Mitsuhashi, N.; Miki, T.; Senbongi, H.; Yokoi, N.; Yano, H.; Miyazaki,

[97503] M.; Nakajima, N.; Iwanaga, T.; Yokoyama, Y.; Shibata, T.; Seino, S.

[97504] : MTABC3, a novel mitochondrial ATP-binding cassette protein involved

[97505] in iron homeostasis. J. Biol. Chem. 275: 17536–17540, 2000.

[97506] 16929.

[97507] Bai, C.; Connolly, B.; Metzker, M. L.; Hilliard, C. A.; Liu, X.;

[97508] Sandig, V.; Soderman, A.; Galloway, S. M.; Liu, Q.; Austin, C. P.;

[97509] Caskey, C. T.: Overexpression of M68/DCR3 in human gastrointestinal

[97510] tract tumors independent of gene amplification and its location in

[97511] a four-gene cluster. Proc. Nat. Acad. Sci. 97: 1230–1235, 2000.

[97512] 16930.

- [97513] Pitti, R. M.; Marsters, S. A.; Lawrence, D. A.; Roy, M.; Kischkel,
- [97514] F. C.; Dowd, P.; Huang, A.; Donahue, C. J.; Sherwood, S. W.; Baldwin,
- [97515] D. T.; Godowski, P. J.; Wood, W. I.; Gurney, A. L.; Hillan, K. J.;
- [97516] Cohen, R. L.; Goddard, A. D.; Botstein, D.; Ashkenazi, A.: Genomic
- [97517] amplification of a decoy receptor for Fas ligand in lung and colon
- [97518] cancer. *Nature* 396: 699–703, 1998.
- [97519] 16931.
- [97520] Ringstad, N.; Nemoto, Y.; De Camilli, P.: The SH3p4/Sh3p8/SH3p13
- [97521] protein family: binding partners for synaptojanin and dynamin via
- [97522] a Grb2-like Src homology 3 domain. *Proc. Nat. Acad. Sci.* 94: 8569–8574,
- [97523] 1997.
- [97524] 16932.
- [97525] Sittler, A.; Walter, S.; Wedemeyer, N.; Hasenbank, R.; Scherzinger,

- [97526] E.; Eickhoff, H.; Bates, G. P.; Lehrach, H.; Wanker, E. E.:
SH3GL3
- [97527] associates with the huntington exon 1 protein and pro-
motes the formation
- [97528] of polyglIn-containing protein aggregates. Molec. Cell 2:
427-436,
- [97529] 1998.
- [97530] 16933.
- [97531] Sparks, A. B.; Hoffman, N. G.; McConnell, S. J.; Fowlkes, D.
M.;
- [97532] Kay, B. K.: Cloning of ligand targets: systematic isolation
of SH3
- [97533] domain-containing proteins. Nature Biotech. 14:
741-744, 1996.
- [97534] 16934.
- [97535] Fracchiolla, N. S.; Colombo, G.; Finelli, P.; Maiolo, A. T.;
Neri,
- [97536] A.: EHT, a new member of the MTG8/ETO gene family,
maps on 20q11
- [97537] region and is deleted in acute myeloid leukemias. (Letter)
Blood 92:
- [97538] 3481-3484, 1998.

- [97539] 16935.
- [97540] Kitabayashi, I.; Ida, K.; Morohoshi, F.; Yokoyama, A.; Mit-
suhashi,
- [97541] N.; Shimizu, K.; Nomura, N.; Hayashi, Y.; Ohki, M.: The
AML1-MTG8
- [97542] leukemic fusion protein forms a complex with a novel
member of the
- [97543] MTG8(ETO/CDR) family, MTGR1. Molec. Cell. Biol. 18:
846-858, 1998.
- [97544] 16936.
- [97545] Jensen, D. E.; Proctor, M.; Marquis, S. T.; Gardner, H. P.;
Ha,
- [97546] S. I.; Chodosh, L. A.; Ishov, A. M.; Tommerup, N.; Vissing,
H.; Sekido,
- [97547] Y.; Minna, J.; Borodovsky, A.; Schultz, D. C.; Wilkinson, K.
D.; Maul,
- [97548] G. G.; Barlev, N.; Berger, S. L.; Prendergast, G. C.;
Rauscher, F.
- [97549] J., III: BAP1: a novel ubiquitin hydrolase which binds to the
BRCA1
- [97550] RING finger and enhances BRCA1-mediated cell growth
suppression. Oncogene 16:

[97551] 1097–1112, 1998.

[97552] 16937.

[97553] Nelson, H.; Mandiyan, S.; Noumi, T.; Moriyama, Y.; Miedel, M. C.;

[97554] Nelson, N.: Molecular cloning of cDNA encoding the C subunit of H(+)-ATPase

[97555] from bovine chromaffin granules. J. Biol. Chem. 265: 20390–20393,

[97556] 1990.

[97557] 16938.

[97558] van Hille, B.; Vanek, M.; Richener, H.; Green, J. R.; Bilbe, G.

[97559] : Cloning and tissue distribution of subunits C, D, and E of the human

[97560] vacuolar H(+)-ATPase. Biochem. Biophys. Res. Commun. 197: 15–21,

[97561] 1993.

[97562] 16939.

[97563] Yasuda, T.; Yamaguchi, N.; Kobayashi, K.; Nishi, I.; Hori-nouchi,

[97564] H.; Jalil, M. A.; Li, M. X.; Ushikai, M.; Iijima, M.; Kondo, I.; Saheki,

- [97565] T.: Identification of two novel mutations in the SLC25A13 gene and
- [97566] detection of seven mutations in 102 patients with adult-onset type
- [97567] II citrullinemia. Hum. Genet. 107: 537–545, 2000.
- [97568] 16940.
- [97569] Kondoh, N.; Schweinfest, C. W.; Henderson, K. W.; Papas, T. S.
- [97570] : Differential expression of S19 ribosomal protein, laminin-binding
- [97571] protein, and human lymphocyte antigen class I messenger RNAs associated
- [97572] with colon carcinoma progression and differentiation. Cancer Res. 52:
- [97573] 791–796, 1992.
- [97574] 16941.
- [97575] Matsson, H.; Klar, J.; Draptchinskaia, N.; Gustavsson, P.; Carlsson,
- [97576] B.; Bowers, D.; de Bont, E.; Dahl, N.: Truncating ribosomal protein
- [97577] S19 mutations and variable clinical expression in Diamond-Blackfan

- [97578] anemia. Hum. Genet. 105: 496–500, 1999.
- [97579] 16942.
- [97580] Tentler, D.; Gustavsson, P.; Elinder, G.; Eklof, O.; Gordon, L.;
- [97581] Mandel, A.; Dahl, N.: A microdeletion in 19q13.2 associated with
- [97582] mental retardation, skeletal malformations, and Diamond–Blackfan anaemia
- [97583] suggests a novel contiguous gene syndrome. J. Med. Genet. 37: 128–131,
- [97584] 2000.
- [97585] 16943.
- [97586] Hoornaert, I.; Marynen, P.; Baens, M.: CREBL2, a novel transcript
- [97587] from the chromosome 12 region flanked by ETV6 and CDKN1B. Genomics 51:
- [97588] 154–157, 1998.
- [97589] 16944.
- [97590] Nishiu, J.; Tanaka, T.; Nakamura, Y.: Identification of a novel
- [97591] gene (ECM2) encoding a putative extracellular matrix pro-

tein expressed

[97592] predominantly in adipose and female-specific tissues and its chromosomal

[97593] localization to 9q22.3. Genomics 52: 378-381, 1998.

[97594] 16945.

[97595] Anderson, P.; Nagler-Anderson, C.; O'Brien, C.; Levine, H.; Watkins,

[97596] S.; Slayter, H. S.; Blue, M. L.; Schlossman, S. F.: A monoclonal

[97597] antibody reactive with a 15-kDa cytoplasmic granule-associated protein

[97598] defines a subpopulation of CD8+ T lymphocytes. J. Immun. 144: 574-582,

[97599] 1990.

[97600] 16946.

[97601] Forch, P.; Puig, O.; Kedersha, N.; Martinez, C.; Granneman, S.;

[97602] Seraphin, B.; Anderson, P.; Valcarcel, J.: The apoptosis-promoting

[97603] factor TIA-1 is a regulator of alternative pre-mRNA splicing. Molec.

[97604] Cell 6: 1089-1098, 2000.

- [97605] 16947.
- [97606] Kawakami, A.; Tian, Q.; Streuli, M.; Poe, M.; Edelhoff, S.;
Disteche,
- [97607] C. M.; Anderson, P.: Intron-exon organization and chro-
mosomal localization
- [97608] of the human TIA-1 gene. J. Immun. 152: 4937-4945,
1994.
- [97609] 16948.
- [97610] Tian, Q.; Streuli, M.; Saito, H.; Schlossman, S. F.; Ander-
son,
- [97611] P.: A polyadenylate binding protein localized to the gran-
ules of
- [97612] cytolytic lymphocytes induces DNA fragmentation in tar-
get cells. Cell 67:
- [97613] 629-639, 1991.
- [97614] 16949.
- [97615] Neubauer, G.; King, A.; Rappsilber, J.; Calvio, C.; Watson,
M.;
- [97616] Ajuh, P.; Sleeman, J.; Lamond, A.; Mann, M.: Mass spec-
trometry and
- [97617] EST-database searching allows characterisation of the
multi-protein

- [97618] spliceosome complex. *Nature Genet.* 20: 46–50, 1998.
- [97619] 16950.
- [97620] Talbot, K.; Miguel–Aliaga, I.; Mohaghegh, P.; Ponting, C. P.; Davies, P.; Davies, K. E.: Characterization of a gene encoding survival motor neuron (SMN)–related protein, a constituent of the spliceosome complex. *Hum. Molec. Genet.* 7: 2149–2156, 1998.
- [97624] 16951.
- [97625] Sillekens, P. T. G.; Beijer, R. P.; Habets, W. J.; van Venrooij, W. J.: Molecular cloning of the cDNA for the human U2 snRNA–specific A–prime protein. *Nucleic Acids Res.* 17: 1893–1906, 1989.
- [97628] 16952.
- [97629] Zheng, B.; Albrecht, U.; Kaasik, K.; Sage, M.; Lu, W.; Vaishnav, S.; Li, Q.; Sun, Z. S.; Eichele, G.; Bradley, A.; Lee, C. C.: Nonredundant roles of the mPer1 and mPer2 genes in the mammalian

circadian clock. Cell 105:

[97632] 683–694, 2001.

[97633] 16953.

[97634] Mattei, M.–G.; Roeckel, N.; Olsen, B. R.; Apte, S. S.: Genes of

[97635] the membrane–type matrix metalloproteinase (MT–MMP) gene family, MMP14,

[97636] MMP15, and MMP16, localize to human chromosomes 14, 16, and 8, respectively. Genomics 40:

[97637] 168–169, 1997.

[97638] 16954.

[97639] Sato, H.; Tanaka, M.; Takino, T.; Inoue, M.; Seiki, M.: Assignment

[97640] of the human genes for membrane–type–1, –2, and –3 matrix metalloproteinases

[97641] (MMP14, MMP15, and MMP16) to 14q12.2, 16q12.2–q21, and 8q21, respectively,

[97642] by in situ hybridization. Genomics 39: 412–413, 1997.

[97643] 16955.

[97644] Takino, T.; Sato, H.; Shinagawa, A.; Seiki, M.: Identification

[97645] of the second membrane–type matrix metalloproteinase

(MT-MMP-2) gene

[97646] from a human placenta cDNA library: MT-MMPs form a unique membrane-type

[97647] subclass in the MMP family. *J. Biol. Chem.* 270: 23013-23020, 1995.

[97648] 16956.

[97649] Will, H.; Hinzmann, B.: cDNA sequence and mRNA tissue distribution

[97650] of a novel human matrix metalloproteinase with a potential transmembrane

[97651] segment. *Europ. J. Biochem.* 231: 602-608, 1995.

[97652] 16957.

[97653] Alliel, P. M.; Perin, J.-P.; Jolles, P.; Bonnet, F. J.: Testican,

[97654] a multidomain testicular proteoglycan resembling modulators of cell

[97655] social behaviour. *Europ. J. Biochem.* 214: 347-350, 1993.

[97656] 16958.

[97657] Charbonnier, F.; Perin, J.-P.; Mattei, M.-G.; Camuzat, A.; Bonnet,

[97658] F.; Gressin, L.; Alliel, P. M.: Genomic organization of the human

- [97659] SPOCK gene and its chromosomal localization to 5q31.
Genomics 48:
[97660] 377–380, 1998.
- [97661] 16959.
- [97662] Cheng, J.-F.; Boyartchuk, V.; Zhu, Y.: Isolation and mapping of
[97663] human chromosome 21 cDNA: progress in constructing a
chromosome 21
[97664] expression map. Genomics 23: 75–84, 1994.
- [97665] 16960.
- [97666] Sirotkin, H.; O'Donnell, H.; DasGupta, R.; Halford, S.; St.
Jore,
[97667] B.; Puech, A.; Parimoo, S.; Morrow, B.; Skoultchi, A.;
Weissman, S.
[97668] M.; Scambler, P.; Kucherlapati, R.: Identification of a new
human
[97669] catenin gene family member (ARVCF) from the region
deleted in velo–cardio–facial
[97670] syndrome. Genomics 41: 75–83, 1997.
- [97671] 16961.
- [97672] Andrews, N. C.: Mining copper transport genes. Proc. Nat.

Acad.

[97673] Sci. 98: 6543–6545, 2001.

[97674] 16962.

[97675] Boultonwood, J.; Strickson, A. J.; Jabs, E. W.; Cheng, J.-F.; Fidler,

[97676] C.; Wainscoat, J. S.: Physical mapping of the human ATX1
homologue

[97677] (HAH1) to the critical region of the 5q- syndrome within
5q32, and

[97678] immediately adjacent to the SPARC gene. Hum. Genet.
106: 127–129,

[97679] 2000.

[97680] 16963.

[97681] Hamza, I.; Faisst, A.; Prohaska, J.; Chen, J.; Gruss, P.;
Gitlin,

[97682] J. D.: The metallochaperone Atox1 plays a critical role in
perinatal

[97683] copper homeostasis. Proc. Nat. Acad. Sci. 98: 6848–6852,
2001.

[97684] 16964.

[97685] Klomp, L. W. J.; Lin, S.-J.; Yuan, D. S.; Klausner, R. D.; Cu-

lotta,

[97686] V. C.; Gitlin, J. D.: Identification and functional expression of

[97687] HAH1, a novel human gene involved in copper homeostasis. J. Biol.

[97688] Chem. 272: 9221–9226, 1997.

[97689] 16965.

[97690] Castro, A. M.; Snyder, L. M.: G6PD San Jose: a new variant characterized

[97691] by NADPH inhibition studies. Humangenetik 21: 361–363, 1974.

[97692] 16966.

[97693] Cayanis, E.; Gomperts, E. D.; Balinsky, D.; Disler, P.; Myers,

[97694] A.: G6PD Hillbrow: a new variant of glucose–6–phosphate dehydrogenase

[97695] associated with drug–induced haemolytic anaemia. Brit. J. Haemat. 30:

[97696] 343–350, 1975.

[97697] 16967.

[97698] Cayanis, E.; Lane, A. B.; Jenkins, T.; Nurse, G. T.; Balinsky,

[97699] D.: Glucose–6–phosphate dehydrogenase Porbandar: a

new slow variant

[97700] with slightly reduced activity in a South African family of Indian

[97701] descent. *Biochem. Genet.* 15: 765–773, 1977.

[97702] 16968.

[97703] Cederbaum, A. I.; Beutler, E.: Nonspherocytic hemolytic anemia

[97704] due to G6PD Grand Prairie. *I.R.C.S.* 3: 579, 1975.

[97705] 16969.

[97706] Chan, T. K.; Todd, D.: Characteristics and distribution of glucose–6–phosphate

[97707] dehydrogenase–deficient variants in South China. *Am. J. Hum. Genet.* 24:

[97708] 475–484, 1972.

[97709] 16970.

[97710] Chan, T. K.; Todd, D.; Lai, M. C. S.: Glucose 6–phosphate dehydrogenase:

[97711] identity of erythrocyte and leukocyte enzyme with report of a new

[97712] variant in Chinese. *Biochem. Genet.* 6: 119–124, 1972.

[97713] 16971.

- [97714] Chang, J.-G.; Chiou, S.-S.; Perng, L.-I.; Chen, T.-C.; Liu, T.-C.;
- [97715] Lee, L.-S.; Chen, P.-H.; Tang, T. K.: Molecular characterization
- [97716] of glucose-6-phosphate dehydrogenase (G6PD) deficiency by natural
- [97717] and amplification created restriction sites: five mutations account
- [97718] for most G6PD deficiency cases in Taiwan. Blood 80: 1079-1082, 1992.
- [97719] 16972.
- [97720] Chen, E. Y.; Cheng, A.; Lee, A.; Kuang, W.-J.; Hillier, L.; Green,
- [97721] P.; Schlessinger, D.; Ciccodicola, A.; D'Urso, M.: Sequence of human
- [97722] glucose-6-phosphate dehydrogenase cloned in plasmids and a yeast artificial
- [97723] chromosome. Genomics 10: 792-800, 1991.
- [97724] 16973.
- [97725] Chen, H.-L.; Huang, M.-J.; Huang, C.-S.; Tang, T. K.: G6PD NanKang
- [97726] (517 T-to-C; 173 phe-to-leu): a new Chinese G6PD vari-

ant associated

[97727] with neonatal jaundice. Hum. Hered. 46: 201–204, 1996.

[97728] 16974.

[97729] Chernyak, N. B.; Batischev, A. I.; Lanzina, N. V.; Tokarev, Y.

[97730] N.; Alexeyev, G. A.: Electrophoretic and kinetic properties of glucose-6-phosphate

[97731] dehydrogenase from erythrocytes or patients with hemolytic anemia,

[97732] related to deficiency of the enzyme activity. Vopr. Med. Khim. 23:

[97733] 166–171, 1977.

[97734] 16975.

[97735] Childs, B.; Zinkham, W.; Browne, E. A.; Kimbro, E. L.; Torbert,

[97736] J. V.: A genetic study of a defect in glutathione metabolism of the

[97737] erythrocytes. Bull. Johns Hopkins Hosp. 102: 21–37, 1958.

[97738] 16976.

[97739] Chiu, D. T. Y.; Zuo, L.; Chao, L.; Chen, E.; Louie, E.; Lubin,

- [97740] B.; Liu, T. Z.; Du, C. S.: Molecular characterization of glucose-6-phosphate
- [97741] dehydrogenase (G6PD) deficiency in patients of Chinese descent and
- [97742] identification of new base substitutions in the human G6PD gene. Blood 81:
- [97743] 2150-2154, 1993.
- [97744] 16977.
- [97745] Chockkalingam, K.; Board, P. G.: Further evidence for heterogeneity
- [97746] of glucose-6-phosphate dehydrogenase deficiency in Papua New Guinea. Hum.
- [97747] Genet. 56: 209-212, 1980.
- [97748] 16978.
- [97749] Chockkalingam, K.; Board, P. G.; Breguet, G.: Glucose-6-phosphate
- [97750] dehydrogenase variants of Bali Island (Indonesia). Hum. Genet. 60:
- [97751] 60-62, 1982.
- [97752] 16979.
- [97753] Chockkalingam, K.; Board, P. G.; Nurse, G. T.: Glucose-

6-phosphate

- [97754] dehydrogenase deficiency in Papua New Guinea: the description of 13
- [97755] new variants. Hum. Genet. 60: 189-192, 1982.
- [97756] 16980.
- [97757] Chuanshu, D.; Yankang, X.; Lin, W. Q. R.; Xiaoyun, H.: Studies
- [97758] on erythrocyte glucose-6-phosphate dehydrogenase variants in Chinese.
- [97759] I. Gd(-) Lizu-Baisha. Acta Acad. Med. Zhong Shan 2: 649-658, 1981.
- [97760] 16981.
- [97761] Cocco, P.; Todde, P.; Fornera, S.; Manca, M. B.; Manca, P. Sias,
- [97762] A. R.: Mortality in a cohort of men expressing the glucose-6-phosphate
- [97763] dehydrogenase deficiency. Blood 91: 706-709, 1998.
- [97764] 16982.
- [97765] Cohn, J.; Carter, N.; Warburg, M.: Glucose-6-phosphate dehydrogenase
- [97766] deficiency in a native Danish family: a new variant. Scand.

J. Haemat. 23:

[97767] 403–406, 1979.

[97768] 16983.

[97769] Colonna–Romano, S.; Iolascon, A.; Lippo, S.; Pinto, L.; Cutillo,

[97770] S.; Battistuzzi, G.: Genetic heterogeneity at the glucose–6-phosphate

[97771] dehydrogenase locus in southern Italy: a study on the population of

[97772] Naples. Hum. Genet. 69: 228–232, 1985.

[97773] 16984.

[97774] Cooper, D. W.; Johnston, P. G.; Murtagh, C. E.; Sharman, G. B.;

[97775] Vandeberg, J. L.; Poole, W. E.: Sex-linked isozymes and sex-chromosome

[97776] evolution and inactivation in kangaroos. In: Markeit, C. L.: Isozymes.

[97777] Developmental Biology. New York: Academic Press (pub.) III: 1975.

[97778] Pp. 559–573.

[97779] 16985.

- [97780] Cooper, M. R.; Dechatelet, L. R.; McCall, C. E.; Lavia, M. F.;
- [97781] Spurr, C. L.; Baehner, R. L.: Complete deficiency of leukocyte glucose-6-phosphate
- [97782] dehydrogenase with defective bactericidal activity. J. Clin. Invest. 51:
- [97783] 769-778, 1972.
- [97784] 16986.
- [97785] Corash, L.; Spielberg, S.; Bartsocas, C.; Boxer, L.; Steiner,
- [97786] R.; Sheetz, M.; Egan, M.; Schlesselman, J.; Schulman, J. D.: Reduced
- [97787] chronic hemolysis during high-dose vitamin E administration in Mediterranean-type
- [97788] glucose-6-phosphate dehydrogenase deficiency. New Eng. J. Med. 303:
- [97789] 416-420, 1980.
- [97790] 16987.
- [97791] Corcoran, C. M.; Calabro, V.; Tamagnini, G.; Town, M.; Haidar,
- [97792] B.; Vulliamy, T. J.; Mason, P. J.; Luzzatto, L.: Molecular heterogeneity

- [97793] underlying the G6PD Mediterranean phenotype. Hum. Genet. 88: 688–690, 1992.
- [97794] 1992.
- [97795] 16988.
- [97796] Costa, E.; Cabeda, J. M.; Vieira, E.; Pinto, R.; Pereira, S. A.; Ferraz, L.; Santos, R.; Barbot, J.: Glucose-6-phosphate dehydrogenase
- [97798] Aveiro: a de novo mutation associated with chronic non-spherocytic
- [97799] hemolytic anemia. Blood 95: 1499–1501, 2000.
- [97800] 16989.
- [97801] Crookston, J. H.; Yoshida, A.; Lin, M.; Booser, D. J.: G6PD Toronto. Biochem. Genet. 8: 259–265, 1973.
- [97802] 16990.
- [97803] Csepregthy, M.; Hall, M. K.; Berkow, R. L.; Jackson, S.; Prchal, J. T.: Characterization of a new G6PD variant: G6PD Titusville. Am. J. Med. Sci. 297: 114–117, 1989.

- [97807] 16991.
- [97808] Csepregyhy, M.; Yeilding, A.; Lilly, M.; Hall, K.; Scott, C. W.;
- [97809] Prchal, J. T.: Characterization of a new glucose-
6-phosphate dehydrogenase
- [97810] variant: G6PD Central City. Am. J. Hemat. 28: 61-62,
1988.
- [97811] 16992.
- [97812] D'Urso, M.; Luzzatto, L.; Perroni, L.; Ciccodicola, A.; Gen-
tile,
- [97813] G.; Peluso, I.; Persico, M. G.; Pizzella, T.; Toniolo, D.; Vul-
liamy,
- [97814] T. J.: An extensive search for RFLP in the human glucose-
6-phosphate
- [97815] dehydrogenase locus has revealed a silent mutation in the
coding sequence. Am.
- [97816] J. Hum. Genet. 42: 735-741, 1988.
- [97817] 16993.
- [97818] Roessler, B. J.; Palella, T. D.; Heidler, S.; Becker, M. A.:
- [97819] Identification of distinct PRPS1 mutations in two patients
with X-linked
- [97820] phosphoribosylpyrophosphate synthetase superactivity.

(Abstract) Clin.

[97821] Res. 39: 267A, 1991.

[97822] 16994.

[97823] Moreno Fuenmayor, H.; Roldan-Paris, L.; Bermudez, H.:
Ectodermal

[97824] dysplasia in females and inversion of chromosome 9. J.
Med. Genet. 18:

[97825] 214-217, 1981.

[97826] 16995.

[97827] Nakata, M.; Koshiba, H.; Eto, K.; Nance, W. E.: A genetic
study

[97828] of anodontia in X-linked hypohidrotic ectodermal dysplasia. Am. J.

[97829] Hum. Genet. 32: 908-919, 1980.

[97830] 16996.

[97831] Passarge, E.; Fries, E.: X-chromosome inactivation in X-linked

[97832] hypohidrotic ectodermal dysplasia. Nature N.B. 245:
58-59, 1973.

[97833] 16997.

[97834] Pinheiro, M.; Freire-Maia, N.: Christ-Siemens-Touraine

syndrome--a

[97835] clinical and genetic analysis of a large Brazilian kindred. I.

Affected

[97836] females. II. Affected males. III. Carrier detection. Am. J.

Med.

[97837] Genet. 4: 113-134, 1979.

[97838] 16998.

[97839] Pinheiro, M.; Ideriha, M. T.; Chautard-Freire-Maia, E. A.;

Freire-Maia,

- [97840] N.; Primo-Parmo, S. L.: Christ-Siemens-Touraine syndrome: investigations
- [97841] on two large Brazilian kindreds with a new estimate of the manifestation
- [97842] rate among carriers. Hum. Genet. 57: 428-431, 1981.
- [97843] 16999.
- [97844] Reed, W. B.; Lopez, D. A.; Landing, B. H.: Clinical spectrum
- [97845] of anhidrotic ectodermal dysplasia. Arch. Derm. 102: 134-143, 1970.
- [97846] 17000.
- [97847] Richards, W.; Kaplan, M.: Anhidrotic ectodermal dysplasia: an
- [97848] unusual case of pyrexia in the newborn. Am. J. Dis. Child. 117: 597-598, 1969.
- [97849] 17001.
- [97851] Roberts, E.: The inheritance of anhidrosis associated with anodontia. J.A.M.A. 93: 277-279, 1929.
- [97852] 17002.
- [97853]

- [97854] Saksena, S. S.; Bixler, D.: Facial morphometrics in the identification
- [97855] of gene carriers of X-linked hypohidrotic ectodermal dysplasia. Am.
- [97856] J. Med. Genet. 35: 105–114, 1990.
- [97857] 17003.
- [97858] Simpson, J. L.; Allen, F. H., Jr.; New, M.; German, J.: Absence
- [97859] of close linkage between the locus for Xg and the locus for anhidrotic
- [97860] ectodermal dysplasia. Vox Sang. 17: 465–467, 1969.
- [97861] 17004.
- [97862] Singh, A.; Jolly, S. S.; Kaur, S.: Hereditary ectodermal dysplasia. Brit.
- [97863] J. Derm. 74: 34–37, 1962.
- [97864] 17005.
- [97865] Soderholm, A.-L.; Kaitila, I.: Expression of X-linked hypohidrotic
- [97866] ectodermal dysplasia in six males and in their mothers. Clin. Genet. 28:
- [97867] 136–144, 1985.

- [97868] 17006.
- [97869] Srivastava, A. K.; Durmowicz, M. C.; Hartung, A. J.; Hudson, J.;
- [97870] Ouzts, L. V.; Donovan, D. M.; Cui, C.-Y.; Schlessinger, D.:
Ectodysplasin-A1
- [97871] is sufficient to rescue both hair growth and sweat glands
in Tabby
- [97872] mice. Hum. Molec. Genet. 10: 2973-2981, 2001.
- [97873] 17007.
- [97874] Srivastava, A. K.; Montonen, O.; Saarialho-Kere, U.; Chen,
E.;
- [97875] Baybayan, P.; Pispá, J.; Limon, J.; Schlessinger, D.; Kere, J.:
Fine
- [97876] mapping of the EDA gene: a translocation breakpoint is
associated
- [97877] with a CpG island that is transcribed. Am. J. Hum. Genet.
58: 126-132,
- [97878] 1996.
- [97879] 17008.
- [97880] Srivastava, A. K.; Pispá, J.; Hartung, A. J.; Du, Y.; Ezer, S.;
- [97881] Jenks, T.; Shimada, T.; Pekkanen, M.; Mikkola, M. L.; Ko,

M. S. H.;

- [97882] Thesleff, I.; Kere, J.; Schlessinger, D.: The Tabby phenotype is
[97883] caused by mutation in a mouse homologue of the EDA
gene that reveals
[97884] novel mouse and human exons and encodes a protein
(ectodysplasin-A)
[97885] with collagenous domains. Proc. Nat. Acad. Sci. 94:
13069–13074,
[97886] 1997.
[97887] 17009.
- [97888] Thadani, K. I.: The toothless men of Sind. J. Hered. 25:
483–484,
[97889] 1934.
[97890] 17010.
- [97891] Thurnam, J.: Two cases in which the skin, hair and teeth
were
[97892] very imperfectly developed. Proc. R. M. Chir. Soc. London
31: 71–82,
[97893] 1848.
[97894] 17011.

- [97895] Turleau, C.; Niaudet, P.; Cabanis, M.-O.; Plessis, G.; Cau, D.;
- [97896] de Grouchy, J.: X-linked hypohidrotic ectodermal dysplasia and t(X;12)
- [97897] in a female. Clin. Genet. 35: 462–466, 1989.
- [97898] 17012.
- [97899] Vargas, G. A.; Fantino, E.; George-Nascimento, C.; Gargus, J.
- [97900] J.; Haigler, H. T.: Reduced epidermal growth factor receptor expression
- [97901] in hypohidrotic ectodermal dysplasia and Tabby mice. J. Clin. Invest. 97:
- [97902] 2426–2432, 1996.
- [97903] 17013.
- [97904] Vincent, M. C.; Biancalana, V.; Ginisty, D.; Mandel, J. L.; Calvas,
- [97905] P.: Mutational spectrum of the ED1 gene in X-linked hypohidrotic
- [97906] ectodermal dysplasia. Europ. J. Hum. Genet. 9: 355–363, 2001.
- [97907] 17014.

- [97908] WPA, Federal Writers' Project: Mississippi: A Guide to the
[97909] Magnolia State. New York: Hastings House (pub.) 1938.
Pp. 359
[97910] only.
[97911] 17015.
[97912] Yotsumoto, S.; Fukumaru, S.; Matsushita, S.; Oku, T.;
Kobayashi,
[97913] K.; Saheki, T.; Kanzaki, T.: A novel point mutation of the
EDA gene
[97914] in a Japanese family with anhidrotic ectodermal dysplasia.
(Letter) J.
[97915] Invest. Derm. 111: 1246–1247, 1998.
[97916] 17016.
[97917] Ballabio, A.: Personal Communication. Houston, Texas
3/11/1993.
[97918] 17017.
[97919] Bardin, C. W.; Ross, G. T.; Rifkind, A. B.; Cargille, C. M.;
Lipsett,
[97920] M. B.: Studies of the pituitary Leydig cell axis in young
men with
[97921] hypogonadotropic hypogonadism and hyposmia: compar-

ison with normal

[97922] men, prepubertal boys, and hypo-pituitary patients. J.

Clin. Invest. 48:

[97923] 2046-2056, 1969.

[97924] 17018.

[97925] Bick, D.; Franco, B.; Sherins, R. J.; Heye, B.; Pike, L.; Crawford,

[97926] J.; Maddalena, A.; Incerti, B.; Pragliola, A.; Meitinger, T.; Ballabio,

[97927] A.: Brief report: intragenic deletion of the KALIG-1 gene in Kallmann's

[97928] syndrome. New Eng. J. Med. 326: 1752-1755, 1992.

[97929] 17019.

[97930] Bick, D. P.; Schwanzel-Fukuda, M.; Pfaff, D. W.; Schorderet, D.

[97931] F.; Price, P. A.; Campbell, L.; Huff, R. W.; Moore, C. M.: Prenatal

[97932] diagnosis and investigation of a fetus with chondrodysplasia punctata,

[97933] ichthyosis and Kallmann syndrome due to an Xp deletion: evidence for

[97934] a neuronal migration defect in Kallmann syndrome.

(Abstract) Am.

[97935] J. Hum. Genet. 45 (suppl.): A252, 1989.

[97936] 17020.

[97937] Birnbacher, R.; Wandl-Vergesslich, K.; Frisch, H.: Diagnosis of

[97938] X-recessive Kallmann syndrome in early infancy: evidence of hypoplastic

[97939] rhinencephalon. Europ. J. Pediat. 153: 245-247, 1994.

[97940] 17021.

[97941] Bulow, H. E.; Berry, K. L.; Topper, L. H.; Peles, E.; Hobert, O.

[97942] : Heparan sulfate proteoglycan-dependent induction of axon branching

[97943] and axon misrouting by the Kallmann syndrome gene kal-1. Proc. Nat.

[97944] Acad. Sci. 99: 6346-6351, 2002.

[97945] 17022.

[97946] del Castillo, I.; Cohen-Salmon, M.; Blanchard, S.; Lutfalla, G.;

[97947] Petit, C.: Structure of the X-linked Kallmann syndrome gene and its

- [97948] homologous pseudogene on the Y chromosome. *Nature Genet.* 2: 305–310, 1992.
- [97949] 17023.
- [97950] 17023.
- [97951] De Morsier, G.: Etudes sur les dysraphies cranio-encephaliques.
- [97952] I. Agenesie des lobes olfactifs (telencephaloschizis lateral) et des
- [97953] commissures calleuse et anterieure (telencephaloschizis median): la
- [97954] dysplasie olfacto-genitale. *Schweiz. Arch. Neurol. Psychiat.* 74: 309–361, 1954.
- [97955] 17024.
- [97956] 17024.
- [97957] Basse, F.; Stout, J. G.; Sims, P. J.; Wiedmer, T.: Isolation of
- [97958] an erythrocyte membrane protein that mediates Ca^{2+} -dependent transbilayer
- [97959] movement of phospholipid. *J. Biol. Chem.* 271: 17205–17210, 1996.
- [97960] 17025.
- [97961] Garrett, R. M.; Johnson, J. L.; Graf, T. N.; Feigenbaum, A.;

Rajagopalan,

- [97962] K. V.: Human sulfite oxidase R160Q: identification of the mutation
[97963] in a sulfite oxidase-deficient patient and expression and characterization
[97964] of the mutant enzyme. Proc. Nat. Acad. Sci. 95: 6394-6398, 1998.
[97965] 17026.
- [97966] Kisker, C.; Schindelin, H.; Pacheco, A.; Wehbi, W. A.; Garrett,
[97967] R. M.; Rajagopalan, K. V.; Enemark, J. H.; Rees, D. C.: Molecular
[97968] basis of sulfite oxidase deficiency from the structure of sulfite
[97969] oxidase. Cell 91: 973-983, 1997.
[97970] 17027.
- [97971] Naylor, J.; Brinke, A.; Hassock, S.; Green, P. M.; Giannelli,
[97972] F.: Characteristic mRNA abnormality found in half the patients with
[97973] severe haemophilia A is due to large DNA inversions. Hum. Molec.
[97974] Genet. 2: 1773-1778, 1993.

- [97975] 17028.
- [97976] Renieri, A.; Seri, M.; Myers, J. C.; Pihlajaniemi, T.; Massella,
[97977] L.; Rizzoni, G.; De Marchi, M.: De novo mutation in the
COL4A5 gene
[97978] converting glycine 325 to glutamic acid in Alport syn-
drome. Hum.
[97979] Molec. Genet. 1: 127-129, 1992.
- [97980] 17029.
- [97981] Corral, J.; Forster, A.; Thompson, S.; Lampert, F.; Kaneko,
Y.;
- [97982] Slater, R.; Kroes, W. G.; van der Schoot, C. E.; Ludwig,
W.-D.; Karpas,
[97983] A.; Pocock, C.; Cotter, F.; Rabbitts, T. H.: Acute leukemias
of different
[97984] lineages have similar MLL gene fusions encoding related
chimeric proteins
[97985] resulting from chromosomal translocation. Proc. Nat.
Acad. Sci. 90:
[97986] 8538-8542, 1993.
- [97987] 17030.
- [97988] Knebelmann, B.; Deschenes, G.; Gros, F.; Hors, M.-C.;

Grunfeld,

- [97989] J.-P.; Tryggvason, K.; Gubler, M.-C.; Antignac, C.: Substitution
[97990] of arginine for glycine 325 in the collagen alpha-5 (IV)
chain associated
[97991] with X-linked Alport syndrome: characterization of the
mutation by
[97992] direct sequencing of PCR-amplified lymphoblast cDNA
fragments. Am.
[97993] J. Hum. Genet. 51: 135-142, 1992.
[97994] 17031.
[97995] Renieri, A.; Seri, M.; Myers, J. C.; Pihlajaniemi, T.; Sessa,
[97996] A.; Rizzoni, G.; De Marchi, M.: Alport syndrome caused by
a 5-prime
[97997] deletion within the COL4A5 gene. Hum. Genet. 89:
120-121, 1992.
[97998] 17032.
[97999] Smeets, H. J. M.; Melenhorst, J. J.; Lemmink, H. H.;
Schroder,
[98000] C. H.; Nelen, M. R.; Zhou, J.; Hostikka, S. L.; Tryggvason,
K.; Ropers,
[98001] H.-H.; Jansweijer, M. C. E.; Monnens, L. A. H.; Brunner, H.

G.; van

[98002] Oost, B. A.: Different mutations in the COL4A5 collagen gene in two

[98003] patients with different features of Alport syndrome. Kidney Int. 42:

[98004] 83–88, 1992.

[98005] 17033.

[98006] Thorner, P. S.; Zheng, K.; Kalluri, R.; Jacobs, R.; Hudson, B.

[98007] G.: Coordinate gene expression of the alpha-3, alpha-4, and alpha-5

[98008] chains of collagen type IV. J. Biol. Chem. 271: 13821–13828, 1996.

[98009] 17034.

[98010] Turco, A. E.; Rossetti, S.; Biasi, M. O.; Rizzoni, G.; Massella,

[98011] L.; Saarinen, N. H.; Renieri, A.; Pignatti, P. F.; De Marchi, M.:

[98012] A novel missense mutation in exon 3 of the COL4A5 gene associated

[98013] with late-onset Alport syndrome. Clin. Genet. 48: 261–263, 1995.

[98014] 17035.

- [98015] Vetrie, D.; Boye, E.; Flinter, F.; Bobrow, M.; Harris, A.: DNA
[98016] rearrangements in the alpha-5(IV) collagen gene (COL4A5)
of individuals
[98017] with Alport syndrome: further refinement using pulsed-
field gel electrophoresis. Genomics 14:
[98018] 624-633, 1992.
[98019] 17036.
[98020] Vetrie, D.; Flinter, F.; Bobrow, M.; Harris, A.: Long-range
mapping
[98021] of the gene for the human alpha-5(IV) collagen chain at
Xq22-q23. Genomics 12:
[98022] 130-138, 1992.
[98023] 17037.
[98024] Zheng, K.; Thorner, P. S.; Marrano, P.; Baumal, R.;
McInnes, R.
[98025] R.: Canine X chromosome-linked hereditary nephritis: a
genetic model
[98026] for human X-linked hereditary nephritis resulting from a
single base
[98027] mutation in the gene encoding the alpha-5 chain of colla-
gen type IV. Proc.

- [98028] Nat. Acad. Sci. 91: 3989–3993, 1994.
- [98029] 17038.
- [98030] Zhou, J.; Barker, D. F.; Hostikka, S. L.; Gregory, M. C.;
Atkin,
- [98031] C. L.; Tryggvason, K.: Single base mutation in alpha-5(IV)
collagen
- [98032] chain gene converting a conserved cysteine to serine in
Alport syndrome. Genomics 9:
- [98033] 10–18, 1991.
- [98034] 17039.
- [98035] Zhou, J.; Hertz, J. M.; Leinonen, A.; Tryggvason, K.: Com-
plete
- [98036] amino acid sequence of the human alpha-5(IV) collagen
chain and identification
- [98037] of a single-base mutation in exon 23 converting glycine
521 in the
- [98038] collagenous domain to cysteine in an Alport syndrome
patient. J.
- [98039] Biol. Chem. 267: 12475–12481, 1992.
- [98040] 17040.
- [98041] Zhou, J.; Hostikka, S. L.; Chow, L. T.; Tryggvason, K.:

Characterization

[98042] of the 3-prime half of the human type IV collagen alpha-5 gene that

[98043] is affected in the Alport syndrome. Genomics 9: 1-9, 1991.

[98044] 17041.

[98045] Oohashi, T.; Sugimoto, M.; Mattei, M.-G.; Ninomiya, Y.: Identification

[98046] of a new collagen IV chain, alpha-6(IV), by cDNA isolation and assignment

[98047] of the gene to chromosome Xq22, which is the same locus for COL4A5. J.

[98048] Biol. Chem. 269: 7520-7526, 1994.

[98049] 17042.

[98050] Oohashi, T.; Ueki, Y.; Sugimoto, M.; Ninomiya, Y.: Isolation and

[98051] structure of the COL4A6 gene encoding the human alpha-6(IV) collagen

[98052] chain and comparison with other type IV collagen genes. J. Biol.

[98053] Chem. 270: 26863-26867, 1995.

[98054] 17043.

- [98055] Sugimoto, M.; Oohashi, T.; Ninomiya, Y.: The genes COL4A5 and
- [98056] COL4A6, coding for basement membrane collagen chains alpha-5(IV) and
- [98057] alpha-6(IV), are located head-to-head in close proximity on human
- [98058] chromosome Xq22 and COL4A6 is transcribed from two alternative promoters. Proc.
- [98059] Nat. Acad. Sci. 91: 11679-11683, 1994.
- [98060] 17044.
- [98061] Parry, P.; Wei, Y.; Evans, G.: Cloning and characterization of
- [98062] the t(X;11) breakpoint from a leukemic cell line identify a new member
- [98063] of the forkhead gene family. Genes Chromosomes Cancer 11: 79-84,
- [98064] 1994.
- [98065] 17045.
- [98066] Peters, U.; Haberhausen, G.; Kostrzewa, M.; Nolte, D.; Muller,
- [98067] U.: AFX1 and p54(nrb): fine mapping, genomic structure, and exclusion

- [98068] as candidate genes of X-linked dystonia parkinsonism.
Hum. Genet. 100:
- [98069] 569–572, 1997.
- [98070] 17046.
- [98071] Bohme, B.; VandenBos, T.; Cerretti, D. P.; Park, L. S.;
Holtrich,
- [98072] U.; Rubsamen-Waigmann, H.; Strebhardt, K.: Cell-cell ad-
hesion mediated
- [98073] by binding of membrane-anchored ligand LERK-2 to the
EPH-related receptor
- [98074] human embryonal kinase 2 promotes tyrosine kinase ac-
tivity. J. Biol.
- [98075] Chem. 271: 24747–24752, 1996.
- [98076] 17047.
- [98077] Fletcher, F. A.; Huebner, K.; Shaffer, L. G.; Fairweather, N.
D.;
- [98078] Monaco, A. P.; Muller, U.; Druck, T.; Simoneaux, D. K.;
Chelly, J.;
- [98079] Belmont, J. W.; Beckmann, M. P.; Lyman, S. D.: Assignment
of the
- [98080] gene (EPLG2) encoding a high-affinity binding protein for
the receptor

[98081] tyrosine kinase Elk to a 200–kilobasepair region in human chromosome

[98082] Xq12. Genomics 25: 334–335, 1995.

[98083] 17048.

[98084] Fletcher, F. A.; Renshaw, B.; Hollingsworth, T.; Baum, P.; Lyman,

[98085] S. D.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Davison, B.

[98086] L.: Genomic organization and chromosomal localization of mouse Eplg2,

[98087] a gene encoding a binding protein for the receptor tyrosine kinase

[98088] Elk. Genomics 24: 127–132, 1994.

[98089] 17049.

[98090] Palmer, A.; Zimmer, M.; Erdmann, K. S.; Eulenburg, V.; Porthin,

[98091] A.; Heumann, R.; Deutsch, U.; Klein, R.: EphrinB phosphorylation

[98092] and reverse signaling: regulation by Src kinases and PTP-BL phosphatase. Molec.

[98093] Cell 9: 725–737, 2002.

- [98094] 17050.
- [98095] Gregor, P.; Nash, S. R.; Caron, M. G.; Seldin, M. F.; Warren, S.
- [98096] T.: Assignment of the creatine transporter gene (SLC6A8) to human
- [98097] chromosome Xq28 telomeric to G6PD. *Genomics* 25: 332–333, 1995.
- [98098] 17051.
- [98099] Hahn, K. A.; Salomons, G. S.; Tackels–Horne, D.; Wood, T. C.; Taylor,
- [98100] H. A.; Schroer, R. J.; Lubs, H. A.; Jakobs, C.; Olson, R. L.; Holden,
- [98101] K. R.; Stevenson, R. E.; Schwartz, C. E.: X-linked mental retardation
- [98102] with seizures and carrier manifestations is caused by a mutation in
- [98103] the creatine–transporter gene (SLC6A8) located in Xq28. *Am. J. Hum.*
- [98104] *Genet.* 70: 1349–1356, 2002.
- [98105] 17052.
- [98106] Item, C. B.; Stockler–Ipsiroglu, S.; Stromberger, C.; Muhl, A.;

- [98107] Alessandri, M. G.; Bianchi, M. C.; Tosetti, M.; Fornai, F.; Cioni,
[98108] G.: Arginine:glycine amidinotransferase deficiency: the third inborn
[98109] error of creatine metabolism in humans. Am. J. Hum. Genet. 69: 1127–33,
[98110] 2001.
[98111] 17053.
[98112] Iyer, G. S.; Krahe, R.; Goodwin, L. A.; Doggett, N. A.; Siciliano,
[98113] M. J.; Funanage, V. L.; Proujansky, R.: Identification of a testis-expressed
[98114] creatine transporter gene at 16p11.2 and confirmation of the X-linked
[98115] locus to Xq28. Genomics 34: 143–146, 1996.
[98116] 17054.
[98117] Nash, S. R.; Giros, B.; Kingsmore, S. F.; Rochelle, J. M.; Suter,
[98118] S. T.; Gregor, P.; Seldin, M. F.; Caron, M. G.: Cloning, pharmacological
[98119] characterization, and genomic localization of the human creatine transporter. Receptors

- [98120] Channels 2: 165–174, 1994.
- [98121] 17055.
- [98122] Salomons, G. S.; van Dooren, S. J. M.; Verhoeven, N. M.; Cecil,
- [98123] K. M.; Ball, W. S.; Degrauw, T. J.; Jakobs, C.: X-linked creatine-transporter
- [98124] gene (SLC6A8) defect: a new creatine-deficiency syndrome. Am. J. Hum. Genet. 68: 1497–1500, 2001.
- [98125] 17056.
- [98126] Sandoval, N.; Bauer, D.; Brenner, V.; Coy, J. F.; Drescher, B.;
- [98127] Kioschis, P.; Korn, B.; Nyakatura, G.; Poustka, A.; Reichwald, K.;
- [98128] Rosenthal, A.; Platzer, M.: The genomic organization of a human creatine
- [98129] transporter (CRTR) gene located in Xq28. Genomics 35: 383–385, 1996.
- [98130] 17057.
- [98131] Stockler, S.; Hanefeld, F.; Frahm, J.: Creatine replacement therapy

- [98133] in guanidinoacetate methyltransferase deficiency, a novel
inborn error
- [98134] of metabolism. Lancet 348: 789–90, 1996.
- [98135] 17058.
- [98136] Brzustowicz, L. M.; Farrell, S.; Khan, M. B.; Weksberg, R.:
Mapping
- [98137] of a new SGBS locus to chromosome Xp22 in a family with
a severe form
- [98138] of Simpson–Golabi–Behmel syndrome. Am. J. Hum. Genet.
65: 779–783,
- [98139] 1999.
- [98140] 17059.
- [98141] Filmus, J.; Church, J. G.; Buick, R. N.: Isolation of a cDNA
corresponding
- [98142] to a developmentally regulated transcript in rat intestine.
Molec.
- [98143] Cell Biol. 4243–4249, 1988.
- [98144] 17060.
- [98145] Gelb, B. D.; Shi, G.–P.; Chapman, H. A.; Desnick, R. J.: Pyc-
nodysostosis,
- [98146] a lysosomal disease caused by cathepsin K deficiency. Sci–

ence 273:

[98147] 1236–1238, 1996.

[98148] 17061.

[98149] Gelb, B. D.; Willner, J. P.; Dunn, T. M.; Kardon, N. B.; Verloes,

[98150] A.; Poncin, J.; Desnick, R. J.: Paternal uniparental disomy for chromosome

[98151] 1 revealed by molecular analysis of a patient with pycnodysostosis. Am.

[98152] J. Hum. Genet. 62: 848–854, 1998.

[98153] 17062.

[98154] Polymeropoulos, M. H.; Ortiz De Luna, R. I.; Ide, S. E.; Torres,

[98155] R.; Rubenstein, J.; Francomano, C. A.: The gene for pycnodysostosis

[98156] maps to human chromosome 1cen–q21. Nature Genet. 10: 238–239, 1995.

[98157] 17063.

[98158] Bourke, E.; Frindt, G.; Flynn, P.; Schreiner, G. E.: Primary hyperoxaluria

[98159] with normal alpha–ketoglutarate:glyoxylate carboligase

activity: treatment

[98160] with isocarboxazid. Ann. Intern. Med. 76: 279–284, 1972.

[98161] 17064.

[98162] Danpure, C. J.: Personal Communication. Middlesex, England

[98163] 6/16/1988.

[98164] 17065.

[98165] Danpure, C. J.: Primary hyperoxaluria type 1 and peroxisome-to-mitochondrion

[98166] mistargeting of alanine:glyoxylate aminotransferase. Biochimie 75:

[98167] 309–315, 1993.

[98168] 17066.

[98169] Danpure, C. J.; Jennings, P. R.: Peroxisomal alanine:glyoxylate

[98170] aminotransferase deficiency in primary hyperoxaluria type I. FEBS

[98171] Lett. 201: 20–24, 1986.

[98172] 17067.

[98173] Danpure, C. J.; Jennings, P. R.; Watts, R. W. E.: Enzymolog–

ical

[98174] diagnosis of primary hyperoxaluria type 1 by measurement of hepatic

[98175] alanine:glyoxylate aminotransferase activity. Lancet I: 289-291,

[98176] 1987.

[98177] 17068.

[98178] Brunelli, S.; Faiella, A.; Capra, V.; Nigro, V.; Simeone, A.; Cama,

[98179] A.; Boncinelli, E.: Germline mutation in the homeobox gene EMX2 in

[98180] patients with severe schizencephaly. Nature Genet. 12: 94-96, 1996.

[98181] 17069.

[98182] Burg, J.; Conzelmann, E.; Sandhoff, K.; Solomon, E.; Swallow, D.

[98183] M.: Mapping of the gene coding for the human GM2 activator protein

[98184] to chromosome 5. Ann. Hum. Genet. 49: 41-45, 1985.

[98185] 17070.

[98186] Chen, B.; Rigat, B.; Curry, C.; Mahuran, D. J.: Structure of

the

- [98187] **GM2A gene: identification of an exon 2 nonsense mutation and a naturally occurring transcript with an in-frame deletion of exon 2. Am. J.**
- [98189] **Hum. Genet. 65: 77–87, 1999.**
- [98190] **17071.**
- [98191] **Conzelmann, E.; Sandhoff, K.: AB variant of infantile Gm2–gangliosidosis:**
- [98192] **deficiency of a factor necessary for stimulation of hexosaminidase**
- [98193] **A–catalyzed degradation of ganglioside Gm2 and glycolipid Ga2. Proc.**
- [98194] **Nat. Acad. Sci. 75: 3979–3983, 1978.**
- [98195] **17072.**
- [98196] **Hechtman, P.; Gordon, B. A.; Ng Ying Kin, N. M. K.: Deficiency**
- [98197] **of the hexosaminidase A activator protein in a case of GM2 gangliosidosis;**
- [98198] **variant AB. Pediat. Res. 16: 217–222, 1982.**
- [98199] **17073.**

- [98200] Heng, H. H. Q.; Xie, B.; Shi, X.-M.; Tsui, L.-C.; Mahuran, D. J.
- [98201] : Refined mapping of the GM2 activator protein (GM2A) locus to 5q31.3-q33.1,
- [98202] distal to the spinal muscular atrophy locus. *Genomics* 18: 429-431,
- [98203] 1993.
- [98204] 17074.
- [98205] Kleyn, P. W.; Brzustowicz, L. M.; Wilhelmsen, K. C.; Freimer, N.
- [98206] B.; Miller, J. M.; Munsat, T. L.; Gilliam, T. C.: Spinal muscular
- [98207] atrophy is not the result of mutations at the beta-hexosaminidase
- [98208] or GM(2)-activator locus. *Neurology* 41: 1418-1422, 1991.
- [98209] 17075.
- [98210] Li, S. C.; Hirabayashi, Y.; Li, Y. T.: A new variant of type-AB
- [98211] GM2-gangliosidosis. *Biochem. Biophys. Res. Commun.* 101: 479-485,
- [98212] 1981.

[98213] 17076.

[98214] Barton, D. E.; Yang-Feng, T. L.; Francke, U.: The human tyrosine

[98215] aminotransferase gene mapped to the long arm of chromosome 16 (region

[98216] 16q22-q24) by somatic cell hybrid analysis and in situ hybridization. Hum.

[98217] Genet. 72: 221-224, 1986.

[98218] 17077.

[98219] Beinfang, D. C.; Kuwabara, T.; Pueschel, S. M.: The Richner-Hanhart

[98220] syndrome: report of a case with associated tyrosinemia. Arch. Ophthal. 94:

[98221] 1133-1137, 1976.

[98222] 17078.

[98223] Bohnert, A.; Anton-Lamprecht, I.: Richner-Hanhart's syndrome:

[98224] ultrastructural abnormalities of epidermal keratinization indicating

[98225] a causal relationship to high intracellular tyrosine levels. J. Invest.

[98226] Derm. 79: 68–74, 1982.

[98227] 17079.

[98228] Buist, N.: Phenylketonuria and related problems. In: Nyhan, W.

[98229] L.: Amino Acid Metabolism and Genetic Variation. New York: McGraw-Hill

[98230] (pub.) 1967. Pp. 117 only.

[98231] 17080.

[98232] Chitayat, D.; Balbul, A.; Hani, V.; Mamer, O. A.; Clow, C.; Scriver,

[98233] C. R.: Hereditary tyrosinaemia type II in a consanguineous Ashkenazi

[98234] Jewish family: intrafamilial variation in phenotype; absence of parental

[98235] phenotype effects on the fetus. J. Inherit. Metab. Dis. 15: 198–203,

[98236] 1992.

[98237] 17081.

[98238] Crovato, F.; Desirello, G.; Gatti, R.; Babbini, N.; Rebora, A.

[98239] : Richner–Hanhart syndrome spares a plantar autograft. Arch. Derm. 121:

[98240] 539–540, 1985.

[98241] 17082.

[98242] Fellman, J. H.; Vanbellinghen, P. J.; Jones, R. T.; Koler, R. D.

[98243] : Soluble and mitochondrial forms of tyrosine aminotransferase: relationship

[98244] to human tyrosinemia. *Biochemistry* 8: 615–622, 1969.

[98245] 17083.

[98246] Garibaldi, L. R.; Siliato, F.; De Martini, I.; Scarsi, M. R.; Romano,

[98247] C.: Oculocutaneous tyrosinosis: report of two cases in the same family. *Helv.*

[98248] *Paediat. Acta* 32: 173–180, 1977.

[98249] 17084.

[98250] Goldsmith, L. A.; Kang, E. S.; Bienfang, D. C.; Jimbow, K.; Gerald,

[98251] P. S.; Baden, H. P.: Tyrosinemia with plantar and palmar keratosis

[98252] and keratitis. *J. Pediat.* 83: 798–805, 1973.

[98253] 17085.

- [98254] Hanhart, E.: Neue Sonderformen von Keratosis palmo-plantaris,
- [98255] u.a. eine regelmaessig-dominante mit systematisierten Lipomen, ferner
- [98256] 2 einfach-rezessive mit Schwachsinn und z.T. mit Hornhautveraenderungen
- [98257] des Auges (Ektodermatosyndrom). Dermatologica 94: 286-308, 1947.
- [98258] 17086.
- [98259] Hunziker, N.: Richner-Hanhart syndrome and tyrosinemia type II. Dermatologica 160:
- [98260] 180-189, 1980.
- [98261] 17087.
- [98262] Kennaway, N. G.; Buist, N. R. M.: Metabolic studies in a patient
- [98263] with hepatic cytosol tyrosine aminotransferase deficiency. Pediat.
- [98264] Res. 5: 287-297, 1971.
- [98265] 17088.
- [98266] Muller, G.; Scherer, G.; Zentgraf, H.; Ruppert, S.; Hermann,

- [98267] B.; Lehrach, H.; Schutz, G.: Isolation, characterization and chromosomal
- [98268] mapping of the mouse tyrosine aminotransferase gene. J. Molec. Biol. 184:
- [98269] 367–373, 1985.
- [98270] 17089.
- [98271] Natt, E.; Kao, F.-T.; Rettenmeier, R.; Scherer, G.: Assignment
- [98272] of the human tyrosine aminotransferase gene to chromosome 16. Hum.
- [98273] Genet. 72: 225–228, 1986.
- [98274] 17090.
- [98275] Kim, E. K.; Yoo, O. J.; Song, K. Y.; Yoo, H. W.; Choi, S. Y.;
- [98276] Cho, S. W.; Hahn, S. H.: Identification of three novel mutations
- [98277] and a high frequency of the arg778-to-leu mutation in Korean patients
- [98278] with Wilson disease. Hum. Mutat. 11: 275–278, 1998.
- [98279] 17091.
- [98280] Kuo, Y.-M.; Gitschier, J.; Packman, S.: Developmental expression

- [98281] of the mouse mottled and toxic milk genes suggests distinct functions
- [98282] for the Menkes and Wilson disease copper transporters. Hum. Molec.
- [98283] Genet. 6: 1043–1049, 1997.
- [98284] 17092.
- [98285] Loudianos, G.; Dessi, V.; Lovicu, M.; Angius, A.; Figus, A.; Lilliu,
- [98286] F.; De Virgiliis, S.; Nurchi, A. M.; Deplano, A.; Moi, P.; Pirastu,
- [98287] M.; Cao, A.: Molecular characterization of Wilson disease in the
- [98288] Sardinian population—evidence of a founder effect. Hum. Mutat. 14:
- [98289] 294–303, 1999.
- [98290] 17093.
- [98291] Loudianos, G.; Dessi, V.; Lovicu, M.; Angius, A.; Nurchi, A.;
- [98292] Sturniolo, G. C.; Marcellini, M.; Zancan, L.; Bragetti, P.; Akar,
- [98293] N.; Yagci, R.; Vegnente, A.; Cao, A.; Pirastu, M.: Further delineation

- [98294] of the molecular pathology of Wilson disease in the
Mediterranean
- [98295] population. Hum. Mutat. 12: 89–94, 1998.
- [98296] 17094.
- [98297] Okada, T.; Shiono, Y.; Hayashi, H.; Satoh, H.; Sawada, T.;
Suzuki,
- [98298] A.; Takeda, Y.; Yano, M.; Michitaka, K.; Onji, M.; Mabuchi,
H.: Mutational
- [98299] analysis of ATP7B and genotype–phenotype correlation in
Japanese with
- [98300] Wilson's disease. Hum. Mutat. 15: 454–462, 2000.
- [98301] 17095.
- [98302] Sasaki, N.; Hayashizaki, Y.; Muramatsu, M.; Matsuda, Y.;
Ando,
- [98303] Y.; Kuramoto, T.; Serikawa, T.; Azuma, T.; Naito, A.; Agui,
T.; Yamashita,
- [98304] T.; Miyoshi, I.; Takeichi, N.; Kasai, N.: The gene responsi-
ble for
- [98305] LEC hepatitis, located on rat chromosome 16, is the ho-
molog to the
- [98306] human Wilson disease gene. Biochem. Biophys. Res. Com-
mun. 202: 512–518,

- [98307] 1994.
- [98308] 17096.
- [98309] McKusick, V. A.: On the X Chromosome of Man. Washington: Am.
- [98310] Inst. Biol. Sci. (pub.) 1964.
- [98311] 17097.
- [98312] Ichinose, H.; Ohye, T.; Suzuki, T.; Sumi-Ichinose, C.; Nomura,
- [98313] T.; Hagino, Y.; Nagatsu, T.: Molecular cloning of the human Nurr1
- [98314] gene: characterization of the human gene and cDNAs. Gene 230: 233-239,
- [98315] 1999.
- [98316] 17098.
- [98317] Law, S. W.; Conneely, O. M.; DeMayo, F. J.; O'Malley, B. W.: Identification
- [98318] of a new brain-specific transcription factor, NURR1. Molec. Endocr. 6:
- [98319] 2129-2135, 1992.
- [98320] 17099.

- [98321] Mages, H. W.; Rilke, O.; Bravo, R.; Senger, G.; Kroczeck, R. A.
- [98322] : NOT, a human immediate-early response gene closely related to the
- [98323] steroid/thyroid hormone receptor NAK1/TR3. *Molec. Endocr.* 8: 1583-1591,
- [98324] 1994.
- [98325] 17100.
- [98326] McEvoy, A. N.; Murphy, E. A.; Ponnio, T.; Conneely, O. M.; Bresnihan,
- [98327] B.; FitzGerald, O.; Murphy, E. P.: Activation of nuclear orphan receptor
- [98328] NURR1 transcription by NF-kappaB and cyclic adenosine 5-prime-monophosphate
- [98329] response element-binding protein in rheumatoid arthritis synovial
- [98330] tissue. *J. Immun.* 168: 2979-2987, 2002.
- [98331] 17101.
- [98332] Okabe, T.; Takayanagi, R.; Imasaki, K.; Haji, M.; Nawata, H.; Watanabe,
- [98333] T.: cDNA cloning of a NGFI-B/nur77-related transcription factor from

- [98334] an apoptotic human T cell line. *J. Immun.* 154: 3871–3879, 1995.
- [98335] 17102.
- [98336] Torii, T.; Kawai, T.; Nakamura, S.; Kawakami, H.: Organization
- [98337] of the human orphan nuclear receptor Nurr1 gene. *Gene* 230: 225–232,
- [98338] 1999.
- [98339] 17103.
- [98340] Xu, P.-Y.; Liang, R.; Jankovic, J.; Hunter, C.; Zeng, Y.-X.; Ashizawa,
- [98341] T.; Lai, D.; Le, W.-D.: Association of homozygous 7048G7049 variant
- [98342] in the intron six of Nurr1 gene with Parkinson's disease. *Neurology* 58:
- [98343] 881–884, 2002.
- [98344] 17104.
- [98345] Hanahan, D.: Signaling vascular morphogenesis and maintenance. *Science* 277:
- [98346] 48–50, 1997.
- [98347] 17105.

- [98348] Wong, S. M. E.; Lowe, A. W.: Sequence of the cDNA encoding human
- [98349] GP-2, the major membrane protein in the secretory granule of the exocrine
- [98350] pancreas. *Gene* 171: 311–312, 1996.
- [98351] 17106.
- [98352] Miyoshi, J.; Higashi, T.; Mukai, H.; Ohuchi, T.; Kakunaga, T.:
- [98353] Structure and transforming potential of the human cot oncogene encoding
- [98354] a putative protein kinase. *Molec. Cell. Biol.* 11: 4088–4096, 1991.
- [98355] 17107.
- [98356] Ohara, R.; Miyoshi, J.; Aoki, M.; Toyoshima, K.: The murine cot
- [98357] proto-oncogene: genome structure and tissue-specific expression. *Jpn.*
- [98358] *J. Cancer Res.* 84: 518–525, 1993.
- [98359] 17108.
- [98360] Ishihara, H.; Shibasaki, Y.; Kizuki, N.; Wada, T.; Yazaki, Y.;
- [98361] Asano, T.; Oka, Y.: Type I phosphatidylinositol–

4-phosphate 5-kinases:

- [98362] cloning of the third isoform and deletion/substitution analysis of
- [98363] members of this novel lipid kinase family. *J. Biol. Chem.* 273: 8741-8748,
- [98364] 1998.
- [98365] 17109.
- [98366] Law, D. J.; Tarle, S. A.; Merchant, J. L.: The human ZBP-89 homolog,
- [98367] located at chromosome 3q21, represses gastrin gene expression. *Mammalian*
- [98368] *Genome* 9: 165-167, 1998.
- [98369] 17110.
- [98370] Merchant, J. L.; Iyer, G. R.; Taylor, B. R.; Kitchen, J. R.; Mortensen,
- [98371] E. R.; Wang, Z.; Flintoft, R. J.; Michel, J. B.; Bassel-Duby, R.:
- [98372] ZBP-89, a Kruppel-like zinc finger protein, inhibits epidermal growth
- [98373] factor induction of the gastrin promoter. *Molec. Cell. Biol.* 16:
- [98374] 6644-6653, 1996.

[98375] 17111.

[98376] Wang, Y.; Kobori, J. A.; Hood, L.: The ht-beta gene encodes a

[98377] novel CACCC box-binding protein that regulates T-cell receptor gene

[98378] expression. Molec. Cell. Biol. 13: 5691-5701, 1993.

[98379] 17112.

[98380] Everett, R. D.; Meredith, M.; Orr, A.; Cross, A.; Kathoria, M.;

[98381] Parkinson, J.: A novel ubiquitin-specific protease is dynamically

[98382] associated with the PML nuclear domain and binds to a herpesvirus

[98383] regulatory protein. EMBO J. 16: 1519-1530, 1997. Note: Corrected

[98384] republication of EMBO J.:16: 566-577, 1997.

[98385] 17113.

[98386] Li, M.; Chen, D.; Shiloh, A.; Luo, J.; Nikolaev, A. Y.; Qin, J.;

[98387] Gu, W.: Deubiquitination of p53 by HAUSP is an important pathway

[98388] for p53 stabilization. Nature 416: 648-653, 2002.

- [98389] 17114.
- [98390] Robinson, P. A.; Lomonte, P.; Leek, J. P.; Markham, A. F.;
Everett,
- [98391] R. D.: Assignment of herpesvirus-associated ubiquitin-specific protease
- [98392] gene HAUSP to human chromosome band 16p13.3 by in situ hybridization. Cytogenet.
- [98393] Cell Genet. 83: 100 only, 1998.
- [98394] 17115.
- [98395] Zhou, G.; Bao, Z. Q.; Dixon, J. E.: Components of a new human
- [98396] protein kinase signal transduction pathway. J. Biol. Chem. 270:
- [98397] 12665-12669, 1995.
- [98398] 17116.
- [98399] Lee, J.-D.; Ulevitch, R. J.; Han, J.: Primary structure of BMK1:
- [98400] a new mammalian MAP kinase. Biochem. Biophys. Res. Commun. 213:
- [98401] 715-724, 1995.
- [98402] 17117.

- [98403] Purandare, S. M.; Lee, J.-D.; Patel, P. I.: Assignment of big
- [98404] MAP kinase (PRKM7) to human chromosome 17 band
p11.2 with somatic
- [98405] cell hybrids. Cytogenet. Cell Genet. 83: 258–259, 1998.
- [98406] 17118.
- [98407] Regan, C. P.; Li, W.; Boucher, D. M.; Spatz, S.; Su, M. S.;
Kuida,
- [98408] K.: Erk5 null mice display multiple extraembryonic vascu-
lar and embryonic
- [98409] cardiovascular defects. Proc. Nat. Acad. Sci. 99:
9248–9253, 2002.
- [98410] 17119.
- [98411] Brennan, T. M. H.; Landau, D.; Shalev, H.; Lamb, F.;
Schutte, B.
- [98412] C.; Walder, R. Y.; Mark, A. L.; Carmi, R.; Sheffield, V. C.:
Linkage
- [98413] of infantile Bartter syndrome with sensorineural deafness
to chromosome
- [98414] 1p. Am. J. Hum. Genet. 62: 355–361, 1998.
- [98415] 17120.
- [98416] Birkenhager, R.; Otto, E.; Schurmann, M. J.; Vollmer, M.;

Ruf,

[98417] E.-M.; Maier-Lutz, I.; Beekmann, F.; Fekete, A.; Omran, H.;
Feldmann,

[98418] D.; Milford, D. V.; Jeck, N.; Konrad, M.; Landau, D.; Knoers,
N. V.

[98419] A. M.; Antignac, C.; Sudbrak, R.; Kispert, A.; Hildebrandt,
F.: Mutation

[98420] of BSND causes Bartter syndrome with sensorineural deaf-
ness and kidney

[98421] failure. *Nature Genet.* 29: 310–314, 2001.

[98422] 17121.

[98423] Barlow, G. M.; Chen, X.-N.; Shi, Z. Y.; Lyons, G. E.; Kurnit,
D.

[98424] M.; Celle, L.; Spinner, N. B.; Zackai, E.; Pettenati, M. J.; Van
Riper,

[98425] A. J.; Vekemans, M. J.; Mjaatvedt, C. H.; Korenberg, J. R.:
Down

[98426] syndrome congenital heart disease: a narrowed region and
a candidate

[98427] gene. *Genet. Med.* 3: 91–101, 2001.

[98428] 17122.

[98429] Schmucker, D.; Clemens, J. C.; Shu, H.; Worby, C. A.; Xiao,

J.;

[98430] Muda, M.; Dixon, J. E.; Zipursky, S. L.: *Drosophila Dscam is an axon*

[98431] *guidance receptor exhibiting extraordinary molecular diversity. Cell* 101:

[98432] 671–684, 2000.

[98433] 17123.

[98434] Yamakawa, K.; Huo, Y.-K.; Haendel, M. A.; Hubert, R.; Chen, X.-N.;

[98435] Lyons, G. E.; Korenberg, J. R.: *DSCAM: a novel member of the immunoglobulin*

[98436] *superfamily maps in a Down syndrome region and is involved in the*

[98437] *development of the nervous system. Hum. Molec. Genet.* 7: 227–237,

[98438] 1998.

[98439] 17124.

[98440] Rowles, J.; Scherer, S. W.; Xi, T.; Majer, M.; Nickle, D. C.; Rommens,

[98441] J. M.; Popov, K. M.; Harris, R. A.; Riebow, N. L.; Xia, J.; Tsui,

[98442] L.-C.; Bogardus, C.; Prochazka, M.: *Cloning and charac-*

terization

[98443] of PDK4 on 7q21.3 encoding a fourth pyruvate dehydrogenase kinase

[98444] isoenzyme in human. J. Biol. Chem. 271: 22376–22382, 1996.

[98445] 17125.

[98446] Dode, C.; Weil, D.; Levilliers, J.; Crozet, F.; Chaib, H.; Levi-Acobas,

[98447] F.; Guilford, P.; Petit, C.: Sequence characterization of a newly

[98448] identified human alpha-tubulin gene (TUBA2). Genomics 47: 125–130,

[98449] 1998.

[98450] 17126.

[98451] Bertolino, E.; Reimund, B.; Wildt-Perinic, D.; Clerc, R. G.: A

[98452] novel homeobox protein which recognizes a TGT core and functionally

[98453] interferes with a retinoid-responsive motif. J. Biol. Chem. 270:

[98454] 31178–31188, 1995.

[98455] 17127.

- [98456] Bertolino, E.; Wildt, S.; Richards, G.; Clerc, R. G.: Expression
of a novel murine homeobox gene in the developing cerebellar external
granular layer during its proliferation. Dev. Dyn. 205:
410–420,
1996.
- [98459] 17128.
- [98460] 17128.
- [98461] Gripp, K. W.; Edwards, M. C.; Mowat, D.; Meinecke, P.;
Richieri-Costa,
A.; Zackai, E. H.; Elledge, S.; Muenke, M.: Mutations in the
transcription
factor TGIF in holoprosencephaly. (Abstract) Am. J. Hum.
Genet. 63:
A32, 1998.
- [98464] 17129.
- [98465] 17129.
- [98466] Sulik, K. K.; Dehart, D. B.; Rogers, J. M.; Chernoff, N.: Teratogenicity
of low doses of all-trans retinoic acid in presomite mouse
embryos. Teratology 51:
398–403, 1995.

- [98469] 17130.
- [98470] Wotton, D.; Lo, R. S.; Lee, S.; Massague, J.: A Smad transcriptional
- [98471] corepressor. Cell 97: 29–39, 1999.
- [98472] 17131.
- [98473] Zhu, J.; Petersen, S.; Tessarollo, L.; Nussenzweig, A.: Targeted
- [98474] disruption of the Nijmegen breakage syndrome gene NBS1 leads to early
- [98475] embryonic lethality in mice. Curr. Biol. 11: 105–109, 2001.
- [98476] 17132.
- [98477] Cooper, P. R.; Smilnich, N. J.; Day, C. D.; Nowak, N. J.; Reid,
- [98478] L. H.; Pearsall, R. S.; Reece, M.; Prawitt, D.; Landers, J.; Housman,
- [98479] D. E.; Winterpacht, A.; Zabel, B. U.; Pelletier, J.; Weissman, B.
- [98480] E.; Shows, T. B.; Higgins, M. J.: Divergently transcribed overlapping
- [98481] genes expressed in liver and kidney and located in the 11p15.5 imprinted

- [98482] domain. Genomics 49: 38–51, 1998.
- [98483] 17133.
- [98484] Kershaw, D. B.; Beck, S. G.; Wharram, B. L.; Wiggins, J. E.;
Goyal,
- [98485] M.; Thomas, P. E.; Wiggins, R. C.: Molecular cloning and
characterization
- [98486] of human podocalyxin-like protein: orthologous relation-
ship to rabbit
- [98487] PCLP1 and rat podocalyxin. J. Biol. Chem. 272:
15708–15714, 1997.
- [98488] 17134.
- [98489] Kershaw, D. B.; Wiggins, J. E.; Wharram, B. L.; Wiggins, R.
C.
- [98490] : Assignment of the human podocalyxin-like protein
(PODXL) gene to
- [98491] 7q32–q33. Genomics 45: 239–240, 1997.
- [98492] 17135.
- [98493] Chan, K. K.; Tsui, S. K. W.; Lee, S. M. Y.; Luk, S. C. W.;
Liew,
- [98494] C. C.; Fung, K. P.; Waye, M. M. Y.; Lee, C. Y.: Molecular
cloning

- [98495] and characterization of FHL2, a novel LIM domain protein preferentially
- [98496] expressed in human heart. *Gene* 210: 345–350, 1998.
- [98497] 17136.
- [98498] Genini, M.; Schwalbe, P.; Scholl, F. A.; Remppis, A.; Mattei, M.-G.;
- [98499] Schafer, B. W.: Subtractive cloning and characterization of DRAL,
- [98500] a novel LIM-domain protein down-regulated in rhabdomyosarcoma. *DNA*
- [98501] *Cell Biol.* 16: 433–442, 1997.
- [98502] 17137.
- [98503] Prols, F.; Liehr, T.; Rinke, R.; Rautenstrauss, B.: Assignment
- [98504] of the microvascular endothelial differentiation gene 1 (MDG1) to
- [98505] human chromosome band 14q24.2–q24.3 by fluorescence in situ hybridization. *Cytogenet.*
- [98506] *Cell Genet.* 79: 149–150, 1997.
- [98507] 17138.
- [98508] Claverie-Martin, F.; Wang, M.; Cohen, S. N.: ARD-1 cDNA

from human

[98509] cells encodes a site-specific single-strand endoribonuclease that

[98510] functionally resembles Escherichia coli RNase E. J. Biol. Chem. 272:

[98511] 13823-13828, 1997.

[98512] 17139.

[98513] Van Eynde, A.; Wera, S.; Beullens, M.; Torrekens, S.; Van Leuven,

[98514] F.; Stalmans, W.; Bollen, M.: Molecular cloning of NIPP-1, a nuclear

[98515] inhibitor of protein phosphatase-1, reveals homology with polypeptides

[98516] involved in RNA processing. J. Biol. Chem. 270: 28068-28074, 1995.

[98517] 17140.

[98518] Tekin, M.; Dogu, F.; Tacyildiz, N.; Akar, E.; Ikinciogullari,

[98519] A.; Ogur, G.; Yavuz, G.; Babacan, E.; Akar, N.: 657del5 mutation

[98520] in the NBS1 gene is associated with Nijmegen breakage syndrome in

[98521] a Turkish family. Clin. Genet. 62: 84-88, 2002.

[98522] 17141.

[98523] Varon, R.; Reis, A.; Henze, G.; Einsiedel, H. G.; Sperling, K.;

[98524] Seeger, K.: Mutations in the Nijmegen breakage syndrome
gene (NBS1)

[98525] in childhood acute lymphoblastic leukemia (ALL). Cancer
Res. 61:

[98526] 3570–3572, 2001.

[98527] 17142.

[98528] Wilda, M.; Demuth, I.; Concannon, P.; Sperling, K.;
Hameister,

[98529] H.: Expression pattern of the Nijmegen breakage syn-
drome gene, Nbs1,

[98530] during murine development. Hum. Molec. Genet. 9:
1739–1744, 2000.

[98531] 17143.

[98532] Wu, X.; Ranganathan, V.; Weisman, D. S.; Heine, W. F.; Cic-
cone,

[98533] D. N.; O'Neill, T. B.; Crick, K. E.; Pierce, K. A.; Lane, W. S.;
Rathbun,

[98534] G.; Livingston, D. M.; Weaver, D. T.: ATM phosphorylation
of Nijmegen

- [98535] breakage syndrome protein is required in a DNA damage response. Nature 405:
- [98536] 477–482, 2000.
- [98537] 17144.
- [98538] Zhao, S.; Weng, Y.–C.; Yuan, S.–S. F.; Lin, Y.–T.; Hsu, H.–C.;
- [98539] Lin, S.–C. J.; Gerbino, E.; Song, M.; Zdzenicka, M. Z.; Gatti, R.
- [98540] A.; Shay, J. W.; Ziv, Y.; Shiloh, Y.; Lee, E. Y.–H. P.: Functional
- [98541] link between ataxia–telangiectasia and Nijmegen breakage syndrome
- [98542] gene products. Nature 405: 473–477, 2000.
- [98543] 17145.
- [98544] Mankodi, A.; Urbinati, C. R.; Yuan, Q.–P.; Moxley, R. T.; Sansone,
- [98545] V.; Krym, M.; Henderson, D.; Schalling, M.; Swanson, M. S.; Thornton,
- [98546] C. A.: Muscleblind localizes to nuclear foci of aberrant RNA in myotonic
- [98547] dystrophy types 1 and 2. Hum. Molec. Genet. 10: 2165–2170, 2001.

- [98548] 17146.
- [98549] Hittner, H. M.; Kretzer, F. L.; Antoszyk, J. H.; Ferrell, R. E.;
- [98550] Mehta, R. S.: Variable expressivity of autosomal dominant
anterior
- [98551] segment mesenchymal dysgenesis in six generations. Am.
J. Ophthalm. 93:
- [98552] 57–70, 1982.
- [98553] 17147.
- [98554] Semina, E. V.; Ferrell, R. E.; Mintz–Hittner, H. A.; Bitoun, P.;
- [98555] Alward, W. L. M.; Reiter, R. S.; Funkhauser, C.; Daack–
Hirsch, S.;
- [98556] Murray, J. C.: A novel homeobox gene PITX3 is mutated in
families
- [98557] with autosomal–dominant cataracts and ASMD. Nature
Genet. 19: 167–170,
- [98558] 1998.
- [98559] 17148.
- [98560] Semina, E. V.; Murray, J. C.; Reiter, R.; Hrstka, R. F.; Graw,
[98561] J.: Deletion in the promoter region and altered expression
of Pitx3

- [98562] homeobox gene in aphakia mice. Hum. Molec. Genet. 9: 1575–1585, 2000.
- [98563] 17149.
- [98564] Semina, E. V.; Reiter, R. S.; Murray, J. C.: Isolation of a new homeobox gene belonging to the Pitx/Rieg family: expression during lens development and mapping to the aphakia region on mouse chromosome 19. Hum. Molec. Genet. 6: 2109–2116, 1997. 17150.
- [98569] Lin, C.-S.; Shen, W.; Chen, Z. P.; Tu, Y.-H.; Matsudaira, P.: Identification of I-plastin, a human fimbrin isoform expressed in intestine and kidney. Molec. Cell. Biol. 14: 2457–2467, 1994. 17151.
- [98573] Kent, J.; Lee, M.; Schedl, A.; Boyle, S.; Fantes, J.; Powell, M.; Rushmere, N.; Abbott, C.; van Heyningen, V.; Bickmore, W. A.: The

- [98576] reticulocalbin gene maps to the WAGR region in human and to the Small
- [98577] eye Harwell deletion in mouse. *Genomics* 42: 260–267, 1997.
- [98578] 17152.
- [98579] Ozawa, M.: Cloning of a human homologue of mouse reticulocalbin
- [98580] reveals conservation of structural domains in the novel endoplasmic
- [98581] reticulum resident Ca^{2+} -binding protein with multiple EF-hand motifs. *J.*
- [98582] *Biochem.* 117: 1113–1119, 1995.
- [98583] 17153.
- [98584] Ozawa, M.; Muramatsu, T.: Reticulocalbin, a novel endoplasmic
- [98585] reticulum resident Ca^{2+} -binding protein with multiple EF-hand motifs
- [98586] and a carboxyl-terminal HDEL sequence. *J. Biol. Chem.* 268: 699–705,
- [98587] 1993.
- [98588] 17154.

- [98589] Yan, W. L.; Lerner, T. J.; Haines, J. L.; Gusella, J. F.: Sequence
[98590] analysis and mapping of a novel human mitochondrial
ATP synthase subunit
[98591] 9 cDNA (ATP5G3). *Genomics* 24: 375–377, 1994.
[98592] 17155.
[98593] Aho, S.; McLean, W. H. I.; Li, K.; Uitto, J.: cDNA cloning,
mRNA
[98594] expression, and chromosomal mapping of human and
mouse periplakin
[98595] genes. *Genomics* 48: 242–247, 1998.
[98596] 17156.
[98597] Luo, G.; Leroy, E.; Kozak, C. A.; Polymeropoulos, M. H.;
Horowitz,
[98598] R.: Mapping of the gene (NRAP) encoding N-RAP in the
mouse and human
[98599] genomes. *Genomics* 45: 229–232, 1997.
[98600] 17157.
[98601] Luo, G.; Zhang, J. Q.; Nguyen, T.-P.; Herrera, A. H.; Pater-
son,
[98602] B.; Horowitz, R.: Complete cDNA sequence and tissue lo-

calization

[98603] of N-RAP, a novel nebulin-related protein of striated muscle. Cell

[98604] Motil. Cytoskeleton 38: 75-90, 1997.

[98605] 17158.

[98606] Ando-Akatsuka, Y.; Saitou, M.; Hirase, T.; Kishi, M.; Sakakibara,

[98607] A.; Itoh, M.; Yonemura, S.; Furuse, M.; Tsukita, S.: Inter-species

[98608] diversity of the occludin sequence: cDNA cloning of human, mouse,

[98609] dog, and rat-kangaroo homologues. J. Cell Biol. 133: 43-47, 1996.

[98610] 17159.

[98611] Strahm, B.; Rittweiler, K.; Duffner, U.; Brandau, O.; Orłowska-Volk,

[98612] M.; Karajannis, M. A.; zur Stadt, U.; Tiemann, M.; Reiter, A.; Brandis,

[98613] M.; Meindl, A.; Niemeyer, C. M.: Recurrent B-cell non-Hodgkin's lymphoma

[98614] in two brothers with X-linked lymphoproliferative disease without

- [98615] evidence for Epstein-Barr virus infection. *Brit. J. Haemat.* 108:
- [98616] 377-382, 2000.
- [98617] 17160.
- [98618] Sylla, B. S.; Murphy, K.; Cahir-McFarland, E.; Lane, W. S.; Mosialos,
- [98619] G.; Kieff, E.: The X-linked lymphoproliferative syndrome gene product
- [98620] SH2D1A associates with p62(dok) (Dok1) and activates NF-kappa-beta. *Proc.*
- [98621] *Nat. Acad. Sci.* 97: 7470-7475, 2000.
- [98622] 17161.
- [98623] Sylla, B. S.; Wang, Q.; Hayoz, D.; Lathrop, G. M.; Lenoir, G.
- [98624] M.: Multipoint linkage mapping of the Xq25-q26 region in a family
- [98625] affected by the X-linked lymphoproliferative syndrome. *Clin. Genet.* 36:
- [98626] 359-462, 1989.
- [98627] 17162.
- [98628] Tangye, S. G.; Lazetic, S.; Woollatt, E.; Sutherland, G. R.; Lanier,

- [98629] L. L.; Phillips, J. H.: Cutting edge: human 2B4, an activating NK
- [98630] cell receptor, recruits the protein tyrosine phosphatase SHP-2 and
- [98631] the adaptor signaling protein SAP. J. Immun. 162: 6981-6985, 1999.
- [98632] 17163.
- [98633] Tangye, S. G.; Phillips, J. H.; Lanier, L. L.; Nichols, K. E.
- [98634] : Cutting edge: functional requirement for SAP in 2B4-mediated activation
- [98635] of human natural killer cells as revealed by the X-linked lymphoproliferative
- [98636] syndrome. J. Immun. 165: 2932-2936, 2000.
- [98637] 17164.
- [98638] Thorley-Lawson, D. A.; Schooley, R. T.; Bhan, A. K.; Nadler, L.
- [98639] M.: Epstein-Barr virus superinduces a new human B cell differentiation
- [98640] antigen (B-LAST 1) expressed on transformed lymphoblasts. Cell 30:
- [98641] 415-425, 1982.

- [98642] 17165.
- [98643] Vowels, M. R.; Lam-Po-Tang, R.; Berdoukas, V.; Ford, D.;
Thierry,
- [98644] D.; Purtilo, D.; Gluckman, E.: Correction of X-linked lym-
phoproliferative
- [98645] disease by transplantation of cord-blood stem cells. New
Eng. J.
- [98646] Med. 329: 1623-1625, 1993.
- [98647] 17166.
- [98648] Williams, L. L.; Rooney, C. M.; Conley, M. E.; Brenner, M.
K.;
- [98649] Krance, R. A.; Heslop, H. E.: Correction of Duncan's syn-
drome by
- [98650] allogeneic bone marrow transplantation. Lancet 342:
587-588, 1993.
- [98651] 17167.
- [98652] Wu, C.; Nguyen, K. B.; Pien, G. C.; Wang, N.; Gullo, C.;
Duncan,
- [98653] H.; Sosa, M. R.; Edwards, M. J.; Borrow, P.; Satoskar, A. R.;
Sharpe,
- [98654] A. H.; Biron, C. A.; Terhorst, C.: SAP controls T cell re-
sponses

- [98655] to virus and terminal differentiation of T(H)2 cells. *Nature Immun.* 2:
410–414, 2001.
- [98656] 17168.
- [98657] Wyandt, H. E.; Skare, J. C.; Grierson, H. L.; Purtilo, D. T.;
- [98658] Milunsky, A.: Detection of a chromosomal deletion of Xq25 in an affected male with X-linked lymphoproliferative disease. (Abstract) *Am. J. Hum. Genet.* 45 (suppl.): A108 only, 1989.
- [98659] 17169.
- [98660] Yin, L.; Ferrand, V.; Lavoue, M.-F.; Hayoz, D.; Philippe, N.;
- [98661] Souillet, G.; Seri, M.; Giacchino, R.; Castagnola, E.; Hodgson, S.;
- [98662] Sylla, B. S.; Romeo, G.: SH2D1A mutation analysis for diagnosis of XLP in typical and atypical patients. *Hum. Genet.* 105: 501–505, 1999.
- [98663] 17170.

[98669] Yin, L.; Tocco, T.; Pauly, S.; Lenoir, G. M.; Romeo, G.: Absence

[98670] of SH2D1A point mutation in 62 Burkitts lymphoma cell lines. Am.

[98671] J. Hum. Genet. 65 (suppl. 1868): A331 only, 1999.

[98672] 17171.

[98673] Berglund, E. O.; Ranscht, B.: Molecular cloning and in situ localization

[98674] of the human contactin gene (CNTN1) on chromosome 12q11-q12. Genomics 21:

[98675] 571-582, 1994.

[98676] 17172.

[98677] Laxminarayan, K. M.; Chan, B. K.; Tetaz, T.; Bird, P. I.; Mitchell,

[98678] C. A.: Characterization of a cDNA encoding the 43-kDa membrane-associated

[98679] inositol-polyphosphate 5-phosphatase. J. Biol. Chem. 269: 17305-17310,

[98680] 1994.

[98681] 17173.

[98682] Fink, J. K.; Jones, S. M.; Esposito, C.; Wilkowski, J.: Human

- [98683] microtubule-associated protein 1a (MAP1A) gene: genomic organization,
- [98684] cDNA sequence, and developmental-and tissue-specific expression. *Genomics* 35:
- [98685] 577-585, 1996.
- [98686] 17174.
- [98687] Quinton, R.; Duke, V. M.; de Zoysa, P. A.; Platts, A. D.; Valentine,
- [98688] A.; Kendall, B.; Pickman, S.; Kirk, J. M. W.; Besser, G. M.; Jacobs,
- [98689] H. S.; Bouloux, P. M. G.: The neuroradiology of Kallmann's syndrome:
- [98690] a genotypic and phenotypic analysis. *J. Clin. Endocr. Metab.* 81:
- [98691] 3010-3017, 1996.
- [98692] 17175.
- [98693] Rogol, A.; Mittal, K.; White, B. J.; McGinnis, M. H.; Liebllich,
- [98694] J. M.; Rosen, S. W.: HLA-compatible paternity in two 'fertile eunuchs'
- [98695] with congenital hypogonadotrophic hypogonadism and anosmia (the Kallmann

- [98696] syndrome). J. Clin. Endocr. Metab. 51: 275–279, 1980.
- [98697] 17176.
- [98698] Befort, K.; Filliol, D.; Decaillot, F. M.; Gaveriaux–Ruff, C.;
- [98699] Hoehe, M. R; Kieffer, B. L.: A single nucleotide polymorphic mutation
- [98700] in the human mu–opioid receptor severely impairs receptor signaling. J.
- [98701] Biol. Chem. 276: 3130–3137, 2001.
- [98702] 17177.
- [98703] Bond, C.; LaForge, K. S.; Tian, M.; Melia, D.; Zhang, S.; Borg,
- [98704] L.; Gong, J.; Schluger, J.; Strong, J. A.; Leal, S. M.; Tischfield,
- [98705] J. A.; Kreek, M. J.; Yu, L.: Single–nucleotide polymorphism in the
- [98706] human mu opioid receptor gene alters beta–endorphin binding and activity:
- [98707] possible implications for opiate addiction. Proc. Nat. Acad. Sci. 95:
- [98708] 9608–9613, 1998.
- [98709] 17178.

- [98710] Chen, Y.; Mestek, A.; Liu, J.; Hurley, J. A.; Yu, L.: Molecular
[98711] cloning and functional expression of a mu-opioid recep-
tor from rat
[98712] brain. *Molec. Pharm.* 44: 8–12, 1993.
[98713] 17179.
- [98714] He, L.; Fong, J.; von Zastrow, M.; Whistler, J. L.: Regulation
[98715] of opioid receptor trafficking and morphine tolerance by
receptor
[98716] oligomerization. *Cell* 108: 271–282, 2002.
[98717] 17180.
- [98718] Hoehe, M. R.; Kopke, K.; Wendel, B.; Rohde, K.; Flach-
meier, C.;
[98719] Kidd, K. K.; Berrettini, W. H.; Church, G. M.: Sequence vari-
ability
[98720] and candidate gene analysis in complex disease: associa-
tion of mu
[98721] opioid receptor gene variation with substance depen-
dence. *Hum. Molec.*
[98722] *Genet.* 9: 2895–2908, 2000.
[98723] 17181.

- [98724] Holtt, V.: A polymorphism (A118G) in the mu-opioid receptor gene
- [98725] affects the response to morphine-6-glucuronide in humans. Pharmacogenetics 12:
- [98726] 1-2, 2002.
- [98727] 17182.
- [98728] LaForge, K. S.; Yuferov, V.; Kreek, M. J.: Opioid receptor and
- [98729] peptide gene polymorphisms: potential implications for addictions. Europ.
- [98730] J. Pharm. 410: 249-268, 2000.
- [98731] 17183.
- [98732] Lotsch, J.; Skarke, C.; Grosch, S.; Darimont, J.; Schmidt, H.;
- [98733] Geisslinger, G.: The polymorphism A118G of the human mu-opioid receptor
- [98734] gene decreases the pupil constrictory effect of morphine-6-glucuronide
- [98735] but not that of morphine. Pharmacogenetics 12: 3-9, 2002.
- [98736] 17184.
- [98737] Mansour, A.; Fox, C. A.; Burke, S.; Akil, H.; Watson, S. J.:

- [98738] Immunohistochemical localization of the clone mu opioid receptor in
- [98739] the rat CNS. *J. Chem. Neuroanat.* 8: 283–305, 1995.
- [98740] 17185.
- [98741] Meyer, U. A.: Pharmacogenetics and adverse drug reactions. *Lancet* 356:
- [98742] 1667–1671, 2000.
- [98743] 17186.
- [98744] Mogil, J. S.: The genetic mediation of individual differences
- [98745] in sensitivity to pain and its inhibition. *Proc. Nat. Acad. Sci.* 96:
- [98746] 7744–7751, 1999.
- [98747] 17187.
- [98748] Uhl, G. R.; Sora, I.; Wang, Z.: The mu opiate receptor as a candidate
- [98749] gene for pain: polymorphisms, variations in expression, nociception,
- [98750] and opiate responses. *Proc. Nat. Acad. Sci.* 96:
- [98751] 7752–7755, 1999.
- [98751] 17188.

- [98752] Wang, J.-B.; Johnson, P. S.; Persico, A. M.; Hawkins, A. L.; Griffin,
[98753] C. A.; Uhl, G. R.: Human mu opiate receptor: cDNA and
genomic clones,
[98754] pharmacologic characterization and chromosomal assign-
ment. FEBS Lett. 338:
[98755] 217-222, 1994.
[98756] 17189.
- [98757] Wu, D.; Kang, Y.-S.; Bickel, U.; Pardridge, W. M.: Blood-
brain
[98758] barrier permeability to morphine-6-glucuronide is
markedly reduced
[98759] compared with morphine. Drug Metab. Dispos. 25:
768-771, 1997.
[98760] 17190.
- [98761] Zubietta, J.-K.; Smith, Y. R.; Bueller, J. A.; Xu, Y.; Kilbourn,
[98762] M. R.; Jewett, D. M.; Meyer, C. R.; Koeppe, R. A.; Stohler, C.
S.
[98763] : Regional mu opioid receptor regulation of sensory and
affective
[98764] dimensions of pain. Science 293: 311-315, 2001.

[98765] 17191.

[98766] McBride, O. W.; Yi, H. F.; Srivastava, M.: The human cytochrome

[98767] b561 gene (CYB561) is located at 17q11-qter. Genomics 21: 662-663,

[98768] 1994.

[98769] 17192.

[98770] Srivastava, M.: Genomic structure and expression of the human

[98771] gene encoding cytochrome b(561), an integral protein of the cromaffin

[98772] granule membrane. J. Biol. Chem. 270: 22714-22720, 1995.

[98773] 17193.

[98774] Albarosa, R.; DiDonato, S.; Finocchiaro, G.: Redefinition of the

[98775] coding sequence of the MXI1 gene and identification of a polymorphic

[98776] repeat in the 3-prime non-coding region that allows the detection

[98777] of loss of heterozygosity of chromosome 10q25 in glioblastomas. Hum.

- [98778] Genet. 95: 709–711, 1995.
- [98779] 17194.
- [98780] Mitchell, C. A.; Speed, C. J.; Nicholl, J.; Sutherland, G. R.:
- [98781] Chromosomal mapping of the gene (INPP5A) encoding the
43-kDa membrane-associated
- [98782] inositol polyphosphate 5-phosphatase to 10q26.3 by flu-
orescence in
- [98783] situ hybridization. Genomics 31: 139–140, 1996.
- [98784] 17195.
- [98785] Phillips, D. R.; Agin, R. P.: Platelet membrane defects in
Glanzmann's
- [98786] thrombasthenia: evidence for decreased amounts of two
major glycoproteins. J.
- [98787] Clin. Invest. 60: 535–545, 1977.
- [98788] 17196.
- [98789] Pittman, M. A., Jr.; Graham, J. B.: Glanzmann's thrombopa-
thy:
- [98790] an autosomal recessive trait in one family. Am. J. Med. Sci.
247:
- [98791] 293–303, 1964.

- [98792] 17197.
- [98793] Poncz, M.; Rifat, S.; Collier, B. S.; Newman, P. J.; Shattil, S. J.; Parrella, T.; Fortina, P.; Bennett, J. S.: Glanzmann thrombasthenia secondary to a gly273-to-aspartate mutation adjacent to the first calcium-binding domain of platelet glycoprotein IIb. *J. Clin. Invest.* 93: 172-179, 1994.
- [98797] 17198.
- [98798] 17198.
- [98799] Prandini, M. H.; Denarier, E.; Frachet, P.; Uzan, G.; Marguerie, G.: Isolation of the human platelet glycoprotein IIb gene and characterization of the 5-prime flanking region. *Biochem. Biophys. Res. Commun.* 156: 595-601, 1988.
- [98802] 595-601, 1988.
- [98803] 17199.
- [98804] Pytela, R.; Pierschbacher, M. D.; Ginsberg, M. H.; Plow, E. F.;
- [98805] Ruoslahti, E.: Platelet membrane glycoprotein IIb/IIIa:

member of

[98806] a family of arg-gly-asp-specific adhesion receptors. Science 231:

[98807] 1559-1562, 1986.

[98808] 17200.

[98809] Ruan, J.; Peyruchaud, O.; Alberio, L.; Valles, G.; Clemetson,

[98810] K.; Bourre, F.; Nurden, A. T.: Double heterozygosity of the GPIIb

[98811] gene in a Swiss patient with Glanzmann's thrombasthenia. Brit. J.

[98812] Haemat. 102: 918-925, 1998.

[98813] 17201.

[98814] Russell, M. E.; Seligsohn, U.; Coller, B. S.; Ginsberg, M. H.;

[98815] Skoglund, P.; Quertermous, T.: Structural integrity of the glycoprotein

[98816] IIb and IIIa genes in Glanzmann thrombasthenia patients from Israel. Blood 72:

[98817] 1833-1836, 1988.

[98818] 17202.

[98819] Schlegel, N.; Gayet, O.; Morel-Kopp, M.-C.; Wyler, B.; Hur-

taud-Roux,

- [98820] M.-F.; Kaplan, C.; McGregor, J.: The molecular genetic basis of Glanzmann's
- [98821] thrombasthenia in a gypsy population in France: identification of
- [98822] a new mutation on the alpha(IIB) gene. Blood 86: 977-982, 1995.
- [98823] 17203.
- [98824] Seligsohn, U.; Mibashan, R. S.; Rodeck, C. H.; Nicolaides, K.
- [98825] H.; Millar, D. S.; Coller, B. S.: Prenatal diagnosis of Glanzmann's
- [98826] thrombasthenia. (Letter) Lancet II: 1419 only, 1985.
- [98827] 17204.
- [98828] Sosnoski, D.; Emanuel, B. S.; Hawkins, A. L.; van Tuinen, P.;
- [98829] Ledbetter, D. H.; Nussbaum, R. L.; Kaos, F.-T.; Schwartz, E.; Phillips,
- [98830] D.; Bennett, J. S.; Fitzgerald, L. A.; Poncz, M.: Chromosomal localization
- [98831] of the genes for the vitronectin and fibronectin receptors alpha-subunits

- [98832] and for platelet glycoproteins IIb and IIIa. J. Clin. Invest. 81:
- [98833] 1993–1998, 1988.
- [98834] 17205.
- [98835] Tongio, M. M.; Lutz, P.; Hauptmann, G.; Rodier, L.; Levy, J.-M.;
- [98836] Mayer, S.; Cazenave, J.-P.: Type I Glanzmann's thrombasthenia segregates
- [98837] independently of Ss and Duffy systems and the A, B, C, factor B, C2
- [98838] and C4 loci of the HLA complex. Tissue Antigens 20: 22–27, 1982.
- [98839] 17206.
- [98840] Uzan, G.; Van Cong, N.; Gross, M. S.; Jegou-Foubert, C.; Frachet,
- [98841] P.; Boucheix, C.; Margerie, G.; Frezal, J.: Assignment of a gene
- [98842] coding for human platelet GPIIb (GP2B) to 17q21.1–q21.3 by in situ
- [98843] hybridization. (Abstract) Cytogenet. Cell Genet. 46: 707 only, 1987.
- [98844] 17207.

- [98845] Van Cong, N.; Uzan, G.; Gross, M. S.; Jegou-Foubert, C.; Frchet,
- [98846] P.; Boucheix, C.; Marguerie, G.; Frezal, J.: Assignment of human
- [98847] platelet GP2B (GPIIb) gene to chromosome 17, region q21.1–q21.3. Hum.
- [98848] Genet. 80: 389–392, 1988.
- [98849] 17208.
- [98850] von dem Borne, A. E. G. K.; von Riesz, E.; Verheugt, F. W. A.;
- [98851] ten Cate, J. W.; Koppe, J. G.; Engelfriet, C. P.; Nijenhuis, L. E.
- [98852] : Bak(a), a new platelet-specific antigen involved in neonatal allo-immune
- [98853] thrombocytopenia. Vox Sang. 39: 113–120, 1980.
- [98854] 17209.
- [98855] Waller, H. D.; Gross, R.: Genetische Enzymdefecte als Ursache
- [98856] von Thrombocytopathien. Verh. Dtsch. Ges. Inn. Med. 70: 476–494,
- [98857] 1964.

[98858] 17210.

- [98859] Wilcox, D. A.; Gill, J.; Newman, P. J.: Glanzmann thrombasthenia
- [98860] resulting from a single amino acid substitution flanking the fibrinogen
- [98861] gamma-chain dodecapeptide binding domain on GPIIb. (Abstract) Blood 82:
- [98862] 210a only, 1993.
- [98863] 17211.
- [98864] Wilcox, D. A.; Paddock, C. M.; Lyman, S.; Gill, J. C.; Newman,
- [98865] P. J.: Glanzmann thrombasthenia resulting from a single amino acid
- [98866] substitution between the second and third calcium-binding domains
- [98867] of GPIIb: role of the GPIIb amino terminus in integrin subunit association. J.
- [98868] Clin. Invest. 95: 1553 only, 1995.
- [98869] 17212.
- [98870] Wilcox, D. A.; Wautier, J. L.; Pidard, D.; Newman, P. J.: A single
- [98871] amino acid substitution flanking the fourth calcium binding domain

- [98872] of alpha-IIb prevents maturation of the alpha-IIb-beta-3 integrin
- [98873] complex. J. Biol. Chem. 269: 4450-4457, 1994.
- [98874] 17213.
- [98875] Zaizov, R.; Cohen, I.; Matoth, Y.: Thrombasthenia: a study of
- [98876] two siblings. Acta Paediat. Scand. 57: 522-526, 1968.
- [98877] 17214.
- [98878] Dreyer, B.; Tranebjaerg, L.; Rosenberg, T.; Weston, M. D.; Kimberling,
- [98879] W. J.; Nilssen, O.: Identification of novel USH2A mutations: implications
- [98880] for the structure of USH2A protein. Europ. J. Hum. Genet. 8: 500-506,
- [98881] 2000.
- [98882] 17215.
- [98883] Eudy, J. D.; Weston, M. D.; Yao, S.; Hoover, D. M.; Rehm, H. L.;
- [98884] Ma-Edmonds, M.; Yan, D.; Ahmad, I.; Cheng, J. J.; Ayuso, C.; Cremers,
- [98885] C.; Davenport, S.; Moller, C.; Talmadge, C. B.; Beisel, K. W.;

Tamayo,

[98886] M.; Morton, C. C.; Swaroop, A.; Kimberling, W. J.; Sumegi, J.: Mutation

[98887] of a gene encoding a protein with extracellular matrix motifs in Usher

[98888] syndrome type IIa. *Science* 280: 1753–1757, 1998.

[98889] 17216.

[98890] Huang, D.; Eudy, J. D.; Uzvolgyi, E.; Davis, J. R.; Talmadge, C.

[98891] B.; Pretto, D.; Weston, M. D.; Lehman, J. E.; Zhou, M.; Seemayer,

[98892] T. A.; Ahmad, I.; Kimberling, W. J.; Sumegi, J.: Identification of

[98893] the mouse and rat orthologs of the gene mutated in Usher syndrome

[98894] type IIA and the cellular source of USH2A mRNA in retina, a target

[98895] tissue of the disease. *Genomics* 80: 195–203, 2002.

[98896] 17217.

[98897] Kimberling, W. J.; Weston, M. D.; Moller, C.; Davenport, S. L.

[98898] H.; Shugart, Y. Y.; Priluck, I. A.; Martini, A.; Milani, M.;

Smith,

[98899] R. J.: Localization of Usher syndrome type II to chromosome 1q. Genomics 7:

[98900] 245–249, 1990.

[98901] 17218.

[98902] Kimberling, W. J.; Weston, M. D.; Moller, C.; van Aarem, A.; Cremers,

[98903] C. W. R. J.; Sumegi, J.; Ing, P. S.; Connolly, C.; Martini, A.; Milani,

[98904] M.; Tamayo, M. L.; Bernal, J.; Greenberg, J.; Ayuso, C.: Gene mapping

[98905] of Usher syndrome type IIa: localization of the gene to a 2.1-cM segment

[98906] on chromosome 1q41. Am. J. Hum. Genet. 56: 216–223, 1995.

[98907] 17219.

[98908] Lewis, R. A.; Otterud, B.; Stauffer, D.; Lalouel, J.-M.; Lerpert,

[98909] M.: Mapping recessive ophthalmic diseases: linkage of the locus for

[98910] Usher syndrome type II to a DNA marker on chromosome 1q. Genomics 7:

- [98911] 250–256, 1990.
- [98912] 17220.
- [98913] Liu, X.–Z.; Hope, C.; Liang, C. Y.; Zou, J. M.; Xu, L. R.; Cole,
[98914] T.; Mueller, R. F.; Bunday, S.; Nance, W.; Steel, K. P.;
Brown, S.
[98915] D. M.: A mutation (2314delG) in the Usher syndrome type
IIA gene:
[98916] high prevalence and phenotypic variation. (Letter) Am. J.
Hum. Genet. 64:
[98917] 1221–1225, 1999.
- [98918] 17221.
- [98919] Pieke–Dahl, S.; van Aarem, A.; Dobin, A.; Cremers, C. W. R.
J.;
- [98920] Kimberling, W. J.: Genetic heterogeneity of Usher syn-
drome type II
[98921] in a Dutch population. J. Med. Genet. 33: 753–757, 1996.
- [98922] 17222.
- [98923] Pieke Dahl, S.; Weston, M. D.; Kimberling, W. J.; Gorin, M.
B.;
- [98924] Shugart, Y. Y.; Kenyon, J. B.: Possible genetic heterogene-
ity of

- [98925] Usher syndrome type 2: a family unlinked to chromosome 1q markers.
- [98926] (Abstract) Am. J. Hum. Genet. 49 (suppl.): 200 only, 1991.
- [98927] 17223.
- [98928] Rivolta, C.; Sweklo, E. A.; Berson, E. L.; Dryja, T. P.: Mis-sense
- [98929] mutation in the USH2A gene: association with recessive retinitis pigmentosa
- [98930] without hearing loss. Am. J. Hum. Genet. 66: 1975–1978, 2000.
- [98931] 17224.
- [98932] Robbins, J. H.; Scudiero, D. A.; Otsuka, F.; Tarone, R. E.; Brumback,
- [98933] R. A.; Wirtschafter, J. D.; Polinsky, R. J.; Barrett, S. F.; Moshell,
- [98934] A. N.; Scarpinato, R. G.; Ganges, M. B.; Nee, L. E.; Meyer, S. A.;
- [98935] Clatterbuck, B. E.: Hypersensitivity to DNA-damaging agents in cultured
- [98936] cells from patients with Usher's syndrome and Duchenne muscular dystrophy. J.
- [98937] Neurol. Neurosurg. Psychiat. 47: 391–398, 1984.

- [98938] 17225.
- [98939] Smith, R. J. H.; Lee, E. C.; Kimberling, W. J.; Daiger, S. P.;
- [98940] Pelias, M. Z.; Keats, B. J. B.; Jay, M.; Bird, A.; Reardon, W.;
Guest,
- [98941] M.; Ayyagari, R.; Hejtmancik, J. F.: Localization of two
genes for
- [98942] Usher syndrome type I to chromosome 11. *Genomics* 14:
995–1002, 1992.
- [98943] 17226.
- [98944] Weston, M. D.; Eudy, J. D.; Fujita, S.; Yao, S.-F.; Usami, S.;
- [98945] Cremers, C.; Greenburg, J.; Ramesar, R.; Martini, A.;
Moller, C.;
- [98946] Smith, R. J.; Sumegi, J.; Kimberling, W. J.: Genomic struc-
ture and
- [98947] identification of novel mutations in usherin, the gene re-
sponsible
- [98948] for Usher syndrome type IIa. *Am. J. Hum. Genet.* 66:
1199–1210, 2000.
- [98949] Note: Erratum: *Am. J. Hum. Genet.* 66: 2020 only, 2000.
- [98950] 17227.
- [98951] Gasparini, P.; De Fazio, A.; Croce, A. I.; Stanziale, P.; Ze-

lante,

[98952] L.: Usher syndrome type III (USH3) linked to chromosome 3q in an

[98953] Italian family. J. Med. Genet. 35: 666–667, 1998.

[98954] 17228.

[98955] Joensuu, T.; Hamalainen, R.; Yuan, B.; Johnson, C.; Tegelberg,

[98956] S.; Gasparini, P.; Zelante, L.; Pirvola, U.; Parakinen, L.; Lehesjoki,

[98957] A.-E.; de la Chapelle, A.; Sankila, E.-M.: Mutations in a novel gene

[98958] with transmembrane domains underlie Usher syndrome type 3. Am. J.

[98959] Hum. Genet. 69: 673–684, 2001. Note: Erratum: Am. J. Hum. Genet.

[98960] 69: 1160 only, 2001.

[98961] 17229.

[98962] Sankila, E.-M.; Pakarinen, L.; Kaariainen, H.; Aittomaki, K.; Karjalainen,

[98963] S.; Sistonen, P.; de la Chapelle, A.: Assignment of an Usher syndrome

[98964] type III (USH3) gene to chromosome 3q. Hum. Molec.

Genet. 4: 93–98,

[98965] 1995.

[98966] 17230.

[98967] Anantharaman, V.; Koonin, E. V.; Aravind, L.: Peptide–
N-glycanases

[98968] and DNA repair proteins, Xp-C/Rad4, are, respectively,
active and

[98969] inactivated enzymes sharing a common transglutaminase
fold. Hum.

[98970] Molec. Genet. 10: 1627–1630, 2001.

[98971] 17231.

[98972] Cheo, D. L.; Burns, D. K.; Meira, L. B.; Houle, J. F.; Fried–
berg,

[98973] E. C.: Mutational inactivation of the xeroderma pigmento–
sum group

[98974] C gene confers predisposition to
2-acetylaminofluorene-induced liver

[98975] and lung cancer and to spontaneous testicular cancer in
Trp53 –/–

[98976] mice. Cancer Res. 59: 771–775, 1999.

[98977] 17232.

- [98978] Cleaver, J. E.: DNA repair in human xeroderma pigmentosum group
- [98979] C cells involves a different distribution of damaged sites in confluent
- [98980] and growing cells. Nucleic Acids Res. 14: 8155–8165, 1986.
- [98981] 17233.
- [98982] Emmert, S.; Kobayashi, N.; Khan, S. G.; Kraemer, K. H.: The xeroderma
- [98983] pigmentosum group C gene leads to selective repair of cyclobutane
- [98984] pyrimidine dimers rather than 6–4 photoproducts. Proc. Nat. Acad.
- [98985] Sci. 97: 2151–2156, 2000.
- [98986] 17234.
- [98987] Halley, D. J. J.; Keijzer, W.; Jaspers, N. G. J.; Niermeijer, M.
- [98988] F.; Kleijer, W. J.; Boue, J.; Boue, A.; Bootsma, D.: Prenatal diagnosis
- [98989] of xeroderma pigmentosum (group C) using assays of unscheduled DNA
- [98990] synthesis and postreplication repair. Clin. Genet. 16: 137–146,

[98991] 1979.

[98992] 17235.

[98993] Hananian, J.; Cleaver, J. E.: Xeroderma pigmentosum ex-
hibiting

[98994] neurological disorders and systemic lupus erythematosus.
Clin. Genet. 17:

[98995] 39-45, 1980.

[98996] 17236.

[98997] Kaur, G. P.; Athwal, R. S.: Identification of a DNA repair
gene

[98998] on human chromosome 5 that complements xeroderma
pigmentosum cells

[98999] of group C. (Abstract) Am. J. Hum. Genet. 51 (suppl.):
A51, 1992.

[99000] 17237.

[99001] Khan, S. G.; Levy, H. L.; Legerski, R.; Quackenbush, E.;
Reardon,

[99002] J. T.; Emmert, S.; Sancar, A.; Li, L.; Schneider, T. D.;
Cleaver,

[99003] J. E.; Kraemer, K. H.: Xeroderma pigmentosum group C
splice mutation

- [99004] associated with autism and hypoglycinemia. J. Invest. Derm. 111:
- [99005] 791–796, 1998.
- [99006] 17238.
- [99007] Legerski, R.; Peterson, C.: Expression cloning of a human DNA
- [99008] repair gene involved in xeroderma pigmentosum group C. Nature 359:
- [99009] 70–73, 1992.
- [99010] 17239.
- [99011] Legerski, R. J.; Liu, P.; Li, L.; Peterson, C. A.; Zhao, Y.; Leach,
- [99012] R. J.; Naylor, S. L.; Siciliano, M. J.: Assignment of xeroderma pigmentosum
- [99013] group C (XPC) gene to chromosome 3p25. Genomics 21: 266–269, 1994.
- [99014] 17240.
- [99015] Li, L.; Bales, E. S.; Peterson, C. A.; Legerski, R. J.: Characterization
- [99016] of molecular defects in xeroderma pigmentosum group C. Nature Genet. 5:

- [99017] 413–417, 1993.
- [99018] 17241.
- [99019] Lynch, H. T.; Fusaro, R. M.; Johnson, J. A.: Xeroderma pigmentosum:
- [99020] complementation group C and malignant melanoma. Arch. Derm. 120:
- [99021] 175–179, 1984.
- [99022] 17242.
- [99023] Masutani, C.; Sugasawa, K.; Yanagisawa, J.; Sonoyama, T.; Ui,
- [99024] M.; Enomoto, T.; Takio, K.; Tanaka, K.; van der Spek, P.; Bootsma,
- [99025] D.; Hoeijmakers, J. H. J.; Hanaoka, F.: Purification and cloning
- [99026] of a nucleotide excision repair complex involving the xeroderma pigmentosum
- [99027] group C protein and a human homologue of yeast RAD23. EMBO J. 13:
- [99028] 1831–1843, 1994.
- [99029] 17243.
- [99030] Peterson, C.; Legerski, R.: High-frequency transformation

of

[99031] human repair-deficient cell lines by an Epstein-Barr virus-based cDNA

[99032] expression vector. *Gene* 107: 279-284, 1991.

[99033] 17244.

[99034] Sands, A. T.; Abuin, A.; Sanchez, A.; Conti, C. J.; Bradley, A.

[99035] : High susceptibility to ultraviolet-induced carcinogenesis in mice

[99036] lacking XPC. *Nature* 377: 162-165, 1995.

[99037] 17245.

[99038] Seguin, L. R.; Tarone, R. E.; Liao, K.; Robbins, J. H.: Ultraviolet

[99039] light-induced chromosomal aberrations in cultured cells from Cockayne

[99040] syndrome and complementation group C xeroderma pigmentosum patients:

[99041] lack of correlation with cancer susceptibility. *Am. J. Hum. Genet.* 42:

[99042] 468-475, 1988.

[99043] 17246.

- [99044] Slor, H.; Batko, S.; Khan, S. G.; Sobe, T.; Emmert, S.; Khadavi,
- [99045] A.; Frumkin, A.; Busch, D. B.; Albert, R. B.; Kraemer, K. H.: Clinical,
- [99046] cellular, and molecular features of an Israeli xeroderma pigmentosum
- [99047] family with a frameshift mutation in the XPC gene: sun protection
- [99048] prolongs life. J. Invest. Derm. 115: 974–980, 2000.
- [99049] 17247.
- [99050] Sugasawa, K.; Ng, J. M. Y.; Masutani, C.; Iwai, S.; van der Spek,
- [99051] P. J.; Eker, A. P. M.; Hanaoka, F.; Bootsma, D.; Hoeijmakers, J. H.
- [99052] J.: Xeroderma pigmentosum group C protein complex is the initiator
- [99053] of global genome nucleotide excision repair. Molec. Cell 2: 223–232,
- [99054] 1998.
- [99055] 17248.
- [99056] Suzuki, T.; Park, H.; Hollingsworth, N. M.; Sternglanz, R.; Lennarz,

- [99057] W. L.: PNG1, a yeast gene encoding a highly conserved peptide:N-glycanase. J.
- [99058] Cell Biol. 149: 1039–1052, 2000.
- [99059] 17249.
- [99060] Teitz, T.; Naiman, T.; Avissar, S. S.; Bar, S.; Okayama, H.; Canaani, D.: Complementation of the UV-sensitive phenotype of a xeroderma pigmentosum human cell line by transfection with a cDNA clone library. Proc.
- [99063] Nat. Acad. Sci. 84: 8801–8804, 1987.
- [99064] 17250.
- [99065] van der Spek, P. J.; Smit, E. M. E.; Beverloo, H. B.; Sugawara, K.; Masutani, C.; Hanaoka, F.; Hoeijmakers, J. H. J.; Hage-meijer, A.: Chromosomal localization of three repair genes: the xeroderma pigmentosum group C gene and two human homologs of yeast RAD23. Genomics 23: 651–658, 1994.

[99070] 17251.

[99071] van der Spek, P. J.; Visser, C. E.; Hanaoka, F.; Smit, B.;
Hagemeyer,

[99072] A.; Bootsma, D.; Hoeijmakers, J. H. J.: Cloning, comparative mapping,

[99073] and RNA expression of the mouse homologues of the *Saccharomyces cerevisiae*

[99074] nucleotide excision repair gene RAD23. *Genomics* 31:
20-27, 1996.

[99075] 17252.

[99076] Moser, H. W.: Personal Communication. Baltimore, Md.
6/29/1989.

[99077] 17253.

[99078] Moser, H. W.: Lorenzo's Oil. (Film Review). *Lancet* 341:
544,

[99079] 1993.

[99080] 17254.

[99081] Moser, H. W.: Personal Communication. Baltimore, Md.
1980.

[99082] 17255.

[99083] Moser, H. W.: Personal Communication. Baltimore, Md.

10/13/1987.

[99084] 17256.

[99085] Moser, H. W.; Kok, F.; Neumann, S.; Borel, J.; Bergin, A.;
Mostafa,

[99086] S. D.; Panoscha, R.; Davoli, C. T.; Shankroff, J.; Smith, K.
D.:

[99087] Adrenoleukodystrophy update: genetics and effect of
Lorenzo's oil

[99088] therapy in asymptomatic patients. Int. Pediat. 9: 196–204,
1994.

[99089] 17257.

[99090] Moser, H. W.; Moser, A. B.; Frayer, K. K.; Chen, W.; Schul-
man,

[99091] J. D.; O'Neill, B. P.; Kishimoto, Y.: Adrenoleukodystrophy:
increased

[99092] plasma content of saturated very long chain fatty acids.
Neurology 31:

[99093] 1241–1249, 1981.

[99094] 17258.

[99095] Moser, H. W.; Moser, A. B.; Kawamura, N.; Murphy, J.;
Suzuki,

[99096] K.; Schaumburg, H.; Kishimoto, Y.: Adrenoleukodystrophy:

elevated

[99097] C26 fatty acid in cultured skin fibroblasts. *Ann. Neurol.* 7:
542–549,

[99098] 1980.

[99099] 17259.

[99100] Moser, H. W.; Moser, A. B.; Naidu, S.; Bergin, A.: Clinical
aspects

[99101] of adrenoleukodystrophy and adrenomyeloneuropathy.
Dev. Neurosci. 13:

[99102] 254–261, 1991.

[99103] 17260.

[99104] Moser, H. W.; Moser, A. E.; Singh, I.; O'Neill, B. P.:
Adrenoleukodystrophy:

[99105] survey of 303 cases: biochemistry, diagnosis, and therapy.
Ann. Neurol. 16:

[99106] 628–641, 1984.

[99107] 17261.

[99108] Moser, H. W.; Smith, K. D.; Moser, A. B.: X-linked
adrenoleukodystrophy. In:

[99109] Scriver, C. R.; Beaudet, A. L.; Sly, W. S.; Valle, D. (eds): *The*

[99110] *Metabolic and Molecular Bases of Inherited Disease*. New

York: McGraw-Hill

[99111] 1995. Pp. 2325–2349.

[99112] 17262.

[99113] Moser, H. W.; Tutschka, P. J.; Brown, F. R., III; Moser, A. E.;

[99114] Yeager, A. M.; Singh, I.; Mark, S. A.; Kumar, A. A. J.; Mc-
Donnell,

[99115] J. M.; White, C. L., III; Maumenee, I. H.; Green, W. R.; Pow-
ers, J.

[99116] M.; Santos, G. W.: Bone marrow transplant in
adrenoleukodystrophy. *Neurology* 34:

[99117] 1410–1417, 1984.

[99118] 17263.

[99119] Mosser, J.; Douar, A.–M.; Sarde, C.–O.; Kioschis, P.; Feil,
R.;

[99120] Moser, H.; Poustka, A.–M.; Mandel, J.–L.; Aubourg, P.: Pu-
tative X-linked

[99121] adrenoleukodystrophy gene shares unexpected homology
with ABC transporters. *Nature* 361:

[99122] 726–730, 1993.

[99123] 17264.

[99124] Mosser, J.; Lutz, Y.; Stoeckel, M. E.; Sarde, C. O.; Kretz, C.;

- [99125] Douar, A. M.; Lopez, J.; Aubourg, P.; Mandel, J. L.: The gene responsible
[99126] for adrenoleukodystrophy encodes a peroxisomal membrane protein. Hum.
[99127] Molec. Genet. 3: 265–271, 1994.
[99128] 17265.
- [99129] Naidu, S.; Bresnan, M. J.; Griffin, D.; O'Toole, S.; Moser, H.
[99130] W.: Childhood adrenoleukodystrophy: failure of intensive immunosuppression
[99131] to arrest neurologic progression. Arch. Neurol. 45: 846–848, 1988.
[99132] 17266.
- [99133] Netik, A.; Forss-Petter, S.; Holzinger, A.; Molzer, B.; Unterrainer,
[99134] G.; Berger, J.: Adrenoleukodystrophy-related protein can compensate
[99135] functionally for adrenoleukodystrophy protein deficiency (X-ALD):
[99136] implications for therapy. Hum. Molec. Genet. 8: 907–913, 1999.
[99137] 17267.

- [99138] Noetzel, M. J.; Landau, W. M.; Moser, H. W.:
Adrenoleukodystrophy
- [99139] carrier state presenting as a chronic nonprogressive spinal
cord disorder. Arch.
- [99140] Neurol. 44: 566–567, 1987.
- [99141] 17268.
- [99142] O'Neill, B. P.; Marmion, L. C.; Feringa, E. R.: The
adrenoleukomyeloneuropathy
- [99143] complex: expression in four generations. Neurology 31:
151–156,
- [99144] 1981.
- [99145] 17269.
- [99146] O'Neill, B. P.; Moser, H. W.; Marmion, L. C.:
Adrenoleukodystrophy:
- [99147] elevated C26 fatty acid in cultured skin fibroblasts and
correlation
- [99148] with disease expression in three generations of a kindred.
Neurology 32:
- [99149] 540–542, 1982.
- [99150] 17270.
- [99151] O'Neill, B. P.; Moser, H. W.; Marmion, L. C.: The

adrenoleukomyeloneuropathy

[99152] (ALMN) complex: elevated C26 fatty acid in cultured skin fibroblasts

[99153] and correlation with disease expression in three generations of a

[99154] kindred. (Abstract) Neurology 30: 352 only, 1980.

[99155] 17271.

[99156] O'Neill, B. P.; Moser, H. W.; Saxena, K. M.: Familial X-linked

[99157] Addison disease as an expression of adrenoleukodystrophy (ALD): elevated

[99158] C26 fatty acid in cultured skin fibroblasts. Neurology 32: 543-547,

[99159] 1982.

[99160] 17272.

[99161] O'Neill, B. P.; Swanson, J. W.; Brown, F. R., III; Griffin, J.

[99162] W.; Moser, H. W.: Familial spastic paraparesis: an adrenoleukodystrophy

[99163] phenotype? Neurology 35: 1233-1235, 1985.

[99164] 17273.

[99165] Peckham, R. S.; Marshall, M. C., Jr.; Rosman, P. M.; Farag,

A.;

- [99166] Kabadi, U.; Wallace, E. Z.: A variant of adrenomyeloneu-
ropathy with
[99167] hypothalamic-pituitary dysfunction and neurologic remis-
sion after
[99168] glucocorticoid replacement therapy. Am. J. Med. 72:
173-176, 1982.
[99169] 17274.
- [99170] Penman, R. W. B.: Addison's disease in association with
spastic
[99171] paraplegia. Brit. Med. J. 1: 402, 1960.
[99172] 17275.
- [99173] Poulos, A.; Gibson, R.; Sharp, P.; Beckman, K.; Grattan-
Smith,
[99174] P.: Very long chain fatty acids in X-linked
adrenoleukodystrophy
[99175] brain after treatment with Lorenzo's oil. Ann. Neurol. 36:
741-746,
[99176] 1994.
[99177]
- [99178] 100. Powers, J. M.; Schaumburg, H. H.: The testis in

adreno-leukodystrophy. Am.

[99179] J. Path. 102: 90-98, 1981.

[99180]

[99181] 101. Powers, J. M.; Schaumburg, H. H.: A fatal cause of sexual inadequacy

[99182] in men: adreno-leukodystrophy. J. Urol. 124: 583-585, 1980.

[99183]

[99184] 102. Powers, J. M.; Schaumburg, H. H.: Adreno-leukodystrophy (sex-linked

[99185] Schilder's disease): a pathogenic hypothesis based on ultrastructural

[99186] lesions in adrenal cortex, peripheral nerve and testis. Am. J. Path. 76:

[99187] 481-500, 1974.

[99188]

[99189] 103. Powers, J. M.; Schaumburg, H. H.; Gaffney, C. L.: Kluver-Bucy

[99190] syndrome caused by adreno-leukodystrophy. Neurology 30: 1131-1132,

[99191] 1980.

[99192]

[99193] 104. Probst, A.; Ulrich, J.; Heitz, P. U.; Herschkowitz, N.:
Adrenomyeloneuropathy:

[99194] a protracted, pseudosystematic variant of
adrenoleukodystrophy. Acta

[99195] Neuropath. 49: 105–115, 1980.

[99196]

[99197] 105. Restuccia, D.; Di Lazzaro, V.; Valeriani, M.; Oliviero,
A.; Le

[99198] Pera, D.; Barba, C.; Cappa, M.; Bertini, E.; Di Capua, M.;
Tonali,

[99199] P.: Neurophysiologic follow-up of long-term dietary treat-
ment in

[99200] adult-onset adrenoleukodystrophy. Neurology 52:
810–816, 1999.

[99201]

[99202] 106. Ropers, H.–H.; Burmeister, P.; von Petrykowski, W.;
Schindera,

[99203] F.: Leukodystrophy, skin hyperpigmentation, and adrenal
atrophy:

[99204] Siemerling–Creutzfeldt disease: transmission through
several generations

[99205] in two families. Am. J. Hum. Genet. 27: 547–553, 1975.

[99206]

[99207] 107. Ropers, H.–H.; Zimmermann, J.; Wienker, T.:

Adrenoleukodystrophy

[99208] (Siemerling–Creutzfeldt disease): heterozygote with two
clonal fibroblast

[99209] populations. Clin. Genet. 11: 114–118, 1977.

[99210]

[99211] 108. Rosen, N. L.; Lechtenberg, R.; Wisniewski, K.;

Pullarkat, R.;

[99212] Bennett, H. S.: Adrenoleukomyeloneuropathy with onset in
early childhood. Ann.

[99213] Neurol. 17: 311–312, 1985.

[99214]

[99215] 109. Ruiz, M.; Coll, M. J.; Pampols, T.; Giros, M.: X-linked
adrenoleukodystrophy:

[99216] phenotype distribution and expression of ALDP in Spanish
kindreds. Am.

[99217] J. Med. Genet. 76: 424–427, 1998.

[99218]

- [99219] 110. Sack, G. H., Jr.; Alpern, M.; Webster, T.; Feil, R. P.; Morrell, J. C.; Chen, G.; Chen, W.; Caskey, C. T.; Moser, H. W.: Chromosomal rearrangement segregating with adrenoleukodystrophy: a molecular analysis. Proc. Nat. Acad. Sci. 90: 9489–9493, 1993.
- [99223]
- [99224] 111. Sack, G. H., Jr.; Morrell, J. C.: Adrenoleukodystrophy: overlapping deletions point to a gene location in Xq28. Biochem. Biophys. Res. Commun. 191: 955–960, 1993.
- [99227]
- [99228] 112. Sack, G. H., Jr.; Raven, M. B.; Moser, H. W.: Color vision defects in adrenomyeloneuropathy. Am. J. Hum. Genet. 44: 794–798, 1989.
- [99230]
- [99231] 113. Sadeghi-Nejad, A.; Senior, B.: Adrenomyeloneuropathy presenting

[99232] as Addison's disease in childhood. New Eng. J. Med. 322: 13–16,

[99233] 1990.

[99234]

[99235] 114. Sarde, C.-O.; Mosser, J.; Kioschis, P.; Kretz, C.; Vicaire, S.;

[99236] Aubourg, P.; Poustka, A.; Mandel, J.-L.: Genomic organization of

[99237] the adrenoleukodystrophy gene. Genomics 22: 13–20, 1994.

[99238]

[99239] 115. Sarde, C.-O.; Thomas, J.; Sadoulet, H.; Garnier, J.-M.; Mandel,

[99240] J.-L.: cDNA sequence of Aldgh, the mouse homolog of the X-linked

[99241] adrenoleukodystrophy gene. Mammalian Genome 5: 810–813, 1994.

[99242]

[99243] 116. Schaumburg, H. H.; Powers, J. M.; Raine, C. S.; Spencer, P. S.;

[99244] Griffin, J. W.; Prineas, J. W.; Boehme, D. M.: Adrenomyeloneuropathy:

- [99245] a probable variant of adrenoleukodystrophy. II. General pathologic,
- [99246] neuropathologic, and biochemical aspects. Neurology 27: 1114–1119,
- [99247] 1977.
- [99248]
- [99249] 117. Schaumburg, H. H.; Powers, J. M.; Raine, C. S.; Suzuki, K.; Richardson,
- [99250] E. P., Jr.: Adrenoleukodystrophy: a clinical and pathological study
- [99251] of 17 cases. Arch. Neurol. 32: 577–591, 1975.
- [99252]
- [99253] 118. Schaumburg, H. H.; Richardson, E. P., Jr.; Johnson, P. C.; Cohen,
- [99254] R. B.; Powers, J. M.; Raine, C. S.: Schilder's disease: sex-linked
- [99255] recessive transmission with specific adrenal changes. Arch. Neurol. 27:
- [99256] 458–460, 1972.
- [99257]
- [99258] 119. Schilder, P.: Zur Frage der Encephalitis Periaxialis

Diffusa

[99259] (Sogenannte Diffuse Sklerose). Z. ges Neurol. Psychiat. 15: 359–376,

[99260] 1913.

[99261]

[99262] 120. Seneca, S.; Lissens, W.: DNA diagnosis of X-linked adrenoleukodystrophy. J.

[99263] Inherit. Metab. Dis. 18 (suppl. 1): 34–44, 1995.

[99264]

[99265] 121. Shani, N.; Watkins, P. A.; Valle, D.: PXA1, a possible *Saccharomyces*

[99266] *cerevisiae* ortholog of the human adrenoleukodystrophy gene. Proc.

[99267] Nat. Acad. Sci. 92: 6012–6016, 1995.

[99268]

[99269] 122. Siemerling, E.; Creutzfeldt, H. G.: Bronzekrankheit und sklerosierende

[99270] Encephalomyelitis. Arch. Psychiat. Nervkrankh. 68: 217–244, 1923.

[99271]

[99272] 123. Singh, I.; Lazo, O.; Contreras, M.; Hashmi, M.; Stan-

ley, W. S.

[99273] : Peroxisomal lignoceroyl-CoA ligase deficiency in X-linked adrenoleukodystrophy.

[99274] (Abstract) Am. J. Hum. Genet. 43: A16, 1988.

[99275]

[99276] 124. Sobue, G.; Ueno-Natsukari, I.; Okamoto, H.; Connell, T. A.; Aizawa,

[99277] I.; Mizoguchi, K.; Honma, M.; Ishikawa, G.; Mitsuma, T.; Natsukari,

[99278] N.: Phenotypic heterogeneity of an adult form of adrenoleukodystrophy

[99279] in monozygotic twins. Ann. Neurol. 36: 912-915, 1994.

[99280]

[99281] 125. Spira, T. J.; Adam, A.; Goodman, R. M.; Berger, A.: Recombination

[99282] between cerebral sclerosis--Addison's disease and the Xg blood-groups.

[99283] (Letter) Lancet II: 820-821, 1971.

[99284]

[99285] 126. Takano, H.; Koike, R.; Onodera, O.; Sasaki, R.; Tsuji, S.: Mutational

- [99286] analysis and genotype-phenotype correlation of 29 unrelated Japanese
- [99287] patients with X-linked adrenoleukodystrophy. Arch. Neurol. 56: 295-300,
- [99288] 1999.
- [99289]
- [99290] 127. Turkington, R. W.; Stempfel, R. S., Jr.: Adrenocortical atrophy
- [99291] and diffuse cerebral sclerosis (Addison-Schilder's disease). J. Pediat. 69:
- [99292] 406-412, 1966.
- [99293]
- [99294] 128. Ulrich, J.; Herschkowitz, N.; Heitz, P.; Sigrist, T. H.; Baerlocher,
- [99295] P.: Adrenoleukodystrophy: preliminary report of a congenital case;
- [99296] light- and electronmicroscopical, immunohistochemical and biochemical
- [99297] findings. Acta Neuropath. 43: 77-83, 1978.
- [99298]
- [99299] 129. Unterrainer, G.; Molzer, B.; Forss-Petter, S.; Berger,

J.: Co-expression

[99300] of mutated and normal adrenoleukodystrophy protein reduces protein

[99301] function: implications for gene therapy of X-linked adrenoleukodystrophy. Hum.

[99302] Molec. Genet. 9: 2609–2616, 2000.

[99303]

[99304] 130. Uyama, E.; Iwagoe, H.; Maeda, J.; Nakamura, M.; Terasaki, T.;

[99305] Ando, M.: Presenile-onset cerebral adrenoleukodystrophy presenting

[99306] as Balint's syndrome and dementia. Neurology 43: 1249–1251, 1993.

[99307]

[99308] 131. Valle, D.; Gartner, J.: Penetrating the peroxisome. Nature 361:

[99309] 682–683, 1993.

[99310]

[99311] 132. van den Bosch, H.; Schutgens, R. B. H.; Wanders, R. J. A.; Tager,

[99312] J. M.: Biochemistry of peroxisomes. Annu. Rev. Biochem.

61: 157-197,

[99313] 1992.

[99314]

[99315] 133. van Geel, B. M.; Assies, J.; Weverling, G. J.; Barth, P.
G.:

[99316] Predominance of the adrenomyeloneuropathy phenotype
of X-linked adrenoleukodystrophy

[99317] in the Netherlands: a survey of 30 kindreds. Neurology 44:
2343-2346,

[99318] 1994.

[99319]

[99320] 134. van Oost, B. A.; van Zandvoort, P.; Hoogeboom, A.;
Bakkeren,

[99321] J.; Hamel, B.; Brunner, H.; Knoers, N.; Ropers, H. H.: Tight
linkage

[99322] between adrenoleukodystrophy and DXS 52. (Abstract)
Cytogenet. Cell

[99323] Genet. 46: 708 only, 1987.

[99324]

[99325] 135. van Oost, B. A.; van Zandvoort, P. M.; Tunte, W.;
Brunner, H.

- [99326] G.; Hoogeboom, A. J. M.; Maaswinkel-Mooy, P. D.;
Bakkeren, J.; Hamel,
- [99327] B.; Ropers, H. H.: Linkage analysis in X-linked
adrenoleukodystrophy
- [99328] and application in post- and prenatal diagnosis. Hum.
Genet. 86:
- [99329] 404-407, 1991.
- [99330]
- [99331] 136. Walsh, P. J.: Adrenoleukodystrophy: report of two
cases with
- [99332] relapsing and remitting courses. Arch. Neurol. 37:
448-450, 1980.
- [99333]
- [99334] 137. Wanders, R. J. A.; van Roermund, C. W. T.; van Wij-
land, M. J.
- [99335] A.; Nijenhuis, A. A.; Tromp, A.; Schutgens, R. B. H.;
Brouwer-Kelder,
- [99336] E. M.; Schram, A. W.; Tager, J. M.; van den Bosch, H.;
Schalkwijk,
- [99337] C.: X-linked adrenoleukodystrophy: defective peroxisomal
oxidation
- [99338] of very long chain fatty acids but not of very long chain

fatty acyl-CoA

[99339] esters. Clin. Chim. Acta 165: 321-329, 1987.

[99340]

[99341] 138. Wanders, R. J. A.; van Roermund, C. W. T.; van Wijland, M. J.

[99342] A.; Schutgens, R. B. H.; Heikoop, J.; van den Bosch, H.; Schram, A.

[99343] W.; Tager, J. M.: Peroxisomal fatty acid beta-oxidation in relation

[99344] to the accumulation of very long chain fatty acids in cultured skin

[99345] fibroblasts from patients with Zellweger syndrome and other peroxisomal

[99346] disorders. J. Clin. Invest. 80: 1778-1783, 1987.

[99347]

[99348] 139. Wanders, R. J. A.; van Roermund, C. W. T.; van Wijland, M. J.

[99349] A.; Schutgens, R. B. H.; van den Bosch, H.; Schram, A. W.; Tager,

[99350] J. M.: Direct demonstration that the deficient oxidation of very

[99351] long chain fatty acids in X-linked adrenoleukodystrophy is

due to

[99352] an impaired ability of peroxisomes to activate very long chain fatty

[99353] acids. *Biochem. Biophys. Res. Commun.* 153: 618–624, 1988.

[99354]

[99355] 140. Watkiss, E.; Webb, T.; Bunday, S.: Is skewed X inactivation

[99356] responsible for symptoms in female carriers for adrenoleucodystrophy? *J.*

[99357] *Med. Genet.* 30: 651–654, 1993.

[99358]

[99359] 141. Wilichowski, E.; Ohlenbusch, A.; Korenke, G. C.; et al.: Identical

[99360] mitochondrial DNA in monozygotic twins with discordant adrenoleukodystrophy

[99361] phenotype. *Ann. Neurol.* 43: 835–836, 1998.

[99362]

[99363] 142. Willems, P. J.; Vits, L.; Wanders, R. J. A.; Coucke, P. J.; Van

[99364] der Auwera, B. J.; Van Elsen, A. F.; Raeymaekers, P.; Van

Broeckhoven,

[99365] C.; Schutgens, R. B. H.; Dacremont, G.; Leroy, J. G.; Martin, J.-J.;

[99366] Dumon, J. E.: Linkage of DNA markers at Xq28 to adrenoleukodystrophy

[99367] and adrenomyeloneuropathy present within the same family. Arch. Neurol. 47:

[99368] 665-669, 1990.

[99369]

[99370] 143. Wray, S. H.; Cogan, D. G.; Kuwabara, T.; Schaumburg, H. H.; Powers,

[99371] J. M.: Adrenoleukodystrophy with disease of the eye and optic nerve. Am.

[99372] J. Ophthalm. 82: 480-485, 1976.

[99373] 17276.

[99374] Dinulos, M. B.; Bassi, M. T.; Rugarli, E. I.; Chapman, V.; Ballabio,

[99375] A.; Disteche, C. M.: A new region of conservation is defined between

[99376] human and mouse X chromosomes. Genomics 35: 244-247, 1996.

[99377] 17277.

- [99378] Schiaffino, M. V.; Bassi, M. T.; Rugarli, E. I.; Renieri, A.; Galli,
L.; Ballabio, A.: Cloning of a human homologue of the
Xenopus laevis
APX gene from the ocular albinism type 1 critical region.
Hum. Molec. Genet. 4: 373–382, 1995.
17278.
- [99382] 17278.
- [99383] Bachner, D.; Sedlacek, Z.; Korn, B.; Hameister, H.; Poustka,
A.
: Expression patterns of two human genes coding for dif-
ferent rab
GDP-dissociation inhibitors (GDIs), extremely conserved
proteins involved
in cellular transport. Hum. Molec. Genet. 4: 701–708,
1995.
17279.
- [99387] 17279.
- [99388] Black, G. C. M.; Perveen, R.; Bonshek, R.; Cahill, M.; Clay-
ton-Smith,
J.; Lloyd, I. C.; McLeod, D.: Coats' disease of the retina
(unilateral
retinal telangiectasis) caused by somatic mutation in the

NDP gene:

- [99391] a role for norrin in retinal angiogenesis. Hum. Molec. Genet. 8: 2031–2035, 1999.
- [99392] 17280.
- [99393] Walpole, S. M.; Hiriyana, K. T.; Nicolaou, A.; Bingham, E. L.;
- [99394] Durham, J.; Vaudin, M.; Ross, M. T.; Yates, J. R. W.; Sieving, P.
- [99395] A.; Trump, D.: Identification and characterization of the human homologue
- [99396] (RAI2) of a mouse retinoic acid-induced gene in Xp22. Genomics 55: 275–283, 1999.
- [99397] 17281.
- [99398] Mao, M.; Fu, G.; Wu, J.-S.; Zhang, Q.-H.; Zhou, J.; Kan, L.-X.;
- [99399] Huang, Q.-H.; He, K.-L.; Gu, B.-W.; Han, Z.-G.; Shen, Y.; Gu, J.;
- [99400] Yu, Y.-P.; Xu, S.-H.; Wang, Y.-X.; Chen, S.-J.; Chen, Z.: Identification
- [99401] of genes expressed in human CD34+ hematopoietic

stem/progenitor cells

[99404] by expressed sequence tags and efficient full-length
cDNA cloning. Proc.

[99405] Nat. Acad. Sci. 95: 8175–8180, 1998.

[99406] 17282.

[99407] Pearce, W. G.; Sanger, R.; Race, R. R.: Ocular albinism and
Xg. Lancet I:

[99408] 1282–1283, 1968.

[99409] 17283.

[99410] Rosenberg, T.; Schwartz, M.: X-linked ocular albinism:
prevalence

[99411] and mutations—a national study. Europ. J. Hum. Genet. 6:
570–577,

[99412] 1998.

[99413] 17284.

[99414] Schiaffino, M. V.; Bassi, M. T.; Galli, L.; Renieri, A.; Bruttini,

[99415] M.; De Nigris, F.; Bergen, A. A. B.; Charles, S. J.; Yates, J. R.

[99416] W., Meindl, A.; Lewis, R. A.; King, R. A.; Ballabio, A.: Anal-
ysis

[99417] of the OA1 gene reveals mutations in only one-third of

patients with

[99418] X-linked ocular albinism. Hum. Molec. Genet. 4:
2319-2325, 1995.

[99419] 17285.

[99420] Schiaffino, M. V.; d'Addio, M.; Alloni, A.; Baschiroto, C.;
Valetti,

[99421] C.; Cortese, K.; Puri, C.; Bassi, M. T.; Colla, C.; De Luca,
M.; Tacchetti,

[99422] C.; Ballabio, A.: Ocular albinism: evidence for a defect in
an intracellular

[99423] signal transduction system. Nature Genet. 23: 108-112,
1999.

[99424] 17286.

[99425] Schnur, R. E.; Gao, M.; Wick, P. A.; Keller, M.; Benke, P. J.;

[99426] Edwards, M. J.; Grix, A. W.; Hockey, A.; Jung, J. H.; Kidd, K.
K.;

[99427] Kistenmacher, M.; Levin, A. V.; and 11 others: OA1 muta-
tions and

[99428] deletions in X-linked ocular albinism. Am. J. Hum. Genet.
62: 800-809,

[99429] 1998.

[99430] 17287.

- [99431] Schnur, R. E.; Trask, B. J.; van den Engh, G.; Punnett, H. H.;
- [99432] Kistenmacher, M.; Tomeo, M. A.; Naidu, R. E.; Nussbaum, R. L.: An
- [99433] Xp22 microdeletion associated with ocular albinism and ichthyosis:
- [99434] approximation of breakpoints and estimation of deletion size by using
- [99435] cloned DNA probes and flow cytometry. Am. J. Hum. Genet. 45: 706–720,
- [99436] 1989.
- [99437] 17288.
- [99438] Schnur, R. E.; Wick, P. A.; Bailey, C.; Rebbeck, T.; Weleber,
- [99439] R. G.; Wagstaff, J.; Grix, A. W.; Pagon, R. A.; Hockey, A.; Edwards,
- [99440] M. J.: Phenotypic variability in X-linked ocular albinism: relationship
- [99441] to linkage genotypes. Am. J. Hum. Genet. 55: 484–496, 1994.
- [99442] 17289.
- [99443] Vogt, A.: Die Iris: Albinismus solum bulbi. Atlas Spalt-Lampen-Mikroskopie

[99444] (pub.) (3rd ed.): 1942.

[99445] 17290.

[99446] Waardenburg, P. J.; Van den Bosch, J.: X-chromosomal ocular albinism

[99447] in Dutch family. Ann. Hum. Genet. 21: 101–122, 1956.

[99448] 17291.

[99449] Zhang, Y.; McMahon, R.; Charles, S. J.; Green, J. S.; Moore, A.

[99450] T.; Barton, D. E.; Yates, J. R. W.: Genetic mapping of the Kallmann

[99451] syndrome and X linked ocular albinism gene loci. J. Med. Genet. 30:

[99452] 923–925, 1993.

[99453] 17292.

[99454] Barth, P. G.; Van Wijngaarden, G. K.; Bethlem, J.: X-linked myotubular

[99455] myopathy with fatal neonatal asphyxia. Neurology 25: 531–536, 1975.

[99456] 17293.

[99457] Blondeau, F.; Laporte, J.; Bodin, S.; Superti-Furga, G.; Payastre,

- [99458] B.; Mandel, J.-L.: Myotubularin, a phosphatase deficient in myotubular
- [99459] myopathy, acts on phosphatidylinositol 3-kinase and phosphatidylinositol
- [99460] 3-phosphate pathway. Hum. Molec. Genet. 9: 2223-2229, 2000.
- [99461] 17294.
- [99462] Braga, S. E.; Gerber, A.; Meier, C.; Weiersmuller, A.; Zimmermann,
- [99463] A.; Herrmann, U.; Liechti, S.; Moser, H.: Severe neonatal asphyxia
- [99464] due to X-linked centronuclear myopathy. Europ. J. Pediat. 150: 132-135,
- [99465] 1990.
- [99466] 17295.
- [99467] Breningstall, G. N.; Grover, W. D.; Marks, H. G.: Maternal muscle
- [99468] biopsy in X-linked recessive centronuclear (myotubular) myopathy. Am.
- [99469] J. Med. Genet. 39: 13-18, 1991.
- [99470] 17296.

- [99471] Verga, V.; Hall, B. K.; Wang, S.; Johnson, S.; Higgins, J. V.;
- [99472] Glover, T. W.: Localization of the translocation breakpoint
in a
- [99473] female with Menkes syndrome to Xq13.2-q13.3 proximal
to PGK-1. Am.
- [99474] J. Hum. Genet. 48: 1133-1138, 1991.
- [99475] 17297.
- [99476] Seltzer, W. K.; Firminger, H.; Klein, J.; Pike, A.; Fennessey,
- [99477] P.; McCabe, E. R. B.: Adrenal dysfunction in glycerol kinase
deficiency. Biochem.
- [99478] Med. 33: 189-199, 1985.
- [99479] 17298.
- [99480] Cianfriglia, M.; Miggiano, V. C.; Meo, T.; Muller, H. J.;
- Muller,
- [99481] E.; Battistuzzi, G.: Evidence for syntenic between the rabbit
gene
- [99482] loci coding for HPRT, PGK and G6PD in mouse-rabbit so-
matic cell hybrids.
- [99483] (Abstract) Cytogenet. Cell Genet. 25: 142, 1979.
- [99484] 17299.

- [99485] Goss, S. J.; Harris, H.: Gene transfer by means of cell fusion.
- [99486] I. Statistical mapping of the human X-chromosome by analysis of radiation-induced
- [99487] gene segregation. J. Cell Sci. 25: 17–37, 1977.
- [99488] 17300.
- [99489] Gill, P.; Ivanov, P. L.; Kimpton, C.; Piercy, R.; Benson, N.;
- [99490] Tully, G.; Evett, I.; Hagelberg, E.; Sullivan, K.: Identification
- [99491] of the remains of the Romanov family by DNA analysis. Nature Genet. 6:
- [99492] 130–135, 1994.
- [99493] 17301.
- [99494] Gitschier, J.: Maternal duplication associated with gene deletion
- [99495] in sporadic hemophilia. Am. J. Hum. Genet. 43: 274–279, 1988.
- [99496] 17302.
- [99497] Gitschier, J.; Drayna, D.; Tuddenham, E. G. D.; White, R. L.;
- [99498] Lawn, R. M.: Genetic mapping and diagnosis of haemophilia A achieved

- [99499] through a BclI polymorphism in the factor VIII gene. Nature 314:
- [99500] 738–740, 1985.
- [99501] 17303.
- [99502] Gitschier, J.; Wood, W. I.; Shuman, M. A.; Lawn, R. M.: Identification
- [99503] of a missense mutation in the factor VIII gene of a mild hemophiliac. Science 232:
- [99504] 1415–1416, 1986.
- [99505] 17304.
- [99506] Gitschier, J.; Wood, W. I.; Tuddenham, E. G. D.; Shuman, M. A.;
- [99507] Goralka, T. M.; Chen, E. Y.; Lawn, R. M.: Detection and sequence
- [99508] of mutations in the factor VIII gene of haemophiliacs. Nature 315:
- [99509] 427–430, 1985.
- [99510] 17305.
- [99511] Graham, J. B.; Green, P. P.; McGraw, R. A.; Davis, L. M.: Application
- [99512] of molecular genetics to prenatal diagnosis and carrier

detection

- [99513] in the hemophilias: some limitations. Blood 66: 759-764, 1985.
- [99514] 17306.
- [99515] Graham, J. B.; McLendon, W. W.; Brinkhous, K. M.: Mild hemophilia:
- [99516] an allelic form of the disease. Am. J. Med. Sci. 225: 46-53, 1953.
- [99517] 17307.
- [99518] Green, P. M.; Montandon, A. J.; Bentley, D. R.; Giannelli, F.
- [99519] : Genetics and molecular biology of haemophilias A and B. Blood Coagul.
- [99520] Fibrinolysis 2: 539-565, 1991.
- [99521] 17308.
- [99522] Grover, H.; Phillips, M. A.; Lillicrap, D. P.; Giles, A. R.; Garvey,
- [99523] M. B.; Teitel, J.; Rivard, G.; Blanchette, V.; White, B. N.; Holden,
- [99524] J. J. A.: Carrier deletion of haemophilia A using DNA markers in
- [99525] families with an isolated affected male. Clin. Genet. 32:

- 10-19,
[99526] 1987.
- [99527] 17309.
- [99528] Grozdea, J.; Colombies, P.; Bierme, R.; Ducos, J.:
Myeloperoxidases
[99529] and genetics of haemophilia A. (Letter) Lancet II: 220 only,
1969.
- [99530] 17310.
- [99531] Haldane, J. B. S.; Smith, C. A. B.: A new estimate of the
linkage
[99532] between the genes for colour-blindness and haemophilia
in man. Ann.
[99533] Eugen. 14: 10-31, 1947.
- [99534] 17311.
- [99535] Harper, K.; Winter, R. M.; Pembrey, M. E.; Hartley, D.;
Davies,
[99536] K. E.; Tuddenham, E. G. D.: A clinically useful DNA probe
closely
[99537] linked to haemophilia A. Lancet II: 6-8, 1984.
- [99538] 17312.
- [99539] Hemker, H. C.; Muller, A. D.; Hermens, W. T.; Zwaal, R. F.

A.

- [99540] : Oral treatment of hemophilia A by gastrointestinal absorption of
- [99541] factor VIII entrapped in liposomes. Lancet I: 70–71, 1980.
- [99542] 17313.
- [99543] Hermann, J.: Der Einfluss des Zeugungsalters auf die Mutationen
- [99544] zu Haemophilie A. Humangenetik 3: 1–16, 1966.
- [99545] 17314.
- [99546] Higuchi, M.; Antonarakis, S. E.; Kasch, L.; Oldenburg, J.; Economou-Petersen,
- [99547] E.; Olek, K.; Arai, M.; Inaba, H.; Kazazian, H. H., Jr.: Molecular
- [99548] characterization of mild-to-moderate hemophilia A: detection of the
- [99549] mutation in 25 of 29 patients by denaturing gradient gel electrophoresis. Proc.
- [99550] Nat. Acad. Sci. 88: 8307–8311, 1991.
- [99551] 17315.
- [99552] Higuchi, M.; Kazazian, H. H., Jr.; Kasch, L.; Warren, T. C.; McGinniss,

- [99553] M. J.; Phillips, J. A., III; Kasper, C., III; Janco, R.; Antonarakis,
- [99554] S. E.: Molecular characterization of severe hemophilia A suggests
- [99555] that about half the mutations are not within the coding regions and
- [99556] splice junctions of the factor VIII gene. Proc. Nat. Acad. Sci. 88:
- [99557] 7405–7409, 1991.
- [99558] 17316.
- [99559] Higuchi, M.; Kochhan, L.; Olek, K.: A somatic mosaic for haemophilia
- [99560] A detected at the DNA level. Molec. Biol. Med. 5: 23–27, 1988.
- [99561] 17317.
- [99562] Higuchi, M.; Kochhan, L.; Schwaab, R.; Egli, H.; Brackmann, H.
- [99563] H.; Horst, J.; Olek, K.: Molecular defects in hemophilia A: identification
- [99564] and characterization of mutations in the factor VIII gene and family
- [99565] analysis. Blood 74: 1045–1051, 1989.

[99566] 17318.

[99567] Higuchi, M.; Wong, C.; Kochhan, L.; Olek, K.; Aronis, S.;
Kasper,

[99568] C. K.; Kazazian, H. H., Jr.; Antonarakis, S. E.: Characteriza-
tion

[99569] of mutations in the factor VIII gene by direct sequencing
of amplified

[99570] genomic DNA. Genomics 6: 65-71, 1990.

[99571] 17319.

[99572] Howard, P. L.; Hoag, J. B.; Bovill, E. G.; Heintz, N. H.: Spon-
taneous

[99573] mutation in the male gamete as a cause of hemophilia A:
clarification

[99574] of a case using DNA probes. Am. J. Hemat. 28: 167-169,
1988.

[99575] 17320.

[99576] Hoyer, L. W.; Breckenridge, R. T.: Two forms of
haemophilia?

[99577] (Letter) Lancet II: 457 only, 1968.

[99578] 17321.

[99579] Inaba, H.; Fujimaki, M.; Kazazian, H. H., Jr.; Antonarakis, S.

- [99580] E.: Mild hemophilia A resulting from arg-to-leu substitution in exon
- [99581] 26 of the factor VIII gene. Hum. Genet. 81: 335-338, 1989.
- [99582] 17322.
- [99583] Ivanov, P. L.; Wadhams, M. J.; Roby, R. K.; Holland, M. M.; Weedn,
- [99584] V. W.; Parsons, T. J.: Mitochondrial DNA sequence heteroplasmy in
- [99585] the Grand Duke of Russia Georgij Romanov establishes the authenticity
- [99586] of the remains of Tsar Nicholas II. Nature Genet. 12: 417-420, 1996.
- [99587] 17323.
- [99588] Jaffe, E. A.; Nachman, R. L.: Subunit structure of factor VIII
- [99589] antigen synthesized by cultured human endothelial cells. J. Clin.
- [99590] Invest. 56: 698-702, 1975.
- [99591] 17324.
- [99592] Janco, R. L.; Phillips, J. A., III; Orlando, P. J.; Woodard, M.
- [99593] J.; Wion, K. L.; Lawn, R. M.: Detection of hemophilia A car-

riers

[99594] using intragenic factor VIII:C DNA polymorphisms. Blood
69: 1539–1541,

[99595] 1987.

[99596] 17325.

[99597] Johnson, D. J.; Pemberton, S.; Acquila, M.; Mori, P. G.;
Tuddenham,

[99598] E. G.; O'Brien, D. P.: Factor VIII S373L: mutation at P1' site
confers

[99599] thrombin cleavage resistance, causing mild haemophilia
A. Thromb.

[99600] Haemost. 71: 428–433, 1994.

[99601] 17326.

[99602] Jonsdottir, S.; Diamond, C.; Levinson, B.; Magnusson, S.;
Jensson,

[99603] O.; Gitschier, J.: Missense mutations causing mild
hemophilia A in

[99604] Iceland detected by denaturing gradient gel electrophore-
sis. Hum.

[99605] Mutat. 1: 506–508, 1992.

[99606] 17327.

- [99607] Holterhus, P. M.; Sinnecker, G. H. G.; Wollmann, H. A.;
Struve,
[99608] D.; Homburg, N.; Kruse, K.; Hiort, O.: Expression of two
functionally
[99609] different androgen receptors in a patient with androgen
insensitivity. Europ.
[99610] J. Pediat. 158: 702–706, 1999.
[99611] 17328.
[99612] Shimmin, L. C.; Chang, B.–H.; Li, W.–H.: Male-driven evo-
lution
[99613] of DNA sequences. Nature 362: 745–747, 1993.
[99614] 17329.
[99615] Warburg, M.: Personal Communication. Copenhagen, Den-
mark 1966.
[99616] 17330.
[99617] Imasaki, K.; Okabe, T.; Murakami, H.; Tanaka, Y.; Haji, M.;
Takayanagi,
[99618] R.; Nawata, H.: Androgen insensitivity syndrome due to
new mutations
[99619] in the DNA-binding domain of the androgen receptor.
Molec. Cell.

- [99620] Endocr. 120: 15–24, 1996.
- [99621] 17331.
- [99622] Jakubiczka, S.; Nedel, S.; Werder, E. A.; Schleiermacher, E.;
- [99623] Theile, U.; Wolff, G.; Wieacker, P.: Mutations of the andro-
gen receptor
- [99624] gene in patients with complete androgen insensitivity.
Hum. Mutat. 9:
- [99625] 57–61, 1997.
- [99626] 17332.
- [99627] Kang, H.-Y.; Yeh, S.; Fujimoto, N.; Chang, C.: Cloning and
characterization
- [99628] of human prostate coactivator ARA54, a novel protein that
associates
- [99629] with the androgen receptor. J. Biol. Chem. 274:
8570–8576, 1999.
- [99630] 17333.
- [99631] Kazemi-Esfarjani, P.; Beitel, L. K.; Trifiro, M.; Kaufman, M.;
- [99632] Rennie, P.; Sheppard, P.; Matusik, R.; Pinsky, L.: Substitu-
tion of
- [99633] valine-865 by methionine or leucine in the human andro-
gen receptor

- [99634] causes complete or partial androgen insensitivity, respectively with
- [99635] distinct androgen receptor phenotypes. *Molec. Endocr.* 7: 37–46,
- [99636] 1993.
- [99637] 17334.
- [99638] Kittles, R. A.; Young, D.; Weinrich, S.; Hudson, J.; Argypoulos,
- [99639] G.; Ukoli, F.; Adams–Campbell, L.; Dunston, G. M.: Extent of linkage
- [99640] disequilibrium between the androgen receptor gene CAG and GGC repeats
- [99641] in human populations: implications for prostate cancer risk. *Hum.*
- [99642] *Genet.* 109: 253–261, 2001.
- [99643] 17335.
- [99644] Klocker, H.; Kaspar, F.; Eberle, J.; Uberreiter, S.; Radmayr, C.; Bartsch, G.: Point mutation in the DNA binding domain of the
- [99646] androgen receptor in two families with Reifenstein syndrome. *Am.*

[99647] J. Med. Genet. 50: 1318–1327, 1992.

[99648] 17336.

[99649] Knoke, I.; Allera, A.; Wieacker, P.: Significance of the CAG

[99650] repeat length in the androgen receptor gene (AR) for the
transactivation

[99651] function of an M780I mutant AR. Hum. Genet. 104:
257–261, 1999.

[99652] 17337.

[99653] Kobayashi, Y.; Miwa, S.; Merry, D. E.; Kume, A.; Mei, L.;
Doyu,

[99654] M.; Sobue, G.: Caspase–3 cleaves the expanded androgen
receptor protein

[99655] of spinal and bulbar muscular atrophy in a polyglutamine
repeat length–dependent

[99656] manner. Biochem. Biophys. Res. Commun. 252: 145–150,
1998.

[99657] 17338.

[99658] Koivisto, P. A.; Schleutker, J.; Helin, H.; Ehren–van Eekelen,

[99659] C.; Kallioniemi, O.–P.; Trapman, J.: Androgen receptor
gene alterations

[99660] and chromosomal gains and losses in prostate carcinomas

appearing

[99661] during finasteride treatment for benign prostatic hyperplasia. Clin.

[99662] Cancer Res. 5: 3578–3582, 1999.

[99663] 17339.

[99664] Kooy, R. F.; Reyniers, E.; Storm, K.; Vits, L.; van Velzen, D.;

[99665] de Ruiter, P. E.; Brinkmann, A. O.; de Paepe, A.; Willems, P. J.:

[99666] CAG repeat contraction in the androgen receptor gene in three brothers

[99667] with mental retardation. Am. J. Med. Genet. 85: 209–213, 1999.

[99668] 17340.

[99669] La Spada, A.; Fischbeck, K. H.: Androgen receptor gene defect

[99670] in X-linked spinal and bulbar muscular atrophy. (Abstract) Am. J.

[99671] Hum. Genet. 49 (suppl.): 20, 1991.

[99672] 17341.

[99673] Lewis, M.; Kaita, H.; Giblett, E. R.; Anderson, J.; Philipps,

[99674] S.; Steinberg, A. G.; McAlpine, P. J.: Multiplicity of genetic

polymorphisms

[99675] of blood in the Schmiedeleut Hutterites. Am. J. Med.

Genet. 22:

[99676] 477–485, 1985.

[99677] 17342.

[99678] Lim, H. N.; Chen, H.; McBride, S.; Dunning, A. M.; Nixon,

R. M.;

[99679] Hughes, I. A.; Hawkins, J. R.: Longer polyglutamine tracts

in the

[99680] androgen receptor are associated with moderate to severe

undermasculinized

[99681] genitalia in XY males. Hum. Molec. Genet. 9: 829–834,

2000.

[99682] 17343.

[99683] Lobaccaro, J.-M.; Lumbroso, S.; Belon, C.; Galtier-Dereure,

F.;

[99684] Bringer, J.; Lesimple, T.; Namer, M.; Cutuli, B. F.; Pujol, H.;

Sultan,

[99685] C.: Androgen receptor gene mutation in male breast can-

cer. Hum.

[99686] Molec. Genet. 2: 1799–1802, 1993.

[99687] 17344.

- [99688] Lubahn, D. B.; Brown, T. R.; Simental, J. A.; Higgs, H. N.; Migeon,
- [99689] C. J.; Wilson, E. M.; French, F. S.: Sequence of the intron/
exon
- [99690] junctions of the coding region of the human androgen re-
ceptor gene
- [99691] and identification of a point mutation in a family with
complete androgen
- [99692] insensitivity. Proc. Nat. Acad. Sci. 86: 9534–9538, 1989.
- [99693] 17345.
- [99694] Lubahn, D. B.; Joseph, D. R.; Sar, M.; Tan, J.; Higgs, H. N.;
- [99695] Larson, R. E.; French, F. S.; Wilson, E. M.: The human an-
drogen receptor:
- [99696] complementary deoxyribonucleic acid cloning, sequence
analysis and
- [99697] gene expression in prostate. Molec. Endocr. 2:
1265–1275, 1988.
- [99698] 17346.
- [99699] Lubahn, D. B.; Joseph, D. R.; Sullivan, P. M.; Willard, H. F.;
- [99700] French, F. S.; Wilson, E. M.: Cloning of human androgen
receptor

- [99701] complementary DNA and localization to the X chromosome. Science 240:
- [99702] 327-330, 1988.
- [99703] 17347.
- [99704] Lumbroso, S.; Lobaccaro, J. M.; Georget, V.; Leger, J.; Poul,
- [99705] N.; Terouanne, B.; Evain-Brion, D.; Czernichow, P.; Sultan, C.: A
- [99706] novel substitution (leu707-to-arg) in exon 4 of the androgen receptor
- [99707] gene causes complete androgen resistance. J. Clin. Endocr. Metab. 81:
- [99708] 1984-1988, 1996.
- [99709] 17348.
- [99710] Stuppia, L.; Calabrese, G.; Franchi, P. G.; Mingarelli, R.; Gatta,
- [99711] V.; Palka, G.; Dallapiccola, B.: Widening of a Y-chromosome interval-6
- [99712] deletion transmitted from a father to his infertile son accounts for
- [99713] an oligozoospermia critical region distal to the RBM1 and DAZ genes.

- [99714] (Letter) *Am. J. Hum. Genet.* 59: 1393–1395, 1996.
- [99715] 17349.
- [99716] Brown, G. M.; Furlong, R. A.; Sargent, C. A.; Erickson, R. P.;
- [99717] Longepied, G.; Mitchell, M.; Jones, M. H.; Hargreave, T. B.;
Cooke,
- [99718] H. J.; Affara, N. A.: Characterisation of the coding se-
quence and
- [99719] fine mapping of the human DFFRY gene and comparative
expression analysis
- [99720] and mapping to the Sxr-b interval of the mouse Y chro-
mosome of the
- [99721] Dffry gene. *Hum. Molec. Genet.* 7: 97–107, 1998.
- [99722] 17350.
- [99723] Ferlin, A.; Moro, E.; Garolla, A.; Foresta, C.: Human male
infertility
- [99724] and Y chromosome deletions: role of the AZF-candidate
genes DAZ, RBM
- [99725] and DFFRY. *Hum. Reprod.* 14: 1710–1716, 1999.
- [99726] 17351.
- [99727] Foresta, C.; Ferlin, A.; Moro, E.: Deletion and expression
analysis

- [99728] of AZFa genes on the human Y chromosome revealed a major role for
- [99729] DBY in male infertility. *Hum. Molec. Genet.* 9: 1161–1169, 2000.
- [99730] 17352.
- [99731] Sargent, C. A.; Boucher, C. A.; Kirsch, S.; Brown, G.; Weiss, B.; Trundley, A.; Burgoyne, P.; Saut, N.; Durand, C.; Levy, N.; Terriou, P.; Hargreave, T.; Cooke, H.; Mitchell, M.; Rappold, G. A.; Affara, N. A.: The critical region of overlap defining the AZFa male infertility interval of proximal Yq contains three transcribed sequences. *J. Med. Genet.* 36: 670–677, 1999.
- [99732] 17353.
- [99733] Shen, P.; Wang, F.; Underhill, P. A.; Franco, C.; Yang, W.-H.; Roxas, A.; Sung, R.; Lin, A. A.; Hyman, R. W.; Vollrath, D.; Davis, R. W.; Cavalli-Sforza, L. L.; Oefner, P. J.: Population genetic implications

- [99741] from sequence variation in four Y chromosome genes.
Proc. Nat. Acad.
- [99742] Sci. 97: 7354–7359, 2000.
- [99743] 17354.
- [99744] Sun, C.; Skaletsky, H.; Birren, B.; Devon, K.; Tang, Z.; Silber,
ber,
- [99745] S.; Oates, R.; Page, D. C.: An azoospermic man with a de
novo point
- [99746] mutation in the Y-chromosomal gene USP9Y. Nature
Genet. 23: 429–432,
- [99747] 1999.
- [99748] 17355.
- [99749] Thomson, R.; Pritchard, J. K.; Shen, P.; Oefner, P. J.; Feld-
man,
- [99750] M. W.: Recent common ancestry of human Y chromo-
somes: evidence from
- [99751] DNA sequence data. Proc. Nat. Acad. Sci. 97: 7360–7365,
2000.
- [99752] 17356.
- [99753] Schiebel, K.; Winkelmann, M.; Mertz, A.; Xu, X.; Page, D.
C.; Weil,

- [99754] D.; Petit, C.; Rappold,, G. A.: Abnormal XY interchange between a
- [99755] novel isolated protein kinase gene, PRKY, and its homologue, PRKX,
- [99756] accounts for one third of all (Y+)XX males and (Y-)XY females. Hum.
- [99757] Molec. Genet. 6: 1985-1989, 1997.
- [99758] 17357.
- [99759] Scott, D.; Addey, C.; Ellis, P; James, E.; Mitchell, M. J.; Saut,
- [99760] N.; Jurcevic, S.; Simpson, E.: Dendritic cells permit identification
- [99761] of genes encoding MHC class II-restricted epitopes of transplantation
- [99762] antigens. Immunity 12: 711-720, 2000.
- [99763] 17358.
- [99764] Wu, G.; Chai, J.; Suber, T. L.; Wu, J.-W.; Du, C.; Wang, X.; Shi,
- [99765] Y.: Structural basis of IAP recognition by Smac/DIABLO. Nature 408:
- [99766] 1008-1012, 2000.
- [99767] 17359.

- [99768] Shih, V. E.: Urea cycle disorders and other congenital hyperammonemic
- [99769] syndromes. In: Stanbury, J. B.; Wyngaarden, J. B.; Fredrickson, D.
- [99770] S.: The Metabolic Basis of Inherited Disease. New York: McGraw-Hill
- [99771] (pub.) (4th ed.): 1978. Pp. 362–386.
- [99772]
- [99773] 100. Shih, V. E.; Berson, E. L.; Mandell, R.; Schmidt, S. Y.: Ornithine
- [99774] ketoacid transaminase deficiency in gyrate atrophy of the choroid
- [99775] and retina. Am. J. Hum. Genet. 30: 174–179, 1978.
- [99776]
- [99777] 101. Short, E. M.; Conn, H. O.; Snodgrass, P. J.; Campbell, A. G.
- [99778] M.; Rosenberg, L. E.: Evidence for X-linked dominant inheritance
- [99779] of ornithine transcarbamylase deficiency. New Eng. J. Med. 288:
- [99780] 7–12, 1973.

[99781]

[99782] 102. Snodgrass, P. J.; Wappner, R. S.; Brandt, I. K.: White cell

[99783] ornithine transcarbamylase activity cannot detect the liver enzyme

[99784] deficiency. (Letter) *Pediat. Res.* 12: 873, 1978.

[99785]

[99786] 103. Stoll, C.; Bieth, R.; Dreyfus, J.; Flori, E.; Lutz, P.; Levy,

[99787] J.-M.: Une nouvelle famille avec mutation du gene de structure de

[99788] l'ornithine carbamyltransferase humaine. *Arch. Franc. Pe-diat.* 35:

[99789] 512-518, 1978.

[99790]

[99791] 104. Strautnieks, S.; Rutland, P.; Malcolm, S.: Arginine 109 to glutamine

[99792] mutation in a girl with ornithine carbamoyl transferase deficiency. *J.*

[99793] *Med. Genet.* 28: 871-874, 1991.

[99794]

- [99795] 105. Suess, P. J.; Tsai, M. Y.; Holzknecht, R. A.; Horowitz, M.; Tuchman, M.: Screening for gene deletions and known mutations in 13 patients with ornithine transcarbamylase deficiency. *Biochem. Med. Metab.* Biol. 47: 250–259, 1992.
- [99799]
- [99800] 106. Sunshine, P.; Lindenbaum, J. E.; Levy, H. L.; Freeman, J. M. : Hyperammonemia due to a defect in hepatic ornithine transcarbamylase. *Pediatrics* 50: 100–111, 1972.
- [99803]
- [99804] 107. Thaler, M. M.; Hoogenraad, N. J.; Boswell, M.: Reye's syndrome due to a novel protein-tolerant variant of ornithine-transcarbamylase deficiency. *Lancet* II: 438–440, 1974.
- [99807]
- [99808] 108. Tripp, J. H.; Hargreaves, T.; Anthony, P. P.; Searle, J.

F.;

[99809] Miller, P.; Leonard, J. V.; Patrick, A. D.; Oberholzer, V. G.:

Sodium

[99810] valproate and ornithine carbamyl transferase deficiency.

(Letter) Lancet I:

[99811] 1165–1166, 1981.

[99812]

[99813] 109. Tuchman, M.: Mutations and polymorphisms in the human ornithine

[99814] transcarbamylase gene. Hum. Mutat. 2: 174–178, 1993.

[99815]

[99816] 110. Tuchman, M.; Holzknecht, R. A.; Gueron, A. B.; Berry, S. A.;

[99817] Tsai, M. Y.: Six new mutations in the ornithine transcarbamylase

[99818] gene detected by single-strand conformational polymorphism. Pediat.

[99819] Res. 32: 600–604, 1992.

[99820]

[99821] 111. Tuchman, M.; Jaleel, N.; Morizono, H.; Sheehy, L.; Lynch, M.

- [99822] G.: Mutations and polymorphisms in the human ornithine transcarbamylase
- [99823] gene. Hum. Mutat. 19: 93–107, 2002.
- [99824]
- [99825] 112. Tuchman, M.; Matsuda, I.; Munnich, A.; Malcolm, S.; Strautnieks,
- [99826] S.; Briede, T.: Proportions of spontaneous mutations in males and
- [99827] females with ornithine transcarbamylase deficiency. Am. J. Med. Genet. 55:
- [99828] 67–70, 1995.
- [99829]
- [99830] 113. Tuchman, M.; Morizono, H.; Reish, O.; Yuan, X.; Allewell, N.
- [99831] M.: The molecular basis of ornithine transcarbamylase deficiency:
- [99832] modelling the human enzyme and the effects of mutations. J. Med.
- [99833] Genet. 32: 680–688, 1995.
- [99834]
- [99835] 114. Tuchman, M.; Plante, R. J.; Garcia-Perez, M. A.; Ru-

bio, V.:

[99836] Relative frequency of mutations causing ornithine transcarbamylase

[99837] deficiency in 78 families. Hum. Genet. 97: 274–276, 1996.

[99838]

[99839] 115. Tuchman, M.; Plante, R. J.; McCann, M. T.; Qureshi, A. A.: Seven

[99840] new mutations in the human ornithine transcarbamylase gene. Hum.

[99841] Mutat. 4: 57–60, 1994.

[99842]

[99843] 116. Veres, G.; Gibbs, R. A.; Scherer, S. E.; Caskey, C. T.: The

[99844] molecular basis of the sparse fur mouse mutation. Science 237: 415–417,

[99845] 1987.

[99846]

[99847] 117. Wareham, K. A.; Lyon, M. F.; Glenister, P. H.; Williams, E. D.

[99848] : Age related reactivation of an X-linked gene. Nature 327: 725–727,

[99849] 1987.

[99850]

[99851] 118. Wettke-Schafer, R.; Kantner, G.: X-linked dominant inherited

[99852] diseases with lethality in hemizygous males. Hum. Genet. 64: 1-23,

[99853] 1983.

[99854]

[99855] 119. Wilson, C. J.; Lee, P. J.; Leonard, J. V.: Plasma glutamine

[99856] and ammonia concentrations in ornithine carbamoyltransferase deficiency

[99857] and citrullinaemia. J. Inherit. Metab. Dis. 24: 691-695, 2001.

[99858]

[99859] 120. Winter, R. M.; Donnai, D.; Burn, J.; Tuckers, S. M.: Fetal valproate

[99860] syndrome: is there a recognisable phenotype?. J. Med. Genet. 24:

[99861] 692-695, 1987.

[99862]

- [99863] 121. Yorifuji, T.; Muroi, J.; Uematsu, A.; Tanaka, K.; Kiwaki, K.;
- [99864] Endo, F.; Matsuda, I.; Nagasaka, H.; Furusho, K.: X-inactivation
- [99865] pattern in the liver of a manifesting female with ornithine transcarbamylase
- [99866] (OTC) deficiency. Clin. Genet. 54: 349–353, 1998.
- [99867]
- [99868] 122. Yudkoff, M.; Daikhin, Y.; Nissim, I.; Jawad, A.; Wilson, J.;
- [99869] Batshaw, M.: In vivo nitrogen metabolism in ornithine transcarbamylase
- [99870] deficiency. J. Clin. Invest. 98: 2167–2173, 1996.
- [99871]
- [99872] 123. Yudkoff, M.; Yang, W.; Snodgrass, P. J.; Segal, S.: Ornithine
- [99873] transcarbamylase deficiency in a boy with normal development. J.
- [99874] Pediat. 96: 441–443, 1980.
- [99875] 17360.
- [99876] Macke, J. P.; Hu, N.; Hu, S.; Bailey, M.; King, V. L.; Brown,

[99877] T.; Hamer, D.; Nathans, J.: Sequence variation in the androgen receptor

- [99878] gene is not a common determinant of male sexual orientation. Am.
- [99879] J. Hum. Genet. 53: 844–852, 1993.
- [99880] 17361.
- [99881] Nakamura, H.; Izumoto, Y.; Kambe, H.; Kuroda, T.; Mori, T.; Kawamura,
- [99882] K.; Yamamoto, H.; Kishimoto, T.: Molecular cloning of complementary
- [99883] DNA for a novel human hepatoma–derived growth factor: its homology
- [99884] with high mobility group–1 protein. J. Biol. Chem. 269: 25143–25149,
- [99885] 1994.
- [99886] 17362.
- [99887] Wanschura, S.; Schoenmakers, E. F. P. M.; Huysmans, C.; Bartnitzke,
- [99888] S.; Van de Ven, W. J. M.; Bullerdiek, J.: Mapping of the gene encoding
- [99889] the human hepatoma–derived growth factor (HDGF) with homology to the
- [99890] high–mobility group (HMG)–1 protein to Xq25. Genomics 32: 298–300,

- [99891] 1996.
- [99892] 17363.
- [99893] Gedeon, A. K.; Glass, I. A.; Connor, J. M.; Mulley, J. C.: Genetic
- [99894] localisation of MRX27 to Xq24-26 defines another discrete gene for
- [99895] non-specific X-linked mental retardation. Am. J. Med. Genet. 64:
- [99896] 121-124, 1996.
- [99897] 17364.
- [99898] Amir, J.; Alpert, G.; Statter, M.; Gutman, A.; Reisner, S. H.:
- [99899] Intracranial haemorrhage in siblings and ornithine transcarbamylase
- [99900] deficiency. Acta Paediat. Scand. 71: 671-673, 1982.
- [99901] 17365.
- [99902] Anadiotis, G.; Ierardi-Curto, L.; Kaplan, P. B.; Berry, G. T.:
- [99903] Ornithine transcarbamylase deficiency and pancreatitis. J. Pediat. 138:
- [99904] 123-124, 2001.
- [99905] 17366.

- [99906] Ardinger, H. H.; Atkin, J. F.; Blackston, R. D.; Elsas, L. J.;
- [99907] Clarren, S. K.; Livingstone, S.; Flannery, D. B.; Pellock, J. M.;
- [99908] Harrod, M. J.; Lammer, E. J.; Majewski, F.; Schinzel, A.; Toriello,
- [99909] H. V.; Hanson, J. W.: Verification of the fetal valproate syndrome
- [99910] phenotype. *Am. J. Med. Genet.* 29: 171–185, 1988.
- [99911] 17367.
- [99912] McMurtrie, E. B.; Barbosa, M. D. F. S.; Zerial, M.; Kingsmore,
- [99913] S. F.: Rab17 and Rab18, small GTPases with specificity for polarized
- [99914] epithelial cells: genetic mapping in the mouse. *Genomics* 45: 623–625,
- [99915] 1997.
- [99916] 17368.
- [99917] Schafer, U.; Seibold, S.; Schneider, A.; Neugebauer, E.: Isolation
- [99918] and characterisation of the human rab18 gene after stimulation of

- [99919] endothelial cells with histamine. FEBS Lett. 466: 148–154, 2000.
- [99920] 17369.
- [99921] Doupnik, C. A.; Davidson, N.; Lester, H. A.: The inward rectifier
- [99922] potassium channel family. Curr. Opin. Neurobiol. 5: 268–277, 1995.
- [99923] 17370.
- [99924] Tada, Y.; Horio, Y.; Takumi, T.; Terayama, M.; Tsuji, L.; Copeland,
- [99925] N. G.; Jenkins, N. A.; Kurachi, Y.: Assignment of the glial inwardly
- [99926] rectifying potassium channel K(AB)–2/Kir4.1 (Kcnj10) gene to the distal
- [99927] region of mouse chromosome 1. Genomics 45: 629–630, 1997.
- [99928] 17371.
- [99929] Takumi, T.; Ishii, T.; Horio, Y.; Morishige, K.–I.; Takahashi,
- [99930] N.; Yamada, M.; Yamashita, T.; Kiyama, H.; Sohmiya, K.; Nakanishi,
- [99931] S.; Kurachi, Y.: A novel ATP–dependent inward rectifier potassium

- [99932] channel expressed predominantly in glial cells. J. Biol. Chem. 270: 16339–16436, 1995.
- [99933] 17372.
- [99934] Schenker, T.; Trueb, B.: Assignment of the gene for a developmentally regulated GTP-binding protein (DRG2) to human chromosome bands 17p13–p12 by in situ hybridization. Cytogenet. Cell Genet. 79: 274–275, 1997.
- [99935] 17373.
- [99936] Heikkinen, J.; Risteli, M.; Wang, C.; Latvala, J.; Rossi, M.; Valtavaara, M.; Myllyla, R.: Lysyl hydroxylase 3 is a multifunctional protein possessing collagen glucosyltransferase activity. J. Biol. Chem. 275: 36158–36163, 2000.
- [99937] 17374.
- [99938] Passoja, K.; Rautavuoma, K.; Ala-Kokko, L.; Kosonen, T.; Kivirikko,

- [99945] K. I.: Cloning and characterization of a third human lysyl hydroxylase
- [99946] isoform. Proc. Nat. Acad. Sci. 95: 10482–10486, 1998.
- [99947] 17375.
- [99948] Valtavaara, M.; Szpirer, C.; Szpirer, J.; Myllyla, R.: Primary
- [99949] structure, tissue distribution, and chromosomal localization of a
- [99950] novel isoform of lysyl hydroxylase (lysyl hydroxylase 3). J. Biol.
- [99951] Chem. 273: 12881–12886, 1998.
- [99952] 17376.
- [99953] Ishibashi, T.; Bottaro, D. P.; Michieli, P.; Kelley, C. A.; Aaronson,
- [99954] S. A.: A novel dual specificity phosphatase induced by serum stimulation
- [99955] and heat shock. J. Biol. Chem. 269: 29897–29902, 1994.
- [99956] 17377.
- [99957] Kwak, S. P.; Dixon, J. E.: Multiple dual specificity protein tyrosine
- [99958] phosphatases are expressed and regulated differentially in liver cell

- [99959] lines. J. Biol. Chem. 270: 1156–1160, 1995.
- [99960] 17378.
- [99961] Kovalenko, O. V.; Golub, E. I.; Bray–Ward, P.; Ward, D. C.;
Radding,
- [99962] C. M.: A novel nucleic acid–binding protein that interacts
with human
- [99963] Rad51 recombinase. Nucleic Acids Res. 25: 4946–4953,
1997.
- [99964] 17379.
- [99965] Mizuta, R.; LaSalle, J. M.; Cheng, H.–L.; Shinohara, A.;
Ogawa,
- [99966] H.; Copeland, N.; Jenkins, N. A.; Lalande, M.; Alt, F. W.:
RAB22
- [99967] and RAB163/mouse BRCA2: proteins that specifically in–
teract with the
- [99968] RAD51 protein. Proc. Nat. Acad. Sci. 94: 6927–6932,
1997.
- [99969] 17380.
- [99970] Blasius, R.; Weber, R. G.; Lichter, P.; Ogilvie, A.: A novel
orphan
- [99971] G protein–coupled receptor primarily expressed in the
brain is localized

- [99972] on human chromosomal band 2q21. J. Neurochem. 70: 1357–1365, 1998.
- [99973] 17381.
- [99974] Raport, C. J.; Schweickart, V. L.; Chantry, D.; Eddy, R. L., Jr.;
- [99975] Shows, T. B.; Godiska, R.; Gray, P. W.: New members of the chemokine
- [99976] receptor gene family. J. Leukoc. Biol. 59: 18–23, 1996.
- [99977] 17382.
- [99978] Shindo, M.; Nakano, H.; Kuroyanagi, H.; Shirasawa, T.; Mi-hara,
- [99979] M.; Gilbert, D. J.; Jenkins, N. A.; Copeland, N. G.; Yagita, H.; Okumura,
- [99980] K.: cDNA cloning, expression, subcellular localization, and chromosomal
- [99981] assignment of mammalian aurora homologues, aurora-related kinase (ARK)
- [99982] 1 and 2. Biochem. Biophys. Res. Commun. 244: 285–292, 1998.
- [99983] 17383.
- [99984] Presky, D. H.; Yang, H.; Minetti, L. J.; Chua, A. O.; Nabavi, N.;

- [99985] Wu, C.-Y.; Gately, M. K.; Gubler, U.: A functional interleukin 12
- [99986] receptor complex is composed of two beta-type cytokine receptor subunits. Proc.
- [99987] Nat. Acad. Sci. 93: 14002-14007, 1996.
- [99988] 17384.
- [99989] Yamamoto, K.; Kobayashi, H.; Miura, O.; Hirose, S.; Miyasaka,
- [99990] N.: Assignment of IL12RB1 and IL12RB2, interleukin-12 receptor beta-1
- [99991] and beta-2 chains, to human chromosome 19 band p13.1 and chromosome
- [99992] 1 band p31.2, respectively, by in situ hybridization. Cytogenet.
- [99993] Cell Genet. 77: 257-258, 1997.
- [99994] 17385.
- [99995] Turelli, P.; Doucas, V.; Craig, E.; Mangeat, B.; Klages, N.; Evans,
- [99996] R.; Kalpana, G.; Trono, D.: Cytoplasmic recruitment of INI1 and PML
- [99997] on incoming HIV preintegration complexes: interference with early

[99998] steps of viral replication. *Molec. Cell* 7: 1245–1254, 2001.

[99999] 17386.

[100000] Lafreniere, R. G.; Rochefort, D. L.; Kibar, Z.; Fon, E. A.;
Han,

[100001] F.-Y.; Cochius, J.; Kang, X.; Baird, S.; Korneluk, R. G.; An-
dermann,

[100002] E.; Rommens, J. M.; Rouleau, G. A.: Isolation and charac-
terization

[100003] of GT335, a novel human gene conserved in *Escherichia*
coli and mapping

[100004] to 21q22.3. *Genomics* 38: 264–272, 1996.

[100005] 17387.

[100006] Nagamine, K.; Kudoh, J.; Minoshima, S.; Kawasaki, K.;
Asakawa,

[100007] S.; Ito, F.; Shimizu, N.: Isolation of cDNA for a novel hu-
man protein

[100008] KNP-I that is homologous to the *E. coli* SCRP-27A protein
from the

[100009] autoimmune polyglandular disease type I (APECED) region
of chromosome

[100010] 21q22.3. *Biochem. Biophys. Res. Commun.* 225: 608–616,
1996.

[100011]17388.

[100012]Scott, H. S.; Chen, H.; Rossier, C.; Lalioti, M. D.; Antonarakis,

[100013]S. E.: Isolation of a human gene (HES1) with homology to an Escherichia

[100014]coli and a zebrafish protein that maps to chromosome 21q22.3. Hum.

[100015]Genet. 99: 616–623, 1997.

[100016]17389.

[100017]Bernier-Villamor, V.; Sampson, D. A.; Matunis, M. J.; Lima, C.

[100018]D.: Structural basis for E2-mediated SUMO conjugation revealed by

[100019]a complex between ubiquitin-conjugating enzyme Ubc9 and RanGAP1. Cell 108:

[100020]345–356, 2002.

[100021]17390.

[100022]Shi, Y.; Zou, M.; Farid, N. R.; Paterson, M. C.: Association of

[100023]FHIT (fragile histidine triad), a candidate tumour suppressor gene,

- [100024]with the ubiquitin-conjugating enzyme hUBC9. *Biochem. J.* 352: 443-448,
- [100025]2000.
- [100026]17391.
- [100027]Tachibana, M.; Iwata, N.; Watanabe, A.; Nobukuni, Y.; Plo-
plis,
- [100028]B.; Kajigaya, S.: Assignment of the gene for a ubiquitin-
conjugating
- [100029]enzyme (UBE2I) to human chromosome band 16p13.3 by
in situ hybridization. *Cytogenet.*
- [100030]*Cell Genet.* 75: 222-223, 1996.
- [100031]17392.
- [100032]Wang, Z.-Y.; Qiu, Q.-Q.; Seufert, W.; Taguchi, T.; Testa, J.
R.;
- [100033]Whitmore, S. A.; Callen, D. F.; Welsh, D.; Shenk, T.; Deuel,
T. F.
- [100034]: Molecular cloning of the cDNA and chromosome local-
ization of the
- [100035]gene for human ubiquitin-conjugating enzyme 9. *J. Biol.*
Chem. 271:
- [100036]24811-24816, 1996.

[100037]17393.

[100038]Yasugi, T.; Howley, P. M.: Identification of the structural
and

[100039]functional human homolog of the yeast ubiquitin conjugating enzyme

[100040]UBC9. Nucleic Acids Res. 24: 2005–2010, 1996.

[100041]17394.

[100042]Bowen, M. A.; Patel, D. D.; Li, X.; Modrell, B.; Malacko, A.
R.;

[100043]Wang, W.-C.; Marquardt, H.; Neubauer, M.; Pesando, J. M.;
Francke,

[100044]U.; Haynes, B. F.; Aruffo, A.: Cloning, mapping, and characterization

[100045]of activated leukocyte–cell adhesion molecule (ALCAM), a
CD6 ligand. J.

[100046]Exp. Med. 181: 2213–2220, 1995.

[100047]17395.

[100048]Skonier, J. E.; Bowen, M. A.; Emswiler, J.; Aruffo, A.; Bajorath,

[100049]J.: Recognition of diverse proteins by members of the immunoglobulin

[100050]superfamily: delineation of the receptor binding site in the human

[100051]CD6 ligand ALCAM. Biochemistry 35: 12287–12291, 1996.

[100052]17396.

[100053]Bord, S.; Horner, A.; Beavan, S.; Compston, J.: Estrogen receptors

[100054]alpha and beta are differentially expressed in developing human bone. J.

[100055]Clin. Endocr. Metab. 86: 2309–2314, 2001.

[100056]17397.

[100057]Watanabe, T. K.; Fujiwara, T.; Kawai, A.; Shimizu, F.; Takami,

[100058]S.; Hirano, H.; Okuno, S.; Ozaki, K.; Takeda, S.; Shimada, Y.; Nagata,

[100059]M.; Takaichi, A.; Takahashi, E.; Nakamura, Y.; Shin, S.: Cloning,

[100060]expression, and mapping of UBE2I, a novel gene encoding a human homologue

[100061]of yeast ubiquitin-conjugating enzymes which are critical for regulating

[100062]the cell cycle. Cytogenet. Cell Genet. 72: 86–89, 1996.

[100063]17398.

[100064]Chaidarun, S. S.; Swearingen, B.; Alexander, J. M.: Differential

[100065]expression of estrogen receptor-beta (ER-beta) in human
pituitary

[100066]tumors: functional interactions with ER-alpha and a tumor-specific

[100067]splice variant. J. Clin. Endocr. Metab. 83: 3308-3315,
1998.

[100068]17399.

[100069]Chu, S.; Mamers, P.; Burger, H. G.; Fuller, P. J.: Estrogen
receptor

[100070]isoform gene expression in ovarian stromal and epithelial
tumors. J.

[100071]Clin. Endocr. Metab. 85: 1200-1205, 2000.

[100072]17400.

[100073]Enmark, E.; Peltö-Huikko, M.; Grandien, K.; Lagercrantz,
S.; Lagercrantz,

[100074]J.; Fried, G.; Nordenskjöld, M.; Gustafsson, J.-A.: Human
estrogen

[100075]receptor beta-gene structure, chromosomal localization,
and expression

[100076]pattern. J. Clin. Endocr. Metab. 82: 4258–4265, 1997.

[100077]17401.

[100078]Esmaeli, B.; Harvey, J. T.; Hewlett, B.: Immunohistochemical evidence

[100079]for estrogen receptors in meibomian glands. Ophthalmology 107: 180–184,

[100080]2000.

[100081]17402.

[100082]Forsell, C.; Enmark, E.; Axelman, K.; Blomberg, M.; Wahlund, L.-O.;

[100083]Gustafsson, J.-A.; Lannfelt, L.: Investigations of a CA repeat in

[100084]the oestrogen receptor beta gene in patients with Alzheimer's disease. Europ.

[100085]J. Hum. Genet. 9: 802–804, 2001.

[100086]17403.

[100087]Korach, K. S.; Couse, J. F.; Curtis, S. W.; Washburn, T. F.; Lindzey,

[100088]J.; Kimbro, K. S.; Eddy, E. M.; Migliaccio, S.; Snedeker, S. M.; Lubahn,

[100089]D. B.; Schomberg, D. W.; Smith, E. P.: Estrogen receptor

gene disruption:

[100090]molecular characterization and experimental and clinical phenotypes. Recent

[100091]Prog. Horm. Res. 51: 159–188, 1996.

[100092]17404.

[100093]Krege, J. H.; Hodgin, J. B.; Couse, J. F.; Enmark, E.; Warner,

[100094]M.; Mahler, J. F.; Sar, M.; Korach, K. S.; Gustafsson, J.-A.; Smithies,

[100095]O.: Generation and reproductive phenotypes of mice lacking estrogen

[100096]receptor beta. Proc. Nat. Acad. Sci. 95: 15677–15682, 1998.

[100097]17405.

[100098]Krezel, W.; Dupont, S.; Krust, A.; Chambon, P.; Chapman, P. F.

[100099]: Increased anxiety and synaptic plasticity in estrogen receptor beta-deficient

[100100]mice. Proc. Nat. Acad. Sci. 98: 12278–12282, 2001.

[100101]17406.

[100102]Kuiper, G. G. J. M.; Enmark, E.; Peltö-Huikko, M.; Nilsson, S.;

[100103]Gustafsson, J.-A.: Cloning of a novel estrogen receptor expressed

[100104]in rat prostate and ovary. Proc. Nat. Acad. Sci. 93: 5925–5930,

[100105]1996.

[100106]17407.

[100107]Mosialos, G.; Birkenbach, M.; Yalamanchili, R.; VanArsdale, T.;

[100108]Ware, C.; Kieff, E.: The Epstein–Barr virus transforming protein

[100109]LMP1 engages signaling proteins for the tumor necrosis factor receptor

[100110]family. Cell 80: 389–399, 1995.

[100111]17408.

[100112]Rothe, M.; Wong, S. C.; Henzel, W. J.; Goeddel, D. V.: A novel

[100113]family of putative signal transducers associated with the cytoplasmic

[100114]domain of the 75 kDa tumor necrosis factor receptor. Cell 78: 681–692,

[100115]1994.

[100116]17409.

[100117]Siemienski, K.; Peters, N.; Scheurich, P.; Wajant, H.: Organization

[100118]of the human tumour necrosis factor receptor-associated factor 1 (TRAF1)

[100119]gene and mapping to chromosome 9q33-34. Gene 195: 35-39, 1997.

[100120]17410.

[100121]Tsitsikov, E. N.; Laouini, D.; Dunn, I. F.; Sannikova, T. Y.; Davidson,

[100122]L.; Alt, F. W.; Geha, R. S.: TRAF1 is a negative regulator of TNF

[100123]signaling: enhanced TNF signaling in TRAF1-deficient mice. Immunity 15:

[100124]647-657, 2001.

[100125]17411.

[100126]Miyamoto, K.; Kesterson, R. A.; Yamamoto, H.; Taketani, Y.; Nishiwaki,

[100127]E.; Tatsumi, S.; Inoue, Y.; Morita, K.; Takeda, E.; Pike, J. W.:

[100128]Structural organization of the human vitamin D receptor chromosomal

[100129]gene and its promoter. Molec. Endocr. 11: 1165–1179, 1997.

[100130]17412.

[100131]Pedeutour, F.; Merscher, S.; Durieux, E.; Montgomery, K.; Krauter,

[100132]K.; Clevy, J.-P.; Barcelo, G.; Kucherlapati, R.; Gaudray, P.; Turc-Carel,

[100133]C.: Mapping of the 12q12–q22 region with respect to tumor translocation

[100134]breakpoints. Genomics 22: 512–518, 1994.

[100135]17413.

[100136]Riggs, B. L.: Vitamin D-receptor genotypes and bone density.

[100137](Editorial) New Eng. J. Med. 337: 125–126, 1997.

[100138]17414.

[100139]Ritchie, H. H.; Hughes, M. R.; Thompson, E. T.; Malloy, P. J.;

[100140]Hochberg, Z.; Feldman, D.; Pike, J. W.; O'Malley, B. W.: An ochre

[100141]mutation in the vitamin D receptor gene causes hereditary 1,25-dihydroxyvitamin

[100142]D3-resistant rickets in three families. Proc. Nat. Acad. Sci.

86:

[100143]9783–9787, 1989.

[100144]17415.

[100145]Saijo, T.; Ito, M.; Takeda, E.; Mahbubul Huq, A. H. M.;

Naito,

[100146]E.; Yokota, I.; Sone, T.; Pike, J. W.; Kuroda, Y.: A unique

mutation

[100147]in the vitamin D receptor gene in three Japanese patients

with vitamin

[100148]D-dependent rickets type II: utility of single-strand con-

formation

[100149]polymorphism analysis for heterozygous carrier detection.

Am. J.

[100150]Hum. Genet. 49: 668–673, 1991.

[100151]17416.

[100152]Sainz, J.; Van Tornout, J. M.; Loro, L.; Sayre, J.; Roe, T. F.;

[100153]Gilsanz, V.: Vitamin D-receptor gene polymorphisms and

bone density

[100154]in prepubertal American girls of Mexican descent. New

Eng. J. Med. 337:

[100155]77–82, 1997.

[100156]17417.

[100157]Sone, T.; Marx, S. J.; Liberman, U. A.; Pike, J. W.: A unique

[100158]point mutation in the human vitamin D receptor chromo-
somal gene confers

[100159]hereditary resistance to 1,25-dihydroxyvitamin D(3).

Molec. Endocr. 4:

[100160]623–631, 1990.

[100161]17418.

[100162]Suarez, F.; Zeghoud, F.; Rossignol, C.; Walrant, O.;

Garabedian,

[100163]M.: Association between vitamin D receptor gene poly-
morphism and

[100164]sex-dependent growth during the first two years of life. J.
Clin.

[100165]Endocr. Metab. 82: 2966–2970, 1997.

[100166]17419.

[100167]Tao, C.; Yu, T.; Garnett, S.; Briody, J.; Knight, J.; Wood-
head,

[100168]H.; Cowell, C. T.: Vitamin D receptor alleles predict growth
and

[100169]bone density in girls. Arch. Dis. Child. 79: 488–494, 1998.

[100170]17420.

[100171]Uitterlinden, A. G.; Burger, H.; Huang, Q.; Odding, E.; van
Duijn,

[100172]C. M.; Hofman, A.; Birkenhager, J. C.; van Leeuwen, J. P. T.
M.; Pols,

[100173]H. A. P.: Vitamin D receptor genotype is associated with
radiographic

[100174]osteoarthritis at the knee. J. Clin. Invest. 100: 259–263,
1997.

[100175]17421.

[100176]Uitterlinden, A. G.; Weel, A. E. A. M.; Burger, H.; Fang, Y.;

[100177]Van Duijn, C. M.; Hofman, A.; Van Leeuwen, J. P. T. M.;
Pols, H. A.

[100178]P.: Interaction between the vitamin D receptor gene and
collagen

[100179]type I- α -1 gene in susceptibility for fracture. J. Bone
Miner.

[100180]Res. 16: 379–385, 2001.

[100181]17422.

[100182]Whitfield, G. K.; Selznick, S. H.; Haussler, C. A.; Hsieh,

J.-C.;

[100183]Galligan, M. A.; Jurutka, P. W.; Thompson, P. D.; Lee, S. M.;
Zerwekh,

[100184]J. E.; Haussler, M. R.: Vitamin D receptors from patients
with resistance

[100185]to 1,25-dihydroxyvitamin D(3): point mutations confer re-
duced transactivation

[100186]in response to ligand and impaired interaction with the
retinoid X

[100187]receptor heterodimeric partner. Molec. Endocr. 10:
1617-1631, 1996.

[100188]17423.

[100189]Yoshizawa, T.; Handa, Y.; Uematsu, Y.; Takeda, S.; Sekine,
K.;

[100190]Yoshihara, Y.; Kawakami, T.; Arioka, K.; Sato, H.;
Uchiyama, Y.; Masushige,

[100191]S.; Fukamizu, A.; Matsumoto, T.; Kato, S.: Mice lacking the
vitamin

[100192]D receptor exhibit impaired bone formation, uterine hy-
poplasia and

[100193]growth retardation after weaning. Nature Genet. 16:
391-396, 1997.

[100194]17424.

[100195]Zhu, W.; Malloy, P. J.; Delvin, E.; Chabot, G.; Feldman, D.:

[100196]Hereditary 1,25-dihydroxyvitamin D-resistant rickets due to an opal

[100197]mutation causing premature termination of the vitamin D receptor. J.

[100198]Bone Miner. Res. 13: 259–264, 1998.

[100199]17425.

[100200]Bejjani, B. A.; Lewis, R. A.; Tomey, K. F.; Anderson, K. L.; Dueker,

[100201]D. K.; Jabak, M.; Astle, W. F.; Otterud, B.; Leppert, M.; Lupski,

[100202]J. R.: Mutations in CYP1B1, the gene for cytochrome P4501B1, are

[100203]the predominant cause of primary congenital glaucoma in Saudi Arabia. Am.

[100204]J. Hum. Genet. 62: 325–333, 1998.

[100205]17426.

[100206]Bejjani, B. A.; Stockton, D. W.; Lewis, R. A.; Tomey, K. F.; Dueker,

[100207]D. K.; Jabak, M.; Astle, W. F.; Lupski, J. R.: Multiple CYP1B1 mutations

[100208]and incomplete penetrance in an inbred population segregating primary

[100209]congenital glaucoma suggest frequent de novo events and a dominant

[100210]modifier locus. Hum. Molec. Genet. 9: 367–374, 2000.

[100211]17427.

[100212]Lin, J.; Arnold, H. B.; Della-Fera, M. A.; Azain, M. J.; Hartzell,

[100213]D. L.; Baile, C. A.: Myostatin knockout in mice increases myogenesis

[100214]and decreases adipogenesis. Biochem. Biophys. Res. Commun. 291:

[100215]701–706, 2002.

[100216]17428.

[100217]McPherron, A. C.; Lawler, A. M.; Lee, S.-J.: Regulation of skeletal

[100218]muscle mass in mice by a new TGF- β superfamily member. Nature 387:

[100219]83–90, 1997.

[100220]17429.

[100221]Strautnieks, S. S.; Bull, L. N.; Knisely, A. S.; Kocoshis, S. A.;

[100222]Dahl, N.; Arnell, H.; Sokal, E.; Dahan, K.; Childs, S.; Ling, V.;

[100223]Tanner, M. S.; Kagalwalla, A. F.; Nemeth, A.; Pawlowska, J.; Baker,

[100224]A.; Mieli-Vergani, G.; Freimer, N. B.; Gardiner, R. M.; Thompson,

[100225]R. J.: A gene encoding a liver-specific ABC transporter is mutated

[100226]in progressive familial intrahepatic cholestasis. Nature Genet. 20:

[100227]233-238, 1998.

[100228]17430.

[100229]Roldan-Arjona, T.; Wei, Y.-F.; Carter, K. C.; Klungland, A.; Anselmino,

[100230]C.; Wang, R.-P.; Augustus, M.; Lindahl, T.: Molecular cloning and

[100231]functional expression of a human cDNA encoding the antitumor enzyme

[100232]8-hydroxyguanine-DNA glycosylase. Proc. Nat. Acad. Sci. 94: 8016-8020,

[100233]1997.

[100234]17431.

[100235]Goldman, M. J.; Anderson, G. M.; Stolzenberg, E. D.; Kari,
U. P.;

[100236]Zasloff, M.; Wilson, J. M.: Human beta-defensin-1 is a
salt-sensitive

[100237]antibiotic in lung that is inactivated in cystic fibrosis. Cell
88:

[100238]553-560, 1997.

[100239]17432.

[100240]Guo, Z.; Turner, C.; Castle, D.: Relocation of the t-SNARE
SNAP-23

[100241]from lamellipodia-like cell surface projections regulates
compound

[100242]exocytosis in mast cells. Cell 94: 537-548, 1998.

[100243]17433.

[100244]Lazo, P. A.; Nadal, M.; Ferrer, M.; Area, E.; Hernandez-Tor-
res,

[100245]J.; Nabokina, S. M.; Mollinedo, F.; Estivill, X.: Genomic or-
ganization,

[100246]chromosomal localization, alternative splicing, and iso-
forms of the

[100247]human synaptosome-associated protein-23 gene impli-
cated in vesicle-membrane

[100248]fusion processes. Hum. Genet. 108: 211–215, 2001.

[100249]17434.

[100250]Mollinedo, F.; Lazo, P. A.: Identification of two isoforms of

[100251]the vesicle–membrane fusion protein SNAP–23 in human
neutrophils and

[100252]HL–60 cells. Biochem. Biophys. Res. Commun. 231:
808–812, 1997.

[100253]17435.

[100254]Ravichandran, V.; Chawla, A.; Roche, P. A.: Identification
of

[100255]a novel syntaxin– and synaptobrevin/VAMP–binding pro-
tein, SNAP–23,

[100256]expressed in non–neuronal tissues. J. Biol. Chem. 271:
13300–13303,

[100257]1996.

[100258]17436.

[100259]Shukla, A.; Corydon, T. J.; Nielsen, S.; Hoffmann, H. J.;
Dahl,

[100260]R.: Identification of three new splice variants of the SNARE
protein

[100261]SNAP–23. Biochem. Biophys. Res. Commun. 285:

320–327, 2001.

[100262]17437.

[100263]Mugita, N.; Kumura, Y.; Ogawa, M.; Saya, H.; Nakao, M.:

Identification

[100264]of a novel, tissue-specific calpain htra-3; a human homologue of the

[100265]Caenorhabditis elegans sex determination gene. Biochem. Biophys.

[100266]Res. Commun. 239: 845–850, 1997.

[100267]17438.

[100268]Bohme, B.; Holtrich, U.; Wolf, G.; Luzius, H.; Grzeschik, K.-H.;

[100269]Strebhardt, K.; Rubsamen-Waigmann, H.: PCR mediated detection of

[100270]a new human receptor-tyrosine-kinase, HEK 2. Oncogene 8: 2857–2862,

[100271]1993.

[100272]17439.

[100273]Ruiz, J. C.; Conlon, F. L.; Robertson, E. J.: Identification of

[100274]novel protein kinases expressed in the myocardium of the developing

[100275]mouse heart. Mech. Dev. 48: 153–164, 1994.

[100276]17440.

[100277]Ashbourne, K. J.; Byth, B. C.; Meijers, J. C. M.; Cox, D. W.:

[100278]Polymorphism of the protein C inhibitor (PCI) gene on
chromosome 14. Hum.

[100279]Molec. Genet. 2: 92 only, 1993.

[100280]17441.

[100281]Sadler, J. E.: Combined factors V and VIII deficiency climbs
onto

[100282]the map. (Editorial) J. Clin. Invest. 99: 555–556, 1997.

[100283]17442.

[100284]Yasuda, T.; Nadano, D.; Iida, R.; Tanaka, Y.; Nakanaga, M.;
Kishi,

[100285]K.: Discovery of a genetic polymorphism of human plasma
protein C

[100286]inhibitor (PCI): genetic survey utilizing isoelectric focusing
followed

[100287]by immunoblotting, immunological and biochemical char-
acterization. Hum.

[100288]Genet. 89: 265–269, 1992.

[100289]17443.

[100290]Arturi, F.; Russo, D.; Schlumberger, M.; du Villard, J.-A.;
Caillou,

[100291]B.; Vigneri, P.; Wicker, R.; Chiefari, E.; Suarez, H. G.; Filetti,

[100292]S.: Iodide symporter gene expression in human thyroid
tumors. J.

[100293]Clin. Endocr. Metab. 83: 2493–2496, 1998.

[100294]17444.

[100295]Cho, J.-Y.; Leveille, R.; Kao, R.; Rousset, B.; Parlow, A. F.;

[100296]Burak, W. E., Jr.; Mazzaferri, E. L.; Jhiang, S. M.: Hormonal
regulation

[100297]of radioiodide uptake activity and Na⁺/I⁻ symporter ex-
pression in

[100298]mammary glands. J. Clin. Endocr. Metab. 85: 2936–2943,
2000.

[100299]17445.

[100300]Couch, R. M.; Dean, H. J.; Winter, J. S.: Congenital hy-
pothyroidism

[100301]caused by defective iodide transport. J. Pediat. 106:
950–953, 1985.

[100302]17446.

[100303]Li, Z.; Matsuoka, S.; Hryshko, L. V.; Nicoll, D. A.; Bersohn, M.

[100304]M.; Burke, E. P.; Lifton, R. P.; Philipson, K. D.: Cloning of the

[100305]NCX2 isoform of the plasma membrane Na(+)-Ca(2+) exchanger. J. Biol.

[100306]Chem. 269: 17434-17439, 1994.

[100307]17447.

[100308]Detter, J. C.; Zhang, Q.; Mules, E. H.; Novak, E. K.; Mishra, V.

[100309]S.; Li, W.; McMurtrie, E. B.; Tchernev, V. T.; Wallace, M. R.; Seabra,

[100310]M. C.; Swank, R. T.; Kingsmore, S. F.: Rab geranylgeranyl transferase

[100311]alpha mutation in the gunmetal mouse reduces Rab prenylation and platelet

[100312]synthesis. Proc. Nat. Acad. Sci. 97: 4144-4149, 2000.

[100313]17448.

[100314]Novak, E. K.; Reddington, M.; Zhen, L.; Stenberg, P. E.; Jackson,

[100315]C. W.; McGarry, M. P.; Swank, R. T. :Blood 85: 1781-1789, 1995.

[100316]17449.

[100317]Swank, R. T.; Jiang, S. Y.; Reddington, M.; Conway, J.;
Stephenson,

[100318]D.; McGarry, M. P.; Novak, E. K. :Blood 81: 2626–2635,
1993.

[100319]17450.

[100320]Bui, T. D.; Rankin, J.; Smith, K.; Huguet, E. L.; Ruben, S.;
Strachan,

[100321]T.; Harris, A. L.; Lindsay, S.: A novel human Wnt gene,
WNT10B, maps

[100322]to 12q13 and is expressed in human breast carcinomas.
Oncogene 14:

[100323]1249–1253, 1997.

[100324]17451.

[100325]Hardiman, G.; Kastelein, R. A.; Bazan, J. F.: Isolation, char-
acterization

[100326]and chromosomal localization of human WNT10B. Cyto-
genet. Cell Genet. 77:

[100327]278–282, 1997.

[100328]17452.

[100329]Ross, S. E.; Hemati, N.; Longo, K. A.; Bennett, C. N.; Lucas,

P.

[100330]C.; Erickson, R. L.; MacDougald, O. A.: Inhibition of adipogenesis

[100331]by Wnt signaling. Science 289: 950–953, 2000.

[100332]17453.

[100333]Meyerhardt, J. A.; Look, A. T.; Bigner, S. H.; Fearon, E. R.:

[100334]Identification and characterization of neogenin, a DCC-related gene. Oncogene 14:

[100335]1129–1136, 1997.

[100336]17454.

[100337]Vielmetter, J.; Chen, X.-N.; Miskevich, F.; Lane, R. P.; Yamakawa,

[100338]K.; Korenberg, J. R.; Dreyer, W. J.: Molecular characterization of

[100339]human neogenin, a DCC-related protein, and the mapping of its gene

[100340](NEO1) to chromosomal position 15q22.3–q23. Genomics 41: 414–421,

[100341]1997.

[100342]17455.

[100343]Vielmetter, J.; Kayyem, J. F.; Roman, J. M.; Dreyer, W. J.:

Neogenin,

[100344]an avian cell surface protein expressed during terminal neuronal differentiation,

[100345]is closely related to the human tumor suppressor molecule deleted

[100346]in colorectal cancer. *J. Cell Biol.* 127: 2009–2020, 1994.

[100347]17456.

[100348]Rosenquist, T. A.; Zharkov, D. O.; Grollman, A. P.: Cloning and

[100349]characterization of a mammalian 8-oxoguanine DNA glycosylase. *Proc.*

[100350]*Nat. Acad. Sci.* 94: 7429–7434, 1997.

[100351]17457.

[100352]Tani, M.; Shinmura, K.; Kohno, T.; Shiroishi, T.; Wakama, S.;

[100353]Kim, S.-R.; Nohmi, T.; Kasai, H.; Takenoshita, S.; Nagamachi, Y.;

[100354]Yokota, J.: Genomic structure and chromosomal localization of the

[100355]mouse Ogg1 gene that is involved in the repair of 8-hydroxyguanine

[100356]in DNA damage. *Mammalian Genome* 9: 32–37, 1998.

[100357]17458.

[100358]Lim, H. N.; Hawkins, J. R.; Hughes, I. A.: Genetic evidence
to

[100359]exclude the androgen receptor co-factor, ARA70 (NCOA4)
as a candidate

[100360]gene for the causation of undermasculinised genitalia.
(Letter) Clin.

[100361]Genet. 59: 284–286, 2001.

[100362]17459.

[100363]Santoro, M.; Dathan, N. A.; Berlingieri, M. T.; Bongarzone,
I.;

[100364]Paulin, C.; Grieco, M.; Pierotti, M. A.; Vecchio, G.; Fusco,
A.:

[100365]Molecular characterization of RET/PTC3: a novel rear-
ranged version

[100366]of the RET proto-oncogene in a human thyroid papillary
carcinoma. Oncogene 9:

[100367]509–516, 1994.

[100368]17460.

[100369]Yeh, S.; Chang, C.: Cloning and characterization of a spe-
cific

[100370]coactivator, ARA-70, for the androgen receptor in human prostate cells. Proc.

[100371]Nat. Acad. Sci. 93: 5517-5521, 1996.

[100372]17461.

[100373]Huttner, K. M.; Kozak, C. A.; Bevins, C. L.: The mouse genome

[100374]encodes a single homolog of the antimicrobial peptide human beta-defensin

[100375]1. FEBS Lett. 413: 45-49, 1997.

[100376]17462.

[100377]Jia, H. P.; Starner, T.; Ackermann, M.; Kirby, P.; Tack, B. F.;

[100378]McCray, P. B., Jr.: Abundant human beta-defensin-1 expression in

[100379]milk and mammary gland epithelium. J. Pediat. 138: 109-112, 2001.

[100380]17463.

[100381]Morrison, G. M.; Davidson, D. J.; Kilanowski, F. M.; Borthwick,

[100382]D. W.; Crook, K.; Maxwell, A. I.; Govan, J. R. W.; Dorin, J. R.:

[100383]Mouse beta defensin-1 is a functional homolog of human

beta defensin-1. Mammalian

[100384]Genome 9: 453-457, 1998.

[100385]17464.

[100386]Singh, P. K.; Jia, H. P.; Wiles, K.; Hesselberth, J.; Liu, L.;

[100387]Conway, B.-A. D.; Greenberg, E. P.; Valore, E. V.; Welsh, M.
J.; Ganz,

[100388]T.; Tack, B. F.; McCray, P. B., Jr.: Production of beta-
defensins

[100389]by human airway epithelia. Proc. Nat. Acad. Sci. 95:
14961-14966,

[100390]1998.

[100391]17465.

[100392]Tunzi, C. R.; Harper, P. A.; Bar-Oz, B.; Valore, E. V.; Sem-
ple,

[100393]J. L.; Watson-MacDonell, J.; Ganz, T.; Ito, S.: Beta-defensin
expression

[100394]in human mammary gland epithelia. Pediat. Res. 48:
30-35, 2000.

[100395]17466.

[100396]Valore, E. V.; Park, C. H.; Quayle, A. J.; Wiles, K. R.; Mc-
Cray,

[100397]P. B., Jr.; Ganz, T.: Human beta-defensin-1: an antimicrobial peptide

[100398]of urogenital tissues. J. Clin. Invest. 101: 1633-1642, 1998.

[100399]17467.

[100400]Nagasawa, A.; Kubota, R.; Imamura, Y.; Nagamine, K.; Wang, Y.;

[100401]Asakawa, S.; Kudoh, J.; Minoshima, S.; Mashima, Y.; Oguchi, Y.; Shimizu,

[100402]N.: Cloning of the cDNA for a new member of the immunoglobulin superfamily

[100403](ISLR) containing leucine-rich repeat (LRR). Genomics 44: 273-279,

[100404]1997.

[100405]17468.

[100406]Paoloni-Giacobino, A.; Chen, H.; Peitsch, M. C.; Rossier, C.; Antonarakis,

[100407]S. E.: Cloning of the TMPRSS2 gene, which encodes a novel serine

[100408]protease with transmembrane, LDLRA, and SRCR domains and maps to 21q22.3. Genomics 44:

[100409]309-320, 1997. Note: Erratum: Genomics 44: 309-320,

1997.

[100410]17469.

[100411]McAllister, G.; Whiting, P.; Hammond, E. A.; Knowles, M.
R.; Atack,

[100412]J. R.; Bailey, F. J.; Maigetter, R.; Ragan, C. I.: cDNA cloning
of

[100413]human and rat brain myo-inositol monophosphatase: ex-
pression and characterization

[100414]of the human recombinant enzyme. Biochem. J. 284:
749-754, 1992.

[100415]17470.

[100416]Sjoholt, G.; Molven, A.; Lovlie, R.; Wilcox, A.; Sikela, J. M.;

[100417]Steen, V. M.: Genomic structure and chromosomal local-
ization of a

[100418]human myo-inositol monophosphatase gene (IMPA). Ge-
nomics 45: 113-122,

[100419]1997.

[100420]17471.

[100421]Steen, V. M.; Gulbrandsen, A. K.; Eiken, H. G.; Berle, J. O.:

[100422]Lack of genetic variation in the coding region of the myo-
inositol

[100423]monophosphatase gene in lithium-treated patients with
manic depressive

[100424]illness. Pharmacogenetics 6: 113–116, 1996.

[100425]17472.

[100426]McDonald, M. T.; Flejter, W.; Sheldon, S.; Putzi, M. J.;
Gorski,

[100427]J. L.: XY sex reversal and gonadal dysgenesis due to 9p24
monosomy. Am.

[100428]J. Med. Genet. 73: 321–326, 1997.

[100429]17473.

[100430]Muroya, K.; Okuyama, T.; Goishi, K.; Ogiso, Y.; Fukuda, S.;
Kameyama,

[100431]J.; Sato, H.; Suzuki, Y.; Terasaki, H.; Gomyo, H.; Wakui, K.;
Fukushima,

[100432]Y.; Ogata, T.: Sex-determining gene(s) on distal 9p: clinical and

[100433]molecular studies in six cases. J. Clin. Endocr. Metab. 85:
3094–3100,

[100434]2000.

[100435]17474.

[100436]Raymond, C. S.; Shamu, C. E.; Shen, M. M.; Seifert, K. J.;

Hirsch,

[100437]B.; Hodgkin, J.; Zarkower, D.: Evidence for evolutionary conservation

[100438]of sex-determining genes. Nature 391: 691-695, 1998.

[100439]17475.

[100440]Shan, Z.; Zabel, B.; Trautmann, U.; Hillig, U.; Ottolenghi, C.;

[100441]Wang, Y.; Haaf, T.: FISH mapping of the sex-reversal region on human

[100442]chromosome 9p in two XY females and in primates. Europ. J. Hum. Genet. 8:

[100443]167-173, 2000.

[100444]17476.

[100445]Shen, M. M.; Hodgkin, J.: mab-3, a gene required for sex-specific

[100446]yolk protein expression and a male-specific lineage in C. elegans. Cell 54:

[100447]1019-1031, 1988.

[100448]17477.

[100449]Smith, C. A.; McClive, P. J.; Western, P. S.; Reed, K. J.; Sinclair,

- [100450]A. H.: Conservation of a sex-determining gene. (Letter)
Nature 402:
[100451]601-602, 1999.
- [100452]17478.
- [100453]Veitia, R.; Nunes, M.; Brauner, R.; Doco-Fenzy, M.; Joanny-
Flinois,
[100454]O.; Jaubert, F.; Lortat-Jacob, S.; Fellous, M.; McElreavey,
K.: Deletions
[100455]of distal 9p associated with 46,XY male to female sex re-
versal: definition
[100456]of the breakpoints at 9p23.3-p24.1. Genomics 41:
271-274, 1997.
- [100457]17479.
- [100458]Takekawa, M.; Posas, F.; Saito, H.: A human homolog of
the yeast
[100459]Ssk2/Ssk22 MAP kinase kinase kinases, MTK1, mediates
stress-induced
[100460]activation of the p38 and JNK pathways. EMBO J. 16:
4973-4982, 1997.
- [100461]17480.
- [100462]Chapman, D. L.; Agulnik, I.; Hancock, S.; Silver, L. M.; Pa-
paioannou,

- [100463]V. E.: Tbx6, a mouse T-box gene implicated in paraxial mesoderm formation at gastrulation. Dev. Biol. 180: 534–542, 1996.
- [100465]17481.
- [100466]Chapman, D. L.; Papaioannou, V. E.: Three neural tubes in mouse embryos with mutations in the T-box gene Tbx6. Nature 391: 695–697, 1998.
- [100468]17482.
- [100469]17482.
- [100470]Papapetrou, C.; Putt, W.; Fox, M.; Edwards, Y. H.: The human TBX6 gene: cloning and assignment to chromosome 16p11.2. Genomics 55: 238–241, 1999.
- [100472]238–241, 1999.
- [100473]17483.
- [100474]Spicer, A. P.; Olson, J. S.; McDonald, J. A.: Molecular cloning and characterization of a cDNA encoding the third putative mammalian hyaluronan synthase. J. Biol. Chem. 272: 8957–8961,

1997.

[100477]17484.

[100478]Bagri, A.; Marin, O.; Plump, A. S.; Mak, J.; Pleasure, S. J.;
Rubenstein,

[100479]J. L. R.; Tessier-Lavigne, M.: Slit proteins prevent midline
crossing

[100480]and determine the dorsoventral position of major axonal
pathways in

[100481]the mammalian forebrain. Neuron 33: 233–248, 2002.

[100482]17485.

[100483]Dallol, A.; Forgacs, E.; Martinez, A.; Sekido, Y.; Walker, R.;

[100484]Kishida, T.; Rabbitts, P.; Maher, E. R.; Minna, J. D.; Latif, F.:

[100485]Tumour specific promoter region methylation of the hu-
man homologue

[100486]of the Drosophila Roundabout gene DUTT1 (ROBO1) in
human cancers. Oncogene 21:

[100487]3020–3028, 2002.

[100488]17486.

[100489]Kidd, T.; Brose, K.; Mitchell, K. J.; Fetter, R. D.; Tessier-
Lavigne,

[100490]M.; Goodman, C. S.; Tear, G.: Roundabout controls axon

crossing of

[100491]the CNS midline and defines a novel subfamily of evolutionarily conserved

[100492]guidance receptors. Cell 92: 205–215, 1998.

[100493]17487.

[100494]Sundaresan, V.; Chung, G.; Heppell–Parton, A.; Xiong, J.; Grundy,

[100495]C.; Roberts, I.; James, L.; Cahn, A.; Bench, A.; Douglas, J.; Minna,

[100496]J.; Sekido, Y.; Lerman, M.; Latif, F.; Bergh, J.; Li, H.; Lowe, N.;

[100497]Ogilvie, D.; Rabbitts, P.: Homozygous deletions at 3p12 in breast

[100498]and lung cancer. Oncogene 17: 1723–1729, 1998.

[100499]17488.

[100500]Zallen, J. A.; Yi, B. A.; Bargmann, C. I.: The conserved immunoglobulin

[100501]superfamily member SAX–3/Robo directs multiple aspects of axon guidance

[100502]in C. elegans. Cell 92: 217–227, 1998.

[100503]17489.

[100504]Faber, P. W.; Barnes, G. T.; Srinidhi, J.; Chen, J.; Gusella, J.
 [100505]F.; MacDonald, M. E.: Huntingtin interacts with a family of
 WW domain
 [100506]proteins. Hum. Molec. Genet. 7: 1463–1474, 1998.
 [100507]17490.
 [100508]Hattula, K.; Peranen, J.: FIP-2, a coiled-coil protein, links
 [100509]huntingtin to Rab8 and modulates cellular morphogene-
 sis. Curr. Biol. 10:
 [100510]1603–1606, 2000.
 [100511]17491.
 [100512]Li, Y.; Kang, J.; Horwitz, M. S.: Interaction of an adenovirus
 [100513]E3 14.7-kilodalton protein with a novel tumor necrosis
 factor alpha-inducible
 [100514]cellular protein containing leucine zipper domains. Molec.
 Cell.
 [100515]Biol. 18: 1601–1610, 1998.
 [100516]17492.
 [100517]Fischer, U.; Heckel, D.; Michel, A.; Janka, M.; Hulsebos, T.;
 Meese,
 [100518]E.: Cloning of a novel transcription factor-like gene am-

plified in

[100519]human glioma including astrocytoma grade I. Hum. Molec.

Genet. 6:

[100520]1817–1822, 1997.

[100521]17493.

[100522]Gracia, E.; Fischer, U.; Elkahouloun, A.; Trent, J. M.; Meese,

E.;

[100523]Meltzer, P. S.: Isolation of genes amplified in human can-

cers by

[100524]microdissection mediated cDNA capture. Hum. Molec.

Genet. 5: 595–600,

[100525]1996.

[100526]17494.

[100527]Gerard, M.; Hernandez, L.; Wevrick, R.; Stewart, C. L.: Dis-

ruption

[100528]of the mouse necdin gene results in early post-natal

lethality. Nature

[100529]Genet. 23: 199–202, 1999.

[100530]17495.

[100531]Hurst, L. D.; McVean, G.; Moore, T.: Imprinted genes have

few

[100532]and small introns. (Letter) Nature Genet. 12: 234–237,
1996.

[100533]17496.

[100534]Jay, P.; Rougeulle, C.; Massacrier, A.; Moncla, A.; Mattei,
M.–G.;

[100535]Malzac, P.; Roeckel, N.; Taviaux, S.; Lefranc, J.–L. B.; Cau,
P.;

[100536]Berta, P.; Lalande, M.; Muscatelli, F.: The human necdin
gene, NDN,

[100537]is maternally imprinted and located in the Prader–Willi
syndrome chromosomal

[100538]region. Nature Genet. 17: 357–360, 1997.

[100539]17497.

[100540]MacDonald, H. R.; Wevrick, R.: The necdin gene is deleted
in Prader–Willi

[100541]syndrome and is imprinted in human and mouse. Hum.
Molec. Genet. 6:

[100542]1873–1878, 1997.

[100543]17498.

[100544]Maruyama, K.; Usami, M.; Aizawa, T.; Yoshikawa, K.: A
novel brain–specific

[100545] mRNA encoding nuclear protein (necdin) expressed in neurally differentiated embryonal carcinoma cells. *Biochem. Biophys. Res. Comm.* 178: 291–296, 1991.

[100547] 17499.

[100548] Muscatelli, F.; Arous, D. N.; Massacrier, A.; Boccaccio, I.; Le Moal, M.; Cau, P.; Cremer, H.: Disruption of the mouse *Necdin* gene results in hypothalamic and behavioral alterations reminiscent of the human Prader–Willi syndrome. *Hum. Molec. Genet.* 9: 3101–3110, 2000.

[100553] 17500.

[100554] Nakada, Y.; Taniura, H.; Uetsuki, T.; Inazawa, J.; Yoshikawa, K. The human chromosomal gene for necdin, a neuronal growth suppressor, in the Prader–Willi syndrome deletion region. *Gene* 213: 65–72, 1998.

[100558]17501.

[100559]Nicholls, R. D.: Incriminating gene suspects, Prader-Willi style. Nature

[100560]Genet. 23: 132-134, 1999.

[100561]17502.

[100562]Tsai, T.-F.; Armstrong, D.; Beaudet, A. L.: Necdin-deficient mice

[100563]do not show lethality or the obesity and infertility of Prader-Willi

[100564]syndrome. (Letter) Nature Genet. 22: 15-16, 1999.

[100565]17503.

[100566]Watrin, F.; Roeckel, N.; Lacroix, L.; Mignon, C.; Mattei, M.-G.;

[100567]Disteche, C.; Muscatelli, F.: The mouse necdin gene is expressed

[100568]from the paternal allele only and lies in the 7C region of the mouse

[100569]chromosome 7, a region of conserved synteny to the human Prader-Willi

[100570]syndrome region. Europ. J. Hum. Genet. 5: 324-332, 1997.

[100571]17504.

[100572]Woodage, T.; Basrai, M. A.; Baxevanis, A. D.; Hieter, P.;
Collins,

[100573]F. S.: Characterization of the CHD family of proteins. Proc.
Nat.

[100574]Acad. Sci. 94: 11472–11477, 1997.

[100575]17505.

[100576]Seelig, H. P.; Renz, M.; Targoff, I. N.; Ge, Q.; Frank, M. B.:

[100577]Two forms of the major antigenic protein of the dermato-
myositis-specific

[100578]Mi-2 autoantigen. (Letter) Arthritis Rheum. 39:
1769–1771, 1996.

[100579]17506.

[100580]Ernstsson, S.; Betz, R.; Lagercrantz, S.; Larsson, C.; Ericks-
son,

[100581]S.; Cederberg, A.; Carlsson, P.; Enerback, S.: Cloning and
characterization

[100582]of freac-9 (FKHL17), a novel kidney-expressed human
forkhead gene

[100583]that maps to chromosome 1p32–p34. Genomics 46:
78–85, 1997.

[100584]17507.

[100585]Ozkaynak, E.; Schnegelsberg, P. N. J.; Jin, D. F.; Clifford, G.

[100586]M.; Warren, F. D.; Drier, E. A.; Oppermann, H.: Osteogenic
protein-2:

[100587]a new member of the transforming growth factor-beta
superfamily expressed

[100588]in early embryogenesis. J. Biol. Chem. 267: 25220-25227,
1992.

[100589]17508.

[100590]Kinoh, H.; Hayashita, H.; Kajita, M.; Okada, A.; Seiki, M.:
Assignment

[100591]of the genes for membrane-type-4 matrix metallopro-
teinase (Mmp17,

[100592]MMP17) to mouse chromosome 5, human chromosome
band 12q24.3 and membrane-type-5

[100593]matrix metalloproteinase (Mmp24, MMP24) to mouse
chromosome 2 and

[100594]human chromosome band 20q11.2-q12, respectively, by
radiation hybrid

[100595]and in situ hybridization. Cytogenet. Cell Genet. 87:
97-98, 1999.

[100596]17509.

[100597]Bingle, C. D.; Gowan, S.: Molecular cloning of the forkhead transcription

[100598]factor HNF-3- α from a human pulmonary adenocarcinoma cell line. Biochim.

[100599]Biophys. Acta 1307: 17-20, 1996.

[100600]17510.

[100601]Hannigan, G. E.; Bayani, J.; Weksberg, R.; Beatty, B.; Pandita,

[100602]A.; Dedhar, S.; Squire, J.: Mapping of the gene encoding the integrin-linked

[100603]kinase, ILK, to human chromosome 11p15.5-p15.4. Genomics 42: 177-179,

[100604]1997.

[100605]17511.

[100606]Hannigan, G. E.; Leung-Hagesteijn, C.; Fitz-Gibbon, L.; Coppolino,

[100607]M. G.; Radeva, G.; Filmus, J.; Bell, J. C.; Dedhar, S.: Regulation

[100608]of cell adhesion and anchorage-dependent growth by a new beta-1-integrin-linked

[100609]protein kinase. Nature 379: 91-96, 1996.

[100610]17512.

[100611]Omeis, I. A.; Hsu, Y.-C.; Perin, M. S.: Mouse and human neuronal

[100612]pentraxin 1 (NPXT1): conservation, genomic structure, and chromosomal

[100613]localization. Genomics 36: 543–545, 1996.

[100614]17513.

[100615]Kubota, N.; Terauchi, Y.; Miki, H.; Tamemoto, H.; Yamauchi, T.;

[100616]Komeda, K.; Satoh, S.; Nakano, R.; Ishii, C.; Sugiyama, T.; Eto, K.;

[100617]Tsubamoto, Y.; and 17 others: PPAR- γ mediates high-fat diet-induced

[100618]adipocyte hypertrophy and insulin resistance. Molec. Cell 4: 597–609,

[100619]1999.

[100620]17514.

[100621]Lapsys, N. M.; Kriketos, A. D.; Lim-Fraser, M.; Poynten, A. M.;

[100622]Lowy, A.; Furler, S. M.; Chisholm, D. J.; Cooney, G. J.: Expression

[100623] of genes involved in lipid metabolism correlate with peroxisome proliferator-activated

[100624] receptor gamma expression in human skeletal muscle. J. Clin. Endocr.

[100625] Metab. 85: 4293–4297, 2000.

[100626] 17515.

[100627] Lehmann, J. M.; Moore, L. B.; Smith–Oliver, T. A.; Wilkison, W.

[100628] O.; Willson, T. M.; Kliewer, S. A.: An antidiabetic thiazolidinedione

[100629] is a high affinity ligand for peroxisome proliferator-activated receptor

[100630] gamma (PPAR gamma). J. Biol. Chem. 270: 12953–12956, 1995.

[100631] 17516.

[100632] Lowell, B. B.: PPAR-gamma: an essential regulator of adipogenesis

[100633] and modulator of fat cell function. Cell 99: 239–242, 1999.

[100634] 17517.

[100635] Martin, G.; Schoonjans, K.; Staels, B.; Auwerx, J.: PPAR-gamma

- [100636]activators improve glucose homeostasis by stimulating fatty acid uptake
- [100637]in the adipocytes. *Atherosclerosis* 137: S75–S80, 1998.
- [100638]17518.
- [100639]Meirhaeghe, A.; Fajas, L.; Helbecque, N.; Cottel, D.; Lebel, P.;
- [100640]Dallongeville, J.; Deeb, S.; Auwerx, J.; Amouyel, P.: A genetic polymorphism
- [100641]of the peroxisome proliferator–activated receptor gamma gene influences
- [100642]plasma leptin levels in obese tumors. *Hum. Molec. Genet.* 7: 435–440,
- [100643]1998.
- [100644]17519.
- [100645]Miles, P. D. G.; Barak, Y.; He, W.; Evans, R. M.; Olefsky, J.
- [100646]M.: Improved insulin–sensitivity in mice heterozygous for PPAR–gamma
- [100647]deficiency. *J. Clin. Invest.* 105: 287–292, 2000.
- [100648]17520.
- [100649]Mueller, E.; Sarraf, P.; Tontonoz, P.; Evans, R. M.; Martin, K.

[100650]J.; Zhang, M.; Fletcher, C.; Singer, S.; Spiegelman, B. M.:

Terminal

[100651]differentiation of human breast cancer through PPAR-
gamma. Molec.

[100652]Cell. 1: 465-470, 1998.

[100653]17521.

[100654]Mueller, E.; Smith, M.; Sarraf, P.; Kroll, T.; Aiyer, A.; Kauf-
man,

[100655]D. S.; Oh, W.; Demetri, G.; Figg, W. D.; Zhou, X.-P.; Eng,
C.; Spiegelman,

[100656]B. M.; Kantoff, P. W.: Effects of ligand activation of peroxi-
some

[100657]proliferator-activated receptor gamma in human prostate
cancer. Proc.

[100658]Nat. Acad. Sci. 97: 10990-10995, 2000.

[100659]17522.

[100660]Nagy, L.; Tontonoz, P.; Alvarez, J. G. A.; Chen, H.; Evans,
R.

[100661]M.: Oxidized LDL regulates macrophage gene expression
through ligand

[100662]activation of PPAR-gamma. Cell 93: 229-240, 1998.

[100663]17523.

[100664]Oh, E. Y.; Min, K. M.; Chung, J. H.; Min, Y.-K.; Lee, M.-S.;
Kim,

[100665]K.-W.; Lee, M.-K.: Significance of pro12ala mutation in
peroxisome

[100666]proliferator-activated receptor-gamma2 in Korean dia-
betic and obese

[100667]subjects. J. Clin. Endocr. Metab. 85: 1801-1804, 2000.

[100668]17524.

[100669]Pritchard, J. K.; Rosenberg, N. A.: Use of unlinked genetic
markers

[100670]to detect population stratification in association studies.
Am. J.

[100671]Hum. Genet. 65: 220-228, 1999.

[100672]17525.

[100673]Ristow, M.; Muller-Wieland, D.; Pfeiffer, A.; Krone, W.;
Kahn,

[100674]C. R.: Obesity associated with a mutation in a genetic reg-
ulator

[100675]of adipocyte differentiation. New Eng. J. Med. 339:
953-959, 1998.

[100676]17526.

[100677]Rocchi, S.; Picard, F.; Vamecq, J.; Gelman, L.; Potier, N.; Zeyer,

[100678]D.; Dubuquoy, L.; Bac, P.; Champy, M.-F.; Plunket, K. D.; Leesnitzer,

[100679]L. M.; Blanchard, S. G.; Desreumaux, P.; Moras, D.; Renaud, J.-P.;

[100680]Auwerx, J.: A unique PPAR- γ ligand with potent insulin-sensitizing

[100681]yet weak adipogenic activity. *Molec. Cell* 8: 737-747, 2001.

[100682]17527.

[100683]Rosen, E. D.; Sarraf, P.; Troy, A. E.; Bradwin, G.; Moore, K.;

[100684]Milstone, D. S.; Spiegelman, B. M.; Mortensen, R. M.: PPAR- γ

[100685]is required for the differentiation of adipose tissue in vivo and

[100686]in vitro. *Molec. Cell* 4: 611-617, 1999.

[100687]17528.

[100688]Ricote, M.; Huang, J.; Fajas, L.; Li, A.; Welch, J.; Najib, J.;

[100689]Witztum, J. L.; Auwerx, J.; Palinski, W.; Glass, C. K.: Ex-

pression

[100690]of the peroxisome proliferator-activated receptor gamma
(PPAR-gamma)

[100691]in human atherosclerosis and regulation in macrophages
by colony stimulating

[100692]factors and oxidized low density lipoprotein. Proc. Nat.
Acad. Sci. 95:

[100693]7614-7619, 1998.

[100694]17529.

[100695]Savage, D. B.; Agostini, M.; Barroso, I.; Gurnell, M.; Luan,
J.;

[100696]Meirhaeghe, A.; Harding, A.-H.; Ihrke, G.; Rajanayagam,
O.; Soos,

[100697]M. A.; George, S.; Berger, D.; and 9 others: Digenic inheri-
tance

[100698]of severe insulin resistance in a human pedigree. Nature
Genet. 31:

[100699]379-384, 2002. Note: Erratum: Nature Genet. 32: 211
only, 2002.

[100700]17530.

[100701]Smith, W. M.; Zhou, X.-P.; Kurose, K.; Gao, X.; Latif, F.;
Kroll,

[100702]T.; Sugano, K.; Cannistra, S. A.; Clinton, S. K.; Maher, E. R.;
Prior,
[100703]T. W.; Eng, C.: Opposite association of two PPARG variants
with cancer:
[100704]overrepresentation of H449H in endometrial carcinoma
cases and underrepresentation
[100705]of P12A in renal cell carcinoma cases. Hum. Genet. 109:
146–151,
[100706]2001.
[100707]17531.
[100708]Tarrade, A.; Schoonjans, K.; Pavan, L.; Auwerx, J.; Ro-
chette-Egly,
[100709]C.; Evain-Brion, D.; Fournier, T.: PPAR-gamma/RXR-alpha
heterodimers
[100710]control human trophoblast invasion. J. Clin. Endocr.
Metab. 86:
[100711]5017–5024, 2001.
[100712]17532.
[100713]Tontonoz, P.; Hu, E.; Devine, J.; Beale, E. G.; Spiegelman,
B.
[100714]M.: PPAR gamma 2 regulates adipose expression of the
phosphoenolpyruvate

[100715]carboxykinase gene. Molec. Cell. Biol. 15: 351–357, 1995.

[100716]17533.

[100717]Tontonoz, P.; Hu, E.; Graves, R. A.; Budavari, A. I.; Spiegelman,

[100718]B. M.: mPPAR gamma 2: tissue-specific regulator of an adipocyte enhancer. Genes

[100719]Dev. 8: 1224–1234, 1994.

[100720]17534.

[100721]Tontonoz, P.; Nagy, L.; Alvarez, J. G. A.; Thomazy, V. A.; Evans,

[100722]R. M.: PPAR-gamma promotes monocyte/macrophage differentiation and

[100723]uptake of oxidized LDL. Cell 93: 241–252, 1998.

[100724]17535.

[100725]Wang, M.: Isolation and characterization of a human gene encoding

[100726]a single-strand-specific endoribonuclease. Ph.D. Thesis: Stanford

[100727]Univ. , 1995.

[100728]17536.

- [100729]Wang, M.; Cohen, S. N.: ard-1: a human gene that re-
verses the
- [100730]effects of temperature-sensitive and deletion mutations in
the Escherichia
- [100731]coli rne gene and encodes an activity producing RNase E-
like cleavages. Proc.
- [100732]Nat. Acad. Sci. 91: 10591-10595, 1994.
- [100733]17537.
- [100734]Amendt, B. A.; Sutherland, L. B.; Semina, E. V.; Russo, A.
F.:
- [100735]The molecular basis of Rieger syndrome: analysis of PITX2
homeodomain
- [100736]protein activities. J. Biol. Chem. 273: 20066-20072, 1998.
- [100737]17538.
- [100738]Arakawa, H.; Nakamura, T.; Zhadanov, A. B.; Fidanza, V.;
Yano,
- [100739]T.; Bullrich, F.; Shimizu, M.; Blechman, J.; Mazo, A.;
Canaani, E.;
- [100740]Croce, C. M.: Identification and characterization of the
ARP1 gene,
- [100741]a target for the human acute leukemia ALL1 gene. Proc.
Nat. Acad.

[100742]Sci. 95: 4573–4578, 1998.

[100743]17539.

[100744]Cushman, L. J.; Camper, S. A.: Molecular basis of pituitary dysfunction

[100745]in mouse and human. Mammalian Genome 12: 485–494, 2001.

[100746]17540.

[100747]Doward, W.; Perveen, R.; Lloyd, I. C.; Ridgway, A. E. A.; Wilson,

[100748]L.; Black, G. C. M.: A mutation in the RIEG1 gene associated with

[100749]Peters' anomaly. J. Med. Genet. 36: 152–155, 1999.

[100750]17541.

[100751]Flomen, R. H.; Vatcheva, R.; Gorman, P. A.; Baptista, P. R.; Groet,

[100752]J.; Barisic, I.; Ligutic, I.; Nizetic, D.: Construction and analysis

[100753]of a sequence-ready map in 4q25: Rieger syndrome can be caused by

[100754]haploinsufficiency of RIEG, but also by chromosome breaks approximately

[100755]90 kb upstream of this gene. Genomics 47: 409–413,
1998.

[100756]17542.

[100757]Gage, P. J.; Camper, S. A.: Pituitary homeobox 2, a novel
member

[100758]of the bicoid-related family of homeobox genes, is a po-
tential regulator

[100759]of anterior structure formation. Hum. Molec. Genet. 6:
457–464,

[100760]1997.

[100761]17543.

[100762]Kozak, M.: Interpreting cDNA sequences: some insights
from studies

[100763]on translation. Mammalian Genome 7: 563–574, 1996.

[100764]17544.

[100765]Kozlowski, K.; Walter, M. A.: Variation in residual PITX2
activity

[100766]underlies the phenotypic spectrum of anterior segment
developmental

[100767]disorders. Hum. Molec. Genet. 9: 2131–2139, 2000.

[100768]17545.

[100769]Lin, C. R.; Kioussi, C.; O'Connell, S.; Briata, P.; Szeto, D.;

[100770]Liu, F.; Izpisua-Belmonte, J. C.; Rosenfeld, M. G.: Pitx2
regulates

[100771]lung asymmetry, cardiac positioning and pituitary and
tooth morphogenesis. Nature 401:

[100772]279-282, 1999.

[100773]17546.

[100774]Logan, M.; Pagan-Westphal, S. M.; Smith, D. M.; Paganessi,
L.;

[100775]Tabin, C. J.: The transcription factor Pitx2 mediates situs-
specific

[100776]morphogenesis in response to left-right asymmetric sig-
nals. Cell 94:

[100777]307-317, 1998.

[100778]17547.

[100779]Lu, M.-F.; Pressman, C.; Dyer, R.; Johnson, R. L.; Martin, J.

[100780]F.: Function of Rieger syndrome gene in left-right asym-
metry and

[100781]craniofacial development. Nature 401: 276-278, 1999.

[100782]17548.

[100783]Martin, D. M.; Probst, F. J.; Fox, S. E.; Schimmenti, L. A.;
Semina,

[100784]E. V.; Hefner, M. A.; Belmont, J. W.; Camper, S. A.: Exclu-
sion of

[100785]PITX2 mutations as a major cause of CHARGE association.
Am. J. Med.

[100786]Genet. 111: 27–30, 2002.

[100787]17549.

[100788]Pellegrini-Bouiller, I.; Manrique, C.; Gunz, G.; Grino, M.;
Zamora,

[100789]A. J.; Figarella-Branger, D.; Grisoli, F.; Jaquet, P.; Enjalbert,

[100790]A.: Expression of the members of the Ptx family of tran-
scription

[100791]factors in human pituitary adenomas. J. Clin. Endocr.
Metab. 84:

[100792]2212–2220, 1999.

[100793]17550.

[100794]Piedra, M. E.; Icardo, J. M.; Albajar, M.; Rodriguez-Rey, J.
C.;

[100795]Ros, M. A.: Pitx2 participates in the late phase of the
pathway controlling

[100796]left-right asymmetry. Cell 94: 319–324, 1998.

[100797]17551.

[100798]Priston, M.; Kozlowski, K.; Gill, D.; Letwin, K.; Buys, Y.;
Levin,

[100799]A. V.; Walter, M. A.; Heon, E.: Functional analyses of two
newly

[100800]identified PITX2 mutants reveal a novel molecular mecha-
nism for Axenfeld-Rieger

[100801]syndrome. Hum. Molec. Genet. 10: 1631–1638, 2001.

[100802]17552.

[100803]Bayever, E.; Ladisch, S.; Philippart, M.; Brill, N.; Nuwer, M.;

[100804]Sparkes, R. S.; Feig, S. A.: Bone-marrow transplantation
for metachromatic

[100805]leucodystrophy. Lancet II: 471–473, 1985.

[100806]17553.

[100807]Beratis, N. G.; Danesino, C.; Hirschhorn, K.: Detection of
homozygotes

[100808]and heterozygotes for metachromatic leukodystrophy in
lymphoid cell

[100809]lines and peripheral leukocytes. Ann. Hum. Genet. 38:
485–493, 1975.

[100810]17554.

[100811]Berger, J.; Gmach, M.; Mayr, U.; Molzer, B.; Bernheimer, H.:

[100812]Coincidence of two novel arylsulfatase A alleles and mutation 459+1G→A

[100813]within a family with metachromatic leukodystrophy: molecular basis

[100814]of phenotypic heterogeneity. Hum. Mutat. 13: 61–68, 1999.

[100815]17555.

[100816]Betts, T. A.; Smith, W. T.; Kelly, R. E.: Adult metachromatic

[100817]leukodystrophy (sulphatide lipidosis) simulating acute schizophrenia:

[100818]report of a case. Neurology 18: 1140–1142, 1968.

[100819]17556.

[100820]Black, J. W.; Cumings, J. N.: Infantile metachromatic leukodystrophy. J.

[100821]Neurol. Neurosurg. Psychiat. 24: 233–239, 1961.

[100822]17557.

[100823]Bohne, W.; von Figura, K.; Gieselmann, V.: An 11-bp deletion

[100824]in the arylsulfatase A gene of a patient with late infantile metachromatic

[100825]leukodystrophy. Hum. Genet. 87: 155–158, 1991.

[100826]17558.

[100827]Bosch, E. P.; Hart, M. N.: Late adult-onset metachromatic leukodystrophy:

[100828]dementia and polyneuropathy in a 63-year-old man. Arch. Neurol. 35:

[100829]475–477, 1978.

[100830]17559.

[100831]Bruns, G. A. P.; Mintz, B. J.; Leary, A. C.; Regina, V. M.; Gerald,

[100832]P. S.: Expression of human arylsulfatase A in man-hamster somatic

[100833]cell hybrids. Cytogenet. Cell Genet. 22: 182–185, 1978.

[100834]17560.

[100835]Butterworth, J.; Broadhead, D. M.; Keay, A. J.: Low arylsulphatase

[100836]A activity in a family without metachromatic leukodystrophy. Clin.

[100837]Genet. 14: 213–218, 1978.

[100838]17561.

[100839]Caillaud, C.; Blanchor, C.; Akli, S.; Crosnier, J. M.; Puech,

[100840]J. P.; Kahn, A.; Poenaru, L.: Molecular basis of late infantile metachromatic

[100841]leukodystrophy in France. (Abstract) ESGLD Workshop Delphi, and

[100842]Greece, and Oct, 1993.

[100843]17562.

[100844]Chabas, A.; Castellvi, S.; Bayes, M.; Balcells, S.; Grinberg,

[100845]D.; Vilageliu, L.; Marfany, G.; Lissens, W.; Gonzalez-Duarte, R.:

[100846]Frequency of the arylsulphatase A pseudodeficiency allele in the Spanish

[100847]population. Clin. Genet. 44: 320-323, 1993.

[100848]17563.

[100849]Chang, P. L.; Davidson, R. G.: Pseudo arylsulphatase-A deficiency

[100850]in healthy individuals: genetic and biochemical relationship to metachromatic

[100851]leukodystrophy. Proc. Nat. Acad. Sci. 80: 7323-7327, 1983.

[100852]17564.

[100853]Chang, P. L.; Davidson, R. G.: Complementation of arylsul-
fatase

[100854]A in somatic hybrids of metachromatic leukodystrophy
and multiple

[100855]sulfatase deficiency disorder fibroblasts. Proc. Nat. Acad.
Sci. 77:

[100856]6166-6170, 1980.

[100857]17565.

[100858]Chang, P. L.; Rosa, N. E.; Davidson, R. G.: Somatic cell hy-
bridization

[100859]studies on the genetic regulation and allelic mutations in
metachromatic

[100860]leukodystrophy. Hum. Genet. 61: 231-235, 1982.

[100861]17566.

[100862]Coulter-Mackie, M. B.; Rip, J.; Ludman, M. D.; Beis, J.;
Cole,

[100863]D. E. C.: Metachromatic leucodystrophy (MLD) in a patient
with a

[100864]constitutional ring chromosome 22. J. Med. Genet. 32:
787-791, 1995.

[100865]17567.

[100866]Cravioto, H.; O'Brien, J.; Lockwood, R.; Kasten, F. H.;
Booker,

[100867]J.: Metachromatic leukodystrophy (sulfatide lipidoses) cul-
tured in

[100868]vitro. Science 156: 243-245, 1967.

[100869]17568.

[100870]DeLuca, C.; Brown, J. A.; Shows, T. B.: Lysosomal arylsul-
fatase

[100871]deficiencies in humans: chromosome assignment of aryl-
sulfatase A and

[100872]B. Proc. Nat. Acad. Sci. 76: 1957-1961, 1979.

[100873]17569.

[100874]DeLuca, C.; Champion, M. J.; Shows, T. B.: Arylsulfatase-A
(ARSA)

[100875]synteny with beta-glucuronidase (BGUS) indicates assign-
ment to human

[100876]chromosome 7 in man-Chinese hamster hybrids.
(Abstract) Winnipeg

[100877]Gene Mapping Conf. , 1977.

[100878]17570.

[100879] Draghia, R.; Letourneur, F.; Drugan, C.; Manicom, J.; Blanchot,

[100880] C.; Kahn, A.; Poenaru, L.; Caillaud, C.: Metachromic leukodystrophy:

[100881] identification of the first deletion in exon 1 and of nine novel point

[100882] mutations in the arylsulfatase gene. Hum. Mutat. 9: 234-242, 1997.

[100883] 17571.

[100884] Dubois, G.; Harzer, K.; Baumann, N.: Very low arylsulfatase A

[100885] and cerebroside sulfatase activities in leukocytes of healthy members

[100886] of metachromatic leukodystrophy family. Am. J. Hum. Genet. 29: 191-194,

[100887] 1977.

[100888] 17572.

[100889] Eto, Y.; Tahara, T.; Koda, N.; Yamaguchi, S.; Ito, F.; Okuno, A.: Prenatal diagnosis of metachromatic leukodystrophy: a diagnosis

[100891] by amniotic fluid and its confirmation. Arch. Neurol. 39: 29-32,

[100892]1982.

[100893]17573.

[100894]Farrell, D. F.: Heterozygote detection in MLD: allelic mutations

[100895]at the ARA locus. Hum. Genet. 59: 129–134, 1981.

[100896]17574.

[100897]Farrell, D. F.; MacMartin, M. P.; Clark, A. F.: Multiple
molecular
[100898]forms of arylsulfatase A in different forms of metachro-
matic leukodystrophy
[100899](MLD). Neurology 29: 16–20, 1979.
[100900]17575.

[100901]Felice, K. J.; Gomez Lira, M.; Natowicz, M.; Grunnet, M. L.;
Tsongalis,
[100902]G. J.; Sima, A. A. F.; Kaplan, R. F.: Adult-onset MLD: a
gene mutation
[100903]with isolated polyneuropathy. Neurology 55: 1036–1039,
2000.
[100904]17576.

[100905]Fluharty, A. L.; Fluharty, C. B.; Bohne, W.; von Figura, K.;
Gieselmann,
[100906]V.: Two new arylsulfatase A (ARSA) mutations in a juvenile
metachromatic
[100907]leukodystrophy (MLD) patient. Am. J. Hum. Genet. 49:
1340–1350,
[100908]1991.
[100909]17577.

[100910] Francke, U.; Tetri, P.; Taggart, R. T.; Oliver, N.: Conserved
[100911] autosomal syntenic group on mouse (MMU) chromosome
15 and human (HSA)
[100912] chromosome 22: assignment of a gene for arylsulfatase A
to MMU 15
[100913] and regional mapping of DIA1, ARSA, and ACO2 on
HSA22. Cytogenet.
[100914] Cell Genet. 31: 58–69, 1981.
[100915] 17578.
[100916] Geurts van Kessel, A. H. M.; Westerveld, A.; de Groot, P.
G.;
[100917] Meera Khan, P.; Hagemeijer, A.: Regional localization of
the genes
[100918] coding for human ACO2, ARSA, and NAGA on chromo-
some 22. Cytogenet.
[100919] Cell Genet. 28: 169–172, 1980.
[100920] 17579.
[100921] Gieselmann, V.: An assay for the rapid detection of the
arylsulfatase
[100922] A pseudodeficiency allele facilitates diagnosis and genetic
counseling

[100923]for metachromatic leukodystrophy. Hum. Genet. 86:
251–255, 1991.

[100924]17580.

[100925]Gieselmann, V.; Fluharty, A. L.; Tonnesen, T.; Von Figura, K.

[100926]: Mutations in the arylsulfatase A pseudodeficiency allele
causing

[100927]metachromatic leukodystrophy. Am. J. Hum. Genet. 49:
407–413, 1991.

[100928]17581.

[100929]Gieselmann, V.; Polten, A.; Kreysing, J.; von Figura, K.:
Arylsulfatase

[100930]A pseudodeficiency: loss of a polyadenylation (sic) signal
and N-glycosylation

[100931]site. Proc. Nat. Acad. Sci. 86: 9436–9440, 1989.

[100932]17582.

[100933]Gieselmann, V.; Zlotogora, J.; Harris, A.; Wenger, D. A.;
Morris,

[100934]C. P.: Molecular genetics of metachromatic leukodystro-
phy. Hum.

[100935]Mutat. 4: 233–242, 1994.

[100936]17583.

[100937]Goebel, H. H.; Pilz, H.; Argyrakis, A.: Adult metachromatic leukodystrophy.

[100938]II. Ultrastructural findings in peripheral nerve and skeletal muscle. Europ.

[100939]Neurol. 15: 308–317, 1977.

[100940]17584.

[100941]Beck, M.; Glossl, J.; Grubisic, A.; Spranger, J.: Heterogeneity

[100942]of Morquio disease. Clin. Genet. 29: 325–331, 1986.

[100943]17585.

[100944]Maroteaux, P.; Stanescu, V.; Stanescu, R.; Kresse, H.; Hors-Cayla,

[100945]M. C.: Heterogeneite des formes frustes de la maladie de Morquio. Arch.

[100946]Franc. Pediat. 39: 761–765, 1982.

[100947]17586.

[100948]Saunders, M.; Sweetman, L.; Robinson, B.; Roth, K.; Kohn, S.;

[100949]Sherwood, G.; Gravel, R.: Multiple carboxylase defects and complementation

[100950] studies in biotin responsive organic aciduria. (Abstract)
Am. J. Hum.

[100951] Genet. 31: 61A, 1979.

[100952] 17587.

[100953] Saunders, M. E.; Sherwood, W. G.; Duthie, M.; Surh, L.;
Gravel,

[100954] R. A.: Evidence for a defect of holocarboxylase synthetase
activity

[100955] in cultured lymphoblasts from a patient with biotin-
responsive multiple

[100956] carboxylase deficiency. Am. J. Hum. Genet. 34: 590-601,
1982.

[100957] 17588.

[100958] Suormala, T.; Fowler, B.; Duran, M.; Burtcher, A.;
Fuchshuber,

[100959] A.; Tratzmuller, R.; Lenze, M. J.; Raab, K.; Baur, B.; Wick,
H.; Baumgartner,

[100960] R.: Five patients with a biotin-responsive defect in holo-
carboxylase

[100961] formation: evaluation of responsiveness to biotin therapy
in vivo

[100962] and comparative biochemical studies in vitro. Pediat. Res.

41: 666-673,

[100963]1997.

[100964]17589.

[100965]Suzuki, Y.; Aoki, Y.; Ishida, Y.; Chiba, Y.; Iwamatsu, A.;
Kishino,

[100966]T.; Niikawa, N.; Matsubara, Y.; Narisawa, K.: Isolation and
characterization

[100967]of mutations in the human holocarboxylase synthetase
cDNA. Nature

[100968]Genet. 8: 122-128, 1994.

[100969]17590.

[100970]Sweetman, L.; Surh, L.; Baker, H.; Peterson, R. M.; Nyhan,
W.

[100971]L.: Clinical and metabolic abnormalities in a boy with di-
etary deficiency

[100972]of biotin. Pediatrics 68: 553-558, 1981.

[100973]17591.

[100974]Thoene, J.; Baker, H.; Yoshino, M.; Sweetman, L.: Biotin-
responsive

[100975]carboxylase deficiency associated with subnormal plasma
and urinary

[100976]biotin. New Eng. J. Med. 304: 817–820, 1981.

[100977]17592.

[100978]Thoene, J.; Sweetman, L.; Yoshino, M.: Biotin–responsive
multiple

[100979]carboxylase deficiency. (Abstract) Am. J. Hum. Genet. 31:
64A, 1979.

[100980]17593.

[100981]Wolf, B.; Grier, R. E.; Parker, W. D.; Goodman, S. I.; Allen,

[100982]R. J.: Deficient biotinidase activity in late–onset multiple
carboxylase

[100983]deficiency. (Letter) New Eng. J. Med. 308: 161, 1983.

[100984]17594.

[100985]Yang, X.; Aoki, Y.; Li, X.; Sakamoto, O.; Hiratsuka, M.; Gib-
son,

[100986]K. M.; Kure, S.; Narisawa, K.; Matsubara, Y.; Suzuki, Y.:
Haplotype

[100987]analysis suggests that the two predominant mutations in
Japanese patients

[100988]with holocarboxylase synthetase deficiency are founder
mutations. J.

[100989]Hum. Genet. 45: 358–362, 2000.

[100990]17595.

[100991]Yang, X.; Aoki, Y.; Li, X.; Sakamoto, O.; Hiratsuka, M.;
Kure,

[100992]S.; Taheri, S.; Christensen, E.; Inui, K.; Kubota, M.; Ohira,
M.;

[100993]Ohki, M.; and 10 others: Structure of human holocarboxy-
lase synthetase

[100994]gene and mutation spectrum of holocarboxylase syn-
thetase deficiency. Hum.

[100995]Genet. 109: 526-534, 2001.

[100996]17596.

[100997]Zhang, X. X.; Leon-Del-Rio, A.; Gravel, R. A.; Eydoux, P.:
Assignment

[100998]of holocarboxylase synthetase gene (HLCS) to human
chromosome band

[100999]21q22.1 and to mouse chromosome band 16C4 by in situ
hybridization. Cytogenet.

[101000]Cell Genet. 76: 179 only, 1997.

[101001]17597.

[101002]Mangos, J. A.; Opitz, J. M.; Lobeck, C. C.; Cookson, D. V.:
Familial

[101003]juvenile nephronophthisis: an unrecognized renal disease
in the United

[101004]States. Pediatrics 34: 337-345, 1964.

[101005]17598.

[101006]McKusick, V. A.: Mendelian Inheritance in Man. A Catalog
of Human

[101007]Genes and Genetic Disorders. (12th ed.) Baltimore: Johns
Hopkins

[101008]Univ. Press , 1998. Pp. I only. Note: Table 15.

[101009]17599.

[101010]Medhioub, M.; Cherif, D.; Benessy, F.; Silbermann, F.;
Gubler,

[101011]M. C.; Le Paslier, D.; Cohen, D.; Weissenbach, J.; Beck-
mann, J.; Antignac,

[101012]C.: Refined mapping of a gene (NPH1) causing familial ju-
venile nephronophthisis

[101013]and evidence for genetic heterogeneity. Genomics 22:
296-301, 1994.

[101014]17600.

[101015]Meier, D. A.; Hess, J. W.: Familial nephropathy with retini-
tis

[101016]pigmentosa: a new oculorenal syndrome in adults. Am. J. Med. 39:

[101017]58–69, 1965.

[101018]17601.

[101019]Mongeau, J. G.; Worthen, H. G.: Nephronophthisis and medullary

[101020]cystic disease. Am. J. Med. 43: 345–355, 1967.

[101021]17602.

[101022]Saunier, S.; Calado, J.; Benessy, F.; Silbermann, F.; Heilig,

[101023]R.; Weissenbach, J.; Antignac, C.: Characterization of the NPHP1

[101024]locus: mutational mechanism involved in deletions in familial juvenile

[101025]nephronophthisis. Am. J. Hum. Genet. 66: 778–789, 2000.

[101026]17603.

[101027]Saunier, S.; Calado, J.; Heilig, R.; Silbermann, F.; Benessy,

[101028]F.; Morin, G.; Konrad, M.; Broyer, M.; Gubler, M.–C.; Weissenbach,

[101029]J.; Antignac, C.: A novel gene that encodes a protein with a putative

[101030]src homology 3 domain is a candidate gene for familial juvenile nephronophthisis. Hum.

[101031]Molec. Genet. 6: 2317-2323, 1997.

[101032]17604.

[101033]Sherman, F. E.; Studnicki, F. M.; Fetterman, G. H.: Renal lesions

[101034]of familial juvenile nephronophthisis examined by microdissection. Am.

[101035]J. Clin. Path. 55: 391-400, 1971.

[101036]17605.

[101037]Simon, E. A.; Cook, S.; Davisson, M. T.; D'Eustachio, P.; Guay-Woodford,

[101038]L. M.: The mouse congenital polycystic kidney (cpk) locus maps within

[101039]1.3 cM of the chromosome 12 marker D12Nyu2. Genomics 21: 415-418,

[101040]1994.

[101041]17606.

[101042]Steele, B. T.; Lirenman, D. S.; Beattie, G. W.: Nephronophthisis. Am.

[101043]J. Med. 68: 531-538, 1980.

[101044]17607.

[101045]Strauss, M. B.; Sommers, S. C.: Medullary cystic disease
and

[101046]familial juvenile nephronophthisis: clinical and pathologi-
cal identity. New

[101047]Eng. J. Med. 277: 863-864, 1967.

[101048]17608.

[101049]Sworn, M. J.; Eisinger, A. J.: Medullary cystic disease and
juvenile

[101050]nephronophthisis in separate members of the same fam-
ily. Arch. Dis.

[101051]Child. 47: 278-281, 1972.

[101052]17609.

[101053]Von Sydow, G.; Ranstrom, S.: Familial juvenile
nephronophthisis. Acta

[101054]Paediat. 51: 561-574, 1962.

[101055]17610.

[101056]Waldherr, R.; Lennert, T.; Weber, H. P.; Fodisch, H. J.;
Scharer,

[101057]K.: The nephronophthisis complex: a clinicopathologic
study in children. Virchows

[101058]Arch. A Path. Anat. Histol. 394: 235–254, 1982.

[101059]17611.

[101060]Bolk, S.; Puffenberger, E. G.; Hudson, J.; Morton, D. H.;
Chakravarti,

[101061]A.: Elevated frequency and allelic heterogeneity of con-
genital nephrotic

[101062]syndrome, Finnish type, in the Old Order Mennonites.
(Letter) Am.

[101063]J. Hum. Genet. 65: 1785–1790, 1999.

[101064]17612.

[101065]Anikster, Y.; Kleta, R.; Shaag, A.; Gahl, W. A.; Elpeleg, O.:

[101066]Type III 3-methylglutaconic aciduria (optic atrophy plus
syndrome,

[101067]or Costeff optic atrophy syndrome): identification of the
OPA3 gene

[101068]and its founder mutation in Iraqi Jews. Am. J. Hum. Genet.
69: 1218–1224,

[101069]2001.

[101070]17613.

[101071]Bourdon, V.; Philippe, C.; Labrune, O.; Amsallem, D.;
Arnould,

[101072]C.; Jonveaux, P.: A detailed analysis of the MECP2 gene:
prevalence

[101073]of recurrent mutations and gross DNA rearrangements in
Rett syndrome

[101074]patients. Hum. Genet. 108: 43–50, 2001.

[101075]17614.

[101076]Brinke, A.; Green, P. M.; Giannelli, F.: Characterization of
the

[101077]gene (VBP1) and transcript for the von Hippel–Lindau
binding protein

[101078]and isolation of the highly conserved murine homologue.
Genomics 45:

[101079]105–112, 1997.

[101080]17615.

[101081]Brinke, A.; Tagliavacca, L.; Naylor, J.; Green, P.; Gian-
grande,

[101082]P.; Giannelli, F.: Two chimaeric transcription units result
from

[101083]an inversion breaking intron 1 of the factor VIII gene and a
region

[101084]reportedly affected by reciprocal translocations in T-cell
leukaemia. Hum.

- [101085]Molec. Genet. 5: 1945–1951, 1996.
- [101086]17616.
- [101087]Clifford, S. C.; Walsh, S.; Hewson, K.; Green, E. K.; Brinke, A.;
- [101088]Green, P. M.; Gianelli, F.; Eng, C.; Maher, E. R.: Genomic organization
- [101089]and chromosomal localization of the human CUL2 gene and the role of
- [101090]von Hippel–Lindau tumor suppressor–binding protein (CUL2 and VBP1)
- [101091]mutation and loss in renal–cell carcinoma development. Genes Chromosomes
- [101092]Cancer 26: 20–28, 1999.
- [101093]17617.
- [101094]Hemberger, M.; Himmelbauer, H.; Neumann, H. P. H.; Plate, K. H.;
- [101095]Schwarzkopf, G.; Fundele, R.: Expression of the von Hippel–Lindau–binding
- [101096]protein–1 (Vbp1) in fetal and adult mouse tissues. Hum. Molec. Genet. 8:
- [101097]229–236, 1999.

[101098]17618.

[101099]Tsuchiya, H.; Iseda, T.; Hino, O.: Identification of a novel protein

[101100](VBP-1) binding to the von Hippel-Lindau (VHL) tumor suppressor gene

[101101]product. Cancer Res. 56: 2881-2885, 1996.

[101102]17619.

[101103]Aoyama, T.; Tsushima, K.; Souri, M.; Kamijo, T.; Suzuki, Y.; Shimozawa,

[101104]N.; Orii, T.; Hashimoto, T.: Molecular cloning and functional expression

[101105]of a human peroxisomal acyl-coenzyme A oxidase. Biochem. Biophys.

[101106]Res. Commun. 198: 1113-1118, 1994.

[101107]17620.

[101108]Fan, C.-Y.; Pan, J.; Chu, R.; Lee, D.; Kluckman, K. D.; Usuda,

[101109]N.; Singh, I.; Yeldandi, A. V.; Rao, M. S.; Maeda, N.; Reddy, J. K.

[101110]: Hepatocellular and hepatic peroxisomal alterations in mice with

[101111]a disrupted peroxisomal fatty acyl-coenzyme A oxidase gene. J. Biol.

[101112]Chem. 271: 24698–24710, 1996.

[101113]17621.

[101114]Fournier, B.; Saudubray, J.-M.; Benichou, B.; Lyonnet, S.; Munnich,

[101115]A.; Clevers, H.; Poll-The, B. T.: Large deletion of the peroxisomal

[101116]acyl-CoA oxidase gene in pseudoneonatal adrenoleukodystrophy. J.

[101117]Clin. Invest. 94: 526–531, 1994.

[101118]17622.

[101119]Goldfischer, S. L.; Collins, J.; Rapin, I.; Neumann, P.; Neglia,

[101120]W.; Spiro, A. J.; Ishii, T.; Roels, F.; Vamecq, F.; Vamecq, J.; Van

[101121]Hoof, F.: Deficiencies in several peroxisomal oxidative activities. J.

[101122]Pediat. 108: 25–32, 1986.

[101123]17623.

[101124]Moghrabi, N. N.; Dawson, D. B.; Bennett, M. J.: Assignment

of

[101125]the human peroxisomal palmitoyl-CoA oxidase gene to
chromosome 17q23-qter

[101126]by PCR technique. J. Inherit. Metab. Dis. 18: 705-709,
1995.

[101127]17624.

[101128]Naidu, S.; Hoefler, G.; Watkins, P. A.; Chen, W. W.; Moser,
A.

[101129]B.; Hoefler, S.; Rance, N. E.; Powers, J. M.; Beard, M.;
Green, W.

[101130]R.; Hashimoto, T.; Moser, H. W.: Neonatal seizures and re-
tardation

[101131]in a female with biochemical features of X-linked
adrenoleukodystrophy:

[101132]a possible new peroxisomal disease entity. Neurology 38:
1100-1107,

[101133]1988.

[101134]17625.

[101135]Poll-The, B. T.; Roels, F.; Ogier, H.; Scotto, J.; Vamecq, J.;

[101136]Schutgens, R. B. H.; Wanders, R. J. A.; van Roermund, C. W.
T.; van

[101137]Wiljand, M. J. A.; Schram, A. W.; Tager, J. M.; Saudubray,

J.-M.:

[101138]A new peroxisomal disorder with enlarged peroxisomes and a specific

[101139]deficiency of acyl-CoA oxidase (pseudo-neonatal adrenoleukodystrophy). Am.

[101140]J. Hum. Genet. 42: 422-434, 1988.

[101141]17626.

[101142]Reddy, J. K.; Rao, M. S.: Peroxisome proliferators and cancer:

[101143]mechanisms and implications. Trends Pharm. Sci. 7: 438-443, 1986.

[101144]17627.

[101145]Varanasi, U.; Chu, R.; Chu, S.; Espinosa, R.; LeBeau, M. M.; Reddy,

[101146]J. K.: Isolation of the human peroxisomal acyl-CoA oxidase gene:

[101147]organization, promoter analysis, and chromosomal localization. Proc.

[101148]Nat. Acad. Sci. 91: 3107-3111, 1994.

[101149]17628.

[101150]Watkins, P. A.; McGuinness, M. C.; Raymond, G. V.; Hicks,

B. A.;

[101151]Sisk, J. M.; Moser, A. B.; Moser, H. W.: Distinction between peroxisomal

[101152]bifunctional enzyme and acyl-CoA oxidase deficiencies.

Ann. Neurol. 38:

[101153]472-477, 1995.

[101154]17629.

[101155]Buyse, I. M.; Fang, P.; Hoon, K. T.; Amir, R. E.; Zoghbi, H.

Y.;

[101156]Roa, B. B.: Diagnostic testing for Rett syndrome by DHPLC and direct

[101157]sequencing analysis of the MECP2 gene: identification of several novel

[101158]mutations and polymorphisms. Am. J. Hum. Genet. 67: 1428-1436, 2000.

[101159]17630.

[101160]Cheadle, J. P.; Gill, H.; Fleming, N.; Maynard, J.; Kerr, A.; Leonard,

[101161]H.; Krawczak, M.; Cooper, D. N.; Lynch, S.; Thomas, N.; Hughes, H.;

[101162]Hulten, M.; Ravine, D.; Sampson, J. R.; Clarke, A.: Long-read sequence

- [101163]analysis of the MECP2 gene in Rett syndrome patients:
correlation
[101164]of disease severity with mutation type and location. Hum.
Molec.
[101165]Genet. 9: 1119–1129, 2000.
[101166]17631.
- [101167]Chen, R. Z.; Akbarian, S.; Tudor, M.; Jaenisch, R.: Defi-
ciency
[101168]of methyl–CpG binding protein–2 in CNS neurons results
in a Rett–like
[101169]phenotype in mice. Nature Genet. 27: 327–331, 2001.
[101170]17632.
- [101171]Claes, S.; Devriendt, K.; D'Adamo, P.; Meireleire, J.; Raey-
maekers,
[101172]P.; Toniolo, D.; Cassiman, J.–J.; Fryns, J.–P.: X–linked se-
vere mental
[101173]retardation and a progressive neurological disorder in a
Belgian family:
[101174]clinical and genetic studies. Clin. Genet. 52: 155–161,
1997.
[101175]17633.

[101176]Clayton-Smith, J.; Watson, P.; Ramsden, S.; Black, G. C. M.:
[101177]Somatic mutation in MECP2 as a non-fatal neurodevelopmental disorder
[101178]in males. Lancet 356: 830-832, 2000.
[101179]17634.
[101180]Couvert, P.; Bienvenu, T.; Aquaviva, C.; Poirier, K.;
Moraine,
[101181]C.; Gendrot, C.; Verloes, A.; Andres, C.; Le Fevre, A. C.;
Souville,
[101182]J.; Steffann, J.; des Portes, V.; Ropers, H.-H.; Yntema, H.
G.; Fryns,
[101183]J.-P.; Briault, S.; Chelly, J.; Cherif, B.: MECP2 is highly mutated
[101184]in X-linked mental retardation. Hum. Molec. Genet. 10:
941-946,
[101185]2001.
[101186]17635.
[101187]D'Esposito, M.; Quaderi, N. A.; Ciccodicola, A.; Bruni, P.;
Esposito,
[101188]T.; D'Urso, M.; Brown, S. D. M.: Isolation, physical mapping, and

- [101189]Northern analysis of the X-linked human gene encoding methyl CpG-binding
- [101190]protein, MECP2. Mammalian Genome 7: 533–535, 1996.
- [101191]17636.
- [101192]De Bona, C.; Zappella, M.; Hayek, G.; Meloni, I.; Vitelli, F.;
- [101193]Bruttini, M.; Cusano, R.; Loffredo, P.; Longo, I.; Renieri, A.:
Preserved
- [101194]speech variant is allelic of classic Rett syndrome. Europ. J.
Hum.
- [101195]Genet. 8: 325–330, 2000.
- [101196]17637.
- [101197]Dragich, J.; Houwink–Manville, I.; Schanen, C.: Rett syn-
drome:
- [101198]a surprising result of mutation in MECP2. Hum. Molec.
Genet. 9:
- [101199]2365–2375, 2000.
- [101200]17638.
- [101201]Gendrot, C.; Ronce, N.; Raynaud, M.; Ayrault, A.–D.;
- Dourlens,
- [101202]J.; Castelnau, P.; Muh, J.–P.; Chelly, J.; Moraine, C.: X-
linked

[101203]nonspecific mental retardation (MRX16) mapping to distal
Xq28: linkage

[101204]study and neuropsychological data in a large family. Am.
J. Med.

[101205]Genet. 83: 411–418, 1999.

[101206]17639.

[101207]Guy, J.; Hendrich, B.; Holmes, M.; Martin, J. E.; Bird, A.: A

[101208]mouse Mecp2-null mutation causes neurological symptoms that mimic
toms that mimic

[101209]Rett syndrome. Nature Genet. 27: 322–326, 2001.

[101210]17640.

[101211]Hagberg, B.: Rett's syndrome: prevalence and impact on
progressive

[101212]severe mental retardation in girls. Acta Pediat. Scand. 74:
405–408,

[101213]1985.

[101214]17641.

[101215]Heilstedt, H. A.; Shahbazian, M. D.; Lee, B.: Infantile hypotonia
tonia

[101216]as a presentation of Rett syndrome. Am. J. Med. Genet.
111: 238–242,

[101217]2002.

[101218]17642.

[101219]Hoffbuhr, K.; Devaney, J. M.; LaFleur, B.; Sirianni, N.;
Scacheri,

[101220]C.; Giron, J.; Schuette, J.; Innis, J.; Marino, M.; Philippart,
M.;

[101221]Narayanan, V.; Umansky, R.; Kronn, D.; Hoffman, E. P.;
Naidu, S.:

[101222]MeCP2 mutations in children with and without the pheno-
type of Rett

[101223]syndrome. Neurology 56: 1486–1495, 2001.

[101224]17643.

[101225]Huppke, P.; Laccone, F.; Kramer, N.; Engel, W.; Hanefeld,
F.:

[101226]Rett syndrome: analysis of MECP2 and clinical characteri-
zation of

[101227]31 patients. Hum. Molec. Genet. 9: 1369–1375, 2000.

[101228]17644.

[101229]Imessaoudene, B.; Bonnefont, J.-P.; Royer, G.; Cormier-
Daire,

[101230]V.; Lyonnet, S.; Lyon, G.; Munnich, A.; Amiel, J.: MECP2

mutation

[101231]in non-fatal, non-progressive encephalopathy in a male. J.

Med. Genet. 38:

[101232]171-174, 2001.

[101233]17645.

[101234]Jones, P. L.; Veenstra, G. J. C.; Wade, P. A.; Vermaak, D.;

Kass,

[101235]S. U.; Landsberger, N.; Strouboulis, J.; Wolffe, A. P.:

Methylated

[101236]DNA and MeCP2 recruit histone deacetylase to repress

transcription. Nature

[101237]Genet. 19: 187-191, 1998.

[101238]17646.

[101239]Klauck, S. M.; Lindsay, S.; Beyer, K. S.; Splitt, M.; Burn, J.;

[101240]Poustka, A.: A mutation hot spot for nonspecific X-linked

mental

[101241]retardation in the MECP2 gene causes the PPM-X syn-

drome. Am. J. Hum.

[101242]Genet. 70: 1034-1037, 2002.

[101243]17647.

[101244]Kleefstra, T.; Yntema, H. G.; Oudakker, A. R.; Romein, T.;

Sisternans,

[101245]E.; Nillesen, W.; van Bokhoven, H.; de Vries, B. B. A.;
Hamel, B.

[101246]C. J.: De novo MECP2 frameshift mutation in a boy with
moderate mental

[101247]retardation, obesity and gynaecomastia. Clin. Genet. 61:
359–362,

[101248]2002.

[101249]17648.

[101250]LaSalle, J. M.; Goldstine, J.; Balmer, D.; Greco, C. M.:
Quantitative

[101251]localization of heterogeneous methyl–CpG-binding pro-
tein 2 (MeCP2)

[101252]expression phenotypes in normal and Rett syndrome brain
by laser scanning

[101253]cytometry. Hum. Molec. Genet. 10: 1729–1740, 2001.

[101254]17649.

[101255]Lewis, J. D.; Meehan, R. R.; Henzel, W. J.; Maurer–Fogy, I.;
Jeppesen,

[101256]P.; Klein, F.; Bird, A.: Purification, sequence, and cellular
localization

[101257]of a novel chromosomal protein that binds to methylated

DNA. Cell 69:

[101258]905–914, 1992.

[101259]17650.

[101260]Meloni, I.; Bruttini, M.; Longo, I.; Mari, F.; Rizzolio, F.;

D'Adamo,

[101261]P.; Denvriendt, K.; Fryns, J.-P.; Toniolo, D.; Renieri, A.: A

mutation

[101262]in the Rett syndrome gene, MECP2, causes X-linked men-

tal retardation

[101263]and progressive spasticity in males. Am. J. Hum. Genet.

67: 982–985,

[101264]2000. Note: Erratum: Am. J. Hum. Genet. 67: 1365 only,

2000.

[101265]17651.

[101266]Moncla, A.; Kpebe, A.; Missirian, C.; Mancini, J.; Villard, L.

[101267]: Polymorphisms in the C-terminal domain of MECP2 in

mentally handicapped

[101268]boys: implications for genetic counselling. Europ. J. Hum.

Genet. 10:

[101269]86–89, 2002.

[101270]17652.

[101271]Nan, X.; Campoy, F. J.; Bird, A.: MeCP2 is a transcriptional
 [101272]repressor with abundant binding sites in genomic chromatin. Cell 88:
 [101273]471–481, 1997.
 [101274]17653.

[101275]Reuber, B. E.; Germain-Lee, E.; Collins, C. S.; Morrell, J. C.;
 [101276]Ameritunga, R.; Moser, H. W.; Valle, D.; Gould, S. J.: Mutations
 in PEX1 are the most common cause of peroxisome biogenesis disorders. Nature
 [101278]Genet. 17: 445–448, 1997.
 [101279]17654.

[101280]Corbaz, A.; ten Hove, T.; Herren, S.; Graber, P.; Schwartzburd,
 [101281]B.; Belzer, I.; Harrison, J.; Plitz, T.; Kosco-Vilbois, M. H.; Kim,
 [101282]S.-H.; Dinarello, C. A.; Novick, D.; van Deventer, S.; Chvatchko,
 [101283]Y.: IL-18-binding protein expression by endothelial cells and macrophages
 [101284]is up-regulated during active Crohn's disease. J. Immun.

168: 3608–3616,

[101285]2002.

[101286]17655.

[101287]Turner, N.; Mason, P. J.; Brown, R.; Fox, M.; Povey, S.;
Rees,

[101288]A.; Pusey, C. D.: Molecular cloning of the human Good-
pasture antigen

[101289]demonstrates it to be the alpha-3 chain of type IV colla-
gen. J. Clin.

[101290]Invest. 89: 592–601, 1992.

[101291]17656.

[101292]Alexander, R. W.: Hypertension and the pathogenesis of
atherosclerosis.

[101293]Oxidative stress and the mediation of arterial inflamma-
tory response:

[101294]a new perspective. Hypertension 25: 155–161, 1995.

[101295]17657.

[101296]Bu-Ghanim, H. N.; Casimir, C. M.; Povey, S.; Segal, A. W.:
The

[101297]alpha subunit of cytochrome b(-245) mapped to chromo-
some 16. Genomics 8:

[101298]568–570, 1990.

[101299]17658.

[101300]Clark, R. A.; Malech, H. L.; Gallin, J. I.; Nuno, H.; Volpp, B.

[101301]D.; Pearson, D. W.; Nauseef, W. M.; Curnutte, J. T.: Genetic variants

[101302]of chronic granulomatous disease: prevalence of deficiencies of two

[101303]cytosolic components of the NADPH oxidase system. New Eng. J. Med. 321:

[101304]647–652, 1989.

[101305]17659.

[101306]Abbott, M. H.; Folstein, S. E.; Abbey, H.; Pyeritz, R. E.: Psychiatric

[101307]manifestations of homocystinuria due to cystathionine beta-synthase

[101308]deficiency: prevalence, natural history, and relationship to neurologic

[101309]impairment and vitamin B(6)–responsiveness. Am. J. Med. Genet. 26:

[101310]959–969, 1987.

[101311]17660.

[101312]Almgren, B.; Eriksson, H.; Hemmingsson, A.; Hillerdal, G.;
Larsson,

[101313]E.; Aberg, H.: Abdominal aortic aneurysm in homocystin-
uria. Acta

[101314]Chir. Scand. 144: 545-546, 1978.

[101315]17661.

[101316]Aral, B.; Coude, M.; London, J.; Aupetit, J.; Chasse, J.-F.;
Zabot,

[101317]M.-T.; Chadeaux-Vekemans, B.; Kamoun, P.: Two novel
mutations (K384E

[101318]and L539S) in the C-terminal moiety of the cystathionine
beta-synthase

[101319]protein in two French pyridoxine-responsive homocystin-
uria patients. Hum.

[101320]Mutat. 9: 81-82, 1997.

[101321]17662.

[101322]Avramopoulos, D.; Cox, T.; Kraus, J. P.; Chakravarti, A.;
Antonarakis,

[101323]S. E.: Linkage mapping of the cystathionine beta-synthase
(CBS) gene

[101324]on human chromosome 21 using a DNA polymorphism in
the 3-prime untranslated

[101325]region. Hum. Genet. 90: 566–568, 1993.

[101326]17663.

[101327]Barber, G. W.; Spaeth, G. L.: Pyridoxine therapy in homocystinuria.

[101328](Letter) Lancet I: 337 only, 1967.

[101329]17664.

[101330]Bass, H. N.; LaGrave, D.; Mardach, R.; Cederbaum, S. D.; Fuster,

[101331]C. D.; Chetty, M.: Spontaneous pneumothorax in association with pyridoxine-responsive

[101332]homocystinuria. J. Inherit. Metab. Dis. 20: 831–832, 1997.

[101333]17665.

[101334]Boers, G. H. J.; Fowler, B.; Smals, A. G. H.; Trijbels, F. J. M.;

[101335]Leermakers, A. I.; Kleijer, W. J.; Kloppenborg, P. W. C.: Improved

[101336]identification of heterozygotes for homocystinuria due to cystathionine

[101337]synthase deficiency by the combination of methionine loading and enzyme

[101338]determination in cultured fibroblasts. Hum. Genet. 69: 164–169,

[101339]1985.

[101340]17666.

[101341]Boers, G. H. J.; Smals, A. G. H.; Trijbels, F. J. M.; Fowler, B.;

[101342]Bakkeren, J. A. J. M.; Schoonderwaldt, H. C.; Kleijer, W. J.;
Kloppenborg,

[101343]P. W. C.: Heterozygosity for homocystinuria in premature
peripheral

[101344]and cerebral occlusive arterial disease. New Eng. J. Med.
313: 709-715,

[101345]1985.

[101346]17667.

[101347]Bross, P.; Corydon, T. J.; Andresen, B. S.; Jorgensen, M. M.;
Bolund,

[101348]L.; Gregersen, N.: Protein misfolding and degradation in
genetic

[101349]diseases. Hum. Mutat. 14: 186-198, 1999.

[101350]17668.

[101351]Burke, J. P.; O'Keefe, M.; Bowell, R.; Naughten, E. R.: Ocular

[101352]complications in homocystinuria: early and late treated.
Brit. J.

[101353]Ophthal. 73: 427-431, 1989.

[101354]17669.

[101355]Carey, M. C.; Donovan, D. E.; Fitzgerald, O.; McAuley, F.
D.:

[101356]Homocystinuria: a clinical and pathological study of nine
subjects

[101357]in six families. Am. J. Med. 45: 7-25, 1968.

[101358]17670.

[101359]Carey, M. C.; Fennelly, J. J.; Fitzgerald, O.: Homocystinuria.

[101360]II. Subnormal serum folate levels, increased folate clear-
ance and

[101361]effects of folic acid therapy. Am. J. Med. 45: 26-31, 1968.

[101362]17671.

[101363]Carson, N. A. J.; Carre, I. J.: Treatment of homocystinuria
with

[101364]pyridoxine: a preliminary study. Arch. Dis. Child. 44:
387-392,

[101365]1969.

[101366]17672.

[101367]Carson, N. A. J.; Cusworth, D. C.; Dent, C. E.; Field, C. M.

B.;

[101368]Neill, D. W.; Westall, R. G.: Homocystinuria: a new inborn error

[101369]of metabolism associated with mental deficiency. Arch. Dis. Child. 38:

[101370]425-436, 1963.

[101371]17673.

[101372]Carson, N. A. J.; Neill, D. W.: Metabolic abnormalities detected

[101373]in a survey of mentally backward individuals in Northern Ireland. Arch.

[101374]Dis. Child. 37: 505-513, 1962.

[101375]17674.

[101376]Chadefaux, B.; Rethore, M. O.; Raoul, O.; Ceballos, I.; Poissonnier,

[101377]M.; Gilgenkrantz, S.; Allard, D.: Cystathionine beta synthase: gene

[101378]dosage effect in trisomy 21. Biochem. Biophys. Res. Commun. 128:

[101379]40-44, 1985.

[101380]17675.

[101381]Chao, C.-L.; Tsai, H.-H.; Lee, C.-M.; Hsu, S.-M.; Kao, J.-T.;

[101382]Chien, K.-L.; Sung, F.-C.; Lee, Y.-T.: The graded effect of
hyperhomocysteinemia

[101383]on the severity and extent of coronary atherosclerosis.
Atherosclerosis 147:

[101384]379-386, 1999.

[101385]17676.

[101386]Hoffmann, M. M.; Jacob, S.; Luft, D.; Schmulling, R.-M.;
Rett,

[101387]K.; Marz, W.; Haring, H.-U.; Matthaei, S.: Type I hyper-
lipoproteinemia

[101388]due to a novel loss of function mutation of lipoprotein li-
pase, cys239-trp,

[101389]associated with recurrent severe pancreatitis. J. Clin. En-
docr. Metab. 85:

[101390]4795-4798, 2000.

[101391]17677.

[101392]Holt, L. E., Jr.; Aylward, F. X.; Timbers, H. G.: Idiopathic

[101393]familial lipemia. Bull. Johns Hopkins Hosp. 64: 279-314,
1939.

[101394]17678.

[101395]Ishimura-Oka, K.; Faustinella, F.; Kihara, S.; Smith, L. C.; Oka, K.; Chan, L.: A missense mutation (trp86-to-arg) in exon 3 of the lipoprotein lipase gene: a cause of familial chylomicronemia. Am. J. Hum. Genet. 50: 1275-1280, 1992.

[101399]17679.

[101400]Kastelein, J. J. P.; Groenemeyer, B. E.; Hallman, D. M.; Henderson, H.; Reymer, P. W. A.; Gagne, S. E.; Jansen, H.; Seidell, J. C.; Kromhout, D.; Jukema, J. W.; Bruschke, A. V. G.; Boerwinkle, E.; Hayden, M. R.; The Regress Study Group: The asn9 variant of lipoprotein lipase is associated with the -93G promoter mutation and an increased risk of coronary artery disease. Clin. Genet. 53: 27-33, 1998.

[101406]17680.

[101407]Kirchgessner, T. G.; Chuat, J.-C.; Heinzmann, C.; Etienne, J.;

[101408]Guilhot, S.; Svenson, K.; Ameis, D.; Pilon, C.; d'Auriol, L.;
Andalibi,
[101409]A.; Schotz, M. C.; Galibert, F.; Lusi, A. J.: Organization of
the
[101410]human lipoprotein lipase gene and evolution of the lipase
gene family. Proc.
[101411]Nat. Acad. Sci. 86: 9647-9651, 1989.
[101412]17681.
[101413]Kirchgessner, T. G.; Svenson, K. L.; Lusi, A. J.; Schotz, M.
[101414]C.: The sequence of cDNA encoding lipoprotein lipase: a
member of
[101415]a lipase gene family. J. Biol. Chem. 262: 8463-8466,
1987.
[101416]17682.
[101417]Kobayashi, J.; Nishida, T.; Ameis, D.; Stahnke, G.; Schotz,
M.
[101418]C.; Hashimoto, H.; Fukamachi, I.; Shirai, K.; Saito, Y.;
Yoshida,
[101419]S.: A heterozygous mutation (the codon for ser477 to a
stop codon)
[101420]in lipoprotein lipase contributes to a defect in lipid inter-
face recognition

[101421]in a case with type I hyperlipidemia. Biochem. Biophys.

Res. Commun. 182:

[101422]70-77, 1992.

[101423]17683.

[101424]Kobayashi, J.; Sasaki, N.; Tashiro, J.; Inadera, H.; Saito, Y.;

[101425]Yoshida, S.: A missense mutation (ala334-to-thr) in exon
7 of the

[101426]lipoprotein lipase gene in a case with type I hyperlipi-
demia. Biochem.

[101427]Biophys. Res. Commun. 191: 1046-1054, 1993.

[101428]17684.

[101429]Langlois, S.; Deeb, S.; Brunzell, J.; Kastelein, J. J.; Hayden,

[101430]M. R.: A unique insertion accounts for a significant pro-
portion of

[101431]the mutations in the lipoprotein lipase (LPL)
gene.(Abstract) Am.

[101432]J. Hum. Genet. 43: A191, 1988.

[101433]17685.

[101434]Langlois, S.; Deeb, S.; Brunzell, J. D.; Kastelein, J. J.; Hay-
den,

[101435]M. R.: A major insertion accounts for a significant proportion of
[101436]mutations underlying human lipoprotein lipase deficiency.
Proc. Nat.
[101437]Acad. Sci. 86: 948–952, 1989.
[101438]17686.
[101439]Levak–Frank, S.; Radner, H.; Walsh, A.; Stollberger, R.;
Knipping,
[101440]G.; Hoefler, G.; Sattler, W.; Weinstock, P. H.; Breslow, J. L.;
Zechner,
[101441]R.: Muscle–specific overexpression of lipoprotein lipase
causes a
[101442]severe myopathy characterized by proliferation of mitochondria and
[101443]peroxisomes in transgenic mice. J. Clin. Invest. 96:
976–986, 1995.
[101444]17687.
[101445]Li, S.; Oka, K.; Galton, D.; Stocks, J.: Pvu–II RFLP at the human
[101446]lipoprotein lipase (LPL) gene locus. Nucleic Acids Res. 16:
2358,
[101447]1988.

[101448]17688.

[101449]Lo, J. Y.; Smith, L. C.; Chan, L.: Lipoprotein lipase: role of

[101450]intramolecular disulfide bonds in enzyme catalysis.

Biochem. Biophys.

[101451]Res. Commun. 206: 266-271, 1995.

[101452]17689.

[101453]Ma, Y.; Henderson, H. E.; Julien, P.; Roederer, G.; Brunzell,

[101454]J.; Hayden, M. R.: A missense mutation (pro-to-leu207) in

[101455]the human lipoprotein lipase gene is the major cause of
type I hyperlipoproteinemia

[101456]in French Canadians. (Series) Miami Short Reports. Ad-
vances

[101457]in Gene Technology: The Molecular Biology of Human Ge-
netic Disease.

[101458]New York: IRL Press (pub.) 1: 1991. Pp. 34 only.

[101459]17690.

[101460]Ma, Y.; Henderson, H. E.; Ven Murthy, M. R.; Roederer, G.;
Monsalve,

[101461]M. V.; Clarke, L. A.; Normand, T.; Julien, P.; Gagne, C.;
Lambert,

[101462]M.; Davignon, J.; Lupien, P. J.; Brunzell, J.; Hayden, M. R.: A
mutation

[101463]in the human lipoprotein lipase gene as the most common
cause of familial

[101464]chylomicronemia in French Canadians. New Eng. J. Med.
324: 1761–1766,

[101465]1991.

[101466]17691.

[101467]Ma, Y.; Liu, M.–S.; Chitayat, D.; Bruin, T.; Beisiegel, U.;
Benlian,

[101468]P.; Foubert, L.; De Gennes, J. L.; Funke, H.; Forsythe, I.;
Blaichman,

[101469]S.; Papanikolaou, M.; Erkelens, D. W.; Kastelein, J.; Brun-
zell, J.

[101470]D.; Hayden, M. R.: Recurrent missense mutations at the
first and

[101471]second base of codon arg243 in human lipoprotein lipase
in patients

[101472]of different ancestries. Hum. Mutat. 3: 52–58, 1994.

[101473]17692.

[101474]Ma, Y.; Liu, M.–S.; Ginzinger, D.; Frohlich, J.; Brunzell, J.

[101475]D.; Hayden, M. R.: Gene–environment interaction in the
conversion
[101476]of a mild–to–severe phenotype in a patient homozygous
for a ser172–to–cys
[101477]mutation in the lipoprotein lipase gene. J. Clin. Invest. 91:
1953–1958,
[101478]1993.
[101479]17693.
[101480]Ma, Y.; Wilson, B. I.; Bijvoet, S.; Henderson, H. E.; Cramb,
E.;
[101481]Roederer, G.; Ven Murthy, M. R.; Julien, P.; Bakker, H. D.;
Kastelein,
[101482]J. J. P.; Brunzell, J. D.; Hayden, M. R.: A missense mutation
(asp250–to–asn)
[101483]in exon 6 of the human lipoprotein lipase gene causes
chylomicronemia
[101484]in patients of different ancestries. Genomics 13: 649–653,
1992.
[101485]17694.
[101486]Mattei, M. G.; Etienne, J.; Chuat, J. C.; Nguyen, V. C.;
Brault,
[101487]D.; Bernheim, A.; Galibert, F.: Assignment of the human

lipoprotein

[101488]lipase (LPL) gene to chromosome band 8p22. Cytogenet.

Cell Genet. 63:

[101489]45–46, 1993.

[101490]17695.

[101491]Miesenbock, G.; Holzl, B.; Foger, B.; Brandstatter, E.;

Paulweber,

[101492]B.; Sandhofer, F.; Patsch, J. R.: Heterozygous lipoprotein

lipase

[101493]deficiency due to a missense mutation as the cause of im-
paired triglyceride

[101494]tolerance with multiple lipoprotein abnormalities. J. Clin.

Invest. 91:

[101495]448–455, 1993.

[101496]17696.

[101497]Nevin, D. N.; Brunzell, J. D.; Deeb, S. S.: The LPL gene in
individuals

[101498]with familial combined hyperlipidemia and decreased LPL
activity. Arteriosclerosis

[101499]Thromb. 14: 869–873, 1994.

[101500]17697.

- [101501]Nevin, N. C.; Slack, J.: Hyperlipidaemic xanthomatosis II: mode
[101502]of inheritance in 55 families with essential hyperlipi-
daemia and xanthomatosis. J.
[101503]Med. Genet. 5: 9-28, 1968.
[101504]17698.
- [101505]Nickerson, D. A.; Taylor, S. L.; Weiss, K. M.; Clark, A. G.;
Hutchinson,
[101506]R. G.; Stengard, J.; Salomaa, V.; Vartiainen, E.; Boerwinkle,
E.;
[101507]Sing, C. F.: DNA sequence diversity in a 9.7-kb region of
the human
[101508]lipoprotein lipase gene. Nature Genet. 19: 233-240,
1998.
[101509]17699.
- [101510]Jacobson, J. H.; Stephens, G.: Hereditary choroidoretinal
degeneration:
[101511]study of a family including electroretinography and adap-
tometry. Arch.
[101512]Ophthalmol. 67: 321-335, 1962.
[101513]17700.

- [101514]Jay, M.: Personal Communication. London, England
6/18/1987.
- [101515]17701.
- [101516]Kaplan, J.; Bonneau, D.; Frezal, J.; Munnich, A.; Dufier, J. L.
- [101517]: Clinical and genetic heterogeneity in retinitis pigmentosa. Hum.
- [101518]Genet. 85: 635–642, 1990.
- [101519]17702.
- [101520]Kaplan, J.; Pelet, A.; Martin, C.; Delrieu, O.; Ayme, S.; Bonneau,
- [101521]D.; Briard, M. L.; Hanauer, A.; Larget-Piet, L.; Lefrancois, P.; Michel-Awad,
- [101522]A.; Plauchu, H.; Dufier, J. L.; Frezal, J.; Munnich, A.: Phenotype-genotype
- [101523]correlations in X linked retinitis pigmentosa. J. Med. Genet. 29:
- [101524]615–623, 1992.
- [101525]17703.
- [101526]Keats, B.; Ott, J.; Conneally, M.: Report of the committee
- [101527]on linkage and gene order. Cytogenet. Cell Genet. 51:

459–502, 1989.

[101528]17704.

[101529]Klein, D.; Franceschetti, A.; Hussels, I.; Race, R. R.; Sanger,

[101530]R.: X-linked retinitis pigmentosa and linkage studies with
the Xg

[101531]blood-groups. Lancet I: 974–975, 1967.

[101532]17705.

[101533]Litt, M.; Weleber, R. G.; Lovrien, E. W.; Litt, R. H.; Payne,

[101534]N.; Murphey, W.; Dubay, C.; Willard, H. F.: A family study
of X-linked

[101535]retinitis pigmentosa. (Abstract) Cytogenet. Cell Genet. 46:
651

[101536]only, 1987.

[101537]17706.

[101538]McQuarrie, M. D.: Two pedigrees of hereditary blindness
in man. J.

[101539]Genet. 30: 147–153, 1935.

[101540]17707.

[101541]Mears, A. J.; Gieser, L.; Yan, D.; Chen, C.; Fahrner, S.;
Hiriyanna,

[101542]S.; Fujita, R.; Jacobson, S. G.; Sieving, P. A.; Swaroop, A.:
Protein-truncation

[101543]mutations in the RP2 gene in a North American cohort of
families with

[101544]X-linked retinitis pigmentosa. (Letter) Am. J. Hum. Genet.
64: 897-900,

[101545]1999.

[101546]17708.

[101547]Meitinger, T.; Fraser, N. A.; Lorenz, B.; Zrenner, E.;
Murken,

[101548]J.; Craig, I. W.: Linkage of X-linked retinitis pigmentosa to
the

[101549]hypervariable DNA marker M27-beta (DXS255). Hum.
Genet. 81: 283-286,

[101550]1989.

[101551]17709.

[101552]Miano, M. G.; Testa, F.; Filippini, F.; Trujillo, M.; Conte, I.;

[101553]Lanzara, C.; Millan, J. M.; De Bernardo, C.; Grammatico, B.;
Mangino,

[101554]M.; Torrente, I.; Carrozzo, R.; Simonelli, F.; Rinaldi, E.;
Ventruto,

- [101555]V.; D'Urso, M.; Ayuso, C.; Ciccodicola, A.: Identification of novel
- [101556]RP2 mutations in a subset of X-linked retinitis pigmentosa families
- [101557]and prediction of new domains. Hum. Mutat. 18: 109–119, 2001.
- [101558]17710.
- [101559]Mukai, S.; Dryja, T. P.; Bruns, G. A. P.; Aldridge, J. F.; Berson,
- [101560]E. L.: Linkage between the X-linked retinitis pigmentosa locus and
- [101561]the L1.28 locus. Am. J. Ophthal. 100: 225–229, 1985.
- [101562]17711.
- [101563]Schwahn, U.; Lenzner, S.; Dong, J.; Feil, S.; Hinzmann, B.; van
- [101564]Duijnhoven, G.; Kirschner, R.; Hemberger, M.; Bergen, A. A. B.; Rosenberg,
- [101565]T.; Pinckers, A. J. L. G.; Fundele, R.; Rosenthal, A.; Cremers, F.
- [101566]P. M.; Ropers, H.-H.; Berger, W.: Positional cloning of the gene
- [101567]for X-linked retinitis pigmentosa 2. Nature Genet. 19:

327-332,

[101568]1998.

[101569]17712.

[101570]Schwahn, U.; Paland, N.; Techritz, S.; Lenzner, S.; Berger, W.

[101571]: Mutations in the X-linked RP2 gene cause intracellular misrouting

[101572]and loss of the protein. Hum. Molec. Genet. 10: 1177-1183, 2001.

[101573]17713.

[101574]Sharon, D.; Bruns, G. A. P.; McGee, T. L.; Sandberg, M. A.; Berson,

[101575]E. L.; Dryja, T. P.: X-linked retinitis pigmentosa: mutation spectrum

[101576]of the RPGR and RP2 genes and correlation with visual function. Invest.

[101577]Ophthal. Vis. Sci. 41: 2712-2721, 2000.

[101578]17714.

[101579]Spence, M. A.; Elston, R. C.; Cederbaum, S. D.: Pedigree analysis

[101580]to determine the mode of inheritance in a family with re-

tinitis pigmentosa. Clin.

[101581]Genet. 5: 338-343, 1974.

[101582]17715.

[101583]Thiselton, D. L.; Hampson, R. M.; Nayudu, M.; Van
Maldergem, L.;

[101584]Wolf, M. L.; Saha, B. K.; Bhattacharya, S. S.; Hardcastle, A.
J.:

[101585]Mapping the RP2 locus for X-linked retinitis pigmentosa
on proximal

[101586]Xp: a genetically defined 5-cM critical region and exclu-
sion of candidate

[101587]genes by physical mapping. Genome Res. 6: 1093-1102,
1996.

[101588]17716.

[101589]Usher, C. H.: Bowman lecture on a few hereditary eye af-
fections. Trans.

[101590]Ophthal. Soc. U.K. 55: 164-245, 1935.

[101591]17717.

[101592]Waardenburg, P. J.: Das menschliche Auge und seine Er-
banlagean.

[101593]S-Gravenhage: Martinus Nijhoff (pub.) 1932.

[101594]17718.

[101595]Waardenburg, P. J.; Franceschetti, A.; Klein, D.: Genetics
and

[101596]Ophthalmology. Oxford: Blackwell (pub.) 1: 1961. Pp. 799
only.

[101597]17719.

[101598]Warburg, M.; Simonsen, S. E.: Sex-linked recessive retinitis

[101599]pigmentosa: a preliminary study of the carriers. Acta Oph-
thal. 46:

[101600]494-499, 1968.

[101601]17720.

[101602]Wright, A. F.; Bhattacharya, S. S.; Aldred, M. A.; Jay, M.;
Carothers,

[101603]A. D.; Thomas, N. S. T.; Bird, A. C.; Jay, B.; Evans, H. J.: Ge-
netic

[101604]localisation of the RP2 type of X linked retinitis pigmen-
tosa in a

[101605]large kindred. J. Med. Genet. 28: 453-457, 1991.

[101606]17721.

[101607]Wright, A. F.; Bhattacharya, S. S.; Clayton, J. F.; Dempster,

[101608]M.; Tippett, P.; McKeown, C. M. E.; Jay, M.; Jay, B.; Bird, A. C.

[101609]; Linkage relationships between X-linked retinitis pigmentosa and

[101610]nine short-arm markers: exclusion of the disease locus from Xp21 and

[101611]localization to between DXS7 and DXS14. Am. J. Hum. Genet. 41: 635-644,

[101612]1987.

[101613]17722.

[101614]Wright, A. F.; Bhattacharya, S. S.; Clayton, J. F.; Jay, M.; Jay,

[101615]B.; Bird, A. C.; Evans, H. J.: Linkage relationships about X-linked

[101616]retinitis pigmentosa (RP2). (Abstract) Cytogenet. Cell Genet. 46:

[101617]720 only, 1987.

[101618]17723.

[101619]Lahn, B. T.; Page, D. C.: Retroposition of autosomal mRNA yielded

[101620]testis-specific gene family on human Y chromosome. Nature Genet. 21:

[101621]429–433, 1999.

[101622]17724.

[101623]Yen, P. H.: A long-range restriction map of deletion interval

[101624]6 of the human Y chromosome: a region frequently deleted in azoospermic

[101625]males. Genomics 54: 5–12, 1998.

[101626]17725.

[101627]MacLean, H. E.; Chu, S.; Warne, G. L.; Zajac, J. D.: Related

[101628]individuals with different androgen receptor gene deletions. J. Clin.

[101629]Invest. 91: 1123–1128, 1993.

[101630]17726.

[101631]Madgar, I.; Green, L.; Kent-First, M.; Weissenberg, R.; Gershoni-Baruch,

[101632]R.; Goldman, B.; Friedman, E.: Genotyping of Israeli infertile men

[101633]with idiopathic oligozoospermia. Clin. Genet. 62: 203–207, 2002.

[101634]17727.

[101635] Marcelli, M.; Tilley, W. D.; Wilson, C. M.; Griffin, J. E.; Wilson,
son,
[101636] J. D.; McPhaul, M. J.: Definition of the human androgen re-
ceptor
[101637] gene structure permits the identification of mutations that
cause
[101638] androgen resistance: premature termination of the recep-
tor protein
[101639] at amino acid residue 588 causes complete androgen re-
sistance. Molec.
[101640] Endocr. 4: 1105-1116, 1990.
[101641] 17728.
[101642] Marcelli, M.; Tilley, W. D.; Wilson, C. M.; Wilson, J. D.; Grif-
fin,
[101643] J. E.; McPhaul, M. J.: A single nucleotide substitution intro-
duces
[101644] a premature termination codon into the androgen recep-
tor gene of a
[101645] patient with receptor-negative androgen resistance. J.
Clin. Invest. 85:
[101646] 1522-1528, 1990.
[101647] 17729.

[101648]Marcelli, M.; Tilley, W. D.; Zoppi, S.; Griffin, J. E.; Wilson,
[101649]J. D.; McPhaul, M. J.: Androgen resistance associated with
a mutation
[101650]of the androgen receptor at amino acid 772 (arg-to-cys)
results from
[101651]a combination of decreased messenger ribonucleic acid
levels and impairment
[101652]of receptor function. J. Clin. Endocr. 73: 318–325, 1991.
[101653]17730.

[101654]McC Campbell, A.; Taylor, J. P.; Taye, A. A.; Robitschek, J.; Li,
[101655]M.; Walcott, J.; Merry, D.; Chai, Y.; Paulson, H.; Sobue, G.;
Fischbeck,
[101656]K. H.: CREB-binding protein sequestration by expanded
polyglutamine. Hum.
[101657]Molec. Genet. 9: 2197–2202, 2000.
[101658]17731.

[101659]McPhaul, M. J.; Griffin, J. E.: Male pseudohermaphroditism
caused
[101660]by mutations of the human androgen receptor. J. Clin. En-
docr. Metab. 84:
[101661]3435–3441, 1999.

[101662]17732.

[101663]McPhaul, M. J.; Marcelli, M.; Tilley, W. D.; Griffin, J. E.;
Isidro-Gutierrez,

[101664]R. F.; Wilson, J. D.: Molecular basis of androgen resistance
in a

[101665]family with a qualitative abnormality of the androgen re-
ceptor and

[101666]responsive to high-dose androgen therapy. J. Clin. Invest.
87: 1413-1421,

[101667]1991.

[101668]17733.

[101669]McPhaul, M. J.; Marcelli, M.; Zoppi, S.; Griffin, J. E.; Wilson,

[101670]J. D.: Genetic basis of endocrine disease 4: the spectrum
of mutations

[101671]in the androgen receptor gene that causes androgen re-
sistance. J.

[101672]Clin. Endocr. Metab. 76: 17-23, 1993.

[101673]17734.

[101674]McPhaul, M. J.; Marcelli, M.; Zoppi, S.; Wilson, C. M.; Grif-
fin,

[101675]J. E.; Wilson, J. D.: Mutations in the ligand-binding domain

of the

[101676] androgen receptor gene cluster in two regions of the
gene. J. Clin.

[101677] Invest. 90: 2097–2101, 1992.

[101678] 17735.

[101679] McPhaul, M. J.; Schweikert, H.-U.; Allman, D. R.: Assess-
ment

[101680] of androgen receptor function in genital skin fibroblasts
using a

[101681] recombinant adenovirus to deliver an androgen-respon-
sive reporter

[101682] gene. J. Clin. Endocr. Metab. 82: 1944–1948, 1997.

[101683] 17736.

[101684] Mifsud, A.; Ramirez, S.; Yong, E. L.: Androgen receptor
gene

[101685] CAG trinucleotide repeats in anovulatory infertility and
polycystic

[101686] ovaries. J. Clin. Endocr. Metab. 85: 3484–3488, 2000.

[101687] 17737.

[101688] Migeon, B. R.; Brown, T. R.; Axelman, J.; Migeon, C. J.:
Studies

[101689] of the locus for androgen receptor: localization on the human X and

[101690] evidence for homology with the Tfm locus in the mouse.
Proc. Nat.

[101691] Acad. Sci. 78: 6339–6343, 1981.

[101692] 17738.

[101693] Mongan, N. P.; Jaaskelainen, J.; Green, K.; Schwabe, J. W.;
Shimura,

[101694] N.; Dattani, M.; Hughes, I. A.: Two de novo mutations in
the AR gene

[101695] cause the complete androgen insensitivity syndrome in a
pair of monozygotic

[101696] twins. J. Clin. Endocr. Metab. 87: 1057–1061, 2002.

[101697] 17739.

[101698] Mononen, N.; Syrjakoski, K.; Matikainen, M.; Tammela, T.
L. J.;

[101699] Schleutker, J.; Kallioniemi, O.-P.; Trapman, J.; Koivisto, P.
A.:

[101700] Two percent of Finnish prostate cancer patients have a
germ-line mutation

[101701] in the hormone-binding domain of the androgen receptor
gene. Cancer

[101702]Res. 60: 6479–6481, 2000.

[101703]17740.

[101704]Muroño, K.; Mendonça, B. B.; Arnhold, I. J. P.; Rigon, A. C.
M.

[101705]M.; Migeon, C. J.; Brown, T. R.: Human androgen insensi-
tivity due

[101706]to point mutations encoding amino acid substitutions in
the androgen

[101707]receptor steroid-binding domain. Hum. Mutat. 6:
152–162, 1995.

[101708]17741.

[101709]Nakao, R.; Haji, M.; Yanase, T.; Ogo, A.; Takayanagi, R.;
Katsube,

[101710]T.; Fukumaki, Y.; Nawata, H.: A single amino acid substi-
tution (met786-to-val)

[101711]in the steroid-binding domain of human androgen recep-
tor leads to

[101712]complete androgen insensitivity syndrome. J. Clin. Endocr.
Metab. 74:

[101713]1152–1157, 1992.

[101714]17742.

- [101715]Newmark, J. R.; Hardy, D. O.; Tonb, D. C.; Carter, B. S.; Epstein,
stein,
[101716]J. I.; Isaacs, W. B.; Brown, T. R.; Barrack, E. R.: Androgen
receptor
[101717]gene mutations in human prostate cancer. Proc. Nat.
Acad. Sci. 89:
[101718]6319-6323, 1992.
[101719]17743.
[101720]Nguyen, D.; Steinberg, S. V.; Rouault, E.; Chagnon, S.;
Gottlieb,
[101721]B.; Pinsky, L.; Trifiro, M.; Mader, S.: A G577R mutation in
the human
[101722]AR P box results in selective decreases in DNA binding
and in partial
[101723]androgen insensitivity syndrome. Molec. Endocr. 15:
1790-1802, 2001.
[101724]17744.
[101725]Ohno, S.: The Y-linked antigen locus and the X-linked
Tfm locus
[101726]as major regulatory genes of the mammalian sex deter-
mining mechanism. J.
[101727]Steroid Biochem. 8: 585-592, 1977.

[101728]

[101729]100. Ohno, S.: Simplicity of mammalian regulatory systems inferred

[101730]by single gene determination of sex phenotypes. Nature
234: 134–137,

[101731]1971.

[101732]

[101733]101. Ong, Y. C.; Wong, H. B.; Adaikan, G.; Yong, E. L.: Directed

[101734]pharmacological therapy of ambiguous genitalia due to an
androgen

[101735]receptor gene mutation. (Letter) Lancet 354: 1444–1445,
1999.

[101736]

[101737]102. Patterson, M. N.; Hughes, I. A.; Gottlieb, B.; Pinsky,
L.: The

[101738]androgen receptor gene mutations database. Nucleic
Acids Res. 22:

[101739]3560–3562, 1994.

[101740]

[101741]103. Pinsky, L.; Kaufman, M.; Killinger, D. W.; Burko, B.;

Shatz,

[101742]D.; Volpe, R.: Human minimal androgen insensitivity with normal dihydrotestosterone-binding

[101743]capacity in cultured genital skin fibroblasts: evidence for an androgen-selective

[101744]qualitative abnormality of the receptor. Am. J. Hum. Genet. 36:

[101745]965-978, 1984.

[101746]

[101747]104. Pinsky, L.; Kaufman, M.; Levitsky, L. L.: Partial androgen resistance

[101748]due to a distinctive qualitative defect of the androgen receptor. Am.

[101749]J. Med. Genet. 27: 459-466, 1987.

[101750]

[101751]105. Pinsky, L.; Kaufman, M.; Summitt, R. L.: Congenital androgen

[101752]insensitivity due to a qualitatively abnormal androgen receptor. Am.

[101753]J. Med. Genet. 10: 91-99, 1981.

[101754]

- [101755]106. Pinsky, L.; Trifiro, M.; Sebbaghatian, N.; Kaufman, M.;
Chang,
[101756]C.; Trapman, J.; Brinkmann, A. O.; Kuiper, G. G. J. M.; Ris,
C. J.;
[101757]Brown, C. J.; Willard, H. F.; Sergovich, F.: A deletional al-
teration
[101758]of the androgen receptor (AR) gene in a sporadic patient
with complete
[101759]androgen insensitivity (CAI) who is mentally retarded.
(Abstract) Am.
[101760]J. Hum. Genet. 45 (suppl.): A212, 1989.
[101761]
[101762]107. Prior, L.; Bordet, S.; Trifiro, M. A.; Mhatre, A.; Kauf-
man, M.;
[101763]Pinsky, L.; Wrogeman, K.; Belsham, D. D.; Pereira, F.;
Greenberg,
[101764]C.; Trapman, J.; Brinkman, A. O.; Chang, C.; Liao, S.: Re-
placement
[101765]of arginine 773 by cysteine or histidine in the human an-
drogen receptor
[101766]causes complete androgen insensitivity with different re-
ceptor phenotypes. Am.

[101767]J. Hum. Genet. 51: 143–155, 1992.

[101768]

[101769]108. Quigley, C. A.; Friedman, K. J.; Johnson, A.; Lafreniere, R.

[101770]G.; Silverman, L. M.; Lubahn, D. B.; Brown, T. R.; Wilson, E. M.;

[101771]Willard, H. F.; French, F. S.: Complete deletion of the androgen

[101772]receptor gene: definition of the null phenotype of the androgen insensitivity

[101773]syndrome and determination of carrier status. J. Clin. Endocr. Metab. 74:

[101774]927–933, 1992.

[101775]

[101776]109. Ris-Stalpers, C.; Kuiper, G. G. J. M.; Faber, P. W.; Schweikert,

[101777]H. U.; van Rooij, H. C. J.; Zegers, N. D.; Hodgins, M. B.; Degenhart,

[101778]H. J.; Trapman, J.; Brinkmann, A. O.: Aberrant splicing of androgen

[101779]receptor mRNA results in synthesis of a nonfunctional receptor protein

[101780]in a patient with androgen insensitivity. Proc. Nat. Acad. Sci. 87:

[101781]7866–7870, 1990.

[101782]

[101783]110. Rodien, P.; Mebarki, F.; Mowszowicz, I.; Chaussain, J.-L.; Young,

[101784]J.; Morel, Y.; Schaison, G.: Different phenotypes in a family with

[101785]androgen insensitivity caused by the same M780I point mutation in

[101786]the androgen receptor gene. J. Clin. Endocr. Metab. 81: 2994–2998,

[101787]1996.

[101788]

[101789]111. Sai, T.; Seino, S.; Chang, C.; Trifiro, M.; Pinsky, L.; Mhatre,

[101790]A.; Kaufman, M.; Lambert, B.; Trapman, J.; Brinkmann, A. O.; Rosenfield,

[101791]R. L.; Liao, S.: An exonic point mutation of the androgen receptor

[101792]gene in a family with complete androgen insensitivity. Am. J. Hum.

[101793]Genet. 46: 1095–1100, 1990.

[101794]

[101795]112. Sammarco, I.; Grimaldi, P.; Rossi, P.; Cappa, M.;
Moretti, C.;

[101796]Frajese, G.; Geremia, R.: Novel point mutation in the splice
donor

[101797]site of exon–intron junction 6 of the androgen receptor
gene in a

[101798]patient with partial androgen insensitivity syndrome. J.
Clin. Endocr.

[101799]Metab. 85: 3256–3261, 2000.

[101800]

[101801]113. Schoenberg, M. P.; Hakimi, J. M.; Wang, S.; Bova, G.
S.; Epstein,

[101802]J. I.; Fischbeck, K. H.; Isaacs, W. B.; Walsh, P. C.; Barrack, E.

[101803]R.: Microsatellite mutation (CAG (24–to–18)) in the andro-
gen receptor

[101804]gene in human prostate cancer Biochem. Biophys. Res.
Commun. 198:

[101805]74–80, 1994.

[101806]

[101807]114. Shang, Y.; Myers, M.; Brown, M.: Formation of the androgen receptor

[101808]transcription complex. Molec. Cell 9: 601–610, 2002.

[101809]

[101810]115. Simeoni, S.; Mancini, M. A.; Stenoien, D. L.; Marcelli, M.; Weigel,

[101811]N. L.; Zanisi, M.; Martini, L.; Poletti, A.: Motoneuronal cell death

[101812]is not correlated with aggregate formation of androgen receptors containing

[101813]an elongated polyglutamine tract. Hum. Molec. Genet. 9: 133–144,

[101814]2000.

[101815]

[101816]116. Sullivan, D. A.; Sullivan, B. D.; Ullman, M. D.; Rocha, E. M.;

[101817]Krenzer, K. L.; Cermak, J. M.; Toda, I.; Doane, M. G.; Evans, J. E.;

[101818]Wickham, L. A.: Androgen influence on the meibomian gland. Invest.

[101819]Ophthal. Vis. Sci. 41: 3732–3742, 2000.

[101820]

[101821]117. Sultan, C.; Lumbroso, S.; Poujol, N.; Belon, C.;
Boudon, C.;

[101822]Lobaccaro, J.-M.: Mutations of androgen receptor gene in
androgen

[101823]insensitivity syndromes. J. Steroid Biochem. Molec. Biol.
46: 519-530,

[101824]1993.

[101825]

[101826]118. Sutherland, R. W.; Wiener, J. S.; Hicks, J. P.; Marcelli,
M.;

[101827]Gonzales, E. T.; Roth, D. R.; Lamb, D. J.: Androgen recep-
tor gene

[101828]mutations are rarely associated with isolated penile hy-
pospadias. J.

[101829]Urol. 156: 828-831, 1996.

[101830]

[101831]119. Taplin, M.-E.; Bubley, G. J.; Shuster, T. D.; Frantz, M.
E.;

[101832]Spooner, A. E.; Ogata, G. K.; Keer, H. N.; Balk, S. P.: Muta-
tion

[101833]of the androgen-receptor gene in metastatic androgen-independent prostate

[101834]cancer. New Eng. J. Med. 332: 1393-1398, 1995.

[101835]

[101836]120. Tilley, W. D.; Marcelli, M.; Wilson, J. D.; McPhaul, M. J.:

[101837]Characterization and expression of a cDNA encoding the human androgen

[101838]receptor. Proc. Nat. Acad. Sci. 86: 327-331, 1989.

[101839]

[101840]121. Trifiro, M.; Prior, L.; Pinsky, L.; Kaufman, M.; Chang, C.; Trapman,

[101841]J.; Brinkmann, A. O.; Kuiper, G. G. J. M.; Ris, C.: A single transition

[101842]at an exonic CpG site apparently abolishes androgen receptor (AR)-binding

[101843]activity in a family with complete androgen insensitivity (CAI). (Abstract) Am.

[101844]J. Hum. Genet. 45 (suppl.): A225, 1989.

[101845]

[101846]122. Trifiro, M.; Prior, R. L.; Sabbaghian, N.; Pinsky, L.;

Kaufman,

[101847]M.; Nylén, E. G.; Belsham, D. D.; Greenberg, C. R.; Wrogermann, K.

[101848]: Amber mutation creates a diagnostic Mael site in the androgen receptor

[101849]gene of a family with complete androgen insensitivity. Am. J. Med.

[101850]Genet. 40: 493–499, 1991.

[101851]

[101852]123. Tut, T. G.; Ghadessy, F. J.; Trifiro, M. A.; Pinsky, L.; Yong,

[101853]E. L.: Long polyglutamine tracts in the androgen receptor are associated

[101854]with reduced trans-activation, impaired sperm production, and male

[101855]infertility. J. Clin. Endocr. Metab. 82: 3777–3782, 1997.

[101856]

[101857]124. Visakorpi, T.; Hyytinen, E.; Koivisto, P.; Tanner, M.; Keinanen,

[101858]R.; Palmberg, C.; Palotie, A.; Tammela, T.; Isola, J.; Kallioniemi,

[101859]O.-P.: In vivo amplification of the androgen receptor gene

and progression

[101860] of human prostate cancer. *Nature Genet.* 9: 401–406,
1995.

[101861]

[101862] 125. Von Eckardstein, S.; Syska, A.; Gromoll, J.; Kamischke, A.; Simoni,

[101863] M.; Nieschlag, E.: Inverse correlation between sperm concentration

[101864] and number of androgen receptor CAG repeats in normal men. *J. Clin.*

[101865] *Endocr. Metab.* 86: 2585–2590, 2001.

[101866]

[101867] 126. Wang, Q.; Ghadessy, F. J.; Trounson, A.; de Kretser, D.; McLachlan,

[101868] R.; Ng, S. C.; Yong, E. L.: Azoospermia associated with a mutation

[101869] in the ligand-binding domain of an androgen receptor displaying normal

[101870] ligand binding, but defective trans-activation. *J. Clin. Endocr.*

[101871] *Metab.* 83: 4303–4309, 1998.

[101872]

[101873]127. Weidemann, W.; Peters, B.; Romalo, G.; Spindler, K.-D.; Schweikert, H.-U.: Response to androgen treatment in a patient with partial androgen insensitivity and a mutation in the deoxyribonucleic acid-binding domain of the androgen receptor. J. Clin. Endocr. Metab. 83: 1173-1176, 1998.

[101878]

[101879]128. Welch, W. J.; Diamond, M. I.: Glucocorticoid modulation of androgen receptor nuclear aggregation and cellular toxicity is associated with distinct forms of soluble expanded polyglutamine protein. Hum. Molec. Genet. 10: 3063-3074, 2001.

[101883]

[101884]129. Westberg, L.; Baghaei, F.; Rosmond, R.; Hellstrand, M.; Landen, M.; Jansson, M.; Holm, G.; Bjorntorp, P.; Eriksson, E.: Polymorphisms

[101886] of the androgen receptor gene and the estrogen receptor
beta gene

[101887] are associated with androgen levels in women. J. Clin. Endocr. Metab. 86:

[101888] 2562–2568, 2001.

[101889]

[101890] 130. Wieacker, P.; Breckwoldt, M.; Gal, A.: Testicular feminization:

[101891] diagnosis and search for closely linked restriction fragment length

[101892] polymorphism. Dis. Markers 3: 213–218, 1985.

[101893]

[101894] 131. Wieacker, P.; Griffin, J. E.; Wienker, T.; Lopez, J. M.; Wilson,

[101895] J. D.; Breckwoldt, M.: Linkage analysis with RFLPs in families with

[101896] androgen resistance syndromes: evidence for close linkage between

[101897] the androgen receptor locus and the DXS1 segment. Hum. Genet. 76:

[101898] 248–252, 1987.

[101899]

[101900]132. Wilson, C. M.; McPhaul, M. J.: A and B forms of the
androgen

[101901]receptor are present in human genital skin fibroblasts.
Proc. Nat.

[101902]Acad. Sci. 91: 1234–1238, 1994.

[101903]

[101904]133. Wilson, J. D.: The promiscuous receptor: prostate
cancer comes

[101905]of age. (Editorial) New Eng. J. Med. 332: 1440–1441,
1995.

[101906]

[101907]134. Wilson, J. D.; Carlson, B. R.; Weaver, D. D.; Kovacs, W.
J.;

[101908]Griffin, J. E.: Endocrine and genetic characterization of
cousins

[101909]with male pseudohermaphroditism: evidence that the Lubs
phenotype

[101910]can result from a mutation that alters the structure of the
androgen

[101911]receptor. Clin. Genet. 26: 363–370, 1984.

[101912]

[101913]135. Wooster, R.; Mangion, J.; Eeles, R.; Smith, S.;
Dowsett, M.;

[101914]Averill, D.; Barrett-Lee, P.; Easton, D. F.; Ponder, B. A. J.;
Stratton,

[101915]M. R.: A germline mutation in the androgen receptor gene
in two brothers

[101916]with breast cancer and Reifenstein syndrome. Nature
Genet. 2: 132–134,
[101917]1992.

[101918]

[101919]136. Zhang, L.; Leeflang, E. P.; Yu, J.; Arnheim, N.: Study-
ing human
[101920]mutations by sperm typing: instability of CAG trinucleotide repeats

[101921]in the human androgen receptor gene. Nature Genet. 7:
531–535, 1994.

[101922]

[101923]137. Zhu, Y.-S.; Cai, L.-Q.; Cordero, J. J.; Canovatchel, W.
J.; Katz,

[101924]M. D.; Imperato-McGinley, J.: A novel mutation in the CAG
triplet

[101925]region of exon 1 of androgen receptor gene causes complete androgen

[101926]insensitivity syndrome in a large kindred. J. Clin. Endocr.
Metab. 84:

[101927]1590–1594, 1999.

[101928]

[101929]138. Zoppi, S.; Wilson, C. M.; Harbison, M. D.; Griffin, J. E.;
Wilson,

[101930]J. D.; McPhaul, M. J.; Marcelli, M.: Complete testicular fem-
inization

[101931]caused by an amino-terminal truncation of the androgen
receptor with

[101932]downstream initiation. J. Clin. Invest. 91: 1105–1112,
1993.

[101933]17745.

[101934]Konig, A.; Happle, R.; Fink-Puches, R.; Soyer, H. P.; Born-
holdt,

[101935]D.; Engel, H.; Grzeschik, K.-H.: A novel missense mutation
of NSDHL

[101936]in an unusual case of CHILD syndrome showing bilateral,
almost symmetric

[101937]involvement. J. Am. Acad. Derm. 46: 594–596, 2002.

[101938]17746.

[101939]Labit-Le Bouteiller, C.; Jamme, M. F.; David, M.; Silve, S.;
Lanau,

[101940]C.; Dhers, C.; Picard, C.; Rahier, A.; Taton, M.; Loison, G.;
Caput,

[101941]D.; Ferrara, P.; Lupker, J.: Antiproliferative effects of

SR31747A

[101942]in animal cell lines are mediated by inhibition of cholesterol biosynthesis

[101943]at the sterol isomerase step. *Europ. J. Biochem.* 256: 342–349, 1998.

[101944]17747.

[101945]Liu, X. Y.; Dangel, A. W.; Kelley, R. I.; Zhao, W.; Denny, P.;

[101946]Botcherby, M.; Cattanach, B.; Peters, J.; Hunsicker, P. R.; Mallon,

[101947]A.–M.; Strivens, M. A.; Bate, R.; Miller, W.; Rhodes, M.; Brown, S.

[101948]D. M.; Herman, G. E.: The gene mutated in bare patches and striated

[101949]mice encodes a novel 3- β -hydroxysteroid dehydrogenase. *Nature*

[101950]*Genet.* 22: 182–187, 1999.

[101951]17748.

[101952]Schindelhauer, D.; Hellebrand, H.; Grimm, L.; Bader, I.; Meitinger,

[101953]T.; Wehnert, M.; Ross, M.; Meindl, A.: Long-range map of a 3.5-Mb

[101954]region in Xp11.23–22 with a sequence-ready map from a

1.1-Mb gene-rich

[101955]interval. *Genome Res.* 6: 1056–1069, 1996.

[101956]17749.

[101957]Silve, S.; Dupuy, P. H.; Labit-Lebouteiller, C.; Kaghad, M.;
Chalon,

[101958]P.; Rahier, A.; Taton, M.; Lupker, J.; Shire, D.; Loison, G.:
Emopamil-binding

[101959]protein, a mammalian protein that binds a series of struc-
turally diverse

[101960]neuroprotective agents, exhibits delta(8)-delta(7) sterol
isomerase

[101961]activity in yeast. *J. Biol. Chem.* 271: 22434–22440, 1996.

[101962]17750.

[101963]Traupe, H.; Muller, D.; Atherton, D.; Kalter, D. C.; Cremers,

[101964]F. P. M.; van Oost, B. A.; Ropers, H.-H.: Exclusion mapping
of the

[101965]X-linked dominant chondrodysplasia punctata/
ichthyosis/cataract/short

[101966]stature (Happle) syndrome: possible involvement of an
unstable pre-mutation. *Hum.*

[101967]*Genet.* 89: 659–665, 1992.

[101968]17751.

[101969]Jin, H.; Gardner, R. J.; Viswesvariah, R.; Muntoni, F.;
Roberts,

[101970]R. G.: Two novel members of the interleukin-1 receptor
gene family,

[101971]one deleted in Xp22.1-Xp21.3 mental retardation. *Europ.*
J. Hum. Genet. 8:

[101972]87-94, 2000.

[101973]17752.

[101974]Montini, E.; Buchner, G.; Spalluto, C.; Andolfi, G.; Caruso,
A.;

[101975]den Dunnen, J. T.; Trump, D.; Rocchi, M.; Ballabio, A.;
Franco, B.

[101976]: Identification of SCML2, a second human gene homolo-
gous to the *Drosophila*

[101977]sex comb on midleg (Scm): a new gene cluster on Xp22.
Genomics 58:

[101978]65-72, 1999.

[101979]17753.

[101980]Goto, M.; Imamura, O.; Kuromitsu, J.; Matsumoto, T.;
Yamabe, Y.;

- [101981]Tokutake, Y.; Suzuki, N.; Mason, B.; Drayna, D.; Sugawara, M.; Sugimoto, M.; Furuichi, Y.: Analysis of helicase gene mutations in Japanese Werner's syndrome patients. Hum. Genet. 99: 191–193, 1997.
- [101982]17754.
- [101983]Gray, M. D.; Shen, J.–C.; Kamath–Loeb, A. S.; Blank, A.; Sopher, B. L.; Martin, G. M.; Oshima, J.; Loeb, L. A.: The Werner syndrome protein is a DNA helicase. Nature Genet. 17: 100–103, 1997.
- [101984]17755.
- [101985]Imamura, O.; Ichikawa, K.; Yamabe, Y.; Goto, M.; Sugawara, M.; Furuichi, Y.: Cloning of a mouse homologue of the human Werner syndrome gene and assignment to 8A4 by fluorescence in situ hybridization. Genomics 41: 298–300, 1997.
- [101986]17756.

- [101994]Ishikawa, Y.; Sugano, H.; Matsumoto, T.; Furuichi, Y.;
Miller,
[101995]R. W.; Goto, M.: Unusual features of thyroid carcinomas in
Japanese
[101996]patients with Werner syndrome and possible genotype-
phenotype relations
[101997]to cell type and race. Cancer 85: 1345-1352, 1999.
[101998]17757.
- [101999]Marciniak, R. A.; Lombard, D. B.; Johnson, F. B.; Guarente,
L.
[102000]: Nucleolar localization of the Werner syndrome protein in
human cells. Proc.
[102001]Nat. Acad. Sci. 95: 6887-6892, 1998.
[102002]17758.
- [102003]Boyle, S.; Gilchrist, S.; Bridger, J. M.; Mahy, N. L.; Ellis, J.
[102004]A.; Bickmore, W. A.: The spatial organization of human
chromosomes
[102005]within the nuclei of normal and emerin-mutant cells.
Hum. Molec.
[102006]Genet. 10: 211-219, 2001.
[102007]17759.

- [102008]Clarke, A.: Hypohidrotic ectodermal dysplasia. J. Med. Genet. 24: 659–663, 1987.
- [102009]17760.
- [102010]17760.
- [102011]Cartegni, L.; Raffaele di Barletta, M.; Barresi, R.; Squarzoni, S.; Sabatelli, P.; Maraldi, N.; Mora, M.; Di Blasi, C.; Corne-lio,
- [102012]F.; Merlini, L.; Villa, A.; Cobianchi, F.; Toniolo, D.: Heart-specific
- [102013]localization of emerin: new insights into Emery–Dreifuss
- [102014]muscular
- [102015]dystrophy. Hum. Molec. Genet. 6: 2257–2264, 1997.
- [102016]17761.
- [102017]Consalez, G. G.; Thomas, N. S. T.; Stayton, C. L.; Knight, S. J.
- [102018]L.; Johnson, M.; Hopkins, L. C.; Harper, P. S.; Elsas, L. J.; Warren,
- [102019]S. T.: Assignment of Emery–Dreifuss muscular dystrophy to the distal
- [102020]region of Xq28: the results of a collaborative study. Am. J. Hum.

[102021]Genet. 48: 468–480, 1991.

[102022]17762.

[102023]Ellis, J. A.; Yates, J. R. W.; Kendrick-Jones, J.; Brown, C. A.

[102024]: Changes at P183 of emerin weaken its protein–protein interactions

[102025]resulting in X-linked Emery–Dreifuss muscular dystrophy. Hum. Genet. 104:

[102026]262–268, 1999.

[102027]17763.

[102028]Hodgson, S. V.; Boswinkel, E.; Walker, A.; Bobrow, M.; Davies,

[102029]K.; Dubowitz, V.; Granata, G.; Merlini, L.: Linkage analysis using

[102030]nine DNA polymorphisms along the length of the X chromosome locates

[102031]the gene for Emery–Dreifuss muscular dystrophy to distal Xq. (Abstract) J.

[102032]Med. Genet. 23: 169–170, 1986.

[102033]17764.

[102034]Klauck, S.; Wilgenbus, P.; Yates, J. R. W.; Muller, C. R.; Poustka,

[102035]A.: Identification of novel mutations in three families with
Emery–Dreifuss

[102036]muscular dystrophy. Hum. Mol. Genet. 4: 1853–1857,
1995.

[102037]17765.

[102038]Manilal, S.; Nguyen thi Man; Sewry, C. A.; Morris, G. E.:
The

[102039]Emery–Dreifuss muscular dystrophy protein, emerin, is a
nuclear membrane

[102040]protein. Hum. Molec. Genet. 5: 801–808, 1996.

[102041]17766.

[102042]Manilal, S.; Recan, D.; Sewry, C. A.; Hoeltzenbein, M.;
Llense,

[102043]S.; Leturcq, F.; Deburgrave, N.; Barbot, J.–C.; Nguyen thi
Man; Muntoni,

[102044]F.; Wehnert, M.; Kaplan, J.–C.; Morris, G. E.: Mutations in
Emery–Dreifuss

[102045]muscular dystrophy and their effects on emerin protein
expression. Hum.

[102046]Molec. Genet. 7: 855–864, 1998.

[102047]17767.

- [102048]Nevo, Y.; Al-Lozi, M.; Parsadanian, A. S.; Elliott, J. L.; Connolly,
muscular
[102049]A. M.; Pestronk, A.: Mutation analysis in Emery–Dreifuss
[102050]dystrophy. *Pediat. Neurol.* 21: 456–459, 1999.
[102051]17768.
- [102052]Romeo, G.; Roncuzzi, L.; Sangiorgi, S.; Giacanelli, M.;
Liguori,
[102053]M.; Tessarolo, D.; Rocchi, M.: Mapping of the Emery–
Dreifuss gene
[102054]through reconstruction of crossover points in two Italian
pedigrees. *Hum.*
[102055]*Genet.* 80: 59–62, 1988.
[102056]17769.
- [102057]Thomas, N. S. T.; Williams, H.; Elsas, L. J.; Hopkins, L. C.;
[102058]Sarfarazi, M.; Harper, P. S.: Localisation of the gene for
Emery–Dreifuss
[102059]muscular dystrophy to the distal long arm of the X chromosome. *J.*
[102060]*Med. Genet.* 23: 596–598, 1986.
[102061]17770.

- [102062]Wulff, K.; Parrish, J. E.; Herrmann, F. H.; Wehnert, M.: Six
[102063]novel mutations in the emerin gene causing X-linked
Emery–Dreifuss
[102064]muscular dystrophy. Hum. Mutat. 9: 526–530, 1997.
[102065]17771.
- [102066]Yamada, T.; Kobayashi, T.: A novel emerin mutation in a
Japanese
[102067]patient with Emery–Dreifuss muscular dystrophy. Hum.
Genet. 97:
[102068]693–694, 1996.
[102069]17772.
- [102070]Yates, J. R. W.; Affara, N. A.; Jamieson, D. M.; Ferguson–
Smith,
[102071]M. A.; Hausmanowa–Petrusewicz, I.; Zaremba, J.;
Borkowska, J.; Johnston,
[102072]A. W.; Kelly, K.: Emery–Dreifuss muscular dystrophy: lo-
calisation
[102073]to Xq27.3–qter confirmed by linkage to the factor VIII
gene. J. Med.
[102074]Genet. 23: 587–590, 1986.
[102075]17773.

- [102076]Yates, J. R. W.; Bagshaw, J.; Aksmanovic, V. M. A.;
Coomber, E.;
- [102077]McMahon, R.; Whittaker, J. L.; Morrison, P. J.; Kendrick-
Jones, J.;
- [102078]Ellis, J. A.: Genotype-phenotype analysis in X-linked
Emery-Dreifuss
[102079]muscular dystrophy and identification of a missense mu-
tation associated
[102080]with a milder phenotype. Neuromusc. Disord. 9: 159-165,
1999.
- [102081]17774.
- [102082]Yates, J. R. W.; Warner, J. P.; Smith, J. A.; Deymeer, F.;
Azulay,
[102083]J.-P.; Hausmanowa-Petrusewicz, I.; Zaremba, J.;
Borkowska, J.; Affara,
[102084]N. A.; Ferguson-Smith, M. A.: Emery-Dreifuss muscular
dystrophy:
[102085]linkage to markers in distal Xq28. J. Med. Genet. 30:
108-111, 1993.
- [102086]17775.
- [102087]Yorifuji, H.; Tadano, Y.; Tsuchiya, Y.; Ogawa, M.; Goto, K.;
Umetani,

[102088]A.; Asaka, Y.; Arahata, K.: Emerin, deficiency of which causes Emery–Dreifuss

[102089]muscular dystrophy, is localized at the inner nuclear membrane. Neurogenetics 1:

[102090]135–140, 1997.

[102091]17776.

[102092]Allen, R. C.; Armitage, R. J.; Conley, M. E.; Rosenblatt, H.; Jenkins,

[102093]N. A.; Copeland, N. G.; Bedell, M. A.; Edelhoff, S.; Disteche, C.

[102094]M.; Simoneaux, D. K.; Fanslow, W. C.; Belmont, J.; Spriggs, M. K.

[102095]: CD40 ligand gene defects responsible for X-linked hyper-IgM syndrome. Science 259:

[102096]990–993, 1993.

[102097]17777.

[102098]Aruffo, A.; Farrington, M.; Hollenbaugh, D.; Li, X.; Milatovich,

[102099]A.; Nonoyama, S.; Bajorath, J.; Grosmaire, L. S.; Stenkamp, R.; Neubauer,

[102100]M.; Roberts, R. L.; Noelle, R. J.; Ledbetter, J. A.; Francke, U.;

- [102101] Ochs, H. D.: The CD40 ligand, gp39, is defective in activated T cells
from patients with X-linked hyper-IgM syndrome. Cell 72: 291-300,
1993.
- [102103] 17778.
- [102104] 17778.
- [102105] DiSanto, J. P.; Bonnefoy, J. Y.; Gauchat, J. F.; Fischer, A.; de
Saint Basile, G.: CD40 ligand mutations in X-linked immunodeficiency
with hyper-IgM. Nature 361: 541-543, 1993.
- [102107] 17779.
- [102108] 17779.
- [102109] Gauchat, J.-F.; Aubry, J.-P.; Mazzei, G.; Life, P.; Jomotte, T.;
- [102110] Elson, G.; Bonnefoy, J.-Y.: Human CD40-ligand: molecular cloning,
cellular distribution and regulation of expression by factors controlling
IgE production. FEBS Lett. 315: 259-266, 1993.
- [102112] 17780.
- [102113] 17780.
- [102114] Graf, D.; Korthauer, U.; Mages, H. W.; Senger, G.; Kroczeck, R.

[102115]A.: Cloning of TRAP, a ligand for CD40 on human T cells.
Europ.

[102116]J. Immun. 22: 3191–3194, 1992.

[102117]17781.

[102118]Hollenbaugh, D.; Grosmaire, L. S.; Kullas, C. D.; Chalupny,
N.

[102119]J.; Braesch-Andersen, S.; Noelle, R. J.; Stamenkovic, I.;
Ledbetter,

[102120]J. A.; Aruffo, A.: The human T cell antigen gp39, a mem-
ber of the

[102121]TNF gene family, is a ligand for the CD40 receptor: ex-
pression of

[102122]a soluble form of gp39 with B cell co-stimulatory activity.
EMBO

[102123]J. 11: 4313–4321, 1992.

[102124]17782.

[102125]Hors-Cayla, M. C.; Heuertz, S.; Van Cong, N.; Benne, F.:
Cattle

[102126]gene mapping by somatic cell hybridization. (Abstract)
Cytogenet.

[102127]Cell Genet. 25: 165–166, 1979.

[102128]17783.

[102129]Biamonti, G.; Ruggiu, M.; Saccone, S.; Della Valle, G.; Riva, S.

[102130]: Two homologous genes, originated by duplication, encode the human

[102131]hnRNP proteins A2 and A1. Nucleic Acids Res. 22: 1996–2002, 1994.

[102132]17784.

[102133]Clarke, A.; Burn, J.: Sweat testing to identify female carriers

[102134]of X linked hypohidrotic ectodermal dysplasia. J. Med. Genet. 28:

[102135]330–333, 1991.

[102136]17785.

[102137]Clarke, A.; Sarfarazi, M.; Thomas, N. S. T.; Roberts, K.; Harper,

[102138]P. S.: X-linked hypohidrotic ectodermal dysplasia: DNA probe linkage

[102139]analysis and gene localization. Hum. Genet. 75: 378–380, 1987.

[102140]17786.

[102141]Clarke, A.; Sarfarazi, M.; Zonana, J.; Thomas, N. S. T.;
Roberts,

[102142]K.; Harper, P. S.: Regional localization of X-linked hypo-
hidrotic

[102143]ectodermal dysplasia (EDA). (Abstract) Cytogenet. Cell
Genet. 46:

[102144]594, 1987.

[102145]17787.

[102146]Cohen, M. M.; Lin, C. C.; Sybert, V.; Orecchio, E. J.: Two
human

[102147]X-autosome translocations identified by autoradiography
and fluorescence. Am.

[102148]J. Hum. Genet. 24: 583-597, 1972.

[102149]17788.

[102150]Crawford, P. J. M.; Aldred, M. J.; Clarke, A.: Clinical and ra-
diographic

[102151]dental findings in X linked hypohidrotic ectodermal dys-
plasia. J.

[102152]Med. Genet. 28: 181-185, 1991.

[102153]17789.

[102154]Darwin, C.: The Variation of Animals and Plants under Do-

mestication.

[102155]London: John Murray (pub.) (2nd ed.): 1875. Pp. 319 only.

[102156]17790.

[102157]de la Chapelle, A.: The map of the human X chromosome. In: Crosignani,

[102158]P. G.; Fraccaro, M.; Rubin, B. L.: Genetic Control of Gamete Production

[102159]and Function. New York: Grune and Stratton (pub.) 1982. Pp. 33–48.

[102160]17791.

[102161]de la Chapelle, A.: Personal Communication. Helsinki, Finland

[102162]9/18/1990.

[102163]17792.

[102164]Durmowicz, M. C.; Cui, C.-Y.; Schlessinger, D.: The EDA gene

[102165]is a target of, but does not regulate Wnt signaling. Gene 285: 203–211,

[102166]2002.

[102167]17793.

[102168]Elomaa, O.; Pulkkinen, K.; Hannelius, U.; Mikkola, M.;

Saarialho-Kere,

[102169]U.; Kere, J.: Ectodysplasin is released by proteolytic shedding and

[102170]binds to the EDAR protein. Hum. Molec. Genet. 10: 953–962, 2001.

[102171]17794.

[102172]Ezer, S.; Bayes, M.; Elomaa, O.; Schlessinger, D.; Kere, J.:

[102173]Ectodysplasin is a collagenous trimeric type II membrane protein with

[102174]a tumor necrosis factor-like domain and co-localizes with cytoskeletal

[102175]structures at lateral and apical surfaces of cells. Hum. Molec. Genet. 8:

[102176]2079–2086, 1999.

[102177]17795.

[102178]Familusi, J. B.; Jaiyesimi, F.; Ojo, C. O.; Attah, E. B.: Hereditary

[102179]anhidrotic ectodermal dysplasia: studies in a Nigerian family. Arch.

[102180]Dis. Child. 50: 642–647, 1975.

[102181]17796.

- [102182]Ferguson, B. M.; Brockdorff, N.; Formstone, E.; Ngyuen, T.;
Kronmiller,
[102183]J. E.; Zonana, J.: Cloning of Tabby, the murine homolog of
the human
[102184]EDA gene: evidence for a membrane-associated protein
with a short
[102185]collagenous domain. Hum. Molec. Genet. 6: 1589–1594,
1997.
[102186]17797.
- [102187]Ferguson, B. M.; Thomas, N. S. T.; Munoz, F.; Morgan, D.;
Clarke,
[102188]A.; Zonana, J.: Scarcity of mutations detected in families
with X
[102189]linked hypohidrotic ectodermal dysplasia: diagnostic im-
plications. J.
[102190]Med. Genet. 35: 112–115, 1998.
[102191]17798.
- [102192]Filippi, G.; Rinaldi, A.; Crisponi, G. L.; Siniscalco, M.: X-
mapping
[102193]in man: evidence against measurable linkage between an-
hidrotic ectodermal
[102194]dysplasia and G6PD deficiency. J. Med. Genet. 16:

223-226, 1979.

[102195]17799.

[102196]Fox, L. A.: X-linked anhidrotic ectodermal dysplasia manifesting

[102197]in a female. Birth Defects Orig. Art. Ser. X(4): 319-320, 1974.

[102198]17800.

[102199]Freire-Maia, N.; Pinheiro, M.: So-called 'anhidrotic ectodermal

[102200]dysplasia.'. Int. J. Derm. 19: 455-456, 1980.

[102201]17801.

[102202]Gerald, P. S.; Brown, J. A.: Report of the committee on the genetic

[102203]constitution of the X chromosome. Cytogenet. Cell Genet. 13: 29-34,

[102204]1974.

[102205]17802.

[102206]Gilgenkrantz, S.; Blanchet-Bardon, C.; Nazzaro, V.; Formiga, L.;

[102207]Mujica, P.; Alembik, Y.: Hypohidrotic ectodermal dysplasia: clinical

[102208]study of a family of 30 over three generations. Hum.

Genet. 81:

[102209]120-122, 1989.

[102210]17803.

[102211]Grant, R.; Falls, H. F.: Anodontia: report of a case associated

[102212]with ectodermal dysplasia of the anhidrotic type. Am. J. Orthodont. 30:

[102213]661-672, 1944.

[102214]17804.

[102215]Graves, R.: The Whitaker Negroes. In: Encounters: An anthropology

[102216]from the first ten years of Encounter Magazine.: Spender, S.; Kristol,

[102217]J.; Lasky, M. J., eds. New York: Basic Books, Inc. (pub.) Pp.

[102218]92-105: 1963.

[102219]17805.

[102220]Halperin, S. L.; Curtis, G. M.: Anhidrotic ectodermal dysplasia

[102221]associated with mental deficiency. Am. J. Ment. Defic. 46: 459-463,

[102222]1942.

[102223]17806.

[102224]Hanauer, A.; Alembik, Y.; Arveiler, B.; Formiga, L.;

Gilgenkrantz,

[102225]S.; Mandel, J. L.: Genetic mapping of hypohidrotic ecto-

dermal dysplasia:

[102226]linkage to DXS159 and DXYS1. (Abstract) Cytogenet. Cell

Genet. 46:

[102227]626 only, 1987.

[102228]17807.

[102229]Hanauer, A.; Alembik, Y.; Arveiler, B.; Formiga, L.;

Gilgenkrantz,

[102230]S.; Mandel, J. L.: Genetic mapping of anhidrotic ectoder-

mal dysplasia:

[102231]DXS159, a closely linked proximal marker. Hum. Genet.

80: 177-180,

[102232]1988.

[102233]17808.

[102234]Happle, R.: Association of pigmentary anomalies with

chromosomal

[102235]and genetic mosaicism and chimerism. (Letter) Am. J.

Hum. Genet. 48:

[102236]1013-1014, 1991.

[102237]17809.

[102238]Happle, R.; Frosch, P. J.: Manifestation of the lines of
Blaschko

[102239]in women heterozygous for X-linked hypohidrotic ecto-
dermal dysplasia. Clin.

[102240]Genet. 27: 468-471, 1985.

[102241]17810.

[102242]Harper, P. S.: Personal Communication. Cardiff, Wales
6/24/1986.

[102243]17811.

[102244]Hutt, F. B.: An earlier record of the toothless men of Sind.
J.

[102245]Hered. 26: 65-66, 1935.

[102246]17812.

[102247]Jespersen, H. G.: Hereditary ectodermal dysplasia of an-
hidrotic

[102248]type. Acta Paediat. 51: 712-720, 1962.

[102249]17813.

[102250]Kapalanga, J.; Blecher, S. R.: Effect of the X-linked gene
Tabby

[102251](Ta) on eyelid opening and incisor eruption in neonatal
mice is opposite

[102252]to that of epidermal growth factor. Development 108:
349–355, 1990.

[102253]17814.

[102254]Kere, J.; Srivastava, A. K.; Montonen, O.; Zonana, J.;
Thomas,

[102255]N.; Ferguson, B; Munoz, F.; Morgan, D.; Clarke, A.; Bay-
bayan, P.;

[102256]Chen, E. Y.; Ezer, S.; Saarialho-Kere, U.; de la Chapelle, A.;
Schlessinger,

[102257]D.: X-linked anhidrotic (hypohidrotic) ectodermal dyspla-
sia is caused

[102258]by mutation in a novel transmembrane protein. Nature
Genet. 13:

[102259]409–416, 1996.

[102260]17815.

[102261]Kerr, C. B.; Wells, R. S.; Cooper, K. E.: Gene effect in carri-
ers

[102262]of anhidrotic ectodermal dysplasia. J. Med. Genet. 3:

169–176, 1966.

[102263]17816.

[102264]Kleinebrecht, J.; Degenhardt, K.-H.; Grubisic, A.; Gunther, E.;

[102265]Svejcar, J.: Sweat pore counts in ectodermal dysplasias. Hum. Genet. 57:

[102266]437–439, 1981.

[102267]17817.

[102268]Kline, A. H.; Sidbury, J. B., Jr.; Richter, C. P.: The occurrence

[102269]of ectodermal dysplasia and corneal dysplasia in one family. J. Pediat. 55:

[102270]355–366, 1959.

[102271]17818.

[102272]Kobiela, K.; Kobiela, A.; Roszkiewicz, J.; Wierzba, J.; Limon,

[102273]J.; Trzeciak, W. H.: Mutations in the EDA gene in three unrelated

[102274]families reveal no apparent correlation between phenotype and genotype

[102275]in the patients with an X-linked anhidrotic ectodermal

dysplasia. Am.

[102276]J. Med. Genet. 100: 191–197, 2001.

[102277]17819.

[102278]Stoddart, K. L.; Jermak, C.; Nagaraja, R.; Schlessinger, D.;
Bech–Hansen,

[102279]N. T.: Physical map covering a 2 Mb region in human
Xp11.3 distal

[102280]to DX6849. Gene 227: 111–116, 1999.

[102281]17820.

[102282]Swanson, D. A.; Freund, C. L.; Ploder, L.; McInnes, R. R.;
Valle,

[102283]D.: A ubiquitin C–terminal hydrolase gene on the proximal
short arm

[102284]of the X chromosome: implications for X–linked retinal
disorders. Hum.

[102285]Molec. Genet. 5: 533–538, 1996.

[102286]17821.

[102287]Olinsky, S.; Loop, B. T.; DeKosky, A.; Ripepi, B.; Weng, W.;
Cummins,

[102288]J.; Wenger, S. L.; Yan, Y.; Lagenaur, C.; Narayanan, V.:
Chromosomal

[102289]mapping of the human M6 genes. Genomics 33: 532–536, 1996.

[102290]17822.

[102291]Yan, Y.; Lagenaur, C.; Narayanan, V.: Molecular cloning of M6:

[102292]identification of a PLP/DM20 gene family. Neuron 11: 423–431, 1993.

[102293]17823.

[102294]Ballabio, A.; Parenti, G.; Napolitano, E.; Di Natale, P.; Andria,

[102295]G.: Genetic complementation of steroid sulphatase after somatic cell

[102296]hybridization of X-linked ichthyosis and multiple sulphatase deficiency. Hum.

[102297]Genet. 70: 315–317, 1985.

[102298]17824.

[102299]Nan, X.; Ng, H.–H.; Johnson, C. A.; Laherty, C. D.; Turner, B.

[102300]M.; Eisenman, R. N.; Bird, A.: Transcriptional repression by the

[102301]methyl–CpG-binding protein MeCP2 involves a histone deacetylase complex. Nature 393:

[102302]386-389, 1998.

[102303]17825.

[102304]Nan, X.; Tate, P.; Li, E.; Bird, A.: DNA methylation specifies

[102305]chromosomal localization of MeCP2. Molec. Cell. Biol. 16:
414-421,

[102306]1996.

[102307]17826.

[102308]Arn, P.; Hauser, B.; Maestri, N.; Herman, G.; Thomas, G.
H.; Brusilow,

[102309]S. W.: Hyperammonemic coma in adult females heterozy-
gous at the ornithine

[102310]transcarbamylase locus (OTCDH). (Abstract) Am. J. Hum.
Genet. 45

[102311](suppl.): A3, 1989.

[102312]17827.

[102313]Andersson, M.; Page, D. C.; Pettay, D.; Subrt, I.; Turleau,
C.;

[102314]de Grouchy, J.; de la Chapelle, A.: Y;autosome transloca-
tions and

[102315]mosaicism in the aetiology of 45,X maleness: assignment
of fertility

[102316]factor to distal Yq11. Hum. Genet. 79: 2–7, 1988.

[102317]17828.

[102318]Chandley, A. C.; Gosden, J. R.; Hargreave, T. B.; Spowart, G.;

[102319]Speed, R. M.; McBeath, S.: Deleted Yq in the sterile son of a man

[102320]with a satellited Y chromosome (Yqs). J. Med. Genet. 26: 145–153,

[102321]1989.

[102322]17829.

[102323]Robledo, R.; Melis, P.; Schillinger, E.; Casciano, I.; Balazs,

[102324]I.; Rinaldi, A.; Siniscalco, M.; Filippi, G.: X-linked ichthyosis

[102325]without STS deficiency: clinical, genetical, and molecular studies. Am.

[102326]J, Med. Genet. 59: 143–148, 1995.

[102327]17830.

[102328]Franco, B.; Meroni, G.; Parenti, G.; Levilliers, J.; Bernard, L.;

[102329]Gebbia, M.; Cox, L.; Maroteaux, P.; Sheffield, L.; Rappold, G. A.;

[102330]Andria, G.; Petit, C.; Ballabio, A.: A cluster of sulfatase genes

[102331]on Xp22.3: mutations in chondrodysplasia punctata (CDPX) and implications

[102332]for warfarin embryopathy. Cell 81: 1–20, 1995.

[102333]17831.

[102334]Meroni, G.; Franco, B.; Archidiacono, N.; Messali, S.; Andolfi,

[102335]G.; Rocchi, M.; Ballabio, A.: Characterization of a cluster of sulfatase

[102336]genes on Xp22.3 suggests gene duplications in an ancestral pseudoautosomal

[102337]region. Hum. Molec. Genet. 5: 423–431, 1996.

[102338]17832.

[102339]Franco, B.; Meroni, G.; Parenti, G.; Levilliers, J.; Bernard, L.;

[102340]Gebbia, M.; Cox, L.; Maroteaux, P.; Sheffield, L.; Rappold, G. A.;

[102341]Andria, G.; Petit, C.; Ballabio, A.: A cluster of sulfatase genes

[102342]on Xp22.3: mutations in chondrodysplasia punctata (CDPX) and implications

[102343]for warfarin embryopathy. Cell 81: 15–20, 1995.

[102344]17833.

[102345]Puca, A. A.; Zollo, M.; Repetto, M.; Andolfi, G.; Guffanti, A.;

[102346]Simon, G.; Ballabio, A.; Franco, B.: Identification by shot–
gun sequencing,

[102347]genomic organization, and functional analysis of a fourth
arylsulfatase

[102348]gene (ARSF) from the Xp22.3 region. Genomics 42:
192–199, 1997.

[102349]17834.

[102350]Adler, D. A.; Quaderi, N. A.; Brown, S. D. M.; Chapman, V.
M.;

[102351]Moore, J.; Tate, P.; Disteché, C. M.: The X-linked methy–
lated DNA

[102352]binding protein, Mecp2, is subject to X inactivation in the
mouse. Mammalian

[102353]Genome 6: 491–492, 1995.

[102354]17835.

[102355]Amano, K.; Nomura, Y.; Segawa, M.; Yamakawa, K.: Muta–
tional analysis

[102356]of the MECP2 gene in Japanese patients with Rett syn–

drome. J. Hum.

[102357]Genet. 45: 231–236, 2000.

[102358]17836.

[102359]Amir, R. E.; Van den Veyver, I. B.; Wan, M.; Tran, C. Q.;
Francke,

[102360]U.; Zoghbi, H. Y.: Rett syndrome is caused by mutations in
X-linked

[102361]MECP2, encoding methyl–CpG–binding protein 2. Nature
Genet. 23:

[102362]185–188, 1999.

[102363]17837.

[102364]Ballestar, E.; Yusufzai, T. M.; Wolffe, A. P.: Effects of Rett

[102365]syndrome mutations of the methyl–CpG binding domain
of the transcriptional

[102366]repressor MeCP2 on selectivity for association with
methylated DNA. Biochemistry 39:

[102367]7100–7106, 2000.

[102368]17838.

[102369]Balmer, D.; Arredondo, J.; Samaco, R. C.; LaSalle, J. M.:
MECP2

[102370]mutations in Rett syndrome adversely affect lymphocyte

growth, but

[102371]do not affect imprinted gene expression in blood or brain.

Hum. Genet. 110:

[102372]545–552, 2002.

[102373]17839.

[102374]Bienvenu, T.; Carrie, A.; de Roux, N.; Vinet, M.-C.; Jon-
veaux,

[102375]P.; Couvert, P.; Villard, L.; Arzimanoglou, A.; Beldjord, C.;
Fontes,

[102376]M.; Tardieu, M.; Chelly, J.: MECP2 mutations account for
most cases

[102377]of typical forms of Rett syndrome. Hum. Molec. Genet. 9:
1377–1384,

[102378]2000.

[102379]17840.

[102380]van de Vosse, E.; Walpole, S. M.; Nicolaou, A.; van der
Bent, P.;

[102381]Cahn, A.; Vaudin, M.; Ross, M. T.; Durham, J.; Pavitt, R.;
Wilkinson,

[102382]J.; Grafham, D.; Bergen, A. A. B.; van Ommen, G.-J. B.;
Yates, J.

[102383]R. W.; den Dunnen, J. T.; Trump, D.: Characterization of

SCML1, a

[102384]new gene in Xp22, with homology to developmental poly-comb genes. Genomics 49:

[102385]96–102, 1998.

[102386]17841.

[102387]Ballabio, A.; Bardoni, B.; Carrozzo, R.; Andria, G.; Bick, D.;

[102388]Campbell, L.; Hamel, B.; Ferguson-Smith, M. A.; Gimelli, G.; Fraccaro,

[102389]M.; Maraschio, P.; Zuffardi, O.; Guioli, S.; Camerino, G.: Contiguous

[102390]gene syndromes due to deletions in the distal short arm of the human

[102391]X chromosome. Proc. Nat. Acad. Sci. 86: 10001–10005, 1989.

[102392]17842.

[102393]Fujikawa-Adachi, K.; Nishimori, I.; Taguchi, T.; Onishi, S.: Human

[102394]mitochondrial carbonic anhydrase VB: cDNA cloning, mRNA expression,

[102395]subcellular localization, and mapping to chromosome X. J. Biol. Chem. 274:

[102396]21228–21233, 1999.

[102397]17843.

[102398]Shah, G. N.; Hewett-Emmett, D.; Grubb, J. H.; Migas, M. C.; Fleming,

[102399]R. E.; Waheed, A.; Sly, W. S.: Mitochondrial carbonic anhydrase CA

[102400]VB: differences in tissue distribution and pattern of evolution from

[102401]those of CA VA suggest distinct physiological roles. Proc. Nat. Acad.

[102402]Sci. 97: 1677-1682, 2000.

[102403]17844.

[102404]Nagase, T.; Seki, N.; Ishikawa, K.; Ohira, M.; Kawarabayashi, Y.;

[102405]Ohara, O.; Tanaka, A.; Kotani, H.; Miyajima, N.; Nomura, N.: Prediction

[102406]of the coding sequences of unidentified human genes. VI. The coding

[102407]sequences of 80 new genes (KIAA0201-KIAA0280) deduced by analysis

[102408]of cDNA clones from cell line KG-1 and brain. DNA Res. 3: 321-329,

[102409]1996. Note: Supplement: DNA Res. 3: 341-354, 1996.

[102410]17845.

[102411]Bowling, F.; McGown, I.; McGill, J.; Cowley, D.; Tuchman, M.:

[102412]Maternal gonadal mosaicism causing ornithine transcarbamylase deficiency. Am.

[102413]J. Med. Genet. 85: 452-454, 1999.

[102414]17846.

[102415]Bruton, C. J.; Corsellia, J. A. N.; Russell, A.: Hereditary hyperammonemia. Brain 93:

[102416]423-434, 1970.

[102417]17847.

[102418]Burdakin, J. H.; Norum, R. A.: Recombination between loci for

[102419]ornithine transcarbamylase (OTC) deficiency and G6PD. (Abstract) Am.

[102420]J. Hum. Genet. 33: 38A, 1981.

[102421]17848.

[102422]Asao, H.; Okuyama, C.; Kumaki, S.; Ishii, N.; Tsuchiya, S.; Foster,

[102423]D.; Sugamura, K.: Cutting edge: the common gamma-chain is an indispensable

[102424]subunit of the IL-21 receptor complex. J. Immun. 167:
1-5, 2001.

[102425]17849.

[102426]Arn, P. H.; Hauser, E. R.; Thomas, G. H.; Herman, G.; Hess,
D.;

[102427]Brusilow, S. W.: Hyperammonemia in women with a muta-
tion at the ornithine

[102428]carbamoyltransferase locus: a cause of postpartum coma.
New Eng.

[102429]J. Med. 322: 1652-1655, 1990.

[102430]17850.

[102431]Batshaw, M. L.; Roan, Y.; Jung, A. L.; Rosenberg, L. A.;
Brusilow,

[102432]S. W.: Cerebral dysfunction in asymptomatic carriers of
ornithine

[102433]transcarbamylase deficiency. New Eng. J. Med. 302:
482-485, 1980.

[102434]17851.

[102435]Bonaiti-Pellie, C.; Pelet, A.; Ogier, H.; Nelson, J.-R.;
Largilliere,

[102436]C.; Berthelot, J.; Saudubray, J.-M.; Munnich, A.: A probable
sex

[102437] difference in mutation rates in ornithine transcarbamylase deficiency. Hum.

[102438] Genet. 84: 163–166, 1990.

[102439] 17852.

[102440] Campbell, A. G. M.; Rosenberg, L. E.; Snodgrass, P. J.;
Nuzum,

[102441] C. T.: Lethal neonatal hyperammonaemia due to complete
ornithine–transcarbamylase

[102442] deficiency. (Letter) Lancet II: 217–218, 1971.

[102443] 17853.

[102444] Campbell, A. G. M.; Rosenberg, L. E.; Snodgrass, P. J.;
Nuzum,

[102445] C. T.: Ornithine transcarbamylase deficiency: a cause of
lethal neonatal

[102446] hyperammonemia in males. New Eng. J. Med. 288: 1–6,
1973.

[102447] 17854.

[102448] Carstens, R. P.; Fenton, W. A.; Rosenberg, L. R.: Identifica-
tion

[102449] of RNA splicing errors resulting in human ornithine tran-
scarbamylase

[102450]deficiency. Am. J. Hum. Genet. 48: 1105–1114, 1991.

[102451]17855.

[102452]Cathelineau, L.; Saudubray, J.–M.; Polonovski, C.: Hetero-
geneous

[102453]mutations of the structural gene of human ornithine car-
bamyltransferase

[102454]as observed in five personal cases. Enzyme 18: 103–113,
1974.

[102455]17856.

[102456]Cavard, C.; Grimber, G.; Dubois, N.; Chasse, J.–F.; Ben-
noun, M.;

[102457]Minet–Thuriaux, M.; Kamoun, P.; Briand, P.: Correction of
mouse ornithine

[102458]transcarbamylase deficiency by gene transfer into the
germ line. Nucleic

[102459]Acids Res. 16: 2099–2110, 1988.

[102460]17857.

[102461]Clayton–Smith, J.; Donnai, D.: Fetal valproate syndrome. J.

[102462]Med. Genet. 32: 724–727, 1995.

[102463]17858.

[102464]Corey, L.; Rubin, R. J.; Hattwick, M. A. W.; Noble, G. R.;
Cassidy,

[102465]E.: A nationwide outbreak of Reye's syndrome: its epidemiologic relationship

[102466]to influenza B. Am. J. Med. 61: 615-625, 1976.

[102467]17859.

[102468]DeMars, R.; LeVan, S. L.; Trend, B. L.; Russell, L. B.: Abnormal

[102469]ornithine carbamyltransferase in mice having the sparse-fur mutation. Proc.

[102470]Nat. Acad. Sci. 73: 1693-1697, 1976.

[102471]17860.

[102472]Drogari, E.; Leonard, J. V.: Late onset ornithine carbamoyl transferase

[102473]deficiency in males. Arch. Dis. Child. 63: 1363-1367, 1988.

[102474]17861.

[102475]Faiella, A.; Wernig, M.; Consalez, G. G.; Hostick, U.; Hofmann,

[102476]C.; Hustert, E.; Boncinelli, E.; Balling, R.; Nadeau, J. H.: A mouse

- [102477]model for valproate teratogenicity: parental effects,
homeotic transformations,
[102478]and altered HOX expression. Hum. Molec. Genet. 9:
227–236, 2000.
[102479]17862.
- [102480]Feldmann, D.; Rozet, J.–M.; Pelet, A.; Hentzen, D.; Briand,
P.;
[102481]Hubert, P.; Largilliere, C.; Rabier, D.; Farriaux, J.–P.;
Munnich,
[102482]A.: Site specific screening for point mutations in ornithine
transcarbamylase
[102483]deficiency. J. Med. Genet. 29: 471–475, 1992.
[102484]17863.
- [102485]Finkelstein, J. E.; Francomano, C. A.; Brusilow, S. W.;
Traystman,
[102486]M. D.: Use of denaturing gradient gel electrophoresis for
detection
[102487]of mutation and prospective diagnosis in late onset or–
nithine transcarbamylase
[102488]deficiency. Genomics 7: 167–172, 1990.
[102489]17864.

[102490]Finkelstein, J. E.; Hauser, E.; Brusilow, S. W.: Late onset ornithine

[102491]transcarbamylase deficiency (OTCD) in males. (Abstract)
Am. J. Hum.

[102492]Genet. 45 (suppl.): A5, 1989.

[102493]17865.

[102494]Finkelstein, J. E.; Hauser, E. R.; Leonard, C. O.; Brusilow, S.

[102495]W.: Late-onset ornithine transcarbamylase deficiency in
male patients. J.

[102496]Pediat. 117: 897-902, 1990.

[102497]17866.

[102498]Fox, J. E.; Hack, A. M.; Fenton, W. A.; Golbus, M. S.; Winter,

[102499]S.; Kalousek, F.; Rozen, R.; Brusilow, S. W.; Rosenberg, L.
E.: Prenatal

[102500]diagnosis of ornithine transcarbamylase deficiency with
use of DNA

[102501]polymorphisms. New Eng. J. Med. 315: 1205-1208, 1986.

[102502]17867.

[102503]Fox, J. E.; Hack, A. M.; Fenton, W. A.; Rosenberg, L. E.:
Identification

[102504]and application of additional restriction fragment length polymorphisms

[102505]at the human ornithine transcarbamylase locus. Am. J. Hum. Genet. 38:

[102506]841–847, 1986.

[102507]17868.

[102508]Garcia-Perez, M. A.; Sanjurjo, P.; Rubio, V.: Demonstration of

[102509]the spf-ash mutation in Spanish patients with ornithine transcarbamylase

[102510]deficiency of moderate severity. Hum. Genet. 95: 183–186, 1995.

[102511]17869.

[102512]Gelehrter, T. D.; Rosenberg, L. E.: Ornithine transcarbamylase

[102513]deficiency: unsuccessful therapy of neonatal hyperammonemia with N-carbamyl-L-glutamate

[102514]and L-arginine. New Eng. J. Med. 292: 351–352, 1975.

[102515]17870.

[102516]Genet, S.; Cranston, T.; Middleton-Price, H. R.: Mutation detection

[102517]in 65 families with a possible diagnosis of ornithine carbamoyltransferase
[102518]deficiency including 14 novel mutations. J. Inherit. Metab. Dis. 23:
[102519]669–676, 2000.
[102520]17871.
[102521]Gilbert–Dussardier, B.; Segues, B.; Rozet, J.–M.; Rabier, D.;
[102522]Calvas, P.; de Lumley, L.; Bonnefond, J.–P.; Munnich, A.:
Partial
[102523]duplication [dup. TCAC (178)] and novel point mutations
(T125M, G188R,
[102524]A209V, and H302L) of the ornithine transcarbamylase
gene in congenital
[102525]hyperammonemia. Hum. Mutat. 8: 74–76, 1996.
[102526]17872.
[102527]Gilchrist, J. M.; Coleman, R. A.: Ornithine transcarbamylase
[102528]deficiency: adult onset of severe symptoms. Ann. Intern. Med. 106:
[102529]556–558, 1987.
[102530]17873.

- [102531]Glick, T. H.; Likosky, W. H.; Levitt, L. P.; Mellin, H.;
Reynolds,
[102532]D. W.: Reye's syndrome: an epidemiologic approach. *Pedi-*
iatrics 46:
[102533]371–377, 1970.
[102534]17874.
- [102535]Grompe, M.; Caskey, C. T.; Fenwick, R. G.: Improved
molecular
[102536]diagnostics for ornithine transcarbamylase deficiency. *Am.*
J. Hum.
[102537]*Genet.* 48: 212–222, 1991.
[102538]17875.
- [102539]Grompe, M.; Muzny, D. M.; Caskey, C. T.: Scanning detec-
tions
[102540]of mutations in human ornithine transcarbamylase defi-
ciency (OTC)
[102541]by chemical mismatch cleavage. (Abstract) *Am. J. Hum.*
Genet. 45
[102542](suppl.): A192, 1989.
[102543]17876.
- [102544]Ogata, T.; Wakui, K.; Muroya, K.; Ohashi, H.; Matsuo, N.;

Brown,

[102545]D. M.; Ishii, T.; Fukushima, Y.: Microphthalmia with linear skin

[102546]defects syndrome in a mosaic female infant with monosomy for the Xp22

[102547]region: molecular analysis of the Xp22 breakpoint and the X-inactivation

[102548]pattern. Hum. Genet. 103: 51-56, 1998.

[102549]17877.

[102550]Lankes, W.; Griesmacher, A.; Grunwald, J.; Schwartz-Albiez, R.;

[102551]Keller, R.: A heparin-binding protein involved in inhibition of smooth-muscle

[102552]cell proliferation. Biochem. J. 251: 831-842, 1988.

[102553]17878.

[102554]Lankes, W. T.; Furthmayr, H.: Moesin: a member of the protein

[102555]4.1-talin-ezrin family of proteins. Proc. Nat. Acad. Sci. 88: 8297-8301,

[102556]1991.

[102557]17879.

- [102558] Shcherbina, A.; Bretscher, A.; Rosen, F. S.; Kenney, D. M.; Remold-O'Donnell, E.: The cytoskeletal linker protein moesin: decreased levels in Wiskott-Aldrich syndrome platelets and identification of a cleavage pathway in normal platelets. *Brit. J. Haemat.* 106: 216-223, 1999.
- [102562] 17880.
- [102563] Wilgenbus, K. K.; Hsieh, C.-L.; Lankes, W. T.; Milatovich, A.; Francke, U.; Furthmayr, H.: Structure and localization on the X chromosome of the gene coding for the human filopodial protein moesin (MSN). *Genomics* 19: 326-333, 1994.
- [102567] 17881.
- [102568] Deng, L.; Wang, C.; Spencer, E.; Yang, L.; Braun, A.; You, J.; Slaughter, C.; Pickart, C.; Chen, Z. J.: Activation of the I-kappa-B complex by TRAF6 requires a dimeric ubiquitin-conjugating enzyme complex

[102571]and a unique polyubiquitin chain. Cell 103: 351–361,
2000.

[102572]17882.

[102573]Nakano, H.; Oshima, H.; Chung, W.; Williams–Abbott, L.;
Ware, C.

[102574]F.; Yagita, H.; Okumura, K.: TRAF5, an activator of NF–
kappaB and

[102575]putative signal transducer for the lymphotoxin–beta re–
ceptor. J.

[102576]Biol. Chem. 271: 14661–14664, 1996.

[102577]17883.

[102578]Nakano, H.; Sakon, S.; Koseki, H.; Takemori, T.; Tada, K.;
Matsumoto,

[102579]M.; Munechika, E.; Sakai, T.; Shirasawa, T.; Akiba, H; Ko–
bata, T.;

[102580]Santee, S. M.; Ware, C. F.; Renner, P. D.; Taniguchi, M.;
Yagita,

[102581]H.; Okumura, K.: Targeted disruption of Traf5 gene causes
defects

[102582]in CD40– and CD27–mediated lymphocyte activation.
Proc. Nat. Acad.

[102583]Sci. 96: 9803–9808, 1999.

[102584]17884.

[102585]Nakano, H.; Shindo, M.; Yamada, K.; Yoshida, M. C.; Santee, S.

[102586]M.; Ware, C. F.; Jenkins, N. A.; Gilbert, D. J.; Yagita, H.; Copeland,

[102587]N. G.; Okumura, K.: Human TNF receptor-associated factor 5 (TRAF5):

[102588]cDNA cloning, expression and assignment of the TRAF5 gene to chromosome

[102589]1q32. Genomics 42: 26–32, 1997.

[102590]17885.

[102591]Augarten, A.; Kerem, B.-S.; Yahav, Y.; Noiman, S.; Rivlin, Y.;

[102592]Tal, A.; Blau, H.; Ben-Tur, L.; Szeinberg, A.; Kerem, E.; Gazit, E.

[102593]: Mild cystic fibrosis and normal or borderline sweat test in patients

[102594]with the 3849 + 10 kb C-to-T mutation. Lancet 342: 25–26, 1993.

[102595]17886.

[102596]Bal, J.; Stuhmann, M.; Schloesser, M.; Schmidtke, J.; Reiss, J.

- [102597]: A cystic fibrosis patient homozygous for the nonsense mutation R553X. J.
- [102598]Med. Genet. 28: 715–717, 1991.
- [102599]17887.
- [102600]Ballabio, A.; Gibbs, R. A.; Caskey, C. T.: PCR test for cystic
- [102601]fibrosis deletion. (Letter) Nature 343: 220 only, 1990.
- [102602]17888.
- [102603]Baylin, S. B.; Rosenstein, B. J.; Marton, L. J.; Lockwood, D.
- [102604]H.: Age-related abnormalities of circulating polyamines and diamine
- [102605]oxidase activity in cystic fibrosis heterozygotes and homozygotes. Pediat.
- [102606]Res. 14: 921–925, 1980.
- [102607]17889.
- [102608]Bienvenu, T.; Beldjord, C.; Adjiman, M.; Kaplan, J. C.: Male
- [102609]infertility as the only presenting sign of cystic fibrosis when homozygous
- [102610]for the mild mutation R117H. (Letter) J. Med. Genet. 30: 797 only,
- [102611]1993.

[102612]17890.

[102613]Bobadilla, J. L.; Macek, M., Jr.; Fine, J. P.; Farrell, P. M.

[102614]: Cystic fibrosis: a worldwide analysis of CFTR mutations–
–correlation

[102615]with incidence data and application to screening. Hum.
Mutat. 19:

[102616]575–606, 2002.

[102617]17891.

[102618]Broackes–Carter, F. C.; Mouchel, N.; Gill, D.; Hyde, S.; Bas–
sett,

[102619]J.; Harris, A.: Temporal regulation of CFTR expression
during ovine

[102620]lung development: implications for CF gene therapy. Hum.
Molec. Genet. 11:

[102621]125–131, 2002.

[102622]17892.

[102623]Brown, C. R.; Hong–Brown, L. Q.; Biwersi, J.; Verkman, A.
S.;

[102624]Welsh, W. J.: Chemical chaperones correct the mutant
phenotype of

[102625]the deltaF508 cystic fibrosis transmembrane conductance

regulator

[102626]protein. Cell Stress Chaperones 1: 117–125, 1996.

[102627]17893.

[102628]Burger, J.; Macek, M., Jr.; Stuhmann, M.; Reis, A.;

Krawczak,

[102629]M.; Schmidtke, J.: Genetic influences in the formation of

nasal polyps.

[102630](Letter) Lancet 337: 974, 1991.

[102631]17894.

[102632]Callen, A.; Diener–West, M.; Zeitlin, P. L.; Rubenstein, R. C.

[102633]: A simplified cyclic adenosine monophosphate–mediated

sweat rate

[102634]test for quantitative measure of cystic fibrosis transmem–

brane regulator

[102635](CFTR) function. J. Pediat. 137: 849–855, 2000.

[102636]17895.

[102637]Casals, T.; Ramos, M. D.; Gimenez, J.; Larriba, S.; Nunes,

V.;

[102638]Estivill, X.: High heterogeneity for cystic fibrosis in Span–

ish families

[102639]: 75 mutations account for 90% of chromosomes. Hum.

Genet. 101:

[102640]365–370, 1997.

[102641]17896.

[102642]Casals, T.; Vazquez, C.; Lazaro, C.; Girbau, E.; Gimenez, F.
J.;

[102643]Estivill, X.: Cystic fibrosis in the Basque country: high frequency

[102644]of mutation $\Delta F508$ in patients of Basque origin. Am. J.
Hum. Genet. 50:

[102645]404–410, 1992.

[102646]17897.

[102647]Castaldo, G.; Rippa, E.; Salvatore, D.; Sibillo, R.; Raia, V.;

[102648]de Ritis, G.; Salvatore, F.: Severe liver impairment in a cystic

[102649]fibrosis-affected child homozygous for the G542X mutation. Am. J.

[102650]Med. Genet. 69: 155–158, 1997.

[102651]17898.

[102652]Chalkley, G.; Harris, A.: A cystic fibrosis patient who is homozygous

[102653]for the G85E mutation has very mild disease. J. Med.

Genet. 28:

[102654]875–877, 1991.

[102655]17899.

[102656]Chalkley, G.; Harris, A.: Lymphocyte mRNA as a resource
for detection

[102657]of mutations and polymorphism in the CF gene. J. Med.

Genet. 28:

[102658]777–780, 1991.

[102659]17900.

[102660]Chang, X.; Cui, L.; Hou, Y.; Jensen, T. J.; Aleksandrov, A.
A.;

[102661]Mengos, A.; Riordan, J. R.: Removal of multiple arginine-
framed trafficking

[102662]signals overcomes misprocessing of delta-F508 CFTR
present in most

[102663]patients with cystic fibrosis. Molec. Cell 4: 137–142,
1999.

[102664]17901.

[102665]Chanson, M.; Scerri, I.; Suter, S.: Defective regulation of
gap

[102666]junctional coupling in cystic fibrosis pancreatic duct cells.

J.

[102667]Clin. Invest. 103: 1677–1684, 1999.

[102668]17902.

[102669]Chehab, F. F.; Johnson, J.; Louie, E.; Goossens, M.;
Kawasaki,

[102670]E.; Erlich, H.: A dimorphic 4-bp repeat in the cystic fibro-
sis gene

[102671]is in absolute disequilibrium with the delta-F508 muta-
tion: implications

[102672]for prenatal diagnosis and mutation origin. Am. J. Hum.
Genet. 48:

[102673]223–226, 1991.

[102674]17903.

[102675]Cheng, S. H.; Gregory, R. J.; Marshall, J.; Paul, S.; Souza, D.

[102676]W.; White, G. A.; O'Riordan, C. R.; Smith, A. E.: Defective
intracellular

[102677]transport and processing of CFTR is the molecular basis of
most cystic

[102678]fibrosis. Cell 63: 827–834, 1990.

[102679]17904.

[102680]Chillon, M.; Casals, T.; Gimenez, J.; Nunes, V.; Estivill, X.

- [102681]: A cystic fibrosis patient homozygous for the new
frameshift mutation
- [102682]936delTA: description and clinical data. J. Med. Genet. 31:
369–370,
[102683]1994.
[102684]17905.
- [102685]Choi, J. Y.; Muallem, D.; Kiselyov, K.; Lee, M. G.; Thomas,
P.
[102686]J.; Muallem, S.: Aberrant CFTR-dependent HCO(–3) trans-
port in mutations
[102687]associated with cystic fibrosis. Nature 410: 94–97, 2001.
[102688]17906.
- [102689]Chu, C.–S.; Trapnell, B. C.; Curristin, S.; Cutting, G. R.;
Crystal,
[102690]R. G.: Genetic basis of variable exon 9 skipping in cystic
fibrosis
[102691]transmembrane conductance regulator mRNA. Nature
Genet. 3: 151–156,
[102692]1993.
[102693]17907.
- [102694]Yost, C.; Farr, G. H., III; Pierce, S. B.; Ferkey, D. M.; Chen,

[102695]M. M.; Kimelman, D.: GBP, an inhibitor of GSK-3, is implicated in

[102696]Xenopus development and oncogenesis. Cell 93:
1031-1041, 1998.

[102697]17908.

[102698]Fernandez-Valle, C.; Tang, Y.; Ricard, J.; Rodenas-Ruano,
A.; Taylor,

[102699]A.; Hackler, E.; Biggerstaff, J.; Iacovelli, J.: Paxillin binds
schwannomin

[102700]and regulates its density-dependent localization and effect on cell

[102701]morphology. Nature Genet. 31: 354-362, 2002.

[102702]17909.

[102703]Glenney, J. R., Jr.; Zokas, L.: Novel tyrosine kinase substrates

[102704]from Rous sarcoma virus-transformed cells are present in
the membrane

[102705]skeleton. J. Cell Biol. 108: 2401-2408, 1989.

[102706]17910.

[102707]Mazaki, Y.; Hashimoto, S.; Sabe, H.: Monocyte cells and
cancer

[102708]cells express novel paxillin isoforms with different binding properties

[102709]to focal adhesion proteins. J. Biol. Chem. 272: 7437–7444, 1997.

[102710]17911.

[102711]Salgia, R.; Li, J.-L.; Lo, S. H.; Brunkhorst, B.; Kansas, G. S.;

[102712]Sobhany, E. S.; Sun, Y.; Pisick, E.; Hallek, M.; Ernst, T.; Tantravahi,

[102713]R.; Chen, L. B.; Griffin, J. D.: Molecular cloning of human paxillin,

[102714]a focal adhesion protein phosphorylated by P210(BCR/ABL). J. Biol.

[102715]Chem. 270: 5039–5047, 1995.

[102716]17912.

[102717]Turner, C. E.; Glenney, J. R., Jr.; Burridge, K.: Paxillin: a

[102718]new vinculin-binding protein present in focal adhesions. J. Cell

[102719]Biol. 111: 1059–1068, 1990.

[102720]17913.

[102721]Nguyen, V. T.; Kiss, T.; Michels, A. A.; Bensaude, O.: 7SK small

[102722]nuclear RNA binds to and inhibits the activity of
CDK9/cyclin T complexes. Nature 414:

[102723]322–325, 2001.

[102724]17914.

[102725]Peng, J.; Zhu, Y.; Milton, J. T.; Price, D. H.: Identification

[102726]of multiple cyclin subunits of human P-TEFb. Genes Dev.
12: 755–762,

[102727]1998.

[102728]17915.

[102729]Yang, Z.; Zhu, Q.; Luo, K.; Zhou, Q.: The 7SK small nuclear
RNA

[102730]inhibits the CDK9/cyclin T1 kinase to control transcrip-
tion. Nature 414:

[102731]317–322, 2001.

[102732]17916.

[102733]Adachi, H.; Tsujimoto, M.; Hattori, M.; Arai, H.; Inoue, K.:
Differential

[102734]tissue distribution of the beta- and gamma-subunits of
human cytosolic

[102735]platelet-activating factor acetylhydrolase (isoform 1).
Biochem.

[102736]Biophys. Res. Commun. 233: 10–13, 1997.

[102737]17917.

[102738]Moro, F.; Arrigo, G.; Fogli, A.; Bernard, L.; Carrozzo, R.:

The

[102739]beta and gamma subunits of the human platelet-activating factor acetyl

[102740]hydrolase isoform Ib (PAFAH1B2 and PAFAH1B3) map to chromosome 11q23

[102741]and 19q13.1, respectively. Genomics 51: 157–159, 1998.

[102742]17918.

[102743]Daigo, Y.; Isomura, M.; Nishiwaki, T.; Tamari, M.;

Ishikawa, S.;

[102744]Kai, M.; Murata, Y.; Takeuchi, K.; Yamane, Y.; Hayashi, R.; Minami,

[102745]M.; Fujino, M. A.; Hojo, Y.; Uchiyama, I.; Takagi, T.; Nakamura, Y.

[102746]: Characterization of a 1200-kb genomic segment of chromosome 3p22–p21.3. DNA

[102747]Res. 6: 37–44, 1999.

[102748]17919.

[102749]Erlich, R.; Gleeson, P. A.; Campbell, P.; Dietzsch, E.; Toh,

B.-H.

[102750] Molecular characterization of trans-Golgi p230: a human peripheral

[102751] membrane protein encoded by a gene on chromosome 6p12-22 contains

[102752] extensive coiled-coil alpha-helical domains and a granin motif. J.

[102753] Biol. Chem. 271: 8328-8337, 1996.

[102754] 17920.

[102755] Barr, F. A.; Nakamura, N.; Warren, G.: Mapping the interaction

[102756] between GRASP65 and GM130, components of a protein complex involved

[102757] in the stacking of Golgi cisternae. EMBO J. 17: 3258-3268, 1998.

[102758] 17921.

[102759] Boguski, M. S.; Lowe, T. M. J.; Tolstochev, C. M.: dbEST:database

[102760] for 'expressed sequence tags.' (Letter) Nature Genet. 4: 332-333,

[102761] 1993.

[102762] 17922.

[102763]Izón, D. J.; Aster, J. C.; He, Y.; Weng, A.; Karnell, F. G.; Patriub,

[102764]V.; Xu, L.; Bakkour, S.; Rodriguez, C.; Allman, D.; Pear, W. S.:

[102765]Deltex1 redirects lymphoid progenitors to the B cell lineage by antagonizing

[102766]Notch1. *Immunity* 16: 231–243, 2002.

[102767]17923.

[102768]Lee, L.; Dowhanick–Morrissette, J.; Katz, A.; Jukofsky, L.; Krantz,

[102769]J. D.: Chromosomal localization, genomic characterization, and mapping

[102770]to the Noonan syndrome critical region of the human Deltex (DTX1)

[102771]gene. *Hum. Genet.* 107: 577–581, 2000.

[102772]17924.

[102773]Matsuno, K.; Eastman, D.; Mitsiades, T.; Quinn, A. M.; Cancian,

[102774]M. L.; Ordentlich, P.; Kadesch, T.; Artavanis–Tsakonas, S.: Human

[102775]deltex is a conserved regulator of Notch signalling. *Nature Genet.* 19:

[102776]74–78, 1998.

[102777]17925.

[102778]Chen, J. J.; Reid, C. E.; Band, V.; Androphy, E. J.: Interaction

[102779]of papillomavirus E6 oncoproteins with a putative calcium-binding

[102780]protein. *Science* 269: 529–531, 1995.

[102781]17926.

[102782]Imai, T.; Matsuda, K.; Shimojima, T.; Hashimoto, T.; Masuhiro,

[102783]Y.; Kitamoto, T.; Sugita, A.; Suzuki, K.; Matsumoto, H.; Masushige,

[102784]S.; Nogi, Y.; Muramatsu, M.; Handa, H.; Kato, S.: ERC-55, a binding

[102785]protein for the papilloma virus E6 oncoprotein, specifically interacts

[102786]with vitamin D receptor among nuclear receptors.

Biochem. Biophys.

[102787]*Res. Commun.* 233: 765–769, 1997.

[102788]17927.

[102789]Wang, J. Y.; Zhen, D. K.; Bianchi, D. W.; Androphy, E. J.; Chen,

[102790]J. J.: Assignment of the gene for ERC-55 (RCN2) to human chromosome

[102791]band 15q22.33-q24.1 by in situ hybridization. Cytogenet. Cell Genet. 79:

[102792]60-61, 1997.

[102793]17928.

[102794]Weis, K.; Griffiths, G.; Lamond, A. I.: The endoplasmic reticulum

[102795]calcium-binding protein of 55 kDa is a novel EF-hand protein retained

[102796]in the endoplasmic reticulum by a carboxyl-terminal His-Asp-Glu-Leu

[102797]motif. J. Biol. Chem. 269: 19142-19150, 1994.

[102798]17929.

[102799]Chan, A. M.-L.; Chedid, M.; McGovern, E. S.; Popescu, N. C.; Miki,

[102800]T.; Aaronson, S. A.: Expression cDNA cloning of a serine kinase transforming

[102801]gene. Oncogene 8: 1329-1333, 1993.

[102802]17930.

[102803]Bahn, R. S.; Heufelder, A. E.; Dutton, C. M.: A point muta-

tion

[102804]of the TSH receptor in retro-ocular fibroblasts from a patient with

[102805]Graves' ophthalmopathy. (Abstract) J. Endocr. Invest. 16 (suppl.

[102806]2-6): 30 only, 1993.

[102807]17931.

[102808]Moreland, R. J.; Dresser, M. E.; Rodgers, J. S.; Roe, B. A.; Conaway,

[102809]J. W.; Conaway, R. C.; Hanas, J. S.: Identification of a transcription

[102810]factor IIIA-interacting protein. Nucleic Acids Res. 28: 1986-1993,

[102811]2000.

[102812]17932.

[102813]Biebermann, H.; Schoneberg, T.; Hess, C.; Germak, J.; Gundermann,

[102814]T.; Gruters, A.: The first activating TSH receptor mutation in transmembrane

[102815]domain 1 identified in a family with nonautoimmune hyperthyroidism. J.

[102816]Clin. Endocr. Metab. 86: 4429-4433, 2001.

[102817]17933.

[102818]de Roux, N.; Polak, M.; Couet, J.; Leger, J.; Czernichow, P.;
Milgrom,

[102819]E.; Misrahi, M.: A neomutation of the thyroid-stimulating
hormone

[102820]receptor in a severe neonatal hyperthyroidism. J. Clin. En-
docr. Metab. 81:

[102821]2023-2026, 1996.

[102822]17934.

[102823]Feliciello, A.; Porcellini, A.; Ciullo, I.; Bonavolonta, G.;
Avvedimento,

[102824]E. V.; Fenzi, G.: Expression of thyrotropin-receptor mRNA
in healthy

[102825]and Graves' disease retro-orbital tissue. Lancet 342:
337-338, 1993.

[102826]17935.

[102827]Fuhrer, D.; Holzapfel, H.-P.; Wonerow, P.; Scherbaum, W.
A.; Paschke,

[102828]R.: Somatic mutations in the thyrotropin receptor gene
and not in

[102829]the Gs-alpha protein gene in 31 toxic thyroid nodules. J.
Clin. Endocr.

[102830]Metab. 82: 3885–3891, 1997.

[102831]17936.

[102832]Fuhrer, D.; Wonerow, P.; Willgerodt, H.; Paschke, R.: Identification

[102833]of a new thyrotropin receptor germline mutation
(leu629phe) in a family

[102834]with neonatal onset of autosomal dominant nonautoimmune hyperthyroidism. J.

[102835]Clin. Endocr. Metab. 82: 4234–4238, 1997.

[102836]17937.

[102837]Gabriel, E. M.; Bergert, E. R.; Grant, C. S.; van Heerden, J.

[102838]A.; Thompson, G. B.; Morris, J. C.: Germline polymorphism
of codon

[102839]727 of human thyroid-stimulating hormone receptor is
associated with

[102840]toxic multinodular goiter. J. Clin. Endocr. Metab. 84:
3328–3335,

[102841]1999.

[102842]17938.

[102843]Graves, P.; Pritsker, A.; Davies, T. F.: Post-translational
processing

[102844] of the natural human thyrotropin receptor: demonstration
of more than

[102845] two cleavage sites. J. Clin. Endocr. Metab. 84: 2177–2181,
1999.

[102846] 17939.

[102847] Gruters, A.; Schoneberg, T.; Biebermann, H.; Krude, H.;
Kron,

[102848] H. P.; Dralle, H.; Gudermann, T.: Severe congenital hyper-
thyroidism

[102849] caused by a germ-line neo mutation in the extracellular
portion of

[102850] the thyrotropin receptor. J. Clin. Endocr. Metab. 83:
1431–1436,

[102851] 1998.

[102852] 17940.

[102853] Heldin, N.-E.; Gustavsson, B.; Westermark, K.; Wester-
mark, B.

[102854]: A somatic point mutation in a putative ligand binding
domain of

[102855] the TSH receptor in a patient with autoimmune hyperthy-
roidism. J.

[102856] Clin. Endocr. Metab. 73: 1374–1376, 1991.

[102857]17941.

[102858]Holzapfel, H.-P.; Wonerow, P.; von Petrykowski, W.; Henschen,

[102859]M.; Scherbaum, W. A.; Paschke, R.: Sporadic congenital hyperthyroidism

[102860]due to a spontaneous germline mutation in the thyrotropin receptor

[102861]gene. J. Clin. Endocr. Metab. 82: 3879-3884, 1997.

[102862]17942.

[102863]Khoo, D. H. C.; Parma, J.; Rajasoorya, C.; Ho, S. C.; Vassart,

[102864]G.: A germline mutation of the thyrotropin receptor gene associated

[102865]with thyrotoxicosis and mitral valve prolapse in a Chinese family. J.

[102866]Clin. Endocr. Metab. 84: 1459-1462, 1999.

[102867]17943.

[102868]Kopp, P.; Muirhead, S.; Jourdain, N.; Gu, W.-X.; Jameson, J. L.;

[102869]Rodd, C.: Congenital hyperthyroidism caused by a solitary toxic adenoma

[102870]harboring a novel somatic mutation
(serine281-isoleucine) in the extracellular
[102871]domain of the thyrotropin receptor. J. Clin. Invest. 100:
1634-1639,
[102872]1997.
[102873]17944.
[102874]Kopp, P.; van Sande, J.; Parma, J.; Duprez, L.; Gerber, H.;
Joss,
[102875]E.; Jameson, J. L.; Dumont, J. E.; Vassart, G.: Congenital
hyperthyroidism
[102876]caused by a mutation in the thyrotropin-receptor gene.
New Eng. J.
[102877]Med. 332: 150-154, 1995.
[102878]17945.
[102879]Libert, F.; Lefort, A.; Gerard, C.; Parmentier, M.; Perret, J.;
[102880]Ludgate, M.; Dumont, J. E.; Vassart, G.: Cloning, sequenc-
ing and
[102881]expression of the human thyrotropin (TSH) receptor: evi-
dence for binding
[102882]of autoantibodies. Biochem. Biophys. Res. Commun. 165:
1250-1255,

[102883]1989.

[102884]17946.

[102885]Libert, F.; Passage, E.; Lefort, A.; Vassart, G.; Mattei, M.-G.

[102886]: Localization of human thyrotropin receptor gene to
chromosome region

[102887]14q31 by in situ hybridization. Cytogenet. Cell Genet. 54:
82-83,

[102888]1990.

[102889]17947.

[102890]Loosfelt, H.; Pichon, C.; Jolivet, A.; Misrahi, M.; Caillou, B.;

[102891]Jamous, M.; Vannier, B.; Milgrom, E.: Two-subunit struc-
ture of the

[102892]human thyrotropin receptor. Proc. Nat. Acad. Sci. 89:
3765-3769,

[102893]1992.

[102894]17948.

[102895]Muhlberg, T.; Herrmann, K.; Joba, W.; Kirchberger, M.;
Heberling,

[102896]H.-J.; Heufelder, A. E.: Lack of association of nonautoim-
mune hyperfunctioning

- [102897] thyroid disorders and a germline polymorphism of codon 727 of the
- [102898] human thyrotropin receptor in a European Caucasian population. J.
- [102899] Clin. Endocr. Metab. 85: 2640–2643, 2000.
- [102900] 17949.
- [102901] Garcia-Barcelo, M.; Law, P. T. W.; Tsui, S. K. W.; Fung, K. P.;
- [102902] Lee, C. Y.; Waye, M. M. Y.: Mapping of the human ribosomal large
- [102903] subunit protein gene RPL29 to human chromosome 3q29–qter. Genomics 46:
- [102904] 148–151, 1997.
- [102905] 17950.
- [102906] Kirn-Safran, C. B.; Dayal, S.; Martin-DeLeon, P. A.; Carson, D.
- [102907] D.: Cloning, expression, and chromosome mapping of the murine Hip/Rpl29
- [102908] gene. Genomics 68: 210–219, 2000.
- [102909] 17951.
- [102910] Law, P. T. W.; Tsui, S. K. W.; Lam, W. Y.; Luk, S. C. W.;

Hwaang,

[102911]D. M.; Liew, C. C.; Lee, C. Y.; Fung, K. P.; Waye, M. M. Y.: A novel

[102912]cDNA encoding a human homologue of ribosomal protein L29. *Biochim.*

[102913]*Biophys. Acta* 1305: 105–108, 1996.

[102914]17952.

[102915]Liu, S.; Smith, S. E.; Julian, J.; Rohde, L. H.; Karin, N. J.;

[102916]Carson, D. D.: cDNA cloning and expression of HIP, a novel cell surface

[102917]heparan sulfate/heparin-binding protein of human uterine epithelial

[102918]cells and cell lines. *J. Biol. Chem.* 271: 11817–11823, 1996.

[102919]17953.

[102920]Rohde, L. H.; Julian, J.; Babaknia, A.; Carson, D. D.: Cell surface

[102921]expression of HIP, a novel heparin/heparan sulfate-binding protein,

[102922]of human uterine epithelial cells and cell lines. *J. Biol. Chem.* 271:

[102923]11824–11830, 1996.

[102924]17954.

[102925]Giachino, C.; Lantelme, E.; Lanzetti, L.; Saccone, S.; Della Valle,

[102926]G.; Migone, N.: A novel SH3-containing human gene family preferentially

[102927]expressed in the central nervous system. Genomics 41: 427-434, 1997.

[102928]17955.

[102929]So, C. W.; Caldas, C.; Liu, M.-M.; Chen, S.-J.; Huang, Q.-H.; Gu,

[102930]L.-J.; Sham, M. H.; Wiedemann, L. M.; Chan, L. C.: EEN encodes for

[102931]a member of a new family of proteins containing an Src homology 3

[102932]domain and is the third gene located on chromosome 19p13 that fuses

[102933]to MLL in human leukemia. Proc. Nat. Acad. Sci. 94: 2563-2568, 1997.

[102934]17956.

[102935]Carling, T.; Kindmark, A.; Hellman, P.; Holmberg, L.; Akerstrom,

[102936]G.; Rastad, J.: Vitamin D receptor alleles b, a, and T: risk factors

[102937]for sporadic primary hyperparathyroidism (HPT) but not HPT of uremia

[102938]or MEN1. Biochem. Biophys. Res. Commun. 231: 329-332, 1997.

[102939]17957.

[102940]Carling, T.; Rastad, J.; Akerstrom, G.; Westin, G.: Vitamin D

[102941]receptor (VDR) and parathyroid hormone messenger ribonucleic acid

[102942]levels correspond to polymorphic VDR alleles in human parathyroid

[102943]tumors. J. Clin. Endocr. Metab. 83: 2255-2259, 1998.

[102944]17958.

[102945]Carling, T.; Ridefelt, P.; Hellman, P.; Rastad, J.; Akerstrom,

[102946]G.: Vitamin D receptor polymorphisms correlate to parathyroid cell

[102947]function in primary hyperparathyroidism. J. Clin. Endocr. Metab. 82:

[102948]1772–1775, 1997.

[102949]17959.

[102950]Correa, P.; Rastad, J.; Schwarz, P.; Westin, G.; Kindmark, A.;

[102951]Lundgren, E.; Akerstrom, G.; Carling, T.: The vitamin D receptor

[102952](VDR) start codon polymorphism in primary hyperparathyroidism and

[102953]parathyroid VDR messenger ribonucleic acid levels. J. Clin. Endocr.

[102954]Metab. 84: 1690–1694, 1999.

[102955]17960.

[102956]Ensrud, K. E.; Stone, K.; Cauley, J. A.; White, C.; Zmuda, J. M.;

[102957]Nguyen, T. V.; Eisman, J. A.; Cummings, S. R.: Vitamin D receptor

[102958]gene polymorphisms and the risk of fractures in older women. J. Bone

[102959]Miner. Res. 14: 1637–1645, 1999.

[102960]17961.

[102961]Ferrari, S.; Manen, D.; Bonjour, J.-P.; Slosman, D.; Rizzoli,

R.

[102962]; Bone mineral mass and calcium and phosphate metabolism in young

[102963]men: relationships with vitamin D receptor allelic polymorphisms. J.

[102964]Clin. Endocr. Metab. 84: 2043–2048, 1999.

[102965]17962.

[102966]Gennari, L.; Becherini, L.; Masi, L.; Mansani, R.; Gonnelli, S.;

[102967]Cepollaro, C.; Martini, S.; Montagnani, A.; Lentini, G.; Becorpi,

[102968]A. M.; Brandi, M. L.: Vitamin D and estrogen receptor allelic variants

[102969]in Italian postmenopausal women: evidence of multiple gene contribution

[102970]to bone mineral density. J. Clin. Endocr. Metab. 83: 939–944, 1998.

[102971]17963.

[102972]Kristjansson, K.; Rut, A. R.; Hewison, M.; O'Riordan, J. L. H.;

[102973]Hughes, M. R.: Two mutations in the hormone binding domain of the

[102974] vitamin D receptor cause tissue resistance to 1,25 dihydroxyvitamin

[102975] D(3). J. Clin. Invest. 92: 12-16, 1993.

[102976] 17964.

[102977] Labuda, M.; Ross, M. V.; Fujiwara, T. M.; Morgan, K.; Ledbetter,

[102978] D.; Hughes, M. R.; Glorieux, F. H.: Two hereditary defects related

[102979] to vitamin D metabolism map to the same region of human chromosome

[102980] 12q.11(Abtract) Cytogenet. Cell Genet. 58: 1978, 1991.

[102981] 17965.

[102982] Jurutka, P. W.; Remus, L. S.; Whitfield, G. K.; Thompson, P. D.;

[102983] Hsieh, J.-C.; Zitzer, H.; Tavakkoli, P.; Galligan, M. A.; Dang, H.

[102984] T. L.; Haussler, C. A.; Haussler, M. R.: The polymorphic N terminus

[102985] in human vitamin D receptor isoforms influences transcriptional activity

[102986] by modulating interaction with transcription factor IIB. Molec. Endocr. 14:

[102987]401–420, 2000.

[102988]17966.

[102989]Lin, U.-T.; Malloy, P. J.; Sakati, N.; Al-Ashwal, A.; Feldman,

[102990]D.: A novel mutation in the deoxyribonucleic acid-binding
domain

[102991]of the vitamin D receptor causes hereditary
1,25-dihydroxyvitamin

[102992]D-resistant rickets. J. Clin. Endocr. Metab. 81:
2564–2569, 1996.

[102993]17967.

[102994]Lorentzon, M.; Lorentzon, R.; Nordstrom, P.: Vitamin D re-
ceptor

[102995]gene polymorphism is associated with birth height,
growth to adolescence,

[102996]and adult stature in healthy Caucasian men: a cross-
sectional and

[102997]longitudinal study. J. Clin. Endocr. Metab. 85: 1666–1671,
2000.

[102998]17968.

[102999]Makishima, M.; Lu, T. T.; Xie, W.; Whitfield, G. K.; Domoto,
H.;

[103000]Evans, R. M.; Haussler, M. R.; Mangelsdorf, D. J.: Vitamin D receptor

[103001]as an intestinal bile acid sensor. Science 296: 1313–1316, 2002.

[103002]17969.

[103003]Mechica, J. B.; Leite, M. O. R.; Mendonca, B. B.; Frazzatto, E.

[103004]S. T.; Borelli, A.; Latronico, A. C.: A novel nonsense mutation in

[103005]the first zinc finger of the vitamin D receptor causing hereditary

[103006]1,25-dihydroxyvitamin D(3)-resistant rickets. J. Clin. Endocr. Metab. 82:

[103007]3892–3894, 1997.

[103008]17970.

[103009]Furlan, M.; Robles, R.; Galbusera, M.; Remuzzi, G.; Kyrle, P. A.;

[103010]Brenner, B.; Krause, M.; Scharrer, I.; Aumann, V.; Mittler, U.; Solenthaler,

[103011]M.; Lammle, B.: Von Willebrand factor-cleaving protease in thrombotic

[103012]thrombocytopenic purpura and the hemolytic-uremic syn-

drome. New Eng.

[103013]J. Med. 339: 1578–1584, 1998.

[103014]17971.

[103015]Furlan, M.; Robles, R.; Lammle, B.: Partial purification and characterization

[103016]of a protease from human plasma cleaving von Willebrand factor to

[103017]fragments produced by in vivo proteolysis. Blood 87: 4223–4234,

[103018]1996.

[103019]17972.

[103020]Nichols, A. F.; Ong, P.; Linn, S.: Mutations specific to the xeroderma

[103021]pigmentosum group E Ddb– phenotype. J. Biol. Chem. 271: 24317–24320,

[103022]1996.

[103023]17973.

[103024]Itoh, T.; Linn, S.; Kamide, R.; Tokushige, H.; Katori, N.; Hosaka,

[103025]Y.; Yamaizumi, M.: Xeroderma pigmentosum variant heterozygotes show

[103026]reduced levels of recovery of replicative DNA synthesis in
the presence
[103027]of caffeine after ultraviolet irradiation. J. Invest. Derm.
115:
[103028]981–985, 2000.
[103029]17974.
[103030]Shisheva, A.; Sudhof, T. C.; Czech, M. P.: Cloning, charac-
terization,
[103031]and expression of a novel GDP dissociation inhibitor iso-
form from
[103032]skeletal muscle. Molec. Cell. Biol. 14: 3459–3468, 1994.
[103033]17975.
[103034]Fathi, Z.; Corjay, M. H.; Shapira, H.; Wada, E.; Benya, R.;
Jensen,
[103035]R.; Viallet, J.; Sausville, E. A.; Battey, J. F.: BRS-3: a novel
[103036]bombesin receptor subtype selectively expressed in testis
and lung
[103037]carcinoma cells. J. Biol. Chem. 268: 5979–5984, 1993.
[103038]17976.
[103039]Gorbulev, V.; Akhundova, A.; Grzeschik, K. H.; Fahrenholz,
F.:

- [103040] Organization and chromosomal localization of the gene
for the human
- [103041] bombesin receptor subtype expressed in pregnant uterus.
FEBS Lett. 340:
- [103042] 260–264, 1994.
- [103043] 17977.
- [103044] Ohki–Hamazaki, H.; Watase, K.; Yamamoto, K.; Ogura, H.;
Yamano,
- [103045] M.; Yamada, K.; Maeno, H.; Imaki, J.; Kikuyama, S.; Wada,
E.; Wada,
- [103046] K.: Mice lacking bombesin receptor subtype–3 develop
metabolic defects
- [103047] and obesity. Nature 390: 166–169, 1997.
- [103048] 17978.
- [103049] Banfi, S.; Borsani, G.; Bulfone, A.; Ballabio, A.: Drosophila–re-
lated
- [103050] expressed sequences. Hum. Molec. Genet. 6: 1745–1753,
1997.
- [103051] 17979.
- [103052] Castrillon, D. H.; Wasserman, S. A.: Diaphanous is re-
quired for

[103053]cytokinesis in *Drosophila* and shares domains of similarity with the

[103054]products of the limb deformity gene. *Development* 120: 3367–3377,

[103055]1994.

[103056]17980.

[103057]Philippe, C.; Cremers, F. P. M.; Chery, M.; Bach, I.; Abbadi, N.;

[103058]Ropers, H. H.; Gilgenkrantz, S.: Physical mapping of DNA markers

[103059]in the q13–q22 region of the human X chromosome. *Genomics* 17: 147–152,

[103060]1993.

[103061]17981.

[103062]Bione, S.; Sala, C.; Manzini, C.; Arrigo, G.; Zuffardi, O.; Banfi,

[103063]S.; Borsani, G.; Jonveaux, P.; Philippe, C.; Zuccotti, M.; Bal-labio,

[103064]A.; Toniolo, D.: A human homologue of the *Drosophila melanogaster*

[103065]diaphanous gene is disrupted in a patient with premature ovarian failure:

[103066]evidence for conserved function in oogenesis and implications for

[103067]human sterility. *Am. J. Hum. Genet.* 62: 533–541, 1998.

[103068]17982.

[103069]Sala, C.; Arrigo, G.; Torri, G.; Martinazzi, F.; Riva, P.; Larizza,

[103070]L.; Philippe, C.; Jonveaux, P.; Sloan, F.; Labella, T.; Toniolo, D.

[103071]: Eleven X chromosome breakpoints associated with premature ovarian

[103072]failure (POF) map to a 15-Mb YAC contig spanning Xq21. *Genomics* 40:

[103073]123–131, 1997.

[103074]17983.

[103075]Jagla, W. M.; Jagle, H.; Hayashi, T.; Sharpe, L. T.; Deeb, S.

[103076]S.: The molecular basis of dichromatic color vision in males with

[103077]multiple red and green visual pigment genes. *Hum. Molec. Genet.* 11:

[103078]23–32, 2002.

[103079]17984.

- [103080]Huijing, F.; Fernandez, J.: X-chromosomal inheritance of
liver
[103081]glycogenosis with phosphorylase kinase deficiency. Am. J.
Hum. Genet. 21:
[103082]275–284, 1969.
[103083]17985.
- [103084]Franco, B.; Guioli, S.; Pragliola, A.; Incerti, B.; Bardoni, B.;
[103085]Tonlorenzi, R.; Carrozzo, R.; Maestrini, E.; Pieretti, M.;
Taillon–Miller,
[103086]P.; Brown, C. J.; Willard, H. F.; Lawrence, C.; Persico, M. G.;
Camerino,
[103087]G.; Ballabio, A.: A gene deleted in Kallmann's syndrome
shares homology
[103088]with neural cell adhesion and axonal path–finding
molecules. Nature 353:
[103089]529–536, 1991.
[103090]17986.
- [103091]Lachner, M.; O'Carroll, D.; Rea, S.; Mechtler, K.; Jenuwein,
T.
[103092]Methylation of histone H3 lysine 9 creates a binding site
for HP1

[103093]proteins. Nature 410: 116–120, 2001.

[103094]17987.

[103095]Melcher, M.; Schmid, M.; Aagaard, L.; Selenko, P.; Laible, G.;

[103096]Jenuwein, T.: Structure–function analysis of SUV39H1 reveals a dominant

[103097]role in heterochromatin organization, chromosome segregation, and

[103098]mitotic progression. Molec. Cell Biol. 20: 3728–3841, 2000.

[103099]17988.

[103100]Nielsen, S. J.; Schneider, R.; Bauer, U.–M.; Bannister, A. J.;

[103101]Morrison, A.; O'Carroll, D.; Firestein, R.; Cleary, M.; Jenuwein,

[103102]T.; Herrera, R. E.; Kouzarides, T.: Rb targets histone H3 methylation

[103103]and HP1 to promoters. Nature 412: 561–565, 2001.

[103104]17989.

[103105]Peters, A. H. F. M.; O'Carroll, D.; Scherthan, H.; Mechtler, K.;

[103106]Sauer, S.; Schofer, C.; Weipoltshammer, K.; Pagani, M.;

Lachner, M.;

[103107]Kohlmaier, A.; Opravil, S.; Doyle, M.; Sibilia, M.; Jenuwein, T.:

[103108]Loss of the Suv39h histone methyltransferases impairs mammalian heterochromatin

[103109]and genome stability. Cell 107: 323–337, 2001.

[103110]17990.

[103111]Rea, S.; Eisenhaber, F.; O'Carroll, D.; Strahl, B. D.; Sun, Z.-W.;

[103112]Schmid, M.; Opravil, S.; Mechtler, K.; Ponting, C. P.; Allis, C. D.;

[103113]Jenuwein, T.: Regulation of chromatin structure by site-specific

[103114]histone H3 methyltransferases. Nature 406: 593–599, 2000.

[103115]17991.

[103116]Scott, A. F.: Personal Communication. Baltimore, Md. 8/7/2000.

[103117]17992.

[103118]Haltiwanger, R. S.; Blomberg, M. A.; Hart, G. W.: Glycosylation

[103119]of nuclear and cytoplasmic proteins: purification and

characterization

[103120] of a uridine diphospho-N-acetylglucosamine:polypeptide
beta-N-acetylglucosaminyltransferase. J.

[103121] Biol. Chem. 267: 9005-9013, 1992.

[103122] 17993.

[103123] Kreppel, L. K.; Blomberg, M. A.; Hart, G. W.: Dynamic gly-
cosylation

[103124] of nuclear and cytosolic proteins: cloning and characteri-
zation of

[103125] a unique O-GlcNAc transferase with multiple tetratri-
copeptide repeats. J.

[103126] Biol. Chem. 272: 9308-9315, 1997.

[103127] 17994.

[103128] Lubas, W. A.; Frank, D. W.; Krause, M.; Hanover, J. A.: O-
linked

[103129] GlcNAc transferase is a conserved nucleocytoplasmic pro-
tein containing

[103130] tetratricopeptide repeats. J. Biol. Chem. 272: 9316-9324,
1997.

[103131] 17995.

[103132] Shafi, R.; Iyer, S. P. N.; Ellies, L. G.; O'Donnell, N.; Marek,

- [103133]K. W.; Chui, D.; Hart, G. W.; Marth, J. D.: The O-GlcNAc transferase
- [103134]gene resides on the X chromosome and is essential for embryonic stem
- [103135]cell viability and mouse ontogeny. Proc. Nat. Acad. Sci. 97: 5735–5739,
- [103136]2000.
- [103137]17996.
- [103138]Yang, X.; Zhang, F.; Kudlow, J. E.: Recruitment of O-GlcNAc transferase
- [103139]to promoters by corepressor mSin3A: coupling protein O-GlcNAcylation
- [103140]to transcriptional repression. Cell 110: 69–80, 2002.
- [103141]17997.
- [103142]Danon, M. J.; Oh, S. J.; DiMauro, S.; Manaligod, J. R.; Eastwood,
- [103143]A.; Naidu, S.; Schliselfeld, L. H.: Lysosomal glycogen storage disease
- [103144]with normal acid maltase. Neurology 31: 51–57, 1981.
- [103145]17998.
- [103146]Dworzak, F.; Casazza, F.; Mora, C. M.; De Maria, R.;

Gronda, E.;

[103147]Baroldi, G.; Rimoldi, M.; Morandi, L.; Cornelio, F.: Lysosomal glycogen

[103148]storage with normal acid maltase: a familial study with successful

[103149]heart transplant. Neuromusc. Disord. 4: 243–247, 1994.

[103150]17999.

[103151]Liu, Y.; Hoffmann, A.; Grinberg, A.; Westphal, H.; McDonald, M.

[103152]P.; Miller, K. M.; Crawley, J. N.; Sandhoff, K.; Suzuki, K.; Proia,

[103153]R. L.: Mouse model of GM2 activator deficiency manifests cerebellar

[103154]pathology and motor impairment. Proc. Nat. Acad. Sci. 94: 8138–8143,

[103155]1997.

[103156]18000.

[103157]Mahuran, D. J.: Personal Communication. Toronto, Ontario, Canada

[103158]12/1/1994.

[103159]18001.

[103160]O'Neill, B.; Butler, A. B.; Young, E.; Falk, P. M.; Bass, N. H.

[103161]: Adult-onset Gm2-gangliosidosis: seizures, dementia,
and normal pressure

[103162]hydrocephalus associated with glycolipid storage in the
brain and

[103163]arachnoid granulation. Neurology 28: 1117-1123, 1978.

[103164]18002.

[103165]Phillips, J. A., III: Personal Communication. Baltimore, Md.

[103166]8/11/1983.

[103167]18003.

[103168]Sakuraba, H.; Itoh, K.; Shimmoto, M.; Utsumi, K.; Kase, R.;
Hashimoto,

[103169]Y.; Ozawa, T.; Ohwada, Y.; Imataka, G.; Eguchi, M.; Fu-
rukawa, T.;

[103170]Schepers, U.; Sandhoff, K.: GM2 gangliosidosis AB variant:
clinical

[103171]and biochemical studies of a Japanese patient. Neurology
52: 372-377,

[103172]1999.

[103173]18004.

[103174]Schepers, U.; Glombitza, G.; Lemm, T.; Hoffmann, A.;
Chabas, A.;

[103175]Ozand, P.; Sandhoff, K.: Molecular analysis of a
GM2-activator deficiency

[103176]in two patients with GM2-gangliosidosis AB variant. Am. J.
Hum. Genet. 59:

[103177]1048-1056, 1996.

[103178]18005.

[103179]Schroder, M.; Schnabel, D.; Hurwitz, R.; Young, E.; Suzuki,
K.;

[103180]Sandhoff, K.: Molecular genetics of GM2-gangliosidosis
AB variant:

[103181]a novel mutation and expression in BHK cells. Hum.
Genet. 92: 437-440,

[103182]1993.

[103183]18006.

[103184]Schroder, M.; Schnabel, D.; Suzuki, K.; Sandhoff, K.: A mu-
tation

[103185]in the gene of a glycolipid-binding protein (GM2 activator)
that causes

[103186]GM2-gangliosidosis variant AB. FEBS Lett. 290: 1-3, 1991.

[103187]18007.

[103188]Swallow, D. M.; Islam, I.; Fox, M. F.; Povey, S.; Klima, H.;
Schepers,

[103189]U.; Sandhoff, K.: Regional localization of the gene coding
for the

[103190]GM2 activator protein (GM2A) to chromosome 5q32-33
and confirmation

[103191]of the assignment of GM2AP to chromosome 3. Ann. Hum.
Genet. 57:

[103192]187-193, 1993.

[103193]18008.

[103194]Valentine, C. R.; Heflich, R. H.: The association of non-
sense

[103195]mutations with exon-skipping in hprt mRNA of Chinese
hamster ovary

[103196]cells results from an artifact of RT-PCR. RNA 3: 660-676,
1997.

[103197]18009.

[103198]Xie, B.; Kennedy, J. L.; McInnes, B.; Auger, D.; Mahuran, D.:

[103199]Identification of a processed pseudogene related to the
functional

[103200] gene encoding the G-M2 activator protein: localization of the pseudogene

[103201] to human chromosome 3 and the functional gene to human chromosome

[103202] 5. Genomics 14: 796-798, 1992.

[103203] 18010.

[103204] Xie, B.; Wang, W.; Mahuran, D. J.: A cys138-to-arg substitution

[103205] in the G-M2 activator protein is associated with the AB variant form

[103206] of G-M2 gangliosidosis. Am. J. Hum. Genet. 50: 1046-1052, 1992.

[103207] 18011.

[103208] Yamanaka, S.; Johnson, O. N.; Lyu, M. S.; Kozak, C. A.; Proia,

[103209] R. L.: The mouse gene encoding the G(M2) activator protein (Gm2a):

[103210] cDNA sequence, expression, and chromosome mapping. Genomics 24:

[103211] 601-604, 1994.

[103212] 18012.

[103213]Ainsworth, P. J.; Coulter-Mackie, M. B.: A double mutation
in
[103214]exon 6 of the beta-hexosaminidase alpha subunit in a pa-
tient with
[103215]the B1 variant of Tay-Sachs disease. Am. J. Hum. Genet.
51: 802-809,
[103216]1992.
[103217]18013.
[103218]Akalin, N.; Shi, H.-P.; Vavougios, G.; Hechtman, P.; Lo, W.;
Scriber,
[103219]C. R.; Mahuran, D.; Kaplan, F.: Novel Tay-Sachs disease
mutations
[103220]from China. Hum. Mutat. 1: 40-46, 1992.
[103221]18014.
[103222]Akerman, B. R.; Zielenski, J.; Triggs-Raine, B. L.; Prencz, E.
[103223]M.; Natowicz, M. R.; Lim-Steele, J. S. T.; Kaback, M. M.;
Miles, E.
[103224]H.; Thomas, G. H.; Clarke, J. T. R.; Gravel, R. A.: A muta-
tion common
[103225]in non-Jewish Tay-Sachs disease: frequency and RNA
studies. Hum.

[103226]Mutat. 1: 303–309, 1992.

[103227]18015.

[103228]Akli, S.; Chelly, J.; Lacorte, J.; Poenaru, L.; Kahn, A.: Seven

[103229]novel Tay–Sachs mutations detected by chemical mismatch cleavage of

[103230]PCR–amplified cDNA fragments. Genomics 11: 124–134, 1991.

[103231]18016.

[103232]Akli, S.; Chomel, J.–C.; Lacorte, J.–M.; Bachner, L.; Poenaru,

[103233]A.; Poenaru, L.: Ten novel mutations in the HEXA gene in non–Jewish

[103234]Tay–Sachs patients. Hum. Molec. Genet. 2: 61–67, 1993.

[103235]18017.

[103236]Lyons, L. B.; Cox, R. P.; Dancis, J.: Complementation analysis

[103237]of maple syrup urine disease in heterokaryons derived from cultured

[103238]human fibroblasts. Nature 243: 533–535, 1973.

[103239]18018.

[103240]Singh, S.; Willers, I.; Goedde, H. W.: Heterogeneity in
maple

[103241]syrup urine disease: aspects of cofactor requirement and
complementation

[103242]in cultured fibroblasts. Clin. Genet. 11: 277–284, 1977.

[103243]18019.

[103244]Ohno, K.; Brengman, J.; Tsujino, A.; Engel, A. G.: Human
endplate

[103245]acetylcholinesterase deficiency caused by mutations in the
collagen-like

[103246]tail subunit (ColQ) of the asymmetric enzyme. Proc. Nat.
Acad. Sci. 95:

[103247]9654–9659, 1998.

[103248]18020.

[103249]Schwartz, J. M.; Reiss, A. L.; Jaffe, E. R.: Hereditary methemoglobinemia

[103250]with deficiency of NADH cytochrome b5 reductase. In:
Stanbury, J. B.;

[103251]Wyngaarden, J. B.; Fredrickson, D. S.; Goldstein, J. L.;
Brown, M.

[103252]S.: The Metabolic Basis of Inherited Disease. New York:
McGraw-Hill

[103253](pub.) (5th ed.): 1983. Pp. 1654–1665.

[103254]18021.

[103255]Scott, E. M.: The relationship of diaphorase of human erythrocytes

[103256]to inheritance of methemoglobinemia. J. Clin. Invest. 39: 1176–1179,

[103257]1960.

[103258]18022.

[103259]Scott, E. M.; Lewis, M.; Kaita, H.; Chown, B.; Giblett, E. R.

[103260]: The absence of close linkage of methemoglobinemia and blood group

[103261]loci. Am. J. Hum. Genet. 15: 493–494, 1963.

[103262]18023.

[103263]Scott, E. M.; Wright, R. C.: The absence of close linkage of

[103264]methemoglobinemia and other loci. Am. J. Hum. Genet. 21: 194–195,

[103265]1969.

[103266]18024.

[103267]Shirabe, K.: Personal Communication. Oita, Japan

7/21/1997.

[103268]18025.

[103269]Shirabe, K.; Fujimoto, Y.; Yubisui, T.; Takeshita, M.: An in-frame

[103270]deletion of codon 298 of the NADH-cytochrome b(5) reductase gene results

[103271]in hereditary methemoglobinemia type II (generalized type): a functional

[103272]implication for the role of the COOH-terminal region of the enzyme. J.

[103273]Biol. Chem. 269: 5952-5957, 1994.

[103274]18026.

[103275]Shirabe, K.; Landi, M. T.; Takeshita, M.; Uziel, G.; Fedrizzi,

[103276]E.; Borgese, N.: A novel point mutation in a 3-prime splice site

[103277]of the NADH-cytochrome b5 reductase gene results in immunologically

[103278]undetectable enzyme and impaired NADH-dependent ascorbate regeneration

[103279]in cultured fibroblasts of a patient with type II hereditary methemoglobinemia. Am.

[103280]J. Hum. Genet. 57: 302-310, 1995.

[103281]18027.

[103282]Shirabe, K.; Yubisui, T.; Borgese, N.; Tang, C.; Hultquist,
D.

[103283]E.; Takeshita, M.: Enzymatic instability of NADH-cy-
tochrome b5 reductase

[103284]as a cause of hereditary methemoglobinemia type I (red
cell type). J.

[103285]Biol. Chem. 267: 20416-20421, 1992.

[103286]18028.

[103287]Tanishima, K.; Matsuki, T.; Fukuda, N.; Takeshita, M.;
Yoneyama,

[103288]Y.: NADH-cytochrome b5 reductase in platelets and
leukocytes with

[103289]special reference to normal levels and to levels in carriers
of hereditary

[103290]methemoglobinemia with or without neurological symp-
toms. Acta Haemat. 63:

[103291]7-12, 1980.

[103292]18029.

[103293]Tanishima, K.; Tanimoto, K.; Tomoda, A.; Mawatari, K.;
Matsukawa,

- [103294]S.; Yoneyama, Y.; Ohkuwa, H.; Takazakura, E.: Hereditary methemoglobinemia due to cytochrome b(5) reductase deficiency in blood cells without associated neurologic and mental disorders. Blood 66: 1288–1291, 1985.
- [103297]18030.
- [103298]18030.
- [103299]Tomatsu, S.; Kobayashi, Y.; Fukumaki, Y.; Yubisui, T.; Orii, T.;
- [103300]Sakaki, Y.: The organization and the complete nucleotide sequence of the human NADH–cytochrome b5 reductase gene. Gene 80: 353–361, 1989.
- [103302]18031.
- [103303]18031.
- [103304]Townes, P. L.; Morrison, M.: Investigation of the defect in a variant of hereditary methemoglobinemia. Blood 19: 60–74, 1962.
- [103306]18032.

[103307]Trost, C.: The Blue People of Troublesome Creek. Science
82

[103308](Nov.): 35–39, 1982.

[103309]18033.

[103310]Vieira, L. M.; Kaplan, J.-C.; Kahn, A.; Leroux, A.: Four new
[103311]mutations in the NADH-cytochrome b5 reductase gene
from patients with

[103312]recessive congenital methemoglobinemia type II. Blood
85: 2254–2262,

[103313]1995.

[103314]18034.

[103315]Vives-Corróns, J. L.; Pujades, A.; Vela, E.; Corretger, J. M.;

[103316]Leroux, A.; Kaplan, J. C.: Congenital methemoglobin-reduc-
tase (cytochrome

[103317]b5 reductase) deficiency associated with mental retarda-
tion in a Spanish

[103318]girl. Acta Haemat. 59: 348–353, 1978.

[103319]18035.

[103320]Waller, H. D.: Inherited methemoglobinemia (enzyme defi-
ciencies). Humangenetik 9:

[103321]217–218, 1970.

[103322]18036.

[103323]Wang, Y.; Wu, Y.-S.; Zheng, P.-Z.; Yang, W.-X.; Fang, G.-A.; Tang,

[103324]Y.-C.; Xie, F.; Lan, F.-H.; Zhu, Z.-Y.: A novel mutation in the NADH-cytochrome

[103325]b5 reductase gene of a Chinese patient with recessive congenital methemoglobinemia. Blood 95:

[103326]3250–3255, 2000.

[103327]18037.

[103328]West, C. A.; Gomperts, B. D.; Huehns, E. R.; Kessel, I.; Ashby,

[103329]J. R.: Demonstration of an enzyme variant in a case of congenital

[103330]methaemoglobinaemia. Brit. Med. J. 4: 212–214, 1967.

[103331]18038.

[103332]Worster-Drought, C.; White, J. C.; Sargent, F.: Familial, idiopathic

[103333]methaemoglobinaemia associated with mental deficiency and neurological

[103334]abnormalities. Brit. Med. J. 2: 114–118, 1953.

[103335]18039.

[103336]Wu, Y.-S.; Huang, C.-H.; Wan, Y.; Huang, Q.-J.; Zhu, Z.-Y.:

Identification

[103337]of a novel point mutation (leu72-to-pro) in the NADH-cytochrome b5

[103338]reductase gene of a patient with hereditary methaemoglobinaemia type

[103339]J. Brit. J. Haemat. 102: 575-577, 1998.

[103340]18040.

[103341]Yubisui, T.; Miyata, T.; Iwanaga, S.; Tamura, M.; Takeshita, M.

[103342]: Complete amino acid sequence of NADH-cytochrome b5 reductase purified

[103343]from human erythrocytes. J. Biochem. 99: 407-422, 1986.

[103344]18041.

[103345]Yubisui, T.; Naitoh, Y.; Zenno, S.; Tamura, M.; Takeshita, M.;

[103346]Sakaki, Y.: Molecular cloning of cDNAs of human liver and placenta

[103347]NADH-cytochrome b(5) reductase. Proc. Nat. Acad. Sci. 84: 3609-3613,

[103348]1987.

[103349]18042.

[103350]Yubisui, T.; Shirabe, K.; Takeshita, M.; Kobayashi, Y.;
Fukumaki,

[103351]Y.; Sakaki, Y.; Takano, T.: Structural role of serine 127 in
the

[103352]NADH-binding site of human NADH-cytochrome b5 re-
ductase. J. Biol.

[103353]Chem. 266: 66-70, 1991.

[103354]18043.

[103355]Alvarez, L.; Corrales, F.; Martin-Duce, A.; Mato, J. M.:
Characterization

[103356]of a full-length cDNA encoding human liver S-
adenosylmethionine synthetase:

[103357]tissue-specific gene expression and mRNA levels in hep-
atopathies. Biochem.

[103358]J. 293: 481-486, 1993.

[103359]18044.

[103360]Blom, H. J.; Davidson, A. J.; Finkelstein, J. D.; Luder, A. S.;

[103361]Bernardini, I.; Martin, J. J.; Tangerman, A.; Trijbels, J. M. F.;

- [103362]Mudd, S. H.; Goodman, S. I.; Gahl, W. A.: Persistent hypermethioninaemia
[103363]with dominant inheritance. J. Inherit. Metab. Dis. 15:
188–197,
[103364]1992.
[103365]18045.
- [103366]Cabrero, C.; Puerta, J.; Alemany, S.: Purification and comparison
[103367]of two forms of S-adenosyl-L-methionine synthetase
from rat liver. Europ.
[103368]J. Biochem. 170: 299–304, 1987.
[103369]18046.
- [103370]Chamberlin, M. E.; Ubagai, T.; Mudd, S. H.; Levy, H. L.;
Chou,
[103371]J. Y.: Dominant inheritance of isolated hypermethionine-
mia is associated
[103372]with a mutation in the human methionine adenosyltrans-
ferase 1A gene. Am.
[103373]J. Hum. Genet. 60: 540–546, 1997.
[103374]18047.
- [103375]Chamberlin, M. E.; Ubagai, T.; Mudd, S. H.; Thomas, J.;

Pao, V.

[103376]Y.; Nguyen, T. K.; Levy, H. L.; Greene, C.; Freehauf, C.;
Chou, J.

[103377]Y.: Methionine adenosyltransferase I/III deficiency: novel
mutations

[103378]and clinical variations. Am. J. Hum. Genet. 66: 347–355,
2000.

[103379]18048.

[103380]Chamberlin, M. E.; Ubagai, T.; Mudd, S. H.; Wilson, W. G.;
Leonard,

[103381]J. V.; Chou, J. Y.: Demyelination of the brain is associated
with

[103382]methionine adenosyltransferase I/III deficiency. J. Clin. In-
vest. 98:

[103383]1021–1027, 1996.

[103384]18049.

[103385]Finkelstein, J. D.; Kyle, W. E.; Martin, J. J.: Abnormal me-
thionine

[103386]adenosyltransferase in hypermethioninemia. Biochem.
Biophys. Res.

[103387]Commun. 66: 1491–1497, 1975.

[103388]18050.

- [103389]Lim, L. E.; Duclos, F.; Broux, O.; Bourg, N.; Sunada, Y.; Al-
lamand,
[103390]V.; Meyer, J.; Richard, I.; Moomaw, C.; Slaughter, C.; Tome,
F. M.
[103391]S.; Fardeau, M.; Jackson, C. E.; Beckmann, J. S.; Campbell,
K. P.
[103392]; Beta-sarcoglycan: characterization and role in limb-
girdle muscular
[103393]dystrophy linked to 4q12. Nature Genet. 11: 257-265,
1995.
[103394]18051.
[103395]Passos-Bueno, M. R.; Vainzof, M.; Moreira, E. S.; Zatz, M.:
Seven
[103396]autosomal recessive limb-girdle muscular dystrophies in
the Brazilian
[103397]population: from LGMD2A to LGMD2G. Am. J. Med. Genet.
82: 392-398,
[103398]1999.
[103399]18052.
[103400]Jarvela, I.: Infantile neuronal ceroid lipofuscinosis (CLN1):
[103401]linkage disequilibrium in the Finnish population and evi-
dence that

[103402]variant late infantile form (variant CLN2) represents a nonallelic

[103403]locus. Genomics 10: 333–337, 1991.

[103404]18053.

[103405]Nishiyama, K.; Funai, T.; Katafuchi, R.; Hattori, F.; Onoyama,

[103406]K.; Ichiyama, A.: Primary hyperoxaluria type I due to a point mutation

[103407]of T to C in the coding region of the serine:pyruvate aminotransferase

[103408]gene. Biochem. Biophys. Res. Commun. 176: 1093–1099, 1991.

[103409]18054.

[103410]Pirulli, D.; Puzzer, D.; Ferri, L.; Crovella, S.; Amoroso, A.;

[103411]Ferrettini, C.; Marangella, M.; Mazzola, G.; Florian, F.: Molecular

[103412]analysis of hyperoxaluria type 1 in Italian patients reveals eight

[103413]new mutations in the alanine:glyoxylate aminotransferase gene. Hum.

[103414]Genet. 104: 523–525, 1999.

[103415]18055.

[103416]Watts, R. W. E.; Calne, R. Y.; Rolles, K.; Danpure, C. J.;
Morgan,

[103417]S. H.; Mansell, M. A.; Williams, R.; Purkiss, P.: Successful
treatment

[103418]of primary hyperoxaluria type I by combined hepatic and
renal transplantation. Lancet II:

[103419]474–475, 1987.

[103420]18056.

[103421]Argoff, C. E.; Kaneski, C. R.; Blanchette–Mackie, E. J.;
Comly,

[103422]M.; Dwyer, N. K.; Brown, A.; Brady, R. O.; Pentchev, P. G.:
Type

[103423]C Niemann–Pick disease: documentation of abnormal LDL
processing in

[103424]lymphocytes. Biochem. Biophys. Res. Commun. 171:
38–45, 1990.

[103425]18057.

[103426]Natt, E.; Kida, K.; Odievre, M.; Di Rocco, M.; Scherer, G.:
Point

[103427]mutations in the tyrosine aminotransferase gene in ty-
rosinemia type

[103428]II. Proc. Nat. Acad. Sci. 89: 9297–9301, 1992.

[103429]18058.

[103430]Brenner, V.; Nyakatura, G.; Rosenthal, A.; Platzer, M.: Ge-
nomic

[103431]organization of two novel genes on human Xq28: compact
head to head

[103432]arrangement of IDH-gamma and TRAP-delta is conserved
in rat and mouse. Genomics 44:

[103433]8–14, 1997.

[103434]18059.

[103435]Achen, M. G.; Jeltsch, M.; Kukk, E.; Makinen, T.; Vitali, A.;
Wilks,

[103436]A. F.; Alitalo, K.; Stacker, S. A.: Vascular endothelial
growth factor

[103437]D (VEGF-D) is a ligand for the tyrosine kinases VEGF re-
ceptor 2 (Flk1)

[103438]and VEGF receptor 3 (Flt4). Proc. Nat. Acad. Sci. 95:
548–553, 1998.

[103439]18060.

[103440]Stacker, S. A.; Caesar, C.; Baldwin, M. E.; Thornton, G. E.;
Williams,

[103441]R. A.; Prevo, R.; Jackson, D. G.; Nishikawa, S.; Kubo, H.;
Achen,

[103442]M. G.: VEGF-D promotes the metastatic spread of tumor
cells via the

[103443]lymphatics. Nature Med. 7: 186–191, 2001.

[103444]18061.

[103445]Yamada, Y.; Nezu, J.; Shimane, M.; Hirata, Y.: Molecular
cloning

[103446]of a novel vascular endothelial growth factor, VEGF-D. Ge-
nomics 42:

[103447]483–488, 1997.

[103448]18062.

[103449]Davies, P. A.; Hanna, M. C.; Hales, T. G.; Kirkness, E. F.:
Insensitivity

[103450]to anaesthetic agents conferred by a class of GABA(A) re-
ceptor subunit. Nature 385:

[103451]820–823, 1997.

[103452]18063.

[103453]Sinkkonen, S. T.; Hanna, M. C.; Kirkness, E. F.; Korpi, E. R.:

[103454]GABA-A receptor epsilon and theta subunits display un-
usual structural

[103455]variation between species and are enriched in the rat locus ceruleus. J.

[103456]Neurosci. 20: 3588–3595, 2000.

[103457]18064.

[103458]Wilke, K.; Gaul, R.; Klauck, S. M.; Poustka, A.: A gene in human

[103459]chromosome band Xq28 (GABRE) defines a putative new subunit class

[103460]of the GABA(A) neurotransmitter receptor. Genomics 45: 1–10, 1997.

[103461]18065.

[103462]Mazzarella, R.; Ciccodicola, A.; Esposito, T.; Arcucci, A.; Migliaccio,

[103463]C.; Jones, C.; Schlessinger, D.; D'Urso, M.; D'Esposito, M.: Human

[103464]protein kinase C iota gene (PRKCI) is closely linked to the BTK gene

[103465]in Xq21.3. Genomics 26: 629–631, 1995.

[103466]18066.

[103467]Lafreniere, R. G.; Carrel, L.; Willard, H. F.: A novel trans-membrane

[103468] transporter encoded by the XPCT gene in Xq13.2. Hum.

Molec. Genet. 3:

[103469] 1133–1139, 1994.

[103470] 18067.

[103471] Baric, I.; Sarnavka, V.; Fumic, K.; Maradin, M.; Begovic, D.;

Ruiter,

[103472] J. P. N.; Wanders, R. J. A.: A new case of succinyl–

CoA:acetoacetate

[103473] transferase deficiency: favourable course despite very low
residual

[103474] activity. J. Inherit. Metab. Dis. 24: 81–82, 2001.

[103475] 18068.

[103476] Berry, G. T.; Fukao, T.; Mitchell, G. A.; Mazur, A.; Ciafre,

M.;

[103477] Gibson, J.; Kondo, N.; Palmieri, M. J.: Neonatal hypogly–

caemia in

[103478] severe succinyl–CoA:3–oxoacid CoA–transferase defi–
ciency. J. Inherit.

[103479] Metab. Dis. 24: 587–595, 2001.

[103480] 18069.

[103481] Fukao, T.; Mitchell, G. A.; Song, X.–Q.; Nakamura, H.;

Kassovska-Bratinova,

[103482]S.; Orii, K. E.; Wraith, J. E.; Besley, G.; Wanders, R. J. A.;

Niezen-Koning,

[103483]K. E.; Berry, G. T.; Palmieri, M.; Kondo, N.: Succinyl-

CoA:3-ketoacid

[103484]CoA transferase (SCOT): cloning of the human SCOT gene,
tertiary structural

[103485]modeling of the human SCOT monomer, and characteri-
zation of three

[103486]pathogenic mutations. Genomics 68: 144-151, 2000.

[103487]18070.

[103488]Fukao, T.; Song, X.-Q.; Watanabe, H.; Hirayama, K.;
Sakazaki, H.;

[103489]Shintaku, H.; Imanaka, M.; Orii, T.; Kondo, N.: Prenatal di-
agnosis

[103490]of succinyl-coenzyme A:3-ketoacid coenzyme A trans-
ferase deficiency. Prenatal

[103491]Diag. 16: 471-474, 1996.

[103492]18071.

[103493]Kassovska-Bratinova, S.; Fukao, T.; Song, X.-Q.; Duncan,
A. M.

[103494]Y.; Chen, H. S.; Robert, M.-F.; Perez-Cerda, C.; Ugarte, M.;

Chartrand,

[103495]C.; Vobecky, S.; Kondo, N.; Mitchell, G. A.: Succinyl

CoA:3-oxoacid

[103496]CoA transferase (SCOT): human cDNA cloning, human
chromosomal mapping

[103497]to 5p13, and mutation detection in a SCOT-deficient pa-
tient. Am.

[103498]J. Hum. Genet. 59: 519-528, 1996.

[103499]18072.

[103500]Middleton, B.; Day, R.; Lombes, A.; Saudubray, J. M.: In-
fantile

[103501]ketoacidosis associated with decreased activity of suc-
cinyl-CoA:3-ketoacid

[103502]CoA-transferase. J. Inherit. Metab. Dis. 10 (suppl. 2):
273-275,

[103503]1987.

[103504]18073.

[103505]Mitchell, G. A.; Kassovska-Bratinova, S.; Boukaftane, Y.;
Robert,

[103506]M.-F.; Wang, S. P.; Ashmarina, L.; Lambert, M.; Lapierre,
P.; Potier,

[103507]E.: Medical aspects of ketone body metabolism. Clin. In-

vest. Med. 18:

[103508]193–216, 1995.

[103509]18074.

[103510]Niezen-Koning, K. E.; Wanders, R. J. A.; Ruiter, J. P. N.;
Ijlst,

[103511]L.; Visser, G.; Reitsma-Bierens, W. C. C.; Heymans, H. S.
A.; Reijngoud,

[103512]D. J.; Smit, G. P. A.: Succinyl-CoA:acetoacetate transferase
deficiency:

[103513]Identification of a new patient with a neonatal onset and
review of

[103514]the literature. *Europ. J. Pediatr.* 156: 870–873, 1997.

[103515]18075.

[103516]Perez-Cerda, C.; Merinero, B.; Sanz, P.; Jimenez, A.; Her-
nandez,

[103517]C.; Garcia, M. J.; Ugarte, M.: A new case of succinyl-
CoA:acetoacetate

[103518]transferase deficiency. *J. Inherit. Metab. Dis.* 15: 371–373,
1992.

[103519]18076.

[103520]Sakazaki, H.; Hirayama, K.; Murakami, S.; Yonezawa, S.;

Shintaku,

[103521]H.; Sawada, Y.; Fukao, T.; Watanabe, H.; Orii, T.; Isshiki, G.: A

[103522]new Japanese case of succinyl-CoA:3-ketoacid CoA-transferase deficiency. J.

[103523]Inherit. Metab. Dis. 18: 323-325, 1995.

[103524]18077.

[103525]Snyderman, S. E.; Sansaricq, C.; Middleton, B.: Succinyl-CoA:3-ketoacid

[103526]CoA-transferase deficiency. Pediatrics 101: 709-711, 1998.

[103527]18078.

[103528]D'Adamo, P.; Fassone, L.; Gedeon, A.; Janssen, E. A. M.; Bione,

[103529]S.; Bolhuis, P. A.; Barth, P. G.; Wilson, M.; Haan, E.; Orstavik,

[103530]K. H.; Patton, M. A.; Green, A. J.; Zammarchi, E.; Donati, M. A.;

[103531]Toniolo, D.: The X-linked gene G4.5 is responsible for different

[103532]infantile dilated cardiomyopathies. Am. J. Hum. Genet. 61: 862-867,

[103533]1997.

[103534]18079.

[103535]Gedeon, A. K.; Wilson, M. J.; Colley, A. C.; Sillence, D. O.;
Mulley,

[103536]J. C.: X linked fatal infantile cardiomyopathy maps to Xq28
and is

[103537]possibly allelic to Barth syndrome. J. Med. Genet. 32:
383–388,

[103538]1995.

[103539]18080.

[103540]Gecz, J.; Baker, E.; Donnelly, A.; Ming, J. E.; McDonald–
McGinn,

[103541]D. M.; Spinner, N. B.; Zackai, E. H.; Sutherland, G. R.; Mul–
ley, J.

[103542]C.: Fibroblast growth factor homologous factor 2 (FHF2):
gene structure,

[103543]expression and mapping to the Borjeson–Forss–
man–Lehmann syndrome region

[103544]in Xq26 delineated by a duplication breakpoint in a BFLS–
like patient. Hum.

[103545]Genet. 104: 56–63, 1999.

[103546]18081.

[103547]Lovec, H.; Hartung, H.; Verdier, A.-S.; Mattei, M.-G.; Birnbaum,

[103548]D.; Goldfarb, M.; Coulier, F.: Assignment of FGF13 to human chromosome

[103549]band Xq21 by in situ hybridization. Cytogenet. Cell Genet. 76: 183-184,

[103550]1997.

[103551]18082.

[103552]Smallwood, P. M.; Munoz-Sanjuan, I.; Tong, P.; Macke, J. P.; Hendry,

[103553]S. H. C.; Gilbert, D. J.; Copeland, N. G.; Jenkins, N. A.; Nathans,

[103554]J.: Fibroblast growth factor (FGF) homologous factors: new members

[103555]of the FGF family implicated in nervous system development. Proc.

[103556]Nat. Acad. Sci. 93: 9850-9857, 1996.

[103557]18083.

[103558]Jones, M. H.; Furlong, R. A.; Burkin, H.; Chalmers, I. J.; Brown,

[103559]G. M.; Khwaja, O.; Affara, N. A.: The Drosophila developmental gene

[103560]fat facets has a human homologue in Xp11.4 which escapes X-inactivation

[103561]and has related sequences on Yq11.2. Hum. Molec. Genet. 5: 1695-1701,

[103562]1996.

[103563]18084.

[103564]Caldas, C.; Hahn, S. A.; da Costa, L. T.; Redston, M. S.; Schutte,

[103565]M.; Seymour, A. B.; Weinstein, C. L.; Hruban, R. H.; Yeo, C. J.; Kern,

[103566]S. E.: Frequent somatic mutations and homozygous deletions of the

[103567]p16 (MTS1) gene in pancreatic adenocarcinoma. Nature Genet. 8: 27-32,

[103568]1994.

[103569]18085.

[103570]Liu, Q.; Yan, Y.-X.; McClure, M.; Nakagawa, H.; Fujimura, F.;

[103571]Rustgi, A. K.: MTS-1 (CDKN2) tumor suppressor gene deletions are

[103572]a frequent event in esophagus squamous cancer and pancreatic adenocarcinoma

[103573]cell lines. *Oncogene* 10: 619–622, 1995.

[103574]18086.

[103575]Whelan, A. J.; Bartsch, D.; Goodfellow, P. J.: Brief report:

[103576]a familial syndrome of pancreatic cancer and melanoma with a mutation

[103577]in the CDKN2 tumor-suppressor gene. *New Eng. J. Med.* 333: 975–977,

[103578]1995.

[103579]18087.

[103580]Schutte, M.; da Costa, L. T.; Hahn, S. A.; Moskaluk, C.; Hoque,

[103581]A. T. M. S.; Rozenblum, E.; Weinstein, C. L.; Bittner, M.; Meltzer,

[103582]P. S.; Trent, J. M.; Yeo, C. J.; Hruban, R. H.; Kern, S. E.: Identification

[103583]by representational difference analysis of a homozygous deletion in

[103584]pancreatic carcinoma that lies with the BRCA2 region. *Proc. Nat.*

[103585]Acad. Sci. 92: 5950–5954, 1995.

[103586]18088.

[103587]Stoffers, D. A.; Zinkin, N. T.; Stanojevic, V.; Clarke, W. L.;

[103588]Habener, J. F.: Pancreatic agenesis attributable to a single nucleotide

[103589]deletion in the human IPF1 gene coding sequence. Nature Genet. 15:

[103590]106–110, 1997.

[103591]18089.

[103592]Tacconi, P.; Ferrigno, P.; Cocco, L.; Cannas, A.; Tamburini, G.;

[103593]Bergonzi, P.; Giagheddu, M.: Sclerosteosis: report of a case in a

[103594]black African man. Clin. Genet. 53: 497–501, 1998.

[103595]18090.

[103596]Tanzi, R. E.; Petrukhin, K.; Chernov, I.; Pellequer, J. L.; Wasco,

[103597]W.; Ross, B.; Romano, D. M.; Parano, E.; Pavone, L.; Brzustowicz,

[103598]L. M.; Devoto, M.; Peppercorn, J.; Bush, A. I.; Sternlieb, I.; Pirastu,

[103599]M.; Gusella, J. F.; Evgrafov, O.; Penchaszadeh, G. K.;
Honig, B.;

[103600]Edelman, I. S.; Soares, M. B.; Scheinberg, I. H.; Gilliam, T.
C.:

[103601]The Wilson disease gene is a copper transporting ATPase
with homology

[103602]to the Menkes disease gene. Nature Genet. 5: 344–350,
1993.

[103603]18091.

[103604]Terada, K.; Nakako, T.; Yang, X.-L.; Iida, M.; Aiba, N.; Mi-
namiya,

[103605]Y.; Nakai, M.; Sakaki, T.; Miura, N.; Sugiyama, T.: Restora-
tion of

[103606]holoceruloplasmin synthesis in LEC rat after infusion of
recombinant

[103607]adenovirus bearing WND cDNA. J. Biol. Chem. 273:
1815–1820, 1998.

[103608]18092.

[103609]Thomas, G. R.; Forbes, J. R.; Roberts, E. A.; Walshe, J. M.;
Cox,

[103610]D. W.: The Wilson disease gene: spectrum of mutations
and their consequences. Nature

- [103611]Genet. 9: 210–217, 1995. Note: Correction. Nature Genet.
9: 451 only,
[103612]1995.
[103613]18093.
- [103614]Thomas, G. R.; Jensson, O.; Gudmundsson, G.;
Thorsteinsson, L.;
[103615]Cox, D. W.: Wilson disease in Iceland: a clinical and ge-
netic study. Am.
[103616]J. Hum. Genet. 56: 1140–1146, 1995.
[103617]18094.
- [103618]Thomas, G. R.; Roberts, E. A.; Walshe, J. M.; Cox, D. W.:
Haplotypes
[103619]and mutations in Wilson disease. Am. J. Hum. Genet. 56:
1315–1319,
[103620]1995.
[103621]18095.
- [103622]Beutler, E.; Kuhl, W.; Comings, D.: Hexosaminidase
isozyme in
[103623]type O Gm2 gangliosidosis (Sandhoff–Jatzkewitz disease).
Am. J. Hum.
[103624]Genet. 27: 628–638, 1975.

[103625]18096.

[103626]Boustany, R.-M. N.; Tanaka, A.; Nishimoto, J.; Suzuki, K.:
Genetic

[103627]cause of a juvenile form of Tay-Sachs disease in a
Lebanese child. Ann.

[103628]Neurol. 29: 104-107, 1991.

[103629]18097.

[103630]Brown, C. A.; Mahuran, D. J.: Beta-hexosaminidase
isozymes from

[103631]cells cotransfected with alpha and beta cDNA constructs:
analysis

[103632]of the alpha-subunit missense mutation associated with
the adult form

[103633]of Tay-Sachs disease. Am. J. Hum. Genet. 5: 497-508,
1993.

[103634]18098.

[103635]Chern, C. J.; Beutler, E.; Kuhl, W.; Gilbert, F.; Mellman, W.

[103636]J.; Croce, C. M.: Characterization of heteropolymeric hex-
osaminidase

[103637]A in human x mouse hybrid cells. Proc. Nat. Acad. Sci. 73:
3637-3640,

[103638]1976.

[103639]18099.

[103640]Chern, C. J.; Kennett, R.; Engel, E.; Mellman, W. J.; Croce, C.

[103641]M.: Assignment of the structural genes for the alpha sub-unit of hexosaminidase

[103642]A, mannosephosphate isomerase and pyruvate kinase to the region of

[103643]22-qter of human chromosome 15. Somat. Cell Genet. 3: 553-560, 1977.

[103644]18100.

[103645]Coulter-Mackie, M. B.: Molecular characterization of both alleles

[103646]in an unusual Tay-Sachs disease B1 variant. (Letter) Am. J. Hum.

[103647]Genet. 54: 1126-1127, 1994.

[103648]18101.

[103649]De Braekeleer, M.; Hechtman, P.; Andermann, E.; Kaplan, F.: The

[103650]French Canadian Tay-Sachs disease deletion mutation: identification

[103651] of probable founders. Hum. Genet. 89: 83–87, 1992.

[103652] 18102.

[103653] De Gasperi, R.; Gama Sosa, M. A.; Battistini, S.; Yeretsian, J.;

[103654] Raghavan, S.; Zelnik, N.; Leshinsky, E.; Kolodny, E. H.:
Late-onset

[103655] G(M2)–gangliosidosis: Ashkenazi Jewish family with an
exon 5 mutation

[103656] (tyr180–to–his) in the Hex A alpha–chain gene. Neurology
47: 547–552,

[103657] 1996.

[103658] 18103.

[103659] dos Santos, M. R.; Tanaka, A.; sa Miranda, M. C.; Ribeiro, M.

[103660] G.; Maia, M.; Suzuki, K.: G(M2)–gangliosidosis B1 variant:
analysis

[103661] of beta–hexosaminidase alpha gene mutations in 11 pa-
tients from a

[103662] defined region in Portugal. Am. J. Hum. Genet. 49:
886–890, 1991.

[103663] 18104.

[103664]Drucker, L.; Navon, R.: Tay-Sachs disease in an Israeli Arab

[103665]family: trp26-to-stop in the alpha-subunit of hexosaminidase A. Hum.

[103666]Mutat. 2: 415-417, 1993.

[103667]18105.

[103668]Drucker, L.; Proia, R. L.; Navon, R.: Identification and rapid
[103669]detection of three Tay-Sachs mutations in the Moroccan Jewish population. Am.

[103670]J. Hum. Genet. 51: 371-377, 1992.

[103671]18106.

[103672]Fernandes, M.; Kaplan, F.; Natowicz, M.; Prence, E.;
Kolodny,

[103673]E.; Kaback, M.; Hechtman, P.: A new Tay-Sachs disease B1
allele in

[103674]exon 7 in two compound heterozygotes each with a second novel mutation. Hum.

[103675]Molec. Genet. 1: 759-761, 1992.

[103676]18107.

[103677]Fernandes, M. J. G.; Hechtman, P.; Boulay, B.; Kaplan, F.: A

[103678]chronic GM(2) gangliosidosis variant with a HEXA splicing defect:

[103679]quantitation of HEXA mRNAs in normal and mutant fibroblasts. *Europ.*

[103680]*J. Hum. Genet.* 5: 129–136, 1997.

[103681]18108.

[103682]Formiga, L. de F.; Poenaru, L.; Couronne, F.; Flori, E.; Eibel,

[103683]J. L.; Deminatti, M. M.; Savary, J. B.; Lai, J. L.; Gilgenkrantz,

[103684]S.; Pierson, M.: Interstitial deletion of chromosome 15: two cases. *Hum.*

[103685]*Genet.* 80: 401–404, 1988.

[103686]18109.

[103687]Goebel, H. H.; Stolte, G.; Kustermann-Kuhn, B.; Harzer, K.: B(1)

[103688]variant of G(M2) gangliosidosis in a 12-year-old patient. *Pediat.*

[103689]*Res.* 25: 89–93, 1989.

[103690]18110.

[103691]Gordon, B. A.; Gordon, K. E.; Hinton, G. G.; Cadera, W.; Feleki,

[103692]V.; Bayleran, J.; Hechtman, P.: Tay Sachs disease: B1 variant. *Pediat.*

[103693]*Neurol.* 4: 54–57, 1988.

[103694]18111.

[103695]Greenberg, D. A.; Kaback, M. M.: Estimation of the frequency

[103696]of hexosaminidase A variant alleles in the American Jewish population. *Am.*

[103697]*J. Hum. Genet.* 34: 444–451, 1982.

[103698]18112.

[103699]Natt, E.; Westphal, E.-M.; Toth-Fejel, S. E.; Magenis, R. E.;

[103700]Buist, N. R. M.; Rettenmeier, R.; Scherer, G.: Inherited and de novo

[103701]deletion of the tyrosine aminotransferase gene locus at 16q22.1–q22.3

[103702]in a patient with tyrosinemia type II. *Hum. Genet.* 77: 352–358,

[103703]1987.

[103704]18113.

[103705]Pelet, B.; Antener, I.; Faggioni, R.; Spahr, A.; Gautier, E.:

- [103706]Tyrosinemia without liver or renal damage with plantar and palmar
- [103707]keratosis and keratitis (hypertyrosinemia type II). *Helv. Paediat.*
- [103708]*Acta* 34: 177–183, 1979.
- [103709]18114.
- [103710]Rehak, A.; Selim, M. M.; Yadav, G.: Richner–Hanhart syndrome
- [103711](tyrosinaemia–II) (report of four cases without ocular involvement). *Brit.*
- [103712]*J. Derm.* 104: 469–475, 1981.
- [103713]18115.
- [103714]Rettenmeier, R.; Natt, E.; Zentgraf, H.; Scherer, G.: Isolation
- [103715]and characterization of the human tyrosine aminotransferase gene. *Nucleic*
- [103716]*Acids Res.* 18: 3853–3861, 1990.
- [103717]18116.
- [103718]Richner, H.: Hornhautaffektion bei Keratoma palmare et plantare
- [103719]hereditarium. *Klin. Mbl. Augenheilk.* 100: 580–588, 1938.

[103720]18117.

[103721]Schmid, W.; Muller, G.; Schutz, G.; Gluecksohn-Waelsch, S.: Deletions

[103722]near the albino locus on chromosome 7 of the mouse affect the level

[103723]of tyrosine aminotransferase mRNA. Proc. Nat. Acad. Sci. 82: 2866-2869,

[103724]1985.

[103725]18118.

[103726]Tallab, T.M.: Richner-Hanhart syndrome: importance of early diagnosis

[103727]and early intervention. J. Am. Acad. Dermatol. 35: 857-859, 1996.

[103728]18119.

[103729]Ventura, G.; Biasini, G.; Petrozzi, M.: Cheratomia palmo-plantare

[103730]dissipatum associato a lesioni corneali in due fratelli. Boll. Oculist. 44:

[103731]497-510, 1965.

[103732]18120.

[103733]Waardenburg, P. J.; Franceschetti, A.; Klein, D.: Genetics

and

[103734]Ophthalmology. Springfield, Ill.: Charles C Thomas (pub.)

1:

[103735]1961. Pp. 515-517.

[103736]18121.

[103737]Westphal, E.-M.; Natt, E.; Grimm, T.; Odievre, M.; Scherer,
G.

[103738]: The human tyrosine aminotransferase gene: characteri-
zation of restriction

[103739]fragment length polymorphisms and haplotype analysis in
a family with

[103740]tyrosinemia type II. Hum. Genet. 79: 260-264, 1988.

[103741]18122.

[103742]Perheentupa, J.: Autoimmune polyendocrinopathy-candido-
sis-ectodermal

[103743]dystrophy (APECED).In: Eriksson, A. W.; Forsius, H. R.;
Nevanlinna,

[103744]H. R.; Workman, P. L.; Norio, R. K.: Population Structure
and Genetic

[103745]Disorders. New York: Academic Press (pub.) 1980. Pp.
583-587.

[103746]18123.

[103747]Prader, A.: Personal Communication. Zurich, Switzerland
1972.

[103748]18124.

[103749]Ramsay, C.; Winqvist, O.; Puhakka, L.; Halonen, M.; Moro,
A.;

[103750]Kampe, O.; Eskelin, P.; Pelto-Huikko, M.; Peltonen, L.: Aire
deficient

[103751]mice develop multiple features of APECED phenotype and
show altered

[103752]immune response. Hum. Molec. Genet. 11: 397-409,
2002.

[103753]18125.

[103754]Rinderle, C.; Christensen, H.-M.; Schweiger, S.; Lehrach,
H.;

[103755]Yaspo, M.-L.: AIRE encodes a nuclear protein co-localizing
with cytoskeletal

[103756]filaments: altered sub-cellular distribution of mutants
lacking PHD

[103757]zinc fingers. Hum. Molec. Genet. 8: 277-290, 1999.

[103758]18126.

[103759]Rosatelli, M. C.; Meloni, A.; Meloni, A.; Devoto, M.; Cao,
A.;

- [103760] Scott, H. S.; Peterson, P.; Heino, M.; Krohn, K. J.;
Nagamine, K.;
- [103761] Kudoh, J.; Shimizu, N.; Antonarakis, S. E.: A common mu-
tation in
- [103762] Sardinian autoimmune polyendocrinopathy-candidia-
sis-ectodermal dystrophy
- [103763] patients. Hum. Genet. 103: 428-434, 1998.
- [103764] 18127.
- [103765] Saenger, P.: Primary hypoaldosteronism due to zona
glomerulosa
- [103766] defect. (Letter) New Eng. J. Med. 310: 1394 only, 1984.
- [103767] 18128.
- [103768] Saenger, P.; Levine, L. S.; Irvine, W. J.; Gottesdiener, K.;
- Rauh,
- [103769] W.; Sonino, N.; Chow, D.; New, M. I.: Progressive adrenal
failure
- [103770] in polyglandular autoimmune disease. J. Clin. Endocr.
Metab. 54:
- [103771] 863-868, 1982.
- [103772] 18129.
- [103773] Scott, H. S.; Heino, M.; Peterson, P.; Mittaz, L.; Lalioti, M.

[103774]D.; Betterle, C.; Cohen, A.; Seri, M.; Lerone, M.; Romeo, G.; Collin,
 [103775]P.; Salo, M.; and 9 others: Common mutations in autoimmune polyendocrinopathy–candidiasis–ectodermal
 [103776]dystrophy patients of different origins. *Molec. Endocr.* 12: 1112–1119,
 [103777]1998.
 [103778]18130.

[103779]Shapiro, M. S.; Zamir, R.; Weiss, E.; Radnay, J.; Shenkman, L.
 [103780]: The polyglandular deficiency syndrome: a new variant in Persian
 [103781]Jews. *J. Endocr. Invest.* 10: 1–7, 1987.
 [103782]18131.

[103783]Spinner, M. W.; Blizzard, R. M.; Childs, B.: Clinical and genetic
 [103784]heterogeneity in idiopathic Addison's disease and hypoparathyroidism. *J.*
 [103785]*Clin. Endocr.* 28: 795–804, 1968.
 [103786]18132.

[103787]Sweetnam, W. P.: Juvenile familial endocrinopathy. *Lancet*

I:

[103788]463–465, 1966.

[103789]18133.

[103790]Wang, C.-Y.; Davoodi-Semiromi, A.; Huang, W.; Connor, E.; Shi,

[103791]J.-D.; She, J.-X.: Characterization of mutations in patients with

[103792]autoimmune polyglandular syndrome type 1 (APS1). Hum. Genet. 103:

[103793]681–685, 1998.

[103794]18134.

[103795]Whitaker, J. A.; Landing, B. H.; Esselborn, V. M.; Williams, R.

[103796]R.: Syndrome of familial juvenile hypoadrenocorticism, hypoparathyroidism

[103797]and superficial moniliasis. J. Clin. Endocr. 16: 1374–1387, 1956.

[103798]18135.

[103799]Wirfalt, A.: Genetic heterogeneity in autoimmune polyglandular

[103800]failure. Acta Med. Scand. 210: 7–13, 1981.

[103801]18136.

[103802]Zlotogora, J.; Shapiro, M. S.: Polyglandular autoimmune syndrome

[103803]type I among Iranian Jews. J. Med. Genet. 29: 824–826, 1992.

[103804]18137.

[103805]Hediger, M. A.: New view at C. Nature Med. 8: 445–446, 2002.

[103806]18138.

[103807]Sotiriou, S.; Gispert, S.; Cheng, J.; Wang, Y.; Chen, A.; Hoogstraten–Miller,

[103808]S.; Miller, G. F.; Kwon, O.; Levine, M.; Guttentag, S. H.; Nussbaum,

[103809]R. L.: Ascorbic–acid transporter Slc23a1 is essential for vitamin

[103810]C transport into the brain and for perinatal survival. Nature Med. 8:

[103811]514–517, 2002.

[103812]18139.

[103813]Chuang, D. T.; Fisher, C. W.; Lau, K. S.; Griffin, T. A.; Wynn,

[103814]R. M.; Cox, R. P.: Maple syrup urine disease: domain structure, mutations
[103815]and exon skipping in the dihydrolipoyl transacylase (E2) component
[103816]of the branched-chain alpha-keto acid dehydrogenase complex. Molec.
[103817]Biol. Med. 8: 49-63, 1991.
[103818]18140.
[103819]Chuang, J. L.; Cox, R. P.; Chuang, D. T.: E2 transacylase-deficient
[103820](type II) maple syrup urine disease: aberrant splicing of E2 mRNA
[103821]caused by internal intronic deletions and association with thiamine-responsive
[103822]phenotype. J. Clin. Invest. 100: 736-744, 1997.
[103823]18141.
[103824]Fisher, C. W.; Fisher, C. R.; Chuang, J. L.; Lau, K. S.; Chuang,
[103825]D. T.; Cox, R. P.: Occurrence of a 2-bp (AT) deletion allele and
[103826]nonsense (G-to-T) mutant allele at the E2 (DBT) locus of six patients

[103827]with maple syrup urine disease: multiple-exon skipping as a secondary

[103828]effect of the mutations. Am. J. Hum. Genet. 52: 414-424, 1993.

[103829]18142.

[103830]Fisher, C. W.; Lau, K. S.; Fisher, C. R.; Wynn, R. M.; Cox, R.

[103831]P.; Chuang, D. T.: A 17-bp insertion and a phe215-to-cys missense

[103832]mutation in the dihydrolipoyl transacylase (E2) mRNA from a thiamine-responsive

[103833]maple syrup urine disease patient WG-34. Biochem. Biophys. Res. Commun. 174:

[103834]804-809, 1991.

[103835]18143.

[103836]Herring, W. J.; Litwer, S.; Weber, J. L.; Danner, D. J.: Molecular

[103837]genetic basis of maple syrup urine disease in a family with two defective

[103838]alleles for branched chain acyltransferase and localization of the

[103839]gene to human chromosome 1. Am. J. Hum. Genet. 48: 342-350, 1991.

[103840]18144.

[103841]Hummel, K. B.; Litwer, S.; Bradford, A. P.; Aitken, A.; Dan-
ner,

[103842]D. J.; Yeaman, S. J.: Nucleotide sequence of a cDNA for
branched

[103843]chain acyltransferase with analysis of the deduced protein
structure. J.

[103844]Biol. Chem. 263: 6165–6168, 1988.

[103845]18145.

[103846]Indo, Y.; Akaboshi, I.; Nobukuni, Y.; Endo, F.; Matsuda, I.:
Maple

[103847]syrup urine disease: a possible biochemical basis for the
clinical

[103848]heterogeneity. Hum. Genet. 80: 6–10, 1988.

[103849]18146.

[103850]Lau, K. S.; Eddy, R. L.; Shows, T. B.; Fisher, C. W.; Chuang,

[103851]D. T.; Cox, R. P.: Localization of the dihydrolipoamide
branched–chain

[103852]transacylase gene (DBT) of the human branched–chain
keto acid dehydrogenase

[103853]complex to chromosome 1. Cytogenet. Cell Genet. 56:

33–35, 1991.

[103854]18147.

[103855]Lau, K. S.; Griffin, T. A.; Hu, C.-W. C.; Chuang, D. T.: Conservation

[103856]of primary structure in the lipoyl-bearing and dihydrolipoyl dehydrogenase

[103857]binding domains of mammalian branched-chain alpha-keto acid dehydrogenase

[103858]complex: molecular cloning of human and bovine transacylase (E2) cDNAs. *Biochemistry* 27:

[103859]1972–1981, 1988.

[103860]18148.

[103861]Lau, K. S.; Herring, W. J.; Chuang, J. L.; McKean, M.; Danner,

[103862]D. J.; Cox, R. P.; Chuang, D. T.: Structure of the gene encoding

[103863]dihydrolipoyl transacylase (E2) component of human branched chain

[103864]alpha-keto acid dehydrogenase and characterization of an E2 pseudogene. *J.*

[103865]*Biol. Chem.* 267: 24090–24096, 1992.

[103866]18149.

[103867]Lebo, R. V.; Shapiro, L. R.; Fenerci, E. Y.; Hoover, J. M.;
Chuang,
[103868]J. L.; Chuang, D. T.; Kronn, D. F.: Rare etiology of autosomal recessive
[103869]disease in a child with noncarrier parents. Am. J. Hum.
Genet. 67:
[103870]750–754, 2000.
[103871]18150.
[103872]Litwer, S.; Danner, D. J.: Identification of a cDNA clone in
[103873]lambda-gt11 for the transacylase component of branched
chain ketoacid
[103874]dehydrogenase. Biochem. Biophys. Res. Commun. 131:
961–967, 1985.
[103875]18151.
[103876]Litwer, S.; Danner, D. J.: Mitochondrial import and processing
[103877]of an in vitro synthesized human prebranched chain acyltransferase
[103878]fragment. Am. J. Hum. Genet. 43: 764–769, 1988.
[103879]18152.
[103880]Mitsubuchi, H.; Nobukuni, Y.; Akaboshi, I.; Indo, Y.; Endo,

F.;

[103881]Matsuda, I.: Maple syrup urine disease caused by a partial deletion

[103882]in the inner E2 core domain of the branched chain alpha-keto acid

[103883]dehydrogenase complex due to aberrant splicing: a single base deletion

[103884]at a 5-prime-splice donor site of an intron of the E2 gene disrupts

[103885]the consensus sequence in this region. J. Clin. Invest. 87: 1207-1211,

[103886]1991.

[103887]18153.

[103888]Tsuruta, M.; Mitsubuchi, H.; Mardy, S.; Miura, Y.; Hayashida,

[103889]Y.; Kinugasa, A.; Ishitsu, T.; Matsuda, I.; Indo, Y.: Molecular basis

[103890]of intermittent maple syrup urine disease: novel mutations in the

[103891]E2 gene of the branched-chain alpha-keto acid dehydrogenase complex. J.

[103892]Hum. Genet. 43: 91-100, 1998.

[103893]18154.

[103894]Zneimer, S. M.; Lau, K. S.; Eddy, R. L.; Shows, T. B.;
Chuang,

[103895]J. L.; Chuang, D. T.; Cox, R. P.: Regional assignment of two
genes

[103896]of the human branched-chain alpha-keto acid dehydro-
genase complex:

[103897]the E1-beta gene (BCKDHB) to chromosome 6p21-22 and
the E2 gene (DBT)

[103898]to chromosome 1p31. Genomics 10: 740-747, 1991.

[103899]18155.

[103900]Gahl, W.; Finkelstein, J.; Martin, J.; Mullen, K.; Hoofnagle,
J.;

[103901]Backlund, P.; Bernar, J.; Bernardini, I.: Hepatic methionine
adenosyltransferase

[103902]deficiency in a 31-year-old man. (Abstract) Am. J. Hum.
Genet. 37:

[103903]A8 only, 1985.

[103904]18156.

[103905]Gahl, W. A.; Bernardini, I.; Finkelstein, J. D.; Tangerman,
A.;

- [103906]Martin, J. J.; Blom, H. J.; Mullen, K. D.; Mudd, S. H.:
Transsulfuration
[103907]in an adult with hepatic methionine adenosyltransferase
deficiency. J.
[103908]Clin. Invest. 81: 390–397, 1988.
[103909]18157.
[103910]Gahl, W. A.; Finkelstein, J. D.; Mullen, K. D.; Bernardini, I.;
[103911]Martin, J. J.; Backlund, P.; Ishak, K. G.; Hoofnagle, J. H.;
Mudd,
[103912]S. H.: Hepatic methionine adenosyltransferase deficiency
in a 31-year-old
[103913]man. Am. J. Hum. Genet. 40: 39–49, 1987.
[103914]18158.
[103915]Gaull, G. E.; Bender, A. N.; Vulovic, D.; Tallan, H. H.;
Schaffner,
[103916]F.: Methioninemia and myopathy: a new disorder. Ann.
Neurol. 9:
[103917]423–432, 1981.
[103918]18159.
[103919]Gaull, G. E.; Tallan, H. H.: Methionine adenosyltransferase
deficiency:

- [103920]new enzymatic defect associated with hypermethionine-
mia. Science 186:
- [103921]59-60, 1974.
- [103922]18160.
- [103923]Gaull, G. E.; Tallan, H. H.; Lonsdale, D.; Przyrembel, H.;
Schaffner,
- [103924]F.; von Bassewitz, D. B.: Hypermethioninemia associated
with methionine
- [103925]adenosyltransferase deficiency: clinical, morphologic, and
biochemical
- [103926]observations in four patients. J. Pediat. 98: 734-741,
1981.
- [103927]18161.
- [103928]Gout, J.-P.; Serre, J.-C.; Dieterlen, M.; Antener, I.; Frappat,
- [103929]P.; Bost, M.; Beaudoin, A.: Une nouvelle cause
d'hypermethioninemie
- [103930]de l'enfant: le deficit en S-adenosyl-
methionine-synthetase. Arch.
- [103931]Franc. Pediat. 34: 416-423, 1977.
- [103932]18162.
- [103933]Hazelwood, S.; Bernardini, I.; Shotelersuk, V.; Tangerman,

A.;

[103934]Guo, J.; Mudd, H.; Gahl, W. A.: Normal brain myelination in a patient

[103935]homozygous for a mutation that encodes a severely truncated methionine

[103936]adenosyltransferase I/III. Am. J. Med. Genet. 75: 395-400, 1998.

[103937]18163.

[103938]Horikawa, S.; Tsukada, K.: Molecular cloning and nucleotide sequence

[103939]of cDNA encoding the human liver S-adenosylmethionine synthetase. Biochem.

[103940]Int. 25: 81-90, 1991.

[103941]18164.

[103942]Hug, G.; Cussen, L. J.; Schubert, W. K.; Chuck, G.: 'Fingerprints'

[103943]in the liver: loss of methionine-activating enzyme. J. Clin. Invest. 47:

[103944]49A-50A, 1968.

[103945]18165.

[103946]Meny, R. G.; Gutberlet, R. L.; Ozand, P.; Morris, C.; Kim, C.

[103947]H.: Hypermethioninemia in an infant. Am. J. Dis. Child.
132: 261–262,

[103948]1978.

[103949]18166.

[103950]Mudd, S. H.; Levy, H. L.; Tangerman, A.; Boujet, C.; Buist,
N.;

[103951]Davidson–Mundt, A.; Hudgins, L.; Oyanagi, K.; Nagao, M.;
Wilson, W.

[103952]G.: Isolated persistent hypermethioninemia. Am. J. Hum.
Genet. 57:

[103953]882–892, 1995.

[103954]18167.

[103955]Natori, Y.: Studies on ethionine. VI. Sex-dependent behavior

[103956]of methionine and ethionine in rats. J. Biol. Chem. 238: 2075–2080,

[103957]1963.

[103958]18168.

[103959]Sakata, S. F.; Shelly, L. L.; Ruppert, S.; Schutz, G.; Chou, J.

[103960]Y.: Cloning and expression of murine S-adenosylmethionine synthetase. J.

[103961]Biol. Chem. 268: 13978–13986, 1993.

[103962]18169.

[103963]Surtees, R.; Leonard, J.; Austin, S.: Association of demyelination

[103964]with deficiency of cerebrospinal-fluid S-adenosylmethionine in inborn

[103965]errors of methyl-transfer pathway. Lancet 338: 1550–1554, 1991.

[103966]18170.

[103967]Ubagai, T.; Lei, K.-J.; Huang, S.; Mudd, S. H.; Levy, H. L.;

Chou,

[103968]J. Y.: Molecular mechanisms of an inborn error of methionine pathway:

[103969]methionine adenosyltransferase deficiency. J. Clin. Invest. 96:

[103970]1943–1947, 1995.

[103971]18171.

[103972]Cahane, M.; Treister, G.; Abraham, F. A.; Melamed, S.:
Glaucoma

[103973]in siblings with Morquio syndrome. Brit. J. Ophthal. 74:
382–383,

[103974]1990.

[103975]18172.

[103976]DiFerrante, N. M.; Ginsburg, L. C.; Donnelly, P. V.; DiFerrante,

[103977]D. T.; Caskey, C. T.: Deficiencies of glucosamine–6–sulfate
or galactosamine–6–sulfate

[103978]sulfatases are responsible for different mucopolysacchari-
doses. Science 199:

[103979]79–81, 1978.

[103980]18173.

- [103981]Fujimoto, A.; Horwitz, A. L.: Biochemical defect of non-keratan-sulfate-excreting
- [103982]Morquio syndrome. Am. J. Med. Genet. 15: 265-273, 1983.
- [103983]18174.
- [103984]Fukuda, S.; Tomatsu, S.; Masue, M.; Sukegawa, K.; Iwata, H.; Ogawa,
- [103985]T.; Nakashima, Y.; Hori, T.; Yamagishi, A.; Hanyu, Y.; Morooka, K.;
- [103986]Kiman, T.; Hashimoto, T.; Orie, T.: Mucopolysaccharidosis type IVA:
- [103987]N-acetyl-galactosamine-6-sulfate sulfatase exonic point mutations in
- [103988]classical Morquio and mild cases. J. Clin. Invest. 90: 1049-1053,
- [103989]1992.
- [103990]18175.
- [103991]Fukuda, S.; Tomatsu, S. Masuno, M.; Ogawa, T.; Yamagishi, A.;
- [103992]Rezvi, G. M. M.; Sukegawa, K.; Shimozawa, N.; Suzuki, Y.; Kondo, N.;
- [103993]Imaizumi, K.; Kuroki, Y.; Okabe, T.; Orie, T.: Mucopolysac-

charidosis

[103994]IVA: submicroscopic deletion of 16q24.3 and a novel
R386C mutation

[103995]of N-acetylgalactosamine-6-sulfate sulfatase gene in a
classical Morquio

[103996]disease. Hum. Mutat. 7: 123-134, 1996.

[103997]18176.

[103998]Gadbois, P.; Moreau, J.; Laberge, C.: La maladie de
Morquio dans

[103999]la province de Quebec. Un. Med. Canada 102: 602-607,
1973.

[104000]18177.

[104001]Gibson, G. J.; Saccone, G. T. P.; Brooks, D. A.; Clements, P.

[104002]R.; Hopwood, J. J.: Human N-acetyl-
galactosamine-4-sulphate sulphatase:

[104003]purification, monoclonal antibody production and native
and subunit

[104004]M(r) values. Biochem. J. 248: 755-764, 1987.

[104005]18178.

[104006]Glossl, J.; Kresse, H.; Mendla, K.; Cantz, M.; Rosenkranz,
W.

[104007]: Partial deficiency of glycoprotein neuraminidase in some patients

[104008]with Morquio disease type A. *Pediat. Res.* 18: 302–305, 1984.

[104009]18179.

[104010]Glossl, J.; Lembeck, K.; Gamse, G.; Kresse, H.: Morquio's disease

[104011]type A: absence of material cross reacting with antibodies against

[104012]N-acetylglactosamine-6-sulfate sulfatase. *Hum. Genet.* 54: 87–91,

[104013]1980.

[104014]18180.

[104015]Greenberg, A. D.: Atlantoaxial dislocations. *Brain* 91: 655–684,

[104016]1968.

[104017]18181.

[104018]Guiney, W. B., Jr.; Stevenson, R. E.: Morquio disease (MPS IV)

[104019]with survival to age 67 years. *Proc. Greenwood Genet. Center* 1:

[104020]84–87, 1982.

[104021]18182.

[104022]Hecht, J. T.; Scott, C. I., Jr.; Smith, T. K.; Williams, J. C.

[104023]: Mild manifestations of the Morquio syndrome. (Letter)
Am. J. Med.

[104024]Genet. 18: 369–371, 1984.

[104025]18183.

[104026]Holzgreve, W.; Grobe, H.; von Figura, K.; Kresse, H.; Beck,
H.;

[104027]Mattei, J. F.: Morquio syndrome: clinical findings in 11 pa-
tients

[104028]with MPS IVA and 2 patients with MPS IVB. Hum. Genet.
57: 360–365,

[104029]1981.

[104030]18184.

[104031]Hori, T.; Tomatsu, S.; Nakashima, Y.; Uchiyama, A.;
Fukuda, S.;

[104032]Sukegawa, K.; Shimozawa, N.; Suzuki, Y.; Kondo, N.; Hori-
uchi, T.;

[104033]Ogura, S.; Orii, T.: Mucopolysaccharidosis type IVA: com-
mon double

[104034]deletion in the N-acetylgalactosamine-6-sulfatase gene
(GALNS). Genomics 26:

[104035]535-542, 1995.

[104036]18185.

[104037]Hussels, I.: Morquio syndrome in a woman with two normal children.In:

[104038]Bergsma, D.: Skeletal Dysplasias. Amsterdam: Excerpta
Medica (pub.)

[104039]1974. Pp. 465-466.

[104040]18186.

[104041]Kato, Z.; Fukuda, S.; Tomatsu, S.; Vega, H.; Yasunaga, T.;
Yamagishi,

[104042]A.; Yamada, N.; Valencia, A.; Barrera, L. A.; Sukegawa, K.;
Orie,

[104043]T.; Kondo, N.: A novel common missense mutation G301C
in the N-acetylgalactosamine-6-sulfate

[104044]sulfatase gene in mucopolysaccharidosis IVA. Hum. Genet.
101: 97-101,

[104045]1997.

[104046]18187.

[104047]Langer, L. O., Jr.; Carey, L. S.: The roentgenographic fea-

tures

[104048]of the KS mucopolysaccharidosis of Morquio

(Morquio-Brailsford's disease). Am.

[104049]J. Roentgen. 97: 1-20, 1966.

[104050]18188.

[104051]Levin, L. S.; Jorgenson, R. J.; Salinas, C. F.: Oral findings

[104052]in the Morquio syndrome (mucopolysaccharidosis IV). Oral

Surg. Oral

[104053]Med. Oral Path. 39: 390-395, 1975.

[104054]18189.

[104055]Linker, A.; Evans, L. R.; Langer, L. O., Jr.: Morquio's dis-

ease

[104056]and mucopolysaccharide excretion. J. Pediat. 77:

1039-1047, 1970.

[104057]18190.

[104058]Maroteaux, P.; Lamy, M.: Opacités corneennes et troubles

metaboliques

[104059]dans la maladie de Morquio. Rev. Franc. Etud. Clin. Biol. 6:

481-483,

[104060]1961.

[104061]18191.

[104062]Masuno, M.; Tomatsu, S.; Nakashima, Y.; Hori, T.; Fukuda, S.;

[104063]Masue, M.; Sukegawa, K.; Oorii, T.: Mucopolysaccharidosis
IV A: assignment

[104064]of the human N-acetylgalactosamine-6-sulfate sulfatase
(GALNS) gene

[104065]to chromosome 16q24. Genomics 16: 777-778, 1993.

[104066]18192.

[104067]Matalon, R.; Arbogast, B.; Dorfman, A.: Morquio's syn-
drome: a

[104068]deficiency of chondroitin sulfate N-acetylhexosamine sul-
fate sulfatase.

[104069](Abstract) Pediat. Res. 8: 436, 1974.

[104070]18193.

[104071]McKusick, V. A.: Osler as a medical geneticist. Johns Hop-
kins

[104072]Med. J. 139: 163-174, 1976.

[104073]18194.

[104074]McKusick, V. A.: Heritable Disorders of Connective Tissue.
Non-keratan-sulfate-excreting

[104075]Morquio syndrome. St. Louis: C. V. Mosby (pub.) 1972. Pp.

600-604.

[104076]Note: Fig. 11-30...

[104077]18195.

[104078]Morquio, L.: Sur une forme de dystrophie osseuse familiale. Bull.

[104079]Soc. Pediat. Paris 27: 145-152, 1929.

[104080]18196.

[104081]Morris, C. P.; Guo, X.-H.; Apostolou, S.; Hopwood, J. J.; Scott,

[104082]H. S.: Morquio A syndrome: cloning, sequence, and structure of the

[104083]human N-acetylgalactosamine 6-sulfatase (GALNS) gene. Genomics 22:

[104084]652-654, 1994.

[104085]18197.

[104086]Nakashima, Y.; Tomatsu, S.; Hori, T.; Fukuda, S.; Sukegawa, K.;

[104087]Kondo, N.; Suzuki, Y.; Shimozawa, N.; Orii, T.: Mucopolysaccharidosis

[104088]IV A: molecular cloning of the human N-acetylgalactosamine-6-sulfatase

[104089]gene (GALNS) and analysis of the 5-prime-flanking region. Genomics 20:

[104090]99-104, 1994.

[104091]18198.

[104092]Nelson, J.; Broadhead, D.; Mossman, J.: Clinical findings in

[104093]12 patients with MPS IV A (Morquio's disease): further evidence for

[104094]heterogeneity. Part I: clinical and biochemical findings. Clin. Genet. 33:

[104095]111-120, 1988.

[104096]18199.

[104097]Nelson, J.; Kinirons, M.: Clinical findings in 12 patients with

[104098]MPS IV A (Morquio's disease): further evidence for heterogeneity.

[104099]Part II: dental findings. Clin. Genet. 33: 121-125, 1988.

[104100]18200.

[104101]Nelson, J.; Thomas, P. S.: Clinical findings in 12 patients with

[104102]MPS IV A (Morquio's disease): further evidence for heterogeneity.

- [104103]Part III: odontoid dysplasia. Clin. Genet. 33: 126–130,
1988.
- [104104]18201.
- [104105]Osler, W.: Sporadic cretinism in America. Am. J. Med. Sci.
114:
[104106]377–401, 1897.
- [104107]18202.
- [104108]Nauseef, W. M.; Brigham, S.; Cogley, M.: Hereditary
myeloperoxidase
[104109]deficiency due to a missense mutation of arginine 569 to
tryptophan. J.
[104110]Biol. Chem. 269: 1212–1216, 1994.
- [104111]18203.
- [104112]Bartholomew, L. G.; Baggenstoss, A. H.; Morlock, C. G.;
Comfort,
[104113]M. W.: Primary atrophy and lipomatosis of the pancreas.
Gastroenterology 36:
[104114]563–572, 1959.
- [104115]18204.
- [104116]Bodian, M.; Sheldon, W.; Lightwood, R.: Congenital hy-
poplasia

[104117]of the exocrine pancreas. Acta Paediat. 53: 282-293,
1964.

[104118]18205.

[104119]Cipolli, M.; D'Orazio, C.; Delmarco, A.; Marchesini, C.; Mi-
ano,

[104120]A.; Mastella, G.: Shwachman's syndrome: pathomorphosis
and long-term

[104121]outcome. J. Pediat. Gastroent. Nutr. 29: 265-272, 1999.

[104122]18206.

[104123]Danks, D. M.; Haslam, R. H. A.; Mayne, V.; Kaufmann, H. J.;
Holtzapple,

[104124]P. G.: Metaphyseal chondrodysplasia, neutropenia, and
pancreatic

[104125]insufficiency presenting with respiratory distress in the
neonatal

[104126]period. Arch. Dis. Child. 51: 697-701, 1976.

[104127]18207.

[104128]Davie, A. M.: The 'singles' method for segregation analysis
under

[104129]incomplete ascertainment. Ann. Hum. Genet. 42:
507-512, 1979.

[104130]18208.

[104131]Dokal, I.; Rule, S.; Chen, F.; Potter, M.; Goldman, J.: Adult

[104132]onset of acute myeloid leukaemia (M6) in patients with
Shwachman–Diamond

[104133]syndrome. Brit. J. Haemat. 99: 171–173, 1997.

[104134]18209.

[104135]Dror, Y.; Freedman, M. H.: Shwachman–Diamond syn-
drome. Brit.

[104136]J. Haemat. 118: 701–713, 2002.

[104137]18210.

[104138]Dror, Y.; Freedman, M. H.: Shwachman–Diamond syn-
drome: an inherited

[104139]preleukemic bone marrow failure disorder with aberrant
hematopoietic

[104140]progenitors and faulty marrow microenvironment. Blood
94: 3048–3054,

[104141]1999.

[104142]18211.

[104143]Dror, Y.; Freedman, M. H.: Shwachman–Diamond syn-
drome marrow

[104144]cells show abnormally increased apoptosis mediated through the Fas

[104145]pathway. Blood 97: 3011–3016, 2001.

[104146]18212.

[104147]Dror, Y.; Ginzberg, H.; Dalal, I.; Cherepanov, V.; Downey, G.;

[104148]Durie, P.; Roifman, C. M.; Freedman, M. H.: Immune function in patients

[104149]with Shwachman–Diamond syndrome. Brit. J. Haemat. 114: 712–717,

[104150]2001.

[104151]18213.

[104152]Fraccaro, M.; Scappaticci, S.; Arico, M.: Shwachman syndrome

[104153]and chromosome breakage. (Letter) Hum. Genet. 79: 194, 1988.

[104154]18214.

[104155]Genieser, N. B.; Halac, E. R.; Greco, M. A.; Selvarani Richards,

[104156]H. M.: Shwachman–Bodian syndrome. J. Comput. Assist. Tomogr. 6:

[104157]1191–1192, 1982.

[104158]18215.

[104159]Ginzberg, H.; Shin, J.; Ellis, L.; Goobie, S.; Morrison, J.;
Corey,

[104160]M.; Durie, P. R.; Rommens, J. M.: Segregation analysis in
Shwachman–Diamond

[104161]syndrome: evidence for recessive inheritance. Am. J. Hum.
Genet. 66:

[104162]1413–1416, 2000.

[104163]18216.

[104164]Ginzberg, H.; Shin, J.; Ellis, L.; Morrison, J.; Ip, W.; Dror,

[104165]Y.; Freedman, M.; Heitlinger, L. A.; Belt, M. A.; Corey, M.;
Rommens,

[104166]J. M.; Durie, P. R.: Shwachman syndrome: phenotypic
manifestations

[104167]of sibling sets and isolated cases in a large patient cohort
are similar. J.

[104168]Pediat. 135: 81–88, 1999.

[104169]18217.

[104170]Goldstein, R.: Congenital lipomatosis of the pancreas.
Malabsorption,

[104171]dwarfism, leukopenia with relative granulocytopenia and thrombocytopenia. Clin.

[104172]Pediat. 7: 419–422, 1968.

[104173]18218.

[104174]Goobie, S.; Morrison, J.; Ginzberg, H.; Ellis, L.; Corey, M.;

[104175]Masuno, M.; Imaizumi, K.; Kuroki, Y.; Fujiwara, T. M.; Morgan, K.;

[104176]Durie, P. R.; Rommens, J. M.: Exclusion of linkage of Shwachman–Diamond

[104177]syndrome to chromosome regions 6q and 12q implicated by a de novo

[104178]translocation. Am. J. Med. Genet. 85: 171–174, 1999.

[104179]18219.

[104180]Li, C. C.; Mantel, N.: A simple method of estimating the segregation

[104181]ratio under complete ascertainment. Am. J. Hum. Genet. 20: 61–81,

[104182]1968.

[104183]18220.

[104184]Mack, D. R.; Forstner, G. G.; Wilschanski, M.; Freedman, M. H.;

[104185]Durie, P. R.: Shwachman syndrome: exocrine pancreatic dysfunction

[104186]and variable phenotypic expression. Gastroenterology 111: 1593–1602,

[104187]1996.

[104188]18221.

[104189]Masuno, M.; Imaizumi, K.; Nishimura, G.; Nakamura, M.; Saito,

[104190]I.; Akagi, K.; Kuroki, Y.: Shwachman syndrome associated with de

[104191]novo reciprocal translocation t(6;12)(q16.2;q21.2). J. Med. Genet. 32:

[104192]894–895, 1995.

[104193]18222.

[104194]McLennan, T. W.; Steinbach, H. L.: Shwachman's syndrome: the

[104195]broad spectrum of bony abnormalities. Radiology 112: 167–173, 1974.

[104196]18223.

[104197]Goobie, S.; Popovic, M.; Morrison, J.; Ellis, L.; Ginzberg, H.;

[104198]Boocock, G. R. B.; Ehtesham, N.; Betard, C.; Brewer, C. G.;

Roslin,

[104199]N. M.; Hudson, T. J.; Morgan, K.; Fujiwara, T. M.; Durie, P.
R.; Rommens,

[104200]J. M.: Shwachman–Diamond syndrome with exocrine pan-
creatic dysfunction

[104201]and bone marrow failure maps to the centromeric region
of chromosome

[104202]7. Am. J. Hum. Genet. 68: 1048–1054, 2001.

[104203]18224.

[104204]Nezelof, C.; Watchi, M.: L'hypoplasie congenitale lipoma-
teuse

[104205]du pancreas exocrine chez l'enfant (Deux observations et
revue de

[104206]la litterature). Arch. Franc. Pediat. 18: 1135–1172, 1961.

[104207]18225.

[104208]Pringle, E. M.; Young, W. F.; Haworth, E. M.: Syndrome of
pancreatic

[104209]insufficiency, blood dyscrasia and metaphyseal dysplasia.
Proc. Roy.

[104210]Soc. Med. 61: 776–777, 1968.

[104211]18226.

[104212]Rothbaum, R. J.; Williams, D. A.; Daugherty, C. C.: Unusual surface

[104213]distribution of concanavalin A reflects a cytoskeletal defect in neutrophils

[104214]in Shwachman's syndrome. Lancet II: 800–801, 1982.

[104215]18227.

[104216]Saint-Martin, J.; Fournet, J. P.; Charlas, J.; Schaison, G.; Nodot,

[104217]A.; Meyer, B.; Vialatte, J.: Insuffisance pancreatique externe avec

[104218]granulopenie chronique. Arch. Franc. Pediat. 26: 861–871, 1969.

[104219]18228.

[104220]Saunders, E. F.; Gall, G.; Freedman, M. H.: Granulopoiesis in

[104221]Shwachman's syndrome (pancreatic insufficiency and bone marrow dysfunction). Pediatrics 64:

[104222]515–519, 1979.

[104223]18229.

[104224]Cazeneuve, C.; Sarkisian, T.; Pecheux, C.; Dervichian, M.; Nedelec,

[104225]B.; Reinert, P.; Ayvazyan, A.; Kouyoumdjian, J.-C.;

Ajrapetyan, H.;

[104226]Delpech, M.; Goossens, M.; Dode, C.; Grateau, G.; Amse-
lem, S.: MEFV-gene

[104227]analysis in Armenian patients with familial Mediterranean
fever: diagnostic

[104228]value and unfavorable renal prognosis of the M694V ho-
mozygous genotype--genetic

[104229]and therapeutic implications. Am. J. Hum. Genet. 65:
88-97, 1999.

[104230]18230.

[104231]Centola, M.; Wood, G.; Frucht, D. M.; Galon, J.; Aringer, M.;

[104232]Farrell, C.; Kingma, D. W.; Horwitz, M. E.; Mansfield, E.;

Holland,

[104233]S. M.; O'Shea, J. J.; Rosenberg, H. F.; Malech, H. L.; Kast-
ner, D.

[104234]L.: The gene for familial Mediterranean fever, MEFV, is ex-
pressed

[104235]in early leukocyte development and is regulated in re-
sponse to inflammatory

[104236]mediators. Blood 95: 3223-3231, 2000.

[104237]18231.

[104238]Carpenter, S.; Karpati, G.; Andermann, F.; Jakob, J. C.; Andermann,

[104239]E.: The ultrastructural characteristics of the abnormal cytosomes

[104240]in Batten-Kuf's disease. Brain 100: 137-156, 1977.

[104241]18232.

[104242]Klockars, T.; Savukoski, M.; Isosomppi, J.; Laan, M.; Jarvela,

[104243]I.; Petrukhin, K.; Palotie, A.; Peltonen, L.: Efficient construction

[104244]of a physical map by fiber-fish of the CLN5 region: refined assignment

[104245]and long-range contig covering the critical region on 13q22. Genomics 35:

[104246]71-78, 1996.

[104247]18233.

[104248]Santavuori, P.; Rapola, J.; Nuutila, A.; Raininko, R.; Lappi, M.;

[104249]Launes, J.; Herva, R.; Sainio, K.: The spectrum of Jansky-Bielschowsky

[104250]disease. Neuropediatrics 22: 92-96, 1991.

[104251]18234.

[104252]Santavuori, P.; Rapola, J.; Sainio, K.; Raitta, C.: A variant
[104253]of Jansky-Bielschowsky disease. Neuropediatrics 13:
135–141, 1982.

[104254]18235.

[104255]Savukoski, M.; Kestila, M.; Williams, R.; Jarvela, I.; Sharp, J.;
[104256]Harris, J.; Santavuori, P.; Gardiner, M.; Peltonen, L.: De-
fined chromosomal

[104257]assignment of CLN5 demonstrates that at least four loci
are involved

[104258]in the pathogenesis of human ceroid lipofuscinoses. Am.
J. Hum. Genet. 55:

[104259]695–701, 1994.

[104260]18236.

[104261]Savukoski, M.; Klockars, T.; Holmberg, V.; Santavuori, P.;
Lander,

[104262]E. S.; Peltonen, L.: CLN5, a novel gene encoding a putative
transmembrane

[104263]protein mutated in Finnish variant late infantile neuronal
ceroid

[104264]lipofuscinosis. Nature Genet. 19: 286–288, 1998.

[104265]18237.

[104266]Varilo, T.; Savukoski, M.; Norio, R.; Santavuori, P.; Peltonen,

[104267]L.; Jarvela, I.: The age of human mutation: genealogical and linkage

[104268]disequilibrium analysis of the CLN5 mutation in the Finnish population. Am.

[104269]J. Hum. Genet. 58: 506–512, 1996.

[104270]18238.

[104271]Williams, R.; Santavuori, P.; Peltonen, L.; Gardiner, R. M.; Jarvela,

[104272]I.: A variant form of late infantile neuronal ceroid lipofuscinosis

[104273](CLN5) is not an allelic form of Batten (Spielmeyer–Vogt–Sjogren,

[104274]CLN3) disease: exclusion of linkage to the CLN3 region of chromosome

[104275]16. Genomics 20: 289–290, 1994.

[104276]18239.

[104277]Busard, B. L. S. M.; Renier, W. O.; Gabreels, F. J. M.; Jaspar,

[104278]H. H. J.; van Haelst, U. J. G.; Slooff, J. L.: Lafora's disease:

[104279]comparison of inclusion bodies in skin and in brain. Arch.
Neurol. 43:

[104280]296-299, 1986.

[104281]18240.

[104282]Busard, H. L. S. M.; Gabreels-Festen, A. A. W. M.; Renier,
W. O.;

[104283]Gabreels, F. J. M.; Stadhouders, A. M.: Axilla skin biopsy: a
reliable

[104284]test for the diagnosis of Lafora's disease. Ann. Neurol. 21:
599-601,

[104285]1987.

[104286]18241.

[104287]Fluharty, A. L.; Porter, M. T.; Hirsh, G. A.; Pevida, E.; Ki-
hara,

[104288]H.: Metachromasia in fibroblasts from a patient with
Lafora's disease.

[104289](Letter) Lancet II: 109-110, 1970.

[104290]18242.

[104291]Ganesh, S.; Agarwala, K. L.; Ueda, K.; Akagi, T.; Shoda, K.;
Usui,

[104292]T.; Hashikawa, T.; Osada, H.; Delgado-Escueta, A. V.; Ya-

makawa, K.

[104293] Laforin, defective in the progressive myoclonus epilepsy of Lafora

[104294] type, is a dual-specificity phosphatase associated with polyribosomes. Hum.

[104295] Molec. Genet. 9: 2251–2261, 2000.

[104296] 18243.

[104297] Gomez–Garre, P.; Sanz, Y.; Rodriguez de Cordoba, S.; Ser-ratosa,

[104298] J. M.: Mutational spectrum of the EPM2A gene in progressive myoclonus

[104299] epilepsy of Lafora: high degree of allelic heterogeneity and prevalence

[104300] of deletions. Europ. J. Hum. Genet. 8: 946–954, 2000.

[104301] 18244.

[104302] Harriman, D. G. F.; Millar, J. H. D.: Progressive familial myoclonic

[104303] epilepsy in 3 families: its clinical features and pathological basis. Brain 78:

[104304] 325–349, 1955.

[104305] 18245.

- [104306]Janeway, R.; Ravens, J. R.; Pearce, L. A.; Odor, D. L.; Suzuki,
Suzuki,
[104307]K.: Progressive myoclonus epilepsy with Lafora inclusion
bodies.
[104308]I. Clinical, genetic, histopathologic and biochemical as-
pects. Arch.
[104309]Neurol. 16: 565–582, 1967.
[104310]18246.
[104311]Lehesjoki, A.–E.; Koskiniemi, M.; Pandolfo, M.; Antonelli,
A.;
[104312]Kyllerman, M.; Wahlstrom, J.; Nergardh, A.; Burmeister, M.;
Sistonen,
[104313]P.; Norio, R.; de la Chapelle, A.: Linkage studies in pro-
gressive
[104314]myoclonus epilepsy: Unverricht–Lundborg and Lafora’s
diseases. Neurology 42:
[104315]1545–1550, 1992.
[104316]18247.
[104317]Maddox, L. O.; Descartes, M.; Collins, J.; Keating, J.;
Rosenfeld,
[104318]S.; Palmer, C.; Carroll, A. J.; Kuzniecky, R.: Identification of

[104319]a recombination event narrowing the Lafora disease gene region. J.

[104320]Med. Genet. 34: 590–591, 1997.

[104321]18248.

[104322]Harding, B. N.; Leonard, J. V.; Erdohazi, M.: Ornithine carbamoyl

[104323]transferase deficiency: a neuropathological study. Europ. J. Pediat. 141:

[104324]215–220, 1984.

[104325]18249.

[104326]Amadou, C.; Ribouchon, M. T.; Mattei, M. G.; Jenkins, N. A.; Gilbert,

[104327]D. J.; Copeland, N. G.; Avoustin, P.; Pontarotti, P.: Localization

[104328]of new genes and markers to the distal part of the human major histocompatibility

[104329]complex (MHC) region and comparison with the mouse: new insights into

[104330]the evolution of mammalian genomes. Genomics 26: 9–20, 1995.

[104331]18250.

- [104332]Solinas-Toldo, S.; Lengauer, C.; Fries, R.: Comparative genome map of human and cattle. Genomics 27: 489–496, 1995.
- [104333]18251.
- [104334]Olives, B.; Martial, S.; Mattei, M.–G.; Matassi, G.; Rousset, G.; Ripoche, P.; Cartron, J.–P.; Bailly, P.: Molecular characterization of a new urea transporter in the human kidney. FEBS Lett. 386: 156–160, 1996.
- [104335]18252.
- [104336]Ranade, K.; Wu, K.–W.; Hwu, C.–M.; Ting, C.–T.; Pei, D.; Pesich, R.; Hebert, J.; Chen, Y.–D. I.; Pratt, R.; Olshen, R.; Masaki, K.; Risch, N.; Cox, D. R.; Botstein, D.: Genetic variation in the human urea transporter–2 is associated with variation in blood pressure. Hum. Molec. Genet. 10: 2157–2164, 2001.

[104345]18253.

[104346]Galko, M. J.; Tessier-Lavigne, M.: Function of an axonal chemoattractant

[104347]modulated by metalloprotease activity. Science 289: 1365–1367, 2000.

[104348]18254.

[104349]Stein, E.; Tessier-Lavigne, M.: Hierarchical organization of guidance

[104350]receptors: silencing of netrin attraction by Slit through a Robo/DCC

[104351]receptor complex. Science 291: 1928–1938, 2001.

[104352]18255.

[104353]Stein, E.; Zou, Y.; Poo, M.; Tessier-Lavigne, M.: Binding of

[104354]DCC by netrin-1 to mediate axon guidance independent of adenosine

[104355]A2B receptor activation. Science 291: 1976–1982, 2001.

[104356]18256.

[104357]Orr, H. T.; Zoghbi, H. Y.: SCA1 molecular genetics: a history

[104358]of a 13 year collaboration against glutamines. Hum. Molec. Genet. 10:

[104359]2307–2311, 2001.

[104360]18257.

[104361]Lubahn, D. B.; Moyer, J. S.; Golding, T. S.; Couse, J. F.; Ko-
rach,

[104362]K. S.; Smithies, O.: Alteration of reproductive function but
not

[104363]prenatal sexual development after insertional disruption
of the mouse

[104364]estrogen receptor gene. Proc. Nat. Acad. Sci. 90:
11162–11166, 1993.

[104365]18258.

[104366]Hanna, I. H.; Dawling, S.; Roodi, N.; Guengerich, F. P.; Parl,

[104367]F. F.: Cytochrome P450 1B1 (CYP1B1) pharmacogenetics:
association

[104368]of polymorphisms with functional differences in estrogen
hydroxylation

[104369]activity. Cancer Res. 60: 3440–3444, 2000.

[104370]18259.

[104371]Dai, G.; Levy, O.; Carrasco, N.: Cloning and characteriza-
tion

[104372]of the thyroid iodide transporter. Nature 379: 458–460,

1996.

[104373]18260.

[104374]Costanzo, A.; Merlo, P.; Pediconi, N.; Fulco, M.; Sartorelli, V.;

[104375]Cole, P. A.; Fontemaggi, G.; Fanciulli, M.; Schiltz, L.; Blandino,

[104376]G.; Balsano, C.; Levrero, M.: DNA damage-dependent acetylation of

[104377]p73 dictates the selective activation of apoptotic target genes. Molec.

[104378]Cell 9: 175–186, 2002.

[104379]18261.

[104380]Flores, E. R.; Tsai, K. Y.; Crowley, D.; Sengupta, S.; Yang, A.;

[104381]McKeon, F.; Jacks, T.: p63 and p73 are required for p53-dependent

[104382]apoptosis in response to DNA damage. Nature 416: 560–564, 2002.

[104383]18262.

[104384]Jost, C. A.; Marin, M. C.; Kaelin, W. G., Jr.: p73 is a human

[104385]p53-related protein that can induce apoptosis. Nature

389: 191–194,

[104386]1997.

[104387]18263.

[104388]Kaghad, M.; Bonnet, H.; Yang, A.; Creancier, L.; Biscan, J.-C.;

[104389]Valent, A.; Minty, A.; Chalon, P.; Lelias, J.-M.; Dumont, X.; Ferrara,

[104390]P.; McKeon, F.; Caput, D.: Monoallelically expressed gene related

[104391]to p53 at 1p36, a region frequently deleted in neuroblastoma and other

[104392]human cancers. Cell 90: 809–819, 1997.

[104393]18264.

[104394]Marin, M. C.; Jost, C. A.; Brooks, L. A.; Irwin, M. S.; O'Nions,

[104395]J.; Tidy, J. A.; James, N.; McGregor, J. M.; Harwood, C. A.; Yulug,

[104396]I. G.; Vousden, K. H.; Allday, M. J.; Gusterson, B.; Ikawa, S.; Hinds,

[104397]P. W.; Crook, T.; Kaelin, W. G., Jr. A common polymorphism acts

[104398]as an intragenic modifier of mutant p53 behaviour. Nature

Genet. 25:

[104399]47–54, 2000.

[104400]18265.

[104401]Morrison, R. S.; Kinoshita, Y.: p73--guilt by association?

Science 289:

[104402]257–258, 2000.

[104403]18266.

[104404]Pozniak, C. D.; Radinovic, S.; Yang, A.; McKeon, F.; Kaplan,
D.

[104405]R.; Miller, F. D.: An anti-apoptotic role for the p53 family
member,

[104406]p73, during developmental neuron death. Science 289:
304–306, 2000.

[104407]18267.

[104408]Yang, A.; Walker, N.; Bronson, R.; Kaghad, M.; Ooster-
wegel, M.;

[104409]Bonnin, J.; Vagner, C.; Bonnet, H.; Dikkes, P.; Sharpe, A.;
McKeon,

[104410]F.; Caput, D.: p73-deficient mice have neurological,
pheromonal and

[104411]inflammatory defects but lack spontaneous tumours. Na-

ture 404: 99–103,

[104412]2000.

[104413]18268.

[104414]Moore, J. T.; McKee, D. D.; Slentz–Kesler, K.; Moore, L. B.;
Jones,

[104415]S. A.; Horne, E. L.; Su, J.–L.; Kliewer, S. A.; Lehmann, J. M.;
Willson,

[104416]T. M.: Cloning and characterization of human estrogen re-
ceptor beta

[104417]isoforms. *Biochem. Biophys. Res. Commun.* 247: 75–78,
1998.

[104418]18269.

[104419]Mosselman, S.; Polman, J.; Dijkema, R.: ER–beta: identifi-
cation

[104420]and characterization of a novel human estrogen receptor.
FEBS Lett. 392:

[104421]49–53, 1996.

[104422]18270.

[104423]Ogawa, S.; Emi, M.; Shiraki, M.; Hosoi, T.; Ouchi, Y.; Inoue,

[104424]S.: Association of estrogen receptor beta (ESR2) gene
polymorphism

[104425]with blood pressure. J. Hum. Genet. 45: 327–330, 2000.

[104426]18271.

[104427]Ogawa, S.; Inoue, S.; Watanabe, T.; Orimo, A.; Hosoi, T.;
Ouchi,

[104428]Y.; Muramatsu, M.: Molecular cloning and characterization
of human

[104429]estrogen receptor beta-cx: a potential inhibitor of estrogen
action

[104430]in human. Nucleic Acids Res. 26: 3505–3512, 1998.

[104431]18272.

[104432]Osterlund, M. K.; Gustafsson, J. A.; Keller, E.; Hurd, Y. L.:

[104433]Estrogen receptor beta (ER-beta) messenger ribonucleic
acid (mRNA)

[104434]expression within the human forebrain: distinct distribution
pattern

[104435]to ER-alpha mRNA. J. Clin. Endocr. Metab. 85: 3840–3846,
2000.

[104436]18273.

[104437]Pasquali, D.; Rossi, V.; Esposito, D.; Abbondanza, C.; Puca,
G.

[104438]A.; Bellastella, A.; Sinisi, A. A.: Loss of estrogen receptor

beta

[104439]expression in malignant human prostate cells in primary cultures and

[104440]in prostate cancer tissues. J. Clin. Endocr. Metab. 86: 2051–2055,

[104441]2001.

[104442]18274.

[104443]Petersen, D. N.; Tkalcevic, G. T.; Koza–Taylor, P. H.; Turi, T.

[104444]G.; Brown, T. A.: Identification of estrogen receptor beta–2, a functional

[104445]variant of estrogen receptor beta expressed in normal rat tissues. Endocrinology 139:

[104446]1082–1092, 1998.

[104447]18275.

[104448]Rissman, E. F.; Heck, A. L.; Leonard, J. E.; Shupnik, M. A.; Gustafsson,

[104449]J.–A.: Disruption of estrogen receptor beta gene impairs spatial

[104450]learning in female mice. Proc. Nat. Acad. Sci. 99: 3996–4001, 2002.

[104451]18276.

[104452]Rosenkranz, K.; Hinney, A.; Ziegler, A.; Hermann, H.;
Fichter,
[104453]M.; Mayer, H.; Siegfried, W.; Young, J. K.; Remschmidt, H.;
Hebebrand,
[104454]J.: Systematic mutation screening of the estrogen receptor
beta gene
[104455]in probands of different weight extremes: identification of
several
[104456]genetic variants. J. Clin. Endocr. Metab. 83: 4524-4527,
1998.
[104457]18277.
[104458]Speirs, V.; Adams, I. P.; Walton, D. S.; Atkin, S. L.: Identifi-
cation
[104459]of wild-type and exon 5 deletion variants of estrogen re-
ceptor beta
[104460]in normal human mammary gland. J. Clin. Endocr. Metab.
85: 1601-1605,
[104461]2000.
[104462]18278.
[104463]Tsukamoto, K.; Inoue, S.; Hosoi, T.; Orimo, H.; Emi, M.:
Isolation
[104464]and radiation hybrid mapping of dinucleotide repeat poly-

morphism at

[104465]the human estrogen receptor beta locus. J. Hum. Genet.

43: 73–74,

[104466]1998.

[104467]18279.

[104468]Weihua, Z.; Saji, S.; Makinen, S.; Cheng, G.; Jensen, E. V.;

Warner,

[104469]M.; Gustafsson, J.–A.: Estrogen receptor (ER)–beta, a mod–

ulator of

[104470]ER–alpha in the uterus. Proc. Nat. Acad. Sci. 97:

5936–5941, 2000.

[104471]18280.

[104472]Westberg, L.; Baghaei, F.; Rosmond, R.; Hellstrand, M.;

Landen,

[104473]M.; Jansson, M.; Holm, G.; Bjorntorp, P.; Eriksson, E.: Poly–

morphisms

[104474]of the androgen receptor gene and the estrogen receptor

gene are associated

[104475]with androgen levels in women. J. Clin. Endocr. Metab. 86:

2562–2568,

[104476]2001.

[104477]18281.

[104478]Zhu, Y.; Bian, Z.; Lu, P.; Karas, R. H.; Bao, L.; Cox, D.; Hodgins,

[104479]J.; Shaul, P. W.; Thoren, P.; Smithies, O.; Gustafsson, J.-A.; Mendelsohn,

[104480]M. E.: Abnormal vascular function and hypertension in mice deficient

[104481]in estrogen receptor beta. *Science* 295: 505–508, 2002.

[104482]18282.

[104483]DesGroseillers, L.; Lemieux, N.: Localization of a human double-stranded

[104484]RNA-binding protein gene (STAU) to band 20q13.1 by fluorescence in

[104485]situ hybridization. *Genomics* 36: 527–529, 1996.

[104486]18283.

[104487]Marion, R. M.; Fortes, P.; Beloso, A.; Dotti, C.; Ortin, J.: A

[104488]human sequence homologue of staufen is an RNA-binding protein that

[104489]is associated with polysomes and localizes to the rough endoplasmic

[104490]reticulum. *Molec. Cell. Biol.* 19: 2212–2219, 1999.

[104491]18284.

[104492]Wickham, L.; Duchaine, T.; Luo, M.; Nabi, I. R.; DesGro-
seillers,

[104493]L.: Mammalian stauferin is a double-stranded-RNA- and
tubulin-binding

[104494]protein which localizes to the rough endoplasmic reticu-
lum. Molec.

[104495]Cell. Biol. 19: 2220-2230, 1999.

[104496]18285.

[104497]Lee, J. E.; Hollenberg, S. M.; Snider, L.; Turner, D. L.; Lip-
nick,

[104498]N.; Weintraub, H.: Conversion of *Xenopus* ectoderm into
neurons by

[104499]NeuroD, a basic helix-loop-helix protein. Science 268:
836-844,

[104500]1995.

[104501]18286.

[104502]Plasilova, M.; Stoilov, I.; Sarfarazi, M.; Kadasi, L.; Ferakova,

[104503]E.; Ferak, V.: Identification of a single ancestral CYP1B1
mutation

[104504]in Slovak Gypsies (Roms) affected with primary congenital
glaucoma. J.

[104505]Med. Genet. 36: 290–294, 1999.

[104506]18287.

[104507]Schwartzman, M. L.; Balazy, M.; Masferrer, J.; Abraham, N.
G.;

[104508]McGiff, J. C.; Murphy, R. C.: 12(R)–hydroxyicosatetraenoic
acid:

[104509]a cytochrome P450–dependent arachidonate metabolite
that inhibits

[104510]Na⁺,K⁺–ATPase in the cornea. Proc. Nat. Acad. Sci. 84:
8125–8129,

[104511]1987.

[104512]18288.

[104513]Stoilov, I.; Akarsu, A. N.; Alozie, I.; Child, A.; Barsoum–
Homsy,

[104514]M.; Turacli, M. E.; Or, M.; Lewis, R. A.; Ozdemir, N.; Brice,
G.;

[104515]Aktan, S. G.; Chevrette, L.; Coca–Prados, M.; Sarfarazi, M.:
Sequence

[104516]analysis and homology modeling suggest that primary
congenital glaucoma

[104517]on 2p21 results from mutations disrupting either the
hinge region

- [104518] or the conserved core structures of cytochrome P4501B1.
Am. J. Hum.
- [104519] Genet. 62: 573–584, 1998.
- [104520] 18289.
- [104521] Stoilov, I.; Akarsu, A. N.; Sarfarazi, M.: Identification of
three
[104522] different truncating mutations in cytochrome P4501B1
(CYP1B1) as the
[104523] principal cause of primary congenital glaucoma
(buphthalmos) in families
[104524] linked to the GLC3A locus on chromosome 2p21. Hum.
Molec. Genet. 6:
[104525] 641–647, 1997.
- [104526] 18290.
- [104527] Sutter, T. R.; Tang, Y. M.; Hayes, C. L.; Wo, Y.-Y. P.; Jabs,
E.
- [104528] W.; Li, X.; Yin, H.; Cody, C. W.; Greenlee, W. F.: Complete
cDNA
[104529] sequence of a human dioxin-inducible mRNA identifies a
new gene subfamily
[104530] of cytochrome P450 that maps to chromosome 2. J. Biol.
Chem. 269:

[104531]13092–13099, 1994.

[104532]18291.

[104533]Tang, Y. M.; Wo, Y.-Y. P.; Stewart, J.; Hawkins, A. L.; Griffin,
fin,

[104534]C. A.; Sutter, T. R.; Greenlee, W. F.: Isolation and characterization
terization

[104535]of the human cytochrome P450 CYP1B1 gene. J. Biol.
Chem. 271: 28324–28330,

[104536]1996.

[104537]18292.

[104538]Vincent, A.; Billingsley, G.; Priston, M.; Williams–Lyn, D.;
Sutherland,

[104539]J.; Glaser, T.; Oliver, E.; Walter, M. A.; Heathcote, G.; Levin,
A.;

[104540]Heon, E.: Phenotypic heterogeneity of CYP1B1: mutations
in a patient

[104541]with Peters' anomaly. J. Med. Genet. 38: 324–326, 2001.

[104542]18293.

[104543]Chen, H. T.; Bhandoola, A.; Difilippantonio, M. J.; Zhu, J.;
Brown,

[104544]M. J.; Tai, X.; Rogakou, E. P.; Brotz, T. M.; Bonner, W. M.;

Ried,

[104545]T.; Nussenzweig, A.: Response to RAG-mediated V(D)J cleavage by NBS1

[104546]and gamma-H2AX. Science 290: 1962–1964, 2000.

[104547]18294.

[104548]Petersen, S.; Casellas, R.; Reina-San-Martin, B.; Chen, H. T.;

[104549]Difilippantonio, M. J.; Wilson, P. C.; Hanitsch, L.; Celeste, A.;

[104550]Muramatsu, M.; Pilch, D. R.; Redon, C.; Ried, T.; Bonner, W. M.; Honjo,

[104551]T.; Nussenzweig, M. C.; Nussenzweig, A.: AID is required to initiate

[104552]Nbs1/gamma-H2AX focus formation and mutations at sites of class switching. Nature 414:

[104553]660–665, 2001.

[104554]18295.

[104555]Lan, M. S.; Lu, J.; Goto, Y.; Notkins, A. L.: Molecular cloning

[104556]and identification of a receptor-type protein tyrosine phosphatase,

[104557]IA-2, from human insulinoma. DNA and Cell Biol. 13:

505-514, 1994.

[104558]18296.

[104559]Leiter, E. H.; Tsumura, H.; Serreze, D. V.; Chapman, H. D.;
Rabin,

[104560]D. U.; Lan, M. S.; Notkins, A. L.: Mapping to chromosomes
1 and 12

[104561]of mouse homologs of human protein tyrosine phos-
phatase, receptor-type,

[104562]related genes encoding pancreatic beta cell autoantigens.
Mammalian

[104563]Genome 8: 949-950, 1997.

[104564]18297.

[104565]Rabin, D. U.; Pleasic, S. M.; Shapiro, J. A.; Yoo-Warren, H.;
Oles,

[104566]J.; Hicks, J. M.; Goldstein, D. E.; Rae, P. M.: Islet cell anti-
gen

[104567]512 is a diabetes-specific islet autoantigen related to pro-
tein tyrosine

[104568]phosphatases. J. Immun. 152: 3183-3188, 1994.

[104569]18298.

[104570]Xie, J.; Zhang, B.; Lan, M. S.; Notkins, A. L.: Genomic

structure

[104571]and promoter sequence of the insulin-dependent diabetes mellitus autoantigen,

[104572]IA-2 (PTPRN). Genomics 54: 338-343, 1998.

[104573]18299.

[104574]Demetriou, M.; Granovsky, M.; Quaggin, S.; Dennis, J. W.:

Negative

[104575]regulation of T-cell activation and autoimmunity by Mgat5 N-glycosylation. Nature 409:

[104576]733-739, 2001.

[104577]18300.

[104578]van den Maagdenberg, A. M. J. M.; Weghuis, D. O.; Rijss,

J.; van

[104579]de Wetering, R. A. C.; Wieringa, B.; van Kessel, A. G.; Hendriks,

[104580]W. J. A. J.: Assignment of the human gene for receptor-type protein

[104581]tyrosine phosphatase IA-2 (PTPRN) to chromosome region 2q35-q36.1

[104582]and identification of an intragenic genetic marker. Cytogenet. Cell

[104583]Genet. 73: 145-148, 1996.

[104584]18301.

[104585]Granovsky, M.; Fata, J.; Pawling, J.; Muller, W. J.; Khokha, R.;

[104586]Dennis, J. W.: Suppression of tumor growth and metastasis in Mgat5-deficient

[104587]mice. Nature Med. 6: 306-12, 2000.

[104588]18302.

[104589]Saito, H.; Nishikawa, A.; Gu, J.; Ihara, Y.; Soejima, H.; Wada,

[104590]Y.; Sekiya, C.; Niikawa, N.; Taniguchi, N.: cDNA cloning and chromosomal

[104591]mapping of human N-acetylglucosaminyltransferase V+. Biochem. Biophys.

[104592]Res. Commun. 198: 318-327, 1994.

[104593]18303.

[104594]Morasso, M. I.; Yonescu, R.; Griffin, C. A.; Sargent, T. D.: Localization

[104595]of human DLX8 to chromosome 17q21.3-q22 by fluorescence in situ hybridization. Mammalian

[104596]Genome 8: 302-303, 1997.

[104597]18304.

[104598]Quinn, L. M.; Johnson, B. V.; Nicholl, J.; Sutherland, G. R.;
Kalionis,
[104599]B.: Isolation and identification of homeobox genes from
the human
[104600]placenta including a novel member of the Distal-less fam-
ily, DLX4. Gene 187:
[104601]55-61, 1997.
[104602]18305.
[104603]Boddy, M. N.; Howe, K.; Etkin, L. D.; Solomon, E.;
Freemont, P.
[104604]S.: PIC 1, a novel ubiquitin-like protein which interacts
with the
[104605]PML component of a multiprotein complex that is dis-
rupted in acute
[104606]promyelocytic leukaemia. Oncogene 13: 971-982, 1996.
[104607]18306.
[104608]Desterro, J. M. P.; Rodriguez, M. S.; Hay, R. T.: SUMO-1
modification
[104609]of I-kappa-B-alpha inhibits NF-kappa-B activation. Molec.
Cell 2:
[104610]233-239, 1998.

[104611]18307.

[104612]Lapenta, V.; Chiurazzi, P.; van der Spek, P.; Pizzuti, A.;
Hanaoka,

[104613]F.; Brahe, C.: SMT3A, a human homologue of the *S. cere-*
visiae SMT3

[104614]gene, maps to chromosome 21qter and defines a novel
gene family. *Genomics* 40:

[104615]362–366, 1997.

[104616]18308.

[104617]Mao, Y.; Sun, M.; Desai, S. D.; Liu, L. F.: SUMO–1 conjuga-
tion

[104618]to topoisomerase I: a possible repair response to topoiso-
merase-mediated

[104619]DNA damage. *Proc. Nat. Acad. Sci.* 97: 4046–4051, 2000.

[104620]18309.

[104621]Okura, T.; Gong, L.; Kamitani, T.; Wada, T.; Okura, I.; Wei,
C.-F.;

[104622]Chang, H.-M.; Yeh, E. T. H.: Protection against Fas/
APO–1– and tumor

[104623]necrosis factor-mediated cell death by a novel protein,
sentrin. *J.*

[104624]Immun. 157: 4277–4281, 1996.

[104625]18310.

[104626]Shen, Z.; Pardington–Purtymun, P. E.; Comeaux, J. C.;
Moyzis, R.

[104627]K.; Chen, D. J.: UBL1, a human ubiquitin–like protein asso-
ciating

[104628]with human RAD51/RAD52 proteins. Genomics 36:
271–279, 1996.

[104629]18311.

[104630]Bengtsson, E.; Neame, P. J.; Heinegard, D.; Sommarin, Y.:
The

[104631]primary structure of a basic leucine–rich repeat protein,
PRELP, found

[104632]in connective tissues. J. Biol. Chem. 270: 25639–25644,
1995.

[104633]18312.

[104634]Grover, J.; Chen, X.–N.; Korenberg, J. R.; Recklies, A. D.;
Roughley,

[104635]P. J.: The gene organization, chromosome location, and
expression

[104636]of a 55–kDa matrix protein (PRELP) of human articular
cartilage. Genomics 38:

[104637]109–117, 1996.

[104638]18313.

[104639]Bigg, H. F.; Shi, Y. E.; Liu, Y. E.; Steffensen, B.; Overall, C.

[104640]M.: Specific, high affinity binding of tissue inhibitor of metalloproteinases–4

[104641](TIMP–4) to the COOH–terminal hemopexin–like domain of human gelatinase

[104642]A. J. Biol. Chem. 272: 15496–15500, 1997.

[104643]18314.

[104644]Greene, J.; Wang, M.; Liu, Y. E.; Raymond, L. A.; Rosen, C.; Shi,

[104645]Y. E.: Molecular cloning and characterization of human tissue inhibitor

[104646]of metalloproteinase 4. J. Biol. Chem. 271: 30375–30380, 1996.

[104647]18315.

[104648]Leco, K. J.; Apte, S. S.; Taniguchi, G. T.; Hawkes, S. P.; Khokha,

[104649]R.; Schultz, G. A.; Edwards, D. R.: Murine tissue inhibitor of metalloproteinases–4

[104650](Timp–4): cDNA isolation and expression in adult mouse

tissues. FEBS

[104651]Lett. 401: 213–217, 1997.

[104652]18316.

[104653]Olson, T. M; Hirohata, S.; Ye, J.; Leco, K.; Seldin, M. F.;
Apte,

[104654]S. S.: Cloning of the human tissue inhibitor of metallopro-
teinase-4

[104655]gene (TIMP4) and localization of the TIMP4 and Timp4
gene to human

[104656]chromosome 3p25 and mouse chromosome 6, respec-
tively. Genomics 51:

[104657]148–151, 1998.

[104658]18317.

[104659]Wang, M.; Liu, Y. E.; Greene, J.; Sheng, S.; Fuchs, A.;
Rosen,

[104660]E. M.; Shi, Y. E.: Inhibition of tumor growth and metastasis
of human

[104661]breast cancer cells transfected with tissue inhibitor of
metalloproteinase

[104662]4. Oncogene 14: 2767–2774, 1997.

[104663]18318.

- [104664]Hsu, L. C.; Chang, W.-C.: Sequencing and expression of the human
- [104665]ALDH8 encoding a new member of the aldehyde dehydrogenase family. Gene 174:
- [104666]319–322, 1996.
- [104667]18319.
- [104668]Hsu, L. C.; Chang, W.-C.; Lin, S. W.; Yoshida, A.: Cloning and
- [104669]characterization of genes encoding four additional human aldehyde
- [104670]dehydrogenase isozymes. Adv. Exp. Med. Biol. 372: 159–168, 1995.
- [104671]18320.
- [104672]Fujiwara, H.; Tatsumi, K.-I.; Miki, K.; Harada, T.; Okada, S.;
- [104673]Nose, O.; Kodama, S.; Amino, N.: Recurrent T354P mutation of the
- [104674]Na⁺/I⁻ symporter in patients with iodide transport defect. J. Clin.
- [104675]Endocr. Metab. 83: 2940–2943, 1998.
- [104676]18321.
- [104677]Kosugi, S.; Bhayana, S.; Dean, H. J.: A novel mutation in

the

[104678]sodium/iodide symporter gene in the largest family with
iodide transport

[104679]defect. J. Clin. Endocr. Metab. 84: 3248–3253, 1999.

[104680]18322.

[104681]Kosugi, S.; Inoue, S.; Matsuda, A.; Jhiang, S. M.: Novel,
missense

[104682]and loss-of-function mutations in the sodium/iodide
symporter gene

[104683]causing iodide transport defect in three Japanese patients.
J. Clin.

[104684]Endocr. Metab. 83: 3373–3376, 1998.

[104685]18323.

[104686]Kosugi, S.; Sato, Y.; Matsuda, A.; Ohyama, Y.; Fujieda, K.;
Inomata,

[104687]H.; Kameya, T.; Isozaki, O.; Jhiang, S. M.: High prevalence
of T354P

[104688]sodium/iodide symporter gene mutation in Japanese pa-
tients with iodide

[104689]transport defect who have heterogeneous clinical pictures.
J. Clin.

[104690]Endocr. Metab. 83: 4123–4129, 1998.

[104691]18324.

[104692]Levy, O.; Ginter, C. S.; De la Vieja, A.; Levy, D.; Carrasco,

[104693]N.: Identification of a structural requirement for thyroid
Na(+)/I(-)

[104694]symporter (NIS) function from analysis of a mutation that
causes human

[104695]congenital hypothyroidism. FEBS Lett. 429: 36-40, 1998.

[104696]18325.

[104697]Matsuda, A.; Kosugi, S.: A homozygous missense muta-
tion of the

[104698]sodium/iodide symporter gene causing iodide transport
defect. J.

[104699]Clin. Endocr. Metab. 82: 3966-3971, 1997.

[104700]18326.

[104701]Ohmori, M.; Endo, T.; Harii, N.; Onaya, T.: A novel thyroid
transcription

[104702]factor is essential for thyrotropin-induced up-regulation
of Na+ /I-

[104703]symporter gene expression. Molec. Endocr. 12: 727-736,
1998.

[104704]18327.

- [104705]Pohlenz, J.; Medeiros-Neto, G.; Gross, J. L.; Silveiro, S. P.;
- [104706]Knobel, M.; Refetoff, S.: Hypothyroidism in a Brazilian kindred due
- [104707]to iodide trapping defect caused by a homozygous mutation in the sodium/iodide
- [104708]symporter gene. *Biochem. Biophys. Res. Commun.* 240: 488–491, 1997.
- [104709]18328.
- [104710]Pohlenz, J.; Rosenthal, I. M.; Weiss, R. E.; Jhiang, S. M.; Barrant,
- [104711]C.; Refetoff, S.: Congenital hypothyroidism due to mutations in the
- [104712]sodium/iodide symporter: identification of a nonsense mutation producing
- [104713]a downstream cryptic 3-prime splice site. *J. Clin. Invest.* 101: 1028–1035, 1998.
- [104714]18329.
- [104715]18329.
- [104716]Smanik, P. A.; Liu, Q.; Furminger, T. L.; Ryu, K.; Xing, S.; Mazzaferri,
- [104717]E. L.; Jhiang, S. M.: Cloning of the human sodium iodide symporter. *Biochem.*

[104718]Biophys. Res. Commun. 226: 339–345, 1996.

[104719]18330.

[104720]Smanik, P. A.; Ryu, K.-Y.; Theil, K. S.; Mazzaferri, E. L.; Jhi-
ang,

[104721]S. M.: Expression, exon–intron organization, and chromo-
some mapping

[104722]of the human sodium iodide symporter. Endocrinology
138: 3555–3558,

[104723]1997.

[104724]18331.

[104725]Spitzweg, C.; Joba, W.; Schriever, K.; Goellner, J. R.; Morris,

[104726]J. C.; Heufelder, A. E.: Analysis of human sodium iodide
symporter

[104727]immunoreactivity in human exocrine glands. J. Clin. En-
docr. Metab. 84:

[104728]4178–4184, 1999.

[104729]18332.

[104730]Venkataraman, G. M.; Yatin, M.; Marcinek, R.; Ain, K. B.:
Restoration

[104731]of iodide uptake in dedifferentiated thyroid carcinoma: re-
lationship

- [104732]to human Na(+)/I(-) symporter gene methylation status. J. Clin. Endocr.
- [104733]Metab. 84: 2449–2457, 1999.
- [104734]18333.
- [104735]Lyon, M. F.; Peters, J.; Glenister, P. H.; Ball, S.; Wright, E.
- [104736]: The scurfy mouse mutant has previously unrecognized hematological abnormalities and resembles Wiskott–Aldrich syndrome. Proc. Nat.
- [104738]Acad. Sci. 87: 2433–2437, 1990.
- [104739]18334.
- [104740]Rawson, R. B.; Zelenski, N. G.; Nijhawan, D.; Ye, J.; Sakai, J.;
- [104741]Hasan, M. T.; Chang, T. Y.; Brown, M. S.; Goldstein, J. L.: Complementation
- [104742]cloning of S2P, a gene encoding a putative metalloprotease required
- [104743]for intramembrane cleavage of SREBPs. Molec. Cell 1: 47–57, 1997.
- [104744]18335.
- [104745]Ye, J.; Rawson, R. B.; Komuro, R.; Chen, X.; Dave, U. P.;

Prywes,

[104746]R.; Brown, M. S.; Goldstein, J. L.: ER stress induces cleavage of

[104747]membrane-bound ATF6 by the same proteases that process SREBPs. Molec.

[104748]Cell 6: 1355–1364, 2000.

[104749]18336.

[104750]Zelenski, N. G.; Rawson, R. B.; Brown, M. S.; Goldstein, J. L.

[104751]: Membrane topology of S2P, a protein required for intramembraneous

[104752]cleavage of sterol regulatory element-binding proteins. J. Biol.

[104753]Chem. 274: 21973–21980, 1999.

[104754]18337.

[104755]Baytel, D.; Shalom, S.; Madgar, I.; Weissenberg, R.; Don, J.:

[104756]The human Pim-2 proto-oncogene and its testicular expression. Biochim.

[104757]Biophys. Acta 1442: 274–285, 1998.

[104758]18338.

[104759]Cocchia, M.; Huber, R.; Pantano, S.; Chen, E. Y.; Ma, P.;

Forabosco,

[104760]A.; Ko, M. S. H.; Schlessinger, D.: PLAC1, an Xq26 gene
with placenta-specific

[104761]expression. Genomics 68: 305–312, 2000.

[104762]18339.

[104763]Kim, V. N.; Kataoka, N.; Dreyfuss, G.: Role of the non-
sense-mediated

[104764]decay factor hUpf3 in the splicing-dependent exon-exon
junction complex. Science 293:

[104765]1832–1836, 2001.

[104766]18340.

[104767]Lykke-Andersen, J.; Shu, M.-D.; Steitz, J. A.: Human Upf
proteins

[104768]target an mRNA for nonsense-mediated decay when
bound downstream of

[104769]a termination codon. Cell 103: 1121–1131, 2000.

[104770]18341.

[104771]Scott, A. F.: Personal Communication. Baltimore, Md.
2/9/2001.

[104772]18342.

[104773]Serin, G.; Gersappe, A.; Black, J. D.; Aronoff, R.; Maquat, L.

- [104774]E.: Identification and characterization of human orthologues to *Saccharomyces*
- [104775]*cerevisiae* Upf2 protein and Upf3 protein (*Caenorhabditis elegans* SMG-4). *Molec.*
- [104776]*Cell. Biol.* 21: 209–223, 2001.
- [104777]18343.
- [104778]Kelley, W. N.; Rosenbloom, F. M.; Henderson, J. F.; Seeg-
- millar,
- [104779]J. E.: A specific enzyme defect in gout associated with
- overproduction
- [104780]of uric acid. *Proc. Nat. Acad. Sci.* 57: 1735–1739, 1967.
- [104781]18344.
- [104782]McDonald, J. A.; Kelley, W. N.: Lesch–Nyhan syndrome: al-
- tered
- [104783]kinetic properties of mutant enzyme. *Science* 171:
- 689–691, 1971.
- [104784]18345.
- [104785]de Leeuw, B.; Balemans, M.; Geurts van Kessel, A.: A novel
- Kruppel-associated
- [104786]box containing the SSX gene (SSX3) on the human X chro-
- mosome is not

[104787] implicated in t(X;18)-positive synovial sarcomas. Cyto-genet. Cell

[104788] Genet. 73: 179-183, 1996.

[104789] 18346.

[104790] Gure, A. O.; Tureci, O.; Sahin, U.; Tsang, S.; Scanlan, M. J.;

[104791] Jager, E.; Knuth, A.; Pfreundschuh, M.; Old, L. J.; Chen, Y.-T.:

[104792] SSX: a multigene family with several members transcribed in normal

[104793] testis and human cancer. Int. J. Cancer 72: 965-971, 1997.

[104794] 18347.

[104795] Sossey-Alaoui, K.; Lyon, J. A.; Jones, L.; Abidi, F. E.; Hartung,

[104796] A. J.; Hane, B.; Schwartz, C. E.; Stevenson, R. E.; Srivastava, A.

[104797] K.: Molecular cloning and characterization of TRPC5 (HTRP5), the

[104798] human homologue of a mouse brain receptor-activated capacitative Ca(2+)

[104799] entry channel. Genomics 60: 330-340, 1999.

[104800]18348.

[104801]Tipnis, S. R.; Hooper, N. M.; Hyde, R.; Karran, E.; Christie, G.;

[104802]Turner, A. J.: A human homolog of angiotensin-converting enzyme:

[104803]cloning and functional expression as a captopril-insensitive carboxypeptidase. J.

[104804]Biol. Chem. 275: 33238-33243, 2000.

[104805]18349.

[104806]Johnson, R. E.; Kondratick, C. M.; Prakash, S.; Prakash, L.: hRAD30

[104807]mutations in the variant form of xeroderma pigmentosum. Science 285:

[104808]263-265, 1999.

[104809]18350.

[104810]Masutani, C.; Kusumoto, R.; Yamada, A.; Dohmae, N.; Yokoi, M.;

[104811]Yuasa, M.; Araki, M.; Iwai, S.; Takio, K.; Hanaoka, F.: The XPV (xeroderma

[104812]pigmentosum variant) gene encodes human DNA polymerase eta. Nature 399:

[104813]700–704, 1999.

[104814]18351.

[104815]Neitz, M.; Neitz, J.: Numbers and ratios of visual pigment genes

[104816]for normal red–green color vision. Science 267: 1013–1016, 1995.

[104817]18352.

[104818]Reyniers, E.; Van Thienen, M.–N.; Meire, F.; De Boulle, K.; Devries,

[104819]K.; Kestelijn, P.; Willems, P. J.: Gene conversion between red and

[104820]defective green opsin gene in blue cone monochromacy. Genomics 29:

[104821]323–328, 1995.

[104822]18353.

[104823]Ruberg, F. L.; Skene, D. J.; Hanifin, J. P.; Rollag, M. D.; English,

[104824]J.; Arendt, J.; Brainard, G. C.: Melatonin regulation in humans with

[104825]color vision deficiencies. J. Clin. Endocr. Metab. 81: 2980–2985,

[104826]1996.

[104827]18354.

[104828]Ueyama, H.; Kuwayama, S.; Imai, H.; Tanabe, S.; Oda, S.;
Nishida,

[104829]Y.; Wada, A.; Shichida, Y.; Yamade, S.: Novel missense
mutations

[104830]in red/green opsin genes in congenital color-vision defi-
ciencies. Biochem.

[104831]Biophys. Res. Commun. 294: 205-209, 2002.

[104832]18355.

[104833]Brunner, W.: Ueber den Vererbungsmodus der verschiede-
nen Typen

[104834]der angeborenen Rotgruenblindheit. Albrecht von Graefes
Arch. Ophthal. 124:

[104835]1-52, 1932.

[104836]18356.

[104837]Crone, R. A.: Spectral sensitivity in color-defective sub-
jects

[104838]and heterozygous carriers. Am. J. Ophthal. 48: 231-238,
1959.

[104839]18357.

[104840]Deeb, S. S.; Lindsey, D. T.; Hibiya, Y.; Sanocki, E.; Winder-
ickx,

[104841]J.; Teller, D. Y.; Motulsky, A. G.: Genotype-phenotype re-
lationships

[104842]in human red/green color-vision defects: molecular and
psychophysical

[104843]studies. Am. J. Hum. Genet. 51: 687-700, 1992.

[104844]18358.

[104845]Drummond-Borg, M.; Deeb, S.; Motulsky, A. G.: Molecular
basis

[104846]of abnormal red-green color vision: a family with three
types of color

[104847]vision defects. Am. J. Hum. Genet. 43: 675-683, 1988.

[104848]18359.

[104849]Emmerson, B. T.; Thompson, L.; Wallace, D. C.; Spence, M.
A.:

[104850]Absence of measurable linkage between the loci for hy-
poxanthine-guanine

[104851]phosphoribosyltransferase and deutan color blindness.
Am. J. Hum.

[104852]Genet. 26: 78-82, 1974.

[104853]18360.

[104854]Keating, J. P.; Brown, B. I.; White, N. H.; DiMauro, S.: X-linked

[104855]glycogen storage disease: a cause of hypotonia, hyperuricemia, and

[104856]growth retardation. Am. J. Dis. Child. 139: 609-613, 1985.

[104857]18361.

[104858]Krebs, E. G.; Love, D. S.; Bratvold, G. E.; Trayser, K. A.; Meyer,

[104859]W. L.; Fischer, E. H.: Purification and properties of rabbit skeletal

[104860]muscle phosphorylase B kinase. Biochemistry 3: 1022-1033, 1964.

[104861]18362.

[104862]Kumatori, A.; Faizunnessa, N. N.; Suzuki, S.; Moriuchi, T.; Kurozumi,

[104863]H.; Nakamura, M.: Nonhomologous recombination between the cytochrome

[104864]b-558 heavy chain gene (CYBB) and LINE-1 causes an X-linked chronic

[104865]granulomatous disease. Genomics 53: 123-128, 1998.

[104866]18363.

[104867]Lyon, J. B., Jr.; Porter, J.; Robertson, M.: Phosphorylase B

[104868]kinase inheritance in mice. Science 155: 1550–1551,
1967.

[104869]18364.

[104870]Malthus, R.; Clark, D. G.; Watts, C.; Sneyd, J. G. T.: Glyco-
gen-storage

[104871]disease in rats, a genetically determined deficiency of liver
phosphorylase

[104872]kinase. Biochem. J. 188: 99–106, 1980.

[104873]18365.

[104874]McAdams, A. J.; Hug, G.; Bove, K. E.: Glycogen storage
disease,

[104875]type I to X: criteria for morphologic diagnosis. Hum. Path.
5: 463–487,

[104876]1974.

[104877]18366.

[104878]Migeon, B. R.; Huijing, F.: Glycogen-storage disease asso-
ciated

[104879]with phosphorylase kinase deficiency: evidence for X inac-
tivation. Am.

[104880]J. Hum. Genet. 26: 360–368, 1974.

[104881]18367.

[104882]Nagai, T.; Matsuo, N.; Tsuchiya, Y.; Cho, H.; Hasegawa, Y.; Igarashi,

[104883]Y.: Proximal renal tubular acidosis associated with glycogen storage

[104884]disease, type 9. Acta Paediat. Scand. 77: 460–463, 1988.

[104885]18368.

[104886]Ryder–Cook, A. S.; Derry, J. M. J.; Barnard, P. J.: Localization

[104887]of the phosphorylase kinase alpha subunit gene on the mouse X chromosome.

[104888](Abstract) Cytogenet. Cell Genet. 51: 1071–1072, 1989.

[104889]18369.

[104890]Schimke, R. N.; Zakheim, R. M.; Corder, R. C.; Hug, G.: Glycogen

[104891]storage disease type IX: benign glycogenosis of liver and hepatic

[104892]phosphorylase kinase deficiency. J. Pediat. 83: 1031–1034, 1973.

[104893]18370.

[104894]Schneider, A.; Davidson, J. J.; Wullrich, A.; Kilimann, M. W.

[104895]Phosphorylase kinase deficiency in I-strain mice is associated with

[104896]a frameshift mutation in the alpha-subunit muscle isoform. Nature

[104897]Genet. 5: 381-385, 1993.

[104898]18371.

[104899]van den Berg, I. E. T.; van Beurden, E. A. C. M.; Malingre, H.

[104900]E. M.; Ploos van Amstel, H. K.; Poll-The, B. T.; Smeitink, J. A. M.;

[104901]Lamers, W. H.; Berger, R.: X-linked liver phosphorylase kinase deficiency

[104902]is associated with mutations in the human liver phosphorylase kinase

[104903]alpha subunit. Am. J. Hum. Genet. 56: 381-387, 1995.

[104904]18372.

[104905]Varsanyi, M.; Vrbica, A.; Heilmeyer, L. M. G., Jr.: X-linked

[104906]dominant inheritance of partial phosphorylase kinase deficiency in

[104907]mice. Biochem. Genet. 18: 247-261, 1980.

[104908]18373.

[104909]Wallis, P. G.; Sidbury, J. B., Jr.; Harris, R. C.: Hepatic phosphorylase

[104910]defect. Studies on peripheral blood. Am. J. Dis. Child. 111: 278–282,

[104911]1966.

[104912]18374.

[104913]Wauters, J. G.; Bossuyt, P. J.; Davidson, J.; Hendrickx, J.; Kilimann,

[104914]M. W.; Willems, P. J.: Regional mapping of a liver alpha-subunit

[104915]gene of phosphorylase kinase (PHKA) to the distal region of human

[104916]chromosome Xp. Cytogenet. Cell Genet. 60: 194–196, 1992.

[104917]18375.

[104918]Willems, P. J.; Gerver, W. J. M.; Berger, R.; Fernandes, J.:

[104919]The natural history of liver glycogenosis due to phosphorylase kinase

[104920]deficiency: a longitudinal study of 41 patients. Europ. J. Pediat. 149:

[104921]268–271, 1990.

[104922]18376.

[104923]Willems, P. J.; Hendrickx, J.; Van der Auwera, B. J.; Vits, L.;

[104924]Raeymaekers, P.; Coucke, P. J.; Van den Bergh, I.; Berger, R.; Smit,

[104925]G. P. A.; Van Broeckhoven, C.; Kilimann, M. W.; Van Elsen, A. F.;

[104926]Fernandes, J. F.: Mapping of the gene for X-linked liver glycogenosis

[104927]due to phosphorylase kinase deficiency to human chromosome region

[104928]Xp22. Genomics 9: 565–569, 1991.

[104929]18377.

[104930]Williams, H. E.; Field, J. B.: Low leukocyte phosphorylase in

[104931]hepatic phosphorylase deficient glycogen storage disease. J. Clin.

[104932]Invest. 40: 1841–1845, 1961.

[104933]18378.

[104934]Zander, N. F.; Meyer, H. E.; Hoffmann–Posorske, E.; Crabb, J.

- [104935]W.; Heilmeyer, L. M. G., Jr.; Kilimann, M. W.: cDNA cloning and complete
[104936]primary structure of skeletal muscle phosphorylase kinase (alpha subunit). Proc.
[104937]Nat. Acad. Sci. 85: 2929–2933, 1988.
[104938]18379.
- [104939]Berta, P.; Hawkins, J. R.; Sinclair, A. H.; Taylor, A.; Griffiths,
[104940]B. L.; Goodfellow, P. N.; Fellous, M.: Genetic evidence equating
[104941]SRY and the testis-determining factor. Nature 348: 448–450, 1990.
[104942]18380.
- [104943]Disteche, C. M.; Casanova, M.; Saal, H.; Friedman, C.; Sybert,
[104944]V.; Graham, J.; Thuline, H.; Page, D.; Fellous, M.: Small deletions
[104945]of the short arm of the Y chromosome in 46,XY females. Proc. Nat.
[104946]Acad. Sci. 83: 7841–7844, 1986.
[104947]18381.
- [104948]Georgopoulos, N. A.; Pralong, F. P.; Seidman, C. E.; Seid-

man,

[104949]J. G.; Crowley, W. F., Jr.; Vallejo, M.: Genetic heterogeneity evidenced

[104950]by low incidence of KAL-1 gene mutations in sporadic cases of gonadotropin-releasing

[104951]hormone deficiency. J. Clin. Endocr. Metab. 82: 213-217, 1997.

[104952]18382.

[104953]Guioli, S.; Incerti, B.; Zanaria, E.; Bardoni, B.; Franco, B.;

[104954]Taylor, K.; Ballabio, A.; Camerino, G.: Kallmann syndrome due to

[104955]a translocation resulting in an X/Y fusion gene. Nature Genet. 1:

[104956]337-340, 1992.

[104957]18383.

[104958]Hardelin, J.-P.; Levilliers, J.; Blanchard, S.; Carel, J.-C.;

[104959]Leutenegger, M.; Pinard-Bertelletto, J.-P.; Bouloux, P.; Petit, C.

[104960]: Heterogeneity in the mutations responsible for X chromosome-linked

[104961]Kallmann syndrome. Hum. Molec. Genet. 2: 373-377,

1993.

[104962]18384.

[104963]Hardelin, J.-P.; Levilliers, J.; del Castillo, I.; Cohen-Salmon,

[104964]M.; Legouis, R.; Blanchard, S.; Compain, S.; Bouloux, P.;

Kirk, J.;

[104965]Moraine, C.; Chaussain, J.-L.; Weissenbach, J.; Petit, C.: X

chromosome-linked

[104966]Kallmann syndrome: stop mutations validate the candi-

date gene. Proc.

[104967]Nat. Acad. Sci. 89: 8190-8194, 1992.

[104968]18385.

[104969]Henkin, R. I.: Abnormalities of taste and olfaction in pa-

tients

[104970]with chromatin negative gonadal dysgenesis. J. Clin. En-

docr. 27:

[104971]1436-1440, 1967.

[104972]18386.

[104973]Hermanussen, M.; Sippell, W. G.: Heterogeneity of Kallmann's

[104974]syndrome. Clin. Genet. 28: 106–111, 1985.

[104975]18387.

[104976]Hipkin, L. J.; Casson, I. F.; Davis, J. C.: Identical twins discordant

[104977]for Kallmann's syndrome. J. Med. Genet. 27: 198–199, 1990.

[104978]18388.

[104979]Incerti, B.; Guioli, S.; Pragliola, A.; Zanaria, E.; Borsani,

[104980]G.; Tonlorenzi, R.; Bardoni, B.; Franco, B.; Wheeler, D.; Balabio,

[104981]A.; Camerino, G.: Kallmann syndrome gene on the X and Y chromosomes:

[104982]implications for evolutionary divergence of human sex chromosomes. Nature

[104983]Genet. 2: 311–314, 1992.

[104984]18389.

[104985]Kirk, J. M. W.; Grant, D. B.; Besser, G. M.; Shalet, S.; Quinton,

[104986]R.; Smith, C. S.; White, M.; Edwards, O.; Bouloux, P.–M. G.:

Unilateral

[104987]renal aplasia in X-linked Kallmann's syndrome. Clin.

Genet. 46:

[104988]260-262, 1994.

[104989]18390.

[104990]Krams, M.; Quinton, R.; Ashburner, J.; Friston, K. J.; Frack-
owiak,

[104991]R. S. J.; Bouloux, P.-M. G.; Passingham, R. E.: Kallmann's
syndrome:

[104992]mirror movements associated with bilateral corticospinal
tract hypertrophy. Neurology 52:

[104993]816-822, 1999.

[104994]18391.

[104995]Legouis, R.; Cohen-Salmon, M.; del Castillo, I.; Levilliers,
J.;

[104996]Capy, L.; Mornon, J.-P.; Petit, C.: Characterization of the
chicken

[104997]and quail homologues of the human gene responsible for
the X-linked

[104998]Kallmann syndrome. Genomics 17: 516-518, 1993.

[104999]18392.

- [105000]Legouis, R.; Hardelin, J.-P.; Levilliers, J.; Claverie, J.-M.;
- [105001]Compain, S.; Wunderle, V.; Millasseau, P.; Le Paslier, D.;
Cohen,
- [105002]D.; Caterina, D.; Bougueleret, L.; Delemarre-Van de Waal,
H.; Lutfalla,
- [105003]G.; Weissenbach, J.; Petit, C.: The candidate gene for the
X-linked
- [105004]Kallmann syndrome encodes a protein related to adhesion
molecules. Cell 67:
- [105005]423-435, 1991.
- [105006]18393.
- [105007]Males, J. L.; Townsend, J. L.; Schneider, R. A.: Hypogo-
nadotropic
- [105008]hypogonadism with anosmia--Kallmann's syndrome: a
disorder of olfactory
- [105009]and hypothalamic function. Arch. Intern. Med. 131:
501-507, 1973.
- [105010]18394.
- [105011]Maya-Nunez, G.; Covarrubias, S. C.; Zenteno, J. C.; Ulloa-
Aguirre,
- [105012]A.; Kofman-Alfaro, S.; Mendez, J. P.: Contiguous gene
syndrome due

[105013] to deletion of the first three exons of the Kallmann gene
and complete

[105014] deletion of the steroid sulphatase gene. Clin. Endocr. 48:
713–718,

[105015] 1998.

[105016] 18395.

[105017] Maya–Nunez, G.; Zenteno, J. C.; Ulloa–Aguirre, A.; Kof–
man–Alfaro,

[105018] S.; Mendez, J. P.: A recurrent missense mutation in the
KAL gene

[105019] in patients with X–linked Kallmann's syndrome. J. Clin. En–
docr. Metab. 83:

[105020] 1650–1653, 1998.

[105021] 18396.

[105022] Meitinger, T.; Heye, B.; Petit, C.; Levilliers, J.; Golla, A.;

[105023] Moraine, C.; Dalla Piccola, B.; Sippell, W. G.; Murken, J.;

Ballabio,

[105024] A.: Definitive localization of X–linked Kallmann syndrome
(hypogonadotropic

[105025] hypogonadism and anosmia) to Xp22.3: close linkage to
the hypervariable

- [105026]repeat sequence CRI-S232. Am. J. Hum. Genet. 47:
664-669, 1990.
- [105027]18397.
- [105028]Nagata, K.; Yamamoto, T.; Chikumi, H.; Ikeda, T.; Yamamoto, H.;
- [105029]Hashimoto, K.; Yoneda, K.; Nanba, E.; Ninomiya, H.; Ishitobi, K.:
- [105030]A novel interstitial deletion of KAL1 in a Japanese family with Kallmann
- [105031]syndrome. J. Hum. Genet. 45: 237-240, 2000.
- [105032]18398.
- [105033]Nass, R.: Mirror movement asymmetries in congenital hemiparesis:
- [105034]the inhibition hypothesis revisited. Neurology 35:
1059-1062, 1985.
- [105035]18399.
- [105036]Nowakowski, H.; Lenz, W.: Genetic aspects in male hypogonadism. Recent
- [105037]Prog. Horm. Res. 17: 53-95, 1961.
- [105038]18400.
- [105039]Oppermann, D.; Happ, J.; Mayr, W. R.: Stimulation of sper-

matogenesis

[105040]and biological paternity by intranasal (low dose) gonadotropin-releasing

[105041]hormone (GnRH) in a male with Kallmann's syndrome: intra-individual

[105042]comparison of GnRH and gonadotropins for stimulation of spermatogenesis. J.

[105043]Clin. Endocr. Metab. 65: 1060-1066, 1987.

[105044]18401.

[105045]Parenti, G.; Rizzolo, M. G.; Ghezzi, M.; Di Maio, S.; Sperandio,

[105046]M. P.; Incerti, B.; Franco, B.; Ballabio, A.; Andria, G.: Variable

[105047]penetrance of hypogonadism in a sibship with Kallmann syndrome due

[105048]to a deletion of the KAL gene. Am. J. Med. Genet. 57: 476-478, 1995.

[105049]18402.

[105050]Pawlowitzki, I. H.; Diekstall, P.; Schadel, A.; Miny, P.: Estimating

[105051]frequency of Kallmann syndrome among hypogonadic and among anosmic

[105052]patients. Am. J. Med. Genet. 26: 473–479, 1987.

[105053]18403.

[105054]Petit, C.; Levilliers, J.; Weissenbach, J.: Long-range restriction

[105055]map of the terminal part of the short arm of the human X chromosome. Proc.

[105056]Nat. Acad. Sci. 87: 3680–3684, 1990.

[105057]18404.

[105058]Pittman, J.: Personal Communication. Boston, Mass. 1966.

[105059]18405.

[105060]Prager, D.; Braunstein, G. D.: X-chromosome-linked Kallmann's

[105061]syndrome: pathology at the molecular level. J. Clin. Endocr. Metab. 76:

[105062]824–826, 1993.

[105063]18406.

[105064]Imperato-McGinley, J.; Peterson, R. E.; Leshin, M.; Griffin, J.

[105065]E.; Cooper, G.; Draghi, S.; Berenyi, M.; Wilson, J. D.: Steroid

[105066]alpha-reductase deficiency in a 65-year-old male pseudohermaphrodite:

[105067]the natural history, ultrastructure of the testes, and evidence for

[105068]inherited enzyme heterogeneity. J. Clin. Endocr. Metab. 50: 15-22,

[105069]1980.

[105070]18407.

[105071]Jenkins, E. P.; Andersson, S.; Imperato-McGinley, J.; Wilson,

[105072]J. D.; Russell, D. W.: Genetic and pharmacological evidence for more

[105073]than one human steroid 5-alpha-reductase. J. Clin. Invest. 89: 293-300,

[105074]1992.

[105075]18408.

[105076]Francke, U.; Taggart, R. T.: Assignment of the gene for cytoplasmic

[105077]superoxide dismutase (Sod-1) to a region of chromosome 16 and HPRT

[105078]to a region of the X-chromosome in the mouse. Proc. Nat. Acad. Sci. 76:

[105079]5230-5233, 1979.

[105080]18409.

[105081]Francke, U.; Taggart, R. T.: Regional mapping of SOD-1 on
mouse

[105082]chromosome 16, and of HPRT and alpha-GAL (Ags) on the
mouse X, using

[105083]Chinese hamster-mouse T(X;16)16H somatic cell hybrids.
(Abstract) Cytogenet.

[105084]Cell Genet. 25: 155-156, 1979.

[105085]18410.

[105086]Fujimori, S.; Davidson, B. L.; Kelley, W. N.; Palella, T. D.:

[105087]Identification of a single nucleotide change in the hypox-
anthine-guanine

[105088]phosphoribosyltransferase gene (HPRT-Yale) responsible
for Lesch-Nyhan

[105089]syndrome. J. Clin. Invest. 83: 11-13, 1989.

[105090]18411.

[105091]Fujimori, S.; Hidaka, Y.; Davidson, B. L.; Palella, T. D.; Kel-
ley,

[105092]W. N.: Identification of a single nucleotide change in a
mutant gene

- [105093]for hypoxanthine-guanine phosphoribosyltransferase (HPRT-Ann Arbor). Hum. Genet. 79: 39-43, 1988.
- [105094]18412.
- [105095]18412.
- [105096]Fujimori, S.; Kamatani, N.; Nishida, Y.; Ogasawara, N.; Akaoka, I.: Hypoxanthine guanine phosphoribosyltransferase deficiency: nucleotide substitution causing Lesch-Nyhan syndrome identified for the first time among Japanese. Hum. Genet. 84: 483-486, 1990.
- [105097]18413.
- [105098]18413.
- [105099]Fujimori, S.; Tagaya, T.; Kamatani, N.; Akaoka, I.: A germ line mutation within the coding sequence for the putative 5-phosphoribosyl-1-pyrophosphate binding site of hypoxanthine-guanine phosphoribosyltransferase (HPRT) in a Lesch-Nyhan patient: missense mutations within a functionally important region probably cause disease. Hum. Genet. 90: 385-388,

[105106]1992.

[105107]18414.

[105108]Fujimori, S.; Tagaya, T.; Yamaoka, N.; Kamatani, N.;
Akaoka, I.

[105109]: Identification of mutations leading to the Lesch–Nyhan
syndrome

[105110]in two unrelated Japanese patients. (Abstract) Jpn. J. Hum.
Genet. 36:

[105111]58 only, 1991.

[105112]18415.

[105113]Gellin, J.; Benne, F.; Renard, C.; Vaiman, M.; Hors–Cayla,
M.

[105114]C.; Gillois, M.: Pig gene mapping: syntenic, attempt to as-
sign the

[105115]histocompatibility complex (SLA). (Abstract) Cytogenet.
Cell Genet. 25:

[105116]159 only, 1979.

[105117]18416.

[105118]Gibbs, R. A.; Caskey, C. T.: Identification and localization

[105119]of mutations at the Lesch–Nyhan locus by ribonuclease A
cleavage. Science 236:

[105120]303–305, 1987.

[105121]18417.

[105122]Gibbs, R. A.; Nguyen, P.–N.; Edwards, A.; Civitello, A. B.;
Caskey,

[105123]C. T.: Multiplex DNA deletion detection and exon se-
quencing of the

[105124]hypoxanthine phosphoribosyltransferase gene in Lesch-
Nyhan families. *Genomics* 7:

[105125]235–244, 1990.

[105126]18418.

[105127]Gibbs, R. A.; Nguyen, P.–N.; McBride, L. J.; Koepf, S. M.;
Caskey,

[105128]C. T.: Identification of mutations leading to the Lesch-
Nyhan syndrome

[105129]by automated direct DNA sequencing of in vitro amplified
cDNA. *Proc.*

[105130]*Nat. Acad. Sci.* 86: 1919–1923, 1989.

[105131]18419.

[105132]Gordon, R. B.; Dawson, P. A.; Sculley, D. G.; Emmerson, B.
T.;

[105133]Caskey, C. T.; Gibbs, R. A.: The molecular characterisation

of HPRT(Chermside)

[105134]and HPRT(Coorparoo): two Lesch–Nyhan patients with reduced amounts

[105135]of mRNA. Gene 108: 299–304, 1991.

[105136]18420.

[105137]Gordon, R. B.; Sculley, D. G.; Dawson, P. A.; Beacham, I. R.;

[105138]Emmerson, B. T.: Identification of a single nucleotide substitution

[105139]in the coding sequence of in vitro amplified cDNA from a patient with

[105140]partial HPRT deficiency (HPRT–Brisbane). J. Inherit. Metab. Dis. 13:

[105141]692–700, 1990.

[105142]18421.

[105143]Greene, M. L.: Clinical features of patients with the 'partial'

[105144]deficiency of the X-linked uricaciduria enzyme. Arch. Intern. Med. 130:

[105145]193–198, 1972.

[105146]18422.

[105147]Gutensohn, W.; Jahn, H.: Partial deficiency of hypoxan–

thine-phosphoribosyltransferase:

[105148]evidence for a structural mutation in a patient with gout.

Europ.

[105149]J. Clin. Invest. 9: 43-47, 1979.

[105150]18423.

[105151]Hashmi, S.; Miller, O. J.: Further evidence of X-linkage of hypoxanthine

[105152]phosphoribosyl-transferase in the mouse. Cytogenet. Cell Genet. 17:

[105153]35-41, 1976.

[105154]18424.

[105155]Anguiano, A.; Oates, R. D.; Amos, J. A.; Dean, M.; Gerrard, B.;

[105156]Stewart, C.; Maher, T. A.; White, M. B.; Milunsky, A.: Congenital

[105157]bilateral absence of the vas deferens: a primarily genital form of

[105158]cystic fibrosis. J.A.M.A. 267: 1794-1797, 1992.

[105159]18425.

[105160]Chillon, M.; Casals, T.; Mercier, B.; Bassas, L.; Lissens, W.;

[105161]Silber, S.; Romey, M.-C.; Ruiz-Romero, J.; Verlingue, C.;

Claustres,

[105162]M.; Nunes, V.; Ferec, C.; Estivill, X.: Mutations in the cystic fibrosis

[105163]gene in patients with congenital absence of the vas deferens. New

[105164]Eng. J. Med. 332: 1475–1480, 1995.

[105165]18426.

[105166]Dumur, V.; Gervais, R.; Rigot, J.-M.; Delomel-Vinner, E.; Decaestecker,

[105167]B.; Lafitte, J.-J.; Roussel, P.: Congenital bilateral absence of

[105168]the vas deferens (CBAVD) and cystic fibrosis transmembrane regulator

[105169](CFTR): correlation between genotype and phenotype. Hum. Genet. 97:

[105170]7–10, 1996.

[105171]18427.

[105172]Andersson, M.; Page, D. C.; de la Chapelle, A.: Chromosome Y-specific

[105173]DNA is transferred to the short arm of X chromosome in human XX males. Science 233:

[105174]786–788, 1986.

[105175]18428.

[105176]Ferguson-Smith, M. A.: X-Y chromosomal interchange in
the aetiology

[105177]of true hermaphroditism and of XX Klinefelter's syndrome.
Lancet II:

[105178]475-476, 1966.

[105179]18429.

[105180]Tiepolo, L.; Zuffardi, O.: Localization of factors controlling

[105181]spermatogenesis in the nonfluorescent portion of the hu-
man Y chromosome

[105182]long arm. Hum. Genet. 34: 119-124, 1976.

[105183]18430.

[105184]Sertie, A. L.; Sossi, V.; Camargo, A. A.; Zatz, M.; Brahe, C.;

[105185]Passos-Bueno, M. R.: Collagen XVIII, containing an en-
dogenous inhibitor

[105186]of angiogenesis and tumor growth, plays a critical role in
the maintenance

[105187]of retinal structure and in neural tube closure (Knobloch
syndrome). Hum.

[105188]Molec. Genet. 9: 2051-2058, 2000.

[105189]18431.

[105190]Fujiwara, H.; Tatsumi, K.; Miki, K.; Harada, T.; Miyai, K.;
Takai,

[105191]S.; Amino, N.: Congenital hypothyroidism caused by a
mutation in

[105192]the Na(+)/I(-) symporter. (Letter) Nature Genet. 16:
124-125, 1997.

[105193]18432.

[105194]Muroi, J.; Yorifuji, T.; Uematsu, A.; Shigematsu, Y.; Oni-
gata,

[105195]K.; Maruyama, H.; Nobutoki, T.; Kitamura, A.; Nakahata,
T.: Molecular

[105196]and clinical analysis of Japanese patients with
3-hydroxy-3-methylglutaryl

[105197]CoA lyase (HL) deficiency. Hum. Genet. 107: 320-326,
2000.

[105198]18433.

[105199]Ozand, P. T.; Al Aqeel, A.; Gascon, G.; Brismar, J.; Thomas,
E.;

[105200]Gleispach, H.: 3-Hydroxy-3-methylglutaryl-coenzyme A
(HMG-CoA) lyase

[105201]deficiency in Saudi Arabia. J. Inherit. Metab. Dis. 14:

174-188,

[105202]1991.

[105203]18434.

[105204]Ozand, P. T.; Devol, E. B.; Gascon, G. G.: Neurometabolic diseases

[105205]at a national referral center: five years experience at the King Faisal

[105206]Specialist Hospital and Research Centre. J. Child Neurol. 7 (suppl.):

[105207]S4-S11, 1992.

[105208]18435.

[105209]Ribes, A.; Briones, P.; Vilaseca, M. A.; Baraibar, R.; Gairi,

[105210]J. M.: Sudden death in an infant with 3-hydroxy-3-methylglutaryl-CoA

[105211]lyase deficiency. J. Inherit. Metab. Dis. 13: 752-753, 1990.

[105212]18436.

[105213]Robinson, B. H.; Oei, J.; Sherwood, W. G.; Slyper, A. H.; Heininger,

[105214]J.; Mamer, O. A.: Hydroxymethylglutaryl CoA lyase deficiency: features

[105215]resembling Reye syndrome. Neurology 30: 714-718,

1980.

[105216]18437.

[105217]Roe, C. R.; Millington, D. S.; Maltby, D. A.: Identification

[105218]of 3-methylglutaryl carnitine: a new diagnostic metabolite
of 3-hydroxy-3-methylglutaryl-coenzyme

[105219]A lyase deficiency. J. Clin. Invest. 77: 1391-1394, 1986.

[105220]18438.

[105221]Shilkin, R.; Wilson, G.; Owles, E.:

3-Hydroxy-3-methylglutaryl

[105222]coenzyme A lyase deficiency: follow-up of first described
case. Acta

[105223]Paediat. Scand. 70: 265-268, 1981.

[105224]18439.

[105225]Shutgens, R. B. H.; Haymans, H.; Ketel, A.: Lethal hypo-
glycemia

[105226]in a child with a deficiency of 3-hydroxy-3-methylglutaryl
coenzyme

[105227]A lyase. J. Pediat. 94: 89-91, 1979.

[105228]18440.

[105229]Sovik, O.; Sweetman, L.; Gibson, K. M.; Nyhan, W. L.: Ge-

netic

[105230] complementation analysis of

3-hydroxy-3-methylglutaryl-coenzyme A

[105231] lyase deficiency in cultured fibroblasts. Am. J. Hum.

Genet. 36:

[105232] 791-801, 1984.

[105233] 18441.

[105234] Wanders, R. J. A.; Schutgens, R. B. H.; Zoeters, P. H. M.:

3-Hydroxy-3-methylglutaryl-CoA

[105235] lyase in human skin fibroblasts: study of its properties
and deficient

[105236] activity in 3-hydroxy-3-methylglutaric aciduria patients
using a simple

[105237] spectrophotometric method. Clin. Chim. Acta 171:
95-102, 1988.

[105238] 18442.

[105239] Wang, S.; Nadeau, J. H.; Duncan, A.; Robert, M.-F.;
Fontaine,

[105240] G.; Schappert, K.; Johnson, K. R.; Zietkiewicz, E.; Hruz, P.;
Miziorko,

[105241] H.; Mitchell, G. A.: 3-Hydroxy-3-methylglutaryl coenzyme
A lyase

[105242](HL): cloning and characterization of a mouse liver HL
cDNA and subchromosomal

[105243]mapping of the human and mouse HL genes. Mammalian
Genome 4: 382-387,

[105244]1993.

[105245]18443.

[105246]Wang, S. P.; Marth, J. D.; Oligny, L. L.; Vachon, M.; Robert,

[105247]M.-F.; Ashmarina, L.; Mitchell, G. A.:

3-Hydroxy-3-methylglutaryl-CoA

[105248]lyase (HL): gene targeting causes prenatal lethality in HL-
deficient

[105249]mice. Hum. Molec. Genet. 7: 2057-2062, 1998.

[105250]18444.

[105251]Wang, S. P.; Robert, M.-F.; Gibson, K. M.; Wanders, R. J. A.;

[105252]Mitchell, G. A.: 3-Hydroxy-3-methylglutaryl CoA lyase
(HL): mouse

[105253]and human HL gene (HMGCL) cloning and detection of
large gene deletions

[105254]in two unrelated HL-deficient patients. Genomics 33:
99-104, 1996.

[105255]18445.

[105256]Wilson, W. G.; Cass, M. B.; Sovik, O.; Gibson, K. M.; Sweetman,

[105257]L.: A child with acute pancreatitis and recurrent hypoglycemia due

[105258]to 3-hydroxy-3-methylglutaryl-CoA lyase deficiency. Europ. J. Pediat. 142:

[105259]289-291, 1984.

[105260]18446.

[105261]Wysocki, S. J.; Hahnel, R.: 3-Hydroxy-3-methylglutaric aciduria:

[105262]3-hydroxy-3-methylglutaryl-coenzyme A lyase levels in leucocytes. Clin.

[105263]Chim. Acta 73: 373-375, 1976.

[105264]18447.

[105265]Wysocki, S. J.; Hahnel, R.:

3-Hydroxy-3-methylglutaryl-coenzyme

[105266]A lyase deficiency: a review. J. Inherit. Metab. Dis. 9: 225-233,

[105267]1986.

[105268]18448.

[105269]Laporte, J.; Kioschis, P.; Hu, L.-J.; Kretz, C.; Carlsson, B.;

[105270]Poustka, A.; Mandel, J. L.; Dahl, N.: Cloning and characterization
[105271]of an alternatively spliced gene in proximal Xq28 deleted
[105272]in two patients
[105273]with intersexual genitalia and myotubular myopathy. Genomics 41:
[105274]458-462, 1997.
[105275]18449.
[105276]Caspi, M.; Atlas, R.; Kantor, A.; Sapir, T.; Reiner, O.: Interaction
[105277]between LIS1 and doublecortin, two lissencephaly gene
[105278]products. Hum. Molec. Genet. 9: 2205-2213, 2000.
[105279]18450.
[105280]des Portes, V.; Francis, F.; Pinard, J.-M.; Desguerre, I.;
[105281]Moutard,
[105282]M.-L.; Snoeck, I.; Meiners, L. C.; Capron, F.; Cusmai, R.;
[105283]Ricci,
[105284]S.; Motte, J.; Echenne, B.; Ponsot, G.; Dulac, O.; Chelly, J.;
[105285]Beldjord,
[105286]C.: Doublecortin is the major gene causing X-linked sub-
[105287]cortical laminar

[105283]heterotopia (SCLH). Hum. Molec. Genet. 7: 1063–1070, 1998.

[105284]18451.

[105285]Gleeson, J. G.; Lin, P. T.; Flanagan, L. A.; Walsh, C. A.: Doublecortin

[105286]is a microtubule-associated protein and is expressed widely by migrating

[105287]neurons. Neuron 23: 257–271, 1999.

[105288]18452.

[105289]Gleeson, J. G.; Minnerath, S.; Kuzniecky, R. I.; Dobyns, W. B.;

[105290]Young, I. D.; Ross, M. E.; Walsh, C. A.: Somatic and germline mosaic

[105291]mutations in the doublecortin gene are associated with variable phenotypes. Am.

[105292]J. Hum. Genet. 67: 574–581, 2000.

[105293]18453.

[105294]Gleeson, J. G.; Minnerath, S. R.; Fox, J. W.; Allen, K. M.; Luo,

[105295]R. F.; Hong, S. E.; Berg, M. J.; Kuzniecky, R.; Reitnauer, P. J.;

- [105296]Borgatti, R.; Puche Mira, A.; Guerrini, R.; and 14 others:
Characterization
[105297]of mutations in the gene doublecortin in patients with
double cortex
[105298]syndrome. *Ann. Neurol.* 45: 146–153, 1999.
[105299]18454.
- [105300]Pilz, D. T.; Kuc, J.; Matsumoto, N.; Bodurtha, J.; Bernadi, B.;
[105301]Tassinari, C. A.; Dobyns, W. B.; Ledbetter, D. H.: Subcorti-
cal band
[105302]heterotopia in rare affected males can be caused by mis-
sense mutations
[105303]in DCX (XLIS) or LIS1. *Hum. Molec. Genet.* 8: 1757–1760,
1999.
[105304]18455.
- [105305]Sapir, T.; Horesh, D.; Caspi, M.; Atlas, R.; Burgess, H. A.;
Wolf,
[105306]S. G.; Francis, F.; Chelly, J.; Elbaum, M.; Pietrokovski, S.;
Reiner,
[105307]O.: Doublecortin mutations cluster in evolutionarily con-
served functional
[105308]domains. *Hum. Molec. Genet.* 9: 703–712, 2000.

[105309]18456.

[105310]Shmueli, O.; Gdalyahu, A.; Sorokina, K.; Nevo, E.; Avivi, A.;

[105311]Reiner, O.: DCX in PC12 cells: CREB-mediated transcription and neurite

[105312]outgrowth. Hum. Molec. Genet. 10: 1061–1070, 2001.

[105313]18457.

[105314]Sossey-Alaoui, K.; Hartung, A. J.; Guerrini, R.; Manchester, D.

[105315]K.; Posar, A.; Puche-Mira, A.; Andermann, E.; Dobyns, W. B.; Srivastava,

[105316]A. K.: Human doublecortin (DCX) and the homologous gene in mouse

[105317]encode a putative Ca(2+)-dependent signaling protein which is mutated

[105318]in human X-linked neuronal migration defects. Hum. Molec. Genet. 7:

[105319]1327–1332, 1998.

[105320]18458.

[105321]Yourc'h, P.; Petit, E.; Muh, J. P.; Andres, C.; Bienvenu, T.;

[105322]Beldjord, C.; Chelly, J.; Barthelemy, C.: Exclusion of the coding

- [105323] sequence of the doublecortin gene as a susceptibility locus in autistic disorder. (Letter) Am. J. Med. Genet. 108: 164–167, 2002.
- [105324] 18459.
- [105326] Farnsworth, D.: The Farnsworth–Munsell 100 hue and dichotomy tests for colour vision. J. Ophthal. Soc. Am. 33: 568–578, 1943.
- [105328] 18460.
- [105329] Franceschetti, A.; Klein, D.: Two families with parents of different types of red–green blindness. Acta Genet. Statist. Med. 7: 255–259, 1957.
- [105331] 18461.
- [105332] 18461.
- [105333] Fraser, G. R.: Estimation of the recombination fraction between the protan and deutan loci. Am. J. Hum. Genet. 21: 593–599, 1969.
- [105335] 18462.
- [105336] Kalmus, H.: Diagnosis and Genetics of Defective Colour Vision.

[105337]Oxford: Pergamon Press (pub.) 1965.

[105338]18463.

[105339]Kondo, T.: Untersuchungen bei angeborenen Farbensinn-
Anomalien.

[105340]Ueber das Zustandekommen und Wesen der angeborenen
Farbensinn-Anomalien. Acta

[105341]Soc. Ophthal. Jpn. 45: 659, 1941.

[105342]18464.

[105343]Nemoto, H.; Murao, M.: A genetic study of colorblindness.
Jpn.

[105344]J. Hum. Genet. 6: 165-173, 1961.

[105345]18465.

[105346]Schmidt, I.: A sign of manifest heterozygosity in carriers
of

[105347]color deficiency. Am. J. Optom. 32: 404-408, 1955.

[105348]18466.

[105349]Siniscalco, M.; Filippi, G.; Latte, B.: Recombination be-
tween

[105350]protan and deutan genes: data on their relative positions
in respect

[105351]of the G6PD locus. Nature 204: 1062–1064, 1964.

[105352]18467.

[105353]Sloan, L. L.: Evaluation of the Tokyo Medical College color vision

[105354]test. Am. J. Ophthal. 52: 650–659, 1961.

[105355]18468.

[105356]Thuline, H. C.; Hodgkin, W. E.; Fraser, G. R.; Motulsky, A. G.

[105357]: Genetics of protan and deutan color–vision anomalies: an instructive

[105358]family. Am. J. Hum. Genet. 21: 581–592, 1969.

[105359]18469.

[105360]Vanderdonck, R.; Verriest, G.: Femme protanomale et heterozygote

[105361]mixte (genes de la protanomalie et de la deuteranopie en position

[105362]de repulsion) ayant deux fils deuteranopes, un fils protanomal et

[105363]deux fils normaux. Biotypologie 21: 110–120, 1960.

[105364]18470.

[105365] von Kries, J.; Nagel, W.: Ueber den Einfluss von Licht-
staerke

[105366] und Adaptation auf das Sehen des Dichromaten
(Gruenblinden). Ztschr.

[105367] Psychol. Physiol. Sinnesorg. 12: 1–38, 1896.

[105368] 18471.

[105369] von Planta, P.: Die Haeufigkeit der angeborenen Far-
bensinnstoerungen

[105370] bei Knaben und Maedchen und ihre Feststellung durch die
ueblichen

[105371] klinischen Proben. Albrecht von Graefes Arch. Klin. Exp.
Ophthal. 120:

[105372] 253–281, 1928.

[105373] 18472.

[105374] Waaler, G. H.: Ueber die Erblichkeitsverhaeltnisse der ver-
schieden

[105375] Arten von angeborener Rotgruenblindheit. Ztschr. F. In-
dukt. Abstammungs-u.

[105376] Vererbungsbl. 45: 279–333, 1927.

[105377] 18473.

[105378] Winderickx, J.; Lindsey, D. T.; Sanocki, E.; Teller, D. Y.;

Motulsky,

[105379]A. G.; Deeb, S. S.: Polymorphism in red photopigment underlies variation

[105380]in colour matching. *Nature* 356: 431–433, 1992.

[105381]18474.

[105382]Bordarier, C.; Robain, O.; Rethore, M.–O.; Dulac, O.; Dhellemmes,

[105383]C.: Inverted neurons in agyria: a Golgi study of a case with abnormal

[105384]chromosome 17. *Hum. Genet.* 73: 374–378, 1986.

[105385]18475.

[105386]Chong, S. S.; Lo Nigro, C.; Roschke, A. V.; Tanigami, A.; Pack,

[105387]S. D.; Smith, A. C. M.; Carrozzo, R.; Dobyns, W. B.; Ledbetter, D.

[105388]H.: Point mutations and an intragenic deletion in three ILS patients

[105389]confirm LIS1 as the lissencephaly causative gene in isolated lissencephaly

[105390]sequence and Miller–Dieker syndrome. (Abstract) *Am. J. Hum. Genet.* 59

[105391](suppl.): A23 only, 1996.

[105392]18476.

[105393]Hanahan, D. J. A.: Platelet activating factor: a biologically

[105394]active phosphoglyceride. Annu. Rev. Biochem. 55:

483–509, 1986.

[105395]18477.

[105396]Hattori, M.; Adachi, H.; Tsujimoto, M.; Arai, H.; Inoue, K.:

[105397]Miller–Dieker lissencephaly gene encodes a subunit of

brain platelet–activating

[105398]factor. Nature 370: 216–218, 1994.

[105399]18478.

[105400]Hattori, M.; Arai, H.; Inoue, K.: Purification and characteri-
zation

[105401]of bovine brain platelet–activating factor acetylhydrolase.

J. Biol.

[105402]Chem. 268: 18748–18753, 1993.

[105403]18479.

[105404]Ye, L.; Nakura, J.; Mitsuda, N.; Fujioka, Y.; Kamino, K.;

Ohta,

[105405]T.; Jinno, Y.; Niikawa, N.; Miki, T.; Ogihara, T.: Genetic as-

sociation

[105406]between chromosome 8 microsatellite (MS8-134) and
Werner syndrome

[105407](WRN): chromosome microdissection and homozygosity
mapping. Genomics 28:

[105408]566-569, 1995.

[105409]18480.

[105410]Yu, C.-E.; Oshima, J.; Fu, Y.-H.; Wijsman, E. M.; Hisama,
F.;

[105411]Alisch, R.; Matthews, S.; Nakura, J.; Miki, T.; Quais, S.;
Martin,

[105412]G. M.; Mulligan, J.; Schellenberg, G. D.: Positional cloning
of the

[105413]Werner's syndrome gene. Science 272: 258-262, 1996.

[105414]18481.

[105415]Yu, C.-E.; Oshima, J.; Goddard, K. A. B.; Miki, T.; Nakura,
J.;

[105416]Ogihara, T.; Poot, M.; Hoehn, H.; Fraccaro, M.; Piussan, C.;
Martin,

[105417]G. M.; Schellenberg, G. D.; Wijsman, E. M.: Linkage dise-
quilibrium

[105418]and haplotype studies of chromosome 8p11.1-21.1
markers and Werner

- [105419] syndrome. Am. J. Hum. Genet. 55: 356–364, 1994.
- [105420] 18482.
- [105421] Yu, C.-E.; Oshima, J.; Wijsman, E. M.; Nakura, J.; Miki, T.;
Piussan,
[105422] C.; Matthews, S.; Fu, Y.-H.; Mulligan, J.; Martin, G. M.;
Schellenberg,
[105423] G. D.; Werner's Syndrome Collaborative Group: Mutations
in the
[105424] consensus helicase domains of the Werner syndrome
gene. Am. J. Hum.
[105425] Genet. 60: 330–341, 1997.
- [105426] 18483.
- [105427] Gibbons, R. J.; Brueton, L.; Buckle, V. J.; Burn, J.; Clayton-
Smith,
[105428] J.; Davison, B. C. C.; Gardner, R. J. M.; Homfray, T.; Kear-
ney, L.;
[105429] Kingston, H. M.; Newbury-Ecob, R.; Porteous, M. E. P.;
Wilkie, A.
[105430] O. M.; Higgs, D. R.: Clinical and hematologic aspects of
the X-linked
[105431] alpha-thalassemia/mental retardation syndrome (ATR-X).
Am. J. Med.

[105432]Genet. 55: 288–299, 1995.

[105433]18484.

[105434]Marchese, A.; Sawzdargo, M.; Nguyen, T.; Cheng, R.;
Heng, H. H.

[105435]Q.; Nowak, T.; Im, D-S.; Lynch, K. R.; George, S. R.;
O'Dowd, B. F.

[105436]: Discovery of three novel orphan G-protein-coupled re-
ceptors. Genomics 56:

[105437]12–21, 1999.

[105438]18485.

[105439]Sanchis, D.; Fleury, C.; Chomiki, N.; Gubern, M.; Huang,
Q.; Neverova,

[105440]M.; Gregoire, F.; Easlick, J.; Raimbault, S.; Levi-Meyrueis,
C.; Miroux,

[105441]B.; Collins, S.; Seldin, M.; Richard, D.; Warden, C.; Bouil-
laud, F.;

[105442]Ricquier, D.: BMCP1, a novel mitochondrial carrier with
high expression

[105443]in the central nervous system of humans and rodents, and
respiration

[105444]uncoupling activity in recombinant yeast. J. Biol. Chem.
273: 34611–34615,

[105445]1998.

[105446]18486.

[105447]Blanco, P.; Sargent, C. A.; Boucher, C. A.; Mitchell, M.; Af-
fara,

[105448]N. A.: Conservation of PCDHX in mammals; expression of
X/Y genes

[105449]predominantly in brain. Mammalian Genome 11: 906–914,
2000.

[105450]18487.

[105451]Ciccodicola, A.; D'Esposito, M.; Esposito, T.; Gian-
francesco, F.;

[105452]Migliaccio, C.; Miano, M. G.; Matarazzo, M. R.; Vacca, M.;
Franze,

[105453]A.; Cuccurese, M.; Cocchia, M.; Curci, A.; and 9 others:
Differentially

[105454]regulated and evolved genes in the fully sequenced Xq/Yq
pseudoautosomal

[105455]region. Hum. Molec. Genet. 9: 395–401, 2000.

[105456]18488.

[105457]Mumm, S.; Molini, B.; Terrell, J.; Srivastava, A.; Sch-
lessinger,

[105458]D.: Evolutionary features of the 4-Mb Xq21.3 XY homology region revealed
[105459]by a map at 60-kb resolution. *Genome Res.* 7: 307–314, 1997.
[105460]18489.
[105461]Schwartz, A.; Chan, D. C.; Brown, L. G.; Alagappan, R.; Pettay,
[105462]D.; Disteche, C.; McGillivray, B.; de la Chapelle, A.; Page, D. C.
[105463]: Reconstructing hominid Y evolution: X-homologous block, created
[105464]by X–Y transposition, was disrupted by Yp inversion through LINE
[105465]recombination. *Hum. Molec. Genet.* 7: 1–11, 1998.
[105466]18490.
[105467]Tilford, C. A.; Kuroda-Kawaguchi, T.; Skaletsky, H.; Rozen, S.;
[105468]Brown, L. G.; Rosenberg, M.; McPherson, J. D.; Wylie, K.; Sekhon,
[105469]M.; Kucaba, T. A.; Waterston, R. H.; Page, D. C.: A physical map
[105470]of the human Y chromosome. *Nature* 409: 943–945,

2001.

[105471]18491.

[105472]des Portes, V.; Pinard, J. M.; Billuart, P.; Vinet, M. C.;

Koulakoff,

[105473]A.; Carrie, A.; Gelot, A.; Dupuis, E.; Motte, J.; Berwald-Net-
ter,

[105474]Y.; Catala, M.; Kahn, A.; Beldjord, C.; Chelly, J.: A novel
CNS gene

[105475]required for neuronal migration and involved in X-linked
subcortical

[105476]laminal heterotopia and lissencephaly syndrome. Cell 92:
51-61,

[105477]1998.

[105478]18492.

[105479]Gleeson, J. G.; Allen, K. M.; Fox, J. W.; Lamperti, E. D.;
Berkovic,

[105480]S.; Scheffer, I.; Cooper, E. C.; Dobyns, W. B.; Minnerath, S.
R.;

[105481]Ross, M. E.; Walsh, C. A.: Doublecortin, a brain-specific
gene mutated

[105482]in human X-linked lissencephaly and double cortex syn-
drome, encodes

[105483]a putative signaling protein. Cell 92: 63–72, 1998.

[105484]18493.

[105485]Matsumoto, N.; Leventer, R. J.; Kuc, J. A.; Mewborn, S. K.;
Dudliceck,

[105486]L. L.; Ramocki, M. B.; Pilz, D. T.; Mills, P. L.; Das, S.; Ross,
M.

[105487]E.; Ledbetter, D. H.; Dobyns, W. B.: Mutation analysis of
the DCX

[105488]gene and genotype/phenotype correlation in subcortical
band heterotopia. Europ.

[105489]J. Hum. Genet. 9: 5–12, 2001.

[105490]18494.

[105491]Batch, J. A.; Evans, B. A. J.; Hughes, I. A.; Patterson, M. N.

[105492]: Mutations of the androgen receptor gene identified in
perineal hypospadias. J.

[105493]Med. Genet. 30: 198–201, 1993.

[105494]18495.

[105495]Gibbons, R. J.; McDowell, T. L.; Raman, S.; O'Rourke, D. M.;
Garrick,

[105496]D.; Ayyub, H.; Higgs, D. R.: Mutations in ATRX, encoding a
SWI/SNF-like

[105497]protein, cause diverse changes in the pattern of DNA methylation. Nature

[105498]Genet. 24: 368–371, 2000.

[105499]18496.

[105500]Gibbons, R. J.; Picketts, D. J.; Villard, L.; Higgs, D. R.: Mutations

[105501]in a putative global transcriptional regulator cause X-linked mental

[105502]retardation with alpha-thalassemia (ATR-X syndrome). Cell 80: 837–845,

[105503]1995.

[105504]18497.

[105505]Kiesewetter, S.; Macek, M., Jr.; Davis, C.; Curristin, S. M.;

[105506]Chu, C.-S.; Graham, C.; Shrimpton, A. E.; Cashman, S. M.; Tsui, L.-C.;

[105507]Mickle, J.; Amos, J.; Highsmith, W. E.; Shuber, A.; Witt, D. R.; Crystal,

[105508]R. G.; Cutting, G. R.: A mutation in CFTR produces different phenotypes

[105509]depending on chromosomal background. Nature Genet. 5: 274–278, 1993.

[105510]18498.

[105511]Lossi, A. M.; Millan, J. M.; Villard, L.; Orellana, C.; Cardoso,

[105512]C.; Prieto, F.; Fontes, M.; Martinez, F.: Mutation of the
XNP/ATR-X

[105513]gene in a family with severe mental retardation, spastic
paraplegia

[105514]and skewed pattern of X inactivation: demonstration that
the mutation

[105515]is involved in the inactivation bias. (Letter) Am. J. Hum.
Genet. 65:

[105516]558-562, 1999.

[105517]18499.

[105518]Martinez, F.; Tomas, M.; Millan, J. M.; Fernandez, A.; Palau,

[105519]F.; Prieto, F.: Genetic localisation of mental retardation
with spastic

[105520]diplegia to the pericentromeric region of the X chromo-
some: X inactivation

[105521]in female carriers. J. Med. Genet. 35: 284-287, 1998.

[105522]18500.

[105523]Mattei, J. F.; Collignon, P.; Ayme, S.; Giraud, F.: X-linked

[105524]mental retardation, growth retardation, deafness and microgenitalism:

[105525]a second familial report. Clin. Genet. 23: 70-74, 1983.

[105526]18501.

[105527]McDowell, T. L.; Gibbons, R. J.; Sutherland, H.; O'Rourke, D.

[105528]M.; Bickmore, W. A.; Pombo, A.; Turley, H.; Gatter, K.; Picketts,

[105529]D. J.; Buckle, V. J.; Chapman, L.; Rhodes, D.; Higgs, D. R.: Localization

[105530]of a putative transcriptional regulator (ATRX) at pericentromeric

[105531]heterochromatin and the short arms of acrocentric chromosomes. Proc.

[105532]Nat. Acad. Sci. 96: 13983-13988, 1999.

[105533]18502.

[105534]Pask, A.; Renfree, M. B.; Graves, J. A. M.: The human sex-reversing

[105535]ATRX gene has a homologue on the marsupial Y chromosome, ATRY: implications

[105536]for the evolution of mammalian sex determination. Proc. Nat. Acad.

[105537]Sci. 97: 13198–13202, 2000.

[105538]18503.

[105539]Picketts, D. J.; Higgs, D. R.; Bachoo, S.; Blake, D. J.; Quarrell,

[105540]O. W. J.; Gibbons, R. J.: ATRX encodes a novel member of the SNF2

[105541]family of proteins: mutations point to a common mechanism underlying

[105542]the ATR-X syndrome. Hum. Molec. Genet. 5: 1899–1907, 1996.

[105543]18504.

[105544]Picketts, D. J.; Tastan, A. O.; Higgs, D. R.; Gibbons, R. J.:

[105545]Comparison of the human and murine ATRX gene identifies highly conserved,

[105546]functionally important domains. Mammalian Genome 9: 400–403, 1998.

[105547]18505.

[105548]Reardon, W.; Gibbons, R. J.; Winter, R. M.; Baraitser, M.: Male

[105549]pseudohermaphroditism in sibs with the alpha-thalassemia/mental retardation

- [105550](ATR-X) syndrome. Am. J. Med. Genet. 55: 285–287, 1995.
- [105551]18506.
- [105552]Stayton, C. L.; Dabovic, B.; Gulisano, M.; Gecz, J.; Broccoli, V.; Giovanazzi, S.; Bossolasco, M.; Monaco, L.; Rastan, S.; Boncinelli, E.; Bianchi, M. E.; Consalez, G. G.: Cloning and characterization of a new human Xq13 gene, encoding a putative helicase. Hum. Molec. Genet. 3: 1957–1964, 1994.
- [105556]18507.
- [105558]Villard, L.; Bonino, M.-C.; Abidi, F.; Ragusa, A.; Belougne, J.; Lossi, A.-M.; Seaver, L.; Bonnefont, J.-P.; Romano, C.; Fichera, M.; Lacombe, D.; Hanauer, A.; Philip, N.; Schwartz, C.; Fontes, M.: Evaluation of a mutation screening strategy for sporadic cases of ATR-X syndrome. J. Med. Genet. 36: 183–186, 1999.

[105563]18508.

[105564]Villard, L.; Fontes, M.; Ades, L. C.; Gecz, J.: Identification
[105565]of a mutation in the XNP/ATR-X gene in a family reported
as Smith-Fineman-Myers
[105566]syndrome. (Letter) Am. J. Med. Genet. 91: 83-85, 2000.

[105567]18509.

[105568]Villard, L.; Gecz, J.; Mattei, J. F.; Fontes, M.; Saugier-Weber,
[105569]P.; Munnich, A.; Lyonnet, S.: XNP mutation in a large fam-
ily with
[105570]Juberg-Marsidi syndrome. (Letter) Nature Genet. 12:
359-360, 1996.

[105571]18510.

[105572]Villard, L.; Lacombe, D.; Fontes, M.: A point mutation in
the
[105573]XNP gene, associated with an ATR-X phenotype without
alpha-thalassemia. Europ.

[105574]J. Hum. Genet. 4: 316-320, 1996.

[105575]18511.

[105576]Villard, L.; Lossi, A.-M.; Cardoso, C.; Proud, V.; Chiaroni,
P.;

[105577]Colleaux, L.; Schwartz, C.; Fontes, M.: Determination of the genomic structure of the XNP/ATRX gene encoding a potential zinc finger helicase. Genomics 43: 149–155, 1997.

[105580]18512.

[105581]Villard, L.; Toutain, A.; Lossi, A.-M.; Gecz, J.; Houdayer, C.; Moraine, C.; Fontes, M.: Splicing mutation in the ATR-X gene can lead to a dysmorphic mental retardation phenotype without alpha-thalassemia. Am. J. Hum. Genet. 58: 499–505, 1996.

[105585]18513.

[105586]Wada, T.; Kubota, T.; Fukushima, Y.; Saitoh, S.: Molecular genetic study of Japanese patients with X-linked alpha-thalassemia/mental retardation syndrome (ATR-X). Am. J. Med. Genet. 94: 242–248, 2000.

[105589]18514.

[105590]Wang, L. H.; Collins, A.; Lawrence, S.; Keats, B. J.; Morton,

[105591]N. E.: Integration of gene maps: chromosome X. Genomics
22: 590-604,

[105592]1994.

[105593]18515.

[105594]Yntema, H. G.; Poppelaars, F. A.; Derksen, E.; Oudakker, A.
R.;

[105595]van Roosmalen, T.; Jacobs, A.; Obbema, H.; Brunner, H. G.;
Hamel,

[105596]B. C. J.; van Bokhoven, H.: Expanding phenotype of XNP
mutations:

[105597]mild to moderate mental retardation. Am. J. Med. Genet.
110: 243-247,

[105598]2002.

[105599]18516.

[105600]Borkhardt, A.; Repp, R.; Haas, O. A.; Leis, T.; Harbott, J.;
Kreuder,

[105601]J.; Hammermann, J.; Henn, T.; Lampert, F.: Cloning and
characterization

[105602]of AFX, the gene that fuses to MLL in acute leukemias with
a t(X;11)(q13;q23). Oncogene 14:

[105603]195-202, 1997.

[105604]18517.

[105605]Katz, F.; Hinshelwood, S.; Rutland, P.; Jones, A.; Kinnon, C.;

[105606]Morgan, G.: Mutation analysis in CD40 ligand deficiency leading to

[105607]X-linked hypogammaglobulinemia with hyper IgM syndrome. Hum. Mutat. 8:

[105608]223–228, 1996.

[105609]18518.

[105610]Kolvraa, S.; Kruse, T. A.; Jensen, P. K. A.; Linde, K. H.; Vestergaard,

[105611]S. R.; Bolund, L.: Close linkage between X-linked ectodermal dysplasia

[105612]and a cloned DNA sequence detecting a two allele restriction fragment

[105613]length polymorphism in the region Xp11–q12. Hum. Genet. 74: 284–287,

[105614]1986.

[105615]18519.

[105616]Korthauer, U.; Graf, D.; Mages, H. W.; Briere, F.; Padayachee,

- [105617]M.; Malcolm, S.; Ugazio, A. G.; Notarangelo, L. D.; Levin-sky, R. J.;
- [105618]Kroczek, R. A.: Defective expression of T-cell CD40 ligand causes
- [105619]X-linked immunodeficiency with hyper-IgM. Nature 361: 539-541, 1993.
- [105620]18520.
- [105621]Kraakman, M. E. M.; de Weers, M.; Espanol, T.; Schuurman, R. K.
- [105622]B.; Hendriks, R. W.: Identification of a CD40L gene mutation and
- [105623]genetic counselling in a family with immunodeficiency with hyperimmunoglobulinemia
- [105624]M. Clin. Genet. 48: 46-48, 1995.
- [105625]18521.
- [105626]Kroczek, R. A.; Graf, D.; Brugnani, D.; Giliani, S.; Korthauer,
- [105627]U.; Ugazio, A.; Senger, G.; Mages, H. W.; Villa, A.; Notarangelo,
- [105628]L. D.: Defective expression of CD40 ligand on T cell causes 'X-linked
- [105629]immunodeficiency with hyper-IgM (HIGM1)'. Immun. Rev.

138: 39–59,

[105630]1994.

[105631]18522.

[105632]Lin, Q.; Rohrer, J.; Allen, R. C.; Larche, M.; Greene, J. M.;

[105633]Shigeoka, A. O.; Gatti, R. A.; Derauf, D. C.; Belmont, J. W.;

Conley,

[105634]M. E.: A single strand conformation polymorphism study
of CD40 ligand:

[105635]efficient mutation analysis and carrier detection of X-
linked hyper

[105636]IgM syndrome. J. Clin. Invest. 97: 196–201, 1996.

[105637]18523.

[105638]Nonoyama, S.; Shimadzu, M.; Toru, H.; Seyama, K.; Nunoi,
H.; Neubauer,

[105639]M.; Yata, J.; Och, H. D.: Mutations of the CD40 ligand gene
in 13

[105640]Japanese patients with X-linked hyper-IgM syndrome.
Hum. Genet. 99:

[105641]624–627, 1997.

[105642]18524.

[105643]Padayachee, M.; Feighery, C.; Finn, A.; McKeown, C.;

Levinsky,

[105644]R. J.; Kinnon, C.; Malcolm, S.: Mapping of the X-linked
form of hyper-IgM

[105645]syndrome (HIGM1) to Xq26 by close linkage to HPRT. Ge-
nomics 14:

[105646]551-553, 1992.

[105647]18525.

[105648]Macchi, P.; Villa, A.; Strina, D.; Sacco, M. G.; Morali, F.;
Brugnoni,

[105649]D.; Giliani, S.; Mantuano, E.; Fasth, A.; Andersson, B.;
Zegers, B.

[105650]J. M.; Cavagni, G.; Reznick, I.; Levy, J.; Zan-Bar, I.; Porat,
Y.;

[105651]Airo, P.; Plebani, A.; Vezzoni, P.; Notarangelo, L. D.: Char-
acterization

[105652]of nine novel mutations in the CD40 ligand gene in pa-
tients with X-linked

[105653]hyper IgM syndrome of various ancestry. Am. J. Hum.
Genet. 56: 898-906,

[105654]1995.

[105655]18526.

[105656]Padayachee, M.; Levinsky, R. J.; Kinnon, C.; Finn, A.; McKe-

own,

[105657]C.; Feighery, C.; Notarangelo, L. D.; Hendriks, R. W.; Read, A. P.;

[105658]Malcolm, S.: Mapping of the X linked form of hyper IgM syndrome (HIGM1). J.

[105659]Med. Genet. 30: 202–205, 1993.

[105660]18527.

[105661]Pilia, G.; Porta, B.; Padayachee, M.; Malcolm, S.; Zucchi, I.;

[105662]Villa, A.; Macchi, P.; Vezzoni, P.; Schlessinger, D.: Human CD40L

[105663]gene maps between DXS144E and DXS300 in Xq26. Genomics 22: 249–251,

[105664]1994.

[105665]18528.

[105666]Ramesh, N.; Fuleihan, R.; Geha, R.: Molecular pathology of X-linked

[105667]immunoglobulin deficiency with normal or elevated IgM (HIGMX-1). Immun.

[105668]Rev. 138: 87–104, 1994.

[105669]18529.

[105670]Seyama, K.; Kira, S.; Ishidoh, K.; Souma, S.; Miyakawa, T.;

Kominami,

[105671]E.: Genomic structure and PCR-SSCP analysis of the human CD40 ligand

[105672]gene: its application to prenatal screening for X-linked hyper-IgM

[105673]syndrome. Human Genet. 97: 180-185, 1996.

[105674]18530.

[105675]Seyama, K.; Nonoyama, S.; Gangsaas, I.; Hollenbaugh, D.; Pabst,

[105676]H. F.; Aruffo, A.; Ochs, H. D.: Mutations of the CD40 ligand and gene

[105677]and its effect on CD40 ligand expression in patients with X-linked

[105678]hyper IgM syndrome. Blood 92: 2421-2434, 1998.

[105679]18531.

[105680]Tan, J.; Town, T.; Crawford, F.; Mori, T.; DelleDonne, A.; Crescentini,

[105681]R.; Obregon, D.; Flavell, R. A.; Mullan, M. J.: Role of CD40 ligand

[105682]in amyloidosis in transgenic Alzheimer's mice. Nature Neurosci. 28Oct,

[105683]2002. Note: Advance Electronic Publication.

[105684]18532.

[105685]Chatterjee, A.; Faust, C. J.; Herman, G. E.: Genetic and physical

[105686]mapping of the biglycan gene on the mouse X chromosome. Mammalian

[105687]Genome 4: 33–36, 1993.

[105688]18533.

[105689]Das, S.; Metzenberg, A.; Pai, G. S.; Gitschier, J.: Mutational

[105690]analysis of the biglycan gene excludes it as a candidate for X-linked

[105691]dominant chondrodysplasia punctata, dyskeratosis congenita, and incontinentia

[105692]pigmenti. (Letter) Am. J. Hum. Genet. 54: 922–925, 1994.

[105693]18534.

[105694]Fisher, L. W.; Heegaard, A.-M.; Vetter, U.; Vogel, W.; Just, W.;

[105695]Termine, J. D.; Young, M. F.: Human biglycan gene: putative promoter,

[105696]intron–exon junctions, and chromosomal localization. J. Biol. Chem. 266:

[105697]14371–14377, 1991.

[105698]18535.

[105699]Fisher, L. W.; Termine, J. D.; Young, M. F.: Deduced-protein sequence

[105700]of bone small proteoglycan I (biglycan) shows homology with proteoglycan

[105701]II (decorin) and several nonconnective tissue proteins in a variety

[105702]of species. J. Biol. Chem. 264: 4571-4576, 1989.

[105703]18536.

[105704]Geerkens, C.; Vetter, U.; Just, W.; Fedarko, N. S.; Fisher, L.

[105705]W.; Young, M. F.; Termine, J. D.; Gehron Robey, P.; Wohrle, D.; Vogel,

[105706]W.: The X-chromosomal human biglycan gene BGN is subject to X inactivation

[105707]but is transcribed like an X-Y homologous gene. Hum. Genet. 96:

[105708]44-52, 1995.

[105709]18537.

[105710]McBride, O. W.; Fisher, L. W.; Young, M. F.: Localization of PGI

[105711](biglycan, BGN) and PGII (decorin, DCN, PG-40) genes on

human chromosomes

[105712]Xq13-qter and 12q, respectively. Genomics 6: 219-225,
1990.

[105713]18538.

[105714]Traupe, H.; van den Ouweland, A. M. W.; van Oost, B. A.;
Vogel,

[105715]W.; Vetter, U.; Warren, S. T.; Rocchi, M.; Darlison, M. G.;
Ropers,

[105716]H.-H.: Fine mapping of the human biglycan (BGN) gene
within the Xq28

[105717]region employing a hybrid cell panel. Genomics 13:
481-483, 1992.

[105718]18539.

[105719]Wegrowski, Y.; Pillarisetti, J.; Danielson, K. G.; Suzuki, S.;

[105720]Iozzo, R. V.: The murine biglycan: complete cDNA cloning,
genomic

[105721]organization, promoter function, and expression. Ge-
nomics 30: 8-17,

[105722]1995.

[105723]18540.

[105724]Xu, T.; Bianco, P.; Fisher, L. W.; Longenecker, G.; Smith, E.;

[105725]Goldstein, S.; Bonadio, J.; Boskey, A.; Heegaard, A.-M.;
Sommer, B.;

[105726]Satomura, K.; Dominguez, P.; Zhao, C.; Kulkarni, A. B.;
Robey, P.

[105727]G.; Young, M. F.: Targeted disruption of the biglycan gene
leads

[105728]to an osteoporosis-like phenotype in mice. Nature Genet.
20: 78-82,

[105729]1998.

[105730]18541.

[105731]Sandhoff, K.; Harzer, K.; Wasse, W.; Jatzkewitz, H.: En-
zyme

[105732]alterations and lipid storage in three variants of Tay-Sachs
disease. J.

[105733]Neurochem. 18: 2469-2489, 1971.

[105734]18542.

[105735]Bowcock, A. M.; Farrer, L. A.; Hebert, J. M.; Agger, M.;
Sternlieb,

[105736]I.; Scheinberg, I. H.; Buys, C. H. C. M.; Scheffer, H.; Fryd-
man, M.;

[105737]Chajek-Saul, T.; Bonne-Tamir, B.; Cavalli-Sforza, L. L.:
Eight closely

[105738]linked loci place the Wilson disease locus within
13q14-q21. Am.

[105739]J. Hum. Genet. 43: 664-674, 1988.

[105740]18543.

[105741]Yoshida, K.; Sugano, S.: Identification of a novel protocad-
herin

[105742]gene (PCDH11) on the human XY homology region in
Xq21.3. Genomics 62:

[105743]540-543, 1999.

[105744]18544.

[105745]de la Chapelle, A.; Sankila, E.-M.; Lindlof, M.; Aula, P.; No-
rio,

[105746]R.: Norrie disease caused by a gene deletion allowing car-
rier detection

[105747]and prenatal diagnosis. Clin. Genet. 28: 317-320, 1985.

[105748]18545.

[105749]Denney, R. M.; Fritz, R. R.; Patel, N. T.; Abell, C. W.: Hu-
man

[105750]liver MAO-A and MAO-B separated by immunoaffinity
chromatography with

[105751]MAO-B-specific monoclonal antibody. Science 215:

1400–1403, 1982.

[105752]18546.

[105753]Kochersperger, L. M.; Parker, E. L.; Siciliano, M.; Darling-
ton,

[105754]G. J.; Denney, R. M.: Assignment of genes for human
monoamine oxidases

[105755]A and B to the X chromosome. J. Neurosci. Res. 16:
601–616, 1986.

[105756]18547.

[105757]Sims, K. B.; de la Chapelle, A.; Norio, R.; Sankila, E.–M.;
Hsu,

[105758]Y.–P. P.; Rinehart, W. B.; Corey, T. J.; Ozelius, L.; Powell, J.
F.;

[105759]Bruns, G.; Gusella, J. F.; Murphy, D. L.; Breakefield, X. O.:
Monoamine

[105760]oxidase deficiency in males with an X chromosome dele-
tion. Neuron 2:

[105761]1069–1076, 1989.

[105762]18548.

[105763]Binda, C.; Newton–Vinson, P.; Hubalek, F.; Edmondson, D.
E.; Mattevi,

[105764]A.: Structure of human monoamine oxidase B, a drug tar-

get for the

[105765]treatment of neurological disorders. Nature Struct. Biol. 9:
22–26,

[105766]2002.

[105767]18549.

[105768]Checkoway, H.; Franklin, G. M.; Costa-Mallen, P.; Smith-
Weller,

[105769]T.; Dilley, J.; Swanson, P. D.; Costa, L. G.: A genetic poly-
morphism

[105770]of MAO-B modifies the association of cigarette smoking
and Parkinson's

[105771]disease. Neurology 50: 1458–1461, 1998.

[105772]18550.

[105773]Fowler, J. S.; Volkow, N. D.; Wang, G.-J.; Pappas, N.; Lo-
gan, J.;

[105774]MacGregor, R.; Alexoff, D.; Shea, C.; Schlyer, D.; Wolf, A.
P.; Warner,

[105775]D.; Zezulkova, I.; Cilento, R.: Inhibition of monoamine oxi-
dase B

[105776]in the brains of smokers. Nature 379: 733–736, 1996.

[105777]18551.

- [105778] Hamano, Y.; Kodama, H.; Fujikawa, Y.; Tanaka, Y.;
Nishimura, K.;
- [105779] Yanagisawa, M.: Use of immunocytochemical analysis of a
duodenal
- [105780] biopsy specimen to identify a carrier of ornithine tran-
scarbamylase
- [105781] deficiency. New Eng. J. Med. 318: 1521-1523, 1988.
- [105782] 18552.
- [105783] Bull, P. C.; Thomas, G. R.; Rommens, J. M.; Forbes, J. R.;
- Cox,
- [105784] D. W.: The Wilson disease gene is a putative copper trans-
porting
- [105785] P-type ATPase similar to the Menkes gene. Nature Genet.
5: 327-337,
- [105786] 1993.
- [105787] 18553.
- [105788] Forbes, J. R.; Cox, D. W.: Copper-dependent trafficking of
Wilson
- [105789] disease mutant ATP7B proteins. Hum. Molec. Genet. 9:
1927-1935,
- [105790] 2000.

[105791]18554.

[105792]Ha-Hao, D.; Hefter, H.; Stremmel, W.; Castaneda-Guillot, C.; Hernandez,

[105793]A. H.; Cox, D. W.; Auburger, G.: His1069-to-gln and six novel Wilson

[105794]disease mutations: analysis of relevance for early diagnosis and phenotype. Europ.

[105795]J. Hum. Genet. 6: 616-623, 1998.

[105796]18555.

[105797]Grino, P. B.; Griffin, J. E.; Cushard, W. G., Jr.; Wilson, J.

[105798]D.: A mutation of the androgen receptor associated with partial androgen

[105799]resistance, familial gynecomastia, and fertility. J. Clin. Endocr.

[105800]Metab. 66: 754-761, 1988.

[105801]18556.

[105802]Holterhus, P.-M.; Sinnecker, G. H. G.; Hiort, O.: Phenotypic

[105803]diversity and testosterone-induced normalization of mutant L712F androgen

[105804]receptor function in a kindred with androgen insensitivity. J. Clin.

[105805]Endocr. Metab. 85: 3245–3250, 2000.

[105806]18557.

[105807]Hughes, I. A.; Evans, B. A. J.: Complete androgen insensitivity

[105808]syndrome characterized by increased concentration of a normal androgen

[105809]receptor in genital skin fibroblasts. J. Clin. Endocr. Metab. 63:

[105810]309–315, 1986.

[105811]18558.

[105812]Imperato-McGinley, J.; Ip, N. Y.; Gautier, T.; Neuweiler, J.;

[105813]Gruenspan, H.; Liao, S.; Chang, C.; Balazs, I.: DNA linkage analysis

[105814]and studies of the androgen receptor gene in a large kindred with

[105815]complete androgen insensitivity. Am. J. Med. Genet. 36: 104–108,

[105816]1990.

[105817]18559.

[105818]Jukier, L.; Kaufman, M.; Pinsky, L.; Peterson, R. E.: Partial

- [105819] androgen resistance associated with secondary
5-alpha-reductase deficiency:
- [105820] identification of a novel qualitative androgen receptor de-
fect and
- [105821] clinical implications. J. Clin. Endocr. Metab. 59: 679-688,
1984.
- [105822] 18560.
- [105823] Kaufman, M.; Pinsky, L.; Bowin, A.; Au, M. W. S.: Familial
external
- [105824] genital ambiguity due to a transformation defect of an-
drogen-receptor
- [105825] complexes that is expressed with
- 5-alpha-dihydrotestosterone and the
- [105826] synthetic androgen methyltrienolone. Am. J. Med. Genet.
18: 493-507,
- [105827] 1984.
- [105828] 18561.
- [105829] Kaufman, M.; Straisfeld, C.; Pinsky, L.: Male pseudo-
hermaphroditism
- [105830] presumably due to target organ unresponsiveness to an-
drogens: deficient
- [105831] 5-alpha-dihydrotestosterone binding in cultured skin fi-

broblasts. J.

[105832]Clin. Invest. 58: 345-350, 1976.

[105833]18562.

[105834]Keenan, B. S.; Meyer, W. J., III; Hadjian, A. J.; Jones, H. W.;

[105835]Migeon, C. J.: Syndrome of androgen insensitivity in man:
absence

[105836]of 5- α -dihydrotestosterone binding protein in skin fi-
broblasts. J.

[105837]Clin. Endocr. 38: 1143-1146, 1974.

[105838]18563.

[105839]Liao, S.; Witte, D.: Autoimmune anti-androgen-receptor
antibodies

[105840]in human serum. Proc. Nat. Acad. Sci. 82: 8345-8348,
1985.

[105841]18564.

[105842]Lin, S.-Y.; Ohno, S.: The binding of androgen receptor to
DNA

[105843]and RNA. Biochim. Biophys. Acta 654: 181-186, 1981.

[105844]18565.

[105845]Aradhya, S.; Bardaro, T.; Galgoczy, P.; Yamagata, T.; Es-

posito,

[105846]T.; Patlan, H.; Ciccodicola, A.; Munnich, A.; Kenwrick, S.;

Platzer,

[105847]M.; D'Urso, M.; Nelson, D. L.: Multiple pathogenic and be-

nign genomic

[105848]rearrangements occur at a 35 kb duplication involving the

NEMO and

[105849]LAGE2 genes. Hum. Molec. Genet. 10: 2557-2567, 2001.

[105850]18566.

[105851]Aradhya, S.; Courtois, G.; Rajkovic, A.; Lewis, R. A.; Levy,

M.;

[105852]Israel, A.; Nelson, D. L.: Atypical forms of incontinentia

pigmenti

[105853]in male individuals result from mutations of a cytosine

tract in exon

[105854]10 of NEMO (IKK-gamma). Am. J. Hum. Genet. 68:

765-771, 2001.

[105855]18567.

[105856]Aradhya, S.; Woffendin, H.; Jakins, T.; Bardaro, T.; Espos-

ito,

[105857]T.; Smahi, A.; Shaw, C.; Levy, M.; Munnich, A.; D'Urso, M.;

Lewis,

[105858]R. A.; Kenwrick, S.; Nelson, D. L.: A recurrent deletion in
the ubiquitously
[105859]expressed NEMO (IKK- γ) gene accounts for the vast
majority of
[105860]incontinentia pigmenti mutations. Hum. Molec. Genet. 10:
2171–2179,
[105861]2001.
[105862]18568.
[105863]Doffinger, R.; Smahi, A.; Bessia, C.; Geissmann, F.; Fein-
berg,
[105864]J.; Durandy, A.; Bodemer, C.; Kenwrick, S.; Dupuis-Girod,
S.; Blanche,
[105865]S.; Wood, P.; Rabia, S. H.; and 16 others: X-linked an-
hidrotic
[105866]ectodermal dysplasia with immunodeficiency is caused by
impaired NF- κ B
[105867]signaling. Nature Genet. 27: 277–285, 2001.
[105868]18569.
[105869]Jain, A.; Ma, C. A.; Liu, S.; Brown, M.; Cohen, J.; Strober, W.
[105870]: Specific missense mutations in NEMO result in hyper-IgM
syndrome

[105871]with hypohydrotic (sic) ectodermal dysplasia. Nature Immun. 2: 223–228,

[105872]2001.

[105873]18570.

[105874]Jin, D. Y.; Jeang, K. T.: Isolation of full-length cDNA and chromosomal

[105875]localization of human NF- κ B modulator NEMO to Xq28. J. Biomed.

[105876]Sci. 6: 115–120, 1999.

[105877]18571.

[105878]Kosaki, K.; Shimasaki, N.; Fukushima, H.; Hara, M.; Ogata, T.;

[105879]Matsuo, N.: Female patient showing hypohidrotic ectodermal dysplasia

[105880]and immunodeficiency (HED-ID). (Letter) Am. J. Hum. Genet. 69: 664–665,

[105881]2001.

[105882]18572.

[105883]Li, Q.; Van Antwerp, D.; Mercurio, F.; Lee, K.-F.; Verma, I. M.

[105884]: Severe liver degeneration in mice lacking the I- κ B

kinase 2

[105885]gene. Science 284: 321–325, 1999.

[105886]18573.

[105887]Li, Y.; Kang, J.; Friedman, J.; Tarassishin, L.; Ye, J.; Kovalenko,

[105888]A.; Wallach, D.; Horwitz, M. S.: Identification of a cell protein

[105889](FIP-3) as a modulator of NF-kappa-B activity and as a target of an

[105890]adenovirus inhibitor of tumor necrosis factor alpha-induced apoptosis. Proc.

[105891]Nat. Acad. Sci. 96: 1042–1047, 1999.

[105892]18574.

[105893]Makris, C.; Godfrey, V. L.; Krahn-Senftleben, G.; Takahashi, T.;

[105894]Roberts, J. L.; Schwarz, T.; Feng, L.; Johnson, R. S.; Karin, M.:

[105895]Female mice heterozygous for IKK-gamma/NEMO deficiencies develop a

[105896]dermatopathy similar to the human X-linked disorder incontinentia

[105897]pigmenti. Molec. Cell 5: 969–979, 2000.

[105898]18575.

[105899]May, M. J.; D'Acquisto, F.; Madge, L. A.; Glockner, J.;
Poher,

[105900]J. S.; Ghosh, S.: Selective inhibition of NF-kappa-B activa-
tion by

[105901]a peptide that blocks the interaction of NEMO with the I-
kappa-B kinase

[105902]complex. Science 289: 1550-1554, 2000.

[105903]18576.

[105904]Roberts, J. L.; Morrow, B.; Vega-Rich, C.; Salafia, C. M.; Ni-
towsky,

[105905]H. M.: Incontinentia pigmenti in a newborn male infant
with DNA confirmation. Am.

[105906]J. Med. Genet. 75: 159-163, 1998.

[105907]18577.

[105908]Rothwarf, D. M.; Zandi, E.; Natoli, G.; Karin, M.: IKK-
gamma

[105909]is an essential regulatory subunit of the I-kappa-B kinase
complex. Nature 395:

[105910]297-300, 1998.

[105911]18578.

[105912]Rudolph, D.; Yeh, W.-C.; Wakeham, A.; Rudolph, B.; Nal-
lainathan,
[105913]D.; Potter, J.; Elia, A. J.; Mak, T. W.: Severe liver degenera-
tion
[105914]and lack of NF-kappa-B activation in NEMO/
IKK-gamma-deficient mice. Genes
[105915]Dev. 14: 854-862, 2000.
[105916]18579.
[105917]Schmidt-Supprian, M.; Bloch, W.; Courtois, G.; Addicks, K.;
Israel,
[105918]A.; Rajewsky, K.; Pasparakis, M.: NEMO/
IKK-gamma-deficient mice model
[105919]incontinentia pigmenti. Molec. Cell 5: 981-992, 2000.
[105920]18580.
[105921]The International Incontinentia Pigmenti Consortium: Ge-
nomic
[105922]rearrangement in NEMO impairs NF-kappa-B activation
and is a cause
[105923]of incontinentia pigmenti. Nature 405: 466-472, 2000.
[105924]18581.
[105925]The International IP Consortium: Survival of male patients

[105926]with incontinentia pigmenti carrying a lethal mutation can
be explained

[105927]by somatic mosaicism or Klinefelter syndrome. Am. J.
Hum. Genet. 69:

[105928]1210–1217, 2001.

[105929]18582.

[105930]Yamaoka, S.; Courtois, G.; Bessia, C.; Whiteside, S. T.;
Weil,

[105931]R.; Agou, F.; Kirk, H. E.; Kay, R. J.; Israel, A.: Complemen-
tation

[105932]cloning of NEMO, a component of the I-kappa-B kinase
complex essential

[105933]for NF-kappa-B activation. Cell 93: 1231–1240, 1998.

[105934]18583.

[105935]Zonana, J.; Elder, M. E.; Schneider, L. C.; Orlow, S. J.; Moss,

[105936]C.; Golabi, M.; Shapira, S. K.; Farndon, P. A.; Wara, D. W.;
Emmal,

[105937]S. A.; Ferguson, B. M.: A novel X-linked disorder of im-
mune deficiency

[105938]and hypohidrotic ectodermal dysplasia is allelic to inconti-
nencia

[105939] pigmenti and due to mutations in IKK-gamma (NEMO).

Am. J. Hum. Genet. 67:

[105940] 1555–1562, 2000.

[105941] 18584.

[105942] Bauer, M. F.; Gempel, K.; Reichert, A. S.; Rappold, G. A.;

Lichtner,

[105943] P.; Gerbitz, K.-D.; Neupert, W.; Brunner, M.; Hofmann, S.:

Genetic

[105944] and structural characterization of the human mitochondrial inner membrane

[105945] translocase. J. Molec. Biol. 289: 69–82, 1999.

[105946] 18585.

[105947] Kruse, T. A.; Kolvraa, S.; Bolund, L.; Kaitila, I.; Soderholm,

[105948] A. L.; de la Chapelle, A.; MacDermot, K. D.: X-linked anhidrotic

[105949] ectodermal dysplasia (EDA): multipoint linkage analysis.

(Abstract) Cytogenet.

[105950] Cell Genet. 51: 1026, 1989.

[105951] 18586.

[105952] Limon, J.; Filipiuk, J.; Nedoszytko, B.; Mrozek, K.; Castren,

[105953]M.; Larramendy, M.; Roszkiewicz, J.: X-linked anhidrotic
ectodermal
[105954]dysplasia and de novo t(X;1) in a female. Hum. Genet. 87:
338–340,
[105955]1991.
[105956]18587.
[105957]MacDermot, K. D.; Hulten, M.: Female with hypohidrotic
ectodermal
[105958]dysplasia and de novo (X;9) translocation: clinical docu-
mentation
[105959]of the AnLy cell line case. Hum. Genet. 84: 577–579,
1990.
[105960]18588.
[105961]MacDermot, K. D.; Malcolm, S.: X linked hypohidrotic ec-
todermal
[105962]dysplasia (CST syndrome): clinical and molecular study
using X chromosome
[105963]markers. (Abstract) J. Med. Genet. 23: 167, 1986.
[105964]18589.
[105965]MacDermot, K. D.; Morgan, S. H.; Cheshire, J. K.; Wilson, T.
M.;

- [105966] Kaitila, I.; Soderholm, L.; de la Chapelle, A.; Kolvraa, S.; Bolund, L.; Kruse, T. A.: X-linked anhidrotic ectodermal dysplasia (EDA): two point and multipoint linkage analysis using 7 polymorphic DNA markers. (Abstract) Cytogenet. Cell Genet. 46: 653-654, 1987.
- [105970] 18590.
- [105971] MacDermot, K. D.; Winter, R. M.; Malcolm, S.: Gene localization of X-linked hypohidrotic ectodermal dysplasia (C-S-T syndrome). Hum. Genet. 74: 172-173, 1986.
- [105974] 18591.
- [105975] Malagon, V.; Taveras, J. E.: Congenital anhidrotic ectodermal and mesodermal dysplasia. Arch. Derm. 74: 253-258, 1956.
- [105977] 18592.
- [105978] Monreal, A. W.; Zonana, J.; Ferguson, B.: Identification of a

- [105979] new splice form of the EDA1 gene permits detection of nearly all X-linked
- [105980] hypohidrotic ectodermal dysplasia mutations. Am. J. Hum. Genet. 63:
- [105981] 380–389, 1998.
- [105982] 18593.
- [105983] Pedrini, V.; Lennzi, L.; Zamtotti, V.: Isolation and identification
- [105984] of keratosulphate in urine of patients affected by Morquio–Ullrich
- [105985] disease. Proc. Soc. Exp. Biol. Med. 110: 847–849, 1962.
- [105986] 18594.
- [105987] Riedner, E. D.; Levin, L. S.: Hearing patterns in Morquio's syndrome
- [105988] (mucopolysaccharidosis IV). Arch. Otolaryng. 103: 518–520, 1977.
- [105989] 18595.
- [105990] Robins, M. M.; Stevens, H. F.; Linker, A.: Morquio's disease:
- [105991] an abnormality of mucopolysaccharide metabolism. J. Pediat. 62:

[105992]881–889, 1963.

[105993]18596.

[105994]Singh, J.; DiFerrante, N. M.; Niebes, P.; Tavella, D.: N-acetylgalactosamine-6-sulfate

[105995]sulfatase in man: absence of the enzyme in Morquio disease. J. Clin.

[105996]Invest. 57: 1036–1040, 1976.

[105997]18597.

[105998]Sukegawa, K.; Nakamura, H.; Kato, Z.; Tomatsu, S.; Montano, A.

[105999]M.; Fukao, T.; Toietta, G.; Tortora, P.; Orii, T.; Kondo, N.: Biochemical

[106000]and structural analysis of missense mutations in N-acetylgalactosamine-6-sulfate

[106001]sulfatase causing mucopolysaccharidosis IVA phenotypes. Hum. Molec.

[106002]Genet. 9: 1283–1290, 2000.

[106003]18598.

[106004]Tomatsu, S.; Fukuda, S.; Cooper, A.; Wraith, J. E.; Ferreira,

[106005]P.; Di Natale, P.; Tortora, P.; Fujimoto, A.; Kato, Z.; Ya-

mada, N.;

[106006]Isogai, K.; Yamagishi, A.; Sukegawa, K.; Suzuki, Y.; Shimozawa, N.;

[106007]Kondo, N.; Sly, W. S.; Orii, T.: Fourteen novel mucopolysaccharidosis

[106008]IVA producing mutations in GALNS gene. Hum. Mutat. 10: 368-375,

[106009]1997.

[106010]18599.

[106011]Tomatsu, S.; Fukuda, S.; Cooper, A.; Wraith, J. E.; Rezvi, G.

[106012]M. M.; Yamagishi, A.; Yamada, N.; Kato, Z.; Isogai, K.; Sukegawa,

[106013]K.; Kondo, N.; Suzuki, Y.; Shimozawa, N.; Orii, T.: Mucopolysaccharidosis

[106014]IVA: identification of a common missense mutation I113F in the N-acetylgalactosamine-6-sulfate

[106015]sulfatase gene. Am. J. Hum. Genet. 57: 556-563, 1995.

[106016]18600.

[106017]Tomatsu, S.; Fukuda, S.; Cooper, A.; Wraith, J. E.; Yamada, N.;

[106018]Isogai, K.; Kato, Z.; Sukegawa, K.; Kondo, N.; Suzuki, Y.;

Shimozawa,

[106019]N.; Orii, T.: Two new mutations, Q473X and N487S, in a
Caucasian

[106020]patient with mucopolysaccharidosis IVA (Morquio disease).
Hum. Mutat. 6:

[106021]195–196, 1995.

[106022]18601.

[106023]Tomatsu, S.; Fukuda, S.; Masue, M.; Sukegawa, K.; Fukao,
T.; Yamagishi,

[106024]A.; Hori, T.; Iwata, H.; Ogawa, T.; Nakashima, Y.; Hanyu,
Y.; Hashimoto,

[106025]T.; Titani, K.; Oyama, R.; Suzuki, M.; Yagi, K.; Hayashi, Y.;
Orii,

[106026]T.: Morquio disease: isolation, characterization and ex-
pression of

[106027]full-length cDNA for human N-acetyl-
galactosamine-6-sulfate sulfatase. Biochem.

[106028]Biophys. Res. Commun. 181: 677–683, 1991.

[106029]18602.

[106030]Tomatsu, S.; Fukuda, S.; Masue, M.; Sukegawa, K.; Ma-
suno, M.;

[106031]Orii, T.: Mucopolysaccharidosis type IVA: characterization

and chromosomal

[106032] localization of N-acetylgalactosamine-6-sulfate sulfatase gene and

[106033] genetic heterogeneity. (Abstract) Am. J. Hum. Genet. 51 (suppl.):

[106034] A178, 1992.

[106035] 18603.

[106036] Tomatsu, S.; Fukuda, S.; Yamagishi, A.; Cooper, A.; Wraith, J.

[106037] E.; Hori, T.; Kato, Z.; Yamada, N.; Isogai, K.; Sukegawa, K.; Kondo,

[106038] N.; Suzuki, Y.; Shimozawa, N.; Orii, T.: Mucopolysacchari-
dosis IVA:

[106039] four new exonic mutations in patients with N-
acetylgalactosamine-6-sulfate

[106040] sulfatase deficiency. Am. J. Hum. Genet. 58: 950-962,
1996.

[106041] 18604.

[106042] Tylki-Szymanska, A.; Czartoryska, B.; Bunge, S.; van
Diggelen,

[106043] O. P.; Kleijer, W. J.; Poorthuis, B. J. H. M.; Huijmans, J. G.
M.;

[106044]Gorska, D.: Clinical, biochemical and molecular findings in
a two-generation

[106045]Morquio A family. Clin. Genet. 53: 369–374, 1998.

[106046]18605.

[106047]Von Noorden, G. K.; Zellweger, H.; Ponseti, I. V.: Ocular
findings

[106048]in Morquio–Ullrich's disease. Arch. Ophthal. 64: 585–591,
1960.

[106049]18606.

[106050]Wang, L.; Ou, X.; Sebesta, I.; Vondrak, K.; Krijt, J.; Elleder,

[106051]M.; Poupetova, H.; Ledvinova, J.; Zeman, J.; Simmonds, H.
A.; Tischfield,

[106052]J. A.; Sahota, A.: Combined adenine phosphoribosyltrans-
ferase and

[106053]N-acetylgalactosamine-6-sulfate sulfatase deficiency.
Molec. Genet.

[106054]Metab. 68: 78–85, 1999.

[106055]18607.

[106056]Wiedemann, H.–R.: The pioneers of pediatric medicine:
Luis Morquio

[106057](1867–1935). Europ. J. Pediat. 151: 549, 1992.

[106058]18608.

[106059]Yamada, N.; Fukuda, S.; Tomatsu, S.; Muller, V.; Hopwood, J. J.;

[106060]Nelson, J.; Kato, Z.; Yamagishi, A.; Sukegawa, K.; Kondo, N.; Orii,

[106061]T.: Molecular heterogeneity in mucopolysaccharidosis IVA in Australia

[106062]and Northern Ireland: nine novel mutations including T312S, a common

[106063]allele that confers a mild phenotype. Hum. Mutat. 11: 202-208, 1998.

[106064]18609.

[106065]Yuen, M.; Fensom, A. H.: Diagnosis of classical Morquio's disease:

[106066]N-acetylgalactosamine 6-sulphate sulphatase activity in cultured fibroblasts,

[106067]leukocytes, amniotic cells and chorionic villi. J. Inherit. Metab.

[106068]Dis. 8: 80-86, 1985.

[106069]18610.

[106070]Zellweger, H.; Ponseti, I. V.; Pedrini, V.; Stamler, F. S.; Von

[106071]Noorden, G. K.: Morquio-Ullrich's disease: report of 2 cases. J.

[106072]Pediat. 59: 549-561, 1961.

[106073]18611.

[106074]Shmerling, D. H.; Prader, A.; Hitzig, W. H.; Giedion, A.; Hadorn,

[106075]B.; Kuhni, M.: The syndrome of exocrine pancreatic insufficiency,

[106076]neutropenia, metaphyseal dysostosis and dwarfism. Helv. Paediat.

[106077]Acta 24: 547-575, 1969.

[106078]18612.

[106079]Moller, D. E.; Xia, C. H.; Tang, W.; Zhu, A. X.; Jakubowski, M.

[106080]; Human rsk isoforms: cloning and characterization of tissue-specific

[106081]expression. Am. J. Physiol. 266: C351-C359, 1994.

[106082]18613.

[106083]Trivier, E.; De Cesare, D.; Jacquot, S.; Pannetier, S.; Zackai,

[106084]E.; Young, I.; Mandel, J.-L.; Sassone-Corsi, P.; Hanauer, A.: Mutations

[106085]in the kinase Rsk-2 associated with Coffin-Lowry syndrome. Nature 384:

[106086]567-570, 1996.

[106087]18614.

[106088]Zhao, Y.; Bjorbaek, C.; Weremowicz, S.; Morton, C. C.; Moller,

[106089]D. E.: RSK3 encodes a novel pp90rsk isoform with a unique N-terminal

[106090]sequence: growth factor-stimulated kinase function and nuclear translocation. Molec.

[106091]Cell. Biol. 15: 4353-4363, 1995.

[106092]18615.

[106093]Liu, Z.; Sun, C.; Olejniczak, E. T.; Meadows, R. P.; Betz, S. F.;

[106094]Oost, T.; Herrmann, J.; Wu, J. C.; Fesik, S. W.: Structural basis

[106095]for binding of Smac/DIABLO to the XIAP BIR3 domain. Nature 408:

[106096]1004-1008, 2000.

[106097]18616.

[106098]Srinivasula, S. M.; Hegde, R.; Saleh, A.; Datta, P.; Shiozaki,

[106099]E.; Chai, J.; Lee, R.-A.; Robbins, P. D.; Fernandes-Alnemri, T.; Shi, Y.; Alnemri, E. S.: A conserved XIAP-interaction motif in caspase-9 and Smac/DIABLO regulates caspase activity and apoptosis. *Nature* 410: 112-116, 2001.

[106102]18617.

[106103]Shwachman, H.; Diamond, L. K.; Oski, F. A.; Khaw, K. T.: The syndrome of pancreatic insufficiency and bone marrow dysfunction. *J. Pediat.* 65: 645-663, 1964.

[106106]18618.

[106107]Shwachman, H.; Holsclaw, D.: Some clinical observations on the Shwachman syndrome (pancreatic insufficiency and bone marrow hypoplasia). *Birth Defects Orig. Art. Ser.* 8(3): 46-49, 1972.

[106110]18619.

[106111]Smith, O. P.; Chan, M. Y.; Evans, J.; Veys, P.: Shwachman-Di-

among

[106113] syndrome and matched unrelated donor BMT. Bone Marrow Transplant. 16:

[106114] 717–718, 1995.

[106115] 18620.

[106116] Smith, O. P.; Hann, I. M.; Chessells, J. M.; Reeves, B. R.; Milla,

[106117] P.: Haematological abnormalities in Shwachman–Diamond syndrome. Brit.

[106118] J. Haemat. 94: 279–284, 1996.

[106119] 18621.

[106120] Tada, H.; Ri, T.; Yoshida, H.; Ishimoto, K.; Kaneko, M.; Yamashiro,

[106121] Y.; Shinohara, T.: A case of Shwachman syndrome with increased spontaneous

[106122] chromosome breakage. Hum. Genet. 77: 289–291, 1987.

[106123] 18622.

[106124] Taybi, H.; Mitchell, A. D.; Friedman, G. D.: Metaphyseal dysostosis

[106125] and associated syndrome of pancreatic insufficiency and blood disorders. Radiology 93:

[106126]563–571, 1969.

[106127]18623.

[106128]Woods, W. G.; Roloff, J. S.; Lukens, J. N.; Krivit, W.: The occurrence

[106129]of leukemia in patients with the Shwachman syndrome. J. Pediat. 99:

[106130]425–428, 1981.

[106131]18624.

[106132]Wulfeck, B.; Trauner, D.; Marsden, D.; Nyhan, W.; Tallal, P.:

[106133]Neurobehavioral profiles in two sisters with Shwachman–Diamond syndrome. Dysmorph.

[106134]Clin. Genet. 5: 15–22, 1991.

[106135]18625.

[106136]Britton, C. H.; Schultz, R. A.; Zhang, B.; Esser, V.; Foster, D.

[106137]W.; McGarry, J. D.: Human liver mitochondrial carnitine palmitoyltransferase

[106138]: characterization of its cDNA and chromosomal localization and partial

[106139]analysis of the gene. Proc. Nat. Acad. Sci. 92: 1984–1988, 1995.

[106140]18626.

[106141]Song, X.-Q.; Fukao, T.; Watanabe, H.; Shintaku, H.; Hirayama,

[106142]K.; Kassovska-Bratinova, S.; Kondo, N.; Mitchell, G. A.: Succinyl-CoA:3-ketoacid

[106143]CoA transferase (SCOT) deficiency: two pathogenic mutations, V133E

[106144]and C456F, in Japanese siblings. Hum. Mutat. 12: 83-88, 1998.

[106145]18627.

[106146]Spence, M. W.; Murphy, M. G.; Cook, H. W.; Ripley, B. A.; Embil,

[106147]J. A.: Succinyl CoA:3-ketoacid CoA transferase deficiency: a 'new'

[106148]phenotype? (Abstract) Pediat. Res. 7: 394 only, 1973.

[106149]18628.

[106150]Tildon, J. T.; Cornblath, M.: Succinyl-CoA: 3-ketoacid CoA-transferase

[106151]deficiency. A cause for ketoacidosis in infancy. J. Clin. Invest. 51:

[106152]493-498, 1972.

[106153]18629.

[106154]Cannizzaro, L. A.; Chen, Y. Q.; Rafi, M. A.; Wenger, D. A.:
Regional

[106155]mapping of the human galactocerebrosidase gene (GALC)
to 14q31 by

[106156]in situ hybridization. Cytogenet. Cell Genet. 66: 244–245,
1994.

[106157]18630.

[106158]De Gasperi, R.; Sosa, M. A. G.; Sartorato, E. L.; Battistini,
S.;

[106159]MacFarlane, H.; Gusella, J. F.; Krivit, W.; Kolodny, E. H.:
Molecular

[106160]heterogeneity of late-onset forms of globoid-cell
leukodystrophy. Am.

[106161]J. Hum. Genet. 59: 1233–1242, 1996.

[106162]18631.

[106163]Duchen, L. W.; Eicher, E. M.; Jacobs, J. M.; Scaravilli, F.;
Teixeira,

[106164]F.: Hereditary leucodystrophy in the mouse: the new mu-
tant twitcher. Brain 103:

[106165]695–710, 1980.

[106166]18632.

[106167]Trauner, D. A.; Page, T.; Greco, C.; Sweetman, L.; Kulovich, S.;

[106168]Nyhan, W. L.: Progressive neurodegenerative disorder in a patient

[106169]with nonketotic hyperglycinemia. J. Pediat. 98: 272–275, 1981.

[106170]18633.

[106171]Minassian, B. A.; Lee, J. R.; Herbrick, J.–A.; Huizenga, J.; Soder,

[106172]S.; Mungall, A. J.; Dunham, I.; Gardner, R.; Fong, C. G.; Carpenter,

[106173]S.; Jardim, L.; Satishchandra, P.; Andermann, E.; Snead, O. C., III;

[106174]Lopes–Cendes, I.; Tsui, L.–C.; Delgado–Escueta, A. V.; Rouleau, G.

[106175]A.; Scherer, S. W.: Mutations in a gene encoding a novel protein

[106176]tyrosine phosphatase cause progressive myoclonus epilepsy. Nature

[106177]Genet. 20: 171–174, 1998.

[106178]18634.

- [106179]Berger, R.; Smit, G. P. A.; Schierbeek, H.; Bijsterveld, K.; le
[106180]Coutre, R.: Mevalonic aciduria: an inborn error of cholesterol biosynthesis? Clin.
[106181]Chim. Acta 152: 219–222, 1985.
[106182]18635.
- [106183]Cuisset, L.; Drenth, J. P. H.; Simon, A.; Vincent, M. F.; van
der
[106184]Velde Visser, S.; van der Meer, J. W. M.; Grateau, G.;
Delpech, M.;
[106185]International Hyper-IgD Study Group: Molecular analysis
of MVK
[106186]mutations and enzymatic activity in hyper-IgD and periodic fever syndrome. Europ.
[106187]J. Hum. Genet. 9: 260–266, 2001.
[106188]18636.
- [106189]Drenth, J. P. H.; Cuisset, L.; Grateau, G.; Vasseur, C.; van
de
[106190]Velde-Visser, S. D.; de Jong, J. G. N.; Beckmann, J. S.; van
der Meer,
[106191]J. W. M.; Delpech, M.; International Hyper-IgD Study
Group: Mutations

[106192]in the gene encoding mevalonate kinase cause hyper-IgD
and periodic

[106193]fever syndrome. *Nature Genet.* 22: 178–181, 1999.

[106194]18637.

[106195]Drenth, J. P. H.; Haagsma, C. J.; van der Meer, J. W. M.; In-
ternational

[106196]Hyper-IgD Study Group: Hyperimmunoglobulinemia D and
periodic fever

[106197]syndrome: the clinical spectrum in a series of 50 patients.
Medicine 73:

[106198]133–144, 1994.

[106199]18638.

[106200]Gibson, K. M.; Hoffmann, G.; Nyhan, W. L.; Sweetman, L.;
Berger,

[106201]R.; le Coultre, R.; Smit, G. P. A.: Mevalonate kinase defi-
ciency

[106202]in a child with cerebellar ataxia hypotonia and mevalonic
aciduria. *Europ.*

[106203]*J. Pediat.* 148: 250–252, 1988.

[106204]18639.

[106205]Gibson, K. M.; Hoffmann, G.; Nyhan, W. L.; Sweetman, L.;

Brandt,

[106206]J. K.; Wappner, R. S.; Bader, P. I.: Mevalonic aciduria: family studies

[106207]in mevalonate kinase deficiency, an inborn error of cholesterol biosynthesis. J.

[106208]Inherit. Metab. Dis. 10 (suppl. 2): 282–285, 1987.

[106209]18640.

[106210]Gibson, K. M.; Hoffmann, G. F.; Tanaka, R. D.; Bishop, R. W.; Chambliss,

[106211]K. L.: Mevalonate kinase map position 12q24. Chromosome Res. 5:

[106212]150 only, 1997.

[106213]18641.

[106214]Hinson, D. D.; Rogers, Z. R.; Hoffmann, G. F.; Schachtele, M.;

[106215]Fingerhut, R.; Kohlschutter, A.; Kelley, R. I.; Gibson, K. M.: Hematological

[106216]abnormalities and cholestatic liver disease in two patients with mevalonate

[106217]kinase deficiency. Am. J. Med. Genet. 78: 408–412, 1998.

[106218]18642.

- [106219]Hinson, D. D.; Ross, R. M.; Krisans, S.; Shaw, J. L.; Kozich, V.;
- [106220]Rolland, M.-O.; Divry, P.; Mancini, J.; Hoffmann, G. F.; Gibson, K.
- [106221]M.: Identification of a mutation cluster in mevalonate kinase deficiency,
- [106222]including a new mutation in a patient of Mennonite ancestry. Am.
- [106223]J. Hum. Genet. 65: 327-335, 1999.
- [106224]18643.
- [106225]Hoffmann, G.; Gibson, K. M.; Brandt, I. K.; Bader, P. I.; Wappner,
- [106226]R. S.; Sweetman, L.: Mevalonic aciduria--an inborn error of cholesterol
- [106227]and nonsterol isoprene biosynthesis. New Eng. J. Med. 314: 1610-1614,
- [106228]1986.
- [106229]18644.
- [106230]Hoffmann, G. F.; Charpentier, C.; Mayatepek, E.; Mancini, J.;
- [106231]Leichsenring, M.; Gibson, K. M.; Divry, P.; Hrebicek, M.; Lehnert,

[106232]W.; Sartor, K.; Trefz, F. K.; Rating, D.; Bremer, H. J.; Nyhan, W.

[106233]L.: Clinical and biochemical phenotype in 11 patients with mevalonic

[106234]aciduria. *Pediatrics* 91: 915–921, 1993.

[106235]18645.

[106236]Houten, S. M; Koster, J.; Romeijn, G.-J.; Frenkel, J.; Di Rocco,

[106237]M.; Caruso, U.; Landrieu, P.; Kelley, R. I.; Kuis, W.; Poll-The, B.

[106238]T.; Gibson, K. M.; Wanders, R. J. A.; Waterham, H. R.: Organization

[106239]of the mevalonate kinase (MVK) gene and identification of novel mutations

[106240]causing mevalonic aciduria and hyperimmunoglobulinemia D and periodic

[106241]fever syndrome. *Europ. J. Hum. Genet.* 9: 253–259, 2001.
Note: Erratum:

[106242]*Europ. J. Med. Genet.* 9: 651 only, 2001.

[106243]18646.

[106244]Krisans, S. K.; Ericsson, J.; Edwards, P. A.; Keller, G. A.:

[106245]Farnesyl-diphosphate synthase is localized in peroxisomes. J. Biol.

[106246]Chem. 269: 14165-14169, 1994.

[106247]18647.

[106248]Mancini, J.; Philip, N.; Chabrol, B.; Divry, P.; Rolland, M.-O.;

[106249]Pinsard, N.: Mevalonic aciduria in 3 siblings: a new recognizable

[106250]metabolic encephalopathy. Pediat. Neurol. 9: 243-246, 1993.

[106251]18648.

[106252]Schafer, B. L.; Bishop, R. W.; Kratunis, V. J.; Kalinowski, S.

[106253]S.; Mosley, S. T.; Gibson, K. M.; Tanaka, R. D.: Molecular cloning

[106254]of human mevalonate kinase and identification of a missense mutation

[106255]in the genetic disease mevalonic aciduria. J. Biol. Chem. 267: 13229-13238,

[106256]1992.

[106257]18649.

[106258]van der Meer, J. W. M.; Vossen, J. M.; Radl, J.; van

Nieuwkoop,

[106259]J. A.; Meyer, C. J. L. M.; Lobatto, S.; van Furth, R.: Hyper-immunoglobulinaemia

[106260]D and periodic fever: a new syndrome. Lancet I: 1087-1090, 1984.

[106261]18650.

[106262]Houten, S. M.; Kuis, W.; Duran, M.; de Koning, T. J.; van Royen-Kerkhof,

[106263]A.; Romeijn, G. J.; Frenkel, J.; Dorland, L.; de Barse, M. M. J.;

[106264]Huijbers, W. A. R.; Rijkers, G. T.; Waterham, H. R.; Wanders, R. J.

[106265]A.; Poll-The, B. T.: Mutations in MVK, encoding mevalonate kinase,

[106266]cause hyperimmunoglobulinaemia D and periodic fever syndrome. Nature

[106267]Genet. 22: 175-177, 1999.

[106268]18651.

[106269]Houten, S. M.; Romeijn, G. J.; Koster, J.; Gray, R. G. F.; Darbyshire,

[106270]P.; Smit, G. P. A.; de Klerk, J. B. C.; Duran, M.; Gibson, K. M.;

- [106271]Wanders, R. J. A.; Waterham, H. R.: Identification and characterization
[106272]of three novel missense mutations in mevalonate kinase
cDNA causing
[106273]mevalonic aciduria, a disorder of isoprene biosynthesis.
Hum. Molec.
[106274]Genet. 8: 1523–1528, 1999.
[106275]18652.
- [106276]Norio, R.; Koskiniemi, M.: Progressive myoclonus epilepsy:
genetic
[106277]and nosological aspects with special reference to 107
Finnish patients. Clin.
[106278]Genet. 15: 382–398, 1979.
[106279]18653.
- [106280]Ortiz-Hidalgo, C.: The man behind Lafora's bodies. Am. J.
Surg.
[106281]Path. 10: 358–361, 1986.
[106282]18654.
- [106283]Sainz, J.; Minassian, B. A.; Serratosa, J. M.; Gee, M. N.;
Sakamoto,
[106284]L. M.; Iranmanesh, R.; Bohlega, S.; Baumann, R. J.; Ryan,

S.; Sparkes,

[106285]R. S.; Delgado-Escueta, A. V.: Lafora progressive myoclonus epilepsy:

[106286]narrowing the chromosome 6q24 locus by recombinations and homozygosities.

[106287](Letter) Am. J. Hum. Genet. 61: 1205-1209, 1997.

[106288]18655.

[106289]Sarlin, M. B.; Kloepper, H. W.; Mickle, W. A.; Heath, R. G.:

[106290]The detection of carriers in hereditary myoclonic epilepsy. Acta

[106291]Genet. Med. Gemellol. 9: 466-471, 1960.

[106292]18656.

[106293]Schwarz, G. A.; Yanoff, M.: Lafora's disease, distinct clinico-pathologic

[106294]form of Unverricht's syndrome. Arch. Neurol. 12: 172-188, 1965.

[106295]18657.

[106296]Serratos, J. M.; Delgado-Escueta, A. V.; Posada, I.; Shih, S.;

[106297]Drury, I.; Berciano, J.; Zabala, J. A.; Antunez, M. C.; Sparkes, R.

- [106298]S.: The gene for progressive myoclonus epilepsy of the Lafora type
[106299]maps to chromosome 6q. Hum. Molec. Genet. 4:
1657–1663, 1995.
[106300]18658.
- [106301]Serratos, J. M.; Gomez–Garre, P.; Gallardo, M. E.; Anta, B.;
[106302]Beltran–Valero de Bernabe, D.; Lindhout, D.; Augustijn, P.
B.; Tassinari,
[106303]C. A.; Michelucci, R.; Malafosse, A.; Topcu, M.; Grid, D.;
Dravet,
[106304]C.; Berkovic, S. F.; Rodriguez de Cordoba, S.: A novel pro-
tein tyrosine
[106305]phosphatase gene is mutated in progressive myoclonus
epilepsy of the
[106306]Lafora type (EPM2). Hum. Molec. Genet. 8: 345–352,
1999.
[106307]18659.
- [106308]Yanoff, M.; Schwarz, G. A.: Lafora's disease: a distinct ge-
netically
[106309]determined form of Unverricht's syndrome. Genet. Hum.
14: 235–244,
[106310]1965.

[106311]18660.

[106312]Yokoi, S.; Austin, J.; Witmer, F.; Sakai, M.: Studies in myoclonus

[106313]epilepsy (Lafora body forms). I. Isolation and preliminary characterization

[106314]of Lafora bodies in two cases. Arch. Neurol. 19: 15–33, 1968.

[106315]18661.

[106316]Mowat, D. R.; Croaker, G. D. H.; Cass, D. T.; Kerr, B. A.; Chaitow,

[106317]J.; Ades, L. C.; Chia, N. L.; Wilson, M. J.: Hirschsprung disease,

[106318]microcephaly, mental retardation, and characteristic facial features:

[106319]delineation of a new syndrome and identification of a locus at chromosome

[106320]2q22–q23. J. Med. Genet. 35: 617–623, 1998.

[106321]18662.

[106322]Wakamatsu, N.; Yamada, Y.; Yamada, K.; Ono, T.; Nomura, N.; Taniguchi,

[106323]H.; Kitoh, H.; Mutoh, N.; Yamanaka, T.; Mushiaki, K.; Kato, K.; Sonta,

[106324]S.; Nagaya, M.: Mutations in SIP1, encoding Smad interacting protein-1,
[106325]cause a form of Hirschsprung disease. Nature Genet. 27:
369-370,
[106326]2001.
[106327]18663.
[106328]Zweier, C.; Albrecht, B.; Mitulla, B.; Behrens, R.; Beese, M.;
[106329]Gillesen-Kaesbach, G.; Rott, H.-D.; Rauch, A.: 'Mowat-
Wilson' syndrome
[106330]with and without Hirschsprung disease is a distinct, recognizable
[106331]multiple congenital anomalies-mental retardation syndrome caused by
[106332]mutations in the zinc finger homeo box 1B gene. Am. J. Med. Genet. 108:
177-181, 2002.
[106333]18664.
[106334]18664.
[106335]Yamada, K.; Yamada, Y.; Nomura, N.; Miura, K.; Wakako, R.; Hayakawa,
[106336]C.; Matsumoto, A.; Kumagai, T.; Yoshimura, I.; Miyazaki, S.; Kato,

[106337]K.; Sonta, S.; Ono, H.; Yamanaka, T.; Nagaya, M.; Wakamatsu, N.:

[106338]Nonsense and frameshift mutations in ZFX1B, encoding Smad-interacting

[106339]protein 1, cause a complex developmental disorder with a great variety

[106340]of clinical features. Am. J. Hum. Genet. 69: 1178–1185, 2001.

[106341]18665.

[106342]Alm, J.; Holmgren, G.; Larsson, A.; Schimpfessel, L.: Histidinaemia

[106343]in Sweden: report on a neonatal screening programme. Clin. Genet. 20:

[106344]229–233, 1981.

[106345]18666.

[106346]Anakura, M.; Matsuda, I.; Arashima, S.; Fukushima, N.; Oka, Y.

[106347]: Histidinemia--classical and atypical form in siblings. Am. J. Dis.

[106348]Child. 129: 858–861, 1975.

[106349]18667.

- [106350]Auerbach, V. H.; DiGeorge, A. M.; Baldrige, R. C.;
Tourtellotte,
[106351]C. D.; Brigham, M. P.: Histidinemia: a deficiency in histi-
dase resulting
[106352]in the urinary excretion of histidine and of imidazolepyru-
vic acid. J.
[106353]Pediat. 60: 487-497, 1962.
[106354]18668.
- [106355]Bruckman, C.; Berry, H. K.; Dasenbrock, R. J.: Histidinemia
in
[106356]two successive generations. Am. J. Dis. Child. 119:
221-227, 1970.
[106357]18669.
- [106358]Coulombe, J. T.; Kammerer, B. L.; Levy, H. L.; Hirsch, B. Z.;
Sriver,
[106359]C. R.: Histidinaemia. Part III: Impact; a prospective study. J.
[106360]Inherit. Metab. Dis. 6: 58-61, 1983.
[106361]18670.
- [106362]Kacser, H. K.; Bulfield, G.; Wallace, M. E.: Histidinaemia
mutant
[106363]in the mouse. Nature 244: 77-79, 1973.

[106364]18671.

[106365]Kappelman, M.; Thomas, G. H.; Howell, R. R.: Histidinemia
in a

[106366]Negro child. Am. J. Dis. Child. 122: 212–214, 1971.

[106367]18672.

[106368]La Du, B. N.: Histidinemia: current status. Am. J. Dis.
Child. 113:

[106369]88–92, 1967.

[106370]18673.

[106371]La Du, B. N.: Histidinemia.In: Stanbury, J. B.; Wyngaarden,
J.

[106372]B.; Fredrickson, D. S.: Metabolic Basis of Inherited Disease.
New

[106373]York: McGraw–Hill (pub.) (4th ed.): 1978. Pp. 317–327.

[106374]18674.

[106375]La Du, B. N.; Howell, R. R.; Jacoby, G. A.; Seegmiller, J. E.;

[106376]Sober, E. K.; Zannoni, V. G.; Canby, J. P.; Ziegler, L. K.:
Clinical

[106377]and biochemical studies on two cases of histidinemia. Pe-
diatrics 32:

[106378]216–227, 1963.

[106379]18675.

[106380]Lam, W. K.; Cleary, M. A.; Wraith, J. E.; Walter, J. H.: Histid-
inaemia:

[106381]a benign metabolic disorder. Arch. Dis. Child. 74:
343–346, 1996.

[106382]18676.

[106383]Lemieux, B.; Auray-Blais, C.; Giguere, R.; Shapcott, D.;
Scriver,

[106384]C. R.: Newborn urine screening experience with over one
million infants

[106385]in the Quebec network of genetic medicine. J. Inherit.
Metab. Dis. 11:

[106386]45–55, 1988.

[106387]18677.

[106388]Levy, H. L.; Shin, V. E.; Madigan, P. M.: Routine newborn
screening

[106389]for histidinemia: clinical and biochemical results. New
Eng. J. Med. 291:

[106390]1214–1219, 1974.

[106391]18678.

[106392]Neville, B. G. R.; Bentovim, A.; Clayton, B. E.; Sheperd, J.:

[106393]Histidinaemia: study of relation between clinical and biological findings

[106394]in 7 subjects. Arch. Dis. Child. 47: 190-200, 1972.

[106395]18679.

[106396]Neville, B. G. R.; Harris, R. F.; Stern, D. J.; Stern, J.: Maternal

[106397]histidinaemia. Arch. Dis. Child. 46: 119-121, 1971.

[106398]18680.

[106399]Popkin, J. S.; Clow, C. L.; Scriver, C. R.; Grove, J.: Is hereditary

[106400]histidinemia harmful? Lancet I: 721-722, 1974.

[106401]18681.

[106402]Rosenblatt, D. S.; Mohyuddin, F.; Scriver, C. R.: Histidinemia

[106403]discovered by urine screening after renal transplantation. Pediatrics 46:

[106404]47-53, 1970.

[106405]18682.

[106406]Barton, R. W.; Neufeld, E. F.: A distinct biochemical deficit

- [106407]in the Maroteaux–Lamy syndrome (mucopolysaccharidosis VI). J. Pediat. 80: 114–116, 1972.
- [106408]18683.
- [106409]Hechtman, P.; Boulay, B.; De Braekeleer, M.; Andermann, E.; Melancon, S.; Larochelle, J.; Prevost, C.; Kaplan, F.: The intron 7 donor splice site transition: a second Tay–Sachs disease mutation in French Canada. Hum. Genet. 90: 402–406, 1992.
- [106410]18684.
- [106411]Hechtman, P.; Kaplan, F.; Bayleran, J.; Boulay, B.; Andermann, E.; de Braekeleer, M.; Melancon, S.; Lambert, M.; Potier, M.; Gagne, R.; Kolodny, E.; Clow, C.; Capua, A.; Prevost, C.; Sriver, C.: More than one mutant allele causes infantile Tay–Sachs disease in French–Canadians. Am. J. Hum. Genet. 47: 815–822, 1990.

[106420]18685.

[106421]Hellkuhl, B.; Mayr, W. R.; Grzeschik, K.-H.: Localization of

[106422]MPI, PK-M2, IDH-M, and the alpha subunit of hex-
osaminidase (HEX-A)

[106423]to the q21-qter region of human chromosome 15. Cyto-
genet. Cell Genet. 22:

[106424]503-505, 1978.

[106425]18686.

[106426]Hou, Y.; Vavougios, G.; Hinek, A.; Wu, K. K.; Hechtman, P.;
Kaplan,

[106427]F.; Mahuran, D. J.: The val192-to-leu mutation in the al-
pha-subunit

[106428]of beta-hexosaminidase A is not associated with the
B1-variant form

[106429]of Tay-Sachs disease. Am. J. Hum. Genet. 59: 52-58,
1996.

[106430]18687.

[106431]Kappler, J.; Gieselmann, V.; Propping, P.: Hexosaminidase-
-pseudodeficiency?.

[106432](Letter) Am. J. Hum. Genet. 47: 880-881, 1990.

[106433]18688.

[106434]Keats, B. J. B.; Elston, R. C.; Andermann, E.: Pedigree discriminant

[106435]analysis of two French Canadian Tay-Sachs families.
Genet. Epidemiol. 4:

[106436]77-85, 1987.

[106437]18689.

[106438]Kelly, T. E.; Chase, G. A.; Kaback, M. M.; Kumor, K.; McKusick,

[106439]V. A.: Tay-Sachs disease: high gene frequency in a non-Jewish population. Am.

[106440]J. Hum. Genet. 27: 287-291, 1975.

[106441]18690.

[106442]Kelly, T. E.; Reynolds, L. W.; O'Brien, J. S.: Segregation within

[106443]a family to two mutant alleles for hexosaminidase A. Clin.
Genet. 9:

[106444]540-543, 1976.

[106445]18691.

[106446]Lalley, P. A.; Rattazzi, M. C.; Shows, T. B.: Human beta-D-N-acetylhexosaminidases

[106447]A and B: expression and linkage relationships in somatic

cell hybrids. Proc.

[106448]Nat. Acad. Sci. 71: 1569–1573, 1975.

[106449]18692.

[106450]Korneluk, R. G.; Mahuran, D. J.; Neote, K.; Klavins, M. H.;
O'Dowd,

[106451]B. F.; Tropak, M.; Willard, H. F.; Anderson, M.-J.; Lowden,
J. A.;

[106452]Gravel, R. A.: Isolation of cDNA clones coding for the al-
pha-subunit

[106453]of human beta-hexosaminidase: extensive homology be-
tween the alpha-

[106454]and beta-subunits and studies on Tay-Sachs disease. J.
Biol. Chem. 261:

[106455]8407–8413, 1986.

[106456]18693.

[106457]Landels, E. C.; Green, P. M.; Ellis, I. H.; Fensom, A. H.; Bo-
brow,

[106458]M.: Beta-hexosaminidase splice site mutation has a high
frequency

[106459]among non-Jewish Tay-Sachs disease carriers from the
British Isles. J.

[106460]Med. Genet. 29: 563–567, 1992.

[106461]18694.

[106462]Landels, E. C.; Green, P. M.; Ellis, I. H.; Fensom, A. H.;
Kaback,

[106463]M. M.; Lim-Steele, J.; Zeiger, K.; Levy, N.; Bobrow, M.: Fur-
ther

[106464]investigation of the HEXA gene intron 9 donor splice site
mutation

[106465]frequently found in non-Jewish Tay-Sachs disease pa-
tients from the

[106466]British Isles. J. Med. Genet. 30: 479-481, 1993.

[106467]18695.

[106468]Lau, M. M. H.; Neufeld, E. F.: A frameshift mutation in a
patient

[106469]with Tay-Sachs disease causes premature termination and
defective

[106470]intracellular transport of the alpha-subunit of beta-
hexosaminidase. J.

[106471]Biol. Chem. 264: 21376-21380, 1989.

[106472]18696.

[106473]McDowell, G. A.; Mules, E. H.; Fabacher, P.; Shapira, E.;
Blitzer,

- [106474]M. G.: The presence of two different infantile Tay-Sachs disease mutations in a Cajun population. Am. J. Hum. Genet. 51: 1071-1077, 1992.
- [106476]18697.
- [106477]18697.
- [106478]Lyerla, T. A.; Konola, J. T.; Skiba, M. C.; Raghavan, S.: Galactocerebrosidase activity in somatic cell hybrids derived from twitcher mouse/control human fibroblasts is associated with human chromosome 17. Am. J. Hum. Genet. 44: 198-207, 1989.
- [106480]18698.
- [106481]18698.
- [106482]Sakai, N.; Inui, K.; Fujii, N.; Fukushima, H.; Nishimoto, J.; Yanagihara, I.; Isegawa, Y.; Iwamatsu, A.; Okada, S.: Krabbe disease: isolation and characterization of a full-length cDNA for human galactocerebrosidase. Biochem. Biophys. Res. Commun. 198: 485-491, 1994.
- [106486]18699.

[106488]Sweet, H.: Twitcher (twi) is on chromosome 12. Mouse
Newsletter 75:

[106489]30, 1986.

[106490]18700.

[106491]Black, S. H.; Pelias, M. Z.; Miller, J. B.; Blitzer, M. G.;
Shapira,

[106492]E.: Maroteaux-Lamy syndrome in a large consanguineous
kindred: biochemical

[106493]and immunological studies. Am. J. Med. Genet. 25:
273-279, 1986.

[106494]18701.

[106495]Brooks, D. A.; McCourt, P. A. G.; Gibson, G. J.; Ashton, L.
J.;

[106496]Shutter, M.; Hopwood, J. J.: Analysis of N-
acetylgalactosamine-4-sulfatase

[106497]protein and kinetics in mucopolysaccharidosis type VI pa-
tients. Am.

[106498]J. Hum. Genet. 48: 710-719, 1991.

[106499]18702.

[106500]Brooks, D. A.; McCourt, P. A. G.; Gibson, G. J.; Hopwood, J.
J.

[106501]: Immunoquantification of the low abundance lysosomal enzyme N-acetylgalactosamine

[106502]4-sulphatase. J. Inherit. Metab. Dis. 13: 108-120, 1990.

[106503]18703.

[106504]Cantor, L. B.; Disseler, J. A.; Wilson, F. M., II: Glaucoma in

[106505]the Maroteaux-Lamy syndrome. Am. J. Ophthal. 108: 426-430, 1989.

[106506]18704.

[106507]Crawley, A. C.; Brooks, D. A.; Muller, V. J.; Petersen, B. A.;

[106508]Isaac, E. L.; Bielicki, J.; King, B. M.; Boulter, C. D.; Moore, A.

[106509]J.; Fazzalari, N. L.; Anson, D. S.; Byers, S.; Hopwood, J. J.: Enzyme

[106510]replacement therapy in a feline model of Maroteaux-Lamy syndrome. J.

[106511]Clin. Invest. 97: 1864-1873, 1996.

[106512]18705.

[106513]Crawley, A. C.; Niedzielski, K. H.; Isaac, E. L.; Davey, R. C.

[106514]A.; Byers, S.; Hopwood, J. J.: Enzyme replacement therapy from birth

- [106515]in a feline model of mucopolysaccharidosis type VI. J. Clin. Invest. 99: 651-662, 1997.
- [106517]18706.
- [106518]Crawley, A. C.; Yogalingam, G.; Muller, V. J.; Hopwood, J. J.: Two mutations within a feline mucopolysaccharidosis type VI colony cause three different clinical phenotypes. J. Clin. Invest. 101: 109-119, 1998.
- [106522]18707.
- [106523]DeLuca, C.; Brown, J. A.; Shows, T. B.: Lysosomal arylsulfatase deficiencies in humans: chromosome assignment for arylsulfatase A and B. Proc. Nat. Acad. Sci. 76: 1957-1961, 1979.
- [106526]18708.
- [106527]DiFerrante, N. M.; Hyman, B. H.; Kish, W.; Donnelly, P. V.; Nichols, B. L., Jr.; Dutton, R. V.: Mucopolysaccharidosis VI

(Maroteaux-Lamy

[106529]disease) clinical and biochemical study of a mild variant case. Johns

[106530]Hopkins Med. J. 135: 42-53, 1974.

[106531]18709.

[106532]Dudin, G.; Alexander, D.; Talj, F.; Deeb, M.; Musallam, S.;
Der

[106533]Kaloustian, V. M.: Interstitial deletion of band q12 of
chromosome

[106534]5. Clin. Genet. 25: 455-458, 1984.

[106535]18710.

[106536]Evers, M.; Saftig, P.; Schmidt, P.; Hafner, A.; McLoughlin, D.

[106537]B.; Schmahl, W.; Hess, B.; von Figura, K.; Peters, C.: Tar-
geted disruption

[106538]of the arylsulfatase B gene results in mice resembling the
phenotype

[106539]of mucopolysaccharidosis VI. Proc. Nat. Acad. Sci. 93:
8214-8219,

[106540]1996.

[106541]18711.

[106542]Fidzianska, E.; Abramowicz, T.; Czartoryska, B.; Glo-

gowska, I.;

[106543]Gorska, D.; Mazurozak, T.; Rodo, M.; Wehr, H.; Witkowska,
J.: Localization

[106544]of the human arylsulfatase B gene locus to chromosome 5
by use of

[106545]hamster-man somatic cell hybrids.(Abstract) 7th Int.
Cong. Hum. Genet.,

[106546]Berlin 611-612, 1986.

[106547]18712.

[106548]Fidzianska, E.; Abramowicz, T.; Czartoryska, B.; Glo-
owska, I.;

[106549]Gorska, D.; Rodo, M.: Assignment of the gene for human
arylsulfatase

[106550]B, ARSB, to chromosome region 5p11-5qter. Cytogenet.
Cell Genet. 38:

[106551]150-151, 1984.

[106552]18713.

[106553]Fox, M. F.; DuToit, D. L.; Warnich, L.; Retief, A. E.: Regional

[106554]localization of alpha-galactosidase (GLA) to Xpter-q22,
hexosaminidase

[106555]B (HEXB) to 5q13-qter, and arylsulfatase B (ARSB) to

5pter-q13. Cytogenet.

[106556]Cell Genet. 38: 45-49, 1984.

[106557]18714.

[106558]Gitzelmann, R.: Personal Communication. Zurich, Switzerland

[106559]4/27/1990.

[106560]18715.

[106561]Goldberg, M. F.; Scott, C. I., Jr.; McKusick, V. A.: Hydrocephalus

[106562]and papilledema in the Maroteaux-Lamy syndrome (mucopolysaccharidosis

[106563]type VI). Am. J. Ophthal. 69: 969-975, 1970.

[106564]18716.

[106565]Haskins, M. E.; Jezyk, P. F.; Patterson, D. F.: Mucopolysaccharide

[106566]storage disease in three families of cats with arylsulfatase B deficiency:

[106567]leukocyte studies and carrier identification. Pediat. Res. 13: 1203-1210,

[106568]1979.

[106569]18717.

[106570]Hellkuhl, B.; Grzeschik, K.-H.: Assignment of a gene for
arylsulfatase

[106571]B to human chromosome 5 (EC 3.1.6.1) using human-
mouse somatic cell

[106572]hybrids. Cytogenet. Cell Genet. 22: 203-206, 1978.

[106573]18718.

[106574]Hopwood, J. J.: Personal Communication. North Adelaide,
South

[106575]Australia 5/9/1991.

[106576]18719.

[106577]Isbrandt, D.; Hopwood, J. J.; von Figura, K.; Peters, C.: Two

[106578]novel frameshift mutations causing premature stop
codons in a patient

[106579]with the severe form of Maroteaux-Lamy syndrome. Hum.
Mutat. 7:

[106580]361-363, 1996.

[106581]18720.

[106582]Jezyk, P. F.; Haskins, M. E.; Patterson, D. F.; Mellman, W. J.;

[106583]Greenstein, M.: Mucopolysaccharidosis in a cat with aryl-
sulfatase

[106584]B deficiency: a model of Maroteaux–Lamy syndrome. Science 198: 834–836,
[106585]1977.
[106586]18721.
[106587]Jin, W.–D.; Jackson, C. E.; Desnick, R. J.; Schuchman, E. H.:
[106588]Mucopolysaccharidosis type VI: identification of three mutations in
[106589]the arylsulfatase B gene of patients with the severe and mild phenotypes
[106590]provides molecular evidence for genetic heterogeneity. Am. J. Hum.
[106591]Genet. 50: 795–800, 1992.
[106592]18722.
[106593]Krivit, W.; Pierpont, M. E.; Ayaz, K.; Tsai, M.; Ramsay, N. K.
[106594]C.; Kersey, J. H.; Weisdorf, S.; Sibley, R.; Snover, D.; McGovern,
[106595]M. M.; Schwartz, M. F.; Desnick, R. J.: Bone–marrow transplantation
[106596]in the Maroteaux–Lamy syndrome (mucopolysaccharidosis type VI): biochemical
[106597]and clinical status 24 months after transplantation. New

Eng. J.

[106598]Med. 311: 1606–1611, 1984.

[106599]18723.

[106600]Kunieda, T.; Simonaro, C. M.; Yoshida, M.; Ikadai, H.;
Levan,

[106601]G.; Desnick, R. J.; Schuchman, E. H.: Mucopolysaccharido-
sis type

[106602]VI in rats: isolation of cDNAs encoding arylsulfatase B,
chromosomal

[106603]localization of the gene, and identification of the muta-
tion. Genomics 29:

[106604]582–587, 1995.

[106605]18724.

[106606]Levy, L. A.; Lewis, J. C.; Sumner, T. E.: Ultrastructures of

[106607]Reilly bodies (metachromatic granules) in the Maroteaux–
Lamy syndrome

[106608](mucopolysaccharidosis VI): a histochemical study. Am. J.
Clin. Path. 73:

[106609]416–422, 1980.

[106610]18725.

[106611]Litjens, T.; Baker, E. G.; Beckmann, K. R.; Morris, C. P.;

Hopwood,

[106612]J. J.; Callen, D. F.: Chromosomal localization of ARSB, the
gene

[106613]for human N-acetylgalactosamine-4-sulphatase. Hum.
Genet. 82: 67-68,

[106614]1989.

[106615]18726.

[106616]Litjens, T.; Brooks, D. A.; Peters, C.; Gibson, G. J.; Hop-
wood,

[106617]J. J.: Identification, expression, and biochemical character-
ization

[106618]of N-acetylgalactosamine-4-sulfatase mutations and re-
lationship with

[106619]clinical phenotype in MPS-VI patients. Am. J. Hum. Genet.
58: 1127-1134,

[106620]1996.

[106621]18727.

[106622]Litjens, T.; Hopwood, J. J.: Mucopolysaccharidosis type VI:
structural

[106623]and clinical implications of mutations in N-
acetylgalactosamine-4-sulfatase. Hum.

[106624]Mutat. 18: 282-295, 2001.

[106625]18728.

[106626]Litjens, T.; Morris, C. P.; Robertson, E. F.; Peters, C.; von

[106627]Figura, K.; Hopwood, J. J.: An N-acetyl-

galactosamine-4-sulfatase

[106628]mutation (delta-G-238) results in a severe Maroteaux-

Lamy phenotype. Hum.

[106629]Mutat. 1: 397-402, 1992.

[106630]18729.

[106631]Taroni, F.; Verderio, E.; Dworzak, F.; Willems, P. J.; Cava-
dini,

[106632]P.; DiDonato, S.: Identification of a common mutation in
the carnitine

[106633]palmitoyltransferase II gene in familial recurrent myo-
globinuria patients. Nature

[106634]Genet. 4: 314-320, 1993.

[106635]18730.

[106636]Gellera, C.; Verderio, E.; Floridia, G.; Finocchiaro, G.; Mon-
termini,

[106637]L.; Cavadini, P.; Zuffardi, O.; Taroni, F.: Assignment of the
human

[106638]carnitine palmitoyltransferase II gene (CTP1) to chromo-

some 1p32. Genomics 24:

[106639]195–197, 1994.

[106640]18731.

[106641]Agmon, D.; Green, J.; Platau, E.; Better, O. S.: Isolated
adrenal

[106642]mineralocorticoid deficiency due to amyloidosis associ-
ated with familial

[106643]Mediterranean fever. Am. J. Med. Sci. 288: 40–43, 1984.

[106644]18732.

[106645]Akarsu, A. N.; Saatci, U.; Ozen, S.; Bakkaloglu, A.; Besbas,
N.;

[106646]Sarfarazi, M.: Genetic linkage study of familial Mediter-
ranean fever

[106647](FMF) to 16p13.3 and evidence for genetic heterogeneity
in the Turkish

[106648]population. J. Med. Genet. 34: 573–578, 1997.

[106649]18733.

[106650]Lukong, K. E.; Elsliger, M.–A.; Chang, Y.; Richard, C.;
Thomas,

[106651]G.; Carey, W.; Tyłki-Szymanska, A.; Czartoryska, B.; Buch-
holz, T.;

[106652]Rodriguez Criado, G.; Palmeri, S.; Pshezhetsky, A. V.:
Characterization

[106653]of the sialidase molecular defects in sialidosis patients
suggests

[106654]the structural organization of the lysosomal multienzyme
complex. Hum.

[106655]Molec. Genet. 9: 1075–1085, 2000.

[106656]18734.

[106657]Fernandes, M.; Poirier, C.; Lespinasse, F.; Carle, G. F.: The

[106658]mouse homologs of human GIF, DDB1, and CFL1 genes
are located on chromosome

[106659]19. Mammalian Genome 9: 339–342, 1998.

[106660]18735.

[106661]Aminoff, M.; Carter, J. E.; Chadwick, R. B.; Johnson, C.;
Grasbeck,

[106662]R.; Abdelaal, M. A.; Broch, H.; Jenner, L. B.; Verroust, P. J.;
Moestrup,

[106663]S. K.; de la Chapelle, A.; Krahe, R.: Mutations in CUBN, en-
coding

[106664]the intrinsic factor–vitamin B12 receptor, cubilin, cause
hereditary

[106665]megaloblastic anaemia 1. Nature Genet. 21: 309–313, 1999.

[106666]18736.

[106667]Aminoff, M.; Tahvanainen, E.; Grasbeck, R.; Weissenbach, J.; Broch,

[106668]H.; de la Chapelle, A.: Selective intestinal malabsorption of vitamin

[106669]B12 displays recessive mendelian inheritance: assignment of a locus

[106670]to chromosome 10 by linkage. Am. J. Hum. Genet. 57: 824–831, 1995.

[106671]18737.

[106672]Kozyraki, R.; Kristiansen, M.; Silahtaroglu, A.; Hansen, C.; Jacobsen,

[106673]C.; Tommerup, N.; Verroust, P. J.; Moestrup, S. K.: The human intrinsic

[106674]factor–vitamin B12 receptor, cubilin: molecular characterization and

[106675]chromosomal mapping of the gene to 10p within the autosomal recessive

[106676]megaloblastic anemia (MGA1) region. Blood 91: 3593–3600, 1998.

[106677]18738.

[106678]Carney, J. P.; Maser, R. S.; Olivares, H.; Davis, E. M.; Le
Beau,

[106679]M.; Yates, J. R., III; Hays, L.; Morgan, W. F.; Petrini, J. H. J.

[106680]: The hMre11/hRad50 protein complex and Nijmegen
breakage syndrome:

[106681]linkage of double-strand break repair to the cellular DNA
damage response. Cell 93:

[106682]477-486, 1998.

[106683]18739.

[106684]Matsuo, T.; Egawa, I.; Okada, S.; Suetsugu, M.; Yamamoto,
K.;

[106685]Watanabe, M.: Sialidosis type 2 in Japan: clinical study in
two siblings'

[106686]cases and review of literature. J. Neurol. Sci. 58: 45-55,
1983.

[106687]18740.

[106688]Miyatake, T.; Atsumi, T.; Obayashi, T.; Mizuno, Y.; Ando,
S.;

[106689]Ariga, T.; Matsui-Nakamura, K.; Yamada, T.: Adult type
neuronal storage

- [106690]disease with neuraminidase deficiency. Ann. Neurol. 6:
232–243,
[106691]1979.
- [106692]18741.
- [106693]Mueller, O. T.; Henry, W. M.; Haley, L. L.; Byers, M. G.;
Eddy,
[106694]R. L.; Shows, T. B.: Sialidosis and galactosialidosis: chro-
mosomal
[106695]assignment of two genes associated with neuraminidase
deficiency disorders. Proc.
[106696]Nat. Acad. Sci. 83: 1817–1821, 1986.
- [106697]18742.
- [106698]Naganawa, Y.; Itoh, K.; Shimmoto, M.; Takiguchi, K.; Doi,
H.;
- [106699]Nishizawa, Y.; Kobayashi, T.; Kamei, S.; Lukong, K. E.;
- Pshezhetsky,
[106700]A. V.; Sakuraba, H.: Molecular and structural studies of
Japanese
[106701]patients with sialidosis type 1. J. Hum. Genet. 45:
241–249, 2000.
- [106702]18743.

[106703] Nakamura, Y.; Takahashi, Y.; Yamaguchi, S.; Omiya, S.;
Orii, T.;

[106704] Yara, A.; Gushiken, M.: Severe infantile sialidosis--the
characteristics

[106705] of oligosaccharides isolated from the urine and the ab-
dominal ascites. Tohoku

[106706] J. Exp. Med. 166: 407-415, 1992.

[106707] 18744.

[106708] O'Brien, J. S.: Neuraminidase deficiency in the cherry red
spot-myoclonus

[106709] syndrome. Biochem. Biophys. Res. Commun. 79:
1136-1141, 1977.

[106710] 18745.

[106711] O'Brien, J. S.: The cherry red spot-myoclonus syndrome: a
newly

[106712] recognized inherited lysosomal storage disease due to
acid neuraminidase

[106713] deficiency. Clin. Genet. 14: 55-60, 1978.

[106714] 18746.

[106715] O'Brien, J. S.; Warner, T. G.: Sialidosis: delineation of sub-
types

[106716]by neuraminidase assay. Clin. Genet. 17: 35–38, 1980.

[106717]18747.

[106718]Oohira, T.; Nagata, N.; Akaboshi, I.; Matsuda, I.; Naito, S.:

[106719]The infantile form of sialidosis type II associated with
congenital

[106720]adrenal hyperplasia: possible linkage between HLA and
the neuraminidase

[106721]deficiency gene. Hum. Genet. 70: 341–343, 1985.

[106722]18748.

[106723]Oohira, T.; Yoshida, M. C.; Matsuda, I.: Additional evi-
dence

[106724]for the location of the alpha-neuraminidase gene on
chromosome 6. Jpn.

[106725]J. Hum. Genet. 31: 309–310, 1986.

[106726]18749.

[106727]Orii, T.; Minami, R.; Sukegawa, K.; Sato, S.; Tsugawa, S.;
Horino,

[106728]K.; Miura, R.; Nakao, T.: A new type of mucopolidosis with
beta-galactosidase

[106729]deficiency and glycopeptiduria. Tohoku J. Exp. Med. 107:
303–315,

[106730]1972.

[106731]18750.

[106732]Paschke, E.; Trinkl, G.; Erwa, W.; Pavelka, M.; Mutz, I.;
Roscher,

[106733]A.: Infantile type of sialic acid storage disease with
sialuria. Clin.

[106734]Genet. 29: 417-424, 1986.

[106735]18751.

[106736]Peters, J.; Swallow, D. M.; Andrews, S. J.; Evans, L.: A gene

[106737](Neu-1) on chromosome 17 of the mouse affects acid al-
pha-glucosidase

[106738]and codes for neuraminidase. Genet. Res. Camb. 38:
47-55, 1981.

[106739]18752.

[106740]Pincus, J. H.; Rossi, J. P.; Daroff, R. B.: Delayed develop-
ment

[106741]of disturbed mucopolysaccharide metabolism in a Hurler
variant. Arch.

[106742]Neurol. 16: 244-253, 1967.

[106743]18753.

[106744]Potier, M.; Beauregard, G.; Gelisle, M.; Mameli, L.; Hong, V.

[106745]N.; Melancon, S. B.; Dallaire, L.: Neuraminidase activity in
the

[106746]mucopolysaccharidoses (types I, II and III) and the cherry-red spot
myoclonus

[106747]syndrome. Clin. Chim. Acta 99: 97-105, 1979.

[106748]18754.

[106749]Pshchetsky, A. V.; Richard, C.; Michaud, L.; Ignotz, S.;
Wang,

[106750]S.; Elsliger, M.-A.; Qu, J.; Leclerc, D.; Gravel, R.; Dallaire,
L.;

[106751]Potier, M.: Cloning, expression and chromosomal map-
ping of human

[106752]lysosomal sialidase and characterization of mutations in
sialidosis. Nature

[106753]Genet. 15: 316-320, 1997.

[106754]18755.

[106755]Rottier, R. J.; Bonten, E.; d'Azzo, A.: A point mutation in
the

[106756]neu-1 locus causes the neuraminidase defect in the SM/J
mouse. Hum.

[106757]Molec. Genet. 7: 313–321, 1998.

[106758]18756.

[106759]Sanfilippo, S. J.; Yunis, J. J.; Worthen, H. G.: An unusual storage

[106760]disease resembling the Hurler–Hunter syndrome.
(Abstract) Am. J.

[106761]Dis. Child. 104: 553 only, 1962.

[106762]18757.

[106763]Sasagasako, N.; Miyahara, S.; Saito, N.; Shinnoh, N.;
Kobayashi,

[106764]T.; Goto, I.: Prenatal diagnosis of congenital sialidosis.
Clin.

[106765]Genet. 44: 8–11, 1993.

[106766]18758.

[106767]Sergi, C.; Penzel, R.; Uhl, J.; Zoubaa, S.; Dietrich, H.;
Decker,

[106768]N.; Rieger, P.; Kopitz, J.; Otto, H. F.; Kiessling, M.; Cantz,
M.

[106769]: Prenatal diagnosis and fetal pathology in a Turkish family harboring

[106770]a novel nonsense mutation in the lysosomal alpha–

N-acetyl-neuraminidase

[106771](sialidase) gene. Hum. Genet. 109: 421-428, 2001.

[106772]18759.

[106773]Shows, T. B.: Personal Communication. Buffalo, N. Y.

10/9/1985.

[106774]18760.

[106775]Spranger, J.: Advances in bone dysplasias. (Abstract) Sixth

[106776]Int. Cong. Hum. Genet., Jerusalem , 1981.

[106777]18761.

[106778]Spranger, J.: Mucopolidosis I: phenotype and nosology.

Perspect.

[106779]Inherit. Metab. Dis. 4: 303-315, 1981.

[106780]18762.

[106781]Spranger, J.; Cantz, M.: Mucopolidosis I, the cherry red

spot-myoclonus

[106782]syndrome and neuraminidase deficiency. Birth Defects

Orig. Art. Ser. XIV(6B):

[106783]105-112, 1978.

[106784]18763.

[106785]Spranger, J. W.: Personal Communication. Kiel, Germany
1972.

[106786]18764.

[106787]Spranger, J. W.; Gehler, J.; Cantz, M.: Mucopolidosis I--a
sialidosis. Am.

[106788]J. Med. Genet. 1: 21-29, 1977.

[106789]18765.

[106790]Spranger, J. W.; Wiedemann, H. R.: The genetic mucolipi-
doses.

[106791]Diagnosis and differential diagnosis. Humangenetik 9:
113-139, 1970.

[106792]18766.

[106793]Spranger, J. W.; Wiedemann, H. R.; Tolksdorf, M.; Graucob,
E.;

[106794]Caesar, R.: Lipomucopolysaccharidose: eine neue Spe-
icherkrankheit. Z.

[106795]Kinderheilk. 103: 285-306, 1968.

[106796]18767.

[106797]Steinmann, L.; Tharp, B. R.; Dorfman, L. J.; Forno, L. S.;
Sogg,

[106798]R. L.; Kelts, K. A.; O'Brien, J. S.: Peripheral neuropathy in

the

[106799]cherry red spot–myoclonus syndrome (sialidosis type I).

Ann. Neurol. 7:

[106800]450–456, 1980.

[106801]18768.

[106802]Stevenson, R. E.; Lubinsky, M.; Taylor, H. A.; Wenger, D.

A.;

[106803]Schroer, R. J.; Olmstead, P. M.: Sialic acid storage disease

with

[106804]sialuria: clinical and biochemical features in the severe in-

fantile

[106805]type. Pediatrics 72: 441–449, 1983.

[106806]18769.

[106807]Swallow, D. M.; Evans, L.; Stewart, G.; Thomas, P. K.;

Abrams,

[106808]J. D.: Sialidosis type I: cherry red spot–myoclonus syn-

drome with

[106809]sialidase deficiency and altered electrophoretic mobility of

some

[106810]enzymes known to be glycoproteins. II. Enzyme studies.

Ann. Hum.

[106811]Genet. 43: 27–35, 1979.

[106812]18770.

[106813]Swallow, D. M.; Hoogeveen, A. T.; Verheijen, F. W.; Galjaard,

[106814]H.: Complementation analysis of human sialidase deficiency using

[106815]natural substrates. Ann. Hum. Genet. 45: 105–112, 1981.

[106816]18771.

[106817]Kitagawa, T.: An animal model of human acid sphingomyelinase

[106818]deficiency (Niemann–Pick disease) and the study of its enzyme replacement. Jpn.

[106819]J. Hum. Genet. 32: 55–69, 1987.

[106820]18772.

[106821]Kurimasa, A.; Ohno, K.; Oshimura, M.: Restoration of the cholesterol

[106822]metabolism in 3T3 cell lines derived from the sphingomyelinosis mouse

[106823](spm/spm) by transfer of a human chromosome 18. Hum. Genet. 92:

[106824]157–162, 1993.

[106825]18773.

- [106826]Miyawaki, S.; Yoshida, H.; Mitsuoka, S.; Enomoto, H.; Ikehara,
hara,
[106827]S.: A mouse model for Niemann–Pick disease: influence of
genetic
[106828]background on disease expression in spm/spm mice. J.
Hered. 77:
[106829]379–384, 1986.
[106830]18774.
[106831]Mailly, F.; Palmen, J.; Muller, D. P. R.; Gibbs, T.; Lloyd, J.;
[106832]Brunzell, J.; Durrington, P.; Mitropoulos, K.; Betteridge, J.;
Watts,
[106833]G.; Lithell, H.; Angelico, F.; Humphries, S. E.; Talmud, P. J.:
Familial
[106834]lipoprotein lipase (LPL) deficiency: a catalogue of LPL gene
mutations
[106835]identified in 20 patients from the UK, Sweden, and Italy.
Hum. Mutat. 10:
[106836]465–473, 1997.
[106837]18775.
[106838]Normand, T.; Bergeron, J.; Fernandez–Margallo, T.;
Bharucha, A.;

[106839]Ven Murthy, M. R.; Julien, P.; Gagne, C.; Dionne, C.; De
Braekeleer,
[106840]M.; Ma, R.; Hayden, M. R.; Lupien, P. J.: Geographic distri-
bution
[106841]and genealogy of mutation 207 of the lipoprotein lipase
gene in the
[106842]French Canadian population of Quebec. Hum. Genet. 89:
671-675, 1992.
[106843]18776.

[106844]Monsalve, M. V.; Henderson, H.; Roederer, G.; Julien, P.;
Deeb,
[106845]S.; Kastelein, J. J. P.; Peritz, L.; Devlin, R.; Bruin, T.; Murthy,
[106846]M. R. V.; Gagne, C.; Davignon, J.; Lupien, P. J.; Brunzell, J.
D.;
[106847]Hayden, M. R.: A missense mutation at codon 188 of the
human lipoprotein
[106848]lipase gene is a frequent cause of lipoprotein lipase defi-
ciency in
[106849]persons of different ancestries. J. Clin. Invest. 86:
728-734, 1990.
[106850]18777.

[106851]Campbell, C.; Cucci, R. A.; Prasad, S.; Green, G. E.; Edeal, J.

[106852]B.; Galer, C. E.; Karniski, L. P.; Sheffield, V. C.; Smith, R. J.
[106853]H.: Pendred syndrome, DFNB4, and PDS/SLC26A4 identification of eight
[106854]novel mutations and possible genotype-phenotype correlations. Hum.
[106855]Mutat. 17: 403-411, 2001.
[106856]18778.
[106857]Everett, L. A.; Glaser, B.; Beck, J. C.; Idol, J. R.; Buchs, A.;
[106858]Heyman, M.; Adawi, F.; Hazani, E.; Nassir, E.; Baxevanis, A.
D.; Sheffield,
[106859]V. C.; Green, E. D.: Pendred syndrome is caused by mutations in a
[106860]putative sulphate transporter gene (PDS). Nature Genet.
17: 411-422,
[106861]1997.
[106862]18779.
[106863]Fugazzola, L.; Mannavola, D.; Cerutti, N.; Maghnie, M.;
Pagella,
[106864]F.; Bianchi, P.; Weber, G.; Persani, L.; Beck-Peccoz, P.:
Molecular
[106865]analysis of the Pendred's syndrome gene and magnetic

resonance imaging

[106866]studies of the inner ear are essential for the diagnosis of
true Pendred's

[106867]syndrome. J. Clin. Endocr. Metab. 85: 2469–2475, 2000.

[106868]18780.

[106869]Kopp, P.; Karamanoglu Arseven, O.; Sabacan, L.; Kotlar, T.;
Dupuis,

[106870]J.; Cavaliere, H.; Santos, C. L. S.; Jameson, J. L.; Medeiros–
Neto,

[106871]G.: Phenocopies for deafness and goiter development in a
large inbred

[106872]Brazilian kindred with Pendred's syndrome associated with
a novel

[106873]mutation in the PDS gene. J. Clin. Endocr. Metab. 84:
336–341, 1999.

[106874]18781.

[106875]Masmoudi, S.; Charfedine, I.; Hmani, M.; Grati, M.; Ghor–
bel, A.

[106876]M.; Elgaied–Boulila, A.; Drira, M.; Hardelin, J.–P.; Ayadi, M.:
Pendred

[106877]syndrome: phenotypic variability in two families carrying
the same

[106878]PDS missense mutation. Am. J. Med. Genet. 90: 38–44,
2000.

[106879]18782.

[106880]Boughman, J. A.; Conneally, P. M.; Nance, W. E.: Population
genetic

[106881]studies of retinitis pigmentosa. Am. J. Hum. Genet. 32:
223–235,

[106882]1980.

[106883]18783.

[106884]Gomez–Lira, M.; Perusi, C.; Mottes, M.; Pignatti, P. F.;
Manfredi,

[106885]M.; Rizzuto, N.; Salviati, A.: Molecular genetic characteri-
zation

[106886]of two metachromatic leukodystrophy patients who carry
the T799G mutation

[106887]and show different phenotypes; description of a novel
null-type mutation. Hum.

[106888]Genet. 102: 459–463, 1998.

[106889]18784.

[106890]Neer, E. J.; Schmidt, C. J.; Smith, T.: LIS is more. Nature

[106891]Genet. 5: 3–4, 1993.

[106892]18785.

[106893]Reiner, O.; Carrozzo, R.; Shen, Y.; Wehnert, M.;
Faustinella,

[106894]F.; Dobyns, W. B.; Caskey, C. T.; Ledbetter, D. H.: Isolation
of

[106895]a Miller-Dieker lissencephaly gene containing G protein
beta-subunit-like

[106896]repeats. Nature 364: 717-721, 1993.

[106897]18786.

[106898]Greene, H.; Hug, G.; Schubert, W. K.: Arylsulfatase A in the
[106899]urine and metachromatic leukodystrophy. J. Pediat. 71:
709-711,

[106900]1967.

[106901]18787.

[106902]Greenfield, J. G.: Form of progressive cerebral sclerosis in
[106903]infants associated with primary degeneration of interfascicular
glia. Proc.

[106904]Roy. Soc. Med. 26: 690-697, 1933.

[106905]18788.

[106906]Gustavson, K.-H.; Hagberg, B.: The incidence and genetics

of

[106907]metachromatic leukodystrophy in northern Sweden. Acta
Paediat. Scand. 60:

[106908]585–590, 1971.

[106909]18789.

[106910]Hagberg, B.; Sourander, P.; Svennerholm, L.: Sulfatide lipi-
dosis

[106911]in childhood. Am. J. Dis. Child. 104: 644–656, 1962.

[106912]18790.

[106913]Haltia, T.; Palo, J.; Haltia, M.; Icen, A.: Juvenile metachro-
matic

[106914]leukodystrophy: clinical, biochemical, and neuropatho-
logic studies

[106915]in nine new cases. Arch. Neurol. 37: 42–46, 1980.

[106916]18791.

[106917]Harvey, J. S.; Carey, W. F.; Morris, C. P.: Importance of the

[106918]glycosylation and polyadenylation variants in metachro-
matic leukodystrophy

[106919]pseudodeficiency phenotype. Hum. Molec. Genet. 7:
1215–1219, 1998.

[106920]18792.

- [106921]Harvey, J. S.; Nelson, P. V.; Carey, W. F.; Robertson, E. F.;
- [106922]Morris, C. P.: An arylsulfatase A (ARSA) missense mutation
(T274M)
- [106923]causing late-infantile metachromatic leukodystrophy.
Hum. Mutat. 2:
[106924]261–267, 1993.
- [106925]18793.
- [106926]Hasegawa, Y.; Kawame, H.; Eto, Y.: Mutations in the aryl-
sulfatase
- [106927]A gene of Japanese patients with metachromatic
leukodystrophy. DNA
[106928]Cell Biol. 12: 493–498, 1993.
- [106929]18794.
- [106930]Hasegawa, Y.; Kawame, H.; Ida, H.; Ohashi, T.; Eto, Y.: Sin-
gle
- [106931]exon mutation in arylsulfatase A gene has two effects:
loss of enzyme
- [106932]activity and aberrant splicing. Hum. Genet. 93: 415–420,
1994.
- [106933]18795.
- [106934]Heinisch, U.; Zlotogora, J.; Kafert, S.; Gieselmann, V.: Mul-

tiple

[106935] mutations are responsible for the high frequency of
metachromatic

[106936] leukodystrophy in a small geographic area. Am. J. Hum.
Genet. 56:

[106937] 51–57, 1995.

[106938] 18796.

[106939] Herz, B.; Bach, G.: Arylsulfatase A in pseudodeficiency.
Hum.

[106940] Genet. 66: 147–150, 1984.

[106941] 18797.

[106942] Hess, B.; Saftig, P.; Hartmann, D.; Coenen, R.; Lullmann–
Rauch,

[106943] R.; Goebel, H. H.; Evers, M.; von Figura, K.; D'Hooge, R.;
Nagels,

[106944] G.; De Deyn, P.; Peters, C.; Gieselmann, V.: Phenotype of
arylsulfatase

[106945] A–deficient mice: relationship to human metachromatic
leukodystrophy. Proc.

[106946] Nat. Acad. Sci. 93: 14821–14826, 1996.

[106947] 18798.

[106948]Hohenschutz, C.; Eich, P.; Friedl, W.; Waheed, A.; Conzelmann,

[106949]E.; Propping, P.: Pseudodeficiency of arylsulfatase A: a common genetic

[106950]polymorphism with possible disease implications. Hum. Genet. 82:

[106951]45–48, 1989.

[106952]18799.

[106953]Hohenschutz, C.; Friedl, W.; Schlor, K.–H.; Waheed, A.; Conzelmann,

[106954]E.; Sandhoff, K.; Propping, P.: Probable metachromatic leukodystrophy/pseudodeficiency

[106955]compound heterozygote at the arylsulfatase A locus with neurological

[106956]and psychiatric symptomatology. Am. J. Med. Genet. 31: 169–175,

[106957]1988.

[106958]18800.

[106959]Holve, S.; Hu, D.; McCandless, S. E.: Metachromatic leukodystrophy

[106960]in the Navajo: fallout of the American–Indian Wars of the nineteenth

[106961]century. Am. J. Med. Genet. 101: 203–208, 2001.

[106962]18801.

[106963]Honke, K.; Kobayashi, T.; Fujii, T.; Gasa, S.; Xu, M.; Takamaru,

[106964]Y.; Kondo, R.; Tsuji, S.; Makita, A.: An adult-type metachromatic

[106965]leukodystrophy caused by substitution of serine for glycine-122 in

[106966]arylsulfatase A. Hum. Genet. 92: 451–456, 1993.

[106967]18802.

[106968]Hors-Cayla, M. C.; Heuertz, S.; Van Cong, N.; Weil, D.; Frezal,

[106969]J.: Confirmation of the assignment of the gene for arylsulfatase

[106970]A to chromosome 22 using somatic cell hybrids. Hum. Genet. 49: 33–39,

[106971]1979.

[106972]18803.

[106973]Jervis, G. A.: Infantile metachromatic leukodystrophy (Greenfield's

[106974]disease). J. Neuropath. Exp. Neurol. 19: 323–341, 1960.

[106975]18804.

[106976]Kaback, M. M.; Howell, R. R.: Infantile metachromatic leukodystrophy:

[106977]heterozygote detection in skin fibroblasts and possible applications

[106978]to intrauterine diagnosis. New Eng. J. Med. 282: 1336–1340, 1970.

[106979]18805.

[106980]Kappler, J.; Sommerlade, H. J.; von Figura, K.; Gieselmann, V.

[106981]: Complex arylsulfatase A alleles causing metachromatic leukodystrophy. Hum.

[106982]Mutat. 4: 119–127, 1994.

[106983]18806.

[106984]Kappler, J.; von Figura, K.; Gieselmann, V.: Late-onset metachromatic

[106985]leukodystrophy: molecular pathology in two siblings. Ann. Neurol. 31:

[106986]256–261, 1992.

[106987]18807.

[106988]Kihara, H.: Genetic heterogeneity in metachromatic

leukodystrophy. Am.

[106989]J. Hum. Genet. 34: 171-181, 1982.

[106990]18808.

[106991]Kihara, H.; Fluharty, A. L.; O'Brien, J. S.; Fish, C. H.:

Metachromatic

[106992]leukodystrophy caused by a partial cerebroside sulfatase defect. Clin.

[106993]Genet. 21: 253-261, 1982.

[106994]18809.

[106995]Kihara, H.; Meek, W. E.; Fluharty, A. L.: Attenuated activities

[106996]and structural alterations of arylsulfatase A in tissues from subjects

[106997]with pseudo arylsulfatase A deficiency. Hum. Genet. 74: 59-62, 1986.

[106998]18810.

[106999]Kohn, H.; Manowitz, P.; Miller, M.; Kling, A.: Neuropsychological

[107000]deficits in obligatory heterozygotes for metachromatic leukodystrophy. Hum.

[107001]Genet. 79: 8-12, 1988.

[107002]18811.

[107003]Kondo, R.; Wakamatsu, N.; Yoshino, H.; Fukuhara, N.; Miy-
atake,

[107004]T.; Tsuji, S.: Identification of a mutation in the arylsulfa-
tase

[107005]A gene of a patient with adult-type metachromatic
leukodystrophy. Am.

[107006]J. Hum. Genet. 48: 971-978, 1991.

[107007]18812.

[107008]Kreysing, J.; Bohne, W.; Bosenberg, C.; Marchesini, S.;
Turpin,

[107009]J. C.; Baumann, N.; von Figura, K.; Gieselmann, V.: High
residual

[107010]aryl sulfatase A (ARSA) activity in a patient with late-
infantile metachromatic

[107011]leukodystrophy. Am. J. Hum. Genet. 53: 339–346, 1993.

[107012]18813.

[107013]Krivit, W.; Shapiro, E.; Kennedy, W.; Lipton, M.; Lockman, L.;

[107014]Smith, S.; Summers, C. G.; Wenger, D. A.; Tsai, M. Y.; Ramsay, N.

[107015]K. C.; Kersey, J. H.; Yao, J. K.; Kaye, E.: Treatment of late infantile

[107016]metachromatic leukodystrophy by bone marrow transplantation. New

[107017]Eng. J. Med. 322: 28–32, 1990.

[107018]18814.

[107019]Langenbeck, U.; Dunker, P.; Heipertz, R.; Pilz, H.: Inheritance

[107020]of metachromatic leukodystrophy. (Letter) Am. J. Hum. Genet. 29:

[107021]639–640, 1977.

[107022]18815.

[107023]Li, Z. G.; Waye, J. S.; Chang, P. L.: Diagnosis of arylsulphatase

[107024]A deficiency. Am. J. Med. Genet. 43: 976–982, 1992.

[107025]18816.

[107026]Lugowska, A.; Berger, J.; Tylki-Szymanska, A.; Czartowska, B.;

[107027]Loschl, B.; Molzer, B.: High prevalence of I179S mutation in patients

[107028]with late-onset metachromatic leukodystrophy. (Letter) Clin. Genet. 61:

[107029]389–390, 2002.

[107030]18817.

[107031]Lyon, G.; Arthiu, M.; Thieffry, S.: Leucodystrophie metachromatique

[107032]infantile familiale: etude de deux observations, dont une avec examen

[107033]anatomique et chimique. Rev. Neurol. 104: 508–533, 1961.

[107034]18818.

[107035]Masters, P. L.; MacDonald, W. B.; Ryan, M. M. P.; Cumings, J.

[107036]N.: Familial leucodystrophy. Arch. Dis. Child. 39: 345–355, 1964.

[107037]18819.

- [107038] Moser, H. W.: Sulfatide lipidosis: metachromatic leukodys-
trophy. In:
- [107039] Stanbury, J. B.; Wyngaarden, J. B.; Fredrickson, D. S.: The
Metabolic
- [107040] Basis of Inherited Disease. New York: McGraw-Hill (pub.)
(3rd
- [107041] ed.): 1972. Pp. 688-729.
- [107042] 18820.
- [107043] Kestila, M.; Lenkkeri, U.; Mannikko, M.; Lamerdin, J.; Mc-
Cready,
- [107044] P.; Putaala, H.; Ruotsalainen, V.; Morita, T.; Nissinen, M.;
Herva,
- [107045] R.; Kashtan, C. E.; Peltonen, L.; Holmberg, C.; Olsen, A.;
Tryggvason,
- [107046] K.: Positionally cloned gene for a novel glomerular pro-
tein--nephrin--is
- [107047] mutated in congenital nephrotic syndrome. Molec. Cell 1:
575-582,
- [107048] 1998.
- [107049] 18821.
- [107050] Muller, D.; Pilz, H.; Ter Meulen, V.: Studies on adult
metachromatic

[107051]leukodystrophy. I. Clinical, morphological and histochemical observations

[107052]in two cases. J. Neurol. Sci. 9: 567–584, 1969.

[107053]18822.

[107054]Narahara, K.; Takahashi, Y.; Murakami, M.; Tsuji, K.; Yokoyama,

[107055]Y.; Murakami, R.; Ninomiya, S.; Seino, Y.: Terminal 22q deletion

[107056]associated with a partial deficiency of arylsulphatase A. J. Med.

[107057]Genet. 29: 432–433, 1992.

[107058]18823.

[107059]Nelson, P. V.; Carey, W. F.; Morris, C. P.: Population frequency

[107060]of the arylsulphatase A pseudo-deficiency allele. Hum. Genet. 87:

[107061]87–88, 1991.

[107062]18824.

[107063]Nicholls, R. G.; Roy, A. G.: Arylsulfatases. In: Boyer, P. D.

[107064]: The Enzymes. New York: Academic Press (pub.) 5: 1971. Pp. 21–41.

[107065]18825.

[107066]Ott, R.; Waye, J. S.; Chang, P. L.: Evolutionary origins of
two

[107067]tightly linked mutations in arylsulfatase-A pseudodeficiency. Hum.

[107068]Genet. 101: 135–140, 1997.

[107069]18826.

[107070]Pastor-Soler, N. M.; Rafi, M. A.; Hoffman, J. D.; Hu, D.;
Wenger,

[107071]D. A.: Metachromatic leukodystrophy in the Navajo Indian
population:

[107072]a splice site mutation in intron 4 of the arylsulfatase A
gene. Hum.

[107073]Mutat. 4: 199–207, 1994.

[107074]18827.

[107075]Percy, A. K.; Brady, R. O.: Metachromatic leukodystrophy:
diagnosis

[107076]with samples of venous blood. Science 161: 594–595,
1968.

[107077]18828.

[107078]Percy, A. K.; Kaback, M. M.: Infantile and adult-onset

metachromatic

[107079]leukodystrophy: biochemical comparisons and predictive diagnosis. New

[107080]Eng. J. Med. 285: 785-787, 1971.

[107081]18829.

[107082]Percy, A. K.; Kaback, M. M.; Herndon, R. M.: Metachromatic leukodystrophy:

[107083]comparison of early- and late-onset forms. Neurology 27: 933-941,

[107084]1977.

[107085]18830.

[107086]Pilz, H.; Duensing, I.; Heipertz, R.; Seidel, D.; Lowitsch, K.;

[107087]Hopf, H. C.; Goebel, H. H.: Adult metachromatic leukodystrophy. I.

[107088]Clinical manifestation in a female aged 44 years, previously diagnosed

[107089]in the preclinical state. Europ. Neurol. 15: 301-307, 1977.

[107090]18831.

[107091]Pilz, H.; Muller, D.: Studies on adult metachromatic leukodystrophy.

[107092]II. Biochemical aspects of adult cases of metachromatic

leukodystrophy. J.

[107093]Neurol. Sci. 9: 585–595, 1969.

[107094]18832.

[107095]Poenaru, L.; Castelnau, L.; Besancon, A.–M.; Nicolesco, H.;
Akli,

[107096]S.; Theophil, D.: First trimester prenatal diagnosis of
metachromatic

[107097]leukodystrophy on chorionic villi by 'immunoprecipitation–elec-
trophoresis.'. J.

[107098]Inherit. Metab. Dis. 11: 123–130, 1988.

[107099]18833.

[107100]Polten, A.; Fluharty, A. L.; Fluharty, C. B.; Kappler, J.; von

[107101]Figura, K.; Gieselmann, V.: Molecular basis of different
forms of

[107102]metachromatic leukodystrophy. New Eng. J. Med. 324:
18–22, 1991.

[107103]18834.

[107104]Porter, M. T.; Fluharty, A. L.; Kihara, H.: Correction of ab-
normal

[107105]cerebroside sulfate metabolism in cultured metachromatic
leukodystrophy

[107106]fibroblasts. Science 172: 1263–1265, 1971.

[107107]18835.

[107108]Porter, M. T.; Fluharty, A. L.; Trammell, J.; Kihara, H.: A correlation

[107109]of intracellular cerebroside sulfatase activity in fibroblasts with

[107110]latency in metachromatic leukodystrophy. Biochem. Biophys. Res. Commun. 44:

[107111]660–666, 1971.

[107112]18836.

[107113]Propping, P.; Friedl, W.; Huschka, M.; Schlor, K.–H.; Reimer,

[107114]F.; Lee–Vaupel, M.; Conzelmann, E.; Sandhoff, K.: The influence of

[107115]low arylsulfatase A activity on neuropsychiatric morbidity: a large–scale

[107116]screening in patients. Hum. Genet. 74: 244–248, 1986.

[107117]18837.

[107118]Quigley, H. A.; Green, W. R.: Clinical and ultrastructural ocular

[107119]histopathologic studies of adult–onset metachromatic

leukodystrophy. Am.

[107120]J. Ophthalmol. 82: 472-479, 1976.

[107121]18838.

[107122]Regis, S.; Corsolini, F.; Stroppiano, M.; Cusano, R.; Filo-
carno,

[107123]M.: Contribution of arylsulfatase A mutations located on
the same

[107124]allele to enzyme activity reduction and metachromatic
leukodystrophy

[107125]severity. Hum. Genet. 110: 351-355, 2002.

[107126]18839.

[107127]Ricketts, M. H.; Poretz, R. D.; Manowitz, P.: The R496H
mutation

[107128]of arylsulfatase A does not cause metachromatic
leukodystrophy. Hum.

[107129]Mutat. 12: 238-239, 1998.

[107130]18840.

[107131]Sanguinetti, N.; Marsh, J.; Jackson, M.; Fensom, A. H.;
Warren,

[107132]R. C.; Rodeck, C. H.: The arylsulphatases of chorionic villi:
potential

- [107133] problems in the first-trimester diagnosis of metachromatic leucodystrophy
- [107134] and Maroteaux-Lamy disease. Clin. Genet. 30: 302-308, 1986.
- [107135] 18841.
- [107136] Schaap, T.; Zlotogora, J.; Elian, E.; Barak, Y.; Bach, G.: The
- [107137] genetics of the aryl sulfatase A locus. Am. J. Hum. Genet. 33: 531-539, 1981.
- [107138]
- [107139]
- [107140] 100. Schutta, H. S.; Pratt, R. T. C.; Metz, H.; Evans, K. A.; Carter,
- [107141] C. O.: A family study of the late infantile and juvenile forms of
- [107142] metachromatic leukodystrophy. J. Med. Genet. 3: 86-91, 1966.
- [107143]
- [107144] 101. Shen, N.; Li, Z.-G.; Waye, J. S.; Francis, G.; Chang, P. L.:
- [107145] Complications in the genotypic molecular diagnosis of pseudo arylsulfatase

[107146]A deficiency. Am. J. Med. Genet. 45: 631–637, 1993.

[107147]

[107148]102. Sourander, P.; Svennerholm, L.: Sulphatide lipidosis
in the

[107149]adult with the clinical picture of progressive organic de-
mentia with

[107150]epileptic seizures. Acta Neuropath. 1: 384–396, 1962.

[107151]

[107152]103. Stein, C.; Gieselmann, V.; Kreysing, J.; Schmidt, B.;
Pohlmann,

[107153]R.; Waheed, A.; Meyer, H. E.; O'Brien, J. S.; von Figura, K.:
Cloning

[107154]and expression of human arylsulfatase A. J. Biol. Chem.
264: 1252–1259,

[107155]1989.

[107156]

[107157]104. Stumpf, D. A.; Austin, J.: Metachromatic leukodystro-
phy (MLD).

[107158]IX. Qualitative and quantitative differences in urinary aryl-
sulfatase

[107159]A in different forms of MLD. Arch. Neurol. 24: 117–124,

1971.

[107160]

[107161]105. Tonnesen, T.; Bro, P. V.; Nielsen, K. B.; Lykkelund, C.:

Metachromatic

[107162]leukodystrophy and pseudoarylsulfatase A deficiency in a

Danish family. Acta

[107163]Paediat. Scand. 72: 175–178, 1983.

[107164]

[107165]106. Tonnesen, T.; Vrang, C.; Wiesmann, U. N.; Christom-

anou, H.; Lou,

[107166]H. O.: Atypical metachromatic leukodystrophy? Problems
with the biochemical

[107167]diagnosis. Hum. Genet. 67: 170–173, 1984.

[107168]

[107169]107. Van Bogaert, L. V.; Dewulf, A.: Diffuse progressive

leukodystrophy

[107170]in the adult with production of metachromatic degenerative
products

[107171](Alzheimer-Baroncini). Arch. Neurol. Psychiat. 42:

1083–1097, 1939.

[107172]

[107173]108. von Figura, K.; Steckel, F.; Conary, J.; Hasilik, A.; Shaw, E.

[107174]: Heterogeneity in late-onset metachromatic leukodystrophy: effect

[107175]of inhibitors of cysteine proteinases. Am. J. Hum. Genet. 39: 371-382,

[107176]1986.

[107177]

[107178]109. von Figura, K.; Steckel, F.; Hasilik, A.: Juvenile and adult

[107179]metachromatic leukodystrophy: partial restoration of aryl-sulfatase

[107180]A (cerebroside sulfatase) activity by inhibitors of thiol proteinases. Proc.

[107181]Nat. Acad. Sci. 80: 6066-6070, 1983.

[107182]

[107183]110. Waheed, A.; Steckel, F.; Hasilik, A.; von Figura, K.: Two allelic

[107184]forms of human arylsulfatase A with different numbers of asparagine-linked

[107185]oligosaccharides. Am. J. Hum. Genet. 35: 228-233, 1983.

[107186]

[107187]111. Waltz, G.; Harik, S. I.; Kaufman, B.: Adult metachromatic leukodystrophy:

[107188]value of computed tomographic scanning and magnetic resonance imaging

[107189]of the brain. Arch. Neurol. 44: 225–227, 1987.

[107190]

[107191]112. Yatziv, S.; Russell, A.: An unusual form of metachromatic leukodystrophy

[107192]in three siblings. Clin. Genet. 19: 222–227, 1981.

[107193]

[107194]113. Zlotogora, J.; Bach, G.: Deficiency of lysosomal hydrolases

[107195]in apparently healthy individuals. Am. J. Med. Genet. 14: 73–80,

[107196]1983.

[107197]

[107198]114. Zlotogora, J.; Bach, G.; Barak, Y.; Elian, E.: Metachromatic

[107199]leukodystrophy in the Habbanite Jews: high frequency in a genetic

[107200]isolate and screening for heterozygotes. Am. J. Hum.
Genet. 32:

[107201]663–669, 1980.

[107202]

[107203]115. Zlotogora, J.; Furman–Shaharabani, Y.; Harris, A.;
Barth, M.

[107204]L.; von Figura, K.; Gieselmann, V.: A single origin for the
most

[107205]frequent mutation causing late infantile metachromatic
leucodystrophy. J.

[107206]Med. Genet. 31: 672–674, 1994.

[107207]

[107208]116. Zlotogora, J.; Gieselman, V.; von Figura, K.; Zeigler,
M.; Bach,

[107209]G.: Late infantile metachromatic leukodystrophy in Israel.
Biomed.

[107210]Pharmacother. 48: 347–350, 1994.

[107211]18842.

[107212]Rozet, J.–M.; Gerber, S.; Ghazi, I. Perrault, I.; Ducroq, D.;

[107213]Souied, E.; Cabot, A.; Dufier, J.–L.; Munnich, A.; Kaplan, J.:
Mutations

[107214] of the retinal specific ATP binding transporter gene (ABCR)
in a single
[107215] family segregating both autosomal recessive retinitis pig-
mentosa RP19
[107216] and Stargardt disease: evidence of clinical heterogeneity
at this
[107217] locus. J. Med. Genet. 36: 447-451, 1999.
[107218] 18843.
[107219] Rozet, J.-M.; Gerber, S.; Souied, E.; Perrault, I.; Chatelin,
[107220] S.; Ghazi, I.; Leowski, C.; Dufier, J.-L.; Munnich, A.; Ka-
plan, J.
[107221]: Spectrum of ABCR gene mutations in autosomal reces-
sive macular dystrophies. Europ.
[107222] J. Hum. Genet. 6: 291-295, 1998.
[107223] 18844.
[107224] Coppa, G. V.; Giorgi, P. L.; Felici, L.; Gabrielli, O.; Donti,
[107225] E.; Bernasconi, S.; Kresse, H.; Paschke, E.; Mastropaolo, C.:
Clinical
[107226] heterogeneity in Sanfilippo disease
(mucopolysaccharidosis III) type
[107227] D: presentation of two new cases. Europ. J. Pediat. 140:

130–133,

[107228]1983.

[107229]18845.

[107230]Dhoot, G. K.; Gustafsson, M. K.; Ai, X.; Sun, W.; Standiford, D.

[107231]M.; Emerson, C. P., Jr.: Regulation of Wnt signaling and embryo patterning

[107232]by an extracellular sulfatase. Science 293: 1663–1666, 2001.

[107233]18846.

[107234]Freeman, C.; Clements, P. R.; Hopwood, J. J.: Human liver N-acetylglucosamine-6-sulphate

[107235]sulphatase: purification and characterization. Biochem. J. 246:

[107236]347–354, 1987.

[107237]18847.

[107238]Freeman, C.; Hopwood, J. J.: Human liver N-acetylglucosamine-6-sulphate

[107239]sulphatase: catalytic properties. Biochem. J. 246: 355–365, 1987.

[107240]18848.

[107241]Gatti, R.; Borrone, C.; Durand, P.; De Virgiliis, S.; Sanna, G.;

[107242]Cao, A.; von Figura, K.; Kresse, H.; Paschke, E.: Sanfilippo type

[107243]D disease: clinical findings in two patients with a new variant of

[107244]mucopolysaccharidosis III. Europ. J. Pediat. 138: 168–171, 1982.

[107245]18849.

[107246]Kaplan, P.; Wolfe, L. S.: Sanfilippo syndrome type D. J. Pediat. 110:

[107247]267–271, 1987.

[107248]18850.

[107249]Kresse, H.; Paschke, E.; von Figura, K.; Gilberg, W.; Fuchs, W.

[107250]: Sanfilippo disease type D: deficiency of N-acetylglucosamine-6-sulfate

[107251]sulfatase required for heparan sulfate degradation. Proc. Nat. Acad.

[107252]Sci. 77: 6822–6826, 1980.

[107253]18851.

- [107254]Robertson, D. A.; Callen, D. F.; Baker, E. G.; Morris, C. P.; Hopwood, J. J.: Chromosomal localization of the gene for human glucosamine-6-sulphatase to 12q14. Hum. Genet. 79: 175-178, 1988.
- [107257]18852.
- [107258]Robertson, D. A.; Freeman, C.; Nelson, P. V.; Morris, C. P.; Hopwood, J. J.: Human glucosamine-6-sulfatase cDNA reveals homology with steroid sulfatase. Biochem. Biophys. Res. Commun. 157: 218-224, 1988.
- [107261]18853.
- [107262]Siciliano, L.; Fiumara, A.; Pavone, L.; Freeman, C.; Robertson, D.; Morris, C. P.; Hopwood, J. J.; Di Natale, P.; Musumeci, S.; Horwitz, A. L.: Sanfilippo syndrome type D in two adolescent sisters. J. Med. Genet. 28: 402-405, 1991.
- [107266]18854.

[107267]Thompson, J. N.; Jones, M. Z.; Dawson, G.; Huffman, P. S.:

N-acetylglucosamine

[107268]6-sulphatase deficiency in a Nubian goat: a model of San-filippo syndrome

[107269]type D (mucopolysaccharidosis IIID). J. Inherit. Metab. Dis.
15:

[107270]760-768, 1992.

[107271]18855.

[107272]Baker, E.; Guo, X.-H.; Orsborn, A. M.; Sutherland, G. R.;
Callen,

[107273]D. F.; Hopwood, J. J.; Morris, C. P.: The Morquio A syn-drome (mucopolysaccharidosis

[107274]IVA) gene maps to 16q24.3. Am. J. Hum. Genet. 52:
96-98, 1993.

[107275]18856.

[107276]Beck, M.; Braun, S.; Coerdts, W.; Merz, E.; Young, E.; Sewell,
A.

[107277]C.: Fetal presentation of Morquio disease type A. Prenatal
Diag. 12:

[107278]1019-1029, 1992.

[107279]18857.

[107280]Blaw, M. E.; Langer, L. O., Jr.: Spinal cord compression in
Morquio–Brailsford's

[107281]disease. J. Pediat. 74: 593–600, 1969.

[107282]18858.

[107283]Brailsford, J. F.: Chondro–osteo–dystrophy: roentgeno–
graphic and

[107284]clinical features of child with dislocation of vertebrae. Am.
J.

[107285]Surg. 7: 404–410, 1929.

[107286]18859.

[107287]Bunge, S.; Kleijer, W. J.; Tylki–Szymanska, A.; Steglich, C.;
Beck,

[107288]M.; Tomatsu, S.; Fukuda, S.; Poorthuis, B. J. H. M.; Czarto–
ryska,

[107289]B.; Orii, T.; Gal, A.: Identification of 31 novel mutations in
the

[107290]N–acetylgalactosamine–6–sulfatase gene reveals exces–
sive allelic heterogeneity

[107291]among patients with Morquio A syndrome. Hum. Mutat.
10: 223–232,

[107292]1997.

[107293]18860.

[107294]Engel, A. G.; Lambert, E. H.; Gomez, M. R.: A new myasthenic syndrome

[107295]with end-plate acetylcholinesterase deficiency, small nerve terminals,

[107296]and reduced acetylcholine release. Ann. Neurol. 1: 315-330, 1977.

[107297]18861.

[107298]Faiella, A.; Brunelli, S.; Granata, T.; D'Incerti, L.; Cardini,

[107299]R.; Lenti, C.; Battaglia, G.; Boncinelli, E.: A number of schizencephaly

[107300]patients including 2 brothers are heterozygous for germline mutations

[107301]in the homeobox gene EMX2. Europ. J. Hum. Genet. 5: 186-190, 1997.

[107302]18862.

[107303]Li, Y.-T.; Hirabayashi, Y.; Li, S.-C.: Differentiation of two

[107304]variants of type-AB Gm-2-gangliosidosis using chromogenic substrates. Am.

[107305]J. Hum. Genet. 35: 520-522, 1983.

[107306]18863.

- [107307]Rousseau-Merck, M.-F.; Duro, D.; Berger, R.; Thiesen, H. J.: Chromosomal
- [107308]localization of two KOX zinc finger genes on chromosome bands 7q21-q22. Ann.
- [107309]Genet. 38: 81-84, 1995.
- [107310]18864.
- [107311]Tommerup, N.; Vissing, H.: Isolation and fine mapping of 16 novel
- [107312]human zinc finger-encoding cDNAs identify putative candidate genes
- [107313]for developmental and malignant disorders. Genomics 27: 259-264,
- [107314]1995.
- [107315]18865.
- [107316]Canman, C. E.; Radany, E. H.; Parsels, L. A.; Davis, M. A.; Lawrence,
- [107317]T. S.; Maybaum, J.: Induction of resistance to fluoro-deoxyuridine
- [107318]cytotoxicity and DNA damage in human tumor cells by expression of
- [107319]Escherichia coli deoxyuridinetriphosphatase. Cancer Res. 54: 2296-2298,

[107320]1994.

[107321]18866.

[107322]Canman, C. E.; Tang, H. Y.; Normolle, D. P.; Lawrence, T. S.; Maybaum,

[107323]J.: Variations in patterns of DNA damage induced in human colorectal

[107324]tumor cells by 5-fluorodeoxyuridine: implications for mechanisms of

[107325]resistance and cytotoxicity. Proc. Nat. Acad. Sci. 89: 10474-10478,

[107326]1992.

[107327]18867.

[107328]Chu, R.; Lin, Y.; Rao, M. S.; Reddy, J. K.: Cloning and identification

[107329]of rat deoxyuridine triphosphatase as an inhibitor of peroxisome proliferator-activated

[107330]receptor alpha. J. Biol. Chem. 271: 27670-27676, 1996.

[107331]18868.

[107332]Cohen, D.; Heng, H. H. Q.; Shi, X.-M.; McIntosh, E. M.; Tsui, L.-C.;

[107333]Pearlman, R. E.: Assignment of the human dUTPase gene

(DUT) to chromosome

[107334]15q15-q21.1 by fluorescence in situ hybridization. Genomics 40:

[107335]213-215, 1997.

[107336]18869.

[107337]el-Hajj, H. H.; Zhang, H.; Weiss, B.: Lethality of a dut (deoxyuridine

[107338]triphosphatase) mutation in Escherichia coli. J. Bacteriol. 170:

[107339]1069-1075, 1988.

[107340]18870.

[107341]Ladner, R. D.; Caradonna, S. J.: The human dUTPase gene encodes

[107342]both nuclear and mitochondrial isoforms: differential expression of

[107343]the isoforms and characterization of a cDNA encoding the mitochondrial

[107344]species. J. Biol. Chem. 272: 19072-19080, 1997.

[107345]18871.

[107346]Ladner, R. D.; McNulty, D. E.; Carr, S. A.; Roberts, G. D.; Caradonna,

[107347]S. J.: Characterization of distinct nuclear and mitochondrial forms
[107348]of human deoxyuridine triphosphate nucleotidohydrolase.
J. Biol.
[107349]Chem. 271: 7745–7751, 1996.
[107350]18872.
[107351]Lindahl, T.: DNA repair enzymes. Annu. Rev. Biochem. 51:
61–87,
[107352]1982.
[107353]18873.
[107354]McIntosh, E. M.; Ager, D. D.; Gadsden, M. H.; Haynes, R.
H.: Human
[107355]dUTP pyrophosphatase: cDNA sequence and potential biological importance
[107356]of the enzyme. Proc. Nat. Acad. Sci. 89: 8020–8024,
1992. Note:
[107357]Erratum: Proc. Nat. Acad. Sci.:90: 4328 only, 1993.
[107358]18874.
[107359]Boring, L.; Gosling, J.; Cleary, M.; Charo, I. F.: Decreased
lesion
[107360]formation in CCR2 –/– mice reveals a role for chemokines

in the initiation

[107361]of atherosclerosis. Nature 394: 894–897, 1998.

[107362]18875.

[107363]Charo, I. F.; Myers, S. J.; Herman, A.; Franci, C.; Connolly, A.

[107364]J.; Coughlin, S. R.: Molecular cloning and functional expression

[107365]of two monocyte chemoattractant protein 1 receptors reveals alternative

[107366]splicing of the carboxyl-terminal tails. Proc. Nat. Acad. Sci. 91:

[107367]2752–2756, 1994.

[107368]18876.

[107369]Sun, D.–S.; Chang, A. C.; Jenkins, N. A.; Gilbert, D. J.; Copeland,

[107370]N. G.; Chang, N.–C. A.: Identification, molecular characterization,

[107371]and chromosomal localization of the cDNA encoding a novel leucine

[107372]zipper motif-containing protein. Genomics 36: 54–62, 1996.

[107373]18877.

[107374]Barrett, T. E.; Savva, R.; Panayotou, G.; Barlow, T.; Brown, T.;

[107375]Jiricny, J.; Pearl, L. H.: Crystal structure of a G:T/U mismatch-specific

[107376]DNA glycosylase: mismatch recognition by complementary-strand interactions. Cell 92:

[107377]117-129, 1998.

[107378]18878.

[107379]De Gregorio, L.; Gallinari, P.; Gariboldi, M.; Manenti, G.; Pierotti,

[107380]M. A.; Jiricny, J.; Dragani, T. A.: Genetic mapping of thymine DNA

[107381]glycosylase (Tdg) gene and of one pseudogene in the mouse. Mammalian

[107382]Genome 7: 909-910, 1996.

[107383]18879.

[107384]Lindahl, T.: DNA repair enzymes. Ann. Rev. Biochem. 51: 61-87,

[107385]1982.

[107386]18880.

[107387]Neddermann, P.; Gallinari, P.; Lettieri, T.; Schmid, D.;

Truong,

[107388]O.; Hsuan, J. J.; Wiebauer, K.; Jiricny, J.: Cloning and expression

[107389]of human G/T mismatch-specific thymine-DNA glycosylase. J. Biol.

[107390]Chem. 271: 12767-12774, 1996.

[107391]18881.

[107392]Sard, L.; Tornielli, S.; Gallinari, P.; Minoletti, F.; Jiricny,

[107393]J.; Lettieri, T.; Pierotti, M. A.; Sozzi, G.; Radice, P.: Chromosomal

[107394]localizations and molecular analysis of TDG gene-related sequences. Genomics 44:

[107395]222-226, 1997.

[107396]18882.

[107397]DiMarco, S. P.; Glover, T. W.; Miller, D. E.; Reines, D.; Warren,

[107398]S. T.: Transcription elongation factor SII (TCEA) maps to human chromosome

[107399]3p22-p21.3. Genomics 36: 185-188, 1996.

[107400]18883.

[107401]Park, H.; Baek, K.; Jeon, C.; Agarwal, K.; Yoo, O.: Charac-

terization

[107402]of the gene encoding the human transcriptional elongation factor TFIIIS. Gene 139:

[107403]263–267, 1994.

[107404]18884.

[107405]Reines, D.: Nascent RNA cleavage by transcription elongation complexes. In:

[107406]Conaway, R. C.; Conaway, J. W. (eds.): Transcription: Mechanisms

[107407]and Regulation. New York: Raven Press 1994. Pp. 263–278.

[107408]18885.

[107409]Thomas, M. J.; Platas, A. A.; Hawley, D. K.: Transcriptional fidelity

[107410]and proofreading by RNA polymerase II. Cell 93: 627–637, 1998.

[107411]18886.

[107412]Applequist, S. E.; Selg, M.; Raman, C.; Jack, H.–M.: Cloning and

[107413]characterization of HUPF1, a human homolog of the Saccharomyces cerevisiae

[107414]nonsense mRNA–reducing UPF1 protein. Nucleic Acids

Res. 25: 814–821,

[107415]1997.

[107416]18887.

[107417]Medghalchi, S. M.; Frischmeyer, P. A.; Mendell, J. T.; Kelly, A.

[107418]G.; Lawler, A. M.; Dietz, H. C.: Rent1, a trans-effector of nonsense-mediated

[107419]mRNA decay, is essential for mammalian embryonic viability. Hum.

[107420]Molec. Genet. 10: 99–105, 2001.

[107421]18888.

[107422]Mendell, J. T.; ap Rhys, C. M. J.; Dietz, H. C.: Separable roles

[107423]for rent1/hUpf1 in altered splicing and decay of nonsense transcripts. Science 298:

[107424]419–371, 2002.

[107425]18889.

[107426]Perlick, H. A.; Medghalchi, S. M.; Spencer, F. A.; Dietz, H. C.

[107427]: Cloning and characterization of a human regulator of nonsense transcript

[107428]stability. (Abstract) Am. J. Hum. Genet. 59 (suppl.): A32

only,

[107429]1996.

[107430]18890.

[107431]Perlick, H. A.; Medghalchi, S. M.; Spencer, F. A.; Kendzior,
R.

[107432]J., Jr.; Dietz, H. C.: Mammalian orthologues of a yeast reg-
ulator

[107433]of nonsense transcript stability. Proc. Nat. Acad. Sci. 93:
10928-10932,

[107434]1996.

[107435]18891.

[107436]Sun, X.; Perlick, H. A.; Dietz, H. C.; Maquat, L. E.: A mu-
tated

[107437]human homologue to yeast Upf1 protein has a dominant-
negative effect

[107438]on the decay of nonsense-containing mRNAs in mam-
malian cells. Proc.

[107439]Nat. Acad. Sci. 95: 10009-10014, 1998.

[107440]18892.

[107441]Eng, C.; Myers, S. M.; Kogon, M. D.; Sanicola, M.; Hession,

C.;

- [107442]Cate, R. L.; Mulligan, L. M.: Genomic structure and chromosomal localization
[107443]of the human GDNFR- α gene. *Oncogene* 16: 597-601, 1998.
[107444]18893.
- [107445]GFR- α Nomenclature Committee: Nomenclature of GPI-linked
[107446]receptors for the GDNF ligand family. *Neuron* 19: 485 only, 1997.
[107447]18894.
- [107448]Gorodinsky, A.; Zimonjic, D. B.; Popescu, N. C.; Milbrandt, J.
[107449]: Assignment of the GDNF family receptor α -1 (GFRA1) to human
[107450]chromosome band 10q26 by in situ hybridization. *Cytogenet. Cell Genet.* 78:
[107451]289-290, 1997.
[107452]18895.
- [107453]Jing, S.; Wen, D.; Yu, Y.; Holst, P. L.; Luo, Y.; Fang, M.; Tamir,
[107454]R.; Antonio, L.; Hu, Z.; Cupples, R.; Louis, J.-C.; Hu, S.; Al-

trock,

[107455]B. W.; Fox, G. M.: GDNF-induced activation of the Ret protein tyrosine

[107456]kinase is mediated by GDNFR-alpha, a novel receptor for GDNF. Cell 85:

[107457]1113-1124, 1996.

[107458]18896.

[107459]Paratcha, G.; Ledda, F.; Baars, L.; Couplier, M.; Besset, V.; Anders,

[107460]J.; Scott, R.; Ibanez, C. F.: Released GFR-alpha-1 potentiates downstream

[107461]signaling, neuronal survival, and differentiation via a novel mechanism

[107462]of recruitment of c-Ret to lipid rafts. Neuron 29: 171-184, 2001.

[107463]18897.

[107464]Puliti, A.; Cinti, R.; Seri, M.; Ceccherini, I.; Romeo, G.: Assignment

[107465]of mouse Gfra1, the homologue of a new human HSCR candidate gene,

[107466]to the telomeric region of mouse chromosome 19. Cytogenet. Cell Genet. 78:

[107467]291–294, 1997.

[107468]18898.

[107469]Shefelbine, S. E.; Khorana, S.; Schultz, P. N.; Huang, E.;
Thobe,

[107470]N.; Hu, Z. J.; Fox, G. M.; Jing, S.; Cote, G. J.; Gagel, R. F.:
Mutational

[107471]analysis of the GDNF/RET–GDNFR–alpha signaling com-
plex in a kindred

[107472]with vesicoureteral reflux. Hum. Genet. 102: 474–478,
1998.

[107473]18899.

[107474]Bardelli, A.; Longati, P.; Albero, D.; Goruppi, S.; Schneider,

[107475]C.; Ponzetto, C.; Comoglio, P. M.: HGF receptor associates
with the

[107476]anti-apoptotic protein BAG-1 and prevents cell death.

EMBO J. 15:

[107477]6205–6212, 1996.

[107478]18900.

[107479]Takayama, S.; Kochel, K.; Irie, S.; Inazawa, J.; Abe, T.; Sato,

[107480]T.; Druck, T.; Huebner, K.; Reed, J. C.: Cloning of cDNAs
encoding

[107481]the human BAG1 protein and localization of the human
BAG1 gene to

[107482]chromosome 9p12. Genomics 35: 494–498, 1996.

[107483]18901.

[107484]Takayama, S.; Sato, T.; Krajewski, S.; Kochel, K.; Irie, S.;
Millan,

[107485]J. A.; Reed, J. C.: Cloning and functional analysis of BAG–1:
a novel

[107486]Bcl–2–binding protein with anti–cell death activity. Cell
80: 279–284,

[107487]1995.

[107488]18902.

[107489]Wang, H.–G.; Takayama, S.; Rapp, U. R.; Reed, J. C.: Bcl–2
interacting

[107490]protein, BAG–1, binds to and activates the kinase Raf–1.
Proc. Nat.

[107491]Acad. Sci. 93: 7063–7068, 1996.

[107492]18903.

[107493]Yang, X.; Pater, A.; Tang, S.–C.: Cloning and characteriza-
tion

[107494]of the human BAG–1 gene promoter: upregulation by tu–

mor-derived p53

[107495]mutants. *Oncogene* 18: 4546–4553, 1999.

[107496]18904.

[107497]Chen, H.; Chrast, R.; Rossier, C.; Gos, A.; Antonarakis, S.
E.;

[107498]Kudoh, J.; Yamaki, A.; Shindoh, N.; Maeda, H.; Minoshima,
S.; Shimizu,

[107499]N.: Single-minded and Down syndrome? (Letter) *Nature*
Genet. 10:

[107500]9–10, 1995.

[107501]18905.

[107502]Chrast, R.; Scott, H. S.; Madani, R.; Huber, L.; Wolfer, D. P.;

[107503]Prinz, M.; Aguzzi, A.; Lipp, H.-P.; Antonarakis, S. E.: Mice
trisomic

[107504]for a bacterial artificial chromosome with the single-
minded 2 gene

[107505](Sim2) show phenotypes similar to some of those present
in the partial

[107506]trisomy 16 mouse models of Down syndrome. *Hum.*
Molec. Genet. 9:

[107507]1853–1864, 2000.

[107508]18906.

[107509]Dahmane, N.; Charron, G.; Lopes, C.; Yaspo, M.-L.; Maunoury, C.;

[107510]Decorte, L.; Sinet, P.-M.; Bloch, B.; Delabar, J.-M.: Down syndrome-critical

[107511]region contains a gene homologous to *Drosophila* sim expressed during

[107512]rat and human central nervous system development. Proc. Nat. Acad.

[107513]Sci. 92: 9191-9195, 1995.

[107514]18907.

[107515]Ema, M.; Ikegami, S.; Hosoya, T.; Mimura, J.; Ohtani, H.; Nakao,

[107516]K.; Inokuchi, K.; Katsuki, M.; Fujii-Kuriyama, Y.: Mild impairment

[107517]of learning and memory in mice overexpressing the mSim2 gene located

[107518]on chromosome 16: an animal model of Down's syndrome. Hum. Molec.

[107519]Genet. 8: 1409-1415, 1999.

[107520]18908.

- [107521]Moffett, P.; Dayo, M.; Reece, M.; McCormick, M. K.; Pelletier,
J.: Characterization of msim, a murine homologue of the
Drosophila
sim transcription factor. Genomics 35: 144–155, 1996.
18909.
- [107525]Muenke, M.; Bone, L. J.; Mitchell, H. F.; Hart, I.; Walton, K.;
Hall-Johnson, K.; Ippel, E. F.; Dietz-Band, J.; Kvaloy, K.;
Fan, C.-M.;
Tessier-Lavigne, M.; Patterson, D.: Physical mapping of
the holoprosencephaly
critical region in 21q22.3, exclusion of SIM2 as a candidate gene
for holoprosencephaly, and mapping of SIM2 to a region
of chromosome
21 important for Down syndrome. Am. J. Hum. Genet. 57:
1074–1079,
1995.
- [107532]18910.
- [107533]Yamaki, A.; Noda, S.; Kudoh, J.; Shindoh, N.; Maeda, H.;
Minoshima,

[107534]S.; Kawasaki, K.; Shimizu, Y.; Shimizu, N.: The mammalian single-minded

[107535](SIM) gene: mouse cDNA structure and diencephalic expression indicate

[107536]a candidate gene for Down syndrome. Genomics 35: 136–143, 1996.

[107537]18911.

[107538]de Winter, J. P.; Leveille, F.; van Berkel, C. G. M.; Rooimans,

[107539]M. A.; van der Weel, L.; Steltenpool, J.; Demuth, I.; Morgan, N. V.;

[107540]Alon, N.; Bosnoyan–Collins, L.; Lightfoot, J.; Leegwater, P. A.; Waisfisz,

[107541]Q.; Komatsu, K.; Arwert, F.; Pronk, J. C.; Mathew, C. G.; Digweed,

[107542]M.; Buchwald, M.; Joenje, H.: Isolation of a cDNA representing the

[107543]Fanconi anemia complementation group E gene. Am. J. Hum. Genet. 67:

[107544]1306–1308, 2000. Note: Erratum: Am. J. Hum. Genet. 67: 1365 only,

[107545]2000.

[107546]18912.

[107547]Hirotune, S.; Fleck, M. W.; Gambello, M. J.; Bix, G. J.;
Chen,

[107548]A.; Clark, G. D.; Ledbetter, D. H.; McBain, C. J.; Wynshaw-
Boris,

[107549]A.: Graded reduction of Pafah1b1 (Lis1) activity results in
neuronal

[107550]migration defects and early embryonic lethality. Nature
Genet. 19:

[107551]333–339, 1998.

[107552]18913.

[107553]Kurahashi, H.; Sakamoto, M.; Ono, J.; Honda, A.; Okada,
S.; Nakamura,

[107554]Y.: Molecular cloning of the chromosomal breakpoint in
the LIS1 gene

[107555]of a patient with isolated lissencephaly and balanced
t(8;17). Hum.

[107556]Genet. 103: 189–192, 1998.

[107557]18914.

[107558]Liu, Z.; Steward, R.; Luo, L.: Drosophila Lis1 is required for

[107559]neuroblast proliferation, dendritic elaboration and axonal

transport. Nature

[107560]Cell Biol. 2: 776–783, 2000.

[107561]18915.

[107562]Lo Nigro, C.; Chong, S. S.; Smith, A. C. M.; Dobyns, W. B.;
Carrozzo,

[107563]R.; Ledbetter, D. H.: Point mutations and an intragenic
deletion

[107564]in LIS1, the lissencephaly causative gene in isolated
lissencephaly

[107565]sequence and Miller–Dieker syndrome. Hum. Molec.
Genet. 6: 157–164,

[107566]1997.

[107567]18916.

[107568]Reiner, O.; Bar–Am, I.; Sapir, T.; Shmueli, O.; Carrozzo, R.;

[107569]Lindsay, E. A.; Baldini, A.; Ledbetter, D. H.; Cahana, A.:
LIS2,

[107570]gene and pseudogene, homologous to LIS1 (lissencephaly
1), located

[107571]on the short and long arms of chromosome 2. Genomics
30: 251–256,

[107572]1995.

[107573]18917.

[107574]Sakamoto, M.; Ono, J.; Okada, S.; Masuno, M.; Nakamura, Y.; Kurahashi,

[107575]H.: Alteration of the LIS1 gene in Japanese patients with isolated

[107576]lissencephaly sequence or Miller–Dieker syndrome. Hum. Genet. 103:

[107577]586–589, 1998.

[107578]18918.

[107579]Smith, D. S.; Niethammer, M.; Ayala, R.; Zhou, Y.; Gambello, M.

[107580]J.; Wynshaw–Boris, A.; Tsai, L.–H.: Regulation of cytoplasmic dynein

[107581]behaviour and microtubule organization by mammalian Lis1. Nature

[107582]Cell Biol. 2: 767–775, 2000.

[107583]18919.

[107584]Sosa–Pineda, B.; Wigle, J. T.; Oliver, G.: Hepatocyte migration

[107585]during liver development requires Prox1. Nature Genet. 25: 254–255,

[107586]2000.

[107587]18920.

[107588]Wigle, J. T.; Chowdhury, K.; Gruss, P.; Oliver, G.: Prox1
function

[107589]is crucial for mouse lens-fibre elongation. Nature Genet.
21: 318-322,

[107590]1999.

[107591]18921.

[107592]Wigle, J. T.; Oliver, G.: Prox1 function is required for the
development

[107593]of the murine lymphatic system. Cell 98: 769-778, 1999.

[107594]18922.

[107595]Zinovieva, R. D.; Duncan, M. K.; Johnson, T. R.; Torres, R.;
Polymeropoulos,

[107596]M. H.; Tomarev, S. I.: Structure and chromosomal localization of
tion of

[107597]the human homeobox gene Prox 1. Genomics 35:
517-522, 1996.

[107598]18923.

[107599]Ladenburger, E. M.; Fackelmayer, F. O.; Hameister, H.;
Knippers,

- [107600]R.: MCM4 and PRKDC, human genes encoding proteins
MCM4 and DNA-PKcs,
[107601]are close neighbours located on chromosome 8q12-q13.
Cytogenet. Cell
[107602]Genet. 77: 268-270, 1997.
[107603]18924.
- [107604]Ma, Y.; Pannicke, U.; Schwarz, K.; Lieber, M. R.: Hairpin
opening
[107605]and overhang processing by an Artemis/DNA-dependent
protein kinase
[107606]complex in nonhomologous end joining and V(D)J recom-
bination. Cell 108:
[107607]781-794, 2002.
[107608]18925.
- [107609]Satoh, T.; Tsuruga, H.; Yabuta, N.; Ishidate, M., Jr.; No-
jima,
[107610]H.: Assignment of the human CDC21 (MCM4) gene to
chromosome 8q11.2. Genomics 46:
[107611]525-526, 1997.
[107612]18926.
- [107613]Barresi, R.; Di Blasi, C.; Negri, T.; Brugnoli, R.; Vitali, A.;

- [107614]Felisari, G.; Salandi, A.; Daniel, S.; Cornelio, F.; Morandi, L.;
- [107615]Mora, M.: Disruption of heart sarcoglycan complex and severe cardiomyopathy
- [107616]caused by beta sarcoglycan mutations. J. Med. Genet. 37: 102–107,
- [107617]2000.
- [107618]18927.
- [107619]Bonnemann, C. G.; Modi, R.; Noguchi, S.; Mizuno, Y.; Yoshida, M.;
- [107620]Gussoni, E.; McNally, E. M.; Duggan, D. J.; Angelini, C.; Hoffman,
- [107621]E. P.; Ozawa, E.; Kunkel, L. M.: Beta-sarcoglycan (A3b) mutations
- [107622]cause autosomal recessive muscular dystrophy with loss of the sarcoglycan
- [107623]complex. Nature Genet. 11: 266–273, 1995. Note: Erratum: Nature
- [107624]Genet. 12: 110 only, 1996.
- [107625]18928.
- [107626]Bonnemann, C. G.; Passos-Bueno, M. R.; McNally, E. M.; Vainzof,

- [107627]M.; Moreira, E. S.; Marie, S. K.; Pavanello, R. C. M.;
Noguchi, S.;
- [107628]Ozawa, E.; Zatz, M.; Kunkel, L. M.: Genomic screening for
beta-sarcoglycan
- [107629]gene mutations: missense mutations may cause severe
limb-girdle muscular
- [107630]dystrophy type 2E (LGMD 2E). Hum. Molec. Genet. 5:
1953-1961, 1996.
- [107631]18929.
- [107632]Bonnemann, C. G.; Wong, J.; Ben Hamida, C.; Ben Hamida,
M.; Hentati,
- [107633]F.; Kunkel, L. M.: LGMD 2E in Tunisia is caused by a ho-
mozygous missense
- [107634]mutation in beta-sarcoglycan exon 3. Neuromusc. Dis. 8:
193-197,
- [107635]1998.
- [107636]18930.
- [107637]Durbeej, M.; Cohn, R. D.; Hrstka, R. F.; Moore, S. A.; Alla-
mand,
- [107638]V.; Davidson, B. L.; Williamson, R. A.; Campbell, K. P.: Dis-
ruption
- [107639]of the beta-sarcoglycan gene reveals pathogenetic com-

plexity of limb-girdle

[107640]muscular dystrophy type 2E. Molec. Cell 5: 141–151,
2000.

[107641]18931.

[107642]Joenje, H.; Lo Ten Foe, J. R.; Oostra, A. B.; van Berkel, C. G.

[107643]M.; Rooimans, M. A.; Schroeder-Kurth, T.; Wegner, R.-D.;
Gille, J.

[107644]J. P.; Buchwald, M.; Arwert, F.: Classification of Fanconi
anemia

[107645]patients by complementation analysis: evidence for a fifth
genetic

[107646]subtype. Blood 86: 2156–2160, 1995.

[107647]18932.

[107648]Fielder, P. J.; Guevara-Aguirre, J.; Rosenbloom, A. L.;
Carlsson,

[107649]L.; Hintz, R. L.; Rosenfeld, R. G.: Expression of serum in-
sulin-like

[107650]growth factors, insulin-like growth factor-binding pro-
teins, and the

[107651]growth hormone-binding protein in heterozygote relatives
of Ecuadorian

[107652]growth hormone receptor deficient patients. J. Clin. En-

docr. Metab. 74:

[107653]743–750, 1992.

[107654]18933.

[107655]Fisker, S.; Kristensen, K.; Rosenfalck, A. M.; Pedersen, S.
B.;

[107656]Ebdrup, L.; Richelsen, B.; Hilsted, J.; Christiansen, J. S.;
Jorgensen,

[107657]J. O. L.: Gene expression of a truncated and the full-
length growth

[107658]hormone (GH) receptor in subcutaneous fat and skeletal
muscle in GH-deficient

[107659]adults: impact of GH treatment. J. Clin. Endocr. Metab. 86:
792–796,

[107660]2001.

[107661]18934.

[107662]Gastier, J. M.; Berg, M. A.; Vesterhus, P.; Reiter, E. O.;
Francke,

[107663]U.: Diverse deletions in the growth hormone receptor
gene cause growth

[107664]hormone insensitivity syndrome. Hum. Mutat. 16:
323–333, 2000.

[107665]18935.

- [107666] Godowski, P. J.; Leung, D. W.; Meacham, L. R.; Galgani, J. P.;
- [107667] Hellmiss, R.; Keret, R.; Rotwein, P. S.; Parks, J. S.; Laron, Z.;
- [107668] Wood, W. I.: Characterization of the human growth hormone receptor gene and demonstration of a partial gene deletion in two patients
- [107669] with Laron-type dwarfism. Proc. Nat. Acad. Sci. 86: 8083–8087, 1989.
- [107671] 18936.
- [107672] Iida, K.; Takahashi, Y.; Kaji, H.; Nose, O.; Okimura, Y.; Abe,
- [107673] H.; Chihara, K.: Growth hormone (GH) insensitivity syndrome with
- [107674] high serum GH-binding protein levels caused by a heterozygous splice
- [107675] site mutation of the GH receptor gene producing a lack of intracellular
- [107676] domain. J. Clin. Endocr. Metab. 83: 531–537, 1998.
- [107677] 18937.
- [107678] Iida, K.; Takahashi, Y.; Kaji, H.; Takahashi, M. O.; Okimura,

[107679]Y.; Nose, O.; Abe, H.; Chihara, K.: Functional characterization of
 truncated growth hormone (GH) receptor-(1-277) causing
 partial GH
 insensitivity syndrome with high GH-binding protein. J.
 Clin. Endocr.
 Metab. 84: 1011-1016, 1999.
 18938.
 Jorge, A. A.; Souza, S. C.; Arnhold, I. J.; Mendonca, B. B.:
 Poor reproducibility of IGF-I and IGF binding protein-3
 generation
 test in children with short stature and normal coding re-
 gion of the
 GH receptor gene. J. Clin. Endocr. Metab. 87: 469-472,
 2002.
 18939.
 Kaji, H.; Nose, O.; Tajiri, H.; Takahashi, Y.; Iida, K.; Taka-
 hashi,
 T.; Okimura, Y.; Abe, H.; Chihara, K.: Novel compound
 heterozygous
 mutations of growth hormone (GH) receptor gene in a pa-
 tient with GH

[107692]insensitivity syndrome. J. Clin. Endocr. Metab. 82:
3705–3709, 1997.

[107693]18940.

[107694]Kranzler, J. H.; Rosenbloom, A. L.; Martinez, V.; Guevara-
Aguirre,

[107695]J.: Normal intelligence with severe insulin-like growth fac-
tor I

[107696]deficiency due to growth hormone receptor deficiency: a
controlled

[107697]study in a genetically homogeneous population. J. Clin.
Endocr.

[107698]Metab. 83: 1953–1958, 1998.

[107699]18941.

[107700]Laron, Z.; Klinger, B.; Erster, B.; Silbergeld, A.: Serum GH

[107701]binding protein activities identifies the heterozygous car-
riers for

[107702]Laron type dwarfism. Acta Endocr. 121: 603–608, 1989.

[107703]18942.

[107704]Leung, D. W.; Spencer, S. A.; Cachianes, G.; Hammonds, R.
G.;

[107705]Collins, C.; Henzel, W. J.; Barnard, R.; Waters, M. J.; Wood,

W. I.

[107706] Growth hormone receptor and serum binding protein:
purification,

[107707] cloning and expression. Nature 330: 537–543, 1987.

[107708] 18943.

[107709] Marzullo, P.; Buckway, C.; Pratt, K. L.; Colao, A.; Guevara–
Aguirre,

[107710] J.; Rosenfeld, R. G.: Leptin concentrations in GH defi–
ciency: the

[107711] effect of GH insensitivity. J. Clin. Endocr. Metab. 87:
540–545,

[107712] 2002.

[107713] 18944.

[107714] Menon, R. K.; Cheng, H.; Singh, M.: Identification and
characterization

[107715] of single strand DNA-binding protein that represses
growth hormone

[107716] receptor gene expression. Molec. Endocr. 11: 1291–1304,
1997.

[107717] 18945.

[107718] Metherell, L. A.; Akker, S. A.; Munroe, P. B.; Rose, S. J.;

Caulfield,

[107719]M.; Savage, M. O.; Chew, S. L.; Clark, A. J. L.: Pseudoexon
activation

[107720]as a novel mechanism for disease resulting in atypical
growth-hormone

[107721]insensitivity. Am. J. Hum. Genet. 69: 641-646, 2001.

[107722]18946.

[107723]Rosenbloom, A. L.; Guevara-Aguirre, J.; Berg, M. A.;
Francke,

[107724]U.: Stature in Ecuadorians heterozygous for growth hor-
mone receptor

[107725]gene E180 splice mutation does not differ from that of
homozygous

[107726]normal relatives. J. Clin. Endocr. Metab. 83: 2373-2375,
1998.

[107727]18947.

[107728]Sanchez, J. E.; Perera, E.; Baumbach, L.; Cleveland, W. W.:
Growth

[107729]hormone receptor mutations in children with idiopathic
short stature. J.

[107730]Clin. Endocr. Metab. 83: 4079-4083, 1998.

[107731]18948.

- [107732]Shen, X. Y.; Holt, R. I. G.; Miell, J. P.; Justice, S.; Portmann,
[107733]B.; Postel-Vinay, M.-C.; Ross, R. J. M.: Cirrhotic liver ex-
presses
[107734]low levels of the full-length and truncated growth hor-
mone receptors. J.
[107735]Clin. Endocr. Metab. 83: 2532-2538, 1998.
[107736]18949.
- [107737]Spencer, S. A.; Hammonds, R. G.; Henzel, W. J.; Rodriguez,
H.;
[107738]Waters, M. J.; Wood, W. I.: Rabbit liver growth hormone re-
ceptor
[107739]and serum binding protein: purification, characterization,
and sequence. J.
[107740]Biol. Chem. 263: 7862-7867, 1988.
[107741]18950.
- [107742]Walker, J. L.; Crock, P. A.; Behncken, S. N.; Rowlinson, S.
W.;
[107743]Nicholson, L. M.; Boulton, T. J. C.; Waters, M. J.: A novel
mutation
[107744]affecting the interdomain link region of the growth hor-
mone receptor

[107745]in a Vietnamese girl, and response to long-term treatment with recombinant

[107746]human insulin-like growth factor-I and luteinizing hormone-releasing

[107747]hormone analogue. J. Clin. Endocr. Metab. 83: 2554-2561, 1998.

[107748]18951.

[107749]Denier, C.; Ducros, A.; Durr, A.; Eymard, B.; Chassande, B.; Tournier-Lasserre,

[107750]E.: Missense CACNA1A mutation causing episodic ataxia type 2. Arch.

[107751]Neurol. 58: 292-295, 2001.

[107752]18952.

[107753]Byk, T.; Dobransky, T.; Cifuentes-Diaz, C.; Sobel, A.: Identification

[107754]and molecular characterization of Unc-33-like phosphoprotein (Ulip),

[107755]a putative mammalian homolog of the axonal guidance-associated unc-33

[107756]gene product. J. Neurosci. 16: 688-701, 1996.

[107757]18953.

- [107758]Choe, H.; Farzan, M.; Sun, Y.; Sullivan, N.; Rollins, B.;
Ponath,
[107759]P. D.; Wu, L.; Mackay, C. R.; LaRosa, G.; Newman, W.; Ger-
ard, N.;
- [107760]Gerard, C.; Sodroski, J.: The beta-chemokine receptors
CCR3 and CCR5
[107761]facilitate infection by primary HIV-1 isolates. Cell 85:
1135-1148,
[107762]1996.
[107763]18954.
- [107764]Diriong, S.; Lory, P.; Williams, M. E.; Ellis, S. B.; Harpold,
[107765]M. M.; Taviaux, S.: Chromosomal localization of the hu-
man genes for
[107766]alpha-1A, alpha-1B, and alpha-1E voltage-dependent
Ca(2+) channel
[107767]subunits. Genomics 30: 605-609, 1995.
[107768]18955.
- [107769]Ducros, A.; Denier, C.; Joutel, A.; Cecillon, M.; Lescoat, C.;
- [107770]Vahedi, K.; Darcel, F.; Vicaut, E.; Bousser, M.-G.; Tournier-
Lasserre,
[107771]E.: The clinical spectrum of familial hemiplegic migraine

associated

[107772]with mutations in a neuronal calcium channel. New Eng. J.
Med. 345:

[107773]17–24, 2001.

[107774]18956.

[107775]Dunlap, K.; Luebke, J. I.; Turner, T. J.: Exocytotic Ca(2+)
channels

[107776]in mammalian central neurons. Trends Neurosci. 18:
89–98, 1995.

[107777]18957.

[107778]Fletcher, C. F.; Lutz, C. M.; O'Sullivan, T. N.; Shaughnessy,
J.

[107779]D, Jr.; Hawkes, R.; Frankel, W. N.; Copeland, N. G.; Jenkins,
N. A.

[107780]: Absence epilepsy in tottering mutant mice is associated
with calcium

[107781]channel defects. Cell 87: 607–617, 1996.

[107782]18958.

[107783]Friend, K. L.; Crimmins, D.; Phan, T. G.; Sue, C. M.; Colley,
A.;

[107784]Fung, V. S. C.; Morris, J. G. L.; Sutherland, G. R.; Richards,

R.

[107785]J.: Detection of a novel missense mutation and second recurrent mutation

[107786]in the CACNA1A gene in individuals with EA-2 and FHM.

Hum. Genet. 105:

[107787]261–265, 1999.

[107788]18959.

[107789]Hess, E. J.: Migraines in mice? Cell 87: 1149–1151, 1996.

[107790]18960.

[107791]Hoffman, E. P.; Gardner, K.: Ion channels--molecular diving

[107792]rods hit their clinical mark. New Eng. J. Med. (Editorial)

336:

[107793]1599–1600, 1997.

[107794]18961.

[107795]Jodice, C.; Mantuano, E.; Veneziano, L.; Trettel, F.; Sabbadini,

[107796]G.; Calandriello, L.; Francia, A.; Spadaro, M.; Pierelli, F.;

Salvi,

[107797]F.; Ophoff, R. A.; Frants, R. R.; Frontali, M.: Episodic ataxia type

[107798]2 (EA2) and spinocerebellar ataxia type 6 (SCA6) due to
CAG repeat

[107799]expansion in the CACNA1A gene on chromosome 19p.
Hum. Molec. Genet. 6:

[107800]1973–1978, 1997.

[107801]18962.

[107802]Kim, J.-S.; Yue, Q.; Jen, J. C.; Nelson, S. F.; Baloh, R. W.:

[107803]Familial migraine with vertigo: no mutations found in
CACNA1A. Am.

[107804]J. Med. Genet. 79: 148–151, 1998.

[107805]18963.

[107806]Kraus, R. L.; Sinnegger, M. J.; Glossmann, H.; Hering, S.;
Striessnig,

[107807]J.: Familial hemiplegic migraine mutations change al-
pha(1A)Ca(2+)

[107808]channel kinetics. J. Biol. Chem. 273: 5586–5590, 1998.

[107809]18964.

[107810]Guida, S.; Trettel, F.; Pagnutti, S.; Mantuano, E.; Tottene,
A.;

[107811]Veneziano, L.; Fellin, T.; Spadaro, M.; Stauderman, K. A.;
Williams,

[107812]M. E.; Volsen, S.; Ophoff, R. A.; Frants, R. R.; Jodice, C.;
Frontali,
[107813]M.; Pietrobon, D.: Complete loss of P/Q calcium channel
activity
[107814]caused by a CACNA1A missense mutation carried by pa-
tients with episodic
[107815]ataxia type 2. Am. J. Hum. Genet. 68: 759–764, 2001.
[107816]18965.
[107817]Ishikawa, K.; Tanaka, H.; Saito, M.; Ohkoshi, N.; Fujita, T.;
[107818]Yoshizawa, K.; Ikeuchi, T.; Watanabe, M.; Hayashi, A.;
Takiyama, Y.;
[107819]Nishizawa, M.; Nakano, I.; Matsubayashi, K.; Miwa, M.;
Shoji, S.;
[107820]Kanazawa, I.; Tsuji, S.; Mizusawa, H.: Japanese families
with autosomal
[107821]dominant pure cerebellar ataxia map to chromosome
19p13.1–p13.2 and
[107822]are strongly associated with mild CAG expansions in the
spinocerebellar
[107823]ataxia type 6 gene in chromosome 19p13.1. Am J. Hum.
Genet. 61:
[107824]336–346, 1997.

[107825]18966.

[107826]Kors, E. E.; Terwindt, G. M.; Vermeulen, F. L. M. G.; Fitzsimons,

[107827]R. B.; Jardine, P. E.; Heywood, P.; Love, S.; van den Maagdenberg,

[107828]A. M. J. M.; Haan, J.; Frants, R. R.; Ferrari, M. D.: Delayed cerebral

[107829]edema and fatal coma after minor head trauma: role of the CACNA1A

[107830]calcium channel subunit gene and relationship with familial hemiplegic

[107831]migraine. Ann. Neurol. 49: 753–760, 2001.

[107832]18967.

[107833]Kraus, R. L.; Sinnegger, M. J.; Koschak, A.; Glossmann, H.; Stenirri,

[107834]S.; Carrera, P.; Striessnig, J.: Three new familial hemiplegic migraine

[107835]mutants affect P/Q-type Ca^{2+} channel kinetics. J. Biol. Chem. 275:

[107836]9239–9243, 2000.

[107837]18968.

- [107838]Lehmann-Horn, F.; Jurkat-Rott, K.: Voltage-gated ion channels
[107839]and hereditary disease. *Physiol. Rev.* 79: 1317–1372, 1999.
[107840]18969.
- [107841]Matsuyama, Z.; Kawakami, H.; Maruyama, H.; Izumi, Y.; Komure,
[107842]O.; Udaaka, F.; Kameyama, M.; Nishio, T.; Kuroda, Y.; Nishimura, M.;
[107843]Nakamura, S.: Molecular features of the CAG repeats of spinocerebellar
[107844]ataxia 6 (SCA6). *Hum. Molec. Genet.* 6: 1283–1287, 1997.
[107845]18970.
- [107846]Riess, O.; Schols, L.; Bottger, H.; Nolte, D.; Viera-Saecker, A. M. M.; Schimming, C.; Kreuz, F.; Macek, M., Jr.; Krebsova, A.;
[107848]Macek, M., Sr.; Klockgether, T.; Zuhlke, C.; Laccone, F. A.: SCA6
[107849]is caused by moderate CAG expansion in the α 1A-voltage-dependent
[107850]calcium channel gene. *Hum. Molec. Genet.* 6: 1289–1293, 1997.

[107851]18971.

[107852]Sasaki, H.; Kojima, H.; Yabe, I.; Tashiro, K.; Hamada, T.;
Sawa,

[107853]H.; Hiraga, H.; Nagashima, K.: Neuropathological and
molecular studies

[107854]of spinocerebellar ataxia type 6 (SCA6). Acta. Neuropath.
95: 199–204,

[107855]1998.

[107856]18972.

[107857]Scoggan, K. A.; Chandra, T.; Nelson, R.; Hahn, A. F.; Bul-
man,

[107858]D. E.: Identification of two novel mutations in the
CACNA1A gene

[107859]responsible for episodic ataxia type 2. (Letter) J. Med.
Genet. 38:

[107860]249–253, 2001.

[107861]18973.

[107862]Thibault, O.; Landfield, P. W.: Increase in single L-type
calcium

[107863]channels in hippocampal neurons during aging. Science
272: 1017–1019,

[107864]1996.

[107865]18974.

[107866]Toru, S.; Murakoshi, T.; Ishikawa, K.; Saegusa, H.; Fuji-
gasaki,

[107867]H.; Uchihara, T.; Nagayama, S.; Osanai, M.; Mizusawa, H.;
Tanabe,

[107868]T.: Spinocerebellar ataxia type 6 mutation alters P-type
calcium

[107869]channel function. J. Biol. Chem. 275: 10893–10898, 2000.

[107870]18975.

[107871]Trettel, F.; Mantuano, E.; Calabresi, V.; Veneziano, L.;
Olsen,

[107872]A. S.; Georgescu, A.; Gordon, L.; Sabbadini, G.; Frontali,
M.; Jodice,

[107873]C.: A fine physical map of the CACNA1A gene region on
19p13.1–p13.2

[107874]chromosome. Gene 241: 45–50, 2000.

[107875]18976.

[107876]Vahedi, K.; Denier, C.; Ducros, A.; Bousson, V.; Levy, C.;
Chabriat,

[107877]H.; Haguenau, M.; Tournier-Lasserre, E.; Bousser, M. G.:

CACNA1A

[107878]gene de novo mutation causing hemiplegic migraine, coma, and cerebellar

[107879]atrophy. Neurology 55: 1040–1042, 2000.

[107880]18977.

[107881]Ben-Porath, I.; Benvenisty, N.: Characterization of a tumor-associated

[107882]gene, a member of a novel family of genes encoding membrane glycoproteins. Gene 183:

[107883]69–75, 1996.

[107884]18978.

[107885]Bosch, E. P.; Murphy, M. J.; Cancilla, P. A.: Peripheral neurofibromatosis

[107886]and peroneal muscular atrophy. Neurology 31: 1408–1414, 1981.

[107887]18979.

[107888]Bridges, C. B.: The Bar 'gene' a duplication. Science 83: 210–211,

[107889]1936.

[107890]18980.

[107891]Gabreels-Festen, A. A. W. M.; Joosten, E. M. G.; Gabreels,

F. J.

[107892]M.; Jennekens, F. G. I.; Janssen-van Kempen, T. W.: Early morphological

[107893]features in dominantly inherited demyelinating motor and sensory neuropathy

[107894](HMSN type I). J. Neurol. Sci. 107: 145–154, 1992.

[107895]18981.

[107896]Green, M. C.: Genetic variants and strains of the laboratory mouse.

[107897]New York: Oxford Univ. Press (pub.) 1989.

[107898]18982.

[107899]Hartl, D. L.: The most unkindest cut of all. Nature Genet. 12:

[107900]227–229, 1996.

[107901]18983.

[107902]Hoogendijk, J. E.; Hensels, G. W.; Zorn, I.; Valentijn, L.; Janssen,

[107903]E. A. M.; de Visser, M.; Barker, D. F.; Ongerboer de Visser, B. W.;

[107904]Baas, F.; Bolhuis, P. A.: The duplication in Charcot-Marie-Tooth

[107905]disease type 1a spans at least 1100 kb on chromosome
17p11.2. Hum.

[107906]Genet. 88: 215–218, 1991.

[107907]18984.

[107908]Hoogendijk, J. E.; Janssen, E. A. M.; Gabreels–Festen, A. A.
W.

[107909]M.; Hensels, G. W.; Joosten, E. M. G.; Gabreels, F. J. M.;
Zorn, I.;

[107910]Valentijn, L. J.; Baas, F.; Ongerboer de Visser, B. W.; de
Visser,

[107911]M.; Bolhuis, P. A.: Allelic heterogeneity in hereditary motor
and

[107912]sensory neuropathy type 1a (Charcot–Marie–Tooth disease
type 1a). Neurology 43:

[107913]1010–1015, 1993.

[107914]18985.

[107915]Ikegami, T.; Ikeda, H.; Aoyama, M.; Matsuki, T.; Imota, T.;
Fukuuchi,

[107916]Y.; Amano, T.; Toyoshima, I.; Ishihara, Y.; Endoh, H.;
Hayasaka, K.

[107917]: Novel mutations of the peripheral myelin protein 22
gene in two

[107918]pedigrees with Dejerine–Sottas disease. Hum. Genet. 102:
294–298,
[107919]1998.
[107920]18986.
[107921]Ikegami, T.; Ikeda, H.; Chance, P. F.; Kiyosawa, H.; Ya-
mamoto,
[107922]M.; Sobue, G.; Ohnishi, A.; Tachi, N.; Hayasaka, K.: Facili-
tated
[107923]diagnosis of CMT1A duplication in chromosome
17p11.2–12: analysis
[107924]with a CMT1A–REP repeat probe and photostimulated lu-
minescence imaging. Hum.
[107925]Mutat. 9: 563–566, 1997.
[107926]18987.
[107927]Ionasescu, V. V.; Searby, C.; Greenberg, S. A.: Dejerine–Sot-
tas
[107928]disease with sensorineural hearing loss, nystagmus, and
peripheral
[107929]facial nerve weakness: de novo dominant point mutation
of the PMP22
[107930]gene. J. Med. Genet. 33: 1048–1049, 1996.

[107931]18988.

[107932]Isaacs, A. M.; Davies, K. E.; Hunter, A. J.; Nolan, P. M.; Vi-
zor,

[107933]L.; Peters, J.; Gale, D. G.; Kelsell, D. P.; Latham, I. D.;
Chase,

[107934]J. M.; Fisher, E. M. C.; Bouzyk, M. M.; and 11 others: Iden-
tification

[107935]of two new Pmp22 mouse mutants using large-scale mu-
tagenesis and a

[107936]novel rapid mapping strategy. Hum. Molec. Genet. 9:
1865-1871, 2000.

[107937]18989.

[107938]King, P. H.; Waldrop, R.; Lupski, J. R.; Shaffer, L. G.: Char-
cot-Marie-Tooth

[107939]phenotype produced by a duplicated PMP22 gene as part
of a 17p trisomy-translocation

[107940]to the X chromosome. Clin. Genet. 54: 413-416, 1998.

[107941]18990.

[107942]Kousseff, B. G.; Hadro, T. A.; Treiber, D. L.; Wollner, T.;
Morris,

[107943]C.: Charcot-Marie-Tooth disease with sensorineural hear-
ing loss--an

[107944]autosomal dominant trait. Birth Defects Orig. Art. Ser. 18:
223–228,

[107945]1982.

[107946]18991.

[107947]Kovach, M. J.; Lin, J.–P.; Boyadjiev, S.; Campbell, K.;
Mazzeo,

[107948]L.; Herman, K.; Rimer, L. A.; Frank, W.; Llewellyn, B.; Jabs,
E. W.;

[107949]Gelber, D.; Kimonis, V. E.: A unique point mutation in the
PMP22

[107950]gene is associated with Charcot–Marie–Tooth disease and
deafness. Am.

[107951]J. Hum. Genet. 64: 1580–1593, 1999.

[107952]18992.

[107953]Liehr, T.; Rautenstrauss, B.; Grehl, H.; Bathke, K. D.; Ekici,

[107954]A.; Rauch, A.; Rott, H.–D.: Mosaicism for the Charcot–
Marie–Tooth

[107955]disease type 1A duplication suggests somatic reversion.
Hum. Genet. 98:

[107956]22–28, 1996.

[107957]18993.

[107958]Gaetano, C; Matsuo, T.; Thiele, C. J.: Identification and
characterization
[107959]of a retinoic acid-regulated human homologue of the
unc-33-like phosphoprotein
[107960]gene (hUlip) from neuroblastoma cells. J. Biol. Chem. 272:
12195-12201,
[107961]1997.
[107962]18994.
[107963]Matsuo, T.; Stauffer, J. K.; Walker, R. L.; Meltzer, P.; Thiele,
[107964]C. J.: Structure and promoter analysis of the human unc-
33-like phosphoprotein
[107965]gene: E-box required for maximal expression in neurob-
lastoma and myoblasts. J.
[107966]Biol. Chem. 275: 16560-16568, 2000.
[107967]18995.
[107968]Behm, F. G.; Smith, F. O.; Raimondi, S. C.; Pui, C.-H.;
Bernstein,
[107969]J. D.: Human homologue of the rat chondroitin sulfate
proteoglycan,
[107970]NG2, detected by monoclonal antibody 7.1, identifies
childhood acute

- [107971] lymphoblastic leukemias with t(4;11)(q21;q23) or t(11;19)(q23;p13) and MLL gene rearrangements. Blood 87: 1134–1139, 1996.
- [107972] 18996.
- [107973] Pluschke, G.; Vanek, M.; Evans, A.; Dittmar, T.; Schmid, P.; Itin, P.; Filardo, E. J.; Reisfeld, R. A.: Molecular cloning of a human melanoma-associated chondroitin sulfate proteoglycan. Proc. Nat. Acad. Sci. 93: 9710–9715, 1996.
- [107974] 18997.
- [107975] Rettig, W. J.; Dracopoli, N. C.; Goetzger, T. A.; Spengler, B. A.; Biedler, J. L.; Oettgen, H. F.; Old, L. J.: Somatic cell genetic analysis of human cell surface antigens: chromosomal assignments and regulation of expression in rodent–human hybrid cells. Proc. Nat. Acad. Sci. 81: 6437–6441, 1984.

[107984]18998.

[107985]Rettig, W. J.; Real, F. X.; Spengler, B. A.; Biedler, J. L.; Old,

[107986]L. J.: Human melanoma proteoglycan: expression in hybrids controlled

[107987]by intrinsic and extrinsic signals. Science 231: 1281–1284, 1986.

[107988]18999.

[107989]Smith, F. O.; Rauch, C.; Williams, D. E.; March, C. J.; Arthur,

[107990]D.; Hilden, J.; Lampkin, B. C.; Buckley, J. D.; Buckley, C. V.; Woods,

[107991]W. G.; Dinndorf, P. A.; Sorensen, P.; Kersey, J.; Hammond, D.; Bernstein,

[107992]I. D.: The human homologue of rat NG2, a chondroitin sulfate proteoglycan,

[107993]is not expressed on the cell surface of normal hematopoietic cells

[107994]but is expressed by acute myeloid leukemia blasts from poor-prognosis

[107995]patients with abnormalities of chromosome band 11q23. Blood 87:

[107996]1123–1133, 1996.

[107997]19000.

[107998]Clark, J.; Moore, L.; Krasinskas, A.; Way, J.; Battey, J.;
Tamkun,

[107999]J.; Kahn, R. A.: Selective amplification of additional mem-
bers of

[108000]the ADP-ribosylation factor (ARF) family: cloning of addi-
tional human

[108001]and Drosophila ARF-like genes. Proc. Nat. Acad. Sci. 90:
8952-8956,

[108002]1993.

[108003]19001.

[108004]Gipp, J. J.; Bailey, H. H.; Mulcahy, R. T.: Cloning and se-
quencing

[108005]of the cDNA for the light subunit of human liver gamma-
glutamylcysteine

[108006]synthetase and relative mRNA levels for heavy and light
subunits in

[108007]human normal tissues. Biochem. Biophys. Res. Commun.
206: 584-589,

[108008]1995.

[108009]19002.

- [108010]Huang; C.-S.; Anderson; M. E.; Meister, A.: Amino acid sequence
- [108011]and function of the light subunit of rat kidney gamma-glutamylcysteine
- [108012]synthetase. J. Biol. Chem. 268: 20578-20583, 1993.
- [108013]19003.
- [108014]Sierra-Rivera, E.; Dasouki, M.; Summar, M. L.; Krishnamani, M.
- [108015]R. S.; Meredith, M.; Rao, P. N.; Phillips, J. A., III; Freeman, M.
- [108016]L.: Assignment of the human gene (GLCLR) that encodes the regulatory
- [108017]subunit of gamma-glutamylcysteine synthetase to chromosome 1p21. Cytogenet.
- [108018]Cell Genet. 72: 252-254, 1996.
- [108019]19004.
- [108020]Tsuchiya, K.; Mulcahy, R. T.; Reid, L. L.; Disteché, C. M.; Kavanagh,
- [108021]T. J.: Mapping of the glutamate-cysteine ligase catalytic subunit
- [108022]gene (GLCLC) to human chromosome 6p12 and mouse chromosome 9D-E and

[108023]of the regulatory subunit gene (GLCLR) to human chromosome 1p21–p22

[108024]and mouse chromosome 3H1–3. Genomics 30: 630–632, 1995.

[108025]19005.

[108026]Rozet, J.–M.; Gerber, S.; Perrault, I.; Calvas, P.; Souied, E.;

[108027]Chatelin, S.; Viegas–Pequignot, E.; Molina–Gomez, D.; Munnich, A.;

[108028]Kaplan, J.: Structure and refinement of the physical mapping of the

[108029]gamma–glutamylcysteine ligase regulatory subunit (GLCLR) gene to chromosome

[108030]1pp22.1 within the critically deleted region of human malignant mesothelioma. Cytogenet.

[108031]Cell Genet. 82: 91–94, 1998.

[108032]19006.

[108033]Ellis, J. A.; Luzio, J. P.: Identification and characterization

[108034]of a novel protein (p137) which transcytoses bidirectionally in Caco-2

[108035]cells. J. Biol. Chem. 270: 20717–20723, 1995.

[108036]19007.

[108037]Gessler, M.; Klamt, B.; Tsaoussidou, S.; Ellis, J. A.; Luzio, J.

[108038]P.: The gene encoding the GPI-anchored membrane protein p137(GPI)

[108039](M11S1) maps to human chromosome 11p13 and is highly conserved in

[108040]the mouse. Genomics 32: 169–170, 1996.

[108041]19008.

[108042]Rouault, J.-P; Falette, N.; Guehenneux, F.; Guillot, C.; Rimokh,

[108043]R.; Wang, Q.; Berthet, C.; Moyret-Lalle, C.; Savatier, P.; Pain, B.;

[108044] Shaw, P.; Berger, R.; Samarut, J.; Magaud, J.-P.; Ozturk, M.; Samarut, C.; Puisieux, A.: Identification of BTG2, an antiproliferative p53-dependent component of the DNA damage cellular response pathway. Nature Genet. 14: 482-486, 1996.

[108047] 19009.

[108048] Ring, H. Z.; Vameghi-Meyers, V.; Wang, W.; Crabtree, G. R.; Francke, U.: Five SWI/SNF-related, matrix-associated, actin-dependent regulator of chromatin (SMARC) genes are dispersed in the human genome. Genomics 51: 140-143, 1998.

[108052] 19010.

[108053] Wang, W.; Xue, Y.; Zhou, S.; Kuo, A.; Cairns, B. R.; Crabtree, G. R.: Diversity and specialization of mammalian SWI/SNF complexes. Genes Dev. 10: 2117-2130, 1996.

[108057]19011.

[108058]Kim, B.-T.; Kitagawa, H.; Tamura, J.; Saito, T.; Kusche-Gullberg,

[108059]M.; Lindahl, U.; Sugahara, K.: Human tumor suppressor
EXT gene family

[108060]members EXTL1 and EXTL3 encode alpha-
1,4-N-acetylglucosaminyltransferases

[108061]that likely are involved in heparan sulfate/heparin biosynthesis. Proc.

[108062]Nat. Acad. Sci. 98: 7176-7181, 2001.

[108063]19012.

[108064]Wise, C. A.; Clines, G. A.; Massa, H.; Trask, B. J.; Lovett, M.

[108065]: Identification and localization of the gene for EXTL, a
third member

[108066]of the multiple exostoses gene family. Genome Res. 7:
10-16, 1997.

[108067]19013.

[108068]Moskow, J. J.; Bullrich, F.; Huebner, K.; Daar, I. O.; Buch-
berg,

[108069]A. M.: Meis1, a PBX1-related homeobox gene involved in
myeloid leukemia

[108070]in BXH-2 mice. *Molec. Cell. Biol.* 15: 5434–5443, 1995.

[108071]19014.

[108072]Steelman, S.; Moskow, J. J.; Muzynski, K.; North, C.; Druck, T.;

[108073]Montgomery, J. C.; Huebner, K.; Daar, I. O.; Buchberg, A. M.: Identification

[108074]of a conserved family of Meis1-related homeobox genes. *Genome Res.* 7:

[108075]142–156, 1997.

[108076]19015.

[108077]Thorsteinsdottir, U.; Kroon, E.; Jerome, L.; Blasi, F.; Sauvageau,

[108078]G.: Defining roles for HOX and MEIS1 genes in induction of acute

[108079]myeloid leukemia. *Molec. Cell. Biol.* 21: 224–234, 2001.

[108080]19016.

[108081]Capdevila, J.; Tsukui, T.; Esteban, C. R.; Zappavigna, V.; Belmonte,

[108082]J. C. I.: Control of vertebrate limb outgrowth by the proximal factor

[108083]Meis2 and distal antagonism of BMPs by Gremlin. *Molec.*

- Cell 4: 839–849,
[108084]1999.
- [108085]19017.
- [108086]Nakamura, T.; Jenkins, N. A.; Copeland, N. G.: Identification
of a new family of Pbx-related homeobox genes. *Oncogene* 13: 2235–2242,
[108088]1996.
- [108089]19018.
- [108090]Smith, J. E.; Afonja, O.; Yee, H. T.; Inghirami, G.;
Takeshita,
[108091]K.: Chromosomal mapping to 15q14 and expression analysis of the human
[108092]MEIS2 homeobox gene. *Mammalian Genome* 8: 951–952,
1997.
- [108093]19019.
- [108094]Kobayashi, S.; Uemura, H.; Kohda, T.; Nagai, T.; Chinen,
Y.; Naritomi,
[108095]K.; Kinoshita, E.; Ohashi, H.; Imaizumi, K.; Tsukahara, M.;
Sugio,
[108096]Y.; Tonoki, H.; Kishino, T.; Tanaka, T.; Yamada, M.; Tsut-

sumi, O.;

[108097]Niikawa, N.; Kaneko-Ishino, T.; Ishino, F.: No evidence of
PEG1/MEST

[108098]gene mutations in Silver-Russell syndrome patients. Am. J.
Med. Genet. 104:

[108099]225-231, 2001.

[108100]19020.

[108101]Mizuno, K.; Hasegawa, K.; Katagiri, T.; Ogimoto, M.;
Ichikawa,

[108102]T.; Yakura, H.: MPTP-delta, a putative murine homolog of
HPTP-delta,

[108103]is expressed in specialized regions of the brain and in the
B-cell

[108104]lineage. Molec. Cell. Biol. 13: 5513-5523, 1993.

[108105]19021.

[108106]Uetani, N.; Kato, K.; Ogura, H.; Mizuno, K.; Kawano, K.;
Mikoshiba,

[108107]K.; Yakura, H.; Asano, M.; Iwakura, Y.: Impaired learning
with enhanced

[108108]hippocampal long-term potentiation in PTP-
delta-deficient mice. EMBO

[108109]J. 19: 2775-2785, 2000.

[108110]19022.

[108111]Heighway, J.; Betticher, D. C.; Hoban, P. R.; Altermatt, H. J.;

[108112]Cowen, R.: Coamplification in tumors of KRAS2, type 2 in-
ositol 1,4,5

[108113]triphosphate receptor gene, and a novel human gene,
KRAg. Genomics 35:

[108114]207-214, 1996.

[108115]19023.

[108116]Scott, A. F.; Elizaga, A.; Morrell, J.; Bergen, A.; Penno, M. B.

[108117]: Characterization of a gene coamplified with Ki-ras in Y1
murine

[108118]adrenal carcinoma cells that codes for a putative mem-
brane protein. Genomics 20:

[108119]227-230, 1994.

[108120]19024.

[108121]Monte, D.; Baert, J. L.; Defossez, P. A.; de Launoit, Y.; Ste-
helin,

[108122]D.: Molecular cloning and characterization of human ERM,
a new member

[108123]of the Ets family closely related to mouse PEA3 and ER81
transcription

[108124]factors. *Oncogene* 9: 1397–1406, 1994.

[108125]19025.

[108126]Monte, D.; Coutte, L.; Dewitte, F.; Defossez, P.-A.; Le Coniat,

[108127]M.; Stehelin, D.; Berger, R.; de Launoit, Y.: Genomic organization

[108128]of the human ERM (ETV5) gene, a PEA3 group member of ETS transcription

[108129]factors. *Genomics* 35: 236–240, 1996.

[108130]19026.

[108131]Protopopova, M. V.; Vorobieva, N. V.; Protopopov, A. I.; Gizatullin,

[108132]R. Z.; Kashuba, V. I.; Klein, G.; Zabarovsky, E. R.; Graphodatsky,

[108133]A. S.: Assignment of the ERM gene (ETV5) coding for the ets-related

[108134]protein to human chromosome band 3q28 by in situ hybridization. *Cytogenet.*

[108135]*Cell Genet.* 74: 220 only, 1996.

[108136]19027.

[108137]Bosher, J. M.; Williams, T.; Hurst, H. C.: The developmen-

tally

[108138]regulated transcription factor AP-2 is involved in c-erbB-2 overexpression

[108139]in human mammary carcinoma. Proc. Nat. Acad. Sci. 92: 744-747, 1995.

[108140]19028.

[108141]Brown, M. E.; Pollak, M.; Seidman, C. E.; Seidman, J. G.; Chou,

[108142]Y.-H. W.; Riccardi, D.; Herbert, S. C.: Calcium-ion-sensing cell-surface

[108143]receptors. New Eng. J. Med. 333 -234-240, 1995.

[108144]19029.

[108145]Carling, T.; Szabo, E.; Bai, M.; Ridefelt, P.; Westin, G.; Gustavsson,

[108146]P.; Trivedi, S.; Hellman, P.; Brown, E. M.; Dahl, N.; Rastad, J.:

[108147]Familial hypercalcemia and hypercalciuria caused by a novel mutation

[108148]in the cytoplasmic tail of the calcium receptor. J. Clin. Endocr.

[108149]Metab. 85: 2042-2047, 2000.

[108150]19030.

[108151]Cetani, F.; Pinchera, A.; Pardi, E.; Cianferotti, L.; Vignali,
[108152]E.; Picone, A.; Miccoli, P.; Viacava, P.; Marcocci, C.: No evi-
dence
[108153]for mutations in the calcium-sensing receptor gene in
sporadic parathyroid
[108154]adenomas. J. Bone Miner. Res. 14: 878-882, 1999.
[108155]19031.
[108156]D'Souza-Li, L.; Canaff, L.; Janicic, N.; Cole, D. E. C.; Hendy,
[108157]G. N.: An acceptor splice site mutation in the calcium-
sensing receptor
[108158](CASR) gene in familial hypocalciuric hypercalcemia and
neonatal severe
[108159]hyperparathyroidism. Hum. Mutat. 18: 411-421, 2001.
[108160]19032.
[108161]De Luca, F.; Ray, K.; Mancilla, E. E.; Fan, G.-F.; Winer, K. K.;
[108162]Gore, P.; Spiegel, A. M.; Baron, J.: Sporadic hypoparathy-
roidism
[108163]caused by de novo gain-of-function mutations of the
Ca(2+)-sensing
[108164]receptor. J. Clin. Endocr. Metab. 82: 2710-2715, 1997.

[108165]19033.

[108166]Hendy, G. N.; D'Souza-Li, L.; Yang, B.; Canaff, L.; Cole, D. E.

[108167]C.: Mutations of the calcium-sensing receptor (CASR) in familial

[108168]hypocalciuric hypercalcemia, neonatal severe hyperparathyroidism,

[108169]and autosomal dominant hypocalcemia. Hum. Mutat. 16: 281-296, 2000.

[108170]19034.

[108171]Janicic, N.; Pausova, Z.; Cole, D. E. C.; Hendy, G. N.: Insertion

[108172]of an Alu sequence in the Ca(2+)-sensing receptor gene in familial

[108173]hypocalciuric hypercalcemia and neonatal severe hyperparathyroidism. Am.

[108174]J. Hum. Genet. 56: 880-886, 1995.

[108175]19035.

[108176]Jap, T.-S.; Wu, Y.-C.; Jenq, S.-F.; Won, G.-S.: A novel mutation

[108177]in the calcium-sensing receptor gene in a Chinese subject with persistent

[108178]hypercalcemia and hypocalciuria. J. Clin. Endocr. Metab.
86: 13–15,

[108179]2001.

[108180]19036.

[108181]Kobayashi, M.; Tanaka, H.; Tsuzuki, K.; Tsuyuki, M.; Igaki,
H.;

[108182]Ichinose, Y.; Aya, K.; Nishioka, N.; Seino, Y.: Two novel
missense

[108183]mutations in calcium–sensing receptor gene associated
with neonatal

[108184]severe hyperparathyroidism. J. Clin. Endocr. Metab. 82:
2716–2719,

[108185]1997.

[108186]19037.

[108187]Lienhardt, A.; Bai, M.; Lagarde, J.–P.; Rigaud, M.; Zhang,
Z.;

[108188]Jiang, Y.; Kottler, M.–L.; Brown, E. M.; Garabedian, M.: Ac–
tivating

[108189]mutations of the calcium–sensing receptor: management
of hypocalcemia. J.

[108190]Clin. Endocr. Metab. 86: 5313–5323, 2001.

[108191]19038.

[108192]Lienhardt, A.; Garabedian, M.; Bai, M.; Sinding, C.; Zhang, Z.;

[108193]Lagarde, J.-P.; Boulesteix, J.; Rigaud, M.; Brown, E. M.; Kottler,

[108194]M.-L.: A large homozygous or heterozygous in-frame deletion within

[108195]the calcium-sensing receptor's carboxylterminal cytoplasmic tail that

[108196]causes autosomal dominant hypocalcemia. J. Clin. Endocr. Metab. 85:

[108197]1695-1702, 2000.

[108198]19039.

[108199]Lovlie, R.; Eiken, H. G.; Sorheim, J. I.; Boman, H.: The Ca(2+)-sensing

[108200]receptor gene (PCAR1) mutation T151M in isolated autosomal dominant

[108201]hypoparathyroidism. Hum. Genet. 98: 129-133, 1996.

[108202]19040.

[108203]Okazaki, R.; Chikatsu, N.; Nakatsu, M.; Takeuchi, Y.; Ajima, M.;

- [108204]Miki, J.; Fujita, T.; Arai, M.; Totsuka, Y.; Tanaka, K.; Fukumoto,
moto,
- [108205]S.: A novel activating mutation in calcium-sensing receptor gene
tor gene
- [108206]associated with a family of autosomal dominant hypocalcemia. J. Clin.
cemia. J. Clin.
- [108207]Endocr. Metab. 84: 363–366, 1999.
- [108208]19041.
- [108209]Combadiere, C.; Ahuja, S. K.; Murphy, P. M.: Cloning,
chromosomal
- [108210]localization, and RNA expression of a human beta
chemokine receptor-like
- [108211]gene. DNA Cell Biol. 14: 673–680, 1995.
- [108212]19042.
- [108213]Combadiere, C.; Ahuja, S. K.; Van Damme, J.; Tiffany, H.
L.; Gao,
- [108214]J.-L.; Murphy, P. M.: Monocyte chemoattractant protein-3
is a functional
- [108215]ligand for CC chemokine receptors 1 and 2B. J. Biol.
Chem. 270:
- [108216]29671–29675, 1995.

[108217]19043.

[108218]Doranz, B. J.; Rucker, J.; Yi, Y.; Smyth, R. J.; Samson, M.;
Peiper,

[108219]S. C.; Parmentier, M.; Collman, R. G.; Doms, R. W.: A dual-
tropic

[108220]primary HIV-1 isolate that uses fusin and the beta-
chemokine receptors

[108221]CKR-5, CKR-3, and CKR-2b as fusion cofactors. Cell 85:
1149-1158,

[108222]1996.

[108223]19044.

[108224]Mummidi, S.; Ahuja, S. S.; Gonzalez, E.; Anderson, S. A.;
Santiago,

[108225]E. N.; Stephan, K. T.; Craig, F. E.; O'Connell, P.; Tryon, V.;
Clark,

[108226]R. A.; Dolan, M. J.; Ahuja, S. K.: Genealogy of the CCR5 lo-
cus and

[108227]chemokine system gene variants associated with altered
rates of HIV-1

[108228]disease progression. Nature Med. 4: 786-793, 1998.

[108229]19045.

[108230]Peters, W.; Dupuis, M.; Charo, I. F.: A mechanism for the impaired

[108231]IFN- γ production in C-C chemokine receptor 2 (CCR2) knockout mice:

[108232]role of CCR2 in linking the innate and adaptive immune responses. J.

[108233]Immun. 165: 7072–7077, 2000.

[108234]19046.

[108235]Peters, W.; Scott, H. M.; Chambers, H. F.; Flynn J. L.; Charo,

[108236]I. F.; Ernst, J. D.: Chemokine receptor 2 serves an early and essential

[108237]role in resistance to Mycobacterium tuberculosis. Proc. Nat. Acad.

[108238]Sci. 98: 7958–7963, 2001.

[108239]19047.

[108240]Samson, M.; Labbe, O.; Mollereau, C.; Vassart, G.; Parmentier,

[108241]M.: Molecular cloning and functional expression of a new human CC-chemokine

[108242]receptor gene. Biochemistry 35: 3362–3367, 1996.

[108243]19048.

[108244]Samson, M.; Soularue, P.; Vassart, G.; Parmentier, M.: The
genes
[108245]encoding the human CC-chemokine receptors CC-CCR1
to CC-CCR5 (CMKBR1-CMKBR5)
[108246]are clustered in the p21.3-p24 region of chromosome 3.
Genomics 36:
[108247]522-526, 1996.
[108248]19049.
[108249]Sanders, S. K.; Crean, S. M.; Boxer, P. A.; Kellner, D.;
LaRosa,
[108250]G. J.; Hunt, S. W., III.: Functional differences between
monocyte
[108251]chemotactic protein-1 receptor A and monocyte chemo-
tactic protein-1
[108252]receptor B expressed in a Jurkat T cell. J. Immun. 165:
4877-4883,
[108253]2000.
[108254]19050.
[108255]Smith, M. W.; Dean, M.; Carrington, M.; Winkler, C.; Hutt-
ley,
[108256]G. A.; Lomb, D. A.; Goedert, J. J.; O'Brien, T. R.; Jacobson,
L. P.;

- [108257]Kaslow, R.; Buchbinder, S.; Vittinghoff, E.; Vlahov, D.;
Hoots, K.;
- [108258]Hilgartner, M. W.; Hemophilia Growth and Development
Study (HGDS);
- [108259]Multicenter AIDS Cohort Study (MACS); Multicenter
Hemophilia Cohort
- [108260]Study (MHCS); San Francisco City Cohort (SFCC); ALIVE
Study;
- [108261]O'Brien, S. J.: Contrasting genetic influence of CCR2 and
CCR5 variants
- [108262]on HIV-1 infection and disease progression. *Science* 277:
959-965,
- [108263]1997.
- [108264]19051.
- [108265]Wong, L.-M.; Myers, S. J.; Tsou, C.-L.; Gosling, J.; Arai, H.;
- [108266]Charo, I. F.: Organization and differential expression of
the human
- [108267]monocyte chemoattractant protein 1 receptor gene: evi-
dence for the
- [108268]role of the carboxyl-terminal tail in receptor trafficking. *J.*
Biol.
- [108269]*Chem.* 272: 1038-1045, 1997.

[108270]19052.

[108271]Robertson, N. G.; Khetarpal, U.; Gutierrez-Espeleta, G. A.;
Bieber,

[108272]F. R.; Morton, C. C.: Isolation of novel and known genes
from a human

[108273]fetal cochlear cDNA library using subtractive hybridization
and differential

[108274]screening. Genomics 23: 42-50, 1994.

[108275]19053.

[108276]Rothschild, C. B.; Freedman, B. I.; Hodge, R.; Rao, P. N.;
Pettenati,

[108277]M. J.; Anderson, R. A.; Akots, G.; Qadri, A.; Roh, B.; Fajans,
S.

[108278]S.; Reis, A.; Morris, D. J.; Usala, A.; Hayward, C.; Brock, D.;
Colle,

[108279]E.; Spray, B. J.; Rich, S. S.; Bowden, D. W.: Fructose-
1,6-bisphosphatase:

[108280]genetic and physical mapping to human chromosome
9q22.3 and evaluation

[108281]in non-insulin-dependent diabetes mellitus. Genomics 29:
187-194,

[108282]1995.

[108283]19054.

[108284]Simunovic, M. P.; Regan, B. C.; Mollon, J. D.: Is color vision

[108285]deficiency an advantage under scotopic conditions? Invest.
Ophthalm.

[108286]Vis. Sci. 42: 3357–3364, 2001.

[108287]19055.

[108288]Austen, K. F.: Inborn errors of the complement system of
man. New

[108289]Eng. J. Med. 276: 1363–1367, 1967.

[108290]19056.

[108291]Austen, K. F.; Becker, E. L.; Bero, C. E.; Borsos, T.; Dal-
masso,

[108292]A. P.; Dias Da Silva, D.: Nomenclature of complement. Bull.
WHO 39:

[108293]935–938, 1968.

[108294]19057.

[108295]Awdeh, Z. L.; Raum, D. D.; Glass, D.; Agnello, V.; Schur, P.
H.;

[108296]Johnston, R. B., Jr.; Gelfand, E. W.; Ballow, M.; Yunis, E.;
Alper,

- [108297]C. A.: Complement–human histocompatibility antigen
haplotypes in
- [108298]C2 deficiency. J. Clin. Invest. 67: 581–583, 1981.
- [108299]19058.
- [108300]Belin, D. C.; Bordwell, B. J.; Einarson, M. E.; McLean, R. H.;
- [108301]Weinstein, A.; Yunis, E. J.; Rothfield, N. F.: Familial discoid
lupus
- [108302]erythematosus associated with heterozygote C2 defi-
ciency. Arthritis
- [108303]Rheum. 23: 898–903, 1980.
- [108304]19059.
- [108305]Bentley, D. R.: Primary structure of human complement
component
- [108306]C2: homology to two unrelated protein families. Biochem.
J. 239:
- [108307]339–345, 1986.
- [108308]19060.
- [108309]Bentley, D. R.; Porter, R. R.: Isolation of cDNA clones for
human
- [108310]complement component C2. Proc. Nat. Acad. Sci. 81:
1212–1215, 1984.

[108311]19061.

[108312]Bottger, E. C.; Hoffmann, T.; Hadding, U.; Bitter-Suer-
mann, D.

[108313]: Guinea pigs with inherited deficiencies of complement
components

[108314]C2 or C4 have characteristics of immune complex disease.
J. Clin.

[108315]Invest. 78: 689–695, 1986.

[108316]19062.

[108317]Callen, J. P.; Hodge, S. J.; Kulick, K. B.; Stelzer, G.; Buchino,

[108318]J. J.: Subacute cutaneous lupus erythematosus in multiple
members

[108319]of a family with C2 deficiency. Arch. Derm. 123: 66–70,
1987.

[108320]19063.

[108321]Cole, F. S.; Whitehead, A. S.; Auerbach, H. S.; Lint, T.;
Zeitz,

[108322]H. J.; Kilbridge, P.; Colten, H. R.: The molecular basis for
genetic

[108323]deficiency of the second component of human comple-
ment. New Eng.

[108324]J. Med. 313: 11-16, 1985.

[108325]19064.

[108326]Donald, L. J.; Wang, H. S.; Hamerton, J. L.: Assignment of the

[108327]gene for cystathionase (CYS) to human chromosome 16.
(Abstract) Cytogenet.

[108328]Cell Genet. 32: 268 only, 1982.

[108329]19065.

[108330]Frimpter, G. W.: Cystathioninuria: nature of the defect.
Science 149:

[108331]1095-1096, 1965.

[108332]19066.

[108333]Frimpter, G. W.: Cystathioninuria, sulfite oxidase deficiency,

[108334]and 'beta-mercaptolactate-cysteine disulfiduria.In: Stanbury, J. B.;

[108335]Wyngaarden, J. B.; Fredrickson, D. S.: The Metabolic Basis
of Inherited

[108336]Disease. New York: McGraw-Hill (pub.) (3rd ed.): 1972. Pp.
413-425.

[108337]19067.

- [108338]Frimpter, G. W.; Haymovitz, A.; Horwith, M.: Cystathionin-
uria. New
- [108339]Eng. J. Med. 268: 333–339, 1963.
- [108340]19068.
- [108341]Harris, H.; Penrose, L. S.; Thomas, D. H. H.: Cystathionin-
uria. Ann.
- [108342]Hum. Genet. 23: 442–453, 1959.
- [108343]19069.
- [108344]Lu, Y.; O'Dowd, B. F.; Orrego, H.; Israel, Y.: Cloning and
nucleotide
- [108345]sequence of human liver cDNA encoding for cystathionine
gamma-lyase. Biochem.
- [108346]Biophys. Res. Commun. 189: 749–758, 1992.
- [108347]19070.
- [108348]Lyon, I. C. T.; Procopis, P. G.; Turner, B.: Cystathioninuria
- [108349]in a well baby population. Acta Paediat. Scand. 60:
324–328, 1971.
- [108350]19071.
- [108351]Mongeau, J.-G.; Hilgartner, M.; Worthen, H. G.; Frimpter,
G. W.

[108352]: Cystathioninuria: study of an infant with normal mental-
ity, thrombocytopenia,

[108353]and renal calculi. J. Pediat. 69: 1113–1120, 1967.

[108354]19072.

[108355]Pascal, T. A.; Gaull, G. E.; Beratis, N. G.; Gillam, B. M.; Tal-
lan,

[108356]H. H.: Cystathionase deficiency: evidence for genetic het-
erogeneity

[108357]in primary cystathioninuria. Pediat. Res. 12: 125–133,
1978.

[108358]19073.

[108359]Perry, T. L.; Hardwick, D. F.; Hansen, S.; Love, D. L.; Is-
raels,

[108360]S.: Cystathioninuria in two healthy siblings. New Eng. J.
Med. 278:

[108361]590–592, 1968.

[108362]19074.

[108363]Schneiderman, L. J.: Latent cystathioninuria. J. Med. Genet.
4:

[108364]260–263, 1967.

[108365]19075.

- [108366] Scott, C. R.; Dassell, S. W.; Clark, S. H.; Chiang-Teng, C.; Swedberg, K. R.: Cystathioninemia: a benign genetic condition. J. Pediat. 76: 571-577, 1970.
- [108369] 19076.
- [108370] Shaw, K. N. F.; Lieberman, E.; Koch, R.; Donnell, G. N.: Cystathioninuria. Am. J. Dis. Child. 113: 119-128, 1967.
- [108372] 19077.
- [108373] Tada, K.; Yoshida, T.; Yokoyama, Y.; Sato, T.; Nakagawa, H.; Arakawa, T.: Cystathioninuria not associated with vitamin B6 dependency: a probably new type of cystathioninuria. Tohoku J. Exp. Med. 95: 235-242, 1968.
- [108376] 19078.
- [108377] 19078.
- [108378] Whelan, D. T.; Scriver, C. R.: Cystathioninuria and renal iminoglycinuria in a pedigree: a perspective on counseling. New Eng. J.

Med. 278:

[108380]924-927, 1968.

[108381]19079.

[108382]Abeliovich, D.; Lavon, I. P.; Lerer, I.; Cohen, T.; Springer, C.;

[108383]Avital, A.; Cutting, G. R.: Screening for five mutations de-
fects

[108384]97% of cystic fibrosis (CF) chromosomes and predicts a
carrier frequency

[108385]of 1:29 in the Jewish Ashkenazi population. Am. J. Hum.
Genet. 51:

[108386]951-956, 1992.

[108387]19080.

[108388]Hobart, M. J.; Joysey, V.; Lachmann, P. J.: Inherited struc-
tural

[108389]variation and linkage relationships of C7. J. Immunogenet.
5: 157-163,

[108390]1978.

[108391]19081.

[108392]Jeremiah, S. J.; Abbott, C. M.; Murad, Z.; Povey, S.;
Thomas,

- [108393]H. J.; Solomon, E.; DiScipio, R. G.; Fey, G. H.: The assignment of
the genes coding for human complement components C6
and C7 to chromosome
5. *Ann. Hum. Genet.* 54: 141–147, 1990.
19082.
- [108397]Schulpis, K. H.; Papaconstantinou, E. D.; Koidou, A.;
Michelakakis,
H.; Tzamouranis, J.; Patsouras, A.; Shin, Y.: UDP galac-
tose-4-epimerase
deficiency in a 5.5-year-old girl with a unilateral cataract.
J.
Inherit. Metab. Dis. 16: 903–904, 1993.
19083.
- [108402]Thoden, J. B.; Wohlers, T. M.; Fridovich-Keil, J. L.; Holden,
H. M.: Molecular basis for severe epimerase deficiency
galactosemia:
x-ray structure of the human V94M-substituted UDP-
galactose 4-epimerase. J.
Biol. Chem. 276: 20617–20623, 2001.
19084.

[108407] Wohlers, T. M.; Christacos, N. C.; Harreman, M. T.;
Fridovich-Keil,
[108408] J. L.: Identification and characterization of a mutation, in
the
[108409] human UDP-galactose-4-epimerase gene, associated with
generalized
[108410] epimerase-deficiency galactosemia. Am. J. Hum. Genet.
64: 462-470,
[108411] 1999.
[108412] 19085.
[108413] Cheadle, J.; Al-Jader, L.; Goodchild, M.; Meredith, A. L.:
Mild
[108414] pulmonary disease in a cystic fibrosis child homozygous
for R553X. J.
[108415] Med. Genet. 29: 597, 1992.
[108416] 19086.
[108417] Cheadle, J. P.; Meredith, A. L.; Al-Jader, L. N.: A new mis-
sense
[108418] mutation (R1283M) in exon 20 of the cystic fibrosis trans-
membrane
[108419] conductance regulator gene. Hum. Molec. Genet. 1:
123-125, 1992.

[108420]19087.

[108421]Colledge, W. H.; Abella, B. S.; Southern, K. W.; Ratcliff, R.;

[108422]Jiang, C.; Cheng, S. H.; MacVinish, L. J.; Anderson, J. R.;
Cuthbert,

[108423]A. W.; Evans, M. J.: Generation and characterisation of a
delta-F508

[108424]cystic fibrosis mouse model. *Nature Genet.* 10: 445-452,
1995.

[108425]19088.

[108426]Haworth, J. C.; Gourley, B.; Hadorn, B.; Sumida, C.: Malab-
sorption

[108427]and growth failure due to intestinal enterokinase defi-
ciency. *J.*

[108428]*Pediat.* 78: 481-490, 1971.

[108429]19089.

[108430]Haworth, J. C.; Hadorn, B.; Gourley, B.; Prasad, A.;
Troesch, V.

[108431]: Intestinal enterokinase deficiency: occurrence in two sibs
and age

[108432]dependency of clinical expression. *Arch. Dis. Child.* 50:
277-282,

[108433]1975.

[108434]19090.

[108435]Holzinger, A.; Maier, E. M.; Buck, C.; Mayerhofer, P. U.;
Kappler,

[108436]M.; Haworth, J. C.; Moroz, S. P.; Hadorn, H.-B.; Sadler, J.
E.; Roscher,

[108437]A. A.: Mutations in the proenteropeptidase gene are the
molecular

[108438]cause of congenital enteropeptidase deficiency. Am. J.
Hum. Genet. 70:

[108439]20–25, 2002.

[108440]19091.

[108441]DiGeorge, A. M.; Harley, R. D. L.: The association of
aniridia,

[108442]Wilms tumor and genital abnormalities. Arch. Ophthalm. 75:
796–798,

[108443]1966.

[108444]19092.

[108445]Dressler, G. R.; Douglass, E. C.: Pax-2 is a DNA-binding
protein

[108446]expressed in embryonic kidney and Wilms tumor. Proc.

Nat. Acad. Sci. 89:

[108447]1179–1183, 1992.

[108448]19093.

[108449]Fearon, E. R.; Vogelstein, B.; Feinberg, A. P.: Somatic deletion

[108450]and duplication of genes on chromosome 11 in Wilms' tumours. Nature 309:

[108451]176–178, 1984.

[108452]19094.

[108453]Fitzgerald, W. L.; Hardin, H. C., Jr.: Bilateral Wilms' tumor

[108454]in a Wilms tumor family: case report. J. Urol. 73: 468–474, 1955.

[108455]19095.

[108456]Francke, U.: A gene for Wilms tumour? Nature 343: 692–694,

[108457]1990.

[108458]19096.

[108459]Francke, U.; Holmes, L. B.; Atkins, L.; Riccardi, V. M.: Aniridia–Wilms'

[108460]tumor association: evidence for specific deletion of

11p13. Cytogenet.

[108461]Cell Genet. 24: 185-192, 1979.

[108462]19097.

[108463]Francke, U.; Riccardi, V. M.; Hittner, H. M.; Borges, W.: In-
terstitial

[108464]del(11p) as a cause of the aniridia-Wilms tumor associa-
tion: band

[108465]localization and a heritable basis. (Abstract) Am. J. Hum.
Genet. 30:

[108466]81A, 1978.

[108467]19098.

[108468]Fraumeni, J. F., Jr.; Glass, A. G.: Wilms' tumor and congen-
ital

[108469]aniridia. J.A.M.A. 206: 825-828, 1968.

[108470]19099.

[108471]Haber, D. A.; Buckler, A. J.; Glaser, T.; Call, K. M.; Pelletier,

[108472]J.; Sohn, R. L.; Douglass, E. C.; Housman, D. E.: An internal
deletion

[108473]within an 11p13 zinc finger gene contributes to the de-
velopment of

[108474]Wilms' tumor. Cell 61: 1257-1269, 1990.

[108475]19100.

[108476]Haber, D. A.; Park, S.; Maheswaran, S.; Englert, C.; Re, G.
G.;

[108477]Hazen-Martin, D. J.; Sens, D. A.; Garvin, A. J.:
WT1-mediated growth

[108478]suppression of Wilms tumor cells expressing a WT1 splic-
ing variant. Science 262:

[108479]2057-2059, 1993.

[108480]19101.

[108481]Haber, D. A.; Timmers, H. T. M.; Pelletier, J.; Sharp, P. A.;

[108482]Housman, D. E.: A dominant mutation in the Wilms tumor
gene WT1 cooperates

[108483]with the viral oncogene E1A in transformation of primary
kidney cells. Proc.

[108484]Nat. Acad. Sci. 89: 6010-6014, 1992.

[108485]19102.

[108486]Haning, R. V., Jr.; Chesney, R. W.; Moorthy, A. V.; Gilbert,
E.

[108487]F.: A syndrome of chronic renal failure and XY gonadal
dysgenesis

[108488]in young phenotypic females without genital ambiguity.

Am. J. Kidney

[108489]Dis. 6: 40–48, 1985.

[108490]19103.

[108491]Hastie, N. D.: Dominant negative mutations in the Wilms
tumour

[108492](WT1) gene cause Denys–Drash syndrome--proof that a
tumour-suppressor

[108493]gene plays a crucial role in normal genitourinary develop-
ment. Hum.

[108494]Molec. Genet. 1: 293–295, 1992.

[108495]19104.

[108496]Huff, V.; Meadows, A.; Riccardi, V. M.; Strong, L. C.; Saun-
ders,

[108497]G. F.: Parental origin of de novo constitutional deletions of
chromosomal

[108498]band 11p13. Am. J. Hum. Genet. 47: 155–160, 1990.

[108499]19105.

[108500]Jeanpierre, C.; Antignac, C.; Beroud, C.; Lavedan, C.;
Henry,

[108501]J.; Saunders, G.; Williams, B.; Glaser, T.; Junien, C.: Consti-
tutional

[108502]and somatic deletions of two different regions of maternal chromosome

[108503]11 in Wilms tumor. Genomics 7: 434-438, 1990.

[108504]19106.

[108505]Jolles, B.: Wilms' tumor in father and son. Lancet I: 207,

[108506]1973.

[108507]19107.

[108508]Juberg, R. C.; St. Martin, E. C.; Hundley, J. R.: Familial occurrence

[108509]of Wilms's tumor: nephroblastoma in one of monozygous twins and in

[108510]another sibling. Am. J. Hum. Genet. 27: 155-169, 1975.

[108511]19108.

[108512]Junien, C.; Huerre, C.; Despoisse, S.; Gilgenkrantz, S.; Lenoir,

[108513]G. M.: c-Ha-ras1 is not deleted in del (11p13) Wilms' tumor (WAGR)

[108514]and maps to 11p15.1-11p15.5. (Abstract) Cytogenet. Cell Genet. 37:

[108515]503, 1984.

[108516]19109.

[108517]Junien, C.; Turleau, C.; de Grouchy, J.; Said, R.; Rethore, M.-O.;

[108518]Tenconi, R.; Dufier, J. L.: Regional assignment of catalase (CAT)

[108519]gene to band 11p13: association with the aniridia-Wilms' tumor-gonadoblastoma

[108520](WAGR) complex. Ann. Genet. 23: 165-168, 1980.

[108521]19110.

[108522]Kakati, S.; Xiao, H.; Siddiqui, S. Y.; Sreekantaiah, C.; Weier,

[108523]H.-U. G.; Green, D. M.; Fisher, J. E.; Allen, J. E.: Constitutional

[108524]extra chromosomal element in a family with Wilms' tumor. Hum. Genet. 87:

[108525]183-188, 1991.

[108526]19111.

[108527]Kaneko, Y.; Celida Egues, M.; Rowley, J. D.: Interstitial deletion

[108528]of short arm of chromosome 11 limited to Wilms' tumor cells in a patient

[108529]without aniridia. Cancer Res. 41: 4577-4578, 1981.

[108530]19112.

[108531]Kaufman, R. L.; Vietti, T. J.; Wabner, C. I.: Wilms' tumor in
[108532]father and son. Lancet I: 43, 1973.

[108533]19113.

[108534]Kinberg, J. A.; Angle, C. R.; Wilson, R. B.: Nephropathy-go-
nadal
[108535]dysgenesis, type 2: renal failure in three siblings with XY
dysgenesis

[108536]in one. Am. J. Kidney Dis. 9: 507-510, 1987.

[108537]19114.

[108538]Knudson, A. G., Jr.; Strong, L. C.: Mutation and cancer: a
model

[108539]for Wilms' tumor of the kidney. J. Nat. Cancer Inst. 48:
313-324,

[108540]1972.

[108541]19115.

[108542]Knudson, A. G., Jr.; Strong, L. C.: Familial Wilms's tumor.
(Letter) Am.

[108543]J. Hum. Genet. 27: 809-810, 1975.

[108544]19116.

- [108545]Kolata, G. B.: Genes and cancer: the story of Wilms tumor.
Science 207:
[108546]970–971, 1980.
[108547]19117.
- [108548]Kontras, S. B.; Newton, W. A., Jr.: Familial Wilms' tumor.
Birth
[108549]Defects Orig. Art. Ser. X(4): 187–188, 1974.
[108550]19118.
- [108551]Koufos, A.; Hansen, M. F.; Lampkin, B. C.; Workman, M. L.;
Copeland,
[108552]N. G.; Jenkins, N. A.; Cavenee, W. K.: Loss of alleles at loci
on
[108553]human chromosome 11 during genesis of Wilms' tumour.
Nature 309:
[108554]170–172, 1984.
[108555]19119.
- [108556]Kozman, H. M.; Clarke, J. M.; Little, M. H.; Smith, P. J.:
Molecular
[108557]genetic evidence for common pathogenesis of childhood
and adult Wilms'
[108558]tumor. Cancer Genet. Cytogenet. 38: 121–125, 1989.

[108559]19120.

[108560]Kumar, S.; Harrison, C. J.; Heighway, J.; Marsden, H. B.;
West,

[108561]D. C.; Jones, P. M.: A cell line from Wilms' tumour with
deletion

[108562]in short arm of chromosome 11. Int. J. Cancer 40:
499–504, 1987.

[108563]19121.

[108564]Ladda, R. L.; Atkins, L.; Littlefield, J.: Computer-assisted

[108565]analysis of chromosomal abnormalities: detection of a
deletion in

[108566]aniridia Wilms' tumor syndrome. Science 185: 784–787,
1974.

[108567]19122.

[108568]Berx, G.; Staes, K.; van Hengel, J.; Molemans, F.; Busse-
makers,

[108569]M. J. G.; van Bokhoven, A.; van Roy, F.: Cloning and char-
acterization

[108570]of the human invasion suppressor gene E-cadherin
(CDH1). Genomics 26:

[108571]281–289, 1995.

[108572]19123.

[108573]Boggon, T. J.; Murray, J.; Chappuis-Flament, S.; Wong, E.;
Gumbiner,

[108574]B. M.; Shapiro, L.: C-cadherin ectodomain structure and
implications

[108575]for cell adhesion mechanisms. Science 296: 1308–1313,
2002.

[108576]19124.

[108577]Cano, A.; Perez-Moreno, M. A.; Rodrigo, I.; Locascio, A.;
Blanco,

[108578]M. J.; del Barrio, M. G.; Portillo, F.; Nieto, M. A.: The tran-
scription

[108579]factor Snail controls epithelial–mesenchymal transitions
by repressing

[108580]E-cadherin expression. Nature Cell Biol. 2: 76–83, 2000.

[108581]19125.

[108582]Ceccherini, I.; Romeo, G.; Lawrence, S.; Breuning, M. H.;
Harris,

[108583]P. C.; Himmelbauer, H.; Frischauf, A. M.; Sutherland, G. R.;
Germino,

[108584]G. G.; Reeders, S. T.; Morton, N. E.: Construction of a map
of chromosome

[108585]16 by using radiation hybrids. Proc. Nat. Acad. Sci. 89:
104–108,

[108586]1992.

[108587]19126.

[108588]Chun, Y. S.; Lindor, N. M.; Smyrk, T. C.; Petersen, B. T.;
Burgart,

[108589]L. J.; Guilford, P. J.; Donohue, J. H.: Germline E-cadherin
gene

[108590]mutations: is prophylactic total gastrectomy indicated?
Cancer 92:

[108591]181–187, 2001.

[108592]19127.

[108593]Eistetter, H. R.; Adolph, S.; Ringwald, M.; Simon–Cha-
zottes, D.;

[108594]Schuh, R.; Guenet, J. L.; Kemler, R.: Chromosomal map-
ping of the

[108595]structural gene coding for the mouse cell adhesion
molecule uvomorulin. Proc.

[108596]Nat. Acad. Sci. 85: 3489–3493, 1988.

[108597]19128.

[108598]Gayther, S. A.; Goringe, K. L.; Ramus, S. J.; Huntsman, D.;

Roviello,

[108599]F.; Grehan, N.; Machado, J. C.; Pinto, E.; Seruca, R.; Halling, K.;

[108600]MacLeod, P.; Powell, S. M.; Jackson, C. E.; Ponder, B. A. J.; Caldas,

[108601]C.: Identification of germ-line E-cadherin mutations in gastric cancer

[108602]families of European origin. Cancer Res. 58: 4086–4089, 1998.

[108603]19129.

[108604]Grady, W. M.; Willis, J.; Guilford, P. J.; Dunbier, A. K.; Toro,

[108605]T. T.; Lynch, H.; Wiesner, G.; Ferguson, K.; Eng, C.; Park, J.-G.;

[108606]Kim, S.-J.; Markowitz, S.: Methylation of the CDH1 promoter as the

[108607]second genetic hit in hereditary diffuse gastric cancer. Nature Genet. 26:

[108608]16–17, 2000.

[108609]19130.

[108610]Humar, B.; Toro, T.; Graziano, F.; Muller, H.; Dobbie, Z.; Kwang-Yang,

[108611]H.; Eng, C.; Hampel, H.; Gilbert, D.; Winship, I.; Parry, S.;

Ward,

[108612]R.; Findlay, M.; Christian, A.; Tucker, M.; Tucker, K.; Merri-
man,

[108613]T.; Guilford, P.: Novel germline CDH1 mutations in heredi-
tary diffuse

[108614]gastric cancer families. Hum. Mutat. 19: 518–525, 2002.

[108615]19131.

[108616]Ilyas, M.; Tomlinson, I. P. M.; Hanby, A.; Talbot, I. C.; Bod-
mer,

[108617]W. F.: Allele loss, replication errors and loss of expression
of

[108618]E-cadherin in colorectal cancers. Gut 40: 654–659, 1997.

[108619]19132.

[108620]Jones, E. G.: Familial gastric cancer. New Zealand Med. J.
63:

[108621]287–296, 1964.

[108622]19133.

[108623]Lecuit, M.; Dramsi, S.; Gottardi, C.; Fedor–Chaiken, M.;
Gumbiner,

[108624]B.; Cossart, P.: A single amino acid in E-cadherin respon-
sible for

[108625] host specificity towards the human pathogen *Listeria monocytogenes*. *EMBO*

[108626] *J.* 18: 3956–3963, 1999.

[108627] 19134.

[108628] Guilford, P. J.; Hopkins, J. B. W.; Grady, W. M.; Markowitz, S.

[108629] D.; Willis, J.; Lynch, H.; Rajput, A.; Wiesner, G. L.; Lindor, N.

[108630] M.; Burgart, L. J.; Toro, T. T.; Lee, D.; Limacher, J.-M.; Shaw, D.

[108631] W.; Findlay, M. P. N.; Reeve, A. E.: E-cadherin germline mutations

[108632] define an inherited cancer syndrome dominated by diffuse gastric cancer. *Hum.*

[108633] *Mutat.* 14: 249–255, 1999.

[108634] 19135.

[108635] Lecuit, M.; Vandormael-Pournin, S.; Lefort, J.; Huerre, M.; Gounon,

[108636] P.; Dupuy, C.; Babinet, C.; Cossart, P.: A transgenic model for listeriosis:

[108637] role of internalin in crossing the intestinal barrier. *Science* 292:

[108638]1722–1725, 2001.

[108639]19136.

[108640]Lorber, B.: Listeriosis. Clin. Infect. Dis. 24: 1–11, 1997.

[108641]19137.

[108642]Lynch, H. T.; Grady, W.; Lynch, J. F.; Tsuchiya, K. D.; Wiesner,

[108643]G.; Markowitz, S. D.: E-cadherin mutation-based genetic counseling

[108644]and hereditary diffuse gastric carcinoma. Cancer Genet. Cytogenet. 122:

[108645]1–6, 2000.

[108646]19138.

[108647]Mansouri, A.; Goodfellow, P. N.; Kemler, R.: Molecular cloning

[108648]and chromosomal localization of the human cell adhesion molecule uvomorulin

[108649](UVO). (Abstract) Cytogenet. Cell Genet. 46: 655, 1987.

[108650]19139.

[108651]Mansouri, A.; Spurr, N.; Goodfellow, P. N.; Kemler, R.: Characterization

[108652]and chromosomal localization of the gene encoding the human cell adhesion

[108653]molecule uvomorulin. Differentiation 38: 67–71, 1988.

[108654]19140.

[108655]Natt, E.; Magenis, R. E.; Zimmer, J.; Mansouri, A.; Scherer, G.

[108656]; Regional assignment of the human loci for uvomorulin (UVO) and chymotrypsinogen

[108657]B (CTRB) with the help of two overlapping deletions on the long arm

[108658]of chromosome 16. Cytogenet. Cell Genet. 50: 145–148, 1989.

[108659]19141.

[108660]Oda, T.; Kanai, Y.; Oyama, T.; Yoshiura, K.; Shimoyama, Y.; Birchmeier,

[108661]W.; Sugimura, T.; Hirohashi, S.: E-cadherin gene mutations in human

[108662]gastric carcinoma cell lines. Proc. Nat. Acad. Sci. 91: 1858–1862,

[108663]1994.

[108664]19142.

- [108665]Oliveira, C.; Bordin, M. C.; Grehan, N.; Huntsman, D.; Suriano,
ano,
- [108666]G.; Machado, J. C.; Kiviluoto, T.; Aaltonen, L.; Jackson, C.
E.; Seruca,
- [108667]R.; Caldas, C.: Screening E-cadherin in gastric cancer families reveals
germline mutations only in hereditary diffuse gastric cancer kindred. Hum.
Mutat. 19: 510–517, 2002.
- [108670]19143.
- [108671]Overduin, M.; Harvey, T. S.; Bagby, S.; Tong, K. I.; Yau, P.;
- [108672]Takeichi, M.; Ikura, M.: Solution structure of the epithelial
cadherin
domain responsible for selective cell adhesion. Science
267: 386–389,
1995.
- [108675]19144.
- [108676]Perl, A.-K.; Wilgenbus, P.; Dahl, U.; Semb, H.; Christofori,
G.
- [108677]: A causal role for E-cadherin in the transition from adenoma to carcinoma. Nature 392:

[108678]190–193, 1998.

[108679]19145.

[108680]Joenje, H.: Fanconi anaemia complementation groups in
Germany

[108681]and The Netherlands. Hum. Genet. 97: 280–282, 1996.

[108682]19146.

[108683]Joenje, H.; Oostra, A. B.; Wijker, M.; di Summa, F. M.; van
Berkel,

[108684]C. G. M.; Rooimans, M. A.; Ebell, W.; van Weel, M.; Pronk, J.
C.;

[108685]Buchwald, M.; Arwert, F.: Evidence for at least eight Fan-
coni anemia

[108686]genes. Am. J. Hum. Genet. 61: 940–944, 1997.

[108687]19147.

[108688]Rassool, F. V.; Le Beau, M. M.; Shen, M.–L.; Neilly, M. E.;
Espinosa,

[108689]R., III; Ong, S. T.; Boldog, F.; Drabkin, H.; McCarroll, R.;
McKeithan,

[108690]T. W.: Direct cloning of DNA sequences from the common
fragile site

[108691]region at chromosome band 3p14.2. Genomics 35:

109–117, 1996.

[108692]19148.

[108693]Rudduck, C.; Franzen, G.: A new heritable fragile site on human

[108694]chromosome 3. *Hereditas* 98: 297–299, 1983.

[108695]19149.

[108696]Shi, Y.; Zou, M.; Farid, N. R.; Paterson, M. C.: Association

[108697]of FHIT (fragile histidine triad), a candidate tumour suppressor gene,

[108698]with the ubiquitin-conjugating enzyme hUBC9. *Biochem. J.* 352: 443–448,

[108699]2000.

[108700]19150.

[108701]Shiraishi, T.; Druck, T.; Mimori, K.; Flomenberg, J.; Berk, L.;

[108702]Alder, H.; Miller, W.; Huebner, K.; Croce, C. M.: Sequence conservation

[108703]at human and mouse orthologous common fragile regions, FRA3B/FHIT

[108704]and Fra14A2/Fhit. *Proc. Nat. Acad. Sci.* 98: 5722–5727, 2001.

[108705]19151.

[108706]Sozzi, G.; Veronese, M. L.; Negrini, M.; Baffa, R.; Cotticelli,

[108707]M. G.; Inoue, H.; Tornielli, S.; Pilotti, S.; De Gregorio, L.;
Pastorino,

[108708]U.; Pierotti, M. A.; Ohta, M.; Huebner, K.; Croce, C. M.: The
FHIT

[108709]gene at 3p14.2 is abnormal in lung cancer. Cell 85:
17-26, 1996.

[108710]19152.

[108711]Stein, C. K.; Glover, T. W.; Palmer, J. L.; Glisson, B. S.: Di-
rect

[108712]correlation between FRA3B expression and cigarette
smoking. Genes

[108713]Chromosomes Cancer 34: 333-340, 2002.

[108714]19153.

[108715]Virgilio, L.; Shuster, M.; Gollin, S. M.; Veronese, M. L.;
Ohta,

[108716]M.; Huebner, K.; Croce, C. M.: FHIT gene alterations in
head and

[108717]neck squamous cell carcinomas. Proc. Nat. Acad. Sci. 93:
9770-9775,

[108718]1996.

[108719]19154.

[108720]Wegner, R.-D.: A new inducible fragile site on chromosome 3 (p14.2)

[108721]in human lymphocytes. Hum. Genet. 63: 297-298, 1983.

[108722]19155.

[108723]Wegner, R.-D.: Reply to the letter of A. Markkanen, S. Knuutila,

[108724]and A. de la Chapelle. (Letter) Hum. Genet. 65: 218 only, 1983.

[108725]19156.

[108726]Zanesi, N.; Fidanza, V.; Fong, L. Y.; Mancini, R.; Druck, T.;

[108727]Valtieri, M.; Rudiger, T.; McCue, P. A.; Croce, C. M.; Huebner, K.

[108728]: The tumor spectrum in FHIT-deficient mice. Proc. Nat. Acad. Sci. 98:

[108729]10250-10255, 2001.

[108730]19157.

[108731]Nagase, T.; Seki, N.; Ishikawa, K.; Ohira, M.; Kawarabayashi, Y.;

[108732]Ohara, O.; Tanaka, A.; Kotani, H.; Miyajima, N.; Nomura, N.: Prediction
[108733]of the coding sequences of unidentified human genes. VI. The coding
[108734]sequences of 80 new genes (KIAA0201–KIAA0280) deduced by analysis
[108735]of cDNA clones from cell line KG–1 and brain. DNA Res. 3: 321–329,
[108736]1996.
[108737]19158.
[108738]Lee, E.; Marcucci, M.; Daniell, L.; Pypaert, M.; Weisz, O. A.;
[108739]Ochoa, G.–C.; Farsad, K.; Wenk, M. R.; De Camilli, P.: Amphiphysin
[108740]2 (Bin1) and T-tubule biogenesis in muscle. Science 297: 1193–1196,
[108741]2002.
[108742]19159.
[108743]Negorev, D.; Riethman, H.; Wechsler–Reya, R.; Sakamuro, D.; Prendergast,
[108744]G. C.; Simon, D.: The Bin1 gene localizes to human chromosome 2q14

[108745]by PCR analysis of somatic cell hybrids and fluorescence
in situ hybridization. Genomics 33:

[108746]329–331, 1996.

[108747]19160.

[108748]Sakamuro, D.; Elliott, K. J.; Wechsler-Reya, R.; Prender-
gast, G.

[108749]C.: BIN1 is a novel MYC–interacting protein with features
of a tumour

[108750]suppressor. Nature Genet. 14: 69–77, 1996.

[108751]19161.

[108752]Endo, Y.; Sato, Y.; Matsushita, M.; Fujita, T.: Cloning and
characterization

[108753]of the human lectin P35 gene and its related gene. Ge-
nomics 36:

[108754]515–521, 1996.

[108755]19162.

[108756]Betz, R. C.; Schoser, B. G. H.; Kasper, D.; Ricker, K.;
Ramirez,

[108757]A.; Stein, V.; Torbergesen, T.; Lee, Y.–A.; Nothen, M. M.;
Wienker,

[108758]T. F.; Malin, J.–P.; Propping, P.; Reis, A.; Mortier, W.;

Jentsch,

[108759]T. J.; Vorgerd, M.; Kubisch, C.: Mutations in CAV3 cause mechanical

[108760]hyperirritability of skeletal muscle in rippling muscle disease. Nature

[108761]Genet. 28: 218–219, 2001.

[108762]19163.

[108763]Drumheller, T.; McGillivray, B. C.; Behrner, D.; MacLeod, P.; McFadden,

[108764]D. E.; Roberson, J.; Venditti, C.; Chorney, K.; Chorney, M.; Smith,

[108765]D. I.: Precise localisation of 3p25 breakpoints in four patients

[108766]with the 3p– syndrome. J. Med. Genet. 33: 842–847, 1996.

[108767]19164.

[108768]Galbiati, F.; Volonte, D.; Chu, J. B.; Li, M.; Fine, S. W.; Fu,

[108769]M.; Bermudez, J.; Pedemonte, M.; Weidenheim, K. M.; Pestell, R. G.;

[108770]Minetti, C.; Lisanti, M. P.: Transgenic overexpression of caveolin–3

[108771]in skeletal muscle fibers induces a Duchenne–like muscu–

lar dystrophy

[108772]phenotype. Proc. Nat. Acad. Sci. 97: 9689–9694, 2000.

[108773]19165.

[108774]Wayne, S.; Robertson, N. G.; DeClau, F.; Chen, N.; Verhoeven, K.;

[108775]Prasad, S.; Tranebjarg, L.; Morton, C. C.; Ryan, A. F.; Van Camp,

[108776]G.; Smith, R. J. H.: Mutations in the transcriptional activator EYA4

[108777]cause late-onset deafness at the DFNA10 locus. Hum. Molec. Genet. 10:

[108778]195–200, 2001.

[108779]19166.

[108780]Kanai, Y.; Miura, K.; Uehara, T.; Amagai, M.; Takeda, O.; Tanuma,

[108781]S.; Kurosawa, Y.: Natural occurrence of Nuc in the sera of autoimmune-prone

[108782]MRL/lpr mice. Biochem. Biophys. Res. Commun. 196: 729–736, 1993.

[108783]19167.

[108784]Miura, K.; Hirai, M.; Kanai, Y.; Kurosawa, Y.: Organization

of

[108785]the human gene for nucleobindin (NUC) and its chromosomal assignment

[108786]to 19q13.2–q13.4. Genomics 34: 181–186, 1996.

[108787]19168.

[108788]Miura, K.; Titani, K.; Kurosawa, Y.; Kanai, Y.: Molecular cloning

[108789]of nucleobindin, a novel DNA-binding protein that contains both a

[108790]signal peptide and a leucine zipper structure. Biochem. Biophys.

[108791]Res. Commun. 187: 375–380, 1992.

[108792]19169.

[108793]Brewer, G.: Personal Communication. Winston–Salem, N.C. 3/4/1999.

[108794]19170.

[108795]Brewer, G.: An A+U-rich element RNA-binding factor regulates c-myc

[108796]mRNA stability in vitro. Molec. Cell. Biol. 11: 2460–2466, 1991.

[108797]19171.

[108798]Dempsey, L. A.; Li, M.; DePace, A.; Bray-Ward, P.; Maizels, N.

[108799]: The human HNRPD locus maps to 4q21 and encodes a highly conserved

[108800]protein. Genomics 49: 378–384, 1998.

[108801]19172.

[108802]Wagner, B. J.; DeMaria, C. T.; Sun, Y.; Wilson, G. M.; Brewer,

[108803]G.: Structure and genomic organization of the human AUF1 gene: alternative

[108804]pre-mRNA splicing generates four protein isoforms. Genomics 48:

[108805]195–202, 1998.

[108806]19173.

[108807]Wagner, B. J.; Long, L.; Rao, P. N.; Pettenati, M. J.; Brewer,

[108808]G.: Localization and physical mapping of genes encoding the A+U-rich

[108809]element RNA-binding protein AUF1 to human chromosomes 4 and X. Genomics 34:

[108810]219–222, 1996.

[108811]19174.

[108812]Zhang, W.; Wagner, B. J.; Ehrenman, K.; Schaefer, A. W.;
DeMaria,
[108813]C. T.; Crater, D.; DeHaven, K.; Long, L.; Brewer, G.: Purifi-
cation,
[108814]characterization and cDNA cloning of an AU-rich element
RNA-binding
[108815]protein, AUF1. Molec. Cell. Biol. 13: 7652-7665, 1993.
[108816]19175.
[108817]Connelly, M. A.; Grady, R. C.; Mushinski, J. F.; Marcu, K. B.:
[108818]PANG, a gene encoding a neuronal glycoprotein, is ectopi-
cally activated
[108819]by intracisternal A-type particle long terminal repeats in
murine
[108820]plasmacytomas. Proc. Nat. Acad. Sci. 91: 1337-1341,
1994.
[108821]19176.
[108822]Mock, B. A.; Connelly, M. A.; McBride, O. W.; Kozak, C. A.;
Marcu,
[108823]K. B.: Plasmacytoma-associated neuronal glycoprotein,
Pang, maps
[108824]to mouse chromosome 6 and human chromosome 3. Ge-
nomics 34: 226-228,

[108825]1996.

[108826]19177.

[108827]Bolino, A.; Seri, M.; Caroli, F.; Eubanks, J.; Srinivasan, J.;

[108828]Mandich, P.; Schenone, A.; Quattrone, A.; Romeo, G.; Catterall, W.

[108829]A.; Devoto, M.: Exclusion of the SCN2B gene as candidate for CMT4B. Europ.

[108830]J. Hum. Genet. 6: 629-634, 1998.

[108831]19178.

[108832]Eubanks, J.; Srinivasan, J.; Dinulos, M. B.; Distèche, C. M.; Catterall,

[108833]W. A.: Structure and chromosomal localization of the beta2 subunit

[108834]of the human brain sodium channel. Neuroreport 8: 2775-2779, 1997.

[108835]19179.

[108836]Imai, K.; Kingsley, D. M.: Mouse chromosome 9. Mammalian Genome 5:

[108837]S139-S153, 1994.

[108838]19180.

[108839]Noben–Trauth, K.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N.

[108840]A.; Sonoda, G.; Testa, J. R.; Klempnauer, K.–H.: Mybl2 (Bmyb) maps

[108841]to mouse chromosome 2 and human chromosome 20q13.1. Genomics 35:

[108842]610–612, 1996.

[108843]19181.

[108844]Ando, A.; Kikuti, Y. Y.; Shigenari, A.; Kawata, H.; Okamoto, N.;

[108845]Shiina, T.; Chen, L.; Ikemura, T.; Abe, K.; Kimura, M.; Inoko, H.

[108846]: cDNA cloning of the human homologues of the mouse Ke4 and Ke6 genes

[108847]at the centromeric end of the human MHC region. Genomics 35: 600–602,

[108848]1996.

[108849]19182.

[108850]Aziz, N.; Maxwell, M. M.; Brenner, B. M.: Coordinate regulation

[108851]of 11–beta–HSD and Ke6 gene in cpk mouse: implications for steroid

[108852]metabolic defect in PKD. Am. J. Physiol. 267: F791–F797,
1994.

[108853]19183.

[108854]Aziz, N.; Maxwell, M. M.; St.–Jacques, B.; Brenner, B. M.:
Downregulation

[108855]of Ke 6, a novel gene encoded within the major histocom-
patibility

[108856]complex, in murine polycystic kidney disease. Molec. Cell.
Biol. 13:

[108857]1847–1853, 1993.

[108858]19184.

[108859]Kikuti, Y. Y.; Tamiya, G.; Ando, A.; Chen, L.; Kimura, M.;
Ferreira,

[108860]E.; Tsuji, K.; Trowsdale, J.; Inoko, H.: Physical mapping
220 kb

[108861]centromeric of the human MHC and DNA sequence analy-
sis of the 43–kb

[108862]segment including the RING1, HKE6, and HKE4 genes. Ge-
nomics 42:

[108863]422–435, 1997.

[108864]19185.

[108865]Macalma, T.; Otte, J.; Hensler, M. E.; Bockholt, S. M.; Louis,
[108866]H. A.; Kalff-Suske, M.; Grzeschik, K.-H.; von der Ahe, D.;
Beckerle,
[108867]M. C.: Molecular characterization of human zyxin. J. Biol.
Chem. 271:
[108868]31470-31478, 1996.
[108869]19186.
[108870]Zumbrunn, J.; Trueb, B.: Assignment of the ZYX gene for
the LIM
[108871]protein zyxin to human chromosome bands 7q34-q35 by
in situ hybridization. Cytogenet.
[108872]Cell Genet. 81: 283-284, 1998.
[108873]19187.
[108874]Jacobsen, L.; Madsen, P.; Moestrup, S. K.; Lund, A. H.;
Tommerup,
[108875]N.; Nykjaer, A.; Sottrup-Jensen, L.; Gliemann, J.; Petersen,
C. M.
[108876]: Molecular characterization of a novel human hybrid-type
receptor
[108877]that binds the alpha(2)-macroglobulin receptor-associ-
ated protein. J.

[108878]Biol. Chem. 271: 31379–31383, 1996.

[108879]19188.

[108880]Senechal, K.; Halpern, J.; Sawyers, C. L.: The CRKL adaptor protein

[108881]transforms fibroblasts and functions in transformation by the BCR-ABL

[108882]oncogene. J. Biol. Chem. 38: 23255–23261, 1996.

[108883]19189.

[108884]ten Hoeve, J.; Morris, C.; Heisterkamp, N.; Groffen, J.: Iso-
lation

[108885]and chromosomal localization of CRKL, a human crk-like gene. Oncogene 8:

[108886]2469–2474, 1993.

[108887]19190.

[108888]Cheung, A. H.; Stewart, R. J.; Marsden, P. A.: Endothelial Tie2/Tek

[108889]ligands angiopoietin-1 (ANGPT1) and angiopoietin-2 (ANGPT2): regional

[108890]localization of the human genes to 8q22.3–q23 and 8p23. Genomics 48:

[108891]389–391, 1998.

[108892]19191.

[108893]Davis, S.; Aldrich, T. H.; Jones, P. F.; Acheson, A.; Compton,

[108894]D. L.; Jain, V.; Ryan, T. E.; Bruno, J.; Radziejewski, C.;
Maisonpierre,

[108895]P. C.; Yancopoulos, G. D.: Isolation of angiopoietin-1, a
ligand

[108896]for the TIE2 receptor, by secretion-trap expression
cloning. Cell 87:

[108897]1161-1169, 1996.

[108898]19192.

[108899]Folkman, J.; D'Amore, P. A.: Blood vessel formation: what
is its

[108900]molecular basis? Cell 87: 1153-1155, 1996.

[108901]19193.

[108902]Marziliano, N.; Crovella, S.; Audero, E.; Pecile, V.; Bus-
solino,

[108903]F.; Amoroso, A.; Garagna, S.: Genetic mapping of the
mouse homologue

[108904]of the human angiopoietin-1 gene (Agpt) to mouse chro-
mosome 9E2 by

[108905]in situ hybridization. Cytogenet. Cell Genet. 87: 199–200, 1999.

[108906]19194.

[108907]Sato, T. N.; Tozawa, Y.; Deutsch, U.; Wolburg–Buchholz, K.; Fujiwara,

[108908]Y.; Gendron–Maguire, M.; Gridley, T.; Wolburg, H.; Risau, W.; Qin,

[108909]Y.: Distinct roles of the receptor tyrosine kinases Tie–1 and Tie–2

[108910]in blood vessel formation. Nature 376: 70–73, 1995.

[108911]19195.

[108912]Suri, C.; Jones, P. F.; Patan, S.; Bartunkova, S.; Maison–pierre,

[108913]P. C.; Davis, S.; Sato, T. N.; Yancopoulos, G. D.: Requisite role

[108914]of angiopoietin–1, a ligand for the TIE2 receptor, during embryonic

[108915]angiogenesis. Cell 87: 1171–1180, 1996.

[108916]19196.

[108917]Suri, C.; McClain, J.; Thurston, G.; McDonald, D. M.; Zhou, H.;

[108918]Oldmixon, E. H.; Sato, T. N.; Yancopoulos, G. D.: Increased vascularization

[108919]in mice overexpressing angiopoietin-1. Science 282: 468-471, 1998.

[108920]19197.

[108921]Valenzuela, D. M.; Griffiths, J. A.; Rojas, J.; Aldrich, T. H.;

[108922]Jones, P. F.; Zhou, H.; McClain, J.; Copeland, N. G.; Gilbert, D.

[108923]J.; Jenkins, N. A.; Huang, T.; Papadopoulos, N.; Maison-pierre, P.

[108924]C.; Davis, S.; Yancopoulos, G. D.: Angiopoietins 3 and 4: diverging

[108925]gene counterparts in mice and humans. Proc. Nat. Acad. Sci. 96:

[108926]1904-1909, 1999.

[108927]19198.

[108928]Chestkov, A. V.; Baka, I. D.; Kost, M. V.; Georgiev, G. P.; Buchman,

[108929]V. L.: The d4 gene family in the human genome. Genomics 36: 174-177,

[108930]1996.

[108931]19199.

[108932]Gabig, T. G.; Crean, C. D.; Klenk, A.; Long, H.; Copeland, N. G.;

[108933]Gilbert, D. J.; Jenkins, N. A.; Quincey, D.; Parente, F.; Lespinasse,

[108934]F.; Carle, G. F.; Gaudray, P.; and 13 others: Expression and chromosomal

[108935]localization of the Requiem gene. Mammalian Genome 9: 660–665, 1998.

[108936]19200.

[108937]Gabig, T. G.; Mantel, P. L.; Rosli, R.; Crean, C. D.: Requiem:

[108938]a novel zinc finger gene essential for apoptosis in myeloid cells. J.

[108939]Biol. Chem. 269: 29515–29519, 1994.

[108940]19201.

[108941]Smith, J. E., Jr.; Bollekens, J. A.; Inghirami, G.; Takeshita,

[108942]K.: Cloning and mapping of the MEIS1 gene, the human homolog of a

[108943]murine leukemogenic gene. Genomics 43: 99–103, 1997.

[108944]19202.

[108945]Nakanishi-Matsui, M.; Zheng, Y.-W.; Sulciner, D. J.; Weiss, E.

[108946]J.; Ludeman, M. J.; Coughlin, S. R.: PAR3 is a cofactor for PAR4

[108947]activation by thrombin. Nature 404: 609-613, 2000.

[108948]19203.

[108949]Sambrano, G. R.; Weiss, E. J.; Zheng, Y.-W.; Huang, W.; Coughlin,

[108950]S. R.: Role of thrombin signalling in platelets in haemostasis and

[108951]thrombosis. Nature 413: 74-78, 2001.

[108952]19204.

[108953]Crosnier, C.; Driancourt, C.; Raynaud, N.; Dhorne-Pollet, S.; Pollet,

[108954]N.; Bernard, O.; Hadchouel, M.; Meunier-Rotival, M.: Mutations in

[108955]JAGGED1 gene are predominantly sporadic in Alagille syndrome. Gastroenterology 116:

[108956]1141-1148, 1999.

[108957]19205.

[108958]Liese, J.; Kloos, S.; Jendrossek, V.; Petropoulou, T.; Win-

tergerst,

[108959]U.; Notheis, G.; Gahr, M.; Belohradsky, B. H.: Long-term follow-up

[108960]and outcome of 39 patients with chronic granulomatous disease. J.

[108961]Pediat. 137: 687-693, 2000.

[108962]19206.

[108963]Parkos, C. A.; Allen, R. A.; Cochrane, C. G.; Jesaitis, A. J.

[108964]: Purified cytochrome b from human granulocyte plasma membrane is

[108965]comprised of two polypeptides with relative molecular weights of 91,000

[108966]and 22,000. J. Clin. Invest. 80: 732-742, 1987.

[108967]19207.

[108968]Parkos, C. A.; Dinauer, M. C.; Walker, L. E.; Allen, R. A.; Jesaitis,

[108969]A. J.; Orkin, S. H.: Primary structure and unique expression of the

[108970]22-kilodalton light chain of human neutrophil cytochrome b. Proc.

[108971]Nat. Acad. Sci. 85: 3319-3323, 1988.

[108972]19208.

[108973]Schapiro, B. L.; Newburger, P. E.; Klempner, M. S.; Dinauer, M.

[108974]C.: Chronic granulomatous disease presenting in a 69-year-old man. New

[108975]Eng. J. Med. 325: 1786-1790, 1991.

[108976]19209.

[108977]Stasia, M. J.; Bordigoni, P.; Martel, C.; Morel, F.: A novel

[108978]and unusual case of chronic granulomatous disease in a child with

[108979]a homozygous 36-bp deletion in the CYBA gene (A22-0) leading to the

[108980]activation of a cryptic splice site in intron 4. Hum. Genet. 110:

[108981]444-450, 2002.

[108982]19210.

[108983]Ushio-Fukai, M.; Zafari, A. M.; Fukui, T.; Ishizaka, N.; Griendling,

[108984]K.: p22-phox is a critical component of the superoxide-generating

[108985]NADH/NADPH oxidase system and regulates angiotensin

II-induced hypertrophy

[108986]in vascular smooth muscle cells. J. Biol. Chem. 271:
23317-23321,

[108987]1996.

[108988]19211.

[108989]Weening, R. S.; Corbeel, L.; de Boer, M.; Lutter, R.; van
Zwieten,

[108990]R.; Hamers, M. N.; Roos, D.: Cytochrome b deficiency in an
autosomal

[108991]form of chronic granulomatous disease: a third form of
chronic granulomatous

[108992]disease recognized by monocyte hybridization. J. Clin. In-
vest. 75:

[108993]915-920, 1985.

[108994]19212.

[108995]Yamada, M.; Ariga, T.; Kawamura, N.; Ohtsu, M.; Imajoh-
Ohmi, S.;

[108996]Ohshika, E.; Tatsuzawa, O.; Kobayashi, K.; Sakiyama, Y.:
Genetic

[108997]studies of three Japanese patients with
p22-phox-deficient chronic

[108998]granulomatous disease: detection of a possible common

mutant CYBA

[108999]allele in Japan and a genotype–phenotype correlation in these patients. Brit.

[109000]J. Haemat. 108: 511–517, 2000.

[109001]19213.

[109002]Francke, U.; Hsieh, C.–L.; Foellmer, B. E.; Lomax, K. J.; Malech,

[109003]H. L.; Leto, T. L.: Genes for two autosomal recessive forms of chronic

[109004]granulomatous disease assigned to 1q25 (NCF2) and 7q11.23 (NCF1). Am.

[109005]J. Hum. Genet. 47: 483–492, 1990.

[109006]19214.

[109007]Chasse, J. F.; Paul, V.; Escanez, R.; Kamoun, P.; London, J.:

[109008]Human cystathionine beta–synthase: gene organization and expression

[109009]of different 5–prime alternative splicing. Mammalian Genome 8: 917–921,

[109010]1997.

[109011]19215.

[109012]Paulweber, B.; Wiebusch, H.; Miesenboeck, G.; Funke, H.;

Assmann,

[109013]G.; Hoelzl, B.; Sippl, M. J.; Friedl, W.; Patsch, J. R.; Sand-
hofer,

[109014]F.: Molecular basis of lipoprotein lipase deficiency in two
Austrian

[109015]families with type I hyperlipoproteinemia. Atherosclerosis
86: 239-250,

[109016]1991.

[109017]19216.

[109018]Andersson, S.; Berman, D. M.; Jenkins, E. P.; Russell, D. W.:

[109019]Deletion of steroid 5-alpha-reductase 2 gene in male
pseudohermaphroditism. Nature 354:

[109020]159-161, 1991.

[109021]19217.

[109022]Boudon, C.; Lobaccaro, J. M.; Lumbroso, S.; Ogur, G.; Ocal,
G.;

[109023]Belon, C.; Sultan, C.: A new deletion of the
5-alpha-reductase type

[109024]2 gene in a Turkish family with 5-alpha-reductase defi-
ciency. Clin.

[109025]Endocr. 43: 183-188, 1995.

[109026]19218.

[109027]Cai, L.-Q.; Zhu, Y.-S.; Katz, M. D.; Herrera, C.; Baez, J.;
DeFillo-Ricart,

[109028]M.; Shackleton, C. H. L.; Imperato-McGinley, J.:
5-Alpha-reductase-2

[109029]gene mutations in the Dominican Republic. J. Clin. Endocr.
Metab. 81:

[109030]1730-1735, 1996.

[109031]19219.

[109032]Can, S.; Zhu, Y.-S.; Cai, L.-Q.; Ling, Q.; Katz, M. D.; Ak-
gun,

[109033]S.; Shackleton, C. H. L.; Imperato-McGinley, J.: The identi-
fication

[109034]of 5-alpha-reductase-2 and 17-beta-hydroxysteroid de-
hydrogenase-3

[109035]gene defects in male pseudohermaphrodites from a Turk-
ish kindred. J.

[109036]Clin. Endocr. Metab. 83: 560-569, 1998.

[109037]19220.

[109038]Chavez, B.; Valdez, E.; Vilchis, F.: Uniparental disomy in
steroid

[109039]5-alpha-reductase 2 deficiency. J. Clin. Endocr. Metab. 85:
3147-3150,

[109040]2000.

[109041]19221.

[109042]De Vaal, O. M.: Genital intersexuality in three brothers,
connected

[109043]with consanguineous marriages in the three previous gen-
erations. Acta

[109044]Paediat. 44: 35-39, 1955.

[109045]19222.

[109046]Forti, G.; Falchetti, A.; Santoro, S.; Davis, D. L.; Wilson, J.

[109047]D.; Russell, D. W.: Steroid 5 alpha-reductase 2 deficiency:
virilization

[109048]in early infancy may be due to partial function of mutant
enzyme. Clin.

[109049]Endocr. 44: 477–482, 1996.

[109050]19223.

[109051]Greene, S. A.; Symes, E.; Brook, C. G. D.:

5-Alpha-reductase deficiency

[109052]causing male pseudohermaphroditism. Arch. Dis. Child.

53: 751–753,

[109053]1978.

[109054]19224.

[109055]Hochberg, Z.; Chayen, R.; Reiss, N.; Falik, Z.; Makler, A.;

Munichor,

[109056]M.; Farkas, A.; Goldfarb, H.; Ohana, N.; Hiort, O.: Clinical,

biochemical,

[109057]and genetic findings in a large pedigree of male and fe-

male patients

[109058]with 5-alpha-reductase 2 deficiency. J. Clin. Endocr.

Metab. 81:

[109059]2821–2827, 1996.

[109060]19225.

[109061]Imperato-McGinley, J.; Gautier, T.: Inherited

5-alpha-reductase

[109062]deficiency in man. Trends Genet. 2: 130–133, 1986.

[109063]19226.

[109064]Imperato-McGinley, J.; Guerrero, L.; Gautier, T.; German, J.
L.;

[109065]Peterson, R. E.: Steroid-5-alpha-reductase deficiency in
man. An

[109066]inherited form of male pseudohermaphroditism.In:
Bergsma, D.: Genetic

[109067]Forms of Hypogonadism. New York: National Foundation-
March of Dimes

[109068](pub.) 1975. Pp. 91-103.

[109069]19227.

[109070]Imperato-McGinley, J.; Guerrero, L.; Gautier, T.; Peterson,
R.

[109071]E.: Steroid 5-alpha-reductase deficiency in man: an inher-
ited form

[109072]of male pseudohermaphroditism. Science 186:
1213-1215, 1974.

[109073]19228.

[109074]Imperato-McGinley, J.; Miller, M.; Wilson, J. D.; Peterson,
R.

[109075]E.; Shackleton, C.; Gajdusek, D. C.: A cluster of male
pseudohermaphrodites

[109076]with 5- α -reductase deficiency in Papua New Guinea.

Clin. Endocr. 34:

[109077]293-298, 1991.

[109078]19229.

[109079]Imperato-McGinley, J.; Peterson, R. E.; Gautier, T.; Sturla, E.

[109080]; Androgens and the evolution of male-gender identity among male pseudohermaphrodites

[109081]with 5- α -reductase deficiency. New Eng. J. Med. 300: 1233-1237,

[109082]1979.

[109083]19230.

[109084]Chrzanowska, B. L.; Nitzan, M.; Phillips, L. S.; Schulman, J.

[109085]D.: Homocysteic acid: an examination of its possible growth hormone-like

[109086]activity. Metabolism 28: 80-84, 1979.

[109087]19231.

[109088]Cochran, F. B.; Sweetman, L.; Schmidt, K.; Barsh, G.; Kraus, J.;

[109089]Packman, S.: Pyridoxine-unresponsive homocystinuria with an unusual

[109090]clinical course. Am. J. Med. Genet. 35: 519–522, 1990.

[109091]19232.

[109092]Collins, J. E.; Brenton, D. P.: Pancreatitis and homocystinuria. J.

[109093]Inherit. Metab. Dis. 13: 232–233, 1990.

[109094]19233.

[109095]Di Minno, G.; Davi, G.; Margaglione, M.; Cirillo, F.; Grandone,

[109096]E.; Ciabattini, G.; Catalano, I.; Strisciuglio, P.; Andria, G.; Patrono,

[109097]C.; Mancini, M.: Abnormally high thromboxane biosynthesis in homozygous

[109098]homocystinuria: evidence for platelet involvement and probucol-sensitive

[109099]mechanism. J. Clin. Invest. 92: 1400–1406, 1993.

[109100]19234.

[109101]Falcon, C. R.; Cattaneo, M.; Panzeri, D.; Martinelli, I.; Mannucci,

[109102]P. M.: High prevalence of hyperhomocyst(e)inemia in patients with

[109103]juvenile venous thrombosis. Arteriosclerosis Thromb. 14:

- 1080–1083,
[109104]1994.
- [109105]19235.
- [109106]Field, C. M. B.; Carson, N. A. J.; Cusworth, D. C.; Dent, C.
E.;
- [109107]Neill, D. W.: Homocystinuria, a new disorder of
metabolism. (Abstract) Proc.
- [109108]10th Int. Cong. Pediat., Lisbon 274–275, 1962.
- [109109]19236.
- [109110]Fowler, B.; Kraus, J.; Packman, S.; Rosenberg, L. E.: Homo-
cystinuria:
- [109111]evidence for three distinct classes of cystathionine beta-
synthetase
- [109112]mutants in cultured fibroblasts. J. Clin. Invest. 61:
645–653, 1978.
- [109113]19237.
- [109114]Franco, R. F.; Elion, J.; Lavinha, J.; Krishnamoorthy, R.;
Tavella,
- [109115]M. H.; Zago, M. A.: Heterogeneous ethnic distribution of
the 844ins68
- [109116]in the cystathionine beta-synthase gene. Hum. Hered. 48:

338–342,

[109117]1998.

[109118]19238.

[109119]Franken, D. G.; Boers, G. H. J.; Blom, H. J.; Cruysberg, J. R.

[109120]M.; Trijbels, F. J. M.; Hamel, B. C. J.: Prevalence of familial
mild

[109121]hyperhomocysteinemia. Atherosclerosis 125: 71–80,
1996.

[109122]19239.

[109123]Frimpter, G. W.: Homocystinuria: vitamin B6 dependent or
not?

[109124](Editorial) Ann. Intern. Med. 71: 209–211, 1969.

[109125]19240.

[109126]Fryer, R. H.; Wilson, B. D.; Gubler, D. B.; Fitzgerald, L. A.;

[109127]Rodgers, G. M.: Homocysteine, a risk factor for premature
vascular

[109128]disease and thrombosis, induces tissue factor activity in
endothelial

[109129]cells. Arteriosclerosis Thromb. 13: 1327–1333, 1993.

[109130]19241.

- [109131]Gallagher, P. M.; Naughten, E.; Hanson, N. Q.; Schwichtenberg,
berg,
- [109132]K.; Bignell, M.; Yuan, M.; Ward, P.; Yap, S.; Whitehead, A.
S.; Tsai,
M. Y.: Characterization of mutations in the cystathionine
beta-synthase
gene in Irish patients with homocystinuria. Molec. Genet.
Metab. 65:
298–302, 1998.
- [109135]19242.
- [109136]19242.
- [109137]Gallagher, P. M.; Ward, P.; Tan, S.; Naughten, E.; Kraus, J.
P.;
- [109138]Sellar, G. C.; McConnell, D. J.; Graham, I.; Whitehead, A. S.:
High
frequency (71%) of cystathionine beta-synthase mutation
G307S in Irish
homocystinuria patients. Hum. Mutat. 6: 177–180, 1995.
- [109140]19243.
- [109141]19243.
- [109142]Gaustadnes, M.; Ingerslev, J.; Rutiger, N.: Prevalence of
congenital
homocystinuria in Denmark. (Letter) New Eng. J. Med. 340:
1513 only,

[109144]1999.

[109145]19244.

[109146]Gaustadnes, M.; Rudiger, N.; Rasmussen, K.; Ingerslev, J.:
Intermediate

[109147]and severe hyperhomocysteinemia with thrombosis: a
study of genetic

[109148]determinants. Thromb. Haemost. 83: 554-558, 2000.

[109149]19245.

[109150]Gaustadnes, M.; Wilcken, B.; Oliveriusova, J.; McGill, J.;
Fletcher,

[109151]J.; Kraus, J. P.; Wilcken, D. E.: The molecular basis of cys-
tathionine

[109152]beta-synthase deficiency in Australian patients: genotype-
phenotype

[109153]correlations and response to treatment. Hum. Mutat. 20:
117-126,

[109154]2002.

[109155]19246.

[109156]Gerding, H.: Ocular complications and a new surgical ap-
proach

[109157]to lens dislocation in homocystinuria due to cystathion-

ine-beta-synthetase

[109158]deficiency. Europ. J. Pediat. 157 (suppl. 2): S94-S101,
1998.

[109159]19247.

[109160]Gerritsen, T.; Vaughn, J. G.; Waisman, H. A.: The identifi-
cation

[109161]of homocystine in the urine. Biochem. Biophys. Res. Com-
mun. 9: 493-496,

[109162]1962.

[109163]19248.

[109164]Goldstein, J. L.; Campbell, B. K.; Gartler, S. M.: Homo-
cystinuria:

[109165]heterozygote detection using phytohemagglutinin-stimu-
lated lymphocytes. J.

[109166]Clin. Invest. 52: 218, 1973.

[109167]19249.

[109168]Goldstein, J. L.; Campbell, B. K.; Gartler, S. M.: Cystathion-
ine

[109169]synthetase activity in human lymphocytes: induction by
phytohemagglutinin. J.

[109170]Clin. Invest. 51: 1034-1037, 1972.

[109171]19250.

[109172]Gu, Z.; Ramesh, V.; Kozich, V.; Korson, M. S.; Kraus, J. P.;
Shih,

[109173]V. E.: Identification of a molecular genetic defect in homo-
cystinuria

[109174]due to cystathionine beta-synthase deficiency. (Abstract)
Am. J.

[109175]Hum. Genet. 49: 406 only, 1991.

[109176]19251.

[109177]Guttormsen, A. B.; Ueland, P. M.; Kruger, W. D.; Kim, C. E.;
Ose,

[109178]L.; Folling, I.; Refsum, H.: Disposition of homocysteine in
subjects

[109179]heterozygous for homocystinuria due to cystathionine
beta-synthase

[109180]deficiency: relationship between genotype and phenotype.
Am. J. Med.

[109181]Genet. 100: 204-213, 2001.

[109182]19252.

[109183]Harker, L. A.; Ross, R.; Slichter, S. J.; Scott, C. R.: Homo-
cystine-induced

- [109184]arteriosclerosis: the role of endothelial cell injury and platelet
- [109185]response in its genesis. J. Clin. Invest. 58: 731–741, 1976.
- [109186]19253.
- [109187]Harker, L. A.; Slichter, S. J.; Scott, C. R.; Ross, R.: Homocystinuria:
- [109188]vascular injury and arterial thrombosis. New Eng. J. Med. 291: 537–543,
- [109189]1974.
- [109190]19254.
- [109191]Harrison, D. A.; Mullaney, P. B.; Mesfer, S. A.; Awad, A. H.;
- [109192]Dhindsa, H.: Management of ophthalmic complications of homocystinuria. Ophthalmology 105:
- [109193]1886–1890, 1998.
- [109194]19255.
- [109195]Hooft, C.; Carton, D.; Samyn, W.: Pyridoxine treatment in homocystinuria.
- [109196](Letter) Lancet I: 1384 only, 1967.
- [109197]19256.
- [109198]Hu, F. L.; Gu, Z.; Kozich, V.; Kraus, J. P.; Ramesh, V.; Shih,

[109199]V. E.: Molecular basis of cystathionine beta-synthase deficiency

[109200]in pyridoxine responsive and nonresponsive homocystinuria. Hum. Molec.

[109201]Genet. 2: 1857-1860, 1993.

[109202]19257.

[109203]Janosik, M.; Oliveriusova, J.; Janosikova, B.; Sokolova, J.; Kraus,

[109204]E.; Kraus, J. P.; Kozich, V.: Impaired heme binding and aggregation

[109205]of mutant cystathionine beta-synthase subunits in homocystinuria. Am.

[109206]J. Hum. Genet. 68: 1506-1513, 2001.

[109207]19258.

[109208]Kaeser, A. C.; Rodnight, R.; Ellis, B. A.: Psychiatric and biochemical

[109209]aspects of a case of homocystinuria. J. Neurol. Neurosurg. Psychiat. 32:

[109210]88-93, 1969.

[109211]19259.

[109212]Kang, S.-S.; Wong, P. W. K.; Cook, H. Y.; Norusis, M.;

Messer,

[109213]J. V.: Protein-bound homocyst(e)ine: a possible risk factor
for coronary

[109214]artery disease. J. Clin. Invest. 77: 1482–1486, 1986.

[109215]19260.

[109216]Kery, V.; Poneleit, L.; Meyer, J. D.; Manning, M. C.; Kraus, J.

[109217]P.: Binding of pyridoxal 5-prime-phosphate to the heme
protein human

[109218]cystathionine beta-synthase. Biochemistry 38:
2716–2724, 1999.

[109219]19261.

[109220]Kim, C. E.; Gallagher, P. M.; Guttormsen, A. B.; Refsum, H.;
Ueland,

[109221]P. M.; Ose, L.; Folling, I.; Whitehead, A. S.; Tsai, M. Y.;
Kruger,

[109222]W. D.: Functional modeling of vitamin responsiveness in
yeast: a

[109223]common pyridoxine-responsive cystathionine beta-
synthase mutation

[109224]in homocystinuria. Hum. Molec. Genet. 6: 2213–2221,
1997.

[109225]19262.

[109226]Giralt, M.; Park, E. A.; Gurney, A. L.; Liu, J.; Hakimi, P.;
Hanson,
[109227]R. W.: Identification of a thyroid hormone response ele-
ment in the
[109228]phosphoenolpyruvate carboxykinase (GTP) gene: evidence
for synergistic
[109229]interaction between thyroid hormone and cAMP cis-
regulatory elements. J.
[109230]Biol. Chem. 266: 21991-21996, 1991.
[109231]19263.
[109232]Pilz, A. J.; Willer, E.; Povey, S.; Abbott, C. M.: The genes
coding
[109233]for phosphoenolpyruvate carboxykinase-1 (PCK1) and
neuronal nicotinic
[109234]acetylcholine receptor alpha-4 subunit (CHRNA4) map to
human chromosome
[109235]20, extending the known region of homology with mouse
chromosome 2. Ann.
[109236]Hum. Genet. 56: 289-293, 1992.
[109237]19264.
[109238]Olswang, Y.; Cohen, H.; Papo, O.; Cassuto, H.; Croniger, C.
M.;

[109239]Hakimi, P.; Tilghman, S. M.; Hanson, R. W.; Reshef, L.: A mutation
[109240]in the peroxisome proliferator-activated receptor gamma-binding site
[109241]in the gene for the cytosolic form of phosphoenolpyruvate carboxykinase
[109242]reduces adipose tissue size and fat content in mice. Proc. Nat. Acad.
[109243]Sci. 99: 625-630, 2002.
[109244]19265.
[109245]Stoffel, M.; Xiang, K.; Espinosa, R., III; Cox, N. J.; Le Beau,
[109246]M. M.; Bell, G. I.: cDNA sequence and localization of polymorphic
[109247]human cytosolic phosphoenolpyruvate carboxykinase gene (PCK1) to chromosome
[109248]20, band q13.31: PCK1 is not tightly linked to maturity-onset diabetes
[109249]of the young. Hum. Molec. Genet. 2: 1-4, 1993.
[109250]19266.
[109251]Ting, C.-N.; Burgess, D. L.; Chamberlain, J. S.; Keith, T. P.;
[109252]Falls, K.; Meisler, M. H.: Phosphoenolpyruvate carboxyki-

nase (GTP):

[109253] characterization of the human PCK1 gene and localization
distal to

[109254] MODY on chromosome 20. Genomics 16: 698–706, 1993.

[109255] 19267.

[109256] Vidnes, J.; Sovik, O.: Gluconeogenesis in infancy and
childhood.

[109257] III. Deficiency of the extramitochondrial form of hepatic
phosphoenolpyruvate

[109258] carboxykinase in a case of persistent neonatal hypogly-
caemia. Acta

[109259] Paediat. Scand. 65: 301–312, 1976.

[109260] 19268.

[109261] Yu, H.; Thun, R.; Chandrasekharappa, S.; Trent, J. M.;
Zhang, J.;

[109262] Meisler, M. H.: Human PCK1 encoding phosphoenolpyru-
vate carboxykinase

[109263] is located on chromosome 20q13.2. Genomics 15:
219–221, 1993.

[109264] 19269.

[109265] Abarbanel, J. M.; Bashan, N.; Potashnik, R.; Osimani, A.;

Moses,

[109266]S. W.; Herishanu, Y.: Adult muscle phosphorylase 'b' kinase deficiency. *Neurology* 36:

[109267]560–562, 1986.

[109268]19270.

[109269]Pepe, G.; Chimienti, G.; Resta, F.; Di Perna, V.; Tarricone, C.;

[109270]Lovecchio, M.; Colacicco, A. M.; Capurso, A.: A new Italian case

[109271]of lipoprotein lipase deficiency: a leu365-to-val change resulting

[109272]in loss of enzyme activity. *Biochem. Biophys. Res. Commun.* 199:

[109273]570–576, 1994.

[109274]19271.

[109275]Reymer, P. W. A.; Groenemeyer, B. E.; Gagne, E.; Miao, L.; Appelman,

[109276]E. E. G.; Seidel, J. C.; Kromhout, D.; Bijvoet, S. M.; van de Oever,

[109277]K.; Bruin, T.; Hayden, M. R.; Kastelein, J. J. P.: A frequently occurring

[109278]mutation in the lipoprotein lipase gene (Asn291Ser) con-

tributes to

[109279]the expression of familial combined hyperlipidemia. Hum.

Molec. Genet. 4:

[109280]1543–1549, 1995.

[109281]19272.

[109282]Samuels, M. E.; Forbey, K. C.; Reid, J. E.; Abkevich, V.;

Bulka,

[109283]K.; Wardell, B. R.; Bowen, B. R.; Hopkins, P. N.; Hunt, S. C.;

Ballinger,

[109284]D. G.; Skolnick, M. H.; Wagner, S.: Identification of a com-

mon variant

[109285]in the lipoprotein lipase gene in a large Utah kindred as-

certained

[109286]for coronary heart disease: the -93G/D9N variant predis-

poses to low

[109287]HDL-C/high triglycerides. Clin. Genet. 59: 88–98, 2001.

[109288]19273.

[109289]Schreibman, P. H.; Arons, D. L.; Saudek, C. D.; Arky, R. A.:

[109290]Abnormal lipoprotein lipase in familial exogenous hyper-

triglyceridemia. J.

[109291]Clin. Invest. 52: 2074–2082, 1973.

[109292]19274.

[109293]Sprecher, D. L.; Kobayashi, J.; Rymaszewski, M.; Goldberg, I.

[109294]J.; Harris, B. V.; Bellet, P. S.; Ameis, D.; Yunker, R. L.; Black, D. M.; Stein, E. A.; Schotz, M. C.; Wiginton, D. A.:

Trp64-to-nonsense

[109296]mutation in the lipoprotein lipase gene. J. Lipid Res. 33: 859-866,

[109297]1992.

[109298]19275.

[109299]Sternowsky, H. J.; Gaertner, U.; Stahnkel, N.; Kaukel, E.: Juvenile

[109300]familial hypertriglyceridemia and growth retardation: clinical and

[109301]biochemical observations in three siblings. Europ. J. Pediat. 125:

[109302]59-70, 1977.

[109303]19276.

[109304]Takagi, A.; Ikeda, Y.; Tsutsumi, Z.; Shoji, T.; Yamamoto, A.:

[109305]Molecular studies on primary lipoprotein lipase (LPL) defi-

ciency:

[109306]one base deletion (G-916) in exon 5 of LPL gene causes
no detectable

[109307]LPL protein due to the absence of LPL mRNA transcript. J.
Clin. Invest. 89:

[109308]581-591, 1992.

[109309]19277.

[109310]Weinstock, P. H.; Bisgaier, C. L.; Aalto-Setälä, K.; Radner,
H.;

[109311]Ramakrishnan, R.; Levak-Frank, S.; Essenburg, A. D.;
Zechner, R.;

[109312]Breslow, J. L.: Severe hypertriglyceridemia, reduced high
density

[109313]lipoprotein, and neonatal death in lipoprotein lipase
knockout mice:

[109314]mild hypertriglyceridemia with impaired very low density
lipoprotein

[109315]clearance in heterozygotes. J. Clin. Invest. 96: 2555-2568,
1995.

[109316]19278.

[109317]Wessler, S.; Avioli, L. A.: Familial hyperlipoproteinemia.
J.A.M.A. 207:

[109318]929-937, 1969.

[109319]19279.

[109320]Wilson, D. E.; Edwards, C. Q.; Chan, I.-F.: Phenotypic heterogeneity

[109321]in the extended pedigree of a proband with lipoprotein lipase deficiency. Metabolism 32:

[109322]1107-1114, 1983.

[109323]19280.

[109324]Wilson, D. E.; Emi, M.; Iverius, P.-H.; Hata, A.; Wu, L. L.; Hillas,

[109325]E.; Williams, R. R.; Lalouel, J.-M.: Phenotypic expression of heterozygous

[109326]lipoprotein lipase deficiency in the extended pedigree of a proband

[109327]homozygous for a missense mutation. J. Clin. Invest. 86: 735-750,

[109328]1990.

[109329]19281.

[109330]Wilson, D. E.; Hata, A.; Kwong, L. K.; Lingam, A.; Shuhua, J.;

[109331]Ridinger, D. N.; Yeager, C.; Kaltenborn, K. C.; Iverius,

P.-H.; Lalouel,
[109332]J.-M.: Mutations in exon 3 of the lipoprotein lipase gene
segregating
[109333]in a family with hypertriglyceridemia, pancreatitis, and
non-insulin-dependent
[109334]diabetes. J. Clin. Invest. 92: 203-211, 1993.
[109335]19282.
[109336]Wion, K. L.; Kirchgessner, T. G.; Lusi, A. J.; Schotz, M. C.;
[109337]Lawn, R. M.: Human lipoprotein lipase complementary
DNA sequence. Science 235:
[109338]1638-1641, 1987.
[109339]19283.
[109340]Wittrup, H. H.; Tybjaerg-Hansen, A.; Abildgaard, S.; Steffensen,
[109341]R.; Schnohr, P.; Nordestgaard, B. G.: A common substitution (asn291ser)
[109342]in lipoprotein lipase is associated with increased risk of
ischemic
[109343]heart disease. J. Clin. Invest. 99: 1606-1613, 1997.
[109344]19284.
[109345]Wittrup, H. H.; Tybjaerg-Hansen, A.; Steffensen, R.; Deeb,

S.

- [109346]S.; Brunzell, J. D.; Jensen, G.; Nordestgaard, B. G.: Mutations in
the lipoprotein lipase gene associated with ischemic heart
disease
in men: the Copenhagen City Heart Study. *Arterioscler. Thromb. Vasc. Biol.* 19: 1535–1540, 1999.
- [109349]19285.
- [109351]Wood, S.; Schertzer, M.; Hayden, M.; Ma, Y.: Support for
founder
effect for two lipoprotein lipase (LPL) gene mutations in
French Canadians
by analysis of GT microsatellites flanking the LPL gene. *Hum. Genet.* 91: 312–316, 1993.
- [109355]19286.
- [109356]Burke, V.; Colebatch, J. H.; Anderson, C. M.; Simons, M. J.:
Association
of pancreatic insufficiency and chronic neutropenia in
childhood. *Arch. Dis. Child.* 42: 147–157, 1967.

[109359]19287.

[109360]Ando, T.; Klingberg, W. G.; Ward, A. N.; Rasmussen, K.;
Nyhan,

[109361]W. L.: Isovaleric acidemia presenting with altered
metabolism of

[109362]glycine. *Pediat. Res.* 5: 478-486, 1971.

[109363]19288.

[109364]Ando, T.; Nyhan, W. L.; Bachmann, C.; Rasmussen, K.;
Scott, R.;

[109365]Smith, E. K.: Isovaleric acidemia: identification of iso-
valerylglycine,

[109366]and 3-hydroxyisovalerate in urine of a patient previously
reported

[109367]as having butyric and hexanoic acidemia. *J. Pediat.* 82:
243-248,

[109368]1973.

[109369]19289.

[109370]Budd, M. A.; Tanaka, K. R.; Holmes, L. B.; Efron, M. L.;
Crawford,

[109371]J. D.; Isselbacher, K. J.: Isovaleric acidemia: clinical feature

[109372]of a new genetic defect of leucine metabolism. *New Eng. J.*

Med. 277:

[109373]321-327, 1967.

[109374]19290.

[109375]Cohn, R. M.; Yudkoff, R.; Rothman, R.; Segal, S.: Isovaleric
acidemia:

[109376]Use of glycine therapy in neonates. New Eng. J. Med. 299:
996-999,

[109377]1978.

[109378]19291.

[109379]de Sousa, C.; Chalmers, R. A.; Stacey, T. E.; Tracey, B. M.;
Weaver,

[109380]C. M.; Bradley, D.: The response to L-carnitine and glycine
therapy

[109381]in isovaleric acidaemia. Europ. J. Pediat. 144: 451-456,
1986.

[109382]19292.

[109383]Duran, M.; van Sprang, F. J.; Drewes, J. G.; Bruinvis, L.;
Ketling,

[109384]D.; Wadman, S. K.: Two sisters with isovaleric acidemia,
multiple

[109385]attacks of ketoacidosis and normal development. Europ. J.

Pediat. 131:

[109386]205–211, 1979.

[109387]19293.

[109388]Efron, M. L.: Isovaleric acidemia. Am. J. Dis. Child. 113:
74–76,

[109389]1967.

[109390]19294.

[109391]Kupfer, G. M.; Naf, D.; Suliman, A.; Pulsipher, M.;
D'Andrea, A.

[109392]D.: The Fanconi anaemia proteins, FAA and FAC, interact
to form a

[109393]nuclear complex. Nature Genet. 17: 487–490, 1997.

[109394]19295.

[109395]Kupfer, G. M.; Yamashita, T.; Naf, D.; Suliman, A.; Asano,
S.;

[109396]D'Andrea, A. D.: The Fanconi anemia polypeptide, FAC,
binds to the

[109397]cyclin-dependent kinase, cdc2. Blood 90: 1047–1054,
1997.

[109398]19296.

[109399]Liebetrau, W.; Budde, A.; Savoia, A.; Grummt, F.; Hoehn,

H.:

[109400]p53 activates Fanconi anemia group C gene expression.

Hum. Molec.

[109401]Genet. 6: 277–283, 1997.

[109402]19297.

[109403]Lo Ten Foe, J. R.; Kwee, M. L.; Rooimans, M. A.; Oostra, A.

B.;

[109404]Veermans, A. J. P.; van Weel, M.; Pauli, R. M.; Shahidi, N.

T.; Dokal,

[109405]I.; Roberts, I.; Altay, C.; Gluckman, E.; Gibson, R. A.;

Mathew, C.

[109406]G.; Arwert, F.; Joenje, H.: Somatic mosaicism in Fanconi anemia:

[109407]molecular basis and clinical significance. Europ. J. Hum.

Genet. 5:

[109408]137–148, 1997.

[109409]19298.

[109410]Lo Ten Foe, J. R.; Rooimans, M. A.; Joenje, H.; Arwert, F.:

Novel

[109411]frameshift mutation (1806insA) in exon 14 of the Fanconi anemia C

[109412]gene, FAC. Hum. Mutat. 7: 264–265, 1996.

[109413]19299.

[109414]Murer-Orlando, M.; Llerena, J. C., Jr.; Birjandi, F.; Gibson,

[109415]R. A.; Mathew, C. G.: FACC gene mutations and early pre-natal diagnosis

[109416]of Fanconi's anaemia. (Letter) Lancet 342: 686 only, 1993.

[109417]19300.

[109418]Pace, P.; Johnson, M.; Tan, W. M.; Mosedale, G.; Sng, C.; Hoatlin,

[109419]M.; de Winter, J.; Joenje, H.; Gergely, F.; Patel, K. J.: FANCE:

[109420]the link between Fanconi anaemia complex assembly and activity. EMBO

[109421]J. 21: 3414-3423, 2002.

[109422]19301.

[109423]Pang, Q.; Christianson, T. A.; Keeble, W.; Diaz, J.; Faulkner,

[109424]G. R.; Reifsteck, C.; Olson, S.; Bagby, G. C.: The Fanconi anemia

[109425]complementation group C gene product: structural evidence of multifunctionality. Blood 98:

[109426]1392-1401, 2001.

[109427]19302.

[109428]Pang, Q.; Fagerlie, S.; Christianson, T. A.; Keeble, W.;
Faulkner,

[109429]G.; Diaz, J.; Rathbun, R. K.; Bagby, G. C.: The Fanconi ane-
mia protein

[109430]FANCC binds to and facilitates the activation of STAT1 by
gamma interferon

[109431]and hematopoietic growth factors. Molec. Cell. Biol. 20:
4724-4735,

[109432]2000.

[109433]19303.

[109434]Riley, J.; Butler, R.; Ogilvie, D.; Finniear, R.; Jenner, D.;

[109435]Powell, S.; Anand, R.; Smith, J. C.; Markham, A. F.: A novel,
rapid

[109436]method for the isolation of terminal sequences from yeast
artificial

[109437]chromosome (YAC) clones. Nucleic Acids Res. 18:
2887-2890, 1990.

[109438]19304.

[109439]Roberts, R. G.; Coffey, A. J.; Bobrow, M.; Bentley, D. R.:
Determination

[109440]of the exon structure of the distal portion of the dys-
trophin gene

[109441]by vectorette PCR. Genomics 13: 942–950, 1992.

[109442]19305.

[109443]Savoia, A.; Centra, M.; Ianzano, L.; de Cillis, G. P.; Zelante,

[109444]L.; Buchwald, M.: Characterization of the 5-prime region
of the Fanconi

[109445]anaemia group C (FACC) gene. Hum. Molec. Genet. 4:
1321–1326, 1995.

[109446]19306.

[109447]Segal, G. M.; Magenis, R. E.; Brown, M.; Keeble, W.; Smith,
T.

[109448]D.; Heinrich, M. C.; Bagby, G. C., Jr.: Repression of Fanconi
anemia

[109449]gene (FACC) expression inhibits growth of hematopoietic
progenitor

[109450]cells. J. Clin. Invest. 94: 846–852, 1994.

[109451]19307.

[109452]Strathdee, C. A.; Duncan, A. M. V.; Buchwald, M.: Evidence
for

[109453]at least four Fanconi anaemia genes including FACC on

chromosome 9. Nature

[109454]Genet. 1: 196–198, 1992.

[109455]19308.

[109456]Verlander, P. C.; Kaporis, A.; Liu, Q.; Zhang, Q.; Seligsohn,

[109457]U.; Auerbach, A. D.: Carrier frequency of the IVS4 + 4 A-
to-T mutation

[109458]of the Fanconi anemia gene FAC in the Ashkenazi Jewish
population. Blood 86:

[109459]4034–4038, 1995.

[109460]19309.

[109461]Verlinsky, Y.; Rechitsky, S.; Schoolcraft, W.; Strom, C.;
Kuliev,

[109462]A.: Preimplantation diagnosis for Fanconi anemia com-
bined with HLA

[109463]matching. J.A.M.A. 285: 3130–3133, 2001.

[109464]19310.

[109465]Waisfisz, Q.; Morgan, N. V.; Savino, M.; de Winter, J. P.; van

[109466]Berkel, C. G. M.; Hoatlin, M. E.; Ianzano, L.; Gibson, R. A.;
Arwert,

[109467]F.; Savoia, A.; Mathew, C. G.; Pronk, J. C.; Joenje, H.: Spon-

taneous

[109468]functional correction of homozygous Fanconi anaemia alleles reveals

[109469]novel mechanistic basis for reverse mosaicism. Nature Genet. 22:

[109470]379–383, 1999.

[109471]19311.

[109472]Wevrick, R.; Barker, J. E.; Eppig, J. T.; Nadeau, J. H.; Buchwald,

[109473]M.: Expression of the Fanconi anemia group C gene in the mouse. (Abstract) Am.

[109474]J. Hum. Genet. 51 (suppl.): A137 only, 1992.

[109475]19312.

[109476]Wevrick, R.; Barker, J. E.; Nadeau, J. H.; Szpirer, C.; Buchwald,

[109477]M.: Mapping of the murine and rat Facc genes and assessment of flexed-tail

[109478]as a candidate mouse homolog of Fanconi anemia group C. Mammalian

[109479]Genome 4: 440–444, 1993.

[109480]19313.

[109481]Whitney, M. A.; Jakobs, P.; Kaback, M.; Moses, R. E.;

Grompe,

[109482]M.: The Ashkenazi Jewish Fanconi anemia mutation: incidence among

[109483]patients and carrier frequency in the at-risk population.

Hum. Mutat. 3:

[109484]339–341, 1994.

[109485]19314.

[109486]Whitney, M. A.; Royle, G.; Low, M. J.; Kelly, M. A.; Axthelm,

[109487]M. K.; Reifsteck, C.; Olson, S.; Braun, R. E.; Heinrich, M. C.;

Rathbun,

[109488]R. K.; Bagby, G. C.; Grompe, M.: Germ cell defects and hematopoietic

[109489]hypersensitivity to gamma-interferon in mice with a targeted disruption

[109490]of the Fanconi anemia C gene. Blood 88: 49–58, 1996.

[109491]19315.

[109492]Whitney, M. A.; Saito, H.; Jakobs, P. M.; Gibson, R. A.;

Moses,

[109493]R. E.; Grompe, M.: A common mutation in the FACC gene causes Fanconi

[109494] anaemia in Ashkenazi Jews. Nature Genet. 4: 202–205,
1993.

[109495] 19316.

[109496] Yamashita, T.; Barber, D. L.; Zhu, Y.; Wu, N.; D'Andrea, A.
D.

[109497]: The Fanconi anemia polypeptide FACC is localized to the
cytoplasm. Proc.

[109498] Nat. Acad. Sci. 91: 6712–6716, 1994.

[109499] 19317.

[109500] Yamashita, T.; Wu, N.; Kupfer, G.; Corless, C.; Joenje, H.;
Grompe,

[109501] M.; D'Andrea, A. D.: Clinical variability of Fanconi anemia
(type

[109502] C) results from expression of an amino terminal truncated
Fanconi

[109503] anemia complementation group C polypeptide with partial
activity. Blood 87:

[109504] 4424–4432, 1996.

[109505] 19318.

[109506] Youssoufian, H.: Cytoplasmic localization of FAC is essen-
tial

[109507]for the correction of a prerepair defect in Fanconi anemia group C

[109508]cells. J. Clin. Invest. 97: 2003–2010, 1996.

[109509]19319.

[109510]Yousoufian, H.: Localization of Fanconi anemia C protein to

[109511]the cytoplasm of mammalian cells. Proc. Nat. Acad. Sci. 91: 7975–7979,

[109512]1994.

[109513]19320.

[109514]Yousoufian, H.; Li, Y.; Martin, M. E.; Buchwald, M.: Induction

[109515]of Fanconi anemia cellular phenotype in human 293 cells by overexpression

[109516]of a mutant FAC allele. J. Clin. Invest. 97: 957–962, 1996.

[109517]19321.

[109518]Hodge, S. E.; Anderson, C. E.; Neiswanger, K.; Field, L. L.; Spence,

[109519]M. A.; Sparkes, R. S.; Sparkes, M. C.; Crist, M.; Terasaki, P. I.;

[109520]Rimoin, D. L.; Rotter, J. I.: Close genetic linkage between

diabetes

[109521]mellitus and Kidd blood group. Lancet II: 893–895, 1981.

[109522]19322.

[109523]Marcolongo, P.; Banhegyi, G.; Benedetti, A.; Hinds, C. J.;
Burchell,

[109524]A.: Liver microsomal transport of glucose–6–phosphate,
glucose, and

[109525]phosphate in type 1 glycogen storage diseases. J. Clin.
Endocr. Metab. 83:

[109526]224–229, 1998.

[109527]19323.

[109528]Michels, V. V.; Beaudet, A. L.: Hemorrhagic pancreatitis in
a

[109529]patient with glycogen storage disease type I. Clin. Genet.
17: 220–222,

[109530]1980.

[109531]19324.

[109532]Nakai, K.; Sakamoto, H.: Construction of a novel database
containing

[109533]aberrant splicing mutations of mammalian genes. Gene
141: 171–177,

[109534]1994.

[109535]19325.

[109536]Napole, V. M.; Campbell, W. G., Jr.: Hepatoblastoma in infant

[109537]sister and brother. Cancer 39: 2647–2650, 1977.

[109538]19326.

[109539]Nuoffer, J. M.; Mullis, P. E.; Wiesmann, U. N.: Treatment with

[109540]low-dose diazoxide in two growth-retarded prepubertal girls with glycogen

[109541]storage disease type Ia resulted in catch-up growth. J. Inherit.

[109542]Metab. Dis. 20: 790–798, 1997.

[109543]19327.

[109544]Obara, K.; Saito, T.; Sato, H.; Ogawa, M.; Igarashi, Y.; Yoshinaga,

[109545]K.: Renal histology in two adult patients with type I glycogen storage

[109546]disease. Clin. Nephrol. 39: 59–64, 1993.

[109547]19328.

[109548]Parvari, R.; Lei, K.-J.; Bashan, N.; HersHKovitz, E.; Korman,
[109549]S. H.; Barash, V.; Lerman-Sagie, T.; Mandel, H.; Chou, J. Y.;
Moses,
[109550]S. W.: Glycogen storage disease type 1a in Israel: bio-
chemical, clinical,
[109551]and mutational studies. Am. J. Med. Genet. 72: 286-290,
1997.
[109552]19329.
[109553]Pizzo, C. J.: Type I glycogen storage disease with focal
nodular
[109554]hyperplasia of the liver and vasoconstrictive pulmonary
hypertension. Pediatrics 65:
[109555]341-343, 1980.
[109556]19330.
[109557]Qu, Y.; Abdenur, J. E.; Eng, C. M.; Desnick, R. J.: Molecular
[109558]prenatal diagnosis of glycogen storage disease type Ia.
Prenatal
[109559]Diag. 15: 333-336, 1996.
[109560]19331.
[109561]Reitsma-Bierens, W. C. C.: Renal complications in glyco-
gen storage

[109562]disease type I. *Europ. J. Pediat.* 152 (suppl. 1): S60–S62, 1993.

[109563]19332.

[109564]Rennert, O. M.; Mukhopadhyay, D.: Diazoxide in Von Gierke's disease. *Arch.*

[109565]*Dis. Child.* 43: 358–361, 1968.

[109566]19333.

[109567]Restaino, I.; Kaplan, B. S.; Stanley, C.; Baker, L.: Nephrolithiasis,

[109568]hypocitraturia, and a distal renal tubular acidification defect in

[109569]type 1 glycogen storage disease. *J. Pediat.* 122: 392–396, 1993.

[109570]19334.

[109571]Ryan, I. P.; Havel, R. J.; Laros, R. K., Jr.: Three consecutive

[109572]pregnancies in a patient with glycogen storage disease type IA (von

[109573]Gierke's disease). *Am. J. Obstet. Gynec.* 170: 1687–1691, 1994.

[109574]19335.

[109575]Selby, R.; Starzl, T. E.; Yunis, E.; Todo, S.; Tzakis, A. G.;

[109576]Brown, B. I.; Kendall, R. S.: Liver transplantation for type I
and

[109577]type IV glycogen storage disease. Europ. J. Pediat. 152
(suppl.

[109578]1): S71–S76, 1993.

[109579]19336.

[109580]Senior, B.; Loridan, L.: Liver glycogenoses: metabolism of
intravenously

[109581]administered glycerol. New Eng. J. Med. 279: 958–965,
1968.

[109582]19337.

[109583]Senior, B.; Loridan, L.: Functional differentiation of
glycogenoses

[109584]of the liver with respect to the use of glycerol. New Eng. J.
Med. 279:

[109585]965–970, 1968.

[109586]19338.

[109587]Seydewitz, H. H.; Matern, D.: Molecular genetic analysis of
40

[109588]patients with glycogen storage disease type Ia: 100% mu-
tation detection

[109589]rate and 5 novel mutations. (Abstract) Hum. Mutat. 15:
115–116,

[109590]2000.

[109591]19339.

[109592]Shelly, L. L.; Lei, K.-J.; Pan, C.-J.; Sakata, S. F.; Ruppert,

[109593]S.; Schutz, G.; Chou, J. Y.: Isolation of the gene for murine
glucose-6-phosphatase,

[109594]the enzyme deficient in glycogen storage disease type 1A.
J. Biol.

[109595]Chem. 268: 21482–21485, 1993.

[109596]19340.

[109597]Sidbury, J. B., Jr.: The genetics of the glycogen storage
disease.In:

[109598]Steinberg, A. G.; Bearn, A. G.: Progress in Medical Genet-
ics. New

[109599]York: Grune and Stratton (pub.) 4: 1965. Pp. 32–58.

[109600]19341.

[109601]Smit, G. P. A.: The long-term outcome of patients with
glycogen

[109602]storage disease type Ia. Europ. J. Pediat. 152 (suppl. 1):
S52–S55,

[109603]1993.

[109604]19342.

[109605]Spencer-Peet, J.; Norman, M. E.; Lake, B. D.; McNamara, J.;
Patrick,

[109606]A. D.: Hepatic glycogen storage disease. Clinical and laboratory
findings in 23 cases. Quart. J. Med. 40: 95-114, 1971.

[109607]

[109608]19343.

[109609]Stamm, W. E.; Webb, D. I.: Partial deficiency of hepatic
glucose-6-phosphatase

[109610]in an adult patient. Arch. Intern. Med. 135: 1107-1109,
1975.

[109611]19344.

[109612]Stevenson, R. E.; Ben-Menachem, Y.; Dudrick, S.; Howell,
R. R.

[109613]: Hepatocellular carcinoma in type 1 glycogen storage disease. Proc.

[109614]Greenwood Genet. Center 3: 39-46, 1984.

[109615]19345.

[109616]Talente, G. M.; Coleman, R. A.; Alter, C.; Baker, L.; Brown,
B.

[109617]I.; Cannon, R. A.; Chen, Y.-T.; Crigler, J. F., Jr.; Ferreira, P.;
 [109618]Haworth, J. C.; Herman, G. E.; Issenman, R. M.; Keating, J.
 P.; Linde,
 [109619]R.; Roe, T. F.; Senior, B.; Wolfsdorf, J. I.: Glycogen storage
 disease
 [109620]in adults. Ann. Intern. Med. 120: 218-226, 1994.
 [109621]19346.
 [109622]Trioche, P.; Francoual, J.; Capel, L.; Odievre, M.; Linden-
 baum,
 [109623]A.; Labrune, P.: Apolipoprotein E polymorphism and
 serum concentrations
 [109624]in patients with glycogen storage disease type Ia. J. In-
 herit. Metab.
 [109625]Dis. 107-112, 2000.
 [109626]19347.
 [109627]Ullrich, K.; Smit, G. P. A.: Clinical aspects of glycogen
 storage
 [109628]disease type I: summary of the discussions. Europ. J. Pe-
 diat. 152
 [109629](suppl. 1): S87-S88, 1993.
 [109630]19348.

[109631]Weinstein, D. A.; Somers, M. J. G.; Wolfsdorf, J. I.: De-
creased

[109632]urinary citrate excretion in type 1a glycogen storage dis-
ease. J.

[109633]Pediat. 138: 378–382, 2001.

[109634]19349.

[109635]Wierzbicki, A. S.; Watts, G. F.; Lynas, J.; Winder, A. F.;
Wray,

[109636]R.: Very low-density lipoprotein apolipoprotein B-100
turnover in

[109637]glycogen storage disease type Ia (von Gierke disease). J.
Inherit.

[109638]Metab. Dis. 24: 527–534, 2001.

[109639]19350.

[109640]Zangeneh, F.; Limbeck, G. A.; Brown, B. I.; Emch, J. R.; Ar-
casoy,

[109641]M. M.; Goldenberg, V. E.; Kelley, V. C.: Hepatorenal
glycogenosis

[109642](type I glycogenosis) and carcinoma of the liver. J. Pediat.
74:

[109643]73–83, 1969.

[109644]19351.

[109645]Maroteaux, P.; Lamy, M.: Hurler's disease, Morquio's disease,

[109646]and related mucopolysaccharidoses. J. Pediat. 67: 312-323, 1965.

[109647]19352.

[109648]Adams, P. C.: Intrafamilial variation in hereditary hemochromatosis. Digest.

[109649]Dis. Sci. 37: 361-363, 1992.

[109650]19353.

[109651]Aguilar-Martinez, P.; Bismuth, M.; Picot, M. C.; Thelcide, C.;

[109652]Pageaux, G.-P.; Blanc, F.; Blanc, P.; Schved, J.-F.; Larrey, D.:

[109653]Variable phenotypic presentation of iron overload in H63D homozygotes:

[109654]are genetic modifiers the cause? Gut 48: 836-842, 2001.

[109655]19354.

[109656]Anand, S.; Schade, R. R.; Bendetti, C.; Kelly, R.; Rabin, B. S.;

[109657]Krause, J.; Starzl, T. E.; Iwatsuki, S. I.; Van Thiel, D. H.: Id-

iopathic

[109658]alpha-hemochromatosis (sic) and 1-antitrypsin deficiency: coexistence

[109659]in a family with progressive liver disease in the proband.

Hepatology 3:

[109660]714-718, 1983.

[109661]19355.

[109662]Bahram, S.; Gilfillan, S.; Kuhn, L. C.; Moret, R.; Schulze, J.

[109663]B.; Lebeau, A.; Schumann, K.: Experimental hemochromatosis due to

[109664]MHC class I HFE deficiency: immune status and iron metabolism. Proc.

[109665]Nat. Acad. Sci. 96: 13312-13317, 1999.

[109666]19356.

[109667]Balcerzak, S. P.; Westerman, M. P.; Lee, R. E.; Doyle, A. P.:

[109668]Idiopathic hemochromatosis: a study of three families. Am. J. Med. 40:

[109669]857-873, 1966.

[109670]19357.

[109671]Barton, J. C.; Acton, R. T.: Inheritance of two HFE muta-

tions

[109672]in African Americans: cases with hemochromatosis phenotypes and estimates

[109673]of hemochromatosis phenotype frequency. Genet. Med. 3: 294–300,

[109674]2001.

[109675]19358.

[109676]Barton, J. C.; McDonnell, S. M.; Adams, P. C.; Brissot, P.; Powell,

[109677]L. W.; Edwards, C. Q.; Cook, J. D.; Kowdley, K. V.; Hemochromatosis

[109678]Management Working Group: Management of hemochromatosis. Ann. Intern.

[109679]Med. 129: 932–939, 1998.

[109680]19359.

[109681]Barton, J. C.; Patton, M. A.; Edwards, C. Q.; Griffen, L. M.; Kushner,

[109682]J. P.; Meeks, R. G.; Leggett, R. W.: Blood lead concentrations in

[109683]hereditary hemochromatosis. J. Lab. Clin. Med. 124: 193–198, 1994.

[109684]19360.

[109685]Barton, J. C.; Rothenberg, B. E.; Bertoli, L. F.; Acton, R. T.

[109686]: Diagnosis of hemochromatosis in family members of probands: a comparison

[109687]of phenotyping and HFE genotyping. Genet. Med. 1: 89–93, 1999.

[109688]19361.

[109689]Barton, J. C.; Sawada–Hirai, R.; Rothenberg, B. E.; Acton, R.

[109690]T.: Two novel missense mutations of the HFE gene (I105T and G93R)

[109691]and identification of the S65C mutation in Alabama hemochromatosis

[109692]probands. Blood Cells Molec. Dis. 25: 146–154, 1999.

[109693]19362.

[109694]Bassett, M. L.; Doran, T. J.; Halliday, J. W.; Bashir, H. V.;

[109695]Powell, L. W.: Idiopathic hemochromatosis: demonstration of homozygous–heterozygous

[109696]mating by HLA typing of families. Hum. Genet. 60: 352–356, 1982.

[109697]19363.

[109698]Beaumont, C.; Simon, M.; Fauchet, R.; Hespel, J.–P.; Bris–sot,

[109699]P.; Genetet, B.; Bourel, M.: Serum ferritin as a possible
marker
[109700]of the hemochromatosis allele. New Eng. J. Med. 301:
169–174, 1979.
[109701]19364.
[109702]Beaumont, C.; Simon, M.; Smith, P. M.; Worwood, M.: Hep-
atic and
[109703]serum ferritin concentrations in patients with idiopathic
hemochromatosis. Gastroenterology 79:
[109704]877–883, 1980.
[109705]19365.
[109706]Beckman, L. E.; Saha, N.; Spitsyn, V.; Van Landeghem, G.;
Beckman,
[109707]L.: Ethnic differences in the HFE codon 282 (Cys/Tyr)
polymorphism. Hum.
[109708]Hered. 47: 263–267, 1997.
[109709]19366.
[109710]Beutler, E.: Targeted disruption of the HFE gene. Proc. Nat.
[109711]Acad. Sci. 95: 2033–2034, 1998.
[109712]19367.

[109713]Beutler, E.: The significance of the 187G (H63D) mutation
in

[109714]hemochromatosis. (Letter) Am. J. Hum. Genet. 61:
762–764, 1997.

[109715]19368.

[109716]Beutler, E.; Felitti, V. J.; Koziol, J. A.; Ho, N. J.; Gelbart,

[109717]T.: Penetrance of 845G–A (C282Y) HFE hereditary
haemochromatosis

[109718]mutation in the USA. Lancet 359: 211–218, 2002.

[109719]19369.

[109720]Beutler, E.; Gelbart, T.: HLA–H mutations in the Ashkenazi
Jewish

[109721]population. Blood Cells Molec. Dis. 23: 95–98, 1997.

[109722]19370.

[109723]Beutler, E.; Gelbart, T.; West, C.; Lee, P.; Adams, M.; Black–
stone,

[109724]R.; Pockros, P.; Kosty, M.; Venditti, C. P.; Phatak, P. D.;
Seese,

[109725]N. K.; Chorney, K. A.; Ten Elshof, A. E.; Gerhard, G. S.;
Chorney,

[109726]M.: Mutation analysis in hereditary hemochromatosis.

Blood Cells

[109727]Molec. Dis. 22: 187-194, 1996.

[109728]19371.

[109729]Beutler, E.; West, C.; Gelbart, T.: HLA-H and associated proteins

[109730]in patients with hemochromatosis. Molec. Med. 3: 397-402, 1997.

[109731]19372.

[109732]Bodmer, J. G.; Parham, P.; Albert, E. D.; Marsh, S. G. E.: Putting

[109733]a hold on 'HLA-H.' (Letter) Nature Genet. 15: 234-235, 1997.

[109734]19373.

[109735]Borecki, I. B.; Lathrop, G. M.; Bonney, G. E.; Yaouanq, J.; Rao,

[109736]D. C.: Combined segregation and linkage analysis of genetic hemochromatosis

[109737]using affection status, serum iron, and HLA. Am. J. Hum. Genet. 47:

[109738]542-550, 1990.

[109739]19374.

[109740]Borecki, I. B.; Rao, D. C.; Yaouanq, J.; Lalouel, J. M.: Per-
cent

[109741]transferrin saturation in segregating hemochromatosis.
Am. J. Med.

[109742]Genet. 36: 301-305, 1990.

[109743]19375.

[109744]Borecki, I. B.; Rao, D. C.; Yaouanq, J.; Lalouel, J. M.: Segre-
gation

[109745]of genetic hemochromatosis indexed by latent capacity of
transferrin. Am.

[109746]J. Hum. Genet. 45: 465-470, 1989.

[109747]19376.

[109748]Boretto, J.; Jouanolle, A.-M.; Yaouanq, J.; El Kahloun, A.;
Mauvieux,

[109749]V.; Blayau, M.; Perichon, M.; Le Treut, A.; Clayton, J.; Borot,
N.;

[109750]Le Gall, J.-Y.; Pontarotti, P.; David, V.: Anonymous mark-
ers located

[109751]on chromosome 6 in the HLA-A class I region: allelic dis-
tribution

[109752]in genetic haemochromatosis. Hum. Genet. 89: 33-36,
1992.

[109753]19377.

[109754]Maroteaux, P.; Leveque, B.; Marie, J.; Lamy, M.: Une nouvelle

[109755]dysostose avec elimination urinaire de chondroïtine-sulfate B. Presse

[109756]Med. 71: 1849-1852, 1963.

[109757]19378.

[109758]McGovern, M. M.; Mandell, N.; Haskins, M.; Desnick, R. J.: Animal

[109759]model studies of allelism: characterization of arylsulfatase B mutations

[109760]in homoallelic and heteroallelic (genetic compound) homozygotes with

[109761]feline mucopolysaccharidosis VI. Genetics 110: 733-749, 1985.

[109762]19379.

[109763]Migeon, B. R.; Sprenkle, J. A.; Liebaers, I.; Scott, J. F.; Neufeld,

[109764]E. F.: X-linked Hunter syndrome: the heterozygous phenotype in cell

[109765]culture. Am. J. Hum. Genet. 29: 448-454, 1977.

[109766]19380.

[109767]Neufeld, E. F.; Muenzer, J.: The mucopolysaccharidoses. In:
Scriver,

[109768]C. R.; Beaudet, A. L.; Sly, W. S.; Valle, D. (eds.): The
Metabolic

[109769]and Molecular Bases of Inherited Disease New York: Mc-
Graw-Hill

[109770](7th ed.): 1995. Pp. 2465-2494.

[109771]19381.

[109772]Paterson, D. E.; Rad, M.; Harper, G.; Weston, H. J.; Mattin-
gley,

[109773]J.: Maroteaux-Lamy syndrome, mild form--MPS VIb. Brit.
J. Radiol. 55:

[109774]805-812, 1982.

[109775]19382.

[109776]Peterson, D. I.; Bacchus, H.; Seaich, L.; Kelly, T. E.:
Myelopathy

[109777]associated with Maroteaux-Lamy syndrome. Arch. Neurol.
32: 127-129,

[109778]1975.

[109779]19383.

- [109780]Poser, C. M.; Ojemann, R. G.; Sobel, R. A.: MGH CPC case
44–1983. New
- [109781]Eng. J. Med. 309: 1109–1117, 1983.
- [109782]19384.
- [109783]Quigley, H. A.; Kenyon, K. R.: Ultrastructural and histo-
chemical
- [109784]studies of a newly recognized form of systemic mu-
copolysaccharidosis
- [109785](Maroteaux–Lamy syndrome, mild phenotype). Am. J.
Ophthal. 77: 809–818,
- [109786]1974.
- [109787]19385.
- [109788]Saul, R. A.; Stevenson, R. E.; Taylor, H. A.: Atypical presen-
tation
- [109789]with normal stature in Maroteaux–Lamy syndrome (MPS
VI). Proc. Greenwood
- [109790]Genet. Center 3: 49–52, 1984.
- [109791]19386.
- [109792]Schuchman, E. H.; Jackson, C. E.; Desnick, R. J.: Human
arylsulfatase
- [109793]B: MOPAC cloning, nucleotide sequence of a full-length

cDNA, and regions

[109794] of amino acid identity with arylsulfatases A and C. Genomics 6:

[109795] 149–158, 1990.

[109796] 19387.

[109797] Schwartz, G. P.; Cohen, E. J.: Hydrocephalus in Maroteaux–Lamy

[109798] syndrome. Arch. Ophthalmol. 116: 400 only, 1998.

[109799] 19388.

[109800] Spranger, J. W.; Koch, F.; McKusick, V. A.; Natzschka, J.; Wiedemann,

[109801] H. R.; Zellweger, H.: Mucopolysaccharidosis VI (Maroteaux–Lamy's

[109802] disease). Helv. Paediat. Acta 25: 337–362, 1970.

[109803] 19389.

[109804] Stumpf, D. A.; Austin, J. H.; Crocker, A. C.; LaFrance, M.: Mucopolysaccharidosis

[109805] type VI (Maroteaux–Lamy syndrome). I. Sulfatase B deficiency in tissues. Am.

[109806] J. Dis. Child. 126: 747–755, 1973.

[109807] 19390.

- [109808]Tan, C. T. T.; Schaff, H. V.; Miller, F. A., Jr.; Edwards, W.
[109809]D.; Karnes, P. S.: Valvular heart disease in four patients
with Maroteaux-Lamy
[109810]syndrome. *Circulation* 85: 188-195, 1992.
[109811]19391.
- [109812]Tonnesen, T.; Gregersen, H. N.; Guttler, F.: Normal MPS
excretion,
[109813]but dermatan sulphaturia, combined with a mild
Maroteaux-Lamy phenotype. *J.*
[109814]*Med. Genet.* 28: 499-501, 1991.
[109815]19392.
- [109816]van Dorpe, J.; Moerman, P.; Pecceu, A.; van den Steen, P.;
Fryns,
[109817]J. P.: Non-immune hydrops fetalis caused by beta-
glucuronidase deficiency
[109818](mucopolysaccharidosis VII): study of a family with 3 af-
fected siblings. *Genet.*
[109819]*Counsel.* 7: 105-112, 1996.
[109820]19393.
- [109821]Van Dyke, D. L.; Fluharty, A. L.; Schafer, I. A.; Shapiro, L.

[109822]J.; Kihara, H.; Weiss, L.: Prenatal diagnosis of Maroteaux-Lamy syndrome. Am.

[109823]J. Med. Genet. 8: 235-242, 1981.

[109824]19394.

[109825]Vestermarck, S.; Tonnesen, T.; Schultz Andersen, M.; Guttler, F.

[109826]: Mental retardation in a patient with Maroteaux-Lamy. Clin. Genet. 31:

[109827]114-117, 1987.

[109828]19395.

[109829]Villani, G. R. D.; Balzano, N.; Vitale, D.; Saviano, M.; Pavone,

[109830]V. Di Natale, P.: Maroteaux-Lamy syndrome: five novel mutations and

[109831]their structural localization. Biochim. Biophys. Acta 1453: 185-192,

[109832]1999.

[109833]19396.

[109834]Vine, D. T.; McGovern, M. M.; Haskins, M. E.; Desnick, R. J.:

[109835]Feline mucopolysaccharidosis. VI. Purification and charac-

terization

[109836] of the residual arylsulfatase B activity. Am. J. Hum. Genet.

33:

[109837] 916–927, 1981.

[109838] 19397.

[109839] Weller, P. F.; Austen, K. F.: Human eosinophil arylsulfatase

[109840] B: structure and activity of the purified tetrameric lysosomal hydrolase. J.

[109841] Clin. Invest. 71: 114–123, 1983.

[109842] 19398.

[109843] Wicker, G.; Prill, V.; Brooks, D.; Gibson, G.; Hopwood, J.;
von

[109844] Figura, K.; Peters, C.: Mucopolysaccharidosis VI
(Maroteaux–Lamy

[109845] syndrome): an intermediate clinical phenotype caused by
substitution

[109846] of valine for glycine at position 137 of arylsulfatase B. J.
Biol.

[109847] Chem. 266: 21386–21391, 1991.

[109848] 19399.

[109849] Wilson, C. S.; Mankin, H. T.; Pluth, J. R.: Aortic stenosis

and

[109850] mucopolysaccharidosis. Ann. Intern. Med. 92: 496–498,
1980.

[109851] 19400.

[109852] Yogalingam, G.; Hopwood, J. J.; Crawley, A.; Anson, D. S.:
Mild

[109853] feline mucopolysaccharidosis type VI: identification of an
N-acetylgalactosamine-4-sulfatase

[109854] mutation causing instability and increased specific activ-
ity. J.

[109855] Biol. Chem. 273: 13421–13429, 1998.

[109856] 19401.

[109857] Yogalingam, G.; Litjens, T.; Bielicki, J.; Crawley, A. C.;
Muller,

[109858] V.; Anson, D. S.; Hopwood, J. J.: Feline mucopolysacchari-
dosis type

[109859] VI. J. Biol. Chem. 271: 27259–27265, 1996.

[109860] 19402.

[109861] Yoshida, M.; Noguchi, J.; Ikadaï, H.; Takahashi, M.; Na-
gase, S.

[109862] Arylsulfatase B-deficient mucopolysaccharidosis in rats.

J. Clin.

[109863]Invest. 91: 1099–1104, 1993.

[109864]19403.

[109865]Yoshida, M.; Tachibana, M.; Kobayashi, E.; Ikadai, H.; Kuni-
nieda,

[109866]T.: The locus responsible for mucopolysaccharidosis VI
(Maroteaux–Lamy

[109867]syndrome) is located on rat chromosome 2. Genomics 20:
145–146,

[109868]1994.

[109869]19404.

[109870]Young, R.; Kleinman, G.; Ojemann, R. G.; Kolodny, E.;
Davis, K.;

[109871]Halperin, J.; Zalneraitis, E.; DeLong, G. R.: Compressive
myelopathy

[109872]in Maroteaux–Lamy syndrome: clinical and pathological
findings. Ann.

[109873]Neurol. 8: 336–340, 1980.

[109874]19405.

[109875]Bradley, W. G.; Price, D. L.; Watanabe, C. K.: Familial cen-
tronuclear

[109876]myopathy. J. Neurol. Neurosurg. Psychiat. 33: 687–693,
1970.

[109877]19406.

[109878]Touchman, J. W.; Anikster, Y.; Dietrich, N. L.; Braden
Maduro,

[109879]V. V.; McDowell, G.; Shotelersuk, V.; Bouffard, G. G.; Beck-
strom–Sternberg,

[109880]S. M.; Gahl, W. A.; Green, E. D.: The genomic region en-
compassing

[109881]the nephropathic cystinosis gene (CTNS): complete se-
quencing of a

[109882]200-kb segment and discovery of a novel gene within the
common cystinosis–causing

[109883]deletion. Genome Res. 10: 165–173, 2000.

[109884]19407.

[109885]Shapiro, M. B.; Senapathy, P.: RNA splice junctions of dif-
ferent

[109886]classes of eukaryotes: sequence statistics and functional
implications

[109887]in gene expression. Nucleic Acids Res. 15: 7155–7174,
1987.

[109888]19408.

[109889]Applegarth, D. A.; Toone, J. R.: Nonketotic hyperglycemia (glycine
encephalopathy): laboratory diagnosis. *Molec. Genet. Metab.* 74:
139–146, 2001.

[109892]19409.

[109893]Kure, S.; Kojima, K.; Kudo, T.; Kanno, K.; Aoki, Y.; Suzuki, Y.;

[109894]Shinka, T.; Sakata, Y.; Narisawa, K.; Matsubara, Y.: Chromosomal
localization, structure, single-nucleotide polymorphisms, and expression
of the human H-protein gene of the glycine cleavage system (GCSH),
a candidate gene for nonketotic hyperglycinemia. *J. Hum. Genet.* 46:
378–384, 2001.

[109899]19410.

[109900]Sakata, Y.; Owada, Y.; Sato, K.; Kojima, K.; Hisanaga, K.; Shinka,

[109901]T.; Suzuki, Y.; Aoki, Y.; Satoh, J.; Kondo, H.; Matsubara, Y.; Kure,

[109902]S.: Structure and expression of the glycine cleavage system in rat

[109903]central nervous system. Molec. Brain Res. 94: 119–130, 2001.

[109904]19411.

[109905]Toone, J. R.; Applegarth, D. A.; Coulter–Mackie, M. B.; James,

[109906]E. R.: Recurrent mutations in P– and T–proteins of the glycine cleavage

[109907]complex and a novel T–protein mutation (N145I): a strategy for the

[109908]molecular investigation of patients with nonketotic hyperglycinemia

[109909](NKH). Molec. Genet. Metab. 72: 322–325, 2001.

[109910]19412.

[109911]Bamforth, J. S.; Hughes, I. A.; Lazarus, J. H.; Weaver, C. M.;

[109912]Harper, P. S.: Congenital hypothyroidism, spiky hair, and cleft palate. J.

[109913]Med. Genet. 26: 49–60, 1989.

[109914]19413.

[109915]Jobard, F.; Lefevre, C.; Karaduman, A.; Blanchet–Bardon,

C.; Emre,

[109916]S.; Weissenbach, J.; Ozguc, M.; Lathrop, M.; Prud'homme,
J.-F.; Fischer,

[109917]J.: Lipoxxygenase-3 (ALOXE3) and 12(R)-lipoxxygenase
(ALOX12B) are

[109918]mutated in non-bullous congenital ichthyosiform erythro-
derma (NCIE)

[109919]linked to chromosome 17p13.1. Hum. Molec. Genet. 11:
107-113, 2002.

[109920]19414.

[109921]Dinarello, C. A.; Wolff, S. M.; Goldfinger, S. E.; Dale, D. C.;

[109922]Alling, D. W.: Colchicine therapy for familial Mediterranean
fever:

[109923]a double-blind trial. New Eng. J. Med. 291: 934-937,
1974.

[109924]19415.

[109925]Akaboshi, S.; Yano, T.; Miyawaki, S.; Ohno, K.; Takeshita,
K.:

[109926]A C57BL/KsJ mouse model of Niemann-Pick disease (spm)
belongs to the

[109927]same complementation group as the major childhood type
of Niemann-Pick

[109928]disease type C. Hum. Genet. 99: 350–353, 1997.

[109929]19416.

[109930]Levy, E. N.; Shen, Y.; Kupelian, A.; Kruglyak, L.; Aksentijevich,

[109931]I.; Pras, E.; Balow, J. E., Jr.; Linzer, B.; Chen, X.; Shelton, D.

[109932]A.; Gumucio, D.; Pras, M.; Shohat, M.; Rotter, J. I.; Fischel-Ghodsian,

[109933]N.; Richards, R. I.; Kastner, D. L.: Linkage disequilibrium mapping

[109934]places the gene causing familial Mediterranean fever close to D16S246. Am.

[109935]J. Hum. Genet. 58: 523–534, 1996.

[109936]19417.

[109937]Majeed, H. A.; Barakat, M.: Familial Mediterranean fever (recurrent

[109938]hereditary polyserositis) in children: analysis of 88 cases. Europ.

[109939]J. Pediat. 148: 636–641, 1989.

[109940]19418.

[109941]Varon, R.; Vissinga, C.; Platzer, M.; Cerosaletti, K. M.; Chrzanowska,

[109942]K. H.; Saar, K.; Beckmann, G.; Seemanova, E.; Cooper, P.
R.; Nowak,

[109943]N. J.; Stumm, M.; Weemaes, C. M. R.; Gatti, R. A.; Wilson,
R. K.;

[109944]Digweed, M.; Rosenthal, A.; Sperling, K.; Concannon, P.;
Reis, A.

[109945]Nibrin, a novel DNA double-strand break repair protein,
is mutated

[109946]in Nijmegen breakage syndrome. Cell 93: 467-476, 1998.

[109947]19419.

[109948]Becker, J. A.; Vlach, J.; Raben, N.; Nagaraju, K.; Adams, E.
M.;

[109949]Hermans, M. M.; Reuser, A. J. J.; Brooks, S. S.; Tifft, C. J.;
Hirschhorn,

[109950]R.; Huie, M. L.; Nicolino, M.; Plotz, P. H.: The African origin
of

[109951]the common mutation in African American patients with
glycogen-storage

[109952]disease type II. (Letter) Am. J. Hum. Genet. 62: 991-994,
1998.

[109953]19420.

[109954]Boerkoel, C. F.; Exelbert, R.; Nicastrì, C.; Nichols, R. C.;

Miller,

[109955]F. W.; Plotz, P. H.; Raben, N.: Leaky splicing mutation in the acid

[109956]maltase gene is associated with delayed onset of glycogenosis type

[109957]II. Am. J. Hum. Genet. 56: 887–897, 1995.

[109958]19421.

[109959]Byrne, E.; Dennett, X.; Crotty, B.; Trounce, I.; Sands, J. M.;

[109960]Hawkins, R.; Hammond, J.; Anderson, S.; Haan, E. A.; Pol-lard, A.:

[109961]Dominantly inherited cardioskeletal myopathy with lyso-somal glycogen

[109962]storage and normal acid maltase levels. Brain 109: 523–536, 1986.

[109963]19422.

[109964]Friduss, S. R.; Sadoff, W. I.; Hern, A. E.; Fivenson, D. P.:

[109965]Fatal pyoderma gangrenosum in association with C7 defi-ciency. J.

[109966]Am. Acad. Derm. 27: 356–359, 1992.

[109967]19423.

[109968]Lee, T. J.; Utsinger, P. D.; Snyderman, R.; Yount, W. J.;

Sparling,

[109969]P. F.: Familial deficiency of the seventh component of complement

[109970]associated with recurrent bacteremic infections due to *Neisseria*. J.

[109971]*Infect. Dis.* 138: 359–368, 1978.

[109972]19424.

[109973]Nakamura, S.; Ooue, O.; Abe, K.: Genetic polymorphism of the

[109974]seventh component of complement in a Japanese population. *Hum. Genet.* 66:

[109975]279–281, 1984.

[109976]19425.

[109977]Nishimukai, H.; Tamaki, Y.: Genetic polymorphism of the seventh

[109978]component of complement: a new variant. *Vox Sang.* 51: 60–62, 1986.

[109979]19426.

[109980]Nishizaka, H.; Horiuchi, T.; Zhu, Z. B.; Fukumori, Y.; Volanakis,

[109981]J. E.: Genetic bases of human complement C7 deficiency. J.

Immun. 157:

[109982]4239–4243, 1996.

[109983]19427.

[109984]Nurnberger, W.; Pietsch, H.; Seger, R.; Bufon, T.; Wahn, V.:

[109985]Familial deficiency of the seventh component of complement associated

[109986]with recurrent meningococcal infections. Europ. J. Pediat. 148:

[109987]758–760, 1989.

[109988]19428.

[109989]Petersen, B. H.; Lee, T. J.; Snyderman, R.; Brooks, G. F.:
Neisseria

[109990]meningitidis and Neisseria gonorrhoeae bacteremia associated with

[109991]C6, C7, or C8 deficiency. Ann. Intern. Med. 90: 917–920, 1979.

[109992]19429.

[109993]Rittner, C.; Opferkuch, W.; Wellek, B.; Grosse-Wilde, H.;
Wernet,

[109994]P.: Lack of linkage between gene(s) controlling the synthesis of

[109995]the seventh component of complement and the HLA re-
gion on chromosome

[109996]no. 6 in man. Hum. Genet. 34: 137-142, 1976.

[109997]19430.

[109998]Schlesinger, M.; Nave, Z.; Levy, Y.; Slater, P. E.; Fishelson,

[109999]Z.: Prevalence of hereditary properdin, C7 and C8 defi-
ciencies in

[110000]patients with meningococcal infections. Clin. Exp. Immun.
81: 423-427,

[110001]1990.

[110002]19431.

[110003]Washio, K.; Tokunaga, K.; Omoto, K.; Misawa, S.: Human
C7 polymorphism:

[110004]classification and association analysis with C6. Jpn. J.
Hum. Genet. 31:

[110005]345-352, 1986.

[110006]19432.

[110007]Wurzner, R.; Hobart, M. J.; Orren, A.; Tokunaga, K.; Nitze,
R.;

[110008]Gotze, O.; Lachmann, P. J.: A novel protein polymorphism
of human

[110009]complement C7 detected by a monoclonal antibody. Immunogenetics 35:

[110010]398–402, 1992.

[110011]19433.

[110012]Wurzner, R.; Nitze, R.; Gotze, O.: C7*9, a new frequent C7 allele

[110013]detected by an allotype-specific monoclonal antibody. Complement

[110014]Inflamm. 7: 290–297, 1990.

[110015]19434.

[110016]Zimran, A.; Rudensky, B.; Kramer, M. R.; Tedesco, F.; Ehrenfeld,

[110017]M.; Raz, R.; Greif, Z.; Gelber, M.; Lishner, M.; Golan, E.; Hershko,

[110018]C.: Hereditary complement deficiency in survivors of meningococcal

[110019]disease: high prevalence of C7/C8 deficiency in Sephardic (Moroccan)

[110020]Jews. Quart. J. Med. 63: 349–358, 1987.

[110021]19435.

[110022]Hadano, S.; Hand, C. K.; Osuga, H.; Yanagisawa, Y.;

Otomo, A.;

[110023]Devon, R. S.; Miyamoto, N.; Showguchi-Miyata, J.; Okada, Y.; Singaraja,

[110024]R.; Figlewicz, D. A.; Kwiatkowski, T.; and 9 others: A gene encoding

[110025]a putative GTPase regulator is mutated in familial amyotrophic lateral

[110026]sclerosis 2. Nature Genet. 29: 166-173, 2001. Note: Erratum: Nature

[110027]Genet. 29: 352 only, 2001.

[110028]19436.

[110029]Colten, H. R.; Alper, C. A.; Rosen, F. S.: Current concepts in

[110030]immunology: genetics and biosynthesis of complement proteins. New

[110031]Eng. J. Med. 304: 653-656, 1981.

[110032]19437.

[110033]Howlett, N. G.; Taniguchi, T.; Olson, S.; Cox, B.; Waisfisz, Q.;

[110034]de Die-Smulders, C.; Persky, N.; Grompe, M.; Joenje, H.; Pals, G.;

[110035]Ikeda, H.; Fox, E. A.; D'Andrea, A. D.: Biallelic inactivation

of

[110036]BRCA2 in Fanconi anemia. Science 297: 606–609, 2002.

[110037]19438.

[110038]Mazoyer, S.; Dunning, A. M.; Serova, O.; Dearden, J.;
Puget, N.;

[110039]Healey, C. S.; Gayther, S. A.; Mangion, J.; Stratton, M. R.;
Lynch,

[110040]H. T.; Goldgar, D. E.; Ponder, B. A.; Lenoir, G. M.: A poly-
morphic

[110041]stop codon in BRCA2. Nature Genet. 14: 253–254, 1996.

[110042]19439.

[110043]Lowry, R. B.; Applegarth, D. A.; Toone, J. R.; MacDonald,
E.;

[110044]Thunem, N. Y.: An update on the frequency of mu-
copolysaccharide syndromes

[110045]in British Columbia. Hum. Genet. 85: 389–390, 1990.

[110046]19440.

[110047]ten Hove, T.; Corbaz, A.; Amitai, H.; Aloni, S.; Belzer, I.;
Graber,

[110048]P.; Drillenburger, P.; van Deventer, S. J.; Chvatchko, Y.; Te
Velde,

[110049]A. A.: Blockade of endogenous IL-18 ameliorates TNBS-induced colitis
[110050]by decreasing local TNF-alpha production in mice. Gastroenterology 121:
[110051]1372-1379, 2001.
[110052]19441.
[110053]Roux, A.-F.; Rommens, J.; McDowell, C.; Anson-
Cartwright, L.; Bell,
[110054]S.; Schappert, K.; Fishman, G. A.; Musarella, M.: Identification
[110055]of a gene from Xp21 with similarity to the tctex-1 gene of
the murine
[110056]t complex. Hum. Molec. Genet. 3: 257-263, 1994.
[110057]19442.
[110058]Ishikawa, K.; Nagase, T.; Suyama, M.; Miyajima, N.;
Tanaka, A.;
[110059]Kotani, H.; Nomura, N.; Ohara, O.: Prediction of the coding
sequences
[110060]of unidentified human genes. X. The complete sequences
of 100 new
[110061]cDNA clones from brain which can code for large proteins
in vitro. DNA

[110062]Res. 5: 169–176, 1998.

[110063]19443.

[110064]Kipreos, E. T.; Lander, L. E.; Wing, J. P.; He, W. W.; Hedge-
cock,

[110065]E. M.: *cul-1* is required for cell cycle exit in *C. elegans* and
identifies

[110066]a novel gene family. *Cell* 85: 829–839, 1996.

[110067]19444.

[110068]Rasooly, R. S.: Personal Communication. Baltimore, Md.
9/29/1998.

[110069]19445.

[110070]Wang, P. J.; McCarrey, J. R.; Yang, F.; Page, D. C.: An abundance

[110071]of X-linked genes expressed in spermatogonia. Nature
Genet. 27:

[110072]422-426, 2001.

[110073]19446.

[110074]Herold, A.; Suyama, M.; Rodrigues, J. P.; Braun, I. C.; Kuntay,

[110075]U.; Carmo-Fonseca, M.; Bork, P.; Izaurralde, E.: TAP (NXF1)
belongs

[110076]to a multigene family of putative RNA export factors with
a conserved

[110077]modular architecture. Molec. Cell. Biol. 20: 8996-9008,
2000.

[110078]19447.

[110079]Yang, J.; Bogerd, H. P.; Wang, P. J.; Page, D. C.; Cullen, B.
R.

[110080]: Two closely related human nuclear export factors utilize
entirely

[110081]distinct export pathways. Molec. Cell 8: 397–406, 2001.

[110082]19448.

[110083]Ikeda, M.; Ishida, O.; Hinoi, T.; Kishida, S.; Kikuchi, A.:

Identification

[110084]and characterization of a novel protein interacting with

Ral-binding

[110085]protein 1, a putative effector protein of Ral. J. Biol. Chem.

273:

[110086]814–821, 1998.

[110087]19449.

[110088]Koshiba, S.; Kigawa, T.; Iwahara, J.; Kikuchi, A.; Yokoyama,

S.

[110089]: Solution structure of the Eps15 homology domain of a

human POB1

[110090](partner of RalBP1). FEBS Lett. 442: 138–142, 1999.

[110091]19450.

[110092]Crackower, M. A.; Sarao, R.; Oudit, G. Y.; Yagil, C.; Kozier-

adzki,

[110093]J.; Scanga, S. E.; Oliveira-dos-Santos, A. J.; da Costa, J.;

Zhang,

[110094]L.; Pei, Y.; Scholey, J.; Ferrario, C. M.; Manoukian, A. S.;

Chappell,

[110095]M. C.; Backx, P. H.; Yagil, Y.; Penninger, J. M.: An-
giotensin-converting

[110096]enzyme 2 is an essential regulator of heart function. Na-
ture 417:

[110097]822-828, 2002.

[110098]19451.

[110099]Scanlan, M. J.; Gordan, J. D.; Williamson, B.; Stockert, E.;
Bander,

[110100]N. H.; Jongeneel, V.; Gure, A. O.; Jager, D.; Jager, E.;
Knuth, A.;

[110101]Chen, Y.-T.; Old, L. J.: Antigens recognized by autologous
antibody

[110102]in patients with renal-cell carcinoma. Int. J. Cancer 83:
456-464,

[110103]1999.

[110104]19452.

[110105]Yan, Z.; Fedorov, S. A.; Mumby, M. C.; Williams, R. S.:
PR48,

[110106]a novel regulatory subunit of protein phosphatase 2A, in-
teracts with

[110107]Cdc6 and modulates DNA replication in human cells.

Molec. Cell. Biol. 20:

[110108]1021–1029, 2000.

[110109]19453.

[110110]Gitzelmann, R.: Hereditary galactokinase deficiency, a
newly recognized

[110111]cause of juvenile cataracts. *Pediat. Res.* 1: 14–23, 1967.

[110112]19454.

[110113]Kalaydjieva, L.; Perez–Lezaun, A.; Angelicheva, D.;
Onengut, S.;

[110114]Dye, D.; Bosshard, N. U.; Jordanova, A.; Savov, A.;
Yanakiev, P.;

[110115]Kremensky, I.; Radeva, B.; Hallmayer, J.; Markov, A.; Ned-
kova, V.;

[110116]Tournev, I.; Aneva, L.; Gitzelmann, R.: A founder mutation
in the

[110117]GK1 gene is responsible for galactokinase deficiency in
Roma (Gypsies). *Am.*

[110118]*J. Hum. Genet.* 65: 1299–1307, 1999.

[110119]19455.

[110120]Kim, Y. J.; Rosenberg, L. E.: On the mechanism of pyridox-
ine

- [110121] responsive homocystinuria. II. Properties of normal and mutant cystathionine
- [110122] beta-synthase from cultured fibroblasts. Proc. Nat. Acad. Sci. 71:
- [110123] 4821–4825, 1974.
- [110124] 19456.
- [110125] Kluijtmans, L. A. J.; Boers, G. H. J.; Kraus, J. P.; van den Heuvel,
- [110126] L. P. W. J.; Cruysberg, J. R. M.; Trijbels, F. J. M.; Blom, H. J.
- [110127]: The molecular basis of cystathionine beta-synthase deficiency in
- [110128] Dutch patients with homocystinuria: effect of CBS genotype on biochemical
- [110129] and clinical phenotype and on response to treatment. Am. J. Hum.
- [110130] Genet. 65: 59–67, 1999.
- [110131] 19457.
- [110132] Kluijtmans, L. A. J.; Boers, G. H. J.; Stevens, E. M. B.; Re-
- [110133] nier, W. O.; Kraus, J. P.; Trijbels, F. J. M.; van den Heuvel, L. P. W.

[110134]J.; Blom, H. J.: Defective cystathionine beta-synthase regulation
[110135]by S-adenosylmethionine in a partially pyridoxine responsive homocystinuria
[110136]patient. J. Clin. Invest. 98: 285-289, 1996.
[110137]19458.
[110138]Komrower, G. M.: Dietary treatment of homocystinuria.
Am. J.
[110139]Dis. Child. 113: 98-100, 1967.
[110140]19459.
[110141]Kozich, V.; de Franchis, R.; Kraus, J. P.: Molecular defect in
[110142]a patient with pyridoxine-responsive homocystinuria.
Hum. Molec.
[110143]Genet. 2: 815-816, 1993.
[110144]19460.
[110145]Kozich, V.; Kraus, E.; de Franchis, R.; Fowler, B.; Boers, G.
[110146]H. J.; Graham, I.; Kraus, J. P.: Hyperhomocysteinemia in
premature
[110147]arterial disease: examination of cystathionine beta-synthase alleles

[110148]at the molecular level. Hum. Molec. Genet. 4: 623–629,
1995.

[110149]19461.

[110150]Kozich, V.; Kraus, J. P.: Screening for mutations by ex-
pressing

[110151]patient cDNA segments in E. coli: homocystinuria due to
cystathionine

[110152]beta-synthase deficiency. Hum. Mutat. 1: 113–123, 1992.

[110153]19462.

[110154]Kraus, J.; Packman, S.; Fowler, B.; Rosenberg, L. E.: Purifi-
cation

[110155]and properties of cystathionine beta-synthase from hu-
man liver: evidence

[110156]for identical subunits. J. Biol. Chem. 253: 6523–6528,
1978.

[110157]19463.

[110158]Kluijtmans, L. A. J.; van den Heuvel, L. P. W. J.; Boers, G. H.

[110159]J.; Frosst, P.; Stevens, E. M. B.; van Oost, B. A.; den Heijer,
M.;

[110160]Trijbels, F. J. M.; Rozen, R.; Blom, H. J.: Molecular genetic
analysis

[110161]in mild hyperhomocysteinemia: a common mutation in the methylenetetrahydrofolate

[110162]reductase gene is a genetic risk factor for cardiovascular disease. Am.

[110163]J. Hum. Genet. 58: 35–41, 1996.

[110164]19464.

[110165]Kraus, J. P.: Molecular basis of phenotype expression in homocystinuria. J.

[110166]Inherit. Metab. Dis. 17: 383–390, 1994.

[110167]19465.

[110168]Kraus, J. P.; Janosik, M.; Kozich, V.; Mandell, R.; Shih, V.;

[110169]Sperandeo, M. P.; Sebastio, G.; de Franchis, R.; Andria, G.; Kluijtmans,

[110170]L. A. J.; Blom, H.; Boers, G. H. J.; Gordon, R. B.; Kamoun, P.; Tsai,

[110171]M. Y.; Kruger, W. D.; Koch, H. G.; Ohura, T.; Gaustadnes, M.: Cystathionine

[110172]beta-synthase mutations in homocystinuria. Hum. Mutat. 13: 362–375,

[110173]1999.

[110174]19466.

[110175]Kraus, J. P.; Oliveriusova, J.; Sokolova, J.; Kraus, E.; Vlcek,

[110176]C.; de Franchis, R.; Maclean, K. N.; Bao, L.; Bukovska, G.;
Patterson,

[110177]D.; Paces, V.; Ansorge, W.; Kozich, V.: The human cystathionine beta-synthase

[110178](CBS) gene: complete sequence, alternative splicing, and
polymorphisms. Genomics 52:

[110179]312–324, 1998.

[110180]19467.

[110181]Kruger, W. D.; Cox, D. R.: A yeast assay for functional detection

[110182]of mutations in the human cystathionine beta-synthase
gene. Hum.

[110183]Molec. Genet. 4: 1155–1161, 1995.

[110184]19468.

[110185]Kruger, W. D.; Cox, D. R.: A yeast system for expression of
human

[110186]cystathionine beta-synthase: structural and functional
conservation

[110187]of the human and yeast genes. Proc. Nat. Acad. Sci. 91:
6614–6618,

[110188]1994.

[110189]19469.

[110190]Kurczynski, T. W.; Muir, W. A.; Fleisher, L. D.; Palomaki, J.

[110191]F.; Gaull, G. E.; Rassin, D. K.; Abramowsky, C.: Maternal
homocystinuria:

[110192]studies of an untreated mother and fetus. Arch. Dis. Child.
55:

[110193]721–723, 1980.

[110194]19470.

[110195]Loughridge, L. W.: Renal abnormalities in the Marfan syn-
drome. Quart.

[110196]J. Med. 112: 531–544, 1959.

[110197]19471.

[110198]Lubec, B.; Fang–Kircher, S.; Lubec, T.; Blom, H. J.; Boers,
G.

[110199]H. J.: Evidence for McKusick's hypothesis of deficient col-
lagen cross–linking

[110200]in patients with homocystinuria. Biochim. Biophys. Acta
1315: 159–162,

[110201]1996.

[110202]19472.

[110203]Maclean, K. N.; Gaustadnes, M.; Oliveriusova, J.; Janosik, M.;

[110204]Kraus, E.; Kozich, V.; Kery, V.; Skovby, F.; Rudiger, N.; Ingerslev,

[110205]J.; Stabler, S. P.; Allen, R. H.; Kraus, J. P.: High homocysteine

[110206]and thrombosis without connective tissue disorders are associated

[110207]with a novel class of cystathionine beta-synthase (CBS) mutations. Hum.

[110208]Mutat. 19: 641-655, 2002.

[110209]19473.

[110210]Malinow, M. R.; Stampfer, M. J.: Role of plasma homocyst(e)ine

[110211]in arterial occlusive disease. (Editorial) Clin. Chem. 40: 857-858,

[110212]1994.

[110213]19474.

[110214]Mandel, H.; Brenner, B.; Berant, M.; Rosenberg, N.; Lanir, N.;

- [110215]Jakobs, C.; Fowler, B.; Seligsohn, U.: Coexistence of hereditary
- [110216]homocystinuria and factor V Leiden: effect on thrombosis. New Eng.
- [110217]J. Med. 334: 763–768, 1996.
- [110218]19475.
- [110219]McCully, K. S.; Ragsdale, B. D.: Production of arteriosclerosis
- [110220]by homocystinuria. Am. J. Path. 61: 1–12, 1970.
- [110221]19476.
- [110222]McKusick, V. A.: Heritable Disorders of Connective Tissue St.
- [110223]Louis: C. V. Mosby (pub.) (3rd ed.): 1966. Pp. 155.
- [110224]19477.
- [110225]Mudd, S. H.: Vascular disease and homocysteine metabolism. (Editorial) New
- [110226]Eng. J. Med. 313: 751–753, 1985.
- [110227]19478.
- [110228]Mudd, S. H.; Edwards, W. A.; Loeb, P. M.; Brown, M. S.; Laster,

[110229]L.: Homocystinuria due to cystathionine synthase deficiency: the

[110230]effect of pyridoxine. J. Clin. Invest. 49: 1762–1773, 1970.

[110231]19479.

[110232]Mudd, S. H.; Finkelstein, J. D.; Irreverre, F.; Laster, L.: Homocystinuria:

[110233]an enzymatic defect. Science 143: 1443–1445, 1964.

[110234]19480.

[110235]Mudd, S. H.; Havlik, R.; Levy, H. L.; McKusick, V. A.; Feinleib,

[110236]M.: A study of cardiovascular risk in heterozygotes for homocystinuria. Am.

[110237]J. Hum. Genet. 33: 883–893, 1981.

[110238]19481.

[110239]Mudd, S. H.; Levy, H. L.: Disorders of transulfuration. In: Stanbury,

[110240]J. B.; Wyngaarden, J. B.; Fredrickson, D. S.: Metabolic Basis of

[110241]Inherited Disease. New York: McGraw–Hill (pub.) (4th ed.): 1978.

[110242]pp. 458–503.

[110243]19482.

[110244]Hsieh, C. L.; Leto, T. L.; Lomax, K. J.; Malech, H. L.;
Francke,

[110245]U.: The genes for two neutrophil cytosol factors that are
deficient

[110246]in autosomal forms of chronic granulomatous disease are
on human chromosome

[110247]10 (47 kD), and chromosome 1 cen-q32 (65 kD).

(Abstract) Cytogenet.

[110248]Cell Genet. 51: 1015-1016, 1989.

[110249]19483.

[110250]Karnovsky, M. L.: Steps toward an understanding of
chronic granulomatous

[110251]disease. (Editorial) New Eng. J. Med. 308: 274-275, 1983.

[110252]19484.

[110253]Segal, A. W.: Variations on the theme of chronic granulo-
matous

[110254]disease. Lancet I: 1378-1383, 1985.

[110255]19485.

[110256]Segal, A. W.: The molecular and cellular pathology of
chronic

[110257]granulomatous disease. Europ. J. Clin. Invest. 18:
433–443, 1988.

[110258]19486.

[110259]Aoshima, M.; Nunoi, H.; Shimazu, M.; Shimizu, S.; Tatsuzawa, O.;

[110260]Kenney, R. T.; Kanegasaki, S.: Two-exon skipping due to a
point mutation

[110261]in p67-phox-deficient chronic granulomatous disease.
Blood 88: 1841–1845,

[110262]1996.

[110263]19487.

[110264]Bonizzato, A.; Russo, M. P.; Donini, M.; Dusi, S.: Identification

[110265]of a double mutation (D160V–K161E) (sic) in the p67phox
gene of a

[110266]chronic granulomatous disease patient. Biochem. Biophys.
Res. Commun. 231:

[110267]861–863, 1997.

[110268]19488.

[110269]de Boer, M.; Hilarius–Stokman, P. M.; Hossle, J.–P.; Verhoeven,

- [110270]A. J.; Graf, N.; Kenney, R. T.; Seger, R.; Roos, D.: Autosomal recessive
- [110271]chronic granulomatous disease with absence of the 67-kD cytosolic
- [110272]NADPH oxidase component: identification of mutation and detection
- [110273]of carriers. Blood 83: 531–536, 1994.
- [110274]19489.
- [110275]Dancis, J.; Hutzler, J.; Woody, N. C.; Cox, R. P.: Multiple enzyme
- [110276]defects in familial hyperlysinemia. Pediat. Res. 10: 686–691, 1976.
- [110277]19490.
- [110278]Fischer, A. Q.; Challa, V. R.; Burton, B. K.; McLean, W. T.: Cerebellar
- [110279]hemorrhage complicating isovaleric acidemia: a case report. Neurology 31:
- [110280]746–748, 1981.
- [110281]19491.
- [110282]Sacksteder, K. A.; Biery, B. J.; Morrell, J. C.; Goodman, B. K.;

[110283]Geisbrecht, B. V.; Cox, R. P.; Gould, S. J.; Geraghty, M. T.:

Identification

[110284]of the alpha-amino adipic semialdehyde synthase gene,
which is defective

[110285]in familial hyperlysinemia. Am. J. Hum. Genet. 66:
1736-1743, 2000.

[110286]19492.

[110287]Gerdes, A.-M.; Gregersen, N.; Ludvigsson, P.; Guttler, F.: A

[110288]Scandinavian case of isovaleric acidemia. J. Inherit.
Metab. Dis. 11:

[110289]219-220, 1988.

[110290]19493.

[110291]Ikeda, Y.; Keese, S. M.; Tanaka, K.: Molecular heterogeneity

[110292]of variant isovaleryl-CoA dehydrogenase from cultured
isovaleric acidemia

[110293]fibroblasts. Proc. Nat. Acad. Sci. 82: 7081-7085, 1985.

[110294]19494.

[110295]Kelleher, J. F., Jr.; Yudkoff, M.; Hutchinson, R.; August, C.

[110296]S.; Cohn, R. M.: The pancytopenia of isovaleric acidemia.
Pediatrics 65:

[110297]1023-1027, 1980.

[110298]19495.

[110299]Naglak, M.; Salvo, R.; Madsen, K.; Dembure, P.; Elsas, L.:

The

[110300]treatment of isovaleric acidemia with glycine supplement.

Pediat.

[110301]Res. 24: 9-13, 1988.

[110302]19496.

[110303]Newman, C. G. H.; Wilson, B. D. R.; Callaghan, P.; Young,

L.:

[110304]Neonatal death associated with isovalericacidaemia.

Lancet II: 439-441,

[110305]1967.

[110306]19497.

[110307]Rhead, W. J.; Tanaka, K.: Demonstration of a specific mi-

tochondrial

[110308]isovaleryl-CoA dehydrogenase deficiency in fibroblasts

from patients

[110309]with isovaleric acidemia. Proc. Nat. Acad. Sci. 77:

580-583, 1980.

[110310]19498.

[110311]Sidbury, J. B., Jr.; Smith, E. K.; Harlan, W.: An inborn error
[110312]of short-chain fatty acid metabolism: the odor-
of-sweaty-feet syndrome. J.
[110313]Pediat. 70: 8-15, 1967.
[110314]19499.
[110315]Tanaka, K.; Budd, M. A.; Efron, M. L.; Isselbacher, K. J.:
Isovaleric
[110316]acidemia: a new genetic defect of leucine metabolism.
Proc. Nat.
[110317]Acad. Sci. 56: 236-242, 1966.
[110318]19500.
[110319]Tanaka, K.; Orr, J.; Isselbacher, K. J.: Identification of b-
hydroxyisovaleric
[110320]acid in the urine of a patient with isovaleric acidemia.
Biochim.
[110321]Biophys. Acta 152: 638-641, 1968.
[110322]19501.
[110323]Vockley, J.; Parimoo, B.; Tanaka, K.: Molecular characteri-
zation
[110324]of four different classes of mutations in the isovaleryl-
CoA dehydrogenase

[110325]gene responsible for isovaleric acidemia. Am. J. Hum. Genet. 49:

[110326]147–157, 1991.

[110327]19502.

[110328]Mohandas, T. K.; Chen, X.-N.; Rowe, L. B.; Birkenmeier, E. H.;

[110329]Fanning, A. S.; Anderson, J. M.; Korenberg, J. R.: Localization of

[110330]the tight junction protein gene TJP1 to human chromosome 15q13, distal

[110331]to the Prader-Willi/Angelman region, and to mouse chromosome 7. Genomics 30:

[110332]594–597, 1995.

[110333]19503.

[110334]Willott, E.; Balda, M. S.; Fanning, A. S.; Jameson, B.; Van Itallie,

[110335]C.; Anderson, J. M.: The tight junction protein ZO-1 is homologous

[110336]to the Drosophila discs-large tumor suppressor protein of septate

[110337]junctions. Proc. Nat. Acad. Sci. 90: 7834–7838, 1993.

[110338]19504.

[110339]Blancar, M. A.; Crossley, P. H.; Peters, K. G.; Steingrimsson, E.;

[110340]Copeland, N. G.; Jenkins, N. A.; Martin, G. R.; Rutter, W. J.:
Meso1,

[110341]a basic helix-loop-helix protein involved in mammalian
presomitic

[110342]mesoderm development. Proc. Nat. Acad. Sci. 92:
5870-5874, 1995.

[110343]19505.

[110344]Burgess, R.; Rawls, A.; Brown, D.; Bradley, A.; Olson, E. N.:

[110345]Requirement of the paraxis gene for somite formation and
musculoskeletal

[110346]patterning. Nature 384: 570-573, 1996.

[110347]19506.

[110348]Hidai, H.; Quertermous, E. E.; Espinosa, R., III; Le Beau, M.
M.;

[110349]Quertermous, T.: Genomic organization and chromosomal
localization

[110350]of the gene TCF15 encoding the early mesodermal basic
helix-loop-helix

[110351]factor bHLH-EC2. Genomics 30: 598–601, 1995.

[110352]19507.

[110353]Quertermous, E. E.; Hidai, H.; Blonar, M. A.; Quertermous, T.:

[110354]Cloning and characterization of a basic helix–loop–helix protein expressed

[110355]in early mesoderm and the developing somites. Proc. Nat. Acad. Sci. 91:

[110356]7066–7070, 1994.

[110357]19508.

[110358]Quertermous, T.: Personal Communication. Nashville, Tenn. 1/22/1996.

[110359]19509.

[110360]Clevidence, D. E.; Overdier, D. G.; Peterson, R. S.; Porcella,

[110361]A.; Ye, H.; Paulson, K. E.; Costa, R. H.: Members of the HNF–3/forkhead

[110362]family of transcription factors exhibit distinct cellular expression

[110363]patterns in lung and regulate the surfactant protein B promoter. Dev.

[110364]Biol. 166: 195–209, 1994.

[110365]19510.

[110366]Hellqvist, M.; Mahlapuu, M.; Blixt, A.; Enerback, S.; Carlsson,

[110367]P.: The human forkhead protein FREAC-2 contains two functionally

[110368]redundant activation domains and interacts with TBP and TFIIIB. J.

[110369]Biol. Chem. 273: 23335-23343, 1998.

[110370]19511.

[110371]Hellqvist, M.; Mahlapuu, M.; Samuelsson, L.; Enerback, S.; Carlsson,

[110372]P.: Differential activation of lung-specific genes by two forkhead

[110373]proteins, FREAC-1 and FREAC-2. J. Biol. Chem. 271: 4482-4490, 1996.

[110374]19512.

[110375]Larsson, C.; Hellqvist, M.; Pierrou, S.; White, I.; Enerback, S.;

[110376]Carlsson, P.: Chromosomal localization of six human forkhead genes,

[110377]freac-1 (FKHL5), -3 (FKHL7), -4 (FKHL8), -5 (FKHL9), -6 (FKHL10),

[110378]and -8 (FKHL12). Genomics 30: 464-469, 1995.

[110379]19513.

[110380]Pierrou, S.; Hellqvist, M.; Samuelsson, L.; Enerback, S.;
Carlsson,

[110381]P.: Cloning and characterization of seven human forkhead
proteins:

[110382]binding site specificity and DNA bending. EMBO J. 13:
5002-5012,

[110383]1994.

[110384]19514.

[110385]Ernstsson, S.; Pierrou, S.; Hulander, M.; Cederberg, A.;
Hellqvist,

[110386]M.; Carlsson, P.; Enerback, S.: Characterization of the hu-
man forkhead

[110387]gene FREAC-4. J. Biol. Chem. 271: 21094-21099, 1996.

[110388]19515.

[110389]Lorenzetti, D.; Pareyson, D.; Sghirlanzoni, A.; Roa, B. B.;
Abbas,

[110390]N. E.; Pandolfo, M.; Di Donato, S.; Lupski, J. R.: A 1.5-Mb
deletion

[110391]in 17p11.2-p12 is frequently observed in Italian families

with hereditary

[110392]neuropathy with liability to pressure palsies. Am. J. Hum.

Genet. 56:

[110393]91–98, 1995.

[110394]19516.

[110395]Lupski, J. R.; Garcia, C. A.; Parry, G. J.; Patel, P. I.: Charcot–

Marie–Tooth

[110396]polyneuropathy syndrome: clinical, electrophysiological,

and genetic

[110397]aspects.In: Appel, S.: Current Neurology. Chicago: Mosby–

Yearbook

[110398](pub.) 1991. Pp. 1–25.

[110399]19517.

[110400]Lupski, J. R.; Pentao, L.; Williams, L. L.; Patel, P. I.: Stable

[110401]inheritance of the CMT1A DNA duplication in two patients

with CMT1

[110402]and NF1. Am. J. Med. Genet. 45: 92–96, 1993.

[110403]19518.

[110404]Lupski, J. R.; Wise, C. A.; Kuwano, A.; Pentao, L.; Parke, J.

[110405]T.; Glaze, D. G.; Ledbetter, D. H.; Greenberg, F.; Patel, P.

I.:

[110406]Gene dosage is a mechanism for Charcot-Marie-Tooth
disease type 1A. Nature

[110407]Genet. 1: 29-33, 1992.

[110408]19519.

[110409]MacMillan, J. C.; Upadhyaya, M.; Harper, P. S.: Charcot-
Marie-Tooth

[110410]disease type 1a (CMT1a): evidence for trisomy of the re-
gion p11.2

[110411]of chromosome 17 in south Wales families. J. Med. Genet.
29: 12-13,

[110412]1992.

[110413]19520.

[110414]Manfioletti, G.; Ruaro, M. E.; Del Sal, G.; Philipson, L.;
Schneider,

[110415]C.: A growth arrest-specific (gas) gene codes for a mem-
brane protein. Molec.

[110416]Cell. Biol. 10: 2924-2930, 1990.

[110417]19521.

[110418]Marques, W., Jr.; Thomas, P. K.; Sweeney, M. G.; Carr, L.;
Wood,

- [110419]N. W.: Dejerine-Sottas neuropathy and PMP22 point mutations: a new
base pair substitution and a possible "hot spot" on ser72.
Ann. Neurol. 43:
680-683, 1998.
- [110422]19522.
- [110423]Martinotti, A.; Cariani, C. T.; Melani, C.; Sozzi, G.; Spurr,
N. K.; Pierotti, M. A.; Colombo, M. P.: Isolation and mapping to
17p12-13 of the human homologous of the murine
growth arrest specific
Gas-3 gene. Hum. Molec. Genet. 1: 331-334, 1992.
- [110427]19523.
- [110428]Matise, T. C.; Chakravarti, A.; Patel, P. I.; Lupski, J. R.;
Nelis,
E.; Timmerman, V.; Van Broeckhoven, C.; Weeks, D. E.:
Detection of
tandem duplications and implications for linkage analysis.
Am. J.
Hum. Genet. 54: 1110-1121, 1994.
- [110432]19524.

[110433]Matsunami, N.; Smith, B.; Ballard, L.; Lensch, M. W.;
Robertson,
[110434]M.; Albertsen, H.; Hanemann, C. O.; Muller, H. W.; Bird, T.
D.; White,
[110435]R.; Chance, P. F.: Peripheral myelin protein-22 gene maps
in the
[110436]duplication in chromosome 17p11.2 associated with
Charcot-Marie-Tooth
[110437]1A. Nature Genet. 1: 176-179, 1992.
[110438]19525.
[110439]Nelis, E.; De Jonghe, P.; De Vriendt, E.; Patel, P. I.; Martin,
[110440]J.-J.; Van Broeckhoven, C.: Mutation analysis of the nerve
specific
[110441]promoter of the peripheral myelin protein 22 gene in
CMT1 disease
[110442]and HNPP. J. Med. Genet. 35: 590-593, 1998.
[110443]19526.
[110444]Nicholson, G. A.; Valentijn, L. J.; Cherryson, A. K.; Kenner-
son,
[110445]M. L.; Bragg, T. L.; DeKroon, R. M.; Ross, D. A.; Pollard, J.
D.;

[110446]McLeod, J. G.; Bolhuis, P. A.; Baas, F.: A frame shift mutation in

[110447]the PMP22 gene in hereditary neuropathy with liability to pressure

[110448]palsies. Nature Genet. 6: 263–266, 1994.

[110449]19527.

[110450]Niemann, S.; Sereda, M. W.; Suter, U.; Griffiths, I. R.; Nave,

[110451]K.-A.: Uncoupling of myelin assembly and Schwann cell differentiation

[110452]by transgenic overexpression of peripheral myelin protein 22. J.

[110453]Neurosci. 20: 4120–4128, 2000.

[110454]19528.

[110455]Palau, F.; Lofgren, A.; De Jonghe, P.; Bort, S.; Nelis, E.; Sevilla,

[110456]T.; Martin, J.-J.; Vilchez, J.; Prieto, F.; Van Broeckhoven, C.:

[110457]Origin of the de novo duplication in Charcot–Marie–Tooth disease type

[110458]1A: unequal nonsister chromatid exchange during spermatogenesis. Hum.

[110459]Molec. Genet. 2: 2031–2035, 1993.

[110460]19529.

[110461]Patel, P. I.; Roa, B. B.; Welcher, A. A.; Schoener-Scott, R.;

[110462]Trask, B. J.; Pentao, L.; Snipes, G. J.; Garcia, C. A.; Francke,
U.;

[110463]Shooter, E. M.; Lupski, J. R.; Suter, U.: The gene for the
peripheral

[110464]myelin protein PMP-22 is a candidate for Charcot-
Marie-Tooth disease

[110465]type 1A. Nature Genet. 1: 159-165, 1992.

[110466]19530.

[110467]Pentao, L.; Wise, C. A.; Chinault, A. C.; Patel, P. I.; Lupski,

[110468]J. R.: Charcot-Marie-Tooth type 1A duplication appears to
arise from

[110469]recombination at repeat sequences flanking the 1.5 Mb
monomer unit. Nature

[110470]Genet. 2: 292-300, 1992.

[110471]19531.

[110472]Raeymaekers, P.; Timmerman, V.; Nelis, E.; De Jonghe, P.;
Hoogendijk,

[110473]J. E.; Baas, F.; Barker, D. F.; Martin, J. J.; De Visser, M.; Bol-
huis,

[110474]P. A.; Van Broeckhoven, C.; HMSN Collaborative Research Group:

[110475]Duplication in chromosome 17p11.2 in Charcot-Marie-Tooth neuropathy

[110476]type 1a (CMT 1a). Neuromusc. Disord. 1: 93-97, 1991.

[110477]19532.

[110478]Raeymaekers, P.; Timmerman, V.; Nelis, E.; Van Hul, W.; De Jonghe,

[110479]P.; Martin, J.-J.; Van Broeckhoven, C.; HMSN Collaborative Research

[110480]Group: Estimation of the size of the chromosome 17p11.2 duplication

[110481]in Charcot-Marie-Tooth neuropathy type 1a (CMT1a). J. Med. Genet. 29:

[110482]5-11, 1992.

[110483]19533.

[110484]Reiter, L. T.; Hastings, P. J.; Nelis, E.; De Jonghe, P.; Van

[110485]Broeckhoven, C.; Lupski, J. R.: Human meiotic recombination products

[110486]revealed by sequencing a hotspot for homologous strand exchange in

[110487]multiple HNPP deletion patients. Am. J. Hum. Genet. 62:
1023–1033,

[110488]1998.

[110489]19534.

[110490]Reiter, L. T.; Murakami, T.; Koeuth, T.; Pentao, L.; Muzny,
D.

[110491]M.; Gibbs, R. A.; Lupski, J. R.: A recombination hotspot re-
sponsible

[110492]for two inherited peripheral neuropathies is located near a
mariner

[110493]transposon-like element. Nature Genet. 12: 288–297,
1996. Note:

[110494]Erratum: Nature Genet. 19: 303 only, 1998.

[110495]19535.

[110496]Roa, B. B.; Garcia, C. A.; Pentao, L.; Killian, J. M.; Trask,

[110497]B. J.; Suter, U.; Snipes, G. J.; Ortiz-Lopez, R.; Shooter, E.
M.;

[110498]Patel, P. I.; Lupski, J. R.: Evidence for a recessive PMP22
point

[110499]mutation in Charcot-Marie-Tooth disease type 1A. Nature
Genet. 5:

[110500]189–194, 1993.

[110501]19536.

[110502]Roa, B. B.; Garcia, C. A.; Suter, U.; Kulpa, D. A.; Wise, C. A.;

[110503]Mueller, J.; Welcher, A. A.; Snipes, G. J.; Shooter, E. M.; Patel,
tel,

[110504]P. I.; Lupski, J. R.: Charcot–Marie–Tooth disease type 1A:
association

[110505]with a spontaneous point mutation in the PMP22 gene.
New Eng. J.

[110506]Med. 329: 96–101, 1993.

[110507]19537.

[110508]LeGuern, E.; Gouider, R.; Lopes, J.; Abbas, N.; Gugenheim,
M.;

[110509]Tardieu, S.; Ravise, N.; Leger, J.–M.; Vallat, J.–M.; Bouche,
P.;

[110510]Agid, Y.; Brice, A.; French CMT Collaborative Research
Group: Constant

[110511]rearrangement of the CMT1A–REP sequences in HNPP pa-
tients with a deletion

[110512]in chromosome 17p11.2: a study of 30 unrelated cases.
Hum. Molec.

[110513]Genet. 4: 1673–1674, 1995.

[110514]19538.

[110515]De Smaele, E.; Zazzeroni, F.; Papa, S.; Nguyen, D. U.; Jin, R.;

[110516]Jones, J.; Cong, R.; Franzoso, G.: Induction of gadd45–beta by NF–kappa–B

[110517]downregulates pro–apoptotic JNK signaling. Nature 414: 308–313,

[110518]2001.

[110519]19539.

[110520]Garcia–Cardena, G.; Martasek, P.; Masters, B. S. S.; Skidd, P.

[110521]M.; Couet, J. C.; Li, S.; Lisanti, M. P.; Sessa, W. C.: Dissect– ing

[110522]the interaction between nitric oxide synthase (NOS) and caveolin:

[110523]functional significance of the NOS caveolin binding do– main in vivo. J.

[110524]Biol. Chem. 272: 25437–25440, 1997.

[110525]19540.

[110526]Isom, L. L.; Ragsdale, D. S.; De Jongh, K. S.; Westenbroek,

R.

[110527]E.; Reber, B. F. X.; Scheuer, T.; Catterall, W. A.: Structure
and
[110528]function of the beta-2 subunit of brain sodium channels,
a transmembrane
[110529]glycoprotein with a CAM motif. Cell 83: 433-442, 1995.
[110530]19541.

[110531]Mattei, M. G.; Stoetzel, C.; Perrin-Schmitt, F.: The B-HLH
protein
[110532]encoding the M-twist gene is located by in situ hybridiza-
tion on murine
[110533]chromosome 12. Mammalian Genome 4: 127-128, 1993.
[110534]19542.

[110535]Rose, C. S. P.; Patel, P.; Reardon, W.; Malcolm, S.; Winter,
R.
[110536]M.: The TWIST gene, although not disrupted in Saethre-
Chotzen patients
[110537]with apparently balanced translocations of 7p21, is mu-
tated in familial
[110538]and sporadic cases. Hum. Molec. Genet. 6: 1369-1373,
1997.
[110539]19543.

[110540]Shishido, E.; Higashijima, S.; Emori, Y.; Saigo, K.: Two
FGF-receptor

[110541]homologues of Drosophila: one is expressed in mesoder-
mal primordium

[110542]in early embryos. Development 117: 751-761, 1993.

[110543]19544.

[110544]Struhl, K.: Histone acetylation and transcriptional regula-
tory

[110545]mechanisms. Genes Dev. 12: 599-606, 1998.

[110546]19545.

[110547]Tsuji, K.; Narahara, K.; Yokoyama, Y.; Grzeschik, K.-H.;
Kunz,

[110548]J.: The breakpoint on 7p in a patient with t(6;7) and cran-
iosynostosis

[110549]is spanned by a YAC clone containing the D7S503 locus.
Hum. Genet. 95:

[110550]303-307, 1995.

[110551]19546.

[110552]Yousfi, M.; Lasmoles, F.; El Ghouzzi, V.; Marie, P. J.: Twist

[110553]haploinsufficiency in Saethre-Chotzen syndrome induces
calvarial osteoblast

- [110554]apoptosis due to increased TNF expression and caspase-2 activation. Hum. Molec. Genet. 11: 359-369, 2002.
- [110555]19547.
- [110556]Albrecht, U.; Sutcliffe, J. S.; Cattanach, B. M.; Beechey, C. V.;
- [110557]Armstrong, D.; Eichele, G.; Beaudet, A. L.: Imprinted expression of the murine Angelman syndrome gene, Ube3a, in hippocampal and Purkinje neurons. Nature Genet. 17: 75-78, 1997.
- [110558]19548.
- [110559]Fang, P.; Lev-Lehman, E.; Tsai, T.-F.; Matsuura, T.; Benton, C. S.; Sutcliffe, J. S.; Christian, S. L.; Kubota, T.; Halley, D. J.;
- [110560]Meijers-Heijboer, H.; Langlois, S.; Graham, J. M., Jr.; Beuten, J.;
- [110561]Willems, P. J.; Ledbetter, D. H.; Beaudet, A. L.: The spectrum of mutations in UBE3A causing Angelman syndrome. Hum. Molec. Genet. 8:

[110567]129–135, 1999.

[110568]19549.

[110569]Fung, D. C. Y.; Yu, B.; Cheong, K. F.; Smith, A.; Trent, R. J.

[110570]: UBE3A 'mutations' in two unrelated and phenotypically
different

[110571]Angelman syndrome patients. Hum. Genet. 102: 487–492,
1998.

[110572]19550.

[110573]Huang, L.; Kinnucan, E.; Wang, G.; Beaudenon, S.; Howley,
P. M.;

[110574]Huibregtse, J. M.; Pavletich, N. P.: Structure of an
E6AP–Ubch7 complex:

[110575]insights into ubiquitination by the E2–E3 enzyme cascade.
Science 286:

[110576]1321–1326, 1999.

[110577]19551.

[110578]Huibregtse, J. M.; Scheffner, M.; Howley, P. M.: A cellular
protein

[110579]mediates association of p53 with the E6 oncoprotein of
human papillomavirus

[110580]types 16 or 18. EMBO J. 10: 4129–4135, 1991.

[110581]19552.

[110582]Huibregtse, J. M.; Scheffner, M.; Howley, P. M.: Cloning
and expression

[110583]of the cDNA for E6-AP, a protein that mediates the inter-
action of

[110584]the human papillomavirus E6 oncoprotein with p53.

Molec. Cell. Biol. 13:

[110585]775-784, 1993.

[110586]19553.

[110587]Kishino, T.; Wagstaff, J.: Genomic organization of the
UBE3A/E6-AP

[110588]gene and related pseudogenes. Genomics 47: 101-107,
1998.

[110589]19554.

[110590]Malzac, P.; Webber, H.; Moncla, A.; Graham, J. M., Jr.;
Kukolich,

[110591]M.; Williams, C.; Pagon, R. A.; Ramsdell, L. A.; Kishino, T.;
Wagstaff,

[110592]J.: Mutation analysis of UBE3A in Angelman syndrome pa-
tients. Am.

[110593]J. Hum. Genet. 62: 1353-1360, 1998.

[110594]19555.

[110595]Rougeulle, C.; Glatt, H.; Lalande, M.: The Angelman syndrome

[110596]candidate gene, UBE3A/E6-AP, is imprinted in brain.
(Letter) Nature

[110597]Genet. 17: 14-15, 1997.

[110598]19556.

[110599]Dong, C.; Yang, D. D.; Tournier, C.; Whitmarsh, A. J.; Xu,
J.;

[110600]Davis, R. J.; Flavell, R. A.: JNK is required for effector T-cell
[110601]function but not for T-cell activation. Nature 405: 91-94,
2000.

[110602]19557.

[110603]Gupta, S.; Barrett, T.; Whitmarsh, A. J.; Cavanagh, J.; Sluss,
[110604]H. K.; Derijard, B.; Davis, R. J.: Selective interaction of JNK
protein

[110605]kinase isoforms with transcription factors. EMBO J. 15:
2760-2770,

[110606]1996.

[110607]19558.

[110608]Tournier, C.; Hess, P.; Yang, D. D.; Xu, J.; Turner, T. K.;
Nimnual,
[110609]A.; Bar-Sagi, D.; Jones, S. N.; Flavell, R. A.; Davis, R. J.: Re-
quirement
[110610]of JNK for stress-induced activation of the cytochrome c-
mediated
[110611]death pathway. Science 288: 870-874, 2000.
[110612]19559.
[110613]Daugherty, B. L.; Springer, M. S.: The beta-chemokine re-
ceptor
[110614]genes CCR1 (CMKBR1), CCR2 (CMKBR2), and CCR3
(CMKBR3) cluster within
[110615]285 kb on human chromosome 3p21. Genomics 41:
294-295, 1997.
[110616]19560.
[110617]Ayyanathan, K.; Naylor, S. L.; Kunapuli, S. P.: Structural
characterization
[110618]and fine chromosomal mapping of the human P2Y(1)
purinergic receptor
[110619]gene (P2RY1). Somat. Cell Molec. Genet. 22: 419-424,
1996.
[110620]19561.

[110621]Ayyanathan, K.; Webbs, T. E.; Sandhu, A. K.; Athwal, R. S.;
Barnard,

[110622]E. A.; Kunapuli, S. P.: Cloning and chromosomal localization of the

[110623]human P2Y₁ purinoceptor. Biochem. Biophys. Res. Commun. 218: 783–788,

[110624]1996.

[110625]19562.

[110626]Leon, C.; Hechler, B.; Freund, M.; Eckly, A.; Vial, C.;
Ohlmann,

[110627]P.; Dierich, A.; LeMeur, M.; Cazenave, J.-P.; Gachet, C.:
Defective

[110628]platelet aggregation and increased resistance to thrombosis in purinergic

[110629]P2Y₁ receptor-null mice. J. Clin. Invest. 104: 1731–1737,
1999.

[110630]19563.

[110631]Leon, C.; Hechler, B.; Vial, C.; Leray, C.; Cazenave, J.-P.;
Gachet,

[110632]C.: The P2Y₁(1) receptor is an ADP receptor antagonized by
ATP and

[110633]expressed in platelets and megakaryoblastic cells. FEBS

Letts. 403:

[110634]26–30, 1997.

[110635]19564.

[110636]Leon, C.; Vial, C.; Cazenave, J.-P.; Gachet, C.: Cloning and sequencing

[110637]of a human cDNA encoding endothelial P2Y1 purinoceptor. Gene 171:

[110638]295–297, 1996.

[110639]19565.

[110640]Hagiwara, Y.; Sasaoka, T.; Araishi, K.; Imamura, M.; Yori-fuji,

[110641]H.; Nonaka, I.; Ozawa, E.; Kikuchi, T.: Caveolin-3 deficiency causes

[110642]muscle degeneration in mice. Hum. Molec. Genet. 9: 3047–3054, 2000.

[110643]19566.

[110644]Herrmann, R.; Straub, V.; Blank, M.; Kutzick, C.; Franke, N.; Jacob,

[110645]E. N.; Lenard, H.-G.; Kroger, S.; Voit, T.: Dissociation of the dystroglycan

[110646]complex in caveolin-3-deficient limb girdle muscular dys-

trophy. Hum.

[110647]Molec. Genet. 9: 2335–2340, 2000.

[110648]19567.

[110649]Matsuda, C.; Hayashi, Y. K.; Ogawa, M.; Aoki, M.; Murayama, K.;

[110650]Nishino, I.; Nonaka, I.; Arahata, K.; Brown, R. H., Jr.: The sarcolemmal

[110651]proteins dysferlin and caveolin-3 interact in skeletal muscle. Hum.

[110652]Molec. Genet. 10: 1761–1766, 2001.

[110653]19568.

[110654]McNally, E. M.: Personal Communication. Chicago, Ill. 6/8/1998.

[110655]19569.

[110656]McNally, E. M.; de Sa Moreira, E.; Duggan, D. J.; Bonnemann, C.

[110657]G.; Lisanti, M. P.; Lidov, H. G. W.; Vainzof, M.; Passos-Bueno, M.

[110658]R.; Hoffman, E. P.; Zatz, M.; Kunkel, L. M.: Caveolin-3 in muscular

[110659]dystrophy. Hum. Molec. Genet. 7: 871–877, 1998.

[110660]19570.

[110661]Minetti, C.; Sotgia, F.; Bruno, C.; Scartezzini, P.; Broda, P.;

[110662]Bado, M.; Masetti, E.; Mazzocco, M.; Egeo, A.; Donati, M.

A.; Volonte,

[110663]D.; Galbiati, F.; Cordone, G.; Bricarelli, F. D.; Lisanti, M. P.;

[110664]Zara, F.: Mutations in the caveolin-3 gene cause autosomal dominant
mal dominant

[110665]limb-girdle muscular dystrophy. Nature Genet. 18:
365-368, 1998.

[110666]19571.

[110667]Sotgia, F.; Minetti, C.; Lisanti, M. P.: Localization of the

[110668]human caveolin-3 gene to the D3S18/D3S4163/D3S4539
locus (3p25), in

[110669]close proximity to the human oxytocin receptor gene:
identification

[110670]of the caveolin-3 gene as a candidate for deletion in
3p-syndrome. FEBS

[110671]Lett. 452: 177-180, 1999.

[110672]19572.

[110673]Sunada, Y.; Ohi, H.; Hase, A.; Ohi, H.; Hosono, T.; Arata,
S.;

[110674]Higuchi, S.; Matsumura, K.; Shimizu, T.: Transgenic mice expressing
[110675]mutant caveolin-3 show severe myopathy associated with
increased nNOS
[110676]activity. Hum. Molec. Genet. 10: 173-178, 2001.
[110677]19573.

[110678]Tang, Z.; Scherer, P. E.; Okamoto, T.; Song, K.; Chu, C.;
Kohtz,
[110679]D. S.; Nishimoto, I.; Lodish, H. F.; Lisanti, M. P.: Molecular
cloning
[110680]of caveolin-3, a novel member of the caveolin gene family
expressed
[110681]predominantly in muscle. J. Biol. Chem. 271: 2255-2261,
1996.
[110682]19574.

[110683]Vorgerd, M.; Bolz, H.; Patzold, T.; Kubisch, C.; Malin, J.-P.;
[110684]Mortier, W.: Phenotypic variability in rippling muscle dis-
ease. Neurology 52:
[110685]1453-1459, 1999.
[110686]19575.

[110687]Furlong, R. A.; Zhou, C. Y.; Ferguson-Smith, M. A.; Affara,

N.

[110688]A.: Characterization of a kinesin-related gene ATSV,
within the tuberous

[110689]sclerosis locus (TSC1) candidate region on chromosome
9q34. Genomics 33:

[110690]421-429, 1996.

[110691]19576.

[110692]Keller, M. P.; Seifried, B. A.; Rabin, B. A.; Chance, P. F.:
Mapping

[110693]of the kinesin-related gene ATSV to chromosome 2q37.
Hum. Genet. 104:

[110694]254-256, 1999.

[110695]19577.

[110696]Kikkawa, M.; Okada, Y.; Hirokawa, N.: 15-angstrom reso-
lution model

[110697]of the monomeric kinesin motor, KIF1A. Cell 100:
241-252, 2000.

[110698]19578.

[110699]Kikkawa, M.; Sablin, E. P.; Okada, Y.; Yajima, H.; Fletterick,

[110700]R. J.; Hirokawa, N.: Switch-based mechanism of kinesin
motors. Nature 411:

[110701]439–445, 2001.

[110702]19579.

[110703]Klopfenstein, D. R.; Tomishige, M.; Stuurman, N.; Vale, R.
D.:

[110704]Role of phosphatidylinositol(4,5)bispophosphate organiza-
tion in membrane

[110705]transport by the Unc104 kinesin motor. Cell 109:
347–358, 2002.

[110706]19580.

[110707]Okada, Y.; Yamazaki, H.; Sekine–Aizawa, Y.; Hirokawa, N.:
The

[110708]neuron–specific kinesin superfamily protein KIF1A is a
unique monomeric

[110709]motor for anterograde axonal transport of synaptic vesicle
precursors. Cell 81:

[110710]769–780, 1995.

[110711]19581.

[110712]Yonekawa, Y.; Harada, A.; Okada, Y.; Funakoshi, T.; Kanai,
Y.;

[110713]Takei, Y.; Terada, S.; Noda, T.; Hirokawa, N.: Defect in
synaptic

[110714] vesicle precursor transport and neuronal cell death in
KIF1A motor
[110715] protein-deficient mice. J. Cell Biol. 141: 431–441, 1998.
[110716] 19582.

[110717] Godbout, R.; Squire, J.: Amplification of a DEAD box pro-
tein gene
[110718] in retinoblastoma cell lines. Proc. Nat. Acad. Sci. 90:
7578–7582,
[110719] 1993.
[110720] 19583.

[110721] Jones, J. M.; Meisler, M. H.; Isom, L. L.: Scn2b, a voltage-
gated
[110722] sodium channel beta-2 gene on mouse chromosome 9.
Genomics 34: 258–259,
[110723] 1996.
[110724] 19584.

[110725] Bernard, O.; Burkitt, V.; Webb, G. C.; Bottema, C. D. K.;
Nicholl,
[110726] J.; Sutherland, G. R.; Matthew, P.: Structure and chromoso-
mal localization
[110727] of the genomic locus encoding the Kiz1 LIM-kinase gene.

Genomics 35:

[110728]593–596, 1996.

[110729]19585.

[110730]Mao, X.; Jones, T. A.; Williamson, J.; Gutowski, N. J.;
Proschel,

[110731]C.; Noble, M.; Sheer, D.: Assignment of the human and
mouse LIM-kinase

[110732]genes (LIMK1; Limk1) to chromosome bands 7q11.23 and
5G1, respectively,

[110733]by in situ hybridization. Cytogenet. Cell Genet. 74:
190–191, 1996.

[110734]19586.

[110735]Mizuno, K.; Okano, I.; Ohashi, K.; Nunoue, K.; Kuma, K.;
Miyata,

[110736]T.; Nakamura, T.: Identification of a human cDNA encod-
ing a novel

[110737]protein kinase with two repeats of the LIM/double zinc
finger motif. Oncogene 9:

[110738]1605–1612, 1994.

[110739]19587.

[110740]Okano, I.; Hiraoka, J.; Otera, H.; Nunoue, K.; Ohashi, K.;

Iwashita,

[110741]S.; Hirai, M.; Mizuno, K.: Identification and characterization of

[110742]a novel family of serine/threonine kinases containing two N-terminal

[110743]LIM motifs. J. Biol. Chem. 270: 31321–31330, 1995.

[110744]19588.

[110745]Proschel, C.; Blouin, M. J.; Gutowski, N. J.; Ludwig, R.; Noble,

[110746]M.: Limk1 is predominantly expressed in neural tissues and phosphorylates

[110747]serine, threonine and tyrosine residues in vitro. Oncogene 11: 1271–1281,

[110748]1995.

[110749]19589.

[110750]Sotiropoulos, A.; Gineitis, D.; Copeland, J.; Treisman, R.: Signal-regulated

[110751]activation of serum response factor is mediated by changes in actin

[110752]dynamics. Cell 98: 159–169, 1999.

[110753]19590.

[110754]Tassabehji, M.; Metcalfe, K.; Fergusson, W. D.; Carette, M.
J.

[110755]A.; Dore, J. K.; Donnai, D.; Read, A. P.; Proschel, C.;
Gutowski,

[110756]N. J.; Mao, X.; Sheer, D.: LIM-kinase deleted in Williams
syndrome.

[110757](Letter) Nature Genet. 13: 272–273, 1996.

[110758]19591.

[110759]de Sauvage, F. J.; Camerato, T. R.; Goeddel, D. V.: Primary
structure

[110760]and functional expression of the human receptor for Es-
cherichia coli

[110761]heat-stable enterotoxin. J. Biol. Chem. 266:
17912–17918, 1991.

[110762]19592.

[110763]Mann, E. A.; Swenson, E. S.; Copeland, N. G.; Gilbert, D. J.;
Jenkins,

[110764]N. A.; Taguchi, T.; Testa, J. R.; Giannella, R. A.: Localiza-
tion

[110765]of the guanylyl cyclase C gene to mouse chromosome 6
and human chromosome

[110766]12p12. Genomics 34: 265–267, 1996.

[110767]19593.

[110768]Singh, S.; Singh, G.; Heim, J. M.; Gerzer, R.: Isolation and expression

[110769]of a guanylate cyclase-coupled heat stable enterotoxin receptor cDNA

[110770]from a human colonic cell line. Biochem. Biophys. Res. Commun. 179:

[110771]1455-1463, 1991.

[110772]19594.

[110773]Yamamoto, Y.; Huibregtse, J. M.; Howley, P. M.: The human E6-AP

[110774]gene (UBE3A) encodes three potential protein isoforms generated by

[110775]differential splicing. Genomics 41: 263-266, 1997.

[110776]19595.

[110777]Han, J.; Knops, J. F.; Longshore, J. W.; King, P. H.: Localization

[110778]of human elav-like neuronal protein 1 (Hel-N1) on chromosome 9p21

[110779]by chromosome microdissection polymerase chain reaction and fluorescence

[110780]in situ hybridization. Genomics 36: 189–191, 1996.

[110781]19596.

[110782]King, P. H.: Hel–N2: a novel isoform of Hel–N1 which is conserved

[110783]in rat neural tissue and produced in early embryogenesis. Gene 151:

[110784]261–265, 1994.

[110785]19597.

[110786]King, P. H.; Levine, T. D.; Fremeau, R. T., Jr.; Keene, J. D.:

[110787]Mammalian homologs of Drosophila ELAV localized to a neuronal subset

[110788]can bind in vitro to the 3' UTR of mRNA encoding the Id transcriptional

[110789]repressor. J. Neurosci. 14: 1943–52, 1994.

[110790]19598.

[110791]Abel, K. J.; Brody, L. C.; Valdes, J. M.; Erdos, M. R.; McKinley,

[110792]D. R.; Castilla, L. H.; Merajver, S. D.; Couch, F. J.; Friedman, L.

[110793]S.; Ostermeyer, E. A.; Lynch, E. D.; King, M.–C.; Welcsh, P. L.; Osborne–Lawrence,

[110794]S.; Spillman, M.; Bowcock, A. M.; Collins, F. S.; Weber, B.
L.: Characterization

[110795]of EZH1, a human homolog of Drosophila enhancer of
zeste near BRCA1. Genomics 37:

[110796]161–171, 1996.

[110797]19599.

[110798]Quirk, J.; van den Heuvel, M.; Henrique, D.; Marigo, V.;
Jones,

[110799]T. A.; Tabin, C.; Ingham, P. W.: The Smoothed gene and
Hedgehog

[110800]signal transduction in Drosophila and vertebrate develop-
ment. Cold

[110801]Spring Harbor Symp. Quant. Biol. 62: 217–226, 1997.

[110802]19600.

[110803]Sublett, J. E.; Entrekin, R. E.; Look, A. T.; Reardon, D. A.:

[110804]Chromosomal localization of the human Smoothed
gene (SMOH) to 7q32.3

[110805]by fluorescence in situ hybridization and radiation hybrid
mapping. Genomics 50:

[110806]112–114, 1998.

[110807]19601.

- [110808]Xie, J.; Murone, M.; Luoh, S.-M.; Ryan, A.; Gu, Q.; Zhang, C.;
- [110809]Bonifas, J. M.; Lam, C.-W.; Hynes, M.; Goddard, A.; Rosenthal, A.;
- [110810]Epstein, E. H., Jr.; de Sauvage, F. J.: Activating Smoothed mutations
- [110811]in sporadic basal-cell carcinoma. *Nature* 391: 90-92, 1998.
- [110812]19602.
- [110813]Ferbis, D.; Le Chalony, C.; Prosperi, M.-T.; Muleris, M.; Vincent-Salomon,
- [110814]A.; Goubin, G.: Identification, nuclear localization, and binding
- [110815]activities of OZF, a human protein solely composed of zinc finger
- [110816]motifs. *Europ. J. Biochem.* 236: 991-995, 1996.
- [110817]19603.
- [110818]Le Chalony, C.; Apiou, F.; Pibouin, L.; Dutrillaux, B.; Goubin,
- [110819]G.: Constitutive amplification of a zinc finger protein gene in cattle. *DNA*
- [110820]*Cell Biol.* 15: 83-88, 1996.

[110821]19604.

[110822]Le Chalony, C.; Prosperi, M.-T.; Haluza, R.; Apiou, F.;
Dutrillaux,

[110823]B.; Goubin, G.: The OZF gene encodes a protein consisting
essentially

[110824]of zinc-finger motifs. J. Molec. Biol. 236: 399-404, 1994.

[110825]19605.

[110826]Sereda, M.; Griffiths, I. Puhlhofer, A.; Stewart, H.; Rossner,

[110827]M. J.; Zimmermann, F.; Magyar, J. P.; Schneider, A.; Hund,
E.; Meinck,

[110828]H.-M.; Suter, U.; Nave, K.-A.: A transgenic rat model of
Charcot-Marie-Tooth

[110829]disease. Neuron 16: 1049-1060, 1996.

[110830]19606.

[110831]Silander, K.; Meretoja, P.; Nelis, E.; Timmerman, V.; Van
Broeckhoven,

[110832]C.; Aula, P.; Savontaus, M.-L.: A de novo duplication in
17p11.2

[110833]and a novel mutation in the P(0) gene in two Dejerine-Sot-
tas syndrome

[110834]patients. Hum. Mutat. 8: 304-310, 1996.

[110835]19607.

[110836]Stogbauer, F.; Young, P.; Kerschensteiner, M.; Ringelstein, E.

[110837]B.; Assmann, G.; Funke, H.: Recurrent brachial plexus palsies as

[110838]the only clinical expression of hereditary neuropathy with liability

[110839]to pressure palsies associated with a de novo deletion of the peripheral

[110840]myelin protein-22 gene. Muscle Nerve 21: 1199-1201, 1998.

[110841]19608.

[110842]Suter, U.; Moskow, J. J.; Welcher, A. A.; Snipes, G. J.; Kosaras,

[110843]B.; Sidman, R. L.; Buchberg, A. M.; Shooter, E. M.: A leucine-to-proline

[110844]mutation in the putative first transmembrane domain of the 22-kDa

[110845]peripheral myelin protein in the trembler-J mouse. Proc. Nat. Acad.

[110846]Sci. 89: 4382-4386, 1992.

[110847]19609.

- [110848]Suter, U.; Welcher, A. A.; Ozcelik, T.; Snipes, G. J.; Kosaras,
[110849]B.; Francke, U.; Billings-Gagliardi, S.; Sidman, R. L.;
Shooter, E.
[110850]M.: Trembler mouse carries a point mutation in a myelin
gene. Nature 356:
[110851]241-244, 1992.
[110852]19610.
[110853]Takahashi, E.; Takeda, O.; Himoro, M.; Nanao, K.; Takada,
G.;
[110854]Hayasaka, K.: Localization of PMP-22 gene (candidate
gene for the
[110855]Charcot-Marie-Tooth disease 1A) to band 17p11.2 by di-
rect R-banding
[110856]fluorescence in situ hybridization. Jpn. J. Hum. Genet. 37:
303-306,
[110857]1992.
[110858]19611.
[110859]Timmerman, V.; Nelis, E.; Van Hul, W.; Nieuwenhuijsen, B.
W.;
[110860]Chen, K. L.; Wang, S.; Othman, K. B.; Cullen, B.; Leach, R.
J.; Hanemann,

[110861]C. O.; De Jonghe, P.; Raeymaekers, P.; van Ommen, G.-J. B.; Martin,

[110862]J.-J.; Muller, H. W.; Vance, J. M.; Fischbeck, K. H.; Van Broeckhoven,

[110863]C.: The peripheral myelin protein gene PMP-22 is contained within

[110864]the Charcot-Marie-Tooth disease type 1A duplication. Nature Genet. 1:

[110865]171-175, 1992.

[110866]19612.

[110867]Tobler, A. R.; Liu, N.; Mueller, L.; Shooter, E. M.: Differential

[110868]aggregation of the Trembler and Trembler J mutants of peripheral myelin

[110869]protein 22. Proc. Nat. Acad. Sci. 99: 483-488, 2002.

[110870]19613.

[110871]Upadhyaya, M.; Roberts, S. H.; Farnham, J.; MacMillan, J. C.;

[110872]Clarke, A.; Heath, J. P.; Hodges, I. C. G.; Harper, P. S.: Charcot-Marie-Tooth

[110873]disease 1A (CMT1A) associated with a maternal duplication of chromosome

[110874]17p11.2–12. Hum. Genet. 91: 392–394, 1993.

[110875]19614.

[110876]Valentijn, L. J.; Baas, F.; Wolterman, R. A.; Hoogendijk, J. E.;

[110877]van den Bosch, N. H. A.; Zorn, I.; Gabreels–Festen, A. A. W. M.; de

[110878]Visser, M.; Bolhuis, P. A.: Identical point mutations of PMP–22 in

[110879]Trembler–J mouse and Charcot–Marie–Tooth disease type 1A. Nature

[110880]Genet. 2: 288–291, 1992.

[110881]19615.

[110882]Valentijn, L. J.; Bolhuis, P. A.; Zorn, I.; Hoogendijk, J. E.;

[110883]van den Bosch, N.; Hensels, G. W.; Stanton, V. P., Jr.; Housman, D.

[110884]E.; Fischbeck, K. H.; Ross, D. A.; Nicholson, G. A.; Meer–shoek, E.

[110885]J.; Dauwerse, H. G.; van Ommen, G.–J. B.; Baas, F.: The peripheral

[110886]myelin gene PMP–22/GAS–3 is duplicated in Charcot–Marie–Tooth disease

[110887]type 1A. Nature Genet. 1: 166–170, 1992.

[110888]19616.

[110889]Valentijn, L. J.; Ouvrier, R. A.; van den Bosch, N. H. A.; Bolhuis,

[110890]P. A.; Baas, F.; Nicholson, G. A.: Dejerine–Sottas neuropathy is

[110891]associated with a de novo PMP22 mutation. Hum. Mutat. 5: 76–80,

[110892]1995.

[110893]19617.

[110894]Vance, J. M.: Hereditary motor and sensory neuropathies. J.

[110895]Med. Genet. 28: 1–5, 1991.

[110896]19618.

[110897]Wise, C. A.; Garcia, C. A.; Davis, S. N.; Heju, Z.; Pentao, L.;

[110898]Patel, P. I.; Lupski, J. R.: Molecular analyses of unrelated Charcot–Marie–Tooth

[110899](CMT) disease patients suggest a high frequency of the CMT1A duplication. Am.

[110900]J. Hum. Genet. 53: 853–863, 1993.

[110901]19619.

[110902]Wulfsberg, E. A.; Weaver, R. P.; Cunniff, C. M.; Jones, M. C.;

[110903]Jones, K. L.: Chromosome 10qter deletion syndrome: a review and report

[110904]of three new cases. Am. J. Med. Genet. 32: 364–367, 1989.

[110905]19620.

[110906]Young, P.; Wiebusch, H.; Stogbauer, F.; Ringelstein, B.; Assmann,

[110907]G.; Funke, H.: A novel frameshift mutation in PMP22 accounts for

[110908]hereditary neuropathy with liability to pressure palsies. Neurology 48:

[110909]450–452, 1997.

[110910]19621.

[110911]Angrist, M.; Wells, D. E.; Chakravarti, A.; Pandey, A.: Chromosomal

[110912]localization of the mouse Src-like adapter protein (Slap) gene and

[110913]its putative human homolog SLA. Genomics 30: 623–625, 1995.

[110914]19622.

[110915]Holland, S. J.; Liao, X. C.; Mendenhall, M. K.; Zhou, X.;
Pardo,

[110916]J.; Chu, P.; Spencer, C.; Fu, A.; Sheng, N.; Yu, P.; Pali, E.;
Nagin,

[110917]A.; and 14 others: Functional cloning of Src-like adapter
protein-2

[110918](SLAP-2), a novel inhibitor of antigen receptor signaling. J.
Exp.

[110919]Med. 194: 1263-1276, 2001.

[110920]19623.

[110921]Kratchmarova, I.; Sosinowski, T.; Weiss, A.; Witter, K.; Vin-
cenz,

[110922]C.; Pandey, A.: Characterization of promoter region and
genomic structure

[110923]of the murine and human genes encoding Src like adapter
protein. Gene 262:

[110924]267-273, 2001.

[110925]19624.

[110926]Meijerink, P. H. S.; Yanakiev, P.; Zorn, I.; Grierson, A. J.;
Bikker,

- [110927]H.; Dye, D.; Kalaydjieva, L.; Baas, F.: The gene for the human Src-like
[110928]adaptor protein (hSLAP) is located within the 64-kb intron
of the
[110929]thyroglobulin gene. *Europ. J. Biochem.* 254: 297–303,
1998.
[110930]19625.
- [110931]Sosinowski, T.; Pandey, A.; Dixit, V. M.; Weiss, A.: Src-like
[110932]adaptor protein (SLAP) is a negative regulator of T cell re-
ceptor
[110933]signaling. *J. Exp. Med.* 191: 463–474, 2000.
[110934]19626.
- [110935]Bailey, S. M.; Cornforth, M. N.; Kurimasa, A.; Chen, D. J.;
Goodwin,
[110936]E. H.: Strand-specific postreplicative processing of mam-
malian telomeres. *Science* 293:
[110937]2462–2465, 2001.
[110938]19627.
- [110939]Junghans, R. P.; Anderson, C. L.: The protection receptor
for
[110940]IgG catabolism is the beta-2-microglobulin-containing

neonatal intestinal

[110941]transport receptor. Proc. Nat. Acad. Sci. 93: 5512–5516,
1996.

[110942]19628.

[110943]Junghans, R. P.; Ebralidze, A.; Tiwari, B.: Does (CUG)_n re-
peat

[110944]in DMPK mRNA 'paint' chromosome 19 to suppress distant
genes to create

[110945]the diverse phenotype of myotonic dystrophy? A new hy-
pothesis of long-range

[110946]cis autosomal inactivation. Neurogenetics 3: 59–67, 2001.

[110947]19629.

[110948]Waldmann, T. A.; Strober, W.: Metabolism of immunoglob-
ulins. Prog.

[110949]Allergy 13: 1–110, 1969.

[110950]19630.

[110951]Beddow, A. L.; Richards, S. A.; Orem, N. R.; Macara, I. G.:
The

[110952]Ran/TC4 GTPase-binding domain: identification by ex-
pression cloning

[110953]and characterization of a conserved sequence motif. Proc.

Nat. Acad.

[110954]Sci. 92: 3328–3332, 1995.

[110955]19631.

[110956]Fauser, S.; Aslanukov, A.; Roepman, R.; Ferreira, P. A.: Ge-
nomic

[110957]organization, expression, and localization of murine Ran-
binding protein

[110958]2 (RanBP2) gene. Mammalian Genome 12: 406–415, 2001.

[110959]19632.

[110960]Krebber, H.; Bastians, H.; Hoheisel, J.; Lichter, P.; Ponst-
ingl,

[110961]H.; Joos, S.: Localization of the gene encoding the Ran-
binding protein

[110962]RanBP2 to human chromosome 2q11–q13 by fluorescence
in situ hybridization. Genomics 43:

[110963]247–248, 1997.

[110964]19633.

[110965]Pichler, A.; Gast, A.; Seeler, J. S.; Dejean, A.; Melchior, F.

[110966]: The nucleoporin RanBP2 has SUMO1 E3 ligase activity.
Cell 108:

[110967]109–120, 2002.

[110968]19634.

[110969]Wu, J.; Matunis, M. J.; Kraemer, D.; Blobel, G.; Coutavas, E.:

[110970]Nup358, a cytoplasmically exposed nucleoporin with peptide repeats,

[110971]Ran-GTP binding sites, zinc fingers, a cyclophilin A homologous domain,

[110972]and a leucine-rich region. J. Biol. Chem. 270: 14209-14213, 1995.

[110973]19635.

[110974]Yokoyama, N.; Hayashi, N.; Seki, T.; Pante, N.; Ohba, T.; Nishii,

[110975]K.; Kuma, K.; Hayashida, T.; Miyata, T.; Abei, U.; Fukui, M.; Nishimoto,

[110976]T.: A giant nucleopore protein that binds Ran/TC4. Nature 376:

[110977]184-188, 1995.

[110978]19636.

[110979]Chang, A. C.-M.; Janosi, J.; Hulsbeek, M.; de Jong, D.; Jeffrey,

[110980]K. J.; Noble, J. R.; Reddel, R. R.: A novel human cDNA highly homologous

[110981]to the fish hormone stanniocalcin. *Molec. Cell. Endocr.*
112: 241–247,
[110982]1995.
[110983]19637.
[110984]Chang, A. C.–M.; Jeffrey, K. J.; Tokutake, Y.; Shimamoto,
A.; Neumann,
[110985]A. A.; Dunham, M. A.; Cha, J.; Sugawara, M.; Furuichi, Y.;
Reddel,
[110986]R. R.: Human stanniocalcin (STC): genomic structure,
chromosomal
[110987]localization, and the presence of CAG trinucleotide re-
peats. *Genomics* 47:
[110988]393–398, 1998.
[110989]19638.
[110990]Jellinek, D. A.; Chang, A. C.; Larsen, M. R.; Wang, X.;
Robinson,
[110991]P. J.; Reddel, R. R.: Stanniocalcin 1 and 2 are secreted as
phosphoproteins
[110992]from human fibrosarcoma cells. *Biochem. J.* 350:
453–461, 2000.
[110993]19639.

- [110994]Olsen, H. S.; Cepeda, M. A.; Zhang, Q.-Q.; Rosen, C. A.;
Vozzolo,
[110995]B. L.; Wagner, G. F.: Human stanniocalcin: a possible hor-
monal regulator
[110996]of mineral metabolism. Proc. Nat. Acad. Sci. 93:
1792-1796, 1996.
[110997]19640.
- [110998]Varghese, R.; Wong, C. K. C.; Deol, H.; Wagner, G. F.; Di-
Mattia,
[110999]G. E.: Comparative analysis of mammalian stanniocalcin
genes. Endocrinology 139:
[111000]4714-4725, 1998.
[111001]19641.
- [111002]Wagner, G. F.; Guiraudon, C. C.; Milliken, C.; Copp, D. H.:
Immunological
[111003]and biological evidence for a stanniocalcin-like hormone
in human
[111004]kidney. Proc. Nat. Acad. Sci. 92: 1871-1875, 1995.
[111005]19642.
- [111006]Bartsch, O.; Wagner, A.; Hinkel, G. K.; Lichtner, P.; Murken,
J.;

- [111007]Schuffenhauer, S.: No evidence for chromosomal microdeletions at
the second DiGeorge syndrome locus on 10p near
D10S585. (Letter) Am.
J. Med. Genet. 83: 425–426, 1999.
- [111010]19643.
- [111011]Daw, S. C. M.; Taylor, C.; Kraman, M.; Call, K.; Mao, J.;
Schuffenhauer,
S.; Meitinger, T.; Lipson, T.; Goodship, J.; Scambler, P.: A
common
region of 10p deleted in DiGeorge and velocardiofacial
syndromes. Nature
Genet. 13: 458–461, 1996.
- [111014]19644.
- [111015]19644.
- [111016]Lichtner, P.; Konig, R.; Hasegawa, T.; Van Esch, H.;
Meitinger,
T.; Schuffenhauer, S.: An HDR (hypoparathyroidism, deaf-
ness, renal
dysplasia) syndrome locus maps distal to the DiGeorge
syndrome region
on 10p13/14. J. Med. Genet. 37: 33–37, 2000.

[111020]19645.

[111021]Artzt, K.; Shin, H. S.; Bennett, D.: Gene mapping within the
T/t-complex

[111022]of the mouse: II: anomalous position of the H-2 complex
in t-haplotypes. Cell 28:

[111023]471-476, 1982.

[111024]19646.

[111025]Watanabe, T. K.; Fujiwara, T.; Shimizu, F.; Okuno, S.;
Suzuki,

[111026]M.; Takahashi, E.; Nakamura, Y.; Hirai, Y.: Cloning, ex-
pression,

[111027]and mapping of TCTEL1, a putative human homologue of
murine Tcte1,

[111028]to 6q. Cytogenet. Cell Genet. 73: 153-156, 1996.

[111029]19647.

[111030]Davidson, J. D.; Riley, B.; Burright, E.N.; Duvick, L.A.;
Zoghbi,

[111031]H.Y.; Orr, H. T.: Identification and characterization of an
ataxin-1-interactin

[111032]g protein: A1Up, a ubiquitin-like nuclear protein. Hum.
Molec. Genet. 9:

[111033]2305–2312, 2000.

[111034]19648.

[111035]Genis, D.; Matilla, T.; Volpini, V.; Rosell, J.; Davalos, A.; Ferrer,

[111036]I.; Molins, A.; Estivill, X.: Clinical, neuropathologic, and genetic

[111037]studies of a large spinocerebellar ataxia type 1 (SCA1) kindred: (CAG)_n

[111038]expansion and early premonitory signs and symptoms. Neurology 45:

[111039]24–30, 1995.

[111040]19649.

[111041]Klanke, C. A.; Su, Y. R.; Callen, D. F.; Wang, Z.; Meneton, P.;

[111042]Baird, N.; Kandasamy, R. A.; Orlowski, J.; Otterud, B. E.; Leppert,

[111043]M.; Shull, G. E.; Menton, A. G.: Molecular cloning and physical and

[111044]genetic mapping of a novel human Na⁺/H⁺ exchanger (NHE5/SLC9A5) to

[111045]chromosome 16q22.1. Genomics 25: 615–622, 1995.

[111046]19650.

[111047]Chen, C.-Y.; Gherzi, R.; Ong, S.-E.; Chan, E. L.; Raijmakers, R.;

[111048]Pruijn, G. J. M.; Stoecklin, G.; Moroni, C.; Mann, M.; Karin, M.:

[111049]AU binding proteins recruit the exosome to degrade ARE-containing

[111050]mRNAs. Cell 107: 451-464, 2001.

[111051]19651.

[111052]Doyle, K.; Zhang, Y.; Baer, R.; Bina, M.: Distinguishable patterns

[111053]of protein-DNA interactions involving complexes of basic helix-loop-helix

[111054]proteins. J. Biol. Chem. 269: 12099-12105, 1994.

[111055]19652.

[111056]Hu, J.-S.; Olson, E. N.; Kingston, R. E.: HEB, a helix-loop-helix

[111057]protein related to E2A and ITF2 that can modulate the DNA-binding

[111058]ability of myogenic regulatory factors. Molec. Cell. Biol. 12: 1031-1042,

[111059]1992.

[111060]19653.

[111061]Sawada, S.; Littman, D. R.: A heterodimer of HEB and an
E12-related

[111062]protein interacts with the CD4 enhancer and regulates its
activity

[111063]in T-cell lines. Molec. Cell. Biol. 13: 5620-5628, 1993.

[111064]19654.

[111065]Zhang, Y.; Babin, J.; Feldhaus, A. L.; Singh, H.; Sharp, P. A.;

[111066]Bina, M.: HTF4: a new human helix-loop-helix protein.
Nucleic Acids

[111067]Res. 19: 4555 only, 1991.

[111068]19655.

[111069]Zhang, Y.; Bina, M.: The nucleotide sequence of the hu-
man transcription

[111070]factor HTF4a cDNA. DNA Sequence 2: 397-403, 1992.

[111071]19656.

[111072]Zhang, Y.; Flejter, W. L.; Barcroft, C. L.; Riviere, M.; Szpirer,

[111073]J.; Szpirer, C.; Bina, M.: Localization of the human HTF4

transcription

[111074] factors 4 gene (TCF12) to chromosome 15q21. Cytogenet.

Cell Genet. 68:

[111075] 235–238, 1995.

[111076] 19657.

[111077] Hua, X.; Yokoyama, C.; Wu, J.; Briggs, M. R.; Brown, M. S.;

Goldstein,

[111078] J. L.; Wang, X.: SREBP-2, a second basic-he-

lix-loop-helix-leucine

[111079] zipper protein that stimulates transcription by binding to

a sterol

[111080] regulatory element. Proc. Nat. Acad. Sci. 90:

11603–11607, 1993.

[111081] 19658.

[111082] Miserez, A. R.; Cao, G.; Probst, L. C.; Hobbs, H. H.: Struc-

ture

[111083] of the human gene encoding sterol regulatory element

binding protein

[111084] 2 (SREBF2). Genomics 40: 31–40, 1997.

[111085] 19659.

[111086] Yang, J.; Sato, R.; Goldstein, J. L.; Brown, M. S.: Sterol-

resistant

[111087]transcription in CHO cells caused by gene rearrangement
that truncates

[111088]SREBP-2. Genes Dev. 8: 1910–1919, 1994.

[111089]19660.

[111090]Rowen, L.; Young, J.; Birditt, B.; Kaur, A.; Madan, A.;
Philipps,

[111091]D. L.; Qin, S.; Minx, P.; Wilson, R. K.; Hood, L.; Graveley, B.
R.

[111092]: Analysis of the human neurexin genes: alternative splic-
ing and the

[111093]generation of protein diversity. Genomics 79: 587–597,
2002.

[111094]19661.

[111095]Tabuchi, K.; Sudhof, T. C.: Structure and evolution of
neurexin

[111096]genes: insight into the mechanism of alternative splicing.
Genomics 79:

[111097]849–859, 2002.

[111098]19662.

[111099]Ullrich, B.; Ushkaryov, Y. A.; Sudhof, T. C.: Cartography of
neurexins:

[111100]more than 1000 isoforms generated by alternative splicing and expressed

[111101]in distinct subsets of neurons. Neuron 14: 497–507, 1995.

[111102]19663.

[111103]Ushkaryov, Y. A.; Petrenko, A. G.; Geppert, M.; Sudhof, T. C.

[111104]; Neurexins: synaptic cell surface proteins related to the alpha-latrotoxin

[111105]receptor and laminin. Science 257: 50–56, 1992.

[111106]19664.

[111107]Bergman, L.; Silins, G.; Grimmond, S.; Hummerich, H.; Stewart,

[111108]C.; Little, P.; Hayward, N.: A 500–kb sequence–ready cosmid contig

[111109]and transcript map of the MEN1 region on 11q13. Genomics 55: 49–56,

[111110]1999.

[111111]19665.

[111112]Ushkaryov, Y. A.; Petrenko, A. G.; Geppert, M.; Sudhof, T. C.:

[111113]Neurexins: synaptic cell surface proteins related to the alpha-latrotoxin

[111114]receptor and laminin. Science 257: 50–56, 1992.

[111115]19666.

[111116]Scheiffele, P.; Fan, J.; Choih, J.; Fetter, R.; Serafini, T.:

[111117]Neurologin expressed in nonneuronal cells triggers presynaptic development

[111118]in contacting axons. Cell 101: 657–669, 2000.

[111119]19667.

[111120]Abderrahmani, A.; Steinmann, M.; Plaisance, V.; Niederhauser, G.;

[111121]Haeffliger, J.-A.; Mooser, V.; Bonny, C.; Nicod, P.; Waeber, G.: The

[111122]transcriptional repressor REST determines the cell-specific expression

[111123]of the human MAPK8IP1 gene encoding IB1 (JIP-1). Molec. Cell. Biol. 21:

[111124]7256–7627, 2001.

[111125]19668.

[111126]Cavaloc, Y.; Popielarz, M.; Fuchs, J.-P.; Gattoni, R.; Stevenin,

[111127]J.: Characterization and cloning of the human splicing factor 9G8:

[111128]a novel 35 kDa factor of the serine/arginine protein family. EMBO

[111129]J. 13: 2639–2649, 1994.

[111130]19669.

[111131]Popielarz, M.; Cavaloc, Y.; Mattei, M.–G.; Gattoni, R.; Stevenin,

[111132]J.: The gene encoding human splicing factor 9G8: structure, chromosomal

[111133]localization, and expression of alternatively processed transcripts. J.

[111134]Biol. Chem. 270: 17830–17835, 1995.

[111135]19670.

[111136]Chiang, C.–M.; Roeder, R. G.: Cloning of an intrinsic human TFIIID

[111137]subunit that interacts with multiple transcriptional activators. Science 267:

[111138]531–536, 1995.

[111139]19671.

[111140]Wu, Q.; Zhang, T.; Cheng, J.–F.; Kim, Y.; Grimwood, J.;

Schmutz,

[111141]J.; Dickson, M.; Noonan, J. P.; Zhang, M. Q.; Myers, R. M.;

Maniatis,

[111142]T.: Comparative DNA sequence analysis of mouse and human protocadherin

[111143]gene clusters. *Genome Res.* 11: 389–404, 2001.

[111144]19672.

[111145]Kurahashi, H.; Akagi, K.; Inazawa, J.; Ohta, T.; Niikawa, N.; Kayatani,

[111146]F.; Sano, T.; Okada, S.; Nishisho, I.: Isolation and characterization

[111147]of a novel gene deleted in DiGeorge syndrome. *Hum.*

Molec. Genet. 4:

[111148]541–549, 1995.

[111149]19673.

[111150]Aoki, K.; Inazawa, J.; Takahashi, T.; Nakahara, K.; Kasai, M.:

[111151]Genomic structure and chromosomal localization of the gene encoding

[111152]translin, a recombination hotspot binding protein. *Genomics* 43:

[111153]237–241, 1997.

[111154]19674.

[111155]Aoki, K.; Nakahara, K.; Ikegawa, C.; Seto, M.; Takahashi, T.; Minowada,

[111156]J.; Strominger, J. L.; Maziarz, R. T.; Kasai, M.: Nuclear proteins

[111157]binding to a novel target sequence within the recombination hotspot

[111158]regions of Bcl-2 and the immunoglobulin D(H) gene family. Oncogene 9:

[111159]1109-1115, 1994.

[111160]19675.

[111161]Sprecher, E.; Bergman, R.; Szargel, R.; Friedman-Birnbaum, R.;

[111162]Cohen, N.: Identification of a genetic defect in the hairless gene

[111163]in atrichia with papular lesions: evidence for phenotypic heterogeneity

[111164]among inherited atrichias. Am. J. Hum. Genet. 64: 1323-1329, 1999.

[111165]19676.

[111166]Sprecher, E.; Bergman, R.; Szargel, R.; Raz, T.; Labay, V.; Ramon,

[111167]M.; Baruch-Gershoni, R.; Friedman-Birnbaum, R.; Cohen, N.: Atrichia

[111168]with papular lesions maps to 8p in the region containing the human

[111169]hairless gene. Am. J. Med. Genet. 80: 546-550, 1998.

[111170]19677.

[111171]Shannon, R. S.; Mann, J. R.; Harper, E.; Harnden, D. G.; Morten,

[111172]J. E. N.; Herbert, A.: Wilms's tumour and aniridia: clinical and

[111173]cytogenetic features. Arch. Dis. Child. 57: 685-690, 1982.

[111174]19678.

[111175]Strom, T.: A Wilms' tumor family. Acta Paediat. 46: 601-604,

[111176]1957.

[111177]19679.

[111178]Blair, I. P.; Gibson, R. R.; Bennett, C. L.; Chance, P. F.: Search

[111179]for genes involved in Joubert syndrome: evidence that one or more

[111180]major loci are yet to be identified and exclusion of candi-

date genes

[111181]EN1, EN2, FGF8, and BARHL1. Am. J. Med. Genet. 107:
190–196, 2002.

[111182]19680.

[111183]Sweetman, L.; Nyhan, W. L.: Inheritable biotin-treatable
disorders

[111184]and associated phenomena. Ann. Rev. Nutr. 6: 317–343,
1986.

[111185]19681.

[111186]D'Cruz, D.; Taylor, J.; Ahmed, T.; Asherson, R.;
Khamashta, M.;

[111187]Hughes, G. R. V.: Complement factor 2 deficiency: a clinical and

[111188]serological family study. Ann. Rheum. Dis. 51:
1254–1256, 1992.

[111189]19682.

[111190]Day, N. K.; L'Esperance, R.; Good, R. A.; Michael, A. F.;
Hansen,

[111191]J. A.; Dupont, B.; Jersild, C.: Hereditary C2 deficiency: genetic

[111192]studies and association with the HL-A system. J. Exp. Med.
141:

[111193]1464–1469, 1975.

[111194]19683.

[111195]Day, N. K.; Rubinstein, P.; Case, D. J. A.; Good, R. A.; Walker,

[111196]M. E.; Tulchin, N.; Dupont, B.; Jersild, C.: Linkage of gene for

[111197]C2 deficiency and the major histocompatibility complex MHC in man:

[111198]family study of a further case. Vox Sang. 31: 96–102, 1976.

[111199]19684.

[111200]Dewald, G.; Rittner, C.: Polymorphism of the second component

[111201]of human complement (C2): observation of the rare phenotype C2 (=C2

[111202]B) and data on the localization of the C2 locus in the HLA region. Vox

[111203]Sang. 37: 47–54, 1979.

[111204]19685.

[111205]Efthimiou, J.; D'Cruz, D.; Kaplan, P.; Isenberg, D.: Heterozygous

- [111206]C2 deficiency associated with angioedema, myasthenia gravis, and systemic
- [111207]lupus erythematosus. Ann. Rheum. Dis. 45: 428–430, 1986.
- [111208]19686.
- [111209]Einstein, L. P.; Alper, C. A.; Bloch, K. J.; Herrin, J. T.; Rosen, F. S.; David, J. R.; Colten, H. R.: Biosynthetic defect in monocytes from human beings with genetic deficiency of the second component of complement. New Eng. J. Med. 292: 1169–1171, 1975.
- [111212]of complement. New Eng. J. Med. 292: 1169–1171, 1975.
- [111213]19687.
- [111214]Friend, P. S.; Handwerger, B. S.; Kim, Y.; Michael, A. F.; Yunis, E. J.: C2 deficiency in man. Genetic relationship to a mixed lymphocyte reaction determinant (7a*). Immunogenetics 2: 569–576, 1975.
- [111216]reaction determinant (7a*). Immunogenetics 2: 569–576, 1975.
- [111217]19688.
- [111218]Fu, S. M.; Kunkel, H. G.; Brusman, H. P.; Allen, F. H., Jr.; Fotino,

- [111219]M.: Evidence for linkage between HL-A histocompatibility genes and
- [111220]those involved in the synthesis of the second component of complement. J.
- [111221]Exp. Med. 140: 1108-1111, 1974.
- [111222]19689.
- [111223]Gewurz, A.; Lint, T. F.; Roberts, J. L.; Zeitz, H.; Gewurz, H.
- [111224]: Homozygous C2 deficiency with fulminant lupus erythematosus: severe
- [111225]nephritis via the alternative complement pathway. Arthritis Rheum. 21:
- [111226]28-36, 1978.
- [111227]19690.
- [111228]Gibson, D. J.; Glass, D.; Carpenter, C. B.; Schur, P. H.: Hereditary
- [111229]C2 deficiency: diagnosis and HLA gene complex associations. J. Immunogenet. 116:
- [111230]1065-1070, 1976.
- [111231]19691.
- [111232]Johnson, C. A.; Densen, P.; Hurford, R.; Colten, H. R.; Wetzel,

- [111233]R. A.: Deficiency of human complement C2, type I: a 28bp gene deletion
- [111234]leads to a splicing defect. (Abstract) Clin. Res. 39: 304A, 1991.
- [111235]19692.
- [111236]Johnson, C. A.; Densen, P.; Hurford, R. K., Jr.; Colten, H. R.;
- [111237]Wetsel, R. A.: Type I human complement C2 deficiency: a 28-base pair
- [111238]gene deletion causes skipping of exon 6 during RNA splicing. J. Biol.
- [111239]Chem. 267: 9347-9353, 1992.
- [111240]19693.
- [111241]Johnson, C. A.; Densen, P.; Wetsel, R. A.; Cole, F. S.; Goeken,
- [111242]N. E.; Colten, H. R.: Molecular heterogeneity of C2 deficiency. New
- [111243]Eng. J. Med. 326: 871-874, 1992.
- [111244]19694.
- [111245]Kim, Y.; Friend, P. S.; Dresner, I. G.; Yunis, E. J.; Michael,
- [111246]A. F.: Inherited deficiency of the second component of

complement

[111247](C2) with membranoproliferative glomerulonephritis. Am. J. Med. 62:

[111248]765-771, 1977.

[111249]19695.

[111250]Klemperer, M. R.: Hereditary deficiency of the second component

[111251]of complement in man: an immunochemical study. J. Immunogenet. 102:

[111252]168-171, 1969.

[111253]19696.

[111254]Klemperer, M. R.; Austen, K. F.; Rosen, F. S.: Hereditary deficiency

[111255]of second component of complement (C-prime-2) in man: further observations

[111256]on a second kindred. J. Immunogenet. 98: 72-78, 1967.

[111257]19697.

[111258]Klemperer, M. R.; Woodworth, H. C.; Rosen, F. S.; Austen, K. F.

[111259]: Hereditary deficiency of second component of complement (C-prime-2)

[111260]in man. J. Clin. Invest. 45: 880–890, 1966.

[111261]19698.

[111262]Leddy, J. P.; Griggs, R. C.; Klemperer, M. R.; Frank, M. M.:

[111263]Hereditary complement (C2) deficiency with dermato-
myositis. Am. J.

[111264]Med. 58: 83–91, 1975.

[111265]19699.

[111266]Loirat, C.; Levy, M.; Peltier, A. P.; Broyer, M.; Checoury, A.;

[111267]Mathieu, H.: Deficiency of the second component of com-
plement: its

[111268]occurrence with membranoproliferative glomerulonephri-
tis. Arch. Path.

[111269]Lab. Med. 100: 467–472, 1980.

[111270]19700.

[111271]Mahowald, M. L.; Dalmaso, A. P.; Petzel, R. A.; Yunis, E. J.

[111272]: Linkage relationship of C2 deficiency, HLA and glyox-
alase I loci. Vox

[111273]Sang. 37: 321–328, 1979.

[111274]19701.

- [111275]McCarty, D. J.; Tan, E. M.; Zvaifler, N. J.; Koethe, E.;
Duquesnoy,
[111276]R. J.: Serologic studies in a family with heterozygous C2
deficiency. Am.
[111277]J. Med. 71: 945-948, 1981.
[111278]19702.
[111279]Meo, T.; Atkinson, J.; Bernoco, M.; Bernoco, D.; Ceppellini,
R.
[111280]: Mapping of the HLA locus controlling C2 structural vari-
ants and
[111281]linkage disequilibrium between alleles C2-2 and Bw15.
Europ. J. Immun. 6:
[111282]916-919, 1976.
[111283]19703.
[111284]Mortensen, J. P.; Buskjaer, L.; Lamm, L. U.: Studies on the
C2-deficiency
[111285]gene in man. Immunology 39: 541-549, 1980.
[111286]19704.
[111287]Polley, M. J.: Inherited C-prime-2 deficiency in man: lack
of
[111288]immunochemically detectable C-prime-2 protein in

serums from deficient

[111289] individuals. Science 161: 1149–1151, 1968.

[111290] 19705.

[111291] Provost, T. T.; Arnett, F. C.; Reichlin, M.: Homozygous C2 deficiency,

[111292] lupus erythematosus, and anti-Ro (SSA) antibodies.

Arthritis Rheum. 26:

[111293] 1279–1282, 1983.

[111294] 19706.

[111295] Riggs, J. E.; Griggs, R. C.; Rosenfeld, S. I.; May, A. G.; Penn,

[111296] A. S.: Heterozygous C2–deficiency and myasthenia gravis.

Neurology 30:

[111297] 871–873, 1980.

[111298] 19707.

[111299] Ruddy, S.; Austen, K. F.: Inherited abnormalities of the complement

[111300] system in man. Prog. Med. Genet. 7: 69–95, 1970.

[111301] 19708.

[111302] Seligmann, M.; Brouet, J.–C.; Sasportes, M.: Hereditary C2 deficiency

[111303]associated with common variable immunodeficiency. Ann.
Intern. Med. 91:

[111304]216–217, 1979.

[111305]19709.

[111306]Curtis, A.; Nelson, R.; Porteous, M.; Burn, J.; Bhattacharya,

[111307]S. S.: Association of less common cystic fibrosis mutations
with

[111308]a mild phenotype. J. Med. Genet. 28: 34–37, 1991.

[111309]19710.

[111310]Cutting, G. R.; Curristin, S. M.; Nash, E.; Rosenstein, B. J.;

[111311]Lerer, I.; Abeliovich, D.; Hill, A.; Graham, C.: Analysis of
four

[111312]diverse population groups indicates that a subset of cystic
fibrosis

[111313]mutations occur in common among Caucasians. Am. J.
Hum. Genet. 50:

[111314]1185–1194, 1992.

[111315]19711.

[111316]Devoto, M.; Ronchetto, P.; Fanen, P.; Orriols, J. J. T.;

Romeo,

[111317]G.; Goossens, M.; Ferrari, M.; Magnani, C.; Seia, M.; Cremonesi, L.

[111318]Screening for non-delta-F508 mutations in five exons of the cystic

[111319]fibrosis transmembrane conductance regulator (CFTR) gene in Italy. Am.

[111320]J. Hum. Genet. 48: 1127-1132, 1991.

[111321]19712.

[111322]de Vries, H. G.; Collee, J. M.; de Walle, H. E. K.; van Veldhuizen,

[111323]M. H. R.; Smit Sibinga, C. T.; Scheffer, H.; ten Kate, L. P.: Prevalence

[111324]of delta-F508 cystic fibrosis carriers in The Netherlands: logistic

[111325]regression on sex, age, region of residence and number of offspring. Hum.

[111326]Genet. 99: 74-79, 1997.

[111327]19713.

[111328]Dorin, J. R.; Dickinson, P.; Alton, E. W. F. W.; Smith, S. N.;

[111329]Geddes, D. M.; Stevenson, B. J.; Kimber, W. L.; Fleming, S.; Clarke,

- [111330]A. R.; Hooper, M. L.; Anderson, L.; Beddington, R. S. P.;
Porteous,
- [111331]D. J.: Cystic fibrosis in the mouse by targeted insertional
mutagenesis. *Nature* 359:
- [111332]211–215, 1992.
- [111333]19714.
- [111334]Dork, T.; El-Harith, E.-H. A.; Stuhmann, M.; Macek, M.,
Jr.;
- [111335]Egan, M.; Cutting, G. R.; Tzetzis, M.; Kanavakis, E.; Carles,
S.; Claustres,
- [111336]M.; Padoa, C.; Ramsay, M.; Schmidtke, J.: Evidence for a
common ethnic
- [111337]origin of cystic fibrosis mutation 3120+1G-to-A in di-
verse populations.
- [111338](Letter) *Am. J. Hum. Genet.* 63: 656–662, 1998.
- [111339]19715.
- [111340]Dork, T.; Wulbrand, U.; Richter, T.; Neumann, T.; Wolfes,
H.;
- [111341]Wulf, B.; Maass, G.; Tummeler, B.: Cystic fibrosis with three
mutations
- [111342]in the cystic fibrosis transmembrane conductance regula-
tor gene. *Hum.*

[111343]Genet. 87: 441–446, 1991.

[111344]19716.

[111345]Dumur, V.; Lafitte, J. J.; Gervais, R.; Debaecker, D.;
Kesteloot,

[111346]M.; Lalau, G.; Roussel, P.: Abnormal distribution of cystic
fibrosis

[111347]delta-F508 allele in adults with chronic bronchial hyper-
secretion. Lancet 335:

[111348]1340, 1990.

[111349]19717.

[111350]Ferec, C.; Audrezet, M. P.; Mercier, B.; Guillermit, H.;
Moullier,

[111351]P.; Quere, I.; Verlingue, C.: Detection of over 98% cystic fi-
brosis

[111352]mutations in a Celtic population. Nature Genet. 1:
188–191, 1992.

[111353]19718.

[111354]Kelly, T. E.: Personal Communication. Charlottesville, Va.

[111355]1977.

[111356]19719.

- [111357]Lindor, N. M.; Furuichi, Y.; Kitao, S.; Shimamoto, A.; Arndt,
[111358]C.; Jalal, S.: Rothmund–Thomson syndrome due to RECQ4
helicase mutations:
[111359]report and clinical and molecular comparisons with Bloom
syndrome
[111360]and Werner syndrome. Am. J. Med. Genet. 90: 223–228,
2000.
[111361]19720.
- [111362]Bodzioch, M.; Orso, E.; Klucken, J.; Langmann, T.;
Bottcher, A.;
[111363]Diederich, W.; Drobnik, W.; Barlage, S.; Buchler, C.;
Porsch–Ozcurumez,
[111364]M.; Kaminski, W. E.; Hahmann, H. W.; Oette, K.; Rothe, G.;
Aslanidis,
[111365]C.; Lackner, K. J.; Schmitz, G.: The gene encoding ATP–
binding cassette
[111366]transporter 1 is mutated in Tangier disease. Nature Genet.
22: 347–351,
[111367]1999.
[111368]19721.
- [111369]Brooks–Wilson, A.; Marcil, M.; Clee, S. M.; Zhang, L.–H.;
Roomp,

- [111370]K.; van Dam, M.; Yu, L.; Brewer, C.; Collins, J. A.; Molhuizen, H.
- [111371]O. F.; Loubser, O.; Ouelette, B. F. F.; and 14 others: Mutations
- [111372]in ABC1 in Tangier disease and familial high-density lipoprotein deficiency. Nature
- [111373]Genet. 22: 336–345, 1999.
- [111374]19722.
- [111375]Hoogendijk, J. E.; Hensels, G. W.; Gabreels–Festen, A. A. W. M.;
- [111376]Gabreels, F. J. M.; Janssen, E. A. M.; De Jonghe, P.; Martin, J.–J.;
- [111377]Van Broeckhoven, C.; Valentijn, L. J.; Baas, F.; de Visser, M.; Bolhuis,
- [111378]P. A.: De–novo mutation in hereditary motor and sensory neuropathy
- [111379]type I. Lancet 339: 1081–1082, 1992.
- [111380]19723.
- [111381]Menasche, G.; Pastural, E.; Feldmann, J.; Certain, S.; Ersoy, F.;
- [111382]Dupuis, S.; Wulffraat, N.; Bianchi, D.; Fischer, A.; Le Deist, F.;

[111383] de Saint Basile, G.: Mutations in RAB27A cause Griscelli syndrome associated with haemophagocytic syndrome. *Nature Genet.* 25: 173–176, 2000.

[111385] 19724.

[111386] 19724.

[111387] Pastural, E.; Barrat, F. J.; Dufourcq-Lagelouse, R.; Certain, S.;

[111388] Sanal, O.; Jabado, N.; Seger, R.; Griscelli, C.; Fischer, A.; de Saint

[111389] Basile, G.: Griscelli disease maps to chromosome 15q21 and is associated with mutations in the myosin-Va gene. *Nature Genet.* 16: 289–292, 1997.

[111391] 19725.

[111392] 19725.

[111393] Asamoah, A.; Wilson, A. F.; Elston, R. C.; Dalferes, E., Jr.; Berenson,

[111394] G. S.: Segregation and linkage analyses of dopamine-beta-hydroxylase activity in a six-generation pedigree. *Am. J. Med. Genet.* 27: 613–621,

[111396]1987.

[111397]19726.

[111398]Biaggioni, I.; Goldstein, D. S.; Atkinson, T.; Robertson, D.:

[111399]Dopamine-beta-hydroxylase deficiency in humans. Neu-
rology 40: 370-373,

[111400]1990.

[111401]19727.

[111402]Biaggioni, I.; Robertson, D.: Endogenous restoration of
noradrenaline

[111403]by precursor therapy in dopamine-beta-hydroxylase defi-
ciency. Lancet II:

[111404]1170-1172, 1987.

[111405]19728.

[111406]Craig, S. P.; Buckle, V. J.; Lamouroux, A.; Mallet, J.; Craig,

[111407]I. W.: Localization of the human dopamine beta hydroxy-
lase (DBH)

[111408]gene to chromosome 9q34. Cytogenet. Cell Genet. 48:
48-50, 1988.

[111409]19729.

[111410]Dunnette, J.; Weinshilboum, R.: Human serum dopamine

beta-hydroxylase:

[111411]correlation of enzymatic activity with immunoreactive protein in genetically

[111412]defined samples. Am. J. Hum. Genet. 28: 155-166, 1976.

[111413]19730.

[111414]Dunnette, J.; Weinshilboum, R.: Inheritance of low immunoreactive

[111415]human plasma dopamine-beta-hydroxylase: radioimmunoassay studies. J.

[111416]Clin. Invest. 60: 1080-1087, 1977.

[111417]19731.

[111418]Elston, R. C.; Namboodiri, K. K.; Hames, C. G.: Segregation and

[111419]linkage analysis of dopamine-beta-hydroxylase activity. Hum. Hered. 29:

[111420]284-292, 1979.

[111421]19732.

[111422]Gershon, E. S.; Goldin, L. R.: Segregation and linkage studies

[111423]of plasma dopamine-beta-hydroxylase (DBH), erythrocyte catechol-O-methyltransferase

[111424](COMT) and platelet monoamine oxidase (MAO): possible linkage between

[111425]the ABO locus and a gene controlling DBH activity.

(Abstract) Am.

[111426]J. Hum. Genet. 33: 136A only, 1981.

[111427]19733.

[111428]Goldin, L. R.; Gershon, E. S.; Lake, C. R.; Murphy, D. L.; McGinniss,

[111429]M.; Sparkes, R. S.: Segregation and linkage studies of plasma dopamine-beta-hydroxylase

[111430](DBH), erythrocyte catechol-O-methyltransferase (COMT), and platelet

[111431]monoamine oxidase (MAO): possible linkage between the ABO locus and

[111432]a gene controlling DBH activity. Am. J. Hum. Genet. 34: 250-262,

[111433]1982.

[111434]19734.

[111435]Joh, T. H.; Baetge, E. E.; Reis, D. J.: Evidence for the existence

[111436]of a single gene or linked genes coding for catecholamine biosynthetic

[111437]enzymes. Trans. Assoc. Am. Phys. 96: 38–43, 1983.

[111438]19735.

[111439]Joh, T. H.; Baetge, E. E.; Ross, M. E.; Albert, V. R.; Moon, H.

[111440]M.; Reis, D. J.: Existence of catecholamine biosynthetic
enzyme gene

[111441]family. (Abstract) Clin. Res. 31: 528 only, 1983.

[111442]19736.

[111443]Joh, T. H.; Baetge, E. E.; Ross, M. E.; Reis, D. J.: Biochem-
istry

[111444]and molecular biology of catecholamine neurons: a single
gene or gene

[111445]family hypothesis. Clin. Exp. Hypertension 6A: 11–21,
1984.

[111446]19737.

[111447]Kobayashi, K.; Kurosawa, Y.; Fujita, K.; Nagatsu, T.: Hu-
man dopamine

[111448]beta-hydroxylase gene: two mRNA types having different
3-prime-terminal

[111449]regions are produced through alternative polyadenylation.
Nucleic

[111450]Acids Res. 17: 1089–1102, 1989.

[111451]19738.

[111452]Lamouroux, A.; Vigny, A.; Faucon Biguet, N.; Darmon, M.
C.; Franck,

[111453]R.; Henry, J.-P.; Mallet, J.: The primary structure of human
dopamine-beta-hydroxylase:

[111454]insights into the relationship between the soluble and the
membrane-bound

[111455]forms of the enzyme. EMBO J. 6: 3931-3937, 1987.

[111456]19739.

[111457]Lea, R. A.; Dohy, A.; Jordan, K.; Quinlan, S.; Brimage, P. J.;

[111458]Griffiths, L. R.: Evidence for allelic association of the
dopamine

[111459]beta-hydroxylase gene (DBH) with susceptibility to typical
migraine. Neurogenetics 3:

[111460]35-40, 2000.

[111461]19740.

[111462]Mathias, C. J.; Bannister, R. B.; Cortelli, P.; Heslop, K.; Po-
lak,

[111463]J. M.; Raimbach, S.; Springall, D. R.; Watson, L.: Clinical,
autonomic

[111464]and therapeutic observations in two siblings with postural

hypotension

[111465]and sympathetic failure due to an inability to synthesize
noradrenaline

[111466]from dopamine because of a deficiency of dopamine beta
hydroxylase. Quart.

[111467]J. Med. 75: 617-633, 1990.

[111468]19741.

[111469]McKinney, E. F.; Walton, R. T.; Yudkin, P.; Fuller, A.; Hal-
dar,

[111470]N. A.; Mant, D.; Murphy, M.; Welsh, K. I.; Marshall, S. E.:
Association

[111471]between polymorphisms in dopamine metabolic enzymes
and tobacco consumption

[111472]in smokers. Pharmacogenetics 10: 483-491, 2000.

[111473]19742.

[111474]O'Malley, K. L.; Mauron, A.; Raese, J.; Barchas, J. D.; Kedes,

[111475]L.: Genes for catecholamine biosynthesis: cloning by ex-
pression and

[111476]identification of the cDNA for rat dopamine beta-
hydroxylase. Proc.

[111477]Nat. Acad. Sci. 80: 2161-2165, 1983.

[111478]19743.

[111479]Ogihara, T.; Nugent, C. A., Jr.; Shen, S.-W.; Goldfein, S.:
Serum

[111480]dopamine-beta-hydroxylase activity in parents and chil-
dren. J. Lab.

[111481]Clin. Med. 85: 566-573, 1975.

[111482]19744.

[111483]Perry, S. E.; Summar, M. L.; Phillips, J. A., III; Robertson,

[111484]D.: Linkage analysis of the human dopamine beta-
hydroxylase gene. Genomics 10:

[111485]493-495, 1991.

[111486]19745.

[111487]Robertson, D.; Haile, V.; Perry, S. E.; Robertson, R. M.;
Phillips,

[111488]J. A., III; Biaggioni, I.: Dopamine beta-hydroxylase defi-
ciency:

[111489]a genetic disorder of cardiovascular regulation. Hyperten-
sion 18:

[111490]1-8, 1991.

[111491]19746.

- [111492]Ross, S. B.; Wetterberg, L.; Myrhed, M.: Genetic control of plasma
- [111493]dopamine-beta-hydroxylase. Life Sci. 12: 529-532, 1973.
- [111494]19747.
- [111495]Schanberg, S. M.; Stone, R. A.; Kirshner, N.; Gunnells, J. C.;
- [111496]Robinson, R. R.: Plasma dopamine beta-hydroxylase: a possible aid
- [111497]in the study and evaluation of hypertension. Science 183: 523-525,
- [111498]1974.
- [111499]19748.
- [111500]Hoefsloot, L. H.; van der Ploeg, A. T.; Kroos, M. A.; Hoogeveen-Westerveld,
- [111501]M.; Oostra, B. A.; Reuser, A. J. J.: Adult and infantile glycogenosis
- [111502]type II in one family, explained by allelic diversity. Am. J. Hum.
- [111503]Genet. 46: 45-52, 1990.
- [111504]19749.
- [111505]de Villiers, J. N. P.; Hillermann, R.; Loubser, L.; Kotze, M.

[111506]J.: Spectrum of mutations in the HFE gene implicated in
haemochromatosis

[111507]and porphyria. Hum. Molec. Genet. 8: 1517–1522, 1999.

[111508]19750.

[111509]Marlar, R. A.; Griffin, J. H.: Deficiency of protein C inhibitor

[111510]in combined factor V/VIII deficiency disease. J. Clin. In-
vest. 66:

[111511]1186–1189, 1980.

[111512]19751.

[111513]Meijers, J. C.; Chung, D. W.: Organization of the gene cod-
ing

[111514]for human protein C inhibitor (plasminogen activator in-
hibitor-3):

[111515]assignment of the gene to chromosome 14. J. Biol. Chem.
266: 15028–15034,

[111516]1991.

[111517]19752.

[111518]Nichols, W. C.; Seligsohn, U.; Zivelin, A.; Terry, V. H.;
Arnold,

[111519]N. D.; Siemieniak, D. R.; Kaufman, R. J.; Ginsburg, D.:
Linkage of

[111520]combined factors V and VIII deficiency to chromosome
18q by homozygosity

[111521]mapping. J. Clin. Invest. 99: 596–601, 1997.

[111522]19753.

[111523]Suzuki, K.; Deyashiki, Y.; Nishioka, J.; Kurachi, K.; Akira,
M.;

[111524]Yamamoto, S.; Hashimoto, S.: Characterization of a cDNA
for human

[111525]protein C inhibitor: a new member of the plasma serine
protease inhibitor

[111526]superfamily. J. Biol. Chem. 262: 611–616, 1987.

[111527]19754.

[111528]Abramson, R. D.; Barbosa, P.; Kalumuck, K.; O'Brien, W. E.:
Characterization

[111529]of the human argininosuccinate lyase gene and analysis of
exon skipping. Genomics 10:

[111530]126–132, 1991.

[111531]19755.

[111532]Allan, J. D.; Cusworth, D. C.; Dent, C. E.; Wilson, V. K.: A
disease,

[111533]probably hereditary, characterized by severe mental defi-

ciency and

[111534]a constant gross abnormality of amino acid metabolism.

Lancet I:

[111535]182–187, 1958.

[111536]19756.

[111537]Asai, K.; Ishii, S.; Ohta, S.; Furusho, K.: Fatal hyperam-
monaemia

[111538]in argininosuccinic aciduria following enflurane anaesthe-
sia. (Letter) Europ.

[111539]J. Paediat. 157: 169–170, 1997.

[111540]19757.

[111541]Bohles, H.; Heid, H.; Harms, D.; Schmid, D.; Fekl, W.:
Argininosuccinic

[111542]aciduria: metabolic studies and effects of treatment with
keto-analogues

[111543]of essential amino acids. Europ. J. Pediat. 128: 225–233,
1978.

[111544]19758.

[111545]Brusilow, S. W.; Batshaw, M. L.: Arginine therapy of argini-
nosuccinase

[111546]deficiency. Lancet I: 124–127, 1979.

[111547]19759.

[111548]Cathelineau, L.; Dinh, D. P.; Briand, P.; Kamoun, P.: Studies

[111549]on complementation in argininosuccinate synthetase and
argininosuccinate

[111550]lyase deficiencies in human fibroblasts. Hum. Genet. 57:
282-284,

[111551]1981.

[111552]19760.

[111553]Collins, F. S.; Summer, G. K.; Schwartz, R. P.; Parke, J. C.,
Jr.

[111554]: Neonatal argininosuccinic aciduria--survival after early
diagnosis

[111555]and dietary management. J. Pediat. 96: 429-431, 1980.

[111556]19761.

[111557]Coryell, M. E.; Hall, W. K.; Thevaos, T. G.; Welter, D. A.;
Gatz,

[111558]A. J.; Horton, B. F.; Sisson, B. D.; Looper, J. W., Jr.; Farrow,
R.

[111559]T.: Familial study of human enzyme defect, argininosuc-
cinic aciduria. Biochem.

[111560]Biophys. Res. Commun. 14: 307-312, 1964.

[111561]19762.

[111562]Fleisher, L. D.; Rassin, D. K.; Desnick, R. J.; Salwen, H. R.;

[111563]Rogers, P.; Bean, M.; Gaull, G. E.: Argininosuccinic
aciduria: prenatal

[111564]studies in a family at risk. Am. J. Hum. Genet. 31:
439-445, 1979.

[111565]19763.

[111566]Glick, N. R.; Snodgrass, P. J.; Schafer, I. A.: Neonatal
argininosuccinic

[111567]aciduria with normal brain and kidney but absent liver
argininosuccinate

[111568]lyase activity. Am. J. Hum. Genet. 28: 22-30, 1976.

[111569]19764.

[111570]Goodman, S. I.; Mace, J. W.; Turner, B.; Garrett, W. J.: An-
tenatal

[111571]diagnosis of argininosuccinic aciduria. Clin. Genet. 4:
236-240,

[111572]1973.

[111573]19765.

[111574]Kint, J. A.; Carton, D.: Deficient argininosuccinase activity

[111575]in brain in argininosuccinicaciduria. (Letter) Lancet II: 635
only,
[111576]1968.
[111577]19766.
[111578]Kvedar, J. C.; Baden, H. P.; Baden, L. A.; Shih, V. E.;
Kolodny,
[111579]E. H.: Dietary management reverses grooving and abnormal polarization
[111580]of hair shafts in argininosuccinase deficiency. Am. J. Med.
Genet. 40:
[111581]211-213, 1991.
[111582]19767.
[111583]Levin, B.: Argininosuccinic aciduria. Am. J. Dis. Child. 113:
[111584]162-165, 1967.
[111585]19768.
[111586]Levin, B.; MacKay, H. M.; Oberholzer, V. G.: Argininosuccinic
[111587]aciduria: an inborn error of amino acid metabolism. Arch.
Dis. Child. 36:
[111588]622-632, 1961.

[111589]19769.

[111590]Lewis, P. D.; Miller, A. L.: Argininosuccinic aciduria: case

[111591]report with neuropathological findings. Brain 93:

413-422, 1970.

[111592]19770.

[111593]Matuo, S.; Tatsuno, M.; Kobayashi, K.; Saheki, T.; Miyata,

T.;

[111594]Iwanaga, S.; Amaya, Y.; Mori, M.: Isolation of cDNA clones

of human

[111595]argininosuccinate lyase and corrected amino acid se-

quence. FEBS Lett. 234:

[111596]395-399, 1988.

[111597]19771.

[111598]McInnes, R. R.; Shih, V.; Chilton, S.: Interallelic comple-

mentation

[111599]in an inborn error of metabolism: genetic heterogeneity in

argininosuccinate

[111600]lyase deficiency. Proc. Nat. Acad. Sci. 81: 4480-4484,

1984.

[111601]19772.

[111602]Moser, H. W.; Efron, M. L.; Brown, H.; Diamond, R.; Neu-

mann, C.

[111603]G.: Argininosuccinic aciduria: report of two cases and demonstration

[111604]of intermittent elevation of blood ammonia. Am. J. Med. 42: 9–26,

[111605]1967.

[111606]19773.

[111607]Sinclair, D. A.; Mills, K.; Guarente, L.: Accelerated aging and

[111608]nucleolar fragmentation in yeast sgs1 mutants. Science. 277: 1313–1316,

[111609]1997.

[111610]19774.

[111611]Koster, J. F.; Busch, H. F. M.; Slee, R. G.; van Weerden, T. W.

[111612]; Glycogenosis type II: the infantile– and late–onset acid maltase

[111613]deficiency observed in one family. Clin. Chim. Acta 87: 451–453,

[111614]1978.

[111615]19775.

- [111616] Kroos, M. A.; Van der Kraan, M.; Van Diggelen, O. P.; Kleijer,
[111617] W. J.; Reuser, A. J. J.: Two extremes of the clinical spectrum of
[111618] glycogen storage disease type II in one family: a matter of
genotype. Hum.
[111619] Mutat. 9: 17–22, 1997.
[111620] 19776.
[111621] Kroos, M. A.; Van der Kraan, M.; Van Diggelen, O. P.; Kleijer,
[111622] W. J.; Reuser, A. J. J.; Van den Boogaard, M. J.; Ausems, M.
G. E.
[111623] M.; Ploos van Amstel, H. K.; Poenaru, L.; Nicolino, M.;
Wevers, R.
[111624]: Glycogen storage disease type II: frequency of three
common mutant
[111625] alleles and their associated clinical phenotypes studied in
121 patients. J.
[111626] Med. Genet. 32: 836–837, 1995.
[111627] 19777.
[111628] Loonen, M. C. B.; Schram, A. W.; Koster, J. F.; Niermeijer,
M.

[111629]F.; Busch, H. F. M.; Martin, J. J.; Brouwer-Kelder, B.; Mekes, W.;

[111630]Slee, R. G.; Tager, J. M.: Identification of heterozygotes for glycogenosis

[111631]2 (acid maltase deficiency). Clin. Genet. 19: 55-63, 1981.

[111632]19778.

[111633]Martiniuk, F.; Bodkin, M.; Tzall, S.; Hirschhorn, R.: Identification

[111634]of the base-pair substitution responsible for a human acid alpha glucosidase

[111635]allele with lower 'affinity' for glycogen (GAA 2) and transient gene

[111636]expression in deficient cells. Am. J. Hum. Genet. 47: 440-445, 1990.

[111637]19779.

[111638]Martiniuk, F.; Chen, A.; Mack, A.; Arvanitopoulos, E.; Chen, Y.;

[111639]Rom, W. N.; Codd, W. J.; Hanna, B.; Alcabes, P.; Raben, N.; Plotz,

[111640]P.: Carrier frequency for glycogen storage disease type II in New

[111641]York and estimates of affected individuals born with the

disease.

[111642](Letter) Am. J. Med. Genet. 79: 69–72, 1998.

[111643]19780.

[111644]Diamond, T.; Stiel, D.; Posen, S.: Osteoporosis in
hemochromatosis:

[111645]iron excess, gonadal deficiency, or other factors? Ann. In-
tern. Med. 110:

[111646]430–436, 1989.

[111647]19781.

[111648]Dokal, I.; Lord, D.; Rhodes, D.; Bydder, G.; Cox, T.: Detec-
tion

[111649]of hereditary haemochromatosis in an HLA-identical pedi-
gree showing

[111650]discordance between HLA class I genes and the disease
locus. Hum.

[111651]Genet. 88: 209–214, 1991.

[111652]19782.

[111653]Dugast, I. J.; Papadopoulos, P.; Zappone, E.; Jones, C.;
Theriault,

[111654]K.; Handelman, G. J.; Benarous, R.; Drysdale, J. W.: Identi-
fication

[111655] of two human ferritin H genes on the short arm of chromosome 6. Genomics 6:

[111656] 204–211, 1990.

[111657] 19783.

[111658] Edwards, C. Q.; Carroll, M.; Bray, P. F.; Cartwright, G. E.:

[111659] Hereditary hemochromatosis: diagnosis in siblings and children. New

[111660] Eng. J. Med. 297: 7–13, 1977.

[111661] 19784.

[111662] Edwards, C. Q.; Cartwright, G. E.; Skolnick, M. H.; Amos, D. B.

[111663]: Homozygosity for hemochromatosis: clinical manifestations. Ann.

[111664] Intern. Med. 93: 519–525, 1980.

[111665] 19785.

[111666] Edwards, C. Q.; Griffen, L. M.; Dadone, M. M.; Skolnick, M. H.;

[111667] Kushner, J. P.: Mapping the locus for hereditary hemochromatosis:

[111668] localization between HLA-B and HLA-A. Am. J. Hum. Genet. 38: 805–811,

[111669]1986.

[111670]19786.

[111671]Edwards, C. Q.; Griffen, L. M.; Dadone, M. M.; Skolnick, M.
H.;

[111672]Kushner, J. P.: The locus for hereditary hemochromatosis
maps between

[111673]HLA-A and HLA-B. (Abstract) Cytogenet. Cell Genet. 40:
620, 1985.

[111674]19787.

[111675]Edwards, C. Q.; Griffen, L. M.; Goldgar, D.; Drummond, C.;
Skolnick,

[111676]M. H.; Kushner, J. P.: Prevalence of hemochromatosis
among 11,065

[111677]presumably healthy blood donors. New Eng. J. Med. 318:
1355-1362,

[111678]1988.

[111679]19788.

[111680]Edwards, C. Q.; Skolnick, M. H.; Kushner, J. P.: Hereditary
hemochromatosis:

[111681]contributions of genetic analyses. Prog. Hemat. 12:
43-71, 1981.

[111682]19789.

[111683]Edwards, J. H.: Allelic association in man.In: Eriksson, A.
W.;

[111684]Forsius, H. R.; Nevanlinna, H. R.; Workman, P. L.; Norio, R.
K. (eds.)

[111685]: Population Structure and Genetic Disorders: Seventh
Sigrid Juselius

[111686]Foundation Symposium: Mariehamn, Aland Islands, Fin-
land, August 1978.

[111687]London: Academic Press 1980. Pp. 239-255.

[111688]19790.

[111689]El Kahloun, A.; Chauvel, B.; Mauvieux, V.; Dorval, I.;
Jouanolle,

[111690]A.-M.; Gicquel, I.; Le Gall, J.-Y.; David, V.: Localization of
seven

[111691]new genes around the HLA-A locus. Hum. Molec. Genet.
2: 55-60, 1993.

[111692]19791.

[111693]Elzouki, A.-N. Y.; Hultcrantz, R.; Stal, P.; Befrits, R.; Eriks-
son,

[111694]S.: Increased PiZ gene frequency for alpha(1) antitrypsin in
patients

[111695]with genetic haemochromatosis. Gut 36: 922–926, 1995.

[111696]19792.

[111697]Eriksson, S.; Lindmark, B.; Hanik, L.: A Swedish family with

[111698]alpha-1-antitrypsin deficiency, haemochromatosis,
haemoglobinopathy

[111699]D and early death in liver cirrhosis. J. Hepatol. 2: 65–72,
1986.

[111700]19793.

[111701]Escobar, G. J.; Heyman, M. B.; Smith, W. B.; Thaler, M. M.:
Primary

[111702]hemochromatosis in childhood. Pediatrics 80: 549–554,
1987.

[111703]19794.

[111704]European Haemochromatosis Consortium: Polymorphism
in intron

[111705]4 of HFE does not compromise haemochromatosis muta-
tion results. (Letter) Nature

[111706]Genet. 23: 271 only, 1999.

[111707]19795.

[111708]Feder, J. N.; Penny, D. M.; Irrinki, A.; Lee, V. K.; Lebron, J.

- [111709]A.; Watson, N.; Tsuchihashi, Z.; Sigal, E.; Bjorkman, P. J.;
Schatzman,
- [111710]R. C.: The hemochromatosis gene product complexes with
the transferrin
- [111711]receptor and lowers its affinity for ligand binding. Proc.
Nat. Acad.
- [111712]Sci. 95: 1472–1477, 1998.
- [111713]19796.
- [111714]Feder, J. N.; Tsuchihashi, Z.; Irrinki, A.; Lee, V. K.; Mapa,
- [111715]F. A.; Morikang, E.; Prass, C. E.; Starnes, S. M.; Wolff, R. K.;
- Parkkila,
- [111716]S.; Sly, W. S.; Schatzman, R. C. :J. Biol. Chem. 272:
14025–14028,
- [111717]1997.
- [111718]19797.
- [111719]Feller, E. R.; Pont, A.; Wands, J. R.; Carter, E. A.; Foster,
- [111720]G.; Kourides, I. A.; Isselbacher, K. J.: Familial hemochro-
matosis:
- [111721]physiologic studies in the precirrhotic stage of the dis-
ease. New
- [111722]Eng. J. Med. 296: 1422–1426, 1977.

[111723]19798.

[111724]Felts, J. H.; Nelson, J. R.; Herndon, C. N.; Spurr, C. L.:
Hemochromatosis

[111725]in two young sisters: case studies and a family survey.
Ann. Intern.

[111726]Med. 67: 117–123, 1967.

[111727]19799.

[111728]Feder, J. N.; Gnirke, A.; Thomas, W.; Tsuchihashi, Z.;
Ruddy,

[111729]D. A.; Basava, A.; Dormishian, F.; Domingo, R., Jr.; Ellis, M.
C.;

[111730]Fullan, A.; Hinton, L. M.; Jones, N. L.; Kimmel, B. E.; Kron-
mal, G.

[111731]S.; Lauer, P.; Lee, V. K.; Loeb, D. B.; Mapa, F. A.; McClel-
land, E.;

[111732]Meyer, N. C.; Mintier, G. A.; Moeller, N.; Moore, T.;
Morikang, E.;

[111733]Prass, C. E.; Quintana, L.; Starnes, S. M.; Schatzman, R. C.;
Brunke,

[111734]K. J.; Drayna, D. T.; Risch, N. J.; Bacon, B. R.; Wolff, R. K.: A

[111735]novel MHC class I-like gene is mutated in patients with
hereditary

[111736]haemochromatosis. Nature Genet. 13: 399–408, 1996.

[111737]19800.

[111738]Fleming, R. E.; Holden, C. C.; Tomatsu, S.; Waheed, A.;
Brunt,

[111739]E. M.; Britton, R. S.; Bacon, B. R.; Roopenian, D. C.; Sly, W.
S.

[111740] Mouse strain differences determine severity of iron accu-
mulation

[111741]in Hfe knockout model of hereditary hemochromatosis.
Proc. Nat. Acad.

[111742]Sci. 98: 2707–2711, 2001.

[111743]19801.

[111744]Fleming, R. E.; Migas, M. C.; Zhou, X. Y.; Jiang, J.; Britton,

[111745]R. S.; Brunt, E. M.; Tomatsu, S.; Waheed, A.; Bacon, B. R.;
Sly, W.

[111746]S.: Mechanism of increased iron absorption in murine
model of hereditary

[111747]hemochromatosis: increased duodenal expression of the
iron transporter

[111748]DMT1. Proc. Nat. Acad. Sci. 96: 3143–3148, 1999.

[111749]19802.

[111750]Goei, V. L.; Parimoo, S.; Capossela, A.; Chu, T. W.; Gruen, J.

[111751]R.: Isolation of novel non-HLA gene fragments from the hemochromatosis

[111752]region (6p21.3) by cDNA hybridization selection. Am. J. Hum. Genet. 54:

[111753]244-251, 1994.

[111754]19803.

[111755]Gomez, P. S.; Parks, S.; Ries, R.; Tran, T. C.; Gomez, P. F.;

[111756]Press, R. D.: Polymorphism in intron 4 of HFE does not compromise

[111757]haemochromatosis mutation results (Letter) Nature Genet. 23: 272

[111758]only, 1999.

[111759]19804.

[111760]Gordeuk, V.; Mukiibi, J.; Hasstedt, S.; Samowitz, W.; Edwards,

[111761]C. Q.; West, G.; Ndambire, S.; Emmamial, J.; Nkanza, N.; Chapanduka,

[111762]Z.; Randall, M.; Boone, P.; Romano, P.; Martell, R. W.; Yamashita,

[111763]T.; Effler, P.; Brittenham, G. :New Eng. J. Med. 326:

95–100,

[111764]1992.

[111765]19805.

[111766]Griffiths, W.; Cox, T.: Haemochromatosis: novel gene discovery

[111767]and the molecular pathophysiology of iron metabolism.
Hum. Molec.

[111768]Genet. 9: 2377–2382, 2000.

[111769]19806.

[111770]Grove, J.; Daly, A. K.; Burt, A. D.; Guzail, M.; James, O. F.

[111771]W.; Bassendine, M. F.; Day, C. P.: Heterozygotes for HFE mutations

[111772]have no increased risk of advanced alcoholic liver disease.
Gut 43:

[111773]262–266, 1998.

[111774]19807.

[111775]Halliday, J. W.; Cowlshaw, J. L.; Russo, A. M.; Powell, L. W.

[111776]: Serum–ferritin in diagnosis of haemochromatosis: a study of 43 families. Lancet II:

[111777]621-624, 1977.

[111778]19808.

[111779]Hardy, L.; Hansen, J. L.; Kushner, J. P.; Knisely, A. S.:

Neonatal

[111780]hemochromatosis: genetic analysis of transferrin-receptor, H-apoferritin,

[111781]and L-apoferritin loci and of the human leukocyte antigen class I

[111782]region. Am. J. Path. 137: 149-153, 1990.

[111783]19809.

[111784]Beattie, E. C.; Stellwagen, D.; Morishita, W.; Bresnahan, J. C.;

[111785]Ha, B. K.; Von Zastrow, M.; Beattie, M. S.; Malenka, R. C.: Control

[111786]of synaptic strength by glial TNF-alpha. Science 295: 2282-2285,

[111787]2002.

[111788]19810.

[111789]Beutler, B.; Krochin, N.; Milsark, I. W.; Luedke, C.; Cerami, A.

[111790]: Control of cachectin (tumor necrosis factor) synthesis:

mechanisms

[111791] of endotoxin resistance. Science 232: 977–980, 1986.

[111792] 19811.

[111793] Brenner, D. A.; O'Hara, M.; Angel, P.; Chojkier, M.; Karin, M.

[111794]; Prolonged activation of JUN and collagenase genes by tumour necrosis

[111795] factor- α . Nature 337: 661–663, 1989.

[111796] 19812.

[111797] Broudy, V. C.; Kaushansky, K.; Segal, G. M.; Harlan, J. M.; Adamson,

[111798] J. W.: Tumor necrosis factor type α stimulates human endothelial

[111799] cells to produce granulocyte/macrophage colony-stimulating factor. Proc.

[111800] Nat. Acad. Sci. 83: 7467–7471, 1986.

[111801] 19813.

[111802] Bruce, A. J.; Boling, W.; Kindy, M. S.; Peschon, J.; Kraemer, P.

[111803] J.; Carpenter, M. K.; Holtsberg, F. W.; Mattson, M. P.: Altered neuronal

- [111804]and microglial responses to excitotoxic and ischemic
brain injury
- [111805]in mice lacking TNF receptors. Nature Med. 2: 788–794,
1996.
- [111806]19814.
- [111807]Cabrera, M.; Shaw, M. A.; Sharples, C.; Williams, H.;
Castes, M.;
- [111808]Convit, J.; Blackwell, J. M.: Polymorphism in tumor necro-
sis factor
- [111809]genes associated with mucocutaneous leishmaniasis. J.
Exp. Med. 182:
- [111810]1259–1264, 1995.
- [111811]19815.
- [111812]Conway, D. J.; Holland, M. J.; Bailey, R. L.; Campbell, A. E.;
- [111813]Mahdi, O. S.; Jennings, R.; Mbena, E.; Mabey, D. C.: Scar-
ring trachoma
- [111814]is associated with polymorphism in the tumor necrosis
factor alpha
- [111815](TNF- α) gene promoter and with elevated TNF- α levels in tear
- [111816]fluid. Infect. Immun. 65: 1003–1006, 1997.

[111817]19816.

[111818]Davis, J. M.; Narachi, M. A.; Alton, N. K.; Arakawa, T.:
Structure

[111819]of human tumor necrosis factor alpha derived from re-
combinant DNA. Biochemistry 26:

[111820]1322-1326, 1987.

[111821]19817.

[111822]Escobar-Morreale, H. F.; Calvo, R. M.; Sancho, J.; San Mil-
lan,

[111823]J. L.: TNF-alpha and hyperandrogenism: a clinical, bio-
chemical, and

[111824]molecular genetic study. J. Clin. Endocr. Metab. 86:
3761-3767,

[111825]2001.

[111826]19818.

[111827]Gorman, J. D.; Sack, K. E.; Davis, J. C., Jr.: Treatment of
ankylosing

[111828]spondylitis by inhibition of tumor necrosis factor-alpha.
New Eng.

[111829]J. Med. 346: 1349-1356, 2002.

[111830]19819.

- [111831]Herrmann, S.-M.; Ricard, S.; Nicaud, V.; Mallet, C.;
Arveiler,
[111832]D.; Evans, A.; Ruidavets, J.-B.; Luc, G.; Bara, L.; Parra,
H.-J.;
[111833]Poirier, O.; Cambien, F.: Polymorphisms of the tumour
necrosis factor- α
[111834]gene, coronary heart disease and obesity. *Europ. J. Clin.*
Invest. 28:
[111835]59-66, 1998.
[111836]19820.
[111837]Inoko, H.; Trowsdale, J.: Linkage of TNF genes to the HLA-
B locus. *Nucleic*
[111838]*Acids Res.* 15: 8957-8962, 1987.
[111839]19821.
[111840]Knight, J. C.; Udalova, I.; Hill, A. V. S.; Greenwood, B. M.;
[111841]Peshu, N.; Marsh, K.; Kwiatkowski, D.: A polymorphism
that affects
[111842]OCT-1 binding to the TNF promoter region is associated
with severe
[111843]malaria. *Nature Genet.* 22: 145-150, 1999.
[111844]19822.

[111845]Koss, K.; Satsangi, J.; Fanning, G. C.; Welsh, K. I.; Jewell,

[111846]D. P.: Cytokine (TNF- α , LT- α , and IL-10) polymorphisms in

[111847]inflammatory bowel diseases and normal controls: differential effects

[111848]on production and allele frequencies. Genes Immun. 1: 185-190, 2000.

[111849]19823.

[111850]Marino, M. W.; Dunn, A.; Grail, D.; Inglese, M.; Noguchi, Y.;

[111851]Richards, E.; Jungbluth, A.; Wada, H.; Moore, M.; Williamson, B.;

[111852]Basu, S.; Old, L. J.: Characterization of tumor necrosis factor-deficient

[111853]mice. Proc. Nat. Acad. Sci. 94: 8093-8098, 1997.

[111854]19824.

[111855]McCusker, S. M.; Curran, M. D.; Dynan, K. B.; McCullagh, C. D.;

[111856]Urquhart, D. D.; Middleton, D.; Patterson, C. C.; McIlroy, S. P.;

[111857]Passmore, A. P.: Association between polymorphism in regulatory region

[111858]of gene encoding tumour necrosis factor- α and risk
of Alzheimer's

[111859]disease and vascular dementia: a case-control study.

Lancet 357:

[111860]436-439, 2001.

[111861]19825.

[111862]Remaley, A. T.; Rust, S.; Rosier, M.; Knapper, C.; Naudin,
L.;

[111863]Broccardo, C.; Peterson, K. M.; Koch, C.; Arnould, I.;
Prades, C.;

[111864]Duverger, N.; Funke, H.; Assman, G.; Dinger, M.; Dean, M.;
Chimini,

[111865]G.; Santamarina-Fojo, S.; Fredrickson, D. S.; Deneffe, P.
Brewer,

[111866]H. B., Jr.: Human ATP-binding cassette transporter 1
(ABC1): genomic

[111867]organization and identification of the genetic defect in the
original

[111868]Tangier disease kindred. Proc. Nat. Acad. Sci. 96:
12685-12690,

[111869]1999.

[111870]19826.

- [111871]Rust, S.; Rosier, M.; Funke, H.; Real, J.; Amoura, Z.; Piette, J.-C.; Deleuze, J.-F.; Brewer, H. B.; Duverger, N.; Deneffe, P.; Assmann, G.: Tangier disease is caused by mutations in the gene encoding ATP-binding cassette transporter 1. *Nature Genet.* 22: 352–355, 1999.
- [111872]19827.
- [111873]Young, S. G.; Fielding, C. J.: The ABCs of cholesterol efflux. *Nature Genet.* 22: 316–318, 1999.
- [111874]19828.
- [111875]Garcia-Higuera, I.; Taniguchi, T.; Ganesan, S.; Meyn, M. S.; Timmers, C.; Hejna, J.; Grompe, M.; D'Andrea, A. D.: Interaction of the Fanconi anemia proteins and BRCA1 in a common pathway. *Molec. Cell* 7: 249–262, 2001.
- [111876]19829.
- [111877]Mudd, S. H.; Levy, H. L.; Abeles, R. H.: A derangement in

[111885]metabolism leading to homocystinemia, cystathioninemia and methylmalonic

[111886]aciduria. Biochem. Biophys. Res. Commun. 35: 121–126, 1969.

[111887]19830.

[111888]Mudd, S. H.; Levy, H. L.; Skovby, F.: Disorders of transsulfuration.In:

[111889]Scriver, C. R.; Beaudet, A. L.; Sly, W. S.; Valle, D. (eds.): The

[111890]Metabolic and Molecular Bases of Inherited Disease. Vol. 1. New

[111891]York: McGraw–Hill (7th ed.): 1995. Pp. 1279–1327.

[111892]19831.

[111893]Mudd, S. H.; Skovby, F.; Levy, H. L.; Pettigrew, K. D.; Wilcken,

[111894]B.; Pyeritz, R. E.; Andria, G.; Boers, G. H. J.; Bromberg, I. L.;

[111895]Cerone, R.; Fowler, B.; Grobe, H.; Schmidt, H.; Schweitzer, L.: The

[111896]natural history of homocystinuria due to cystathionine beta-synthase

[111897]deficiency. Am. J. Hum. Genet. 37: 1–31, 1985.

[111898]19832.

- [111899] Munke, M.; Kraus, J.; Watkins, P.; Tanzi, R.; Gusella, J.;
Millington
- [111900] Ward, A.; Watson, M.; Francke, U.: Homocystinuria gene
on human chromosome
- [111901] 21 mapped with cloned cystathionine beta-synthase
probe and in situ
- [111902] hybridization of other chromosome 21 probes. (Abstract)
Cytogenet.
- [111903] Cell Genet. 40: 706-707, 1985.
- [111904] 19833.
- [111905] Munke, M.; Kraus, J. P.; Ohura, T.; Francke, U.: The gene
for
- [111906] cystathionine beta-synthase (CBS) maps to the subtelom-
eric region
- [111907] on human chromosome 21q and to proximal mouse chro-
mosome 17. Am.
- [111908] J. Hum. Genet. 42: 550-559, 1988.
- [111909] 19834.
- [111910] Munnich, A.; Saudubray, J.-M.; Dautzenberg, M.-D.; Parvy,
P.;
- [111911] Ogier, H.; Girot, R.; Manigne, P.; Frezal, J.: Diet-responsive
proconvertin

[111912](factor VII) deficiency in homocystinuria. J. Pediat. 102:
730-734,

[111913]1983.

[111914]19835.

[111915]Nishinaga, M.; Ozawa, T.; Shimada, K.: Homocysteine, a
thrombogenic

[111916]agent, suppresses anticoagulant heparan sulfate expres-
sion in cultured

[111917]porcine aortic endothelial cells. J. Clin. Invest. 92:
1381-1386,

[111918]1993.

[111919]19836.

[111920]Nugent, A.; Hadden, D. R.; Carson, N. A. J.: Long-term
survival

[111921]of homocystinuria: the first case. Lancet 352: 624-625,
1998.

[111922]19837.

[111923]Perry, T. L.; Hansen, S.; Love, D. L.; Crawford, L. E.; Tis-
chler,

[111924]B.: Treatment of homocystinuria with a low-methionine
diet, supplemental

[111925]cystine and a methyl donor. Lancet II: 474–478, 1968.

[111926]19838.

[111927]Peterschmitt, M. J.; Simmons, J. R.; Levy, H. L.: Reduction
of

[111928]false negative results in screening of newborns for homo-
cystinuria. New

[111929]Eng. J. Med. 341: 1572–1576, 1999.

[111930]19839.

[111931]Pogribna, M.; Melnyk, S.; Pogribny, I.; Chango, A.; Yi, P.;
James,

[111932]S. J.: Homocysteine metabolism in children with Down
syndrome: in

[111933]vitro modulation. Am. J. Hum. Genet. 69: 88–95, 2001.

[111934]19840.

[111935]Ratnoff, O. D.: Activation of Hageman factor by L-
homocystine. Science 162:

[111936]1007–1009, 1968.

[111937]19841.

[111938]Reish, O.; Townsend, D.; Berry, S. A.; Tsai, M. Y.; King, R.
A.

- [111939] Tyrosinase inhibition due to interaction of homo-
cyst(e)ine with
- [111940] copper: the mechanism for reversible hypopigmentation in
homocystinuria
- [111941] due to cystathionine beta-synthase deficiency. Am. J.
Hum. Genet. 57:
- [111942] 127-132, 1995.
- [111943] 19842.
- [111944] Rosenquist, T. H.; Ratashak, S. A.; Selhub, J.: Homocys-
teine
- [111945] induces congenital defects of the heart and neural tube:
effect of
- [111946] folic acid. Proc. Nat. Acad. Sci. 93: 15227-15232, 1996.
- [111947] 19843.
- [111948] Saudubray, J. M.: Personal Communication. Paris, France
5/20/1997.
- [111949] 19844.
- [111950] Schimke, R. N.; McKusick, V. A.; Huang, T.; Pollack, A. D.:
Homocystinuria:
- [111951] studies of 20 families with 38 affected members. J.A.M.A.
193: 711-719,

[111952]1965.

[111953]19845.

[111954]Schnyder, G.; Roffi, M.; Pin, R.; Flammer, Y.; Lange, H.;
Eberli,

[111955]F. R.; Meier, B.; Turi, Z. G.; Hess, O. M.: Decreased rate of
coronary

[111956]restenosis after lowering of plasma homocysteine levels.
New Eng.

[111957]J. Med. 345: 1593–1600, 2001.

[111958]19846.

[111959]Sebastio, G.: Personal Communication. Padua, Italy
3/20/1997.

[111960]19847.

[111961]Sebastio, G.; Sperandeo, M. P.; Panico, M.; de Franchis, R.;
Kraus,

[111962]J. P.; Andria, G.: The molecular basis of homocystinuria
due to cystathionine

[111963]beta-synthase deficiency in Italian families, and report of
four novel

[111964]mutations. Am. J. Hum. Genet. 56: 1324–1333, 1995.

[111965]19848.

[111966]Shan, X.; Dunbrack, R. L., Jr.; Christopher, S. A.; Kruger, W.

[111967]D.: Mutations in the regulatory domain of cystathionine
beta-synthase

[111968]can functionally suppress patient-derived mutations in
cis. Hum.

[111969]Molec. Genet. 10: 635–643, 2001.

[111970]19849.

[111971]Shan, X.; Kruger, W. D.: Correction of disease-causing CBS
mutations

[111972]in yeast. Nature Genet. 19: 91–93, 1998.

[111973]19850.

[111974]Taniguchi, T.; Garcia-Higuera, I.; Xu, B.; Andreassen, P. R.;
Gregory,

[111975]R. C.; Kim, S.-T.; Lane, W. S.; Kastan, M. B.; D'Andrea, A.
D.: Convergence

[111976]of the Fanconi anemia and ataxia telangiectasia signaling
pathways. Cell 109:

[111977]459–472, 2002.

[111978]19851.

[111979]Timmers, C.; Taniguchi, T.; Hejna, J.; Reifsteck, C.; Lucas,
L.;

- [111980] Bruun, D.; Thayer, M.; Cox, B.; Olson, S.; D'Andrea, A. D.;
Moses,
- [111981] R.; Grompe, M.: Positional cloning of a novel Fanconi ane-
mia gene,
- [111982] FANCD2. Molec. Cell 7: 241–248, 2001.
- [111983] 19852.
- [111984] Whitney, M.; Thayer, M.; Reifsteck, C.; Olson, S.; Smith, L.;
Jakobs,
- [111985] P. M.; Leach, R.; Naylor, S.; Joenje, H.; Grompe, M.: Micro-
cell mediated
- [111986] chromosome transfer maps the Fanconi anaemia group D
gene to chromosome
- [111987] 3p. Nature Genet. 11: 341–343, 1995.
- [111988] 19853.
- [111989] Hejna, J. A.; Timmers, C. D.; Reifsteck, C.; Bruun, D. A.;
Lucas,
- [111990] L. W.; Jakobs, P. M.; Toth–Fejel, S.; Unsworth, N.; Clemens,
S. L.;
- [111991] Garcia, D. K.; Naylor, S. L.; Thayer, M. J.; Olson, S. B.;
Grompe,
- [111992] M.; Moses, R. E.: Localization of the Fanconi anemia com-
plementation

- [111993]group D gene to a 200-kb region on chromosome 3p25.3.
Am. J. Hum.
- [111994]Genet. 66: 1540–1551, 2000.
- [111995]19854.
- [111996]Buchwald, M.: Complementation groups: one or more per
gene? Nature
- [111997]Genet. 11: 228–230, 1995.
- [111998]19855.
- [111999]Stambolian, D.; Ai, Y.; Sidjanin, D.; Nesburn, K.; Sathe, G.;
- [112000]Rosenberg, M.; Bergsma, D. J.: Cloning of the galactoki-
nase cDNA
- [112001]and identification of mutations in two families with
cataracts. Nature
- [112002]Genet. 10: 307–312, 1995.
- [112003]19856.
- [112004]Alano, A.; Almashanu, S.; Chinsky, J. M.; Costeas, P.;
- Blitzer,
- [112005]M. G.; Wulfsberg, E. A.; Cowan, T. M.: Molecular character-
ization
- [112006]of a unique patient with epimerase-deficiency galac-
tosaemia. J. Inherit.

[112007]Metab. Dis. 21: 341–350, 1998.

[112008]19857.

[112009]Daude, N.; Gallaher, T. K.; Zeschnigk, M.; Starzinski-Powitz, A.;

[112010]Petry, K. G.; Haworth, I. S.; Reichardt, J. K. V.: Molecular cloning,

[112011]characterization, and mapping of a full-length cDNA encoding human

[112012]UDP-galactose 4-prime-epimerase. Biochem. Molec. Med. 56: 1–7, 1995.

[112013]19858.

[112014]Henderson, M. J.; Holton, J. B.; MacFaul, R.: Further observations

[112015]in a case of uridine diphosphate galactose-4-epimerase deficiency

[112016]with a severe clinical presentation. J. Inherit. Metab. Dis. 6:

[112017]17–20, 1983.

[112018]19859.

[112019]Holton, J. B.; Gillett, M. G.; MacFaul, R.; Young, R.: Galactosaemia:

- [112020] a new severe variant due to uridine diphosphate galactose-4-epimerase
- [112021] deficiency. Arch. Dis. Child. 56: 885-887, 1981.
- [112022] 19860.
- [112023] Kingsley, D. M.; Kozarsky, K. F.; Hobbie, L.; Krieger, M.:
Reversible
- [112024] defects in O-linked glycosylation and LDL receptor expression in a
- [112025] UDP-Gal/UDP-GalNAc 4-epimerase deficient mutant. Cell 44: 749-759,
- [112026] 1986.
- [112027] 19861.
- [112028] Maceratesi, P.; Daude, N.; Dallapiccola, B.; Novelli, G.;
Allen,
- [112029] R.; Okano, Y.; Reichardt, J.: Human UDP-galactose
4-prime epimerase
- [112030] (GALE) gene and identification of five missense mutations
in patients
- [112031] with epimerase-deficiency galactosemia. Molec. Genet. Metab. 63:
- [112032] 26-30, 1998.

[112033]19862.

[112034]Piller, F.; Hanlon, M. H.; Hill, R. L.: Co-purification and
characterization

[112035]of UDP-glucose 4-epimerase and UDP-
N-acetylglucosamine 4-epimerase

[112036]from porcine submaxillary glands. J. Biol. Chem. 258:
10774-10778,

[112037]1983.

[112038]19863.

[112039]Quimby, B. B.; Alano, A.; Almashanu, S.; DeSandro, A. M.;
Cowan,

[112040]T. M.; Fridovich-Keil, J. L.: Characterization of two muta-
tions associated

[112041]with epimerase-deficiency galactosemia, by use of a yeast
expression

[112042]system for human UDP-galactose-4-epimerase. Am. J.
Hum. Genet. 61:

[112043]590-598, 1997.

[112044]19864.

[112045]Dubiel, B.; Dabrowski, C.; Wetts, R.; Tanaka, K.: Comple-
mentation

[112046]studies of isovaleric acidemia and glutaric aciduria type II
using

[112047]cultured skin fibroblasts. J. Clin. Invest. 72: 1543–1552,
1983.

[112048]19865.

[112049]McKusick, V. A.: Ellis–van Creveld syndrome and the
Amish. Nature

[112050]Genet. 24: 203–204, 2000.

[112051]19866.

[112052]McKusick, V. A.; Egeland, J. A.; Eldridge, R.; Krusen, D. E.:

[112053]Dwarfism in the Amish. I. The Ellis–van Creveld syndrome.
Bull. Johns

[112054]Hopkins Hosp. 115: 306–336, 1964.

[112055]19867.

[112056]Coppola, G.; De Michele, G.; Cavalcanti, F.; Pianese, L.;
Perretti,

[112057]A.; Santoro, L.; Vita, G.; Toscano, A.; Amboni, M.;
Grimaldi, G.;

[112058]Salvatore, E.; Caruso, G.; Filla, A.: Why do some Friedre-
ich's ataxia

[112059]patients retain tendon reflexes? A clinical, neurophysio-

logical and

[112060]molecular study. J. Neurol. 246: 353–357, 1999.

[112061]19868.

[112062]Delatycki, M. B.; Knight, M.; Koenig, M.; Cossee, M.;
Williamson,

[112063]R.; Forrest, S. M.: G130V, a common FRDA point mutation,
appears

[112064]to have arisen from a common founder. Hum. Genet. 105:
343–346,

[112065]1999.

[112066]19869.

[112067]Delatycki, M. B.; Paris, D. B. B. P.; Gardner, R. J. M.; Nichol-
son,

[112068]G. A.; Nassif, N.; Storey, E.; MacMillan, J. C.; Collins, V.;
Williamson,

[112069]R.; Forrest, S. M.: Clinical and genetic study of Friedreich
ataxia

[112070]in an Australian population. Am. J. Med. Genet. 87:
168–174, 1999.

[112071]19870.

[112072]Durr, A.; Cossee, M.; Agid, Y.; Campuzano, V.; Mignard,

C.; Penet,

[112073]C.; Mandel, J.-L.; Brice, A.; Koenig, M.: Clinical and genetic abnormalities

[112074]in patients with Friedreich's ataxia. New Eng. J. Med. 335: 1169-1175,

[112075]1996.

[112076]19871.

[112077]Filla, A.; De Michele, G.; Cavalcanti, F.; Pianese, L.; Monticelli,

[112078]A.; Campanella, G.; Coccozza, S.: The relationship between trinucleotide

[112079](GAA) repeat length and clinical features in Friedreich ataxia. Am.

[112080]J. Hum. Genet. 59: 554-560, 1996.

[112081]19872.

[112082]Fujita, R.; Hanauer, A.; Vincent, A.; Mandel, J.-L.; Koenig, M.

[112083]: Physical mapping of two loci (D9S5 and D9S15) tightly linked to

[112084]Friedreich ataxia locus (FRDA) and identification of nearby CpG islands

[112085]by pulse-field gel electrophoresis. Genomics 10:

915–920, 1991.

[112086]19873.

[112087]Gray, J. V.; Johnson, K. J.: Waiting for frataxin. *Nature Genet.* 16:

[112088]323–325, 1997.

[112089]19874.

[112090]Kenney, R. T.; Malech, H. L.; Epstein, N. D.; Roberts, R. L.;
Leto,

[112091]T. L.: Characterization of the p67–phox gene: genomic
organization

[112092]and restriction fragment length polymorphism analysis for
prenatal

[112093]diagnosis in chronic granulomatous disease. *Blood* 82:
3739–3744,

[112094]1993.

[112095]19875.

[112096]Noack, D.; Rae, J.; Cross, A. R.; Munoz, J.; Salmen, S.;
Mendoza,

[112097]J. A.; Rossi, N.; Curnutte, J. T.; Heyworth, P. G.: Autosomal
recessive

[112098]chronic granulomatous disease caused by novel mutations

in NCF-2,

[112099]the gene encoding the p67-phox component of phagocyte NADPH oxidase. Hum.

[112100]Genet. 105: 460-467, 1999.

[112101]19876.

[112102]Nunoi, H.; Iwata, M.; Tatsuzawa, S.; Onoe, Y.; Shimizu, S.; Kanegasaki,

[112103]S.; Matsuda, I.: AG dinucleotide insertion in a patient with chronic

[112104]granulomatous disease lacking cytosolic 67-kD protein. Blood 86:

[112105]329-333, 1995.

[112106]19877.

[112107]Okamura, N.; Babior, B. M.; Mayo, L. A.; Peveri, P.; Smith, R.

[112108]M.; Curnutte, J. T.: The p67-phox cytosolic peptide of the respiratory

[112109]burst oxidase from human neutrophils: functional aspects. J. Clin.

[112110]Invest. 85: 1583-1587, 1990.

[112111]19878.

[112112]Patino, P. J.; Rae, J.; Noack, D.; Erickson, R.; Ding, J.; Garcia

[112113]de Olarte, D.; Curnutte, J. T.: Molecular characterization of autosomal

[112114]recessive chronic granulomatous disease caused by a defect of the

[112115]nicotinamide adenine dinucleotide phosphate (reduced form) oxidase

[112116]component p67-phox. Blood 94: 2505-2514, 1999.

[112117]19879.

[112118]Tanugi-Cholley, L. C.; Issartel, J.-P.; Lunardi, J.; Freycon,

[112119]F.; Morel, F.; Vignais, P. V.: A mutation located at the

5-prime

[112120]splice junction sequence of intron 3 in the p67-phox gene causes the

[112121]lack of p67-phox mRNA in a patient with chronic granulomatous disease. Blood 85:

[112122]242-249, 1995.

[112123]19880.

[112124]Forsius, H.; Eriksson, A.; Nuutila, A.; Vainio-Mattila, B.; Krause,

[112125]U.: A genetic study of three rare retinal disorders: dystrophia retinae

[112126]dysacusis syndrome, X-chromosomal retinoschisis and grouped pigments

[112127]of the retina. Birth Defects Orig. Art. Ser. VII(3): 83-98, 1971.

[112128]19881.

[112129]Hassin, G. B.: Amaurotic family idiocy: late infantile type (Bielschowsky)

[112130]with the clinical picture of decerebrate rigidity. Arch. Neurol.

[112131]Psychiat. 16: 708-727, 1926.

[112132]19882.

[112133]Jarvela, I.: Infantile neuronal ceroid lipofuscinosis (CLN1):

[112134]linkage disequilibrium in the Finnish population and evidence that

[112135]variant late infantile form (variant CLN2) represents a non-allelic

[112136]locus. Genomics 10: 333–337, 1991.

[112137]19883.

[112138]Liu, C.-G.; Sleat, D. E.; Donnelly, R. J.; Lobel, P.: Structural

[112139]organization and sequence of CLN2, the defective gene in classical

[112140]late infantile neuronal ceroid lipofuscinosis. Genomics 50: 206–212,

[112141]1998.

[112142]19884.

[112143]MacLeod, P. M.; Dolman, C. L.; Nickel, R. E.; Chang, E.; Nag,

[112144]S.; Zonana, J.; Silvey, K.: Prenatal diagnosis of neuronal ceroid-lipofuscinoses. Am.

[112145]J. Med. Genet. 22: 781–789, 1985.

[112146]19885.

- [112147] MacLeod, P. M.; Dolman, C. L.; Nickel, R. E.; Chang, E.; Zonana,
nana,
- [112148] J.; Silvey, K.: Prenatal diagnosis of neuronal ceroid lipofus-
cinosis
- [112149] (Letter) New Eng. J. Med. 310: 595 only, 1984.
- [112150] 19886.
- [112151] Seitelberger, F.; Vogel, G.; Stepan, H.: Späta infantile
amaurotische
- [112152] Idiotie. Arch. Psychiat. Nervenkr. 196: 154–190, 1957.
- [112153] 19887.
- [112154] Sharp, J.; Savukoski, M.; Wheeler, R. B.; Harris, J.; Jarvela,
Jarvela,
- [112155] J.; Peltonen, L.; Gardiner, M.; Williams, R.: Linkage analysis
of
- [112156] late-infantile neuronal ceroid-lipofuscinosis. Am. J. Med.
Genet. 57:
- [112157] 348–349, 1995.
- [112158] 19888.
- [112159] Sharp, J. D.; Wheeler, R. B.; Lake, B. D.; Savukoski, M.;
Jarvela,
- [112160] J. E.; Peltonen, L.; Gardiner, R. M.; Williams, R. E.: Loci for
classical

- [112161]and a variant late infantile neuronal ceroid lipofuscinosis map to
- [112162]chromosomes 11p15 and 15q21–23. Hum. Molec. Genet. 6: 591–595, 1997.
- [112163]19889.
- [112164]Sleat, D. E.; Donnelly, R. J.; Lackland, H.; Liu, C.–G.; Sohar,
- [112165]J.; Pullarkat, R. K.; Lobel, P.: Association of mutations in a lysosomal
- [112166]protein with classical late–infantile neuronal ceroid lipofuscinosis. Science 277:
- [112167]1802–1805, 1997.
- [112168]19890.
- [112169]Sleat, D. E.; Gin, R. M.; Sohar, I.; Wisniewski, K.; Sklower–Brooks,
- [112170]S.; Pullarkat, R. K.; Palmer, D. N.; Lerner, T. J.; Boustany, R.–M.;
- [112171]Uldall, P.; Siakotos, A. N.; Donnelly, R. J.; Lobel, P.: Mutational
- [112172]analysis of the defective protease in classic late–infantile neuronal
- [112173]ceroid lipofuscinosis, a neurodegenerative lysosomal storage disorder. Am.

[112174]J. Hum. Genet. 64: 1511–1523, 1999.

[112175]19891.

[112176]Sohar, I.; Sleat, D. E.; Jadot, M.; Lobel, P.: Biochemical
characterization

[112177]of a lysosomal protease deficient in classical late infantile
neuronal

[112178]ceroid lipofuscinosis (LINCL) and development of an en-
zyme-based assay

[112179]for diagnosis and exclusion of LINCL in human specimens
and animal

[112180]models. J. Neurochem. 73: 700–711, 1999.

[112181]19892.

[112182]Volk, B. W.; Wallace, B. J.; Schneck, L.; Saifer, A.: Late in-
fantile

[112183]amaurotic idiocy. Ultramicroscopic and histochemical
studies on a

[112184]case. Arch. Path. 78: 483–500, 1964.

[112185]19893.

[112186]Williams, R.; Vesa, J.; Jarvela, I.; McKay, T.; Mitchison, H.;

[112187]Hellsten, E.; Thompson, A.; Callen, D.; Sutherland, G.;
Luna-Battadano,

[112188]D.; Stallings, R.; Peltonen, L.; Gardiner, M.: Genetic heterogeneity
[112189]in neuronal ceroid lipofuscinosis (NCL): evidence that the
late-infantile
[112190]subtype (Jansky-Bielschowsky disease; CLN2) is not an allelic form
[112191]of the juvenile or infantile subtypes. Am. J. Hum. Genet.
53: 931-935,
[112192]1993.
[112193]19894.
[112194]Yan, W.; Boustany, R.-M. N.; Konradi, C.; Ozelius, L.;
Lerner,
[112195]T.; Trofatter, J. A.; Julier, C.; Breakefield, X. O.; Gusella, J.
[112196]F.; Haines, J. L.: Localization of juvenile, but not late-infantile,
[112197]neuronal ceroid lipofuscinosis on chromosome 16. Am. J.
Hum. Genet. 52:
[112198]89-95, 1993.
[112199]19895.
[112200]Zhong, N.; Wisniewski, K. E.; Hartikainen, J.; Ju, W.; Moroziewicz,

- [112201]D. N.; McLendon, L.; Sklower Brooks, S.; Brown, W. T.: Two common
- [112202]mutations in the CLN2 gene underlie late infantile neuronal ceroid
- [112203]lipofuscinosis. Clin. Genet. 54: 234–238, 1998.
- [112204]19896.
- [112205]Harnden, D. G.: Ataxia–telangiectasia syndrome: cytogenetic and
- [112206]cancer aspects.In: German, J.: Chromosomes and Cancer. New York:
- [112207]Wiley (pub.) 1974. Pp. 619–636.
- [112208]19897.
- [112209]Hartwell, L.: Defects in a cell cycle checkpoint may be responsible
- [112210]for the genomic instability of cancer cells. Cell 71: 543–546, 1992.
- [112211]19898.
- [112212]Hawley, R. S.; Friend, S. H.: Strange bedfellows in even stranger
- [112213]places: the role of ATM in meiotic cells, lymphocytes, tumors, and

[112214]its functional links to p53. Genes Dev. 10: 2383–2388, 1996.

[112215]19899.

[112216]Hecht, F.; Koler, R. D.; Rigas, D. A.; Dahnke, G. S.; Case, M.

[112217]P.; Tisdale, V.; Miller, R. W.: Leukemia and lymphocytes in ataxia–telangiectasia.

[112218](Letter) Lancet II: 1193, 1966.

[112219]19900.

[112220]Henderson, L.; Cole, H.; Arlett, C.; James, S. E.; Cole, J.; Lehmann,

[112221]A.; Rosenbloom, L.; Redmond, T.; Meller, S.: Diagnosis of ataxia–telangiectasia

[112222]by T–lymphocyte cloning assay. (Letter) Lancet II: 1242, 1985.

[112223]19901.

[112224]Hernandez, D.; McConville, C. M.; Stacey, M.; Woods, C. G.; Brown,

[112225]M. M.; Shutt, P.; Rysiecki, G.; Taylor, A. M. R.: A family showing

[112226]no evidence of linkage between the ataxia telangiectasia gene and

[112227]chromosome 11q22-23. J. Med. Genet. 30: 135-140,
1993.

[112228]19902.

[112229]Hoar, D. I.; Sargent, P.: Chemical mutagen hypersensitivity
in

[112230]ataxia-telangiectasia. Nature 261: 590-592, 1976.

[112231]19903.

[112232]Hodge, S. E.; Berkel, A. I.; Gatti, R. A.; Boder, E.; Spence,

[112233]M. A.: Ataxia-telangiectasia and xeroderma pigmentosum:
no evidence

[112234]of linkage to HLA. Tissue Antigens 15: 313-317, 1980.

[112235]19904.

[112236]Huang, P. C.; Sheridan, R. B., III: Genetic and biochemical
studies

[112237]with ataxia telangiectasia. Hum. Genet. 59: 1-9, 1981.

[112238]19905.

[112239]Humphreys, M. W.; Nevin, N. C.; Wooldridge, M. A. W.: Cy-
togenetic

[112240]investigations in a family with ataxia telangiectasia. Hum.
Genet. 83:

[112241]79–82, 1989.

[112242]19906.

[112243]Imai, T.: Personal Communication. Chiba, Japan
9/12/1996.

[112244]

[112245]100. Ishiguro, T.; Taketa, K.; Gatti, R. A.: Tissue of origin
of

[112246]elevated alpha-fetoprotein in ataxia-telangiectasia. Dis.
Markers 4:

[112247]293–297, 1986.

[112248]

[112249]101. Jaspers, N. G. J.; Bootsma, D.: Genetic heterogeneity
in ataxia-telangiectasia

[112250]studied by cell fusion. Proc. Nat. Acad. Sci. 79:
2641–2644, 1982.

[112251]

[112252]102. Jaspers, N. G. J.; Gatti, R. A.; Baan, C.; Linssen, P. C.
M.

[112253]L.; Bootsma, D.: Genetic complementation analysis of
ataxia telangiectasia

[112254]and Nijmegen breakage syndrome: a survey of 50 pa-

tients. Cytogenet.

[112255]Cell Genet. 49: 259–263, 1988.

[112256]

[112257]103. Jaspers, N. G. J.; Painter, R. B.; Paterson, M. C.; Kidson, C.;

[112258]Inoue, T.: Complementation analysis of ataxia–telangiectasia.In:

[112259]Gatti, R. A.; Swift, M.: Ataxia–telangiectasia: Genetics, Neuropathology

[112260]and Immunology of a Degenerative Disease of Childhood. New York:

[112261]Alan R. Liss (pub.) 1985. Pp. 147–162.

[112262]

[112263]104. Johnson, J. P.; Gatti, R. A.; Sears, T. S.; White, R. L.: Inverted

[112264]duplication of J(H) associated with chromosome 14 translocation and

[112265]T–cell leukemia in ataxia–telangiectasia. Am. J. Hum. Genet. 39:

[112266]787–796, 1986.

[112267]

- [112268]105. Johnson, J. P.; White, R. L.; Gatti, R. A.: Rearrange-
ment of
[112269]J(H) genes in a patient with ataxia telangiectasia, chromo-
some 14
[112270]translocation, and T-cell leukemia. (Abstract) Am. J. Hum.
Genet. 37:
[112271]A100, 1985.
- [112272]
- [112273]106. Jung, M.; Zhang, Y.; Lee, S.; Dritschilo, A.: Correction
of
[112274]radiation sensitivity in ataxia telangiectasia cells by a
truncated
[112275]I-kappa-B-alpha. Science 268: 1619-1621, 1995.
- [112276]
- [112277]107. Kapp, L. N.; Painter, R. B.; Yu, L.-C.; van Loon, N.;
Richard,
[112278]C. W., III; James, M. R.; Cox, D. R.; Murnane, J. P.: Cloning
of
[112279]a candidate gene for ataxia-telangiectasia group D. Am. J.
Hum. Genet. 51:
[112280]45-54, 1992.

[112281]

[112282]108. Kastan, K.: Clinical implications of basic research:
ataxia-telangiectasia--broad

[112283]implications for a rare disorder. New Eng. J. Med. 333:
662-663,

[112284]1995.

[112285]

[112286]109. Kastan, M. B.; Onyekwere, O.; Sidransky, D.; Vogel-
stein, B.;

[112287]Craig, R. W.: Participation of p53 protein in the cellular re-
sponse

[112288]to DNA damage. Cancer Res. 51: 6304-6311, 1991.

[112289]

[112290]110. Kastan, M. B.; Zhan, Q.; El-Deiry, W. S.; Carrier, F.;

[112291]Jacks,
T.; Walsh, W. V.; Plunkett, B. S.; Vogelstein, B.; Fornace, A.
J.,

[112292]Jr.: A mammalian cell cycle checkpoint pathway utilizing
p53 and

[112293]GADD45 is defective in ataxia-telangiectasia. Cell 71:
587-597,

[112294]1992.

[112295]

[112296]111. Kennaugh, A. A.; Butterworth, S. V.; Hollis, R.; Baer, R.; Rabbitts,

[112297]T. H.; Taylor, A. M. R.: The chromosome breakpoint at 14q32 in an

[112298]ataxia telangiectasia t(14;14) T cell clone is different from the

[112299]14q32 breakpoint in Burkitts and an inv(14) T cell lymphoma. Hum.

[112300]Genet. 73: 254-259, 1986.

[112301]

[112302]112. Khanna, K. K.; Keating, K. E.; Kozlov, S.; Scott, S.; Gatei,

[112303]M.; Hobson, K.; Taya, Y.; Gabrielli, B.; Chan, D.; Lees-Miller, S.

[112304]P.; Lavin, M. F.: ATM associates with and phosphorylates p53: mapping

[112305]the region of interaction. Nature Genet. 20: 398-400, 1998.

[112306]

[112307]113. Klein, C.; Wenning, G. K.; Quinn, N. P.; Marsden, C.
D.: Ataxia

[112308]without telangiectasia masquerading as benign hereditary
chorea. Mov.

[112309]Disord. 11: 217–220, 1996.

[112310]

[112311]114. Kojis, T. L.; Schreck, R. R.; Gatti, R. A.; Sparkes, R. S.:

[112312]Tissue specificity of chromosomal rearrangements in
ataxia–telangiectasia. Hum.

[112313]Genet. 83: 347–352, 1989.

[112314]

[112315]115. Komatsu, K.; Kodama, S.; Okumura, Y.; Koi, M.; Os-
himura, M.:

[112316]Restoration of radiation resistance in ataxia–telangiectasia
cells

[112317]by the introduction of normal human chromosome 11.
Mutat. Res. 235:

[112318]59–63, 1990.

[112319]

[112320]116. Komatsu, K.; Matsuura, S.; Tauchi, H.; Endo, S.; Ko-
dama, S.;

[112321]Smeets, D.; Weemaes, C.; Oshimura, M.: The gene for Ni-jmegen breakage

[112322]syndrome (V2) is not located on chromosome 11. (Letter)
Am. J. Hum.

[112323]Genet. 58: 885–888, 1996.

[112324]

[112325]117. Korein, J.; Steinman, P. A.; Senz, E. H.: Ataxia–telangiectasia:

[112326]report of a case and review of the literature. Arch. Neurol.
4:

[112327]272–280, 1961.

[112328]

[112329]118. Krishna Kumar, G.; Al Saadi, A.; Yang, S. S.; Mc-Caughey, R. S.

[112330]: Ataxia–telangiectasia and hepatocellular carcinoma. Am.
J. Med.

[112331]Sci. 278: 157–160, 1979.

[112332]

[112333]119. Laake, K.; Jansen, L.; Hahnemann, J. M.; Brondum–Nielsen, K.;

[112334]Lonnqvist, T.; Kaariainen, H.; Sankila, R.; Lahdesmaki, A.;

Hammarstrom,

[112335]L.; Yuen, J.; Tretli, S.; Heiberg, A.; Olsen, J. H.; Tucker, M.;
Kleinerman,

[112336]R.; Borresen-Dale, A.-L.: Characterization of ATM muta-
tions in 41

[112337]Nordic families with ataxia telangiectasia. Hum. Mutat. 16:
232-246,

[112338]2000.

[112339]

[112340]120. Laake, K.; Telatar, M.; Geitvik, G. A.; Hansen, R. O.;
Heiberg,

[112341]A.; Andresen, A. M.; Gatti, R.; Borresen-Dale, A.-L.: Identical
mutation

[112342]in 55% of the ATM alleles in 11 Norwegian AT families: evidence for

[112343]a founder effect. Europ. J. Hum. Genet. 6: 235-244, 1998.

[112344]

[112345]121. Lambert, C.; Schultz, R. A.; Smith, M.; Wagner-
McPherson, C.;

[112346]McDaniel, L. D.; Donlon, T.; Stanbridge, E. J.; Friedberg, E.
C.:

[112347]Functional complementation of ataxia-telangiectasia

group D (AT-D)

[112348]cells by microcell-mediated chromosome transfer and mapping of the

[112349]AT-D locus to the region 11q22-23. Proc. Nat. Acad. Sci. 88: 5907-5911,

[112350]1991.

[112351]

[112352]122. Levin, S.; Perlov, S.: Ataxia-telangiectasia in Israel, with

[112353]observations on its relationship to malignant disease. Israel J. rael J.

[112354]Med. Sci. 7: 1535-1541, 1971.

[112355]

[112356]123. Li, A.; Swift, M.: Mutations at the ataxia-telangiectasia locus

[112357]and clinical phenotypes of A-T patients. Am. J. Med. Genet. 92:

[112358]170-177, 2000.

[112359]

[112360]124. Li, S.; Ting, N. S. Y.; Zheng, L.; Chen, P.-L.; Ziv, Y.; Shiloh,

[112361]Y.; Lee, E. Y.–H. P.; Lee, W.–H.: Functional link of BRCA1 and ataxia

[112362]telangiectasia gene product in DNA damage response.
Nature 406:

[112363]210–215, 2000.

[112364]

[112365]125. Lim, D.–S.; Kim, S.–T.; Xu, B.; Maser, R. S.; Lin, J.; Petrini,

[112366]J. H. J.; Kastan, M. B.: ATM phosphorylates p95/nbs1 in an S-phase

[112367]checkpoint pathway. Nature 404: 613–617, 2000.

[112368]

[112369]126. Lim, D.–S.; Kirsch, D. G.; Canman, C. E.; Ahn, J.–H.; Ziv, Y.;

[112370]Newman, L. S.; Darnell, R. B.; Shiloh, Y.; Kastan, M. B.: ATM binds

[112371]to beta-adaptin in cytoplasmic vesicles. Proc. Nat. Acad. Sci. 95:

[112372]10146–10151, 1998.

[112373]

[112374]127. Lipkowitz, S.; Garry, V. F.; Kirsch, I. R.: Interlocus V–J

recombination

[112375]measures genomic instability in agriculture workers at risk
for lymphoid

[112376]malignancies. Proc. Nat. Acad. Sci. 89: 5301–5305, 1992.

[112377]

[112378]128. Lipkowitz, S.; Stern, M.–H.; Kirsch, I. R.: Hybrid T cell
receptor

[112379]genes formed by interlocus recombination in normal and
ataxia–telangiectasia

[112380]lymphocytes. J. Exp. Med. 172: 409–418, 1990.

[112381]

[112382]129. Lisker, R.; Cobo, A.: Chromosome breakage in
ataxia–telangiectasia.

[112383](Letter) Lancet I: 618, 1970.

[112384]

[112385]130. Littlefield, L. G.; Colyer, S. P.; Joiner, E. E.; DuFrain, R.

[112386]J.; Frome, E.; Cohen, M. M.: Chromosomal radiation sensi-
tivity in

[112387]ataxia telangiectasia long–term lymphoblastoid cell lines.
Cytogenet.

[112388]Cell Genet. 31: 203–213, 1981.

[112389]

[112390]131. Llerena, J., Jr.; Murer-Orlando, M.; McGuire, M.; Zahed, L.;

[112391]Sheridan, R. J.; Berry, A. C.; Bobrow, M.: Spontaneous and induced

[112392]chromosome breakage in chorionic villus samples: a cytogenetic approach

[112393]to first trimester prenatal diagnosis of ataxia telangiectasia syndrome. J.

[112394]Med. Genet. 26: 174-178, 1989.

[112395]

[112396]132. Maserati, E.; Ottolini, A.; Veggiotti, P.; Lanzi, G.; Pasquali,

[112397]F.: Ataxia-without-telangiectasia in two sisters with rearrangements

[112398]of chromosomes 7 and 14. Clin. Genet. 34: 283-287, 1988.

[112399]

[112400]133. Matsuda, Y.; Imai, T.; Shiomi, T.; Saito, T.; Yamauchi, M.; Fukao,

[112401]T.; Akao, Y.; Seki, N.; Ito, H.; Hori, T.: Comparative genome mapping

[112402]of the ataxia-telangiectasia region in mouse, rat, and Syrian hamster. Genomics 34:

[112403]347-352, 1996.

[112404]

[112405]134. McCaw, B. K.; Hecht, F.; Harden, D. G.; Teplitz, R. L.:
Somatic

[112406]rearrangement of chromosome 14 in human lymphocytes.
Proc. Nat. Acad.

[112407]Sci. 72: 2071-2075, 1975.

[112408]

[112409]135. McConville, C. M.; Formstone, C. J.; Hernandez, D.;
Thick, J.;

[112410]Taylor, A. M.: Fine mapping of the chromosome 11q22-23
region using

[112411]PFGE, linkage and haplotype analysis; localization of the
gene for

[112412]ataxia telangiectasia to a 5cM region flanked by NCAM/
DRD2 and STMY/CJ52.75,

[112413]phi 2.22. Nucleic Acids Res. 18: 4335-4343, 1990.

[112414]

[112415]136. McConville, C. M.; Stankovic, T.; Byrd, P. J.; McGuire,

G. M.;

[112416] Yao, Q.-Y.; Lennox, G. G.; Taylor, A. M. R.: Mutations associated

[112417] with variant phenotypes in ataxia-telangiectasia. Am. J. Hum. Genet. 59:

[112418] 320-330, 1996.

[112419]

[112420] 137. McConville, C. M.; Woods, C. G.; Farrall, M.; Metcalfe, J. A.;

[112421] Taylor, A. M. R.: Analysis of 7 polymorphic markers at chromosome

[112422] 11q22-23 in 35 ataxia telangiectasia families; further evidence of

[112423] linkage. Hum. Genet. 85: 215-220, 1990.

[112424]

[112425] 138. McFarlin, D. E.; Strober, W.; Waldmann, T. A.: Ataxia-telangiectasia. Medicine 51:

[112426] 281-314, 1972.

[112427]

[112428] 139. McKusick, V. A.; Cross, H. E.: Ataxia-telangiectasia and Swiss-type

[112429] agammaglobulinemia. Two genetic disorders of the immune mechanism

[112430] in related Amish sibships. J.A.M.A. 195: 739–745, 1966.

[112431]

[112432] 140. Meyn, M. S.: High spontaneous intrachromosomal recombination

[112433] rates in ataxia–telangiectasia. Science 260: 1327–1330, 1993.

[112434]

[112435] 141. Miller, M. E.; Chatten, J.: Ovarian changes in ataxia–telangiectasia. Acta

[112436] Paediat. Scand. 56: 559–561, 1967.

[112437]

[112438] 142. Mohamed, R.; Singh, S. P.; Kumar, S.; Lavin, M. F.: A defect

[112439] in DNA topoisomerase II activity in ataxia–telangiectasia cells. Biochem.

[112440] Biophys. Res. Commun. 149: 233–238, 1987.

[112441]

[112442] 143. Morrell, D.; Chase, C. L.; Swift, M.: Cancers in 44 families

[112443]with ataxia-telangiectasia. Cancer Genet. Cytogenet. 50:
119-123,

[112444]1990.

[112445]

[112446]144. Morrow, D. M.; Tagle, D. A.; Shiloh, Y.; Collins, F. S.;
Hieter,

[112447]P.: TEL1, an *S. cerevisiae* homolog of the human gene mu-
tated in ataxia

[112448]telangiectasia, is functionally related to the yeast check-
point gene

[112449]MEC1. Cell 82: 831-840, 1995.

[112450]

[112451]145. Murnane, J. P.; Painter, R. B.: Complementation of the
effects

[112452]in DNA synthesis in irradiated and unirradiated ataxia-
telangiectasia

[112453]cells. Proc. Nat. Acad. Sci. 79: 1960-1963, 1982.

[112454]

[112455]146. Naito, T.; Matsuura, A.; Ishikawa, F.: Circular chro-
mosome formation

[112456]in a fission yeast mutant defective in two ATM homo-

logues. Nature

[112457]Genet. 20: 203–206, 1998.

[112458]

[112459]147. Nguyen, T. D.; Boyd, J. B.; Green, M. M.: Sensitivity of
Drosophila

[112460]mutants to chemical carcinogens. Mutat. Res. 63: 67–77,
1979.

[112461]

[112462]148. Oxelius, V.–A.; Berkel, A. I.; Hanson, L. A.: IgG2 defi-
ciency

[112463]in ataxia–telangiectasia. New Eng. J. Med. 306: 515–517,
1982.

[112464]

[112465]149. Oxford, J. M.; Harnden, D. G.; Parrington, J. M.; Del-
hanty, J.

[112466]D. A.: Specific chromosome aberrations in ataxia–
telangiectasia. J.

[112467]Med. Genet. 12: 251–262, 1975.

[112468]

[112469]150. Pagani, F.; Buratti, E.; Stuani, C.; Bendix, R.; Dork, T.;
Baralle,

[112470]F. E.: A new type of mutation causes a splicing defect in
ATM. Nature

[112471]Genet. 30: 426–429, 2002.

[112472]

[112473]151. Painter, R. B.; Cramer, P.; Howard, R.; Young, B. R.:
Two forms

[112474]of inhibition of DNA replicon initiation in human cells.In:
Harris,

[112475]C. C.; Cerutti, P. C.: Mechanisms of Chemical Carcinogen-
esis. New

[112476]York: Alan R. Liss (pub.) 1982. Pp. 383–386.

[112477]

[112478]152. Painter, R. B.; Young, B. R.: Radiosensitivity in ataxia-
telangiectasia:

[112479]a new explanation. Proc. Nat. Acad. Sci. 77: 7315–7317,
1980.

[112480]

[112481]153. Paterson, M. C.; Smith, B. P.; Knight, P. A.; Anderson,
A. K.

[112482]: Ataxia telangiectasia: an inherited human disease involv-
ing radiosensitivity,

[112483]malignancy and defective DNA repair.In: Castellani, A.
(ed.): Research

[112484]in Photobiology. New York: Plenum (pub.) 1977. Pp.
207–218.

[112485]

[112486]154. Paterson, M. C.; Smith, B. P.; Lohman, P. H. M.; An-
derson, A.

[112487]K.; Fishman, L.: Defective excision repair of gamma-
ray-damaged DNA

[112488]in human (ataxia-telangiectasia) fibroblasts. Nature 260:
444–447,

[112489]1976.

[112490]

[112491]155. Paterson, M. C.; Smith, P. J.: Ataxia telangiectasia: an
inherited

[112492]human disorder involving hypersensitivity to ionizing ra-
diation and

[112493]related DNA-damaging chemicals. Ann. Rev. Genet. 13:
291–318, 1979.

[112494]

[112495]156. Paulovich, A. G.; Hartwell, L. H.: A checkpoint regu-
lates the

[112496]rate of progression through S phase in *S. cerevisiae* in response to

[112497]DNA damage. *Cell* 82: 841–847, 1995.

[112498]

[112499]157. Pecker, I.; Avraham, K. B.; Gilbert, D. J.; Savitsky, K.; Rotman,

[112500]G.; Harnik, R.; Fukao, T.; Schrock, E.; Hirotsume, S.; Tagle, D. A.;

[112501]Collins, F. S.; Wynshaw-Boris, A.; Ried, T.; Copeland, N. G.; Jenkins,

[112502]N. A.; Shiloh, Y.; Ziv, Y.: Identification and chromosomal localization

[112503]of Atm, the mouse homolog of the ataxia-telangiectasia gene. *Genomics* 35:

[112504]39–45, 1996.

[112505]

[112506]158. Peterson, R. D. A.; Funkhouser, J. D.: Speculations on ataxia-telangiectasia:

[112507]defective regulation of the immunoglobulin gene superfamily. *Immun.*

[112508]*Today* 10: 313–315, 1989.

[112509]

[112510]159. Peterson, R. D. A.; Funkhouser, J. D.: Ataxia–telangiectasia:

[112511]an important clue. (Editorial) New Eng. J. Med. 322: 124–125, 1990.

[112512]

[112513]160. Peterson, R. D. A.; Kelly, W. D.; Good, R. A.: Ataxia–telangiectasia:

[112514]its association with a defective thymus, immunological–deficiency

[112515]disease and malignancy. Lancet I: 1189–1193, 1964.

[112516]

[112517]161. Pippard, E. C.; Hall, A. J.; Barker, D. J. P.; Bridges, B. A.

[112518]; Cancer in homozygotes and heterozygotes of ataxia–telangiectasia

[112519]and xeroderma pigmentosum in Britain. Cancer Res. 48: 2929–2932,

[112520]1988.

[112521]

[112522]162. Rary, J. M.; Bender, M. A.; Kelly, T. E.: A 14/14

marker chromosome

[112523] lymphocyte clone in ataxia telangiectasis. J. Hered. 66:
33–35,

[112524] 1975.

[112525]

[112526] 163. Rary, J. M.; Bender, M. A.; Kelly, T. E.: Cytogenetic
studies

[112527] of ataxia–telangiectasia. (Abstract) Am. J. Hum. Genet. 26:
70,

[112528] 1974.

[112529]

[112530] 164. Reye, C.; Mosman, N. S. W.: Ataxia–telangiectasia.
Am. J. Dis.

[112531] Child. 99: 238–247, 1960.

[112532]

[112533] 165. Richkind, K. E.; Boder, E.; Teplitz, R. L.: Fetal proteins
in

[112534] ataxia–telangiectasia. J.A.M.A. 248: 1346–1347, 1982.

[112535]

[112536] 166. Rosen, F. S.; Harris, N. L.: Case records of the Mas–

sachusetts

[112537]General Hospital: a 30-year-old man with ataxia-telangiectasia and

[112538]dysphagia. New Eng. J. Med. 316: 91-100, 1987.

[112539]

[112540]167. Rosin, M. P.; Ochs, H. D.: In vivo chromosomal instability in

[112541]ataxia-telangiectasia homozygotes and heterozygotes. Hum. Genet. 74:

[112542]335-340, 1986.

[112543]

[112544]168. Rosin, M. P.; Ochs, H. D.; Gatti, R. A.; Boder, E.: Heterogeneity

[112545]of chromosomal breakage levels in epithelial tissue of ataxia-telangiectasia

[112546]homozygotes and heterozygotes. Hum. Genet. 83: 133-138, 1989.

[112547]

[112548]169. Russo, G.; Isobe, M.; Gatti, R.; Finan, J.; Batuman, O.; Huebner,

[112549]K.; Nowell, P. C.; Croce, C. M.: Molecular analysis of a

t(14;14)

[112550]translocation in leukemic T-cells of an ataxia telangiectasia patient. Proc.

[112551]Nat. Acad. Sci. 86: 602-606, 1989.

[112552]

[112553]170. Sanal, O.; Lange, E.; Telatar, M.; Sobel, E.; Salazar-Novak,

[112554]J.; Ersoy, F.; Morrison, A.; Concannon, P.; Tolun, A.; Gatti, R. A.

[112555]: Ataxia-telangiectasia: linkage analysis of chromosome 11q22-23 markers

[112556]in Turkish families. FASEB J. 6: 2848-2852, 1992.

[112557]

[112558]171. Sanal, O.; Wei, S.; Foroud, T.; Malhotra, U.; Concannon, P.;

[112559]Charmley, P.; Salser, W.; Lange, K.; Gatti, R. A.: Further mapping

[112560]of an ataxia-telangiectasia locus to the chromosome 11q23 region. Am.

[112561]J. Hum. Genet. 47: 860-866, 1990.

[112562]

[112563]172. Sandoval, N.; Platzer, M.; Rosenthal, A.; Dork, T.;
Bendix, R.;

[112564]Skawran, B.; Stuhmann, M.; Wegner, R.-D.; Sperling, K.;
Banin, S.;

[112565]Shiloh, Y.; Baumer, A.; Bernthaler, U.; Sennefelder, H.;
Brohm, M.;

[112566]Weber, B. H. F.; Schindler, D.: Characterization of ATM
gene mutations

[112567]in 66 ataxia telangiectasia families. Hum. Molec. Genet. 8:
69–79,

[112568]1999.

[112569]

[112570]173. Sasaki, T.; Tian, H.; Kukita, Y.; Inazuka, M.; Tahira,
T.; Imai,

[112571]T.; Yamauchi, M.; Saito, T.; Hori, T.; Hashimoto–Tamaoki,
T.; Komatsu,

[112572]K.; Nikaido, O.; Hayashi, K.: ATM mutations in patients
with ataxia

[112573]telangiectasia screened by a hierarchical strategy. Hum.
Mutat. 12:

[112574]186–195, 1998.

[112575]

[112576]174. Savitsky, K.; Bar-Shira, A.; Gilad, S.; Rotman, G.; Ziv, Y.;

[112577]Vanagaite, L.; Tagle, D. A.; Smith, S.; Uziel, T.; Sfez, S.; Ashkenazi,

[112578]M.; Pecker, I.; and 18 others: A single ataxia telangiectasia gene

[112579]with a product similar to PI-3 kinase. Science 268: 1749-1753, 1995.

[112580]

[112581]175. Savitsky, K.; Sfez, S.; Tagle, D. A.; Ziv, Y.; Sartiell, A.; Collins,

[112582]F. S.; Shiloh, Y.; Rotman, G.: The complete sequence of the coding

[112583]region of the ATM gene reveals similarity to cell cycle regulators

[112584]in different species. Hum. Molec. Genet. 4: 2025-2032, 1995.

[112585]

[112586]176. Saxon, A.; Stevens, R. H.; Golde, D. W.: Helper and suppressor

[112587]T-lymphocyte leukemia in ataxia-telangiectasia. New Eng. J. Med. 300:

[112588]700–704, 1979.

[112589]

[112590]177. Schaffner, C.; Idler, I.; Stilgenbauer, S.; Dohner, H.;
Lichter,

[112591]P.: Mantle cell lymphoma is characterized by inactivation
of the

[112592]ATM gene. Proc. Nat. Acad. Sci. 97: 2773–2778, 2000.

[112593]

[112594]178. Schaffner, C.; Stilgenbauer, S.; Rappold, G. A.;
Dohner, H.;

[112595]Lichter, P.: Somatic ATM mutations indicate a pathogenic
role of

[112596]ATM in B-cell chronic lymphocytic leukemia. Blood 94:
748–753, 1999.

[112597]

[112598]179. Schalch, D. S.; McFarlin, D. E.; Barlow, M. H.: An un-
usual form

[112599]of diabetes mellitus in ataxia–telangiectasia. New Eng. J.
Med. 282:

[112600]1396–1402, 1970.

[112601]

- [112602]180. Scheres, J. M. J. C.; Hustinx, T. W. J.; Weemaes, C. M. R.:
[112603]Chromosome 7 in ataxia-telangiectasia. J. Pediat. 97:
440-441, 1980.
[112604]
- [112605]181. Scott, S. P.; Bendix, R.; Chen, P.; Clark, R.; Dork, T.;
Lavin,
[112606]M. F.: Missense mutations but not allelic variants alter the
function
[112607]of ATM by dominant interference in patients with breast
cancer. Proc.
[112608]Nat. Acad. Sci. 99: 925-930, 2002.
[112609]
- [112610]182. Sedgwick, R. P.; Boder, E.: Ataxia-telangiectasia.In:
Vinken,
[112611]P. J.; Bruyn, G. W. (eds.): Handbook of Clinical Neurology.
Amsterdam:
[112612]North-Holland Publishing Co. (pub.) 14: 1972. Pp.
267-339.
[112613]
- [112614]183. Shaham, M.; Becker, Y.: The ataxia telangiectasia
clastogenic

[112615]factor is a low molecular weight peptide. Hum. Genet. 58:
422–424,

[112616]1981.

[112617]

[112618]184. Shiloh, Y.; Parshad, R.; Frydman, M.; Sanford, K. K.;
Portnoi,

[112619]S.; Ziv, Y.; Jones, G. M.: G(2) chromosomal radiosensitivity
in families

[112620]with ataxia–telangiectasia. Hum. Genet. 84: 15–18, 1989.

[112621]

[112622]185. Shultz, L. D.; Sweet, H. O.; Davisson, M. T.; Coman,
D. R.:

[112623]'Wasted,' a new mutant of the mouse with abnormalities
characteristic

[112624]of ataxia telangiectasia. Nature 297: 402–404, 1982.

[112625]

[112626]186. Shuster, J.; Hart, Z.; Stimson, C. W.; Brough, A. J.;
Poulik,

[112627]M. D.: Ataxia–telangiectasia with cerebellar tumor. Pedi-
atrics 37:

[112628]776–786, 1966.

[112629]

[112630]187. Sobel, E.; Lange, E.; Jaspers, N. G. J.; Chessa, L.;
Sanal, O.;

[112631]Shiloh, Y.; Taylor, A. M. R.; Weemaes, C. M. A.; Lange, K.;
Gatti,

[112632]R. A.: Ataxia-telangiectasia: linkage evidence for genetic
heterogeneity.

[112633](Letter) Am. J. Hum. Genet. 50: 1343-1348, 1992.

[112634]

[112635]188. Sourander, P.; Bonnevier, J. O.; Olsson, Y.: A case of
ataxia-telangiectasia

[112636]with lesions in the spinal cord. Acta Neurol. Scand. 42:
354-366,

[112637]1966.

[112638]

[112639]189. Spring, K.; Ahangari, F.; Scott, S. P.; Waring, P.; Pur-
die, D.

[112640]M.; Chen, P. C.; Hourigan, K.; Ramsay, J.; McKinnon, P. J.;
Swift,

[112641]M.; Lavin, M. F.: Mice heterozygous for mutation in Atm,
the gene

[112642]involved in ataxia-telangiectasia, have heightened susceptibility

[112643]to cancer. *Nature Genet.* 32: 185–190, 2002.

[112644]

[112645]190. Stankovic, T.; Kidd, A. M. J.; Sutcliffe, A.; McGuire, G. M.;

[112646]Robinson, P.; Weber, P.; Bedenham, T.; Bradwell, A. R.; Easton, D.

[112647]F.; Lennox, G. G.; Haites, N.; Byrd, P. J.; Taylor, A. M. R.: ATM

[112648]mutations and phenotypes in ataxia-telangiectasia families in the

[112649]British Isles: expression of mutant ATM and the risk of leukemia,

[112650]lymphoma, and breast cancer. *Am. J. Hum. Genet.* 62: 334–345, 1998.

[112651]

[112652]191. Stern, M.-H.; Zhang, F.; Griscelli, C.; Thomas, G.; Aurias, A.

[112653]: Molecular characterization of different ataxia telangiectasia T-cell

[112654]clones. I. A common breakpoint at the 14q11.2 band

splits the T-cell

[112655]receptor alpha-chain gene. Hum. Genet. 78: 33-36, 1988.

[112656]

[112657]192. Stern, M.-H.; Zhang, F.; Thomas, G.; Griscelli, C.; Aurias, A.

[112658]; Molecular characterization of ataxia telangiectasia T cell clones.

[112659]III. Mapping the 14q32.1 distal breakpoint. Hum. Genet. 81: 18-22,

[112660]1988.

[112661]

[112662]193. Stewart, G. S.; Maser, R. S.; Stankovic, T.; Bressan, D. A.;

[112663]Kaplan, M. I.; Jaspers, N. G. J.; Raams, A.; Byrd, P. J.; Petrini,

[112664]J. H. J.; Taylor, A. M. R.: The DNA double-strand break repair gene

[112665]hMRE11 is mutated in individuals with an ataxia-telangiectasia-like

[112666]disorder. Cell 99: 577-587, 1999.

[112667]

[112668]194. Stilgenbauer, S.; Schaffner, C.; Litterst, A.; Liebisch, P.;

[112669]Gilad, S.; Bar-Shira, A.; James, M. R.; Lichter, P.; Dohner, H.:

[112670]Biallelic mutations in the ATM gene in T-prolymphocytic leukemia. Nature

[112671]Med. 3: 1155–1159, 1997.

[112672]

[112673]195. Stilgenbauer, S.; Winkler, D.; Ott, G.; Schaffner, C.; Leupolt,

[112674]E.; Bentz, M.; Moller, P.; Muller-Hermelink, H.-K.; James, M. R.;

[112675]Lichter, P.; Dohner, H.: Molecular characterization of 11q deletions

[112676]points to a pathogenic role of the ATM gene in mantle cell lymphoma. Blood 94:

[112677]3262–3264, 1999.

[112678]

[112679]196. Sugimoto, T.; Kidowaki, T.; Sawada, T.; Ohtsuka-Urano, T.; Kusunoki,

[112680]T.: Ataxia-telangiectasia associated with non-T, non-B cell acute

[112681]lymphocytic leukemia. Acta Paediat. Scand. 71: 509–510,
1982.

[112682]

[112683]197. Swift, M.: Cancer risk counseling. (Letter) Science
210: 1074,

[112684]1980.

[112685]

[112686]198. Swift, M.; Morrell, D.; Cromartie, E.; Chamberlin, A.
R.; Skolnick,

[112687]M. H.; Bishop, D. T.: The incidence and gene frequency of
ataxia–telangiectasia

[112688]in the United States. Am. J. Hum. Genet. 39: 573–583,
1986.

[112689]

[112690]199. Swift, M.; Morrell, D.; Massey, R. B.; Chase, C. L.: In-
cidence

[112691]of cancer in 161 families affected by ataxia–telangiectasia.
New

[112692]Eng. J. Med. 325: 1831–1836, 1991.

[112693]

[112694]200. Swift, M.; Reitnauer, P. J.; Morrell, D.; Chase, C. L.:

Breast

[112695]and other cancers in families with ataxia–telangiectasia.

New Eng.

[112696]J. Med. 316: 1289–1294, 1987.

[112697]

[112698]201. Swift, M. R.; Sholman, L.; Perry, M.; Chase, C.: Malignant neoplasms

[112699]in the families of patients with ataxia–telangiectasia. Cancer Res. 36:

[112700]209–215, 1976.

[112701]

[112702]202. Tadjoeidin, M. K.; Fraser, F. C.: Heredity of ataxia–telangiectasia

[112703](Louis–Bar syndrome). Am. J. Dis. Child. 110: 64–68, 1965.

[112704]

[112705]203. Taniguchi, T.; Garcia–Higuera, I.; Xu, B.; Andreassen, P. R.;

[112706]Gregory, R. C.; Kim, S.–T.; Lane, W. S.; Kastan, M. B.; D'Andrea,

[112707]A. D.: Convergence of the Fanconi anemia and ataxia

telangiectasia

[112708]signaling pathways. Cell 109: 459–472, 2002.

[112709]

[112710]204. Taylor, A. M. R.; Flude, E.; Laher, B.; Stacey, M.; McKay, E.;

[112711]Watt, J.; Green, S. H.; Harding, A. E.: Variant forms of ataxia telangiectasia. J.

[112712]Med. Genet. 24: 669–677, 1987.

[112713]

[112714]205. Taylor, A. M. R.; Harnden, D. G.; Arlett, C. F.; Harcourt, S.

[112715]A.; Lehmann, A. R.; Stevens, S.; Bridges, B. A.: Ataxia–telangiectasia:

[112716]a human mutation with abnormal radiation sensitivity. Nature 258:

[112717]427–429, 1975.

[112718]

[112719]206. Taylor, A. M. R.; Metcalfe, J. A.; Oxford, J. M.; Harnden, D.

[112720]G.: Is chromatid–type damage in ataxia–telangiectasia after irradiation

[112721]at G(0) a consequence of defective repair? Nature 260:
441–443,

[112722]1976.

[112723]

[112724]207. Tchirkov, A.; Bay, J.–O.; Pernin, D.; Bignon, Y.–J. Rio,
P.;

[112725]Granchó, M.; Kwiatkowski, F.; Giollant, M.; Malet, P.; Ver-
relle, P.

[112726]: Detection of heterozygous carriers of the ataxia–
telangiectasia

[112727](ATM) gene by G(2) phase chromosomal radiosensitivity of
peripheral

[112728]blood lymphocytes. Hum. Genet. 101: 312–316, 1997.

[112729]

[112730]208. Telatar, M.; Teraoka, S.; Wang, Z.; Chun, H. H.; Liang,
T.; Castellvi–Bel,

[112731]S.; Udar, N.; Borresen–Dale, A.–L.; Chessa, L.; Berna–
towska–Matuskiewicz,

[112732]E.; Porras, O.; Watanabe, M.; Junker, A.; Concannon, P.;
Gatti, R.

[112733]A.: Ataxia–telangiectasia: identification and detection of
founder–effect

[112734] mutations in the ATM gene in ethnic populations. Am. J. Hum. Genet. 62:

[112735] 86–97, 1998.

[112736]

[112737] 209. Telatar, M.; Wang, Z.; Castellvi–Bel, S.; Tai, L.–Q.; Sheikhavandi,

[112738] S.; Regueiro, J. R.; Porras, O.; Gatti, R. A.: A model for ATM heterozygote

[112739] identification in a large population: four founder–effect ATM mutations

[112740] identify most of Costa Rican patients with ataxia telangiectasia. Molec.

[112741] Genet. Metab. 64: 36–43, 1998.

[112742]

[112743] 210. Teplitz, R. L.: Ataxia–telangiectasia. Arch. Neurol. 35: 553–554,

[112744] 1978.

[112745]

[112746] 211. Teraoka, S. N.; Telatar, M.; Becker–Catania, S.; Liang, T.; Onengut,

[112747] S.; Tolun, A.; Chessa, L.; Sanal, O.; Bernatowska, E.; Gatti,

R. A.;

[112748]Concannon, P.: Splicing defects in the ataxia-telangiectasia gene,

[112749]ATM: underlying mutations and consequences. Am. J. Hum. Genet. 64:

[112750]1617-1631, 1999.

[112751]

[112752]212. Thibaut, S.; Sass, U.; Khoury, A.; Simonart, J.-M.: Ataxia-telangiectasia

[112753]and necrobiosis lipoidica: an explainable association. Europ. J.

[112754]Derm. 4: 509-513, 1994.

[112755]

[112756]213. Toledano, S. R.; Lang, B. J.: Ataxia-telangiectasia and acute

[112757]lymphoblastic leukemia. Cancer 45: 1675-1678, 1980.

[112758]

[112759]214. Toyoshima, M.; Hara, T.; Zhang, H.; Yamamoto, T.; Akaboshi, S.;

[112760]Nanba, E.; Ohno, K.; Hori, N.; Sato, K.; Takeshita, K.: Ataxia-telangiectasia

[112761] without immunodeficiency: novel point mutations within
and adjacent

[112762] to the phosphatidylinositol 3-kinase-like domain. Am. J.
Med. Genet. 75:

[112763] 141-144, 1998.

[112764]

[112765] 215. Tse, A. G. D.; Barclay, A. N.; Watts, A.; Williams, A. F.:
A

[112766] glycopospholipid tail at the carboxyl terminus of the
Thy-1 glycoprotein

[112767] of neurons and thymocytes. Science 230: 1003-1008,
1985.

[112768]

[112769] 216. Uziel, T.; Savitsky, K.; Platzer, M.; Ziv, Y.; Helbitz, T.;
Nehls,

[112770] M.; Boehm, T.; Rosenthal, A.; Shiloh, Y.; Rotman, G.: Ge-
nomic organization

[112771] of the ATM gene. Genomics 33: 317-320, 1996.

[112772]

[112773] 217. van Belzen, M. J.; Hiel, J. A. P.; Weemaes, C. M. R.;
Gabreels,

[112774]F. J. M.; van Engelen, B. G. M.; Smeets, D. F. C. M.; van den Heuvel,

[112775]L. P. W. J.: A double missense mutation in the ATM gene of a Dutch

[112776]family with ataxia telangiectasia. Hum. Genet. 102: 187–191, 1998.

[112777]

[112778]218. Vincent, R. A., Jr.; Sheridan, R. B., III; Huang, P. C.: DNA

[112779]strand breakage repair in ataxia–telangiectasia fibroblast–like cells. Mutat.

[112780]Res. 33: 357–366, 1975.

[112781]

[112782]219. Vorechovsky, I.; Luo, L.; Dyer, M. J. S.; Catovsky, D.; Amlot,

[112783]P. L.; Yaxley, J. C.; Foroni, L.; Hammarstrom, L.; Webster, A. D.

[112784]B.; Yuille, M. A. R.: Clustering of missense mutations in the ataxia–telangiectasia

[112785]gene in a sporadic T–cell leukaemia. Nature Genet. 17: 96–99, 1997.

[112786]

[112787]220. Vorechovsky, I.; Luo, L.; Lindblom, A.; Negrini, M.; Webster,

[112788]A. D.; Croce, C. M.; Hammarstrom, L.: ATM mutations in cancer families. Cancer

[112789]Res. 56: 4130–4133, 1996.

[112790]

[112791]221. Waldmann, T. A.; McIntire, K. R.: Serum- α -pha-fetoprotein levels

[112792]in patients with ataxia-telangiectasia. Lancet II: 1112–1115, 1972.

[112793]

[112794]222. Waldmann, T. A.; Misiti, J.; Nelson, D. L.; Kraemer, K. H.:

[112795]Ataxia-telangiectasia: a multisystem hereditary disease with immunodeficiency,

[112796]impaired organ maturation, x-ray hypersensitivity, and a high incidence

[112797]of neoplasia. Ann. Intern. Med. 99: 367–379, 1983.

[112798]

[112799]223. Wang, Y.; Cortez, D.; Yazdi, P.; Neff, N.; Elledge, S. J.; Qin,

[112800]J.: BASC, a super complex of BRCA1-associated proteins involved in

[112801]the recognition and repair of aberrant DNA structures.

Genes Dev. 14:

[112802]927–939, 2000.

[112803]

[112804]224. Wang, Y. A.; Elson, A.; Leder, P.: Loss of p21 in-
creases sensitivity

[112805]to ionizing radiation and delays the onset of lymphoma in
atm-deficient

[112806]mice. Proc. Nat. Acad. Sci. 94: 14590–14595, 1997.

[112807]

[112808]225. Watanabe, A.; Hanazono, H.; Sogawa, H.; Takaya, H.:
Stomach

[112809]cancer in a 14-year-old-boy with ataxia-telangiectasia.
Tohoku J.

[112810]Exp. Med. 121: 127–131, 1977.

[112811]

[112812]226. Waterman, M. J. F.; Stavridi, E. S.; Waterman, J. L. F.;
Halazonetis,

[112813]T. D.: ATM-dependent activation of p53 involves dephos-

phorylation

[112814]and association with 14-3-3 proteins. Nature Genet. 19:
175-178,

[112815]1998.

[112816]

[112817]227. Watts, J. A.; Morley, M.; Burdick, J. T.; Fiori, J. L.;
Ewens,

[112818]W. J.; Spielman, R. S.; Cheung, V. G.: Gene expression
phenotype

[112819]in heterozygous carriers of ataxia telangiectasia. Am. J.
Hum. Genet. 71:

[112820]791-800, 2002.

[112821]

[112822]228. Weinstein, S.; Scottolini, A. G.; Loo, S. Y. T.; Caldwell,
P.

[112823]C.; Bhagavan, N. V.: Ataxia telangiectasia with hepatocel-
lular carcinoma

[112824]in a 15-year-old girl and studies of her kindred. Arch.
Path. Lab.

[112825]Med. 109: 1000-1004, 1985.

[112826]

[112827]229. Welshimer, K.; Swift, M.: Congenital malformations and developmental disabilities in ataxia-telangiectasia, Fanconi anemia, and xeroderma pigmentosum families. Am. J. Hum. Genet. 34: 781-793, 1982.

[112830]

[112831]230. Westphal, C. H.; Rowan, S.; Schmaltz, C.; Elson, A.; Fisher, D. E.; Leder, P.: atm and p53 cooperate in apoptosis and suppression of tumorigenesis, but not in resistance to acute radiation toxicity. Nature Genet. 16: 397-401, 1997.

[112835]

[112836]231. Woods, C. G.; Bunday, S. E.; Taylor, A. M. R.: Unusual features in the inheritance of ataxia telangiectasia. Hum. Genet. 84: 555-562, 1990.

[112839]

[112840]232. Woods, C. G.; Taylor, A. M. R.: Ataxia telangiectasia
in the

[112841]British Isles: the clinical and laboratory features of 70 af-
fected

[112842]individuals. Quart. J. Med. 82: 169–179, 1992.

[112843]

[112844]233. Wooster, R.; Ford, D.; Mangion, J.; Ponder, B. A. J.;
Peto, J.;

[112845]Easton, D. F.; Stratton, M. R.: Absence of linkage to the
ataxia

[112846]telangiectasia locus in familial breast cancer. Hum. Genet.
92:

[112847]91–94, 1993.

[112848]

[112849]234. Worgul, B. V.; Smilenov, L.; Brenner, D. J.; Junk, A.;
Zhou,

[112850]W.; Hall, E. J.: Atm heterozygous mice are more sensitive
to radiation-induced

[112851]cataracts than are their wild-type counterparts. Proc. Nat.
Acad.

[112852]Sci. 99: 9836–9839, 2002.

[112853]

[112854]235. Wright, J.; Teraoka, S.; Onengut, S.; Tolun, A.; Gatti, R. A.;

[112855]Ochs, H. D.; Concannon, P.: A high frequency of distinct ATM gene

[112856]mutations in ataxia-telangiectasia. Am. J. Hum. Genet. 59: 839-846,

[112857]1996.

[112858]

[112859]236. Wu, X.; Ranganathan, V.; Weisman, D. S.; Heine, W. F.; Ciccone,

[112860]D. N.; O'Neill, T. B.; Crick, K. E.; Pierce, K. A.; Lane, W. S.; Rathbun,

[112861]G.; Livingston, D. M.; Weaver, D. T.: ATM phosphorylation of Nijmegen

[112862]breakage syndrome protein is required in a DNA damage response. Nature 405:

[112863]477-482, 2000.

[112864]

[112865]237. Xia, Y.-R.; Welch, C. L.; Warden, C. H.; Lange, E.; Fukao, T.;

[112866]Lusis, A. J.; Gatti, R. A.: Assignment of the mouse ataxia-telangiectasia

[112867]gene (Atm) to mouse chromosome 9. Mammalian Genome 7: 554-555, 1996.

[112868]

[112869]238. Xu, Y.; Ashley, T; Brainerd, E. E.; Bronson, R. T.; Meyn, M.

[112870]S.; Baltimore, D.: Targeted disruption of ATM leads to growth retardation,

[112871]chromosomal fragmentation during meiosis, immune defects, and thymic

[112872]lymphoma. Genes Dev. 10: 2411-2422, 1996.

[112873]

[112874]239. Xu, Y.; Baltimore, D.: Dual roles of ATM in the cellular response

[112875]to radiation and in cell growth control. Genes Dev. 10: 2401-2410,

[112876]1996.

[112877]

[112878]240. Ying, K. L.; Decoteau, W. E.: Cytogenetic anomalies in a patient

[112879]with ataxia, immune deficiency, and high alpha-fetoprotein in the

[112880]absence of telangiectasia. Cancer Genet. Cytogenet. 4: 311-317,

[112881]1981.

[112882]

[112883]241. Yount, W. J.: IgG2 deficiency and ataxia-telangiectasia. (Editorial) New

[112884]Eng. J. Med. 306: 541-543, 1982.

[112885]

[112886]242. Zadik, Z.; Levin, S.; Prager-Lewin, R.; Laron, Z.: Gonadal dysfunction

[112887]in patients with ataxia telangiectasia. Acta Paediat. Scand. 67:

[112888]477-479, 1978.

[112889]

[112890]243. Zakian, V. A.: ATM-related genes: what do they tell us about

[112891]functions of the human gene? Cell 82: 685-687, 1995.

[112892]

[112893]244. Zhang, F.; Stern, M.-H.; Thomas, G.; Aurias, A.:

Molecular characterization

[112894]of ataxia telangiectasia T cell clones. II. The clonal inv(14)

in

[112895]ataxia telangiectasia differs from the inv(14) in T cell lym-

phoma. Hum.

[112896]Genet. 78: 316-319, 1988.

[112897]

[112898]245. Zhang, N.; Chen, P.; Khanna, K. K.; Scott, S.; Gatei,

M.; Kozlov,

[112899]S.; Watters, D.; Spring, K.; Yen, T.; Lavin, M. F.: Isolation of

[112900]full-length ATM cDNA and correction of the ataxia-

telangiectasia cellular

[112901]phenotype. Proc. Nat. Acad. Sci. 94: 8021-8026, 1997.

[112902]

[112903]246. Zhao, S.; Weng, Y.-C.; Yuan, S.-S. F.; Lin, Y.-T.; Hsu,

H.-C.;

[112904]Lin, S.-C. J.; Gerbino, E.; Song, M.; Zdzienicka, M. Z.; Gatti,

R.

[112905]A.; Shay, J. W.; Ziv, Y.; Shiloh, Y.; Lee, E. Y.-H. P.: Func-

tional

[112906]link between ataxia–telangiectasia and Nijmegen breakage syndrome

[112907]gene products. Nature 405: 473–477, 2000.

[112908]

[112909]247. Ziv, Y.; Amiel, A.; Jaspers, N. G. J.; Berkel, A. I.; Shiloh,

[112910]Y.: Ataxia–telangiectasia: a variant with altered in vitro phenotype

[112911]of fibroblast cells. Mutat. Res. 210: 211–219, 1989.

[112912]

[112913]248. Ziv, Y.; Frydman, M.; Lange, E.; Zelnik, N.; Rotman, G.; Julier,

[112914]C.; Jaspers, N. G. J.; Dagan, Y.; Abeliovicz, D.; Dar, H.; Borochowitz,

[112915]Z.; Lathrop, M.; Gatti, R. A.; Shiloh, Y.: Ataxia–telangiectasia:

[112916]linkage analysis in highly inbred Arab and Druze families and differentiation

[112917]from an ataxia–microcephaly–cataract syndrome. Hum. Genet. 88: 619–626,

[112918]1992.

[112919]

[112920]249. Ziv, Y.; Rotman, G.; Frydman, M.; Dagan, J.; Cohen, T.; Foroud,

[112921]T.; Gatti, R. A.; Shiloh, Y.: The ATC (ataxia-telangiectasia complementation

[112922]group C) locus localizes to 11q22-q23. Genomics 9: 373-375, 1991.

[112923]19907.

[112924]Date, H.; Onodera, O.; Tanaka, H.; Iwabuchi, K.; Uekawa, K.; Igarashi,

[112925]S.; Koike, R.; Hiroi, T.; Yuasa, T.; Awaya, Y.; Sakai, T.; and 9

[112926]others: Early-onset ataxia with ocular motor apraxia and hypoalbuminemia

[112927]is caused by mutations in a new HIT superfamily gene. Nature Genet. 29:

[112928]184-188, 2001.

[112929]19908.

[112930]Moreira, M.-C.; Barbot, C.; Tachi, N.; Kozuka, N.; Uchida, E.;

[112931]Gibson, T.; Mendonca, P.; Costa, M.; Barros, J.; Yanagisawa, T.; Watanabe,

[112932]M.; Ikeda, Y.; Aoki, M.; Nagata, T.; Coutinho, P.; Sequeiros, J.;

[112933]Koenig, M.: The gene mutated in ataxia–oculomotor apraxia 1 encodes

[112934]the new HIT/Zn–finger protein aprataxin. *Nature Genet.* 29: 189–193,

[112935]2001.

[112936]19909.

[112937]Shelley, W. B.; Rawnsley, H. M.; Morrow, G., III: Pyridoxine–dependent

[112938]hair pigmentation in association with homocystinuria. *Arch. Derm.* 106:

[112939]228–230, 1972.

[112940]

[112941]100. Shih, V. E.; Efron, M. L.: Pyridoxine–unresponsive homocystinuria. *New*

[112942]*Eng. J. Med.* 283: 1206–1208, 1970.

[112943]

[112944]101. Shih, V. E.; Fringer, J. M.; Mandell, R.; Kraus, J. P.; Berry,

[112945]G. T.; Heidenreich, R. A.; Korson, M. S.; Levy, H. L.;

Ramesh, V.

[112946] A missense mutation (I278T) in the cystathionine beta-synthase gene

[112947] prevalent in pyridoxine-responsive homocystinuria and associated with

[112948] mild clinical phenotype. *Am. J. Hum. Genet.* 57: 34–39, 1995.

[112949]

[112950] 102. Shipman, R. T.; Townley, R. R. W.; Danks, D. M.: Homocystinuria,

[112951] Addisonian pernicious anaemia, and partial deletion of a G chromosome. *Lancet* II:

[112952] 693–694, 1969.

[112953]

[112954] 103. Skovby, F.: Homocystinuria: clinical, biochemical and genetic

[112955] aspects of cystathionine beta-synthase and its deficiency in man. *Acta*

[112956] *Paediat. Scand.* 321 (suppl.): 1–21, 1985.

[112957]

[112958] 104. Skovby, F.; Krassikoff, N.; Francke, U.: Assignment of

the gene

[112959]for cystathionine beta-synthase to human chromosome
21 in somatic

[112960]cell hybrids. Hum. Genet. 65: 291-294, 1984.

[112961]

[112962]105. Skovby, F.; Krassikoff, N.; Francke, U.: Assignment of
the gene

[112963]for cystathionine beta-synthase (CBS) to human chromo-
some 21 in somatic

[112964]cell hybrids. (Abstract) Cytogenet. Cell Genet. 37: 585
only, 1984.

[112965]

[112966]106. Skovby, F.; Kraus, J.; Redlich, C.; Rosenberg, L. E.:
Immunochemical

[112967]studies on cultured fibroblasts from patients with homo-
cystinuria

[112968]due to cystathionine beta-synthase deficiency. Am. J.
Hum. Genet. 34:

[112969]73-83, 1982.

[112970]

[112971]107. Skovby, F.; Kraus, J. P.; Rosenberg, L. E.: Homo-

cystinuria:

[112972]biogenesis of cystathionine beta-synthase subunits in
cultured fibroblasts

[112973]and in an in vitro translation system programmed with fi-
broblast messenger

[112974]RNA. Am. J. Hum. Genet. 36: 452-459, 1984.

[112975]

[112976]108. Spaeth, G. L.; Barber, G. W.: Prevalence of homo-
cystinuria among

[112977]the mentally retarded: evaluation of a specific screening
test. Pediatrics 40:

[112978]586-589, 1967.

[112979]

[112980]109. Sperandio, M. P.; de Franchis, R.; Andria, G.; Sebas-
tio, G.:

[112981]A 68-bp insertion found in a homocystinuric patient is a
common variant

[112982]and is skipped by alternative splicing of the cystathionine
beta-synthase

[112983]mRNA. (Letter) Am. J. Hum. Genet. 59: 1391-1393, 1996.

[112984]

- [112985]110. Sperandeo, M. P.; Panico, M.; Pepe, A.; Candito, M.;
de Franchis,
[112986]R.; Kraus, J. P.; Andria, G.; Sebastio, G.: Molecular analysis
of
[112987]patients affected by homocystinuria due to cystathionine
beta-synthase
[112988]deficiency: report of a new mutation in exon 8 and a dele-
tion in intron
[112989]11. J. Inherit. Metab. Dis. 18: 211-214, 1995.
[112990]
[112991]111. Streeten, B. W.: The nature of the ocular zonule.
Trans. Am.
[112992]Ophthal. Soc. 80: 823-854, 1982.
[112993]
[112994]112. Stubbs, L.; Kraus, J.; Lehrach, H.: The alpha-
A-crystallin and
[112995]cystathionine beta-synthase genes are physically very
closely linked
[112996]in proximal mouse chromosome 17. Genomics 7:
284-288, 1990.
[112997]

[112998]113. Tada, K.; Yoshida, T.; Hirono, H.; Arakawa, T.: Homocystinuria:

[112999]amino acid pattern of the liver. Tohoku J. Exp. Med. 92: 325-332,

[113000]1967.

[113001]

[113002]114. Tsai, M. Y.; Bignell, M.; Schwichtenberg, K.; Hanson, N. Q.:

[113003]High prevalence of a mutation in the cystathionine beta-synthase gene. Am.

[113004]J. Hum. Genet. 59: 1262-1267, 1996.

[113005]

[113006]115. Uhlemann, E. R.; TenPas, J. H.; Lucky, A. W.; Schulman, J. D.;

[113007]Mudd, S. H.; Shulman, N. R.: Platelet survival and morphology in

[113008]homocystinuria due to cystathionine synthase deficiency. New Eng.

[113009]J. Med. 295: 1283-1286, 1976.

[113010]

[113011]116. Uhlendorf, B. W.; Mudd, S. H.: Cystathionine synthase

in tissue

[113012]culture derived from human skin: enzyme defect in homocystinuria. Science 160:

[113013]1007–1009, 1968.

[113014]

[113015]117. Visy, J. M.; Le Coz, P.; Chadeaux, B.; Fressinaud, C.; Woimant,

[113016]F.; Marquet, J.; Zittoun, J.; Visy, J.; Vallat, J. M.; Haguenu, M.

[113017]: Homocystinuria due to 5,10-methylenetetrahydrofolate reductase deficiency

[113018]revealed by stroke in adult siblings. Neurology 41: 1313–1315, 1991.

[113019]

[113020]118. Wadman, S. K.; Cats, B. P.; de Bree, P. K.: Sulfite oxidase

[113021]deficiency and the detection of urinary sulfite. (Letter) Europ.

[113022]J. Pediat. 141: 62–63, 1983.

[113023]

[113024]119. Watanabe, M.; Osada, J.; Aratani, Y.; Kluckman, K.;

Reddick,

[113025]R.; Malinow, M. R.; Maeda, N.: Mice deficient in cystathionine beta-synthase:

[113026]animal models for mild and severe homocyst(e)inemia.

Proc. Nat. Acad.

[113027]Sci. 92: 1585-1589, 1995.

[113028]

[113029]120. Wilcken, D. E.; Wilcken, B.: The pathogenesis of coronary artery

[113030]disease: a possible role for methionine metabolism. J.

Clin. Invest. 57:

[113031]1079-1082, 1976.

[113032]

[113033]121. Wilcken, D. E. L.; Dudman, N. P. B.; Tyrrell, P. A.: Homocystinuria

[113034]due to cystathionine beta-synthase deficiency--the effects of betaine

[113035]treatment in pyridoxine-responsive patients. Metabolism 34: 1115-1121,

[113036]1985.

[113037]

[113038]122. Wilcken, D. E. L.; Wilcken, B.; Dudman, N. P. B.; Tyrrell, P.

[113039]A.: Homocystinuria--the effects of betaine in the treatment of patients

[113040]not responsive to pyridoxine. New Eng. J. Med. 309: 448-453, 1983.

[113041]

[113042]123. Wong, P. W. K.; Schwarz, V.; Komrower, G. M.: The biosynthesis

[113043]of cystathionine in patients with homocystinuria. Pediat. Res. 2:

[113044]149-160, 1968.

[113045]

[113046]124. Yaghmai, R.; Kashani, A. H.; Geraghty, M. T.; Okoh, J.; Pomper,

[113047]M.; Tangerman, A.; Wagner, C.; Stabler, S. P.; Allen, R. H.; Mudd,

[113048]S. H.; Braverman, N.: Progressive cerebral edema associated with

[113049]high methionine levels and betaine therapy in a patient with cystathionine

[113050]beta-synthase (CBS) deficiency. Am. J. Med. Genet. 108:

57-63, 2002.

[113051]

[113052]125. Yap, S.; Naughten, E.: Homocystinuria due to cystathionine beta-synthase

[113053]deficiency in Ireland: 25 years' experience of a newborn screened

[113054]and treated population with reference to clinical outcome and biochemical

[113055]control. J. Inherit. Metab. Dis. 21: 738-747, 1998.

[113056]

[113057]126. Yap, S.; Rushe, H.; Howard, P. M.; Naughten, E. R.: The intellectual

[113058]abilities of early-treated individuals with pyridoxine-nonresponsive

[113059]homocystinuria due to cystathionine beta-synthase deficiency. J.

[113060]Inherit. Metab. Dis. 24: 437-447, 2001.

[113061]19910.

[113062]Burchell, A.; Jung, R. T.; Lang, C. C.; Bennet, W.; Shepherd, A.

[113063]N.: Diagnosis of type 1a and type 1c glycogen storage

diseases in

[113064]adults. Lancet I: 1059-1062, 1987.

[113065]19911.

[113066]Burchell, A.; Waddell, I. D.: Diagnosis of a novel glycogen storage

[113067]disease: type 1aSP. J. Inherit. Metab. Dis. 13: 247-249, 1990.

[113068]19912.

[113069]Chen, Y.-T.; Coleman, R. A.; Scheinman, J. I.; Kolbeck, P. C.;

[113070]Sidbury, J. B.: Renal disease in type I glycogen storage disease. New

[113071]Eng. J. Med. 318: 7-11, 1988.

[113072]19913.

[113073]Chen, Y.-T.; Cornblath, M.; Sidbury, J. B.: Cornstarch therapy

[113074]in type I glycogen-storage disease. New Eng. J. Med. 310: 171-175,

[113075]1984.

[113076]19914.

[113077]Chen, Y.-T.; Scheinman, J. I.; Park, H. K.; Coleman, R. A.;

Roe,

[113078]C. R.: Amelioration of proximal renal tubular dysfunction
in type

[113079]glycogen storage disease with dietary therapy. New Eng.
J. Med. 323:

[113080]590–593, 1990.

[113081]19915.

[113082]Chevalier–Porst, F.; Bozon, D.; Bonardot, A.–M.; Bruni, N.;
Mithieux,

[113083]G.; Mathieu, M.; Maire, I.: Mutation analysis in 24 French
patients

[113084]with glycogen storage disease type 1a. J. Med. Genet. 33:
358–360,

[113085]1996.

[113086]19916.

[113087]Chou, J. Y.: Personal Communication. Bethesda, Md.
11/4/1993.

[113088]19917.

[113089]Chou, J.Y.; Mansfield, B.C.: Molecular genetics of type 1
glycogen

[113090]storage diseases. Trends Endocr. Metab. 10: 104–113,

1999.

[113091]19918.

[113092]Cohen, J. L.; Vinik, A.; Faller, J.; Fox, I. H.: Hyperuricemia

[113093]in glycogen storage disease type I: contributions by hypoglycemia

[113094]and hyperglucagonemia to increased urate production. J.

Clin. Invest. 75:

[113095]251–257, 1985.

[113096]19919.

[113097]Cori, G. T.; Cori, C. F.: Glucose-6-phosphatase of the liver

[113098]in glycogen storage disease. J. Biol. Chem. 199: 661–667, 1952.

[113099]19920.

[113100]Emmett, M.; Narins, R. G.: Renal transplantation in type I glycogenosis:

[113101]failure to improve glucose metabolism. J.A.M.A. 239: 1642–1644,

[113102]1978.

[113103]19921.

[113104]Faivre, L.; Houssin, D.; Valayer, J.; Brouard, J.; Hadchouel,

[113105]M.; Bernard, O.: Long-term outcome of liver transplantation in patients

[113106]with glycogen storage disease type Ia. J. Inherit. Metab. Dis. 22:

[113107]723–732, 1999.

[113108]19922.

[113109]Fine, R. N.; Wilson, W. A.; Donnell, G. N.: Retinal changes in

[113110]glycogen storage disease type I. Am. J. Dis. Child. 115: 328–331,

[113111]1968.

[113112]19923.

[113113]Fraumeni, J. F.; Rosen, P. J.; Hull, E. W.; Barth, R. F.; Shapiro,

[113114]S. R.; O'Connor, J. F.: Hepatoblastoma in infant sisters. Cancer 24:

[113115]1086–1090, 1969.

[113116]19924.

[113117]Furukawa, N.; Kinugasa, A.; Inoue, F.; Imashuku, S.; Takamatsu,

[113118]T.; Sawada, T.: Type I glycogen storage disease with vaso-

constrictive

[113119]pulmonary hypertension. J. Inherit. Metab. Dis. 13:
102–107, 1990.

[113120]19925.

[113121]Gibson, Q. H.: The reduction of methaemoglobin in red
blood cells

[113122]and studies on the cause of idiopathic methaemoglobi-
naemia. Biochem.

[113123]J. 42: 13–23, 1948.

[113124]19926.

- [113125]Greene, H. L.; Slonim, A. E.; O'Neill, J. A., Jr.; Burr, I. M.
- [113126]: Continuous nocturnal intragastric feeding for management of type
- [113127]1 glycogen-storage disease. New Eng. J. Med. 294: 423-425, 1976.
- [113128]19927.
- [113129]Howell, R. R.: The glycogen storage diseases.In: Stanbury, J.
- [113130]B.; Wyngaarden, J. B.; Fredrickson, D. S.: The Metabolic Basis of
- [113131]Inherited Disease. New York: McGraw-Hill (pub.) (4th ed.): 1978.
- [113132]Pp. 137-159.
- [113133]19928.
- [113134]Howell, R. R.: The interrelationship of glycogen storage disease
- [113135]and gout. Arthritis Rheum. 8: 780-785, 1965.
- [113136]19929.
- [113137]Howell, R. R.; Stevenson, R. E.; Ben-Menachem, Y.; Phyliky, R.
- [113138]L.; Berry, D. H.: Hepatic adenomata with type I glycogen

storage

[113139]disease. J. Nucl. Med. 19: 354–358, 1978.

[113140]19930.

[113141]Ito, E.; Sato, Y.; Kawauchi, K.; Munakata, H.; Kamata, Y.;
Yodono,

[113142]H.; Yokoyama, M.: Type 1a glycogen storage disease with
hepatoblastoma

[113143]in siblings. Cancer 59: 1776–1780, 1987.

[113144]19931.

[113145]Kajihara, S.; Matsushashi, S.; Yamamoto, K.; Kido, K.; Tsuji,
K.;

[113146]Tanae, A.; Fujiyama, S.; Itoh, T.; Tanigawa, K.; Uchida, M.;
Setoguchi,

[113147]Y.; Motomura, M.; Mizuta, T.; Sakai, T.: Exon redefinition
by a point

[113148]mutation within exon 5 of the glucose–6–phosphatase
gene is the major

[113149]cause of glycogen storage disease type 1a in Japan. Am. J.
Hum. Genet. 57:

[113150]549–555, 1995.

[113151]19932.

- [113152] Kikuchi, M.; Hasegawa, K.; Handa, I.; Watabe, M.; Nari-
sawa, K.;
- [113153] Tada, K.: Chronic pancreatitis in a child with glycogen
storage disease
- [113154] type 1. *Europ. J. Pediat.* 150: 852–853, 1991.
- [113155] 19933.
- [113156] Lee, P. J.; Dixon, M. A.; Leonard, J. V.: Uncooked corn-
starch--efficacy
- [113157] in type I glycogenosis. *Arch. Dis. Child.* 74: 546–547,
1996.
- [113158] 19934.
- [113159] Lee, W.-J.; Lee, H.-M.; Chi, C.-S.; Shu, S.-G.; Lin, L.-Y.;
- Lin,
- [113160] W.-H.: Genetic analysis of the glucose-6-phosphatase
mutation of
- [113161] type 1a glycogen storage disease in a Chinese family. *Clin.*
Genet. 50:
- [113162] 206–211, 1996.
- [113163] 19935.
- [113164] Lei, K.-J.; Chen, H.; Pan, C.-J.; Ward, J. M.; Mosinger, B.,
Jr.;

[113165]Lee, E. J.; Westphal, H.; Mansfield, B. C.; Chou, J. Y.: Glucose-6-phosphatase

[113166]dependent substrate transport in the glycogen storage disease type-1a

[113167]mouse. *Nature Genet.* 13: 203-209, 1996.

[113168]19936.

[113169]Lei, K.-J.; Chen, Y.-T.; Chen, H.; Wong, L.-J. C.; Liu, J.-L.;

[113170]McConkie-Rosell, A.; Van Hove, J. L. K.; Ou, H. C.-Y.; Yeh, N. J.;

[113171]Pan, L. Y.; Chou, J. Y.: Genetic basis of glycogen storage disease

[113172]type 1a: prevalent mutations at the glucose-6-phosphatase locus. *Am.*

[113173]*J. Hum. Genet.* 57: 766-771, 1995.

[113174]19937.

[113175]Lei, K.-J.; Pan, C.-J.; Shelly, L. L.; Liu, J.-L.; Chou, J. Y.

[113176]: Identification of mutations in the gene for glucose-6-phosphatase,

[113177]the enzyme deficient in glycogen storage disease type 1A. *J. Clin.*

[113178]*Invest.* 93: 1994-1999, 1994.

[113179]19938.

[113180]Lei, K.-J.; Shelly, L. L.; Lin, B.; Sidbury, J. B.; Chen, Y.-T.;

[113181]Nordlie, R. C.; Chou, J. Y.: Mutations in the glucose-
6-phosphatase

[113182]gene are associated with glycogen storage disease types
1a and 1aSP

[113183]but not 1b and 1c. J. Clin. Invest. 95: 234-240, 1995.

[113184]19939.

[113185]Lei, K.-J.; Shelly, L. L.; Pan, C.-J.; Sidbury, J. B.; Chou, J.

[113186]Y.: Mutations in the glucose-6-phosphatase gene that
cause glycogen

[113187]storage disease type 1a. Science 262: 580-583, 1993.

[113188]19940.

[113189]Limmer, J.; Fleig, W. E.; Leupold, D.; Bittner, R.;
Ditschuneit,

[113190]H.; Beger, H.-G.: Hepatocellular carcinoma in type I glyco-
gen storage

[113191]disease. Hepatology 8: 531-537, 1988.

[113192]19941.

[113193]Malatack, J. J.; Finegold, D. N.; Iwatsuki, S.; Shaw, B. W.,

Jr.;

[113194]Gartner, J. C.; Zitelli, B. J.; Roe, T.; Starzl, T. E.: Liver trans-
plantation

[113195]for type I glycogen storage disease. Lancet I: 1073–1075,
1983.

[113196]19942.

[113197]Aksentijevich, I.; Gruberg, L.; Pras, E.; Balow, J. E., Jr.;
Kovo,

[113198]M.; Gazit, E.; Dean, M.; Pras, M.; Kastner, D. L.: Evidence
for linkage

[113199]of the gene causing familial Mediterranean fever to chro-
mosome 17q

[113200]in non-Ashkenazi Jewish families: second locus or type I
error? Hum.

[113201]Genet. 91: 527–534, 1993.

[113202]19943.

[113203]Aksentijevich, I.; Pras, E.; Gruberg, L.; Shen, Y.; Holman,
K.;

[113204]Helling, S.; Prosen, L.; Sutherland, G. R.; Richards, R. I.;
Ramsburg,

[113205]M.; Dean, M.; Pras, M.; Amos, C. I.; Kastner, D. L.: Refined
mapping

[113206]of the gene causing familial Mediterranean fever, by linkage and homozygosity

[113207]studies. Am. J. Hum. Genet. 53: 451-461, 1993.

[113208]19944.

[113209]Aksentijevich, I.; Pras, E.; Gruberg, L.; Shen, Y.; Richards, R.

[113210]I.; Prosen, L.; Helling, S.; Pras, M.; Kastner, D. L.: Linkage disequilibrium

[113211]in the familial Mediterranean fever candidate region.
(Abstract) Am.

[113212]J. Hum. Genet. 51 (suppl.): A181, 1992.

[113213]19945.

[113214]Aksentijevich, I.; Torosyan, Y.; Samuels, J.; Centola, M.; Pras,

[113215]E.; Chae, J. J.; Oddoux, C.; Wood, G.; Azzaro, M. P.; Palumbo, G.;

[113216]Giustolisi, R.; Pras, M.; Ostrer, H.; Kastner, D. L.: Mutation and

[113217]haplotype studies of familial Mediterranean fever reveal new ancestral

[113218]relationships and evidence for a high carrier frequency with reduced

[113219]penetrance in the Ashkenazi Jewish population. Am. J.
Hum. Genet. 64:

[113220]949–962, 1999.

[113221]19946.

[113222]Armenian, H. K.: Enrollment bias and variation in clinical
manifestations:

[113223]a review of consecutive cases of familial paroxysmal poly-
serositis. J.

[113224]Chronic Dis. 36: 209–212, 1983.

[113225]19947.

[113226]Armenian, H. K.; Khachadurian, A. K.: Familial paroxysmal
polyserositis:

[113227]clinical and laboratory findings in 120 cases. Leban. Med.
J. 26:

[113228]605–614, 1973.

[113229]19948.

[113230]Ayesh, S. K.; Ferne, M.; Flechner, I.; Babior, B. M.; Matzner,

[113231]Y.: Partial characterization of a C5a-inhibitor in peritoneal
fluid. J.

[113232]Immun. 144: 3066–3070, 1990.

[113233]19949.

[113234]Babior, B. M.; Matzner, Y.: The familial Mediterranean fever

[113235]gene--cloned at last. New Eng. J. Med. 337: 1548-1549, 1997.

[113236]19950.

[113237]Barakat, M. H.; El-Khawad, A. O.; Gumaa, K. A.; El-Sobki, N. I.;

[113238]Fenech, F. F.: Metaraminol provocative test: a specific diagnostic

[113239]test for familial Mediterranean fever. Lancet I: 656-657, 1984.

[113240]19951.

[113241]Barakat, M. H.; El-Sobki, N. I.; El-Khawad, A. O.; Gumma, K. A.;

[113242]Fenech, F. F.: Diagnosing familial Mediterranean fever. (Letter) Lancet II:

[113243]41-42, 1984.

[113244]19952.

[113245]Barakat, M. H.; Karnik, A. M.; Majeed, H. W. A.; El-Sobki, N.

[113246]I.; Fenech, F. F.: Familial Mediterranean fever (recurrent hereditary

[113247]polyserositis) in Arabs--a study of 175 patients and re-view of the

[113248]literature. Quart. J. Med. 60: 837-847, 1986.

[113249]19953.

[113250]Barakat, M. H.; Mustafa, H. T.; Shakir, R. A.: Mollaret's meningitis:

[113251]a variant of recurrent hereditary polyserositis, both provoked by

[113252]metaraminol. Arch. Neurol. 45: 926-927, 1988.

[113253]19954.

[113254]Ben-Chetrit, E.; Scherrmann, J.-M.; Levy, M.: Colchicine in breast

[113255]milk of patients with familial Mediterranean fever. Arthritis Rheum. 39:

[113256]1213-1217, 1996.

[113257]19955.

[113258]Benson, M. D.; Skinner, M.; Cohen, A. S.: Amyloid deposition

[113259]in a renal transplant in familial Mediterranean fever. Ann.

Intern.

[113260]Med. 87: 31–34, 1977.

[113261]19956.

[113262]Bernot, A.; da Silva, C.; Petit, J.–L.; Cruaud, C.; Caloustian,

[113263]C.; Castet, V.; Ahmed–Arab, M.; Dross, C.; Dupont, M.;

Cattan, D.;

[113264]Smaoui, N.; Dode, C.; and 10 others: Non–founder mutations in the

[113265]MEFV gene establish this gene as the cause of familial Mediterranean

[113266]fever (FMF). Hum. Molec. Genet. 7: 1317–1325, 1998.

[113267]19957.

[113268]Booth, D. R.; Booth, S. E.; Gillmore, J. D.; Hawkins, P. N.; Pepys,

[113269]M. B.: SAA1 alleles as risk factors in reactive systemic AA amyloidosis. Amyloid 5:

[113270]262–265, 1998.

[113271]19958.

[113272]Brik, R.; Litmanovitz, D.; Berkowitz, D.; Shamir, R.; Rosenthal,

[113273]E.; Shinawi, M.; Gershoni–Baruch, R.: Incidence of familial

Mediterranean

[113274]fever (FMF) mutations among children of Mediterranean
extraction with

[113275]functional abdominal pain. J. Pediat. 138: 759–762, 2001.

[113276]19959.

[113277]Cattan, D.; Dervichian, M.; Courillon, A.; Nurit, Y.:

Metaraminol

[113278]provocation test for familial Mediterranean fever. (Letter)
Lancet I:

[113279]1130–1131, 1984.

[113280]19960.

[113281]Cattan, D.; Notarnicola, C.; Molinari, N.; Touitou, I.: In-
flammatory

[113282]bowel disease in non-Ashkenazi Jews with familial
Mediterranean fever. Lancet 355:

[113283]378–379, 2000.

[113284]19961.

[113285]Chae, J. J.; Centola, M.; Aksentijevich, I.; Dutra, A.; Tran,

[113286]M.; Wood, G.; Nagaraju, K.; Kingma, D. W.; Liu, P. P.; Kast-
ner, D.

[113287]L.: Isolation, genomic organization, and expression analy-

sis of the

[113288]mouse and rat homologs of MEFV, the gene for familial
Mediterranean

[113289]fever. Mammalian Genome 11: 428–435, 2000.

[113290]19962.

[113291]Chen, X.; Fischel–Ghodsian, N.; Cercek, A.; Hamon, M.;
Ogur, G.;

[113292]Lotan, R.; Danon, Y.; Shohat, M.: Assessment of pyrin
gene mutations

[113293]in Turks with familial Mediterranean fever (FMF). Hum.
Mutat. 11:

[113294]456–460, 1998.

[113295]19963.

[113296]Daniels, M.; Shohat, T.; Brenner–Ullman, A.; Shohat, M.:
Familial

[113297]Mediterranean fever: high gene frequency among the
non–Ashkenazic

[113298]and Ashkenazic Jewish populations in Israel. Am. J. Med.
Genet. 55:

[113299]311–314, 1995.

[113300]19964.

- [113301] Dewalle, M.; Domingo, C.; Rozenbaum, M.; Ben-Chetrit, E.; Cattan,
- [113302] D.; Bernot, A.; Dross, C.; Dupont, M.; Notarnicola, C.; Levy, M.;
- [113303] Rosner, I.; Demaille, J.; Touitou, I.: Phenotype-genotype correlation
- [113304] in Jewish patients suffering from familial Mediterranean fever (FMF). Europ.
- [113305] J. Hum. Genet. 6: 95-97, 1998.
- [113306] 19965.
- [113307] Ito, K.; Suzuki, H.; Hirohashi, T.; Kume, K.; Shimizu, T.; Sugiyama,
- [113308] Y.: Molecular cloning of canalicular multispecific organic anion
- [113309] transporter defective in EHBR. Am. J. Physiol. 272: G16-G22, 1997.
- [113310] 19966.
- [113311] van Kuijk, M. A.; Kool, M.; Merks, G. F. M.; Geurts van Kessel,
- [113312] A.; Bindels, R. J. M.; Deen, P. M. T.; van Os, C. H.: Assignment
- [113313] of the canalicular multispecific organic anion transporter

gene (CMOAT)

[113314] to human chromosome 10q24 and mouse chromosome
19D2 by fluorescent

[113315] in situ hybridization. Cytogenet. Cell Genet. 77: 285–287,
1997.

[113316] 19967.

[113317] Wada, M.; Toh, S.; Taniguchi, K.; Nakamura, T.; Uchiumi,
T.; Kohno,

[113318] K.; Yoshida, I.; Kimura, A.; Sakisaka, S.; Adachi, Y.;
Kuвано, M.

[113319]: Mutations in the canalicular multispecific organic anion
transporter

[113320] (cMOAT) gene, a novel ABC transporter, in patients with
hyperbilirubinemia

[113321] II/Dubin–Johnson syndrome. Hum. Molec. Genet. 7:
203–207, 1998.

[113322] 19968.

[113323] Spranger, J. W.: Personal Communication. Kiel, Germany
1974.

[113324] 19969.

[113325] Victoria, T.; Rafi, M. A.; Wenger, D. A.: Cloning of the ca-
nine

[113326]GALC cDNA and identification of the mutation causing globoid cell

[113327]leukodystrophy in West Highland White and Cairn terriers.

Genomics 33:

[113328]457–462, 1996.

[113329]19970.

[113330]Wenger, D. A.; Rafi, M. A.; Luzi, P.: Molecular genetics of Krabbe

[113331]disease (globoid cell leukodystrophy): diagnostic and clinical implications. Hum.

[113332]Mutat. 10: 268–279, 1997.

[113333]19971.

[113334]Zlotogora, J.; Chakraborty, S.; Knowlton, R. G.; Wenger, D. A.

[113335]; Krabbe disease locus mapped to chromosome 14 by genetic linkage. Am.

[113336]J. Hum. Genet. 47: 37–44, 1990.

[113337]19972.

[113338]Zlotogora, J.; Regev, R.; Zeigler, M.; Iancu, T. C.; Bach, G.

[113339]; Krabbe disease: increased incidence in a highly inbred community. Am.

[113340]J. Med. Genet. 21: 765–770, 1985.

[113341]19973.

[113342]Rushton, A. R.; Dawson, G.: Genetic linkage studies of the
human

[113343]glycosphingolipid beta–galactosidases. Biochem. Genet.
15: 1071–1082,

[113344]1977.

[113345]19974.

[113346]Balciunaite, G.; Keller, M. P.; Balciunaite, E.; Piali, L.;
Zuklys,

[113347]S.; Mathieu, Y. D.; Gill, J.; Boyd, R.; Sussman, D. J.; Hollan-
der,

[113348]G. A.: Wnt glycoproteins regulate the expression of
FoxN1, the gene

[113349]defective in nude mice. Nature Immun. 15Oct, 2002. Note:
Advance

[113350]Electronic Publication.

[113351]19975.

[113352]Frank, J.; Pignata, C.; Panteleyev, A. A.; Prowse, D. M.;
Baden,

[113353]H.; Weiner, L.; Gaetaniello, L.; Ahmad, W.; Pozzi, N.; Caer–

halmi-Friedman,

[113354]P. B.; Aita, V. M.; Uyttendaele, H.; Gordon, D.; Ott, J.; Brissette,

[113355]J. L.; Christiano, A. M.: Exposing the human nude phenotype. *Nature* 398:

[113356]473–474, 1999.

[113357]19976.

[113358]Nehls, M.; Pfeifer, D.; Schorpp, M.; Hedrich, H.; Boehm, T.:
New

[113359]member of the winged-helix protein family disrupted in
mouse and rat

[113360]nude mutations. *Nature* 372: 103–107, 1994.

[113361]19977.

[113362]Pignata, C.; Fiore, M.; Guzzetta, V.; Castaldo, A.; Sebastio,
G.;

[113363]Porta, F.; Guarino, A.: Congenital alopecia and nail dystrophy associated

[113364]with severe functional T-cell immunodeficiency in two
sibs. *Am. J.*

[113365]*Med. Genet.* 65: 167–170, 1996.

[113366]19978.

[113367]Schorpp, M.; Hofmann, M.; Dear, T. N.; Boehm, T.: Characterization
of mouse and human nude genes. Immunogenetics 46:
509–515, 1997.
[113369]19979.

[113370]Segre, J. A.; Nemhauser, J. L.; Taylor, B. A.; Nadeau, J. H.;
Lander,
[113371]E. S.: Positional cloning of the nude locus: genetic, physi-
cal, and
[113372]transcription maps of the region and mutations in the
mouse and rat. Genomics 28:
[113373]549–559, 1995.
[113374]19980.

[113375]Delpire, E.; Lu, J.; England, R.; Dull, C.; Thorne, T.: Deaf-
ness
[113376]and imbalance associated with inactivation of the secre-
tory Na–K–2Cl
[113377]co-transporter. Nature Genet. 22: 192–195, 1999.
[113378]19981.

[113379]Evans, R. L.; Park, K.; Turner, R. J.; Watson, G. E.; Nguyen,
H.–V.;

- [113380]Dennett, M. R.; Hand, A. R.; Flagella, M.; Shull, G. E.;
Melvin, J.
- [113381]E.: Severe impairment of salivation in $\text{Na}^+/\text{K}^+/\text{2Cl}^-$ co-
transporter
- [113382](NKCC1)-deficient mice. J. Biol. Chem. 275:
26720-26726, 2000.
- [113383]19982.
- [113384]Payne, J. A.; Xu, J.-C.; Haas, M.; Lytle, C. Y.; Ward, D.; For-
bush,
- [113385]B., III: Primary structure, functional expression, and chro-
mosomal
- [113386]localization of the bumetanide-sensitive Na-K-Cl cotrans-
porter in
- [113387]human colon. J. Biol. Chem. 270: 17977-17985, 1995.
- [113388]19983.
- [113389]Quaggin, S. E.; Payne, J. A.; Forbush, B., III; Igarashi, P.:
- [113390]Localization of the renal Na-K-Cl cotransporter gene
(Slc12a1) on
- [113391]mouse chromosome 2. Mammalian Genome 6: 557-561,
1995.
- [113392]19984.

[113393] Xu, J.-C.; Lytle, C.; Zhu, T. T.; Payne, J. A.; Benz, E., Jr.;
 [113394] Forbush, B., III: Molecular cloning and functional expres-
 sion of
 [113395] the bumetanide-sensitive Na-K-Cl cotransporter. Proc.
 Nat. Acad.
 [113396] Sci. 91: 2201-2205, 1994.
 [113397] 19985.
 [113398] Sedlacek, Z.; Konecki, D. S.; Korn, B.; Klauck, S. M.;
 Poustka,
 [113399] A.: Evolutionary conservation and genomic organization of
 XAP-4,
 [113400] an Xq28 located gene coding for a human rab GDP-
 dissociation inhibitor
 [113401] (GDI). Mammalian Genome 5: 633-639, 1995.
 [113402] 19986.
 [113403] Sedlacek, Z.; Munstermann, E.; Mincheva, A.; Lichter, P.;
 Poustka,
 [113404] A.: The human rab GDI beta gene with long retroposon-
 rich introns
 [113405] maps to 10p15 and its pseudogene to 7p11-p13. Mam-
 malian Genome 9:

[113406]78–80, 1998.

[113407]19987.

[113408]Desseyn, J.–L.; Aubert, J.–P.; van Seuning, I.; Porchet, N.;

[113409]Laine, A.: Genomic organization of the 3–prime region of
the human

[113410]mucin gene MUC5B. J. Biol. Chem. 272: 16873–16883,
1997.

[113411]19988.

[113412]Desseyn, J.–L.; Guyonnet–Duperat, V.; Porchet, N.; Aubert,
J.–P.;

[113413]Laine, A.: Human mucin gene MUC5B, the 10.7–kb large
central exon

[113414]encodes various alternate subdomains resulting in a su-
per-repeat:

[113415]structural evidence for a 11p15.5 gene family. J. Biol.
Chem. 272:

[113416]3168–3178, 1997.

[113417]19989.

[113418]Gipson, I. K.; Moccia, R.; Spurr–Michaud, S.; Argueso, P.;
Gargiulo,

[113419]A. R.; Hill, J. A., III; Offner, G. D.; Keutmann, H. T.: The

amount

[113420]of MUC5B in cervical mucus peaks at midcycle. J. Clin. Endocr. Metab. 86:

[113421]594–600, 2001.

[113422]19990.

[113423]Keates, A. C.; Nunes, D. P.; Afdhal, N. H.; Troxler, R. F.; Offner,

[113424]G. D.: Molecular cloning of a major human gall bladder mucin: complete

[113425]C-terminal sequence and genomic organization of MUC5B. Biochem. J. 324:

[113426]295–303, 1997.

[113427]19991.

[113428]Troxler, R. F.; Offner, G. D.; Zhang, F.; Iontcheva, I.; Oppenheim,

[113429]F. G.: Molecular cloning of a novel high molecular weight mucin (MG1)

[113430]from human sublingual gland. Biochem. Biophys. Res. Commun. 217:

[113431]1112–1119, 1995.

[113432]19992.

- [113433] Birck, C.; Poch, O.; Romier, C.; Ruff, M.; Mengus, G.; Lavigne,
A.-C.; Davidson, I.; Moras, D.: Human TAFII28 and TAFII18
interact
through a histone fold encoded by atypical evolutionary
conserved
motifs also found in the SPT3 family. Cell 94: 239–249,
1998.
19993.
- [113437] Mengus, G.; May, M.; Jacq, X.; Staub, A.; Tora, L.; Chambon, P.;
Davidson, I.: Cloning and characterization of hTAFII18,
hTAFII20
and hTAFII28: three subunits of the human transcription
factor TFIID. EMBO
J. 14: 1520–1531, 1995.
19994.
- [113442] Waisfisz, Q.; Saar, K.; Morgan, N. V.; Altay, C.; Leegwater,
P.
A.; de Winter, J. P.; Komatsu, K.; Evans, G. R.; Wegner,
R.-D.; Reis,
A.; Joenje, H.; Arwert, F.; Mathew, C. G.; Pronk, J. C.; Dig-

weed,

[113446]M.: The Fanconi anemia group E gene, FANCE, maps to chromosome 6p. Am.

[113447]J. Hum. Genet. 64: 1400–1405, 1999.

[113448]19995.

[113449]Wegner, R.-D.; Henrichs, I.; Joenje, H.; Schroeder-Kurth, T.:

[113450]Fanconi anemia complementation group E: clinical and cytogenetic data

[113451]of the first patient. Clin. Genet. 50: 479–482, 1996.

[113452]19996.

[113453]Fink, J. M.; Hirsch, B.; Zheng, C.; Deitz, G.; Hatten, M. E.; Ross,

[113454]M. E.: The CNS neuronal migration gene, astrotactin, is mapped to

[113455]human chromosome band 1q25 by fluorescence in situ hybridization (FISH).

[113456](Abstract) Am. J. Hum. Genet. 57: A133 only, 1995.

[113457]19997.

[113458]Baker, C. H.; Matsuda, S. P. T.; Liu, D. R.; Corey, E. J.: Molecular

[113459]cloning of the human gene encoding lanosterol synthase
from a liver

[113460]cDNA library. Biochem. Biophys. Res. Commun. 213:
154–160, 1995.

[113461]19998.

[113462]Roessler, E.; Mittaz, L.; Du, Y.; Scott, H. S.; Chang, J.;
Rossier,

[113463]C.; Guipponi, M.; Matsuda, S. P. T.; Muenke, M.; An-
tonarakis, S. E.

[113464]: Structure of the human lanosterol synthase gene and its
analysis

[113465]as a candidate for holoprosencephaly (HPE1). Hum. Genet.
105: 489–495,

[113466]1999.

[113467]19999.

[113468]Young, M.; Chen, H.; Lalioti, M. D.; Antonarakis, S. E.: The
human

[113469]lanosterol synthase gene maps to chromosome 21q22.3.
Hum. Genet. 97:

[113470]620–624, 1996.

[113471]20000.

[113472]Green, P.; Lipman, D.; Hillier, L.; Waterston, R.; States, D.;

[113473]Claverie, J.-M.: Ancient conserved regions in new gene sequences

[113474]and the protein databases. Science 259: 1711–1716, 1993.

[113475]20001.

[113476]Schwartz, F.; Eisenman, R.; Knoll, J.; Gessler, M.; Bruns, G.:

[113477]cDNA sequence, genomic organization, and evolutionary conservation

[113478]of a novel gene from the WAGR region. Genomics 29: 526–532, 1995.

[113479]20002.

[113480]Schwartz, F.; Neve, R.; Eisenman, R.; Gessler, M.; Bruns, G.:

[113481]A WAGR region gene between PAX-6 and FSHB expressed in fetal brain. Hum.

[113482]Genet. 94: 658–664, 1994.

[113483]20003.

[113484]Joseph, R. E.; Walker, J.; Norris, F. A.: Assignment of the inositol

[113485]polyphosphate 4-phosphatase type I gene (INPP4A) to hu-

man chromosome

[113486]band 2q11.2 by in situ hybridization. Cytogenet. Cell
Genet. 87:

[113487]276–277, 1999.

[113488]20004.

[113489]Norris, F. A.; Auethavekiat, V.; Majerus, P. W.: The isolation

[113490]and characterization of cDNA encoding human and rat
brain inositol

[113491]polyphosphate 4–phosphatase. J. Biol. Chem. 270:
16128–16133, 1995.

[113492]20005.

[113493]van den Maagdenberg, A. M. J. M.; van den Hurk, H. H.;
Olde Weghuis,

[113494]D.; Wieringa, B.; Geurts van Kessel, A.; Hendriks, W. J. A.
J.: Assignment

[113495]of the human protein tyrosine phosphatase epsilon
(PTPRE) gene to

[113496]chromosome 10q26 by fluorescence in situ hybridization.
Genomics 30:

[113497]128–129, 1995.

[113498]20006.

[113499]Yue, Q.; Jen, J. C.; Nelson, S. F.; Baloh, R. W.: Progressive
[113500]ataxia due to a missense mutation in a calcium-channel
gene. Am.
[113501]J. Hum. Genet. 61: 1078–1087, 1997.
[113502]20007.
[113503]Wojcik, J.; Berg, M. A.; Esposito, N.; Geffner, M. E.; Sakati,
[113504]N.; Reiter, E. O.; Dower, S.; Francke, U.; Postel-Vinay,
M.–C.; Finidori,
[113505]J.: Four contiguous amino acid substitutions, identified in
patients
[113506]with Laron syndrome, differently affect the binding affinity
and intracellular
[113507]trafficking of the growth hormone receptor. J. Clin. En-
docr. Metab. 83:
[113508]4481–4489, 1998.
[113509]20008.
[113510]Azim, A. C.; Knoll, J. H. M.; Marfatia, S. M.; Peel, D. J.;
Bryant,
[113511]P. J.; Chishti, A. H.: DLG1: chromosome location of the
closest human
[113512]homologue of the Drosophila discs large tumor suppres-

sor gene. Genomics 30:

[113513]613–616, 1995.

[113514]20009.

[113515]Burgess, D. L.; Rafael, J. A.; Meisler, M. H.; Chamberlain, J.

[113516]S.: Dlgh1, a mouse homolog of the Drosophila discs-large gene, is

[113517]located on chromosome 16. Mammalian Genome 7:
623–624, 1996.

[113518]20010.

[113519]Hanada, T.; Lin, L.; Chandy, K. G.; Oh, S. S.; Chishti, A. H.:

[113520]Human homologue of the Drosophila discs large tumor suppressor binds

[113521]to p56lck tyrosine kinase and Shaker type Kv1.3 potassium channel

[113522]in T lymphocytes. J. Biol. Chem. 272: 26899–26904,
1997.

[113523]20011.

[113524]Lue, R. A.; Marfatia, S. M.; Branton, D.; Chishti, A. H.:
Cloning

[113525]and characterization of hdlg: the human homologue of the
Drosophila

[113526]discs large tumor suppressor binds to protein 4.1. Proc.
Nat. Acad.

[113527]Sci. 91: 9818–9822, 1994.

[113528]20012.

[113529]Mori, K.; Iwao, K.; Miyoshi, Y.; Nakagawara, A.; Kofu, K.;
Akiyama,

[113530]T.; Arita, N.; Hayakawa, T.; Nakamura, Y.: Identification of
brain-specific

[113531]splicing variants of the hDLG1 gene and altered splicing in
neuroblastoma

[113532]cell lines. J. Hum. Genet. 43: 123–127, 1998.

[113533]20013.

[113534]Peters, L. L.; Ciciotte, S. L.; Lin, L.; Chishti, A. H.: The
mouse

[113535]homolog of the Drosophila discs large tumor suppressor
gene maps to

[113536]chromosome 16. Mammalian Genome 7: 619–620, 1996.

[113537]20014.

[113538]Woods, K. A.; Dastot, F.; Preece, M. A.; Clark, A. J. L.; Pos-
tel-Vinay,

[113539]M.-C.; Chatelain, P. G.; Ranke, M. B.; Rosenfeld, R. G.;

Amselem,

[113540]S.; Savage, M. O.: Phenotype:genotype relationships in growth hormone

[113541]insensitivity syndrome. J. Clin. Endocr. Metab. 82: 3529–3535, 1997.

[113542]20015.

[113543]Bertaux, F.; Sharp, A. H.; Ross, C. A.; Lehrach, H.; Bates, G.

[113544]P.; Wanker, E.: HAP1–huntingtin interactions do not contribute to

[113545]the molecular pathology in Huntington's disease transgenic mice. FEBS

[113546]Lett. 426: 229–232, 1998.

[113547]20016.

[113548]Engelender, S.; Sharp, A. H.; Colomer, V.; Tokito, M. K.; Lanahan,

[113549]A.; Worley, P.; Holzbaur, E. L. F.; Ross, C. A.: Huntingtin-associated

[113550]protein 1 (HAP1) interacts with the p150(Glued) subunit of dynactin. Hum.

[113551]Molec. Genet. 6: 2205–2212, 1997.

[113552]20017.

[113553]Li, S.-H.; Hosseini, S. H.; Gutekunst, C.-A.; Hersch, S. M.;
Ferrante,
[113554]R. J.; Li, X.-J.: A human HAP1 homologue: cloning, expres-
sion, and
[113555]interaction with huntingtin. J. Biol. Chem. 273:
19220-19227, 1998.
[113556]20018.
[113557]Li, X.-J.; Li, S.-H.; Sharp, A. H.; Nucifora, F. C., Jr.;
Schilling,
[113558]G.; Lanahan, A.; Worley, P.; Snyder, S. H.; Ross, C. A.: A
huntingtin-associated
[113559]protein enriched in brain with implications for pathology.
Nature 378:
[113560]398-402, 1995.
[113561]20019.
[113562]Nasir, J.; Duan, K.; Nichol, K.; Engelender, S.; Ashworth, R.;
[113563]Colomer, V.; Thomas, S.; Distèche, C. M.; Hayden, M. R.;
Ross, C.
[113564]A.: Gene structure and map location of the murine ho-
molog of the
[113565]Huntington-associated protein, Hap1. Mammalian
Genome 9: 565-570,

[113566]1998.

[113567]20020.

[113568]Nasir, J.; Lafuente, M.-J.; Duan, K.; Colomer, V.; Engelen-
der,

[113569]S.; Ingersoll, R.; Margolis, R. L.; Ross, C. A.; Hayden, M. R.:
Human

[113570]huntingtin-associated protein (HAP-1) gene: genomic or-
ganisation and

[113571]an intragenic polymorphism. Gene 254: 181-187, 2000.

[113572]20021.

[113573]Coon, S. L.; Mazuruk, K.; Bernard, M.; Roseboom, P. H.;
Klein,

[113574]D. C.; Rodriguez, I. R.: The human serotonin N-
acetyltransferase

[113575](EC 2.3.1.87) gene (AANAT): structure, chromosomal lo-
calization, and

[113576]tissue expression. Genomics 34: 76-84, 1996.

[113577]20022.

[113578]Coon, S. L.; Roseboom, P. H.; Baler, R.; Weller, J. L.; Nam-
boodiri,

[113579]M. A. A.; Koonin, E. V.; Klein, D. C.: Pineal serotonin N-

acetyltransferase:

[113580]expression cloning and molecular analysis. Science 270:

1681–1683,

[113581]1995.

[113582]20023.

[113583]Joy, J. E.; Johnson, G. S.; Lazar, T.; Ralph, M. R.;

Hochstrasser,

[113584]A. C.; Menaker, M.; Merrill, C. R.: Protein differences in tau

mutant

[113585]hamsters: candidate clock proteins. Brain Res. Mol. Brain

Res. 15:

[113586]8–14, 1992.

[113587]20024.

[113588]Ralph, M. R.; Menaker, M.: A mutation of the circadian

system

[113589]in golden hamsters. Science 241: 1225–1227, 1988.

[113590]20025.

[113591]Tosini, G.; Menaker, M.: Circadian rhythms in cultured

mammalian

[113592]retina. Science 272: 419–421, 1996.

[113593]20026.

- [113594]Yoshimura, T.; Nagabukuro, A.; Matsuda, Y.; Suzuki, T.; Kuroiwa, A.; Iigo, M.; Namikawa, T.; Ebihara, S.: Chromosomal mapping of the gene encoding serotonin N-acetyltransferase to rat chromosome 10q32.3 and mouse chromosome 11E2. Cytogenet. Cell Genet. 79: 172–175, 1997.
- [113598]20027.
- [113599]Broccoli, D.; Chong, L.; Oelmann, S.; Fernald, A. A.; Marziliano, N.; van Steensel, B.; Kipling, D.; Le Beau, M. M.; de Lange, T.: Comparison of the human and mouse genes encoding the telomeric protein, TRF1: chromosomal localization, expression and conserved protein domains. Hum. Molec. Genet. 6: 69–76, 1997.
- [113604]20028.
- [113605]Chong, L.; van Steensel, B.; Broccoli, D.; Erdjument-Bromage, H.; Hanish, J.; Tempst, P.; de Lange, T.: A human telomeric

protein. Science 270:

[113607]1663–1667, 1995.

[113608]20029.

[113609]Fairall, L.; Chapman, L.; Moss, H.; de Lange, T.; Rhodes,
D.:

[113610]Structure of the TRFH dimerization domain of the human
telomeric proteins

[113611]TRF1 and TRF2. Molec. Cell 8: 351–361, 2001.

[113612]20030.

[113613]Griffith, J. D.; Comeau, L.; Rosenfield, S.; Stansel, R. M.;
Bianchi,

[113614]A.; Moss, H.; de Lange, T.: Mammalian telomeres end in a
large duplex

[113615]loop. Cell 97: 503–514, 1999.

[113616]20031.

[113617]Kim, S.; Kaminker, P.; Campisi, J.: TIN2, a new regulator of
telomere

[113618]length in human cells. Nature Genet. 23: 405–412, 1999.

[113619]20032.

[113620]Marcand, S.; Gilson, E.; Shore, D.: A protein–counting

mechanism

[113621] for telomere length regulation in yeast. *Science* 275:
986–990, 1997.

[113622] 20033.

[113623] Okabe, J.; Eguchi, A.; Masago, A.; Hayakawa, T.; Nakan-
ishi, M.

[113624] TRF1 is a critical trans-acting factor required for de novo
telomere

[113625] formation in human cells. *Hum. Molec. Genet.* 9:
2639–2650, 2000.

[113626] 20034.

[113627] Young, A. C.; Chavez, M.; Giambernardi, T. A.; Mattern, V.;
McGill,

[113628] J. R.; Harris, J. M.; Sarosdy, M. F.; Patel, P.; Sakaguchi, A. Y.

[113629] Organization and expression of human telomere repeat
binding factor

[113630] genes. *Somat. Cell Molec. Genet.* 23: 275–286, 1997.

[113631] 20035.

[113632] Zakian, V. A.: Telomeres: beginning to understand the
end. *Science* 270:

[113633] 1601–1607, 1995.

[113634]20036.

[113635]Brodsky, G.; Otterson, G. A.; Parry, B. B.; Hart, I.; Patterson,
son,

[113636]D.; Kaye, F. J.: Localization of STCH to human chromosome 21q11.1. Genomics 30:

[113637]627-628, 1995.

[113638]20037.

[113639]Otterson, G. A.; Flynn, G. C.; Kratzke, R. A.; Coxon, A.; Johnston,

[113640]P. G.; Kaye, F. J.: Stch encodes the 'ATPase core' of a mitochondrial
crosomal

[113641]stress70 protein. EMBO J. 13: 1216-1225, 1994.

[113642]20038.

[113643]Reeves, R. H.; Rue, E.; Yu, J.; Kao, F.-T.: Stch maps to
mouse

[113644]chromosome 16, extending the conserved synteny with
human chromosome

[113645]21. Genomics 49: 156-157, 1998.

[113646]20039.

[113647]Nielsen, P. J.; Rochelle, J. M.; Seldin, M. F.: The functional

[113648] genes for protein synthesis initiation factor 4A1 and 4A2
map to
[113649] mouse chromosomes 11 and 16. Mammalian Genome 4:
185–186, 1993.
[113650] 20040.

[113651] Nielsen, P. J.; Trachsel, H.: The mouse protein synthesis
initiation
[113652] factor 4A gene family includes two related functional
genes which
[113653] are differentially expressed. EMBO J. 7: 2097–2105, 1988.
[113654] 20041.

[113655] Sudo, K.; Takahashi, E.; Nakamura, Y.: Isolation and map-
ping of
[113656] the human EIF4A2 gene homologous to the murine pro-
tein synthesis initiation
[113657] factor 4A-II gene Eif4a2. Cytogenet. Cell Genet. 71:
385–388, 1995.
[113658] 20042.

[113659] Gibson, M. A.; Hatzinikolas, G.; Kumaratilake, J. S.; Sand-
berg,
[113660] L. B.; Nicholl, J. K.; Sutherland, G. R.; Cleary, E. G.: Further

[113661] characterization of proteins associated with elastic fiber
microfibrils

[113662] including the molecular cloning of MAGP-2 (MP25). J. Biol.
Chem. 271:

[113663] 1096-1103, 1996.

[113664] 20043.

[113665] Gelb, B. D.; Willner, J. P.; Verloes, A.; Herens, C.; Desnick,
[113666] R. J.: Mutation analysis of pycnodysostosis reveals uni-
parental disomy

[113667] of chromosome 1. (Abstract) Am. J. Hum. Genet. 61
(suppl.): A28

[113668] only, 1997.

[113669] 20044.

[113670] Gladyshev, V. N.; Jeang, K.-T.; Stadtman, T. C.: Selenocys-
teine,

[113671] identified as the penultimate C-terminal residue in human
T-cell thioredoxin

[113672] reductase, corresponds to TGA in the human placental
gene. Proc.

[113673] Nat. Acad. Sci. 93: 6146-6151, 1996.

[113674] 20045.

- [113675]Zhang, K.; Lindsberg, P. J.; Tatlisumak, T.; Kaste, M.; Olsen,
[113676]H. S.; Andersson, L. C.: Stanniocalcin: a molecular guard
of neurons
[113677]during cerebral ischemia. Proc. Nat. Acad. Sci. 97:
3637–3642, 2000.
[113678]20046.
- [113679]Gillett, G. T.; Fox, M. F.; Rowe, P. S. N.; Casimir, C. M.;
Povey,
[113680]S.: Mapping of human non-muscle type cofilin (CFL1) to
chromosome
[113681]11q13 and muscle-type cofilin (CFL2) to chromosome 14.
Ann. Hum.
[113682]Genet. 60: 201–211, 1996.
[113683]20047.
- [113684]Ogawa, K.; Tashima, M.; Yumoto, Y.; Okuda, T.; Sawada,
H.; Okuma,
[113685]M.; Maruyama, Y.: Coding sequence of human placenta
cofilin cDNA. Nuc.
[113686]Acids Res. 18: 7169 only, 1990.
[113687]20048.

[113688]Ono, S.; Minami, N.; Abe, H.; Obinata, T.: Characterization
of
[113689]a novel cofilin isoform that is predominantly expressed in
mammalian
[113690]skeletal muscle. J. Biol. Chem. 269: 15280–15286, 1994.
[113691]20049.

[113692]Georgakopoulos, T.; Thireos, G.: Two distinct yeast tran-
scriptional
[113693]activators require the function of the GCN5 protein to
promote normal
[113694]levels of transcription. EMBO J. 11: 4145–4152, 1992.
[113695]20050.

[113696]Inoue, M.; Isomura, M.; Ikegawa, S.; Fujiwara, T.; Shin, S.;
Moriya,
[113697]H.; Nakamura, Y.: Isolation and characterization of a hu-
man cDNA
[113698]clone (GCN5L1) homologous to GCN5, a yeast transcrip-
tion factor. Cytogenet.
[113699]Cell Genet. 73: 134–136, 1996.
[113700]20051.

[113701]Emahazion, T.; Beskow, A.; Gyllenstein, U.; Brookes, A. J.:

Intron

[113702]based radiation hybrid mapping of 15 complex I genes of the human

[113703]electron transport chain. Cytogenet. Cell Genet. 82: 115–119, 1998.

[113704]20052.

[113705]Byrd, P. J.; Cooper, P. R.; Stankovic, T.; Kullar, H. S.; Watts,

[113706]G. D. J.; Robinson, P. J.; Taylor, M. R.: A gene transcribed from

[113707]the bidirectional ATM promoter coding for a serine rich protein: amino

[113708]acid sequence, structure and expression studies. Hum. Molec. Genet. 5:

[113709]1785–1791, 1996.

[113710]20053.

[113711]Chen, X.; Yang, L.; Udar, N.; Liang, T.; Uhrhammer, N.; Xu, S.;

[113712]Bay, J.–O.; Wang, Z.; Dandakar, S.; Chiplunkar, S.; Klisak, I.; Telatar,

[113713]M.; Yang, H.; Concannon, P.; Gatti, R. A.: CAND3: a ubiquitously

[113714]expressed gene immediately adjacent and in opposite

transcriptional

[113715]orientation to the ATM gene at 11q23.1. Mammalian
Genome 8: 129–133,

[113716]1997.

[113717]20054.

[113718]Imai, T.: Personal Communication. Chiba, Japan
9/12/1996.

[113719]20055.

[113720]Imai, T.; Sugawara, T.; Nishiyama, A.; Shimada, R.; Ohki,
R.; Seki,

[113721]N.; Sagara, M.; Ito, H.; Yamauchi, M.; Hori, T.: The struc-
ture and

[113722]organization of the human NPAT gene. Genomics 42:
388–392, 1997.

[113723]20056.

[113724]Imai, T.; Yamauchi, M.; Seki, N.; Sugawara, T.; Saito, T.;
Matsuda,

[113725]Y.; Ito, H.; Nagase, T.; Nomura, N.; Hori, T.: Identification
and

[113726]characterization of a new gene physically linked to the
ATM gene. Genome

[113727]Res. 6: 439–447, 1996.

[113728]20057.

[113729]Borowsky, B.; Hoffman, B. J.: Analysis of a gene encoding two

[113730]glycine transporter variants reveals alternative promoter usage and

[113731]a novel gene structure. J. Biol. Chem. 273: 29077–29085, 1998.

[113732]20058.

[113733]Borowsky, B.; Mezey, E.; Hoffman, B. J.: Two glycine transporter

[113734]variants with distinct localization in the CNS and peripheral tissues

[113735]are encoded by a common gene. Neuron 10: 851–863, 1993.

[113736]20059.

[113737]Jones, E. M. C.; Fernald, A.; Bell, G. I.; Le Beau, M. M.: Assignment

[113738]of SLC6A9 to human chromosome band 1p33 by in situ hybridization. Cytogenet.

[113739]Cell Genet. 71: 211, 1995.

[113740]20060.

[113741]Kim, K.-M.; Kingsmore, S. F.; Han, H.; Yang-Feng, T. L.;
Godinot,

[113742]N.; Seldin, M. F.; Caron, M. G.; Giros, B.: Cloning of the
human

[113743]glycine transporter type 1: molecular and pharmacological
characterization

[113744]of novel isoform variants and chromosomal localization of
the gene

[113745]in the human and mouse genomes. Molec. Pharmacol. 45:
608-617, 1994.

[113746]20061.

[113747]Arai, Y.; Hosoda, F.; Kobayashi, H.; Arai, K.; Hayashi, Y.;
Kamada,

[113748]N.; Kaneko, Y.; Ohki, M.: The inv(11)(p15q22) chromo-
some translocation

[113749]of de novo and therapy-related myeloid malignancies re-
sults in fusion

[113750]of the nucleoporin gene, NUP98, with the putative RNA
helicase gene,

[113751]DDX10. Blood 89: 3936-3944, 1997.

[113752]20062.

[113753]Dash, A. B.; Williams, I. R.; Kutok, J. L.; Tomasson, M. H.;

Anastasiadou,

[113754]E.; Lindahl, K.; Li, S.; Van Etten, R. A.; Borrow, J.; Housman, D.;

[113755]Druker, B.; Gilliland, D. G.: A murine model of CML blast crisis

[113756]induced by cooperation between BCR/ABL and NUP98/HOXA9. Proc. Nat.

[113757]Acad. Sci. 99: 7622–7627, 2002.

[113758]20063.

[113759]Enninga, J.; Levy, D. E.; Blobel, G.; Fontoura, B. M. A.: Role

[113760]of nucleoporin induction in releasing an mRNA nuclear export block. Science 295:

[113761]1523–1525, 2002.

[113762]20064.

[113763]Fontoura, B. M.; Blobel, G.; Matunis, M. J.: A conserved biogenesis

[113764]pathway for nucleoporins: proteolytic processing of a 186–kilodalton

[113765]precursor generates Nup98 and the novel nucleoporin, Nup96. J. Cell

[113766]Biol. 144: 1097–1112, 1999.

[113767]20065.

[113768]Hodel, A. E.; Hodel, M. R.; Griffis, E. R.; Hennig, K. A.; Ratner,

[113769]G. A.; Xu, S.; Powers, M. A.: The three-dimensional structure of

[113770]the autoproteolytic, nuclear pore-targeting domain of the human nucleoporin

[113771]Nup98. *Molec. Cell* 10: 347–358, 2002.

[113772]20066.

[113773]Jaju, R. J.; Fidler, C.; Haas, O. A.; Strickson, A. J.; Watkins,

[113774]F.; Clark, K.; Cross, N. C. P.; Cheng, J.-F.; Aplan, P. D.; Kearney,

[113775]L.; Boultonwood, J.; Wainscoat, J. S.: A novel gene, NSD1, is fused

[113776]to NUP98 in the t(5;11)(q35;p15.5) in de novo childhood acute myeloid

[113777]leukemia. *Blood* 98: 1264–1267, 2001.

[113778]20067.

[113779]Jaju, R. J.; Haas, C. A.; Neat, M.; et al: A new recurrent translocation,

[113780]t(5;11)(q35p15.5), associated with del(5q) in childhood

acute myeloid

[113781]leukemia. Blood 94: 773–780, 1999.

[113782]20068.

[113783]Mizuno, T.; Kaibuchi, K.; Yamamoto, T.; Kawamura, M.;
Sakoda,

[113784]T.; Fujioka, H.; Matsuura, Y.; Takai, Y.: A stimulatory GDP/
GTP exchange

[113785]protein for smg p21 is active on the post-translationally
processed

[113786]form of c-Ki-ras p21 and rhoA p21. Proc. Nat. Acad. Sci.
88: 6442–6446,

[113787]1991.

[113788]20069.

[113789]Nakamura, T.; Largaespada, D. A.; Lee, M. P.; Johnson, L.
A.;

[113790]Ohyashiki, K.; Toyama, K.; Chen, S. J.; Willman, C. L.;
Chen, I.-M.;

[113791]Feinberg, A. P.; Jenkins, N. A.; Copeland, N. G.; Shaugh-
nessy, J.

[113792]D., Jr.: Fusion of the nucleoporin gene NUP98 to HOXA9
by the chromosome

[113793]translocation t(7;11)(p15;p15) in human myeloid

leukaemia. Nature

[113794]Genet. 12: 154–158, 1996.

[113795]20070.

[113796]Radu, A.; Moore, M. S.; Blobel, G.: The peptide repeat domain

[113797]of nucleoporin Nup98 functions as a docking site in transport across

[113798]the nuclear pore complex. Cell 81: 215–222, 1995.

[113799]20071.

[113800]Rosati, R.; La Starza, R.; Veronese, A.; Aventin, A.; Schwienbacher,

[113801]C.; Vallespi, T.; Negrini, M.; Martelli, M. F.; Mecucci, C.: NUP98

[113802]is fused to the NSD3 gene in acute myeloid leukemia associated with

[113803]t(8;11)(p11.2;p15). Blood 99: 3857–3860, 2002.

[113804]20072.

[113805]von Kobbe, C.; van Deursen, J. M. A.; Rodrigues, J. P.; Sitterlin,

[113806]D.; Bachi, A.; Wu, X.; Wilm, M.; Carmo-Fonseca, M.; Izaurralde, E.

[113807]: Vesicular stomatitis virus matrix protein inhibits host cell gene

[113808]expression by targeting the nucleoporin Nup98. Molec. Cell 6: 1243–1252,

[113809]2000.

[113810]20073.

[113811]Cheng, Y.; Austin, S. C.; Rocca, B.; Koller, B. H.; Coffman, T.

[113812]M.; Grosser, T.; Lawson, J. A.; FitzGerald, G. A.: Role of prostacyclin

[113813]in the cardiovascular response to thromboxane A2. Science 296: 539–541,

[113814]2002.

[113815]20074.

[113816]Fuse, I.; Mito, M.; Hattori, A.; Higuchi, W.; Shibata, A.; Ushikubi,

[113817]F.; Okuma, M.; Yahata, K.: Defective signal transduction induced

[113818]by thromboxane A2 in a patient with a mild bleeding disorder: impaired

[113819]phospholipase C activation despite normal phospholipase A2 activation. Blood 81:

[113820]994–1000, 1993.

[113821]20075.

[113822]Hirata, M.; Hayashi, Y.; Ushikubi, F.; Yokota, Y.; Kageyama, R.;

[113823]Nakanishi, S.; Narumiya, S.: Cloning and expression of cDNA for a

[113824]human thromboxane A2 receptor. Nature 349: 617–620, 1991.

[113825]20076.

[113826]Hirata, T.; Kakizuka, A.; Ushikubi, F.; Fuse, I.; Okuma, M.; Narumiya,

[113827]S.: Arg60-to-leu mutation of the human thromboxane A2 receptor in

[113828]a dominantly inherited bleeding disorder. J. Clin. Invest. 94: 1662–1667,

[113829]1994.

[113830]20077.

[113831]Watanabe, O.; Maruyama, I.; Arimura, K.; Kitajima, I.; Arimura,

[113832]H.; Hanatani, M.; Matsuo, K.; Arisato, T.; Osame, M.: Over-production

[113833] of vascular endothelial growth factor/vascular permeability factor

[113834] is causative in Crow-Fukase (POEMS) syndrome. Muscle Nerve 21: 1390-1397,

[113835] 1998.

[113836] 20078.

[113837] Wei, M.-H.; Popescu, N. C.; Lerman, M. I.; Merrill, M. J.; Zimonjic,

[113838] D. B.: Localization of the human vascular endothelial growth factor

[113839] gene, VEGF, at chromosome 6p12. Hum. Genet. 97: 794-797, 1996.

[113840] 20079.

[113841] Wong, A. K.; Alfert, M.; Castrillon, D. H.; Shen, Q.; Holash,

[113842] J.; Yancopoulos, G. D.; Chin, L.: Excessive tumor-elaborated VEGF

[113843] and its neutralization define a lethal paraneoplastic syndrome. Proc.

[113844] Nat. Acad. Sci. 98: 7481-7486, 2001.

[113845] 20080.

[113846] Wulff, C.; Wilson, H.; Lague, P.; Duncan, W. C.; Arm-

strong, D.

[113847]G.; Fraser, H. M.: Angiogenesis in the human corpus luteum: localization

[113848]and changes in angiopoietins, Tie-2, and vascular endothelial growth

[113849]factor messenger ribonucleic acid. J. Clin. Endocr. Metab. 85: 4302-4309,

[113850]2000.

[113851]20081.

[113852]Ylikorkala, A.; Rossi, D. J.; Korsisaari, N.; Luukko, K.; Alitalo,

[113853]K.; Henkemeyer, M.; Makela, T. P.: Vascular abnormalities and deregulation

[113854]of VEGF in Lkb1-deficient mice. Science 293: 1323-1326, 2001.

[113855]20082.

[113856]Bodner, M.; Fridkin, M.; Gozes, I.: Coding sequences for vasoactive

[113857]intestinal peptide and PHM-27 peptide are located on two adjacent

[113858]exons in the human genome. Proc. Nat. Acad. Sci. 82: 3548-3551,

[113859]1985.

[113860]20083.

[113861]Delgado, M.; Abad, C.; Martinez, C.; Leceta, J.; Gomariz, R.
P.

[113862]; Vasoactive intestinal peptide prevents experimental
arthritis by

[113863]downregulating both autoimmune and inflammatory com-
ponents of the

[113864]disease. Nature Med. 7: 563–568, 2001.

[113865]20084.

[113866]Gotoh, E.; Yamagami, T.; Yamamoto, H.; Okamoto, H.:
Chromosomal

[113867]assignment of human VIP/PHM-27 gene to 6q26–q27 re-
gion by spot blot

[113868]hybridization and in situ hybridization. Biochem. Int. 17:
555–562,

[113869]1988.

[113870]20085.

[113871]Gozes, I.; Avidor, R.; Yahav, Y.; Katznelson, D.; Croce, C.
M.;

[113872]Huebner, K.: The gene encoding vasoactive intestinal pep-

gene is located

[113873]on human chromosome 6p21-6qter. Hum. Genet. 75:
41-44, 1987.

[113874]20086.

[113875]Gozes, I.; Nakai, H.; Byers, M.; Avidor, R.; Weinstein, Y.;
Shani,

[113876]Y.; Shows, T. B.: Sequential expression in the nervous sys-
tem of

[113877]C-MYB and VIP genes, located in human chromosomal re-
gion 6q24. Somat.

[113878]Cell Molec. Genet. 13: 305-313, 1987.

[113879]20087.

[113880]Heinz-Erian, P.; Dey, R. D.; Flux, M.; Said, S. I.: Deficient

[113881]vasoactive intestinal peptide innervation in sweat glands
of cystic

[113882]fibrosis patients. Science 229: 1407-1408, 1985.

[113883]20088.

[113884]Itoh, N.; Obata, K.; Yanaihara, N.; Okamoto, H.: Human
preprovasoactive

[113885]intestinal polypeptide contains a novel PHI-27-like pep-
tide, PHM-27. Nature 304:

[113886]547-549, 1983.

[113887]20089.

[113888]Linder, S.; Barkhem, T.; Norberg, A.; Persson, H.;
Schalling, M.;

[113889]Hokfelt, T.; Magnusson, G.: Structure and expression of
the gene

[113890]encoding the vasoactive intestinal peptide precursor. Proc.
Nat.

[113891]Acad. Sci. 84: 605-609, 1987.

[113892]20090.

[113893]Omary, M. B.; Kagnoff, M. F.: Identification of nuclear re-
ceptors

[113894]for VIP on a human colonic adenocarcinoma cell line. Sci-
ence 238:

[113895]1578-1581, 1987.

[113896]20091.

[113897]Couvineau, A.; Rouyer-Fessard, C.; Darmoul, D.; Maoret,
J.-J.;

[113898]Carrero, I.; Ogier-Denis, E.; Laburthe, M.: Human intesti-
nal VIP

[113899]receptor: cloning and functional expression of two cDNA

encoding proteins

[113900]with different N-terminal domains. Biochem. Biophys. Res. Commun. 200:

[113901]769–776, 1994.

[113902]20092.

[113903]Hashimoto, H.; Nishino, A.; Shintani, N.; Hagihara, N.; Copeland,

[113904]N. G.; Jenkins, N. A.; Yamamoto, K.; Matsuda, T.; Ishihara, T.; Nagata,

[113905]S.; Baba, A.: Genomic organization and chromosomal location of the

[113906]mouse vasoactive intestinal polypeptide 1 (VPAC-1) receptor. Genomics 58:

[113907]90–93, 1999.

[113908]20093.

[113909]Sreedharan, S. P.; Huang, J.-X.; Cheung, M.-C.; Goetzl, E. J.:

[113910]Structure, expression, and chromosomal localization of the type I

[113911]human vasoactive intestinal peptide receptor gene. Proc. Nat. Acad.

[113912]Sci. 92: 2939–2943, 1995.

[113913]20094.

[113914]Sreedharan, S. P.; Patel, D. R.; Huang, J.-X.; Goetzel, E. J.:

[113915]Cloning and functional expression of a human neuroen-
docrine vasoactive

[113916]intestinal peptide receptor. Biochem. Biophys. Res. Com-
mun. 193:

[113917]546-553, 1993.

[113918]20095.

[113919]Sreedharan, S. P.; Robichon, A.; Peterson, K. E.; Goetzel, E.
J.

[113920]: Cloning and expression of the human vasoactive intesti-
nal peptide

[113921]receptor. Proc. Nat. Acad. Sci. 88: 4986-4990, 1991.

[113922]20096.

[113923]Anderson, S. R.; Geertinger, P.; Larsen, H.-W.; Mikkelsen,
M.;

[113924]Parving, A.; Vestermark, S.; Warburg, M.: Aniridia, cataract
and

[113925]gonadoblastoma in a mentally retarded girl with deletion
of chromosome

[113926]11: a clinicopathological case report. Ophthalmologica

176: 171-177,

[113927]1978.

[113928]20097.

[113929]Babaian, R. J.; Skinner, D. G.; Waisman, J.: Wilms' tumor in
the

[113930]adult patient: diagnosis, management, and review of the
world medical

[113931]literature. Cancer 45: 1713-1719, 1980.

[113932]20098.

[113933]Beckwith, J. B.: Nephrogenic rests and the pathogenesis of
Wilms

[113934]tumor: developmental and clinical considerations. Am. J.
Med. Genet. 79:

[113935]268-273, 1998.

[113936]20099.

[113937]Blanchet, P.; Daloze, P.; Lesage, R.; Papas, S.; van Camp-
enhout,

[113938]J.: XY gonadal dysgenesis with gonadoblastoma discov-
ered after kidney

[113939]transplantation. Am. J. Obstet. Gynec. 129: 221-222,
1977.

[113940]20100.

[113941]Bond, J. V.: Bilateral Wilms' tumor: age at diagnosis, associated

[113942]congenital anomalies, and possible pattern of inheritance.
Lancet II:

[113943]482-484, 1975.

[113944]20101.

[113945]Bonetta, L.; Huang, A.; Gregoris, M.; Yeger, H.; Williams, B.
R.

[113946]G.: Characterization of a homozygous deletion mapping to
the Wilms

[113947]tumor region on 11p13. (Abstract) Cytogenet. Cell Genet.
51: 965,

[113948]1989.

[113949]20102.

[113950]Bove, K. E.: Wilms' tumor and related abnormalities in the
fetus

[113951]and newborn. Semin. Perinatol. 23: 310-318, 1999.

[113952]20103.

[113953]Breslow, N. E.; Beckwith, J. R.: Epidemiological features of
Wilms'

[113954]tumor: results of the national Wilms' tumor study. J. Nat.
Cancer

[113955]Inst. 68: 429-436, 1982.

[113956]20104.

[113957]Brown, W. T.; Puranik, S. R.; Altman, D. H.; Hardin, H. C.,
Jr.

[113958]: Wilms' tumor in three successive generations. Surgery
72: 756-761,

[113959]1972.

[113960]20105.

[113961]Cordero, J. F.; Li, F. P.; Holmes, L. B.; Gerald, P. S.: Wilms

[113962]tumor in five cousins. Pediatrics 66: 716-719, 1980.

[113963]20106.

[113964]Dao, D. D.; Schroeder, W. T.; Chao, L.-Y.; Kikuchi, H.;
Strong,

[113965]L. C.; Riccardi, V. M.; Pathak, S.; Nichols, W. W.; Lewis, W.
H.;

[113966]Saunders, G. F.: Genetic mechanisms of tumor-specific
loss of 11p

[113967]DNA sequences in Wilms tumor. Am. J. Hum. Genet. 41:
202-217, 1987.

[113968]20107.

[113969]de Martinville, B.; Francke, U.: HRAS1, insulin and beta-globin

[113970]map outside of 11p11.2-p14.1. (Abstract) Cytogenet. Cell Genet. 37:

[113971]530, 1984.

[113972]20108.

[113973]Denys, P.; Malvaux, P.; van den Berghe, H.; Tanghe, W.; Proesmans,

[113974]W.: Association d'un syndrome anatomo-pathologique de pseudohermaphrodisme

[113975]masculin, d'une tumeur de Wilms, d'une nephropathie parenchymateuse

[113976]et d'un mosaicisme XX/XY. Arch. Franc. Pediat. 24: 729-739, 1967.

[113977]20109.

[113978]Ueno, S.; Maruki, Y.; Nakamura, M.; Tomemori, Y.; Kamae, K.; Tanabe,

[113979]H.; Yamashita, Y.; Matsuda, S.; Kaneko, S.; Sano, A.: The gene encoding

[113980]a newly discovered protein, chorein, is mutated in chorea-acanthocytosis. Nature

[113981]Genet. 28: 121–122, 2001.

[113982]20110.

[113983]Wolf, B.; Feldman, G. L.: The biotin-dependent carboxylase deficiencies. Am.

[113984]J. Hum. Genet. 34: 699–716, 1982.

[113985]20111.

[113986]Hirata, T.; Ushikubi, F.; Kakizuka, A.; Okuma, M.; Narumiya, S.

[113987]: Two thromboxane A(2) receptor isoforms in human platelets: opposite

[113988]coupling to adenylyl cyclase with different sensitivity to arg60-to-leu

[113989]mutation. J. Clin. Invest. 97: 949–956, 1996.

[113990]20112.

[113991]Nusing, R. M.; Hirata, M.; Kakizuka, A.; Eki, T.; Ozawa, K.; Narumiya,

[113992]S.: Characterization and chromosomal mapping of the human thromboxane

[113993]A2 receptor gene. J. Biol. Chem. 268: 25253–25259, 1993.

[113994]20113.

[113995]Schwengel, D. A.; Nouri, N.; Meyers, D. A.; Levitt, R. C.:
Linkage

[113996]mapping of the human thromboxane A2 receptor
(TBXA2R) to chromosome

[113997]19p13.3 using transcribed 3-prime untranslated DNA se-
quence polymorphisms. Genomics 18:

[113998]212-215, 1993.

[113999]20114.

[114000]Thomas, D. W.; Mannon, R. B.; Mannon, P. J.; Latour, A.;
Oliver,

[114001]J. A.; Hoffman, M.; Smithies, O.; Koller, B. H.; Coffman, T.
M.:

[114002]Coagulation defects and altered hemodynamic responses
in mice lacking

[114003]receptors for thromboxane A(2). J. Clin. Invest. 102:
1994-2001,

[114004]1998.

[114005]20115.

[114006]Unoki, M.; Furuta, S.; Onouchi, Y.; Watanabe, O.; Doi, S.;
Fujiwara,

[114007]H.; Miyatake, A.; Fujita, K.; Tamari, M.; Nakamura, Y.: As-
sociation

- [114008] studies of 33 single nucleotide polymorphisms (SNPs) in
29 candidate
- [114009] genes for bronchial asthma: positive association a T924C
polymorphism
- [114010] in the thromboxane A2 receptor gene. Hum. Genet. 106:
440–446, 2000.
- [114011] 20116.
- [114012] Ushikubi, F.; Nakajima, M.; Hirata, M.; Okuma, M.; Fuji-
wara, M.;
- [114013] Narumiya, S.: Purification of the thromboxane
A2/prostaglandin H2
- [114014] receptor from human blood platelets.. J. Biol. Chem. 264:
16496–16501,
- [114015] 1989.
- [114016] 20117.
- [114017] Ushikubi, F.; Okuma, M.; Kanaji, K.; Sugiyama, T.;
- Ogorochi, T.;
- [114018] Narumiya, S.; Uchino, H.: Hemorrhagic thrombocytopathy
with platelet
- [114019] thromboxane A2 abnormality: defective signal transduc-
tion with normal
- [114020] binding activity. Thromb. Haemost. 57: 158–164, 1987.

[114021]20118.

[114022]Ades, E. W.; Zwerner, R. K.; Acton, R. T.; Balch, C. M.: Iso-
lation

[114023]and partial characterisation of the human homologue of
Thy-1. J.

[114024]Exp. Med. 151: 400-406, 1980.

[114025]20119.

[114026]Bonewald, L.; Ades, E. W.; Tung, E.; Marchalonis, J. J.;
Wang,

[114027]A. C.: Biochemical characterization of human Thy-1. J. Im-
munogenet. 11:

[114028]283-296, 1984.

[114029]20120.

[114030]Gatti, R. A.; Lathrop, G. M.; Salser, W.; Silver, J.; Lalouel,

[114031]J. M.; White, R.: Location of Thy-1 with respect to a pri-
mary linkage

[114032]map of chromosome 11q. (Abstract) Cytogenet. Cell
Genet. 46: 618

[114033]only, 1987.

[114034]20121.

[114035]Gatti, R. A.; Shaked, R.; Wei, S.; Koyama, M.; Salser, W.;
Silver,

[114036]J.: DNA polymorphism in the human Thy-1 gene. Hum.
Immun. 22:

[114037]145-150, 1988.

[114038]20122.

[114039]Greenspan, R. J.; O'Brien, M. C.: Genetic evidence for the
role

[114040]of Thy-1 in neurite outgrowth in the mouse. J. Neuro-
genet. 5: 25-36,

[114041]1989.

[114042]20123.

[114043]Grzeschik, K.-H.; Kazazian, H. H.: Report of the committee
on

[114044]the genetic constitution of chromosomes 10, 11, and 12.
Cytogenet.

[114045]Cell Genet. 40: 179-205, 1985.

[114046]20124.

[114047]Letarte-Muirhead, M.; Barclay, A. N.; Williams, A. F.: Purifi-
cation

[114048]of the Thy-1 molecule, a major cell surface glycoprotein

of rat thymocytes.

[114049]Biochem. J. 151: 685–697, 1975.

[114050]20125.

[114051]Mansour, M. H.; Negm, H. I.; Cooper, E. L.: Thy-1 evolution. Dev.

[114052]Comp. Immun. 11: 3–15, 1987.

[114053]20126.

[114054]McKenzie, J. L.; Fabre, J. W.: Human Thy-1: unusual localization

[114055]and possible functional significance in lymphoid tissues. J. Immun. 126:

[114056]843–850, 1981.

[114057]20127.

[114058]Morris, R.: Thy-1 in developing nervous tissue. Dev. Neurosci. 7:

[114059]133–160, 1985.

[114060]20128.

[114061]Raff, M. C.: Surface antigenic markers for distinguishing T and

[114062]B lymphocytes in mice. Transplant. Rev. 6: 52–80, 1971.

[114063]20129.

[114064]Rettig, W. J.; Dracopoli, N. C.; Chesa, P. G.; Spengler, B. A.;

[114065]Beresford, H. R.; Davies, P.; Biedler, J. L.; Old, L. J.: Role of

[114066]human chromosome 11 in determining surface antigenic
phenotype of

[114067]normal and malignant cells: somatic cell genetic analysis
of eight

[114068]antigens, including putative human Thy-1. J. Exp. Med.
162: 1603-1619,

[114069]1985.

[114070]20130.

[114071]Rettig, W. J.; Dracopoli, N. C.; Silver, J.; Old, L. J.: Human

[114072]THY-1: regional mapping on chromosome 11 and com-
parison with other

[114073]chromosome 11-encoded cell surface glycoproteins.
(Abstract) Cytogenet.

[114074]Cell Genet. 40: 731 only, 1985.

[114075]20131.

[114076]Rettig, W. J.; Dracopoli, N. C.; Spengler, B. A.; Biedler, J.

[114077]L.; Old, L. J.: Somatic cell genetic analysis of human cell

surface

[114078]antigens, including putative human Thy-1: eight distinct antigenic

[114079]systems controlled by chromosome 11. (Abstract) Cytogenet. Cell

[114080]Genet. 40: 732 only, 1985.

[114081]20132.

[114082]Seki, T.; Spurr, N.; Obata, F.; Goyert, S.; Goodfellow, P.; Silver,

[114083]J.: The human Thy-1 gene: structure and chromosomal location. Proc.

[114084]Nat. Acad. Sci. 82: 6657-6661, 1985.

[114085]20133.

[114086]Tse, A. G. D.; Barclay, A. N.; Watts, A.; Williams, A. F.: A

[114087]glycophospholipid tail at the carboxyl terminus of the Thy-1 glycoprotein

[114088]of neurons and thymocytes. Science 230: 1003-1008, 1985.

[114089]20134.

[114090]Chen, Z.; Friedrich, G. A.; Soriano, P.: Transcriptional enhancer

[114091]factor 1 disruption by a retroviral gene trap leads to heart defects

[114092]and embryonic lethality in mice. Genes Dev. 8: 2293–2301, 1994.

[114093]20135.

[114094]Aso, T.; Tsai, P.; Kawaguchi, T.; Menninger, J. C.; Kitajima, S.;

[114095]Yasukochi, Y.; Ward, D. C.; Weissman, S. M.: Assignment of the human

[114096]GTF2F1 gene to chromosome 19p13.3. Genomics 16: 252–253, 1993.

[114097]20136.

[114098]Aso, T.; Vasavada, H. A.; Kawaguchi, T.; Germino, F. J.; Ganguly,

[114099]S.; Kitajima, S.; Weissman, S. M.; Yasukochi, Y.: Characterization

[114100]of cDNA for the large subunit of the transcription initiation factor

[114101]TFIIF. Nature 355: 461–464, 1992.

[114102]20137.

[114103]Finkelstein, A.; Kostrub, C. F.; Li, J.; Chavez, D. P.; Wang, B.

- [114104]Q.; Fang, S. M.; Greenblatt, J.; Burton, Z. F.: A cDNA encoding RAP74,
[114105]a general initiation factor for transcription by RNA polymerase II. Nature 355:
[114106]464–467, 1992.
[114107]20138.
- [114108]Joliot, V.; Demma, M.; Prywes, R.: Interaction with RAP74 subunit
[114109]of TFIIF is required for transcriptional activation by serum response
[114110]factor. Nature 373: 632–635, 1995.
[114111]20139.
- [114112]Sopta, M.; Burton, Z. F.; Greenblatt, J.: Structure and associated
[114113]DNA–helicase activity of a general transcription initiation factor
[114114]that binds to RNA polymerase II. Nature 341: 410–414, 1989.
[114115]20140.
- [114116]Field, S. J.; Tsai, F.–Y.; Kuo, F.; Zubiaga, A. M.; Kaelin, W.
[114117]G., Jr.; Livingston, D. M.; Orkin, S. H.; Greenberg, M. E.:

E2F-1

[114118]functions in mice to promote apoptosis and suppress proliferation. Cell 85:

[114119]549–561, 1996.

[114120]20141.

[114121]Helin, K.; Lees, J. A.; Vidal, M.; Dyson, N.; Harlow, E.; Fataey,

[114122]A.: A cDNA encoding a pRB-binding protein with properties of the

[114123]transcription factor E2F. Cell 70: 337–350, 1992.

[114124]20142.

[114125]Irwin, M.; Marin, M. C.; Phillips, A. C.; Seelan, R. S.; Smith,

[114126]D. I.; Liu, W.; Flores, E. R.; Tsai, K. Y.; Jacks, T.; Vousden, K.

[114127]H.; Kaelin, W. G., Jr.: Role for the p53 homologue p73 in E2F-1-induced

[114128]apoptosis. Nature 407: 645–648, 2000.

[114129]20143.

[114130]Jacks, T.; Fazeli, A.; Schmitt, E. M.; Bronson, R. T.; Goodell,

[114131]M. A.; Weinberg, R. A.: Effects of an Rb mutation in the

mouse. Nature 359:

[114132]295–300, 1992.

[114133]20144.

[114134]Lees, J. A.; Saito, M.; Valentine, M.; Look, T.; Harlow, E.;
Dyson,

[114135]N.; Helin, K.: The retinoblastoma protein binds to a family
of E2F

[114136]transcription factors. Molec. Cell. Biol. 13: 7813–7825,
1993.

[114137]20145.

[114138]Leone, G.; Sears, R.; Huang, E.; Rempel, R.; Nuckolls, F.;
Park,

[114139]C.–H.; Giangrande, P.; Wu, L.; Saavedra, H. I.; Field, S. J.;
Thompson,

[114140]M. A.; Yang, H.; Fujiwara, Y.; Greenberg, M. E.; Orkin, S.;
Smith,

[114141]C.; Nevins, J. R.: Myc requires distinct E2F activities to in-
duce

[114142]S phase and apoptosis. Molec. Cell 8: 105–113, 2001.

[114143]20146.

[114144]Lissy, N. A.; Davis, P. K.; Irwin, M.; Kaelin, W. G.; Dowdy, S.

[114145]F.: A common E2F-1 and p73 pathway mediates cell death
induced by

[114146]TCR activation. *Nature* 407: 642-645, 2000.

[114147]20147.

[114148]Neuman, E.; Sellers, W. R.; McNeil, J. A.; Lawrence, J. B.;
Kaelin,

[114149]W. G., Jr.: Structure and partial genomic sequence of the
human E2F1

[114150]gene. *Gene* 173: 163-169, 1996.

[114151]20148.

[114152]Nevins, J. R.: The Rb/E2F pathway and cancer. *Hum.*
Molec. Genet. 10:

[114153]699-703, 2001.

[114154]20149.

[114155]Nevins, J. R.: E2F: a link between the Rb tumor suppressor
protein

[114156]and viral oncoproteins. *Science* 258: 424-429, 1992.

[114157]20150.

[114158]Ohtani, K.; DeGregori, J.; Nevins, J. R.: Regulation of the

cyclin

[114159]E gene by transcription factor E2F1. Proc. Nat. Acad. Sci.
92: 12146–12150,

[114160]1995.

[114161]20151.

[114162]Arden, K. C.; Boutin, J.–M.; Djiane, J.; Kelly, P. A.; Cavenee,

[114163]W. K.: The receptors for prolactin and growth hormone are
localized

[114164]in the same region of human chromosome 5. Cytogenet.
Cell Genet. 53:

[114165]161–165, 1990.

[114166]20152.

[114167]Arden, K. C.; Cavenee, W. K.; Boutin, J.–M.; Kelly, P. A.:
The

[114168]genes encoding the receptors for prolactin and growth
hormone map

[114169]to human chromosome 5. (Abstract) Am. J. Hum. Genet.
45 (suppl.):

[114170]A129 only, 1989.

[114171]20153.

[114172]Kelly, A.; Powis, S. H.; Glynne, R.; Radley, E.; Beck, S.;

Trowsdale,

[114173]J.: Second proteasome-related gene in the human MHC class II region. Nature 353:

[114174]667-668, 1991.

[114175]20154.

[114176]Martinez, C. K.; Monaco, J. J.: Homology of proteasome subunits

[114177]to a major histocompatibility complex-linked LMP gene. Nature 353:

[114178]664-667, 1991.

[114179]20155.

[114180]Petes, T. D.: Meiotic recombination hot spots and cold spots. Nature

[114181]Rev. Genet. 2: 360-369, 2001.

[114182]20156.

[114183]Van Kaer, L.; Ashton-Rickardt, P. G.; Eichelberger, M.; Gaczynska,

[114184]M.; Nagashima, K.; Rock, K. L.; Goldberg, A. L.; Doherty, P. C.; Tonegawa,

[114185]S.: Altered peptidase and viral-specific T cell response in LMP2

[114186]mutant mice. Immunity 1: 533–541, 1994.

[114187]20157.

[114188]Zhou, P.; Zanelli, E.; Smart, M.; David, C.: Genomic organization

[114189]and tissue expression of mouse proteasome gene Lmp-2. Genomics 16:

[114190]664–668, 1993.

[114191]20158.

[114192]Deng, G. Y.; Muir, A.; Maclaren, N. K.; She, J.-X.: Association

[114193]of LMP2 and LMP7 genes within the major histocompatibility complex

[114194]with insulin-dependent diabetes mellitus: population and family studies. Am.

[114195]J. Hum. Genet. 56: 528–534, 1995.

[114196]20159.

[114197]Bouhassira, E. E.; Schwartz, R. S.; Yawata, Y.; Ata, K.; Kan-

[114198]A.; Qui, J. J.-H.; Nagel, R. L.; Rybicki, A. C.: An alanine-to-threonine

[114199]substitution in protein 4.2 cDNA is associated with a

Japanese form

[114200]of hereditary hemolytic anemia (protein 4.2-Nippon).

Blood 79: 1846-1854,

[114201]1992.

[114202]20160.

[114203]Doolittle, R. F.; Hunkapiller, M. W.; Hood, L. E.; Devare, S.

[114204]G.; Robbins, K. C.; Aaronson, S. A.; Antoniades, H. N.:

Simian sarcoma

[114205]virus onc gene, v-sis, is derived from the gene (or genes)
encoding

[114206]a platelet-derived growth factor. Science 221: 275-277,
1983.

[114207]20161.

[114208]Frolova, L. Y.; Sudomoina, M. A.; Grigorieva, A. Y.; Zi-
novieva,

[114209]O. L.; Kisselev, L. L.: Cloning and nucleotide sequence of
the structural

[114210]gene encoding for human tryptophanyl-tRNA synthetase.
Gene 109:

[114211]291-296, 1991.

[114212]20162.

[114213]Vassart, G.: Personal Communication. Brussels, Belgium
1/15/1992.

[114214]20163.

[114215]Wenger, G. D.: Personal Communication. Columbus, Ohio
8/3/1993.

[114216]20164.

[114217]Abbes, A. P.; Bruggeman, B.; van den Akker, E. L. T.; de
Groot,

[114218]M. R.; Franken, A. A. M.; Drexhage, V. R.; Engel, H.: Identi-
fication

[114219]of two distinct mutations at the same nucleotide position,
concomitantly

[114220]with a novel polymorphism in the vasopressin-neuro-
physin II gene (AVP-NP

[114221]II) in two Dutch families with familial neurohypophyseal
diabetes

[114222]insipidus. Clin. Chem. 46: 1699-1702, 2000.

[114223]20165.

[114224]Bahnsen, U.; Oosting, P.; Swaab, D. F.; Nahke, P.; Richter,
D.;

[114225]Schmale, H.: A missense mutation in the vasopressin-
neurophysin precursor

[114226]gene cosegregates with human autosomal dominant neurohypophyseal diabetes

[114227]insipidus. EMBO J. 11: 19–23, 1992.

[114228]20166.

[114229]Calvo, B.; Bilbao, J. R.; Rodriguez, A.; Rodriguez–Arnao, M. D.;

[114230]Castano, L.: Molecular analysis in familial neurohypophyseal diabetes

[114231]insipidus: early diagnosis of an asymptomatic carrier. J. Clin. Endocr.

[114232]Metab. 84: 3351–3354, 1999.

[114233]20167.

[114234]Calvo, B.; Bilbao, J. R.; Urrutia, I.; Eizaguirre, J.; Gaztam-bide,

[114235]S.; Castano, L.: Identification of a novel nonsense mutation and

[114236]a missense substitution in the vasopressin–neurophysin II gene in

[114237]two Spanish kindreds with familial neurohypophyseal diabetes insipidus. J.

[114238]Clin. Endocr. Metab. 83: 995–997, 1998.

[114239]20168.

[114240]Doherty-Fuller, E.; Copeland, K. C.: Sweat tests in patients
with

[114241]diabetes insipidus. Clin. Pediat. 27: 330-332, 1988.

[114242]20169.

[114243]Evans, D. A. P.; van der Kleij, A. A. M.; Sonnemans, M. A.
F.;

[114244]Burbach, J. P. H.; van Leeuwen, F. W.: Frameshift muta-
tions at two

[114245]hotspots in vasopressin transcripts in post-mitotic neu-
rons. Proc.

[114246]Nat. Acad. Sci. 91: 6059-6063, 1994.

[114247]20170.

[114248]Gagliardi, P. C.; Bernasconi, S.; Repaske, D. R.: Autosomal
dominant

[114249]neurohypophyseal diabetes insipidus associated with a
missense mutation

[114250]encoding gly(23) to val in neurophysin II. J. Clin. Endocr.
Metab. 82:

[114251]3643-3646, 1997.

[114252]20171.

[114253] Ganz, M. B.; Boyarsky, G.; Sterzel, R. B.; Boron, W. F.: Arginine

[114254] Vasopressin enhances pH(i) regulation in the presence of HCO₃⁻ by

[114255] stimulating three acid-base transport systems. Nature 337: 648-651,

[114256] 1989.

[114257] 20172.

[114258] Grant, F. D.; Ahmadi, A.; Hosley, C. M.; Majzoub, J. A.: Two

[114259] novel mutations of the vasopressin gene associated with familial diabetes

[114260] insipidus and identification of an asymptomatic carrier infant. J.

[114261] Clin. Endocr. Metab. 83: 3958-3964, 1998.

[114262] 20173.

[114263] Heppner, C.; Kotzka, J.; Bullmann, C.; Krone, W.; Muller-Wieland,

[114264] D.: Identification of mutations of the arginine vasopressin-neurophysin

[114265] II gene in two kindreds with familial central diabetes insipidus. J.

[114266]Clin. Endocr. Metab. 83: 693–696, 1998.

[114267]20174.

[114268]Ito, M.; Jameson, J. L.; Ito, M.: Molecular basis of autosomal

[114269]dominant neurohypophyseal diabetes insipidus: cellular toxicity caused

[114270]by the accumulation of mutant vasopressin precursors within the endoplasmic

[114271]reticulum. J. Clin. Invest. 99: 1897–1905, 1997.

[114272]20175.

[114273]Ito, M.; Mori, Y.; Oiso, Y.; Saito, H.: A single base substitution

[114274]in the coding region for neurophysin II associated with familial central

[114275]diabetes insipidus. J. Clin. Invest. 87: 725–728, 1991.

[114276]20176.

[114277]Ito, M.; Oiso, Y.; Murase, T.; Kondo, K.; Saito, H.; Chinzei,

[114278]T.; Racchi, M.; Lively, M. O.: Possible involvement of inefficient

[114279]cleavage of preprovasopressin by signal peptidase as a cause for familial

- [114280]central diabetes insipidus. J. Clin. Invest. 91: 2565–2571, 1993.
- [114281]20177.
- [114282]Korbonits, M.; Kaltsas, G.; Perry, L. A.; Putignano, P.; Grossman, A. B.; Besser, G. M.; Trainer, P. J.: The growth hormone secretagogue hexarelin stimulates the hypothalamo–pituitary–adrenal axis via arginine vasopressin. J. Clin. Endocr. Metab. 84: 2489–2495, 1999.
- [114283]20178.
- [114284]Krishnamani, M. R. S.; Phillips, J. A., III; Copeland, K. C.: Detection of a novel arginine vasopressin defect by dideoxy fingerprinting. J. Clin. Endocr. Metab. 77: 596–598, 1993.
- [114285]20179.
- [114286]Land, H.; Schutz, G.; Schmale, H.; Richter, D.: Nucleotide sequence of cloned cDNA encoding bovine arginine vasopressin–neurophysin II precursor. Nature 295: 299–303, 1982.

[114294]20180.

[114295]Mahoney, C. P.; Weinberger, E.; Bryant, C.; Ito, M.; Jameson,

[114296]J. L.; Ito, M.: Effects of aging on vasopressin production in a kindred

[114297]with autosomal dominant neurohypophyseal diabetes insipidus due to

[114298]the delta-E47 neurophysin mutation. J. Clin. Endocr.

Metab. 87:

[114299]870–876, 2002.

[114300]20181.

[114301]McLeod, J. F.; Kovacs, L.; Gaskill, M. B.; Rittig, S.; Bradley,

[114302]G. S.; Robertson, G. L.: Familial neurohypophyseal diabetes insipidus

[114303]associated with a signal peptide mutation. J. Clin. Endocr.

Metab. 77:

[114304]599A–599G, 1993.

[114305]20182.

[114306]Gardner, T. L.; Elston, D. M.; Wotowic, P. J.: A familial dermatofibrosarcoma

[114307]protuberans. J. Am. Acad. Derm. 39: 504–505, 1998.

[114308]20183.

[114309]Groffen, J.; Heisterkamp, N.; Stephenson, J. R.; Geurts van Kessel,

[114310]A.; de Klein, A.; Grosveld, G.; Bootsma, D.: c-sis is translocated

[114311]from chromosome 22 to chromosome 9 in chronic myelocytic leukemia. J.

[114312]Exp. Med. 158: 9-15, 1983.

[114313]20184.

[114314]Hermansson, M.; Nister, M.; Betsholtz, C.; Heldin, C.-H.; Westermark,

[114315]B.; Funa, K.: Endothelial cell hyperplasia in human glioblastoma:

[114316]coexpression of mRNA for platelet-derived growth factor (PDGF) B chain

[114317]and PDGF receptor suggests autocrine growth stimulation. Proc. Nat.

[114318]Acad. Sci. 85: 7748-7752, 1988.

[114319]20185.

[114320]Josephs, S. F.; Dalla-Favera, R.; Gelmann, E. P.; Gallo, R. C.;

[114321]Wong-Staal, F.: 5-prime viral and human cellular sequences corresponding
[114322]to the transforming gene of simian sarcoma virus. Science
219: 503-505,
[114323]1983.
[114324]20186.
[114325]Josephs, S. F.; Guo, C.; Ratner, L.; Wong-Staal, F.: Human
proto-oncogene
[114326]nucleotide sequences corresponding to the transforming
region of simian
[114327]sarcoma virus. Science 223: 487-491, 1984.
[114328]20187.
[114329]Josephs, S. F.; Ratner, L.; Clarke, M. F.; Westin, E. H.; Reitz,
[114330]M. S.; Wong-Staal, F.: Transforming potential of human c-
sis nucleotide
[114331]sequences encoding platelet-derived growth factor. Science 225:
[114332]636-639, 1984.
[114333]20188.
[114334]Kelly, J. D.; Raines, E. W.; Ross, R.; Murray, M. J.: The B
chain

[114335]of PDGF alone is sufficient for mitogenesis. EMBO J. 4:
3399–3405,

[114336]1985.

[114337]20189.

[114338]Kiuru-Kuhlefelt, S.; El-Rifai, W.; Fanburg-Smith, J.; Kere,
J.;

[114339]Miettinen, M.; Knuutila, S.: Concomitant DNA copy number
amplification

[114340]at 17q and 22q in dermatofibrosarcoma protuberans. Cy-
togenet. Cell

[114341]Genet. 92: 192–195, 2001.

[114342]20190.

[114343]Kozak, C. A.; Sears, J. F.; Hoggan, M. D.: Genetic mapping
of

[114344]the mouse proto-oncogene c-sis to chromosome 15. Sci-
ence 221: 867–869,

[114345]1983.

[114346]20191.

[114347]Lindahl, P.; Johansson, B. R.; Leveen, P.; Betsholtz, C.: Per-
icyte

[114348]loss and microaneurysm formation in PDGF-B-deficient

mice. Science 277:

[114349]242–245, 1997.

[114350]20192.

[114351]Owen, A. J.; Pantazis, P.; Antoniades, H. N.: Simian sarcoma

[114352]virus-transformed cells secrete a mitogen identical to platelet-derived

[114353]growth factor. Science 225: 54–56, 1984.

[114354]20193.

[114355]Rao, C. D.; Igarashi, H.; Chiu, I.-M.; Robbins, K. C.; Aaronson,

[114356]S. A.: Structure and sequence of the human c-sis/platelet-derived

[114357]growth factor 2 (SIS/PDGF2) transcriptional unit. Proc. Nat. Acad.

[114358]Sci. 83: 2392–2396, 1986.

[114359]20194.

[114360]Robbins, K. C.; Antoniades, H. N.; Devare, S. G.; Hunkapiller,

[114361]M. W.; Aaronson, S. A.: Structural and immunological similarities

[114362]between simian sarcoma virus gene product(s) and human platelet-derived

[114363]growth factor. *Nature* 305: 605–608, 1983.

[114364]20195.

[114365]Robbins, K. C.; Devare, S. G.; Reddy, E. P.; Aaronson, S. A.:

[114366]In vivo identification of the transforming gene product of simian

[114367]sarcoma virus. *Science* 218: 1131–1133, 1982.

[114368]20196.

[114369]Simon, M.–P.; Navarro, M.; Roux, D.; Pouyssegur, J.: Structural

[114370]and functional analysis of a chimeric protein COL1A1–PDGFB generated

[114371]by the translocation t(17;22)(q22;q13.1) in dermatofibrosarcoma protuberans

[114372](DP). *Oncogene* 20: 2965–2975, 2001.

[114373]20197.

[114374]Smidt, M.; Kirsch, I.; Ratner, L.: Deletion of Alu sequences

[114375]in the fifth c-sis intron in individuals with meningiomas. *J. Clin.*

[114376]Invest. 86: 1151-1157, 1990.

[114377]20198.

[114378]Turc-Carel, C.; Philip, I.; Berger, M. P.; Philip, T.; Lenoir,

[114379]G. M.: Chromosomal translocations in Ewing's sarcoma.

(Letter) New

[114380]Eng. J. Med. 309: 497-498, 1983.

[114381]20199.

[114382]Shimizu, A.; O'Brien, K. P.; Sjoblom, T.; Pietras, K.; Buch-
dunger,

[114383]E.; Collins, V. P.; Heldin, C.-H.; Dumanski, J. P.; Ostman,
A.: The

[114384]dermatofibrosarcoma protuberans-associated collagen
type I-alpha-1/platelet-derived

[114385]growth factor (PDGF) beta-chain fusion gene generates a
transforming

[114386]protein that is processed to functional PDGF-BB. Cancer
Res. 59:

[114387]3719-3723, 1999.

[114388]20200.

[114389]Simon, M.-P.; Pedeutour, F.; Sirvent, N.; Grosgeorge, J.;
Minoletti,

- [114390]F.; Coindre, J.-M.; Terrier-Lacombe, M.-J.; Mandahl, N.; Craver, R.
- [114391]D.; Blin, N.; Sozzi, G.; Turc-Carel, C.; O'Brien, K. P.; Kedra, D.;
- [114392]Fransson, I.; Guilbaud, C.; Dumanski, J. P.: Deregulation of the
- [114393]platelet-derived growth factor B-chain gene via fusion with collagen
- [114394]gene COL1A1 in dermatofibrosarcoma protuberans and giant-cell fibroblastoma. *Nature*
- [114395]*Genet.* 15: 95-98, 1997.
- [114396]20201.
- [114397]Waterfield, M. D.; Scrace, G. T.; Whittle, N.; Stroobant, P.;
- [114398]Johnsson, A.; Wasteson, A.; Westermarck, B.; Heldin, C.-H.; Huang,
- [114399]J. S.; Deuel, T. F.: Platelet-derived growth factor is structurally
- [114400]related to the putative transforming protein p28(sis) of simian sarcoma
- [114401]virus. *Nature* 304: 35-39, 1983.
- [114402]20202.

[114403]Graphodatsky, A.; Frolova, L.; Biltueva, L.; Eremina, V.;
Lushnikova,
[114404]T.; Sudomoina, M.; Zinovieva, O.; Kisselev, L.: Localization
of the
[114405]tryptophanyl tRNA synthetase gene (WARS) on human and
bovine chromosomes
[114406]by in situ hybridization. Mammalian Genome 4: 183–184,
1993.
[114407]20203.
[114408]Jensen, L. L.; Nielsen, M. M.; Justesen, J.; Hansen, L. L.:
Assignment
[114409]of human NADH dehydrogenase (ubiquinone) 1 beta sub-
complex 3 (NDUFB3)
[114410]and of its four pseudogenes to human chromosomes
2q31.3, 1p13.3-p13.1,
[114411]9q32-q34.1, 14q22.3-q23.1 and 14q32.2 by radiation
hybrid mapping. Cytogenet.
[114412]Cell Genet. 93: 147–150, 2001.
[114413]20204.
[114414]Otani, A.; Slike, B. M.; Dorrell, M. I.; Hood, J.; Kinder, K.;
[114415]Ewalt, K. L.; Cheresch, D.; Schimmel, P.; Friedlander, M.: A
fragment

- [114416] of human TrpRS as a potent antagonist of ocular angiogenesis. Proc.
- [114417] Nat. Acad. Sci. 99: 178–183, 2002.
- [114418] 20205.
- [114419] Shimizu, N.; Kucherlapati, R. S.; Ruddle, F. H.: Assignment of
- [114420] a human gene for tryptophanyl-tRNA synthetase to chromosome 14 using
- [114421] human-mouse somatic cell hybrids. Somat. Cell Genet. 2: 345–357,
- [114422] 1976.
- [114423] 20206.
- [114424] Tolstrup, A. B.; Bejder, A.; Fleckner, J.; Justesen, J.: Transcriptional
- [114425] regulation of the interferon-gamma-inducible tryptophanyl-tRNA synthetase
- [114426] includes alternative splicing. J. Biol. Chem. 270: 397–403, 1995.
- [114427] 20207.
- [114428] Turpaev, K. T.; Zakhariev, V. M.; Sokolova, I. V.; Narovlyansky,

- [114429]A. N.; Amchenkova, A. M.; Justesen, J.; Frolova, L. Y.: Alternative
[114430]processing of the tryptophanyl-tRNA synthetase mRNA
from interferon-treated
[114431]human cells. *Europ. J. Biochem.* 240: 732–737, 1996.
[114432]20208.
- [114433]Wakasugi, K.; Slike, B. M.; Hood, J.; Otani, A.; Ewalt, K. L.;
[114434]Friedlander, M.; Cheresch, D. A.; Schimmel, P.: A human
aminoacyl-tRNA
[114435]synthetase as a regulator of angiogenesis. *Proc. Nat. Acad.*
Sci. 99:
[114436]173–177, 2002.
[114437]20209.
- [114438]Comings, D. E.; Comings, B. G.; Dietz, G.; Muhleman, D.:
Localization
[114439]of human tryptophan oxygenase to 4q31: possible relevance to alcoholism,
[114440]depression and Tourette syndrome. (Abstract) *Am. J. Hum.*
Genet. 45
[114441](suppl.): A135 only, 1989.
[114442]20210.

[114443]Comings, D. E.; Dietz, G.; Muhlman, D.: Localization of human
tryptophan oxygenase to 4q25-q31. (Abstract) Cytogenet.
Cell Genet. 51:
[114445]979 only, 1989.
[114446]20211.
[114447]Comings, D. E.; Muhleman, D.; Dietz, G.; Sherman, M.;
Forest, G.
[114448]L.: Sequence of human tryptophan 2,3-dioxygenase
(TDO2): presence
[114449]of a glucocorticoid response-like element composed of a
GTT repeat
[114450]and an intronic CCCCT repeat. Genomics 29: 390-396,
1995.
[114451]20212.
[114452]Comings, D. E.; Muhleman, D.; Dietz, G. W., Jr.; Donlon,
T.: Human
[114453]tryptophan oxygenase localized to 4q31: possible impli-
cations for
[114454]alcoholism and other behavioral disorders. Genomics 9:
301-308,
[114455]1991.

[114456]20213.

[114457]Donlon, T. A.; Muhleman, D.; Dietz, G.; Comings, D. E.;
Spak, D.

[114458]K.: Localization of human tryptophan oxygenase to
4q31-q32 by in

[114459]situ hybridization. (Abstract) Cytogenet. Cell Genet. 51:
992 only,

[114460]1989.

[114461]20214.

[114462]Miller, J. S.; Moxley, G.; Schwartz, L. B.: Cloning and char-
acterization

[114463]of a second complementary DNA for human tryptase. J.
Clin. Invest. 86:

[114464]864-870, 1990.

[114465]20215.

[114466]Miller, J. S.; Westin, E. H.; Schwartz, L. B.: Cloning and
characterization

[114467]of complementary DNA for human tryptase. J. Clin. Invest.
84: 1188-1195,

[114468]1989.

[114469]20216.

- [114470]Vanderslice, P.; Ballinger, S. M.; Tam, E. K.; Goldstein, S. M.;
- [114471]Craig, C. S.; Caughey, G. H.: Human mast cell tryptase: multiple
- [114472]cDNAs and genes reveal a multigene serine protease family. Proc.
- [114473]Nat. Acad. Sci. 87: 3811–3815, 1990.
- [114474]20217.
- [114475]Lewis, W. H.; Yeger, H.: Characterization of the aniridia-Wilms
- [114476]tumor association region of chromosome 11. (Abstract) Cytogenet.
- [114477]Cell Genet. 46: 650, 1987.
- [114478]20218.
- [114479]Mach, B.; Steimle, V.; Reith, W.: MHC class II-deficient combined
- [114480]immunodeficiency: a disease of gene regulation. Immun. Rev. 138:
- [114481]207–221, 1994.
- [114482]20219.
- [114483]Little, M.; Holmes, G.; Bickmore, W.; van Heyningen, V.;

Hastie,

[114484]N.; Wainwright, B.: DNA binding capacity of the WT1 protein is abolished

[114485]by Denys-Drash syndrome WT1 point mutations. Hum. Molec. Genet. 4:

[114486]351-358, 1995.

[114487]20220.

[114488]Little, M. H.; Williamson, K. A.; Mannens, M.; Kelsey, A.; Gosden,

[114489]C.; Hastie, N. D.; van Heyningen, V.: Evidence that WT1 mutations

[114490]in Denys-Drash syndrome patients may act in a dominant-negative fashion. Hum.

[114491]Molec. Genet. 2: 259-264, 1993.

[114492]20221.

[114493]Mannens, M.; Slater, R. M.; Heyting, C.; Blik, J.; de Kraker,

[114494]J.; Coad, N.; de Pagter-Holthuisen, P.; Pearson, P. L.: Molecular

[114495]nature of genetic changes resulting in loss of heterozygosity of chromosome

[114496]11 in Wilms' tumours. Hum. Genet. 81: 41-48, 1988.

[114497]20222.

[114498]Marshall, L. S.; Qureshi, A. R.; DiGeorge, A. M.; Kistener,
macher,

[114499]M. L.; Punnett, H. H.: Aniridia and the 11p13 deletion.
(Abstract) Am.

[114500]J. Hum. Genet. 34: 74A, 1982.

[114501]20223.

[114502]Matsunaga, E.: Genetics of Wilms' tumor. Hum. Genet. 57:
231-246,

[114503]1981.

[114504]20224.

[114505]McCoy, F. E., Jr.; Franklin, W. A.; Aronson, A. J.; Spargo, B.

[114506]H.: Glomerulonephritis associated with male pseudo-
hermaphroditism

[114507]and nephroblastoma. Am. J. Surg. Path. 7: 387-395, 1983.

[114508]20225.

[114509]Meadows, A. T.; Lichtenfeld, J. L.; Koop, C. E.: Wilms' tu-
mor

[114510]in three children of a woman with congenital hemihyper-
trophy. New

[114511]Eng. J. Med. 291: 23-24, 1974.

[114512]20226.

[114513]Michalopoulos, E. E.; Bevilacqua, P. J.; Stokoe, N.; Powers, V.

[114514]E.; Willard, H. F.; Lewis, W. H.: Molecular analysis of gene deletion

[114515]in aniridia--Wilms tumor association. Hum. Genet. 70: 157-162, 1985.

[114516]20227.

[114517]Miller, R. W.; Fraumeni, J. F., Jr.; Manning, M. D.: Association

[114518]of Wilms' tumor with aniridia, hemihypertrophy and other congenital

[114519]malformations. New Eng. J. Med. 270: 922-927, 1964.

[114520]20228.

[114521]Miyagawa, K.; Kent, J.; Moore, A.; Charlieu, J.-P.; Little, M.

[114522]H.; Williamson, K. A.; Kelsey, A.; Brown, K. W.; Hassam, S.; Briner,

[114523]J.; Hayashi, Y.; Hirai, H.; Yazaki, Y.; van Heyningen, V.; Hastie,

[114524]N. D.: Loss of WT1 function leads to ectopic myogenesis

in Wilms'

[114525]tumour. (Letter) Nature Genet. 18: 15–17, 1998.

[114526]20229.

[114527]Nakagome, Y.; Ise, T.; Sakurai, M.; Nakajo, T.; Okamoto,
E.; Takano,

[114528]T.; Nakahori, Y.; Tsuchida, Y.; Nagahara, N.; Takada, Y.;
Ohsawa,

[114529]Y.; Sawaguchi, S.; Toyosaka, A.; Kobayashi, N.; Matsunaga,
E.; Saito,

[114530]S.: High-resolution studies in patients with aniridia-Wilms
tumor

[114531]association, Wilms tumor or related congenital abnormali-
ties. Hum.

[114532]Genet. 67: 245–248, 1984.

[114533]20230.

[114534]Neidhardt, M.: Wilms-tumor und Aniridie--ein genetisch
fixiertes

[114535]Syndrome? Klin. Paediat. 184: 312–316, 1972.

[114536]20231.

[114537]Nordenskjold, A.; Fricke, G.; Anvret, M.: Absence of muta-
tions

[114538]in the WT1 gene in patients with XY gonadal dysgenesis.

Hum. Genet. 96:

[114539]102–104, 1995.

[114540]20232.

[114541]Orkin, S. H.; Goldman, D. S.; Sallan, S. E.: Development of
homozygosity

[114542]for chromosome 11p markers in Wilms' tumour. Nature
309: 172–174,

[114543]1984.

[114544]20233.

[114545]Park, S.; Bernard, A.; Bove, K. E.; Sens, D. A.; Hazen–Mar-
tin,

[114546]D. J.; Garvin, A. J.; Haber, D. A.: Inactivation of WT1 in
nephrogenic

[114547]rests, genetic precursors to Wilms' tumour. Nature Genet.
5: 363–367,

[114548]1993.

[114549]20234.

[114550]Park, S.; Tomlinson, G.; Nisen, P.; Haber, D. A.: Altered
trans–activational

[114551]properties of a mutated WT1 gene product in a WAGR–

associated Wilms'

[114552]tumor. Cancer Res. 53: 4757-4760, 1993.

[114553]20235.

[114554]Puissant, H.; Azoulay, M.; Serre, J.-L.; Piet, L.; Junien, C.

[114555]: Molecular analysis of a reciprocal translocation

t(5;11)(q11;p13)

[114556]in a WAGR patient. Hum. Genet. 79: 280-282, 1988.

[114557]20236.

[114558]Rahman, N.; Arbour, L.; Tonin, P.; Renshaw, J.; Pelletier, J.;

[114559]Baruchel, S.; Pritchard-Jones, K.; Stratton, M. R.; Narod, S.

A.:

[114560]Evidence for a familial Wilms' tumour gene (FWT1) on

chromosome 17q12-q21. Nature

[114561]Genet. 13: 461-463, 1996.

[114562]20237.

[114563]Reeve, A. E.; Housiaux, P. J.; Gardner, R. J. M.; Chewings,

W.

[114564]E.; Grindley, R. M.; Millow, L. J.: Loss of a Harvey ras allele

in

[114565]sporadic Wilms' tumour. Nature 309: 174-176, 1984.

[114566]20238.

[114567]Riccardi, V. M.; Hittner, H. M.; Strong, L. C.; Fernbach, D.
J.;

[114568]Lebo, R.; Ferrell, R. E.: Wilms tumor with aniridia/iris dys-
plasia

[114569]and apparently normal chromosomes. J. Pediat. 100:
574–577, 1982.

[114570]20239.

[114571]Patek, C. E.; Little, M. H.; Fleming, S.; Miles, C.; Charlieu,

[114572]J.-P.; Clarke, A. R.; Miyagawa, K.; Christie, S.; Doig, J.;
Harrison,

[114573]D. J.; Porteous, D. J.; Brookes, A. J.; Hooper, M. L.; Hastie,
N.

[114574]D.: A zinc finger truncation of murine WT1 results in the
characteristic

[114575]urogenital abnormalities of Denys–Drash syndrome. Proc.
Nat. Acad.

[114576]Sci. 96: 2931–2936, 1999.

[114577]20240.

[114578]Riccardi, V. M.; Sujansky, E.; Smith, A. C.; Francke, U.:
Chromosomal

[114579] imbalance in the aniridia--Wilms' tumor association: 11p
interstitial

[114580] deletion. Pediatrics 61: 604-610, 1978.

[114581] 20241.

[114582] Salazar, H.; Kanbour, A.; Burgess, F.: Ultrastructure and
observations

[114583] on the histogenesis of mesotheliomas 'adenomatoid tu-
mors' of the female

[114584] genital tract. Cancer 29: 141-152, 1972.

[114585] 20242.

[114586] Schmickel, R. D.: Chromosomal deletions and enzyme de-
ficiencies. J.

[114587] Pediat. 108: 244-246, 1986.

[114588] 20243.

[114589] Schroeder, W. T.; Chao, L.-Y.; Dao, D. D.; Strong, L. C.;
Pathak,

[114590] S.; Riccardi, V.; Lewis, W. H.; Saunders, G. F.: Nonrandom
loss of

[114591] maternal chromosome 11 alleles in Wilms tumors. Am. J.
Hum. Genet. 40:

[114592] 413-420, 1987.

[114593]20244.

[114594]Schumacher, V.; Schneider, S.; Figge, A.; Wildhardt, G.;
Harms,

[114595]D.; Schmidt, D.; Weirich, A.; Ludwig, R.; Royer-Pokora, B.:
Correlation

[114596]of germ-line mutations and two-hit inactivation of the
WT1 gene with

[114597]Wilms tumors of stromal-predominant histology. Proc.
Nat. Acad. Sci. 94:

[114598]3972-3977, 1997.

[114599]20245.

[114600]Babcock, M.; de Silva, D.; Oaks, R.; Davis-Kaplan, S.; Ji-
ralerspong,

[114601]S.; Montermini, L.; Pandolfo, M.; Kaplan, J.: Regulation of
mitochondrial

[114602]iron accumulation by Yfh1p, a putative homolog of
frataxin. Science 276:

[114603]1709-1712, 1997.

[114604]20246.

[114605]Bidichandani, S. I.; Ashizawa, T.; Patel, P. I.: The GAA
triplet-repeat

[114606] expansion in Friedreich ataxia interferes with transcription and may

[114607] be associated with an unusual DNA structure. Am. J. Hum. Genet. 62:

[114608] 111–121, 1998.

[114609] 20247.

[114610] Bidichandani, S. I.; Ashizawa, T.; Patel, P. I.: Atypical Friedreich

[114611] ataxia caused by compound heterozygosity for a novel missense mutation

[114612] and the GAA triplet-repeat expansion. (Letter) Am. J. Hum. Genet. 60:

[114613] 1251–1256, 1997.

[114614] 20248.

[114615] Cavadini, P.; Gellera, C.; Patel, P. I.; Isaya, G.: Human frataxin

[114616] maintains mitochondrial iron homeostasis in *Saccharomyces cerevisiae*. Hum.

[114617] Molec. Genet. 9: 2523–2530, 2000.

[114618] 20249.

[114619] MacPherson, A. I. S.; Richmond, J.; Donaldson, G. W. K.;

Muir,

[114620]A. R.: The role of the spleen in congenital spherocytosis.

Am. J.

[114621]Med. 50: 35–41, 1971.

[114622]20250.

[114623]Masera, G.; Mieli, G.; Petrone, M.; Porcelli, P.: Transient
aplastic

[114624]crisis in hereditary spherocytosis. Acta Haemat. 63:
28–31, 1980.

[114625]20251.

[114626]Miraglia del Giudice, E.; Perrotta, S.; Nobili, B.; Specchia,

[114627]C.; d'Urzo, G.; Iolascon, A.: Coinheritance of Gilbert syn-
drome increases

[114628]risk for developing gallstones in patients with hereditary
spherocytosis. Blood 94:

[114629]2259–2262, 1999.

[114630]20252.

[114631]Miwa, S.; Tanaka, K. R.; Valentine, W. N.: Enolase activity
of

[114632]erythrocytes in hereditary spherocytosis. Nature 195:
613–614, 1962.

[114633]20253.

[114634]Mohler, D. N.; Wheby, M. S.: Patients with hereditary spherocytosis

[114635]may have clinically significant iron overload when they are also heterozygous

[114636]for hemochromatosis. Trans. Am. Clin. Climatol. Assoc. 96: 34-40,

[114637]1984.

[114638]20254.

[114639]Mohler, D. N.; Wheby, M. S.: Hemochromatosis heterozygotes may

[114640]have significant iron overload when they also have hereditary spherocytosis. Am.

[114641]J. Med. Sci. 292: 320-324, 1986.

[114642]20255.

[114643]Moiseyev, V. S.; Korovina, E. A.; Polotskaya, E. L.; Poliyan-skaya,

[114644]I. S.; Yazdovsky, V. V.: Hypertrophic cardiomyopathy associated with

[114645]hereditary spherocytosis in three generations of one family.(Letter) Lancet II:

[114646]853–854, 1987.

[114647]20256.

[114648]Morton, N. E.; MacKinney, A. A.; Kosower, N. S.; Schilling, R.

[114649]F.; Gray, M. P.: Genetics of spherocytosis. Am. J. Hum. Genet. 14:

[114650]170–184, 1962.

[114651]20257.

[114652]Motulsky, A. G.; Anderson, R.; Sparkes, R. S.; Huestis, R. H.

[114653]Marrow transplantation in newborn mice with hereditary spherocytosis.

[114654]A model system. Trans. Assoc. Am. Phys. 75: 64–72, 1962.

[114655]20258.

[114656]Nakashima, K.; Yamauchi, K.; Miwa, S.; Fujimura, K.; Mizutani,

[114657]A.; Kuramoto, A.: Glutathione reductase deficiency in a kindred with

[114658]hereditary spherocytosis. Am. J. Hemat. 4: 141–150, 1978.

[114659]20259.

[114660]Ng, J.-P.; Cumming, R. L. C.; Horn, E. H.; Hogg, R. B.:
Hereditary

[114661]spherocytosis revealed by human parvovirus infection.(Letter) Brit.

[114662]J. Haemat. 65: 379-380, 1987.

[114663]20260.

[114664]Nozawa, Y.; Noguchi, T.; Iida, H.; Fukushima, H.; Sekiya,
T.;

[114665]Ito, Y.: Erythrocyte membrane of hereditary spherocytosis:
alteration

[114666]in surface ultrastructure and membrane proteins, as inferred by scanning

[114667]electron microscopy and SDS-disc gel electrophoresis.
Clin. Chim.

[114668]Acta 55: 81-86, 1974.

[114669]20261.

[114670]Okamoto, N.; Wada, Y.; Nakamura, Y.; Nakayama, M.;
Chiyo, H.;

[114671]Murayama, K.; Inoue, T.; Kanzaki, A.; Yawata, Y.; Hirono,
A.; Miwa,

[114672]S.: Hereditary spherocytic anemia with deletion of the
short arm
[114673]of chromosome 8. Am. J. Med. Genet. 58: 225–229, 1995.
[114674]20262.

[114675]Peretz, E.; Hallel–Halevy, D.; Grunwald, M. H.; Halevy, S.:
Hereditary
[114676]spherocytosis with leg ulcers which healed after splenec-
tomy. Europ.
[114677]J. Derm. 7: 527–528, 1997.
[114678]20263.

[114679]Rao, K. R. P.; Patel, A. R.; Anderson, M. J.; Hodgson, J.;
Jones,
[114680]S. E.; Pattison, J. R.: Infection with parvovirus–like agent
and
[114681]aplastic crisis in adults with chronic hemolytic anemia.
Ann. Intern.
[114682]Med. 98: 930–932, 1983.
[114683]20264.

[114684]Reznikoff–Etievant, M. F.; Bonaiti, C.; Maigret, P.;
Malvoisin,
[114685]A.; Maynier, M.; Mesnard, G.; Haupman, G.: Hereditary

spherocytosis

[114686]linkage. Brit. J. Haemat. 46: 153–155, 1980.

[114687]20265.

[114688]Sengar, D. P. S.; McLeish, W. A.; Smiley, R. K.; Luke, B.:

HLA

[114689]and hereditary spherocytosis. Vox Sang. 33: 278–279,
1977.

[114690]20266.

[114691]Shohet, S. B.: Reconstitution of spectrin–deficient spherocytic

[114692]mouse erythrocyte membranes. J. Clin. Invest. 64:
483–494, 1979.

[114693]20267.

[114694]Stratton, R. F.; Crudo, D. F.; Varela, M.; Shapira, E.: Deletion

[114695]of the proximal short arm of chromosome 8. Am. J. Med. Genet. 42:

[114696]15–18, 1992.

[114697]20268.

[114698]Tse, W. T.; Meninger, J.; Ward, D.; John, K.; Lux, S. E.; Forget,

[114699]B. G.: Genomic cloning and chromosomal sublocalization
of the human

[114700]ankyrin gene.(Abstract) Clin. Res. 38: 266A, 1990.

[114701]20269.

[114702]Tsukada, T.; Koike, T.; Koike, R.; Sanada, M.; Takahashi,
M.;

[114703]Shibata, A.; Nunoue, T.: Epidemic of aplastic crisis in pa-
tients

[114704]with hereditary spherocytosis in Japan.(Letter) Lancet 1:
1401,

[114705]1985.

[114706]20270.

[114707]White, R.; Barker, J.: Normoblastosis, a mutant mouse with
severe

[114708]hemolytic anemia.(Abstract) Blood 70S: 57a, 1987.

[114709]20271.

[114710]White, R. A.; Birkenmeier, C. S.; Lux, S. E.; Barker, J. E.:

[114711]Ankyrin and the hemolytic anemia mutation, nb, map to
mouse chromosome

[114712]8: presence of the nb allele is associated with a truncated
erythrocyte

[114713]ankyrin. Proc. Nat. Acad. Sci. 87: 3117–3121, 1990.

[114714]20272.

[114715]Wiley, J. S.: Co-ordinated increase of sodium leak and sodium

[114716]pump in hereditary spherocytosis. Brit. J. Haemat. 22: 529–542,

[114717]1972.

[114718]20273.

[114719]Wiley, J. S.; Firkin, B. G.: An unusual variant of hereditary

[114720]spherocytosis. Am. J. Med. 48: 63–71, 1970.

[114721]20274.

[114722]Zail, S. S.; Krawitz, E.; Viljoen, E.; Kramer, S.; Metz, J.:

[114723]Atypical hereditary spherocytosis: biochemical studies and sites of

[114724]erythrocyte destruction. Brit. J. Haemat. 13: 323–334, 1967.

[114725]20275.

[114726]Sukhatme, V. P.; Vollmer, A. C.; Erikson, J.; Isobe, M.; Croce,

[114727]C.; Parnes, J. R.: Gene for the human T cell differentiation

antigen

[114728]Leu-2/T8 is closely linked to the kappa light chain locus
on chromosome

[114729]2. J. Exp. Med. 161: 429-434, 1985.

[114730]20276.

[114731]Traver, D.; Akashi, K.; Manz, M.; Merad, M.; Miyamoto, T.;
Engleman,

[114732]E. G.; Weissman, I. L.: Development of CD8-alpha-positive
dendritic

[114733]cells from a common myeloid progenitor. Science 290:
2152-2154,

[114734]2000.

[114735]20277.

[114736]Weichhold, G. M.; Huber, C.; Parnes, J. R.; Zachau, H. G.:
The

[114737]CD8-alpha locus is located on the telomere side of the
immunoglobulin-kappa

[114738]locus at a distance of 2 Mb. Genomics 16: 512-514, 1993.

[114739]20278.

[114740]Angel, J. M.; Moore, J. L.; Pelphrey, A.; Richie, E. R.: The
mouse

[114741]homolog of the rhombotin (Ttg-1) gene maps on chromosome 7 distal

[114742]to the beta-globin (Hbb) locus. Mammalian Genome 4: 281-282, 1993.

[114743]20279.

[114744]Boehm, T.; Foroni, L.; Kennedy, M.; Rabbitts, T. H.: The rhombotin

[114745]gene belongs to a class of transcriptional regulators with a potential

[114746]novel protein dimerisation motif. Oncogene 5: 1103-1105, 1990.

[114747]20280.

[114748]Chen, Q.; Cheng, J. T.; Tasi, L. H.; Schneider, N.; Buchanan, G.;

[114749]Carroll, A.; Crist, W.; Ozanne, B.; Siciliano, M. J.; Baer, R.: The

[114750]TAL gene undergoes chromosome translocation in T cell leukemia and

[114751]potentially encodes a helix-loop-helix protein. EMBO J. 9: 415-424,

[114752]1990.

[114753]20281.

[114754]Greenberg, J.; Boehm, T.; Surane, A.; Keynes, R.; Sofronew, M.;

[114755]Rabbitts, T. H.: A human T cell translocation involving chromosome

[114756]11p15 identifies a gene expressed in developing rhombomeres of the

[114757]mouse hind brain. (Abstract) Cytogenet. Cell Genet. 51: 1008 only,

[114758]1989.

[114759]20282.

[114760]Honey, K.; Benlagha, K.; Beers, C.; Forbush, K.; Teyton, L.; Kleijmeer,

[114761]M. J.; Rudensky, A. Y.; Bendelac, A.: Thymocyte expression of cathepsin

[114762]L is essential for NKT cell development. Nature Immun. 7Oct, 2002.

[114763]Note: Advance Electronic Publication.

[114764]20283.

[114765]Adkison, L. R.; Taylor, S.; Beamer, W. G.: Mutant gene-induced

[114766]disorders of structure, function and thyroglobulin synthesis in congenital

[114767]goitre (cog/cog) in mice. J. Endocr. 126: 51–58, 1990.

[114768]20284.

[114769]Avvedimento, V. E.; Di Lauro, R.; Monticelli, A.; Bernardi, F.;

[114770]Patracchini, P.; Calzolari, E.; Martini, G.; Varrone, S.: Mapping

[114771]of human thyroglobulin gene on the long arm of chromosome 8 by in

[114772]situ hybridization. Hum. Genet. 71: 163–166, 1985.

[114773]20285.

[114774]Baas, F.; Bikker, H.; Geurts van Kessel, A.; Melsert, R.; Pearson,

[114775]P. L.; de Vijlder, J. J. M.; van Ommen, G.–J. B.: The human thyroglobulin

[114776]gene: a polymorphic marker localized distal to C–MYC on chromosome

[114777]8 band q24. Hum. Genet. 69: 138–143, 1985.

[114778]20286.

[114779]Baas, F.; van Ommen, G.–J. B.; Bikker, H.; Arnberg, A. C.; de Vijlder,

[114780]J. J. M.: The human thyroglobulin gene is over 300 kb long

and contains

[114781]introns of up to 64 kb. Nucleic Acids Res. 14: 5171–5186,
1986.

[114782]20287.

[114783]Beamer, W. G.; Maltais, L. J.; DeBaets, M. H.; Eicher, E. M.:

[114784]Inherited congenital goiter in mice. Endocrinology 120:
838–840,

[114785]1987.

[114786]20288.

[114787]Berge-Lefranc, J.-L.; Cartouzou, G.; Mattei, M.-G.; Pas-
sage, E.;

[114788]Malezet-Desmoulins, C.; Lissitzky, S.: Localization of the
thyroglobulin

[114789]gene by in situ hybridization to human chromosomes.
Hum. Genet. 69:

[114790]28–31, 1985.

[114791]20289.

[114792]Bernardi, F.; Patracchini, P.; Monticelli, A.; Varrone, S.;

Aiello,

[114793]V.; Calzolari, E.; Marchetti, G.; Avvedimento, V. E.: Human
thyroglobulin

[114794] gene is located on the terminal part of the long arm of chromosome

[114795] 8. (Abstract) Cytogenet. Cell Genet. 40: 582–583, 1985.

[114796] 20290.

[114797] Brocas, H.; Szpirer, J.; Lebo, R. V.; Levan, G.; Szpirer, C.; Cheung,

[114798] M. C.; Vassart, G.: The thyroglobulin gene resides on chromosome

[114799] 8 in man and on chromosome 7 in the rat. Cytogenet. Cell Genet. 39:

[114800] 150–153, 1985.

[114801] 20291.

[114802] Cochaux, P.; Ieiri, T.; Targovnik, H.; Suzuki, M.; Shimoda, S.-I.;

[114803] Perret, J.; Vassart, G.: Identification of a splicing mutation responsible

[114804] for a human hereditary goiter with hypothyroidism. (Abstract) Am.

[114805] J. Hum. Genet. 49 (suppl.): 131 only, 1991.

[114806] 20292.

[114807] Hashimoto, K.; Hirai, M.; Kurosawa, Y.: A gene outside the

human

[114808]MHC related to classical HLA class I genes. Science 269:
693–695,

[114809]1995.

[114810]20293.

[114811]Hashimoto, K.; Hirai, M.; Kurosawa, Y.: Identification of a
mouse

[114812]homolog for the human hereditary haemochromatosis
candidate gene. Biochem.

[114813]Biophys. Res. Commun. 230: 35–39, 1997.

[114814]20294.

[114815]Hickman, P. E.; Hourigan, L. F.; Powell, L. W.; Cordingley,
F.;

[114816]Dimeski, G.; Ormiston, B.; Shaw, J.; Ferguson, W.; Johnson,
M.; Ascough,

[114817]J.; McDonell, K.; Pink, A.; Crawford, D. H. G.: Automated
measurement

[114818]of unsaturated iron binding capacity is an effective
screening strategy

[114819]for C282Y homozygous haemochromatosis. Gut 46:
405–409, 2000.

[114820]20295.

[114821]Hill, W. G.; Robertson, A.: Linkage disequilibrium in finite
[114822]populations. Theor. Appl. Genet. 38: 226–231, 1968.
[114823]20296.

[114824]Jazwinska, E. C.; Cullen, L. M.; Busfield, F.; Pyper, W. R.;
Webb,

[114825]S. I.; Powell, L. W.; Morris, C. P.; Walsh T. P.: Haemochro-
matosis

[114826]and HLA-H. (Letter) Nature Genet. 14: 249–251, 1996.

[114827]20297.

[114828]Jazwinska, E. C.; Lee, S. C.; Webb, S. I.; Halliday, J. W.;
Powell,

[114829]L. W.: Localization of the hemochromatosis gene close to
D6S105. Am.

[114830]J. Hum. Genet. 53: 347–352, 1993.

[114831]20298.

[114832]Jazwinska, E. C.; Pyper, W. R.; Burt, M. J.; Francis, J. L.;
Goldwurm,

[114833]S.; Webb, S. I.; Lee, S. C.; Halliday, J. W.; Powell, L. W.:
Haplotype

[114834]analysis in Australian hemochromatosis patients: evidence
for a predominant

- [114835]ancestral haplotype exclusively associated with
hemochromatosis. Am.
- [114836]J. Hum. Genet. 56: 428–433, 1995.
- [114837]20299.
- [114838]Jeffrey, G. P.; Chakrabarti, S.; Hegele, R. A.; Adams, P. C.:
- [114839]Polymorphism in intron 4 of HFE may cause overestima-
tion of C282Y
- [114840]homozygote prevalence in haemochromatosis. (Letter)
Nature Genet. 22:
- [114841]325–326, 1999.
- [114842]20300.
- [114843]Johnson, G. B., Jr.; Frey, W. G., III: Familial aspects of idio-
pathic
- [114844]hemochromatosis. J.A.M.A. 179: 747–751, 1962.
- [114845]20301.
- [114846]Jouanolle, A.-M.; Yaouanq, J.; Blayau, M.; Perichon, M.;
Fauchet,
- [114847]R.; Font, M.-P.; Le Gall, J.-Y.; David, V.: HLA class I gene
polymorphism
- [114848]in genetic hemochromatosis. Hum. Genet. 85: 279–282,
1990.

[114849]20302.

[114850]Jouanolle, A. M.; Fergelot, P.; Gandon, G.; Yaouanq, J.; Le Gall,

[114851]J. Y.; David, V.: A candidate gene for hemochromatosis: frequency

[114852]of the C282Y and H63D mutations. Hum. Genet. 100: 544–547, 1997.

[114853]

[114854]100. Jouanolle, A. M.; Gandon, G.; Jezequel, P.; Blayau, M.; Champion,

[114855]M. L.; Yaouanq, J.; Mosser, J.; Fergelot, P.; Chauvel, B.; Bouric,

[114856]P.; Carn, G.; Andrieux, N.; Gicquel, I.; Le Gall, J.-Y.; David, V.

[114857]: Haemochromatosis and HLA-H. (Letter) Nature Genet. 14: 251–252,

[114858]1996.

[114859]

[114860]101. Kaikov, Y.; Wadsworth, L. D.; Hassall, E.; Dimmick, J. E.; Rogers,

[114861]P. C. J.: Primary hemochromatosis in children: report of three newly

[114862]diagnosed cases and review of the pediatric literature. Pediatrics 90:

[114863]37–42, 1992.

[114864]

[114865]102. Karlsson, M.; Ikkala, E.; Reunanen, A.; Takkunen, H.; Vuori,

[114866]E.; Makinen, J.: Prevalence of hemochromatosis in Finland. Acta

[114867]Med. Scand. 224: 385–390, 1988.

[114868]

[114869]103. Kidd, K. K.: Genetic linkage and hemochromatosis. (Editorial) New

[114870]Eng. J. Med. 301: 209–210, 1979.

[114871]

[114872]104. Kravitz, K.; Skolnick, M.; Cannings, C.; Carmelli, D.; Baty,

[114873]B.; Amos, B.; Johnson, A.; Mendell, N.; Edwards, C.; Cartwright, G.

[114874]: Genetic linkage between hereditary hemochromatosis and HLA. Am.

[114875]J. Hum. Genet. 31: 601–619, 1979.

[114876]

[114877]105. Krikker, M. A.: A foundation for hemochromatosis.
(Letter) Ann.

[114878]Intern. Med. 97: 782-783, 1982.

[114879]

[114880]106. Kuhl, P.; Kaltwasser, J. P.; Seidl, S.: HLA antigens in
patients

[114881]with idiopathic hemochromatosis (IH). Tissue Antigens 12:
398-401,

[114882]1978.

[114883]

[114884]107. Kushner, J. P.; Skolnick, M. H.; Edwards, C. Q.;
Goldgar, D.;

[114885]Griffen, L. M.; Drummond, C.: Advantageous hemochro-
matosis. (Letter) New

[114886]Eng. J. Med. 319: 1156, 1988.

[114887]

[114888]108. Lalouel, J. M.; Le Mignon, L.; Simon, M.; Fauchet, R.;
Bourel,

[114889]M.; Rao, D. C.; Morton, N. E.: Genetic analysis of idiopathic
hemochromatosis

[114890] using both qualitative (disease status) and quantitative
(serum iron)

[114891] information. Am. J. Hum. Genet. 37: 700–718, 1985.

[114892]

[114893] 109. Lamon, J. M.; Marynick, S. P.; Roseblatt, R.; Donnelly,
S.:

[114894] Idiopathic hemochromatosis in a young female: a case
study and review

[114895] of the syndrome in young people. Gastroenterology 76:
178–183, 1979.

[114896]

[114897] 110. Lebron, J. A.; Bennett, M. J.; Vaughn, D. E.; Chirino, A.
J.;

[114898] Snow, P. M.; Mintier, G. A.; Feder, J. N.; Bjorkman, P. J.:
Crystal

[114899] structure of the hemochromatosis protein HFE and char-
acterization

[114900] of its interaction with transferrin receptor. Cell 93:
111–123,

[114901] 1998.

[114902]

[114903]111. Leggett, B. A.; Halliday, J. W.; Brown, N. N.; Bryant, S.;
Powell,

[114904]L. W.: Prevalence of haemochromatosis amongst asymptomatic
Australians. Brit.

[114905]J. Haemat. 74: 525–530, 1990.

[114906]

[114907]112. Le Mignon, L.; Simon, M.; Fauchet, R.; Edan, G.; Le
Reun, M.;

[114908]Brissot, P.; Genetet, B.; Bourel, M.: An HLA–A1 association
with

[114909]the hemochromatosis allele? Clin. Genet. 24: 171–176,
1983.

[114910]

[114911]113. Levy, J. E.; Montross, L. K.; Cohen, D. E.; Fleming, M.
D.; Andrews,

[114912]N. C.: The C282Y mutation causing hereditary hemochro-
matosis does

[114913]not produce a null allele. Blood 94: 9–11, 1999.

[114914]

[114915]114. Lipinski, M.; Hors, J.; Saleun, J.–P.; Saddi, R.; Passa,
P.;

[114916] Lafaurie, S.; Feingold, N.; Dausset, J.: Idiopathic hemochromatosis:

[114917] linkage with HLA. Tissue Antigens 11: 471–474, 1978.

[114918]

[114919] 115. Lonjou, C.; Collins, A.; Ajioka, R. S.; Jorde, L. B.; Kushner,

[114920] J. P.; Morton, N. E.: Allelic association under map error and recombinational

[114921] heterogeneity: a tale of two sites. Proc. Nat. Acad. Sci. 95: 11366–11370,

[114922] 1998.

[114923]

[114924] 116. Looker, A. C.; Johnson, C. J.: Prevalence of elevated serum

[114925] transferrin saturation in adults in the United States. Ann. Intern.

[114926] Med. 129: 940–945, 1998.

[114927]

[114928] 117. Lord, D. K.; Dunham, I.; Campbell, R. D.; Bomford, A.; Strachan,

[114929] T.; Cox, T. M.: Molecular analysis of the human MHC class

I region

[114930]in hereditary haemochromatosis: a study by pulsed-field gel electrophoresis. Hum.

[114931]Genet. 85: 531–536, 1990.

[114932]

[114933]118. Lucotte, G.: Celtic origin of the C282Y mutation of hemochromatosis. Blood

[114934]Cells Molecules Dis. 24: 433–438, 1998.

[114935]

[114936]119. Lucotte, G.; Coulondre, C.: Association between a 10 kb PvuII

[114937]restriction fragment of genomic DNA with the hemochromatosis gene. Exp.

[114938]Clin. Immunogenet. 3: 219–223, 1986.

[114939]

[114940]120. MacSween, R. N. M.; Scott, A. R.: Hepatic cirrhosis: a clinico-pathological

[114941]review of 520 cases. J. Clin. Path. 26: 936–942, 1973.

[114942]

[114943]121. Maddrey, W. C.; Hamilton, S. R.; Belitsos, N. J.: Famil–

ial hemochromatosis. Johns

[114944]Hopkins Med. J. 144: 66–69, 1979.

[114945]

[114946]122. McGill, J. R.; Naylor, S. L.; Sakaguchi, A. Y.; Moore, C.
M.;

[114947]Boyd, D.; Barrett, K. J.; Shows, T. B.; Drysdale, J. W.: Hu-
man ferritin

[114948]H and L sequences lie on ten different chromosomes.
Hum. Genet. 76:

[114949]66–72, 1987.

[114950]

[114951]123. Mercier, B.; Mura, C.; Ferec, C.: Putting a hold on
'HLA-H.'

[114952](Letter) Nature Genet. 15: 234 only, 1997.

[114953]

[114954]124. Merryweather-Clarke, A. T.; Pointon, J. J.; Shearman,
J. D.;

[114955]Robson, K. J. H.: Global prevalence of putative
haemochromatosis

[114956]mutations. J. Med. Genet. 34: 275–278, 1997.

[114957]

- [114958]125. Merryweather-Clarke, A. T.; Simonsen, H.; Shearman, J. D.; Pointon, J. J.; Norgaard-Pedersen, B.; Robson, K. J. H.: A retrospective anonymous pilot study in screening newborns for HFE mutations in Scandinavian populations. Hum. Mutat. 13: 154-159, 1999.
- [114962]
- [114963]126. Meyer, T. E.; Ballot, D.; Bothwell, T. H.; Green, A.; Derman, D. P.; Baynes, R. D.; Jenkins, T.; Jooste, P. L.; du Toit, E. D.; Jacobs, P. J.: The HLA linked iron loading gene in an Afrikaner population. J. Med. Genet. 24: 348-356, 1987.
- [114967]
- [114968]127. Milman, N.: Hereditary haemochromatosis in Denmark 1950-1985: clinical, biochemical and histological features in 179 patients and 13 preclinical cases. Danish Med. Bull. 38: 385-393, 1991.

[114971]

[114972]128. Milman, N.; Eiberg, H.; Thymann, M.; Fenger, K.:
Transferrin

[114973]subtypes in 51 Danish patients with hereditary
haemochromatosis and

[114974]in 847 normal subjects. Hum. Genet. 88: 475–476, 1992.

[114975]

[114976]129. Milman, N.; Graudal, N.; Nielsen, L. S.; Fenger, K.:
HLA determinants

[114977]in 70 Danish patients with idiopathic haemochromatosis.
Clin. Genet. 33:

[114978]286–292, 1988.

[114979]

[114980]130. Milman, N.; Graudal, N.; Nielsen, L. S.; Fenger, K.: An
HLA

[114981]study in 74 Danish haemochromatosis patients and in 21
of their families. Clin.

[114982]Genet. 41: 6–11, 1992.

[114983]

[114984]131. Milman, N.; Graudal, N.; Nielsen, L. S.; Mathiassen,
B.; Tauris,

- [114985]P.; Lund, B.; Kristensen, J. S.; Fenger, K.: Family studies of hereditary hemochromatosis in Denmark and the Faroe Islands. Hum. Genet. 85: 228–232, 1990.
- [114988]
- [114989]132. Muir, W. A.; McLaren, G. D.; Braun, W.; Askari, A.: Evidence for heterogeneity in hereditary hemochromatosis: evaluation of 174 persons in nine families. Am. J. Med. 76: 806–814, 1984.
- [114992]
- [114993]133. Muir, W. A.; McLaren, G. D.; Braun, W. E.; Askari, A. K.: Hereditary hemochromatosis: dominant inheritance and probable linkage to HLA. (Abstract) Am. J. Hum. Genet. 30: 61A, 1978.
- [114996]
- [114997]134. Mura, C.; Le Gac, G.; Scotet, V.; Ragueneas, O.; Mercier, A.-Y.; Ferec, C.: Variation of iron loading expression in C282Y

homozygous

[114999]haemochromatosis probands and sib pairs. (Letter) J. Med. Genet. 38:

[115000]632–636, 2001.

[115001]

[115002]135. Mura, C.; Nousbaum, J.-B.; Verger, P.; Moalic, M.-T.; Raguenees,

[115003]O.; Mercier, A.-Y.; Ferec, C.: Phenotype–genotype correlation in

[115004]haemochromatosis subjects. Hum. Genet. 101: 271–276, 1997.

[115005]

[115006]136. Mura, C.; Raguenees, O.; Ferec, C.: HFE mutations analysis in

[115007]711 hemochromatosis probands: evidence for S65C implication in mild

[115008]form of hemochromatosis. Blood 93: 2502–2505, 1999.

[115009]

[115010]137. Murphy, P. A.: Personal Communication. Baltimore, Md. 12/12/1987.

[115011]

[115012]138. Niederau, C.; Fischer, R.; Sonnenberg, A.; Stremmel, W.; Trampisch,

[115013]H. J.; Strohmeyer, G.: Survival and causes of death in cirrhotic

[115014]and in noncirrhotic patients with primary hemochromatosis. New Eng.

[115015]J. Med. 313: 1256–1262, 1985.

[115016]

[115017]139. Noll, W. W.; Belloni, D. R.; Stenzel, T. T.; Grody, W. W.: Polymorphism

[115018]in intron 4 of HFE does not compromise haemochromatosis mutation results

[115019](Letter) Nature Genet. 23: 271–272, 1999.

[115020]

[115021]140. Nussbaumer, T.; Plattner, H. C.; Rywlin, A. M.: Hemochromatose

[115022]juvenile chez trois soeurs et un frere avec consanguinite des parents:

[115023]etude anatomoclinique et genetique du syndrome endocrinohepato-myocardique. J.

[115024]Genet. Hum. 1: 53–59, 1952.

[115025]

[115026]141. Olsson, K. S.; Eriksson, K.; Ritter, B.; Heedman, P. A.:
Screening
[115027]for iron overload using transferrin saturation. Acta Med.
Scand. 215:
[115028]105–112, 1984.

[115029]

[115030]142. Olsson, K. S.; Ritter, B.; Rosen, U.; Heedman, P. A.;
Staugard,
[115031]F.: Prevalence of iron overload in central Sweden. Acta
Med. Scand. 213:
[115032]145–150, 1983.

[115033]

[115034]143. Olynyk, J. K.; Cullen, D. J.; Aquilia, S.; Rossi, E.; Sum-
merville,
[115035]L.; Powell, L. W.: A population-based study of the clinical
expression
[115036]of the hemochromatosis gene. New Eng. J. Med. 341:
718–724, 1999.

[115037]

[115038]144. Parkkila, S.; Waheed, A.; Britton, R. S.; Bacon, B. R.;

Zhou,

[115039]X. Y.; Tomatsu, S.; Fleming, R. E.; Sly, W. S.: Association of
the

[115040]transferrin receptor in human placenta with HFE, the pro-
tein defective

[115041]in hereditary hemochromatosis. Proc. Nat. Acad. Sci. 94:
13198-13202,

[115042]1997.

[115043]

[115044]145. Parkkila, S.; Waheed, A.; Britton, R. S.; Feder, J. N.;
Tsuchihashi,

[115045]Z.; Schatzman, R. C.; Bacon, B. R.; Sly, W. S.: Immunohis-
tochemistry

[115046]of HLA-H, the protein defective in patients with hereditary
hemochromatosis,

[115047]reveals unique pattern of expression in gastrointestinal
tract. Proc.

[115048]Nat. Acad. Sci. 94: 2534-2539, 1997.

[115049]

[115050]146. Perkins, K. W.; McInnes, I. W. S.; Blackburn, C. R. B.;
Beal,

[115051]R. W.: Idiopathic hemochromatosis in children: report of a

family. Am.

[115052]J. Med. 39: 118–126, 1965.

[115053]

[115054]147. Phatak, P. D.; Sham, R. L.; Raubertas, R. F.; Dunnigan, K.; O'Leary,

[115055]M. T.; Braggins, C.; Cappuccio, J. D.: Prevalence of hereditary hemochromatosis

[115056]in 16,031 primary care patients. Ann. Intern. Med. 129: 954–961,

[115057]1998.

[115058]

[115059]148. Pollycove, M.: Hemochromatosis. In: Stanbury, J. B.; Wyngaarden,

[115060]J. B.; Fredrickson, D. S.: The Metabolic Basis of Inherited Disease.

[115061]New York: McGraw–Hill (pub.) (3rd ed.): 1972. Pp. 1051–1084.

[115062]

[115063]149. Poullis, A.; Moodie, S. J.; Maxwell, J. D.: Clinical haemochromatosis

[115064]in HFE mutation carriers. Lancet 360: 411–412, 2002.

[115065]

[115066]150. Powell, L. W.; Ferluga, J.; Halliday, J. W.; Bassett, M. L.;

[115067]Kohonen-Corish, M.; Serjeantson, S.: Genetic hemochromatosis and

[115068]HLA linkage. Hum. Genet. 77: 55-56, 1987.

[115069]

[115070]151. Powell, L. W.; George, D. K.; McDonnell, S. M.; Kowdley, K. V.

[115071]: Diagnosis of hemochromatosis. Ann. Intern. Med. 129: 925-931,

[115072]1998.

[115073]

[115074]152. Pozzato, G.; Zorat, F.; Nascimben, F.; Gregorutti, M.; Comar,

[115075]C.; Baracetti, S.; Vatta, S.; Bevilacqua, E.; Begnano, A.; Crovella,

[115076]S.; Amoroso, A.: Haemochromatosis gene mutations in a clustered Italian

[115077]population: evidence of high prevalence in people of Celtic ancestry. Europ.

[115078]J. Hum. Genet. 9: 445–451, 2001.

[115079]

[115080]153. Rabinovitz, M.; Gavalier, J. S.; Kelly, R. H.; Van Thiel, D. H.

[115081]: Association between heterozygous alpha-1-antitrypsin deficiency

[115082]and genetic hemochromatosis. Hepatology 16: 145–148, 1992.

[115083]

[115084]154. Raha-Chowdhury, R.; Bowen, D. J.; Worwood, M.: A new highly

[115085]polymorphic marker in the 5-prime untranslated region of HLA-F shows

[115086]strong allelic association with haemochromatosis. Hum. Genet. 97:

[115087]228–231, 1996.

[115088]

[115089]155. Rhodes, D. A.; Raha-Chowdhury, R.; Cox, T. M.; Trowsdale, J.

[115090]: Homozygosity for the predominant Cys282Tyr mutation and absence

[115091]of disease expression in hereditary haemochromatosis. J.

Med. Genet. 34:

[115092]761–764, 1997.

[115093]

[115094]156. Risch, N.: Haemochromatosis, HFE and genetic complexity. (Letter) Nature

[115095]Genet. 17: 375–376, 1997.

[115096]

[115097]157. Ritter, B.; Safwenberg, J.; Olsson, K. S.: HLA as a marker of

[115098]the hemochromatosis gene in Sweden. Hum. Genet. 68: 62–66, 1984.

[115099]

[115100]158. Robson, K. J. H.; Shearman, J. D.; Merryweather–Clarke, A. T.;

[115101]Pointon, J. J.; Rosenberg, W. M.; Walker, A. P.; Dooley, J. S.; Bomford,

[115102]A.; Raha–Chowdhury, R.; Worwood, M.: Haemochromatosis: a gene at

[115103]last? J. Med. Genet. 34: 148–151, 1997.

[115104]

- [115105]159. Rochette, J.; Pointon, J. J.; Fisher, C. A.; Perera, G.; Arambepola, M.; Kodikara Arichchi, D. S.; De Silva, S.; Vandwalle, J. L.; Monti, J. P.; Old, J. M.; Merryweather-Clarke, A. T.; Weatherall, D. J.; Robson, K. J. H.: Multicentric origin of hemochromatosis gene (HFE) mutations. *Am. J. Hum. Genet.* 64: 1056–1062, 1999.
- [115110]
- [115111]160. Roldan, F. P.; Echenagusia, A. A.; Carro, P. G.: Hemochromatosis presenting as acute liver failure after iron supplementation. (Letter) *New Eng. J. Med.* 339: 269–270, 1998.
- [115114]
- [115115]161. Roth, M.-P.; Giraldo, P.; Hariti, G.; Poloni, E. S.; Sanchez-Mazas, A.; De Stefano, G. F.; Dugoujon, J.-M.; Coppin, H.: Absence of the hemochromatosis gene Cys282Tyr mutation in three ethnic groups from

[115118]Algeria (Mzab), Ethiopia, and Senegal. Immunogenetics
46: 222–225,

[115119]1997.

[115120]

[115121]162. Rothenberg, B. E.; Volland, J. R.: Beta-2 knockout
mice develop

[115122]parenchymal iron overload: a putative role for class I
genes of the

[115123]major histocompatibility complex in iron metabolism.
Proc. Nat. Acad.

[115124]Sci. 93: 1529–1534, 1996.

[115125]

[115126]163. Rowe, J. W.; Wands, J. R.; Mezey, S. E.; Waterbury, L.
A.; Wright,

[115127]J. R.; Tobin, J.; Andres, R.: Familial hemochromatosis:
characteristics

[115128]of the precirrhotic stage in a large kindred. Medicine 56:
197–211,

[115129]1977.

[115130]

[115131]164. Roy, C. N.; Andrews, N. C.: Recent advances in disor-

ders of

[115132]iron metabolism: mutations, mechanisms and modifiers.

Hum. Molec.

[115133]Genet. 10: 2181–2186, 2001.

[115134]

[115135]165. Roy, C. N.; Penny, D. M.; Feder, J. N.; Enns, C. A.: The
hereditary

[115136]hemochromatosis protein, HFE, specifically regulates
transferrin-mediated

[115137]iron uptake in HeLa cells. J. Biol. Chem. 274: 9022–9028,
1999.

[115138]

[115139]166. Ryan, E.; O'Keane, C.; Crowe, J.: Hemochromatosis in
Ireland

[115140]and HFE. Blood Cells Molecules Dis. 24: 428–432, 1998.

[115141]

[115142]167. Saddi, R.; Feingold, J.: Idiopathic haemochromatosis:
an autosomal

[115143]recessive disease. Clin. Genet. 5: 234–241, 1974.

[115144]

[115145]168. Sargent, T., III; Lim, T. H.; Jenson, R. L.: Reduced

chromium

[115146]retention in patients with hemochromatosis, a possible
basis of hemochromatotic

[115147]diabetes. Metabolism 28: 70–79, 1979.

[115148]

[115149]169. Sheldon, J. H.: Haemochromatosis. London: Oxford
Univ. Press

[115150](pub.) 1935. Pp. 19 only.

[115151]

[115152]170. Siemons, L. J.; Mahler, C.: Hypogonadotropic hypog-
onadism in

[115153]hemochromatosis: recovery of reproductive function after
iron depletion. J.

[115154]Clin. Endocr. Metab. 65: 585–587, 1987.

[115155]

[115156]171. Simon, M.; Alexandre, J. L.; Bourel, M.; Le Marec, B.;
Scordia,

[115157]C.: Heredity of idiopathic haemochromatosis: a study of
106 families. Clin.

[115158]Genet. 11: 327–341, 1977.

[115159]

[115160]172. Simon, M.; Bourel, M.; Fauchet, R.; Genetet, B.: Association

[115161]of HLA-A3 and HLA-B14 antigens with idiopathic haemochromatosis. Gut 17:

[115162]332-334, 1976.

[115163]

[115164]173. Simon, M.; Bourel, M.; Genetet, B.; Fauchet, R.:

Heredity of

[115165]idiopathic haemochromatosis. (Letter) Lancet I: 706, 1977.

[115166]

[115167]174. Simon, M.; Bourel, M.; Genetet, B.; Fauchet, R.: Idio-
pathic

[115168]hemochromatosis: demonstration of recessive transmis-
sion and early

[115169]detection by family HLA typing. New Eng. J. Med. 297:
1017-1021,

[115170]1977.

[115171]

[115172]175. Simon, M.; Fauchet, R.; Hespel, J. P.; Beaumont, C.;
Brissot,

[115173]P.; Hary, B.; De Nercy, H. Y. H.; Genetet, B.; Bourel, M.: Id-
iopathic

[115174]hemochromatosis: a study of biochemical expression in
247 heterozygous

[115175]members of 63 families: evidence for a single major HLA-
linked gene. Gastroenterology 78:

[115176]703–708, 1980.

[115177]

[115178]176. Simon, M.; Fauchet, R.; Le Gall, J. Y.; Brissot, P.;
Bourel,

[115179]M.: Immunogenetics of idiopathic hemochromatosis and
secondary iron

[115180]overload.In: Farid, N. R.: Immunogenetics of Endocrine
Disorders.

[115181]New York: Alan R. Liss (pub.) 1988. Pp. 345–371.

[115182]

[115183]177. Simon, M.; Le Mignon, L.; Fauchet, R.; Yaouanq, J.;
David, V.;

[115184]Edan, G.; Bourel, M.: A study of 609 HLA haplotypes
marking for the

[115185]hemochromatosis gene: (1) mapping of the gene near the
HLA-A locus

[115186]and characters required to define a heterozygous popula-
tion and (2)

[115187]hypothesis concerning the underlying cause of hemochro-
matosis-HLA

[115188]association. Am. J. Hum. Genet. 41: 89–105, 1987.

[115189]

[115190]178. Skolnick, M.: Personal Communication. Salt Lake City,
Utah

[115191]9/27/1983.

[115192]

[115193]179. Sorsby, A.: Clinical Genetics. St. Louis: C. V. Mosby
(pub.)

[115194]1953. Pp. 206 only.

[115195]

[115196]180. Sproule, T. J.; Jazwinska, E. C.; Britton, R. S.; Bacon,
B. R.;

[115197]Fleming, R. E.; Sly, W. S.; Roopenian, D. C.: Naturally vari-
ant autosomal

[115198]and sex-linked loci determine the severity of iron over-
load in beta-2-microglobulin-deficient

[115199]mice. Proc. Nat. Acad. Sci. 98: 5170-5174, 2001.

[115200]

[115201]181. Stevens, F. M.; Walters, J. M.; Watt, D. W.; McCarthy,
C. F.

[115202]: Inheritance of idiopathic haemochromatosis. (Letter)
Lancet I:

[115203]1107, 1977.

[115204]

[115205]182. Summers, K. M.; Tam, K. S.; Bartley, P. B.; Drysdale, J.; Zoghbi,

[115206]H. Y.; Halliday, J. W.; Powell, L. W.: Fine mapping of a human chromosome

[115207]6 ferritin heavy chain pseudogene: relevance to haemochromatosis. Hum.

[115208]Genet. 88: 175–178, 1991.

[115209]

[115210]183. Summers, K. M.; Tam, K. S.; Halliday, J. W.; Powell, L. W.:

[115211]HLA determinants in an Australian population of hemochromatosis patients

[115212]and their families. Am. J. Hum. Genet. 45: 41–48, 1989.

[115213]

[115214]184. Thenie, A. C.; Gicquel, I. M.; Hardy, S.; Ferran, H.; Fergelot,

[115215]P. Le Gall, J.-Y.; Mosser, J.: Identification of an endogenous RNA

[115216]transcribed from the antisense strand of the HFE gene.

Hum. Molec.

[115217]Genet. 10: 1859–1866, 2001.

[115218]

[115219]185. The UK Haemochromatosis Consortium: A simple genetic test

[115220]identifies 90% of UK patients with haemochromatosis. Gut 41: 841–844,

[115221]1997.

[115222]

[115223]186. Totaro, A.; Rommens, J. M.; Grifa, A.; Lunardi, C.; Carella,

[115224]M.; Huizenga, J. J.; Roetto, A.; Camaschella, C.; De Sandre, G.; Gasparini,

[115225]P.: Hereditary hemochromatosis: generation of a transcription map

[115226]within a refined and extended map of the HLA class I region. Genomics 31:

[115227]319–326, 1996.

[115228]

[115229]187. Townsend, A.; Drakesmith, H.: Role of HFE in iron metabolism,

[115230]hereditary haemochromatosis, anaemia of chronic disease, and secondary

[115231]iron overload. Lancet 359: 786–790, 2002.

[115232]

[115233]188. Trousseau, A.: Clinical Medicine Lectures Delivered at the Hotel-Dieu

[115234]Paris by A. Trousseau. Philadelphia: Blakiston (pub.) II: 1882.

[115235]pp. 313–315. Note: Translation: 3rd revised and enlarged edition by

[115236]Sir John Rose Cormack and P. Victor Bazire.....

[115237]

[115238]189. Trousseau, A.: Clinique Med. de l'Hotel de Paris. (pub.)

[115239]II: 1865. Pp. 663–698.

[115240]

[115241]190. Valberg, L. S.; Lloyd, D. A.; Ghent, C. N.; Flanagan, P. R.;

[115242]Sinclair, N. R.; Stiller, C. R.; Chamberlain, M. J.: Clinical and

[115243]biochemical expression of the genetic abnormality in idio-

pathic hemochromatosis. Gastroenterology 79:

[115244]884–892, 1980.

[115245]

[115246]191. Waheed, A.; Parkkila, S.; Saarnio, J.; Fleming, R. E.;
Zhou,

[115247]X. Y.; Tomatsu, S.; Britton, R. S.; Bacon, B. R.; Sly, W. S.:
Association

[115248]of HFE protein with transferrin receptor in crypt entero-
cytes of human

[115249]duodenum. Proc. Nat. Acad. Sci. 96: 1579–1584, 1999.

[115250]

[115251]192. Waheed, A.; Parkkila, S.; Zhou, X. Y.; Tomatsu, S.;
Tsuchihashi,

[115252]Z.; Feder, J. N.; Schatzman, R. C.; Britton, R. S.; Bacon, B.
R.;

[115253]Sly, W. S.: Hereditary hemochromatosis: effects of C282Y
and H63D

[115254]mutations on association with beta-2-microglobulin, in-
tracellular

[115255]processing, and cell surface expression of the HFE protein
in COS-7

[115256]cells. Proc. Nat. Acad. Sci. 94: 12384–12389, 1997.

[115257]

[115258]193. Wands, J. R.; Rowe, J. A.; Mezey, S. E.; Waterbury, L. A.; Wright,

[115259]J. R.; Halliday, J. W.; Isselbacher, K. J.; Powell, L. W.: Normal

[115260]serum ferritin concentrations in precirrhotic hemochromatosis. New

[115261]Eng. J. Med. 294: 302-305, 1976.

[115262]

[115263]194. Willis, G.; Wimperis, J. Z.; Lonsdale, R.; Fellows, I. W.; Watson,

[115264]M. A.; Skipper, L. M.; Jennings, B. A.: Incidence of liver disease

[115265]in people with HFE mutations. Gut 46: 401-404, 2000.

[115266]

[115267]195. Worwood, M.; Darke, C.; Trenchard, P.: Hereditary haemochromatosis

[115268]and blood donation. (Letter) Brit. Med. J. 302: 593, 1991.

[115269]

[115270]196. Yaouang, J.; Perichon, M.; Chorney, M.; Pontarotti, P.; Le Treut,

- [115271]A.; El Kahloun, A.; Mauvieux, V.; Blayau, M.; Jouanolle, A.
M.; Chauvel,
- [115272]B.; Moirand, R.; Nouel, O.; Le Gall, J. Y.; Feingold, J.; David,
V.
- [115273]: Anonymous marker loci within 400 kb of HLA-A gener-
ate haplotypes
- [115274]in linkage disequilibrium with the hemochromatosis gene
(HFE). Am.
- [115275]J. Hum. Genet. 54: 252-263, 1994.
- [115276]
- [115277]197. Yaouanq, J.; El Kahloun, A.; Chorney, M.; Jouanolle,
A. M.; Mauvieux,
- [115278]V.; Perichon, M.; Blayau, M.; Pontarotti, P.; Le Gall, J. Y.;
David,
- [115279]V.: Familial screening for genetic haemochromatosis by
means of DNA
- [115280]markers. J. Med. Genet. 29: 320-322, 1992.
- [115281]
- [115282]198. Zappone, E.; Dugast, I.; Papadopoulos, P.; Theriault,
K.; David,
- [115283]V.; LeGall, J.-Y.; Summers, K.; Powell, L.; Drysdale, J.:
Polymorphism

[115284]in a ferritin H gene from chromosome 6p. Hum. Genet.

86: 557–561,

[115285]1991.

[115286]

[115287]199. Zhou, X. Y.; Tomatsu, S.; Fleming, R. E.; Parkkila, S.;

Waheed,

[115288]A.; Jiang, J.; Fei, Y.; Brunt, E. M.; Ruddy, D. A.; Prass, C. E.;

[115289]Schatzman, R. C.; O'Neill, R.; Britton, R. S.; Bacon, B. R.;

Sly,

[115290]W. S.: HFE gene knockout produces mouse model of

hereditary hemochromatosis. Proc.

[115291]Nat. Acad. Sci. 95: 2492–2497, 1998.

[115292]

[115293]200. Zoller, H.; Pietrangelo, A.; Vogel, W.; Weiss, G.: Duo-

denal

[115294]metal-transporter (DMT-1, NRAMP-2) expression in pa-

tients with hereditary

[115295]haemochromatosis. Lancet 353: 2120–2123, 1999.

[115296]20303.

[115297]Azarnia, R.; Reddy, S.; Kmiecik, T. E.; Shalloway, D.;

Loewenstein,

- [115298]W. R.: The cellular src gene product regulates junctional cell-to-cell communication. Science 239: 398–401, 1988.
- [115300]20304.
- [115301]Czernilofsky, A. P.; Levinson, A. D.; Varmus, H. E.; Bishop, J.
- [115302]M.; Tischer, E.; Goodman, H.: Correction to the nucleotide sequence of the src gene of Rous sarcoma virus. Nature 301: 736–738, 1983.
- [115304]20305.
- [115305]Gibbs, C. P.; Tanaka, A.; Anderson, S. K.; Radul, J.; Baar, J.;
- [115306]Ridgway, A.; Kung, H.-J.; Fujita, D. J.: Isolation and structural mapping of a human c-src gene homologous to the transforming gene (v-src) of Rous sarcoma virus. J. Virol. 53: 19–24, 1985.
- [115309]20306.
- [115310]Irby, R. B.; Mao, W.; Coppola, D.; Kang, J.; Loubeau, J. M.; Trudeau, W.; Karl, R.; Fujita, D. J.; Jove, R.; Yeatman, T. J.: Activating

[115312]SRC mutation in a subset of advanced human colon cancers. Nature

[115313]Genet. 21: 187-190, 1999.

[115314]20307.

[115315]Le Beau, M. M.; Westbrook, C. A.; Diaz, M. O.; Rowley, J. D.:

[115316]c-src is consistently conserved in the chromosomal deletion (20q)

[115317]observed in myeloid disorders. Proc. Nat. Acad. Sci. 82: 6692-6696,

[115318]1985.

[115319]20308.

[115320]Lowe, C.; Yoneda, T.; Boyce, B. F.; Chen, H.; Mundy, G. R.; Soriano,

[115321]P.: Osteopetrosis in Src-deficient mice is due to an autonomous defect

[115322]of osteoclasts. Proc. Nat. Acad. Sci. 90: 4485-4489, 1993.

[115323]20309.

[115324]Morris, C. M.; Honeybone, L. M.; Hollings, P. E.; Fitzgerald,

[115325]P. H.: Localization of the SRC oncogene to chromosome band 20q11.2

[115326]and loss of this gene with deletion (20q) in two leukemic patients. Blood 74:

[115327]1768–1773, 1989.

[115328]20310.

[115329]Sakaguchi, A. Y.; Mohandas, T.; Naylor, S. L.: A human c–src

[115330]gene resides on the proximal long arm of chromosome 20 (cen–q13.1). Cancer

[115331]Genet. Cytogenet. 18: 123–129, 1985.

[115332]20311.

[115333]Sakaguchi, A. Y.; Naylor, S. L.; Weinberg, R. A.; Shows, T. B.

[115334]: Organization of human proto–oncogenes. (Abstract) Am. J. Hum. Genet. 34:

[115335]175A, 1982.

[115336]20312.

[115337]Sakaguchi, A. Y.; Zabel, B. U.; Grzeschik, K. H.; Law, M. L.;

[115338]Naylor, S. L.: Human proto–oncogene assignments. (Abstract) Cytogenet.

[115339]Cell Genet. 37: 572–573, 1984.

[115340]20313.

[115341]Wong, B. R.; Besser, D.; Kim, N.; Arron, J. R.; Vologodskaia,

[115342]M.; Hanafusa, H.; Choi, Y.: TRANCE, a TNF family member,
activates

[115343]Akt/PKB through a signaling complex involving TRAF6 and
c-Src. Molec.

[115344]Cell 4: 1041–1049, 1999.

[115345]20314.

[115346]Starzl, T. E.; Demetris, A. J.; Trucco, M.; Ricordi, C.; Ild-
stad,

[115347]S.; Terasaki, P. I.; Murase, N.; Kendall, R. S.; Kocova, M.;
Rudert,

[115348]W. A.; Zeevi, A.; Van Thiel, D.: Chimerism after liver trans-
plantation

[115349]for type IV glycogen storage disease and type 1 Gaucher's
disease. New

[115350]Eng. J. Med. 328: 745–749, 1993.

[115351]20315.

[115352]Trend, P. St. J.; Wiles, C. M.; Spencer, G. T.; Morgan-
Hughes,

[115353]J. A.; Lake, B. D.; Patrick, A. D.: Acid maltase deficiency in

adults:

[115354]diagnosis and management in five cases. Brain 108:
845–860, 1985.

[115355]20316.

[115356]Wokke, J. H. J.; Ausems, M. G. E. M.; van den Boogaard,
M.–J.

[115357]H.; Ippel, E. F.; van Diggelen, O.; Kroos, M. A.; Boer, M.;
Jennekens,

[115358]F. G. I.; Reuser, A. J. J.; Ploos van Amstel, H. K.: Genotype–
phenotype

[115359]correlation in adult–onset acid maltase deficiency. Ann.
Neurol. 38:

[115360]450–454, 1995.

[115361]20317.

[115362]Bao, Y.; Dawson, T. L., Jr.; Chen, Y.–T.: Human glycogen
debranching

[115363]enzyme gene (AGL): complete structural organization and
characterization

[115364]of the 5–prime flanking region. Genomics 38: 155–165,
1996.

[115365]20318.

[115366]Brunberg, J. A.; McCormick, W. F.; Schochet, S. S., Jr.: Type

[115367]III glycogenosis. An adult with diffuse weakness and muscle wasting. Arch.

[115368]Neurol. 25: 171-178, 1971.

[115369]20319.

[115370]Chen, Y.-T.; He, J.-K.; Ding, J.-H.; Brown, B. I.: Glycogen debranching

[115371]enzyme: purification, antibody characterization, and immunoblot analyses

[115372]of type III glycogen storage disease. Am. J. Hum. Genet. 41: 1002-1015,

[115373]1987.

[115374]20320.

[115375]Chen, Y. T.; Burchell, A.: Glycogen storage diseases. In: Scriver,

[115376]C. R.; Beaudet, A. L.; Sly, W. S.; Valle, D. (eds.): The Metabolic

[115377]and Molecular Bases of Inherited Disease. New York: McGraw-Hill

[115378](7th ed.): 1995. Pp. 935-965.

[115379]20321.

[115380]Cohen, J.; Friedman, M.: Renal tubular acidosis associated

with

[115381]type III glycogenosis. *Acta Paediat. Scand.* 68: 779–782,
1979.

[115382]20322.

[115383]Cohn, J.; Wang, P.; Hauge, M.; Henningsen, K.; Jensen, B.;
Svejgaard,

[115384]A.: Amylo-1,6-glucosidase deficiency (glycogenosis type
III) in the

[115385]Faroe Island. *Hum. Hered.* 25: 115–126, 1975.

[115386]20323.

[115387]Coleman, R. A.; Winter, H. S.; Wolf, B.; Chen, Y.-T.: Glyco-
gen

[115388]debranching enzyme deficiency: long-term study of
serum enzyme activities

[115389]and clinical features. *J. Inherit. Metab. Dis.* 15: 869–881,
1992.

[115390]20324.

[115391]Confino, E.; Pauzner, D.; Lidor, A.; Yedwab, G.; David, M.:
Pregnancy

[115392]associated with amylo-1,6-glucosidase deficiency (Forbes'
disease):

[115393]case report. *Brit. J. Obstet. Gynaec.* 91: 494–497, 1984.

[115394]20325.

[115395]DiMauro, S.; Hartwig, G. B.; Hays, A.; Eastwood, A. B.;
Franco,

[115396]R.; Olarte, M.; Chang, M.; Roses, A. D.; Fetell, M.; Schoen-
feldt,

[115397]R. S.; Stern, L. Z.: Debrancher deficiency: neuromuscular
disorder

[115398]in 5 adults. Ann. Neurol. 5: 422-436, 1979.

[115399]20326.

[115400]Fellows, I. W.; Lowe, J. S.; Ogilvie, A.; Stevens, A.; Toghill,

[115401]P. J.; Atkinson, M.: Type III glycogenosis presenting as
liver disease

[115402]in adults with atypical histological features. J. Clin. Path.
36:

[115403]431-434, 1983.

[115404]20327.

[115405]Fernandes, J.: The history of the glycogen storage disease.
(Letter) Europ.

[115406]J. Pediat. 154: 423-424, 1995.

[115407]20328.

- [115408]Garancis, J. C.; Panares, R. R.; Good, T. A.; Kuzma, J. F.:
Type
- [115409]3 glycogenosis. A biochemical and electron microscopic
study. Lab.
- [115410]Invest. 22: 468–477, 1970.
- [115411]20329.
- [115412]Hadjigeorgiou, G. M.; Comi, G. P.; Bordoni, A.; Shen, J.;
Chen,
- [115413]Y.–T.; Salani, S.; Toscano, A.; Fortunato, F.; Lucchiari, S.;
Bresolin,
- [115414]N.; Rodolico, C.; Piscaglia, M. G.; Franceschina, L.; Pa-
padimitriou,
- [115415]A.; Scarlato, G.: Novel donor splice site mutations of AGL
gene in
- [115416]glycogen storage disease type IIIa. J. Inherit. Metab. Dis.
22:
- [115417]762–763, 1999.
- [115418]20330.
- [115419]Levin, S.; Moses, S. W.; Chayoth, R.; Jadoga, N.; Steinitz, K.
- [115420]: Glycogen storage disease in Israel. A clinical, biochemi-
cal and

- [115421]genetic study. Israel J. Med. Sci. 3: 397–410, 1967.
- [115422]20331.
- [115423]Lucchiari, S.; Fogh, I.; Prella, A.; Parini, R.; Bresolin, N.;
- [115424]Melis, D.; Fiori, L.; Scarlato, G.; Comi, G. P.: Clinical and
genetic
- [115425]variability of glycogen storage disease type IIIa: seven
novel AGL
- [115426]gene mutations in the Mediterranean area. Am. J. Med.
Genet. 109:
- [115427]183–190, 2002.
- [115428]20332.
- [115429]Markowitz, A. J.; Chen, Y.-T.; Muenzer, J.; Delbuono, E. A.;
- Lucey,
- [115430]M. R.: A man with type III glycogenosis associated with
cirrhosis
- [115431]and portal hypertension. Gastroenterology 105:
1882–1885, 1993.
- [115432]20333.
- [115433]Miranda, A. F.; DiMauro, S.; Antler, A.; Stern, L. Z.; Row-
- land,
- [115434]L. P.: Glycogen debrancher deficiency is reproduced in

muscle culture. Ann.

[115435]Neurol. 9: 283–288, 1981.

[115436]20334.

[115437]Momoi, T.; Sano, H.; Yamanaka, C.; Sasaki, H.; Mikawa, H.:

Glycogen

[115438]storage disease type III with muscle involvement: reap-
praisal of phenotypic

[115439]variability and prognosis. Am. J. Med. Genet. 42: 696–699,
1992.

[115440]20335.

[115441]Moses, S. W.; Wanderman, K. L.; Myroz, A.; Frydman, M.:
Cardiac

[115442]involvement in glycogen storage disease type III. Europ. J.
Pediat. 148:

[115443]764–766, 1989.

[115444]20336.

[115445]Okubo, M.; Aoyama, Y.; Murase, T.: A novel donor splice
site

[115446]mutation in the glycogen debranching enzyme gene is as-
sociated with

[115447]glycogen storage disease type III. Biochem. Biophys. Res.

Commun. 224:

[115448]493–499, 1996.

[115449]20337.

[115450]Okubo, M.; Horinishi, A.; Makamura, N.; Aoyama, Y.;
Hashimoto,

[115451]M.; Endo, Y.; Murase, T.: A novel point mutation in an ac-
ceptor splice

[115452]site of intron 32 (IVS32 A–12–to–G) but no exon 3 muta-
tions in the

[115453]glycogen debranching enzyme gene in a homozygous pa-
tient with glycogen

[115454]storage disease type IIb. Hum. Genet. 102: 1–5, 1998.

[115455]20338.

[115456]Okubo, M.; Horinishi, A.; Suzuki, Y.; Murase, T.; Hayasaka,
K.

[115457]: Compound heterozygous patient with glycogen storage
disease type

[115458]III: identification of two novel AGL mutations, a donor
splice site

[115459]mutation of Chinese origin and a 1–bp deletion of
Japanese origin. Am.

[115460]J. Med. Genet. 93: 211–214, 2000.

[115461]20339.

[115462]Okubo, M.; Horinishi, A.; Takeuchi, M.; Suzuki, Y.; Sakura, N.;

[115463]Hasegawa, Y.; Igarashi, T.; Goto, K.; Tahara, H.; Uchimoto, S.; Omichi,

[115464]K.; Kanno, H.; Hayasaka, K.; Murase, T.: Heterogeneous mutations

[115465]in the glycogen-debranching enzyme gene are responsible for glycogen

[115466]storage disease type IIIa in Japan. Hum. Genet. 106: 108–115, 2000.

[115467]20340.

[115468]Attard-Montalto, S.; Evans, N.; Sherwood, R. A.: Carotenaemia

[115469]with low vitamin A levels and low retinol-binding protein. J. Inherit.

[115470]Metab. Dis. 15: 929–930, 1992.

[115471]20341.

[115472]Chainani, M.; Sampsell, B.; Elliott, R. W.: Localization of the

[115473]gene for plasma retinol binding protein to the distal half of mouse

[115474]chromosome 19. Genomics 9: 376–379, 1991.

[115475]20342.

[115476]Matsuo, T.: Familial retinol-binding-protein deficiency?
(Letter) Lancet II:

[115477]910 only, 1987.

[115478]20343.

[115479]Matsuo, T.; Matsuo, N.; Shiraga, F.; Koide, N.: Familial
retinol-binding-protein

[115480]deficiency. (Letter) Lancet II: 402–403, 1987.

[115481]20344.

[115482]Morgan, F. F.; Canfield, R. E.; Goodman, D. S.: The partial
structure

[115483]of human plasma prealbumin and retinol-binding protein.
Biochim.

[115484]Biophys. Acta 236: 798–801, 1971.

[115485]20345.

[115486]Rask, L.; Anundi, H.; Fohlman, J.; Peterson, P. A.: The
complete

[115487]amino acid sequence of human serum retinol-binding
protein. Upsala

[115488]J. Med. Sci. 92: 115–146, 1987.

[115489]20346.

[115490]Rask, L.; Vahlquist, A.; Peterson, P. A.: Studies on two physiological

[115491]forms of the human retinol-binding protein differing in vitamin A

[115492]and arginine content. J. Biol. Chem. 246: 6638–6646, 1971.

[115493]20347.

[115494]Rocchi, M.; Covone, A.; Romeo, G.; Faraonio, R.; Colantuoni, V.

[115495]: Regional mapping of RBP4 to 10q23–q24 and RBP1 to 3q21–q22 in man. Somat.

[115496]Cell Molec. Genet. 15: 185–190, 1989.

[115497]20348.

[115498]Seeliger, M. W.; Biesalski, H. K.; Wissinger, B.; Gollnick, H.;

[115499]Gielen, S.; Frank, J.; Beck, S.; Zrenner, E.: Phenotype in retinol

[115500]deficiency due to a hereditary defect in retinol binding protein synthesis. Invest.

[115501]Ophthal. Vis. Sci. 40: 3–11, 1999.

[115502]20349.

[115503]Thuluvath, P. J.: Familial retinol-binding-protein deficiency?

[115504](Letter) Lancet II: 910 only, 1987.

[115505]20350.

[115506]Carson, N. L.; Simpson, N. E.: A physical map of 13 markers on

[115507]chromosome 10 from dosage studies on abnormal cell lines. (Abstract) Cytogenet.

[115508]Cell Genet. 51: 974-975, 1989.

[115509]20351.

[115510]Chin, K. S.; Mathew, C. G. P.; Fong, S. L.; Bridges, C. D. B.;

[115511]Ponder, B. A. J.: Styl RFLP recognised by a human IRBP cDNA localised

[115512]to chromosome 10. Nucleic Acids Res. 16: 1645 only, 1988.

[115513]20352.

[115514]Danciger, M.; Kozak, C. A.; Nickerson, J.; Redmond, T. M.; Farber,

[115515]D. B.: Localization of the gene for interphotoreceptor retinoid-binding

[115516]protein to mouse chromosome 14 near Np-1. Genomics
8: 727-731, 1990.

[115517]20353.

[115518]Farrer, L. A.; Castiglione, C. M.; Kidd, J. R.; Myers, S.; Car-
son,

[115519]N.; Simpson, N. E.; Kidd, K. K.: A linkage group of five
DNA markers

[115520]on human chromosome 10. Genomics 3: 72-77, 1988.

[115521]20354.

[115522]Fong, S.-L.; Liou, G. I.; Landers, R. A.; Alvarez, R. A.; Gon-
zalez-Fernandez,

[115523]F.; Glazebrook, P. A.; Lam, D. M. K.; Bridges, C. D. B.:
Characterization,

[115524]localization, and biosynthesis of an interstitial retinol-
binding

[115525]glycoprotein in the human eye. J. Neurochem. 42:
1667-1676, 1984.

[115526]20355.

[115527]Liou, G. I.; Fong, S.-L.; Beattie, W. G.; Cook, R. G.; Leone,
J.;

[115528]Landers, R. A.; Alvarez, R. A.; Wang, C.; Li, Y.; Bridges, C.
D. B.

[115529] Bovine interstitial retinol-binding protein
(IRBP)—isolation and
[115530] sequence analysis of cDNA clones, characterization and in
vitro translation
[115531] of mRNA. Vision Res. 26: 1645–1653, 1986.
[115532] 20356.

[115533] Liou, G. I.; Fong, S.-L.; Gosden, J.; vanTuinen, P.; Ledbet-
ter,
[115534] D. H.; Christie, S.; Rout, D.; Bhattacharya, S.; Cook, R. G.;
Li,
[115535] Y.; Wang, C.; Bridges, C. D. B.: Human interstitial retinol-
binding
[115536] protein (IRBP): cloning, partial sequence, and chromoso-
mal localization. Somat.
[115537] Cell Molec. Genet. 13: 315–323, 1987.
[115538] 20357.

[115539] Liou, G. I.; Li, Y.; Wang, C.; Fong, S.-L.; Bhattacharya, F. S.;
[115540] Bridges, C. D. B.: Bgl II RFLP recognized by a human IRBP
cDNA localized
[115541] to chromosome 10. Nucleic Acids Res. 15: 3196 only,
1987.

[115542]20358.

[115543]McGuire, W.; Hill, A. V. S.; Allsopp, C. E. M.; Greenwood, B.

[115544]M.; Kwiatkowski, D.: Variation in the TNF- α promoter
region associated

[115545]with susceptibility to cerebral malaria. Nature 371:
508-511, 1994.

[115546]20359.

[115547]Bruford, E. A.; Riise, R.; Teague, P. W.; Porter, K.; Thom-
son,

[115548]K. L.; Moore, A. T.; Jay, M.; Warburg, M.; Schinzel, A.;
Tommerup,

[115549]N.; Tornqvist, K.; Rosenberg, T.; Patton, M.; Mansfield, D.
C.; Wright,

[115550]A. F.: Linkage mapping in 29 Bardet-Biedl syndrome fami-
lies confirms

[115551]loci in chromosomal regions 11q13, 15q22.3-q23, and
16q21. Genomics 41:

[115552]93-99, 1997.

[115553]20360.

[115554]Benbrook, D.; Pfahl, M.: A novel thyroid hormone receptor
encoded

[115555]by a cDNA clone from a human testis library. Science 238:
788-791,

[115556]1987.

[115557]20361.

[115558]Bernal, J.; Refetoff, S.; DeGroot, L. J.: Abnormalities of tri-
iodothyronine

[115559]binding to lymphocyte and fibroblast nuclei from a patient
with peripheral

[115560]tissue resistance to thyroid hormone action. J. Clin. En-
docr. Metab. 47:

[115561]1266-1272, 1978.

[115562]20362.

[115563]Dayton, A. I.; Selden, J. R.; Laws, G.; Dorney, D. J.; Finan,
J.;

[115564]Tripputi, P.; Emanuel, B. S.; Rovera, G.; Nowell, P. C.;

Croce, C.

[115565]M.: A human c-erbA oncogene homologue is closely
proximal to the

[115566]chromosome 17 breakpoint in acute promyelocytic
leukemia. Proc. Nat.

[115567]Acad. Sci. 81: 4495-4499, 1984.

[115568]20363.

[115569]Debuire, B.; Henry, C.; Benaissa, M.; Biserte, G.; Claverie, J.

[115570]M.; Saule, S.; Martin, P.; Stehelin, D.: Sequencing the erbA
gene

[115571]of avian erythroblastosis virus reveals a new type of onco-
gene. Science 224:

[115572]1456–1459, 1984.

[115573]20364.

[115574]Ferro, M. T.; San Roman, C.: Constitutional t(15;17). Can-
cer

[115575]Genet. Cytogenet. 4: 89–91, 1981.

[115576]20365.

[115577]Gullberg, H.; Rudling, M.; Forrest, D.; Angelin, B.;
Vennstrom,

[115578]B.: Thyroid hormone receptor beta-deficient mice show
complete loss

[115579]of the normal cholesterol 7- α -hydroxylase (CYP7A)
response to

[115580]thyroid hormone but display enhanced resistance to di-
etary cholesterol. Molec.

[115581]Endocr. 14: 1739–1749, 2000.

[115582]20366.

[115583]Ichikawa, K.; Hughes, I. A.; Horwitz, A. L.; DeGroot, L. J.:

Characterization

[115584]of nuclear thyroid hormone receptors of cultured skin fibroblasts

[115585]from patients with resistance to thyroid hormone.

Metabolism 36:

[115586]392–399, 1987.

[115587]20367.

[115588]Iskaros, J.; Pickard, M.; Evans, I.; Sinha, A.; Hardiman, P.;

Ekins,

[115589]R.: Thyroid hormone receptor gene expression in first trimester human

[115590]fetal brain. J. Clin. Endocr. Metab. 85: 2620–2623, 2000.

[115591]20368.

[115592]Jansson, M.; Philipson, L.; Vennstrom, B.: Isolation and

characterization

[115593]of multiple human genes homologous to the oncogenes of avian erythroblastosis

[115594]virus. EMBO J. 2: 561–565, 1983.

[115595]20369.

[115596]Jhanwar, S. C.; Chaganti, R. S. K.; Croce, C. M.: Germ-line chromosomal

[115597]localization of human c-erb-A oncogene. Somat. Cell Molec. Genet. 11:

[115598]99–102, 1985.

[115599]20370.

[115600]Kaneshige, M.; Suzuki, H.; Kaneshige, K.; Cheng, J.; Wimbrow,

[115601]H.; Barlow, C.; Willingham, M. C.; Cheng, S.: A targeted dominant

[115602]negative mutation of the thyroid hormone alpha-1 receptor causes increased

[115603]mortality, infertility, and dwarfism in mice. Proc. Nat. Acad. Sci. 98:

[115604]15095–15100, 2001.

[115605]20371.

[115606]Laudet, V.; Begue, A.; Henry-Duthoit, C.; Joubel, A.; Martin,

[115607]P.; Stehelin, D.; Saule, S.: Genomic organization of the human thyroid

[115608]hormone receptor alpha (c-erbA-1) gene. Nucleic Acids Res. 19: 1105–1112,

[115609]1991.

[115610]20372.

[115611]Le Beau, M. M.; Westbrook, C. A.; Diaz, M. O.; Rowley, J. D.;

[115612]Oren, M.: Translocation of the p53 gene in t(15;17) in acute promyelocytic

[115613]leukaemia. Nature 316: 826–828, 1985.

[115614]20373.

[115615]Mathieu–Mahul, D.; Xu, D. Q.; Saule, S.; Lidereau, R.; Galibert,

[115616]F.; Berger, R.; Mauchauffe, M.; Larsen, C. J.: An EcoRI restriction

[115617]fragment length polymorphism (RFLP) in the human c-erbA locus. Hum.

[115618]Genet. 71: 41–44, 1985.

[115619]20374.

[115620]McCabe, C. J.; Gittoes, N. J.; Sheppard, M. C.; Franklyn, J. A.

[115621]: Thyroid receptor alpha–1 and alpha–2 mutations in non-functioning

[115622]pituitary tumors. J. Clin. Endocr. Metab. 84: 649–653,

1999.

[115623]20375.

[115624]Menezes-Ferreira, M. M.; Eil, C.; Wortsman, J.; Weintraub, B.

[115625]D.: Decreased nuclear uptake of

[125-I]triiodo-L-thyronine in fibroblasts

[115626]from patients with peripheral thyroid hormone resistance.
J. Clin.

[115627]Endocr. Metab. 59: 1081-1087, 1984.

[115628]20376.

[115629]Mitelman, F.; Manolov, G.; Manolova, Y.; Billstrom, R.;
Heim,

[115630]S.; Kristoffersson, U.; Mandahl, N.; Ferro, M. T.; San Roman, C.:

[115631]High resolution chromosome analysis of constitutional
and acquired

[115632]t(15;17) maps c-erbA to subband 17q11.2. Cancer Genet.
Cytogenet. 22:

[115633]95-98, 1986.

[115634]20377.

[115635]Miyajima, N.; Horiuchi, R.; Shibuya, Y.; Fukushige, S.; Mat-

subara,

[115636]K.; Toyoshima, K.; Yamamoto, T.: Two erbA homologs encoding proteins

[115637]with different T(3) binding capacities are transcribed from opposite

[115638]DNA strands of the same genetic locus. Cell 57: 31-39, 1989.

[115639]20378.

[115640]Nagaya, T.; Nomura, Y.; Fujieda, M.; Seo, H.: Heterodimerization

[115641]preferences of thyroid hormone receptor alpha isoforms. Biochem.

[115642]Biophys. Res. Commun. 226: 426-430, 1996.

[115643]20379.

[115644]Nakai, A.; Seino, S.; Sakurai, A.; Szilak, I.; Bell, G. I.; DeGroot,

[115645]L. J.: Characterization of a thyroid hormone receptor expressed in

[115646]human kidney and other tissues. Proc. Nat. Acad. Sci. 85: 2781-2785,

[115647]1988.

[115648]20380.

- [115649]Ng, L.; Rusch, A.; Amma, L. L.; Nordstrom, K.; Erway, L. C.; Vennstrom, B.; Forrest, D.: Suppression of the deafness and thyroid dysfunction in Thr β -null mice by an independent mutation in the Thra thyroid hormone receptor gene. Hum. Molec. Genet. 10: 2701–2708, 2001.
- [115652]20381.
- [115654]Puzianowska-Kuznicka, M.; Krystyniak, A.; Madej, A.; Cheng, S.-Y.; Nauman, J.: Functionally impaired TR mutants are present in thyroid papillary cancer. J. Clin. Endocr. Metab. 87: 1120–1128, 2002.
- [115657]20382.
- [115658]Refetoff, S.; DeGroot, L. J.; Benard, B.; DeWind, L. T.: Studies of a sibship with apparent hereditary resistance to the intracellular action of thyroid hormone. Metabolism 21: 723–756, 1972.
- [115661]20383.

[115662]Rider, S. H.; Bailey, C. J.; Voss, R.; Sheer, D.; Hiorns, L. R.;

[115663]Solomon, E.: RFLP for the human erb-A1 gene. *Nucleic
Acids Res.* 15:

[115664]863 only, 1987.

[115665]20384.

[115666]Sakurai, A.; Nakai, A.; DeGroot, L. J.: Expression of three
forms

[115667]of thyroid hormone receptor in human tissues. *Molec. En-
docr.* 3:

[115668]392-399, 1989.

[115669]20385.

[115670]Wolf, F. W.; Marks, R. M.; Sarma, V.; Byers, M. G.; Katz, R.
W.;

[115671]Shows, T. B.; Dixit, V. M.: Characterization of a novel tu-
mor necrosis

[115672]factor-alpha-induced endothelial primary response gene.
J. Biol.

[115673]*Chem.* 267: 1317-1326, 1992.

[115674]20386.

[115675]Pandey, A.; Lindberg, R. A.; Dixit, V. M.: Receptor orphans
find

[115676]a family. Curr. Biol. 5: 986–989, 1995.

[115677]20387.

[115678]Aaltonen, L.–M.; Chen, R. W.; Roth, S.; Makitie, A. A.;
Rihkanen,

[115679]H.; Vaheri, A.; Aaltonen, L. A.: Role of TP53 P72R poly-
morphism in

[115680]human papillomavirus associated premalignant laryngeal
neoplasm. J.

[115681]Med. Genet. 38: 327 only, 2001.

[115682]20388.

[115683]Aas, T.; Borresen, A.–L.; Geisler, S.; Smith–Sorenson, B.;
Johnsen,

[115684]H.; Varhaug, J. E.; Akslen, L. A.; Lonning, P. E.: Specific P53
mutations

[115685]are associated with de novo resistance to doxorubicin in
breast cancer

[115686]patients. Nature Med. 2: 811–814, 1996.

[115687]20389.

[115688]Aguilar, F.; Harris, C. C.; Sun, T.; Hollstein, M.; Cerutti, P.

[115689]: Geographic variation of p53 mutational profile in non-
malignant human

[115690]liver. Science 264: 1317–1319, 1994.

[115691]20390.

[115692]Aguilar, F.; Hussain, S. P.; Cerutti, P.: Aflatoxin B1 induces

[115693]the transversion of G-to-T in codon 249 of the p53 tumor
suppressor

[115694]gene in human hepatocytes. Proc. Nat. Acad. Sci. 90:
8586–8590,

[115695]1993.

[115696]20391.

[115697]Ara, S.; Lee, P. S. Y.; Hansen, M. F.; Saya, H.: Codon 72
polymorphism

[115698]of the TP53 gene. Nucleic Acids Res. 18: 4961, 1990.

[115699]20392.

[115700]Arai, M.; Shimizu, S.; Imai, Y.; Nakatsuru, Y.; Oda, H.;
Oohara,

[115701]T.; Ishikawa, T.: Mutations of the Ki-ras, p53 and APC
genes in adenocarcinomas

[115702]of the human small intestine. Int. J. Cancer 70: 390–395,
1997.

[115703]20393.

[115704] Baker, S. J.; Fearon, E. R.; Nigro, J. M.; Hamilton, S. R.;
Preisinger,
[115705] A. C.; Jessup, J. M.; vanTuinen, P.; Ledbetter, D. H.; Barker,
D.
[115706] F.; Nakamura, Y.; White, R.; Vogelstein, B.: Chromosome
17 deletions
[115707] and p53 gene mutations in colorectal carcinomas. Science
244: 217–221,
[115708] 1989.
[115709] 20394.
[115710] Benchimol, S.; Lamb, P.; Crawford, L. V.; Sheer, D.; Shows,
T.
[115711] B.; Bruns, G. A. P.; Peacock, J.: Transformation associated
p53 protein
[115712] is encoded by a gene on human chromosome 17. Somat.
Cell Molec. Genet. 11:
[115713] 505–509, 1985.
[115714] 20395.
[115715] Bernal, J. A.; Luna, R.; Espina, A.; Lazaro, I.; Ramos–
Morales,
[115716] F.; Romero, F.; Arias, C.; Silva, A.; Tortolero, M.; Pintor–
Toro,

[115717]J. A.: Human securin interacts with p53 and modulates
p53-mediated
[115718]transcriptional activity and apoptosis. Nature Genet. 32:
306–311,
[115719]2002.
[115720]20396.
[115721]Beroud, C.; Verdier, F.; Soussi, T.: p53 gene mutation:
software
[115722]and database. Nucleic Acids Res. 24: 147–150, 1996.
[115723]20397.
[115724]Borresen, A.-L.; Hovig, E.; Smith-Sorensen, B.; Malkin, D.;
Lystad,
[115725]S.; Andersen, T. I.; Nesland, J. M.; Isselbacher, K. J.; Friend,
S.
[115726]H.: Constant denaturant gel electrophoresis as a rapid
screening
[115727]technique for p53 mutations. Proc. Nat. Acad. Sci. 88:
8405–8409,
[115728]1991.
[115729]20398.
[115730]Borresen, A. L.; Andersen, T. I.; Garber, J.; Barbier-Piriaux,

[115731]N.; Thorlacius, S.; Eyfjord, J.; Ottestad, L.; Smith-Sorensen, B.;

[115732]Hovig, E.; Malkin, D.; Friend, S. H.: Screening for germ line TP53

[115733]mutations in breast cancer patients. Cancer Res. 52: 3234-3236,

[115734]1992.

[115735]20399.

[115736]Brash, D. E.; Rudolph, J. A.; Simon, J. A.; Lin, A.; McKenna,

[115737]G. J.; Baden, H. P.; Halperin, A. J.; Ponten, J.: A role for sunlight

[115738]in skin cancer: UV-induced p53 mutations in squamous cell carcinoma. Proc.

[115739]Nat. Acad. Sci. 88: 10124-10128, 1991.

[115740]20400.

[115741]Brennan, J. A.; Boyle, J. O.; Koch, W. M.; Goodman, S. N.; Hruban,

[115742]R. H.; Eby, Y. J.; Couch, M. J.; Forastiere, A. A.; Sidransky, D.

[115743]: Association between cigarette smoking and mutation of the p53 gene

[115744]in squamous-cell carcinoma of the head and neck. New
Eng. J. Med. 332:

[115745]712–717, 1995.

[115746]20401.

[115747]Bressac, B.; Kew, M.; Wands, J.; Ozturk, M.: Selective G to T

[115748]mutations of p53 gene in hepatocellular carcinoma from
southern Africa. Nature 350:

[115749]429–431, 1991.

[115750]20402.

[115751]Brodsky, M. H.; Nordstrom, W.; Tsang, G.; Kwan, E.; Rubin,
G.

[115752]M.; Abrams, J. M.: Drosophila p53 binds a damage re-
sponse element

[115753]at the reaper locus. Cell 101: 103–113, 2000.

[115754]20403.

[115755]Buetow, K. H.; Sheffield, V. C.; Zhu, M.; Zhou, T.; Shen, F.;

[115756]Hino, O.; Smith, M.; McMahon, B. J.; Lanier, A. P.; London,
W. T.;

[115757]Redeker, A. G.; Govindarajan, S.: Low frequency of p53
mutations

[115758]observed in a diverse collection of primary hepatocellular carcinomas. Proc.

[115759]Nat. Acad. Sci. 89: 9622–9626, 1992.

[115760]20404.

[115761]Bunz, F.; Dutriaux, A.; Lengauer, C.; Waldman, T.; Zhou, S.; Brown,

[115762]J. P.; Sedivy, J. M.; Kinzler, K. W.; Vogelstein, B.: Requirement

[115763]for p53 and p21 to sustain G2 arrest after DNA damage. Science 282:

[115764]1497–1501, 1998.

[115765]20405.

[115766]Caelles, C.; Helmberg, A.; Karin, M.: p53–dependent apoptosis

[115767]in the absence of transcriptional activation of p53–target genes. Nature 370:

[115768]220–223, 1994.

[115769]20406.

[115770]Colucci–Guyon, E.; Portier, M.–M.; Dunia, I.; Paulin, D.; Pournin,

[115771]S.; Babinet, C.: Mice lacking vimentin develop and repro–

duce without

[115772]an obvious phenotype. Cell 79: 679–694, 1994.

[115773]20407.

[115774]Ferrari, S.; Battini, R.; Kaczmarek, L.; Rittling, S.; Calabretta,

[115775]B.; de Riel, J. K.; Philiponis, V.; Weil, J.-F.; Baserga, R.:
Coding

[115776]sequence and growth regulation of the human vimentin
gene. Molec.

[115777]Cell. Biol. 6: 3614–3620, 1986.

[115778]20408.

[115779]Gieser, L.; Swaroop, A.: Expressed sequence tags and
chromosomal

[115780]localization of cDNA clones from a subtracted retinal pig-
ment epithelium

[115781]library. Genomics 13: 873–876, 1992.

[115782]20409.

[115783]Lilienbaum, A.; Legagneux, V.; Portier, M.-M.; Dellagi, K.;
Paulin,

[115784]D.: Vimentin gene: expression in human lymphocytes and
in Burkitt's

[115785]lymphoma cells. EMBO J. 5: 2809–2814, 1986.

[115786]20410.

[115787]Marcus, E. M.; Smith, B. A.; Telenius, H.; Landsvater, R. M.;
Buys,

[115788]C. H. C. M.; Ferrari, S.; Ponder, B. A. J.; Mathew, C. G. P.:
Bcll

[115789]RFLP for the human vimentin gene. Nucleic Acids Res. 16:
9068 only,

[115790]1988.

[115791]20411.

[115792]Mathew, C. G.; Wakeling, W.; Jones, E.; Easton, D.; Fisher,
R.;

[115793]Strong, C.; Smith, B.; Chin, K.; Little, P.; Nakamura, Y.;
Shows,

[115794]T. B.; Jones, C.; Goodfellow, P. J.; Povey, S.; Ponder, B. A.
J.:

[115795]Regional localization of polymorphic markers on chromo-
some 10 by physical

[115796]and genetic mapping. Ann. Hum. Genet. 54: 121–129,
1990.

[115797]20412.

- [115798]Perreau, J.; Lilienbaum, A.; Vasseur, M.; Paulin, D.: Nu-
cleotide
- [115799]sequence of the human vimentin gene and regulation of
its transcription
- [115800]in tissues and cultured cells. Gene 62: 7-16, 1988.
- [115801]20413.
- [115802]Gimona, M.; Small, J. V.; Moeremans, M.; Van Damme, J.;
Puype,
- [115803]M.; Vandekerckhove, J.: Porcine vinculin and metavinculin
differ
- [115804]by a 68-residue insert located close to the carboxy-ter-
minal part
- [115805]of the molecule. EMBO J. 7: 2329-2334, 1988.
- [115806]20414.
- [115807]Mulligan, L. M.; Gardner, E.; Telenius, H.; Ponder, B. A. J.:
- [115808]Complementary physical and genetic techniques map the
vinculin (VCL)
- [115809]gene on chromosome 10q. Genomics 13: 1347-1349,
1992.
- [115810]20415.
- [115811]Strasser, P.; Gimona, M.; Herzog, M.; Geiger, B.; Small, J.

V.

[115812] Variable and constant regions in the C-terminus of vinculin and

[115813]metavinculin: cloning and expression of fragments in *E. coli*. FEBS

[115814]Lett. 317: 189–194, 1993.

[115815]20416.

[115816]Koteliansky, V. E.; Ogryzko, E. P.; Zhidkova, N. I.; Weller, P.

[115817]A.; Critchley, D. R.; Vancompernelle, K.; Vandekerckhove, J.; Strasser,

[115818]P.; Way, M.; Gimona, M.; Small, J. V.: An additional exon in the

[115819]human vinculin gene specifically encodes meta-vinculin-specific difference

[115820]peptide: cross-species comparison reveals variable and conserved motifs

[115821]in the meta-vinculin insert. *Europ. J. Biochem.* 204: 767–772, 1992.

[115822]20417.

[115823]Turner, C. E.; Burridge, K.: Detection of metavinculin in human

[115824]platelets using a modified talin overlay assay. *Europ. J.*

Cell Biol. 49:

[115825]202–206, 1989.

[115826]20418.

[115827]Weller, P. A.; Ogryzko, E. P.; Corben, E. B.; Zhidkova, N. I.;

[115828]Patel, B.; Price, G. J.; Spurr, N. K.; Koteliensky, V. E.;

Critchley,

[115829]D. R.: Complete sequence of human vinculin and assignment of the

[115830]gene to chromosome 10. Proc. Nat. Acad. Sci. 87: 5667–5671, 1990.

[115831]20419.

[115832]Katsanis, N.; Ansley, S. J.; Badano, J. L.; Eichers, E. R.;

Lewis,

[115833]R. A.; Hoskins, B. E.; Scambler, P. J.; Davidson, W. S.;

Beales, P.

[115834]L.; Lupski, J. R.: Triallelic inheritance in Bardet–Biedl syndrome,

[115835]a mendelian recessive disorder. Science 293: 2256–2259, 2001.

[115836]20420.

[115837]Katsanis, N.; Beales, P. L.; Woods, M. O.; Lewis, R. A.;

Green,

[115838]J. S.; Parfrey, P. S.; Ansley, S. J.; Davidson, W. S.; Lupski, J.

[115839]R.: Mutations in MKKS cause obesity, retinal dystrophy and renal

[115840]malformations associated with Bardet-Biedl syndrome.

Nature Genet. 26:

[115841]67-70, 2000.

[115842]20421.

[115843]Kwitek-Black, A. E.; Carmi, R.; Duyk, G. M.; Buetow, K. H.; Elbedour,

[115844]K.; Parvari, R.; Yandava, C. N.; Stone, E. M.; Sheffield, V. C.:

[115845]Linkage of Bardet-Biedl syndrome to chromosome 16q and evidence for

[115846]non-allelic genetic heterogeneity. Nature Genet. 5: 392-396, 1993.

[115847]20422.

[115848]Ishii, H.; Majerus, P. W.: Thrombomodulin is present in human

[115849]plasma and urine. J. Clin. Invest. 76: 2178-2181, 1985.

[115850]20423.

[115851]Gabreels, B. A. T. F.; Swaab, D. F.; de Kleijn, D. P. V.; Seidah,

[115852]N. G.; Van de Loo, J.-W.; Van de Ven, W. J. M.; Martens, G. J. M.;

[115853]van Leeuwen, F. W.: Attenuation of the polypeptide 7B2, prohormone

[115854]convertase PC2, and vasopressin in the hypothalamus of some Prader-Willi

[115855]patients: indications for a processing defect. J. Clin. Endocr. Metab. 83:

[115856]591-599, 1998.

[115857]20424.

[115858]Bell, K.; Hopper, K. E.; McKenzie, H. A.; Murphy, W. H.; Shaw,

[115859]D. C.: A comparison of bovine alpha-lactalbumin A and B of Droughtmaster. Biochim.

[115860]Biophys. Acta 214: 437-444, 1970.

[115861]20425.

[115862]Bell, K.; McKenzie, H. A.; Murphy, W. H.; Shaw, D. C.: Beta-lactoglobulin

[115863](Droughtmaster): a unique protein variant. Biochim. Biophys. Acta 214:

[115864]427-436, 1970.

[115865]20426.

[115866]Julkunen, M.; Seppala, M.; Janne, O. A.: Complete amino acid sequence

[115867]of human placental protein 14: a progesterone-regulated uterine protein

[115868]homologous to beta-lactoglobulins. Proc. Nat. Acad. Sci. 85: 8845-8849,

[115869]1988.

[115870]20427.

[115871]Morris, H. R.; Dell, A.; Easton, R. L.; Panico, M.; Koistinen,

[115872]H.; Koistinen, R.; Oehninger, S.; Patankar, M. S.; Seppala, M.; Clark,

[115873]G. F.: Gender-specific glycosylation of human glycodefin affects

[115874]its contraceptive activity. J. Biol. Chem. 271: 32159-32167, 1996.

[115875]20428.

[115876]Song, M.; Ramaswamy, S.; Ramachandran, S.; Flowers, L. C.; Horowitz,

[115877]J. R.; Rock, J. A.; Parthasarathy, S.: Angiogenic role for gly-

codelin

[115878]in tumorigenesis. Proc. Nat. Acad. Sci. 98: 9265–9270,
2001.

[115879]20429.

[115880]Van Cong, N.; Vaisse, C.; Gross, M.–S.; Slim, R.; Milgrom,
E.;

[115881]Bernheim, A.: The human placental protein 14 (PP14) gene
is localized

[115882]on chromosome 9q34. Hum. Genet. 86: 515–518, 1991.

[115883]20430.

[115884]Lee, F. S.; Fox, E. A.; Zhou, H.–M.; Strydom, D. J.; Vallee, B.

[115885]L.: Primary structure of human placental ribonuclease in-
hibitor. Biochemistry 27:

[115886]8545–8553, 1988.

[115887]20431.

[115888]Weremowicz, S.; Fox, E. A.; Morton, C. C.; Vallee, B. L.: The

[115889]placental ribonuclease inhibitor (RNH) gene is located on
chromosome

[115890]subband 11p15.5. Genomics 8: 717–721, 1990.

[115891]20432.

[115892]Zneimer, S. M.; Crawford, D.; Schneider, N. R.; Beutler, B.:
Mapping

[115893]of the human ribonuclease inhibitor gene (RNH) to chromosome 11p15

[115894]by in situ hybridization. Genomics 8: 175–178, 1990.

[115895]20433.

[115896]Coughlin, P.; Nicholl, J.; Sun, J.; Salem, H.; Bird, P.; Sutherland,

[115897]G. R.: Chromosomal mapping of the human proteinase inhibitor 6 (PI6)

[115898]gene to 6p25 by fluorescence in situ hybridization. Genomics 26:

[115899]431–433, 1995.

[115900]20434.

[115901]Coughlin, P.; Sun, J.; Cerruti, L.; Salem, H. H.; Bird, P.:
Cloning

[115902]and molecular characterization of a human intracellular serine proteinase

[115903]inhibitor. Proc. Nat. Acad. Sci. 90: 9417–9421, 1993.

[115904]20435.

[115905]Zhang, K.; Westberg, J. A.; Paetau, A.; von Boguslawsky,

K.; Lindsberg,

[115906]P.; Erlander, M.; Guo, H.; Su, J.; Olsen, H. S.; Andersson, L.
C.

[115907]: High expression of stanniocalcin in differentiated brain
neurons. Am.

[115908]J. Path. 153: 439–445, 1998.

[115909]20436.

[115910]Hamilton, S. E.; Hurley, J. B.: A phosphodiesterase in-
hibitor

[115911]specific to a subset of bovine retinal cones. J. Biol. Chem.
265:

[115912]11259–11264, 1990.

[115913]20437.

[115914]Shimizu–Matsumoto, A.; Itoh, K.; Inazawa, J.; Nishida, K.;
Matsumoto,

[115915]Y.; Kinoshita, S.; Matsubara, K.; Okubo, K.: Isolation and
chromosomal

[115916]localization of the human cone cGMP phosphodiesterase
gamma cDNA (PDE6H). Genomics 32:

[115917]121–124, 1996.

[115918]20438.

[115919]Isomura, M.; Okui, K.; Fujiwara, T.; Shin, S.; Nakamura, Y.:

Cloning

[115920]and mapping of a novel human cDNA homologous to

DROER, the enhancer

[115921]of the *Drosophila melanogaster* rudimentary gene. *Ge-*

nomics 32: 125-127,

[115922]1996.

[115923]20439.

[115924]Chen, J.; Engle, S. J.; Seilhamer, J. J.; Tischfield, J. A.:

Cloning

[115925]and recombinant expression of a novel human low molec-

ular weight Ca^{2+} -dependent

[115926]phospholipase A2. *J. Biol. Chem.* 269: 2365-2368, 1994.

[115927]20440.

[115928]Tischfield, J. A.; Xia, Y.-R.; Shih, D. M.; Klisak, I.; Chen, J.;

[115929]Engle, S. J.; Siakotos, A. N.; Winstead, M. V.; Seilhamer, J.

J.;

[115930]Allamand, V.; Gyapay, G.; Lusi, A. J.: Low molecular

weight, calcium-dependent

[115931]phospholipase A(2) genes are linked and map to homolo-

gous chromosome

[115932]regions in mouse and human. Genomics 32: 328–333, 1996.

[115933]20441.

[115934]Hayashi, M.; Fujimoto, S.; Takano, H.; Ushiki, T.; Abe, K.; Ishikura,

[115935]H.; Yoshida, M. C.; Kirchhoff, C.; Ishibashi, T.; Kasahara, M.: Characterization

[115936]of a human glycoprotein with a potential role in sperm-egg fusion:

[115937]cDNA cloning, immunohistochemical localization, and chromosomal assignment

[115938]of the gene (AEGL1). Genomics 32: 367–374, 1996.

[115939]20442.

[115940]Muzio, M.; Bosisio, D.; Polentarutti, N.; D'amico, G.; Stoppacciaro,

[115941]A.; Mancinelli, R.; van't Veer, C.; Penton-Rol, G.; Ruco, L. P.; Allavena,

[115942]P.; Mantovani, A.: Differential expression and regulation of Toll-like

[115943]receptors (TLR) in human leukocytes: selective expression of TLR3

[115944]in dendritic cells. J. Immun. 164: 5998–6004, 2000.

[115945]20443.

[115946]Rock, F. L.; Hardiman, G.; Timans, J. C.; Kastelein, R. A.;
Bazan,

[115947]J. F.: A family of human receptors structurally related to
Drosophila

[115948]Toll. Proc. Nat. Acad. Sci. 95: 588–593, 1998.

[115949]20444.

[115950]Perrimon, N.; Mahowald, A. P.: Multiple functions of seg-
ment polarity

[115951]genes in Drosophila. Dev. Biol. 119: 587–600, 1987.

[115952]20445.

[115953]Pizzuti, A.; Novelli, G.; Mari, A.; Ratti, A.; Colosimo, A.;
Amati,

[115954]F.; Penso, D.; Sangiuolo, F.; Calabrese, G.; Palka, G.; Silani,
V.;

[115955]Gennarelli, M.; Mingarelli, R.; Scarlato, G.; Scambler, P.;
Dallapiccola,

[115956]B.: Human homologue sequences to the Drosophila di-
shevelled segment-polarity

[115957]gene are deleted in the DiGeorge syndrome. Am. J. Hum.
Genet. 58:

[115958]722–729, 1996.

[115959]20446.

[115960]Onyango, P.; Lubyova, B.; Gardellin, P.; Kurzbauer, R.;
Weith,

[115961]A.: Molecular cloning and expression analysis of five novel
genes

[115962]in chromosome 1p36. Genomics 50: 187–198, 1998.

[115963]20447.

[115964]Barbier, M.; Attoub, S.; Calvez, R.; Laffargue, M.; Jarry, A.;

[115965]Mareel, M.; Altruda, F.; Gespach, C.; Wu, D.; Lu, B.; Hirsch,
E.;

[115966]Wymann, M. P.: Weakening link to colorectal cancer? Na-
ture 413:

[115967]796 only, 2001.

[115968]20448.

[115969]Crackower, M. A.; Oudit, G. Y.; Kozieradzki, I.; Sarao, R.;
Sun,

[115970]H.; Sasaki, T.; Hirsch, E.; Suzuki, A.; Shioi, T.; Irie-Sasaki,
J.;

[115971]Sah, R.; Cheng, H.–Y. M.; and 13 others: Regulation of
myocardial

[115972]contractility and cell size by distinct PI3K–PTEN signaling pathways. Cell 110:

[115973]737–749, 2002.

[115974]20449.

[115975]Hirsch, E.; Katanaev, V. L.; Garlanda, C.; Azzolino, O.;
Pirola,

[115976]L.; Silengo, L.; Sozzani, S.; Mantovani, A.; Altruda, F.;
Wymann,

[115977]M. P.: Central role for G protein–coupled phosphoinositide
3–kinase

[115978]gamma in inflammation. Science 287: 1049–1053, 2000.

[115979]20450.

[115980]Jiang, K.; Zhong, B.; Gilvary, D. L.; Corliss, B. C. Hong–
Geller,

[115981]E.; Wei, S; Djeu, J. Y.: Pivotal role of phosphoinositide–3
kinase

[115982]in regulation of cytotoxicity in natural killer cells. Nature
Immun. 1:

[115983]419–425, 2000.

[115984]20451.

[115985]Kikuta, Y.; Kato, M.; Yamashita, Y.; Miyauchi, Y.; Tanaka,

K.;

[115986]Kamada, N.; Kusunose, M.: Human leukotriene B₄ omega-hydroxylase

[115987](CYP4F3) gene: molecular cloning and chromosomal localization. DNA

[115988]Cell Biol. 17: 221-230, 1998.

[115989]20452.

[115990]Kikuta, Y.; Kusunose, E.; Endo, K.; Yamamoto, S.; Sogawa, K.; Fujii-Kuriyama,

[115991]Y.; Kusunose, M.: A novel form of cytochrome P-450 family 4 in human

[115992]polymorphonuclear leukocytes: cDNA cloning and expression of leukotriene

[115993]B₄ omega-hydroxylase. J. Biol. Chem. 268: 9376-9380, 1993.

[115994]20453.

[115995]Hamra, F. K.; Forte, L. R.; Eber, S. L.; Pidhorodeckyj, N. V.;

[115996]Krause, W. J.; Freeman, R. H.; Chin, D. T.; Tompkins, J. A.; Fok,

[115997]K. F.; Smith, C. E.; Duffin, K. L.; Siegel, N. R.; Currie, M. G.:

[115998]Uroguanylin: structure and activity of a second endoge-

nous peptide

[115999] that stimulates intestinal guanylate cyclase. Proc. Nat.

Acad. Sci. 90:

[116000] 10464–10468, 1993.

[116001] 20454.

[116002] Kita, T.; Smith, C. E.; Fok, K. F.; Duffin, K. L.; Moore, W. M.;

[116003] Karabatsos, P. J.; Kachur, J. F.; Hamra, F. K.; Pid-
horodeckyj, N.

[116004] V.; Forte, L. R.; Currie, M. G.: Characterization of human
uroguanylin:

[116005] a member of the guanylin peptide family. Am. J. Physiol.
266: F342–F348,

[116006] 1994.

[116007] 20455.

[116008] Miyazato, M.; Nakazato, M.; Matsukura, S.; Kangawa, K.;

Matsuo,

[116009] H.: Genomic structure and chromosomal localization of
human uroguanylin. Genomics 43:

[116010] 359–365, 1997.

[116011] 20456.

[116012] Whitaker, T. L.; Steinbrecher, K. A.; Copeland, N. G.;

Gilbert,

[116013]D. J.; Jenkins, N. A.; Cohen, M. B.: The uroguanylin gene
(Guca1b)

[116014]is linked to guanylin (Guca2) on mouse chromosome 4.

Genomics 45:

[116015]348–354, 1997.

[116016]20457.

[116017]Haft, C. R.; de la Luz Sierra, M.; Barr, V. A.; Haft, D. H.;
Taylor,

[116018]S. I.: Identification of a family of sorting nexin molecules
and

[116019]characterization of their association with receptors. Molec.
Cell.

[116020]Biol. 18: 7278–7287, 1998.

[116021]20458.

[116022]Kurten, R. C.; Cadena, D. L.; Gill, G. N.: Enhanced degra-
dation

[116023]of EGF receptors by a sorting nexin, SNX1. Science 272:
1008–1010,

[116024]1996.

[116025]20459.

- [116026]Zhong, Q.; Lazar, C. S.; Tronchere, H.; Sato, T.; Meerloo, T.;
- [116027]Yeo, M.; Songyang, Z.; Emr, S. D.; Gill, G. N.: Endosomal localization
- [116028]and function of sorting nexin 1. *Proc. Nat. Acad. Sci.* 99: 6767–6772,
- [116029]2002.
- [116030]20460.
- [116031]Klingensmith, J.; Nusse, R.; Perrimon, N.: The *Drosophila* segment
- [116032]polarity gene *dishevelled* encodes a novel protein required for response
- [116033]to wingless signal. *Genes Dev.* 8: 118–130, 1994.
- [116034]20461.
- [116035]Holmes, S. E.; Riazzi, M. A.; Gong, W.; McDermid, H. E.; Sellinger,
- [116036]B. T.; Hua, A.; Chen, F.; Wang, Z.; Zhang, G.; Roe, B.; Gonzalez,
- [116037]J.; McDonald–McGinn, D. M.; Zackai, E.; Emanuel, B. S.; Budarf, M.
- [116038]L.: Disruption of the clathrin heavy chain–like gene (CLTCL) associated

[116039]with features of DGS/VCFS: a balanced (21;22)(p12;q11) translocation. Hum.

[116040]Molec. Genet. 6: 357–367, 1997.

[116041]20462.

[116042]Kedra, D.; Peyrard, M.; Fransson, I.; Collins, J. E.; Dunham, I.;

[116043]Roe, B. A.; Dumanski, J. P.: Characterization of a second human clathrin

[116044]heavy chain polypeptide gene (CLH-22) from chromosome 22q11. Hum.

[116045]Molec. Genet. 5: 625–631, 1996.

[116046]20463.

[116047]Long, K. R.; Trofatter, J. A.; Ramesh, V.; McCormick, M. K.; Buckler,

[116048]A. J.: Cloning and characterization of a novel human clathrin heavy

[116049]chain gene (CLTCL). Genomics 35: 466–472, 1996.

[116050]20464.

[116051]Lagenaur, C.; Kunemund, V.; Fischer, G.; Fushiki, S.; Schachner,

[116052]M.: Monoclonal M6 antibody interferes with neurite exten-

sion of cultured

[116053]neurons. J. Neurobiol. 23: 71–88, 1992.

[116054]20465.

[116055]Shimizu, F.; Watanabe, T. K.; Fujiwara, T.; Takahashi, E.;
Nakamura,

[116056]Y.; Maekawa, H.: Isolation and mapping of the human gly-
coprotein

[116057]M6 gene (GPM6A) to 4q33–to–q34. Cytogenet. Cell Genet.
74: 138–139,

[116058]1996.

[116059]20466.

[116060]Bui, T. D.; Beier, D. R.; Jonssen, M.; Smith, K.; Dorrington,
S.

[116061]M.; Kaklamanis, L.; Kearney, L.; Regan, R.; Sussman, D. J.;
Harris,

[116062]A. L.: cDNA cloning of a human dishevelled DVL–3 gene,
mapping to

[116063]3q27, and expression in human breast and colon carcino-
mas. Biochem.

[116064]Biophys. Res. Commun. 239: 510–516, 1997.

[116065]20467.

[116066]Scheffner, M.; Huibregtse, J. M.; Vierstra, R. D.; Howley, P.

[116067]M.: The HPV-16 E6 and E6-AP complex functions as a ubiquitin-protein

[116068]ligase in the ubiquitination of p53. Cell 75: 495-505, 1993.

[116069]20468.

[116070]Cheng, Y.-S. E.; Patterson, C. E.; Staeheli, P.: Interferon-induced

[116071]guanylate-binding proteins lack an N(T)KXD consensus motif and bind

[116072]GMP in addition to GDP and GTP. Molec. Cell. Biol. 11: 4717-4725,

[116073]1991.

[116074]20469.

[116075]Kumar, S.; Li, Q.; Dua, A.; Ying, Y.-K.; Bagchi, M. K.; Bagchi,

[116076]I. C.: Messenger ribonucleic acid encoding interferon-inducible guanylate

[116077]binding protein 1 is induced in human endometrium within the putative

[116078]window of implantation. J. Clin. Endocr. Metab. 86: 2420-2427, 2001.

[116079]20470.

[116080]Prochazka, M.; Staeheli, P.; Holmes, R. S.; Haller, O.: Interferon-induced

[116081]guanylate-binding proteins: mapping of the murine Gbp-1 locus to chromosome

[116082]3. Virology 145: 273-279, 1985.

[116083]20471.

[116084]Strehlow, I.; Lohmann-Matthes, M. L.; Decker, T.: The interferon-inducible

[116085]GBP1 gene: structure and mapping to human chromosome 1. Gene 144:

[116086]295-299, 1994.

[116087]20472.

[116088]Scheffner, M.; Werness, B. A.; Huibregtse, J. M.; Levine, A. J.;

[116089]Howley, P. M.: The E6 oncoprotein encoded by human papillomavirus

[116090]types 16 and 18 promotes the degradation of p53. Cell 63: 1129-1136,

[116091]1990.

[116092]20473.

[116093]Tsai, T.-F.; Raas-Rothschild, A.; Ben-Neriah, Z.; Beaudet, A.

[116094]L.: Prenatal diagnosis and carrier detection for a point mutation

[116095]in UBE3A causing Angelman syndrome. (Letter) Am. J. Hum. Genet. 63:

[116096]1561-1563, 1998.

[116097]20474.

[116098]Yu, T. H.; Hoffman, A. R.: Imprinting of the Angelman syndrome

[116099]gene, UBE3A, is restricted to brain. (Letter) Nature Genet. 17:

[116100]12-13, 1997.

[116101]20475.

[116102]Avraham, K. B.; Prezioso, V. R.; Chen, W. S.; Lai, E.; Sladek,

[116103]F. M.; Zhong, W.; Darnell, J. E., Jr.; Jenkins, N. A.; Copeland, N.

[116104]G.: Murine chromosomal location of four hepatocyte-enriched transcription

[116105]factors: HNF-3-alpha, HNF3-beta, HNF-3-gamma, and HFN-4. Genomics 13:

[116106]264–268, 1992.

[116107]20476.

[116108]Chartier, F. L.; Bossu, J.-P.; Laudet, V.; Fruchart, J.-C.:

Cloning

[116109]and sequencing of cDNAs encoding the human hepatocyte
nuclear factor

[116110]4 indicate the presence of two isoforms in human liver.

Gene 147:

[116111]269–272, 1994.

[116112]20477.

[116113]Eeckhoutte, J.; Formstecher, P.; Laine, B.: Maturity-onset
diabetes

[116114]of the young type 1 (MODY1)–associated mutations R154X
and E276Q in

[116115]hepatocyte nuclear factor 4–alpha (HNF4–alpha) gene im-
pair recruitment

[116116]of p300, a key transcriptional coactivator. Molec. Endocr.
15: 1200–1210,

[116117]2001.

[116118]20478.

[116119]Furuta, H.; Iwasaki, N.; Oda, N.; Hinokio, Y.; Horikawa, Y.;

Yamagata,

[116120]K.; Yano, N.; Sugahiro, J.; Ogata, M.; Ohgawara, H.; Omori,
Y.; Iwamoto,

[116121]Y.; Bell, G. I.: Organization and partial sequence of the
hepatocyte

[116122]nuclear factor-4-alpha/MODY1 gene and identification of
a missense

[116123]mutation, R127W, in a Japanese family with MODY. Dia-
betes 46: 1652-1657,

[116124]1997.

[116125]20479.

[116126]Hani, E. H.; Suaud, L.; Boutin, P.; Chevre, J.-C.; Durand, E.;

[116127]Philippi, A.; Demenais, F.; Vionnet, N.; Furuta, H.; Velho,
G.; Bell,

[116128]G. I.; Laine, B.; Froguel, P.: A missense mutation in hepa-
tocyte

[116129]nuclear factor-4-alpha, resulting in a reduced transacti-
vation activity,

[116130]in human late-onset non-insulin-dependent diabetes
mellitus. J. Clin.

[116131]Invest. 101: 521-526, 1998.

[116132]20480.

- [116133] Lausen, J.; Thomas, H.; Lemm, I.; Bulman, M.;
Borgschulze, M.;
- [116134] Lingott, A.; Hattersley, A. T.; Ryffel, G. U.: Naturally occurring
ring
- [116135] mutations in the human HNF4- α gene impair the
function of the
- [116136] transcription factor to a varying degree. Nucleic Acids Res.
28:
- [116137] 430-437, 2000.
- [116138] 20481.
- [116139] Li, J.; Ning, G.; Duncan, S. A.: Mammalian hepatocyte differentiation
ferentiation
- [116140] requires the transcription factor HNF-4- α . Genes Dev.
14: 464-474,
- [116141] 2000.
- [116142] 20482.
- [116143] Stoffel, M.; Duncan, S. A.: The maturity-onset diabetes of
the
- [116144] young (MODY1) transcription factor HNF4- α regulates
expression
- [116145] of genes required for glucose transport and metabolism.
Proc. Nat.

- [116146]Acad. Sci. 94: 13209–13214, 1997.
- [116147]20483.
- [116148]Yamagata, K.; Furuta, H.; Oda, N.; Kaisaki, P. J.; Menzel, S.;
- [116149]Cox, N. J.; Fajans, S. S.; Signorini, S.; Stoffel, M.; Bell, G. I.
- [116150]; Mutations in the hepatocyte nuclear factor-4- α gene
in maturity-onset
- [116151]diabetes of the young (MODY1). Nature 384: 458–460,
1996.
- [116152]20484.
- [116153]Zouali, H.; Hani, E. H.; Philippi, A.; Vionnet, N.; Beckmann,
- [116154]J. S.; Demenais, F.; Froguel, P.: A susceptibility locus for
early-onset
- [116155]non-insulin dependent (type 2) diabetes mellitus maps to
chromosome
- [116156]20q, proximal to the phosphoenolpyruvate carboxykinase
gene. Hum.
- [116157]Molec. Genet. 6: 1401–1408, 1997.
- [116158]20485.
- [116159]Frolova, L.; Le Goff, X.; Rasmussen, H. H.; Cheperegin, S.;
- Druegon,

[116160]G.; Kress, M.; Arman, I.; Haenni, A.-L.; Celis, J. E.; Philippe, M.;

[116161]Justesen, J.; Kisselev, L.: A highly conserved eukaryotic protein

[116162]family possessing properties of polypeptide chain release factor. Nature 372:

[116163]701–703, 1994.

[116164]20486.

[116165]Guenet, L.; Henry, C.; Toutain, B.; Dubourg, C.; Le Gall, J. Y.;

[116166]David, V.; Le Treut, A.: Eukaryotic translation termination factor

[116167]gene (ETF1/eRF1) maps at D5S500 in a commonly deleted region of chromosome

[116168]5q31 in malignant myeloid diseases. Cytogenet. Cell Genet. 88: 82–86,

[116169]2000.

[116170]20487.

[116171]Hansen, L. L.; Jakobsen, C. G.; Justesen, J.: Assignment of the

[116172]human translation termination factor 1 (ETF1) to 5q31.1 and of the

[116173]proximal marker D5S1995 by radiation hybrid mapping.

Cytogenet. Cell

[116174]Genet. 87: 256–257, 1999.

[116175]20488.

[116176]Le Goff, X.; Philippe, M.; Jean-Jean, O.: Overexpression of
human

[116177]release factor 1 alone has an antisuppressor effect in hu-
man cells. Molec.

[116178]Cell Biol. 17: 3164–3172, 1997.

[116179]20489.

[116180]Song, H.; Mugnier, P.; Das, A. K.; Webb, H. M.; Evans, D.
R.; Tuite,

[116181]M. F.; Hemmings, B. A.; Barford, D.: The crystal structure
of human

[116182]eukaryotic release factor eRF1--mechanism of stop codon recognition

[116183]and peptidyl-tRNA hydrolysis. Cell 100: 311-321, 2000.

[116184]20490.

[116185]Liu, Q.; Fischer, U.; Wang, F.; Dreyfuss, G.: The spinal muscular

[116186]atrophy disease gene product, SMN, and its associated protein SIP1

[116187]are in a complex with spliceosomal snRNP proteins. Cell 90: 1013-1021,

[116188]1997.

[116189]20491.

[116190]Meister, G.; Buhler, D.; Laggerbauer, B.; Zobawa, M.; Lottspeich,

[116191]F.; Fischer, U.: Characterization of a nuclear 20S complex containing

[116192]the survival of motor neurons (SMN) protein and a specific subset

[116193]of spliceosomal Sm proteins. Hum. Molec. Genet. 9: 1977-1986, 2000.

[116194]20492.

- [116195] Aramburu, J.; Garcia-Cozar, F.; Raghavan, A.; Okamura, H.; Rao, A.; Hogan, P. G.: Selective inhibition of NFAT activation by a peptide spanning the calcineurin targeting site of NFAT. Molec. Cell 1: 627-637, 1998.
- [116198] 20493.
- [116199] Castigli, E.; Pahwa, R.; Good, R. A.; Geha, R. S.; Chatila, T. A.: Molecular basis of a multiple lymphokine deficiency in a patient with severe combined immunodeficiency. Proc. Nat. Acad. Sci. 90: 4728-4732, 1993.
- [116203] 20494.
- [116205] Chatila, T.; Castigli, E.; Pahwa, R.; Pahwa, S.; Chirmule, N.; Oyaizu, N.; Good, R. A.; Geha, R. S.: Primary combined immunodeficiency resulting from defective transcription of multiple T-cell lymphokine genes. Proc. Nat. Acad. Sci. 87: 10033-10037, 1990.

[116209]20495.

[116210]Horsley, V.; Pavlath, G. K.: NFAT: ubiquitous regulator of cell

[116211]differentiation and adaptation. J. Cell Biol. 156: 771–774, 2002.

[116212]20496.

[116213]Jauliac, S.; Lopez-Rodriguez, C.; Shaw, L. M.; Brown, L. F.; Rao,

[116214]A.; Toker, A.: The role of NFAT transcription factors in integrin-mediated

[116215]carcinoma invasion. Nature Cell Biol. 4: 540–544, 2002.

[116216]20497.

[116217]Li, X.; Ho, S. N.; Luna, J.; Giacalone, J.; Thomas, D. J.; Timmerman,

[116218]L. A.; Crabtree, G. R.; Francke, U.: Cloning and chromosomal localization

[116219]of the human and murine genes for the T-cell transcription factors

[116220]NFATc and NFATp. Cytogenet. Cell Genet. 68: 185–191, 1995.

[116221]20498.

[116222]Northrop, J. P.; Ho, S. N.; Chen, L.; Thomas, D. J.; Timmerman,
[116223]L. A.; Nolan, G. P.; Admon, A.; Crabtree, G. R.: NF-AT
components
[116224]define a family of transcription factors targeted in T-cell
activation. Nature 369:
[116225]497–502, 1994.
[116226]20499.
[116227]Okamura, H.; Aramburu, J.; Garcia-Rodriguez, C.; Viola, J.
P. B.;
[116228]Raghavan, A.; Tashirani, M.; Zhang, X.; Qin, J.; Hogan, P.
G.; Rao,
[116229]A.: Concerted dephosphorylation of the transcription fac-
tor NFAT1
[116230]induces a conformational switch that regulates transcrip-
tional activity. Molec.
[116231]Cell 6: 539–550, 2000.
[116232]20500.
[116233]Pahwa, R.; Chatila, T.; Pahwa, S.; Paradise, C.; Day, N. K.;
Geha,
[116234]R.; Schwartz, S. A.; Slade, H.; Oyaizu, N.; Good, R. A.: Re-
combinant

[116235]interleukin 2 therapy in severe combined immunodeficiency disease. Proc.

[116236]Nat. Acad. Sci. 86: 5069–5073, 1989.

[116237]20501.

[116238]Park, J.; Takeuchi, A.; Sharma, S.: Characterization of a new

[116239]isoform of the NFAT (nuclear factor of activated T cells) gene family

[116240]member NFATc. J. Biol. Chem. 271: 20914–20921, 1996.

[116241]20502.

[116242]Peng, S. L.; Gerth, A. J.; Ranger, A. M.; Glimcher, L. H.: NFATc1

[116243]and NFATc2 together control both T and B cell activation and differentiation. Immunity 14:

[116244]13–20, 2001.

[116245]20503.

[116246]Aoki, K.; Suzuki, K.; Sugano, T.; Tasaka, T.; Nakahara, K.; Kuge,

[116247]O.; Omori, A.; Kasai, M.: A novel gene, 'Translin,' encodes a recombination

[116248]hotspot binding protein associated with chromosomal

translocations. Nature

[116249]Genet. 10: 167–174, 1995.

[116250]20504.

[116251]Badge, R. M.; Yardley, J.; Jeffreys, A. J.; Armour, J. A. L.:

[116252]Crossover breakpoint mapping identifies a subtelomeric
hotspot for

[116253]male meiotic recombination. Hum. Molec. Genet. 9:
1239–1244, 2000.

[116254]20505.

[116255]Hosaka, T.; Kanoe, H.; Nakayama, T.; Murakami, H.; Ya-
mamoto, H.;

[116256]Nakamata, T.; Tsuboyama, T.; Oka, M.; Kasai, M.; Sasaki,
M. S.; Nakamura,

[116257]T.; Toguchida, J.: Translin binds to the sequences adjacent
to the

[116258]breakpoints of the TLS and CHOP genes in liposarcomas
with translocation

[116259]t(12;16). Oncogene 19: 5821–5825, 2000.

[116260]20506.

[116261]Kasai, M.; Aoki, K.; Matsuo, Y.; Minowada, J.; Maziarz, R.
T.;

[116262]Strominger, J. L.: Recombination hotspot associated factors specifically

[116263]recognize novel target sequences at the site of interchromosomal rearrangements

[116264]in T-ALL patients with t(8;14)(q24;q11) and (t(1;14)(p32;q11). *Int.*

[116265]*Immun.* 6: 1017–1025, 1994.

[116266]20507.

[116267]Cirillo, L. A.; Lin, F. R.; Cuesta, I.; Friedman, D.; Jarnik, M.;

[116268]Zaret, K. S.: Opening of compacted chromatin by early developmental

[116269]transcription factors HNF3(FoxA) and GATA-4. *Molec. Cell* 9: 279–289,

[116270]2002.

[116271]20508.

[116272]Laitinen, M. P. E.; Anttonen, M.; Ketola, I.; Wilson, D. B.; Ritvos,

[116273]O.; Butzow, R.; Heikinheimo, M.: Transcription factors GATA-4 and

[116274]GATA-6 and a GATA family cofactor, FOG-2, are expressed in human ovary

[116275]and sex cord-derived ovarian tumors. J. Clin. Endocr.

Metab. 85:

[116276]3476–3483, 2000.

[116277]20509.

[116278]Freyd, G.; Kim, S. K.; Horvitz, H. R.: Novel cysteine-rich
motif

[116279]and homeodomain in the product of the *Caenorhabditis*
elegans cell

[116280]lineage gene *lin-11*. Nature 344: 876–879, 1990.

[116281]20510.

[116282]Mbikay, M.; Tadros, H.; Seidah, N. G.; Simpson, E. M.:
Linkage

[116283]mapping of the gene for the LIM-homeoprotein LIM3
(locus *Lhx3*) to

[116284]mouse chromosome 2. Mammalian Genome 6: 818–819,
1995.

[116285]20511.

[116286]Sharma, K.; Leonard, A. E.; Lettieri, K.; Pfaff, S. L.: Genetic

[116287]and epigenetic mechanisms contribute to motor neuron
pathfinding. Nature 406:

[116288]515–519, 2000.

[116289]20512.

[116290]Sharma, K.; Sheng, H. Z.; Lettieri, K.; Li, H.; Karavanov, A.;

[116291]Potter, S.; Westphal, H.; Pfaff, S. L.: LIM homeodomain factors Lhx3

[116292]and Lhx4 assign subtype identities for motor neurons.

Cell 95: 817–828,

[116293]1998.

[116294]20513.

[116295]Sheng, H. Z.; Zhadanov, A. B.; Mosinger, B., Jr.; Fujii, T.; Bertuzzi,

[116296]S.; Grinberg, A.; Lee, E. J.; Huang, S.–P.; Mahon, K. A.; Westphal,

[116297]H.: Specification of pituitary cell lineages by the LIM homeobox

[116298]gene Lhx3. Science 272: 1004–1007, 1996.

[116299]20514.

[116300]Sloop, K. W.; Meier, B. C.; Bridwell, J. L.; Parker, G. E.; Schiller,

[116301]A. M.; Rhodes, S. J.: Differential activation of pituitary hormone

[116302]genes by human Lhx3 isoforms with distinct DNA binding

properties. Molec.

[116303]Endocr. 13: 2212–2225, 1999.

[116304]20515.

[116305]Sloop, K. W.; Showalter, A. D.; Von Kap–Herr, C.; Pettenati, M.

[116306]J.; Rhodes, S. J.: Analysis of the human LHX3 neuroendocrine transcription

[116307]factor gene and mapping to the subtelomeric region of chromosome 9. Gene 245:

[116308]237–243, 2000.

[116309]20516.

[116310]Sloop, K. W.; Walvoord, E. C.; Showalter, A. D.; Pescovitz, O.

[116311]H.; Rhodes, S. J.: Molecular analysis of LHX3 and PROP–1 in pituitary

[116312]hormone deficiency patients with posterior pituitary ectopia. J.

[116313]Clin. Endocr. Metab. 85: 2701–2708, 2000.

[116314]20517.

[116315]Brenneman, M. A.; Wagener, B. M.; Miller, C. A.; Allen, C.; Nickoloff,

[116316]J. A.: XRCC3 controls the fidelity of homologous recombination: roles
 [116317]for XRCC3 in late stages of recombination. Molec. Cell 10:
 387–395,
 [116318]2002.
 [116319]20518.
 [116320]Liu, N.; Lamerdin, J. E.; Tebbs, R. S.; Schild, D.; Tucker, J.
 [116321]D.; Shen, M. R.; Brookman, K. W.; Siciliano, M. J.; Walter, C.
 A.;
 [116322]Fan, W.; Narayana, L. S.; Zhou, Z.-Q.; Adamson, A. W.;
 Sorensen, K.
 [116323]J.; Chen, D. J.; Jones, N. J.; Thompson, L. H.: XRCC2 and
 XRCC3,
 [116324]new human Rad51-family members, promote chromo-
 some stability and protect
 [116325]against DNA cross-links and other damages. Molec. Cell
 1: 783–793,
 [116326]1998.
 [116327]20519.
 [116328]Tebbs, R. S.; Zhao, Y.; Tucker, J. D.; Scheerer, J. B.; Sicil-
 iano,

[116329]M. J.; Hwang, M.; Liu, N.; Legerski, R. J.; Thompson, L. H.:

Correction

[116330]of chromosomal instability and sensitivity to diverse mutagens by

[116331]a cloned cDNA of the XRCC3 DNA repair gene. Proc. Nat. Acad. Sci. 92:

[116332]6354–6358, 1995.

[116333]20520.

[116334]Winsey, S. L.; Haldar, N. A.; Marsh, H. P.; Bunce, M.; Marshall,

[116335]S. E.; Harris, A. L.; Wojnarowska, F.; Welsh, K. I.: A variant within

[116336]the DNA repair gene XRCC3 is associated with the development of melanoma

[116337]skin cancer. Cancer Res. 60: 5612–5616, 2000.

[116338]20521.

[116339]Hayashida, S.; Yamasaki, K.; Asada, Y.; Soeda, E.; Niikawa, N.;

[116340]Kishino, T.: Construction of a physical and transcript map flanking

[116341]the imprinted MEST/PEG1 region at 7q32. Genomics 66: 221–225, 2000.

[116342]20522.

[116343]de Wind, N.; Dekker, M.; Claij, N.; Jansen, L.; van Klink, Y.;

[116344]Radman, M.; Riggins, G.; van der Valk, M.; van't Wout, K.;
te Riele,

[116345]H.: HNPCC-like cancer predisposition in mice through si-
multaneous

[116346]loss of Msh3 and Msh6 mismatch-repair protein func-
tions. Nature Genet. 23:

[116347]359-362, 1999.

[116348]20523.

[116349]Chen, Y. H.; Hansen, L.; Chen, M. X.; Bjorbaek, C.; Vester-
gaard,

[116350]H.; Hansen, T.; Cohen, P. T. W.; Pedersen, O.: Sequence of
the human

[116351]glycogen-associated regulatory subunit of type 1 protein
phosphatase

[116352]and analysis of its coding region and mRNA level in mus-
cle from patients

[116353]with NIDDM. Diabetes 43: 1234-1241, 1994.

[116354]20524.

[116355]Braun-Dullaes, R. C.; Mann, M. J.; Ziegler, A.; von der

Leyen,

[116356]H. E.; Dzaou, V. J.: A novel role for the cyclin-dependent kinase

[116357]inhibitor p27(Kip1) in angiotensin II-stimulated vascular smooth muscle

[116358]cell hypertrophy. J. Clin. Invest. 104: 815–823, 1999.

[116359]20525.

[116360]Carrano, A. C.; Eytan, E.; Hershko, A.; Pagano, M.: SKP2 is required

[116361]for ubiquitin-mediated degradation of the CDK inhibitor p27. Nature

[116362]Cell Biol. 1: 193–199, 1999.

[116363]20526.

[116364]Di Cristofano, A.; De Acetis, M.; Koff, A.; Cordon-Cardo, C.; Pandolfi,

[116365]P. P.: Pten and p27(KIP1) cooperate in prostate cancer tumor suppression

[116366]in the mouse. Nature Genet. 27: 222–224, 2001.

[116367]20527.

[116368]Fero, M. L.; Rivkin, M.; Tasch, M.; Porter, P.; Carow, C. E.; Firpo,

[116369]E.; Polyak, K.; Tsai, L.-H.; Broudy, V.; Perlmutter, R. M.;
Kaushansky,
[116370]K.; Roberts, J. M.: A syndrome of multiorgan hyperplasia
with features
[116371]of gigantism, tumorigenesis, and female sterility in
p27(Kip1)-deficient
[116372]mice. Cell 85: 733-744, 1996.
[116373]20528.

[116374]Gowen, M.; Lazner, F.; Dodds, R.; Kapadia, R.; Feild, J.;
Tavaria,
[116375]M.; Bertoncello, I.; Drake, F.; Zavarselk, S.; Tellis, I.; Hert-
zog,
[116376]P.; Debouck, C.; Kola, I.: Cathepsin K knockout mice de-
velop osteopetrosis
[116377]due to a deficit in matrix degradation but not demineral-
ization. J.
[116378]Bone Miner. Res. 14: 1654-1663, 1999.
[116379]20529.

[116380]Buyse, I. M.; Shao, G.; Huang, S.: The retinoblastoma pro-
tein
[116381]binds to RIZ, a zinc-finger protein that shares an epitope
with the

[116382]adenovirus E1A protein. Proc. Nat. Acad. Sci. 92:
4467-4471, 1995.

[116383]20530.

[116384]Buyse, I. M.; Takahashi, E.; Huang, S.: Physical mapping of
the

[116385]retinoblastoma interacting zinc finger gene RIZ to D1S228
on chromosome

[116386]1p36. Genomics 34: 119-121, 1996.

[116387]20531.

[116388]Chadwick, R. B.; Jiang, G.-L.; Bennington, G. A.; Yuan, B.;
Johnson,

[116389]C. K.; Stevens, M. W.; Niemann, T. H.; Peltomaki, P.;
Huang, S.; de

[116390]la Chapelle, A.: Candidate tumor suppressor RIZ is fre-
quently involved

[116391]in colorectal carcinogenesis. Proc. Nat. Acad. Sci. 97:
2662-2667,

[116392]2000.

[116393]20532.

[116394]Mock, B. A.; Coleman, M. P.; Huang, S.: Riz maps to distal
chromosome

[116395]4 near genes involved in tumorigenesis and nerve degeneration. Mammalian

[116396]Genome 7: 637 only, 1996.

[116397]20533.

[116398]Poetsch, M.; Dittberner, T.; Woenckhaus, C.: Frameshift mutations

[116399]of RIZ, but no point mutations in RIZ1 exons in malignant melanomas

[116400]with deletions in 1p36. Oncogene 21: 3038–3042, 2002.

[116401]20534.

[116402]Malek, N. P.; Sundberg, H.; McGrew, S.; Nakayama, K.; Kyriakidis,

[116403]T. R.; Roberts, J. M.: A mouse knock-in model exposes sequential

[116404]proteolytic pathways that regulate p27(Kip1) in G1 and S phase. Nature 413:

[116405]323–327, 2001.

[116406]20535.

[116407]Martin, E.; Cacheux, V.; Cave, H.; Lapierre, J. M.; Le\Paslier,

[116408]D.; Grandchamp, B.: Localization of the CDKN4/p27(Kip1)

gene to human

[116409]chromosome 12p12.3. Hum. Genet. 96: 668–670, 1995.

[116410]20536.

[116411]Mitsuhashi, T.; Aoki, Y.; Eksioglu, Y. Z.; Takahashi, T.;

Bhide,

[116412]P. G.; Reeves, S. A.; Caviness, V. S., Jr.: Overexpression of

p27(Kip1)

[116413]lengthens the G1 phase in a mouse model that targets in-

ducible gene

[116414]expression to central nervous system progenitor cells.

Proc. Nat.

[116415]Acad. Sci. 98: 6435–6440, 2001.

[116416]20537.

[116417]Polyak, K.; Lee, M.-H.; Erdjument-Bromage, H.; Koff, A.;

Roberts,

[116418]J. M.; Tempst, P.; Massague, J.: Cloning of p27(Kip1), a cy-

clin-dependent

[116419]kinase inhibitor and a potential mediator of extracellular

antimitogenic

[116420]signals. Cell 78: 59–66, 1994.

[116421]20538.

[116422]Saito, T.; Seki, N.; Hattori, A.; Hayashi, A.; Abe, M.; Araki,

[116423]R.; Fujimori, A.; Fukumura, R.; Kozuma, S.; Matsuda, Y.:

Structure,

[116424]expression profile, and chromosomal location of a mouse
gene homologous

[116425]to human DNA-PK(cs) interacting protein (KIP) gene.

Mammalian Genome 10:

[116426]315–317, 1999.

[116427]20539.

[116428]Sherr, C. J.; Roberts, J. M.: Inhibitors of mammalian G1 cy-
clin-dependent

[116429]kinases. Genes Dev. 9: 1149–1163, 1995.

[116430]20540.

[116431]Toyoshima, H.; Hunter, T.: p27, a novel inhibitor of G1 cy-
clin-Cdk

[116432]protein kinase activity, is related to p21. Cell 78: 67–74,
1994.

[116433]20541.

[116434]Khakh, B. S.; Zhou, X.; Sydes, J.; Galligan, J. J.; Lester, H.

[116435]A.: State-dependent cross-inhibition between transmit-
ter-gated cation

[116436]channels. Nature 406: 405–410, 2000.

[116437]20542.

[116438]Chen, J. D.; Evans, R. M.: A transcriptional co-repressor
that

[116439]interacts with nuclear hormone receptors. Nature 377:
454–457, 1995.

[116440]20543.

[116441]Fischle, W.; Dequiedt, F.; Hendzel, M. J.; Guenther, M. G.;
Lazar,

[116442]M. A.; Voelter, W.; Verdin, E.: Enzymatic activity associated
with

[116443]class II HDACs is dependent on a multiprotein complex
containing HDAC3

[116444]and SMRT/N-CoR. Molec. Cell 9: 45–57, 2002.

[116445]20544.

[116446]Horlein, A. J.; Naar, A. M.; Heinzel, T.; Torchia, J.; Gloss,
B.;

[116447]Kurokawa, R.; Ryan, A.; Kamel, Y.; Soderstrom, M.; Glass,
C. K.; Rosenfeld,

[116448]M. G.: Ligand-independent repression by the thyroid hor-
mone receptor

[116449]mediated by a nuclear receptor co-repressor. Nature 377:
397-403,

[116450]1995.

[116451]20545.

[116452]Hu, X.; Lazar, M. A.: The CoRNR motif controls the recruit-
ment

[116453]of corepressors by nuclear hormone receptors. Nature
402: 93-96,

[116454]1999.

[116455]20546.

[116456]Jiang, Q.; Galiegue-Zouitina, S.; Roumier, C.; Hildebrand,
M. P.;

[116457]Thomas, S.; Coignet, L. J.: Genomic organization and re-
fined mapping

[116458]of the human nuclear corepressor 2 (NCOR2)/ silencing
mediator of

[116459]retinoid and thyroid hormone receptor (SMRT) gene on
chromosome 12q24.3. Cytogenet.

[116460]Cell Genet. 92: 217-220, 2001.

[116461]20547.

[116462]Ordentlich, P.; Downes, M.; Xie, W.; Genin, A.; Spinner, N.

B.;

[116463]Evans, R. M.: Unique forms of human and mouse nuclear
receptor corepressor

[116464]SMRT. Proc. Nat. Acad. Sci. 96: 2639–2644, 1999.

[116465]20548.

[116466]Hansen, L.; Hansen, T.; Vestergaard, H.; BJORBAEK, C.; Ech-
wald,

[116467]S. M.; Clausen, J. O.; Chen, Y. H.; Chen, M. X.; Cohen, P. T.
W.;

[116468]Pedersen, O.: A widespread amino acid polymorphism at
codon 905 of

[116469]the glycogen-associated regulatory subunit of protein
phosphatase-1

[116470]is associated with insulin resistance and hypersecretion of
insulin. Hum.

[116471]Molec. Genet. 4: 1313–1320, 1995.

[116472]20549.

[116473]Hegele, R. A.; Harris, S. B.; Zinman, B.; Wang, J.; Cao, H.;
Hanley,

[116474]A. J. G.; Tsui, L.-C.; Scherer, S. W.: Variation in the
AU(AT)-rich

[116475]element within the 3-prime-untranslated region of

PPP1R3 is associated

[116476]with variation in plasma glucose in aboriginal Canadians.

J. Clin.

[116477]Endocr. Metab. 83: 3980–3983, 1998.

[116478]20550.

[116479]Savage, D. B.; Agostini, M.; Barroso, I.; Gurnell, M.; Luan,

J.;

[116480]Meirhaeghe, A.; Harding, A.-H.; Ihrke, G.; Rajanayagam,

O.; Soos,

[116481]M. A.; George, S.; Berger, D.; and 9 others: Digenic inheritance
tance

[116482]of severe insulin resistance in a human pedigree. Nature

Genet. 31:

[116483]379–384, 2002.

[116484]20551.

[116485]Tang, P. M.; Bondor, J. A.; Swiderek, K. M.; DePaoli-Roach,

A.

[116486]A.: Molecular cloning and expression of the regulatory
(RG1) subunit

[116487]of the glycogen-associated protein phosphatase. J. Biol.

Chem. 266:

[116488]15782–15789, 1991.

[116489]20552.

[116490]International Cystinuria Consortium: Functional analysis of

[116491]mutations in SLC7A9, and genotype-phenotype correlation in non-Type

[116492]I cystinuria. Hum. Molec. Genet. 10: 305-316, 2001.

[116493]20553.

[116494]International Cystinuria Consortium: Non-type I cystinuria caused

[116495]by mutations in SLC7A9, encoding a subunit (b(0,+)-AT) of rBAT. Nature

[116496]Genet. 23: 52-57, 1999.

[116497]20554.

[116498]Garcia, J. G.; Lazar, V.; Gilbert-McClain, L. I.; Gallagher, P.

[116499]J.; Verin, A. D.: Myosin light chain kinase in endothelium: molecular

[116500]cloning and regulation. Am. J. Resp. Cell Molec. Biol. 16: 489-494,

[116501]1997.

[116502]20555.

[116503]Giorgi, D.; Brand-Arpon, V.; Rouquier, S.: The functional

myosin

[116504]light chain kinase (MYLK) gene localizes with marker
D3S3552 on human

[116505]chromosome 3q21 in a greater than 5-Mb yeast artificial
chromosome

[116506]region and is not linked to olfactory receptor genes. Cyto-
genet.

[116507]Cell Genet. 92: 85-88, 2001.

[116508]20556.

[116509]Lazar, V.; Garcia, J. G. N.: A single human myosin light
chain

[116510]kinase gene (MLCK; MYLK) transcribes multiple nonmuscle
isoforms. Genomics 57:

[116511]256-267, 1999.

[116512]20557.

[116513]Potier, M.-C.; Chelot, E.; Pekarsky, Y.; Gardiner, K.;
Rossier,

[116514]J.; Turnell, W. G.: The human myosin light chain kinase
(MLCK) from

[116515]hippocampus: cloning, sequencing, expression, and local-
ization to

[116516]3cen-q21. Genomics 29: 562-570, 1995.

[116517]20558.

[116518]Walker, L. A.; MacDonald, J. A.; Liu, X.; Nakamoto, R. K.;
Haystead,

[116519]T. A. J.; Somlyo, A. V.; Somlyo, A. P.: Site-specific phos-
phorylation

[116520]and point mutations of telokin modulate its
Ca(2+)-desensitizing effect

[116521]in smooth muscle. J. Biol. Chem. 276: 24519-24524,
2001.

[116522]20559.

[116523]Watterson, D. M.; Schavocky, J. P.; Guo, L.; Weiss, C.;
Chlenski,

[116524]A.; Shirinsky, V. P.; Van Eldik, L. J.; Haiech, J.: Analysis of
the

[116525]kinase-related protein gene found at human chromosome
3q21 in a multi-gene

[116526]cluster: organization, expression, alternative splicing, and
polymorphic

[116527]marker. J. Cell. Biochem. 75: 481-491, 1999.

[116528]20560.

[116529]Druck, T.; Gu, Y.; Prabhala. G.; Cannizzaro, L. A.; Park,
S.-H.;

[116530] Huebner, K.; Keen, J. H.: Chromosome localization of human genes
[116531] for clathrin adaptor polypeptides AP2-beta and AP50 and
the clathrin-binding
[116532] protein, VCP. Genomics 30: 94-97, 1995.
[116533] 20561.
[116534] Liu, Q.; Feng, Y.; Forgac, M.: Activity and in vitro reassembly
of the coated vesicle (H⁺)-ATPase requires the 50-kDa
subunit of the
[116535] clathrin assembly complex AP-2. J. Biol. Chem. 269:
31592-31597,
[116537] 1994.
[116538] 20562.
[116539] Thuriereau, C.; Brosius, J.; Burne, C.; Jolles, P.; Keen, J. H.;
[116540] Mattaliano, R. J.; Chow, E. P.; Ramachandran, K. L.; Kirchhausen,
[116541] T.: Molecular cloning and complete amino acid sequence
of AP50, an
[116542] assembly protein associated with clathrin-coated vesicles.
DNA 7:

[116543]663–669, 1988.

[116544]20563.

[116545]Ponnambalam, S.; Robinson, M. S.; Jackson, A. P.; Peiperl, L.;

[116546]Parham, P.: Conservation and diversity in families of coated vesicle

[116547]adaptins. *J. Biol. Chem.* 265: 4814–4820, 1990.

[116548]20564.

[116549]Waelter, S.; Scherzinger, E.; Hasenbank, R.; Nordhoff, E.; Lurz,

[116550]R.; Goehler, H.; Gauss, C.; Sathasivam, K.; Bates, G. P.; Lehrach,

[116551]H.; Wanker, E. E.: The huntingtin interacting protein HIP1 is a clathrin

[116552]and alpha-adaptin-binding protein involved in receptor-mediated endocytosis. *Hum.*

[116553]*Molec. Genet.* 10: 1807–1817, 2001.

[116554]20565.

[116555]Kaneko-Ishino, T.; Kuroiwa, Y.; Miyoshi, N.; Kohda, T.; Suzuki,

[116556]R.; Yokoyama, M.; Viville, S.; Barton, S. C.; Ishino, F.;

Surani,

[116557]M. A.: Peg1/Mest imprinted gene on chromosome 6 identified by cDNA

[116558]subtraction hybridization. Nature Genet. 11: 52–59, 1995.

[116559]20566.

[116560]Kobayashi, S.; Kohda, T.; Miyoshi, N.; Kuroiwa, Y.; Aisaka, K.;

[116561]Tsutsumi, O.; Kaneko-Ishino, T.; Ishino, F.: Human PEG1/MEST, an

[116562]imprinted gene on chromosome 7. Hum. Molec. Genet. 6: 781–786, 1997.

[116563]20567.

[116564]Kosaki, K.; Kosaki, R.; Craigen, W. J.; Matsuo, N.: Isoform-specific

[116565]imprinting of the human PEG1/MEST gene. (Letter) Am. J. Hum. Genet. 66:

[116566]309–312, 2000.

[116567]20568.

[116568]Kotzot, D.; Schmitt, S.; Bernasconi, F.; Robinson, W. P.; Lurie,

[116569]J. W.; Ilyina, H.; Mehes, K.; Hamel, B. C. J.; Otten, B. J.;

Hergersberg,

[116570]M.; Werder, E.; Shoenle, E.; Schinzel, A.: Uniparental disomy 7 in

[116571]Silver–Russell syndrome and primordial growth retardation. Hum. Molec.

[116572]Genet. 4: 583–587, 1995.

[116573]20569.

[116574]Lefebvre, L.; Viville, S.; Barton, S. C.; Ishino, F.; Keverne,

[116575]E. B.; Surani, M. A.: Abnormal maternal behaviour and growth retardation

[116576]associated with loss of the imprinted gene Mest. Nature Genet. 20:

[116577]163–169, 1998.

[116578]20570.

[116579]Lefebvre, L.; Viville, S.; Barton, S. C.; Ishino, F.; Surani, M.

[116580]A.: Genomic structure and parent–of–origin–specific methylation of

[116581]Peg1. Hum. Molec. Genet. 6: 1907–1915, 1997.

[116582]20571.

[116583]Ayala–Madrigal, M. L.; Doerr, S.; Ramirez–Duenas, M. L.;

Hansmann,

[116584]J.: Assignment of karyopherin alpha 1 (KPNA1) to human chromosome

[116585]band 3q21 by in situ hybridization. Cytogenet. Cell Genet. 90: 58-59,

[116586]2000.

[116587]20572.

[116588]Conti, E.; Uy, M.; Leighton, L.; Blobel, G.; Kuriyan, J.: Crystallographic

[116589]analysis of the recognition of a nuclear localization signal by the

[116590]nuclear import factor karyopherin alpha. Cell 94: 193-204, 1998.

[116591]20573.

[116592]Cortes, P.; Ye, Z.-S.; Baltimore, D.: RAG-1 interacts with the

[116593]repeated amino acid motif of the human homologue of the yeast protein

[116594]SRP1. Proc. Nat. Acad. Sci. 91: 7633-7637, 1994.

[116595]20574.

[116596]Aplan, P. D.; Johnson, B. E.; Russell, E.; Chervinsky, D. S.;

Kirsch,

[116597]J. R.: Cloning and characterization of TCTA, a gene located
at the

[116598]site of a t(1;3) translocation. Cancer Res. 55: 1917–1921,
1995.

[116599]20575.

[116600]Ghetti, A.; Pinol-Roma, S.; Michael, W. M.; Morandi, C.;
Dreyfuss,

[116601]G.: HNRNP I, the polypyrimidine tract-binding protein:
distinct nuclear

[116602]localization and association with hnRNAs. Nucleic Acids
Res. 20:

[116603]3671–3678, 1992.

[116604]20576.

[116605]Raimondi, E.; Romanelli, M. G.; Moralli, D.; Gamberi, C.;
Russo,

[116606]M. P.; Morandi, C.: Assignment of the human gene encod-
ing heterogeneous

[116607]nuclear RNA ribonucleoprotein I (PTB) to chromosome
14q23–q24.1. Genomics 27:

[116608]553–555, 1995.

[116609]20577.

[116610]Romanelli, M. G.; Lorenzi, P.; Morandi, C.: Organization of the

[116611]human gene encoding heterogeneous nuclear ribonucleo-
protein type I

[116612](hnRNP I) and characterization of hnRNP I related pseudo-
gene. Gene 255:

[116613]267-272, 2000.

[116614]20578.

[116615]Hirota, H.; Chen, J.; Betz, U. A. K.; Rajewsky, K.; Gu, Y.;
Ross,

[116616]J., Jr.; Muller, W.; Chien, K. R.: Loss of a gp130 cardiac
muscle

[116617]cell survival pathway is a critical event in the onset of
heart failure

[116618]during biochemical stress. Cell 97: 189-198, 1999.

[116619]20579.

[116620]Kidd, V. J.; Nesbitt, J. E.; Fuller, G. M.: Chromosomal local-
ization

[116621]of the IL-6 receptor signal transducing subunit, gp130
(IL6ST). Somat.

[116622]Cell Molec. Genet. 18: 477-483, 1992.

[116623]20580.

[116624]Kishimoto, T.; Taga, T.; Akira, S.: Cytokine signal trans-
duction. Cell 76:

[116625]253–262, 1994.

[116626]20581.

[116627]Rodriguez, C.; Grosgeorge, J.; Nguyen, V. C.; Gaudray, P.;
Theillet,

[116628]C.: Human gp130 transducer chain gene (IL6ST) is local-
ized to chromosome

[116629]band 5q11 and possesses a pseudogene on chromosome
band 17p11. Cytogenet.

[116630]Cell Genet. 70: 64–67, 1995.

[116631]20582.

[116632]Haagerup, A.; Hertz, J. M.; Christensen, M. F.; Binderup,
H.; Kruse,

[116633]T. A.: Cathepsin K gene mutations and 1q21 haplotypes in
patients

[116634]with pycnodysostosis in an outbred population. Europ. J.
Hum. Genet. 8:

[116635]431–436, 2000.

[116636]20583.

- [116637]Ho, N.; Punturieri, A.; Wilkin, D.; Szabo, J.; Johnson, M.;
Whaley,
[116638]J.; Davis, J.; Clark, A.; Weiss, S.; Francomano, C.: Mutations
of
[116639]CTSK result in pycnodysostosis via a reduction in cathep-
sin K protein. J.
[116640]Bone Miner. Res. 14: 1649–1653, 1999.
[116641]20584.
[116642]Inaoka, T.; Bilbe, G.; Ishibashi, O.; Tezuka, K.; Kumegawa,
M.;
[116643]Kokubo, T.: Molecular cloning of human cDNA for cathep-
sin K: novel
[116644]cysteine proteinase predominantly expressed in bone.
Biochem. Biophys.
[116645]Res. Commun. 206: 89–96, 1995.
[116646]20585.
[116647]Johnson, M. R.; Polymeropoulos, M. H.; Vos, H. L.; Ortiz de
Luna,
[116648]R. I.; Francomano, C. A.: A nonsense mutation in the
cathepsin K
[116649]gene observed in a family with pycnodysostosis. Genome
Res. 6: 1050–1055,

[116650]1996.

[116651]20586.

[116652]Hoogenraad, C. C.; Koekkoek, B.; Akhmanova, A.; Krugers, H.; Dortland,

[116653]B.; Miedema, M.; van Alphen, A.; Kistler, W. M.; Jaegle, M.; Koutsourakis,

[116654]M.; Van Camp, N.; Verhoye, M.; van der Linden, A.; Kaverina, I.; Grosveld,

[116655]F.; De Zeeuw, C. I.; Galjart, N.: Targeted mutation of Cyln2 in the

[116656]Williams syndrome critical region links CLIP-115 haploinsufficiency

[116657]to neurodevelopmental abnormalities in mice. Nature Genet. 32: 116–127,

[116658]2002. Note: Erratum: Nature Genet. 32: 331 only, 2002.

[116659]20587.

[116660]Bannerman, R. M.; Edwards, J. A.; Kreimer-Birnbaum, M.; McFarland,

[116661]E.; Russell, E. S.: Hereditary microcytic anaemia in the mouse; studies

[116662]in iron distribution and metabolism. Brit. J. Haemat. 23: 235–245,

[116663]1972.

[116664]20588.

[116665]Silverstein, A. M.: Essential hypocomplementemia: report
of a

[116666]case. Blood 16: 1338–1341, 1960.

[116667]20589.

[116668]Hintner, H.; Wolff, K.: Generalized atrophic benign epider-
molysis

[116669]bullosa. Arch. Derm. 118: 375–384, 1982.

[116670]20590.

[116671]McGrath, J. A.; Gatalica, B.; Christiano, A. M.; Li, K.;
Owaribe,

[116672]K.; McMillan, J. R.; Eady, R. A. J.; Uitto, J.: Mutations in the

[116673]180-kD bullous pemphigoid antigen (BPAG2), a
hemidesmosomal transmembrane

[116674]collagen (COL17A1), in generalized atrophic benign epi-
dermolysis bullosa. Nature

[116675]Genet. 11: 83–86, 1995.

[116676]20591.

[116677]Vincent, J. B.; Herbrick, J.-A.; Gurling, H. M. D.; Bolton, P.

[116678]F.; Roberts, W.; Scherer, S. W.: Identification of a novel
gene on

[116679]chromosome 7q31 that is interrupted by a translocation
breakpoint

[116680]in an autistic individual. Am. J. Hum. Genet. 67: 510–514,
2000.

[116681]20592.

[116682]Vourc'h, P.; Bienvenu, T.; Beldjord, C.; Chelly, J.;
Barthelemy,

[116683]C.; Muh, J. P.; Andres, C.: No mutations in the coding re-
gion of

[116684]the Rett syndrome gene MECP2 in 59 autistic patients. Eu-
rop. J. Hum.

[116685]Genet. 9: 556–558, 2001.

[116686]20593.

[116687]Jaspers, M.; Zhang, Z.; Marynen, P.; Vekemans, S.; Aly, M.
S.;

[116688]Cuppens, H.; Hillicker, C.; Cassiman, J.-J.: Localization of
the

[116689]genes encoding the alpha-2 and alpha-4 subunits of the
human VLA-receptors

[116690]to chromosome 5q23–31 and 2q31–32 respectively.

(Abstract) Cytogenet.

[116691]Cell Genet. 58: 1870 only, 1991.

[116692]20594.

[116693]Kaplan, C.; Morel-Kopp, M. C.; Kroll, H.; Kiefel, V.;
Schlegel,

[116694]N.; Chesnel, N.; Mueller-Eckhardt, C.: HPA-5b (Br-a)
neonatal alloimmune

[116695]thrombocytopenia: clinical and immunological analysis of
39 cases. Brit.

[116696]J. Haemat. 78: 425-429, 1991.

[116697]20595.

[116698]Kiefel, V.; Santoso, S.; Katzmann, B.; Mueller-Eckhardt, C.:
A

[116699]new platelet-specific alloantigen Br(a): report of 4 cases
with neonatal

[116700]alloimmune thrombocytopenia. Vox Sang. 54: 101-106,
1988.

[116701]20596.

[116702]Kiefel, V.; Shechter, Y.; Atias, D.; Kroll, H.; Santoso, S.;
Mueller-Eckhardt,

[116703]C.: Neonatal alloimmune thrombocytopenia due to anti-

Br(b) (HPA-5a):

[116704]report of three cases in two families. Vox Sang. 60:
244-245, 1991.

[116705]20597.

[116706]Kritzik, M.; Savage, B.; Nugent, D. J.; Santoso, S.; Ruggeri,
Z.

[116707]M.; Kunicki, T. J.: Nucleotide polymorphisms in the alpha-
2 gene

[116708]define multiple alleles that are associated with differences
in platelet

[116709]alpha-2/beta-1 density. Blood 92: 2382-2388, 1998.

[116710]20598.

[116711]Kunicki, T. J.; Kritzik, M.; Annis, D. S.; Nugent, D. J.:
Hereditary

[116712]variation in platelet integrin alpha-2-beta-1 density is as-
sociated

[116713]with two silent polymorphisms in the alpha-2 gene coding
sequence. Blood 89:

[116714]1939-1943, 1997.

[116715]20599.

[116716]Mueller-Eckhardt, C.; Kiefel, V.; Grubert, A.; Kroll, H.;

Weisheit,

[116717]M.; Schmidt, S.; Mueller-Eckhardt, G.; Santoso, S.: 348

cases of

[116718]suspected neonatal alloimmune thrombocytopenia. Lancet

I: 363-366,

[116719]1989.

[116720]20600.

[116721]Nieuwenhuis, H. K.; Akkerman, J. W. N.; Houdijk, W. P.;

Sixma,

[116722]J. J.: Human blood platelets showing no response to colla-

gen fail

[116723]to express surface glycoprotein Ia. Nature 318: 470-472,

1985.

[116724]20601.

[116725]Santoso, S.; Amrhein, J.; Hofmann, H. A.; Sachs, U. J. H.;

Walka,

[116726]M. M.; Kroll, H.; Kiefel, V.: A point mutation thr799met on

the alpha-2

[116727]integrin leads to the formation of new human platelet al-

loantigen

[116728]Sit(a) and affects collagen-induced aggregation. Blood 94:

4103-4111,

[116729]1999.

[116730]20602.

[116731]Santoso, S.; Kalb, R.; Walka, M.; Kiefel, V.; Mueller-Eckhardt,

[116732]C.; Newman, P. J.: The human platelet alloantigens Br(a) and Br(b)

[116733]are associated with a single amino acid polymorphism on glycoprotein

[116734]Ia (integrin subunit alpha-2). J. Clin. Invest. 92: 2427-2432, 1993.

[116735]20603.

[116736]Santoso, S.; Kunicki, T. J.; Kroll, H.; Haberbosch, W.; Gardemann,

[116737]A.: Association of the platelet glycoprotein Ia C807T gene polymorphism

[116738]with nonfatal myocardial infarction in younger patients. Blood 93:

[116739]2449-2453, 1999.

[116740]20604.

[116741]Takada, Y.; Hemler, M. E.: The primary structure of the VLA-2/collagen

[116742]receptor alpha-2 subunit (platelet GPIa): homology to
other integrins

[116743]and the presence of a possible collagen-binding domain.

J. Cell Biol. 109:

[116744]397-407, 1989.

[116745]20605.

[116746]von Beckerath, N.; Koch, W.; Mehilli, J.; Bottiger, C.;
Schomig,

[116747]A.; Kastrati, A.: Glycoprotein Ia gene C807T polymorphism
and risk

[116748]for major adverse cardiac events within the first 30 days
after coronary

[116749]artery stenting. Blood 95: 3297-3301, 2000.

[116750]20606.

[116751]Fernandez-Ruiz, E.; Pardo-Manuel de Villena, F.; Rubio, M.
A.;

[116752]Corbi, A. L.; Rodriguez de Cordoba, S.; Sanchez-Madrid,
F.: Mapping

[116753]of the human VLA-alpha-4 gene to chromosome
2q31-q32. Europ. J. Immun. 22:

[116754]587-590, 1992.

[116755]20607.

[116756]Roseblatt, M.; Vuillet-Gaugler, M. H.; Leroy, C.;
Coulombel, L.

[116757]: Coexpression of two fibronectin receptors, VLA-4 and
VLA-5, by immature

[116758]human erythroblastic precursor cells. J. Clin. Invest. 87:
6-11,

[116759]1991.

[116760]20608.

[116761]Zhang, Z.; Vekemans, S.; Aly, M. S.; Jaspers, M.; Marynen,
P.;

[116762]Cassiman, J.-J.: The gene for the alpha-4 subunit of the
VLA-4 integrin

[116763]maps to chromosome 2q31-32. Blood 78: 2396-2399,
1991.

[116764]20609.

[116765]Hiesberger, T.; Trommsdorff, M.; Howell, B. W.; Goffinet,
A.; Mumby,

[116766]M. C.; Cooper, J. A.; Herz, J.: Direct binding of reelin to
VLDL

[116767]receptor and apoE receptor 2 induces tyrosine phosphory-
lation of disabled-1

[116768]and modulates tau phosphorylation. Neuron 24: 481–489,
1999.

[116769]20610.

[116770]Trommsdorff, M.; Gotthardt, M.; Hiesberger, T.; Shelton,
J.; Stockinger,

[116771]W.; Nimpf, J.; Hammer, R. E.; Richardson, J. A.; Herz, J.:
Reeler/Disabled-like

[116772]disruption of neuronal migration in knockout mice lacking
the VLDL

[116773]receptor and ApoE receptor 2. Cell 97: 689–701, 1999.

[116774]20611.

[116775]Archidiacono, N.; Grimaldi, G.; Rocchi, M.; Romeo, G.:
Mapping

[116776]of three sequenced clones coding for Zn(++) finger pro-
teins. Cytogenet.

[116777]Cell Genet. 51: 952 only, 1989.

[116778]20612.

[116779]Miller, J.; McLachlan, A. D.; Klug, A.: Repetitive zinc-
binding

[116780]domains in the protein transcription factor IIIA from
Xenopus oocytes. EMBO

[116781]J. 4: 1609–1614, 1985.

[116782]20613.

[116783]Rocchi, M.; Gentile, E.; Rosati, M.; Grimaldi, G.: The human
KRAB/FPB

[116784]containing zinc finger gene ZNF2 maps to chromosome
2q11.2. Cytogenet.

[116785]Cell Genet. 86: 305–306, 1999.

[116786]20614.

[116787]Lloyd, S. L.; Sargent, C. A.; Chalmers, J.; Lim, E.; Habeebu,
S.

[116788]S. M.; Affara, N. A.: An X-linked zinc finger gene mapping
to Xq21.1–q21.3

[116789]closely related to ZFX and ZFY: possible origins from a
common ancestral

[116790]gene. Nucleic Acids Res. 19: 4835–4841, 1991.

[116791]20615.

[116792]Rousseau-Merck, M. F.; Tunnacliffe, A.; Berger, R.; Ponder,
B.

[116793]A. J.; Thiesen, H. J.: A cluster of expressed zinc finger pro-
tein

[116794]genes in the pericentromeric region of human chromo-

some 10. Genomics 13:

[116795]845–848, 1992.

[116796]20616.

[116797]Huebner, K.; Druck, T.; Croce, C. M.; Thiesen, H. J.:

Twenty-seven

[116798]nonoverlapping zinc finger cDNAs from human T cells
map to nine different

[116799]chromosomes with apparent clustering. Am. J. Hum.
Genet. 48: 726–740,

[116800]1991.

[116801]20617.

[116802]Rousseau-Merck, M.-F.; Hillion, J.; Jonveaux, P.; Berger,
R.; Thiesen,

[116803]H. J.: Chromosomal localization of two human zinc finger
protein

[116804]genes (Kox), to chromosome 17p13p12. (Abstract) Cyto-
genet. Cell Genet. 58:

[116805]2009, 1991.

[116806]20618.

[116807]Rousseau-Merck, M.-F.; Hillion, J.; Jonveaux, P.; Couillin,
P.;

[116808]Seite, P.; Thiesen, H.-J.; Berger, R.: Chromosomal localization of

[116809]9 KOX zinc finger genes: physical linkages suggest clustering of KOX

[116810]genes on chromosomes 12, 16, and 19. Hum. Genet. 92: 583-587, 1993.

[116811]20619.

[116812]Bray, P.; Lichter, P.; Thiesen, H.-J.; Ward, D. C.; Dawid, I. B.

[116813]: Characterization and mapping of human genes encoding zinc finger

[116814]proteins. Proc. Nat. Acad. Sci. 88: 9563-9567, 1991.

[116815]20620.

[116816]Naylor, S. L.; Klebe, R. J.; Shows, T. B.: Argininosuccinic aciduria:

[116817]assignment of the argininosuccinate lyase gene to the pter-q22 region

[116818]of human chromosome 7 by bioautography. Proc. Nat. Acad. Sci. 75:

[116819]6159-6162, 1978.

[116820]20621.

[116821]Pastural, E.; Ersoy, F.; Yalman, N.; Wulffraat, N.; Grillo, E.;

[116822]Ozkinay, F.; Tezcan, I.; Gedikoglu, G.; Philippe, N.; Fischer, A.;

[116823]de Saint Basile, G.: Two genes are responsible for Griscelli syndrome

[116824]at the same 15q21 locus. Genomics 63: 299–306, 2000.

[116825]20622.

[116826]Lehmann, O. J.; El-ashry, M. F.; Ebenezer, N. D.; Ocaka, L.; Francis,

[116827]P. J.; Wilkie, S. E.; Patel, R. J.; Ficker, L.; Jordan, T.; Khaw,

[116828]P. T.; Bhattacharya, S. S.: A novel keratocan mutation causing autosomal

[116829]recessive cornea plana. Invest. Ophthalm. Vis. Sci. 42: 3118–3122,

[116830]2001.

[116831]20623.

[116832]Hayasaka, K.; Tada, K.; Kikuchi, G.; Winter, S.; Nyhan, W. L.:

[116833]Nonketotic hyperglycinemia: two patients with primary defects of P-protein

[116834]and T-protein, respectively, in the glycine cleavage system. Pediat.

[116835]Res. 17: 967-970, 1983.

[116836]20624.

[116837]Kure, S.; Mandel, H.; Rolland, M.-O.; Sakata, Y.; Shinka, T.;
Drugan,

[116838]A.; Boneh, A.; Tada, K.; Matsubara, Y.; Narisawa, K.: A
missense

[116839]mutation (his42arg) in the T-protein gene from a large Is-
raeli-Arab

[116840]kindred with nonketotic hyperglycinemia. Hum. Genet.
102: 430-434,

[116841]1998.

[116842]20625.

[116843]Kure, S.; Shinka, T.; Sakata, Y.; Osamu, N.; Takayanagi, M.;
Tada,

[116844]K.; Matsubara, Y.; Narisawa, K.: A one-base deletion
(183delC) and

[116845]a missense mutation (D276H) in the T-protein gene from
a Japanese

[116846]family with nonketotic hyperglycinemia. J. Hum. Genet.
43: 135-137,

[116847]1998.

[116848]20626.

[116849]Nanao, K.; Okamura-Ikeda, K.; Motokawa, Y.; Danks, D.
M.; Baumgartner,

[116850]E. R.; Takada, G.; Hayasaka, K.: Identification of the muta-
tions

[116851]in the T-protein gene causing typical and atypical nonke-
totic hyperglycinemia. Hum.

[116852]Genet. 93: 655-658, 1994.

[116853]20627.

[116854]Nanao, K.; Takada, G.; Takahashi, E.; Seki, N.; Komatsu,
Y.; Okamura-Ikeda,

[116855]K.; Motokawa, Y.; Hayasaka, K.: Structure and chromoso-
mal localization

[116856]of the aminomethyltransferase gene (AMT). Genomics 19:
27-30, 1994.

[116857]20628.

[116858]Toone, J. R.; Applegarth, D. A.; Coulter-Mackie, M. B.;
James,

[116859]E. R.: Biochemical and molecular investigations of patients
with

[116860]nonketotic hyperglycemia. Molec. Genet. Metab. 70:
116-121, 2000.

[116861]20629.

[116862]Toone, J. R.; Applegarth, D. A.; Coulter-Mackie, M. B.;
James,

[116863]E. R.: Identification of the first reported splice site mutation

[116864](IVS7-1G-A) in the aminomethyltransferase (T-protein) gene (AMT) of

[116865]the glycine cleavage complex in 3 unrelated families with nonketotic

[116866]hyperglycinemia. (Abstract) Hum. Mutat. 17: 76 only, 2000.

[116867]20630.

[116868]Fujiwara, K.; Okamura-Ikeda, K.; Hayasaka, K.; Motokawa, Y.: The

[116869]primary structure of human H-protein of the glycine cleavage system

[116870]deduced by cDNA cloning. Biochem. Biophys. Res. Commun. 176: 711-716,

[116871]1991.

[116872]20631.

[116873]Hiraga, K.; Kochi, H.; Hayasaka, K.; Kikuchi, G.; Nyhan, W. L.

[116874]: Defective glycine cleavage system in non-ketotic hyperglycinemia:

[116875]occurrence of a less active glycine decarboxylase and an abnormal

[116876]aminomethyl carrier protein. J. Clin. Invest. 68: 525-534, 1981.

[116877]20632.

[116878]Hiraga, K.; Kure, S.; Yamamoto, M.; Ishiguro, Y.; Suzuki, T.:

[116879]Cloning of cDNA encoding human H-protein, a constituent of the glycine

[116880]cleavage system. Biochem. Biophys. Res. Commun. 151: 758-762, 1988.

[116881]20633.

[116882]Koyata, H.; Hiraga, K.: The glycine cleavage system: structure

[116883]of a cDNA encoding human H-protein, and partial characterization of

[116884]its gene in patients with hyperglycinemias. Am. J. Hum. Genet. 48:

[116885]351-361, 1991.

[116886]20634.

[116887]O'Brien, W. E.; McInnes, R.; Kalumuck, K.; Adcock, M.:

Cloning

[116888]and sequence analysis of cDNA for human argininosuccinate lyase. Proc.

[116889]Nat. Acad. Sci. 83: 7211-7215, 1986.

[116890]20635.

[116891]Piatigorsky, J.; O'Brien, W. E.; Norman, B. L.; Kalumuck, K.;

[116892]Wistow, G. J.; Borras, T.; Nickerson, J. M.; Wawrousek, E.
F.: Gene

[116893]sharing by delta-crystallin and argininosuccinate lyase.
Proc. Nat.

[116894]Acad. Sci. 85: 3479-3483, 1988.

[116895]20636.

[116896]Piatigorsky, J.; Wistow, G.: The recruitment of crystallins:

[116897]new functions precede gene duplication. Science 252:
1078-1079,

[116898]1991.

[116899]20637.

[116900]Pijpers, L.; Kleijer, W. J.; Reuss, A.; Jahoda, M. G. J.; Los,

[116901]F. J.; Sachs, E. S.; Wladimiroff, J. W.: Transabdominal

chorionic

[116902] villus sampling in a multiple pregnancy at risk of argininosuccinic

[116903] aciduria: a case report. Am. J. Med. Genet. 36: 449–450, 1990.

[116904] 20638.

[116905] Qureshi, I. A.; Letarte, J.; Ouellet, R.; Lemieux, B.: Enzymologic

[116906] and metabolic studies in two families affected by argininosuccinic

[116907] aciduria. Pediat. Res. 12: 256–262, 1978.

[116908] 20639.

[116909] Shih, V. E.; Efron, M. L.: Urea cycle disorders. In: Stanbury, J. B.; Wyngaarden, J. B.; Fredrickson, D. S.: The Metabolic Basis

[116911] of Inherited Disease. New York: McGraw–Hill (pub.) (3rd ed.)

[116912]: 1972. Pp. 370–392.

[116913] 20640.

[116914] Shih, V. E.; Littlefield, J. W.; Moser, H. W.: Argininosuccinase

- [116915]deficiency in fibroblasts cultured from patients with
argininosuccinic
aciduria. *Biochem. Genet.* 3: 81–83, 1969.
- [116917]20641.
- [116918]Simard, L.; O'Brien, W. E.; McInnes, R. R.: Argininosuccinate
lyase deficiency: evidence for heterogeneous structural
gene mutations
by immunoblotting. *Am. J. Hum. Genet.* 39: 38–51, 1986.
- [116921]20642.
- [116922]Todd, S.; McGill, J. R.; McCombs, J. L.; Moore, C. M.; Weider,
J.; Naylor, S. L.: cDNA sequence, interspecies comparison
and gene
mapping analysis of argininosuccinate lyase. *Genomics* 4:
53–59,
1989.
- [116926]20643.
- [116927]Van der Heiden, C.; Gerards, L. J.; van Biervliet, J. P. G. M.;
Desplanque, J.; DeBree, P. K.; Van Sprang, F. J.; Wadman,
S. K.:

[116929] Lethal neonatal argininosuccinate lyase deficiency in four children

[116930] from one sibship. *Helv. Paediat. Acta* 31: 407-417, 1976.

[116931] 20644.

[116932] Walker, D. C.; McCloskey, D. A.; Simard, L. R.; McInnes, R. R.

[116933]: Molecular analysis of human argininosuccinate lyase (ASAL): mutant

[116934] characterization and alternate splicing of the active site. (Abstract) *Am.*

[116935] *J. Hum. Genet.* 45 (suppl.): A227 only, 1989.

[116936] 20645.

[116937] Walker, D. C.; McCloskey, D. A.; Simard, L. R.; McInnes, R. R.

[116938]: Identification of a mutation frequently involved in inter-allelic

[116939] complementation at the human argininosuccinic acid lyase locus. (Abstract) *Am.*

[116940] *J. Hum. Genet.* 47 (suppl.): A169 only, 1990.

[116941] 20646.

[116942] Walker, D. C.; McCloskey, D. A.; Simard, L. R.; McInnes, R.

R.

[116943] Molecular analysis of human argininosuccinate lyase:
mutant characterization

[116944] and alternative splicing of the coding region. Proc. Nat.
Acad. Sci. 87:

[116945] 9625–9629, 1990.

[116946] 20647.

[116947] Widhalm, K.; Koch, S.; Scheibenreiter, S.; Knoll, E.;
Colombo,

[116948] J. P.; Bachmann, C.; Thalhammer, O.: Long-term follow-up
of 12 patients

[116949] with the late-onset variant of argininosuccinic acid lyase
deficiency:

[116950] no impairment of intellectual and psychomotor develop-
ment during therapy. Pediatrics 89:

[116951] 1182–1184, 1992.

[116952] 20648.

[116953] Aloysia, M.; Gelb, A. G.; Fudenberg, H.; Hamper, J.; Tip-
pett, P.;

[116954] Race, R. R.: The expected 'Bombay' group O(H-A1) and
O(H-A2). Transfusion 1:

[116955] 212–217, 1961.

[116956]20649.

[116957]Bhende, Y. M.; Deshpande, C. K.; Bhatia, H. M.; Sanger, R.;
Race,

[116958]R. R.; Morgan, W. T. J.; Watkins, W. M.: A 'new' blood
group character

[116959]related to the ABO system. Lancet I: 903–904, 1952.

[116960]20650.

[116961]Gerard, G.; Vitrac, D.; Le Pendu, J.; Muller, A.; Oriol, R.: H-
deficient

[116962]blood groups (Bombay) of Reunion Island. Am. J. Hum.
Genet. 34:

[116963]937–947, 1982.

[116964]20651.

[116965]Hrubisko, M.; Laluha, J.; Mergancova, O.; Zakovicova, S.:
New

[116966]variants in the ABOH blood group system due to interac-
tion of recessive

[116967]genes controlling the formation of H antigen in erythro-
cytes: the

[116968]'Bombay-like' phenotypes O HM, OB HM, OAB HM. Vox
Sang. 19: 113–122,

[116969]1970.

[116970]20652.

[116971]Kelly, R. J.; Ernst, L. K.; Larsen, R. D.; Bryant, J. G.; Robinson,

[116972]J. S.; Lowe, J. B.: Molecular basis for H blood group deficiency

[116973]in Bombay (Oh) and para-Bombay individuals. Proc. Nat. Acad. Sci. 91:

[116974]5843–5847, 1994.

[116975]20653.

[116976]Koda, Y.; Soejima, M.; Johnson, P. H.; Smart, E.; Kimura, H.:

[116977]Missense mutation of FUT1 and deletion of FUT2 are responsible for

[116978]Indian Bombay phenotype at ABO blood group system. Biochem. Biophys.

[116979]Res. Commun. 238: 21–25, 1997.

[116980]20654.

[116981]Larsen, R. D.; Ernst, L. K.; Nair, R. P.; Lowe, J. B.: Molecular

[116982]cloning, sequence, and expression of a human GDP–L–fucose:beta–D–galactoside

[116983]2-alpha-L-fucosyltransferase cDNA that can form the H blood group

[116984]antigen. Proc. Nat. Acad. Sci. 87: 6674-6678, 1990.

[116985]20655.

[116986]Le Pendu, J.; Cartron, J. P.; Lemieux, R. U.; Oriol, R.: The presence

[116987]of at least two different H-blood-group-related beta-D-Gal alpha-2-L-fucosyltransferases

[116988]in human serum and the genetics of blood group H substances. Am.

[116989]J. Hum. Genet. 37: 749-760, 1985.

[116990]20656.

[116991]Le Pendu, J.; Clamagirand-Mulet, C.; Cartron, J.-P.; Gerard, G.;

[116992]Vitrac, D.; Oriol, R.: H-deficient blood groups of Reunion Island.

[116993]III. Alpha-2-L-fucosyltransferase activity in sera of homozygous and

[116994]heterozygous individuals. Am. J. Hum. Genet. 35: 497-507, 1983.

[116995]20657.

[116996]Le Pendu, J.; Gerard, G.; Vitrac, D.; Juszczak, G.; Liberge, G.;

[116997]Rouger, P.; Salmon, C.; Lambert, F.; Dalix, A.-M.; Oriol, R.:
H-deficient

[116998]blood groups of Reunion Island. II. Differences between
Indians (Bombay

[116999]phenotype) and whites (Reunion phenotype). Am. J. Hum.
Genet. 35:

[117000]484-496, 1983.

[117001]20658.

[117002]Levine, P.; Robinson, E.; Celano, M.; Briggs, O.; Falkinburg,

[117003]L.: Gene interaction resulting in suppression of blood
group substance

[117004]B. Blood 10: 1100-1108, 1955.

[117005]20659.

[117006]Oriol, R.: Personal Communication. Paris, France
2/7/1983.

[117007]20660.

[117008]Sathe, M.; Bhatia, H. M.: Bombay (OH) phenotype in two
generations

[117009]in an Indian family. Vox Sang. 30: 312-314, 1976.

[117010]20661.

[117011]Yunis, E. J.; Svoldal, J. M.; Bridges, R. A.: Genetics of the

[117012]Bombay phenotype. Blood 33: 124–132, 1969.

[117013]20662.

[117014]Lees–Miller, J. P.; Helfman, D. M.: The molecular basis for
tropomyosin

[117015]isoform diversity. BioEssays 13: 429–437, 1991.

[117016]20663.

[117017]MacLeod, A. R.; Gooding, C.: Human hTM–alpha gene: ex-
pression

[117018]in muscle and nonmuscle tissue. Molec. Cell. Biol. 8:
433–440, 1988.

[117019]20664.

[117020]Muthuchamy, M.; Pieples, K.; Rethinasamy, P.; Hoit, B.;
Grupp,

[117021]J. L.; Boivin, G. P.; Wolska, B.; Evans, C.; Solaro, R. J.; Wiece-
zorek,

[117022]D. F.: Mouse model of a familial hypertrophic cardiomy-
opathy mutation

[117023]in alpha–tropomyosin manifests cardiac dysfunction. Circ.
Res. 85:

[117024]47–56, 1999.

[117025]20665.

[117026]Schleef, M.; Werner, K.; Satzger, U.; Kaupmann, K.;
Jockusch, H.

[117027]: Chromosomal localization and genomic cloning of the
mouse alpha-tropomyosin

[117028]gene Tpm-1. Genomics 17: 519–521, 1993.

[117029]20666.

[117030]Thierfelder, L.; Watkins, H.; MacRae, C.; Lamas, R.;
McKenna, W.;

[117031]Vosberg, H.-P.; Seidman, J. G.; Seidman, C. E.: Alpha-
tropomyosin

[117032]and cardiac troponin T mutations cause familial hyper-
trophic cardiomyopathy:

[117033]a disease of the sarcomere. Cell 77: 701–712, 1994.

[117034]20667.

[117035]Watkins, H.; Anan, R.; Coviello, D. A.; Spirito, P.; Seidman,

[117036]J. G.; Seidman, C. E.: A de novo mutation in alpha-
tropomyosin that

[117037]causes hypertrophic cardiomyopathy. Circulation 91:
2302–2305, 1995.

[117038]20668.

[117039]Watkins, H.; McKenna, W. J.; Thierfelder, L.; Suk, H. J.;
Anan,

[117040]R.; O'Donoghue, A.; Spirito, P.; Matsumori, A.; Moravec, C.
S.; Seidman,

[117041]J. G.; Seidman, C. E.: Mutations in the genes for cardiac
troponin

[117042]T and alpha-tropomyosin in hypertrophic cardiomyopa-
thy. New Eng.

[117043]J. Med. 332: 1058-1064, 1995.

[117044]20669.

[117045]Gahlmann, R.; Kedes, L.: Cloning, structural analysis, and
expression

[117046]of the human fast twitch skeletal muscle troponin C gene.
J. Biol.

[117047]Chem. 265: 12520-12528, 1990.

[117048]20670.

[117049]Gahlmann, R.; Wade, R.; Gunning, P.; Kedes, L.: Differen-
tial expression

[117050]of slow and fast skeletal muscle troponin C: slow skeletal
muscle

[117051]troponin C is expressed in human fibroblasts. J. Molec.
Biol. 201:

[117052]379–391, 1988.

[117053]20671.

[117054]Townsend, P. J.; Yacoub, M. H.; Barton, P. J. R.: Assign-
ment of

[117055]the human fast skeletal muscle troponin C gene (TNNC2)
between D20S721

[117056]and GCT10F11 on chromosome 20 by somatic cell hybrid
analysis. Ann.

[117057]Hum. Genet. 61: 457–459, 1997.

[117058]20672.

[117059]Bermingham, N.; Hernandez, D.; Balfour, A.; Gilmour, F.;
Martin,

[117060]J. E.; Fisher, E. M. C.: Mapping TNNC1, the gene that en-
codes cardiac

[117061]troponin I in the human and mouse. Genomics 30:
620–622, 1995.

[117062]20673.

[117063]Kwon, B. S.; Halaban, R.; Chintamaneni, C.: Molecular ba-
sis of

[117064]mouse Himalayan mutation. Biochem. Biophys. Res. Commun. 161: 252-260,

[117065]1989.

[117066]20674.

[117067]DiLella, A. G.: Chromosomal assignment of the human immunophilin

[117068]FKBP-12 gene. Biochem. Biophys. Res. Commun. 179: 1427-1433, 1991.

[117069]20675.

[117070]DiLella, A. G.; Hawkins, A.; Craig, R. J.; Schreiber, S. L.; Griffin,

[117071]C. A.: Chromosomal band assignments of the genes encoding human FKBP12

[117072]and FKBP13. Biochem. Biophys. Res. Commun. 189: 819-823, 1992.

[117073]20676.

[117074]Goebl, M. G.: The peptidyl-prolyl isomerase, FK506-binding protein,

[117075]is most likely the 12 kd endogenous inhibitor 2 of protein kinase

[117076]C. (Letter) Cell 64: 1051-1052, 1991.

[117077]20677.

[117078]Jin, Y.-J.; Albers, M. W.; Lane, W. S.; Bierer, B. E.; Schreiber,

[117079]S. L.; Burakoff, S. J.: Molecular cloning of a membrane-associated

[117080]human FK506- and rapamycin-binding protein, FKBP-13.
Proc. Nat. Acad.

[117081]Sci. 88: 6677-6681, 1991.

[117082]20678.

[117083]Maki, N.; Sekiguchi, F.; Nishimaki, J.; Miwa, K.; Hayano, T.;
Takahashi,

[117084]N.; Suzuki, M.: Complementary DNA encoding the human
T-cell FK506-binding

[117085]protein, a peptidylprolyl cis-trans isomerase distinct from
cyclophilin. Proc.

[117086]Nat. Acad. Sci. 87: 5440-5443, 1990.

[117087]20679.

[117088]Peattie, D. A.; Hsaio, K.; Benasutti, M.; Lippke, J. A.: Three

[117089]distinct messenger RNAs can encode the human immunosuppressant-binding

[117090]protein FKBP12. Gene 150: 251-257, 1994.

[117091]20680.

[117092]Shou, W.; Aghdasi, B.; Armstrong, D. L.; Guo, Q.; Bao, S.;
Charng,

[117093]M.-J.; Mathews, L. M.; Schneider, M. D.; Hamilton, S. L.;
Matzuk,

[117094]M. M.: Cardiac defects and altered ryanodine receptor
function in

[117095]mice lacking FKBP12. Nature 391: 489–492, 1998.

[117096]20681.

[117097]Standaert, R. F.; Galat, A.; Verdine, G. L.; Schreiber, S. L.:

[117098]Molecular cloning and overexpression of the human
FK506-binding protein

[117099]FKBP. Nature 346: 671–674, 1990.

[117100]20682.

[117101]Wang, T.; Donahoe, P. K.; Zervos, A. S.: Specific interaction

[117102]of type I receptors of the TGF-beta family with the im-
munophilin FKBP-12. Science 265:

[117103]674–676, 1994.

[117104]20683.

[117105]Bichi, R.; Shinton, S. A.; Martin, E. S.; Koval, A.; Calin, G.

[117106]A.; Cesari, R.; Russo, G.; Hardy, R. R.; Croce, C. M.: Human chronic
[117107]lymphocytic leukemia modeled in mouse by targeted TCL1
expression. Proc.
[117108]Nat. Acad. Sci. 99: 6955–6960, 2002.
[117109]20684.
[117110]Davey, M. P.; Bertness, V.; Nakahara, K.; Johnson, J. P.;
McBride,
[117111]O. W.; Waldmann, T. A.; Kirsch, I. R.: Juxtaposition of the
T-cell
[117112]receptor alpha-chain locus (14q11) and a region (14q32)
of potential
[117113]importance in leukemogenesis by a 14;14 translocation in
a patient
[117114]with T-cell chronic lymphocytic leukemia and ataxia-
telangiectasia. Proc.
[117115]Nat. Acad. Sci. 85: 9287–9291, 1988.
[117116]20685.
[117117]Hecht, F.; Morgan, R.; Hecht, B. K.-M.; Smith, S. D.: Com-
mon region
[117118]on chromosome 14 in T-cell leukemia and lymphoma.
Science 226: 1445–1447,

[117119]1984.

[117120]20686.

[117121]Laine, J.; Kunstle, G.; Obata, T.; Sha, M.; Noguchi, M.: The protooncogene

[117122]TCL1 is an Akt kinase coactivator. *Molec. Cell* 6: 395–407, 2000.

[117123]20687.

[117124]Mathieu–Mahul, D.; Caubet, J. F.; Bernheim, A.; Mauchauffe, M.;

[117125]Palmer, E.; Berger, R.; Larsen, C.–J.: Molecular cloning of a DNA

[117126]fragment from human chromosome 14(14q11) involved in T–cell malignancies. *EMBO*

[117127]*J.* 4: 3427–3433, 1985.

[117128]20688.

[117129]Mengle–Gaw, L.; Albertson, D. G.; Sherrington, P. D.; Rabbits,

[117130]T. H.: Analysis of a T–cell tumor–specific breakpoint cluster at

[117131]human chromosome 14q32. *Proc. Nat. Acad. Sci.* 85: 9171–9175, 1988.

[117132]20689.

[117133]Howard, H. C.; Mount, D. B.; Rochefort, D.; Byun, N.;
Dupre, N.;

[117134]Lu, J.; Fan, X.; Song, L.; Riviere, J.-B.; Prevost, C.; Horst, J.;

[117135]Simonati, A.; and 12 others: The K-Cl cotransporter KCC3
is mutant

[117136]in a severe peripheral neuropathy associated with agene-
sis of the

[117137]corpus callosum. Nature Genet. 7Oct, 2002. Note: Ad-
vance Electronic

[117138]Publication.

[117139]20690.

[117140]Haire, R. N.; Litman, G. W.: The murine form of TXK, a
novel TEC

[117141]kinase expressed in thymus maps to chromosome 5.
Mammalian Genome 6:

[117142]476-480, 1995.

[117143]20691.

[117144]Haire, R. N.; Ohta, Y.; Lewis, J. E.; Fu, S. M.; Kroisel, P.; Lit-
man,

[117145]G. W.: TXK, a novel human tyrosine kinase expressed in T

cells shares

[117146]sequence identity with Tec family kinases and maps to
4p12. Hum.

[117147]Molec. Genet. 3: 897–901, 1994.

[117148]20692.

[117149]Schneider, H.; Schwartzberg, P. L.; Rudd, C. E.: Resting
lymphocyte

[117150]kinase (Rlk/Txk) phosphorylates the YVKM motif and reg-
ulates PI 3-kinase

[117151]binding to T-cell antigen CTLA-4. Biochem. Biophys. Res.
Commun. 252:

[117152]14–19, 1998.

[117153]20693.

[117154]Takeba, Y.; Nagafuchi, H.; Takeno, M.; Kashiwakura, J.;
Suzuki,

[117155]N.: Txk, a member of nonreceptor tyrosine kinase of Tec
family, acts

[117156]as a Th1 cell-specific transcription factor and regulates
IFN-gamma

[117157]gene transcription. J. Immun. 168: 2365–2370, 2002.

[117158]20694.

[117159]Gasdaska, J. R.; Gasdaska, P. Y.; Gallegos, A.; Powis, G.:
Human

[117160]thioredoxin reductase gene localization to chromosomal
position 12q23-q24.1

[117161]and mRNA distribution in human tissue. Genomics 37:
257-259, 1996.

[117162]20695.

[117163]Gasdaska, P. Y.; Gasdaska, J. R.; Cochran, S.; Powis, G.:
Cloning

[117164]and sequencing of a human thioredoxin reductase. FEBS
Lett. 373:

[117165]5-9, 1995.

[117166]20696.

[117167]Russo, M. W.; Sevetson, B. R.; Milbrandt, J.: Identification
of

[117168]NAB1, a repressor of NGFI-A- and Krox20-mediated tran-
scription. Proc.

[117169]Nat. Acad. Sci. 92: 6873-6877, 1995.

[117170]20697.

[117171]Svaren, J.; Apel, E. D.; Simburger, K. S.; Jenkins, N. A.;
Gilbert,

[117172]D. J.; Copeland, N. A.; Milbrandt, J.: The Nab2 and Stat6 genes share

[117173]a common transcription termination region. Genomics 41: 33–39, 1997.

[117174]20698.

[117175]Swirnoff, A. H.; Apel, E. D.; Svaren, J.; Sevetson, B. R.; Zimonjic,

[117176]D. B.; Popescu, N. C.; Milbrandt, J.: Nab1, a corepressor of NGFI-A

[117177](Egr-1), contains an active transcriptional repression domain. Molec.

[117178]Cell. Biol. 18: 512–524, 1998.

[117179]20699.

[117180]Venken, K.; Di Maria, E.; Bellone, E.; Balestra, P.; Cassandrini,

[117181]D.; Mandich, P.; De Jonghe, P.; Timmerman, V.; Svaren, J.: Search

[117182]for mutation in the EGR2 corepressor proteins, NAB1 and NAB2, in human

[117183]peripheral neuropathies. Neurogenetics 4: 37–41, 2002.

[117184]20700.

- [117185]Nichols, R. C.; Blinder, J.; Pai, S. I.; Ge, Q.; Targoff, I. N.;
- [117186]Plotz, P. H.; Liu, P.: Assignment of two human autoantigen genes:
- [117187]isoleucyl-tRNA synthetase locates to 9q21 and lysyl-tRNA synthetase
- [117188]locates to 16q23-q24. Genomics 36: 210-213, 1996.
- [117189]20701.
- [117190]Nichols, R. C.; Raben, N.; Boerkoel, C. F.; Plotz, P. H.: Human
- [117191]isoleucyl-tRNA synthetase: sequence of the cDNA, alternative mRNA
- [117192]splicing, and the characteristics of an unusually long C-terminal
- [117193]extension. Gene 155: 299-304, 1995.
- [117194]20702.
- [117195]Tommerup, N.; Leffers, H.: Assignment of human KH-box-containing
- [117196]genes by in situ hybridization: HNRNPK maps to 9q21.32-q21.33, PCBP1
- [117197]to 2p12-p13, and PCBP2 to 12q13.12-q13.13, distal to FRA12A. Genomics 32:

[117198]297–298, 1996.

[117199]20703.

[117200]Martell, K. J.; Kwak, S.; Hakes, D. J.; Dixon, J. E.; Trent, J.

- [117201]M.: Chromosomal localization of four human VH1-like protein-tyrosine phosphatases. Genomics 22: 462-464, 1994.
- [117202]20704.
- [117203]Chrast, R.; Chen, H.; Morris, M. A.; Antonarakis, S. E.: Mapping of the human transcription factor GABPA (E4TF1-60) gene to chromosome 21. Genomics 28: 119-122, 1995.
- [117204]20705.
- [117205]Gugneja, S.; Virbasius, J. V.; Scarpulla, R. C.: Four structurally distinct, non-DNA-binding subunits of human nuclear respiratory factor 2 share a conserved transcriptional activation domain. Molec. Cell. Biol. 15: 102-111, 1995.
- [117206]20706.
- [117207]Guo, A.; Nie, F.; Wong-Riley, M.: Human nuclear respiratory factor 2-alpha subunit cDNA: isolation, subcloning, sequencing,

and in situ

[117215]hybridization of transcripts in normal and monocularly
deprived macaque

[117216]visual system. J. Comp. Neurol. 417: 221–232, 2000.

[117217]20707.

[117218]Sawada, J.; Goto, M.; Watanabe, H.; Handa, H.; Yoshida, M.
C.:

[117219]Regional mapping of two subunits of transcription factor
E4TF1 to

[117220]human chromosome. Jpn. J. Cancer Res. 86: 10–12, 1995.

[117221]20708.

[117222]Virbasius, J. V.; Scarpulla, R. C.: Activation of the human
mitochondrial

[117223]transcription factor A gene by nuclear respiratory factors:
a potential

[117224]regulatory link between nuclear and mitochondrial gene
expression

[117225]in organelle biogenesis. Proc. Nat. Acad. Sci. 91:
1309–1313, 1994.

[117226]20709.

[117227]Watanabe, H.; Sawada, J.; Yano, K.-I.; Yamaguchi, K.; Goto,

M.;

[117228]Handa, H.: cDNA cloning of transcription factor E4TF1 subunits with

[117229]Ets and notch motifs. *Molec. Cell. Biol.* 13: 1385–1391, 1993.

[117230]20710.

[117231]Chambrud, B.; Radanyi, C.; Camonis, J. H.; Rajkowski, K.; Schumacher,

[117232]M.; Baulieu, E.–E.: Immunophilins, Refsum disease, and lupus nephritis:

[117233]the peroxisomal enzyme phytanoyl–CoA alpha–hydroxylase is a new FKBP–associated

[117234]protein. *Proc. Nat. Acad. Sci.* 96: 2104–2109, 1999.

[117235]20711.

[117236]Dixon, J.; Loftus, S. K.; Gladwin, A. J.; Scambler, P. J.; Wasmuth,

[117237]J. J.; Dixon, M. J.: Cloning of the human heparan sulfate–N–deacetylase/N–sulfotransferase

[117238]gene from the Treacher Collins syndrome candidate region at 5q32–q33.1. *Genomics* 26:

[117239]239–244, 1995.

[117240]20712.

[117241]Gladwin, A. J.; Dixon, J.; Loftus, S. K.; Wasmuth, J. J.;
Dixon,

[117242]M. J.: Genomic organization of the human heparan sul-
fate-N-deacetylase/N-sulfotransferase

[117243]gene: exclusion from a causative role in the pathogenesis
of Treacher

[117244]Collins syndrome. Genomics 32: 471-473, 1996.

[117245]20713.

[117246]Hashimoto, Y.; Orellana, A.; Gil, G.; Hirschberg, C. B.:
Molecular

[117247]cloning and expression of rat liver N-heparan sulfate sul-
fotransferase. J.

[117248]Biol. Chem. 267: 15744-15750, 1995.

[117249]20714.

[117250]Chen, H.; Antonarakis, S. E.: Localisation of a human ho-
mologue

[117251]of the Drosophila *mnb* and rat *Dyrk* genes to chromosome
21q22.2. Hum.

[117252]Genet. 99: 262-265, 1997.

[117253]20715.

[117254]Patil, N.; Cox, D. R.; Bhat, D.; Faham, M.; Myers, R. M.; Pe-

terson,

[117255]A. S.: A potassium channel mutation in weaver mice implicates membrane

[117256]excitability in granule cell differentiation. *Nature Genet.*

11:

[117257]126–129, 1995.

[117258]20716.

[117259]Shindoh, N.; Kudoh, J.; Maeda, H.; Yamaki, A.; Minoshima, S.; Shimizu,

[117260]Y.; Shimizu, N.: Cloning of a human homolog of the *Drosophila* minibrain/rat

[117261]Dyrk gene from 'the Down syndrome critical region' of chromosome 21. *Biochem.*

[117262]*Biophys. Res. Commun.* 225: 92–99, 1996.

[117263]20717.

[117264]Smith, D. J.; Stevens, M. E.; Sudanagunta, S. P.; Bronson, R. T.;

[117265]Makhinson, M.; Watabe, A. M.; O'Dell, T. J.; Fung, J.; Weier, H.–U.

[117266]G.; Cheng, J.–F.; Rubin, E. M.: Functional screening of 2 Mb of human

[117267]chromosome 21q22.2 in transgenic mice implicates mini–

brain in learning

[117268]defects associated with Down syndrome. Nature Genet.
16: 28–36,

[117269]1997.

[117270]20718.

[117271]Song, W.-J.; Chung, S.-H.; Kurnit, D. M.: The murine Dyrk
protein

[117272]maps to chromosome 16, localizes to the nucleus, and
can form multimers. Biochem.

[117273]Biophys. Res. Commun. 231: 640–644, 1997.

[117274]20719.

[117275]Song, W.-J.; Sternberg, L. R.; Kasten-Sportes, C.; Van
Keuren,

[117276]M. L.; Chung, S.-H.; Slack, A. C.; Miller, D. E.; Glover, T.
W.; Chiang,

[117277]P.-W.; Lou, L.; Kurnit, D. M.: Isolation of human and
murine homologues

[117278]of the Drosophila minibrain gene: human homologue
maps to 21q22.2

[117279]in the Down syndrome 'critical region.' Genomics 38:
331–339, 1996.

[117280]20720.

[117281]Tejedor, F.; Zhu, X. R.; Kaltenbach, E.; Ackermann, A.;
Baumann,
[117282]A.; Canal, I.; Heisenberg, M.; Fischbach, K. F.; Pongs, O.:
Minibrain:
[117283]a new protein kinase family involved in postembryonic
neurogenesis
[117284]in *Drosophila*. *Neuron* 14: 287–301, 1995.
[117285]20721.

[117286]Algar, E.; Brickell, S.; Deeble, G.; Amor, D.; Smith, P.:
Analysis
[117287]of CDKN1C in Beckwith Wiedemann syndrome. *Hum. Mu-
tat.* 15: 497–508,
[117288]2000.
[117289]20722.

[117290]Bhuiyan, Z. A.; Yatsuki, H.; Sasaguri, T.; Joh, K.; Soejima,
H.;
[117291]Zhu, X.; Hatada, I. Morisaki, H.; Morisaki, T.; Mukai, T.:
Functional
[117292]analysis of the p57(KIP2) gene mutation in Beckwith-
Wiedemann syndrome. *Hum.*
[117293]*Genet.* 104: 205–210, 1999.

[117294]20723.

[117295]Fitzpatrick, G. V.; Soloway, P. D.; Higgins, M. J.: Regional
loss

[117296]of imprinting and growth deficiency in mice with a tar-
geted deletion

[117297]of KvDMR1. Nature Genet. 9Sept, 2002. Note: Advance
Electronic Publication.

[117298]20724.

[117299]Hatada, I.; Mukai, T.: Genomic imprinting of p57(KIP2), a
cyclin-dependent

[117300]kinase inhibitor, in mouse. Nature Genet. 11: 204-206,
1995.

[117301]20725.

[117302]Hatada, I.; Nabetani, A.; Morisaki, H.; Xin, Z.; Ohishi, S.;
Tonoki,

[117303]H.; Niikawa, N.; Inoue, M.; Komoto, Y.; Okada, A.; Ste-
ichen, E.; Ohashi,

[117304]H.; Fukushima, Y.; Nakayama, M.; Mukai, T.: New
p57(KIP2) mutations

[117305]in Beckwith-Wiedemann syndrome. Hum. Genet. 100:
681-683, 1997.

[117306]20726.

[117307] Hatada, I.; Ohashi, H.; Fukushima, Y.; Kaneko, Y.; Inoue, M.; Komoto, Y.; Okada, A.; Ohishi, S.; Nabetani, A.; Morisaki, H.; Nakayama, M.; Niikawa, M.; Mukai, T.: An imprinted gene p57(KIP2) is mutated in Beckwith–Wiedemann syndrome. *Nature Genet.* 14: 171–173, 1996.

[117311] 20727.

[117312] John, R. M.; Ainscough, J. F.–X.; Barton, S. C.; Surani, M. A. [117313]: Distant cis–elements regulate imprinted expression of the mouse p57(Kip2) (Cdkn1c) gene: implications for the human disorder, Beckwith–Wiedemann [117314] syndrome. *Hum. Molec. Genet.* 10: 1601–1609, 2001.

[117316] 20728.

[117317] Lee, M.–H.; Reynisdottir, I.; Massague, J.: Cloning of p57(KIP2), [117318] a cyclin–dependent kinase inhibitor with unique domain structure and [117319] tissue distribution. *Genes Dev.* 9: 639–649, 1995.

[117320]20729.

[117321]Zhang, X.; Zhou, J.; Reenders, S. T.; Tryggvason, K.: Structure

[117322]of the human type IV collagen COL4A6 gene, which is mutated in Alport

[117323]syndrome-associated leiomyomatosis. Genomics 33: 473-479, 1996.

[117324]20730.

[117325]Zhou, J.; Ding, M.; Zhao, Z.; Reenders, S. T.: Complete primary

[117326]structure of the sixth chain of human basement membrane collagen,

[117327]alpha-6(IV): isolation of the cDNAs for alpha-6(IV) and comparison

[117328]with five other type IV collagen chains. J. Biol. Chem. 269: 13193-13199,

[117329]1994.

[117330]20731.

[117331]Nathans, J.; Maumenee, I. H.; Zrenner, E.; Sadowski, B.; Sharpe,

[117332]L. T.; Lewis, R. A.; Hansen, E.; Rosenberg, T.; Schwartz, M.; Heckenlively,

[117333]J. R.; Traboulsi, E.; Klingaman, R.; Bech-Hansen, N. T.;
LaRoche,

[117334]G. R.; Pagon, R. A.; Murphey, W. H.; Weleber, R. G.: Genetic
heterogeneity

[117335]among blue-cone monochromats. Am. J. Hum. Genet. 53:
987-1000, 1993.

[117336]20732.

[117337]Bell, J.; Haldane, J. B. S.: The linkage between the genes
for

[117338]colour-blindness and haemophilia in man. Proc. Roy. Soc.
London 123B:

[117339]119-150, 1937.

[117340]20733.

[117341]Filippi, G.; Rinaldi, A.; Palmarino, R.; Seravalli, E.; Sinis-
calco,

[117342]M.: Linkage disequilibrium for two X-linked genes in Sar-
dinia and

[117343]its bearing on the statistical mapping of the human X
chromosome. Genetics 86:

[117344]199-222, 1977.

[117345]20734.

- [117346]Thompson, D. A.; McHenry, C. L.; Li, Y.; Richards, J. E.;
Othman,
[117347]M. I.; Schwinger, E.; Vollrath, D.; Jacobson, S. G.; Gal, A.:
Retinal
[117348]dystrophy due to paternal isodisomy for chromosome 1 or
chromosome
[117349]2, with homoallelism for mutations in RPE65 or MERTK,
respectively. Am.
[117350]J. Hum. Genet. 70: 224–229, 2002.
[117351]20735.
[117352]Aguirre, G. D.; Baldwin, V.; Weeks, K. M.; Acland, G. M.;
Ray,
[117353]K.: Frequency of the codon 807 mutation in the cGMP
phosphodiesterase
[117354]beta-subunit gene in Irish setters and other dog breeds
with hereditary
[117355]retinal degeneration. J. Hered. 90: 143–147, 1999.
[117356]20736.
[117357]Bateman, J. B.; Klisak, I.; Kojis, T.; Mohandas, T.; Sparkes,
R.
[117358]S.; Li, T.; Applebury, M. L.; Bowes, C.; Farber, D. B.: As-
signment

[117359] of the beta-subunit of rod photoreceptor cGMP phosphodiesterase gene

[117360] PDEB (homolog of the mouse rd gene) to human chromosome 4p16. Genomics 12:

[117361] 601-603, 1992.

[117362] 20737.

[117363] Bayes, M.; Giordano, M.; Balcells, S.; Grinberg, D.; Vigeliliu,

[117364] L.; Martinez, I.; Ayuso, C.; Benitez, J.; Ramos-Arroyo, M. A.; Chivelet,

[117365] P.; Solans, T.; Valverde, D.; Amselem, S.; Goossens, M.; Baiget, M.;

[117366] Gonzalez-Duarte, R.; Besmond, C.: Homozygous tandem duplication within

[117367] the gene encoding the beta-subunit of rod phosphodiesterase as a cause

[117368] for autosomal recessive retinitis pigmentosa. Hum. Mutat. 5: 228-234,

[117369] 1995.

[117370] 20738.

[117371] Bennett, J.; Tanabe, T.; Sun, D.; Zeng, Y.; Kjeldbye, H.; Gouras,

[117372]P.; Maguire, A. M.: Photoreceptor cell rescue in retinal degeneration

[117373](rd) mice by in vivo gene therapy. Nature Med. 2: 649, 1996.

[117374]20739.

[117375]Bowes, C.; Danciger, M.; Kozak, C. A.; Farber, D. B.: Isolation

[117376]of a candidate cDNA for the gene causing retinal degeneration in the

[117377]rd mouse. Proc. Nat. Acad. Sci. 86: 9722–9726, 1989.

Note: Erratum:

[117378]Proc. Nat. Acad. Sci 87:1625, 1990...

[117379]20740.

[117380]Bowes, C.; Li, T.; Danciger, M.; Baxter, L. C.; Applebury, M. L.;

[117381]Farber, D. B.: Retinal degeneration in the rd mouse is caused by

[117382]a defect in the beta subunit of rod cGMP–phosphodiesterase. Nature 347:

[117383]677–680, 1990.

[117384]20741.

- [117385]Bowes, C.; Li, T.; Frankel, W. N.; Danciger, M.; Coffin, J. M.;
- [117386]Applebury, M. L.; Farber, D. B.: Localization of a retroviral element within the rd gene coding for the beta subunit of cGMP phosphodiesterase. Proc.
- [117387]Nat. Acad. Sci. 90: 2955–2959, 1993.
- [117388]20742.
- [117389]Cachon–Gonzalez, M. B.; Fenner, S.; Coffin, J. M.; Moran, C.; Best, S.; Stoye, J. P.: Structure and expression of the hairless gene of mice. Proc. Nat. Acad. Sci. 91: 7717–7721, 1994.
- [117390]20743.
- [117391]Clements, P. J. M.; Gregory, C. Y.; Peterson–Jones, S. M.; Sargan, D. R.; Bhattacharya, S. S.: Confirmation of the rod cGMP phosphodiesterase beta subunit (PDE–beta) nonsense mutation in affected rcd–1 Irish setters in the UK and the development of a diagnostic test. Curr.

[117398]Eye Res. 12: 861–866, 1993.

[117399]20744.

[117400]Collins, C.; Hutchinson, G.; Kowbel, D.; Riess, O.; Weber, B.;

[117401]Hayden, M. R.: The human beta-subunit of rod photoreceptor cGMP phosphodiesterase:

[117402]complete retinal cDNA sequence and evidence for expression in brain. Genomics 13:

[117403]698–704, 1992.

[117404]20745.

[117405]Danciger, M.; Bowes, C.; Kozak, C. A.; LaVail, M. M.; Farber,

[117406]D. B.: Fine mapping of a putative rd cDNA and its cosegregation with

[117407]rd expression. Invest. Ophthalmol. Vis. Sci. 31: 1427–1432, 1990.

[117408]20746.

[117409]Farber, D. B.; Danciger, J. S.; Aguirre, G.: The beta subunit
[117410]of cyclic GMP phosphodiesterase mRNA is deficient in canine rod-cone

[117411]dysplasia 1. Neuron 9: 349–356, 1992.

[117412]20747.

[117413]Farber, D. B.; Lolley, R. N.: Enzyme basis for cyclic GMP
accumulation

[117414]in degenerative photoreceptor cells of mouse retina. J.
Cyclic Nucleotide

[117415]Res. 2: 139–148, 1976.

[117416]20748.

[117417]Farber, D. B.; Lolley, R. N.: Cyclic guanosine monophos-
phate:

[117418]elevation in degenerating photoreceptor cells of the C3H
mouse retina. Science 186:

[117419]449–451, 1974.

[117420]20749.

[117421]Gal, A.; Orth, U.; Baehr, W.; Schwinger, E.; Rosenberg, T.:
Heterozygous

[117422]missense mutation in the rod cGMP phosphodiesterase
beta-subunit gene

[117423]in autosomal dominant stationary night blindness. Nature
Genet. 7:

[117424]64–68, 1994. Note: Correction: Nature Genet. 7: 551 only,
1994...

[117425]20750.

[117426]Bogenmann, E.; Lochrie, M. A.; Simon, M. I.: Cone cell-specific

[117427]genes expressed in retinoblastoma. Science 240: 76-78, 1988.

[117428]20751.

[117429]Bonaiti-Pellie, C.; Briard-Guillemot, M. L.: Segregation analysis

[117430]in hereditary retinoblastoma. Hum. Genet. 57: 411-419, 1981.

[117431]20752.

[117432]Nagasaki, H.; Ito, M.; Yuasa, H.; Saito, H.; Fukase, M.; Hamada,

[117433]K.; Ishikawa, E.; Katakami, H.; Oiso, Y.: Two novel mutations in

[117434]the coding region for neurophysin-II associated with familial central

[117435]diabetes insipidus. J. Clin. Endocr. Metab. 80: 1352-1356, 1995.

[117436]20753.

[117437]Bartram, C. R.; de Klein, A.; Hagemeijer, A.; van Agthoven, T.;

[117438]Geurts van Kessel, A.; Bootsma, D.; Grosveld, G.; Ferguson-Smith,
son-Smith,
[117439]M. A.; Davies, T.; Stone, M.; Heisterkamp, N.; Stephenson,
J. R.;
[117440]Groffen, J.: Translocation of c-abl oncogene correlates
with the
[117441]presence of a Philadelphia chromosome in chronic myelo-
cytic leukaemia. Nature 306:
[117442]277-280, 1983.
[117443]20754.
[117444]Rittig, S.; Robertson, G. L.; Siggaard, C.; Kovacs, L.;
Gregersen,
[117445]N.; Nyborg, J.; Pedersen, E. B.: Identification of 13 new
mutations
[117446]in the vasopressin-neurophysin II gene in 17 kindreds
with familial
[117447]autosomal dominant neurohypophyseal diabetes in-
sipidus. Am. J. Hum.
[117448]Genet. 58: 107-117, 1996.
[117449]20755.
[117450]Russell, J. T.; Brownstein, M. J.; Gainer, H.: Trypsin liber-
ates

[117451]an arginine vasopressin-like peptide and neurophysin
from a M(r)20,000

[117452]putative common precursor. Proc. Nat. Acad. Sci. 76:
6086-6090,

[117453]1979.

[117454]20756.

[117455]Sachs, H.; Fawcett, P.; Takabatake, Y.; Portanova, R.:
Biosynthesis

[117456]and release of vasopressin and neurophysin. Recent Prog.
Horm. Res. 25:

[117457]447-491, 1969.

[117458]20757.

[117459]O'Brien, R. O.; Taske, N. L.; Hansbro, P. M.; Matthaei, K. I.;

[117460]Hogan, S. P.; Denborough, M. A.; Foster, P. S.: Exclusion of
defects

[117461]in the skeletal muscle specific regions of the DHPR alpha-
1 subunit

[117462]as frequent causes of malignant hyperthermia. J. Med.
Genet. 32:

[117463]913-914, 1995.

[117464]20758.

[117465]Wichterle, H.; Hanspal, M.; Palek, J.; Jarolim, P.: Combination
[117466]of two mutant alpha spectrin alleles underlies a severe
[117467]spherocytic
[117467]hemolytic anemia. J. Clin. Invest. 98: 2300–2307, 1996.
[117468]20759.
[117469]Nijenhuis, M.; van den Akker, E. L. T.; Zalm, R.; Franken,
A.
[117470]A. M.; Abbes, A. P.; Engel, H.; de Wied, D.; Burbach, J. P.
H.: Familial
[117471]neurohypophysial diabetes insipidus in a large Dutch kin-
dred: effect
[117472]of the onset of diabetes on growth in children and cell bi-
ological
[117473]defects of the mutant vasopressin prohormone. J. Clin.
Endocr. Metab. 86:
[117474]3410–3420, 2001.
[117475]20760.
[117476]Nijenhuis, M.; Zalm, R.; Burbach, J. P. H.: Mutations in the
[117477]vasopressin prohormone involved in diabetes insipidus
impair endoplasmic

[117478]reticulum export but not sorting. J. Biol. Chem. 274:
21200–21208,

[117479]1999.

[117480]20761.

[117481]Olias, G.; Richter, D.; Schmale, H.: Heterologous expres-
sion

[117482]of human vasopressin–neurophysin precursors in a pitu-
itary cell line:

[117483]defective transport of a mutant protein from patients with
familial

[117484]diabetes insipidus. DNA Cell Biol. 15: 929–935, 1996.

[117485]20762.

[117486]Repaske, D. R.; Summar, M. L.; Krishnamani, M. R. S.; Gul-
tekin,

[117487]E. K.; Arriazu, M. C.; Roubicek, M. E.; Blanco, M.; Isaac, G.
B.;

[117488]Phillips, J. A., III: Recurrent mutations in the vasopressin–
neurophysin

[117489]II gene cause autosomal dominant neurohypophyseal dia-
betes insipidus. J.

[117490]Clin. Endocr. Metab. 81: 2328–2334, 1996.

[117491]20763.

[117492]Riddell, D. C.; Mallonee, R.; Phillips, J. A.; Parks, J. S.; Sex-
ton,

[117493]L. A.; Hamerton, J. L.: Chromosomal assignment of human
sequences

[117494]encoding arginine vasopressin-neurophysin II and growth
hormone releasing

[117495]factor. Somat. Cell Molec. Genet. 11: 189-195, 1985.

[117496]20764.

[117497]Repaske, D. R.; Medlej, R.; Gultekin, E. K.; Krishnamani, M.
R.

[117498]S.; Halaby, G.; Findling, J. W.; Phillips, J. A., III: Hetero-
geneity

[117499]in clinical manifestation of autosomal dominant neurohy-
popyseal diabetes

[117500]insipidus caused by a mutation encoding Ala-1 Val in the
signal peptide

[117501]of the arginine vasopressin/neurophysin II/copeptin pre-
cursor. J.

[117502]Clin. Endocr. Metab. 82: 51-56, 1997.

[117503]20765.

[117504]Rutishauser, J.; Boni-Schnetzler, M.; Boni, J.; Wichmann, W.;

[117505]Huisman, T.; Vallotton, M. B.; Froesch, E. R.: A novel point mutation

[117506]in the translation initiation codon of the pre-pro-vasopressin-neurophysin

[117507]II gene: Cosegregation with morphological abnormalities and clinical

[117508]symptoms in autosomal dominant neurohypophyseal diabetes. J. Clin.

[117509]Endocr. Metab. 81: 192-198, 1996.

[117510]20766.

[117511]Sausville, E.; Carney, D.; Battey, J.: The human vasopressin

[117512]gene is linked to the oxytocin gene and is selectively expressed in

[117513]a cultured lung cancer cell line. J. Biol. Chem. 260: 10236-10241,

[117514]1985.

[117515]20767.

[117516]Scherbaum, W. A.; Bottazzo, G. F.: Autoantibodies to vasopressin

[117517]cells in idiopathic diabetes insipidus: evidence for an autoimmune

[117518]variant. *Lancet* I: 897–901, 1983.

[117519]20768.

[117520]Siggaard, C.; Rittig, S.; Corydon, T. J.; Andreasen, P. H.; Jensen,

[117521]T. G.; Andresen, B. S.; Robertson, G. L.; Gregersen, N.; Bolund, L.;

[117522]Pedersen, E. B.: Clinical and molecular evidence of abnormal processing

[117523]and trafficking of the vasopressin preprohormone in a large kindred

[117524]with familial neurohypophyseal diabetes insipidus due to a signal

[117525]peptide mutation. *J. Clin. Endocr. Metab.* 84: 2933–2941, 1999.

[117526]20769.

[117527]Summar, M. L.; Phillips, J. A., III; Battey, J.; Castiglione,

[117528]C. M.; Kidd, K. K.; Maness, K. J.; Weiffenbach, B.; Gravius, T. C.

[117529]: Linkage relationships of human arginine vasopressin-neurophysin-II

[117530]and oxytocin–neurophysin–I to prodynorphin and other
loci on chromosome

[117531]20. Molec. Endocr. 4: 947–950, 1990.

[117532]20770.

[117533]Sunde, D. A.; Sokol, H. W.: Quantification of rat neuro-
physins

[117534]by polyacrylamide gel electrophoresis (page): application
to the rat

[117535]with hereditary hypothalamic diabetes insipidus. Ann. N.Y.
Acad.

[117536]Sci. 248: 345–364, 1975.

[117537]20771.

[117538]Ueta, Y.; Taniguchi, S.; Yoshida, A.; Murakami, I.; Mitani,
Y.;

[117539]Hisatome, I.; Manabe, I.; Sato, R.; Tsuboi, M.; Ohtahara, A.;
Nanba,

[117540]E.; Shigemasa, C.: A new type of familial central diabetes
insipidus

[117541]caused by a single base substitution in the neurophysin II
coding

[117542]region of the vasopressin gene. J. Clin. Endocr. Metab. 81:
1787–1790,

[117543]1996.

[117544]20772.

[117545]Valtin, H.; Sawyer, W. H.; Sokol, H. W.: Neurohypophyseal principles

[117546]in rats homozygous and heterozygous for hypothalamic diabetes insipidus

[117547](Brattleboro strain). Endocrinology 77: 701-706, 1965.

[117548]20773.

[117549]van Leeuwen, F. W.; de Kleijn, D. P. V.; van den Hurk, H. H.;

[117550]Neubauer, A.; Sonnemans, M. A. F.; Sluijs, J. A.; Koycu, S.; Ramdjielal,

[117551]R. D. J.; Salehi, A.; Martens, G. J. M.; Grosveld, F. G.; Burbach,

[117552]J. P. H.; Hol, E. M.: Frameshift mutants of beta-amyloid precursor

[117553]protein and ubiquitin-B in Alzheimer's and Down patients. Science 279:

[117554]242-247, 1998.

[117555]20774.

[117556]Yuasa, H.; Ito, M.; Nagasaki, H.; Oiso, Y.; Miyamoto, S.;

Sasaki,

[117557]N.; Saito, H.: Glu-47, which forms a salt bridge between neurophysin-II

[117558]and arginine vasopressin, is deleted in patients with familial central

[117559]diabetes insipidus. J. Clin. Endocr. Metab. 77: 600-604, 1993.

[117560]20775.

[117561]Oesterling, J. E.; Chan, D. W.; Epstein, J. I.; Kimball, A. W.,

[117562]Jr.; Bruzek, D. J.; Rock, R. C.; Brendler, C. B.; Walsh, P. C.: Prostate

[117563]specific antigen in the preoperative and postoperative evaluation

[117564]of localized prostatic cancer treated with radical prostatectomy. J.

[117565]Urol. 139: 766-772, 1988.

[117566]20776.

[117567]Riegman, P. H. J.; Vlietstra, R. J.; van der Korput, J. A. G. M.;

[117568]Brinkmann, A. O.; Trapman, J.: The promoter of the prostate-specific

[117569]antigen gene contains a functional androgen responsive

element. Molec.

[117570]Endocr. 5: 1921–1930, 1991.

[117571]20777.

[117572]van den Elsen, P.; Bruns, G.; Gerhard, D. S.; Pravtcheva, D.;

[117573]Jones, C.; Housman, D.; Ruddle, F. A.; Orkin, S.; Terhorst,

C.: Assignment

[117574]of the gene coding for the T3–delta subunit of the

T3–T-cell receptor

[117575]complex to the long arm of human chromosome 11 and

to mouse chromosome

[117576]9. Proc. Nat. Acad. Sci. 82: 2920–2924, 1985.

[117577]20778.

[117578]van Rijs, J.; Giguere, V.; Hurst, J.; van Agthoven, T.; Geurts

[117579]van Kessel, A.; Goyert, S.; Grosveld, F.: Chromosomal lo-

calization

[117580]of the human Thy–1 gene. Proc. Nat. Acad. Sci. 82:

5832–5835, 1985.

[117581]20779.

[117582]Williams, A. F.; Gagnon, J.: Neuronal cell Thy–1 glycopro-

tein:

[117583]homology with immunoglobulin. Science 216: 696–703,

1982.

[117584]20780.

[117585]Johansson, M.; Karlsson, A.: Cloning of the cDNA and
chromosome

[117586]localization of the gene for human thymidine kinase 2. J.
Biol. Chem. 272:

[117587]8454–8458, 1997.

[117588]20781.

[117589]Mandel, H.; Szargel, R.; Labay, V.; Elpeleg, O.; Saada, A.;
Shalata,

[117590]A.; Anbinder, Y.; Berkowitz, D.; Hartman, C.; Barak, M.;
Eriksson,

[117591]S.; Cohen, N.: The deoxyguanosine kinase gene is mu-
tated in individuals

[117592]with depleted hepatocerebral mitochondrial DNA. Nature
Genet. 29:

[117593]337–341, 2001. Note: Erratum: Nature Genet. 29: 491
only, 2001.

[117594]20782.

[117595]Saada, A.; Shaag, A.; Mandel, H.; Nevo, Y.; Eriksson, S.;
Elpeleg,

[117596]O.: Mutant mitochondrial thymidine kinase in mitochon-

drial DNA depletion

[117597]myopathy. Nature Genet. 29: 342–344, 2001.

[117598]20783.

[117599]Wang, L.; Munch–Petersen, B.; Herrstrom Sjoberg, A.; Hell–
man, U.;

[117600]Bergman, T.; Jornvall, H.; Eriksson, S.: Human thymidine
kinase 2:

[117601]molecular cloning and characterisation of the enzyme ac–
tivity with

[117602]antiviral and cytostatic nucleoside substrates. FEBS Lett.
443:

[117603]170–174, 1999.

[117604]20784.

[117605]Willecke, K.; Reuber, T.; Kucherlapati, R. S.; Ruddle, F. H.:

[117606]Human mitochondrial thymidine kinase is coded for by a
gene on chromosome

[117607]16 of the nucleus. Somat. Cell Genet. 3: 237–245, 1977.

[117608]20785.

[117609]Klobutcher, L. A.; Ruddle, F. H.: Phenotype stabilisation
and

[117610]integration of transferred material in chromosome–medi–

ated gene transfer. Nature 280:

[117611]657-660, 1979.

[117612]20786.

[117613]Kuo, W.-L.; Hirschhorn, R.; Huie, M. L.; Hirschhorn, K.: Localization

[117614]and ordering of acid alpha-glucosidase (GAA) and thymidine kinase

[117615](TK1) by fluorescence in situ hybridization. Hum. Genet. 97: 404-406,

[117616]1996.

[117617]20787.

[117618]Chang, J. G.; Scarpa, A.; Eddy, R. L.; Byers, M. G.; Harris, A.

[117619]S.; Morrow, J. S.; Watkins, P.; Shows, T. B.; Forget, B. G.: Cloning

[117620]of a portion of the chromosomal gene and cDNA for human beta-fodrin,

[117621]the nonerythroid form of beta-spectrin. Genomics 17: 287-293, 1993.

[117622]20788.

[117623]Watkins, P. C.; Eddy, R.; Forget, B. G.; Chang, J. G.; Rochelle,

[117624]R.; Shows, T. B.: Assignment of a non-erythroid spectrin gene to

[117625]human chromosome 2. (Abstract) Am. J. Hum. Genet. 43: A161, 1988.

[117626]20789.

[117627]Birkenmeier, C. S.; McFarland-Starr, E. C.; Barker, J. E.: Chromosomal

[117628]location of three spectrin genes: relationship to the inherited hemolytic

[117629]anemias of mouse and man. Proc. Nat. Acad. Sci. 85: 8121-8125, 1988.

[117630]20790.

[117631]Agre, P.; Asimos, A.; Casella, J. F.; McMillan, C.: Inheritance

[117632]pattern and clinical response to splenectomy as a reflection of erythrocyte

[117633]spectrin deficiency in hereditary spherocytosis. New Eng. J. Med. 315:

[117634]1579-1583, 1986.

[117635]20791.

[117636]Alloisio, N.; Guetarni, D.; Morle, L.; Pothier, B.; Ducluzeau,

[117637]M. T.; Soun, A.; Colonna, P.; Clerc, M.; Philippe, N.; Delaunay, J.

[117638]: SP-alpha(I/65) hereditary elliptocytosis in North Africa.
Am. J.

[117639]Hemat. 23: 113-122, 1986.

[117640]20792.

[117641]Alloisio, N.; Morle, L.; Pothier, B.; Roux, A.-F.; Marechal, J.;

[117642]Ducluzeau, M.-T.; Benhadji-Zouaoui, Z.; Delaunay, J.:
Spectrin Oran

[117643](alpha II/21), a new spectrin variant concerning the alpha
II domain

[117644]and causing severe elliptocytosis in the homozygous
state. Blood 71:

[117645]1039-1047, 1988.

[117646]20793.

[117647]Alloisio, N.; Wilmotte, R.; Marechal, J.; Texier, P.; Denoroy,

[117648]L.; Feo, C.; Benhadji-Zouaoui, Z.; Delaunay, J.: A splice site
mutation

[117649]of alpha-spectrin gene causing skipping of exon 18 in
hereditary elliptocytosis. Blood 81:

[117650]2791-2798, 1993.

[117651]20794.

[117652]Alloisio, N.; Wilmotte, R.; Morle, L.; Baklouti, F.; Marechal,

[117653]J.; Ducluzeau, M.-T.; Denoroy, L.; Feo, C.; Forget, B. G.;

Kastally,

[117654]R.; Delaunay, J.: Spectrin Jendouba: an alpha-II/31 spectrin variant

[117655]that is associated with elliptocytosis and carries a mutation distant

[117656]from the dimer self-association site. Blood 80: 809-815, 1992.

[117657]20795.

[117658]Baklouti, F.; Marechal, J.; Morle, L.; Alloisio, N.; Wilmotte,

[117659]R.; Pothier, B.; Ducluzeau, M.-T.; Kastally, R.; Delaunay, J.:

Occurrence

[117660]of the alpha-I 22 arg-to-his (CGT-to-CAT) spectrin mutation in Tunisia:

[117661]potential association with severe elliptopoikilocytosis. Brit. J.

[117662]Haemat. 78: 108-113, 1991.

[117663]20796.

[117664]Boulanger, L.; LeCompte, M.-C.; Dhermy, D.; Garbarz, M.:

Fast

[117665]screening methods to detect mutations of spectrin in subjects with

[117666]hereditary elliptocytosis. (Letter) Am. J. Hum. Genet. 51: 440–442,

[117667]1992.

[117668]20797.

[117669]Burke, J. P. W. G.; Van Zyl, D.; Zail, S. S.; Coetzer, T. L.:

[117670]Reduced spectrin–ankyrin binding in a South African hereditary elliptocytosis

[117671]kindred homozygous for spectrin St Claude. (Letter) Blood 92: 2591–2592,

[117672]1998.

[117673]20798.

[117674]Coetzer, T.; Zail, S.: Spectrin tetramer–dimer equilibrium in

[117675]hereditary elliptocytosis. Blood 59: 900–905, 1982.

[117676]20799.

[117677]Coetzer, T. L.; Sahr, K.; Prchal, J.; Blacklock, H.; Peterson,

[117678]L.; Koler, R.; Doyle, J.; Manaster, J.; Palek, J.: Four different

[117679] mutations in codon 28 of alpha spectrin are associated with structurally

[117680] and functionally abnormal spectrin alpha(I/74) in hereditary elliptocytosis. J.

[117681] Clin. Invest. 88: 743-749, 1991.

[117682] 20800.

[117683] Dalla Venezia, N.; Wilmotte, R.; Morle, L.; Forissier, A.; Parquet,

[117684] N.; Garbarz, M.; Rousset, T.; Dhermy, D.; Alloisio, N.; Delaunay,

[117685] J.: An alpha-spectrin mutation responsible for hereditary elliptocytosis

[117686] associated in cis with the alpha-V/41 polymorphism. Hum. Genet. 90:

[117687] 641-644, 1993.

[117688] 20801.

[117689] Delaunay, J.; Dhermy, D.: Mutations involving the spectrin heterodimer

[117690] contact site: clinical expression and alterations in specific function. Seminars

[117691] Hemat. 30: 21-33, 1993.

[117692]20802.

[117693]del Giudice, E. M.; Ducluzeau, M. T.; Alloisio, N.; Wilmotte,

[117694]R.; Delaunay, J.; Perrotta, S.; Cutillo, S.; Iolascon, A.: Al-
pha(I/65)

[117695]hereditary elliptocytosis in southern Italy: evidence for an
African

[117696]origin. Hum. Genet. 89: 553-556, 1992.

[117697]20803.

[117698]Dhermy, D.; Lecomte, M. C.; Garbarz, M.; Bournier, O.;
Galand,

[117699]C.; Gautero, H.; Feo, C.; Alloisio, N.; Delaunay, J.; Boivin,
P.:

[117700]Spectrin beta-chain variant associated with hereditary el-
liptocytosis. J.

[117701]Clin. Invest. 70: 707-715, 1982.

[117702]20804.

[117703]Elgsaeter, A.; Stokke, B. T.; Mikkelsen, A.; Branton, D.: The

[117704]molecular basis of erythrocyte shape. Science 234:
1217-1223, 1986.

[117705]20805.

[117706]Miceli-Richard, C.; Lesage, S.; Rybojad, M.; Prieur, A.-M.;
Manouvrier-Hanu,

[117707]S.; Hafner, R.; Chamailard, M.; Zouali, H.; Thomas, G.;
Hugot, J.-P.

[117708]: CARD15 mutations in Blau syndrome. *Nature Genet.* 29:
19-20, 2001.

[117709]20806.

[117710]Bernards, A.; Rubin, C. M.; Westbrook, C. A.; Paskind, M.;
Baltimore,

[117711]D.: The first intron in the human c-abl gene is at least 200
kilobases

[117712]long and is a target for translocations in chronic myeloge-
nous leukemia. *Molec.*

[117713]*Cell. Biol.* 7: 3231-3236, 1987.

[117714]20807.

[117715]Clark, S. S.; McLaughlin, J.; Crist, W. M.; Champlin, R.;
Witte,

[117716]O. N.: Unique forms of the abl tyrosine kinase distinguish
Ph-1-positive

[117717]CML from Ph-1-positive ALL. *Science* 235: 85-88, 1986.

[117718]20808.

[117719]Collins, S. J.; Groudine, M. T.: Rearrangement and amplification

[117720]of c-abl sequences in the human chronic myelogenous leukemia cell

[117721]line K-562. Proc. Nat. Acad. Sci. 80: 4813-4817, 1983.

[117722]20809.

[117723]Cong, F.; Spencer, S.; Cote, J.-F.; Wu, Y.; Tremblay, M. L.; Lasky,

[117724]L. A.; Goff, S. P.: Cytoskeletal protein PSTPIP1 directs the PEST-type

[117725]protein tyrosine phosphatase to the c-Abl kinase to mediate Abl dephosphorylation. Molec.

[117726]Cell 6: 1413-1423, 2000.

[117727]20810.

[117728]Fainstein, E.; Marcelle, C.; Rosner, A.; Canaani, E.; Gale, R.

[117729]P.; Dreazen, O.; Smith, S. D.; Croce, C. M.: A new fused transcript

[117730]in Philadelphia chromosome positive acute lymphocytic leukaemia. Nature 330:

[117731]386-388, 1987.

[117732]20811.

[117733]Goff, S. P.; D'Eustachio, P.; Ruddle, F. H.; Baltimore, D.:
Chromosomal

[117734]assignment of endogenous protooncogene c-abl. Science
218: 1317-1319,

[117735]1982.

[117736]20812.

[117737]Gong, J.; Costanzo, A.; Yang, H.-Q.; Melino, G.; Kaelin, W.
G.,

[117738]Jr.; Levvero, M.; Wang, J. Y. J.: The tyrosine kinase c-Abl
regulates

[117739]p73 in apoptotic response to cisplatin-induced DNA dam-
age. Nature 399:

[117740]806-809, 1999.

[117741]20813.

[117742]Han, L. Wong, D.; Dhaka, A.; Afar, D.; White, M.; Xie, W.;
Herschman,

[117743]H.; Witte, O.; Colicelli, J.: Protein binding and signaling
properties

[117744]of RIN1 suggest a unique effector function. Proc. Nat.
Acad. Sci. 94:

[117745]4954-4959, 1997.

[117746]20814.

[117747]Heisterkamp, N.; Groffen, J.; Stephenson, J. R.; Spurr, N.
K.;

[117748]Goodfellow, P. N.; Solomon, E.; Carritt, B.; Bodmer, W. F.:
Chromosomal

[117749]localization of human cellular homologues of two viral
oncogenes. Nature 299:

[117750]747–749, 1982.

[117751]20815.

[117752]Heisterkamp, N.; Stephenson, J. R.; Groffen, J.; Hansen, P.
F.;

[117753]de Klein, A.; Bartram, C. R.; Grosveld, G.: Localization of
the c-abl

[117754]oncogene adjacent to a translocation breakpoint in
chronic myelocytic

[117755]leukaemia. Nature 306: 239–242, 1983.

[117756]20816.

[117757]Hoffman–Falk, H.; Einat, P.; Shilo, B.–Z.; Hoffmann, F. M.:
Drosophila

[117758]melanogaster DNA clones homologous to vertebrate
oncogenes: evidence

[117759] for a common ancestor to the src and abl cellular genes.

Cell 32:

[117760] 589–598, 1983.

[117761] 20817.

[117762] Huntly, B. J. P.; Reid, A. G.; Bench, A. J.; Campbell, L. J.;

[117763] Telford, N.; Shepherd, P.; Szer, J.; Prince, H. M.; Turner, P.;

Grace,

[117764] C.; Nacheva, E. P.; Green, A. R.: Deletions of the derivative
chromosome

[117765] 9 occur at the time of the Philadelphia translocation and
provide

[117766] a powerful and independent prognostic indicator in
chronic myeloid

[117767] leukemia. Blood 98: 1732–1738, 2001.

[117768] 20818.

[117769] Jhanwar, S. C.; Neel, B. G.; Hayward, W. S.; Chaganti, R. S.
K.

[117770]: Localization of the cellular oncogenes ABL, SIS, and FES
on human

[117771] germ-line chromosomes. Cytogenet. Cell Genet. 38:
73–75, 1984.

[117772]20819.

[117773]Konopka, J. B.; Watanabe, S. M.; Singer, J. W.; Collins, S. J.;

[117774]Witte, O. N.: Cell lines and clinical isolates derived from
Ph1-positive

[117775]chronic myelogenous leukemia patients express c-abl
proteins with

[117776]a common structural alteration. Proc. Nat. Acad. Sci. 82:
1810-1814,

[117777]1985.

[117778]20820.

[117779]Kurzrock, R.; Shtalrid, M.; Romero, P.; Kloetzer, W. S.; Tal-
pas,

[117780]M.; Trujillo, J. M.; Blick, M.; Beran, M.; Gutterman, J. U.: A
novel

[117781]c-abl protein product in Philadelphia-positive acute lym-
phoblastic

[117782]leukaemia. Nature 325: 631-635, 1987.

[117783]20821.

[117784]Lane, M.-A.; Neary, D.; Cooper, G. M.: Activation of a cel-
lular

[117785]transforming gene in tumours induced by Abelson murine

leukaemia virus. Nature 300:

[117786]659–661, 1982.

[117787]20822.

[117788]Li, B.; Boast, S.; de los Santos, K.; Schieren, I.; Quiroz, M.;

[117789]Teitelbaum, S. L.; Tondravi, M. M.; Goff, S. P.: Mice deficient in

[117790]Abl are osteoporotic and have defects in osteoblast maturation. Nature

[117791]Genet. 24: 304–308, 2000.

[117792]20823.

[117793]Melo, J. V.; Gordon, D. E.; Cross, N. C. P.; Goldman, J. M.:

[117794]The ABL–BCR fusion gene is expressed in chronic myeloid leukemia. Blood 81:

[117795]158–165, 1993.

[117796]20824.

[117797]Pluk, H.; Dorey, K.; Superti-Furga, G.: Autoinhibition of c–Abl. Cell 108:

[117798]247–259, 2002.

[117799]20825.

[117800]Schindler, T.; Bornmann, W.; Pellicena, P.; Miller, W. T.;
Clarkson,

[117801]B.; Kuriyan, J.: Structural mechanism for STI-571 inhibition
of Abelson

[117802]tyrosine kinase. Science 289: 1938–1941, 2000.

[117803]20826.

[117804]Schwartzberg, P. L.; Stall, A. M.; Hardin, J. D.; Bowdish, K.

[117805]S.; Humaran, T.; Boast, S.; Harbison, M. L.; Robertson, E.
J.; Goff,

[117806]S. P.: Mice homozygous for the abl(m1) mutation show
poor viability

[117807]and depletion of selected B and T cell populations. Cell
65: 1165–1175,

[117808]1991.

[117809]20827.

[117810]Scott, M. L.; Van Etten, R. A.; Daley, G. Q.; Baltimore, D.:

[117811]v-abl causes hematopoietic disease distinct from that
caused by bcr-abl. Proc.

[117812]Nat. Acad. Sci. 88: 6506–6510, 1991.

[117813]20828.

- [117814]Bouhassira, E. E.; Schwartz, R. S.; Yawata, Y.; Ata, K.;
Nagel,
[117815]R. L.; Rybicki, A. C.: An alanine to threonine substitution
in protein
[117816]4.2 cDNA is associated with a Japanese form of hereditary
hemolytic
[117817]anemia. (Abstract) Clin. Res. 39: 313A, 1991.
[117818]20829.
[117819]Davies, K. A.; Lux, S. E.: Hereditary disorders of the red
cell
[117820]membrane skeleton. Trends Genet. 5: 222–227, 1989.
[117821]20830.
[117822]Ghanem, A.; Pothier, B.; Marechal, J.; Ducluzeau, M. T.;
Morle,
[117823]L.; Alloisio, N.; Feo, C.; Ben Abdeladhim, A.; Fattoum, S.;
Delaunay,
[117824]J.: A haemolytic syndrome associated with the complete
absence of
[117825]red cell membrane protein 4.2 in two Tunisian siblings.
Brit. J.
[117826]Haemat. 75: 414–420, 1990.

[117827]20831.

[117828]Grenard, P.; Bates, M. K.; Aeschlimann, D.: Evolution of
transglutaminase

[117829]genes: identification of a transglutaminase gene cluster
on human

[117830]chromosome 15q15: structure of the gene encoding
transglutaminase

[117831]X and a novel gene family member, transglutaminase Z. J.
Biol. Chem. 276:

[117832]33066–33078, 2001.

[117833]20832.

[117834]Gwynn, B.; Korsgren, C.; Cohen, C. M.; Ciciotte, S. L.; Pe-
ters,

[117835]L. L.: The gene encoding protein 4.2 is distinct from the
mouse platelet

[117836]storage pool deficiency mutation pallid. Genomics 42:
532–535, 1997.

[117837]20833.

[117838]Hayashi, S.; Koomoto, R.; Yano, A.; Ishigami, S.; Tsujino,
G.;

[117839]Saeki, S.; Tanaka, T.: Abnormality in a specific protein of
the erythrocyte

[117840]membrane in hereditary spherocytosis. Biochem. Biophys.
Res. Commun. 57:

[117841]1038–1044, 1974.

[117842]20834.

[117843]Hayette, S.; Dhermy, D.; dos Santos, M.–E.; Bozon, M.;
Drenckhahn,

[117844]D.; Alloisio, N.; Texier, P.; Delaunay, J.; Morle, L.: A dele-
tional

[117845]frameshift mutation in protein 4.2 gene (allele 4.2 Lisboa)
associated

[117846]with hereditary hemolytic anemia. Blood 85: 250–256,
1995.

[117847]20835.

[117848]Hayette, S.; Morle, L.; Bozon, M.; Ghanem, A.; Risinger, M.;
Korsgren,

[117849]C.; Tanner, M. J. A.; Fattoum, S.; Cohen, C. M.; Delaunay,
J.: A

[117850]point mutation in the protein 4.2 gene (allele 4.2 Tozeur)
associated

[117851]with hereditary haemolytic anaemia. Brit. J. Haemat. 89:
762–770,

[117852]1995.

[117853]20836.

[117854]Ideguchi, H.; Nishimura, J.; Nawata, H.; Hamasaki, N.: A
genetic

[117855]defect of erythrocyte band 4.2 protein associated with
hereditary

[117856]spherocytosis. Brit. J. Haemat. 74: 347–353, 1990.

[117857]20837.

[117858]Iwamoto, S.; Kajii, E.; Omi, T.; Kamesaki, T.; Akifuji, Y.;
Ikemoto,

[117859]S.: Point mutation in the band 4.2 gene associated with
autosomal

[117860]recessively inherited erythrocyte band 4.2 deficiency. Eu-
rop. J.

[117861]Haemat. 50: 286–291, 1993.

[117862]20838.

[117863]Korsgren, C.; Cohen, C. M.: Organization of the gene for
human

[117864]erythrocyte membrane protein 4.2: structural similarities
with the

[117865]gene for the A subunit of factor XIII. Proc. Nat. Acad. Sci.
88:

[117866]4840-4844, 1991.

[117867]20839.

[117868]Korsgren, C.; Cohen, C. M.: cDNA sequence, gene sequence, and

[117869]properties of murine pallidin (band 4.2), the protein implicated in

[117870]the murine pallid mutation. Genomics 21: 478-485, 1994.

[117871]20840.

[117872]Korsgren, C.; Lawler, J.; Lambert, S.; Speicher, D.; Cohen, C.

[117873]M.: Complete amino acid sequence and homologies of human erythrocyte

[117874]membrane protein band 4.2. Proc. Nat. Acad. Sci. 87: 613-617, 1990.

[117875]20841.

[117876]Matsuda, M.; Hatano, N.; Ideguchi, H.; Takahira, H.; Fukumaki,

[117877]Y.: A novel mutation causing an aberrant splicing in the protein

[117878]4.2 gene associated with hereditary spherocytosis (protein 4.2-Notame). Hum.

[117879]Molec. Genet. 4: 1187–1191, 1995.

[117880]20842.

[117881]Najfeld, V.; Ballard, S. G.; Menninger, J.; Ward, D. C.;
Bouhassira,

[117882]E. E.; Schwartz, R. S.; Nagel, R. L.; Rybicki, A. C.: The gene
for

[117883]human erythrocyte protein 4.2 maps to chromosome
15q15. Am. J. Hum.

[117884]Genet. 50: 71–75, 1992.

[117885]20843.

[117886]Nozawa, Y.; Hogushi, T.; Iida, H.; Fukushima, H.; Sekiya,
T.;

[117887]Ito, Y.: Erythrocyte membranes in hereditary spherocyto-
sis: alterations

[117888]in surface ultrastructure and membrane proteins as in-
ferred by scanning

[117889]electron microscopy and SDS gel electrophoresis. Clin.
Chim. Acta 55:

[117890]81–85, 1974.

[117891]20844.

[117892]Perrotta, S.; Iolascon, A.; Polito, R.; d'Urzo, G.; Conte, M.

[117893]L.; Miraglia del Giudice, E.: 4.2 Nippon mutation in a non-Japanese

[117894]patient with hereditary spherocytosis. (Letter) Haematologica 84:

[117895]660-662, 1999.

[117896]20845.

[117897]Rybicki, A. C.; Heath, R.; Wolf, J. L.; Lubin, B.; Schwartz, R.

[117898]S.: Deficiency of protein 4.2 in erythrocytes from a patient with

[117899]a Coombs negative hemolytic anemia: evidence for a role of protein

[117900]4.2 in stabilizing ankyrin on the membrane. J. Clin. Invest. 81:

[117901]893-901, 1988.

[117902]20846.

[117903]Sung, L. A.; Chien, S.; Chang, L.-S.; Lambert, K.; Bliss, S. A.;

[117904]Bouhassira, E. E.; Nagel, R. L.; Schwartz, R. S.; Rybicki, A. C.:

[117905]Molecular cloning of human protein 4.2: a major component of the erythrocyte

[117906]membrane. Proc. Nat. Acad. Sci. 87: 955–959, 1990.

[117907]20847.

[117908]Sung, L. A.; Fan, Y. S.; Lambert, K.; Chien, S.; Lin, C.: Map-
ping

[117909]of gene for human erythrocyte protein 4.2 to chromoso-
mal region 15q15–q21.

[117910](Abstract) Cytogenet. Cell Genet. 58: 1996, 1991.

[117911]20848.

[117912]White, R. A.; Dowler, L. L.; Hummel, G. S.; Adkison, L. R.:
Exclusion

[117913]of Ebp4.2 as a candidate for the mouse mutant pallid.

Mouse News

[117914]Letter 95: 492–494, 1997.

[117915]20849.

[117916]White, R. A.; Peters, L. L.; Adkison, L. R.; Korsgren, C.; Co-
hen,

[117917]C. M.; Lux, S. E.: The murine pallid mutation is a platelet
storage

[117918]pool disease associated with the protein 4.2 (pallidin)
gene. Nature

[117919]Genet. 2: 80–83, 1992.

[117920]20850.

[117921]Blank, V.; Andrews, N. C.: The Maf transcription factors:
regulators

[117922]of differentiation. Trends Biochem. Sci. 22: 437–441,
1997.

[117923]20851.

[117924]Jamieson, R. V.; Perveen, R.; Kerr, B.; Carette, M.; Yardley,
J.;

[117925]Heon, E.; Wirth, M. G.; van Heyningen, V.; Donnai, D.; Mu-
nier, F.;

[117926]Black, G. C. M.: Domain disruption and mutation of the
bZIP transcription

[117927]factor, MAF, associated with cataract, ocular anterior seg-
ment dysgenesis

[117928]and coloboma. Hum. Molec. Genet. 11: 33–42, 2002.

[117929]20852.

[117930]Kim, J. I.; Li, T.; Ho, I. C.; Grusby, M. J.; Glimcher, L. H.:

[117931]Requirement for the c-Maf transcription factor in crystallin
gene

[117932]regulation and lens development. Proc. Nat. Acad. Sci. 96:
3781–3785,

[117933]1999.

[117934]20853.

[117935]Nishizawa, M.; Kataoka, K.; Goto, N.; Fujiwara, K. T.;
Kawai, S.

[117936]: v-maf, a viral oncogene that encodes a 'leucine zipper'
motif. Proc.

[117937]Nat. Acad. Sci. 86: 7711-7715, 1989.

[117938]20854.

[117939]Yoshida, M. C.; Nishizawa, M.; Kataoka, K.; Goto, N.; Fuji-
wara,

[117940]K. T.; Kawai, S.: Localization of the human MAF pro-
tooncogene on

[117941]chromosome 16 to bands q22-q23. (Abstract) Cytogenet.
Cell Genet. 58:

[117942]2003 only, 1991.

[117943]20855.

[117944]Harbers, K.; Kuehn, M.; Delius, H.; Jaenisch, R.: Insertion
of

[117945]retrovirus into the first intron of alpha 1 (I) collagen gene
to embryonic

[117946]lethal mutation in mice. Proc. Nat. Acad. Sci. 81:

1504–1508, 1984.

[117947]20856.

[117948]Howard, I. K.: Clyde Edgar Keeler (1900–1994): geneticist,
artist,

[117949]cultural historian. J. Hered. 86: 489–491, 1995.

[117950]20857.

[117951]Jenkins, N. A.; Copeland, N. G.; Taylor, B. A.; Lee, B. K.:
Dilute

[117952](d) coat colour mutation of DBA/2J mice is associated with
the site

[117953]of integration of an ecotropic MLV genome. Nature 293:
370–374,

[117954]1981.

[117955]20858.

[117956]Keeler, C. E.: The inheritance of a retinal abnormality in
white

[117957]mice. Proc. Nat. Acad. Sci. 10: 329–333, 1924.

[117958]20859.

[117959]Khramtsov, N. V.; Feshchenko, E. A.; Suslova, V. A.;
Shmukler,

[117960]B. E.; Terpufov, B. E.; Rakitina, T. V.; Atabekova, N. V.;

Lipkin,

[117961]V. M.: The human rod photoreceptor cGMP phosphodiesterase beta-subunit:

[117962]structural studies of its cDNA and gene. FEBS Lett. 327: 275-278,

[117963]1993.

[117964]20860.

[117965]Lem, J.; Flannery, J. G.; Li, T.; Applebury, M. L.; Farber, D.

[117966]B.; Simon, M. I.: Retinal degeneration is rescued in transgenic rd

[117967]mice by expression of the cGMP phosphodiesterase beta subunit. Proc.

[117968]Nat. Acad. Sci. 89: 4422-4426, 1992.

[117969]20861.

[117970]McKusick, V. A.: The effect of lithium on the electrocardiogram

[117971]of animals and relation of this effect to the ratio of the intracellular

[117972]and extracellular concentrations of potassium. J. Clin. Invest. 33:

[117973]598-610, 1954.

[117974]20862.

[117975]McLaughlin, M. E.; Ehrhart, T. L.; Berson, E. L.; Dryja, T. P.

[117976]: Mutation spectrum of the gene encoding the beta subunit of rod phosphodiesterase

[117977]among patients with autosomal recessive retinitis pigmentosa. Proc.

[117978]Nat. Acad. Sci. 92: 3249-3253, 1995.

[117979]20863.

[117980]McLaughlin, M. E.; Sandberg, M. A.; Berson, E. L.; Dryja, T. P.

[117981]: Recessive mutations in the gene encoding the beta-subunit of rod

[117982]phosphodiesterase in patients with retinitis pigmentosa. Nature Genet. 4:

[117983]130-134, 1993.

[117984]20864.

[117985]Pittler, S. J.; Baehr, W.: Identification of a nonsense mutation

[117986]in the rod photoreceptor cGMP phosphodiesterase beta-subunit gene

[117987]of the rd mouse. Proc. Nat. Acad. Sci. 88: 8322-8326,

1991.

[117988]20865.

[117989]Pittler, S. J.; Keeler, C. E.; Sidman, R. L.; Baehr, W.: PCR

[117990]analysis of DNA from 70-year-old sections of rodless
retina demonstrates

[117991]identity with the mouse rd defect. Proc. Nat. Acad. Sci. 90:
9616-9619,

[117992]1993.

[117993]20866.

[117994]Ray, K.; Baldwin, V. J.; Acland, G. M.; Blanton, S. H.;
Aguirre,

[117995]G. D.: Cosegregation of codon 807 mutation of the canine
rod cGMP

[117996]phosphodiesterase beta gene and rcd1. Invest. Ophthalm.
Vis. Sci. 35:

[117997]4291-4299, 1994.

[117998]20867.

[117999]Riess, O.; Kozak, C.; Van Ommen, G.-J.: Report of the
fourth

[118000]international workshop on human chromosome 4 map-
ping 1996. Cytogenet.

[118001]Cell Genet. 74: 57–69, 1996.

[118002]20868.

[118003]Riess, O.; Noerremoelle, A.; Collins, C.; Mah, D.; Weber, B.;

[118004]Hayden, M. R.: Exclusion of DNA changes in the beta-subunit of the

[118005]c-GMP phosphodiesterase gene as the cause for Huntington's disease. Nature

[118006]Genet. 1: 104–108, 1992.

[118007]20869.

[118008]Riess, O.; Noerremoelle, A.; Weber, B.; Musarella, M. A.; Hayden,

[118009]M. R.: The search for mutations in the gene for the beta subunit

[118010]of the cGMP phosphodiesterase (PDEB) in patients with autosomal recessive

[118011]retinitis pigmentosa. Am. J. Hum. Genet. 51: 755–762, 1992.

[118012]20870.

[118013]Sidman, R. L.; Green, M. C.: Retinal degeneration in the mouse. J.

[118014]Hered. 56: 23–29, 1965.

[118015]20871.

[118016]Suber, M. L.; Pittler, S. J.; Qin, N.; Wright, G. C.; Holcombe,

[118017]Y.; Lee, R. H.; Craft, C. M.; Lolley, R. N.; Baehr, W.; Hur-
witz, R.

[118018]L.: Irish setter dogs affected with rod/cone dysplasia con-
tain a

[118019]nonsense mutation in the rod cGMP phosphodiesterase
beta-subunit gene. Proc.

[118020]Nat. Acad. Sci. 90: 3968-3972, 1993.

[118021]20872.

[118022]Weber, B.; Collins, C.; Kowbel, D.; Riess, O.; Hayden, M. R.:

[118023]Identification of multiple CpG islands and associated con-
served sequences

[118024]in a candidate region for the Huntington disease gene.

Genomics 11:

[118025]1113-1124, 1991.

[118026]20873.

[118027]Weber, B.; Riess, O.; Hutchinson, G.; Collins, C.; Lin, B.;
Kowbel,

[118028]D.; Andrew, S.; Schappert, K.; Hayden, M. R.: Genomic or-
ganization

[118029]and complete sequence of the human gene encoding the
beta-subunit

[118030]of the cGMP phosphodiesterase and its localisation to
4p16.3. Nucleic

[118031]Acids Res. 19: 6263-6268, 1991.

[118032]20874.

[118033]Bear, J. C.; Nemec, T. F.; Kennedy, J. C.; Marshall, W. H.;
Power,

[118034]A. A.; Kolonel, V. M.; Burke, G. B.: Inbreeding in outport
Newfoundland. Am.

[118035]J. Med. Genet. 29: 649-660, 1988.

[118036]20875.

[118037]Fontaine, B.; Vale-Santos, J.; Jurkat-Rott, K.; Reboul, J.;
Plassart,

[118038]E.; Rime, C.-S.; Elbaz, A.; Heine, R.; Guimaraes, J.; Weis-
senbach,

[118039]J.; Baumann, N.; Fardeau, M.; Lehmann-Horn, F.: Mapping
of the hypokalaemic

[118040]periodic paralysis (HypoPP) locus to chromosome
1q31-32 in three European

[118041]families. Nature Genet. 6: 267-272, 1994.

[118042]20876.

[118043]Jurkat-Rott, K.; Mitrovic, N.; Hang, C.; Kouzmekine, A.;
laizzo,

[118044]P.; Herzog, J.; Lerche, H.; Nicole, S.; Vale-Santos, J.; Chau-
veau,

[118045]D.; Fontaine, B.; Lehmann-Horn, F.: Voltage-sensor
sodium channel

[118046]mutations cause hypokalemic periodic paralysis type 2 by
enhanced

[118047]inactivation and reduced current. Proc. Nat. Acad. Sci. 97:
9549-9554,

[118048]2000.

[118049]20877.

[118050]Ptacek, L. J.; Tawil, R.; Griggs, R. C.; Engel, A. G.; Layzer,

[118051]R. B.; Kwiecinski, H.; McManis, P. G.; Santiago, L.; Moore,
M.; Fouad,

[118052]G.; Bradley, P.; Leppert, M. F.: Dihydropyridine receptor
mutations

[118053]cause hypokalemic periodic paralysis. Cell 77: 863-868,
1994.

[118054]20878.

[118055]Armstrong, F. S.: Hyperkalemic familial periodic paralysis
(adynamia

[118056]episodica hereditaria). Ann. Intern. Med. 57: 455–461,
1962.

[118057]20879.

[118058]Bennani-Baiti, I. M.; Jones, B. K.; Liebhaber, S. A.; Cooke,
N.

[118059]E.: Physical linkage of the human growth hormone gene
cluster and

[118060]the skeletal muscle sodium channel alpha-subunit gene
(SCN4A) on chromosome

[118061]17. Genomics 29: 647–652, 1995.

[118062]20880.

[118063]Borg, K.; Hovmoller, M.; Larsson, L.; Edstrom, L.: Paramy-
otonia

[118064]congenita (Eulenberg): clinical, neurophysiological and
muscle biopsy

[118065]observations in a Swedish family. Acta Neurol. Scand. 87:
37–42,

[118066]1993.

[118067]20881.

[118068]Burama, O. J. S.; Schipperheyn, J. J.: Periodic paralysis.In:

[118069]Vinken, P. J.; Bruyn, G. W.: Handbook of Clinical Neurology. Part

[118070]II, Diseases of Muscle. Amsterdam: Elsevier/North Holland (pub.)

[118071]41: 1979. Pp. 147–174.

[118072]20882.

[118073]Eulenberg, A.: Ueber eine familiäre, durch 6 generationen verfolgbare

[118074]Form congenitaler Paramyotonie. Neurologisches Centralblatt 12:

[118075]265–272, 1886.

[118076]20883.

[118077]Feero, W. G.; Wang, J.; Barany, F.; Zhou, J.; Todorovic, S. M.;

[118078]Conwit, R.; Galloway, G.; Hausmanowa-Petrusewicz, I.; Fidzianska,

[118079]A.; Arahata, K.; Wessel, H. B.; Wadeliuss, C.; Marks, H. G.; Hartlage,

[118080]P.; Hayakawa, H.; Hoffman, E. P.: Hyperkalemic periodic paralysis:

[118081]rapid molecular diagnosis and relationship of genotype to phenotype

[118082]in 12 families. Neurology 43: 668–673, 1993.

[118083]20884.

[118084]Fontaine, B.; Khurana, T. S.; Hoffman, E. P.; Bruns, G. A. P.;

[118085]Haines, J. L.; Trofatter, J. A.; Hanson, M. P.; Rich, J.; McFarlane,

[118086]H.; Yasek, D. M.; Romano, D.; Gusella, J. F.; Brown, R. H., Jr.:

[118087]Hyperkalemic periodic paralysis and the adult muscle sodium channel

[118088]alpha-subunit gene. Science 250: 1000–1002, 1990.

[118089]20885.

[118090]George, A. L., Jr.; Ledbetter, D. H.; Kallen, R. G.; Barchi, R.

[118091]L.: Assignment of a human skeletal muscle sodium channel alpha-subunit

[118092]gene (SCN4A) to 17q23.1–25.3. Genomics 9: 555–556, 1991.

[118093]20886.

[118094]Ammann, A. J.; Cain, W. A.; Ishizaka, K.; Hong, R.; Good, R. A.

[118095]: Immunoglobulin E deficiency in ataxia–telangiectasia.

New Eng.

[118096]J. Med. 281: 469–472, 1969.

[118097]20887.

[118098]Amromin, G. D.; Boder, E.; Teplitz, R.: Ataxia–

telangiectasia

[118099]with a 32–year survival: a clinicopathological report. J.

Neuropath.

[118100]Exp. Neurol. 38: 621–643, 1979.

[118101]20888.

[118102]Athma, P.; Rappaport, R.; Swift, M.: Molecular genotyping

shows

[118103]that ataxia–telangiectasia heterozygotes are predisposed

to breast

[118104]cancer. Cancer Genet. Cytogenet. 92: 130–134, 1996.

[118105]20889.

[118106]Aurias, A.; Croquette, M. F.; Nuyts, J. P.; Griscelli, C.;

Dutrillaux,

[118107]B.: New data on clonal anomalies of chromosome 14 in

ataxia telangiectasia:

[118108]tct(14;14) and inv(14). Hum. Genet. 72: 22–24, 1986.

[118109]20890.

[118110]Aurias, A.; Dutrillaux, B.: A possible new type of chromosome
some

[118111]rearrangement: telomere-centromere translocation (tct)
followed by

[118112]double duplication. Hum. Genet. 72: 25-26, 1986.

[118113]20891.

[118114]Aurias, A.; Dutrillaux, B.: Probable involvement of im-
munoglobulin

[118115]superfamily genes in most recurrent chromosomal rear-
rangements from

[118116]ataxia telangiectasia. Hum. Genet. 72: 210-214, 1986.

[118117]20892.

[118118]Aurias, A.; Dutrillaux, B.; Buriot, D.; Lejeune, J.: High fre-
quencies

[118119]of inversions and translocations of chromosomes 7 and
14 in ataxia-telangiectasia. Mutat.

[118120]Res. 69: 369-374, 1980.

[118121]20893.

[118122]Aurias, A.; Dutrillaux, B.; Griscelli, C.: Tandem transloca-
tion

[118123]t(14;14) in isolated and clonal cells in ataxia telangiectasia
are

[118124]different. Hum. Genet. 63: 320–322, 1983.

[118125]20894.

[118126]Baker, B. S.; Boyd, J. B.; Carpenter, A. T. C.; Green, M. M.;
Nguyen,

[118127]T. D.; Ripoll, P.; Smith, P. D.: Genetic controls of meiotic
recombination

[118128]and somatic DNA metabolism in *Drosophila melanogaster*.
Proc. Nat.

[118129]Acad. Sci. 73: 4140–4144, 1976.

[118130]20895.

[118131]Banga, S. S.; Shenkar, R.; Boyd, J. B.: Hypersensitivity of
Drosophila

[118132]mei-41 mutants to hydroxyurea is associated with re-
duced mitotic chromosome

[118133]stability. Mutat. Res. 163: 157–165, 1986.

[118134]20896.

[118135]Banin, S.; Moyal, L.; Shieh, S.-Y.; Taya, Y.; Anderson, C.
W.;

[118136]Chessa, L.; Smorodinsky, N. I.; Prives, C.; Reiss, Y.; Shiloh,

Y.;

[118137]Ziv, Y.: Enhanced phosphorylation of p53 by ATM in response to DNA

[118138]damage. Science 281: 1674–1677, 1998.

[118139]20897.

[118140]Bao, S.; Tibbetts, R. S.; Brumbaugh, K. M.; Fang, Y.; Richardson,

[118141]D. A.; Ali, A.; Chen, S. M.; Abraham, R. T.; Wang, X.-F.: ATR/ATM-mediated

[118142]phosphorylation of human Rad17 is required for genotoxic stress responses. Nature 411:

[118143]969–974, 2001.

[118144]20898.

[118145]Bar, R. S.; Levis, W. R.; Rechler, M. M.; Harrison, L. C.; Siebert,

[118146]C.; Podskalny, J.; Roth, J.; Muggeo, M.: Extreme insulin resistance

[118147]in ataxia telangiectasia: defect in affinity of insulin receptors. New

[118148]Eng. J. Med. 298: 1164–1171, 1978.

[118149]20899.

[118150]Barlow, C.; Dennerly, P. A.; Shigenaga, M. K.; Smith, M. A.;
Morrow,
[118151]J. D.; Roberts, L. J., II; Wynshaw-Boris, A.; Levine, R. L.:
Loss
[118152]of the ataxia-telangiectasia gene product causes oxidative
damage
[118153]in target organs. Proc. Nat. Acad. Sci. 96: 9915-9919,
1999.
[118154]20900.
[118155]Barlow, C.; Eckhaus, M. A.; Schaffer, A. A.; Wynshaw-Boris,
A.
[118156]: Atm haploinsufficiency results in increased sensitivity to
sublethal
[118157]doses of ionizing radiation in mice. Nature Genet. 21:
359-360,
[118158]1999.
[118159]20901.
[118160]Barlow, C.; Hirotsume, S.; Paylor, R.; Liyanage, M.; Eckhaus,
[118161]M.; Collins, F.; Shiloh, Y.; Crawley, J. N.; Ried, T.; Tagle, D.;
[118162]Wynshaw-Boris, A.: Atm-deficient mice: a paradigm of
ataxia telangiectasia. Cell 86:

[118163]159–171, 1996.

[118164]20902.

[118165]Bay, J.–O.; Uhrhammer, N.; Pernin, D.; Presneau, N.;
Tchirkov,

[118166]A.; Vuillaume, M.; Laplace, V.; Grancho, M.; Verrelle, P.;
Hall, J.;

[118167]Bignon, Y.–J.: High incidence of cancer in a family segregating a

[118168]mutation of the ATM gene: possible role of ATM heterozygosity in cancer. Hum.

[118169]Mutat. 14: 485–492, 1999.

[118170]20903.

[118171]Becker, Y.; Tabor, E.; Asher, Y.: Ataxia–telangiectasia fibroblasts

[118172]have less fibronectin mRNA than control cells but have the same levels

[118173]of integrin and beta–actin mRNA. Hum. Genet. 81:
165–170, 1989.

[118174]20904.

[118175]Bender, M. A.; Rary, J. M.; Kale, R. P.: G(2) chromosomal radiosensitivity

[118176]in ataxia telangiectasia lymphocytes. Mutat.Res. 152:
39-47, 1985.

[118177]20905.

[118178]Bender, M. A.; Rary, J. M.; Kale, R. P.: G(0) chromosomal
radiosensitivity

[118179]in ataxia telangiectasia lymphocytes. Mutat. Res. 150:
277-282,

[118180]1985.

[118181]20906.

[118182]Bernstein, R.; Pinto, M.; Jenkins, T.: Ataxia telangiectasia

[118183]with evolution of monosomy 14 and emergence of
Hodgkin's disease. Cancer

[118184]Genet. Cytogenet. 4: 31-37, 1981.

[118185]20907.

[118186]Bigbee, W. L.; Langlois, R. G.; Swift, M.; Jensen, R. H.: Evi-
dence

[118187]for an elevated frequency of in vivo somatic cell mutations
in ataxia

[118188]telangiectasia. Am. J. Hum. Genet. 44: 402-408, 1989.

[118189]20908.

[118190] Bishop, D. T.; Hopper, J.: AT-tributable risks? Nature
Genet. 15:

[118191] 226 only, 1997.

[118192] 20909.

[118193] Bochkov, N. P.; Lopukhin, Y. M.; Kuleshov, N. P.; Ko-
valchuk, L.

[118194] V.: Cytogenetic study of patients with ataxia-
telangiectasia. Humangenetik 24:

[118195] 115-128, 1974.

[118196] 20910.

[118197] Boder, E.: Ataxia-telangiectasia: an overview. In: Gatti, R.
A.;

[118198] Swift, M.: Ataxia-telangiectasia: Genetics, Neuropathology
and Immunology

[118199] of a Degenerative Disease of Childhood. New York: Alan R.
Liss (pub.)

[118200] 1985. Pp. 1-63.

[118201] 20911.

[118202] Boder, E.: Ataxia-telangiectasia: some historic, clinical and

[118203] pathologic observations. In: Bergsma, D. (ed.): Immunode-
ficiency in

[118204]Man and Animals. New York: National Foundation–March
of Dimes (pub.)

[118205]1975. Pp. 255–300.

[118206]20912.

[118207]Boder, E.; Sedgwick, R. P.: Ataxia–telangiectasia: a familial

[118208]syndrome of progressive cerebellar ataxia, oculocuta-
neous telangiectasia

[118209]and frequent pulmonary infection. Pediatrics 21: 526–554,
1958.

[118210]20913.

[118211]Borghesani, P. R.; Alt, F. W.; Bottaro, A.; Davidson, L.; Ak-
soy,

[118212]S.; Rathbun, G. A.; Roberts, T. M.; Swat, W.; Segal, R. A.;
Gu, Y.

[118213]: Abnormal development of Purkinje cells and lymphocytes
in Atm mutant

[118214]mice. Proc. Nat. Acad. Sci. 97: 3336–3341, 2000.

[118215]20914.

[118216]Boyd, J. B.; Golino, M. D.; Nguyen, T. D.; Green, M. M.: Iso-
lation

[118217]and characterization of X-linked mutants of Drosophila

melanogaster

[118218] which are sensitive to mutagens. Genetics 84: 485–506,
1976.

[118219] 20915.

- [118220] Bridges, B. A.; Arlett, C. F.: Risk of breast cancer in ataxia-telangiectasia.
- [118221] (Letter) New Eng. J. Med. 326: 1357-1361, 1992.
- [118222] 20916.
- [118223] Bridges, B. A.; Harnden, D. G.: Ataxia-telangiectasia: A Cellular
- [118224] and Molecular Link between Cancer, Neuropathology, and Immune Deficiency.
- [118225] New York: John Wiley (pub.) 1982.
- [118226] 20917.
- [118227] Barbosa, M. D. F. S.; Barrat, F. J.; Tchernev, V. T.; Nguyen, Q.
- [118228] A.; Mishra, V. S.; Colman, S. D.; Pastural, E.; Dufourcq-Lagelouse,
- [118229] R.; Fischer, A.; Holcombe, R. F.; Wallace, M. R.; Brandt, S. J.; de
- [118230] Saint Basile, G.; Kingsmore S. F.: Identification of mutations in
- [118231] two major mRNA isoforms of the Chediak-Higashi syndrome gene in human
- [118232] and mouse. Hum. Molec. Genet. 6: 1091-1098, 1997.

[118233]20918.

[118234]Barrat, F. J.; Auloge, L.; Pastural, E.; Dufourcq Lagelouse, R.;

[118235]Vilmer, E.; Cant, A. J.; Weissenbach, J.; Le Paslier, D.; Fischer,

[118236]A.; de Saint Basile, G.: Genetic and physical mapping of the Chediak-Higashi

[118237]syndrome on chromosome 1q42-43. Am. J. Hum. Genet. 59: 625-632,

[118238]1996.

[118239]20919.

[118240]Dufourcq-Lagelouse, R.; Lambert, N.; Duval, M.; Viot, G.; Vilmer,

[118241]E.; Fischer, A.; Prieur, M.; de Saint Basile, G.: Chediak-Higashi

[118242]syndrome associated with maternal uniparental isodisomy of chromosome

[118243]1. Europ. J. Hum. Genet. 7: 633-637, 1999.

[118244]20920.

[118245]Faigle, W.; Raposo, G.; Tenza, D.; Pinet, V.; Vogt, A. B.; Kropshofer,

- [118246]H.; Fischer, A.; de Saint-Basile, G.; Amigorena, S.: Deficient peptide loading and MHC class II endosomal sorting in a human genetic immunodeficiency disease: the Chediak-Higashi syndrome. J. Cell Biol. 141: 1121-1134, 1998.
- [118249]1998.
- [118250]20921.
- [118251]Karim, M. A.; Suzuki, K.; Fukai, K.; Oh, J.; Nagle, D. L.; Moore, K. J.; Barbosa, E.; Falik-Borenstein, T.; Filipovich, A.; Ishida, Y. Kivrikko, S.; Klein, C.; and 8 others: Apparent genotype-phenotype correlation in childhood, adolescent, and adult Chediak-Higashi syndrome. Am. J. Med. Genet. 108: 16-22, 2002.
- [118255]J. Med. Genet. 108: 16-22, 2002.
- [118256]20922.
- [118257]Kunieda, T.; Ide, H.; Nakagiri, M.; Yoneda, K.; Konfortov, B.; Ogawa, H.: Localization of the locus responsible for Chediak-Higashi

[118259]syndrome in cattle to bovine chromosome 28. Anim.

Genet. 31: 87–90,

[118260]2000.

[118261]20923.

[118262]El-Ashry, M. F.; El-Aziz, M. M. A.; Wilkins, S.; Cheetham,

M. E.;

[118263]Wilkie, S. E.; Hardcastle A. J.; Halford, S.; Bayoumi, A. Y.;

Ficker,

[118264]L. A.; Tuft, S.; Bhattacharya, S. S.; Ebenezer, N. D.: Identi-

fication

[118265]of novel mutations in the carbohydrate sulfotransferase

gene (CHST6)

[118266]causing macular corneal dystrophy. Invest. Ophthal. Vis.

Sci. 43:

[118267]377–382, 2002.

[118268]20924.

[118269]Akama, T. O.; Nishida, K.; Nakayama, J.; Watanabe, H.;

Ozaki, K.;

[118270]Nakamura, T.; Dota, A.; Kawasaki, S.; Inoue, Y.; Maeda, N.;

Yamamoto,

[118271]S.; Fujiwara, T.; Thonar, E. J.–M. A.; Shimomura, Y.; Ki-

noshita, S.;

[118272]Tanigami, A.; Fukuda, M. N.: Macular corneal dystrophy
type I and
[118273]type II are caused by distinct mutations in a new sulpho-
transferase
[118274]gene. Nature Genet. 26: 237–241, 2000.
[118275]20925.
[118276]Hasegawa, N.; Torii, T.; Kato, T.; Miyajima, H.; Furuhata,
A.;
[118277]Nakayasu, K.; Kanai, A.; Habuchi, O.: Decreased GlcNAc
6–O–sulfotransferase
[118278]activity in the cornea with macular corneal dystrophy. In-
vest. Ophthalm.
[118279]Vis. Sci. 41: 3670–3677, 2000.
[118280]20926.
[118281]Enomoto, A.; Kimura, H.; Chairoungdua, A.; Shigeta, Y.;
Jutabha,
[118282]P.; Cha, S. H.; Hosoyamada, M.; Takeda, M.; Sekine, T.;
Igarashi,
[118283]T.; Matsuo, H.; Kikuchi, Y.; Oda, T.; Ichida, K.; Hosoya, T.;
Shimokata,
[118284]K.; Niwa, T.; Kanai, Y.; Endou, H.: Molecular identification
of a

[118285]renal urate-anion exchanger that regulates blood urate levels. Nature 417:

[118286]447-452, 2002.

[118287]20927.

[118288]Anderson, S. L.; Coli, R.; Daly, I. W.; Kichula, E. A.; Rork, M.

[118289]J.; Volpi, S. A.; Ekstein, J.; Rubin, B. Y.: Familial dysautonomia

[118290]is caused by mutations of the IKAP gene. Am. J. Hum. Genet. 68:

[118291]753-758, 2001.

[118292]20928.

[118293]Blumenfeld, A.; Slaugenhaupt, S. A.; Axelrod, F. B.; Lucette, D.

[118294]E.; Maayan, C.; Liebert, C. B.; Ozelius, L. J.; Trofatter, J. A.;

[118295]Haines, J. L.; Breakefield, X. O.; Gusella, J. F.: Localization of

[118296]the gene for familial dysautonomia on chromosome 9 and definition

[118297]of DNA markers for genetic diagnosis. Nature Genet. 4: 160-164,

[118298]1993.

[118299]20929.

[118300]Blumenfeld, A.; Slaughter, S. A.; Liebert, C. B.; Temper, V.;

[118301]Maayan, C.; Gill, S.; Lucente, D. E.; Idelson, M.; McCormack, K.;

[118302]Monahan, M. A.; Mull, J.; Leyne, M.; Mendillo, M.; Schiripo, T.; Mishori,

[118303]E.; Breakefield, X.; Axelrod, F. B.; Gusella, J. F.: Precise genetic

[118304]mapping and haplotype analysis of the familial dysautonomia gene on

[118305]human chromosome 9q31. Am. J. Hum. Genet. 64: 1110–1118, 1999.

[118306]20930.

[118307]Xia, F.; Taghian, D. G.; DeFrank, J. S.; Zeng, Z.-C.; Willers,

[118308]H.; Iliakis, G.; Powell, S. N.: Deficiency of human BRCA2 leads to

[118309]impaired homologous recombination but maintains normal nonhomologous

[118310]end joining. Proc. Nat. Acad. Sci. 98: 8644–8649, 2001.

[118311]20931.

[118312]Goedert, M.; Hasegawa, J.; Craxton, M.; Leversha, M. A.;
Clegg,
[118313]S.: Assignment of the human stress-activated protein ki-
nase-3 gene
[118314](SAPK3) to chromosome 22q13.3 by fluorescence in situ
hybridization. Genomics 41:
[118315]501-502, 1997.
[118316]20932.
[118317]Wooster, R.; Bignell, G.; Lancaster, J.; Swift, S.; Seal, S.;
[118318]Mangion, J.; Collins, N.; Gregory, S.; Gumbs, C.; Micklem,
G.; Barfoot,
[118319]R.; Hamoudi, R.; Patel, S.; Rice, C.; Biggs, P.; Hashim, Y.;
Smith,
[118320]A.; Connor, F.; Arason, A.; Gudmundsson, J.; Ficenec, D.;
Keisell,
[118321]D.; Ford, D.; Tonin, P.; Bishop D. T.; Spurr, N. K.; Ponder, B.
A.
[118322]J.; Eeles, R.; Peto, J.; Devilee, P.; Cornelisse, C.; Lynch, H.;
Narod,
[118323]S.; Lenoir, G.; Egilsson, V.; Barkadottir, R. B.; Easton, D. F.;
Bentley,
[118324]D. R.; Futreal, P. A.; Ashworth, A.; Stratton, M. R.: Identifi-

cation

[118325]of the breast cancer susceptibility gene BRCA2. Nature
378: 789–792,

[118326]1995.

[118327]20933.

[118328]Wooster, R.; Neuhausen, S. L.; Mangion, J.; Quirk, Y.; Ford,
D.;

[118329]Collins, N.; Nguyen, K.; Seal, S.; Tran, T.; Averill, D.; Fields,

[118330]P.; Marshall, G.; Narod, S.; Lenoir, G. M.; Lynch, H.; Feun-
teun, J.;

[118331]Devilee, P.; Cornelisse, C. J.; Menko, F. H.; Daly, P. A.;
Ormiston,

[118332]W.; McManus, R.; Pye, C.; Lewis, C. M.; Cannon–Albright,
L. A.; Peto,

[118333]J.; Ponder, B. A. J.; Skolnick, M. H.; Easton, D. F.; Goldgar,
D.

[118334]E.; Stratton, M. R.: Localization of a breast cancer suscep-
tibility

[118335]gene, BRCA2, to chromosome 13q12–13. Science 265:
2088–2090, 1994.

[118336]20934.

[118337]Yang, H.; Jeffrey, P. D.; Miller, J.; Kinnucan, E.; Sun, Y.;

Thoma,

[118338]N. H.; Zheng, N.; Chen, P.-L.; Lee, W.-H.; Pavletich, N. P.:

BRCA2

[118339]function in DNA binding and recombination from a

BRCA2-DSS1-ssDNA

[118340]structure. Science 297: 1837-1848, 2002.

[118341]20935.

[118342]Hartley, D. A.; Preiss, A.; Artavanis-Tsakonas, S.: A deduced

[118343]gene product from the Drosophila neurogenic locus, enhancer of split,

[118344]shows homology to mammalian G-protein beta subunit.

Cell 55: 785-795,

[118345]1988.

[118346]20936.

[118347]Hou, E. W.; Li, S.-L.: Genomic organization and chromosome localization

[118348]to band 19p13.3 of the human AES gene: gene product exhibits strong

[118349]similarity to the N-terminal domain of Drosophila enhancer of split

[118350]Groucho protein. DNA Cell Biol. 17: 911-913, 1998.

[118351]20937.

[118352]Miyasaka, H.; Choudhury, B. K.; Hou, E. W.; Li, S. S.-L.:
Molecular

[118353]cloning and expression of mouse and human cDNA en-
coding AES and ESG

[118354]proteins with strong similarity to Drosophila enhancer of
split groucho

[118355]protein. Europ. J. Biochem. 216: 343-352, 1993.

[118356]20938.

[118357]Avraham, K. B.; Prezioso, V. R.; Chen, W. S.; Lai, E.; Sladek,

[118358]F. M.; Zhong, W.; Darnell, J. E., Jr.; Jenkins, N. A.;
Copeland, N.

[118359]G.: Murine chromosomal location of four hepatocyte-en-
riched transcription

[118360]factors: HNF-3-alpha, HNF-3-beta, HNF-3-gamma, and
HFN-4. Genomics 13:

[118361]264-268, 1992.

[118362]20939.

[118363]Deleuze, J. F.; Dhorne, S.; Hazan, J.; Borghi, E.; Raynaud,
N.;

[118364]Pollet, N.; Meunier-Rotival, M.; Deschatrette, J.; Alagille,

D.; Hadchouel,

[118365]M.: Deleted chromosome 20 from a patient with Alagille syndrome isolated

[118366]in a cell hybrid through leucine transport selection: study of three

[118367]candidate genes. Mammalian Genome 5: 663–669, 1994.

[118368]20940.

[118369]Kaestner, K. H.; Hiemisch, H.; Luckow, B.; Schutz, G.: The HNF-3

[118370]gene family of transcription factors in mice: gene structure, cDNA

[118371]sequence, and mRNA distribution. Genomics 20: 377–385, 1994.

[118372]20941.

[118373]Mincheva, A.; Lichter, P.; Schutz, G.; Kaestner, K. H.: Assignment

[118374]of the human genes for hepatocyte nuclear factor 3-alpha, -beta, and

[118375]-gamma (HNF3A, HNF3B, HNF3G) to 14q12–q13, 20p11, and 19q13.2–q13.4. Genomics 39:

[118376]417–419, 1997.

[118377]20942.

[118378]Adams, R. H.; Porras, A.; Alonso, G.; Jones, M.; Vintersten, K.;

[118379]Panelli, S.; Valladares, A.; Perez, L.; Klein, R.; Nebreda, A. R.

[118380]: Essential role of p38- α MAP kinase in placental but not embryonic

[118381]cardiovascular development. *Molec. Cell* 6: 109–116, 2000.

[118382]20943.

[118383]Ge, B.; Gram, H.; Di Padova, F.; Huang, B.; New, L.; Ulevitch,

[118384]R. J.; Luo, Y.; Han, J.: MAPKK-independent activation of p38- α

[118385]mediated by TAB1-dependent autophosphorylation of p38- α . *Science* 295:

[118386]1291–1294, 2002.

[118387]20944.

[118388]Han, J.; Lee, J.-D.; Bibbs, L.; Ulevitch, R. J.: A MAP kinase

[118389]targeted by endotoxin and hyperosmolarity in mammalian cells. *Science* 265:

[118390]808–811, 1994.

[118391]20945.

[118392]Kim, D. H.; Feinbaum, R.; Alloing, G.; Emerson, F. E.;
Garsin,

[118393]D. A.; Inoue, H.; Tanaka-Hino, M.; Hisamoto, N.; Mat-
sumoto, K.; Tan,

[118394]M.-W.; Ausubel, F. M.: A conserved p38 MAP kinase path-
way in *Caenorhabditis*

[118395]*elegans* innate immunity. *Science* 297: 623-626, 2002.

[118396]20946.

[118397]Kumar, S.; McLaughlin, M. M.; McDonnell, P. C.; Lee, J. C.;
Livi,

[118398]G. P.; Young, P. R.: Human mitogen-activated protein ki-
nase CSBP1,

[118399]but not CSBP2, complements a *hog1* deletion in yeast. *J.*
Biol. Chem. 270:

[118400]29043-29046, 1995.

[118401]20947.

[118402]Lee, J. C.; Laydon, J. T.; McDonnell, P. C.; Gallagher, T. F.;

[118403]Kumar, S.; Green, D.; McNulty, D.; Blumenthal, M. J.; Heys,
J. R.;

[118404]Landvatter, S. W.; Stickler, J. E.; McLaughlin, M. M.;

Siemens, I.

[118405]R.; Fisher, S. M.; Livi, G. P.; White, J. R.; Adams, J. L.;

Young,

[118406]P. R.: A protein kinase involved in the regulation of in-

flammatory

[118407]cytokine biosynthesis. *Nature* 372: 739–746, 1994.

[118408]20948.

[118409]Liao, P.; Georgakopoulos, D.; Kovacs, A.; Zheng, M.;

Lerner, D.;

[118410]Pu, H.; Saffitz, J.; Chien, K.; Xiao, R.-P.; Kass, D. A.; Wang,

Y.

[118411]: The in vivo role of p38 MAP kinases in cardiac remodel-

ing and restrictive

[118412]cardiomyopathy. *Proc. Nat. Acad. Sci.* 98: 12283–12288,

2001.

[118413]20949.

[118414]Maizels, E. T.; Mukherjee, A.; Sithanandam, G.; Peters, C.

A.;

[118415]Cottom, J.; Mayo, K. E.; Hunzicker-Dunn, M.: Develop-

mental regulation

[118416]of mitogen-activated protein kinase-activated kinases-2

and -3 (MAPKAPK-2/-3)

[118417]in vivo during corpus luteum formation in the rat. Molec.

Endocr. 15:

[118418]716–733, 2001.

[118419]20950.

[118420]McDonnell, P. C.; DiLella, A. G.; Lee, J. C.; Young, P. R.:

Localization

[118421]of the human stress responsive MAP kinase-like CSAIDs
binding protein

[118422](CSBP) gene to chromosome 6p21.3/21.2. Genomics 29:
301–302, 1995.

[118423]20951.

[118424]New, L.; Jiang, Y.; Zhao, M.; Liu, K.; Zhu, W.; Flood, L. J.;

[118425]Kato, Y.; Parry, G. C. N.; Han, J.: PRAK, a novel protein ki-
nase

[118426]regulated by the p38 MAP kinase. EMBO J. 17: 3372–3384,
1998.

[118427]20952.

[118428]Ni, H.; Wang, X. S.; Diener, K.; Yao, Z.: MAPKAPK5, a novel
mitogen-activate

[118429]d protein kinase (MAPK)-activated protein kinase, is a
substrate of

[118430]the extracellular-regulated kinase (ERK) and p38 kinase.

Biochem.

[118431]Biophys. Res. Commun. 243: 492–496, 1998.

[118432]20953.

[118433]Takekawa, M.; Maeda, T.; Saito, H.: Protein phosphatase
2C- α

[118434]inhibits the human stress-responsive p38 and JNK MAPK
pathways. EMBO

[118435]J. 17: 4744–4752, 1998.

[118436]20954.

[118437]Tamura, K.; Sudo, T.; Senftleben, U.; Dadak, A. M.; John-
son, R.;

[118438]Karin, M.: Requirement for p38- α in erythropoietin
expression:

[118439]a role for stress kinases in erythropoiesis. Cell 102:
221–231,

[118440]2000.

[118441]20955.

[118442]Haber, N.; Stengel, D.; Defer, N.; Roeckel, N.; Mattei,
M.–G.;

[118443]Hanoune, J.: Chromosomal mapping of human adenylyl

cyclase genes

[118444]type III, type V and type VI. Hum. Genet. 94: 69–73, 1994.

[118445]20956.

[118446]Sinnarajah, S.; Dessauer, C. W.; Srikumar, D.; Chen, J.;

Yuen,

[118447]J.; Yilma, S.; Dennis, J. C.; Morrison, E. E.; Vodyanoy, V.;

Kehrl,

[118448]J. H.: RGS2 regulates signal transduction in olfactory neu-

rons by

[118449]attenuating activation of adenylyl cyclase III. Nature 409:

1051–1055,

[118450]2001.

[118451]20957.

[118452]Thaler, J. P.; Lee, S.-K.; Jurata, L. W.; Gill, G. N.; Pfaff,

[118453]S. L.: LIM factor Lhx3 contributes to the specification of

motor

[118454]neuron and interneuron identity through cell-

type-specific protein-protein

[118455]interactions. Cell 110: 237–249, 2002.

[118456]20958.

[118457]Tebbutt, N. C.; Giraud, A. S.; Inglese, M.; Jenkins, B.; War-

ing,

[118458]P.; Clay, F. J.; Malki, S.; Alderman, B. M.; Grail, D.; Hol-
lande,

[118459]F.; Heath, J. K.; Ernst, M.: Reciprocal regulation of gas-
trointestinal

[118460]homeostasis by SHP2 and STAT-mediated trefoil gene ac-
tivation in gp130

[118461]mutant mice. Nature Med. 8: 1089-1097, 2002.

[118462]20959.

[118463]Brown, K. A.; Leek, J. P.; Lench, N. J.; Moynihan, L. M.;
Markham,

[118464]A. F.; Mueller, R. F.: Human sequences homologous to the
gene for

[118465]the cochlear protein Ocp-II do not map to currently known
non-syndromic

[118466]hearing loss loci. Ann. Hum. Genet. 60: 385-389, 1996.

[118467]20960.

[118468]Chen, H.; Thalmann, I.; Adams, J. C.; Avraham, K. B.;
Copeland,

[118469]N. G.; Jenkins, N. A.; Beier, D. R.; Corey, D. P.; Thalmann,
R.; Duyk,

[118470]G. M.: cDNA cloning, tissue distribution, and chromoso-

mal localization

[118471]of Ocp2, a gene encoding a putative transcription-associated factor

[118472]predominantly expressed in the auditory organs. Genomics 27: 389–398,

[118473]1995.

[118474]20961.

[118475]Demetrick, D. J.; Zhang, H.; Beach, D. H.: Chromosomal mapping

[118476]of the genes for the human CDK2/cyclin A-associated proteins p19 (SKP1A

[118477]and SKP1B) and p45 (SKP2). Cytogenet. Cell Genet. 73: 104–107, 1996.

[118478]20962.

[118479]Liang, Y.; Chen, H.; Asher, J. H., Jr.; Chang, C.-C.; Friedman,

[118480]T. B.: Human inner ear OCP2 cDNA maps to 5q22–5q35.2 with related

[118481]sequences on chromosomes 4p16.2–4p14, 5p13–5q22, 7pter–q22, 10 and

[118482]12p13–12qter. Gene 184: 163–167, 1997.

[118483]20963.

[118484] Sowden, J.; Morrison, K.; Schofield, J.; Putt, W.; Edwards, Y.

[118485] A novel cDNA with homology to an RNA polymerase II
elongation factors

[118486] maps to human chromosome 5q31 (TCEB1L) and to
mouse chromosome 11

[118487] (Tceb1l). Genomics 29: 145–151, 1995.

[118488] 20964.

[118489] Thalmann, I.; Rosenthal, H. L.; Moore, B. W.; Thalmann, R.:
Organ

[118490] of Corti-specific polypeptides: OCP-I and OCP-II. Arch.
Oto-Rhino-Laryngol. 226:

[118491] 123–128, 1980.

[118492] 20965.

[118493] Thalmann, I.; Takahashi, K.; Varghese, J.; Comegys, T. H.;
Thalmann,

[118494] R.: Biochemical features of major organ of Corti proteins
(OCP-I

[118495] and OCP-II) including partial amino acid sequence. Laryn-
goscope 100:

[118496] 99–105, 1990.

[118497] 20966.

[118498]Zhang, H.; Kobayashi, R.; Galaktionov, K.; Beach, D.:

p19Skl

[118499]and p45Skl are essential elements of the cyclin A-CDK2
S phase kinase. Cell 82:

[118500]915-925, 1995.

[118501]20967.

[118502]FitzPatrick, D. R.; Germain-Lee, E.; Valle, D.: Isolation and

[118503]characterization of rat and human cDNAs encoding a
novel putative

[118504]peroxisomal enoyl-CoA hydratase. Genomics 27:
457-466, 1995.

[118505]20968.

[118506]Saijo, M.; Sakai, Y.; Kishino, T.; Niikawa, N.; Matsuura, Y.;
Morino,

[118507]K.; Tamai, K.; Taya, Y.: Molecular cloning of a human pro-
tein that

[118508]binds to the retinoblastoma protein and chromosomal
mapping. Genomics 27:

[118509]511-519, 1995.

[118510]20969.

[118511]Anand, A.; Chada, K.: In vivo modulation of Hmgic reduces

obesity. Nature

[118512]Genet. 24: 377–380, 2000.

[118513]20970.

[118514]Arlotta, P.; Tai, A. K.-F.; Manfioletti, G.; Clifford, C.; Jay,

[118515]G.; Ono, S. J.: Transgenic mice expressing a truncated
form of the

[118516]high mobility group I–C protein develop adiposity and an
abnormally

[118517]high prevalence of lipomas. J. Biol. Chem. 275:
14394–14400, 2000.

[118518]20971.

[118519]Ashar, H. R.; Cherath, L.; Przybysz, K. M.; Chada, K.: Ge-
nomic

[118520]characterization of human HMGIC, a member of the ac-
cessory transcription

[118521]factor family found at translocation breakpoints in lipo-
mas. Genomics 31:

[118522]207–214, 1996.

[118523]20972.

[118524]Ashar, H. R.; Schoenberg Fejzo, M.; Tkachenko, A.; Zhou,
X.; Fletcher,

[118525]J. A.; Weremowicz, S.; Morton, C. C.; Chada, K.: Disruption
of the
[118526]architectural factor HMGI-C: DNA-binding AT hook motifs
fused in lipomas
[118527]to distinct transcriptional regulatory domains. Cell 82:
57-65,
[118528]1995.
[118529]20973.
[118530]Chau, K.-Y.; Patel, U. A.; Lee, K.-L. D.; Lam, H.-Y. P.;
Crane-Robinson,
[118531]C.: The gene for the human architectural transcription
factor HMGI-C
[118532]consists of five exons each coding for a distinct functional
element. Nucleic
[118533]Acids Res. 23: 4262-4266, 1995.
[118534]20974.
[118535]Danforth, E., Jr.: Failure of adipocyte differentiation causes
[118536]type II diabetes mellitus. Nature Genet. 26: 13 only, 2000.
[118537]20975.
[118538]Friedmann, M.; Holth, L. T.; Zoghbi, H. Y.; Reeves, R.: Or-
ganization,

[118539]inducible-expression and chromosome localization of the human HMG-I(Y)

[118540]nonhistone protein gene. Nucleic Acids Res. 21: 4259-4267, 1993.

[118541]20976.

[118542]Ishwad, C. S.; Shriver, M. D.; Lassige, D. M.; Ferrell, R. E.:

[118543]The high mobility group I-C gene (HMGI-C): polymorphism and genetic

[118544]localization. Hum. Genet. 99: 103-105, 1997.

[118545]20977.

[118546]Kazmierczak, B.; Dal Cin, P.; Wanschura, S.; Bartnitzke, S.; Van

[118547]den Berghe, H.; Bullerdiek, J.: Cloning and molecular characterization

[118548]of part of a new gene fused to HMGIC in mesenchymal tumors. Am. J.

[118549]Path. 152: 431-435, 1998.

[118550]20978.

[118551]Kazmierczak, B.; Pohnke, Y.; Bullerdiek, J.: Fusion transcripts

[118552]between the HMGIC gene and RTVL-H-related sequences

in mesenchymal

[118553] tumors without cytogenetic aberrations. Genomics 38:
223-226, 1996.

[118554] 20979.

[118555] Manfioletti, G.; Rustighi, A.; Mantovani, F.; Goodwin, G. H.;

[118556] Giancotti, V.: Isolation and characterization of the gene
coding

[118557] for murine high-mobility-group protein HMGI-C. Gene
167: 249-253,

[118558] 1995.

[118559] 20980.

[118560] Mine, N.; Kurose, K.; Nagai, H.; Doi, D.; Ota, Y.; Yoneyama,
K.;

[118561] Konishi, H.; Araki, T.; Emi, M.: Gene fusion involving
HMGIC is a

[118562] frequent aberration in uterine leiomyomas. J. Hum. Genet.
46: 408-412,

[118563] 2001.

[118564] 20981.

[118565] Nucci, M. R.; Weremowicz, S.; Neskey, D. M.; Sornberger,
K.; Tallini,

[118566]G.; Morton, C. C.; Quade, B. J.: Chromosomal translocation
t(8;12)

[118567]induces aberrant HMGIC expression in aggressive an-
giomyxoma of the

[118568]vulva. Genes Chromosomes Cancer 32: 172–176, 2001.

[118569]20982.

[118570]Oka, J.; Matsumoto, A.; Hosokawa, Y.; Inoue, S.: Molecular
cloning

[118571]of human cytosolic purine 5–prime–nucleotidase.

Biochem. Biophys.

[118572]Res. Commun. 205: 917–922, 1994.

[118573]20983.

[118574]Kingsmore, S. F.; Suh, D.; Seldin, M. F.: Genetic mapping
of the

[118575]glycine receptor alpha–3 subunit on mouse chromosome

8. Mammalian

[118576]Genome 5: 831–832, 1994.

[118577]20984.

[118578]Kuhse, J.; Schmieden, V.; Betz, H.: Identification and func-
tional

[118579]expression of a novel ligand binding subunit of the in-

hibitory glycine

[118580]receptor. J. Biol. Chem. 265: 22317–22320, 1990.

[118581]20985.

[118582]Nikolic, Z.; Laube, B.; Weber, R. G.; Lichter, P.; Kioschis, P.;

[118583]Poustka, A.; Mulhardt, C.; Becker, C.–M.: The human
glycine receptor

[118584]subunit alpha–3: GLRA3 gene structure, chromosomal lo-
calization, and

[118585]functional characterization of alternative transcripts. J.
Biol.

[118586]Chem. 273: 19708–19714, 1998.

[118587]20986.

[118588]Way, J. C.; Chalfie, M.: mec–3, a homeobox–containing
gene that

[118589]specifies differentiation of the touch receptor neurons in
C. elegans. Cell 54:

[118590]5–16, 1988.

[118591]20987.

[118592]Zhadanov, A. B.; Bertuzzi, S.; Taira, M.; Dawid, I. B.; West-
phal,

[118593]H.: Expression pattern of the murine LIM class homeobox

gene Lhx3

[118594]in subsets of neural and neuroendocrine tissues. Dev.

Dyn. 202:

[118595]354–364, 1995.

[118596]20988.

[118597]Zhadanov, A. B.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N.

[118598]A.; Westphal, H.: Genomic structure and chromosomal localization

[118599]of the mouse LIM/homeobox gene Lhx3. Genomics 27: 27–32, 1995.

[118600]20989.

[118601]Seimiya, H.; Sawabe, T.; Inazawa, J.; Tsuruo, T.: Cloning, expression

[118602]and chromosomal localization of a novel gene for protein tyrosine

[118603]phosphatase (PTP-U2) induced by various differentiation-inducing agents. Oncogene 10:

[118604]1731–1738, 1995.

[118605]20990.

[118606]Wiggins, R. C.; Wiggins, J. E.; Goyal, M.; Wharram, B. L.; Thomas,

[118607]P. E.: Molecular cloning of cDNAs encoding human
GLEPP1, a membrane
[118608]protein tyrosine phosphatase: characterization of the
GLEPP1 protein
[118609]distribution in human kidney and assignment of the
GLEPP1 gene to
[118610]human chromosome 12p12-p13. Genomics 27: 174-181,
1995.
[118611]20991.

[118612]Jia, S.; VanDusen, W. J.; Diehl, R. E.; Kohl, N. E.; Dixon, R.
[118613]A. F.; Elliston, K. O.; Stern, A. M.; Friedman, P. A.: cDNA
cloning
[118614]and expression of bovine aspartyl (asparaginy) beta-
hydroxylase. J.
[118615]Biol. Chem. 267: 14322-14327, 1992.
[118616]20992.

[118617]Korioth, F.; Gieffers, C.; Frey, J.: Cloning and characteriza-
tion
[118618]of the human gene encoding aspartyl beta-hydroxylase.
Gene 150:
[118619]395-399, 1994.

[118620]20993.

[118621]Lavaissiere, L.; Jia, S.; Nishiyama, M.; de la Monte, S.;
Stern,

[118622]A. M.; Wands, J. R.; Friedman, P. A.: Overexpression of human aspartyl(asparaginy)-beta-hydroxylase

[118623]in hepatocellular carcinoma and cholangiocarcinoma. J.
Clin. Invest. 98:

[118624]1313-1323, 1996.

[118625]20994.

[118626]Lim, K. Y.; Hong, C.-S.; Kim, D. H.: cDNA cloning and
characterization

[118627]of human cardiac junctin. Gene 255: 35-42, 2000.

[118628]20995.

[118629]Scott, A. F.: Personal Communication. Baltimore, Md.
2/19/2001.

[118630]20996.

[118631]Treves, S.; Feriotto, G.; Moccagatta, L.; Gambari, R.;
Zorzato,

[118632]F.: Molecular cloning, expression, functional characterization,
chromosomal

[118633]localization, and gene structure of junctate, a novel inte-

gral calcium

[118634]binding protein of sarco(endo)plasmic reticulum membrane. J. Biol.

[118635]Chem. 275: 39555–39568, 2000.

[118636]20997.

[118637]Wetzel, G. T.; Ding, S.; Chen, F.: Molecular cloning of junctin

[118638]from human and developing rabbit heart. Molec. Genet. Metab. 69:

[118639]252–258, 2000.

[118640]20998.

[118641]Sato, K.; Mano, H.; Ariyama, T.; Inazawa, J.; Yazaki, Y.; Hirai,

[118642]H.: Molecular cloning and analysis of the human Tec protein-tyrosine

[118643]kinase. Leukemia 8: 1663–1672, 1994.

[118644]20999.

[118645]Zhou, P.; Sun, L. J.; Dotsch, V.; Wagner, G.; Verdine, G. L.:

[118646]Solution structure of the core NFATC1/DNA complex. Cell 92: 687–696,

[118647]1998.

[118648]21000.

[118649]McCaffrey, P. G.; Luo, C.; Kerppola, T. K.; Jain, J.; Badalian,

[118650]T. M.; Ho, A. M.; Burgeon, E.; Lane, W. S.; Lambert, J. N.;

Curran,

[118651]T.; Verdine, G. L.; Rao, A.; Hogan, P. G.: Isolation of the

cyclosporin-sensitive

[118652]T cell transcription factor NFATp. Science 262: 750-754,

1993.

[118653]21001.

[118654]Rengarajan, J.; Tang, B.; Glimcher, L. H.: NFATc2 and

NFATc3 regulate

[118655]TH2 differentiation and modulate TCR-responsiveness of

naive TH cells. Nature

[118656]Immun. 3: 48-54, 2002.

[118657]21002.

[118658]Abrams, W. R.; Ma, R.-I.; Kucich, U.; Bashir, M. M.; Decker,

S.;

[118659]Tsipouras, P.; McPherson, J. D.; Wasmuth, J. J.; Rosen-

bloom, J.:

[118660]Molecular cloning of the microfibrillar protein MFAP3 and

assignment

[118661]of the gene to human chromosome 5q32-q33.2. Ge-
nomics 26: 47-54,

[118662]1995.

[118663]21003.

[118664]Atanasoski, S.; Toldo, S. S.; Malipiero, U.; Schreiber, E.;
Fries,

[118665]R.; Fontana, A.: Isolation of the human genomic brain-
2/N-Oct 3 gene

[118666](POUF3) and assignment to chromosome 6q16. Genomics
26: 272-280,

[118667]1995.

[118668]21004.

[118669]Schreiber, E.; Tobler, A.; Malipiero, U.; Schaffner, W.;
Fontana,

[118670]A.: cDNA cloning of human N-Oct 3, a nervous-system
specific POU

[118671]domain transcription factor binding to the octamer DNA
motif. Nucleic

[118672]Acids Res. 21: 253-258, 1993.

[118673]21005.

[118674]Xia, Y.-R.; Andersen, B.; Mehrabian, M.; Diep, A. T.; War-

den, C.

[118675]H.; Mohandas, T.; McEvilly, R. J.; Rosenfeld, M. G.; Lusic, A.
J.

[118676]Chromosomal organization of mammalian POU domain
factors. Genomics 18:

[118677]126–130, 1993.

[118678]21006.

[118679]Brass, N.; Heckel, D.; Sahin, U.; Pfreundschuh, M.;
Sybrecht, G.

[118680]W.; Meese, E.: Translation initiation factor eIF-4gamma is
encoded

[118681]by an amplified gene and induces an immune response in
squamous cell

[118682]lung carcinoma. Hum. Molec. Genet. 6: 33–39, 1997.

[118683]21007.

[118684]Gradi, A.; Imataka, H.; Svitkin, Y. V.; Rom, E.; Raught, B.;
Morino,

[118685]S.; Sonenberg, N.: A novel functional human eukaryotic
translation

[118686]initiation factor 4G. Molec. Cell. Biol. 18: 334–342, 1998.

[118687]21008.

[118688]Imataka, H.; Gradi, A.; Sonenberg, N.: A newly identified
N-terminal

[118689]amino acid sequence of human eIF4G binds
poly(A)-binding protein and

[118690]functions in poly(A)-dependent translation. EMBO J. 17:
7480-7489,

[118691]1998.

[118692]21009.

[118693]Imataka, H.; Sonenberg, N.: Human eukaryotic translation
initiation

[118694]factor 4G (eIF4G) possesses two separate and independent
binding sites

[118695]for eIF4A. Molec. Cell. Biol. 17: 6940-6947, 1997.

[118696]21010.

[118697]Yan, R.; Rhoads, R. E.: Human protein synthesis initiation
factor

[118698]eIF-4-gamma is encoded by a single gene (EIF4G) that
maps to chromosome

[118699]3q27-qter. Genomics 26: 394-398, 1995.

[118700]21011.

[118701]Yan, R.; Rychlik, W.; Etchison, D.; Rhoads, R. E.: Amino

acid

[118702]sequence of the human protein synthesis initiation factor
eIF-4-gamma. J.

[118703]Biol. Chem. 267: 23226-23231, 1992.

[118704]21012.

[118705]Corrigall, A. V.; Hift, R. J.; Hancock, V.; Meissner, D.;
Davids,

[118706]L.; Kirsch, R. E.; Meissner, P. N.: Identification and charac-
terisation

[118707]of a deletion (537delAT) in the protoporphyrinogen oxi-
dase gene in

[118708]a South African variegate porphyria family. Hum. Mutat.
12: 403-407,

[118709]1998.

[118710]21013.

[118711]Aso, T.; Lane, W. S.; Conaway, J. W.; Conaway, R. C.: Elong-
gin

[118712](SIII): a multisubunit regulator of elongation by RNA poly-
merase II. Science 269:

[118713]1439-1443, 1995.

[118714]21014.

[118715] Baldwin, C. T.; Weiss, S.; Farrer, L.; De Stefano, A.; Adair, R.;

[118716] Franklyn, B.; Kidd, K. K.; Korostishevsky, M.; Bonne-Tamir, B.: Linkage

[118717] of congenital, recessive deafness (DFNB4) to chromosome 7q31 and evidence

[118718] for genetic heterogeneity in the Middle Eastern Druze population. Hum.

[118719] Molec. Genet. 4: 1637–1642, 1995.

[118720] 21015.

[118721] Li, X. C.; Everett, L. A.; Lalwani, A. K.; Desmukh, D.; Friedman,

[118722] T. B.; Green, E. D.; Wilcox, E. R.: A mutation in PDS causes non-syndromic

[118723] recessive deafness. (Letter) Nature Genet. 18: 215–217, 1998.

[118724] 21016.

[118725] Boon, L. M.; Mulliken, J. B.; Vikkula, M.; Watkins, H.; Seidman,

[118726] J.; Olsen, B. R.; Warman, M. L.: Assignment of a locus for dominantly

[118727] inherited venous malformations to chromosome 9p. Hum.

Molec. Genet. 3:

[118728]1583–1587, 1994.

[118729]21017.

[118730]Calvert, J. T.; Riney, T. J.; Kontos, C. D.; Cha, E. H.; Prieto,

[118731]V. G.; Shea, C. R.; Berg, J. N.; Nevin, N. C.; Simpson, S. A.;

Pasyk,

[118732]K. A.; Speer, M. C.; Peters, K. G.; Marchuk, D. A.: Allelic

and locus

[118733]heterogeneity in inherited venous malformations. Hum.

Molec. Genet. 8:

[118734]1279–1289, 1999.

[118735]21018.

[118736]Vikkula, M.; Boon, L. M.; Carraway, K. L., III; Calvert, J. T.;

[118737]Diamonti, A. J.; Goumnerov, B.; Pasyk, K. A.; Marchuk, D.

A.; Warman,

[118738]M. L.; Cantley, L. C.; Mulliken, J. B.; Olsen, B. R.: Vascular

dysmorphogenesis

[118739]caused by an activating mutation in the receptor tyrosine

kinase TIE2. Cell 87:

[118740]1181–1190, 1996.

[118741]21019.

[118742]Iwata, T.; Kogame, K.; Toki, T.; Yokoyama, M.; Yamamoto, M.; Ito,

[118743]E.: Structure and chromosome mapping of the human small maf genes

[118744]MAFG and MAFK. Cytogenet. Cell Genet. 82: 88–90, 1998.
[118745]21020.

[118746]Motohashi, H.; Katsuoka, F.; Shavit, J. A.; Engel, J. D.; Yamamoto,

[118747]M.: Positive or negative MARE-dependent transcriptional regulation

[118748]is determined by the abundance of small Maf proteins. Cell 103:

[118749]865–875, 2000.
[118750]21021.

[118751]Peters, L. L.; Eicher, E. M.: The ubiquitous subunit of the globin

[118752]enhancer-binding protein NF-E2 (Nfe2u) maps to mouse chromosome 5. Genomics 22:

[118753]490–491, 1994.
[118754]21022.

[118755]Shavit, J. A.; Motohashi, H.; Onodera, K.; Akasaka, J.; Ya-

mamoto,

[118756]M.; Engel, J. D.: Impaired megakaryopoiesis and behavioral defects

[118757]in mafG-null mutant mice. Genes Dev. 12: 2164–2174, 1998.

[118758]21023.

[118759]Toki, T.; Itoh, J.; Kitazawa, J.; Arai, K.; Hatakeyama, K.; Akasaka,

[118760]J.; Igarashi, K.; Nomura, N.; Yokoyama, M.; Yamamoto, M.; Ito, E.

[118761]: Human small Maf proteins form heterodimers with CNC family transcription

[118762]factors and recognize the NF-E2 motif. Oncogene 14: 1901–1910, 1997.

[118763]21024.

[118764]Yang, B.; He, B.; Abdel-Halim, S. M.; Tibell, A.; Brendel, M. D.;

[118765]Bretzel, R. G.; Efendic, S.; Hillert, J.: Molecular cloning of a

[118766]full-length cDNA for human type 3 adenylyl cyclase and its expression

[118767]in human islets. Biochem. Biophys. Res. Commun. 254: 548–551, 1999.

[118768]21025.

[118769]Chawengsaksophak, K.; James, R.; Hammond, V. E.; Kontgen, F.; Beck,

[118770]F.: Homeosis and intestinal tumours in Cdx2 mutant mice. Nature 386:

[118771]84–87, 1997.

[118772]21026.

[118773]Drummond, F.; Putt, W.; Fox, M.; Edwards, Y. H.: Cloning and chromosome

[118774]assignment of the human CDX2 gene. Ann. Hum. Genet. 61: 393–400,

[118775]1997.

[118776]21027.

[118777]German, M. S.; Wang, J.; Fernald, A. A.; Espinosa, R., III; Le

[118778]Beau, M. M.; Bell, G. I.: Localization of the genes encoding two

[118779]transcription factors, LMX1 and CDX3, regulating insulin gene expression

[118780]to human chromosomes 1 and 13. Genomics 24: 403–404, 1994.

[118781]21028.

- [118782]Woodford–Richens, K. L.; Halford, S.; Rowan, A.; Bevan, S.; Aaltonen, L. A.; Wasan, H.; Bicknell, D.; Bodmer, W. F.; Houlston, R. S.; Tomlinson, J. P. M.: CDX2 mutations do not account for juvenile polyposis or Peutz–Jeghers syndrome and occur infrequently in sporadic colorectal cancers. *Brit. J. Cancer* 84: 1314–1316, 2001.
- [118787]21029.
- [118788]Hiroi, Y.; Kudoh, S.; Monzen, K.; Ikeda, Y.; Yazaki, Y.; Nagai, R.; Komuro, I.: Tbx5 associates with Nkx2–5 and synergistically promotes cardiomyocyte differentiation. *Nature Genet.* 28: 276–280, 2001.
- [118791]21030.
- [118792]Dubbink, H. J.; Verkaik, N. S.; Faber, P. W.; Trapman, J.; Schroder, F. H.; Romijn, J. C.: Tissue-specific and androgen-regulated expression of human prostate-specific transglutaminase. *Biochem. J.*

315: 901-908,

[118795]1996.

[118796]21031.

[118797]Gentile, V.; Grant, F. J.; Porta, R.; Baldini, A.: Localization

[118798]of the human prostate transglutaminase (type IV) gene
(TGM4) to chromosome

[118799]3p21.33-p22 by fluorescence in situ hybridization. Ge-
nomics 27:

[118800]219-220, 1995.

[118801]21032.

[118802]Grant, F. J.; Taylor, D. A.; Sheppard, P. O.; Mathewes, S. L.;

[118803]Lint, W.; Vanaja, E.; Bishop, P. D.; O'Hara, P. J.: Molecular
cloning

[118804]and characterization of a novel transglutaminase cDNA
from a human

[118805]prostate cDNA library. Biochem. Biophys. Res. Commun.
203: 1117-1123,

[118806]1994.

[118807]21033.

[118808]Takai, S.; Long, J. E.; Yamada, K.; Miki, T.: Chromosomal

localization

[118809]of the human ECT2 proto-oncogene to 3q26.1-q26.2 by
somatic cell analysis

[118810]and fluorescence in situ hybridization. Genomics 27:
220-222, 1995.

[118811]21034.

[118812]Takai, S.; Lorenzi, M. V.; Long, J. E.; Yamada, K.; Miki, T.:

[118813]Assignment of the Ect2 protooncogene to mouse chromo-
some band 3B by

[118814]in situ hybridization. Cytogenet. Cell Genet. 81: 83-84,
1998.

[118815]21035.

[118816]Barker, H. M.; Brewis, N. D.; Street, A. J.; Spurr, N. K.; Co-
hen,

[118817]P. T. W.: Three genes for protein phosphatase 1 map to
different

[118818]human chromosomes: sequence, expression and gene lo-
calisation of protein

[118819]serine/threonine phosphatase 1 beta (PPP1CB). Biochim.
Biophys. Acta 1220:

[118820]212-218, 1994.

[118821]21036.

[118822]Saadat, M.; Kakinoki, Y.; Mizuno, Y.; Kikuchi, K.; Yoshida, M.

[118823]C.: Chromosomal localization of human, rat, and mouse protein phosphatase

[118824]type 1 beta catalytic subunit genes (PPP1CB) by fluorescence in situ

[118825]hybridization. Jpn. J. Genet. 69: 697–700, 1994.

[118826]21037.

[118827]Padilla, C. A.; Bajalica, S.; Lagercrantz, J.; Holmgren, A.: The

[118828]gene for human glutaredoxin (GLRX) is localized to human chromosome

[118829]5q14. Genomics 32: 455–457, 1996.

[118830]21038.

[118831]Padilla, C. A.; Martinez-Galisteo, E.; Barcena, J. A.; Spyrou,

[118832]G.; Holmgren, A.: Purification from placenta, amino acid sequence,

[118833]structure comparisons and cDNA cloning of human glutaredoxin. Europ.

[118834]J. Biochem. 227: 27–34, 1995.

[118835]21039.

[118836]Raghavachari, N.; Krysan, K.; Xing K.; Lou, M. F.: Regulation
of thioltransferase expression in human lens epithelial
cells. Invest.
Ophthalm. Vis. Sci. 42: 1002–1008, 2001.
21040.

[118840]Liang, B. T.; Jacobson, K. A.: A physiological role of the
adenosine
A3 receptor: sustained cardioprotection. Proc. Nat. Acad.
Sci. 95:
6995–6999, 1998.
21041.

[118844]Monitto, C. L.; Levitt, R. C.; DiSilvestre, D.; Holroyd, K. J.
Localization of the A(3) adenosine receptor gene
(ADORA3) to human
chromosome 1p. Genomics 26: 637–638, 1995.
21042.

[118848]Sajjadi, F. G.; Firestein, G. S.: cDNA cloning and sequence
analysis
of the human A3 adenosine receptor. Biochim. Biophys.
Acta 1179:

[118850]105–107, 1993.

[118851]21043.

[118852]Zhao, Z.; Ravid, S.; Ravid, K.: Chromosomal mapping of
the mouse

[118853]A3 adenosine receptor gene, Adora3. Genomics 30:
118–119, 1995.

[118854]21044.

[118855]Zhou, Q.-Y.; Li, C.; Olah, M. E.; Johnson, R. A.; Stiles, G. L.;

[118856]Civelli, O.: Molecular cloning and characterization of an
adenosine

[118857]receptor: the A3 adenosine receptor. Proc. Nat. Acad. Sci.
89: 7432–7436,

[118858]1992.

[118859]21045.

[118860]Corset, V.; Nguyen-Ba-Charvet, K. T.; Forcet, C.; Moyse,
E.; Chedotal,

[118861]A.; Mehlen, P.: Netrin–1–mediated axon outgrowth and
cAMP production

[118862]requires interaction with adenosine A2b receptor. Nature
407: 747–750,

[118863]2000.

[118864]21046.

[118865]Mino, R. P.; Spoerri, P. E.; Caballero, S.; Player, D.; Belardinelli,

[118866]L.; Biaggioni, I.; Grant, M. B.: Adenosine receptor antagonists and

[118867]retinal neovascularization in vivo. Invest. Ophthalm. Vis. Sci. 42:

[118868]3320–3324, 2001.

[118869]21047.

[118870]Stein, E.; Zou, Y.; Poo, M.; Tessier-Lavigne, M.: Binding of DCC

[118871]by netrin-1 to mediate axon guidance independent of adenosine A2B

[118872]receptor activation. Science 291: 1976–1982, 2001.

[118873]21048.

[118874]Townsend-Nicholson, A.; Baker, E.; Sutherland, G. R.; Schofield,

[118875]P. R.: Localization of the adenosine A2b receptor subtype gene (ADORA2B)

[118876]to chromosome 17p11.2–p12 by FISH and PCR screening of somatic cell

[118877]hybrids. Genomics 25: 605–607, 1995.

[118878]21049.

[118879]von Gall, C.; Garabette, M. L.; Kell, C. A.; Frenzel, S.; De-
hghani,

[118880]F.; Schumm–Draeger, P.–M.; Weaver, D. R.; Korf, H.–W.;
Hastings, M.

[118881]H.; Stehle, J. H.: Rhythmic gene expression in pituitary de-
pends

[118882]on heterologous sensitization by the neurohormone mela-
tonin. Nature

[118883]Neurosci. 5: 234–238, 2002.

[118884]21050.

[118885]Hirai, S.; Izawa, M.; Osada, S.; Spyrou, G.; Ohno, S.: Acti-
vation

[118886]of the JNK pathway by distantly related protein kinases,
MEKK and

[118887]MUK. Oncogene 12: 641–650, 1996.

[118888]21051.

[118889]Holzman, L. B.; Merritt, S. E.; Fan, G.: Identification,
molecular

[118890]cloning, and characterization of dual leucine zipper bear-

ing kinase.

[118891]A novel serine/threonine protein kinase that defines a second subfamily

[118892]of mixed lineage kinases. J. Biol. Chem. 269: 30808–30817, 1994.

[118893]21052.

[118894]Mata, M.; Merritt, S. E.; Fan, G.; Yu, G. G.; Holzman, L. B.:

[118895]Characterization of dual leucine zipper-bearing kinase, a mixed lineage

[118896]kinase present in synaptic terminals whose phosphorylation state is

[118897]regulated by membrane depolarization via calcineurin. J. Biol. Chem. 271:

[118898]16888–16896, 1996.

[118899]21053.

[118900]Reddy, U. R.; Nycum, L.; Slavic, I.; Biegel, J. A.: Localization

[118901]of the human zipper protein kinase gene (ZPK) to chromosome 12q13

[118902]by fluorescence in situ hybridization and somatic cell hybrid analysis. Genomics 25:

[118903]597–598, 1995.

[118904]21054.

[118905]Reddy, U. R.; Pleasure, D.: Cloning of a novel putative protein

[118906]kinase having a leucine zipper domain from human brain. Biochem.

[118907]Biophys. Res. Commun. 202: 613–620, 1994.

[118908]21055.

[118909]Becq, F.; Hamon, Y.; Bajetto, A.; Gola, M.; Verrier, B.; Chimini,

[118910]G.: ABC1, an ATP binding cassette transporter required for phagocytosis

[118911]of apoptotic cells, generates a regulated anion flux after expression

[118912]in *Xenopus laevis* oocytes. J. Biol. Chem. 272: 2695–2699, 1997.

[118913]21056.

[118914]Decottignies, A.; Goffeau, A.: Complete inventory of the yeast

[118915]ABC proteins. Nature Genet. 15: 137–145, 1997.

[118916]21057.

[118917]Guo, Z.; Inazu, A.; Yu, W.; Suzumura, T.; Okamoto, M.;

Nohara,

[118918]A.; Higashikata, T.; Sano, R.; Wakasugi, K.; Hayakawa, T.;
Yoshida,

[118919]K.; Suehiro, T.; Schmitz, G.; Mabuchi, H.: Double deletions
and missense

[118920]mutations in the first nucleotide-binding fold of the ATP-
binding

[118921]cassette transporter A1 (ABCA1) gene in Japanese patients
with Tangier

[118922]disease. J. Hum. Genet. 47: 325-329, 2002.

[118923]21058.

[118924]Hong, S. H.; Rhyne, J.; Zeller, K.; Miller, M.: Novel ABCA1
compound

[118925]variant associated with HDL cholesterol deficiency.
Biochim. Biophys.

[118926]Acta 1587: 60-64, 2002.

[118927]21059.

[118928]Huang, W.; Moriyama, K.; Koga, T.; Hua, H.; Ageta, M.;
Kawabata,

[118929]S.; Mawatari, K.; Imamura, T.; Eto, T.; Kawamura, M.; Ter-
amoto, T.;

[118930]Sasaki, J.: Novel mutations in ABCA1 gene in Japanese pa-

tients with

[118931]Tangier disease and familial high density lipoprotein deficiency with

[118932]coronary heart disease. *Biochim. Biophys. Acta* 1537: 71–78, 2001.

[118933]21060.

[118934]Ishii, J.; Nagano, M.; Kujiraoka, T.; Ishihara, M.; Egashira,

[118935]T.; Takada, D.; Tsuji, M.; Hattori, H.; Emi, M.: Clinical variant

[118936]of Tangier disease in Japan: mutation of the ABCA1 gene in hypoalphalipoproteinemia

[118937]with corneal lipidosis. *J. Hum. Genet.* 47: 366–369, 2002.

[118938]21061.

[118939]Jennings, M. W.; Jones, R. W.; Wood, W. G.; Weatherall, D. J.

[118940]: Analysis of an inversion within the human beta globin gene cluster. *Nucleic*

[118941]*Acids Res.* 13: 2897–2906, 1985.

[118942]21062.

[118943]Kulozik, A. E.; Bellan-Koch, A.; Kohne, E.; Kleihauer, E.: A

[118944]deletion/inversion rearrangement of the beta-globin gene

cluster in

[118945]a Turkish family with delta-beta(0)-thalassemia intermedia. Blood 79:

[118946]2455-2459, 1992.

[118947]21063.

[118948]Langmann, T.; Klucken, J.; Reil, M.; Liebisch, G.; Luciani, M.-F.;

[118949]Chimini, G.; Kaminski, W. E.; Schmitz, G.: Molecular cloning of the

[118950]human ATP-binding cassette transporter 1 (hABC1): evidence for sterol-dependent

[118951]regulation in macrophages. Biochem. Biophys. Res. Commun. 257: 29-33,

[118952]1999.

[118953]21064.

[118954]Lawn, R. M.; Wade, D. P.; Garvin, M. R.; Wang, X.; Schwartz, K.;

[118955]Porter, J. G.; Seilhamer, J. J.; Vaughan, A. M.; Oram, J. F.: The

[118956]Tangier disease gene product ABC1 controls the cellular apolipoprotein-mediated

[118957]lipid removal pathway. J. Clin. Invest. 104: R25-R31,

1999.

[118958]21065.

[118959]Luciani, M. F.; Denizot, F.; Savary, S.; Mattei, M. G.; Chi-
mini,

[118960]G.: Cloning of two novel ABC transporters mapping on
human chromosome

[118961]9. Genomics 21: 150–159, 1994.

[118962]21066.

[118963]Lapicka–Bodzioch, K.; Bodzioch, M.; Krull, M.; Kielar, D.;
Probst,

[118964]M.; Kiec, B.; Andrikovics, H.; Bottcher, A.; Hubacek, J.;
Aslanidis,

[118965]C.; Suttorp, N.; Schmitz, G.: Homogeneous assay based on
52 primer

[118966]sets to scan for mutations of the ABCA1 gene and its ap-
plication in

[118967]genetic analysis of a new patient with familial high-
density lipoprotein

[118968]deficiency syndrome. Biochim. Biophys. Acta 1537: 42–48,
2001.

[118969]21067.

[118970]Marcil, M.; Boucher, B.; Krimbou, L.; Solymoss, B. C.; Davi-

gnon,

[118971]J.; Frohlich, J.; Genest, J., Jr.: Severe familial HDL deficiency

[118972]in French-Canadian kindreds: clinical, biochemical, and
molecular

[118973]characterization. *Arterioscler. Thromb. Vasc. Biol.* 15:
1015-1024,

[118974]1995.

[118975]21068.

[118976]Marcil, M.; Yu, L.; Krimbou, L.; Boucher, B.; Oram, J. F.;
Cohn,

[118977]J. S.; Genest, J., Jr.: Cellular cholesterol transport and ef-
flux

[118978]in fibroblasts are abnormal in subjects with familial HDL
deficiency. *Arterioscler.*

[118979]*Thromb. Vasc. Biol.* 19: 159-169, 1999.

[118980]21069.

[118981]McNeish, J.; Aiello, R. J.; Guyot, D.; Turi, T.; Gabel, C.;
Aldinger,

[118982]C.; Hoppe, K. L.; Roach, M. L.; Royer, L. J.; de Wet, J.; Broc-
cardo,

[118983]C.; Chimini, G.; Francone, O. L.: High density lipoprotein

deficiency

[118984]and foam cell accumulation in mice with targeted disruption of ATP-binding

[118985]cassette transporter-1. Proc. Nat. Acad. Sci. 97: 4245-4250, 2000.

[118986]21070.

[118987]Pullinger, C. R.; Hakamata, H.; Duchateau, P. N.; Eng, C.; Aouizerat,

[118988]B. E.; Cho, M. H.; Fielding, C. J.; Kane, J. P.: Analysis of hABC1

[118989]gene 5-prime end: additional peptide sequence, promoter region, and

[118990]four polymorphisms. Biochem. Biophys. Res. Commun. 271: 451-455,

[118991]2000.

[118992]21071.

[118993]Petit, M. M. R.; Schoenmakers, E. F. P. M.; Huysmans, C.; Geurts,

[118994]J. M. W.; Mandahl, N.; Van de Ven, W. J. M.: LHFP, a novel translocation

[118995]partner gene of HMGIC in a lipoma, is a member of a new family of

[118996]LHFP-like genes. Genomics 57: 438–441, 1999.

[118997]21072.

[118998]Nishita, Y.; Yoshida, I.; Sado, T.; Takagi, N.: Genomic imprinting

[118999]and chromosomal localization of the human MEST gene. Genomics 36:

[119000]539–542, 1996.

[119001]21073.

[119002]Kohen, R.; Metcalf, M. A.; Khan, N.; Druck, T.; Huebner, K.; Lachowicz,

[119003]J. E.; Meltzer, H. Y.; Sibley, D. R.; Roth, B. L.; Hamblin, M. W.

[119004]: Cloning, characterization, and chromosomal localization of a human

[119005]5-HT-6 serotonin receptor. J. Neurochem. 66: 47–56, 1996.

[119006]21074.

[119007]Pearce, S. H. S.; Williamson, C.; Kifor, O.; Bai, M.; Coulthard,

[119008]M. G.; Davies, M.; Lewis-Barned, N.; McCredie, D.; Powell, H.; Kendall-Taylor,

- [119009]P.; Brown, E. M.; Thakker, R. V.: A familial syndrome of hypocalcemia with hypercalciuria due to mutations in the calcium-sensing receptor. New Eng. J. Med. 335: 1115–1122, 1996.
- [119012]21075.
- [119013]Frank, J.; Aita, V. M.; Ahmad, W.; Lam, H.; Wolff, C.; Christiano, A. M.: Identification of a founder mutation in the protoporphyrinogen oxidase gene in variegate porphyria patients from Chile. Hum. Hered. 51: 160–168, 2001.
- [119017]21076.
- [119018]Frank, J.; Jugert, F. K.; Breitkopf, C.; Goerz, G.; Merk, H. F.; Christiano, A. M.: Recurrent missense mutation in the protoporphyrinogen oxidase gene underlies variegate porphyria. Am. J. Med. Genet. 79: 22–26, 1998.
- [119022]21077.

- [119023] Orso, E.; Broccardo, C.; Kaminski, W. E.; Bottcher, A.;
Liebisch,
[119024] G.; Drobnik, W.; Gotz, A.; Chambenoit, O.; Diederich, W.;
Langmann,
[119025] T.; Spruss, T.; Luciani, M.-F.; Rothe, G.; Lackner, K. J.; Chi-
mini,
[119026] G.; Schmitz, G.: Transport of lipids from Golgi to plasma
membrane
[119027] is defective in Tangier disease patients and Abc1-deficient
mice. Nature
[119028] Genet. 24: 192-196, 2000.
[119029] 21078.
[119030] Hift, R. J.; Meissner, P. N.; Corrigall, A. V.; Ziman, M. R.;
Petersen,
[119031] L. A.; Meissner, D. M.; Davidson, B. P.; Sutherland, J.; Dai-
ley, H.
[119032] A.; Kirsch, R. E.: Variegate porphyria in South Africa,
1688-1996:
[119033] new developments in an old disease. S. Afr. Med. J. 87:
722-731,
[119034] 1997.
[119035] 21079.

- [119036]Hift, R. J.; Meissner, P. N.; Todd, G.; Kirby, P.; Bilsland, D.;
- [119037]Collins, P.; Ferguson, J.; Moore, M. R.: Homozygous variegate porphyria:
- [119038]an evolving clinical syndrome. *Postgrad. Med. J.* 69: 781–786, 1993.
- [119039]21080.
- [119040]Lam, H.; Dragan, L.; Tsou, H. C.; Merk, H.; Peacocke, M.; Goerz,
- [119041]G.; Sassa, S.; Poh-Fitzpatrick, M.; Bickers, D. R.; Christiano, A.
- [119042]M.: Molecular basis of variegated porphyria: a de novo insertion mutation
- [119043]in the protoporphyrinogen oxidase gene. *Hum. Genet.* 99: 126–129,
- [119044]1997.
- [119045]21081.
- [119046]Puy, H.; Robreau, A.-M.; Rosipal, R.; Nordmann, Y.; Deybach, J.-C.
- [119047]: Protoporphyrinogen oxidase: complete genomic sequence and polymorphisms
- [119048]in the human gene. *Biochem. Biophys. Res. Comm.* 226: 226–230, 1996.

[119049]21082.

[119050]Roberts, A. G.; Puy, H.; Dailey, T. A.; Morgan, R. R.; Whatley,

[119051]S. D.; Dailey, H. A.; Martasek, P.; Nordmann, Y.; Deybach, J.-C.;

[119052]Elder, G. H.: Molecular characterization of homozygous variegate

[119053]porphyria. Hum. Molec. Genet. 7: 1921-1925, 1998.

[119054]21083.

[119055]Francavilla, A.; Hagiya, M.; Porter, K. A.; Polimeno, L.; Ihara,

[119056]J.; Starzl, T. E.: Augmenter of liver regeneration: its place in

[119057]the universe of hepatic growth factors. Hepatology 20: 747-757,

[119058]1994.

[119059]21084.

[119060]Giorda, R.; Hagiya, M.; Seki, T.; Shimonishi, M.; Sakai, H.; Michaelson,

[119061]J.; Francavilla, A.; Starzl, T. E.; Trucco, M.: Analysis of the structure

- [119062]and expression of the augmenter of liver regeneration
(ALR) gene. Molec.
- [119063]Med. 2: 97–108, 1996.
- [119064]21085.
- [119065]Hagiya, M.; Francavilla, A.; Polimeno, L.; Ihara, I.; Sakai, H.;
- [119066]Seki, T.; Shimonishi, M.; Porter, K. A.; Starzl, T. E.: Cloning
and
- [119067]sequence analysis of the rat augmenter of liver regenera-
tion (ALR)
- [119068]gene: expression of biologically active recombinant ALR
and demonstration
- [119069]of tissue distribution. Proc. Nat. Acad. Sci. 91:
8142–8146, 1994.
- [119070]Note: Erratum Proc. Nat. Acad. Sci. 92, p. 3076 only,
1995.
- [119071]21086.
- [119072]Lisowsky, T.; Weinstat–Saslow, D. L.; Barton, N.; Reeders,
S. T.;
- [119073]Schneider, M. C.: A new human gene located in the PKD1
region of
- [119074]chromosome 16 is a functional homologue to ERV1 of
yeast. Genomics 29:

[119075]690-697, 1995.

[119076]21087.

[119077]Honda, H.; Inazawa, J.; Nishida, J.; Yazaki, Y.; Hirai, H.:

Molecular

[119078]cloning, characterization, and chromosomal localization
of a novel

[119079]protein-tyrosine phosphatase, HPTP eta. Blood 84:
4186-4194, 1994.

[119080]21088.

[119081]Ostman, A.; Yang, Q.; Tonks, N. K.: Expression of DEP-1, a
receptor-like

[119082]protein-tyrosine-phosphatase, is enhanced with increas-
ing cell density. Proc.

[119083]Nat. Acad. Sci. 91: 9680-9684, 1994.

[119084]21089.

[119085]Jacoby, A. S.; Webb, G. C.; Liu, M. L.; Kofler, B.; Hort, Y. J.;

[119086]Fathi, Z.; Bottema, C. D. K.; Shine, J.; Iismaa, T. P.: Struc-
tural

[119087]organization of the mouse and human GALR1 galanin re-
ceptor genes (Galnr

[119088]and GALNR) and chromosomal localization of the mouse

gene. Genomics 45:

[119089]496–508, 1997.

[119090]21090.

[119091]Tomoda, T.; Kurashige, T.; Moriki, T.; Yamamoto, H.; Fujimoto,

[119092]S.; Taniguchi, T.: Enhanced expression of
poly(ADP-ribose) synthetase

[119093]gene in malignant lymphoma. Am. J. Hemat. 37: 223–227,
1991.

[119094]21091.

[119095]Vasquez, K. M.; Marburger, K.; Intody, Z.; Wilson, J. H.:
Manipulating

[119096]the mammalian genome by homologous recombination.
Proc. Nat. Acad.

[119097]Sci. 98: 8403–8410, 2001.

[119098]21092.

[119099]Yu, S.-W.; Wang, H.; Poitras, M. F.; Coombs, C.; Bowers, W.
J.;

[119100]Federoff, H. J.; Poirier, G. G.; Dawson, T. M.; Dawson, V. L.:
Mediation

[119101]of poly(ADP-ribose) polymerase-1-dependent cell death

by apoptosis-inducing

[119102]factor. Science 297: 259–263, 2002.

[119103]21093.

[119104]Zabel, B. U.; Herzog, H.; Schneider, R.; Auer, B.; Hirsch-Kauffmann,

[119105]M.; Schweiger, M.: Chromosomal sublocalization of the gene for human

[119106]poly (ADP-ribose) polymerase (NAD⁺ ADP-ribosyltransferase) at 1q41–42.

[119107](Abstract) Cytogenet. Cell Genet. 51: 1115, 1989.

[119108]21094.

[119109]Davidson, M. K.; Le Beau, M. M.; Eddy, R. L.; Shows, T. B.; DiPietro,

[119110]L. A.; Kingzette, M.; Hanly, W. C.: Genetic mapping of the human

[119111]polymeric immunoglobulin receptor gene to chromosome region 1q31–q41. Cytogenet.

[119112]Cell Genet. 48: 107–111, 1988.

[119113]21095.

[119114]Hood, L.; Kronenberg, M.; Hunkapiller, T.: T cell antigen receptors

[119115]and the immunoglobulin supergene family. Cell 40:
225–229, 1985.

[119116]21096.

[119117]Krajci, P.; Gedde–Dahl, T., Jr.; Hoyheim, B.; Rogde, S.;
Olaisen,

[119118]B.; Brandtzaeg, P.: The gene encoding human transmem-
brane secretory

[119119]component (locus PIGR) is linked to D1S58 on chromo-
some 1. Hum. Genet. 90:

[119120]215–219, 1992.

[119121]21097.

[119122]Krajci, P.; Grzeschik, K. H.; Geurts van Kessel, A. H. M.;
Olaisen,

[119123]B.; Brandtzaeg, P.: The human transmembrane secretory
component (poly-Ig

[119124]receptor): molecular cloning, restriction fragment length
polymorphism

[119125]and chromosomal sublocalization. Hum. Genet. 87:
642–648, 1991.

[119126]21098.

[119127]Krajci, P.; Kvale, D.; Tasken, K.; Brandtzaeg, P.: Molecular
cloning

[119128]and exon–intron mapping of the gene encoding human transmembrane secretory

[119129]component (the poly-Ig receptor). *Europ. J. Immun.* 22: 2309–2315,

[119130]1992.

[119131]21099.

[119132]Zhang, J.–R.; Mostov, K. E.; Lamm, M. E.; Nanno, M.; Shimida, S.;

[119133]Ohwaki, M.; Tuomanen, E.: The polymeric immunoglobulin receptor translocates

[119134]pneumococci across human nasopharyngeal epithelial cells. *Cell* 102:

[119135]827–837, 2000.

[119136]21100.

[119137]Bear, J. C.; McManamon, P.; Morgan, J.; Payne, R. H.; Lewis, H.;

[119138]Gault, M. H.; Churchill, D. N.: Age at clinical onset and at ultrasonographic

[119139]detection of adult polycystic kidney disease: data for genetic counselling. *Am.*

[119140]*J. Med. Genet.* 18: 45–53, 1984.

[119141]21101.

[119142]Bear, J. C.; Parfrey, P. S.; Morgan, J. M.; Martin, C. J.;
Cramer,

[119143]B. C.: Autosomal dominant polycystic kidney disease: new
information

[119144]for genetic counselling. Am. J. Med. Genet. 43: 548-553,
1992.

[119145]21102.

[119146]Boichis, H.; Passwell, J.; David, R.; Miller, H.: Congenital
hepatic

[119147]fibrosis and nephronophthisis: a family study. Quart. J.
Med. 42:

[119148]221-233, 1973.

[119149]21103.

[119150]Breuning, M. H.; Snijdwint, F. G. M.; Dauwerse, J. G.;
Saris,

[119151]J. J.; Bakker, E.; Pearson, P. L.; van Ommen, G. J. B.: Two
step

[119152]procedure for early diagnosis of polycystic kidney disease
with polymorphic

[119153]DNA markers on both sides of the gene. J. Med. Genet. 27:
614-617,

[119154]1990.

[119155]21104.

[119156]Breuning, M. H.; Snijdwint, F. G. M.; Brunner, H.; Verwest, A.;

[119157]Ijdo, J. W.; Saris, J. J.; Dauwerse, J. G.; Blondin, L.; Keith, T.;

[119158]Callen, D. F.; Hyland, V. J.; Xiao, G. H.; Scherer, G.; Higgs, D.

[119159]R.; Harris, P.; Bachner, L.; Reeders, S. T.; Germino, G.; Pearson,

[119160]P. L.; van Ommen, G. J. B.: Map of 16 polymorphic loci on the short

[119161]arm of chromosome 16 close to the polycystic kidney disease gene (PKD1). J.

[119162]Med. Genet. 27: 603-613, 1990.

[119163]21105.

[119164]Burn, T. C.; Connors, T. D.; Dackowski, W. R.; Petry L. R.; Van

[119165]Raay, T. J.; Millholland, J. M.; Venet, M.; Miller, G.; Hakim, R.

[119166]M.; Landes, G. M.; Klinger, K. W.; Qian, F.; Onuchic, L. F.; Watnick,

- [119167]T.; Germino, G. G.; Doggett, N. A.: Analysis of the genomic sequence
for the autosomal dominant polycystic kidney disease (PKD1) gene predicts
the presence of a leucine-rich repeat. Hum. Molec. Genet. 4: 575-582,
1995.
- [119170]21106.
- [119171]21106.
- [119172]Chanmugam, D.; Rasaretnam, R.; Karunaratne, K. E. S.: Hereditary
spherocytosis and polycystic disease of the kidneys in four members
of a family. Am. J. Hum. Genet. 23: 66, 1971.
- [119174]21107.
- [119175]21107.
- [119176]Coto, E.; Aguado, S.; Alvarez, J.; Menendez Diaz, M. J.; Lopez-Larrea,
C.: Genetic and clinical studies in autosomal dominant polycystic
kidney disease type 1 (ADPKD1). J. Med. Genet. 29: 243-246, 1992.
- [119177]21108.
- [119179]21108.

[119180] de Villiers, J. N. P.; Hillermann, R.; Loubser, L.; Kotze, M. J.

[119181]: Spectrum of mutations in the HFE gene implicated in
haemochromatosis

[119182] and porphyria. Hum. Molec. Genet. 8: 1517–1522, 1999.

[119183] 21109.

[119184] Deybach, J.-C.; Puy, H.; Robreau, A.-M.; Lamoril, J.; Da
Silva,

[119185] V.; Grandchamp, B.; Nordmann, Y.: Mutations in the pro-
toporphyrinogen

[119186] oxidase gene in patients with variegate porphyria. Hum.
Molec. Genet. 5:

[119187] 407–410, 1996.

[119188] 21110.

[119189] Meissner, P. N.; Dailey, T. A.; Hift, R. J.; Ziman, M.; Corri-
gall,

[119190] A. V.; Roberts, A. G.; Meissner, D. M.; Kirsch, R. E.; Dailey,
H.

[119191] A.: A R59W mutation in human protoporphyrinogen oxi-
dase results in

[119192] decreased enzyme activity and is prevalent in South
Africans with

[119193]variegate porphyria. Nature Genet. 13: 95–97, 1996.

[119194]21111.

[119195]Kalman, K.; Nguyen, A.; Tseng–Crank, J.; Dukes, I. D.;
Chandy,

[119196]G.; Hustad, C. M.; Copeland, N. G.; Jenkins, N. A.; Mohren–
weiser,

[119197]H.; Brandriff, B.; Cahalan, M.; Gutman, G. A.; Chandy, K.
G.: Genomic

[119198]organization, chromosomal localization, tissue distribu–
tion, and biophysical

[119199]characterization of a novel mammalian Shaker–related
voltage–gated

[119200]potassium channel, Kv1.7. J. Biol. Chem. 273: 5851–5857,
1998.

[119201]21112.

[119202]Orkiszewski, K. G.; Tedesco, T. A.; Mellman, W. J.; Croce,
C.

[119203]M.: Linkage relationship between the genes for thymidine
kinase and

[119204]galactokinase in different primates. Somat. Cell Genet. 2:
21–26,

[119205]1976.

[119206]21113.

[119207]Dunham, I.; Collins, J.; Wade, R.; Scambler, P.: Possible
role

[119208]for COMT in psychosis associated with velo-cardio-facial
syndrome.

[119209](Letter) Lancet 340: 1361-1362, 1992.

[119210]21114.

[119211]Pulkka, A.; Ihalainen, R.; Suorsa, A.; Riviere, M.; Szpirer, J.;

[119212]Pajunen, A.: Structures and chromosomal localizations of
two rat

[119213]genes encoding S-adenosylmethionine decarboxylase.
Genomics 16:

[119214]342-349, 1993.

[119215]21115.

[119216]Radford, D. M.; Eddy, R.; Haley, L.; Henry, W. M.; Pegg, A.
E.;

[119217]Pajunen, A.; Shows, T. B.: Gene sequences coding for S-
adenosylmethionine

[119218]decarboxylase are present on human chromosome 6 and
the X and are

[119219]not amplified in colon neoplasia. Cytogenet. Cell Genet.

49: 285–288,

[119220]1989.

[119221]21116.

[119222]Kleymenova, E.; Ibraghimov-Beskrovnaya, O.; Kugoh, H.;
Everitt,

[119223]J.; Xu, H.; Kiguchi, K.; Landes, G.; Harris, P.; Walker, C.:

Tuberin-dependent

[119224]membrane localization of polycystin-1: a functional link
between polycystic

[119225]kidney disease and the TSC2 tumor suppressor gene.

Molec. Cell 7:

[119226]823–832, 2001.

[119227]21117.

[119228]Sandford, R.; Sgotto, B.; Burn, T.; Brenner, S.: The tuberin

[119229](TSC2), autosomal dominant polycystic kidney disease
(PKD1), and somatostatin

[119230]type V receptor (SSTR5) genes form a synteny group in the
Fugu genome. Genomics 38:

[119231]84–86, 1996.

[119232]21118.

[119233]Pedeutour, F.; Szpirer, C.; Nahon, J.–L.: Assignment of the

human

[119234]pro-melanin-concentrating hormone gene (PMCH) to
chromosome 12q23-q24

[119235]and two variant genes (PMCHL1 and PMCHL2) to chromo-
some 5p14 and 5q12-q13.

[119236]Genomics 19: 31-37, 1994.

[119237]21119.

[119238]Chen, S.-J.; Zelent, A.; Tong, J.-H.; Yu, H.-Q.; Wang, Z.-Y.;
Derre,

[119239]J.; Berger, R.; Waxman, S.; Chen, Z.: Rearrangements of the retinoic acid receptor alpha and promyelocytic leukemia zinc finger genes resulting from t(11;17)(q23;q21) in a patient with acute promyelocytic leukemia. J. Clin. Invest. 91: 2260–2267, 1993.

[119243]21120.

[119244]Chen, Z.; Brand, N. J.; Chen, A.; Chen, S.-J.; Tong, J.-H.; Wang, Z.-Y.; Waxman, S.; Zelent, A.: Fusion between a novel Kruppel-like zinc finger gene and the retinoic acid receptor-alpha locus due to a variant t(11;17) translocation associated with acute promyelocytic leukaemia. EMBO J. 12: 1161–1167, 1993.

[119249]21121.

[119250]Chen, Z.; Guidez, F.; Rousselot, P.; Agadir, A.; Chen, S.-J.; Wang, Z.-Y.; Degos, L.; Zelent, A.; Waxman, S.; Chomienne, C.: PLZF-RAR-alpha

- [119252] fusion proteins generated from the variant
t(11;17)(q23;q21) translocation
- [119253] in acute promyelocytic leukemia inhibit ligand-dependent
transactivation
- [119254] of wild-type retinoic acid receptors. Proc. Nat. Acad. Sci.
91:
1178-1182, 1994.
- [119256] 21122.
- [119257] Grignani, F.; De Matteis, S.; Nervi, C.; Tomassoni, L.; Gel-
metti,
[119258] V.; Cioce, M.; Fanelli, M.; Ruthardt, M.; Ferrara, F. F.; Za-
mir, I.;
[119259] Seiser, C.; Grignani, F.; Lazar, M. A.; Minucci, S.; Pelicci, P.
G.
- [119260]: Fusion proteins of the retinoic acid receptor-alpha re-
cruit histone
- [119261] deacetylase in promyelocytic leukaemia. Nature 391:
815-818, 1998.
- [119262] 21123.
- [119263] Lin, R. J.; Nagy, L.; Inoue, S.; Shao, W.; Miller, W. H., Jr.;
- [119264] Evans, R. M.: Role of the histone deacetylase complex in
acute promyelocytic

[119265]leukaemia. Nature 391: 811–814, 1998.

[119266]21124.

[119267]Burstedt, M. S.; Sandgren, O.; Holmgren, G.; Forsman-Semb, K.:

[119268]Bothnia dystrophy caused by mutations in the cellular retinaldehyde-binding

[119269]protein gene (RLBP1) on chromosome 15q26. Invest. Ophthalmol. Vis. Sci. 40:

[119270]995–1000, 1999.

[119271]21125.

[119272]Fournier, C. M.; Nicolas, G.; Gallagher, P. G.; Dhermy, D.; Grandchamp,

[119273]B.; Lecomte, M.-C.: Spectrin St Claude, a splicing mutation of the

[119274]human alpha-spectrin gene associated with severe poikilocytic anemia. Blood 89:

[119275]4584–4590, 1997.

[119276]21126.

[119277]Unger, S.; Korkko, J.; Krakow, D.; Lachman, R. S.; Rimoin, D.

[119278]L.; Cohn, D. H.: Double heterozygosity for pseudoachon-

droplasia and

[119279]spondyloepiphyseal dysplasia congenita. Am. J. Med. Genet. 104:

[119280]140–146, 2001.

[119281]21127.

[119282]Burstedt, M. S. I.; Forsman-Semb, K.; Golovleva, I.; Janunger,

[119283]T.; Wachtmeister, L.; Sandgren, O.: Ocular phenotype of Bothnia dystrophy,

[119284]an autosomal recessive retinitis pigmentosa associated with an R234W

[119285]mutation in the RLBP1 gene. Arch Ophthal. 119: 260–267, 2001.

[119286]21128.

[119287]Crabb, J. W.; Goldflam, S.; Harris, S. E.; Saari, J. C.: Cloning

[119288]of the cDNAs encoding the cellular retinaldehyde-binding protein from

[119289]bovine and human retina and comparison of the protein structures. J.

[119290]Biol. Chem. 263: 18688–18692, 1988.

[119291]21129.

[119292]Eichers, E. R.; Green, J. S.; Stockton, D. W.; Jackman, C. S.;

[119293]Whelan, J.; McNamara, J. A.; Johnson, G. J.; Lupski, J. R.;
Katsanis,

[119294]N.: Newfoundland rod-cone dystrophy, an early-onset
retinal dystrophy,

[119295]is caused by splice-junction mutations in RLBP1. Am. J.
Hum. Genet. 70:

[119296]955-964, 2002.

[119297]21130.

[119298]Granse, L.; Abrahamson, M.; Ponjavic, V.; Andreasson, S.:
Electrophysiological

[119299]findings in two young patients with Bothnia dystrophy and
a mutation

[119300]in the RLBP1 gene. Ophthal. Genet. 22: 97-105, 2001.

[119301]21131.

[119302]Intres, R.; Goldflam, S.; Cook, J. R.; Crabb, J. W.: Molecular

[119303]cloning and structural analysis of the human gene encod-
ing cellular

[119304]retinaldehyde-binding protein. J. Biol. Chem. 269:
25411-25418,

[119305]1994.

[119306]21132.

[119307]Katsanis, N.; Shroyer, N. F.; Lewis, R. A.; Cavender, J. C.;
Al-Rajhi,

[119308]A. A.; Jabak, M.; Lupski, J. R.: Fundus albipunctatus and
retinitis

[119309]punctata albescens in a pedigree with an R150Q mutation
in RLBP1. Clin.

[119310]Genet. 59: 424–429, 2001.

[119311]21133.

[119312]Maw, M. A.; Kennedy, B.; Knight, A.; Bridges, R.; Roth, K.
E.;

[119313]Mani, E. J.; Mukkadan, J. K.; Nancarrow, D.; Crabb, J. W.;
Denton,

[119314]M. J.: Mutation of the gene encoding cellular retinalde-
hyde-binding

[119315]protein in autosomal recessive retinitis pigmentosa. Na-
ture Genet. 17:

[119316]198–200, 1997.

[119317]21134.

[119318]Morimura, H.; Berson, E. L.; Dryja, T. P.: Recessive muta-
tions

[119319]in the RLBP1 gene encoding cellular retinaldehyde-binding protein

[119320]in a form of retinitis punctata albescens. Invest. Ophthalm. Vis.

[119321]Sci. 40: 1000–1004, 1999.

[119322]21135.

[119323]Sparkes, R. S.; Heinzmann, C.; Goldflam, S.; Kojis, T.; Saari,

[119324]J. C.; Mohandas, T.; Klisak, I.; Bateman, J. B.; Crabb, J. W.: Assignment

[119325]of the gene (RLBP1) for cellular retinaldehyde-binding protein (CRALBP)

[119326]to human chromosome 15q26 and mouse chromosome 7. Genomics 12: 58–62,

[119327]1992.

[119328]21136.

[119329]Blanton, S. H.; Heckenlively, J. R.; Cottingham, A. W.; Friedman,

[119330]J.; Sadler, L. A.; Wagner, M.; Friedman, L. H.; Daiger, S. P.: Linkage

[119331]mapping of autosomal dominant retinitis pigmentosa (RP1) to the pericentric

[119332]region of human chromosome 8. Genomics 11: 857–869, 1991.

[119333]21137.

[119334]Farrar, G. J.; McWilliam, P.; Bradley, D. G.; Kenna, P.; Lawler,

[119335]M.; Sharp, E. M.; Humphries, M. M.; Eiberg, H.; Conneally, P. M.;

[119336]Traflet, J. A.; Humphries, P.: Autosomal dominant retinitis pigmentosa:

[119337]linkage to rhodopsin and evidence for genetic heterogeneity. Genomics 8:

[119338]35–40, 1990.

[119339]21138.

[119340]Wang, S.-Q.; Song, L.-S.; Lakatta, E. G.; Cheng, H.: Ca(2+) signalling

[119341]between single L-type Ca(2+) channels and ryanodine receptors in heart

[119342]cells. Nature 410: 592–596, 2001.

[119343]21139.

[119344]Laitinen, P. J.; Brown, K. M.; Piippo, K.; Swan, H.; Devaney, J.

- [119345]M.; Brahmabhatt, B.; Donarum, E. A.; Marino, M.; Tiso, N.; Viitasalo,
[119346]M.; Toivonen, L.; Stephan, D. A.; Kontula, K.: Mutations of
the cardiac
[119347]ryanodine receptor (RyR2) gene in familial polymorphic
ventricular
[119348]tachycardia. Circulation 103: 485–490, 2001.
[119349]21140.
- [119350]Marx, S. O.; Reiken, S.; Hisamatsu, Y.; Jayaraman, T.;
Burkhoff,
[119351]D.; Rosemblyt, N.; Marks, A. R.: PKA phosphorylation dis-
sociates
[119352]FKBP12.6 from the calcium release channel (ryanodine re-
ceptor): defective
[119353]regulation in failing hearts. Cell 101: 365–376, 2000.
[119354]21141.
- [119355]Mattei, M. G.; Giannini, G.; Moscatelli, F.; Sorrentino, V.:
Chromosomal
[119356]localization of murine ryanodine receptor genes RYR1,
RYR2, and RYR3
[119357]by in situ hybridization. Genomics 22: 202–204, 1994.

[119358]21142.

[119359]Otsu, K.; Willard, H. F.; Khanna, V. K.; Zorzato, F.; Green, N.

[119360]M.; MacLennan, D. H.: Molecular cloning of cDNA encoding the Ca^{2+}

[119361]release channel (ryanodine receptor) of rabbit cardiac muscle sarcoplasmic

[119362]reticulum. J. Biol. Chem. 265: 13472-13483, 1990.

[119363]21143.

[119364]Priori, S. G.; Napolitano, C.; Tiso, N.; Memmi, M.; Vignati, G.;

[119365]Bloise, R.; Sorrentino, V.; Danieli, G. A.: Mutations in the cardiac

[119366]ryanodine receptor gene (hRyR2) underlie catecholaminergic polymorphic

[119367]ventricular tachycardia. Circulation 102: r49-r53, 2000.

[119368]21144.

[119369]Zorzato, F.; Fujii, J.; Otsu, K.; Phillips, M.; Green, N. M.; Lai,

[119370]F. A.; Meissner, G.; MacLennan, D. H.: Molecular cloning of cDNA

- [119371] encoding human and rabbit forms of the Ca^{2+} release channel (ryanodine
- [119372] receptor) of skeletal muscle sarcoplasmic reticulum. J. Biol. Chem. 265:
- [119373] 2244–2256, 1990.
- [119374] 21145.
- [119375] Bertocchini, F.; Ovitt, C. E.; Conti, A.; Barone, V.; Scholer, H. R.; Bottinelli, R.; Reggiani, C.; Sorrentino, V.: Requirement
- [119376] for the ryanodine receptor type 3 for efficient contraction in neonatal
- [119377] skeletal muscles. EMBO J. 16: 6956–6963, 1997.
- [119378] 21146.
- [119380] Futatsugi, A.; Kato, K.; Ogura, H.; Li, S.-T.; Nagata, E.; Kuwajima,
- [119381] G.; Tanaka, K.; Itohara, S.; Mikoshiba, K.: Facilitation of NMDAR-independent
- [119382] LTP and spatial learning in mutant mice lacking ryanodine receptor
- [119383] type 3. Neuron 24: 701–713, 1999.
- [119384] 21147.

- [119385] Hakamata, Y.; Nakai, J.; Takeshima, H.; Imoto, K.: Primary structure
and distribution of a novel ryanodine receptor/calcium release channel
from rabbit brain. FEBS Lett. 312: 229–235, 1992.
21148.
- [119389] Leeb, T.; Brenig, B.: cDNA cloning and sequencing of the human
ryanodine receptor type 3 (RYR3) reveals a novel alternative splice
site in the RYR3 gene. FEBS Lett. 423: 367–370, 1998.
21149.
- [119393] Sorrentino, V.; Giannini, G.; Malzac, P.; Mattei, M. G.: Localization
of a novel ryanodine receptor gene (RYR3) to human chromosome 15q14–q15
by in situ hybridization. Genomics 18: 163–165, 1993.
21150.
- [119397] Takeshima, H.; Ikemoto, T.; Nishi, M.; Nishiyama, N.; Shimuta,
M.; Sugitani, Y.; Kuno, J.; Saito, I.; Saito, H.; Endo, M.; Iino,

- [119399]M.; Noda, T.: Generation and characterization of mutant mice lacking
- [119400]ryanodine receptor type 3. J. Biol. Chem. 271: 19649–19652, 1996.
- [119401]21151.
- [119402]Tiso, N.; Stephan, D. A.; Nava, A.; Bagattin, A.; Devaney, J. M.;
- [119403]Stanchi, F.; Larderet, G.; Brahmbhatt, B.; Brown, K.; Bauce, B.; Muriago,
- [119404]M.; Basso, C.; Thiene, G.; Danieli, G. A.; Rampazzo, A.: Identification
- [119405]of mutations in the cardiac ryanodine receptor gene in families affected
- [119406]with arrhythmogenic right ventricular cardiomyopathy type 2 (ARVD2). Hum.
- [119407]Molec. Genet. 10: 189–194, 2001.
- [119408]21152.
- [119409]Gallagher, P. G.; Forget, B. G.: Hematologically important mutations:
- [119410]spectrin and ankyrin variants in hereditary spherocytosis. Blood
- [119411]Cells Molec. Dis. 24: 539–543, 1998.

[119412]21153.

[119413]Gallagher, P. G.; Forget, B. G.: Hematologically important mutations:

[119414]spectrin variants in hereditary elliptocytosis and hereditary pyropoikilocytosis. Blood

[119415]Cells Molec. Dis. 22: 254–258, 1996.

[119416]21154.

[119417]Gallagher, P. G.; Forget, B. G.: Spectrin genes in health and disease. Seminars Hemat. 30: 4–20, 1993.

[119419]21155.

[119420]Gallagher, P. G.; Tse, W. T.; Coetzer, T.; Lecomte, M.–C.; Garbarz,

[119421]M.; Zarkowsky, H. S.; Baruchel, A.; Ballas, S. K.; Dhermy, D.; Palek,

[119422]J.; Forget, B. G.: A common type of the spectrin alpha-I 46–50a–kD

[119423]peptide abnormality in hereditary elliptocytosis and pyropoikilocytosis

[119424]is associated with a mutation distant from the proteolytic cleavage

[119425]site: evidence for the functional importance of the triple

helical

[119426]model of spectrin. J. Clin. Invest. 89: 892–898, 1992.

[119427]21156.

[119428]Gallagher, P. G.; Tse, W. T.; Marchesi, S. L.; Zarkowsky, H. S.;

[119429]Forget, B. G.: A defect in alpha-spectrin mRNA accumulation in hereditary

[119430]pyropoikilocytosis. Trans. Assoc. Am. Phys. 104: 32–39, 1991.

[119431]21157.

[119432]Garbarz, M.; Lecomte, M.-C.; Feo, C.; Devaux, I.; Picat, C.; Lefebvre,

[119433]C.; Galibert, F.; Gautero, H.; Bournier, O.; Galand, C.; Forget, B.

[119434]G.; Boivin, P.; Dhermy, D.: Hereditary pyropoikilocytosis and elliptocytosis

[119435]in a white French family with the spectrin alpha(I/74) variant related

[119436]to a CGT to CAT codon change (arg to his) at position 22 of the spectrin

[119437]alpha-I domain. Blood 75: 1691–1698, 1990.

[119438]21158.

[119439]Garbarz, M.; Lecomte, M. C.; Dhermy, D.; Feo, C.; Chave-
roche,

[119440]I.; Gautero, H.; Bournier, O.; Picat, C.; Goepp, A.; Boivin,
P.:

[119441]Double inheritance of an alpha I/65 spectrin variant in a
child with

[119442]homozygous elliptocytosis. Blood 67: 1661–1667, 1986.

[119443]21159.

[119444]Garbarz, M.; Lecomte, M. C.; Lefebvre, C.; Devaux, I.; Gal-
ibert,

[119445]F.; Boivin, P.; Forget, B. G.; Dhermy, D.: The DNAs from HE
and HPP

[119446]patients in a Caucasian French family with the spectrin al-
pha-I/74-kD

[119447]variant contain a CGT-to-CAT codon change (arg-to-his)
at position

[119448]22 of the alpha-I/80-kD domain. (Abstract) Blood 74:
181A, 1989.

[119449]21160.

[119450]Guetarni, D.; Roux, A.-F.; Alloisio, N.; Morle, F.;
Ducluzeau,

[119451]M. T.; Forget, B. G.; Colonna, P.; Delaunay, J.; Godet, J.:

Evidence

[119452] that expression of Sp-alpha-(I/65) hereditary elliptocytosis is compounded

[119453] by a genetic factor that is linked to the homologous alpha-spectrin

[119454] allele. Hum. Genet. 85: 627-630, 1990.

[119455] 21161.

[119456] Hassoun, H.; Coetzer, T. L.; Vassiliadis, J. N.; Sahr, K. E.;

[119457] Maalouf, G. J.; Saad, S. T. O.; Catanzariti, L.; Palek, J.: A novel

[119458] mobile element inserted in the alpha-spectrin gene: spectrin Dayton--a

[119459] truncated alpha-spectrin associated with hereditary elliptocytosis. J.

[119460] Clin. Invest. 94: 643-648, 1994.

[119461] 21162.

[119462] Huebner, K.; Palumbo, A. P.; Isobe, M.; Kozak, C. A.; Monaco,

[119463] S.; Rovera, G.; Croce, C. M.; Curtis, P. J.: The alpha-spectrin gene

[119464] is on chromosome 1 in mouse and man. Proc. Nat. Acad.

Sci. 82: 3790–3793,

[119465]1985.

[119466]21163.

[119467]Iarocci, T. A.; Wagner, G. M.; Mohandas, N.; Lane, P. A.;
Mentzer,

[119468]W. C.: Hereditary poikilocytic anemia associated with the
co-inheritance

[119469]of two alpha spectrin abnormalities. Blood 71:
1390–1396, 1988.

[119470]21164.

[119471]Knowles, W. J.; Bologna, M. L.; Chasis, J. A.; Marchesi, S.
L.;

[119472]Marchesi, V. T.: Common structural polymorphisms in hu-
man erythrocyte

[119473]spectrin. J. Clin. Invest. 73: 973–979, 1984.

[119474]21165.

[119475]Kotula, L.; Laury-Kleintop, L. D.; Showe, L.; Sahr, K.; Lin-
nenbach,

[119476]A. J.; Forget, B.; Curtis, P. J.: The exon–intron organization
of

[119477]the human erythrocyte alpha–spectrin gene. Genomics 9:

131-140,

[119478]1991.

[119479]21166.

[119480]Lambert, S.; Zail, S.: A new variant of the alpha-subunit of

[119481]spectrin in hereditary elliptocytosis. Blood 69: 473-478,

1987.

[119482]21167.

[119483]Lane, P. A.; Shew, R. L.; Iarocci, T. A.; Mohandas, N.; Hays,

[119484]T.; Mentzer, W. C.: Unique alpha-spectrin mutant in a kin-

dred with

[119485]common hereditary elliptocytosis. J. Clin. Invest. 79:

989-996,

[119486]1987.

[119487]21168.

[119488]Lawler, J.; Coetzer, T. L.; Mankad, V. N.; Moore, R. B.; Pr-

chal,

[119489]J. T.; Palek, J.: Spectrin-alpha(I/61): a new structural vari-

ant

[119490]of alpha-spectrin in a double-heterozygous form of

hereditary pyropoikilocytosis. Blood 72:

[119491]1412-1415, 1988.

[119492]21169.

[119493]Lawler, J.; Coetzer, T. L.; Palek, J.; Jacob, H. S.; Luban, N.

[119494]: Sp alpha(I/65): a new variant of the alpha subunit of spectrin in

[119495]hereditary elliptocytosis. Blood 66: 706–709, 1985.

[119496]21170.

[119497]Lawler, J.; Liu, S.-C.; Palek, J.; Prchal, J.: A molecular defect

[119498]in spectrin with a subset of patients with hereditary elliptocytosis:

[119499]alterations in the alpha-subunit domain involved in spectrin self-association. J.

[119500]Clin. Invest. 73: 1688–1695, 1984.

[119501]21171.

[119502]Lecomte, M.-C.; Dhermy, D.; Garbarz, M.; Feo, C.; Gautero, H.;

[119503]Bournier, O.; Picat, C.; Chaveroche, I.; Ester, A.; Galand, C.; Boivin,

[119504]P.: Pathologic and nonpathologic variants of the spectrin molecule

[119505]in two black families with hereditary elliptocytosis. Hum. Genet. 71:

[119506]351–357, 1985.

[119507]21172.

[119508]Lecomte, M.–C.; Dhermy, D.; Solis, C.; Ester, A.; Feo, C.;
Gautero,

[119509]H.; Bournier, O.; Boivin, P.: A new abnormal variant of
spectrin

[119510]in black patients with hereditary elliptocytosis. Blood 65:
1208–1217,

[119511]1985.

[119512]21173.

[119513]Lecomte, M.–C.; Garbarz, M.; Grandchamp, B.; Feo, C.;
Gautero,

[119514]H.; Devaux, I.; Bournier, O.; Galand, C.; d'Auriol, L.; Galib-
ert,

[119515]F.; Sahr, K. E.; Forget, B. G.; Boivin, P.; Dhermy, D.: Sp-
alpha(I/78):

[119516]a mutation of the alpha-I spectrin domain in a white kin-
dred with

[119517]HE and HPP phenotypes. Blood 74: 1126–1133, 1989.

[119518]21174.

[119519]Lecomte, M. C.; Dhermy, D.; Garbarz, M.; Feo, C.; Gautero,

H.;

[119520]Bournier, O.; Picat, C.; Chaveroche, I.; Galand, C.; Boivin, P.:

[119521]Hereditary pyropoikilocytosis and elliptocytosis in a Caucasian family:

[119522]transmission of the same molecular defect in spectrin through three

[119523]generations with different clinical expression. Hum. Genet. 77:

[119524]329-334, 1987.

[119525]21175.

[119526]Lecomte, M. C.; Dhermy, D.; Gautero, H.; Bournier, O.; Galand,

[119527]C.; Boivin, P.: L'elliptocytose hereditaire en Afrique de l'Ouest:

[119528]frequence et repartition des variants de la spectrine. C. R. Acad.

[119529]Sci. 306: 43-46, 1988.

[119530]21176.

[119531]Lecomte, M. C.; Feo, C.; Gautero, H.; Bournier, O.; Galand, C.;

[119532]Garbarz, M.; Boivin, P.; Dhermy, D.: Severe recessive poik-

ilocytic

[119533] anaemia with a new spectrin alpha chain variant. Brit. J.
Haemat. 74:

[119534] 497-507, 1990.

[119535] 21177.

[119536] Linnenbach, A. J.; Speicher, D. W.; Marchesi, V. T.; Forget,
B.

[119537] G.: Cloning of a portion of the chromosomal gene for hu-
man erythrocyte

[119538] alpha-spectrin by using a synthetic gene fragment. Proc.
Nat. Acad.

[119539] Sci. 83: 2397-2401, 1986.

[119540] 21178.

[119541] Bigler, R. D.; Bushkin, Y.; Chiorazzi, N.: S152 (CD27): a
modulating

[119542] disulfide-linked T cell activation antigen. J. Immun. 141:
21-28,

[119543] 1988.

[119544] 21179.

[119545] Hendriks, J.; Gravestien, L. A.; Tesselaar, K.; van Lier, R. A.

[119546] W.; Schumacher, T. N. M.; Borst, J.: CD27 is required for

generation

[119547]and long-term maintenance of T cell immunity. Nature Immun. 1: 433–440,

[119548]2000.

[119549]21180.

[119550]Prasad, K. V. S.; Ao, Z.; Yoon, Y.; Wu, M. X.; Rizk, M.; Jacquot,

[119551]S.; Schlossman, S. F.: CD27, a member of the tumor necrosis factor

[119552]receptor family, induces apoptosis and binds to Siva, a proapoptotic

[119553]protein. Proc. Nat. Acad. Sci. 94: 6346–6351, 1997.

[119554]21181.

[119555]Moody, D. B.; Ulrichs, T.; Muhlecker, W.; Young, D. C.; Gurcha,

[119556]S. S.; Grant, E.; Rosat, J.-P.; Brenner, M. B.; Costello, C. E.; Besra,

[119557]G. S.; Porcelli, S. A.: CD1c-mediated T-cell recognition of isoprenoid

[119558]glycolipids in Mycobacterium tuberculosis infection. Nature 404:

[119559]884–888, 2000.

[119560]21182.

[119561]Park, S.-H.; Bendelac, A.: CD1-restricted T-cell responses
and

[119562]microbial infection. Nature 406: 788-792, 2000.

[119563]21183.

[119564]Albertson, D. G.; Fishpool, R.; Sherrington, P.; Nacheva, E.;
Milstein,

[119565]C.: Sensitive and high resolution in situ hybridization to
human

[119566]chromosomes using biotin labelled probes: assignment of
the human

[119567]thymocyte CD1 antigen genes to chromosome 1. EMBO J.
7: 2801-2805,

[119568]1988.

[119569]21184.

[119570]Balk, S. P.; Bleicher, P. A.; Terhorst, C.: Isolation and char-
acterization

[119571]of a cDNA and gene coding for a fourth CD1 molecule.
Proc. Nat. Acad.

[119572]Sci. 86: 252-256, 1989.

[119573]21185.

[119574]Calabi, F.; Milstein, C.: A novel family of human major histocompatibility

[119575]complex-related genes not mapping to chromosome 6.
Nature 323: 540-543,

[119576]1986.

[119577]21186.

[119578]Nishimura, K.; Taketani, S.; Inokuchi, H.: Cloning of a human

[119579]cDNA for protoporphyrinogen oxidase by complementation in vivo of

[119580]a hemG mutant of Escherichia coli. J. Biol. Chem. 270:
8076-8080,

[119581]1995.

[119582]21187.

[119583]Roberts, A. G.; Whatley, S. D.; Daniels, J.; Holmans, P.;
Fenton,

[119584]J.; Owen, M. J.; Thompson, P.; Long, C.; Elder, G. H.: Partial
characterization

[119585]and assignment of the gene for protoporphyrinogen oxidase and variegate

[119586]porphyria to human chromosome 1q23. Hum. Molec.
Genet. 4: 2387-2390,

[119587]1995.

[119588]21188.

[119589]Taketani, S.; Inazawa, J.; Abe, T.; Furukawa, T.; Kohno, H.;
Tokunaga,

[119590]R.; Nishimura, K.; Inokuchi, H.: The human protopor-
phyrinogen oxidase

[119591]gene (PPOX): organization and location to chromosome 1.
Genomics 29:

[119592]698–703, 1995.

[119593]21189.

[119594]Warnich, L.; Kotze, M. J.; Groenewald, I. M.; Groenewald, J.
Z.;

[119595]van Brakel, M. G.; van Heerden, C. J.; de Villiers, J. N. P.;
van

[119596]de Ven, W. J. M.; Schoenmakers, E. F. P. M.; Taketani, S.;
Retief,

[119597]A. E.: Identification of three mutations and associated
haplotypes

[119598]in the protoporphyrinogen oxidase gene in South African
families with

[119599]variegate porphyria. Hum. Molec. Genet. 5: 981–984,
1996.

[119600]21190.

[119601]Albrecht, B.; Weber, K.; Pongs, O.: Characterization of a voltage-activated

[119602]K-channel gene cluster on human chromosome 12p13. Receptors Channels 3:

[119603]213-220, 1995.

[119604]21191.

[119605]Klocke, R.; Roberds, S. L.; Tamkun, M. M.; Gronemeier, M.; Augustin,

[119606]A.; Albrecht, B.; Pongs, O.; Jockusch, H.: Chromosomal mapping in

[119607]the mouse of eight K(+)-channel genes representing the four Shaker-like

[119608]subfamilies Shaker, Shab, Shaw, and Shal. Genomics 18: 568-574,

[119609]1993.

[119610]21192.

[119611]Curran, M. E.; Landes, G. M.; Keating, M. T.: Molecular cloning,

[119612]characterization, and genomic localization of a human potassium channel

[119613]gene. Genomics 12: 729–737, 1992.

[119614]21193.

[119615]Al-Chalabi, A.; Andersen, P. M.; Nilsson, P.; Chioza, B.;
Andersson,

[119616]J. L.; Russ, C.; Shaw, C. E.; Powell, J. F.; Leigh, P. N.: Dele-
tions

[119617]of the heavy neurofilament subunit tail in amyotrophic
lateral sclerosis. Hum.

[119618]Molec. Genet. 8: 157–164, 1999.

[119619]21194.

[119620]Collard, J.-F.; Cote, F.; Julien, J.-P.: Defective axonal
transport

[119621]in a transgenic mouse model of amyotrophic lateral scler-
osis. Nature 375:

[119622]61–64, 1995.

[119623]21195.

[119624]Figlewicz, D. A.; Krizus, A.; Martinoli, M. G.; Meiningner, V.;

[119625]Dib, M.; Rouleau, G. A.; Julien, J.-P.: Variants of the heavy
neurofilament

[119626]subunit are associated with the development of amy-
otrophic lateral

- [119627]sclerosis. Hum. Molec. Genet. 3: 1757–1761, 1994.
- [119628]21196.
- [119629]Lees, J. F.; Shneidman, P. S.; Skuntz, S. F.; Carden, M. J.;
Lazzarini,
[119630]R. A.: The structure and organization of the human heavy
neurofilament
[119631]subunit (NF-H) and the gene encoding it. EMBO J. 7:
1947–1955, 1988.
- [119632]21197.
- [119633]Mattei, M.-G.; Dautigny, A.; Pham-Dinh, D.; Passage, E.;
Mattei,
[119634]J.-F.; Jolles, P.: The gene encoding the large human neu-
rofilament
[119635]subunit (NF-H) maps to the q121–q131 region on human
chromosome 22. Hum.
[119636]Genet. 80: 293–295, 1988.
- [119637]21198.
- [119638]Rooke, K.; Figlewicz, D. A.; Han, F.; Rouleau, G. A.: Analy-
sis
[119639]of the KSP repeat of the neurofilament heavy subunit in
familial amyotrophic

[119640]lateral sclerosis. Neurology 46: 789–790, 1996.

[119641]21199.

[119642]Rouleau, G. A.; Merel, P.; Lutchman, M.; Sanson, M.; Zucman, J.;

[119643]Marineau, C.; Hoang-Xuan, K.; Demczuk, S.; Desmaze, C.; Plougastel,

[119644]B.; Pulst, S. M.; Lenoir, G.; Bijlsma, E.; Fashold, R.; Duman-ski,

[119645]J.; de Jong, P.; Parry, D.; Eldrige, R.; Aurias, A.; Delattre, O.;

[119646]Thomas, G.: Alteration in a new gene encoding a putative membrane-organizing

[119647]protein causes neuro-fibromatosis type 2. Nature 363: 515–521, 1993.

[119648]21200.

[119649]Tomkins, J.; Usher, P.; Slade, J. Y.; Ince, P. G.; Curtis, A.;

[119650]Bushby, K.; Shaw, P. J.: Novel insertion in the KSP region of the

[119651]neurofilament heavy gene in amyotrophic lateral sclerosis. Neuroreport 9:

[119652]3967–3970, 1998.

[119653]21201.

[119654]Vechio, J. D.; Bruijn, L. I.; Xu, Z.; Brown, R. H., Jr.; Cleveland,

[119655]D. W.: Sequence variants in human neurofilament proteins: absence

[119656]of linkage to familial amyotrophic lateral sclerosis. Ann. Neurol. 40:

[119657]603–610, 1996.

[119658]21202.

[119659]Watson, C. J.; Gaunt, L.; Evans, G.; Patel, K.; Harris, R.; Strachan,

[119660]T.: A disease-associated germline deletion maps the type 2 neurofibromatosis

[119661](NF2) gene between the Ewing sarcoma region and the leukaemia inhibitory

[119662]factor locus. Hum. Molec. Genet. 2: 701–704, 1993.

[119663]21203.

[119664]Umehara, F.; Kiwaki, T.; Yoshikawa, H.; Nishimura, T.; Nakagawa,

[119665]M.; Matsumoto, W.; Hashimoto, K.; Izumo, S.; Arimura, Y.; Arimura,

- [119666]K.; Kuriyama, M.; Osame, M.: Deletion in chromosome 17p11.2 including the peripheral myelin protein-22 (PMP-22) gene in hereditary neuropathy with liability to pressure palsies. J. Neurol. Sci. 133: 173-176, 1995.
- [119669]21204.
- [119670]21204.
- [119671]Barker, P. E.; Besmer, P.; Ruddle, F. H.: Human c-kit oncogene on human chromosome 4. (Abstract) Am. J. Hum. Genet. 37: A143, 1985.
- [119672]21205.
- [119673]21205.
- [119674]Beghini, A.; Larizza, L.; Cairoli, R.; Morra, E.: c-kit activating mutations and mast cell proliferation in human leukemia. (Letter) Blood 92: 701-702, 1998.
- [119675]701-702, 1998.
- [119676]21206.
- [119677]21206.
- [119678]Beghini, A.; Tibiletti, M.; Roversi, G.; Chiaravalli, A.; Serio, G.; Capella, C.; Larizza, L.: Germline mutation in the jux-

tamembrane

[119680]domain of the KIT gene in a family with gastrointestinal
stromal tumors

[119681]and urticaria pigmentosa. Cancer 92: 657–662, 2001.

[119682]21207.

[119683]Blume-Jensen, P.; Jiang, G.; Hyman, R.; Lee, K.-F.;
O’Gorman, S.;

[119684]Hunter, T.: Kit/stem cell factor receptor–induced activa-
tion of phosphatidylinositol

[119685]3–prime–kinase is essential for male fertility. Nature
Genet. 24:

[119686]157–162, 2000.

[119687]21208.

[119688]Bolognia, J. L.; Pawelek, J. M.: Biology of hypopigmenta-
tion. J.

[119689]Am. Acad. Derm. 19: 217–255, 1988.

[119690]21209.

[119691]Brannan, C. I.; Lyman, S. D.; Williams, D. E.; Eisenman, J.;;
Anderson,

[119692]D. M.; Cosman, D.; Bedell, M. A.; Jenkins, N. A.; Copeland,
N. G.

[119693]: Steel–Dickie mutation encodes a c–kit ligand lacking
transmembrane

[119694]and cytoplasmic domains. Proc. Nat. Acad. Sci. 88:
4671–4674, 1991.

[119695]21210.

[119696]Chabot, B.; Stephenson, D. A.; Chapman, V. M.; Besmer,
P.; Bernstein,

[119697]A.: The proto–oncogene c–kit encoding a transmembrane
tyrosine kinase

[119698]receptor maps to the mouse W locus. Nature 335: 88–89,
1988.

[119699]21211.

[119700]d'Auriol, L.; Mattei, M.–G.; Andre, C.; Galibert, F.: Local–
ization

[119701]of the human c–kit protooncogene on the q11–q12 region
of chromosome

[119702]4. Hum. Genet. 78: 374–376, 1988.

[119703]21212.

[119704]De Miguel, M. P.; Cheng, L.; Holland, E. C.; Federspiel, M.
J.;

[119705]Donovan, P. J.: Dissection of the c–Kit signaling pathway
in mouse

[119706]primordial germ cells by retroviral-mediated gene transfer. Proc.

[119707]Nat. Acad. Sci. 99: 10458–10463, 2002.

[119708]21213.

[119709]Dubreuil, P.; Forrester, L.; Rottapel, R.; Reedijk, M.; Fujita,

[119710]J.; Bernstein, A.: The c-fms gene complements the mitogenic defect

[119711]in mast cells derived from mutant W mice but not mi (microphthalmia)

[119712]mice. Proc. Nat. Acad. Sci. 88: 2341–2345, 1991.

[119713]21214.

[119714]el-Omar, M.; Davies, J.; Gupta, S.; Ross, H.; Thompson, R.: Leiomyosarcoma

[119715]in leiomyomatosis of the small intestine. Postgrad. Med. J. 70:

[119716]661–664, 1994.

[119717]21215.

[119718]Fleischman, R. A.: Human piebald trait resulting from a dominant

[119719]negative mutant allele of the c-kit membrane receptor gene. J. Clin.

[119720]Invest. 89: 1713–1717, 1992.

[119721]21216.

[119722]Fleischman, R. A.; Saltman, D. L.; Stastny, V.; Zneimer, S.:

[119723]Deletion of the c-kit protooncogene in the human developmental defect

[119724]piebald trait. Proc. Nat. Acad. Sci. 88: 10885–10889, 1991.

[119725]21217.

[119726]Fritsche–Polanz, R.; Jordan, J.–H.; Feix, A.; Sperr, W. R.; Sunder–Plassmann,

[119727]G.; Valent, P.; Fodinger, M.: Mutation analysis of C–KIT in patients

[119728]with myelodysplastic syndromes without mastocytosis and cases of systemic

[119729]mastocytosis. Brit. J. Haemat. 113: 357–364, 2001.

[119730]21218.

[119731]Geissler, E. N.; Ryan, M. A.; Housman, D. E.: The dominant–white

[119732]spotting (W) locus of the mouse encodes the c-kit protooncogene. Cell 55:

[119733]185–192, 1988.

[119734]21219.

[119735]Giebel, L. B.; Spritz, R. A.: Mutation of the KIT (mast/stem
[119736]cell growth factor receptor) protooncogene in human
piebaldism. Proc.

[119737]Nat. Acad. Sci. 88: 8696–8699, 1991.

[119738]21220.

[119739]Hirota, S.; Isozaki, K.; Moriyama, Y.; Hashimoto, K.;
Nishida,

[119740]T.; Ishiguro, S.; Kawano, K.; Hanada, M.; Kurata, A.;
Takeda, M.;

[119741]Tunio, G. M.; Matsuzawa, Y.; Kanakura, Y.; Shinomura, Y.;
Kitamura,

[119742]Y.: Gain-of-function mutations of c-kit in human gas-
trointestinal

[119743]stromal tumors. Science 279: 577–580, 1998.

[119744]21221.

[119745]Huizinga, J. D.; Thuneberg, L.; Kluppel, M.; Malysz, J.;
Mikkelsen,

[119746]H. B.; Bernstein, A.: W/kit gene required for interstitial
cells

[119747]of Cajal and for intestinal pacemaker activity. Nature 373:

347–349,

[119748]1995.

[119749]21222.

[119750]Furitsu, T.; Tsujimura, T.; Tono, T.; Ikeda, H.; Kitayama,
H.;

[119751]Koshimizu, U.; Sugahara, H.; Butterfield, J. H.; Ashman, L.
K.; Kanayama,

[119752]Y.; Matsuzawa, Y.; Kitamura, Y.; Kanakura, Y.: Identifica-
tion of

[119753]mutations in the coding sequence of the proto-oncogene
c-kit in a

[119754]human mast cell leukemia cell line causing ligand-
independent activation

[119755]of c-kit product. J. Clin. Invest. 92: 1736–1744, 1993.

[119756]21223.

[119757]Ingram, D. A.; Yang, F.-C.; Travers, J. B.; Wenning, M. J.;
Hiatt,

[119758]K.; New, S.; Hood, A.; Shannon, K.; Williams, D. A.; Clapp,
D. W.

[119759]: Genetic and biochemical evidence that haploinsufficiency
of the

[119760]Nf1 tumor suppressor gene modulates melanocyte and

mast cell fates

[119761]in vivo. J. Exp. Med. 191: 181–187, 2000.

[119762]21224.

[119763]Isozaki, K.; Terris, B.; Belghiti, J.; Schiffmann, S.; Hirota,

[119764]S.; Vanderwinden, J.–M.: Germline–activating mutation in
the kinase

[119765]domain of KIT gene in familial gastrointestinal stromal tu-
mors. Am.

[119766]J. Path. 157: 1581–1585, 2000.

[119767]21225.

[119768]Joensuu, H.; Roberts, P. J.; Sarlomo–Rikala, M.; Andersson,
L.

[119769]C.; Tervahartiala, P.; Tuveson, D.; Silberman, S. L.;
Capdeville,

[119770]R.; Dimitrijevic, S.; Druker, B.; Demetri, G. D.: Effect of the
tyrosine

[119771]kinase inhibitor STI571 in a patient with a metastatic gas-
trointestinal

[119772]stromal tumor. New Eng. J. Med. 344: 1052–1056, 2001.

[119773]21226.

[119774]Johansson Moller, M.; Chaudhary, R.; Hellmen, E.; Hoy–

heim, B.;

[119775]Chowdhary, B.; Andersson, L.: Pigs with the dominant white coat color

[119776]phenotype carry a duplication of the KIT gene encoding the mast/stem

[119777]cell growth factor receptor. Mammalian Genome 7: 822–830, 1996.

[119778]21227.

[119779]Lasota, J.; Jasinski, M.; Sarlomo-Rikala, M.; Miettinen, M.:

[119780]Mutations in exon 11 of c-kit occur preferentially in malignant versus

[119781]benign gastrointestinal stromal tumors and do not occur in leiomyomas

[119782]or leiomyosarcomas. Am. J. Path. 154: 53–60, 1999.

[119783]21228.

[119784]Longley, B. J.; Tyrrell, L.; Lu, S.-Z.; Ma, Y.-S.; Langley, K.;

[119785]Ding, T.; Duffy, T.; Jacobs, P.; Tang, L. H.; Modlin, I.: Somatic

[119786]c-KIT activating mutation in urticaria pigmentosa and aggressive mastocytosis:

[119787]establishment of clonality in a human mast cell neoplasm.

Nature

[119788]Genet. 12: 312–314, 1996.

[119789]21229.

[119790]Longley, B. J., Jr.; Metcalfe, D. D.; Tharp, M.; Wang, X.;
Tyrrell,

[119791]L.; Lu, S.-Z.; Heitjan, D.; Ma, Y.: Activating and dominant
inactivating

[119792]c-KIT catalytic domain mutations in distinct clinical forms
of human

[119793]mastocytosis. Proc. Nat. Acad. Sci. 96: 1609–1614, 1999.

[119794]21230.

[119795]Marklund, S.; Kijas, J.; Rodriguez-Martinez, H.;
Ronnstrand, L.;

[119796]Funa, K.; Moller, M.; Lange, D.; Edfors-Lilja, I.; Andersson,
L.:

[119797]Molecular basis for the dominant white phenotype in the
domestic pig. Genome

[119798]Res. 8: 826–833, 1998.

[119799]21231.

[119800]Ingalls, A. M.; Dickie, M. M.; Snell, G. D.: Obese, a new
mutation

[119801]in the house mouse. J. Hered. 41: 317–318, 1950.

[119802]21232.

[119803]Ioffe, E.; Moon, B.; Connolly, E.; Friedman, J. M.: Abnormal

[119804]regulation of the leptin gene in the pathogenesis of obesity. Proc.

[119805]Nat. Acad. Sci. 95: 11852–11857, 1998.

[119806]21233.

[119807]Isse, N.; Ogawa, Y.; Tamura, N.; Masuzaki, H.; Mori, K.;
Okazaki,

[119808]T.; Satoh, N.; Shigemoto, M.; Yoshimasa, Y.; Nishi, S.;
Hosoda, K.;

[119809]Inazawa, J.; Nakao, K.: Structural organization and chromosomal assignment

[119810]of the human obese gene. J. Biol. Chem. 270:
27728–27733, 1995.

[119811]21234.

[119812]Jaquet, D.; Leger, J.; Levy-Marchal, C.; Oury, J. F.; Czerni-
chow,

[119813]P.: Ontogeny of leptin in human fetuses and newborns:
effect of intrauterine

[119814]growth retardation on serum leptin concentrations. J. Clin.

Endocr.

[119815]Metab. 83: 1243–1246, 1998.

[119816]21235.

[119817]Karvonen, M. K.; Pesonen, U.; Heinonen, P.; Laakso, M.;
Rissanen,

[119818]A.; Naukkarinen, H.; Valve, R.; Uusitupa, M. I. J.; Koulu, M.:
Identification

[119819]of new sequence variants in the leptin gene. J. Clin. En-
docr. Metab. 83:

[119820]3239–3242, 1998.

[119821]21236.

[119822]Kawai, K.; Sugimoto, K.; Nakashima, K.; Miura, H.; Ni-
nomiya, Y.

[119823]: Leptin as a modulator of sweet taste sensitivities in mice.
Proc.

[119824]Nat. Acad. Sci. 97: 11044–11049, 2000.

[119825]21237.

[119826]Kennedy, A.; Gettys, T. W.; Watson, P.; Wallace, P.; Gan-
away,

[119827]E.; Pan, Q.; Garvey, W. T.: The metabolic significance of
leptin

- [119828]in humans: Gender-based differences in relationship to adiposity,
- [119829]insulin sensitivity, and energy expenditure. J. Clin. Endocr. Metab. 82: 1293–1300, 1997.
- [119831]21238.
- [119832]Koistinen, H. A.; Koivisto, V. A.; Andersson, S.; Karonen, S.-L.;
- [119833]Kontula, K.; Oksanen, L.; Teramo, K. A.: Leptin concentration in
- [119834]cord blood correlates with intrauterine growth. J. Clin. Endocr. Metab. 82: 3328–3330, 1997.
- [119835]21239.
- [119837]Li, W.-D.; Reed, D. R.; Lee, J. H.; Xu, W.; Kilker, R. L.; So-
- [119838]dam, B. R.; Price, R. A.: Sequence variants in the 5-prime flanking region
- [119839]of the leptin gene are associated with obesity in women. Ann. Hum. Genet. 63: 227–234, 1999.

[119841]21240.

[119842]Lonnqvist, F.; Arner, P.; Nordfors, L.; Schalling, M.: Over-expression

[119843]of the obese (ob) gene in adipose tissue of human obese subjects. Nature

[119844]Med. 1: 950-953, 1995.

[119845]21241.

[119846]Lord, G. M.; Matarese, G.; Howard, J. K.; Baker, R. J.; Bloom,

[119847]S. R.; Lechler, R. I.: Leptin modulates the T-cell immune response

[119848]and reverses starvation-induced immunosuppression. Nature 394: 897-901,

[119849]1998.

[119850]21242.

[119851]Lucantoni, R.; Ponti, E.; Berselli, M. E.; Savia, G.; Minocci,

[119852]A.; Calo, G.; de Medici, C.; Liuzzi, A.; Di Blasio, A. M.: The A19G

[119853]polymorphism in the 5-prime untranslated region of the human obese

[119854]gene does not affect leptin levels in severely obese pa-

tients. J.

[119855]Clin. Endocr. Metab. 85: 3589–3591, 2000.

[119856]21243.

[119857]Maffei, M.; Fei, H.; Lee, G.-H.; Dani, C.; Leroy, P.; Zhang, Y.;

[119858]Proenca, R.; Negrel, R.; Ailhaud, G.; Friedman, J. M.: Increased

[119859]expression in adipocytes of ob RNA in mice with lesions of the hypothalamus

[119860]and with mutations at the db locus. Proc. Nat. Acad. Sci. 92: 6957–6960,

[119861]1995.

[119862]21244.

[119863]Mammes, O.; Betoulle, D.; Aubert, R.; Giraud, V.; Tuzet, S.; Petiet,

[119864]A.; Colas-Linhart, N.; Fumeron, F.: Novel polymorphisms in the 5-prime

[119865]region of the LEP gene: association with leptin levels and response

[119866]to low-calorie diet in human obesity. Diabetes 47: 487–489, 1998.

[119867]21245.

- [119868]Mammes, O.; Betoulle, D.; Aubert, R.; Herbeth, B.; Siest, G.;
- [119869]Fumeron, F.: Association of the G-2548A polymorphism in the 5-prime
- [119870]region of the LEP gene with overweight. Ann. Hum. Genet. 64: 391-394,
- [119871]2000.
- [119872]21246.
- [119873]Mancuso, P.; Gottschalk, A.; Phare, S. M.; Peters-Golden, M.;
- [119874]Lukacs, N. W.; Huffnagle, G. B.: Leptin-deficient mice exhibit impaired
- [119875]host defense in Gram-negative pneumonia. J. Immun. 168: 4018-4024,
- [119876]2002.
- [119877]21247.
- [119878]Mantzoros, C. S.; Flier, J. S.; Rogol, A. D.: A longitudinal
- [119879]assessment of hormonal and physical alterations during normal puberty
- [119880]in boys. v. rising leptin levels may signal the onset of puberty. J.

- [119881]Clin. Endocr. Metab. 82: 1066–1070, 1997.
- [119882]21248.
- [119883]Masuzaki, H.; Ogawa, Y.; Isse, N.; Satoh, N.; Okazaki, T.;
Shigemoto,
- [119884]M.; Mori, K.; Tamura, N.; Hosoda, K.; Yoshimasa, Y.;
Jingami, H.;
- [119885]Kawada, T.; Nakao, K.: Human obese gene expression:
adipocyte-specific
- [119886]expression and regional differences in the adipose tissue.
Diabetes 44:
- [119887]855–858, 1995.
- [119888]21249.
- [119889]Masuzaki, H.; Ogawa, Y.; Sagawa, N.; Hosoda, K.; Mat-
sumoto, T.;
- [119890]Mise, H.; Nishimura, H.; Yoshimasa, Y.; Tanaka, I.; Mori,
T.; Nakao,
- [119891]K.: Nonadipose tissue production of leptin: leptin as a
novel placenta-derived
- [119892]hormone in humans. Nature Med. 3: 1029–1033, 1997.
- [119893]21250.
- [119894]Matkovic, V.; Ilich, J. Z.; Badenhop, N. E.; Skugor, M.;

Clairmont,

[119895]A.; Klisovic, D.; Landoll, J. D.: Gain in body fat is inversely related

[119896]to the nocturnal rise in serum leptin level in young females. J.

[119897]Clin. Endocr. Metab. 82: 1368–1372, 1997.

[119898]21251.

[119899]Matkovic, V.; Ilich, J. Z.; Skugor, M.; Badenhop, N. E.; Goel,

[119900]P.; Clairmont, A.; Klisovic, D.; Nahhas, R. W.; Landoll, J. D.:
Leptin

[119901]is inversely related to age at menarche in human females.
J. Clin.

[119902]Endocr. Metab. 82: 3239–3245, 1997.

[119903]21252.

[119904]Matsuda, J.; Yokota, I.; Iida, M.; Murakami, T.; Naito, E.;
Ito,

[119905]M.; Shima, K.; Kuroda, Y.: Serum leptin concentration in
cord blood:

[119906]relationship to birth weight and gender. J. Clin. Endocr.
Metab. 82:

[119907]1642–1644, 1997.

[119908]21253.

[119909]Miller, S. G.; De Vos, P.; Guerre-Millo, M.; Wong, K.; Hermann,

[119910]T.; Staels, B.; Briggs, M. R.; Auwerx, J.: The adipocyte specific

[119911]transcription factor C/EBP- α modulates human ob gene expression. Proc.

[119912]Nat. Acad. Sci. 93: 5507-5511, 1996.

[119913]21254.

[119914]Minokoshi, Y.; Kim, Y.-B.; Peroni, O. D.; Fryer, L. G. D.; Muller,

[119915]C.; Carling, D.; Kahn, B. B.: Leptin stimulates fatty-acid oxidation

[119916]by activating AMP-activated protein kinase. Nature 415: 339-343,

[119917]2002.

[119918]21255.

[119919]Moffett, S.; Martinson, J.; Shriver, M. D.; Deka, R.; McGarvey,

[119920]S. T.; Barrantes, R.; Ferrell, R. E.: Genetic diversity and evolution

[119921]of the human leptin locus tetranucleotide repeat. Hum.
Genet. 110:

[119922]412–417, 2002.

[119923]21256.

[119924]Montague, C. T.; Farooqi, I. S.; Whitehead, J. P.; Soos, M.
A.;

[119925]Rau, H.; Wareham, N. J.; Sewter, C. P.; Digby, J. E.; Mo-
hammed, S.

[119926]N.; Hurst, J. A.; Cheetham, C. H.; Earley, A. R.; Barnett, A.
H.;

[119927]Prins, J. B.; O'Rahilly, S.: Congenital leptin deficiency is as-
sociated

[119928]with severe early-onset obesity in humans. Nature 387:
903–908,

[119929]1997.

[119930]21257.

[119931]Muzzin, P.; Eisensmith, R. C.; Copeland, K. C.; Woo, S. L.
C.

[119932]: Correction of obesity and diabetes in genetically obese
mice by

[119933]leptin gene therapy. Proc. Nat. Acad. Sci. 93:
14804–14808, 1996.

[119934]21258.

[119935]Kiefer, M. C.; Tucker, J. E.; Joh, R.; Landsberg, K. E.; Saltman,

[119936]D.; Barr, P. J.: Identification of a second human subtilisin-like

[119937]protease gene in the fes/fps region of chromosome 15.
DNA Cell Biol. 10:

[119938]757-769, 1991.

[119939]21259.

[119940]Amiel, J.; Audollent, S.; Joly, D.; Dureau, P.; Salomon, R.;
Tellier,

[119941]A.-L.; Auge, J.; Bouissou, F.; Antignac, C.; Gubler, M.-C.;
Eccles,

[119942]M. R.; Munnich, A.; Vekemans, M.; Lyonnet, S.; Attie-
Bitach, T.:

[119943]PAX2 mutations in renal-coloboma syndrome: mutational
hotspot and

[119944]germline mosaicism. Europ. J. Hum. Genet. 8: 820-826,
2000.

[119945]21260.

[119946]Dehbi, M.; Ghahremani, M.; Lechner, M.; Dressler, G.; Pel-
letier,

[119947]J.: The paired-box transcription factor, PAX2, positively modulates

[119948]expression of the Wilms' tumor suppressor gene. *Oncogene* 13: 447-453,

[119949]1996.

[119950]21261.

[119951]Devriendt, K.; Matthijs, G.; Van Damme, B.; Van Caesbroeck, D.;

[119952]Eccles, M.; Vanrenterghem, Y.; Fryns, J.-P.; Leys, A.: Mis-sense mutation

[119953]and hexanucleotide duplication in the PAX2 gene in two unrelated families

[119954]with renal-coloboma syndrome (MIM 120330). *Hum. Genet.* 103: 149-153,

[119955]1998.

[119956]21262.

[119957]Dressler, G. R.; Deutsch, U.; Chowdhury, K.; Nornes, H. O.; Gruss,

[119958]P.: Pax2, a new murine paired-box-containing gene and its expression

[119959]in the developing excretory system. *Development* 109: 787-795, 1990.

[119960]21263.

[119961]Ford, B.; Rupps, R.; Lirenman, D.; Van Allen, M. I.; Far-
quharson,

[119962]D.; Lyons, C.; Friedman, J. M.: Renal-coloboma syndrome:
prenatal

[119963]detection and clinical spectrum in a large family. Am. J.
Med. Genet. 99:

[119964]137-141, 2001.

[119965]21264.

[119966]Keller, S. A.; Jones, J. M.; Boyle, A.; Barrow, L. L.; Killen,

[119967]P. D.; Green, D. G.; Kapousta, N. V.; Hitchcock, P. F.;
Swank, R.

[119968]T.; Meisler, M. H.: Kidney and retinal defects (Krd), a
transgene-induced

[119969]mutation with a deletion of mouse chromosome 19 that
includes the

[119970]Pax2 locus. Genomics 23: 309-320, 1994.

[119971]21265.

[119972]Narahara, K.; Baker, E.; Ito, S.; Yokoyama, Y.; Yu, S.; He-
witt,

[119973]D.; Sutherland, G. R.; Eccles, M. R.; Richards, R. I.: Locali-

sation

[119974] of a 10q breakpoint within the PAX2 gene in a patient with
a de novo

[119975] t(10;13) translocation and optic nerve coloboma–renal
disease. J.

[119976] Med. Genet. 34: 213–216, 1997.

[119977] 21266.

[119978] Pilz, A. J.; Povey, S.; Gruss, P.; Abbott, C. M.: Mapping of
the

[119979] human homologs of the murine paired–box–containing
genes. Mammalian

[119980] Genome 4: 78–82, 1993.

[119981] 21267.

[119982] Favor, J.; Sandulache, R.; Neuhauser–Klaus, A.; Pretsch, W.;
Chatterjee,

[119983] B.; Senft, E.; Wurst, W.; Blanquet, V.; Grimes, P.; Sporle, R.;
Schughart,

[119984] K.: The mouse Pax2(1Neu) mutation is identical to a hu–
man PAX2 mutation

[119985] in a family with renal–coloboma syndrome and results in
developmental

[119986] defects of the brain, ear, eye, and kidney. Proc. Nat. Acad.

Sci. 93:

[119987]13870-13875, 1996.

[119988]21268.

[119989]Seawright, A.; Fletcher, J. M.; Fantes, J. A.; Morrison, H.;
Porteous,

[119990]D. J.; Li, S. S.-L.; Hastie, N. D.; Van Heyningen, V.: Analysis
of

[119991]WAGR deletions and related translocations with gene-
specific DNA probes,

[119992]using FACS-selected cell hybrids. Somat. Cell Molec.
Genet. 14:

[119993]21-30, 1988.

[119994]21269.

[119995]Huerre, C.; Despoisse, S.; Gilgenkrantz, S.; Lenoir, G. M.;
Junien,

[119996]C.: c-Ha-ras1 is not deleted in aniridia-Wilms' tumour as-
sociation. Nature 305:

[119997]638-641, 1983.

[119998]21270.

[119999]Liou, G. I.; Ma, D.-P.; Yang, Y.-W.; Geng, L.; Zhu, C.;
Baehr,

- [120000]W.: Human interstitial retinoid-binding protein: gene structure and primary sequence. J. Biol. Chem. 264: 8200–8206, 1989.
- [120001]21271.
- [120002]Nakamura, Y.; Lathrop, M.; Bragg, T.; Leppert, M.; O'Connell, P.; Jones, C.; Lalouel, J. M.; White, R.: An extended genetic linkage map of markers for human chromosome 10. Genomics 3: 389–392, 1988.
- [120003]21272.
- [120004]Heitman, J.; Agre, P.: A new face of the rhesus antigen. Nature Genet. 26: 258–259, 2000.
- [120005]21273.
- [120006]Marini, A.–M.; Matassi, G.; Raynal, V.; Andre, B.; Cartron, J.–P.; Cherif–Zahar, B.: The human Rhesus-associated RhAG protein and a kidney homologue promote ammonium transport in yeast. Nature Genet. 26:

[120013]341–344, 2000.

[120014]21274.

[120015]Sap, J.; Munoz, A.; Damm, K.; Goldberg, Y.; Ghysdael, J.;
Leutz,

[120016]A.; Beug, H.; Vennstrom, B.: The c-erb-A protein is a
high-affinity

[120017]receptor for thyroid hormone. Nature 324: 635–640,
1986.

[120018]21275.

[120019]Carrere, N.; Leblanc, R. M.; Begueret, J.; Blouin, P.; Chey-
rou,

[120020]A.: A new mutation of exon 5 of the P53 gene in breast
cancer. Hum.

[120021]Molec. Genet. 2: 1075, 1993.

[120022]21276.

[120023]Walker, R. L.; Anziano, P.; Meltzer, P. S.: A PAC containing
the

[120024]human mitochondrial DNA polymerase gamma gene
(POLG) maps to chromosome

[120025]15q25. Genomics 40: 376–378, 1997.

[120026]21277.

- [120027]Zullo, S. J.; Butler, L.; Zahorchak, R. J.; Macville, M.; Wilkes,
[120028]C.; Merril, C. R.: Localization by fluorescence in situ hybridization
(FISH) of human mitochondrial polymerase gamma (POLG)
[120029]to human chromosome
[120030]band 15q24-q26, and of mouse mitochondrial polymerase gamma (Polg)
[120031]to mouse chromosome band 7E, with confirmation by direct sequence
[120032]analysis of bacterial artificial chromosomes (BACs). Cytogenet. Cell
[120033]Genet. 78: 281-284, 1997.
[120034]21278.
[120035]Bayliss, W.; Starling, E. H.: The mechanism of pancreatic secretion. J.
[120036]Physiol. (London) 28: 325-353, 1902.
[120037]21279.
[120038]Kopin, A. S.; Wheeler, M. B.; Leiter, A. B.: Secretin: structure
[120039]of the precursor and tissue distribution of the mRNA. Proc. Nat.

[120040]Acad. Sci. 87: 2299–2303, 1990.

[120041]21280.

[120042]Marinelli, R. A.; Pham, L.; Agre, P.; LaRusso, N. F.: Secretin

[120043]promotes osmotic water transport in rat cholangiocytes by
increasing

[120044]aquaporin–1 water channels in plasma membrane. J. Biol.
Chem. 272:

[120045]12984–12988, 1997.

[120046]21281.

[120047]Peel, A. L.; Rao, R. V.; Cottrell, B. A.; Hayden, M. R.;
Ellerby,

[120048]L. M.; Bredesen, D. E.: Double–stranded RNA–dependent
protein kinase,

[120049]PKR, binds preferentially to Huntington's disease (HD)
transcripts

[120050]and is activated in HD tissue. Hum. Molec. Genet. 10:
1531–1538,

[120051]2001.

[120052]21282.

[120053]Squire, J.; Meurs, E. F.; Chong, K. L.; McMillan, N. A. J.;
Hovanessian,

- [120054]A. G.; Williams, B. R. G.: Localization of the human interferon-induced,
[120055]ds-RNA activated p68 kinase gene (PRKR) to chromosome
2p21-p22. Genomics 16:
[120056]768-770, 1993.
[120057]21283.
- [120058]Taylor, D. R.; Shi, S. T.; Romano, P. R.; Barber, G. N.; Lal,
M.
[120059]M. C.: Inhibition of the interferon-inducible protein kinase
PKR
[120060]by HCV E2 protein. Science 285: 107-110, 1999.
[120061]21284.
- [120062]Brott, B. K.; Alessandrini, A.; Largaespada, D. A.;
Copeland, N.
[120063]G.; Jenkins, N. A.; Crews, C. M.; Erikson, R. L.: MEK2 is a
kinase
[120064]related to MEK1 and is differentially expressed in murine
tissues. Cell
[120065]Growth Differ. 4: 921-929, 1993.
[120066]21285.
- [120067]Crews, C. M.; Alessandrini, A.; Erikson, R. L.: The primary

structure

[120068]of MEK, a protein kinase that phosphorylates the ERK gene product. Science 258:

[120069]478–480, 1992.

[120070]21286.

[120071]Favata, M. F.; Horiuchi, K. Y.; Manos, E. J.; Daulerio, A. J.;

[120072]Stradley, D. A.; Feeser, W. S.; Van Dyk, D. E.; Pitts, W. J.; Earl,

[120073]R. A.; Hobbs, F.; Copeland, R. A.; Magolda, R. L.; Scherle, P. A.;

[120074]Trzaskos, J. M.: Identification of a novel inhibitor of mitogen-activated

[120075]protein kinase kinase. J. Biol. Chem. 273: 18623–18632, 1998.

[120076]21287.

[120077]Meloche, S.; Gopalbhai, K.; Beatty, B. G.; Scherer, S. W.; Pellerin,

[120078]J.: Chromosome mapping of the human genes encoding the MAP kinase

[120079]kinase MEK1 (MAP2K1) to 15q21 and MEK2 (MAP2K2) to 7q32. Cytogenet.

[120080]Cell Genet. 88: 249–252, 2000.

[120081]21288.

[120082]Orth, K.; Palmer, L. E.; Bao, Z. Q.; Stewart, S.; Rudolph, A. E.;

[120083]Bliska, J. B.; Dixon, J. E.: Inhibition of the mitogen-activated

[120084]protein kinase kinase superfamily by a Yersinia effector. Science 285:

[120085]1920–1923, 1999.

[120086]21289.

[120087]Perry, R. L. S.; Parker, M. H.; Rudnicki, M. A.: Activated MEK1

[120088]binds the nuclear MyoD transcriptional complex to repress transactivation. Molec.

[120089]Cell 8: 291–301, 2001.

[120090]21290.

[120091]Pleschka, S.; Wolff, T.; Ehrhardt, C.; Hobom, G.; Planz, O.; Rapp,

[120092]U. R.; Ludwig, S.: Influenza virus propagation is impaired by inhibition

[120093]of the Raf/MEK/ERK signalling cascade. Nature Cell Biol. 3: 301–305,

[120094]2001.

[120095]21291.

[120096]Rampoldi, L.; Zimbello, R.; Bortoluzzi, S.; Tiso, N.; Valle, G.;

[120097]Lanfranchi, G.; Danieli, G. A.: Chromosomal localization of four

[120098]MAPK signaling cascade genes: MEK1, MEK3, MEK4 and MEKK5. Cytogenet.

[120099]Cell Genet. 78: 301–303, 1997.

[120100]21292.

[120101]Ryan, K. M.; Ernst, M. K.; Rice, N. R.; Vousden, K. H.: Role of

[120102]NF- κ B in p53-mediated programmed cell death. Nature 404: 892–897,

[120103]2000.

[120104]21293.

[120105]Sebolt-Leopold, J. S.; Dudley, D. T.; Herrera, R.; Van Becelaere,

[120106]K.; Wiland, A.; Gowan, R. C.; Tecle, H.; Barrett, S. D.; Bridges,

[120107]A.; Przybranowski, S.; Leopold, W. R.; Saltiel, A. R.: Block-

ade of

[120108] the MAP kinase pathway suppresses growth of colon tumors in vivo. *Nature*

[120109] *Med.* 5: 810–816, 1999.

[120110] 21294.

[120111] Seger, R.; Krebs, E. G.: The MAPK signaling cascade. *FASEB J.* 9:

[120112] 726–735, 1995.

[120113] 21295.

[120114] Seger, R.; Seger, D.; Lozeman, F. J.; Ahn, N. G.; Graves, L. M.;

[120115] Campbell, J. S.; Ericsson, L.; Harrylock, M.; Jensen, A. M.; Krebs,

[120116] E. G.: Human T-cell mitogen-activated protein kinase kinases are

[120117] related to yeast signal transduction kinases. *J. Biol. Chem.* 267:

[120118] 25628–25631, 1992.

[120119] 21296.

[120120] Zheng, C. F.; Guan, K. L.: Cloning and characterization of two

[120121]distinct human extracellular signal-regulated kinase activator kinases,

[120122]MEK1 and MEK2. J. Biol. Chem. 268: 11435-11439, 1993.

[120123]21297.

[120124]White, P. S.; Maris, J. M.; Beltinger, C.; Sulman, E.; Marshall,

[120125]H. N.; Fujimori, M.; Kaufman, B. A.; Biegel, J. A.; Allen, C.; Hilliard,

[120126]C.; Valentine, M. B.; Look, A. T.; Enomoto, H.; Sakiyama, S.; Brodeur,

[120127]G. M.: A region of consistent deletion in neuroblastoma maps within

[120128]human chromosome 1p36.2-36.3. Proc. Nat. Acad. Sci. 92: 5520-5524,

[120129]1995.

[120130]21298.

[120131]Deng, Z.; Morse, J. H.; Slager, S. L.; Cuervo, N.; Moore, K. J.;

[120132]Venetos, G.; Kalachikov, S.; Cayanis, E.; Fischer, S. G.; Barst, R.

[120133]J.; Hodge, S. E.; Knowles, J. A.: Familial primary pulmonary hypertension

[120134](gene PPH1) is caused by mutations in the bone morpho-
genetic protein

[120135]receptor-II gene. Am. J. Hum. Genet. 67: 737-744, 2000.

[120136]21299.

[120137]International PPH Consortium; Lane, K. B.; Machado, R. D.;
Pauciulo,

[120138]M. W.; Thomson, J. R.; Phillips, J. A., III; Loyd, J. E.; Nichols,

[120139]W. C.; Trembath, R. C.: Heterozygous germline mutations
in BMPR2,

[120140]encoding a TGF-beta receptor, cause familial primary pul-
monary hypertension. Nature

[120141]Genet. 26: 81-84, 2000.

[120142]21300.

[120143]Amin, R. S.; Wert, S. E.; Baughman, R. P.; Tomashevski, J.
F.,

[120144]Jr.; Nogee, L. M.; Brody, A. S.; Hull, W. M.; Whitsett, J. A.:
Surfactant

[120145]protein deficiency in familial interstitial lung disease. J.
Pediat. 139:

[120146]85-92, 2001.

[120147]21301.

- [120148]Glasser, S. W.; Burhans, M. S.; Korfhagen, T. R.; Na, C.-L.;
Sly,
[120149]P. D.; Ross, G. F.; Ikegami, M.; Whitsett, J. A.: Altered sta-
bility
[120150]of pulmonary surfactant in SP-C-deficient mice. Proc. Nat.
Acad.
[120151]Sci. 98: 6366-6371, 2001.
[120152]21302.
[120153]Hussain, S. P.; Raja, K.; Amstad, P. A.; Sawyer, M.; Trudel,
L.
[120154]J.; Wogan, G. N.; Hofseth, L. J.; Shields, P. G.; Billiar, T. R.;
[120155]Trautwein, C.; Hohler, T.; Galle, P. R.; Phillips, D. H.;
Markin,
[120156]R.; Marrogi, A. J.; Harris, C. C.: Increased p53 mutation
load in
[120157]nontumorous human liver of Wilson disease and
hemochromatosis: oxyradical
[120158]overload diseases. Proc. Nat. Acad. Sci. 97: 12770-12775,
2000.
[120159]21303.
[120160]Iavarone, A.; Matthay, K. K.; Steinkirchner, T. M.; Israel, M.

[120161]A.: Germ-line and somatic p53 gene mutations in multifocal osteogenic

[120162]sarcoma. Proc. Nat. Acad. Sci. 89: 4207–4209, 1992.

[120163]21304.

[120164]Iggo, R.; Gatter, K.; Bartek, J.; Lane, D.; Harris, A. L.: Increased

[120165]expression of mutant forms of p53 oncogene in primary lung cancer. Lancet 335:

[120166]675–679, 1990.

[120167]21305.

[120168]Isobe, M.; Emanuel, B. S.; Givol, D.; Oren, M.; Croce, C. M.:

[120169]Localization of gene for human p53 tumour antigen to band 17p13. Nature 320:

[120170]84–85, 1986.

[120171]21306.

[120172]Jeffrey, P. D.; Gorina, S.; Pavletich, N. P.: Crystal structure

[120173]of the tetramerization domain of the p53 tumor suppressor at 1.7 angstroms. Science 267:

[120174]1498–1502, 1995.

[120175]21307.

[120176]Jenkins, J. R.; Rudge, K.; Chumakov, P.; Currie, G. A.: The cellular

[120177]oncogene p53 can be activated by mutagenesis. Nature
317: 816–818,

[120178]1985.

[120179]21308.

[120180]Jimenez, G. S.; Nister, M.; Stommel, J. M.; Beeche, M.; Bar-
carse,

[120181]E. A.; Zhang, X.-Q.; O'Gorman, S.; Wahl, G. M.: A transac-
tivation-deficient

[120182]mouse model provides insights into Trp53 regulation and
function. Nature

[120183]Genet. 26: 37–43, 2000.

[120184]21309.

[120185]Jonkers, J.; Meuwissen, R.; van der Gulden, H.; Peterse, H.;
van

[120186]der Valk, M.; Berns, A.: Synergistic tumor suppressor ac-
tivity of

[120187]BRCA2 and p53 in a conditional mouse model for breast
cancer. Nature

[120188]Genet. 29: 418–425, 2001.

[120189]21310.

[120190]Kaelbling, M.; Burk, R. D.; Atkin, N. B.; Johnson, A. B.;
Klinger,

[120191]H. P.: Loss of heterozygosity on chromosome 17p and
mutant p53 in

[120192]HPV-negative cervical carcinomas. Lancet 340: 140–142,
1992.

[120193]21311.

[120194]Keller, D. M.; Zeng, X.; Wang, Y.; Zhang, Q. H.; Kapoor, M.;
Shu,

[120195]H.; Goodman, R.; Lozano, G.; Zhao, Y.; Lu, H.: A DNA
damage-induced

[120196]p53 serine 392 kinase complex contains CK2, hSpt16, and
SSRP1. Molec.

[120197]Cell 283–292, 2001.

[120198]21312.

[120199]Kemp, C. J.; Wheldon, T.; Balmain, A.: p53-deficient mice
are

[120200]extremely susceptible to radiation-induced tumorigenesis.
Nature

[120201]Genet. 8: 66–69, 1994.

[120202]21313.

[120203]Krawczak, M.; Smith-Sorensen, B.; Schmidtke, J.; Kakkar, V. V.;

[120204]Cooper, D. N.; Hovig, E.: Somatic spectrum of cancer-associated single

[120205]basepair substitutions in the TP53 gene is determined mainly by endogenous

[120206]mechanisms of mutation and by selection. Hum. Mutat. 5: 48-57, 1995.

[120207]21314.

[120208]Lamb, P.; Crawford, L.: Characterization of the human p53 gene. Molec.

[120209]Cell. Biol. 6: 1379-1385, 1986.

[120210]21315.

[120211]Latronico, A. C.; Pinto, E. M.; Domenice, S.; Fragoso, M. C. B.

[120212]V.; Martin, R. M.; Zerbini, M. C.; Lucon, A. M.; Mendonca, B. B.:

[120213]An inherited mutation outside the highly conserved DNA-binding domain

[120214]of the p53 tumor suppressor protein in children and adults with sporadic

[120215]adrenocortical tumors. J. Clin. Endocr. Metab. 86:
4970–4973, 2001.

[120216]21316.

[120217]Lavigueur, A.; Maltby, V.; Mock, D.; Rossant, J.; Pawson,
T.;

[120218]Bernstein, A.: High incidence of lung, bone, and lymphoid
tumors

[120219]in transgenic mice overexpressing mutant alleles of the
p53 oncogene. Molec.

[120220]Cell. Biol. 9: 3982–3991, 1989.

[120221]21317.

[120222]Law, J. C.; Strong, L. C.; Chidambaram, A.; Ferrell, R. E.: A

[120223]germ line mutation in exon 5 of the p53 gene in an ex-
tended cancer

[120224]family. Cancer Res. 51: 6385–6387, 1991.

[120225]21318.

[120226]Lee, J. M.; Bernstein, A.: p53 mutations increase resistance

[120227]to ionizing radiation. Proc. Nat. Acad. Sci. 90: 5742–5746,
1993.

[120228]21319.

[120229] Levine, A. J.: p53, the cellular gatekeeper for growth and division. Cell 88:

[120230] 323–331, 1997.

[120231] 21320.

[120232] Levine, A. J.; Momand, J.; Finlay, C. A.: The p53 tumour suppressor

[120233] gene. Nature 351: 453–456, 1991.

[120234] 21321.

[120235] Li, M.; Chen, D.; Shiloh, A.; Luo, J.; Nikolaev, A. Y.; Qin, J.;

[120236] Gu, W.: Deubiquitination of p53 by HAUSP is an important pathway

[120237] for p53 stabilization. Nature 416: 648–653, 2002.

[120238]

[120239] 100. Lu, H.; Levine, A. J.: Human TAFII31 protein is a transcriptional

[120240] coactivator of the p53 protein. Proc. Nat. Acad. Sci. 92: 5154–5158,

[120241] 1995.

[120242]

[120243] 101. Luo, J.; Nikolaev, A. Y.; Imai, S.; Chen, D.; Su, F.;

Shiloh,

[120244]A.; Guarente, L.; Gu, W.: Negative control of p53 by
Sir2- α promotes

[120245]cell survival under stress. Cell 107: 137–148, 2001.

[120246]

[120247]102. Luo, J.; Su, F.; Chen, D.; Shiloh, A.; Gu, W.: Deacety-
lation

[120248]of p53 modulates its effect on cell growth and apoptosis.
Nature 408:

[120249]377–381, 2000.

[120250]

[120251]103. Malkin, D.; Jolly, K. W.; Barbier, N.; Look, A. T.;
Friend, S.

[120252]H.; Gebhardt, M. C.; Andersen, T. I.; Borresen, A.-L.; Li, F.
P.;

[120253]Garber, J.; Strong, L. C.: Germline mutations of the p53
tumor-suppressor

[120254]gene in children and young adults with second malignant
neoplasms. New

[120255]Eng. J. Med. 326: 1309–1315, 1992.

[120256]

[120257]104. Malkin, D.; Li, F. P.; Strong, L. C.; Fraumeni, J. F., Jr.;
Nelson,

- [120258]C. E.; Kim, D. H.; Kassel, J.; Gryka, M. A.; Bischoff, F. Z.;
Tainsky,
[120259]M. A.; Friend, S. H.: Germ line p53 mutations in a familial
syndrome
[120260]of breast cancer, sarcomas, and other neoplasms. Science
250: 1233–1238,
[120261]1990.
[120262]
[120263]105. Marin, M. C.; Jost, C. A.; Brooks, L. A.; Irwin, M. S.;
O'Nions,
[120264]J.; Tidy, J. A.; James, N.; McGregor, J. M.; Harwood, C. A.;
Yulug,
[120265]J. G.; Vousden, K. H.; Allday, M. J.; Gusterson, B.; Ikawa, S.;
Hinds,
[120266]P. W.; Crook, T.; Kaelin, W. G., Jr. A common polymor-
phism acts
[120267]as an intragenic modifier of mutant p53 behaviour. Nature
Genet. 25:
[120268]47–54, 2000.
[120269]
[120270]106. Marino, S.; Vooijs, M.; van der Gulden, H.; Jonker, J.;
Berns,

[120271]A.: Induction of medulloblastomas in p53-null mutant mice by somatic

[120272]inactivation of Rb in the external granular layer cells of the cerebellum. Genes

[120273]Dev. 14: 994–1004, 2000.

[120274]

[120275]107. Masuda, H.; Miller, C.; Koeffler, H. P.; Battifora, H.; Cline,

[120276]M. J.: Rearrangement of the p53 gene in human osteogenic sarcomas. Proc.

[120277]Nat. Acad. Sci. 84: 7716–7719, 1987.

[120278]

[120279]108. Mathupala, S. P.; Heese, C.; Pedersen, P. L.: Glucose catabolism

[120280]in cancer cells: the type II hexokinase promoter contains functionally

[120281]active response elements for the tumor suppressor p53. J. Biol. Chem. 272:

[120282]22776–22780, 1997.

[120283]

[120284]109. Mazoyer, S.; Lalle, P.; Moyret-Lalle, C.; Marcais, C.;

Schraub,

[120285]S.; Frappaz, D.; Sobol, H.; Ozturk, M.: Two germ-line mutations affecting

[120286]the same nucleotide at codon 257 of p53 gene, a rare site for mutations. Oncogene 9:

[120287]1237-1239, 1994.

[120288]

[120289]110. McBride, O. W.; Merry, D.; Givol, D.: The gene for human p53

[120290]cellular tumor antigen is located on chromosome 17 short arm (17p13). Proc.

[120291]Nat. Acad. Sci. 83: 130-134, 1986.

[120292]

[120293]111. McBride, O. W.; Merry, D. E.; Oren, M.; Givol, D.: Human p53

[120294]cellular tumor antigen is on chromosome 17p13. (Abstract) Cytogenet.

[120295]Cell Genet. 40: 694-695, 1985.

[120296]

[120297]112. Metzger, A. K.; Sheffield, V. C.; Duyk, G.; Daneshvar, L.; Edwards,

[120298]M. S. B.; Cogen, P. H.: Identification of a germ-line mutation in

[120299]the p53 gene in a patient with an intracranial ependy-
moma. Proc.

[120300]Nat. Acad. Sci. 88: 7825-7829, 1991.

[120301]

[120302]113. Miller, C.; Mohandas, T.; Wolf, D.; Prokocimer, M.;
Rotter, V.;

[120303]Koeffler, H. P.: Human p53 gene localized to short arm of
chromosome

[120304]17. Nature 319: 783-784, 1986.

[120305]

[120306]114. Moll, U. M.; Riou, G.; Levine, A. J.: Two distinct
mechanisms

[120307]alter p53 in breast cancer: mutation and nuclear exclu-
sion. Proc.

[120308]Nat. Acad. Sci. 89: 7262-7266, 1992.

[120309]

[120310]115. Monpezat, J. P.; Delattre, O.; Bernard, A.; Grunwald,
D.; Remvikos,

[120311]Y.; Muleris, M.; Salmon, R. J.; Frelat, G.; Dutrillaux, B.;

Thomas,

[120312]G.: Loss of alleles on chromosome 18 and on the short arm of chromosome

[120313]17 in polyploid colorectal carcinomas. *Int. J. Cancer* 41: 404–408,

[120314]1988.

[120315]

[120316]116. Mulligan, L. M.; Matlashewski, G. J.; Scrable, H. J.; Cavenee,

[120317]W. K.: Mechanisms of p53 loss in human sarcomas. *Proc. Nat. Acad.*

[120318]*Sci.* 87: 5863–5867, 1990.

[120319]

[120320]117. Nakamura, T.; Pichel, J. G.; Williams–Simons, L.; Westphal, H.

[120321]: An apoptotic defect in lens differentiation caused by human p53

[120322]is rescued by a mutant allele. *Proc. Nat. Acad. Sci.* 92: 6142–6146,

[120323]1995.

[120324]

[120325]118. Neugut, A. I.; Santos, J.: The association between
cancers of

[120326]the small and large bowel. Cancer Epidemiol. Biomarkers
Prev. 2:

[120327]551–553, 1993.

[120328]

[120329]119. Nicol, C. J.; Harrison, M. L.; Laposa, R. R.;
Gimelshtein, I.

[120330]L.; Wells, P. G.: A teratologic suppressor role for p53 in
benzo[a]pyrene-treated

[120331]transgenic p53-deficient mice. Nature Genet. 10:
181–187, 1995.

[120332]

[120333]120. Nigro, J. M.; Baker, S. J.; Preisinger, A. C.; Jessup, J.
M.;

[120334]Hostetter, R.; Cleary, K.; Bigner, S. H.; Davidson, N.; Baylin,
S.;

[120335]Devilee, P.; Glover, T.; Collins, F. S.; Weston, A.; Modali, R.;

[120336]Harris,
C. C.; Vogelstein, B.: Mutations in the p53 gene occur in
diverse

[120337]human tumour types. Nature 342: 705–708, 1989.

[120338]

[120339]121. Oda, T.; Tsuda, H.; Scarpa, A.; Sakamoto, M.; Hirohashi, S.:

[120340]p53 gene mutation spectrum in hepatocellular carcinoma. Cancer Res. 52:

[120341]6358-6364, 1992.

[120342]

[120343]122. Oliner, J. D.; Kinzler, K. W.; Meltzer, P. S.; George, D. L.;

[120344]Vogelstein, B.: Amplification of a gene encoding a p53-associated

[120345]protein in human sarcomas. Nature 358: 80-83, 1992.

[120346]

[120347]123. Olivier, M.; Eeles, R.; Hollstein, M.; Khan, M. A.; Harris, C.

[120348]C.; Hainaut, P.: The IARC TP53 database: new online mutation analysis

[120349]and recommendations to users. Hum. Mutat. 19: 607-614, 2002.

[120350]

[120351]124. Ollmann, M.; Young, L. M.; Di Como, C. J.; Karim, F.;

Belvin,

[120352]M.; Robertson, S.; Whittaker, K.; Demsky, M.; Fisher, W.
W.; Buchman,

[120353]A.; Duyk, G.; Friedman, L.; Prives, C.; Kopczynski, C.:
Drosophila

[120354]p53 is a structural and functional homolog of the tumor
suppressor

[120355]p53. Cell 101: 91–101, 2000.

[120356]

[120357]125. Olschwang, S.; Laurent–Puig, P.; Vassal, A.; Salmon,
R.–J.; Thomas,

[120358]G.: Characterization of a frequent polymorphism in the
coding sequence

[120359]of the Tp53 gene in colonic cancer patients and a control
population. Hum.

[120360]Genet. 86: 369–370, 1991.

[120361]

[120362]126. Parsa, C. F.; Hoyt, C. S.; Lesser, R. L.; Weinstein, J. M.;
Strother,

[120363]C. M.; Muci–Mendoza, R.; Ramella, M.; Manor, R. S.;
Fletcher, W. A.;

[120364]Repka, M. X.; Garrity, J. A.; Ebner, R. N.; Monteiro, M. L. R.;

McFadzean,

[120365]**R. M.; Rubtsova, I. V.; Hoyt, W. F.: Spontaneous regression
of optic**

[120366]**gliomas: thirteen cases documented by serial neuroimag-
ing. Arch.**

[120367]**Ophthalmol. 119: 516–529, 2001.**

[120368]

[120369]**127. Patel, P.; Stephenson, J.; Scheuer, P. J.; Francis, G. E.:
p53**

[120370]**codon 249–ser mutations in hepatocellular carcinoma pa-
tients with**

[120371]**low aflatoxin exposure. (Letter) Lancet 339: 881, 1992.**

[120372]

[120373]**128. Pavletich, N. P.; Chambers, K. A.; Pabo, C. O.: The
DNA-binding**

[120374]**domain of p53 contains the four conserved regions and
the major mutation**

[120375]**hot spots. Genes Dev. 7: 2556–2564, 1993.**

[120376]

[120377]**129. Pearson, M.; Carbone, R.; Sebastiani, C.; Cioce, M.;
Fagioli,**

[120378]M.; Saito, S.; Higashimoto, Y.; Appella, E.; Minucci, S.;
Pandolfi,

[120379]P. P.; Pelicci, P. G.: PML regulates p53 acetylation and pre-
mature

[120380]senescence induced by oncogenic Ras. Nature 406:
207–210, 2000.

[120381]

[120382]130. Polyak, K.; Xia, Y.; Zweler, J. L.; Kinzler, K. W.; Vogel-
stein,

[120383]B.: A model for p53-induced apoptosis. Nature 389:
300–305, 1997.

[120384]

[120385]131. Potzsch, C.; Schaefer, H.–E.; Lubbert, M.: Familial and
metachronous

[120386]malignant lymphoma: absence of constitutional p53 mu-
tations. Am.

[120387]J. Hemat. 62: 144–149, 1999.

[120388]

[120389]132. Raj, K.; Ogston, P.; Beard, P.: Virus-mediated killing
of cells

[120390]that lack p53 activity. Nature 412: 914–917, 2001. Note:
Addendum:

[120391]Nature 416: 202 only, 2002.

[120392]

[120393]133. Raman, V.; Martensen, S. A.; Reisman, D.; Evron, E.;
Odenwald,

[120394]W. F.; Jaffee, E.; Marks, J.; Sukumar, S.: Compromised
HOXA5 function

[120395]can limit p53 expression in human breast tumours. Nature
405: 974–978,

[120396]2000.

[120397]

[120398]134. Reilly, K. M.; Loisel, D. A.; Bronson, R. T.; McLaugh-
lin, M.

[120399]E.; Jacks, T.: Nf1;Trp53 mutant mice develop glioblastoma
with evidence

[120400]of strain-specific effects. Nature Genet. 26: 109–113,
2000.

[120401]

[120402]135. Reisman, D.; Balint, E.; Loging, W. T.; Rotter, V.; Al-
mon, E.

[120403]: A novel transcript encoded within the 10-kb first intron
of the

[120404]human p53 tumor suppressor gene (D17S2179E) is induced during differentiation

[120405]of myeloid leukemia cells. Genomics 38: 364–370, 1996.

[120406]

[120407]136. Reisman, D.; Greenberg, M.; Rotter, V.: Human p53 oncogene contains

[120408]one promoter upstream of exon 1 and a second, stronger promoter within

[120409]intron 1. Proc. Nat. Acad. Sci. 85: 5146–5150, 1988.

[120410]

[120411]137. Ribeiro, R. C.; Sandrini, F.; Figueiredo, B.; Zambetti, G. P.;

[120412]Michalkiewicz, E.; Lafferty, A. R.; DeLacerda, L.; Rabin, M.; Cadwell,

[120413]C.; Sampaio, G.; Cat, I.; Stratakis, C. A.; Sandrini, R.: An inherited

[120414]p53 mutation that contributes in a tissue-specific manner to pediatric

[120415]adrenal cortical carcinoma. Proc. Nat. Acad. Sci. 98: 9330–9335,

[120416]2001.

[120417]

[120418]138. Rideout, W. M., III; Coetzee, G. A.; Olumi, A. F.; Jones, P.

[120419]A.: 5-Methylcytosine as an endogenous mutagen in the human LDL receptor

[120420]and p53 genes. Science 249: 1288–1290, 1990.

[120421]

[120422]139. Rodrigues, N. R.; Rowan, A.; Smith, M. E. F.; Kerr, I. B.; Bodmer,

[120423]W. F.; Gannon, J. V.; Lane, D. P.: p53 mutations in colorectal cancer. Proc.

[120424]Nat. Acad. Sci. 87: 7555–7559, 1990.

[120425]

[120426]140. Romano, J. W.; Ehrhart, J. C.; Duthu, A.; Kim, C. M.; Appella,

[120427]E.; May, P.: A mutation in the p53 gene of a human osteosarcoma cell

[120428]line. (Abstract) Am. J. Hum. Genet. 45 (suppl.): A32, 1989.

[120429]

[120430]141. Roth, J. A.; Nguyen, D.; Lawrence, D. D.; Kemp, B. L.; Carrasco,

[120431]C. H.; Ferson, D. Z.; Hong, W. K.; Komaki, R.; Lee, J. J.;
Nesbitt,

[120432]J. C.; Pisters, K. M. W.; Putnam, J. B.; Schea, R.; Shin, D. M.;
Walsh,

[120433]G. L.; Dolormente, M. M.; Han, C.-I.; Martin, F. D.; Yen, N.;
Xu,

[120434]K.; Stephens, L. C.; McDonnell, T. J.; Mukhopadhyay, T.;
Cai, D.:

[120435]Retrovirus-mediated wild-type p53 gene transfer to tu-
mors of patients

[120436]with lung cancer. Nature Med. 2: 985-990, 1996.

[120437]

[120438]142. Rowley, J. D.: Personal Communication. Chicago, Ill.
1/3/1986.

[120439]

[120440]143. Rutherford, J.; Chu, C. E.; Duddy, P. M.; Charlton, R.
S.; Chumas,

[120441]P.; Taylor, G. R.; Lu, X.; Barnes, D. M.; Camplejohn, R. S.:
Investigations

[120442]on a clinically and functionally unusual and novel germline
p53 mutation. Brit.

[120443]J. Cancer 86: 1592-1596, 2002.

[120444]

[120445]144. Ryan, K. M.; Ernst, M. K.; Rice, N. R.; Vousden, K. H.:

Role

[120446]of NF- κ B in p53-mediated programmed cell death.

Nature 404:

[120447]892–897, 2000.

[120448]

[120449]145. Sah, V. P.; Arradi, L. D.; Mulligan, G. J.; Williams, B.

O.;

[120450]Bronson, R. T.; Jacks, T.: A subset of p53-deficient em-

bryos exhibit

[120451]exencephaly. Nature Genet. 10: 175–180, 1995.

[120452]

[120453]146. Santibanez-Koref, M. F.; Birch, J. M.; Hartley, A. L.;

Morris

[120454]Jones, P. H.; Craft, A. W.; Eden, T.; Crowther, D.; Kelsey, A.

M.;

[120455]Harris, M.: p53 germline mutations in Li-Fraumeni syn-

drome. Lancet 338:

[120456]1490–1491, 1991.

[120457]

- [120458]147. Schiffer, D.; Cavalla, P.; Di Sapio, A.; Giordana, M. T.;
Mauro,
[120459]A.: Mutations and immunohistochemistry of p53 and pro-
liferation markers
[120460]in astrocytic tumors of childhood. *Child's Nerv. Syst.* 11:
517–522,
[120461]1995.
[120462]
[120463]148. Schmitt, C. A.; Fridman, J. S.; Yang, M.; Lee, S.; Bara-
nov, E.;
[120464]Hoffman, R. M.; Lowe, S. W.: A senescence program con-
trolled by p53
[120465]and p16–INK4a contributes to the outcome of cancer
therapy. *Cell* 109:
[120466]335–346, 2002.
[120467]
[120468]149. Sedlacek, Z.; Kodet, R.; Kriz, V.; Seemanova, E.; Vod-
varka, P.;
[120469]Wilgenbus, P.; Mares, J.; Poustka, A.; Goetz, P.: Two Li-
Fraumeni
[120470]syndrome families with novel germline p53 mutations:
loss of the wild-type

[120471]p53 allele in only 50% of tumours. Brit. J. Cancer 77:
1034–1039,

[120472]1998.

[120473]

[120474]150. Shieh, S.–Y.; Ikeda, M.; Taya, Y.; Prives, C.: DNA
damage–induced

[120475]phosphorylation of p53 alleviates inhibition by MDM2. Cell
91: 325–334,

[120476]1997.

[120477]

[120478]151. Sidransky, D.; Von Eschenbach, A.; Tsai, Y. C.; Jones,
P.; Summerhayes,

[120479]I.; Marshall, F.; Paul, M.; Green, P.; Hamilton, S. R.; Frost,
P.;

[120480]Vogelstein, B.: Identification of p53 gene mutations in
bladder cancers

[120481]and urine samples. Science 252: 706–709, 1991.

[120482]

[120483]152. Smela, M. E.; Hamm, M. L.; Henderson, P. T.; Harris,
C. M.; Harris,

[120484]T. M.; Essigmann, J. M.: The aflatoxin B1 formamidopy–

rimidine adduct

[120485] plays a major role in causing the types of mutations observed in human

[120486] hepatocellular carcinoma. Proc. Nat. Acad. Sci. 99: 6655–6660, 2002.

[120487]

[120488] 153. Smith, H. S.: Personal Communication. San Francisco, Calif.

[120489] 11/16/1993.

[120490]

[120491] 154. Smith–Sorensen, B.; Gebhardt, M. C.; Kloen, P.; McIntyre, J.;

[120492] Aguilar, F.; Cerutti, P.; Borresen, A.–L.: Screening for TP53 mutations

[120493] in osteosarcomas using constant denaturant gel electrophoresis (CDGE). Hum.

[120494] Mutat. 2: 274–285, 1993.

[120495]

[120496] 155. Srivastava, S.; Zou, Z.; Pirollo, K.; Blattner, W.; Chang, E.

[120497] H.: Germ–line transmission of a mutated p53 gene in a

cancer-prone

[120498]family with Li-Fraumeni syndrome. Nature 348: 747-749, 1990.

[120499]

[120500]156. Storey, A.; Thomas, M.; Kalita, A.; Harwood, C.; Gardiol, D.;

[120501]Mantovani, F.; Breuer, J.; Leigh, I. M.; Matlashewski, G.; Banks,

[120502]L.: Role of a p53 polymorphism in the development of human papilloma-virus-associated

[120503]cancer. Nature 393: 229-234, 1998.

[120504]

[120505]157. Sun, Y.; Hegamyer, G.; Cheng, Y.-J.; Hildesheim, A.; Chen, J.-Y.;

[120506]Chen, I.-H.; Cao, Y.; Yao, K.-T.; Colburn, N. H.: An infrequent point

[120507]mutation of the p53 gene in human nasopharyngeal carcinoma. Proc.

[120508]Nat. Acad. Sci. 89: 6516-6520, 1992.

[120509]

[120510]158. Takahashi, T.; D'Amico, D.; Chiba, I.; Buchhagen, D.

L.; Minna,

[120511]J. D.: Identification of intronic point mutations as an alternative

[120512]mechanism for p53 inactivation in lung cancer. J. Clin. Invest. 86:

[120513]363–369, 1990.

[120514]

[120515]159. Takahashi, T.; Nau, M. M.; Chiba, I.; Birrer, M. J.; Rosenberg,

[120516]R. K.; Vinocour, M.; Levitt, M.; Pass, H.; Gazdar, A. F.; Minna, J.

[120517]D.: p53: a frequent target for genetic abnormalities in lung cancer. Science 246:

[120518]491–494, 1989.

[120519]

[120520]160. Toguchida, J.; Yamaguchi, T.; Dayton, S. H.; Beauchamp, R. L.;

[120521]Herrera, G. E.; Ishizaki, K.; Yamamuro, T.; Meyers, P. A.; Little,

[120522]J. B.; Sasaki, M. S.; Weichselbaum, R. R.; Yandell, D. W.: Prevalence

[120523]and spectrum of germline mutations of the p53 gene

among patients

[120524]with sarcoma. New Eng. J. Med. 326: 1301–1308, 1992.

[120525]

[120526]161. Tonisson, N.; Zernant, J.; Kurg, A.; Pavel, H.; Slavin, G.; Roomere,

[120527]H.; Meiel, A.; Hainaut, P.; Metspalu, A.: Evaluating the arrayed
rayed

[120528]primer extension resequencing assay of TP53 tumor suppressor gene. Proc.

[120529]Nat. Acad. Sci. 99: 5503–5508, 2002.

[120530]

[120531]162. Tyner, S. D.; Venkatachalam, S.; Choi, J.; Jones, S.; Ghebranious,

[120532]N.; Igelmann, H.; Lu, X.; Soron, G.; Cooper, B.; Brayton, C.; Park,

[120533]S. H.; Thompson, T.; Karsenty, G.; Bradley, A.; Donehower, L. A.:

[120534]p53 mutant mice that display early ageing-associated phenotypes. Nature 415:

[120535]45–53, 2002.

[120536]

[120537]163. Ueda, H.; Ullrich, S. J.; Gangemi, J. D.; Kappel, C. A.;
Ngo,

[120538]L.; Feitelson, M. A.; Jay, G.: Functional inactivation but not
structural

[120539]mutation of p53 causes liver cancer. Nature Genet. 9:
41–47, 1995.

[120540]

[120541]164. Unger, T.; Nau, M. M.; Segal, S.; Minna, J. D.: p53: a
transdominant

[120542]regulator of transcription whose function is ablated by
mutations

[120543]occurring in human cancer. EMBO J. 11: 1383–1390,
1992.

[120544]

[120545]165. vanTuinen, P.; Ledbetter, D. H.: Construction and
utilization

[120546]of a detailed somatic cell hybrid mapping panel for human
chromosome

[120547]17: localization of an anonymous clone to the critical re-
gion of Miller–Dieker

[120548]syndrome, deletion 17p13. (Abstract) Cytogenet. Cell
Genet. 46:

[120549]708–709, 1987.

[120550]

[120551]166. Varley, J. M.; McGowan, G.; Thorncroft, M.; Tricker, K.
J.; Teare,,

[120552]M. D.; Santibanez–Koref, M. F.; Houlston, R. S.; Martin, J.;
Birch,

[120553]J. M.; Evans, D. G. R.: An extended Li–Fraumeni kindred
with gastric

[120554]carcinoma and a codon 175 mutation in TP53. J. Clin.
Genet. 32:

[120555]942–945, 1995.

[120556]

[120557]167. Varley, J. M.; McGown, G.; Thorncroft, M.; Cochrane,
S.; Morrison,

[120558]P.; Woll, P.; Kelsey, A. M.; Mitchell, E. L. D.; Boyle, J.; Birch,

[120559]J. M.; Evans, D. G. R.: A previously undescribed mutation
within

[120560]the tetramerisation domain of TP53 in a family with Li–
Fraumeni syndrome

[120561](sic). Oncogene 12: 2437–2442, 1996.

[120562]

[120563]168. Vaziri, H.; Dessain, S. K.; Eaton, E. N.; Imai, S.-I.;
Frye,

[120564]R. A.; Pandita, T. K.; Guarente, L.; Weinberg, R. A.:
hSIR2-SIRT1

[120565]functions as an NAD-dependent p53 deacetylase. Cell
107: 149-159,

[120566]2001.

[120567]

[120568]169. Vogelstein, B.; Kinzler, K. W.: X-rays strike p53
again. Nature 370:

[120569]174-175, 1994.

[120570]

[120571]170. Vogelstein, B.; Kinzler, K. W.: p53 function and dys-
function. Cell 70:

[120572]523-526, 1992.

[120573]

[120574]171. Wheeler, J. M. D.; Warren, B. F.; Mortensen, N. J. M.;
Kim, H.

[120575]C.; Biddolph, S. C.; Elia, G.; Beck, N. E.; Williams, G. T.;
Shepherd,

[120576]N. A.; Bateman, A. C.; Bodmer, W. F.: An insight into the

genetic

[120577] pathway of adenocarcinoma of the small intestine. Gut 50:
218–223,

[120578] 2002.

[120579]

[120580] 172. Yamanishi, Y.; Boyle, D. L.; Rosengren, S.; Green, D.
R.; Zvaifler,

[120581] N. J.; Firestein, G. S.: Regional analysis of p53 mutations in
rheumatoid

[120582] arthritis synovium. Proc. Nat. Acad. Sci. 99: 10025–10030,
2002.

[120583]

[120584] 173. Yu, A.; Fan, H.-Y.; Liao, D.; Bailey, A. D.; Weiner, A.
M.:

[120585] Activation of p53 or loss of the Cockayne syndrome group
B repair

[120586] protein causes metaphase fragility of human U1, U2, and
5S genes. Molec.

[120587] Cell 5: 801–810, 2000.

[120588]

[120589] 174. Yu, J. L.; Rak, J. W.; Coomber B. L.; Hicklin, D. J.; Ker-

bel,

[120590]R. S.: Effect of p53 status on tumor response to antiangiogenic therapy. Science 295:

[120591]1526–1528, 2002.

[120592]

[120593]175. Zhang, W.; Funk, W. D.; Wright, W. E.; Shay, J. W.; Deisseroth,

[120594]A. B.: Novel DNA binding of p53 mutants and their role in transcriptional

[120595]activation. Oncogene 8: 2555–2559, 1993.

[120596]

[120597]176. Zhang, Y.; Xiong, Y.: A p53 amino-terminal nuclear export signal

[120598]inhibited by DNA damage-induced phosphorylation. Science 292: 1910–1915,

[120599]2001.

[120600]

[120601]177. Zhu, C.; Mills, K. D.; Ferguson, D. O.; Lee, C.; Manis, J.; Fleming,

[120602]J.; Gao, Y.; Morton, C. C.; Alt, F. W.: Unrepaired DNA breaks in

[120603]p53-deficient cells lead to oncogenic gene amplification
subsequent

[120604]to translocations. Cell 109: 811–821, 2002.

[120605]

[120606]178. Ziegler, A.; Jonason, A. S.; Leffell, D. J.; Simon, J. A.;
Sharma,

[120607]H. W.; Kimmelman, J.; Remington, L.; Jacks, T.; Brash, D.
E.: Sunburn

[120608]and p53 in the onset of skin cancer. Nature 372:
773–776, 1994.

[120609]21322.

[120610]Juppner, H.: Molecular cloning and characterization of a
parathyroid

[120611]hormone/parathyroid hormone-related peptide receptor:
a member of

[120612]an ancient family of G protein-coupled receptors. Curr.
Opin. Nephrol.

[120613]Hypertension 3: 371–378, 1994.

[120614]21323.

[120615]Juppner, H.; Abou-Samra, A.-B.; Freeman, M.; Kong, X. F.;
Schipani,

- [120616]E.; Richards, J.; Kolakowski, L. F., Jr.; Hock, J.; Potts, J. T.,
[120617]Jr.; Kronenberg, H. M.; Segre, G. V.: A G protein-linked re-
ceptor
[120618]for parathyroid hormone and parathyroid hormone-re-
lated peptide. Science 254:
[120619]1024–1026, 1991.
[120620]21324.
[120621]Karperien, M.; van der Harten, H. J.; van Schooten, R.;
Farih-Sips,
[120622]H.; den Hollander, N. S.; Kneppers, S. L. J.; Nijweide, P.;
Papapoulos,
[120623]S. E.; Lowik, C. W. G. M.: A frame-shift mutation in the
type I parathyroid
[120624]hormone (PTH)/PTH-related peptide receptor causing
Blomstrand lethal
[120625]osteochondrodysplasia. J. Clin. Endocr. Metab. 84:
3713–3720, 1999.
[120626]21325.
[120627]Lanske, B.; Karaplis, A. C.; Lee, K.; Luz, A.; Vortkamp, A.;
Pirro,
[120628]A.; Karperien, M.; Defize, L. H. K.; Ho, C.; Mulligan, R. C.;
Abou-Samra,

[120629]A.-B.; Juppner, H.; Segre, G. V.; Kronenberg, H. M.: PTH/
PTHrP receptor

[120630]in early development and Indian hedgehog-regulated
bone growth. Science 273:

[120631]663–666, 1996.

[120632]21326.

[120633]Loshkajian, A.; Roume, J; Stanescu, V.; Delezoide, A.;
Stampf,

[120634]F.; Maroteaux, P.: Familial Blomstrand chondrodysplasia
with advanced

[120635]skeletal maturation: further delineation. Am. J. Med.
Genet. 71:

[120636]283–288, 1997.

[120637]21327.

[120638]Mahon, M. J.; Donowitz, M.; Yun, C. C.; Segre, G. V.:
Na⁺/H⁺

[120639]exchanger regulatory factor 2 directs parathyroid hor-
mone 1 receptor

[120640]signalling. Nature 417: 858–861, 2002.

[120641]21328.

[120642]Manen, D.; Palmer, G.; Bonjour, J.-P.; Rizzoli, R.: Positive

- [120643]and negative control of the expression of parathyroid hormone (PTH)/PTH-related
- [120644]protein receptor via proximal promoter P3 in human osteoblast-like
- [120645]cells. J. Clin. Endocr. Metab. 85: 3376–3382, 2000.
- [120646]21329.
- [120647]McCuaig, K. A.; Clarke, J. C.; White, J. H.: Molecular cloning
- [120648]of the gene encoding the mouse parathyroid hormone/parathyroid hormone-related
- [120649]peptide receptor. Proc. Nat. Acad. Sci. 91: 5051–5055, 1994.
- [120650]21330.
- [120651]Minagawa, M.; Watanabe, T.; Kohno, Y.; Mochizuki, H.; Hendy, G.
- [120652]N.; Goltzman, D.; White, J. H.; Yasuda, T.: Analysis of the P3 promoter
- [120653]of the human parathyroid hormone (PTH)/PTH-related peptide receptor
- [120654]gene in pseudohypoparathyroidism type 1b. J. Clin. Endocr. Metab. 86:
- [120655]1394–1397, 2001.

[120656]21331.

[120657]Minagawa, M.; Yasuda, T.; Watanabe, T.; Minamitani, K.;
Takahashi,

[120658]Y.; Goltzman, D.; White, J. H.; Hendy, G. N.; Kohno, Y.: As-
sociation

[120659]between AAAG repeat polymorphism in the P3 promoter
of the human parathyroid

[120660]hormone (PTH)/PTH-related peptide receptor gene and
adult height,

[120661]urinary pyridinoline excretion, and promoter activity. J.
Clin. Endocr.

[120662]Metab. 87: 1791-1796, 2002.

[120663]21332.

[120664]Parfitt, A. M.; Schipani, E.; Rao, D. S.; Kupin, W.; Han,
Z.-H.;

[120665]Juppner, H.: Hypercalcemia due to constitutive activity of
the parathyroid

[120666]hormone (PTH)/PTH-related peptide receptor: comparison
with primary

[120667]hyperparathyroidism. J. Clin. Endocr. Metab. 81:
3584-3588, 1996.

[120668]21333.

[120669] Schipani, E.; Karga, H.; Karaplis, A. C.; Potts, J. T, Jr.; Kronenberg,

[120670] H. M.; Segre, G. V.; Abou-Samra, A.-B.; Juppner, H.: Identical complementary

[120671] deoxyribonucleic acids encode a human renal and bone parathyroid hormone

[120672] (PTH)/PTH-related peptide receptor. *Endocrinology* 132: 2157-2165,

[120673] 1993.

[120674] 21334.

[120675] Schipani, E.; Langman, C.; Hunzelman, J.; Le Merrer, M.; Loke,

[120676] K. Y.; Dillon, M. J.; Silve, C.; Juppner, H.: A novel parathyroid

[120677] hormone (PTH)/PTH-related peptide receptor mutation in Jansen's metaphyseal

[120678] chondrodysplasia. *J. Clin. Endocr. Metab.* 84: 3052-3057, 1999.

[120679] 21335.

[120680] Schipani, E.; Langman, C. B.; Parfitt, A. M.; Jensen, G. S.; Kikuchi,

[120681] S.; Kooh, S. W.; Cole, W. G.; Juppner, H.: Constitutively ac-

tivated

[120682]receptors for parathyroid hormone and parathyroid hormone-related

[120683]peptide in Jansen's metaphyseal chondrodysplasia. New Eng. J. Med. 335:

[120684]708-714, 1996.

[120685]21336.

[120686]Pausova, Z.; Bourdon, J.; Clayton, D.; Mattei, M.-G.; Seldin,

[120687]M. F.; Janicic, N.; Riviere, M.; Szpirer, J.; Levan, G.; Szpirer,

[120688]C.; Goltzman, D.; Hendy, G. N.: Cloning of a parathyroid hormone/parathyroid

[120689]hormone-related peptide receptor (PTHrP) cDNA from a rat osteosarcoma

[120690](UMR 106) cell line: chromosomal assignment of the gene in the human,

[120691]mouse, and rat genomes. Genomics 20: 20-26, 1994.

[120692]21337.

[120693]Schipani, E.; Weinstein, L. S.; Bergwitz, C.; Iida-Klein, A.;

[120694]Kong, X. F.; Stuhmann, M.; Kruse, K.; Whyte, M. P.; Murray, T.; Schmidtke,

[120695]J.; van Dop, C.; Brickman, A. S.; Crawford, J. D.; Potts, J. T.,

Jr.;

[120696]Kronenberg, H. M.; Abou-Samra, A. B.; Segre, G. V.; Jupp-
ner, H.:

[120697]Pseudohypoparathyroidism type Ib is not caused by muta-
tions in the

[120698]coding exons of the human parathyroid hormone
(PTH)/PTH-related peptide

[120699]receptor gene. J. Clin. Endocr. Metab. 80: 1611-1621,
1995.

[120700]21338.

[120701]Suarez, F.; Lebrun, J. J.; Lecossier, D.; Escoubet, B.;
Coureau,

[120702]C.; Silve, C.: Expression and modulation of the parathyroid
hormone

[120703](PTH)/PTH-related peptide receptor messenger ribonu-
cleic acid in skin

[120704]fibroblasts from patients with type Ib pseudohy-
poparathyroidism. J.

[120705]Clin. Endocr. Metab. 80: 965-970, 1995.

[120706]21339.

[120707]Zhang, P.; Jobert, A.-S.; Couvineau, A.; Silve, C.: A ho-
mozygous

- [120708]inactivating mutation in the parathyroid hormone/
parathyroid hormone-related
- [120709]peptide receptor causing Blomstrand chondrodysplasia. J.
Clin. Endocr.
- [120710]Metab. 83: 3365–3368, 1998.
- [120711]21340.
- [120712]Albright, F.: Case records of the Massachusetts General
Hospital
- [120713](case 27461). New Eng. J. Med. 225: 789–791, 1941.
- [120714]21341.
- [120715]Bakre, M. M.; Zhu, Y.; Yin, H.; Burton, D. W.; Terkeltaub,
R.;
- [120716]Deftos, L. J.; Varner, J. A.: Parathyroid hormone-related
peptide
- [120717]is a naturally occurring, protein kinase A-dependent an-
giogenesis
- [120718]inhibitor. Nature Med. 8: 995–1003, 2002.
- [120719]21342.
- [120720]Broadus, A. E.; Mangin, M.; Ikeda, K.; Insogna, K. L.; Weir,
E.
- [120721]C.; Burtis, W. J.; Stewart, A. F.: Humoral hypercalcemia of

cancer:

[120722]identification of a novel parathyroid hormone-like peptide. New Eng.

[120723]J. Med. 319: 556-563, 1988.

[120724]21343.

[120725]Hammonds, R. G., Jr.; McKay, P.; Winslow, G. A.; Diefenbach-Jagger,

[120726]H.; Grill, V.; Glatz, J.; Rodda, C. P.; Moseley, J. M.; Wood, W. I.;

[120727]Martin, T. J.: Purification and characterization of recombinant human

[120728]parathyroid hormone-related protein. J. Biol. Chem. 264: 14806-14811,

[120729]1989.

[120730]21344.

[120731]Hendy, G. N.; Sakaguchi, A. Y.; Lalley, P. A.; Martinez, L.; Yasuda,

[120732]T.; Banville, D.; Goltzman, D.: Gene for parathyroid hormone-like

[120733]peptide is on mouse chromosome 6. Cytogenet. Cell Genet. 53: 80-82,

[120734]1990.

[120735]21345.

[120736]Hendy, G. N.; Sakaguchi, A. Y.; Lalley, P. A.; Martinez, L.;
Yasuda,

[120737]T.; Banville, D.; Goltzman, D.: Gene for parathyroid hor-
mone-like

[120738]peptide is on mouse chromosome 6. (Abstract) Cytogenet.
Cell Genet. 51:

[120739]1003 only, 1989.

[120740]21346.

[120741]Kim, S. K.; Selleri, L.; Lee, J. S.; Zhang, A. Y.; Gu, X.; Ja-
cobs,

[120742]Y.; Cleary, M. L.: Pbx1 inactivation disrupts pancreas de-
velopment

[120743]and in Ipf1-deficient mice promotes diabetes mellitus.
Nature Genet. 30:

[120744]430-435, 2002.

[120745]21347.

[120746]Mercader, N.; Leonardo, E.; Azpiazu, N.; Serrano, A.;
Morata, G.;

[120747]Martinez-A, C.; Torres, M.: Conserved regulation of proxi-
modistal

[120748]limb axis development by Meis1/Hth. Nature 402:
425–429, 1999.

[120749]21348.

[120750]Monica, K.; Galili, N.; Nourse, J.; Saltman, D.; Cleary, M. L.

[120751]: PBX2 and PBX3, new homeobox genes with extensive homology to the

[120752]human proto-oncogene PBX1. Molec. Cell. Biol. 11:
6149–6157, 1991.

[120753]21349.

[120754]Aguado, B.; Campbell, R. D.: The novel gene G17, located
in the

[120755]human major histocompatibility complex, encodes PBX2, a
homeodomain-containing

[120756]protein. Genomics 25: 650–659, 1995.

[120757]21350.

[120758]Katsanis, N.; Fitzgibbon, J.; Fisher, E. M. C.: Paralogy mapping:

[120759]identification of a region in the human MHC triplicated
onto human

[120760]chromosomes 1 and 9 allows the prediction and isolation
of novel PBX

[120761]and NOTCH loci. Genomics 35: 101–108, 1996.

[120762]21351.

[120763]Eisenhofer, G.; Walther, M. M.; Huynh, T.–T.; Li, S.–T.;
Bornstein,

[120764]S. R.; Vortmeyer, A.; Mannelli, M.; Goldstein, D. S.; Line-
han, W.

[120765]M.; Lenders, J. W. M.; Pacak, K.: Pheochromocytomas in
von Hippel–Lindau

[120766]syndrome and multiple endocrine neoplasia type 2 display
distinct

[120767]biochemical and clinical phenotypes. J. Clin. Endocr.
Metab. 86:

[120768]1999–2008, 2001.

[120769]21352.

[120770]Glasser, S. W.; Korfhagen, T. R.; Weaver, T. E.; Clark, J. C.;

[120771]Pilot–Matias, T.; Meuth, J.; Fox, J. L.; Whitsett, J. A.: cDNA,
deduced

[120772]polypeptide structure and chromosomal assignment of
human pulmonary

[120773]surfactant proteolipid, SPL(pVal). J. Biol. Chem. 263: 9–12,
1988.

[120774]21353.

[120775]Ahmad, S.; Banville, D.; Zhao, Z.; Fischer, E. H.; Shen, S.-H.

[120776]: A widely expressed human protein-tyrosine phosphatase
containing

[120777]src homology 2 domains. Proc. Nat. Acad. Sci. 90:
2197-2201, 1993.

[120778]21354.

[120779]Bastien, L.; Ramachandran, C.; Liu, S.; Adam, M.: Cloning,
expression,

[120780]and mutational analysis of SH-PTP2, human protein-ty-
rosine phosphatase

[120781]2-domains. Biochem. Biophys. Res. Commun. 196:
124-133, 1993.

[120782]21355.

[120783]Dechert, U.; Duncan, A. M. V.; Bastien, L.; Duff, C.; Adam,
M.;

[120784]Jirik, F. R.: Protein-tyrosine phosphatase SH-PTP2
(PTPN11) is localized

[120785]to 12q24.1-24.3. Hum. Genet. 96: 609-615, 1995.

[120786]21356.

[120787]Higashi, H.; Tsutsumi, R.; Muto, S.; Sugiyama, T.; Azuma,

T; Asaka,

[120788]M.; Hatakeyama, M.: SHP-2 tyrosine phosphatase as an intracellular

[120789]target of *Helicobacter pylori* CagA protein. *Science* 295: 683–686,

[120790]2002.

[120791]21357.

[120792]Hof, P.; Pluskey, S.; Dhe-Paganon, S.; Eck, M. J.; Shoelson, S.

[120793]E.: Crystal structure of the tyrosine phosphatase SHP-2. *Cell* 92:

[120794]441–450, 1998.

[120795]21358.

[120796]Isobe, M.; Hinoda, Y.; Imai, K.; Adachi, M.: Chromosomal localization

[120797]of an SH2 containing tyrosine phosphatase (SH-PTP3) gene to chromosome

[120798]12q24.1. *Oncogene* 9: 1751–1753, 1994.

[120799]21359.

[120800]Qu, C.-K.; Yu, W.-M.; Azzarelli, B.; Cooper, S.; Broxmeyer, H.

- [120801]E.; Feng, G.-S.: Biased suppression of hematopoiesis and multiple
developmental defects in chimeric mice containing Shp-2
mutant cells. Molec.
Cell. Biol. 18: 6075-6082, 1998.
[120804]21360.
- [120805]Saxton, T. M.; Ciruna, B. G.; Holmyard, D.; Kulkarni, S.;
Harpal,
[120806]K.; Rossant, J.; Pawson, T.: The SH2 tyrosine phosphatase
Shp2 is
[120807]required for mammalian limb development. Nature Genet.
24: 420-423,
[120808]2000.
[120809]21361.
- [120810]Saxton, T. M.; Henkemeyer, M.; Gasca, S.; Shen, R.; Rossi,
D.
[120811]J.; Shalaby, F.; Feng, G.-S.; Pawson, T.: Abnormal meso-
derm patterning
[120812]in mouse embryos mutant for the SH2 tyrosine phos-
phatase Shp-2. EMBO
[120813]J. 16: 2352-2364, 1997.

[120814]21362.

[120815]Tartaglia, M.; Kalidas, K.; Shaw, A.; Song, X.; Musat, D. L.;

[120816]van der Burgt, I.; Brunner, H. G.; Bertola, D. R.; Crosby, A.;
Ion,

[120817]A.; Kucherlapati, R. S.; Jeffery, S.; Patton, M. A.; Gelb, B.
D.:

[120818]PTPN11 mutations in Noonan syndrome: molecular spec-
trum, genotype-phenotype

[120819]correlation, and phenotypic heterogeneity. Am. J. Hum.
Genet. 70:

[120820]1555-1563, 2002.

[120821]21363.

[120822]Zhao, Z. J.; Zhao, R.: Purification and cloning of PZR, a
binding

[120823]protein and putative physiological substrate of tyrosine
phosphatase

[120824]SHP-2. J. Biol. Chem. 273: 29367-29372, 1998.

[120825]21364.

[120826]Allaart, C. F.; Aronson, D. C.; Ruys, T.; Rosendaal, F. R.;
van

[120827]Bockel, J. H.; Bertina, R. M.; Briet, E.: Hereditary protein S

deficiency

[120828]in young adults with arterial occlusive disease. Thromb.

Haemost. 64:

[120829]206–210, 1990.

[120830]21365.

[120831]Beauchamp, N. J.; Daly, M. E.; Makris, M.; Preston, F. E.;

Peake,

[120832]J. R.: A novel mutation in intron K of the PROS1 gene

causes aberrant

[120833]RNA splicing and is a common cause of protein S defi-

ciency in a UK

[120834]thrombophilia cohort. Thromb. Haemost. 79: 1086–1091,

1998.

[120835]21366.

[120836]Bertina, R. M.: Nomenclature proposal for protein S defi-

ciency XXXVI

[120837]Annual meeting of the Scientific and Standardization

Committee of

[120838]the ISTH Barcelona, and Spain, 1990.

[120839]21367.

[120840]Bertina, R. M.; Ploos van Amstel, H. K.; van Wijngaarden,

- A.; Coenen,
[120841]J.; Leemhuis, M. P.; Deutz-Terlouw, P. P.; van der Linden,
I. K.;
[120842]Reitsma, P. H.: Heerlen polymorphism of protein S, an im-
munologic
[120843]polymorphism due to dimorphism of residue 460. Blood
76: 538-548,
[120844]1990.
[120845]21368.
[120846]Bosson, J. L.; Francois, P.; Pernod, G.; Carpentier, P.: Mal-
adie
[120847]thrombo-embolique veineuse chez l'enfant liee a un
deficit transitoire
[120848]en proteine S au decours d'une varicelle. (Letter) Presse
Med. 24:
[120849]415 only, 1995.
[120850]21369.
[120851]Chafa, O.; Fischer, A. M.; Meriane, F.; Chellali, F.; Rahal, S.;
[120852]Sternberg, C.; Benabadji, M.: A new case of 'type II' inher-
ited protein
[120853]S deficiency. Brit. J. Haemat. 73: 501-505, 1989.

[120854]21370.

[120855]Clark, D. A.; Williams, W. L.; Marlar, R. A.: Mesenteric vein
[120856]thrombosis associated with a familial deficiency of free
protein S. Arch.

[120857]Path. Lab. Med. 115: 617–619, 1991.

[120858]21371.

[120859]Comp, P. C.; Esmon, C. T.: Recurrent venous thromboem-
bolism in

[120860]patients with a partial deficiency of protein S. New Eng. J.
Med. 311:

[120861]1525–1528, 1984.

[120862]21372.

[120863]Lorimer, D. D.; Benya, R. V.: Cloning and quantification of
galanin-1

[120864]receptor expression by mucosal cells lining the human
gastrointestinal

[120865]tract. Biochem. Biophys. Res. Commun. 222: 379–385,
1996.

[120866]21373.

[120867]Nicholl, J.; Kofler, B.; Sutherland, G. R.; Shine, J.; Iismaa,

[120868]T. P.: Assignment of the gene encoding human galanin receptor (GALNR)

[120869]to 18q23 by in situ hybridization. Genomics 30: 629–630, 1995.

[120870]21374.

[120871]Parker, E. M.; Izzarelli, D. G.; Nowak, H. P.; Mahle, C. D.; Iben,

[120872]L. G.; Wang, J.; Goldstein, M. E.: Cloning and characterization of

[120873]the rat GALR1 galanin receptor from Rin14B insulinoma cells. Molec.

[120874]Brain Res. 34: 179–189, 1995.

[120875]21375.

[120876]Simoneaux, D. K.; Leach, R. J.; O'Connell, P.: Galanin receptor

[120877]1 gene (Galnr1) is tightly linked to the myelin basic protein gene

[120878]on chromosome 18 in mouse. Mammalian Genome 8: 875 only, 1997.

[120879]21376.

[120880]Walli, R.; Schafer, H.; Morys-Wortmann, C.; Paetzold, G.; Nustede,

[120881]R.; Schmidt, W. E.: Identification and biochemical characterization
of the human brain galanin receptor. J. Molec. Endocr. 13:
347–356,
1994.
21377.

[120885]Futterer, A.; Kruppa, G.; Kramer, B.; Lemke, H.; Kronke,
M.: Molecular
cloning and characterization of human kinectin. Molec.
Biol. Cell 6:
161–170, 1995.
21378.

[120889]Plitz, T.; Pfeffer, K.: Intact lysosome transport and phago-
some
function despite kinectin deficiency. Molec. Cell. Biol. 21:
6044–6055,
2001.
21379.

[120893]Print, C. G.; Leung, E.; Harrison, J. E. B.; Watson, J. D.;
Krissansen,
G. W.: Cloning of a gene encoding a human leukocyte

protein characterised

[120895]by extensive heptad repeats. Gene 144: 221–228, 1994.

[120896]21380.

[120897]Print, C. G.; Morris, C. M.; Spurr, N. K.; Rooke, L.; Kris-
sansen,

[120898]G. W.: The CG-1 gene, a member of the kinectin and ES/
130 family,

[120899]maps to human chromosome band 14q22. Immunogenet-
ics 43: 227–229,

[120900]1996.

[120901]21381.

[120902]Rao, P. N.; Yu, H.; Hodge, R.; Pettenati, M. J.; Sheetz, M. P.

[120903]: Assignment of the human kinectin gene (KTN1), encod-
ing a kinesin-binding

[120904]protein, to chromosome 14 band q22.1 by in situ hy-
bridization. Cytogenet.

[120905]Cell Genet. 79: 196–197, 1997.

[120906]21382.

[120907]Yu, H.; Nicchitta, C. V.; Kumar, J.; Becker, M.; Toyoshima,
I.;

[120908]Sheetz, M. P.: Characterization of kinectin, a kinesin-

binding protein:

[120909]primary sequence and N-terminal topogenic signal analysis. Molec.

[120910]Biol. Cell 6: 171-183, 1995.

[120911]21383.

[120912]Gordenin, D. A.; Kunkel, T. A.; Resnick, M. A.: Repeat expansion--all

[120913]in a flap? Nature Genet. 16: 116-118, 1997.

[120914]21384.

[120915]Greene, A. L.; Snipe, J. R.; Gordenin, D. A.; Resnick, M. A.:

[120916]Functional analysis of human FEN1 in *Saccharomyces cerevisiae* and

[120917]its role in genome stability. Hum. Molec. Genet. 8: 2263-2273, 1999.

[120918]21385.

[120919]Nakatsuru, S.; Sudo, K.; Nakamura, Y.: Isolation and mapping of

[120920]a human gene (MCM2) encoding a product homologous to yeast proteins

[120921]involved in DNA replication. Cytogenet. Cell Genet. 68: 226-230,

[120922]1995.

[120923]21386.

[120924]Augusseau, S.; Jouk, S.; Jalbert, P.; Prieur, M.: DiGeorge syndrome

[120925]and 22q11 rearrangements. (Letter) Hum. Genet. 74: 206 only, 1986.

[120926]21387.

[120927]Van Esch, H.; Groenen, P.; Daw, S.; Poffyn, A.; Holvoet, M.; Scambler,

[120928]P.; Fryns, J.-P.; Van de Ven, W.; Devriendt, K.: Partial DiGeorge

[120929]syndrome in two patients with a 10p rearrangement. Clin. Genet. 55:

[120930]269-276, 1999.

[120931]21388.

[120932]Wadey, R.; Daw, S.; Taylor, C.; Atif, U.; Kamath, S.; Halford,

[120933]S.; O'Donnell, H.; Wilson, D.; Goodship, J.; Burn, J.; Scambler, P.

[120934]: Isolation of a gene encoding an integral membrane protein from the

[120935]vicinity of a balanced translocation breakpoint associated

with DiGeorge

[120936]syndrome. Hum. Molec. Genet. 4: 1027–1033, 1995.

[120937]21389.

[120938]Zhao, Z.; Lee, C.-C.; Jiralerspong, S.; Juyal, R. C.; Lu, F.;
Baldini,

[120939]A.; Greenberg, F.; Caskey, C. T.; Patel, P. I.: The gene for a
human

[120940]microfibril-associated glycoprotein is commonly deleted
in Smith–Magenis

[120941]syndrome patients. Hum. Molec. Genet. 4: 589–597,
1995.

[120942]21390.

[120943]Contractor, A.; Rogers, C.; Maron, C.; Henkemeyer, M.;
Swanson,

[120944]G. T.; Heinemann, S. F.: Trans-synaptic Eph receptor-
ephrin signaling

[120945]in hippocampal mossy fiber LTP. Science 296: 1864–1869,
2002.

[120946]21391.

[120947]Tang, X. X.; Biegel, J. A.; Nycum, L. M.; Yoshioka, A.;
Brodeur,

[120948]G. M.; Pleasure, D. E.; Ikegaki, N.: cDNA cloning, molecular

characterization,

[120949]and chromosomal localization of NET (EPHT2), a human
EPH-related receptor

[120950]protein-tyrosine kinase gene preferentially expressed in
brain. :Genomics 29:

[120951]426-437, 1995.

[120952]21392.

[120953]Kamp, C.; Hirschmann, P.; Voss, H.; Huellen, K.; Vogt, P.
H.:

[120954]Two long homologous retroviral sequence blocks in prox-
imal Yq11 cause

[120955]AZFa microdeletions as a result of intrachromosomal re-
combination

[120956]events. Hum. Molec. Genet. 9: 2563-2572, 2000.

[120957]21393.

[120958]Krausz, C.; Meyts, E. R.-D.; Frydelund-Larson, L.; Quin-
tana-Murci,

[120959]L.; McElreavey, K.; Skakkebaek, N. E.: Double-blind Y
chromosome

[120960]microdeletion analysis in men with known sperm parame-
ters and reproductive

[120961]hormone profiles: microdeletions are specific for sper-

matogenic failure. J.

[120962]Clin. Endocr. Metab. 86: 2638–2642, 2001.

[120963]21394.

[120964]Simpson, E.; Chandler, P.; Goulmy, E.; Ma, K.; Hargreave, T. B.;

[120965]Chandley, A. C.: Loss of the 'azoospermia factor' (AZF) on Yq in

[120966]man is not associated with loss of HYA. Hum. Molec. Genet. 2: 469–471,

[120967]1993.

[120968]21395.

[120969]Sun, C.; Skaletsky, H.; Rozen, S.; Gromoll, J.; Nieschlag, E.;

[120970]Oates, R.; Page, D. C.: Deletion of azoospermia factor a (AZFa) region

[120971]of human Y chromosome caused by recombination between HERV15 proviruses. Hum.

[120972]Molec. Genet. 9: 2291–2296, 2000.

[120973]21396.

[120974]Krausz, C.; Quintana–Murci, L.; Barbaux, S.; Siffroi, J.–P.; Rouba,

[120975]H.; Delafontaine, D.; Souleyreau–Therville, N.; Arvis, G.;

Antoine,

[120976]J. M.; Erdei, E.; Taar, J. P.; Tar, A.; Jeandidier, E.; Plessis, G.;

[120977]Bourgeron, T.; Dadoune, J.-P.; Fellous, M.; McElreavey, K.:
A high

[120978]frequency of Y chromosome deletions in males with noni-
diopathic infertility. J.

[120979]Clin. Endocr. Metab. 84: 3606–3612, 1999.

[120980]21397.

[120981]Ma, K.; Inglis, J. D.; Sharkey, A.; Bickmore, W. A.; Hill, R.

[120982]E.; Prosser, E. J.; Speed, R. M.; Thomson, E. J.; Jobling, M.;
Taylor,

[120983]K.; Wolfe, J.; Cooke, H. J.; Hargreave, T. B.; Chandley, A.
C.: A

[120984]Y chromosome gene family with RNA-binding protein ho-
mology: candidates

[120985]for the azoospermia factor AZF controlling human sper-
matogenesis. Cell 75:

[120986]1287–1295, 1993.

[120987]21398.

[120988]Vergnaud, G.; Page, D. C.; Simmler, M. C.; Brown, L.;

Rouyer,

[120989]F.; Noel, B.; Botstein, D.; de la Chapelle, A.; Weissenbach,
J.:

[120990]A deletion map of the human Y chromosome based on
DNA hybridization. Am.

[120991]J. Hum. Genet. 38: 109–124, 1986.

[120992]21399.

[120993]Vogt, P.; Chandley, A. C.; Hargreave, T. B.; Keil, R.; Ma, K.;

[120994]Sharkey, A.: Microdeletions in interval 6 of the Y chromo-
some of

[120995]males with idiopathic sterility point to disruption of AZF, a
human

[120996]spermatogenesis gene. Hum. Genet. 89: 491–496, 1992.

[120997]21400.

[120998]Vogt, P. H.; Edelmann, A.; Kirsch, S.; Henegariu, O.;
Hirschmann,

[120999]P.; Kiesewetter, F.; Kohn, F. M.; Schill, W. B.; Farah, S.;
Ramos,

[121000]C.; Hartmann, M.; Hartschuh, W.; Meschede, D.; Behre, H.
M.; Castel,

[121001]A.; Nieschlag, E.; Weidner, W.; Grone, H.-J.; Jung, A.; En-

gel, W.;

[121002]Haidl, G.: Human Y chromosome azoospermia factors
(AZF) mapped to

[121003]different subregions in Yq11. Hum. Molec. Genet. 5:
933-943, 1996.

[121004]21401.

[121005]Hawkins, J. R.: Mutational analysis of SRY in XY females.
Hum.

[121006]Mutat. 2: 347-350, 1993.

[121007]21402.

[121008]Hawkins, J. R.; Taylor, A.; Berta, P.; Levilliers, J.; Van der

[121009]Auwera, B.; Goodfellow, P. N.: Mutational analysis of SRY:
nonsense

[121010]and missense mutations in XY sex reversal. Hum. Genet.
88: 471-474,

[121011]1992.

[121012]21403.

[121013]Rohen, C.; Caselitz, J.; Stern, C.; Wanschura, S.; Schoen-
makers,

[121014]E. F.; Van de Ven, W. J.; Barnitzke, S.; Bullerdiek, J.: A
hamartoma

- [121015] of the breast with an aberration of 12q mapped to the
MAR region by
[121016] fluorescence in situ hybridization. Genes Chromosomes
Cancer 84:
[121017] 82–84, 1995.
[121018] 21404.
- [121019] Schoenmakers, E. F. P. M.; Huysmans, C.; Van de Ven, W. J.
M.
[121020]: Allelic knockout of novel splice variants of human re-
combination
[121021] repair gene RAD51B in t(12;14) uterine leiomyomas. Can-
cer Res. 59:
[121022] 19–23, 1999.
[121023] 21405.
- [121024] Zhou, X.; Benson, K. F.; Ashar, H. R.; Chada, K.: Mutation
responsible
[121025] for the mouse pygmy phenotype in the developmentally
regulated factor
[121026] HMGI-C. Nature 377: 771–774, 1995.
[121027] 21406.
- [121028] Petit, M. M. R.; Mols, R.; Schoenmakers, E. F. P. M.; Man-

dahl,

[121029]N.; Van de Ven, W. J. M.: LPP, the preferred fusion partner
gene

[121030]of HMGIC in lipomas, is a novel member of the LIM protein
gene family. Genomics 36:

[121031]118–129, 1996.

[121032]21407.

[121033]Burgess, D. L.; Kohrman, D. C.; Galt, J.; Plummer, N. W.;
Jones,

[121034]J. M.; Spear, B.; Meisler, M. H.: Mutation of a new sodium
channel

[121035]gene, Scn8a, in the mouse mutant 'motor endplate dis-
ease'. Nature

[121036]Genet. 10: 461–465, 1995.

[121037]21408.

[121038]DeRepentigny, Y.; Cote, P. D.; Pool, M.; Bernier, G.; Girard,
S.;

[121039]Vidal, S. M.; Kothary, R.: Pathological and genetic analysis
of the

[121040]degenerating muscle (dmu) mouse: a new allele of Scn8a.
Hum. Molec.

[121041]Genet. 10: 1819–1827, 2001.

[121042]21409.

[121043]Kohrman, D. C.; Harris, J. B.; Meisler, M. H.: Mutation detection

[121044]in the med and med(j) alleles of the sodium channel
Scn8a: unusual

[121045]splicing due to a minor class AT-AC intron. J. Biol. Chem.
271:

[121046]17576-17581, 1996.

[121047]21410.

[121048]Kohrman, D. C.; Plummer, N. W.; Schuster, T.; Jones, J. M.;
Jang,

[121049]W.; Burgess, D. L.; Galt, J.; Spear, B. T.; Meisler, M. H.: In-
sertional

[121050]mutation of the motor endplate disease (med) locus on
mouse chromosome

[121051]15. Genomics 26: 171-177, 1995.

[121052]21411.

[121053]Kohrman, D. C.; Smith, M. R.; Goldin, A. L.; Harris, J.;
Meisler,

[121054]M. H.: A missense mutation in the sodium channel Scn8a
is responsible

[121055]for cerebellar ataxia in the mouse mutant jolting. J. Neurosci. 16:

[121056]5993–5999, 1996.

[121057]21412.

[121058]Meisler, M. H.; Sprunger, L. K.; Plummer, N. W.; Escayg, A.; Jones,

[121059]J. M.: Ion channel mutations in mouse models of inherited neurological

[121060]disease. Ann. Med. 29: 569–574, 1997.

[121061]21413.

[121062]Plummer, N. W.; Galt, J.; Jones, J. M.; Burgess, D. L.; Sprunger,

[121063]L. K.; Kohrman, D. C.; Meisler, M. H.: Exon organization, coding

[121064]sequence, physical mapping, and polymorphic intragenic markers for

[121065]the human neuronal sodium channel gene SCN8A. Genomics 54: 287–296,

[121066]1998.

[121067]21414.

[121068]Plummer, N. W.; McBurney, M. W.; Meisler, M. H.: Alterna-

tive splicing

[121069]of the sodium channel SCN8A predicts a truncated two-domain protein

[121070]in fetal brain and non-neuronal cells. J. Biol. Chem. 272: 24008–24015,

[121071]1997.

[121072]21415.

[121073]Reid, E.; Escayg, A.; Dearlove, A. M.; Lee, D. D.; Meisler, M.

[121074]H.; Rubinsztein, D. C.: The spastic paraplegia SPG10 locus: narrowing

[121075]of critical region and exclusion of sodium channel gene SCN8A as a

[121076]candidate. J. Med. Genet. 38: 65–67, 2001.

[121077]21416.

[121078]Sprunger, L. K.; Escayg, A.; Tallaksen–Greene, S.; Albin, R. L.;

[121079]Meisler, M. H.: Dystonia associated with mutation of the neuronal

[121080]sodium channel Scn8a and identification of the modifier locus Scnm1

[121081]on mouse chromosome 3. Hum. Molec. Genet. 8:

471-479, 1999.

[121082]21417.

[121083]Casey, J. L.; Di Jeso, B.; Rao, K.; Klausner, R. D.; Harford, J.

[121084]B.: Two genetic loci participate in the regulation by iron of
the

[121085]gene for the human transferrin receptor. Proc. Nat. Acad.
Sci. 85:

[121086]1787-1791, 1988.

[121087]21418.

[121088]Enns, C. A.; Suomalainen, H. A.; Gebhardt, J. E.; Schroder,
J.;

[121089]Sussman, H. H.: Human transferrin receptor: expression of
the receptor

[121090]is assigned to chromosome 3. Proc. Nat. Acad. Sci. 79:
3241-3245,

[121091]1982.

[121092]21419.

[121093]Gareau, R.; Gagnon, M. G.; Thellend, C.; Chenard, C.; Au-
dran, M.;

[121094]Chanal, J.-L.; Ayotte, C.; Brisson, G. R.: Transferrin soluble
receptor:

[121095]a possible probe for detection of erythropoietin abuse by athletes. Horm.

[121096]Metab. Res. 26: 311–312, 1994.

[121097]21420.

[121098]Goodfellow, P. N.; Banting, G.; Sutherland, R.; Greaves, M.; Solomon,

[121099]E.; Povey, S.: Expression of human transferrin receptor is controlled

[121100]by a gene on chromosome 3: assignment using species specificity of

[121101]a monoclonal antibody. Somat. Cell Genet. 8: 197–206, 1982.

[121102]21421.

[121103]Larrick, J. W.; Hyman, E. S.: Acquired iron-deficiency anemia

[121104]caused by an antibody against the transferrin receptor. New Eng.

[121105]J. Med. 311: 214–218, 1984.

[121106]21422.

[121107]Levy, J. E.; Jin, O.; Fujiwara, Y.; Kuo, F.; Andrews, N. C.: Transferrin

[121108]receptor is necessary for development of erythrocytes and the nervous

[121109]system. Nature Genet. 21: 396–399, 1999.

[121110]21423.

[121111]Miller, Y. E.; Jones, C.; Scoggin, C.; Morse, H.; Seligman, P.

[121112]; Chromosome 3q (22–ter) encodes the human transferrin receptor. Am.

[121113]J. Hum. Genet. 35: 573–583, 1983.

[121114]21424.

[121115]Newman, R.; Schneider, C.; Sutherland, R.; Vodinelich, L.; Greaves,

[121116]M.: The transferrin receptor. Trends Biochem. Sci. 7: 397–400,

[121117]1982.

[121118]21425.

[121119]Nikinmaa, B.; Schroder, J.: Two antigens, the transferrin receptor

[121120]and p90 assigned to human chromosome 3, are probably the same protein. Hereditas 107:

[121121]55–58, 1987.

[121122]21426.

[121123]Omary, M. B.; Trowbridge, I. S.: Biosynthesis of the human transferrin

[121124]receptor in cultured cells. J. Biol. Chem. 256: 12888–12892, 1981.

[121125]21427.

[121126]Kashuba, V. I.; Gizatullin, R. Z.; Protopopov, A. I.; Allik-mets,

[121127]R.; Korolev, S.; Li, J.; Boldog, F.; Tory, K.; Zabarovska, V.; Marcsek,

[121128]Z.; Sumegi, J.; Klein, G.; Zabarovsky, E. R.; Kisselev, L.: NotI

[121129]linking/jumping clones of human chromosome 3: mapping of the TFRC,

[121130]RAB7 and HAUSP genes to regions rearranged in leukemia and deleted

[121131]in solid tumors. FEBS Lett. 419: 181–185, 1997.

[121132]21428.

[121133]Rabin, M.; McClelland, A.; Kuhn, L.; Ruddle, F. H.: Regional

[121134]localization of the human transferrin receptor gene to 3q26.2–qter. Am.

[121135]J. Hum. Genet. 37: 1112–1116, 1985.

[121136]21429.

[121137]Schneider, C.; Kurkinen, M.; Greaves, M.: Isolation of cDNA clones

[121138]for the human transferrin receptor. EMBO J. 2:
2259–2263, 1983.

[121139]21430.

[121140]Schneider, C.; Owen, M. J.; Banville, D.; Williams, J. G.: Primary

[121141]structure of human transferrin receptor deduced from the mRNA sequence. Nature 311:

[121142]675–678, 1984.

[121143]21431.

[121144]Valenzuela, C. Y.; Avendano, A.; Harb, Z.: Association between

[121145]Rh and plasma iron binding (transferrin). Hum. Genet. 87:
438–440,

[121146]1991.

[121147]21432.

[121148]Vodinelich, L.; Sutherland, R.; Schneider, C.; Newman, R.;
Greaves,

[121149]M.: Receptor for transferrin may be a 'target' structure for natural

[121150]killer cells. Proc. Nat. Acad. Sci. 80: 835–839, 1983.

[121151]21433.

[121152]de Martinville, B.; Francke, U.: The c-Ha-ras1, insulin and beta-globin

[121153]loci map outside the deletion associated with aniridia–Wilms' tumour. Nature 305:

[121154]641–643, 1983.

[121155]21434.

[121156]Hamm, C. W.; Ravkilde, J.; Gerhardt, W.; Jorgensen, P.; Peheim,

[121157]E.; Ljungdahl, L.; Goldmann, B.; Katus, H. A.: The prognostic value

[121158]of serum troponin T in unstable angina. New Eng. J. Med. 327: 146–150,

[121159]1992.

[121160]21435.

[121161]Carmeliet, P.; Ferreira, V.; Breler, G.; Pollefeyt, S.; Kleckens,

[121162]L.; Gertsenstein, M.; Fahrig, M.; Vandenhoeck, A.; Harpal,

K.; Eberhardt,

[121163]C.; Declercq, C.; Pawling, J.; Moons, L.; Collen, D.; Risau,
W.; Nagy,

[121164]A.: Abnormal blood vessel development and lethality in
embryos lacking

[121165]a single VEGF allele. Nature 380: 435–439, 1996.

[121166]21436.

[121167]Martin, L. H.; Calabi, F.; Lefebvre, F.–A.; Bilsland, C. A. G.;

[121168]Milstein, C.: Structure and expression of the human thymocyte antigens

[121169]CD1a, CD1b, and CD1c. Proc. Nat. Acad. Sci. 84:
9189–9193, 1987.

[121170]21437.

[121171]Martin, L. H.; Calabi, F.; Milstein, C.: Isolation of CD1
genes:

[121172]a family of major histocompatibility complex–related differentiation

[121173]antigens. Proc. Nat. Acad. Sci. 83: 9154–9158, 1986.

[121174]21438.

[121175]Moseley, W. S.; Watson, M. L.; Kingsmore, S. F.; Seldin, M.
F.

- [121176]: CD1 defines conserved linkage group border between human chromosomes
- [121177]1 and mouse chromosomes 1 and 3. Immunogenetics 30: 378–382, 1989.
- [121178]21439.
- [121179]Samson, F.; de Jong, P. J.; Trask, B. J.; Koza–Taylor, P.; Speer,
- [121180]M. C.; Potter, T.; Roses, A. D.; Gilbert, J. R.: Assignment of the
- [121181]human slow skeletal troponin T gene to 19q13.4 using somatic cell
- [121182]hybrids and fluorescence in situ hybridization analysis. Genomics 13:
- [121183]1374–1375, 1992.
- [121184]21440.
- [121185]Samson, F.; Gilbert, J. R.; Lee, J. E.; Potter, T. G.; Koza–Taylor,
- [121186]P.; Speer, M. C.; Bachinski, L. L.; Siciliano, M. J.; Roses, A. D.
- [121187]: Isolation and localization of a human slow troponin T gene on chromosome
- [121188]19q. (Abstract) Cytogenet. Cell Genet. 58: 2025, 1991.

[121189]21441.

[121190]Samson, F.; Lee, J. E.; Hung, W. Y.; Potter, T. G.; Herbstreith,

[121191]M.; Roses, A. D.; Gilbert, J. R.: Isolation and localisation of
a

[121192]slow troponin (TNT) gene on chromosome 19 by subtraction of a cDNA

[121193]muscle library using myotonic muscle cDNA. J. Neurosci.
Res. 27:

[121194]441–451, 1990.

[121195]21442.

[121196]Samson, F.; Mesnard, L.; Mihovilovic, M.; Potter, T. G.;
Mercadier,

[121197]J.-J.; Roses, A. D.; Gilbert, J. R.: A new human slow skeletal
troponin

[121198]T (TnTs) mRNA isoform derived from alternative splicing
of a single

[121199]gene. Biochem. Biophys. Res. Commun. 199: 841–847,
1994.

[121200]21443.

[121201]Johnston, J. J.; Kelley, R. I.; Crawford, T. O.; Morton, D. H.;

[121202]Agarwala, R.; Koch, T.; Schaffer, A. A.; Francomano, C. A.;
Biesecker,

[121203]L. G.: A novel nemaline myopathy in the Amish caused by
a mutation

[121204]in troponin T1. Am. J. Hum. Genet. 67: 814–821, 2000.

[121205]21444.

[121206]Novelli, G.; Gennarelli, M.; Rocchi, M.; Dallapiccola, B.: As-
signment

[121207]of the slow troponin T (TNNT1) gene to chromosome 19
using polymerase

[121208]chain reaction. Hum. Genet. 88: 697–698, 1992.

[121209]21445.

[121210]Novelli, G.; Gennarelli, M.; Zelano, G.; Sangiuolo, F.; Roc-
chi,

[121211]M.; Dallapiccola, B.: Isolation and mapping of the slow
skeletal

[121212]troponin T using the polymerase chain reaction. (Abstract)
Cytogenet.

[121213]Cell Genet. 58: 2023, 1991.

[121214]21446.

[121215]Guenet, J.-L.; Simon-Chazottes, D.; Gravel, M.; Hastings,

K. E.

[121216]M.; Schiaffino, S.: Cardiac and skeletal muscle troponin I isoforms

[121217]are encoded by a dispersed gene family on mouse chromosomes 1 and

[121218]7. Mammalian Genome 7: 13–15, 1996.

[121219]21447.

[121220]Wade, R.; Eddy, R.; Shows, T. B.; Kedes, L.: cDNA sequence, tissue-specific

[121221]expression, and chromosomal mapping of the human slow-twitch skeletal

[121222]muscle isoform of troponin I. Genomics 7: 346–357, 1990.

[121223]21448.

[121224]Bhavsar, P. K.; Brand, N. J.; Yacoub, M. H.; Barton, P. J. R.:

[121225]Isolation and characterization of the human cardiac troponin I gene

[121226](TNNI3). Genomics 35: 11–23, 1996.

[121227]21449.

[121228]Kimura, A.; Harada, H.; Park, J.-E.; Nishi, H.; Satoh, M.; Takahashi,

[121229]M.; Hiroi, S.; Sasaoka, T.; Ohbuchi, N.; Nakamura, T.; Koyanagi, T.;

[121230]Hwang, T.-H.; Choo, J.; Chung, K.-S.; Hasegawa, A.; Nagai, R.; Okazaki,

[121231]O.; Nakamura, H.; Matsuzaki, M.; Sakamoto, T.; Toshima, H.; Koga,

[121232]Y.; Imaizumi, T.; Sasazuki, T.: Mutations in the cardiac troponin

[121233]I gene associated with hypertrophic cardiomyopathy. Nature Genet. 16:

[121234]379–382, 1997.

[121235]21450.

[121236]Labarrere, C. A.; Nelson, D. R.; Cox, C. J.; Pitts, D.; Kirlin,

[121237]P.; Halbrook, H.: Cardiac-specific troponin I levels and risk of

[121238]coronary artery disease and graft failure following heart transplantation. J.A.M.A. 284:

[121239]457–464, 2000.

[121240]21451.

[121241]Lang, R.; Gomes, A. V.; Zhao, J.; Housmans, P. R.; Miller, T.;

- [121242]Potter, J. D.: Functional analysis of a troponin I (R145G) mutation
- [121243]associated with familial hypertrophic cardiomyopathy. J. Biol. Chem. 277: 11670–11678, 2002.
- [121244]21452.
- [121245]MacGeoch, C.; Barton, P. J. R.; Vallins, W. J.; Bhavsar, P.; Spurr, N. K.: The human cardiac troponin I locus: assignment to 19p13.2–19q13.2. Hum. Genet. 88: 101–104, 1991.
- [121246]21453.
- [121247]Mogensen, J.; Kruse, T. A.; Borglum, A. D.: Assignment of the human cardiac troponin I gene (TNNI3) to chromosome 19q13.4 by radiation hybrid mapping. Cytogenet. Cell Genet. 79: 272–273, 1997.
- [121248]21454.
- [121249]Crow, R. S.: Peripheral neuritis in myelomatosis. Brit. Med. J. 2: 802–804, 1956.

[121256]21455.

[121257]Dantz, D.; Bewersdorf, J.; Fruehwald-Schultes, B.; Kern, W.; Jelkmann,

[121258]W.; Born, J.; Fehm, H. L.; Peters, A.: Vascular endothelial growth

[121259]factor: a novel endocrine defensive response to hypoglycemia. J.

[121260]Clin. Endocr. Metab. 87: 835–840, 2002.

[121261]21456.

[121262]Ferrara, N.; Carver-Moore, K.; Chen, H.; Dowd, M.; Lu, L.; O'Shea,

[121263]K. S.; Powell-Braxton, L.; Hillan, K. J.; Moore, M. W.: Heterozygous

[121264]embryonic lethality induced by targeted inactivation of the VEGF gene. Nature 380:

[121265]439–442, 1996.

[121266]21457.

[121267]Folkman, J.: Angiogenesis in cancer, vascular, rheumatoid and

[121268]other disease. Nature Med. 1: 27–31, 1995.

[121269]21458.

[121270]Fukumura, D.; Xavier, R.; Sugiura, T.; Chen, Y.; Park, E.-C.;
Lu,

[121271]N.; Selig, M.; Nielsen, G.; Taksir, T.; Jain, R. K.; Seed, B.:
Tumor

[121272]induction of VEGF promoter activity in stromal cells. Cell
94: 715–725,

[121273]1998.

[121274]21459.

[121275]Gerber, H.-P.; Malik, A. K.; Solar, G. P.; Sherman, D.;
Liang,

[121276]X. H.; Meng, G.; Hong, K.; Marsters, J. C.; Ferrara, N.: VEGF
regulates

[121277]haematopoietic stem cell survival by an internal autocrine loop mechanism. Nature 417: 954–958, 2002.

[121279]21460.

[121280]Gerber, H.-P.; Vu, T. H.; Ryan, A. M.; Kowalski, J.; Werb, Z.; Ferrara, N.: VEGF couples hypertrophic cartilage remodeling, ossification and angiogenesis during endochondral bone formation. Nature Med. 5: 623–628, 1999.

[121284]21461.

[121285]Giordano, F. J.; Gerber, H.-P.; Williams, S.-P.; VanBruggen, N.; Bunting, S.; Ruiz-Lozano, P.; Gu, Y.; Nath, A. K.; Huang, Y.; Hickey, R.; Dalton, N.; Peterson, K. L.; Ross, J., Jr.; Chien, K. R.; Ferrara, N.: A cardiac myocyte vascular endothelial growth factor paracrine pathway is required to maintain cardiac function. Proc. Nat. Acad.

[121290]Sci. 98: 5780–5785, 2001.

[121291]21462.

[121292]Helmlinger, G.; Endo, M.; Ferrara, N.; Hlatky, L.; Jain, R. K.

[121293]: Formation of endothelial cell networks. Nature 405:
139–141, 2000.

[121294]21463.

[121295]Hofman, P.; van Blijswijk, B. C.; Gaillard, P. J.; Vrensen, G.

[121296]F. J. M.; Schlingemann, R. O.: Endothelial cell hypertrophy
induced

[121297]by vascular endothelial growth factor in the retina: new
insights

[121298]into the pathogenesis of capillary nonperfusion. Arch.
Ophthalm. 119:

[121299]861–866, 2001.

[121300]21464.

[121301]Holash, J.; Maisonpierre, P. C.; Compton, D.; Boland, P.;
Alexander,

[121302]C. R.; Zagzag, D.; Yancopoulos, G. D.; Wiegand, S. J.: Ves-
sel cooption,

[121303]regression, and growth in tumors mediated by angiopo-
etins and VEGF. Science 284:

[121304]1994–1998, 1999.

[121305]21465.

[121306]Jin, K.; Zhu, Y.; Sun, Y.; Mao, X. O.; Xie, L.; Greenberg, D.

[121307]A.: Vascular endothelial growth factor (VEGF) stimulates
neurogenesis

[121308]in vitro and in vivo. Proc. Nat. Acad. Sci. 99:
11946–11950, 2002.

[121309]21466.

[121310]Mattei, M.–G.; Borg, J.–P.; Rosnet, O.; Marme, D.; Birn-
baum, D.

[121311]: Assignment of vascular endothelial growth factor (VEGF)
and placenta

[121312]growth factor (PIGF) genes to human chromosome
6p12–p21 and 14q24–q31

[121313]regions, respectively. Genomics 32: 168–169, 1996.

[121314]21467.

[121315]Miralles, G. D.; O'Fallon, J. R.; Talley, N. J.: Plasma-cell

[121316]dyscrasia with polyneuropathy: the spectrum of POEMS
syndrome. New

[121317]Eng. J. Med. 327: 1919–1923, 1992.

[121318]21468.

[121319]Mueller, M. D.; Vigne, J.-L.; Minchenko, A.; Lebovic, D. I.;
Leitman,

[121320]D. C.; Taylor, R. N.: Regulation of vascular endothelial
growth factor

[121321](VEGF) gene transcription by estrogen receptors alpha and
beta. Proc.

[121322]Nat. Acad. Sci. 97: 10972-10977, 2000.

[121323]21469.

[121324]Nakanishi, T.; Sobue, I.; Toyokura, Y.; Nishitani, H.;
Kuroiwa,

[121325]Y.; Satoyoshi, E.; Tsubaki, T.; Igata, A.; Ozaki, Y.: The
Crow-Fukase

[121326]syndrome: a study of 102 cases in Japan. Neurology 34:
712-720,

[121327]1984.

[121328]21470.

[121329]Niimi, H.; Arimura, K.; Jonosono, M.; Hashiguchi, T.;
Kawabata,

[121330]M.; Osame, M.; Kitajima, I.: VEGF is causative for pul-
monary hypertension

[121331]in a patient with Crow–Fukase (POEMS) syndrome. Intern.

Med. 39:

[121332]1101–1104, 2000.

[121333]21471.

[121334]Oosthuysen, B.; Moons, L.; Storkebaum, E.; Beck, H.;

Nuyens, D.;

[121335]Brusselmans, K.; Van Dorpe, J.; Hellings, P.; Gorselink, M.;

Heymans,

[121336]S.; Theilmeier, G.; Dewerchin, M.; and 20 others: Deletion

of the

[121337]hypoxia–response element in the vascular endothelial

growth factor

[121338]promoter causes motor neuron degeneration. Nature

Genet. 28: 131–138,

[121339]2001.

[121340]21472.

[121341]Poltorak, Z.; Cohen, T.; Sivan, R.; Kandelis, Y.; Spira, G.;

Vlodavsky,

[121342]I.; Keshet, E.; Neufeld, G.: VEGF145, a secreted vascular

endothelial

[121343]growth factor isoform that binds to extracellular matrix. J.

Biol.

[121344]Chem. 272: 7151–7158, 1997.

[121345]21473.

[121346]Shimpo, S.: Solitary myeloma causing polyneuritis and endocrine

[121347]disorders. Nippon Rinsho 26: 2444–2456, 1968.

[121348]21474.

[121349]Soker, S.; Takashima, S.; Miao, H. Q.; Neufeld, G.; Klagsbrun,

[121350]M.: Neuropilin-1 is expressed by endothelial and tumor cells as an

[121351]isoform-specific receptor for vascular endothelial growth factor. Cell 92:

[121352]735–745, 1998.

[121353]21475.

[121354]Sone, H.; Kawakami, Y.; Sakauchi, M.; Nakamura, Y.; Takahashi,

[121355]A.; Shimano, H.; Okuda, Y.; Segawa, T.; Suzuki, H.; Yamada, N.: Neutralization

[121356]of vascular endothelial growth factor prevents collagen-induced arthritis

[121357]and ameliorates established disease in mice. Biochem.

Biophys. Res.

[121358]Commun. 281: 562–568, 2001.

[121359]21476.

[121360]Springer, M. L.; Chen, A. S.; Kraft, P. E.; Bednarski, M.;
Blau,

[121361]H. M.: VEGF gene delivery to muscle: potential role for
vasculogenesis

[121362]in adults. Molec. Cell 2: 549–558, 1998.

[121363]21477.

[121364]Thurston, G.; Suri, C.; Smith, K.; McClain, J.; Sato, T. N.;
Yancopoulos,

[121365]G. D.; McDonald, D. M.: Leakage-resistant blood vessels in
mice transgenically

[121366]overexpressing angiopoietin-1. Science 286: 2511–2514,
1999.

[121367]21478.

[121368]Tischer, E.; Mitchell, R.; Hartman, T.; Silva, M.; Gospo-
darowicz,

[121369]D.; Fiddes, J. C.; Abraham, J. A.: The human gene for vas-
cular endothelial

[121370]growth factor: multiple protein forms are encoded

through alternative

[121371]exon splicing. J. Biol. Chem. 266: 11947–11954, 1991.

[121372]21479.

[121373]Ahlgren, A.; Yanase, T.; Simpson, E. R.; Winter, J. S. D.;
Waterman,

[121374]M. R.: Compound heterozygous mutations (arg239-to-ter,
pro342-to-thr)

[121375]in the CYP17 (P45017-alpha) gene lead to ambiguous ex-
ternal genitalia

[121376]in a male patient with partial combined
17-alpha-hydroxylase/17,20-lyase

[121377]deficiency. J. Clin. Endocr. Metab. 74: 667–672, 1992.

[121378]21480.

[121379]Auchus, R. J.; Miller, W. L.: Molecular modeling of human
P450c17

[121380](17-alpha-hydroxylase/17,20-lyase): insights into reac-
tion mechanisms

[121381]and effects of mutations. Molec. Endocr. 13: 1169–1182,
1999.

[121382]21481.

[121383]Biason, A.; Mantero, F.; Scaroni, C.; Simpson, E. R.; Water-

man,

[121384]M. R.: Deletion within the CYP17 gene together with insertion of

[121385]foreign DNA is the cause of combined complete 17-alpha-hydroxylase/17,20-lyase

[121386]deficiency in an Italian patient. Molec. Endocr. 5: 2037-2045, 1991.

[121387]21482.

[121388]Biaison-Lauber, A.; Kempken, B.; Werder, E.; Forest, M. G.; Einaudi,

[121389]S.; Ranke, M. B.; Matsuo, N.; Brunelli, V.; Schonle, E. J.; Zachmann,

[121390]M.: 17-alpha-hydroxylase/17,20-lyase deficiency as a model to study

[121391]enzymatic activity regulation: role of phosphorylation. J. Clin.

[121392]Endocr. Metab. 85: 1226-1231, 2000.

[121393]21483.

[121394]Biaison-Lauber, A.; Leiberman, E.; Zachmann, M.: A single amino

[121395]acid substitution in the putative redox partner-binding site of P450c17

[121396] as cause of isolated 17,20-lyase deficiency. J. Clin. Endocr. Metab. 82:

[121397] 3807–3812, 1997.

[121398] 21484.

[121399] Biglieri, E.G.: A prismatic case: 17- α -hydroxylase deficiency:

[121400] 1963–1966. J. Clin. Endocr. Metab. 82: 48–50, 1997.

[121401] 21485.

[121402] Osborne, L. R.; Martindale, D.; Scherer, S. W.; Shi, X.-M.; Huizenga,

[121403] J.; Heng, H. H. Q.; Costa, T.; Pober, B.; Lew, L.; Brinkman, J.; Rommens,

[121404] J.; Koop, B.; Tsui, L.-C.: Identification of genes from a 500-kb

[121405] region at 7q11.23 that is commonly deleted in Williams syndrome patients. Genomics 36:

[121406] 328–336, 1996.

[121407] 21486.

[121408] Rampoldi, L.; Dobson-Stone, C.; Rubio, J. P.; Danek, A.; Chalmers,

[121409] R. M.; Wood, N. W.; Verellen, C.; Ferrer, X.; Malandrini, A.;

Fabrizi,

[121410]G. M.; Brown, R.; Vance, J.; Pericak-Vance, M.; Rudolf, G.;
Carre,

[121411]S.; Alonso, E.; Manfredi, M.; Nemeth, A. H.; Monaco, A. P.:
A conserved

[121412]sorting-associated protein is mutant in chorea-
acanthocytosis. Nature

[121413]Genet. 28: 119-120, 2001.

[121414]21487.

[121415]Belden, W. J.; Barlowe, C.: Role of Erv29p in collecting sol-
uble

[121416]secretory proteins into ER-derived transport vesicles. Sci-
ence 294:

[121417]1528-1531, 2001.

[121418]21488.

[121419]Duhig, T.; Ruhrberg, C.; Mor, O.; Fried, M.: The human
surfeit

[121420]locus. Genomics 52: 72-78, 1998.

[121421]21489.

[121422]Reeves, J. E.; Fried, M.: The surf-4 gene encodes a novel
30 kDa

- [121423]integral membrane protein. *Molec. Membr. Biol.* 12:
201–208, 1995.
- [121424]21490.
- [121425]Archer, B. T., III; Ozcelik, T.; Jahn, R.; Francke, U.; Sudhof,
[121426]T. C.: Structures and chromosomal localizations of two
human genes
[121427]encoding synaptobrevins 1 and 2. *J. Biol. Chem.* 265:
17267–17273,
[121428]1990.
- [121429]21491.
- [121430]Hu, K.; Carroll, J.; Fedorovich, S.; Rickman, C.; Sukhodub,
A.;
- [121431]Davietov, B.: Vesicular restriction of synaptobrevin sug-
gests a role
[121432]for calcium in membrane fusion. *Nature* 415: 646–650,
2002.
- [121433]21492.
- [121434]Doggen, C. J. M.; Kunz, G.; Rosendaal, F. R.; Lane, D. A.;
Vos,
- [121435]H. L.; Stubbs, P. J.; Cats, V. M.; Ireland, H.: A mutation in
the

[121436]thrombomodulin gene, 127G to A coding for
ala25-to-thr, and the risk
[121437]of myocardial infarction in men. Thromb. Haemost. 80:
743-748, 1998.
[121438]21493.

[121439]Esmon, N. L.: Thrombomodulin. Semin. Thromb. Hemost.
13: 454-463,
[121440]1987.
[121441]21494.

[121442]Espinosa, R., III; Sadler, J. E.; Le Beau, M. M.: Regional lo-
calization
[121443]of the human thrombomodulin gene to 20p12-cen. Ge-
nomics 5: 649-650,
[121444]1989.
[121445]21495.

[121446]Franchi, F.; Biguzzi, E.; Cetin, I.; Facchetti, F.; Radaelli, T.;
[121447]Bozzo, M.; Pardi, G.; Faioni, E. M.: Mutations in the throm-
bomodulin
[121448]and endothelial protein C receptor genes in women with
late fetal
[121449]loss. Brit. J. Haemat. 114: 641-646, 2001.

[121450]21496.

[121451]Jackman, R. W.; Beeler, D. L.; Fritze, L.; Soff, G.; Rosenberg,

[121452]R. D.: Human thrombomodulin gene is intron depleted:
nucleic acid

[121453]sequences of the cDNA and gene predict protein structure
and suggest

[121454]sites of regulatory control. Proc. Nat. Acad. Sci. 84:
6425–6429,

[121455]1987.

[121456]21497.

[121457]Kunz, G.; Ireland, H. A.; Stubbs, P. J.; Kahan, M.; Coulton,
G.

[121458]C.; Lane, D. A.: Identification and characterization of a
thrombomodulin

[121459]gene mutation coding for an elongated protein with re-
duced expression

[121460]in a kindred with myocardial infarction. Blood 95:
569–576, 2000.

[121461]21498.

[121462]Ohlin, A.-K.; Marlar, R. A.: The first mutation identified in

[121463]the thrombomodulin gene in a 45-year-old man presenting with thromboembolic

[121464]disease. *Blood* 85: 330–336, 1995.

[121465]21499.

[121466]Okura, Y.; Kato, K.; Hanawa, H.; Izumi, T.; Kamishima, T.; Yamato,

[121467]Y.; Emura, I.; Shibata, A.: Pericardial mesothelioma secreting thrombomodulin. *Am.*

[121468]*Heart J.* 132: 1309–1311, 1996.

[121469]21500.

[121470]Wen, D.; Dittman, W. A.; Ye, R. D.; Deaven, L. L.; Majerus, P.

[121471]W.; Sadler, J. E.: Human thrombomodulin: complete cDNA sequence and

[121472]chromosome localization of the gene. *Biochemistry* 26: 4350–4357,

[121473]1987.

[121474]21501.

[121475]Grimm, C.; Wenzel, A.; Hafezi, F.; Yu, S.; Redmond, T. M.; Reme,

[121476]C. E.: Protection of Rpe65-deficient mice identifies

rhodopsin as

[121477]a mediator of light-induced retinal degeneration. Nature
Genet. 25:

[121478]63–66, 2000.

[121479]21502.

[121480]Barone, V.; Bertocchini, F.; Bottinelli, R.; Protasi, F.; Allen,

[121481]P. D.; Armstrong, C. F.; Reggiani, C.; Sorrentino, V.: Con-
tractile

[121482]impairment and structural alterations of skeletal muscles
from knockout

[121483]mice lacking type 1 and type 3 ryanodine receptors. FEBS
Lett. 422:

[121484]160–164, 1998.

[121485]21503.

[121486]Cavanna, J. S.; Greenfield, A. J.; Johnson, K. J.; Marks, A.
R.;

[121487]Nadal-Ginard, B.; Brown, S. D. M.: Establishment of the
mouse chromosome

[121488]7 region with homology to the myotonic dystrophy region
of human chromosome

[121489]19q. Genomics 7: 12–18, 1990.

[121490]21504.

[121491]Cheng, H.; Lederer, W. J.; Cannell, M. B.: Calcium sparks:
elementary

[121492]events underlying excitation-contraction coupling in heart
muscle. Science 262:

[121493]740-744, 1993.

[121494]21505.

[121495]Vallins, W. J.; Brand, N. J.; Dabhade, N.; Butler-Browne, G.;

[121496]Yacoub, M. H.; Barton, P. J. R.: Molecular cloning of human
cardiac

[121497]troponin I using polymerase chain reaction. FEBS Lett.
270: 57-61,

[121498]1990.

[121499]21506.

[121500]Vincenti, V.; Cassano, C.; Rocchi, M.; Persico, M. G.: As-
signment

[121501]of the vascular endothelial growth factor gene to human
chromosome

[121502]6p21.3. Circulation 93: 1493-1495, 1996.

[121503]21507.

- [121504]Biglieri, E. G.; Herron, M. A.; Brust, N.: 17-Hydroxylation deficiency
- [121505]in man. J. Clin. Invest. 45: 1946-1954, 1966.
- [121506]21508.
- [121507]Chung, B.-C.; Picado-Leonard, J.; Haniu, M.; Bienkowski, M.; Hall, M.; Hall, P. F.; Shively, J. E.; Miller, W. L.: Cytochrome P450c17 (steroid
- [121508]17 alpha-hydroxylase/17,20 lyase): cloning of human adrenal and testis
- [121509]cDNAs indicates the same gene is expressed in both tissues. Proc.
- [121510]Nat. Acad. Sci. 84: 407-411, 1987.
- [121511]21509.
- [121512]D'Armiento, M.; Reda, G.; Bisignani, G.; Tabolli, S.; Capel-laci,
- [121513]S.; Lulli, P.; Carbonara, A.; Biglieri, E. G.: No linkage between
- [121514]HLA and congenital adrenal hyperplasia due to 17-alpha-hydroxylase
- [121515]deficiency. (Letter) New Eng. J. Med. 308: 970-971, 1983.
- [121516]

[121517]21510.

[121518]Di Cerbo, A.; Biason-Laubert, A.; Savino, M.; Piemontese, M. R.;

[121519]Di Giorgio, A.; Perona, M.; Savoia, A.: Combined
17- α -hydroxylase/17,20-lyase

[121520]deficiency caused by phe93cys mutation in the CYP17
gene. J. Clin.

[121521]Endocr. Metab. 87: 898-905, 2002.

[121522]21511.

[121523]Fan, Y.-S.; Sasi, R.; Lee, C.; Winter, J. S. D.; Waterman, M.

[121524]R.; Lin, C. C.: Localization of the human CYP17 gene
(cytochrome

[121525]p450-17- α) to 10q24.3 by fluorescence in situ hybridization and

[121526]simultaneous chromosome banding. Genomics 14:
1110-1111, 1992.

[121527]21512.

[121528]Geller, D. H.; Auchus, R. J.; Mendonca, B. B.; Miller, W. L.:

[121529]The genetic and functional basis of isolated 17,20-lyase
deficiency. Nature

[121530]Genet. 17: 201-205, 1997.

[121531]21513.

[121532]Goldsmith, O.; Solomon, D. H.; Horton, R.: Hypogonadism
and mineralocorticoid

[121533]excess: 17-hydroxylase deficiency. New Eng. J. Med. 277:
673-677,

[121534]1967.

[121535]21514.

[121536]Gupta, M. K.; Geller, D. H.; Auchus, R. J.: Pitfalls in charac-
terizing

[121537]P450c17 mutations associated with isolated 17,20-lyase
deficiency. J.

[121538]Clin. Endocr. Metab. 86: 4416-4423, 2001.

[121539]21515.

[121540]Hanley, N. A.; Rainey, W. E.; Wilson, D. I.; Ball, S. G.;
Parker,

[121541]K. L.: Expression profiles of SF-1, DAX1, and CYP17 in the
human

[121542]fetal adrenal gland: potential interactions in gene regula-
tion. Molec.

[121543]Endocr. 15: 57-68, 2001.

[121544]21516.

[121545]Heremans, G. F. P.; Moolenaar, A. J.; Van Gelderen, H. H.:

Female

[121546]phenotype in a male child due to 17-alpha-hydroxylase
deficiency. Arch.

[121547]Dis. Child. 51: 721-723, 1976.

[121548]21517.

[121549]Imai, T.; Yanase, T.; Waterman, M. R.; Simpson, E. R.;
Pratt,

[121550]J. J.: Canadian Mennonites and individuals residing in the
Friesland

[121551]region of the Netherlands share the same molecular basis
of 17-alpha-hydroxylase

[121552]deficiency. Hum. Genet. 89: 95-96, 1992.

[121553]21518.

[121554]Jones, K. L.; Freidenberg, G. R.; Buchta, R.; Derenoncourt,
A.

[121555]: Male pseudohermaphroditism resulting from
17-alpha-monooxygenase

[121556](P-450C-17) deficiency in two unrelated Guamanians. Am.
J. Dis. Child. 146:

[121557]592-595, 1992.

[121558]21519.

[121559]Kagimoto, K.; Waterman, M. R.; Kagimoto, M.; Ferreira, P.;
Simpson,

[121560]E. R.; Winter, J. S. D.: Identification of a common molecular
basis

[121561]for combined 17-alpha-hydroxylase/17,20-lyase defi-
ciency in two Mennonite

[121562]families. Hum. Genet. 82: 285-286, 1989.

[121563]21520.

[121564]Kagimoto, M.; Winter, J. S. D.; Kagimoto, K.; Simpson, E.
R.;

[121565]Waterman, M. R.: Structural characterization of normal
and mutant

[121566]human steroid 17-alpha-hydroxylase genes: molecular
basis of one example

[121567]of combined 17-alpha-hydroxylase/17,20 lyase defi-
ciency. Molec. Endocr. 2:

[121568]564-570, 1988.

[121569]21521.

[121570]Krohn, K.; Uiho, R.; Aavik, E.; Peterson, P.; Savilahti, K.:

[121571]Identification of molecular cloning of an autoantigen as-

sociated with

[121572]Addison's disease as steroid 17-alpha-hydroxylase.

Lancet 339: 770-773,

[121573]1992.

[121574]21522.

[121575]Laflamme, N.; Leblanc, J.-F.; Mailloux, J.; Faure, N.; Labrie,

[121576]F.; Simard, J.: Mutation R96W in cytochrome P450c17 gene

causes combined

[121577]17-alpha-hydroxylase/17-20-lyase deficiency in two

French Canadian

[121578]patients. J. Clin. Endocr. Metab. 81: 264-268, 1996.

[121579]21523.

[121580]Lin, C. J.; Martens, J. W. M.; Miller, W. L.: NF-1C, Sp1, and

[121581]Sp3 are essential for transcription of the human gene for

P450c17

[121582](steroid 17-alpha-hydroxylase/17,20 lyase) in human

adrenal NCI-H295A

[121583]cells. Molec. Endocr. 15: 1277-1293, 2001.

[121584]21524.

[121585]Lin, D.; Harikrishna, J. A.; Moore, C. C. D.; Jones, K. L.;

Miller,

[121586]W. L.: Missense mutation serine106-to-proline causes
17-alpha-hydroxylase

[121587]deficiency. J. Biol. Chem. 266: 15992-15998, 1991.

[121588]21525.

[121589]Mallin, S. R.: Congenital adrenal hyperplasia secondary to
17-hydroxylase

[121590]deficiency. Ann. Intern. Med. 70: 69-76, 1969.

[121591]21526.

[121592]Mantero, F.; Scaroni, C.; Pasini, C. V.; Fagiolo, U.: No link-
age

[121593]between HLA and congenital adrenal hyperplasia due to
17-alpha-hydroxylase

[121594]deficiency. (Letter) New Eng. J. Med. 303: 530, 1980.

[121595]21527.

[121596]Matteson, K. J.; Picado-Leonard, J.; Chung, B.-C.; Mohan-
das, T.

[121597]K.; Miller, W. L.: Assignment of the gene for adrenal
P450C17 (steroid

[121598]17-alpha-hydroxylase/17,20 lyase) to human chromo-
some 10. J. Clin.

[121599]Endocr. Metab. 63: 789–791, 1986.

[121600]21528.

[121601]Miller, W. L.; Chung, B.-C.; Matteson, K. J.; Voutilainen, R.;

[121602]Picado–Leonard, J.: Molecular biology of steroid hormone synthesis.

[121603](Abstract) DNA 5: 61, 1986.

[121604]21529.

[121605]Donlon, T. A.; Krensky, A. M.; Clayberger, C.: Localization of

[121606]the human T lymphocyte activation gene 519 (D2S69E) to chromosome

[121607]2p12–q11. Cytogenet. Cell Genet. 53: 230–231, 1990.

[121608]21530.

[121609]Jongstra, J.; Schall, T. J.; Dyer, B. J.; Clayberger, C.; Jorgensen,

[121610]J.; Davis, M. M.; Krensky, A. M.: The isolation and sequence of a

[121611]novel gene from a human functional T cell line. J. Exp. Med. 165:

[121612]601–614, 1987.

[121613]21531.

[121614]Kaufmann, S. H. E.: CD8+ T lymphocytes in intracellular
microbial

[121615]infections. Immun. Today 9: 168–174, 1988.

[121616]21532.

[121617]Pena, S. V.; Hanson, D. A.; Carr, B. A.; Goralski, T. J.; Kren-
sky,

[121618]A. M.: Processing, subcellular localization, and function of
519

[121619](granulysin), a human late T cell activation molecule with
homology

[121620]to small, lytic granule proteins. J. Immun. 158:
2680–2688, 1997.

[121621]21533.

[121622]Stenger, S.; Hanson, D. A.; Teitelbaum, R.; Dewan, P.; Ni-
azi, K.

[121623]R.; Froelich, C. J.; Ganz, T.; Thoma–Uszynski, S.; Melian,
A.; Bogdan,

[121624]C.; Porcelli, S. A.; Bloom, B. R.; Krensky, A. M.; Modlin, R.
L.:

[121625]An antimicrobial activity of cytolytic T cells mediated by
granulysin. Science 282:

[121626]121–125, 1998.

[121627]21534.

[121628]Stenger, S.; Mazzaccaro, R. J.; Uyemura, K.; Cho, S.;
Barnes, P.

[121629]F.; Rosat, J. P.; Sette, A.; Brenner, M. B.; Porcelli, S. A.;
Bloom,

[121630]B. R.; Modlin, R. L.: Differential effects of cytolytic T cell
subsets

[121631]on intracellular infection. Science 276: 1684–1687, 1997.

[121632]21535.

[121633]Wong–Staal, F.; Dalla–Favera, R.; Franchini, G.; Gelmann,
E. P.;

[121634]Gallo, R. C.: Three distinct genes in human DNA related to
the transforming

[121635]genes of mammalian sarcoma retroviruses. Science 213:
226–228, 1981.

[121636]21536.

[121637]Aurias, A.; Rimbaut, C.; Buffe, D.; Dubousset, J.; Maz–
abraud, A.

[121638]: Chromosomal translocations in Ewing's sarcoma. (Letter)
New Eng.

[121639]J. Med. 309: 496–497, 1983.

[121640]21537.

[121641]Bartram, C. R.; de Klein, A.; Hagemeijer, A.; Grosveld, G.;
Heisterkamp,

[121642]N.; Groffen, J.: Localization of the human c-sis oncogene
in Ph-1-positive

[121643]and Ph-1-negative chronic myelocytic leukemia by in situ
hybridization. Blood 63:

[121644]223–225, 1984.

[121645]21538.

[121646]Bechet, J.-M.; Bornkamm, G.; Freese, U.-K.; Lenoir, G. M.:
The

[121647]c-sis oncogene is not activated in Ewing's sarcoma.
(Letter) New

[121648]Eng. J. Med. 310: 393 only, 1984.

[121649]21539.

[121650]Bishop, J. M.: Enemies within: the genesis of retrovirus
oncogenes. Cell 23:

[121651]5–6, 1981.

[121652]21540.

- [121653]Bolger, G. B.; Stamberg, J.; Kirsch, I. R.; Hollis, G. F.;
Schwarz,
[121654]D. F.; Thomas, G. H.: Chromosomal translocation t(14;22)
and oncogene
[121655](c-sis) variant in a pedigree with familial meningioma.
New Eng.
[121656]J. Med. 312: 564-567, 1985.
[121657]21541.
[121658]Cohen, J. B.; Levinson, A. D.: A point mutation in the last
intron
[121659]responsible for increased expression and transforming
activity of
[121660]the c-Ha-ras oncogene. Nature 334: 119-124, 1988.
[121661]21542.
[121662]Collins, T.; Ginsburg, D.; Boss, J. M.; Orkin, S. H.; Pober, J.
[121663]S.: Cultured human endothelial cells express platelet-de-
rived growth
[121664]factor B chain: cDNA cloning and structural analysis. Na-
ture 316:
[121665]748-750, 1985.
[121666]21543.

[121667]Dalla-Favera, R.; Gallo, R. C.; Giallongo, A.; Croce, C.:
Chromosomal

[121668]localization of the human homolog (c-sis) of the simian
sarcoma virus

[121669]onc gene. Science 218: 686-688, 1982.

[121670]21544.

[121671]Dalla-Favera, R.; Gelmann, E. P.; Gallo, R. C.; Wong-Staal,
F.

[121672]: A human onc gene homologous to the transforming
gene (v-sis) of

[121673]simian sarcoma virus. Nature 292: 31-35, 1981.

[121674]21545.

[121675]Deuel, T. F.; Huang, J. S.; Huang, S. S.; Stroobant, P.; Wa-
terfield,

[121676]M. D.: Expression of a platelet-derived growth factor-like
protein

[121677]in simian sarcoma virus transformed cells. Science 221:
1348-1350,

[121678]1983.

[121679]21546.

[121680]Devare, S. G.; Reddy, E. P.; Law, J. D.; Robbins, K. C.;

Aaronson,

[121681]S. A.: Nucleotide sequence of the simian sarcoma virus genome: demonstration

[121682]that its acquired cellular sequences encode the transforming gene

[121683]product p28-sis. Proc. Nat. Acad. Sci. 80: 731-735, 1983.

[121684]21547.

[121685]Charlet-B., N.; Logan, P.; Singh, G.; Cooper, T. A.: Dynamic antagonism

[121686]between ETR-3 and PTB regulates cell type-specific alternative splicing. Molec.

[121687]Cell 9: 649-658, 2002.

[121688]21548.

[121689]Borglum, A. D.; Flint, T.; Tommerup, N.; Fleckner, J.; Justesen,

[121690]J.; Kruse, T. A.: Assignment of the human tryptophanyl-tRNA synthetase

[121691]gene (WARS) to chromosome 14q32.2-q32.32. Cytogenet. Cell Genet. 73:

[121692]99-103, 1996.

[121693]21549.

[121694]Denney, R. M.; Borgaonkar, D.; Ruddle, F. H.: Order of
genes for

[121695]NP and TRPRS on chromosome 14. Cytogenet. Cell Genet.
22: 493-497,

[121696]1978.

[121697]21550.

[121698]Francke, U.; Denney, R. M.; Ruddle, F. H.: Intrachromoso-
mal gene

[121699]mapping in man: the gene for tryptophanyl-tRNA syn-
thetase maps in

[121700]region q21-qter of chromosome 14. Somat. Cell Genet. 3:
381-389,

[121701]1977.

[121702]21551.

[121703]Sack, G. H.; Huether, C. A.; Garg, N.: Familial spastic para-
plegia--clinical

[121704]and pathologic studies in a large kindred. Johns Hopkins
Med. J. 143:

[121705]117-121, 1978.

[121706]21552.

[121707]Schady, W.; Dick, J. P. R.; Sheard, A.; Crampton, S.: Central

[121708]motor conduction studies in hereditary spastic paraplegia.

J. Neurol.

[121709]Neurosurg. Psychiat. 54: 775–779, 1991.

[121710]21553.

[121711]Schady, W.; Sheard, A.: A quantitative study of sensory
function

[121712]in hereditary spastic paraplegia. Brain 113: 709–720,
1990.

[121713]21554.

[121714]Schady, W.; Smith, C. M. L.: Sensory neuropathy in heredi-
tary

[121715]spastic paraplegia. J. Neurol. Neurosurg. Psychiat. 57:
693–698,

[121716]1994.

[121717]21555.

[121718]Scheltens, P.; Bruyn, R. P. M.; Hazenberg, G. J.: A Dutch
family

[121719]with autosomal dominant pure spastic paraparesis
(Strumpell's disease). Acta

[121720]Neurol. Scand. 82: 169–173, 1990.

[121721]21556.

- [121722]Schwarz, G. A.; Liu, C. N.: Hereditary (familial) spastic paraplegia:
[121723]further clinical and pathologic observations. Arch. Neurol. Psychiat. 75:
[121724]144–162, 1956.
[121725]21557.
- [121726]Skre, H.: Hereditary spastic paraplegia in Western Norway. Clin. Genet. 6: 165–183, 1974.
[121728]21558.
- [121729]Strumpell, A.: Die primaere Seitenstrangsklerose (spastische Spinalparalyse). Dtsch. Z. Nervenheilk. 27: 291–339, 1904.
[121731]21559.
- [121732]Thurmon, T. F.; He, C.; Haskell, C.; Thorpe, P.; Thurmon, S. G.;
[121733]Rosen, D. R.: Genetic anticipation in a large family with pure autosomal
[121734]dominant hereditary spastic paraplegia. Am. J. Med. Genet. 83: 392–396,

[121735]1999.

[121736]21560.

[121737]Thurmon, T. F.; Walker, B. A.: Two distinct types of autosomal

[121738]dominant spastic paraplegia. Birth Defects Orig. Art. Ser. VII(1):

[121739]216–218, 1971.

[121740]21561.

[121741]Van Bogaert, L.: Etude genetique sur les paraplegies spasmotiques

[121742]familiales. J. Genet. Hum. 1: 6–23, 1952.

[121743]21562.

[121744]Zhao, X.; Alvarado, D.; Rainier, S.; Lemons, R.; Hedera, P.; Weber,

[121745]C. H.; Tükel, T.; Apak, M.; Heiman-Patterson, T.; Ming, L.; Bui, M.;

[121746]Fink, J. K.: Mutations in a newly identified GTPase gene cause autosomal

[121747]dominant hereditary spastic paraplegia. Nature Genet. 29: 326–331,

[121748]2001.

[121749]21563.

[121750]Benson, K. F.; Horwitz, M.; Wolff, J.; Friend, K.; Thompson, E.;

[121751]White, S.; Richards, R. I.; Raskind, W. H.; Bird, T. D.: CAG repeat

[121752]expansion in autosomal dominant familial spastic paraparesis: novel

[121753]expansion in a subset of patients. Hum. Molec. Genet. 7: 1779–1786,

[121754]1998.

[121755]21564.

[121756]Fonknechten, N.; Mavel, D.; Byrne, P.; Davoine, C.–S.; Cruaud,

[121757]C.; Boentsch, D.; Samson, D.; Coutinho, P.; Hutchinson, M.; McMonagle,

[121758]P.; Burgunder, J.–M.; Tartaglione, A.; and 10 others: Spectrum

[121759]of SPG4 mutations in autosomal dominant spastic paraplegia. Hum.

[121760]Molec. Genet. 9: 637–644, 2000.

[121761]21565.

- [121762]Nielsen, J. E.; Koefoed, P.; Abell, K.; Hasholt, L.; Eiberg, H.;
- [121763]Fenger, K.; Niebuhr, E.; Sorensen, S. A.: CAG repeat expansion in
- [121764]autosomal dominant pure spastic paraplegia linked to chromosome 2p21-p24. Hum.
- [121765]Molec. Genet. 6: 1811-1816, 1997.
- [121766]21566.
- [121767]Svenson, I. K.; Ashley-Koch, A. E.; Gaskell, P. C.; Riney, T.
- [121768]J.; Cumming, W. J. K.; Kingston, H. M.; Hogan, E. L.; Boustany, R.-M.
- [121769]N.; Vance, J. M.; Nance, M. A.; Pericak-Vance, M. A.; Marchuk, D.
- [121770]A.: Identification and expression analysis of spastin gene mutations
- [121771]in hereditary spastic paraplegia. Am. J. Hum. Genet. 68: 1077-1085,
- [121772]2001.
- [121773]21567.
- [121774]Ala-Kapee, M.; Nevanlinna, H.; Mali, M.; Jalkanen, M.; Schroder,
- [121775]J.: Localization of gene for human syndecan, an integral

membrane

[121776]proteoglycan and a matrix receptor, to chromosome 2.

Somat. Cell

[121777]Molec. Genet. 16: 501–505, 1990.

[121778]21568.

[121779]Alexander, C. M.; Reichsman, F.; Hinkes, M. T.; Lincecum, J.; Becker,

[121780]K. A.; Cumberledge, S.; Bernfield, M.: Syndecan–1 is required for

[121781]Wnt–1–induced mammary tumorigenesis in mice. Nature Genet. 25: 329–332,

[121782]2000.

[121783]21569.

[121784]Mali, M.; Jaakkola, P.; Arvilommi, A.–M.; Jalkanen, M.: Sequence

[121785]of human syndecan indicates a novel gene family of integral membrane

[121786]proteoglycans. J. Biol. Chem. 265: 6884–6889, 1990.

[121787]21570.

[121788]Oettinger, H. F.; Streeter, H.; Lose, E.; Copeland, N. G.; Gilbert,

[121789]D. J.; Justice, M. J.; Jenkins, N. A.; Mohandas, T.; Bernfield, M.

[121790]: Chromosome mapping of the murine syndecan gene. Genomics 11: 334–338,

[121791]1991.

[121792]21571.

[121793]Reizes, O.; Lincecum, J.; Wang, Z.; Goldberger, O.; Huang, L.;

[121794]Kaksonen, M.; Ahima, R.; Hinkes, M. T.; Barsh, G. S.; Rauvala, H.;

[121795]Bernfield, M.: Transgenic expression of syndecan–1 uncovers a physiological

[121796]control of feeding behavior by syndecan–3. Cell 106: 105–116, 2001.

[121797]21572.

[121798]Sanderson, R. D.; Lalor, P.; Bernfield, M.: B lymphocytes express

[121799]and lose syndecan at specific stages of differentiation. Cell Regulation 1:

[121800]27–35, 1989.

[121801]21573.

- [121802] Westman, P.; Hsieh, C.-L.; Mali, M.; Jalkanen, M.; Francke, U.;
- [121803] Schroder, J.: Assignment of the human syndecan (SDC) gene to short arm of chromosome 2. (Abstract) Cytogenet. Cell Genet. 58: 1873-1874, 1991.
- [121805] 1991.
- [121806] 21574.
- [121807] Asch, A. S.; Barnwell, J.; Silverstein, R. L.; Nachman, R. L.: Isolation of the thrombospondin membrane receptor. J. Clin. Invest. 79: 1054-1061, 1987.
- [121809] 1054-1061, 1987.
- [121810] 21575.
- [121811] de Fraipont, F.; El Atifi, M.; Gicquel, C.; Bertagna, X.; Chambaz, E. M.; Feige, J. J.: Expression of the angiogenesis markers vascular endothelial growth factor-A, thrombospondin-1, and platelet-derived endothelial cell growth factor in human sporadic adrenocortical tumors:

[121815]correlation with genotypic alterations. J. Clin. Endocr.
Metab. 85:

[121816]4734–4741, 2000.

[121817]21576.

[121818]Dixit, V. M.; Hennessy, S. W.; Grant, G. A.; Rotwein, P.;
Frazier,

[121819]W. A.: Characterization of a cDNA encoding the heparin
and collagen

[121820]binding domains of human thrombospondin. Proc. Nat.
Acad. Sci. 83:

[121821]5449–5453, 1986.

[121822]21577.

[121823]Frazier, W. A.: Thrombospondin: a modular adhesive gly-
coprotein

[121824]of platelets and nucleated cells. J. Cell Biol. 105: 625–632,
1987.

[121825]21578.

[121826]Jaffe, E.; Bornstein, P.; Disteché, C. M.: Mapping of the
thrombospondin

[121827]gene to human chromosome 15 and mouse chromosome
2 by in situ hybridization. Genomics 7:

[121828]123–126, 1990.

[121829]21579.

[121830]Lawler, J.; Sunday, M.; Thibert, V.; Duquette, M.; George, E. L.;

[121831]Rayburn, H.; Hynes, R. O.: Thrombospondin–1 is required for normal

[121832]murine pulmonary homeostasis and its absence causes pneumonia. J.

[121833]Clin. Invest. 101: 982–992, 1998.

[121834]21580.

[121835]Rodriguez–Manzaneque, J. C.; Lane, T. F.; Ortega, M. A.; Hynes,

[121836]R. O.; Lawler, J.; Iruela–Arispe, M. L.: Thrombospondin–1 suppresses

[121837]spontaneous tumor growth and inhibits activation of matrix metalloproteinase–9

[121838]and mobilization of vascular endothelial growth factor. Proc. Nat.

[121839]Acad. Sci. 98: 12485–12490, 2001.

[121840]21581.

[121841]Wolf, F. W.; Eddy, R. L.; Shows, T. B.; Dixit, V. M.: Structure

[121842]and chromosomal localization of the human thrombospondin gene. Genomics 6:

[121843]685–691, 1990.

[121844]21582.

[121845]Imai, K.; Sarker, A. H.; Akiyama, K.; Ikeda, S.; Yao, M.; Tsutsui,

[121846]K.; Shohmori, T.; Seki, S.: Genomic structure and sequence of a human

[121847]homologue (NTHL1/NTH1) of Escherichia coli endonuclease III with those

[121848]of the adjacent parts of TSC2 and SLC9A3R2 genes. Gene 222: 287–295,

[121849]1998.

[121850]21583.

[121851]Mira, J.-P.; Cariou, A.; Grall, F.; Delclaux, C.; Losser, M.-R.;

[121852]Heshmati, F.; Cheval, C.; Monchi, M.; Teboul, J.-L.; Riche, F.; Leleu,

[121853]G.; Arbibe, L.; Mignon, A.; Delpech, M.; Dhainaut, J.-F.: Association

[121854]of TNF2, a TNF- α promoter polymorphism, with septic shock susceptibility

[121855]and mortality: a multicenter study. J.A.M.A. 282: 561–568, 1999.

[121856]21584.

[121857]Moffatt, M. F.; Cookson, W. O. C. M.: Tumour necrosis factor

[121858]haplotypes and asthma. Hum. Molec. Genet. 6: 551–554, 1997.

[121859]21585.

[121860]Moraes, M. O.; Duppre, N. C.; Suffys, P. N.; Santos, A. R.; Almeida,

[121861]A. S.; Nery, J. A. C.; Sampaio, E. P.; Sarno, E. N.: Tumor necrosis

[121862]factor–alpha promoter polymorphism TNF2 is associated with a stronger

[121863]delayed–type hypersensitivity reaction in the skin of borderline tuberculoid

[121864]leprosy patients. Immunogenetics 53: 45–47, 2001.

[121865]21586.

[121866]Mulcahy, B.; Waldron–Lynch, F.; McDermott, M. F.; Adams, C.; Amos,

[121867]C. I.; Zhu, D. K.; Ward, R. H.; Clegg, D. O.; Shanahan, F.; Molloy,

[121868]M. G.; O'Gara, F.: Genetic variability in the tumor necrosis factor-lymphotoxin

[121869]region influences susceptibility to rheumatoid arthritis.
Am. J.

[121870]Hum. Genet. 59: 676-683, 1996.

[121871]21587.

[121872]Muller, U.; Jongeneel, C. V.; Nedospasov, S. A.; Lindahl, K.
F.;

[121873]Steinmetz, M.: Tumour necrosis factor and lymphotoxin
genes map close

[121874]to H-2D in the mouse major histocompatibility complex.
Nature 325:

[121875]265-267, 1987.

[121876]21588.

[121877]Nadel, S.; Newport, M. J.; Booy, R.; Levin, M.: Variation in

[121878]the tumor necrosis factor-alpha gene promoter region
may be associated

[121879]with death from meningococcal disease. J. Infect. Dis.
174: 878-880,

[121880]1996.

[121881]21589.

[121882]Nedospasov, S. A.; Hirt, B.; Shakhov, A. N.; Dobrynin, V.
N.;

[121883]Kawashima, E.; Accolla, R. S.; Jongeneel, C. V.: The genes
for tumor

[121884]necrosis factor (TNR-alpha) and lymphotoxin (TNR-beta)
are tandemly

[121885]arranged on chromosome 17 of the mouse. Nucleic Acids
Res. 14: 7713-7725,

[121886]1986.

[121887]21590.

[121888]Nedwin, G. E.; Naylor, S. L.; Sakaguchi, A. Y.; Smith, D.;
Jarrett-Nedwin,

[121889]J.; Pennica, D.; Goeddel, D. V.; Gray, P. W.: Human lym-
photoxin and

[121890]tumor necrosis factor genes: structure, homology and
chromosomal localization. Nucleic

[121891]Acids Res. 13: 6361-6373, 1985.

[121892]21591.

[121893]Norman, R. A.; Bogardus, C.; Ravussin, E.: Linkage be-
tween obesity

[121894]and a marker near the tumor necrosis factor-alpha locus
in Pima Indians. J.

[121895]Clin. Invest. 96: 158–162, 1995.

[121896]21592.

[121897]Obayashi, H.; Hasegawa, G.; Fukui, M.; Kamiuchi, K.; Kitamura,

[121898]A.; Ogata, M.; Kanaitzuka, T.; Shigeta, H.; Kitagawa, Y.; Nakano,

[121899]K.; Nishimura, M.; Ohta, M.; Nakamura, N.: Tumor necrosis factor

[121900]microsatellite polymorphism influences the development of insulin

[121901]dependency in adult-onset diabetes patients with the DRB1*1502–DQB1*0601

[121902]allele and anti-glutamic acid decarboxylase antibodies. J. Clin.

[121903]Endocr. Metab. 85: 3348–3351, 2000.

[121904]21593.

[121905]Old, L. J.: Tumor necrosis factor (TNF). Science 230: 630–632,

[121906]1985.

[121907]21594.

[121908]Pennica, D.; Nedwin, G. E.; Hayflick, J. S.; Seeburg, P. H.;

Derynck,

[121909]R.; Palladino, M. A.; Kohr, W. J.; Aggarwal, B. B.; Goeddel, D. V.

[121910]; Human tumour necrosis factor: precursor structure, expression and

[121911]homology to lymphotoxin. Nature 312: 724-729, 1984.

[121912]21595.

[121913]Ragoussis, J.; Bloemer, K.; Weiss, E. H.; Ziegler, A.: Localization

[121914]of the genes for tumor necrosis factor and lymphotoxin between the

[121915]HLA class I and III regions by field inversion gel electrophoresis. Immunogenetics 27:

[121916]66-69, 1988.

[121917]21596.

[121918]Rasmussen, S. K.; Urhammer, S. A.; Jensen, J. N.; Hansen, T.;

[121919]Borch-Johnsen, K.; Pedersen, O.: The -238 and -308 G6A polymorphisms

[121920]of the tumor necrosis factor alpha gene promoter are not associated

[121921]with features of the insulin resistance syndrome or altered

birth

[121922]weight in Danish Caucasians. J. Clin. Endocr. Metab. 85:
1731–1734,

[121923]2000.

[121924]21597.

[121925]Rosmond, R.; Chagnon, M.; Bouchard, C.; Bjorntorp, P.: G–
308A

[121926]polymorphism of the tumor necrosis factor alpha gene
promoter and

[121927]salivary cortisol secretion. J. Clin. Endocr. Metab. 86:
2178–2180,

[121928]2001.

[121929]21598.

[121930]Roy, S.; McGuire, W.; Mascie–Taylor, C. G.; Saha, B.; Hazra,
S.

[121931]K.; Hill, A. V.; Kwiatkowski, D.: Tumor necrosis factor pro-
moter

[121932]polymorphism and susceptibility to lepromatous leprosy.
J. Infect.

[121933]Dis. 176: 530–532, 1997.

[121934]21599.

[121935]Ruuls, S. R.; Sedgwick, J. D.: Unlinking tumor necrosis factor

[121936]biology from the major histocompatibility complex: lessons from human

[121937]genetics and animal models. Am. J. Hum. Genet. 65: 294-301, 1999.

[121938]21600.

[121939]Spies, T.; Morton, C. C.; Nedospasov, S. A.; Fiers, W.; Pious,

[121940]D.; Strominger, J. L.: Genes for the tumor necrosis factors alpha

[121941]and beta are linked to the human major histocompatibility complex. Proc.

[121942]Nat. Acad. Sci. 83: 8699-8702, 1986.

[121943]21601.

[121944]van Hensbroek, M. B.; Palmer, A.; Onyiorah, E.; Schneider, G.;

[121945]Jaffar, S.; Dolan, G.; Memming, H.; Frenkel, J.; Enwere, G.; Bennett,

[121946]S.; Kwiatkowski, D.; Greenwood, B.: The effect of a monoclonal antibody

[121947]to tumor necrosis factor on survival from childhood cere-

bral malaria. J.

[121948] Infect. Dis. 174: 1091–1097, 1996.

[121949] 21602.

[121950] Sashio, H.; Tamura, K.; Ito, R.; Yamamoto, Y.; Bamba, H.;
Kosaka,

[121951] T.; Fukui, S.; Sawada, K.; Fukuda, Y.; Tamura, K.; Satomi,
M.; Shimoyama,

[121952] T.; Furuyama, J.: Polymorphisms of the TNF gene and the
TNF receptor

[121953] superfamily member 1B gene are associated with suscep-
tibility to ulcerative

[121954] colitis and Crohn's disease, respectively. Immunogenetics
53: 1020–1027,

[121955] 2002.

[121956] 21603.

[121957] Van Ostade, X.; Vandenabeele, P.; Everaerd, B.; Loetscher,
H.;

[121958] Gentz, R.; Brockhaus, M.; Lesslauer, W.; Tavernier, J.;
Brouckaert,

[121959] P.; Fiers, W.: Human TNF mutants with selective activity on
the p55

[121960] receptor. Nature 361: 266–269, 1993.

[121961]21604.

[121962]Wang, A. M.; Creasey, A. A.; Ladner, M. B.; Lin, L. S.;
Strickler,

[121963]J.; Van Arsdel, J. N.; Yamamoto, R.; Mark, D. F.: Molecular
cloning

[121964]of the complementary DNA for human tumor necrosis fac-
tor. Science 228:

[121965]149-154, 1985.

[121966]21605.

[121967]Wilson, A. G.; Symons, J. A.; McDowell, T. L.; et al: Effects

[121968]of a polymorphism in the human tumor necrosis factor al-
pha promoter

[121969]on transcriptional activation. Proc. Nat. Acad. Sci. 94:
3195-3199,

[121970]1997.

[121971]21606.

[121972]Winchester, E. C.; Millwood, I. Y.; Rand, L.; Penny, M. A.;
Kessling,

[121973]A. M.: Association of the TNF-alpha-308 (G-A) polymor-
phism with self-reported

[121974]history of childhood asthma. Hum. Genet. 107: 591-596,

2000.

[121975]21607.

[121976]Witte, J. S.; Palmer, L. J.; O'Connor, R. D.; Hopkins, P. J.;

[121977]Hall, J. M.: Relation between tumour necrosis factor polymorphism

[121978]TNF- α -308 and risk of asthma. *Europ. J. Hum. Genet.* 10: 82-85,

[121979]2002.

[121980]21608.

[121981]Peter, D.; Finn, J. P.; Klisak, I.; Liu, Y.; Kojis, T.; Heinzmann,

[121982]C.; Roghani, A.; Sparkes, R. S.; Edwards, R. H.: Chromosomal localization

[121983]of the human vesicular amine transporter genes. *Genomics* 18: 720-723,

[121984]1993.

[121985]21609.

[121986]Roghani, A.; Welch, C.; Xia, Y.-R.; Liu, Y.; Peter, D.; Finn, J.

[121987]P.; Edwards, R. H.; Lusis, A. J.: Assignment of the mouse vesicular

[121988]monoamine transporter genes, *Slc18a1* and *Slc18a2*, to

chromosomes 8

[121989]and 19 by linkage analysis. Mammalian Genome 7:
393–394, 1996.

[121990]21610.

[121991]Lania, L.; Donti, E.; Pannuti, A.; Pascucci, A.; Pengue, G.;
Felicciello,

[121992]I.; La Mantia, G.; Lanfranccone, L.; Pelicci, P.-G.: cDNA iso-
lation,

[121993]expression analysis, and chromosomal localization of two
human zinc

[121994]finger genes. Genomics 6: 333–340, 1990.

[121995]21611.

[121996]Han, Z.-G.; Zhang, Q.-H.; Ye, M.; Kan, L.-X.; Gu, B.-W.;
He, K.-L.;

[121997]Shi, S.-L.; Zhou, J.; Fu, G.; Mao, M.; Chen, S.-J.; Yu, L.;
Chen,

[121998]Z.: Molecular cloning of six novel Kruppel-like zinc finger
genes

[121999]from hematopoietic cells and identification of a novel
transregulatory

[122000]domain KRN. J. Biol. Chem. 274: 35741–35748, 1999.

[122001]21612.

[122002]Rousseau-Merck, M. F.; Huebner, K.; Berger, R.; Thiesen, H. J.

[122003]: Chromosomal localization of two human zinc finger protein genes,

[122004]ZNF24 (KOX17) and ZNF29 (KOX26), to 18q12 and 17p13-p12, respectively. Genomics 9:

[122005]154-161, 1991.

[122006]21613.

[122007]Seite, P.; Huebner, K.; Rousseau-Merck, M. F.; Berger, R.; Thiesen,

[122008]H. J.: Two human genes encoding zinc finger proteins, ZNF12 (KOX3)

[122009]and ZNF26 (KOX20), map to chromosomes 7p22-p21 and 12q24.33, respectively. Hum.

[122010]Genet. 86: 585-590, 1991.

[122011]21614.

[122012]King, R. A.: Personal Communication. Minneapolis, Minn. 12/31/1992.

[122013]21615.

[122014]Corral, J.; Martin, C.; Perez, R.; Sanchez, I.; Mories, M. T.;

[122015]San Millan, J. L.; Miralles, J. M.; Gonzalez-Sarmiento, R.:

Thyroglobulin

[122016]gene point mutation associated with non-endemic simple goitre. Lancet 341:

[122017]462-464, 1993.

[122018]21616.

[122019]Zinman, B.; Hanley, A. J. G.; Harris, S. B.; Kwan, J.; Fantus,

[122020]J. G.: Circulating tumor necrosis factor-alpha concentrations in

[122021]a Native Canadian population with high rates of type 2 diabetes mellitus. J.

[122022]Clin. Endocr. Metab. 84: 272-278, 1999.

[122023]21617.

[122024]de Vijlder, J. J. M.; Baas, F.; Koch, C. A. M.; Kok, K.; Gons,

[122025]M. H.: Autosomal dominant inheritance of a thyroglobulin abnormality

[122026]suggests cooperation of subunits in hormone formation. Ann. Endocr. 44:

[122027]36 only, 1983.

[122028]21618.

[122029]Hishinuma, A.; Takamatsu, J.; Ohyama, Y.; Yokozawa, T.;

Kanno,

[122030]Y.; Kuma, K.; Yoshida, S.; Matsuura, N.; Ieiri, T.: Two novel
cysteine

[122031]substitutions (C1263R and C1995S) of thyroglobulin
cause a defect

[122032]in intracellular transport of thyroglobulin in patients with
congenital

[122033]goiter and the variant type of adenomatous goiter. J. Clin.
Endocr.

[122034]Metab. 84: 1438–1444, 1999.

[122035]21619.

[122036]Ieiri, T.; Cochaux, P.; Targovnik, H. M.; Suzuki, M.; Shi-
moda,

[122037]S.-I.; Perret, J.; Vassart, G.: A 3-prime splice site mutation
in

[122038]the thyroglobulin gene responsible for congenital goiter
with hypothyroidism. J.

[122039]Clin. Invest. 88: 1901–1905, 1991.

[122040]21620.

[122041]Kim, P. S.; Ding, M.; Menon, S.; Jung, C.-G.; Cheng, J.-M.;
Miyamoto,

[122042]T.; Li, B.; Furudate, S.; Agui, T.: A missense mutation

G2320R in

[122043]the thyroglobulin gene causes non-goitrous congenital
primary hypothyroidism

[122044]in the WIC-rdw rat. Molec. Endocr. 14: 1944-1953, 2000.

[122045]21621.

[122046]Kim, P. S.; Hossain, S. A.; Park, Y.-N.; Lee, I.; Yoo, S.-E.;

[122047]Arvan, P.: A single amino acid change in the acetyl-
cholinesterase-like

[122048]domain of thyroglobulin causes congenital goiter with hy-
pothyroidism

[122049]in the cog/cog mouse: a model of human endoplasmic
reticulum storage

[122050]disease. Proc. Nat. Acad. Sci. 95: 9909-9913, 1998.

[122051]21622.

[122052]Kok, K.; van Dijk, J. E.; Sterk, A.; Baas, F.; van Ommen,
G.-J.

[122053]B.; de Vijlder, J. J. M.: Autosomal recessive inheritance of
goiter

[122054]in Dutch goats. J. Hered. 78: 298-300, 1987.

[122055]21623.

[122056]Landegent, J. E.; Jansen in de Wal, N.; van Ommen, G.-J.

B.; Baas,

[122057]F.; de Vijlder, J. J. M.; van Duijn, P.; van der Ploeg, M.:

Chromosomal

[122058]localization of a unique gene by non-autoradiographic in situ hybridization. Nature 317:

[122059]175-177, 1985.

[122060]21624.

[122061]Lazar, V.; Bidart, J.-M.; Caillou, B.; Mahe, C.; Lacroix, L.;

[122062]Filetti, S.; Schlumberger, M.: Expression of the Na(+)/I(-) symporter

[122063]gene in human thyroid tumors: a comparison study with other thyroid-specific

[122064]genes. J. Clin. Endocr. Metab. 84: 3228-3234, 1999.

[122065]21625.

[122066]Malthiery, Y.; Lissitzky, S.: Primary structure of human thyroglobulin

[122067]deduced from the sequence of its 8448-base complementary DNA. Europ.

[122068]J. Biochem. 165: 491-498, 1987.

[122069]21626.

[122070]Medeiros-Neto, G.; Bunduki, V.; Tomimori, E.; Gomes, S.;

Knobel,

[122071]M.; Martin, R. T.; Zugaib, M.: Prenatal diagnosis and treatment of

[122072]dyshormonogenetic fetal goiter due to defective thyroglobulin synthesis. J.

[122073]Clin. Endocr. Metab. 82: 4239–4242, 1997.

[122074]21627.

[122075]Ricketts, M. H.; Pohl, V.; de Martynoff, G.; Boyd, C. D.; Bester,

[122076]A. J.; Van Jaarsveld, P. P.; Vassart, G.: Defective splicing of thyroglobulin

[122077]gene transcripts in the congenital goitre of the Afrikaner cattle. EMBO

[122078]J. 4: 731–737, 1985.

[122079]21628.

[122080]Ricketts, M. H.; Simons, M. J.; Parma, J.; Mercken, L.; Dong,

[122081]Q.; Vassart, G.: A nonsense mutation causes hereditary goitre in

[122082]the Afrikaner cattle and unmasks alternative splicing of thyroglobulin

[122083]transcripts. Proc. Nat. Acad. Sci. 84: 3181–3184, 1987.

[122084]21629.

[122085]Savagner, F.; Rodien, P.; Reynier, P.; Rohmer, V.; Bigorgne, J.-C.;

[122086]Malthiery, Y.: Analysis of Tg transcripts by real-time RT-PCR in

[122087]the blood of thyroid cancer patients. J. Clin. Endocr. Metab. 87:

[122088]635–639, 2002.

[122089]21630.

[122090]Targovnik, H.; Propato, F.; Varela, V.; Wajchenberg, B.; Knobel,

[122091]M.; D'Abronzio, H. F.; Medeiros-Neto, G.: Low levels of thyroglobulin

[122092]messenger ribonucleic acid in congenital goitrous hypothyroidism with

[122093]defective thyroglobulin synthesis. J. Clin. Endocr. Metab. 69: 1137–1147,

[122094]1989.

[122095]21631.

[122096]Targovnik, H. M.; Frechtel, G. D.; Mendive, F. M.; Vono, J.; Cochaux,

- [122097]P.; Vassart, G.; Medeiros-Neto, G.: Evidence for the segregation
of three different mutated alleles of the thyroglobulin
gene in a
Brazilian family with congenital goiter and hypothyroidism. *Thyroid* 8:
291-297, 1998.
- [122100]21632.
- [122101]21632.
- [122102]Targovnik, H. M.; Medeiros-Neto, G.; Varela, V.; Cochaux, P.;
- [122103]Wajchenberg, B. L.; Vassart, G.: A nonsense mutation causes human
hereditary congenital goiter with preferential production
of a 171-nucleotide-deleted
thyroglobulin ribonucleic acid messenger. *J. Clin. Endocr. Metab.* 77:
210-215, 1993.
- [122106]21633.
- [122107]21633.
- [122108]Targovnik, H. M.; Vono, J.; Billerbeck, A. E. C.; Cerrone, G.
- [122109]E.; Varela, V.; Mendive, F.; Wajchenberg, B. L.; Medeiros-Neto, G.

[122110]: A 138-nucleotide deletion in the thyroglobulin ribonucleic acid

[122111]messenger in a congenital goiter with defective thyroglobulin synthesis. J.

[122112]Clin. Endocr. Metab. 80: 3356-3360, 1995.

[122113]21634.

[122114]Taylor, B. A.; Rowe, L.: The congenital goiter mutation is linked

[122115]to the thyroglobulin gene in the mouse. Proc. Nat. Acad. Sci. 84:

[122116]1986-1990, 1987.

[122117]21635.

[122118]van de Graaf, S. A. R.; Ris-Stalpers, C.; Veenboer, G. J. M.;

[122119]Cammenga, M.; Santos, C.; Targovnik, H. M.; de Vijlder, J. J. M.;

[122120]Medeiros-Neto, G.: A premature stopcodon in thyroglobulin messenger

[122121]RNA results in familial goiter moderate hypothyroidism. J. Clin.

[122122]Endocr. Metab. 84: 2537-2542, 1999.

[122123]21636.

[122124]Van Ommen, G.-J. B.; Baas, F.; Arnberg, A. C.; Pearson, P. L.;

[122125]De Vijlder, J. J. M.: Chromosome mapping and polymorphism study of

[122126]the human thyroglobulin gene. (Abstract) Cytogenet. Cell Genet. 37:

[122127]562 only, 1984.

[122128]21637.

[122129]Van Ommen, G. B.: Merging autosomal dominance and recessivity.

[122130](Letter) Am. J. Hum. Genet. 41: 689-691, 1987.

[122131]21638.

[122132]Yoshida, S.; Takamatsu, J.; Kuma, K.; Murakami, Y.; Sakane, S.;

[122133]Katayama, S.; Tarutani, O.; Ohsawa, N.: A variant of adenomatous

[122134]goiter with characteristic histology and possible hereditary thyroglobulin

[122135]abnormality. J. Clin. Endocr. Metab. 81: 1961-1966, 1996.

[122136]21639.

[122137]Spurr, N. K.; Goodfellow, P. N.; Sheer, D.; Bodmer, W. F.;

Vennstrom,

[122138]B.: Mapping of cellular oncogenes: ERBA1 is on chromosome 17. (Abstract) Cytogenet.

[122139]Cell Genet. 37: 591 only, 1984.

[122140]21640.

[122141]Thompson, C. C.; Weinberger, C.; Lebo, R.; Evans, R. M.: Identification

[122142]of a novel thyroid hormone receptor expressed in the mammalian central

[122143]nervous system. Science 237: 1610-1614, 1987.

[122144]21641.

[122145]Weinberger, C.; Thompson, C. C.; Ong, E. S.; Lebo, R.; Gruol,

[122146]D. J.; Evans, R. M.: The c-erb-A gene encodes a thyroid hormone receptor. Nature 324:

[122147]641-646, 1986.

[122148]21642.

[122149]Zabel, B. U.; Fournier, R. E. K.; Lalley, P. A.; Naylor, S. L.;

[122150]Sakaguchi, A. Y.: Cellular homologs of the avian erythroblastosis

[122151]virus erb-A and erb-B genes are syntenic in mouse but

asynthetic in

[122152]man. Proc. Nat. Acad. Sci. 81: 4874–4878, 1984.

[122153]21643.

[122154]Elchebly, M.; Payette, P.; Michaliszyn, E.; Cromlish, W.;
Collins,

[122155]S.; Loy, A. L.; Normandin, D.; Cheng, A.; Himms-Hagen, J.;
Chan, C.-C.;

[122156]Ramachandran, C.; Gresser, M. J.; Tremblay, M. L.;
Kennedy, B. P.

[122157]: Increased insulin sensitivity and obesity resistance in
mice lacking

[122158]the protein tyrosine phosphatase-1B gene. Science 283:
1544–1548,

[122159]1999.

[122160]21644.

[122161]Forsell, P. K. A. L.; Boie, Y.; Montalibet, J.; Collins, S.;
Kennedy,

[122162]B. P.: Genomic characterization of the human and mouse
protein tyrosine

[122163]phosphatase-1B genes. Gene 260: 145–153, 2000.

[122164]21645.

[122165]Gu, H. F.; Almgren, P.; Lindholm, E.; Frittitta, L.; Pizzuti, A.;

[122166]Trischitta, V.; Groop, L. C.: Association between the human glycoprotein

[122167]PC-1 gene and elevated glucose and insulin levels in paired-sibling

[122168]analysis. Diabetes 49: 1601-1603, 2000.

[122169]21646.

[122170]Haj, F. G.; Verveer, P. J.; Squire, A.; Neel, B. G.; Bastiaens,

[122171]P. I. H.: Imaging sites of receptor dephosphorylation by PTP1B on

[122172]the surface of the endoplasmic reticulum. Science 295: 1708-1711,

[122173]2002.

[122174]21647.

[122175]Jia, Z.; Barford, D.; Flint, A. J.; Tonks, N. K.: Structural

[122176]basis for phosphotyrosine peptide recognition by protein tyrosine

[122177]phosphatase 1B. Science 268: 1754-1758, 1995.

[122178]21648.

- [122179] Kennedy, B. P.; Ramachandran, C.: Protein tyrosine phosphatase-1B
[122180] in diabetes. *Biochem. Pharm.* 60: 877-883, 2000.
[122181] 21649.
- [122182] Mok, A.; Cao, H.; Zinman, B.; Hanley, A. J. G.; Harris, S. B.;
[122183] Kennedy, B. P.; Hegele, R. A.: A single nucleotide polymorphism in
[122184] protein tyrosine phosphatase PTP-1B is associated with
protection
[122185] from diabetes or impaired glucose tolerance in Oji-Cree. *J.*
Clin.
[122186] *Endocr. Metab.* 87: 724-727, 2002.
[122187] 21650.
- [122188] Tonks, N. K.; Diltz, C. D.; Fischer, E. H.: Purification of the
[122189] major protein-tyrosine-phosphatases of human placenta.
J. Biol. Chem. 263:
[122190] 6722-6730, 1988.
[122191] 21651.
- [122192] Barnea, G.; Silvennoinen, O.; Shaanan, B.; Honegger, A. M.;
Canoll,

[122193]P. D.; D'Eustachio, P.; Morse, B.; Levy, J. B.; Laforgia, S.; Huebner,

[122194]K.; Musacchio, J. M.; Sap, J.; Schlessinger, J.: Identification of

[122195]a carbonic anhydrase-like domain in the extracellular region of RPTP-gamma

[122196]defines a new subfamily of receptor tyrosine phosphatases. Molec.

[122197]Cell. Biol. 13: 1497-1506, 1993.

[122198]21652.

[122199]Kastury, K.; Ohta, M.; Lasota, J.; Moir, D.; Dorman, T.; LaForgia,

[122200]S.; Druck, T.; Huebner, K.: Structure of the human receptor tyrosine

[122201]phosphatase gamma gene (PTPRG) and relation to the familial RCC t(3;8)

[122202]chromosome translocation. Genomics 32: 225-235, 1996.

[122203]21653.

[122204]LaForgia, S.; Morse, B.; Levy, J.; Barnea, G.; Cannizzaro, L. A.;

[122205]Li, F.; Nowell, P. C.; Boghosian-Sell, L.; Glick, J.; Weston, A.;

[122206]Harris, C. C.; Drabkin, H.; Patterson, D.; Croce, C. M.; Schlessinger,
[122207]J.; Huebner, K.: Receptor protein-tyrosine phosphatase
gamma is a
[122208]candidate tumor suppressor gene at human chromosome
region 3p21. Proc.
[122209]Nat. Acad. Sci. 88: 5036-5040, 1991.
[122210]21654.
[122211]Latif, F.; Tory, K.; Modi, W.; Geil, L.; LaForgia, S.; Huebner,
[122212]K.; Zbar, B.; Lerman, M. I.: A MspI polymorphism and linkage
mapping
[122213]of the human protein-tyrosine phosphatase G (PTPRG)
gene. Hum. Molec.
[122214]Genet. 2: 91, 1993.
[122215]21655.
[122216]Cool, D. E.; Tonks, N. K.; Charbonneau, H.; Walsh, K. A.;
Fischer,
[122217]E. H.; Krebs, E. G.: cDNA isolated from a human T-cell library
encodes
[122218]a member of the protein-tyrosine-phosphatase family.
Proc. Nat. Acad.

[122219]Sci. 86: 5257–5261, 1989.

[122220]21656.

[122221]Johnson, C. V.; Cool, D. E.; Glaccum, M. B.; Green, N.; Fischer,

[122222]E. H.; Bruskin, A.; Hill, D. E.; Lawrence, J. B.: Isolation and mapping

[122223]of human T-cell protein tyrosine phosphatase sequences: localization

[122224]of genes and pseudogenes discriminated using fluorescence hybridization

[122225]with genomic versus cDNA probes. Genomics 16: 619–629, 1993.

[122226]21657.

[122227]Sakaguchi, A. Y.; Sylvia, V. L.; Martinez, L.; Lalley, P. A.; Shows,

[122228]T. B.; Han, E. S.; Smith, E. A.; Ghosh Choudhury, G.: Assignment

[122229]of tyrosine-specific T-cell phosphatase to conserved syntenic groups

[122230]on human chromosome 18 and mouse chromosome 18. (Abstract) Cytogenet.

[122231]Cell Genet. 58: 2014–2015, 1991.

[122232]21658.

[122233]Sakaguchi, A. Y.; Sylvia, V. L.; Martinez, L.; Smith, E. A.;
Han,

[122234]E. S.; Lalley, P. A.; Shows, T. B.; Ghosh Choudhury, G.: As-
signment

[122235]of tyrosine-specific T-cell phosphatase to conserved syn-
tenic groups

[122236]on human chromosome 18 and mouse chromosome 18.
Genomics 12: 151-154,

[122237]1992.

[122238]21659.

[122239]Adachi, M.; Sekiya, M.; Arimura, Y.; Takekawa, M.; Itoh, F.;
Hinoda,

[122240]Y.; Imai, K.; Yachi, A.: Protein-tyrosine phosphatase ex-
pression

[122241]in pre-B cell NALM-6. Cancer Res. 52: 737-740, 1992.

[122242]21660.

[122243]Adachi, M.; Sekiya, M.; Isobe, M.; Kumura, Y.; Ogita, Z.;
Hinoda,

[122244]Y.; Imai, K.; Yachi, A.: Molecular cloning and chromosomal
mapping

[122245]of a human protein-tyrosine phosphatase LC-PTP.

Biochem. Biophys.

[122246]Res. Commun. 186: 1607-1615, 1992.

[122247]21661.

[122248]Zanke, B.; Suzuki, H.; Kishihara, K.; Mizzen, L.; Minden, M.; Pawson,

[122249]A.; Mak, T. W.: Cloning and expression of an inducible lymphoid-specific,

[122250]protein tyrosine phosphatase (HePTPase). Europ. J. Immun. 22: 235-239,

[122251]1992.

[122252]21662.

[122253]Schols, L.; Szymanski, S.; Peters, S.; Przuntek, H.; Epplen, J.

[122254]T.; Hardt, C.; Riess, O.: Genetic background of apparently idiopathic

[122255]sporadic cerebellar ataxia. Hum. Genet. 107: 132-137, 2000.

[122256]21663.

[122257]Virgilio, L.; Isobe, M.; Narducci, M. G.; Carotenuto, P.; Camerini,

[122258]B.; Kurosawa, N.; ar-Rushdi, A.; Croce, C. M.; Russo, G.:
Chromosome

[122259]walking on the TCL1 locus involved in T-cell neoplasia.
Proc. Nat.

[122260]Acad. Sci. 90: 9275–9279, 1993.

[122261]21664.

[122262]Virgilio, L.; Lazzeri, C.; Bichi, R.; Nibu, K.; Narducci, M. G.;

[122263]Russo, G.; Rothstein, J. L.; Croce, C. M.: Deregulated ex-
pression

[122264]of TCL1 causes T cell leukemia in mice. Proc. Nat. Acad.
Sci. 95:

[122265]3885–3889, 1998.

[122266]21665.

[122267]Virgilio, L.; Narducci, M. G.; Isobe, M.; Billips, L. G.;
Cooper,

[122268]M. D.; Croce, C. M.; Russo, G.: Identification of the TCL1
gene involved

[122269]in T-cell malignancies. Proc. Nat. Acad. Sci. 91:
12530–12534, 1994.

[122270]21666.

[122271]Comp, P. C.; Nixon, R. R.; Cooper, M. R.; Esmon, C. T.: Fa-

milial

[122272]protein S deficiency is associated with recurrent thrombo-
sis. J.

[122273]Clin. Invest. 74: 2082-2088, 1984.

[122274]21667.

[122275]Cooper, D. N.: Personal Communication. London, England
9/17/1993.

[122276]21668.

[122277]D'Angelo, A.; Della Valle, P.; Crippa, L.; Pattarini, E.;
Grimaldi,

[122278]L. M. E.; D'Angelo, S. V.: Autoimmune protein S deficiency
in a boy

[122279]with severe thromboembolic disease. New Eng. J. Med.
328: 1753-1757,

[122280]1993.

[122281]21669.

[122282]Dahlback, B.: Protein S and C4b-binding protein: compo-
nents involved

[122283]in the regulation of the protein C anticoagulant system.
Thromb.

[122284]Haemost. 66: 49-61, 1991.

[122285]21670.

[122286]Dahlback, B.; Stenflo, J.: High molecular weight complex in human

[122287]plasma between vitamin K dependent protein S and complement component

[122288]C4b-binding protein. Proc. Nat. Acad. Sci. 78: 2512-2516, 1981.

[122289]21671.

[122290]Engesser, L.; Broekmans, A. W.; Briet, E.; Brommer, E. J. P.;

[122291]Bertina, R. M.: Hereditary protein S deficiency: clinical manifestations. Ann.

[122292]Intern. Med. 106: 677-682, 1987.

[122293]21672.

[122294]Espinosa-Parrilla, Y.; Morell, M.; Souto, J. C.; Tirado, I.; Fontcuberta,

[122295]J.; Estivill, X.; Sala, N.: Protein S gene analysis reveals the presence

[122296]of a cosegregating mutation in most pedigrees with type I
but not

[122297]type III PS deficiency. Hum. Mutat. 14: 30–39, 1999.

[122298]21673.

[122299]Fulton, A. B.; Hansen, R. M.; Petersen, R. A.; Vanderveen,
D.

[122300]K.: The rod photoreceptors in retinopathy of prematurity:
an electroretinographic

[122301]study. Arch. Ophthalm. 119: 499–505, 2001.

[122302]21674.

[122303]Girolami, A.; Simioni, P.; Lazzaro, A. R.; Pontara, E.; Ruzza,

[122304]G.: Heterozygous protein-S deficiency: a study of a large
kindred. Acta

[122305]Haemat. 84: 162–168, 1990.

[122306]21675.

[122307]Golub, B. M.; Sibony, P. A.; Coller, B. S.: Protein S defi-
ciency

[122308]associated with central retinal artery occlusion. Arch.
Ophthalm. 108:

[122309]918 only, 1990.

[122310]21676.

[122311]Griffin, J. H.; Gruber, A.; Fernandez, J. A.: Reevaluation of

[122312]total, free, and bound protein S and C4b-binding protein
levels in

[122313]plasma anticoagulated with citrate or hirudin. Blood 79:
3203-3211,

[122314]1992.

[122315]21677.

[122316]Hayashi, T.; Nishioka, J.; Shigekiyo, T.; Saito, S.; Suzuki, K.

[122317]: Protein S Tokushima: abnormal molecule with a substi-
tution of glu

[122318]for lys-155 in the second epidermal growth factor-like
domain of protein

[122319]S. Blood 83: 683-690, 1994.

[122320]21678.

[122321]Heeb, M. J.; Rosing, J.; Bakker, H. M.; Fernandez, J. A.;
Tans,

[122322]G.; Griffin, J. H.: Protein S binds to and inhibits factor Xa.
Proc.

[122323]Nat. Acad. Sci. 91: 2728-2732, 1994.

[122324]21679.

[122325]Hoskins, J.; Norman, D. K.; Beckmann, R. J.; Long, G. L.:
Cloning

[122326]and characterization of human liver cDNA encoding a pro-
tein S precursor. Proc.

[122327]Nat. Acad. Sci. 84: 349–353, 1987.

[122328]21680.

[122329]Kamiya, T.; Sugihara, T.; Ogata, K.; Saito, H.; Suzuki, K.;
Nishioka,

[122330]J.; Hashimoto, S.; Yamagata, K.: Inherited deficiency of
protein

[122331]S in a Japanese family with recurrent venous thrombosis: a
study of

[122332]three generations. Blood 67: 406–410, 1986.

[122333]21681.

[122334]Koller, H.; Stoll, G.; Sitzler, M.; Burk, M.; Schottler, B.; Fre-
und,

[122335]H-J.: Deficiency of both protein C and protein S in a family
with

[122336]ischemic strokes in young adults. Neurology 44:
1238–1240, 1994.

[122337]21682.

[122338]Lundwall, A.; Dackowski, W.; Cohen, E.; Shaffer, M.; Mahr, A.;

[122339]Dahlback, B.; Stenflo, J.; Wydro, R.: Isolation and sequence of the

[122340]cDNA for human protein S, a regulator of blood coagulation. Proc.

[122341]Nat. Acad. Sci. 83: 6716-6720, 1986.

[122342]21683.

[122343]Mahasandana, C.; Suvatte, V.; Chuansumrit, A.; Marlar, R. A.;

[122344]Manco-Johnson, M. J.; Jacobson, L. J.; Hathaway, W. E.: Homozygous

[122345]protein S deficiency in an infant with purpura fulminans. J. Pediat. 117:

[122346]750-753, 1990.

[122347]21684.

[122348]Maillard, C.; Berruyer, M.; Serre, C. M.; Dechavanne, M.; Delmas,

[122349]P. D.: Protein-S, a vitamin K-dependent protein, is a bone matrix

[122350]component synthesized and secreted by osteoblasts. Endocrinology 130:

[122351]1599–1604, 1992.

[122352]21685.

[122353]Makris, M.; Leach, M.; Beauchamp, N. J.; Daly, M. E.;
Cooper,

[122354]P. C.; Hampton, K. K.; Bayliss, P.; Peake, I. R.; Miller, G. J.;
Preston,

[122355]F. E.: Genetic analysis, phenotypic diagnosis, and risk of
venous

[122356]thrombosis in families with inherited deficiencies of protein S. Blood 95:

[122357]1935–1941, 2000.

[122358]21686.

[122359]Malnick, S. D. H.; Sthoeger, Z. M.: Autoimmune protein S
deficiency. New

[122360]Eng. J. Med. 329: 1898 only, 1993.

[122361]21687.

[122362]Mannucci, P. M.; Valsecchi, C.; Krachmalnicoff, A.; Faioni,
E.

[122363]M.; Tripodi, A.: Familial dysfunction of protein S. Thromb.

Haemost. 62:

[122364]763-766, 1989.

[122365]21688.

[122366]Mintz-Hittner, H. A.; Miyashiro, M. J.; Knight-Nanan, D.
M.; O'Malley,

[122367]R. E.; Marlar, R. A.: Vitreoretinal findings similar to
retinopathy

[122368]of prematurity in infants with compound heterozygous
protein S deficiency. Ophthalmology 106:

[122369]1525-1530, 1999.

[122370]21689.

[122371]Naylor, S. L.; Marshall, A.; Solomon, A.; Long, G.: Mapping
human

[122372]protein C and protein S to chromosomes 2 and 3, respec-
tively. (Abstract) Cytogenet.

[122373]Cell Genet. 46: 669 only, 1987.

[122374]21690.

[122375]Pan, E. Y.; Gomperts, E. D.; Millen, R.; Gilsanz, V.: Bone
mineral

[122376]density and its association with inherited protein S defi-
ciency. Thromb.

[122377]Res. 58: 221-231, 1990.

[122378]21691.

[122379]Phillips, W. G.; Marsden, J. R.; Hill, F. G.: Purpura fulminans

[122380]due to protein S deficiency following chickenpox. Brit. J. Derm. 127:

[122381]30-32, 1992.

[122382]21692.

[122383]Ploos van Amstel, H. K.; Huisman, M. V.; Reitsma, P. H.; ten Cate,

[122384]J.; Bertina, R. M.: Partial protein S gene deletion in a family with

[122385]hereditary thrombophilia. Blood 73: 479-483, 1989.

[122386]21693.

[122387]Ploos van Amstel, H. K.; Reitsma, P. H.; Bertina, R. M.: The

[122388]human protein S locus: identification of the PS-alpha gene as a site

[122389]of liver protein S messenger RNA synthesis. Biochem. Biophys. Res.

[122390]Commun. 157: 1033-1038, 1988.

[122391]21694.

[122392]Ploos van Amstel, J. K.; van der Zanden, A. L.; Bakker, E.;
Reitsma,

[122393]P. H.; Bertina, R. M.: Two genes homologous with human
protein S

[122394]cDNA are located on chromosome 3. Thromb. Haemost.
58: 982–987,

[122395]1987.

[122396]21695.

[122397]Reitsma, P. H.; Ploos van Amstel, H. K.; Bertina, R. M.:
Three

[122398]novel mutations in five unrelated subjects with hereditary
protein

[122399]S deficiency type I. J. Clin. Invest. 93: 486–492, 1994.

[122400]21696.

[122401]Sacco, R. L.; Owen, J.; Mohr, J. P.; Tatemichi, T. K.; Gross-
man,

[122402]B. A.: Free protein S deficiency: a possible association with
cerebrovascular

[122403]occlusion. Stroke 20: 1657–1661, 1989.

[122404]21697.

[122405] Sas, G.; Blasko, G.; Petro, I.; Griffin, J. H.: A protein S deficient

[122406] family with portal vein thrombosis. (Letter) *Thromb. Haemost.* 54:

[122407] 724 only, 1985.

[122408] 21698.

[122409] Chan, J. Y.; Cheung, M.-C.; Moi, P.; Chan, K.; Kan, Y. W.: Chromosomal

[122410] localization of the human NF-E2 family of bZIP transcription factors

[122411] by fluorescence in situ hybridization. *Hum. Genet.* 95: 265-269,

[122412] 1995.

[122413] 21699.

[122414] Chan, J. Y.; Han, X.-L.; Kan, Y. W.: Cloning of Nrf1, an NF-E2-related

[122415] transcription factor, by genetic selection in yeast. *Proc. Nat. Acad.*

[122416] *Sci.* 90: 11371-11375, 1993.

[122417] 21700.

[122418] Chan, J. Y.; Kwong, M.; Lu, R.; Chang, J.; Wang, B.; Yen, T.

S.

[122419]B.; Kan, Y. W.: Targeted disruption of the ubiquitous CNC-
bZIP transcription

[122420]factor, Nrf-1, results in anemia and embryonic lethality in
mice. EMBO

[122421]J. 17: 1779-1787, 1998.

[122422]21701.

[122423]Luna, L.; Johnsen, O.; Skartlien, A.; Pedeutour, F.; Turc-
Carel,

[122424]C.; Prydz, H.; Kolsto, A.-B.: Molecular cloning of a putative
novel

[122425]human bZIP transcription factor on chromosome 17q22.
Genomics 22:

[122426]553-562, 1994.

[122427]21702.

[122428]Luna, L.; Johnsen, O.; Skartlien, A. H.; Pedeutour, F.; Turc-
Carel,

[122429]C.; Prydz, H.; Kolsto, A.-B.: Molecular cloning of a putative
novel

[122430]human bZIP transcription factor on chromosome 17q22.
Genomics 22:

[122431]553-562, 1994.

[122432]21703.

[122433]McKie, J.; Johnstone, K.; Mattei, M.-G.; Scambler, P.:

Cloning

[122434]and mapping of murine Nfe2l1. Genomics 25: 716–719,
1995.

[122435]21704.

[122436]McKie, J.; Scambler, P. J.: The Nfe2l1 gene maps to distal
mouse

[122437]chromosome 11. Mammalian Genome 7: 89–90, 1996.

[122438]21705.

[122439]Dryja, T. P.; Berson, E. L.; Rao, V. R.; Oprian, D. D.: Het-
erozygous

[122440]missense mutation in the rhodopsin gene as a cause of
congenital stationary

[122441]night blindness. Nature Genet. 4: 280–283, 1993.

[122442]21706.

[122443]Gal, A.; Xu, S.; Piczenik, Y.; Eiberg, H.; Duvigneau, C.;
Schwinger,

[122444]E.; Rosenberg, T.: Gene for autosomal dominant congeni-
tal stationary

[122445]night blindness maps to the same region as the gene for

the beta-subunit
 of the rod photoreceptor cGMP phosphodiesterase (PDEB)
 in chromosome
 4p16.3. Hum. Molec. Genet. 3: 323-325, 1994.
 21707.
 Alfano, J. E.; Berger, J. P.: Retinitis pigmentosa, ophthalmoplegia,
 and spastic quadriplegia. Am. J. Ophthalm. 43: 231-240,
 1957.
 21708.
 Anttinen, A.; Nikoskelainen, E.; Marttila, R. J.; Grenman, R.;
 Falck, B.; Aarnisalo, E.; Kalimo, H.: Familial olivopontocerebellar
 atrophy with macular degeneration: a separate entity
 among the olivopontocerebellar
 atrophies. Acta Neurol. Scand. 73: 180-190, 1986.
 21709.
 Benomar, A.; Krols, L.; Stevanin, G.; Cancel, G.; LeGuern,
 E.;
 David, G.; Ouhabi, H.; Martin, J.-J.; Durr, A.; Zaim, A.;
 Ravise,

- [122459]N.; Busque, C.; Penet, C.; Van Regemorter, N.; Weissenbach, J.; Yahyaoui, M.; Chkili, T.; Agid, Y.; Van Broeckhoven, C.; Brice, A.: The gene for autosomal dominant cerebellar ataxia with pigmentary macular dystrophy maps to chromosome 3p12-p21.1. *Nature Genet.* 10: 84-88, 1995.
- [122463]21710.
- [122464]Benomar, A.; Le Guern, E.; Durr, A.; Ouhabi, H.; Stevanin, G.; Yahyaoui, M.; Chkili, T.; Agid, Y.; Brice, A.: Autosomal-dominant cerebellar ataxia with retinal degeneration (ADCA type II) is genetically different from ADCA type I. *Ann. Neurol.* 35: 439-444, 1994.
- [122468]21711.
- [122469]Bjork, A.; Lindblom, U.; Wadensten, L.: Retinal degeneration in hereditary ataxia. *J. Neurol. Neurosurg. Psychiatr.* 19: 186-193,

[122471]1956.

[122472]21712.

[122473]Carpenter, S.; Schumacher, G. A.: Familial infantile cerebellar

[122474]atrophy associated with retinal degeneration. Arch. Neurol. 14:

[122475]82-94, 1966.

[122476]21713.

[122477]Cooles, P.; Michaud, R.; Best, P. V.: A dominantly inherited progressive

[122478]disease in a black family characterized by cerebellar and retinal

[122479]degeneration, external ophthalmoplegia and abnormal mitochondria. J.

[122480]Neurol. Sci. 87: 275-288, 1988.

[122481]21714.

[122482]Cruysberg, J. R. M.; Eerola, K. U.; Vrijland, H. R.; Aandeker,

[122483]A. L.; Kremer, H. P. H.; Deutman, A. F.: Autosomal recessive cerebellar

[122484]ataxia with bull's-eye macular dystrophy. Am. J. Ophthal.

133: 410–413,

[122485]2002.

[122486]21715.

[122487]David, G.; Abbas, N.; Stevanin, G.; Durr, A.; Yvert, G.; Cancel,

[122488]G.; Weber, C.; Imbert, G.; Saudou, F.; Antoniou, E.; Drabkin, H.;

[122489]Gemmill, R.; Giunti, P.; Benomar, A.; Wood, N.; Ruberg, M.; Agid,

[122490]Y.; Mandel, J.-L.; Brice, A.: Cloning of the SCA7 gene reveals a

[122491]highly unstable CAG repeat expansion. *Nature Genet.* 17: 65–70, 1997.

[122492]21716.

[122493]David, G.; Giunti, P.; Abbas, N.; Coullin, P.; Stevanin, G.; Horta,

[122494]W.; Gemmell, R.; Weissenbach, J.; Wood, N.; Cunha, S.; Drabkin, H.;

[122495]Harding, A. E.; Agid, Y.; Brice, A.: The gene for autosomal dominant

[122496]cerebellar ataxia type II is located in a 5–cM region in 3p12–p13:

[122497]genetic and physical mapping of the SCA7 locus. Am. J. Hum. Genet. 59:

[122498]1328–1336, 1996.

[122499]21717.

[122500]Duinkerke–Eerola, K. U.; Cruysberg, J. R. M.; Deutman, A. F.:

[122501]Atrophic maculopathy associated with hereditary ataxia. Am. J. Ophthal. 90:

[122502]597–603, 1980.

[122503]21718.

[122504]Enevoldson, T. P.; Sanders, M. D.; Harding, A. E.: Autosomal

[122505]dominant cerebellar ataxia with pigmentary macular dystrophy: a clinical

[122506]and genetic study of eight families. Brain 117: 445–460, 1994.

[122507]21719.

[122508]Foster, J. B.; Ingram, T. T. S.: Familial cerebro–macular degeneration

[122509]and ataxia. J. Neurol. Neurosurg. Psychiat. 25: 63–68, 1962.

[122510]21720.

[122511]Froment, J.; Bonnet, P.; Colrat, A.: Heredo-degenerations
retinienne

[122512]et spino-cerebelleuse: variantes ophtalmoscopiques et
neurologiques

[122513]presentees par trois generations successives. J. Med. Lyon
153-163,

[122514]1937.

[122515]21721.

[122516]Giunti, P.; Stevanin, G.; Worth, P. F.; David, G.; Brice, A.;

[122517]Wood, N. W.: Molecular and clinical study of 18 families
with ADCA

[122518]type II: evidence for genetic heterogeneity and de novo
mutation. Am.

[122519]J. Hum. Genet. 64: 1594-1603, 1999.

[122520]21722.

[122521]Gouw, L. G.; Digre, K. B.; Harris, C. P.; Haines, J. H.;

Ptacek,

[122522]L. J.: Autosomal dominant cerebellar ataxia with retinal
degeneration:

[122523]clinical, neuropathologic, and genetic analysis of a large

kindred. Neurology 44:

[122524]1441-1447, 1994.

[122525]21723.

[122526]Gouw, L. G.; Kaplan, C. D.; Haines, J. H.; Digre, K. B.; Rutledge,

[122527]S. L.; Matilla, A.; Leppert, M.; Zoghbi, H. Y.; Ptacek, L. J.:
Retinal

[122528]degeneration characterizes a spinocerebellar ataxia mapping to chromosome

[122529]3p. Nature Genet. 10: 89-93, 1995.

[122530]21724.

[122531]Gu, W.; Wang, Y.; Liu, X.; Zhou, B.; Zhou, Y.; Wang, G.:
Molecular

[122532]and clinical study of spinocerebellar ataxia type 7 in Chinese kindreds. Arch.

[122533]Neurol. 57: 1513-1518, 2000.

[122534]21725.

[122535]Halsey, J. H., Jr.; Scott, T. R.; Farmer, T. W.: Adult hereditary

[122536]cerebelloretinal degeneration. Neurology 17: 87-90,
1967.

[122537]21726.

[122538]Sobetzko, D.; Eich, G.; Kalff-Suske, M.; Grzeschik, K.-H.;
Superti-Furga,

[122539]A.: Boy with syndactylies, macrocephaly, and severe skeletal dysplasia:

[122540]not a new syndrome, but two dominant mutations (GLI3
E543X and COL2A1

[122541]G973R) in the same individual. Am. J. Med. Genet. 90:
239-242, 2000.

[122542]21727.

[122543]Stroschein, S. L.; Wang, W.; Zhou, S.; Zhou, Q.; Luo, K.:
Negative

[122544]feedback regulation of TGF-beta signaling by the SnoN
oncoprotein. Science 286:

[122545]771-774, 1999.

[122546]21728.

[122547]Goldblum, S. E.; Ding, X.; Funk, S. E.; Sage, E. H.: SPARC
(secreted

[122548]protein acidic and rich in cysteine) regulates endothelial
cell shape

[122549]and barrier function. Proc. Nat. Acad. Sci. 91: 3448-3452,
1994.

[122550]21729.

[122551]Lossie, A. C.; Gordon, D. F.; Camper, S. A.: Localization of thyrotropin-releasing

[122552]hormone receptor and thyrotroph embryonic factor on mouse chromosome

[122553]15. Mammalian Genome 4: 621-623, 1993.

[122554]21730.

[122555]Bongarzone, I.; Butti, M. G.; Coronelli, S.; Borrello, M. G.; Santoro,

[122556]M.; Mondellini, P.; Pilotti, S.; Fusco, A.; Della Porta, G.; Pierotti,

[122557]M. A.: Frequent activation of ret protooncogene by fusion with a

[122558]new activating gene in papillary thyroid carcinomas. Cancer Res. 54:

[122559]2979-2985, 1994.

[122560]21731.

[122561]Bongarzone, I.; Butti, M. G.; Fugazzola, L.; Pacini, F.; Pinchera,

[122562]A.; Vorontsova, T. V.; Demidchik, E. P.; Pierotti, M. A.: Comparison

[122563]of the breakpoint regions of ELE1 and RET genes involved
in the generation

[122564]of RET/PTC3 oncogene in sporadic and in radiation-as-
sociated papillary

[122565]thyroid carcinomas. Genomics 42: 252–259, 1997.

[122566]21732.

[122567]Klein, M.; Vignaud, J.-M.; Hennequin, V.; Toussaint, B.;
Bresler,

[122568]L.; Plenat, F.; Leclere, J.; Duprez, A.; Weryha, G.: Increased
expression

[122569]of the vascular endothelial growth factor is a pejorative
prognosis

[122570]marker in papillary thyroid carcinoma. J. Clin. Endocr.
Metab. 86:

[122571]656–658, 2001.

[122572]21733.

[122573]Katyal, S. L.; Singh, G.; Locker, J.: Characterization of a
second

[122574]human pulmonary surfactant-associated protein SP-A
gene. Am. J. Resp.

[122575]Cell Molec. Biol. 6: 446–452, 1992.

[122576]21734.

[122577]Ramet, M.; Lofgren, J.; Albo, O.-P.; Hallman, M.: Surfactant protein-A

[122578]gene locus associated with recurrent otitis media. J. Pediat. 138:

[122579]266-268, 2001.

[122580]21735.

[122581]Mignon, C.; Okada, A.; Mattei, M. G.; Basset, P.: Assignment of

[122582]the human membrane-type matrix metalloproteinase (MMP14) gene to 14q11-q12

[122583]by in situ hybridization. Genomics 28: 360-361, 1995.

[122584]21736.

[122585]Wilson, C. L.; Ouellette, A. J.; Satchell, D. P.; Ayabe, T.; Lopez-Boado,

[122586]Y. S.; Stratman, J. L.; Hultgren, S. J.; Matrisian, L. M.; Parks,

[122587]W. C.: Regulation of intestinal alpha-defensin activation by the

[122588]metalloproteinase matrilysin in innate host defense. Science 286:

[122589]113-117, 1999.

[122590]21737.

[122591]Inglehearn, C. F.; Bashir, R.; Lester, D. H.; Jay, M.; Bird, A.

[122592]C.; Bhattacharya, S. S.: A 3-bp deletion in the rhodopsin
gene in

[122593]a family with autosomal dominant retinitis pigmentosa.
Am. J. Hum.

[122594]Genet. 48: 26-30, 1991.

[122595]21738.

[122596]Inglehearn, C. F.; Keen, T. J.; Bashir, R.; Jay, M.; Fitzke, F.;

[122597]Bird, A. C.; Crombie, A.; Bhattacharya, S.: A completed
screen for

[122598]mutations of the rhodopsin gene in a panel of patients
with autosomal

[122599]dominant retinitis pigmentosa. Hum. Molec. Genet. 1:
41-45, 1992.

[122600]21739.

[122601]Inglehearn, C. F.; Lester, D. H.; Bashir, R.; Atif, U.; Keen,

[122602]T. J.; Sertedaki, A.; Lindsey, J.; Jay, M.; Bird, A. C.; Farrar,
G.

[122603]J.; Humphries, P.; Bhattacharya, S. S.: Recombination be-
tween rhodopsin

[122604]and locus D3S47 (C17) in rhodopsin retinitis pigmentosa families. Am.

[122605]J. Hum. Genet. 50: 590–597, 1992.

[122606]21740.

[122607]Keen, T. J.; Inglehearn, C. F.; Lester, D. H.; Bashir, R.; Jay,

[122608]M.; Bird, A. C.; Jay, B.; Bhattacharya, S. S.: Autosomal dominant

[122609]retinitis pigmentosa: four new mutations in rhodopsin, one of them

[122610]in the retinal attachment site. Genomics 11: 199–205, 1991.

[122611]21741.

[122612]Khorana, H. G.: Rhodopsin, photoreceptor of the rod cell: an

[122613]emerging pattern for structure and function. J. Biol. Chem. 267:

[122614]1–4, 1992.

[122615]21742.

[122616]Kijas, J. W.; Cideciyan, A. V.; Aleman, T. S.; Pianta, M. J.;

[122617]Pearce–Kelling, S. E.; Miller, B. J.; Jacobson, S. G.; Aguirre, G.

[122618]D.; Acland, G. M.: Naturally occurring rhodopsin mutation
in the

[122619]dog causes retinal dysfunction and degeneration mimick-
ing human dominant

[122620]retinitis pigmentosa. Proc. Nat. Acad. Sci. 99: 6328–6333,
2002.

[122621]21743.

[122622]Kiselev, A.; Socolich, M.; Vinos, J.; Hardy, R. W.; Zuker, C.

[122623]S.; Ranganathan, R.: A molecular pathway for light-
dependent photoreceptor

[122624]apoptosis in Drosophila. Neuron 28: 139–152, 2000.

[122625]21744.

[122626]Kranich, H.; Bartkowski, S.; Denton, M. J.; Krey, S.; Dickin-
son,

[122627]P.; Duvigneau, C.; Gal, A.: Autosomal dominant 'sector' re-
tinitis

[122628]pigmentosa due to a point mutation predicting an
asn15-to-ser substitution

[122629]of rhodopsin. Hum. Molec. Genet. 2: 813–814, 1993.

[122630]21745.

[122631]Kumar-Singh, R.; Wang, H.; Humphries, P.; Farrar, G. J.:

Autosomal

[122632]dominant retinitis pigmentosa: no evidence for nonallelic genetic

[122633]heterogeneity on 3q. Am. J. Hum. Genet. 52: 319–326, 1993.

[122634]21746.

[122635]Kumaramanickavel, G.; Maw, M.; Denton, M. J.; John, S.; Srisailapathy

[122636]Srikumari, C. R.; Orth, U.; Oehlmann, R.; Gal, A.: Missense rhodopsin

[122637]mutation in a family with recessive RP. (Letter) Nature Genet. 8:

[122638]10–11, 1994.

[122639]21747.

[122640]Lem, J.; Krasnoperova, N. V.; Calvert, P. D.; Kosaras, B.; Cameron,

[122641]D. A.; Nicolo, M.; Makino, C. L.; Sidman, R. L.: Morphological, physiological,

[122642]and biochemical changes in rhodopsin knockout mice. Proc. Nat. Acad.

[122643]Sci. 96: 736–741, 1999.

[122644]21748.

[122645]Li, T.; Sandberg, M. A.; Pawlyk, B. S.; Rosner, B.; Hayes, K.

[122646]C.; Dryja, T. P.; Berson, E. L.: Effect of vitamin A supplementation

[122647]on rhodopsin mutants threonine-17-to-methionine and proline-347-to-serine

[122648]in transgenic mice and in cell cultures. Proc. Nat. Acad. Sci. 95:

[122649]11933-11938, 1998.

[122650]21749.

[122651]Li, T.; Snyder, W. K.; Olsson, J. E.; Dryja, T. P.: Transgenic

[122652]mice carrying the dominant rhodopsin mutation P347S: evidence for

[122653]defective vectorial transport of rhodopsin to the outer segments. Proc.

[122654]Nat. Acad. Sci. 93: 14176-14181, 1996.

[122655]21750.

[122656]Macke, J. P.; Davenport, C. M.; Jacobson, S. G.; Hennessey, J.

[122657]C.; Gonzalez-Fernandez, F.; Conway, B. P.; Heckenlively, J.; Palmer,

[122658]R.; Maumenee, I. H.; Sieving, P.; Gouras, P.; Good, W.;

Nathans, J.

[122659] Identification of novel rhodopsin mutations responsible for retinitis

[122660] pigmentosa: implications for the structure and function of rhodopsin. Am.

[122661] J. Hum. Genet. 53: 80–89, 1993.

[122662] 21751.

[122663] Mendez, A.; Burns, M. E.; Roca, A.; Lem, J.; Wu, L.-W.; Simon,

[122664] M. I.; Baylor, D. A.; Chen, J.: Rapid and reproducible deactivation

[122665] of rhodopsin requires multiple phosphorylation sites.

Neuron 28:

[122666] 153–164, 2000.

[122667] 21752.

[122668] Naash, M. I.; Hollyfield, J. G.; Al-Ubaidi, M. R.; Baehr, W.:

[122669] Simulation of human autosomal dominant retinitis pigmentosa in transgenic

[122670] mice expressing a mutated murine opsin gene. Proc. Nat. Acad. Sci. 90:

[122671] 5499–5503, 1993.

[122672]21753.

[122673]Nathans, J.; Hogness, D. S.: Isolation and nucleotide sequence

[122674]of the gene encoding human rhodopsin. Proc. Nat. Acad. Sci. 81:

[122675]4851–4855, 1984.

[122676]21754.

[122677]Nathans, J.; Piantanida, T. P.; Eddy, R. L.; Shows, T. B.; Hogness,

[122678]D. S.: Molecular genetics of inherited variation in human color vision. Science 232:

[122679]203–210, 1986.

[122680]21755.

[122681]Nathans, J.; Thomas, D.; Hogness, D. S.: Molecular genetics of

[122682]human color vision: the genes encoding blue, green, and red pigments. Science 232:

[122683]193–202, 1986.

[122684]21756.

[122685]Nie, Z.; Chen, S.; Kumar, R.; Zack, D. J.: RER, an evolutionarily

[122686] conserved sequence upstream of the rhodopsin gene, has enhancer activity. J.

[122687] Biol. Chem. 271: 2667–2675, 1996.

[122688] 21757.

[122689] Oh, K. T.; Weleber, R. G.; Lotery, A.; Oh, D. M.; Billingslea,

[122690] A. M.; Stone, E. M.: Description of a new mutation in rhodopsin,

[122691] pro23 to ala, and comparison with electroretinographic and clinical

[122692] characteristics of the pro23-to-his mutation. Arch. Ophthalmol. 118:

[122693] 1269–1276, 2000.

[122694] 21758.

[122695] Olsson, J. E.; Gordon, J. W.; Pawlyk, B. S.; Roof, D.; Hayes,

[122696] A.; Molday, R. S.; Mukai, S.; Cowley, G. S.; Berson, E. L.; Dryja,

[122697] T. P.: Transgenic mice with a rhodopsin mutation (Pro23His): a mouse

[122698] model of autosomal dominant retinitis pigmentosa. Neuron 9: 815–830,

[122699] 1992.

[122700]21759.

[122701]Palczewski, K.; Kumasaka, T.; Hori, T.; Behnke, C. A.; Mo-
toshima,

[122702]H.; Fox, B. A.; Le Trong, I.; Teller, D. C.; Okada, T.;
Stenkamp,

[122703]R. E.; Yamamoto, M.; Miyano, M.: Crystal structure of
rhodopsin:

[122704]a G protein-coupled receptor. Science 289: 739-745,
2000.

[122705]21760.

[122706]Rao, V. R.; Cohen, G. B.; Oprian, D. D.: Rhodopsin muta-
tion G90D

[122707]and a molecular mechanism for congenital night blind-
ness. Nature 367:

[122708]639-642, 1994.

[122709]21761.

[122710]Robinson, P. R.; Cohen, G. B.; Zhukovsky, E. A.; Oprian, D.
D.

[122711]: Constitutively active mutants of rhodopsin. Neuron 9:
719-725,

[122712]1992.

[122713]21762.

[122714]Rosenfeld, P. J.; Cowley, G. S.; Hahn, L. B.; Sandberg, M. A.;

[122715]Berson, E. L.; Dryja, T. P.: A null mutation within the rhodopsin

[122716]gene in a family with autosomal recessive retinitis pigmentosa. (Abstract) Invest.

[122717]Ophthalmol. Vis. Sci. 33: 1397, 1992.

[122718]21763.

[122719]Mason, I. J.; Murphy, D.; Munke, M.; Francke, U.; Elliott, R. W.;

[122720]Hogan, B. L. M.: Developmental and transformation-sensitive expression

[122721]of the SPARC gene on mouse chromosome 11. EMBO J. 5: 1831-1837,

[122722]1986.

[122723]21764.

[122724]Mason, I. J.; Taylor, A.; Williams, J. G.; Sage, H.; Hogan, B.

[122725]L. M.: Evidence from molecular cloning that SPARC, a major product

[122726]of mouse embryo parietal endoderm, is related to an en-

endothelial cell

[122727] 'culture shock' glycoprotein of Mr 43,000. EMBO J. 5:

1465–1472,

[122728] 1986.

[122729] 21765.

[122730] Naylor, S. L.; Helen-Davis, D.; Villarreal, X. C.; Long, G. L.

[122731]: The human osteonectin gene on chromosome 5 is polymorphic. (Abstract) Cytogenet.

[122732] Cell Genet. 51: 1051 only, 1989.

[122733] 21766.

[122734] Schwartz, R. C.; Young, M. F.; Tsiouras, P.: Two RFLPs in the

[122735] 5-prime end of the human osteonectin (ON) gene. Nucleic Acids Res. 16:

[122736] 9076 only, 1988.

[122737] 21767.

[122738] Stenner, D. D.; Tracy, R. P.; Riggs, B. L.; Mann, K. G.: Human

[122739] platelets contain and secrete osteonectin, a major protein of mineralized

[122740] bone. Proc. Nat. Acad. Sci. 83: 6892–6896, 1986.

[122741]21768.

[122742]Swaroop, A.; Francke, U.: Molecular cloning, cDNA sequence, and

[122743]expression of human SPARC (osteonectin). (Abstract) Am. J. Hum. Genet. 41:

[122744]A240 only, 1987.

[122745]21769.

[122746]Swaroop, A.; Hogan, B. L. M.; Francke, U.: Molecular analysis

[122747]of the cDNA for human SPARC/osteonectin/BM-40: sequence, expression,

[122748]and localization of the gene to chromosome 5q31-q33. Genomics 2:

[122749]37-47, 1988.

[122750]21770.

[122751]Termine, J. D.; Kleinman, H. K.; Whitson, S. W.; Conn, K. M.;

[122752]McGarvey, M. L.; Martin, G. R.: Osteonectin, a bone-specific protein

[122753]linking mineral to collagen. Cell 26: 99-105, 1981.

[122754]21771.

[122755]Thony, B.; Heizmann, C. W.; Mattei, M.-G.: Human GTP-cyclohydrolase

[122756] gene and sepiapterin reductase gene map to region 14q21-q22 and

[122757]2p14-p12, respectively, by in situ hybridization. Genomics 26: 168-170,

[122758]1995.

[122759]21772.

[122760]Jin, H.; Oksenberg, D.; Ashkenazi, A.; Peroutka, S. J.; Duncan,

[122761]A. M. V.; Rozmahel, R.; Yang, Y.; Mengod, G.; Palacios, J. M.; O'Dowd,

[122762]B. F.: Characterization of the human 5-hydroxytryptamine(1B) receptor. J.

[122763]Biol. Chem. 267: 5735-5738, 1992.

[122764]21773.

[122765]Levy, F. O.; Gudermann, T.; Birnbaumer, M.; Kaumann, A. J.; Birnbaumer,

[122766]L.: Molecular cloning of a human gene (S31) encoding a novel serotonin

[122767]receptor mediating inhibition of adenylyl cyclase. FEBS Lett. 296:

[122768]201–206, 1992.

[122769]21774.

[122770]Levy, F. O.; Holtgreve-Grez, H.; Tasken, K.; Solberg, R.;
Ried,

[122771]T.; Gudermann, T.: Assignment of the gene encoding the
5-HT-1E serotonin

[122772]receptor (S31) (locus HTR1E) to human chromosome
6q14–q15. Genomics 22:

[122773]637–640, 1994.

[122774]21775.

[122775]McAllister, G.; Charlesworth, A.; Snodin, C.; Beer, M. S.;
Noble,

[122776]A. J.; Middlemiss, D. N.; Iversen, L. L.; Whiting, P.: Molecu-
lar

[122777]cloning of a serotonin receptor from human brain
(5HT1E): a fifth

[122778]5HT1-like subtype. Proc. Nat. Acad. Sci. 89: 5517–5521,
1992.

[122779]21776.

[122780]Weinshank, R. L.; Zgombick, J. M.; Macchi, M. J.; Branchek,
T.

[122781]A.; Hartig, P. R.: Human serotonin 1D receptor is encoded by a subfamily

[122782]of two distinct genes: 5-HT(1D-alpha) and 5-HT(1D-beta). Proc. Nat.

[122783]Acad. Sci. 89: 3630-3634, 1992.

[122784]21777.

[122785]Pabst, T.; Mueller, B. U.; Zhang, P.; Radomska, H. S.; Nar-ravula,

[122786]S.; Schnittger, S.; Behre, G.; Hiddemann, W.; Tenen, D. G.: Dominant-negative

[122787]mutations of CEBPA, encoding CCAAT/enhancer binding protein-alpha

[122788](C/EBP-alpha), in acute myeloid leukemia. Nature Genet. 27: 263-270,

[122789]2001.

[122790]21778.

[122791]Chen, Z.-Y.; Battinelli, E. M.; Fielder, A.; Bunday, S.; Sims,

[122792]K.; Breakefield, X. O.; Craig, I. W.: A mutation in the Norrie disease

[122793]gene (NDP) associated with X-linked familial exudative vitreoretinopathy. Nature

[122794]Genet. 5: 180–183, 1993.

[122795]21779.

[122796]Dudgeon, J.: Familial exudative vitreo–retinopathy. Trans.
Ophthalm.

[122797]Soc. U.K. 99: 45–49, 1979.

[122798]21780.

[122799]Matsuda, I.; Arashima, S.; Nambu, H.; Takekoshi, Y.;
Anakura,

[122800]M.: Hyperammonemia due to a mutant enzyme of or-
nithine transcarbamylase. Pediatrics 48:

[122801]595–600, 1971.

[122802]21781.

[122803]Yang–Feng, T. L.; DeGennaro, L. J.; Francke, U.: Genes for
synapsin

[122804]I, a neuronal phosphoprotein, map to conserved regions
of human and

[122805]murine X chromosomes. Proc. Nat. Acad. Sci. 83:
8679–8683, 1986.

[122806]21782.

[122807]Matsuda, I.; Nagata, N.; Matsuura, T.; Oyanagi, K.; Tada,
K.;

[122808]Narisawa, K.; Kitagawa, T.; Sakiyama, T.; Yamashita, F.;
Yoshino,
[122809]M.: Retrospective survey of urea cycle disorders: Part 1.
Clinical
[122810]and laboratory observations of thirty-two Japanese male
patients with
[122811]ornithine transcarbamylase deficiency. Am. J. Med. Genet.
38: 85-89,
[122812]1991.
[122813]21783.
[122814]Mavinakere, M.; Morizono, H.; Shi, D.; Allewell, N. M.;
Tuchman,
[122815]M.: The clinically variable R40H mutant ornithine car-
bamoyltransferase
[122816]shows cytosolic degradation of the precursor protein in
CHO cells. J.
[122817]Inherit. Metab. Dis. 24: 614-622, 2001.
[122818]21784.
[122819]Michels, V. V.; Potts, E.; Walser, M.; Beaudet, A. L.: Or-
nithine
[122820]transcarbamylase deficiency: long-term survival. Clin.
Genet. 22:

[122821]211-214, 1982.

[122822]21785.

[122823]Matsuda, I.; Matsuura, T.; Nishiyori, A.; Komaki, S.;
Hoshide,

[122824]R.; Matsumoto, T.; Funakoshi, M.; Kiwaki, K.; Endo, F.;
Hata, A.;

[122825]Shimadzu, M.; Yoshino, M.: Phenotypic variability in male
patients

[122826]carrying the mutant ornithine transcarbamylase (OTC) al-
lele, arg40his,

[122827]ranging from a child with an unfavourable prognosis to an
asymptomatic

[122828]older adult. J. Med. Genet. 33: 645-648, 1996.

[122829]21786.

[122830]Morsy, M. A.; Zhao, J. Z.; Ngo, T. T.; Warman, A. W.;
O'Brien,

[122831]W. E.; Graham, F. L.; Caskey, C. T.: Patient selection may
affect

[122832]gene therapy success: dominant negative effects observed
for ornithine

[122833]transcarbamylase in mouse and human hepatocytes. J.
Clin. Invest. 97:

[122834]826-832, 1996.

[122835]21787.

[122836]Nagata, N.; Matsuda, I.; Oyanagi, K.: Estimated frequency
of
[122837]urea cycle enzymopathies in Japan. (Letter) Am. J. Med.
Genet. 39:

[122838]228-229, 1991.

[122839]21788.

[122840]Nussbaum, R. L.; Boggs, B. A.; Beaudet, A. L.; Doyle, S.;
Potter,

[122841]J. L.; O'Brien, W. E.: New mutation and prenatal diagnosis
in ornithine

[122842]transcarbamylase deficiency. Am. J. Hum. Genet. 38:
149-158, 1986.

[122843]21789.

[122844]Ohtake, A.; Takayanagi, M.; Yamamoto, S.; Kakinuma, H.;
Nakajima,

[122845]H.; Tatibana, M.; Mori, M.: Molecular basis of ornithine
transcarbamylase

[122846]deficiency in spf and spf-ash mutant mice. J. Inherit.
Metab. Dis. 9:

[122847]289–291, 1986.

[122848]21790.

[122849]Oizumi, J.; Ng, W. G.; Koch, R.; Shaw, K. N. F.; Sweetman, L.;

[122850]Velazquez, A.; Donnell, G. N.: Partial ornithine transcarbamylase

[122851]deficiency associated with recurrent hyperammonemia, lethargy and

[122852]depressed sensorium. Clin. Genet. 25: 538–542, 1984.

[122853]21791.

[122854]Old, J. M.; Briand, P. L.; Purvis-Smith, S.; Howard, N. J.; Wilcken,

[122855]B.; Hammond, J.; Pearson, P.; Cathelineau, L.; Williamson, R.; Davies,

[122856]K. E.: Prenatal exclusion of ornithine transcarbamylase deficiency

[122857]by direct gene analysis. Lancet I: 73–75, 1985.

[122858]21792.

[122859]Oppliger Leibundgut, E.; Liechti-Gallati, S.; Colombo, J.-P.;

[122860]Wermuth, B.: Ornithine transcarbamylase deficiency: ten new mutations

[122861]and high proportion of de novo mutations in heterozygous females. Hum.

[122862]Mutat. 9: 409–411, 1997.

[122863]21793.

[122864]Oppliger Leibundgut, E.; Liechti-Gallati, S.; Colombo, J. P.;

[122865]Wermuth, B.: Ornithine transcarbamylase deficiency: new sites with

[122866]increased probability of mutation. Hum. Genet. 95: 191–196, 1995.

[122867]21794.

[122868]Oppliger Leibundgut, E.; Wermuth, B.; Colombo, J.-P.; Liechti-Gallati,

[122869]S.: Ornithine transcarbamylase deficiency: characterization of gene

[122870]mutations and polymorphisms. Hum. Mutat. 8: 333–339, 1996.

[122871]21795.

[122872]Pelet, A.; Rotig, A.; Bonaiti-Pellie, C.; Rabier, D.; Cormier,

[122873]V.; Toumas, E.; Hentzen, D.; Saudubray, J.-M.; Munnich, A.: Carrier

[122874]detection in a partially dominant X-linked disease: or-

ornithine transcarbamylase

[122875] deficiency. Hum. Genet. 84: 167–171, 1990.

[122876] 21796.

[122877] Pembrey, M. E.; Old, J. M.; Leonard, J. V.; Rodeck, C. H.;
Warren,

[122878] R.; Davies, K. E.: Prenatal diagnosis of ornithine carbamoyl
transferase

[122879] deficiency using a gene specific probe. J. Med. Genet. 22:
462–465,

[122880] 1985.

[122881] 21797.

[122882] Ploechl, E.; Ploechl, W.; Stoeckler-Ipsiroglu, S.; Pokorny,
H.;

[122883] Wermuth, B.: Late-onset ornithine transcarbamylase defi-
ciency in

[122884] two families with different mutations in the same codon.
Clin. Genet. 59:

[122885] 111–114, 2001.

[122886] 21798.

[122887] Qureshi, I. A.; Letarte, J.; Ouellet, R.: Ornithine transcar-
bamylase

[122888]deficiency in mutant mice. I. Studies on the characterization of enzyme

[122889]defect and suitability as animal model of human disease. *Pediat.*

[122890]*Res.* 13: 807-811, 1979.

[122891]21799.

[122892]Ricciuti, F. C.; Gelehrter, T. D.; Rosenberg, L. E.: X-chromosome

[122893]inactivation in human liver: confirmation of X-linkage of ornithine

[122894]transcarbamylase. *Am. J. Hum. Genet.* 28: 332-338, 1976.

[122895]21800.

[122896]Rodeck, C. H.; Patrick, A. D.; Pembrey, M. E.; Tzannatos, C.;

[122897]Whitfield, A. E.: Fetal liver biopsy for prenatal diagnosis of ornithine

[122898]carbamyl transferase deficiency. *Lancet* II: 297-300, 1982.

[122899]21801.

[122900]Rosenberg, L. E.; Kalousek, F.; Orsulak, M. D.: Biogenesis of

[122901]ornithine transcarbamylase in *spf(ash)* mutant mice: two

cytoplasmic

[122902]precursors, one mitochondrial enzyme. Science 222:
426-428, 1983.

[122903]21802.

[122904]Rowe, P. C.; Newman, S. L.; Brusilow, S. W.: Natural history
[122905]of symptomatic partial ornithine transcarbamylase defi-
ciency. New

[122906]Eng. J. Med. 314: 541-547, 1986.

[122907]21803.

[122908]Rozen, R.; Fox, J.; Fenton, W. A.; Horwich, A. L.; Rosen-
berg,

[122909]L. E.: Gene deletion and restriction fragment length poly-
morphisms

[122910]at the human ornithine transcarbamylase locus. Nature
313: 815-817,

[122911]1985.

[122912]21804.

[122913]Russell, A.; Levin, B.; Oberholzer, V. G.; Sinclair, L.: Hyper-
ammonaemia.

[122914]A new instance of an inborn enzymatic defect of the
biosynthesis of

[122915]urea. Lancet II: 699–700, 1962.

[122916]21805.

[122917]Schwartz, M.; Christensen, E.; Christensen, N. C.; Skovby, F.;

[122918]Davies, K. E.; Old, J. M.: Detection and exclusion of carriers of

[122919]ornithine transcarbamylase deficiency by RFLP analysis. Clin. Genet. 29:

[122920]449–452, 1986.

[122921]21806.

[122922]Scott, C. R.; Chiang–Teng, C.; Goodman, S. I.; Greensher, A.;

[122923]Mace, J. W.: X-linked transmission of ornithine–transcarbamylase

[122924]deficiency. (Letter) Lancet II: 1148, 1972.

[122925]21807.

[122926]Shapiro, J. M.; Schaffner, F.; Tallan, H. H.; Gaull, G. E.: Mitochondrial

[122927]abnormalities of liver in primary ornithine transcarbamylase deficiency. Pediat.

[122928]Res. 14: 735–759, 1980.

[122929]21808.

[122930]Sinke, R. J.; de Leeuw, B.; Janssen, H. A. P.; Olde Weghuis, D.;

[122931]Suijkerbuijk, R. F.; Meloni, A. M.; Gilgenkrantz, S.; Berger, W.;

[122932]Ropers, H. H.; Sandberg, A. A.; Geurts van Kessel, A.: Localization

[122933]of X chromosome short arm markers relative to synovial sarcoma- and

[122934]renal adenocarcinoma-associated translocation break-points. Hum. Genet. 92:

[122935]305-308, 1993.

[122936]21809.

[122937]Brennan, T. J.; Seeley, W. W.; Kilgard, M.; Schreiner, C. E.; Tecott,

[122938]L. H.: Sound-induced seizures in serotonin 5-HT-2C receptor mutant

[122939]mice. Nature Genet. 16: 387-390, 1997.

[122940]21810.

[122941]Gurevich, I.; Tamir, H.; Arango, V.; Dwork, A. J.; Mann, J. J.;

[122942]Schmauss, C.: Altered editing of serotonin 2C receptor

pre-mRNA in

[122943]the prefrontal cortex of depressed suicide victims. *Neuron*
34: 349–356,

[122944]2002.

[122945]21811.

[122946]Hall, C. S.: Genetic differences in fatal audiogenic
seizures:

[122947]between two inbred strains of house mice. *J. Hered.* 38:
3–6, 1947.

[122948]21812.

[122949]Lappalainen, J.; Zhang, L.; Dean, M.; Oz, M.; Ozaki, N.; Yu,
D.;

[122950]Virkkunen, M.; Weight, F.; Linnoila, M.; Goldman, D.: Iden-
tification,

[122951]expression, and pharmacology of a cys(23)–ser(23) sub-
stitution in

[122952]the human 5-HT(2C) receptor gene (HTR2C). *Genomics*
27: 274–279,

[122953]1995.

[122954]21813.

[122955]Milatovich, A.; Hsieh, C.–L.; Bonaminio, G.; Tecott, L.;

Julius,

[122956]D.; Francke, U.: Serotonin receptor 1c gene assigned to X chromosome

[122957]in human (band q24) and mouse (bands D-F4). Hum.

Molec. Genet. 1:

[122958]681-684, 1992.

[122959]21814.

[122960]Tecott, L. H.; Sun, L. M.; Akana, S. F.; Strack, A. M.;

Lowenstein,

[122961]D. H.; Dallman, M. F.; Julius, D.: Eating disorder and epilepsy in

[122962]mice lacking 5-HT_{2C} serotonin receptors. Nature 374: 542-546, 1995.

[122963]21815.

[122964]Blaschke, R. J.; Monaghan, A. P.; Schiller, S.; Schechinger, B.;

[122965]Rao, E.; Padilla-Nash, H.; Ried, T.; Rappold, G. A.: SHOT, a SHOX-related

[122966]homeobox gene, is implicated in craniofacial, brain, heart, and limb

[122967]development. Proc. Nat. Acad. Sci. 95: 2406-2411, 1998.

[122968]21816.

[122969]Castillo, S.; Youlton, R.; Be, C.: Dyschondrosteosis is controlled

[122970]by X and Y linked loci. (Abstract) Cytogenet. Cell Genet.
40: 601-602,

[122971]1985.

[122972]21817.

[122973]Clement-Jones, M.; Schiller, S.; Rao, E.; Blaschke, R. J.; Zuniga,

[122974]A.; Zeller, R.; Robson, S. C.; Binder, G.; Glass, I.; Strachan,
T.;

[122975]Lindsay, S.; Rappold, G. A.: The short stature homeobox
gene SHOX

[122976]is involved in skeletal abnormalities in Turner syndrome.
Hum. Molec.

[122977]Genet. 9: 695-702, 2000.

[122978]21818.

[122979]Ellison, J. W.; Wardak, Z.; Webster, M.; Chiong, W.: PHOG, a
candidate

[122980]gene for involvement in the short stature of Turner syndrome. (Abstract) Am.

[122981]J. Hum. Genet. 59 (suppl.): A32, 1996.

[122982]21819.

[122983]Ellison, J. W.; Wardak, Z.; Young, M. F.; Robey, P. G.; Webster,
ster,

[122984]M.; Chiong, W.: PHOG, a candidate gene for involvement in
the short

[122985]stature of Turner syndrome. Hum. Molec. Genet. 6:
1341–1347, 1997.

[122986]21820.

[122987]Guichet, A.; Briault, S.; Le Merrer, M.; Moraine, C.: Are
t(X;Y)

[122988](p22;q11) translocations in females frequently associated
with Madelung

[122989]deformity? Clin. Dysmorphol. 6: 341–345, 1997.

[122990]21821.

[122991]Henke, A.; Wapenaar, M.; van Ommen, G.-J.; Maraschio,
P.; Camerino,

[122992]G.; Rappold, G.: Deletions within the pseudoautosomal re-
gion help

[122993]map three new markers and indicate a possible role of
this region

[122994]in linear growth. Am. J. Hum. Genet. 49: 811–819, 1991.

[122995]21822.

[122996]Affara, N. A.; Chalmers, I. J.; Ferguson-Smith, M. A.: Analysis

[122997]of the SRY gene in 22 sex-reversed XY females identifies four new

[122998]point mutations in the conserved DNA binding domain. Hum. Molec.

[122999]Genet. 2: 785-789, 1993.

[123000]21823.

[123001]Behlke, M. A.; Bogan, J. S.; Beer-Romero, P.; Page, D. C.: Evidence

[123002]that the SRY protein is encoded by a single exon on the human Y chromosome. Genomics 17:

[123003]736-739, 1993.

[123004]21824.

[123005]Berkovitz, G. D.; Fechner, P. Y.; Zacur, H. W.; Rock, J. A.; Snyder,

[123006]H. M., III; Migeon, C. J.; Perlman, E. J.: Clinical and pathologic

[123007]spectrum of 46,XY gonadal dysgenesis: its relevance to the understanding

[123008]of sex differentiation. *Medicine* 70: 375–383, 1991.

[123009]21825.

[123010]Bowles, J.; Cooper, L.; Berkman, J.; Koopman, P.: Sry re-
quires

[123011]a CAG repeat domain for male sex determination in *Mus*
musculus. *Nature*

[123012]*Genet.* 22: 405–408, 1999.

[123013]21826.

[123014]Braun, A.; Kammerer, S.; Cleve, H.; Lohrs, U.; Schwarz,
H.-P.;

[123015]Kuhnle, U.: True hermaphroditism in a 46,XY individual,
caused by

[123016]a postzygotic somatic point mutation in the male gonadal
sex-determining

[123017]locus (SRY): molecular genetics and histological findings
in a sporadic

[123018]case. *Am. J. Hum. Genet.* 52: 578–585, 1993.

[123019]21827.

[123020]Brown, S.; Yu, C. C.; Lanzano, P.; Heller, D.; Thomas, L.;
Warburton,

[123021]D.; Kitajewski, J.; Stadtmauer, L.: A de novo mutation

(gln2stop)

[123022]at the 5-prime end of the SRY gene leads to sex reversal
with partial

[123023]ovarian function. (Letter) Am. J. Hum. Genet. 62: 189-192,
1998.

[123024]21828.

[123025]Canto, P.; de la Chesnaye, E.; Lopez, M.; Cervantes, A.;
Chavez,

[123026]B.; Vilchis, F.; Reyes, E.; Ulloa-Aguirre, A.; Kofman-Alfaro,
S.;

[123027]Mendez, J. P.: A mutation in the 5-prime non-high mobil-
ity group

[123028]box region of the SRY gene in patients with Turner syn-
drome and Y

[123029]mosaicism. J. Clin. Endocr. Metab. 85: 1908-1911, 2000.

[123030]21829.

[123031]Cherfas, J.: Sex and the single gene. Science 252: 782,
1991.

[123032]21830.

[123033]Clepet, C.; Schafer, A. J.; Sinclair, A. H.; Palmer, M. S.;
Lovell-Badge,

[123034]R.; Goodfellow, P. N.: The human SRY transcript. Hum.

Molec. Genet. 2:

[123035]2007–2012, 1993.

[123036]21831.

[123037]Davis, R. M.: Localisation of male determining factors in man:

[123038]a thorough review of structural anomalies of the Y chromosome. J.

[123039]Med. Genet. 18: 161–195, 1981.

[123040]21832.

[123041]Dork, T.; Stuhrmann, M.; Miller, K.; Schmidtke, J.: Independent

[123042]observation of SRY mutation I90M in a patient with complete gonadal

[123043]dysgenesis. (Letter) Hum. Mutat. 11: 90–91, 1998.

[123044]21833.

[123045]Eicher, E. M.; Washburn, L. L.: Genetic control of primary sex

[123046]determination in mice. Annu. Rev. Genet. 20: 327–360, 1986.

[123047]21834.

[123048]Foster, J. W.; Brennan, F. E.; Hampikian, G. K.; Goodfellow,

P.

[123049]N.; Sinclair, A. H.; Lovell-Badge, R.; Selwood, L.; Renfree, M. B.;

[123050]Cooper, D. W.; Graves, J. A. M.: Evolution of sex determination and

[123051]the Y chromosome: SRY-related sequences in marsupials. Nature 359:

[123052]531-533, 1992.

[123053]21835.

[123054]Fuqua, J. S.; McLaughlin, J.; Perlman, E. J.; Berkowitz, G. D.

[123055]: Analysis of the SRY gene in gonadal tissue of subjects with 46,XY

[123056]gonadal dysgenesis (Letter) J. Clin. Endocr. Metab. 82: 701-702,

[123057]1997.

[123058]21836.

[123059]Domenice, S.; Nishi, M. Y.; Billerbeck, A. E. C.; Latronico, A.

[123060]C.; Medeiros, M. A.; Russell, A. J.; Vass, K.; Carvalho, F. M.; Frade,

[123061]E. M. C.; Arnhold, I. J. P.; Mendonca, B. B.: A novel mis-

sense mutation

[123062](S18N) in the 5-prime non-HMG box region of the SRY gene in a patient

[123063]with partial gonadal dysgenesis and his normal male relatives. Hum.

[123064]Genet. 102: 213-215, 1998.

[123065]21837.

[123066]Giese, K.; Pagel, J.; Grosschedl, R.: Distinct DNA-binding properties

[123067]of the high mobility group domain of murine and human SRY sex-determining

[123068]factors. Proc. Nat. Acad. Sci. 91: 3368-3372, 1994.

[123069]21838.

[123070]Goodfellow, P. N.; Lovell-Badge, R.: SRY and sex determination

[123071]in mammals. Annu. Rev. Genet. 27: 71-92, 1993.

[123072]21839.

[123073]Gubbay, J.; Collignon, J.; Koopman, P.; Capel, B.; Economou, A.;

[123074]Munsterberg, A.; Vivian, N.; Goodfellow, P. N.; Lovell-Badge, R.:

[123075]A gene mapping to the sex-determining region of the mouse Y chromosome

[123076]is a member of a novel family of embryonically expressed genes. Nature 346:

[123077]245-250, 1990.

[123078]21840.

[123079]Haqq, C. M.; King, C.-Y.; Ukiyama, E.; Falsafi, S.; Haqq, T. N.;

[123080]Donahoe, P. K.; Weiss, M. A.: Molecular basis of mammalian sexual

[123081]determination: activation of Mullerian inhibiting substance gene expression

[123082]by SRY. Science 266: 1494-1500, 1994.

[123083]21841.

[123084]Ramulu, P.; Kennedy, M.; Xiong, W.-H.; Williams, J.; Cowan, M.;

[123085]Blesh, D.; Yau, K.-W.; Hurley, J. B.; Nathans, J.: Normal light response,

[123086]photoreceptor integrity, and rhodopsin dephosphorylation in mice lacking

[123087]both protein phosphatases with EF hands (PPEF-1 and PPEF-2). Molec.

[123088]Cell. Biol. 21: 8605–8614, 2001.

[123089]21842.

[123090]van de Vosse, E.; Franco, B.; van der Bent, P.; Montini, E.;
Orth,

[123091]U.; Hanauer, A.; Tijmes, N.; van Ommen, G.–J. B.; Ballabio,
A.; den

[123092]Dunnen, J. T.; Bergen, A. A. B.: Exclusion of PPEF as the
gene causing

[123093]X-linked juvenile retinoschisis. Hum. Genet. 101:
235–237, 1997.

[123094]21843.

[123095]Spranger, S.; Schiller, S.; Jauch, A.; Wolff, K.; Rauterberg–Ru-
land,

[123096]I.; Hager, D.; Tariverdian, G.; Troger, J.; Rappold, G.: Leri-
Weill

[123097]syndrome as part of a contiguous gene syndrome at
Xp22.3. Am. J.

[123098]Med. Genet. 83: 367–371, 1999.

[123099]21844.

[123100]Stern, M.–H.; Soulier, J.; Rosenzweig, M.; Nakahara, K.;
Canki–Klain,

[123101]N.; Aurias, A.; Sigaux, F.; Kirsch, I. R.: MTCP-1: a novel gene on

[123102]the human chromosome Xq28 translocated to the T cell receptor alpha/delta

[123103]locus in mature T cell proliferations. *Oncogene* 8: 2475-2483, 1993.

[123104]21845.

[123105]Watanabe, T. K.; Fujiwara, T.; Nakamura, Y.; Hirai, Y.; Maekawa,

[123106]H.; Takahashi, E.: Cloning, expression pattern and mapping to Xq

[123107]of NAP1L3, a gene encoding a peptide homologous to human and yeast

[123108]nucleosome assembly proteins. *Cytogenet. Cell Genet.* 74: 281-285,

[123109]1996.

[123110]21846.

[123111]Prakash, S. K.; Paylor, R.; Jenna, S.; Lamarche-Vane, N.; Armstrong,

[123112]D. L.; Xu, B.; Mancini, M. A.; Zoghbi, H. Y.: Functional analysis

[123113]of ARHGAP6, a novel GTPase-activating protein for RhoA.

Hum. Molec.

[123114]Genet. 9: 477–488, 2000.

[123115]21847.

[123116]Schaefer, L.; Prakash, S.; Zoghbi, H. Y.: Cloning and characterization

[123117]of a novel rho-type GTPase-activating protein gene (ARHGAP6) from

[123118]the critical region for microphthalmia with linear skin defects. Genomics 46:

[123119]268–277, 1997.

[123120]21848.

[123121]Aman, M. J.; Tayebi, N.; Obiri, N. I.; Puri, R. K.; Modi, W. S.;

[123122]Leonard, W. J.: cDNA cloning and characterization of the human interleukin

[123123]13 receptor alpha chain. J. Biol. Chem. 271: 29265–29270, 1996.

[123124]21849.

[123125]Guo, J.; Apiou, F.; Mellerin, M.-P.; Lebeau, B.; Jacques, Y.;

[123126]Minvielle, S.: Chromosome mapping and expression of the human interleukin-13

[123127]receptor. Genomics 42: 141–5, 1997.

[123128]21850.

[123129]Hilton, D. J.; Zhang, J.-G.; Metcalf, D.; Alexander, W. S.; Nicola,

[123130]N. A.; Willson, T. A.: Cloning and characterization of a binding

[123131]subunit of the interleukin 13 receptor that is also a component of

[123132]the interleukin 4 receptor. Proc. Nat. Acad. Sci. 93: 497-501, 1996.

[123133]21851.

[123134]Riggs, J. E.; Schochet, S. S., Jr.; Gutmann, L.; Shanske, S.;

[123135]Neal, W. A.; DiMauro, S.: Lysosomal glycogen storage disease without

[123136]acid maltase deficiency. Neurology 33: 873-877, 1983.

[123137]21852.

[123138]Verloes, A.; Massin, M.; Lombet, J.; Grattagliano, B.; Soyeur,

[123139]D.; Rigo, J.; Koulischer, L.; Van Hoof, F.: Nosology of lysosomal

[123140]glycogen storage diseases without in vitro acid maltase deficiency:

[123141]delineation of a neonatal form. Am. J. Med. Genet. 72:
135–142,

[123142]1997.

[123143]21853.

[123144]Kaye, F. J.; Modi, S.; Ivanovska, I.; Koonin, E. V.; Thress, K.;

[123145]Kubo, A.; Kornbluth, S.; Rose, M. D.: A family of ubiquitin-
like

[123146]proteins binds the ATPase domain of Hsp70-like Stch.
FEBS Lett. 467:

[123147]348–352, 2000.

[123148]21854.

[123149]Kaye, F. J.; Shows, T. B.: Assignment of ubiquilin 2
(UBQLN2)

[123150]to human chromosome xp11.23–p11.1 by GeneBridge ra-
diation hybrids. Cytogenet.

[123151]Cell Genet. 89: 116–117, 2000.

[123152]21855.

[123153]Kleijnen, M. F.; Shih, A. H.; Zhou, P.; Kumar, S.; Soccio, R.
E.;

[123154]Kedersha, N. L.; Gill, G.; Howley, P. M.: The hPLIC proteins
may

[123155]provide a link between the ubiquitination machinery and the proteasome. Molec.

[123156]Cell 6: 409–419, 2000.

[123157]21856.

[123158]Lee, D. K.; Nguyen, T.; Lynch, K. R.; Cheng, R.; Vanti, W. B.;

[123159]Arkhitko, O.; Lewis, T.; Evans, J. F.; George, S. R.; O'Dowd, B. F.

[123160]: Discovery and mapping of ten novel G protein–coupled receptor genes. Gene 275:

[123161]83–91, 2001.

[123162]21857.

[123163]Ades, L. C.; Gedeon, A. K.; Wilson, M. J.; Latham, M.; Partington,

[123164]M. W.; Mulley, J. C.; Nelson, J.; Lui, K.; Sillence, D. O.: Barth

[123165]syndrome: clinical features and confirmation of gene localisation

[123166]to distal Xq28. Am. J. Med. Genet. 45: 327–334, 1993.

[123167]21858.

[123168]Barth, P. G.; Scholte, J. A.; Berden, J. A.; Van der Klei–Van Moorsel,

- [123169]J. M.; Luyt-Houwen, I. E. M.; Van't Veer-Korthof, E. T.; Van der Harten,
- [123170]J. J.; Sobotka-Plojhar, M. A.: An X-linked mitochondrial disease
- [123171]affecting cardiac muscle, skeletal muscle and neutrophil leucocytes. J.
- [123172]Neurol. Sci. 62: 327-355, 1983.
- [123173]21859.
- [123174]Barth, P. G.; Van't Veer-Korthof, E. T.; Van Delden, L.; Van Dam,
- [123175]K.; Van der Harten, J. J.; Kuipers, J. R. G.: An X-linked mitochondrial
- [123176]disease affecting cardiac muscle, skeletal muscle and neutrophil leukocytes.In:
- [123177]Busch, H. F. M.; Jennekens, F. G. I.; Schotte, H. R.: Mitochondria
- [123178]and Muscular Diseases. Beetsterzwaag, The Netherlands: Mefar (pub.)
- [123179]1981. Pp. 161-164.
- [123180]21860.
- [123181]Bione, S.; D'Adamo, P.; Maestrini, E.; Gedeon, A. K.; Bolhuis,

- [123182]P. A.; Toniolo, D.: A novel X-linked gene, G4.5.(sic) is responsible
for Barth syndrome. Nature Genet. 12: 385–389, 1996.
- [123184]21861.
- [123185]Bolhuis, P. A.; Hensels, G. W.; Hulsebos, T. J. M.; Baas, F.; Barth,
P. G.: Mapping of the locus for X-linked cardioskeletal myopathy
with neutropenia and abnormal mitochondria (Barth syndrome) to Xq28. Am. J. Hum. Genet. 48: 481–485, 1991.
- [123188]21862.
- [123190]Chen, Y.-T.; Rettig, W. J.; Yenamandra, A. K.; Kozak, C. A.; Chaganti,
R. S. K.; Posner, J. B.; Old, L. J.: Cerebellar degeneration-related
antigen: a highly conserved neuroectodermal marker mapped to chromosomes
X in human and mouse. Proc. Nat. Acad. Sci. 87: 3077–3081, 1990.
- [123194]21863.

[123195]Dropcho, E. J.; Chen, Y. T.; Posner, J. B.; Old, L. J.: Cloning
[123196]of a brain protein identified by autoantibodies from a pa-
tient with
[123197]paraneoplastic cerebellar degeneration. Proc. Nat. Acad.
Sci. 84:
[123198]4552-4556, 1987.
[123199]21864.

[123200]Furieux, H. M.; Rosenblum, M. K.; Dalmau, J.; Wong, E.;
Woodruff,
[123201]P.; Graus, F.; Posner, J. B.: Selective expression of Purk-
inje-cell
[123202]antigens in tumor tissue from patients with paraneoplastic
cerebellar
[123203]degeneration. New Eng. J. Med. 322: 1844-1851, 1990.
[123204]21865.

[123205]Furieux, H. M.; Wong, E.; Posner, J. B.: Isolation of cDNA
clones
[123206]encoding the major Yo paraneoplastic antigen. (Abstract)
Neurology 40
[123207](suppl. 1): 166 only, 1990.
[123208]21866.

[123209]Hirst, M. C.; Bell, M. V.; MacKinnon, R. N.; Watson, J. E. V.;

[123210]Callen, D.; Sutherland, G.; Dahl, N.; Patterson, M. N.;

Schwartz,

[123211]C.; Ledbetter, D.; Ledbetter, S.; Davies, K. E.: Mapping of a
cerebellar

[123212]degeneration related protein and DXS304 around the
fragile site. Am.

[123213]J. Med. Genet. 38: 354–356, 1991.

[123214]21867.

[123215]Rettig, W. J.; Chen, Y. T.; Yenamandra, A. K.; Chaganti, R.
S.

[123216]K.; Dropcho, E. J.; Posner, J. B.; Old, L. J.: Assignment of
the

[123217]cerebellar degeneration–related autoantigen CDR to hu-
man chromosome

[123218]X. (Abstract) Cytogenet. Cell Genet. 46: 681 only, 1987.

[123219]21868.

[123220]Siniscalco, M.; Oberle, I.; Melis, P.; Alhadeff, B.; Murray, J.;

[123221]Filippi, G.; Mattioni, T.; Chen, Y.–T.; Furneaux, H.; Old, L.
J.;

[123222]Posner, J. B.: Physical and genetic mapping of the CDR

gene with

[123223]particular reference to its position with respect to the
FRAXA site. Am.

[123224]J. Med. Genet. 38: 357–362, 1991.

[123225]21869.

[123226]Siniscalco, M.; Alhadeff, B.; Kekish, O.; Melis, P.; Maltioni,

[123227]T.; Filippi, G.; Murray, J. C.; Oberle, I.; Chen, Y. T.; Old, L.
J.;

[123228]Posner, J. B.: The CDR gene, located at the upper bound-
ary of the

[123229]FRAXA, is tightly linked to DXS98 and DXS105 RFLPs,
moderately to

[123230]the FRA(X)-syndrome and loosely to the cluster of
DXS15–DXS52 RFLPs.

[123231](Abstract) Cytogenet. Cell Genet. 51: 1081 only, 1989.

[123232]21870.

[123233]Knight, J. C.; Renwick, P. J.; Downing, J. R.; Okuda, T.:
Physical

[123234]linkage of the cdc2-related gene (PCTK1) and the ubiqui-
tin-activating

[123235]enzyme E1 gene (UBE1) on human Xp11.3. Cytogenet. Cell

Genet. 71:

[123236]155–157, 1995.

[123237]21871.

[123238]Aarskog, D.: A familial syndrome of short stature associated with

[123239]facial dysplasia and genital anomalies. J. Pediat. 77:
856–861,

[123240]1970.

[123241]21872.

[123242]Baldellou, A.; Galve, L.; Bassecourt, M.: Risk of medullary
damage

[123243]in Aarskog–Scott syndrome. (Abstract) Clin. Genet. 23:
225 only,

[123244]1983.

[123245]21873.

[123246]Bawle, E.; Tyrkus, M.; Lipman, S.; Bozimowski, D.: Aarskog
syndrome:

[123247]full male and female expression associated with an X–
autosome translocation. Am.

[123248]J. Med. Genet. 17: 595–602, 1984.

[123249]21874.

[123250]Berman, P. A.; Desjardins, C.; Fraser, F. C.: Inheritance of the

[123251]Aarskog syndrome. Birth Defects Orig. Art. Ser. X(7): 151-159, 1974.

[123252]21875.

[123253]Buechner, M.; Hall, D. H.; Bhatt, H.; Hedgecock, E. M.: Cystic

[123254]canal mutants in *Caenorhabditis elegans* are defective in the apical

[123255]membrane domain of the renal (excretory) cell. Dev. Biol. 214: 227-241,

[123256]1999.

[123257]21876.

[123258]Casteels, M.; Samain, H.; Penninckx, F.; Coremans, G.; Beirinckx,

[123259]J.; Fryns, J. P.: Megadolichosigmoid in a young male with Aarskog

[123260]syndrome. Genet. Counsel. 5: 81-83, 1994.

[123261]21877.

[123262]Estrada, L.; Caron, E.; Gorski, J. L.: Fgd1, the Cdc42 guanine

- [123263]nucleotide exchange factor responsible for faciogenital dysplasia,
- [123264]is localized to the subcortical actin cytoskeleton and Golgi membrane. Hum.
- [123265]Molec. Genet. 10: 485–495, 2001.
- [123266]21878.
- [123267]Fernandez, I.; Tsukahara, M.; Mito, H.; Yoshii, H.; Uchida, M.;
- [123268]Matsuo, K.; Kajii, T.: Congenital heart defects in Aarskog syndrome. Am.
- [123269]J. Med. Genet. 50: 318–322, 1994.
- [123270]21879.
- [123271]Fryns, J.–P.: Dolichomegasigmoid in Aarskog syndrome. (Letter) Am.
- [123272]J. Med. Genet. 45: 122 only, 1993.
- [123273]21880.
- [123274]Fryns, J. P.: Aarskog syndrome: the changing phenotype with age. Am.
- [123275]J. Med. Genet. 43: 420–427, 1992.
- [123276]21881.

[123277]Fryns, J. P.; Macken, J.; Vinken, L.; Igodt-Ameye, L.; van
den

[123278]Berghe, H.: The Aarskog syndrome. Hum. Genet. 42:
129-135, 1978.

[123279]21882.

[123280]Funderburk, S. J.; Crandall, B. F.: The Aarskog syndrome in
three

[123281]brothers. Clin. Genet. 6: 119-124, 1974.

[123282]21883.

[123283]Furukawa, C. T.; Hall, B. D.; Smith, D. W.: The Aarskog
syndrome. J.

[123284]Pediat. 81: 1117-1122, 1972.

[123285]21884.

[123286]Gao, J.; Estrada, L.; Cho, S.; Ellis, R. E.; Gorski, J. L.: The

[123287]Caenorhabditis elegans homolog of FGD1, the human
Cdc42 GEF gene responsible

[123288]for faciogenital dysplasia, is critical for excretory cell mor-
phogenesis. Hum.

[123289]Molec. Genet. 10: 3049-3062, 2001.

[123290]21885.

- [123291]Glover, T. W.; Verga, V.; Rafael, J.; Barcroft, C.; Gorski, J.
[123292]L.; Bawle, E. V.; Higgins, J. V.: Translocation breakpoint in
Aarskog
[123293]syndrome maps to Xp11.21 between ALAS2 and DXS323.
Hum. Molec. Genet. 2:
[123294]1717–1718, 1993.
[123295]21886.
[123296]Grier, R. E.; Farrington, F. H.; Kendig, R.; Mamunes, P.:
Autosomal
[123297]dominant inheritance of the Aarskog phenotype.
(Abstract) Am. J.
[123298]Hum. Genet. 33: 64A only, 1981.
[123299]21887.
[123300]Hoo, J. J.: The Aarskog (facio–digito–genital) syndrome.
Clin.
[123301]Genet. 16: 269–276, 1979.
[123302]21888.
[123303]Kodama, M.; Fujimoto, S.; Namikawa, T.; Matsuda, I.:
Aarskog
[123304]syndrome with isolated growth hormone deficiency. Eu-
rop. J. Pediat. 135:

[123305]273–276, 1981.

[123306]21889.

[123307]Lebel, R. R.; May, M.; Pouls, S.; Lubs, H. A.; Stevenson, R. E.;

[123308]Schwartz, C. E.: Non-syndromic X-linked mental retardation associated

[123309]with a missense mutation (P312L) in the FGD1 gene. Clin. Genet. 61:

[123310]139–145, 2002.

[123311]21890.

[123312]Logie, L. J.; Porteous, M. E. M.: Intelligence and development

[123313]in Aarskog syndrome. Arch. Dis. Child. 79: 359–360, 1998.

[123314]21891.

- [123315] Mikelsaar, R. V.-A.; Lurie, I. W.: Atypical case of Aarskog syndrome. J. Med. Genet. 29: 349-350, 1992.
- [123316] 21892.
- [123317] Nielsen, K. B.: Aarskog syndrome in a Danish family: an illustration of the need for dysmorphology in paediatrics. Clin. Genet. 33: 315-317, 1988.
- [123318] 21893.
- [123319] Oberiter, V.; Lovrencic, M. K.; Schmutzer, L.; Kraus, O.: The Aarskog syndrome. Acta Paediat. Scand. 69: 567-570, 1980.
- [123320] 21894.
- [123321] Orrico, A.; Galli, L.; Falciani, M.; Bracci, M.; Cavaliere, M. L.; Rinaldi, M. M.; Musacchio, A.; Sorrentino, V.: A mutation in the pleckstrin homology (PH) domain of the FGD1 gene in an Italian family with faciogenital dysplasia (Aarskog-Scott syn-

drome). FEBS

[123329]Lett. 478: 216–220, 2000.

[123330]21895.

[123331]Pasteris, N. G.; Cadle, A.; Logie, L. J.; Porteous, M. E. M.;

[123332]Schwartz, C. E.; Stevenson, R. E.; Glover, T. W.; Wilroy, R.
S.; Gorski,

[123333]J. L.: Isolation and analysis of the faciogenital dysplasia
(Aarskog–Scott

[123334]syndrome) gene: a putative, rho/rac guanine nucleotide
exchange factor. Cell 79:

[123335]669–678, 1994.

[123336]21896.

[123337]Pasteris, N. G.; de Gouyon, B.; Cadle A. B.; Campbell, K.;
Herman,

[123338]G. E.; Gorski, J. L.: Cloning and regional localization of the
mouse

[123339]faciogenital dysplasia (Fgd1) gene. Mammalian Genome 6:
658–661,

[123340]1995.

[123341]21897.

[123342]Pedersen, J. C.; Fryns, J. P.; Bracke, P.; Geeraert, M.; Van

Den

[123343]Berghe, H.: The Aarskog syndrome. Ann. Genet. 23: 108–110, 1980.

[123344]21898.

[123345]Porteous, M. E. M.; Goudie, D. R.: Aarskog syndrome. J. Med.

[123346]Genet. 28: 44–47, 1991.

[123347]21899.

[123348]Rafael, J.; Verga, V.; Hall, B.; Burright, E.; Gorski, J.; Bawle,

[123349]J.; Higgins, J. V.; Glover, T. W.: Assignment of the translocation

[123350]breakpoint in a patient with Aarskog syndrome to Xp11.21. (Abstract) Am.

[123351]J. Hum. Genet. 51 (suppl.): A116 only, 1992.

[123352]21900.

[123353]Schwartz, C. E.; Gillessen-Kaesbach, G.; May, M.; Cappa, M.; Gorski,

[123354]J.; Steindl, K.; Neri, G.: Two novel mutations confirm FGD1 is responsible

[123355]for the Aarskog syndrome. Europ. J. Hum. Genet. 8: 869–874, 2000.

[123356]21901.

[123357]Scott, C. I., Jr.: Unusual facies, joint hypermobility, genital
[123358]anomaly and short stature: a new dysmorphic syndrome.

Birth Defects

[123359]Orig. Art. Ser. VII(6): 240–246, 1971.

[123360]21902.

[123361]Asghar, M.; Nevin, N. C.; Beattie, E. D.; McManus, D.;
Roberts,

[123362]G. M. A.; Phillips, J. A.: Contiguous gene deletion syn-
drome involving

[123363]glycerol kinase and Duchenne muscular dystrophy loci. J.
Inherit.

[123364]Metab. Dis. 22: 933–935, 1999.

[123365]21903.

[123366]Balducci, R.; Municchi, G.; Toscano, V.; Mangiantini, A.;
Sabatini,

[123367]R.; Caiola, S.; Tedeschi, B.; Orlandi, L.; Boscherini, B.:
Complex

[123368]glycerol kinase deficiency: an unusual cause of salt-
wasting in males. Clin.

[123369]Endocr. 42: 437–439, 1995.

[123370]21904.

[123371]Bartley, J. A.; Patil, S.; Davenport, S.; Goldstein, D.; Pickens,

[123372]J.: Duchenne muscular dystrophy, glycerol kinase deficiency, and

[123373]adrenal insufficiency associated with Xp21 interstitial deletion. J.

[123374]Pediat. 108: 189-192, 1986.

[123375]21905.

[123376]Borresen, A.-L.; Hellerud, C.; Moller, P.; Sovik, O.; Berg, K.

[123377]: Prenatal diagnosis of glycerol-kinase deficiency associated with

[123378]a DNA deletion on the short arm of the X chromosome. Clin. Genet. 32:

[123379]254-259, 1987.

[123380]21906.

[123381]Darras, B. T.; Francke, U.: Myopathy in complex glycerol kinase

[123382]deficiency patients is due to 3-prime deletions of the dystrophin

[123383]gene. Am. J. Hum. Genet. 43: 126-130, 1988.

[123384]21907.

[123385]Davies, K. E.; Patterson, M. N.; Kenwrick, S. J.; Bell, M. V.;

[123386]Sloan, H. R.; Westman, J. A.; Elsas, L. J., II; Mahan, J.: Fine
mapping

[123387]of glycerol kinase deficiency and congenital adrenal hy-
poplasia within

[123388]Xp21 on the short arm of the human X chromosome. Am.
J. Med. Genet. 29:

[123389]557-564, 1988.

[123390]21908.

[123391]Dipple, K. M.; Zhang, Y.-H.; Huang, B.-L.; McCabe, L. L.;
Dallongeville,

[123392]J.; Inokuchi, T.; Kimura, M.; Marx, H. J.; Roederer, G. O.;
Shih,

[123393]Y.; Yamaguchi, S.; Yoshida, I.; McCabe, E. R. B.: Glycerol
kinase

[123394]deficiency: evidence for complexity in a single gene disor-
der. Hum.

[123395]Genet. 109: 55-62, 2001.

[123396]21909.

[123397]Arias, J.; Alberts, A. S.; Brindle, P.; Claret, F. X.; Smeal, T.;

[123398]Karin, M.; Feramisco, J.; Montminy, M.: Activation of cAMP and mitogen

[123399]responsive genes relies on a common nuclear factor. Nature 370:

[123400]226–229, 1994.

[123401]21910.

[123402]Blough, R. I.; Petrij, F.; Dauwerse, J. G.; Milatovich–Cherry,

[123403]A.; Weiss, L.; Saal, H. M.; Rubinstein, J. H.: Variation in microdeletions

[123404]of the cyclic AMP–responsive element–binding protein gene at chromosome

[123405]band 16p13.3 in the Rubinstein–Taybi syndrome. Am. J. Med. Genet. 90:

[123406]29–34, 2000.

[123407]21911.

[123408]Chen, X.–N.; Korenberg, J. R.: Localization of human CREBBP (CREB

[123409]binding protein) to 16p13.3 by fluorescence in situ hybridization. Cytogenet.

[123410]Cell Genet. 71: 56–57, 1995.

[123411]21912.

- [123412]Chrivia, J. C.; Kwok, R. P. S.; Lamb, N.; Hagiwara, M.;
Montminy,
[123413]M. R.; Goodman, R. H.: Phosphorylated CREB binds specif-
ically to
[123414]the nuclear protein CBP. Nature 365: 855–859, 1993.
[123415]21913.
- [123416]Borrow, J.; Stanton, V. P., Jr.; Andresen, J. M.; Becher, R.;
Behm,
[123417]F. G.; Chaganti, R. S. K.; Civin, C. I.; Distech, C.; Dube, I.;
Frischauf,
[123418]A. M.; Horsman, D.; Mitelman, F.; Volinia, S.; Watmore, A.
E.; Housman,
[123419]D. E.: The translocation t(8;16)(p11;p13) of acute myeloid
leukaemia
[123420]fuses a putative acetyltransferase to the CREB-binding
protein. Nature
[123421]Genet. 14: 33–41, 1996.
[123422]21914.
- [123423]Demarest, S. J.; Martinez–Yamout, M.; Chung, J.; Chen, H.;
Xu,
[123424]W.; Dyson, H. J.; Evans, R. M.; Wright, P. E.: Mutual syner-
gistic

- [123425]folding in recruitment of CBP/p300 by p160 nuclear receptor coactivators. *Nature* 415:
[123426]549–553, 2002.
- [123427]21915.
- [123428]Eckner, R.; Ewen, M. E.; Newsome, D.; Gerdes, M.; De-
Caprio, J.
[123429]A.; Lawrence, J. B.; Livingston, D. M.: Molecular cloning
and functional
[123430]analysis of the adenovirus E1A-associated 300-kD protein
(p300) reveals
[123431]a protein with properties of a transcriptional adaptor.
Genes Dev. 15:
[123432]869–884, 1994.
- [123433]21916.
- [123434]Giles, R. H.; Dauwerse, H. G.; van Ommen, G.-J. B.; Breun-
ing, M.
[123435]H.: Do human chromosomal bands 16p13 and 22q11–13
share ancestral
[123436]origins? (Letter) *Am. J. Hum. Genet.* 63: 1240–1242, 1998.
- [123437]21917.
- [123438]Giles, R. H.; Petrij, F.; Dauwerse, H. G.; den Hollander, A.

I.;

[123439]Lushnikova, T.; van Ommen, G.-J. B.; Goodman, R. H.;
Deaven, L. L.;

[123440]Doggett, N. A.; Peters, D. J.; Breuning, M. H.: Construction
of a

[123441]1.2-Mb contig surrounding, and molecular analysis of, the
human CREB-binding

[123442]protein (CBP/CREBBP) gene on chromosome 16p13.3. Ge-
nomics 42: 96-114,

[123443]1997.

[123444]21918.

[123445]Gonzalez, G. A.; Montminy, M. R.: Cyclic AMP stimulates
somatostatin

[123446]gene transcription by phosphorylation of CREB at serine
133. Cell 59:

[123447]675-680, 1989.

[123448]21919.

[123449]Kung, A. L.; Rebel, V. I.; Bronson, R. T.; Ch'ng, L.-E.; Sieff,

[123450]C. A.; Livingston, D. M.; Yao, T.-P.: Gene dose-dependent
control

[123451]of hematopoiesis and hematologic tumor suppression by

CBP. Genes

[123452]Dev. 14: 272–277, 2000.

[123453]21920.

[123454]Kwok, R. P. S.; Lundblad, J. R.; Chrivia, J. C.; Richards, J.

[123455]P.; Bachinger, H. P.; Brennan, R. G.; Roberts, S. G. E.;

Green, M.

[123456]R.; Goodman, R. H.: Nuclear protein CBP is a coactivator
for the

[123457]transcription factor CREB. Nature 370: 223–226, 1994.

[123458]21921.

[123459]Lin, C. H.; Hare, B. J.; Wagner, G.; Harrison, S. C.; Maniatis,

[123460]T.; Fraenkel, E.: A small domain of CBP/p300 binds di-
verse proteins:

[123461]solution structure and functional studies. Molec. Cell 8:
581–590,

[123462]2001.

[123463]21922.

[123464]Lundblad, J. R.; Kwok, R. P. S.; Lurance, M. E.; Harter, M.

L.;

[123465]Goodman, R. H.: Adenoviral E1A-associated protein p300

as a functional

[123466]homologue of the transcriptional co-activator CBP. Nature
374: 85–88,

[123467]1995.

[123468]21923.

[123469]Murata, T.; Kurokawa, R.; Krones, A.; Tatsumi, K.; Ishii, M.;

[123470]Taki, T.; Masuno, M.; Ohashi, H.; Yanagisawa, M.; Rosen-
feld, M. G.;

[123471]Glass, C. K.; Hayashi, Y.: Defect of histone acetyltrans-
ferase activity

[123472]of the nuclear transcriptional coactivator CBP in Rubin-
stein–Taybi

[123473]syndrome. Hum. Molec. Genet. 10: 1071–1076, 2001.

[123474]21924.

[123475]Nordheim, A.: CREB takes CBP to tango. Nature 370:
177–178,

[123476]1994.

[123477]21925.

[123478]Ohno, S.: Patterns in genome evolution. Curr. Opin. Genet.
Dev. 3:

[123479]911–914, 1993.

[123480]21926.

[123481]Oike, Y.; Hata, A.; Mamiya, T.; Kaname, T.; Noda, Y.;
Suzuki,

[123482]M.; Yasue, H.; Nabeshima, T.; Araki, K.; Yamamura, K.:
Truncated

[123483]CBP protein leads to classical Rubinstein–Taybi syndrome
phenotypes

[123484]in mice: implications for a dominant–negative mechanism.
Hum. Molec.

[123485]Genet. 8: 387–396, 1999.

[123486]21927.

[123487]Kamb, A.; Shattuck–Eidens, D.; Eeles, R.; Liu, Q.; Gruis, N.
A.;

[123488]Ding, W.; Hussey, C.; Tran, T.; Miki, Y.; Weaver–Feldhaus,
J.; McClure,

[123489]M.; Aitken, J. F.; Anderson, D. E.; Bergman, W.; Frants, R.;
Goldgar,

[123490]D. E.; Green, A.; MacLennan, R.; Martin, N. G.; Meyer, L. J.;
Youl,

[123491]P.; Zone, J. J.; Skolnick, M. H.; Cannon–Albright, L. A.:
Analysis

[123492]of the p16 gene (CDKN2) as a candidate for the chromo–

some 9p melanoma

[123493]susceptibility locus. Nature Genet. 8: 22–26, 1994.

[123494]21928.

[123495]Riesewijk, A. M.; Blagitko, N.; Schinzel, A. A.; Hu, L.;
Schulz,

[123496]U.; Hamel, B. C. J.; Ropers, H.–H.; Kalscheuer, V. M.: Evi-
dence against

[123497]a major role of PEG1/MEST Silver–Russell syndrome. Eu-
rop. J. Hum.

[123498]Genet. 6: 114–120, 1998.

[123499]21929.

[123500]Riesewijk, A. M.; Hu, L.; Schulz, U.; Tariverdian, G.;
Hoglund,

[123501]P.; Kere, J.; Ropers, H.–H.; Kalscheuer, V. M.: Monoallelic
expression

[123502]of human PEG1/MEST is paralleled by parent–specific
methylation in

[123503]fetuses. Genomics 42: 236–244, 1997.

[123504]21930.

[123505]Sado, T.; Nakajima, N.; Tada, M.; Takagi, N.: A novel
mesoderm–specific

[123506]cDNA isolated from a mouse embryonal carcinoma cell
line. Dev. Growth

[123507]Differ. 35: 551–560, 1993.

[123508]21931.

[123509]Kamijo, T.; Zindy, F.; Roussel, M. F.; Quelle, D. E.; Down-
ing,

[123510]J. R.; Ashmun, R. A.; Grosveld, G.; Sherr, C. J.: Tumor sup-
pression

[123511]at the mouse INK4a locus mediated by the alternative
reading frame

[123512]product p19(ARF). Cell 91: 649–659, 1997.

[123513]21932.

[123514]Koh, J.; Enders, G. H.; Dynlacht, B. D.; Harlow, E.: Tumour-
derived

[123515]p16 alleles encoding proteins defective in cell-cycle inhi-
bition. Nature 375:

[123516]506–510, 1995.

[123517]21933.

[123518]Komiya, A.; Suzuki, H.; Aida, S.; Yatani, R.; Shimazaki, J.:

[123519]Mutational analysis of CDKN2 (CDK4I/MTS1) gene in tis-
sues and cell

[123520]lines of human prostate cancer. Jpn. J. Cancer Res. 86:
622–625,

[123521]1995.

[123522]21934.

[123523]Krimpenfort, P.; Quon, K. C.; Mooi, W. J.; Loonstra, A.;
Berns,

[123524]A.: Loss of p16(Ink4a) confers susceptibility to metastatic
melanoma

[123525]in mice. Nature 413: 83–85, 2001.

[123526]21935.

[123527]Labarriere, N.; Selvais, P. L.; Lemaigre, F. P.; Michel, A.;
Maiter,

[123528]D. M.; Rousseau, G. G.: A novel transcriptional activator
originating

[123529]from an upstream promoter in the human growth hor-
mone gene. J. Biol.

[123530]Chem. 270: 19205–19208, 1995.

[123531]21936.

[123532]Little, M.; Wainwright, B.: Methylation and p16: suppress-
ing

[123533]the suppressor. Nature Med. 1: 633–634, 1995.

[123534]21937.

[123535]Liu, L.; Dilworth, D.; Gao, L.; Monzon, J.; Summers, A.;
Lassam,

[123536]N.; Hogg, D.: Mutation of the CDKN2A 5-prime UTR cre-
ates an aberrant

[123537]initiation codon and predisposes to melanoma. Nature
Genet. 21:

[123538]128–132, 1999.

[123539]21938.

[123540]Liu, L.; Lassam, N. J.; Slingerland, J. M.; Bailey, D.; Cole,

[123541]D.; Jenkins, R.; Hogg, D.: Germline p16(INK4A) mutation
and protein

[123542]dysfunction in a family with inherited melanoma. Onco-
gene 11: 405–412,

[123543]1995.

[123544]21939.

[123545]Lukas, J.; Parry, D.; Aagaard, L.; Mann, D. J.; Bartkova, J.;

[123546]Strauss, M.; Peters, G.; Bartek, J.: Retinoblastoma-pro-
tein-dependent

[123547]cell-cycle inhibition by the tumour suppressor p16. Na-
ture 375:

[123548]503–506, 1995.

[123549]21940.

[123550]MacGeoch, C.; Bishop, J. A.; Bataille, V.; Bishop, D. T.;
Frischauf,

[123551]A. M.; Meloni, R.; Cuzick, J.; Pinney, E.; Spurr, N. K.: Ge-
netic

[123552]heterogeneity in familial malignant melanoma. Hum.
Molec. Genet. 3:

[123553]2195–2200, 1994.

[123554]21941.

[123555]MacKie, R. M.; Andrew, N.; Lanyon, W. G.; Connor, J. M.:
CDKN2A

[123556]germline mutations in U.K. patients with familial
melanoma and multiple

[123557]primary melanomas. J. Invest. Derm. 111: 269–272, 1998.

[123558]21942.

[123559]Mantelli, M.; Barile, M.; Ciotti, P.; Ghiorzo, P.; Lantieri, F.;

[123560]Pastorino, L.; Catricala, C.; Della Torre, G.; Folco, U.;
Grammatico,

[123561]P.; Padovani, L.; Pasini, B.; and 9 others: High prevalence
of

- [123562]the G101W germline mutation in the CDKN2A (P16ink4a) gene in 62 Italian
- [123563]malignant melanoma families. Am. J. Med. Genet. 107: 214–221, 2002.
- [123564]21943.
- [123565]Merlo, A.; Herman, J. G.; Mao, L.; Lee, D. J.; Gabrielson, E.;
- [123566]Burger, P. C.; Baylin, S. B.; Sidransky, D.: 5-prime CpG island methylation
- [123567]is associated with transcriptional silencing of the tumour suppressor
- [123568]p16/CDKN2/MTS1 in human cancers. Nature Med. 1: 686–692, 1995.
- [123569]21944.
- [123570]Monzon, J.; Liu, L.; Brill, H.; Goldstein, A. M.; Tucker, M. A.;
- [123571]From, L.; McLaughlin, J.; Hogg, D.; Lassam, N. J.: CDKN2A mutations
- [123572]in multiple primary melanomas. New Eng. J. Med. 338: 879–887, 1998.
- [123573]21945.
- [123574]Mori, T.; Miura, K.; Aoki, T.; Nishihira, T.; Mori, S.; Nakamura,

[123575]Y.: Frequent somatic mutation of the MTS1/CDK41
(multiple tumor suppressor/cyclin-dependent
[123576]kinase 4 inhibitor) gene in esophageal squamous cell car-
cinoma. Cancer
[123577]Res. 54: 3396–3397, 1994.
[123578]21946.

[123579]Nobori, T.; Miura, K.; Wu, D. J.; Lois, A.; Takabayashi, K.;
Carson,
[123580]D. A.: Deletions of the cyclin-dependent kinase-4 in-
hibitor gene
[123581]in multiple human cancers. Nature 368: 753–756, 1994.
[123582]21947.

[123583]Ogawa, S.; Hirano, N.; Sato, N.; Takahashi, T.; Hangaishi,
A.;
[123584]Tanaka, K.; Kurokawa, M.; Tanaka, T.; Mitani, K.; Yazaki,
Y.; Hirai,
[123585]H.: Homozygous loss of the cyclin-dependent kinase
4-inhibitor (p16)
[123586]gene in human leukemias. Blood 84: 2431–2435, 1994.
[123587]21948.

[123588]Ohhara, M.; Esumi, M.; Kurosu, Y.: Activation but not inac-

tivation

[123589] of the MTS1 gene is associated with primary colorectal carcinomas. Biochem.

[123590] Biophys. Res. Commun. 226: 791–795, 1996.

[123591] 21949.

[123592] Okuda, T.; Shurtleff, S. A.; Valentine, M. B.; Raimondi, S. C.;

[123593] Head, D. R.; Behm, F.; Curcio-Brint, A. M.; Liu, Q.; Pui, C.-H.; Sherr,

[123594] C. J.; Beach, D.; Look, A. T.; Downing, J. R.: Frequent deletion

[123595] of p16(INK4a)/MTS1 and p15(INK4b)/MTS2 in pediatric acute lymphoblastic

[123596] leukemia. Blood 85: 2321–2330, 1995.

[123597] 21950.

[123598] Petronzelli, F.; Sollima, D.; Coppola, G.; Martini-Neri, M. E.;

[123599] Neri, G.; Genuardi, M.: CDKN2A germline splicing mutation affecting

[123600] both P16(ink4) and P14(arf) RNA processing in a melanoma/neurofibroma

[123601] kindred. Genes Chromosomes Cancer 31: 398–401, 2001.

[123602]21951.

[123603]Petty, E. M.; Gibson, L. H.; Fountain, J. W.; Bolognia, J. L.;

[123604]Yang-Feng, T. L.; Housman, D. E.; Bale, A. E.: Molecular
definition

[123605]of a chromosome 9p21 germ-line deletion in a woman
with multiple melanomas

[123606]and a plexiform neurofibroma: implications for 9p tumour
suppressor

[123607]gene(s). Am. J. Hum. Genet. 53: 96–104, 1993.

[123608]21952.

[123609]Pilon, C.; Pistorello, M.; Moscon, A.; Altavilla, G.; Pagotto,

[123610]U.; Boscaro, M.; Fallo, F.: Inactivation of the p16 tumor
suppressor

[123611]gene in adrenocortical tumors. J. Clin. Endocr. Metab. 84:
2776–2779,

[123612]1999.

[123613]21953.

[123614]Pomerantz, J.; Schreiber-Agus, N.; Liegeois, N. J.; Silver-
man,

[123615]A.; Alland, L.; Chin, L.; Potes, J.; Chen, K.; Orlow, I.; Lee,
H.-W.;

[123616]Cordon-Cardo, C.; DePinho, R. A.: The Ink4a tumor suppressor gene

[123617]product, p19(Arf), interacts with MDM2 and neutralizes MDM2's inhibition

[123618]of p53. Cell 92: 713-723, 1998.

[123619]21954.

[123620]Jayakumar, A.; Chirala, S. S.; Chinault, A. C.; Baldini, A.; Abu-Elheiga,

[123621]L.; Wakil, S. J.: Isolation and chromosomal mapping of genomic clones

[123622]encoding the human fatty acid synthase gene. Genomics 23: 420-424,

[123623]1994.

[123624]21955.

[123625]Jayakumar, A.; Tai, M.-H.; Huang, W.-Y.; Al-Feel, W.; Hsu, M.;

[123626]Abu-Elheiga, L.; Chirala, S. S.; Wakil, S. J.: Human fatty acid synthase:

[123627]properties and molecular cloning. Proc. Nat. Acad. Sci. 92: 8695-8699,

[123628]1995.

[123629]21956.

[123630]Loftus, T. M.; Jaworsky, D. E.; Frehywot, G. L.; Townsend, C. A.;

[123631]Ronnett, G. V.; Lane, M. D.; Kuhajda, F. P.: Reduced food intake

[123632]and body weight in mice treated with fatty acid synthase inhibitors. Science 288:

[123633]2379–2381, 2000.

[123634]21957.

[123635]Wakil, S. J.: Fatty acid synthase, a proficient multifunctional

[123636]enzyme. Biochemistry 28: 4523–4530, 1989.

[123637]21958.

[123638]Ye, Q.; Chung, L. W. K.; Li, S.; Zhau, H. E.: Identification of

[123639]a novel FAS/ER- α fusion transcript expressed in human cancer cells. Biochim.

[123640]Biophys. Acta 1493: 373–377, 2000.

[123641]21959.

[123642]Argeson, A. C.; Druck, T.; Veronese, M. L.; Knopf, J. L.; Buchberg,

- [123643]A. M.; Huebner, K.; Siracusa, L. D.: Phospholipase C gamma-2 (Plcg2) and phospholipase C gamma-1 (Plcg1) map to distinct regions in the human and mouse genomes. *Genomics* 25: 29-35, 1995.
- [123646]21960.
- [123647]Hernandez, D.; Egan, S. E.; Yulug, I. G.; Fisher, E. M. C.: Mapping the gene that encodes phosphatidylinositol-specific phospholipase C-gamma-2 in the human and the mouse. *Genomics* 23: 504-507, 1994.
- [123650]21961.
- [123651]Kang, J. S.; Kohlhuber, F.; Hug, H.; Marme, D.; Eick, D.; Ueffing, M.: Cloning and functional analysis of the hematopoietic cell-specific phospholipase C-gamma-2 promoter. *FEBS Lett.* 399: 14-20, 1996.
- [123654]21962.
- [123655]Dumont, D. J.; Anderson, L.; Breitman, M. L.; Duncan, A. M. V.

- [123656]: Assignment of the endothelial-specific protein receptor tyrosine kinase gene (TEK) to human chromosome 9p21. Genomics 23: 512-513, 1994.
- [123659]21963.
- [123660]Dumont, D. J.; Yamaguchi, T. P.; Conlon, R. A.; Rossant, J.; Breitman, M. L.: tek, a novel tyrosine kinase gene located on mouse chromosome 4, is expressed in endothelial cells and their presumptive precursors. Oncogene 7: 1471-1480, 1992.
- [123664]21964.
- [123665]Delot, E.; King, L. M.; Briggs, M. D.; Wilcox, W. R.; Cohn, D. H.: Trinucleotide expansion mutations in the cartilage oligomeric matrix protein (COMP) gene. Hum. Molec. Genet. 8: 123-128, 1999.
- [123668]21965.
- [123669]Oliner, J. D.; Andresen, J. M.; Hansen, S. K.; Zhou, S.; Tjian,

[123670]R.: SREBP transcriptional activity is mediated through an interaction

[123671]with the CREB-binding protein. *Genes Dev.* 10: 2903–2911, 1996.

[123672]21966.

[123673]Panagopoulos, I.; Fioretos, T.; Isaksson, M.; Samuelsson, U.;

[123674]Billstrom, R.; Strombeck, B.; Mitelman, F.; Johansson, B.: Fusion

[123675]of the MORF and CBP genes in acute myeloid leukemia with the t(10;16)(q22;p13). *Hum.*

[123676]*Molec. Genet.* 10: 395–404, 2001.

[123677]21967.

[123678]Petrij, F.; Dorsman, J. C.; Dauwerse, H. G.; Giles, R. H.; Peeters,

[123679]T.; Hennekam, R. C. M.; Breuning, M. H.; Peters, D. J. M.: Rubinstein–Taybi

[123680]syndrome caused by a de novo reciprocal translocation t(2;16)(q36.3;p13.3). *Am.*

[123681]*J. Med. Genet.* 92: 47–52, 2000.

[123682]21968.

- [123683]Sobulo, O. M.; Borrow, J.; Tomek, R.; Reshmi, S.; Harden, A.;
- [123684]Schlegelberger, B.; Housman, D.; Doggett, N. A.; Rowley, J. D.; Zeleznik-Le, N. J.: MLL is fused to CBP, a histone acetyltransferase, in therapy-related acute myeloid leukemia with a t(11;16)(q23;p13.3). Proc. Nat. Acad. Sci. 94: 8732-8737, 1997.
- [123688]21969.
- [123689]Taine, L.; Goizet, C.; Wen, Z. Q.; Petrij, F.; Breuning, M. H.; Ayme, S.; Saura, R.; Arveiler, B.; Lacombe, D.: Submicroscopic deletion of chromosome 16p13.3 in patients with Rubinstein-Taybi syndrome. Am. J. Med. Genet. 78: 267-270, 1998.
- [123693]21970.
- [123694]Tini, M.; Benecke, A.; Um, S.-J.; Torchia, J.; Evans, R. M.; Chambon, P.: Association of CBP/p300 acetylase and thymine DNA glycosylase

[123696]links DNA repair and transcription. Molec. Cell 9:
265–277, 2002.

[123697]21971.

[123698]Weaver, B. K.; Kumar, K. P.; Reich, N. C.: Interferon regula-
tory

[123699]factor 3 and CREB-binding protein/p300 are subunits of
double-stranded

[123700]RNA-activated transcription factor DRAF1. Molec. Cell.
Biol. 18:

[123701]1359–1368, 1998.

[123702]21972.

[123703]Wydner, K. L.; Bhattacharya, S.; Eckner, R.; Lawrence, J. B.;

[123704]Livingston, D. M.: Localization of human CREB-binding
protein gene

[123705](CREBBP) to 16p13.2–p13.3 by fluorescence in situ hy-
bridization. Genomics 30:

[123706]395–396, 1995.

[123707]21973.

[123708]Yamauchi, T.; Oike, Y.; Kamon, J.; Waki, H.; Komeda, K.;
Tsuchida,

[123709]A.; Date, Y.; Li, M.–X.; Miki, H.; Akanuma, Y.; Nagai, R.;

Kimura,

[123710]S.; Saheki, T.; Nakazato, M.; Naitoh, T.; Yamamura, K.;
Kadowaki,

[123711]T.: Increased insulin sensitivity despite lipodystrophy in
Crebbp

[123712]heterozygous mice. Nature Genet. 30: 221–226, 2002.

[123713]21974.

[123714]Zanger, K.; Radovick, S.; Wondisford, F. E.: CREB binding
protein

[123715]recruitment to the transcription complex requires growth
factor-dependent

[123716]phosphorylation of its GF box. Molec. Cell 7: 551–558,
2001.

[123717]21975.

[123718]Hansen, J. J.; Durr, A.; Cournu-Rebeix, I.; Georgopoulos,
C.; Ang,

[123719]D.; Nielsen, M. N.; Davoine, C.-S.; Brice, A.; Fontaine, B.;
Gregersen,

[123720]N.; Bross, P.: Hereditary spastic paraplegia SPG13 is asso-
ciated

[123721]with a mutation in the gene encoding the mitochondrial
chaperonin

[123722]Hsp60. Am. J. Hum. Genet. 70: 1328–1332, 2002.

[123723]21976.

[123724]Todd, M. J.; Viitanen, P. V.; Lorimer, G. H.: Dynamics of the

[123725]chaperonin ATPase cycle: implications for facilitated protein folding. Science 265:

[123726]659–666, 1994.

[123727]21977.

[123728]Hellevuo, K.; Berry, R.; Sikela, J. M.; Tabakoff, B.: Localization

[123729]of the gene for a novel human adenylyl cyclase (ADCY7) to chromosome

[123730]16. Hum. Genet. 95: 197–200, 1995.

[123731]21978.

[123732]Hellevuo, K.; Yoshimura, M.; Kao, M.; Hoffman, P. L.; Cooper, D.

[123733]M. F.; Tabakoff, B.: A novel adenylyl cyclase sequence cloned from

[123734]the human erythroleukemia cell line. Biochem. Biophys. Res. Commun. 192:

[123735]311–318, 1993.

[123736]21979.

[123737]Moerman, P.; Fryns, J.-P.; Sastrowijoto, S. H.; Vandenberghe, K.;

[123738]Lauweryns, J. M.: Hereditary renal adysplasia: new observations and

[123739]hypotheses. *Pediat. Path.* 14: 405-410, 1994.

[123740]21980.

[123741]Kagawa, W.; Kurumizaka, H.; Ishitani, R.; Fukai, S.; Nureki, O.;

[123742]Shibata, T.; Yokoyama, S.: Crystal structure of the homologous-pairing

[123743]domain from the human Rad52 recombinase in the undecameric form. *Molec.*

[123744]*Cell* 10: 359-371, 2002.

[123745]21981.

[123746]Shen, Z.; Denison, K.; Lobb, R.; Gatewood, J. M.; Chen, D. J.:

[123747]The human and mouse homologs of the yeast RAD52 gene: cDNA cloning,

[123748]sequence analysis, assignment to human chromosome 12p12.2-p13, and

[123749]mRNA expression in mouse tissues. Genomics 25:
199–206, 1995.

[123750]21982.

[123751]Stapleton, D.; Woollatt, E.; Mitchelhill, K. I.; Nicholl, J. K.;

[123752]Fernandez, C. S.; Michell, B. J.; Witters, L. A.; Power, D. A.;
Sutherland,

[123753]G. R.; Kemp, B. E.: AMP-activated protein kinase isoen-
zyme family:

[123754]subunit structure and chromosomal location. FEBS Lett.
409: 452–456,

[123755]1997.

[123756]21983.

[123757]Cox, G. A.; Mahaffey, C. L.; Frankel, W. N.: Identification of

[123758]mouse neuromuscular degeneration gene and mapping of
a second site

[123759]suppressor allele. Neuron 21: 1327–1337, 1998.

[123760]21984.

[123761]Fukita, Y.; Mituza, T.; Shirozu, M.; Ozawa, K.; Shimuzu, A.;
Honjo,

[123762]T.: The human S-mu-bp2, a DNA-binding protein specific
to the single-stranded

[123763]guanine-rich sequence related to the immunoglobulin mu chain switch

[123764]region. J. Biol. Chem. 268: 17463-17470, 1993.

[123765]2 1985.

[123766]Grohmann, K.; Schuelke, M.; Diers, A.; Hoffmann, K.; Lucke, B.;

[123767]Adams, C.; Bertini, E.; Leonhardt-Horti, H.; Muntoni, F.; Ouvrier,

[123768]R.; Pfeufer, A.; Rossi, R.; Van Maldergem, L.; Wilmshurst, J. M.;

[123769]Wienker, T. F.; Sendtner, M.; Rudnik-Schoneborn, S.; Zerres, K.; Hubner,

[123770]C.: Mutations in the gene encoding immunoglobulin mu-binding protein

[123771]2 cause spinal muscular atrophy with respiratory distress type 1. Nature

[123772]Genet. 29: 75-77, 2001.

[123773]2 1986.

[123774]McBride, K.; Robitaille, L.; Tremblay, S.; Argentin, S.; Nemer,

[123775]M.: Fos/jun repression of cardiac specific transcription is targeted

[123776]at a tissue-specific cis element. Mol. Cell. Biol. 13:
600-612,
[123777]1993.
[123778]21987.
[123779]Sebastiani, G.; Durocher, D.; Gros, P.; Nemer, M.; Malo, D.:
Localization
[123780]of the Catf1 transcription factor gene to mouse chromo-
some 19. Mammalian
[123781]Genome 6: 147-148, 1995.
[123782]21988.
[123783]Zonana, J.; Jones, M.; Browne, D.; Litt, M.; Kramer, P.;
Becker,
[123784]H. W.; Brockdorff, N.; Rastan, S.; Davies, K. P.; Clarke, A.;
Thomas,
[123785]N. S. T.: High-resolution mapping of the X-linked hypo-
hidrotic ectodermal
[123786]dysplasia (EDA) locus. Am. J. Hum. Genet. 51: 1036-1046,
1992.
[123787]21989.
[123788]Zonana, J.; Roberts, S. H.; Thomas, N. S. T.; Harper, P. S.:
[123789]Recognition and reanalysis of a cell line from a manifest-

ing female

[123790]with X linked hypohidrotic ectodermal dysplasia and an
X;autosome

[123791]balanced translocation. J. Med. Genet. 25: 383-386, 1988.

[123792]21990.

[123793]Zonana, J.; Schnizel, A.; Upadhyaya, M.; Thomas, N. S. T.;
Anton-Lamprecht,

[123794]I.; Harper, P. S.: Prenatal diagnosis of X-linked hypo-
hidrotic ectodermal

[123795]dysplasia by linkage analysis. Am. J. Med. Genet. 35:
132-135, 1990.

[123796]21991.

[123797]Derry, J. M.; Barnard, P. J.: The gene for tissue inhibitor of

[123798]metalloproteinases (TIMP) is located within an intron of
the synapsin

[123799]I gene on the X chromosome. (Abstract) Cytogenet. Cell
Genet. 58:

[123800]2061-2062, 1991.

[123801]21992.

[123802]Gal, A.; Wieringa, B.; Smeets, D. F. C. M.; Bleeker-
Wagemakers,

[123803]L.; Ropers, H. H.: Submicroscopic interstitial deletion of the X chromosome explains a complex genetic syndrome dominated by Norrie disease. Cytogenet. Cell Genet. 42: 219–224, 1986.

[123806]21993.

[123807]Rugarli, E. I.; Ghezzi, C.; Valsecchi, V.; Ballabio, A.: The Kallmann syndrome gene product expressed in COS cells is cleaved on the cell surface to yield a diffusible component. Hum. Molec. Genet. 5: 1109–1115, 1996.

[123811]21994.

[123812]Schroffner, W. G.; Furth, E. D.: Hypogonadotropic hypogonadism with anosmia (Kallmann's syndrome) unresponsive to clomiphene citrate. J. Clin. Endocr. 31: 267–270, 1970.

[123815]21995.

[123816]Schwanzel-Fukuda, M.; Bick, D.; Pfaff, D. W.: Luteinizing hormone-releasing

[123817]hormone (LHRH)-expressing cells do not migrate normally
in an inherited

[123818]hypogonadal (Kallmann) syndrome. Brain Res. Molec. Brain
Res. 6:

[123819]311–326, 1989.

[123820]21996.

[123821]Soussi-Yanicostas, N.; de Castro, F.; Julliard, A. K.; Perfet-
tini,

[123822]I.; Chedotal, A.; Petit, C.: Anosmin-1, defective in the X-
linked

[123823]form of Kallmann syndrome, promotes axonal branch for-
mation from olfactory

[123824]bulb output neurons. Cell 109: 217–228, 2002.

[123825]21997.

[123826]Sparkes, R. S.; Simpson, R. W.; Paulsen, C. A.: Familial hy-
pogonadotropic

[123827]hypogonadism with anosmia. Arch. Intern. Med. 121:
534–538, 1968.

[123828]21998.

[123829]Wegenke, J. D.; Vehling, D. T.; Wear, J. B., Jr.; Gordon, E.
S.;

[123830]Bargman, G. J.; Deacon, J. S. R.; Herrmann, J. P. R.; Opitz, J. M.

[123831]: Familial Kallmann syndrome with unilateral renal aplasia. Clin.

[123832]Genet. 7: 368–381, 1975.

[123833]21999.

[123834]Soussi-Yanicostas, N.; Hardelin, J.-P.; del Mar Arroyo-Jimenez,

[123835]M.; Ardouin, O.; Legouis, R.; Levilliers, J.; Traincard, F.; Betton,

[123836]J.-M.; Cabanie, L.; Petit, C.: Initial characterization of anosmin-1,

[123837]a putative extracellular matrix protein synthesized by definite neuronal

[123838]cell populations in the central nervous system. J. Cell Sci. 109:

[123839]1749–1757, 1996.

[123840]22000.

[123841]Woods-Samuels, P.; Kazazian, H. H., Jr.; Antonarakis, S. E.:

[123842]Nonhomologous recombination in the human genome: deletions in the

[123843]human factor VIII gene. Genomics 10: 94–101, 1991.

[123844]22001.

[123845]Wray, S.; Grant, P.; Gainer, H.: Evidence that cells expressing
ing

[123846]lutinizing hormone–releasing hormone mRNA in the
mouse are derived

[123847]from progenitor cells in the olfactory placode. Proc. Nat.
Acad.

[123848]Sci. 86: 8132–8136, 1989.

[123849]22002.

[123850]Hata, A.; Tsuzuki, T.; Shimada, K.; Takiguchi, M.; Mori, M.;
Matsuda,

[123851]I.: Structure of the human ornithine transcarbamylase
gene. J. Biochem. 103:

[123852]302–308, 1988.

[123853]22003.

[123854]Hata, A.; Tsuzuki, T.; Shimada, K.; Takiguchi, M.; Mori, M.;
Matsuda,

[123855]I.: Isolation and characterization of the human ornithine
transcarbamylase

[123856]gene: structure of the 5–prime–end region. J. Biochem.

100: 717–725,

[123857]1986.

[123858]22004.

[123859]Hauser, E. R.; Finkelstein, J. E.; Valle, D.; Brusilow, S. W.

[123860]; Allopurinol-induced orotidinuria: a test for mutations at
the ornithine

[123861]carbamoyltransferase locus in women. New Eng. J. Med.
322: 1641–1645,

[123862]1990.

[123863]22005.

[123864]Hentzen, D.; Pelet, A.; Feldman, D.; Rabier, D.; Berthelot,
J.;

[123865]Munnich, A.: Fatal hyperammonemia resulting from a C-
to-T mutation

[123866]at a MspI site of the ornithine transcarbamylase gene.
Hum. Genet. 88:

[123867]153–156, 1991.

[123868]22006.

[123869]Herrin, J. T.; McCredie, D. A.: Peritoneal dialysis in the re-
duction

[123870]of blood ammonia levels in a case of hyperammonaemia.

Arch. Dis.

[123871]Child. 44: 149–151, 1969.

[123872]22007.

[123873]Hjelm, M.; de Silva, L. V. K.; Seakins, J. W. T.; Oberholzer,

[123874]V. G.; Rolles, C. J.: Evidence of inherited urea cycle defect
in

[123875]a case of fatal valproate toxicity. Brit. Med. J. 292: 23–24,
1986.

[123876]22008.

[123877]Hokanson, J. T.; O'Brien, W. E.; Idemoto, J.; Schafer, I. A.:

[123878]Carrier detection in ornithine transcarbamylase deficiency.
J. Pediat. 93:

[123879]75–78, 1978.

[123880]22009.

[123881]Holmes, A. K.; Fowler, B.; Sardharwalla, I. B.: Late-onset
ornithine

[123882]carbamyl transferase deficiency in a male patient: detailed
enzyme

[123883]studies. J. Inherit. Metab. Dis. 10 (suppl. 2): 299–301,
1987.

[123884]22010.

- [123885] Holzgreve, W.; Golbus, M. S.: Prenatal diagnosis of ornithine
- [123886] transcarbamylase deficiency utilizing fetal liver biopsy. Am. J.
- [123887] Hum. Genet. 36: 320–328, 1984.
- [123888] 22011.
- [123889] Honeycutt, D.; Callahan, K.; Rutledge, L.; Evans, B.: Heterozygote
- [123890] ornithine transcarbamylase deficiency presenting as symptomatic hyperammonemia
- [123891] during initiation of valproate therapy. Neurology 42: 666–668, 1992.
- [123892] 22012.
- [123893] Hoogenraad, N.; de Martinis, M. L.; Danks, D. M.: Immunological
- [123894] evidence for an ornithine transcarbamylase lesion resulting in the
- [123895] formation of enzyme with smaller protein subunits. J. Inher. Metab.
- [123896] Dis. 6: 149–152, 1983.
- [123897] 22013.

[123898]Hopkins, I. J.; Connelly, J. F.; Dawson, A. G.; Hird, F. J.;
Maddison,

[123899]T. G.: Hyperammonaemia due to ornithine transcarbamylase deficiency. Arch.

[123900]Dis. Child. 44: 143–148, 1969.

[123901]22014.

[123902]Horwich, A. L.; Fenton, W. A.; Williams, K. R.; Kalousek, F.;

[123903]Kraus, J. P.; Doolittle, R. F.; Konigsberg, W.; Rosenberg, L.
E.:

[123904]Structure and expression of a complementary DNA for the
nuclear coded

[123905]precursor of human mitochondrial ornithine transcarbamylase. Science 224:

[123906]1069–1074, 1984.

[123907]22015.

[123908]Horwich, A. L.; Kalousek, F.; Fenton, W. A.; Pollock, R. A.;
Rosenberg,

[123909]L. E.: Targeting of pre-ornithine transcarbamylase to mitochondria:

[123910]definition of critical regions and residues in the leader
peptide. Cell 44:

[123911]451–459, 1986.

[123912]22016.

[123913]Horwich, A. L.; Kalousek, F.; Mellman, I.; Rosenberg, L. E.:

[123914]A leader peptide is sufficient to direct mitochondrial import of a

[123915]chimeric protein. EMBO J. 4: 1129–1135, 1985.

[123916]22017.

[123917]Horwich, A. L.; Kraus, J. P.; Williams, K.; Kalousek, F.;
Konigsberg,

[123918]W.; Rosenberg, L. E.: Molecular cloning of the cDNA coding for rat

[123919]ornithine transcarbamoylase. Proc. Nat. Acad. Sci. 80:
4258–4262,

[123920]1983.

[123921]22018.

[123922]Hurt, E. C.; van Loon, A. P. G. M.: How proteins find mitochondria

[123923]and intramitochondrial compartments. Trends Biochem.
Sci. 11: 204–207,

[123924]1986.

[123925]22019.

[123926]Kang, E. S.; Solomon, S. S.; Gates, R. E.; Schaeffer, S. J.:

[123927]Red blood cell insulin binding studies in Reye's syndrome
survivors

[123928]and families. Endocr. Res. Commun. 9: 121–133, 1982.

[123929]22020.

[123930]Kay, J. D. S.; Hilton-Jones, D.; Hyman, N.: Valproate toxic-
ity

[123931]and ornithine carbamoyltransferase deficiency. (Letter)
Lancet II:

[123932]1283–1284, 1986.

[123933]22021.

[123934]Kodama, H.; Ohtake, A.; Mori, M.; Okabe, I.; Tatibana, M.;
Kamoshita,

[123935]S.: Ornithine transcarbamylase deficiency: a case with a
truncated

[123936]enzyme precursor and a case with undetectable mRNA ac-
tivity. J. Inherit.

[123937]Metab. Dis. 9: 175–185, 1986.

[123938]22022.

[123939]Komaki, S.; Matsuura, T.; Oyanagi, K.; Hoshide, R.; Kiwaki, K.;

[123940]Endo, F.; Shimadzu, M.; Matsuda, I.: Familial lethal inheritance

[123941]of a mutated paternal gene in females causing X-linked ornithine transcarbamylase

[123942](OTC) deficiency. Am. J. Med. Genet. 69: 177–181, 1997.

[123943]2023.

[123944]Kornfeld, M.; Woodfin, B. M.; Papile, L.; Davis, L. E.; Bernard,

[123945]L. R.: Neuropathology of ornithine carbamyl transferase deficiency. Acta

[123946]Neuropath. 65: 261–264, 1985.

[123947]2024.

[123948]Korson, M. S.; Lillehei, C. W.; Vacanti, J. P.; Levy, H. L.:

[123949]Liver transplantation for ornithine transcarbamylase deficiency (OTCD).

[123950](Abstract) Am. J. Hum. Genet. 45 (suppl.): A8, 1989.

[123951]2025.

[123952]Krieger, I.; Snodgrass, P. J.; Roskamp, J.: Atypical clinical

[123953]course of ornithine transcarbamylase deficiency due to a new mutant

[123954](comparison with Reye's disease). J. Clin. Endocr. Metab. 48: 388-392,

[123955]1979.

[123956]22026.

[123957]Lee, J. T.; Nussbaum, R. L.: An arginine to glutamine mutation

[123958]in residue 109 of human ornithine transcarbamylase completely abolishes

[123959]enzymatic activity in COS1 cells. J. Clin. Invest. 84: 1762-1766,

[123960]1989.

[123961]22027.

[123962]Legius, E.; Baten, E.; Stul, M.; Marynen, P.; Cassiman, J.-J.

[123963]: Sporadic late onset ornithine transcarbamylase deficiency in a boy

[123964]with somatic mosaicism for an intragenic deletion. Clin. Genet. 38:

[123965]155-159, 1990.

[123966]22028.

[123967]Levin, B.; Abraham, J. M.; Oberholzer, V. G.; Burgess, E. A.:

[123968]Hyperammonaemia: a deficiency of liver ornithine transcarbamylase.

[123969]Occurrence in mother and child. Arch. Dis. Child. 44: 152-161, 1969.

[123970]22029.

[123971]Lindgren, V.; de Martinville, B.; Horwich, A. L.; Rosenberg, L.

[123972]E.; Francke, U.: Human ornithine transcarbamylase locus mapped to

[123973]band Xp21.1 near Duchenne muscular dystrophy locus. Science 226:

[123974]698-700, 1984.

[123975]22030.

[123976]Maddalena, A.; Sosnoski, D. M.; Berry, G. T.; Nussbaum, R. L.

[123977]: Mosaicism for an intragenic deletion in a boy with mild ornithine

[123978]transcarbamylase deficiency. New Eng. J. Med. 319: 999-1003, 1988.

[123979]22031.

[123980]Maddalena, A.; Spence, J. E.; O'Brien, W. E.; Nussbaum, R. L.

[123981]: Characterization of point mutations in the same arginine codon in

[123982]three unrelated patients with ornithine transcarbamylase deficiency. J.

[123983]Clin. Invest. 82: 1353-1358, 1988.

[123984]22032.

[123985]Maestri, N. E.; Brusilow, S. W.; Clissold, D. B.; Bassett, S.

[123986]S.: Long-term treatment of girls with ornithine transcarbamylase

[123987]deficiency. New Eng. J. Med. 335: 855-859, 1996.

[123988]22033.

[123989]Maestri, N. E.; Hauser, E. R.; Bartholomew, D.; Brusilow, S. W.

[123990]: Prospective treatment of urea cycle disorders. J. Pediat. 119:

[123991]923-928, 1991.

[123992]22034.

[123993]Alitalo, T.; Karna, J.; Forsius, H.; de la Chapelle, A.: X-linked

[123994]retinoschisis is closely linked to DXS41 and DXS16 but not DXS85. Clin.

[123995]Genet. 32: 192–195, 1987.

[123996]22035.

[123997]Alitalo, T.; Kruse, T. A.; de la Chapelle, A.: Refined localization

[123998]of the gene causing X-linked juvenile retinoschisis. Genomics 9:

[123999]505–510, 1991.

[124000]22036.

[124001]Arden, G. B.; Gorin, M. B.; Polkinghorne, P. J.; Jay, M.; Bird,

[124002]A. C.: Detection of the carrier state of X-linked retinoschisis. Am.

[124003]J. Ophthalm. 105: 590–595, 1988.

[124004]22037.

[124005]Byrne, J.; Warburton, D.: Male excess among anatomically normal

[124006]fetuses in spontaneous abortions. Am. J. Med. Genet. 26: 605–611,

[124007]1987.

[124008]22038.

[124009]Dahl, N.; Goonewardena, P.; Chotai, J.; Anvret, M.; Pettersson,

[124010]U.: DNA linkage of X-linked retinoschisis. Hum. Genet. 78: 228-232,

[124011]1988.

[124012]22039.

[124013]Dahl, N.; Goonewardena, P.; Chotai, J.; Wadelius, C.; Lindsten,

[124014]J.; Pettersson, U.: DNA linkage analysis of X-chromosome linked retinoschisis.

[124015](Abstract) Cytogenet. Cell Genet. 46: 602, 1987.

[124016]22040.

[124017]de Jong, P. T. V. M.; Zrenner, E.; van Meel, G. J.; Keunen, J.

[124018]E. E.; van Norren, D.: Mizuo phenomenon in X-linked retinoschisis:

[124019]pathogenesis of the Mizuo phenomenon. Arch. Ophthalmol. 109: 1104-1108,

[124020]1991.

[124021]22041.

[124022]de la Chapelle, A.; Alitalo, T.; Forsius, H.: X-linked juvenile
[124023]retinoschisis. In: Wright, A.; Jay, B. (eds.): Molecular Genet-
ics
[124024]of Inherited Eye Disorders. Chur: Harwood Academic Publ.
(1st
[124025]ed.): 1994.
[124026]22042.
[124027]Eksandh, L. C.; Ponjavic, V.; Ayyagari, R.; Bingham, E. L.;
Hiriyanna,
[124028]K. T.; Andreasson, S.; Ehinger, B.; Sieving, P. A.: Pheno-
typic expression
[124029]of juvenile X-linked retinoschisis in Swedish families with
different
[124030]mutations in the XLR51 gene. Arch. Ophthal. 118:
1098-1104, 2000.
[124031]22043.
[124032]Eriksson, A. W.; Vainio-Mattila, B.; Krause, U.; Fellman, J.;
[124033]Forsius, H.: Secondary sex ratio in families with X-
chromosomal disorders. Hereditas 57:
[124034]373-381, 1967.
[124035]22044.

- [124036] Forsius, H.; Eriksson, A.; Vainio-Mattila, B.: Geschlechts-
gebundene,
[124037] erbliche Retinoschisis in zwei Familien in Finnland. Klin.
Mbl. Augenheilk. 143:
[124038] 806–816, 1963.
[124039] 22045.
- [124040] Forsius, H.; Krause, U.; Helve, J.; Vuopala, V.; Mustonen,
E.;
[124041] Vainio-Mattila, B.; Fellman, J.; Eriksson, A. W.: Visual acu-
ity in
[124042] 183 cases of X-chromosomal retinoschisis. Canad. J.
Ophthal. 8:
[124043] 385–393, 1973.
[124044] 22046.
- [124045] Forsius, H.; Vainio-Mattila, B.; Eriksson, A.: X-linked
hereditary
[124046] retinoschisis. Brit. J. Ophthal. 46: 678–681, 1962.
[124047] 22047.
- [124048] Forsius, H. R.: Personal Communication. Oulu, Finland
1/1977.
[124049] 22048.

- [124050] Forsius, H. R.; Eriksson, A. W.: Retinoschisis X-chromosomal. In:
- [124051] Eriksson, A. W.; Forsius, H. R.; Nevanlinna, H. R.; Workman, P. L.;
- [124052] Norio, R. K. (eds.): Population Structure and Genetic Disorders.
- [124053] New York: Academic Press (pub.) 1980. Pp. 673–676.
- [124054] 22049.
- [124055] Gehrig, A.; Weber, B. H. F.; Lorenz, B.; Andrassi, M.: First
- [124056] molecular evidence for a de novo mutation in RS1 (XLR1) associated
- [124057] with X-linked juvenile retinoschisis. J. Med. Genet. 36: 932–934,
- [124058] 1999.
- [124059] 22050.
- [124060] Gellert, G.; Petersen, J.; Krawczak, M.; Zoll, B.: Linkage relationship
- [124061] between retinoschisis and four marker loci. Hum. Genet. 79: 382–384,
- [124062] 1988.
- [124063] 22051.

[124064]George, N. D. L.; Yates, J. R. W.; Bradshaw, K.; Moore, A. T.

[124065]: Infantile presentation of X linked retinoschisis. Brit. J.

Ophthal 79:

[124066]653–657, 1995.

[124067]22052.

[124068]George, N. D. L.; Yates, J. R. W.; Moore, A. T.: X linked
retinoschisis. Brit.

[124069]J. Ophthal. 79: 697–702, 1995.

[124070]22053.

[124071]Gieser, E. P.; Falls, H. F.: Hereditary retinoschisis. Am. J.

[124072]Ophthal. 51: 1193–1200, 1961.

[124073]22054.

[124074]Grayson, C.; Reid, S. N. M.; Ellis, J. A.; Rutherford, A.; Sow-
den,

[124075]J. C.; Yates, J. R. W.; Farber, D. B.; Trump, D.:

Retinoschisin,

[124076]the X-linked retinoschisis protein, is a secreted photore-
ceptor protein,

[124077]and is expressed and released by Weri-Rb1 cells. Hum.
Molec. Genet. 9:

[124078]1873–1879, 2000.

[124079]22055.

[124080]Haas, J.: Ueber das Zusammenvorkommen von Veränderungen der Retina

[124081]und Chorioidea. Arch. Augenheilk. 37: 343–348, 1898.

[124082]22056.

[124083]Huopaniemi, L.; Rantala, A.; Forsius, H.; Somer, M.; de la Chapelle,

[124084]A.; Alitalo, T.: Three widespread founder mutations contribute to

[124085]high incidence of X-linked juvenile retinoschisis in Finland. Europ.

[124086]J. Hum. Genet. 7: 368–376, 1999.

[124087]22057.

[124088]Huopaniemi, L.; Rantala, A.; Tahvanainen, E.; de la Chapelle,

[124089]A.; Alitalo, T.: Linkage disequilibrium and physical mapping of X-linked

[124090]juvenile retinoschisis. Am. J. Hum. Genet. 60: 1139–1149, 1997.

[124091]22058.

- [124092]Inoue, Y.; Yamamoto, S.; Okada, M.; Tsujikawa, M.; Inoue, T.;
- [124093]Okada, A. A.; Kusaka, S.; Saito, Y.; Wakabayashi, K.; Miyake, Y.;
- [124094]Fujikado, T.; Tano, Y.: X-linked retinoschisis with point mutations
- [124095]in the XLR51 gene. Arch. Ophthal. 118: 93-96, 2000.
- [124096]22059.
- [124097]Ives, E. J.; Ewing, C. C.; Innes, R.: X-linked juvenile retinoschisis
- [124098]and Xg linkage in five families. (Abstract) Am. J. Hum. Genet. 22:
- [124099]17A-18A, 1970.
- [124100]22060.
- [124101]Kaplan, J.; Pelet, A.; Hentati, H.; Jeanpierre, M.; Briard, M.
- [124102]L.; Journal, H.; Munnich, A.; Dufier, J. L.: Contribution to carrier
- [124103]detection and genetic counselling in X linked retinoschisis. J. Med.
- [124104]Genet. 28: 383-388, 1991.
- [124105]22061.

[124106]Kato, K.; Miyake, Y.; Kachi, S.; Suzuki, T.; Terasaki, H.;
Kawase,

[124107]Y.; Kanda, T.: Axial length and refractive error in X-linked
retinoschisis. Am.

[124108]J. Ophthal. 131: 812–814, 2001.

[124109]22062.

[124110]Kleinert, H.: Eine recessiv-geschlechtsgebundene Form
der idiopathischen

[124111]Netzhautspaltung bei nichtmyopen Jugendlichen. Albrecht
von Graefes

[124112]Arch. Klin. Exp. Ophthal. 154: 295–305, 1953.

[124113]22063.

[124114]Mann, I.; MacRae, A.: Congenital vascular veils in the vit-
reous. Brit.

[124115]J. Ophthal. 22: 1–10, 1938.

[124116]22064.

[124117]Manschot, W. A.: Pathology of hereditary juvenile
retinoschisis. Arch.

[124118]Ophthal. 88: 131–138, 1972.

[124119]22065.

- [124120]Newton, M. S.; Collyer, S.; Phillips, C. I.: Male proband with
- [124121]X linked retinoschisis apparently inherited from his father's family.
- [124122](Letter) J. Med. Genet. 28: 213–214, 1991.
- [124123]22066.
- [124124]Pagenstecher, H. E.: Ueber eine unter dem Bilde der Netzhautablosung
- [124125]verlaufende, erbliche Erkrankung der Retina. Graefes Arch. Ophthal. 86:
- [124126]457–462, 1913.
- [124127]22067.
- [124128]Pawar, H.; Bingham, E. L.; Lunetta, K. L.; Segal, M.; Richards,
- [124129]J. E.; Boehnke, M.; Sieving, P. A.: Refined genetic mapping of juvenile
- [124130]X-linked retinoschisis. Hum. Hered. 45: 206–210, 1995.
- [124131]22068.
- [124132]Retinoschisis Consortium: Functional implications of the spectrum
- [124133]of mutations found in 234 cases with X-linked juvenile retinoschisis

[124134](XLRs). Hum. Molec. Genet. 7: 1185–1192, 1998.

[124135]22069.

[124136]Sauer, C. G.; Gehrig, A.; Warneke–Wittstock, R.; Marquardt, A.;

[124137]Ewing, C. C.; Gibson, A.; Lorenz, B.; Jurklies, B.; Weber, B. H. F.

[124138] Positional cloning of the gene associated with X-linked juvenile

[124139]retinoschisis. Nature Genet. 17: 164–170, 1997.

[124140]22070.

[124141]Sieving, P. A.; Bingham, E. L.; Roth, M. S.; Young, M. R.; Boehnke,

[124142]M.; Kuo, C.–Y.; Ginsburg, D.: Linkage relationship of X-linked juvenile

[124143]retinoschisis with Xp22.1–p22.3 probes. Am. J. Hum. Genet. 47: 616–621,

[124144]1990.

[124145]22071.

[124146]Sorsby, A.; Klein, M.; Gann, J. H.; Siggins, G.: Unusual retinal

[124147]detachment, possibly sex linked. Brit. J. Ophthal. 35:

1-10, 1951.

[124148]22072.

[124149]Springer, W. R.; Cooper, D. N.; Barondes, S. H.: Discoidin I

[124150]is implicated in cell-substratum attachment and ordered
cell migration

[124151]of Dictyostelium discoideum and resembles fibronectin.

Cell 39:

[124152]557-564, 1984.

[124153]22073.

[124154]Vainio-Mattila, B.; Eriksson, A. W.; Forsius, H.: X-
chromosomal

[124155]recessive retinoschisis in the region of Pori: an oph-
thalgo-genetical

[124156]analysis of 103 cases. Acta Ophthal. 47: 1135-1148,
1969.

[124157]22074.

[124158]Van de Vosse, E.; Bergen, A. A. B.; Meershoek, E. J.; Oost-
erwijk,

[124159]J. C.; Gregory, S.; Bakker, B.; Weissenbach, J.; Coffey, A. J.;
van

[124160]Ommen, G.-J. B.; Den Dunnen, J. T.: An Xp22.1-p22.2

YAC contig encompassing

[124161]the disease loci for RS, KFSD, CLS, HYP and RP15: refined
localization

[124162]of RS. *Europ. J. Hum. Genet.* 4: 101–104, 1996.

[124163]22075.

[124164]Weber, B. H. F.; Schrewe, H.; Molday, L. L.; Gehrig, A.;
White,

[124165]K. L.; Seeliger, M. W.; Jaissle, G. B.; Friedburg, C.; Tamm,
E.; Molday,

[124166]R. S.: Inactivation of the murine X-linked juvenile
retinoschisis

[124167]gene, Rs1h, suggests a role of retinoschisin in retinal cell
layer

[124168]organization and synaptic structure. *Proc. Nat. Acad. Sci.*
99: 6222–6227,

[124169]2002.

[124170]22076.

[124171]Wieacker, P.; Wienker, T. F.; Dallapiccola, B.; Bender, K.;
Davies,

[124172]K. E.; Ropers, H. H.: Linkage relationships between
retinoschisis,

[124173]Xg, and a cloned DNA sequence from the distal short arm

of the X chromosome. Hum.

[124174]Genet. 64: 143–145, 1983.

[124175]22077.

[124176]Yanoff, M.; Kertesz Rahn, E.; Zimmerman, L. E.:

Histopathology

[124177]of juvenile retinoschisis. Arch. Ophthal. 79: 49–53, 1968.

[124178]22078.

[124179]Hiraoka, M.; Trese, M. T.; Shastri, B. S.: X-linked juvenile

[124180]retinoschisis associated with a 4–base pair insertion at
codon 55

[124181]of the XLR51 gene. Biochem. Biophys. Res. Commun. 268:
370–372,

[124182]2000.

[124183]22079.

[124184]Hiriyanna, K. T.; Bingham, E. L.; Yashar, B. M.; Ayyagari,
R.;

[124185]Fishman, G.; Small, K. W.; Weinberg, D. V.; Weleber, R. G.;
Lewis,

[124186]R. A.; Andreasson, S.; Richards, J. E.; Sieving, P. A.: Novel
mutations

[124187]in XLR51 causing retinoschisis, including first evidence of

putative

[124188]leader sequence change. Hum. Mutat. 14: 423–427, 1999.

[124189]22080.

[124190]Newman, J. H.; Wheeler, L.; Lane, K. B.; Loyd, E.; Gaddipati, R.;

[124191]Phillips, J. A., III; Loyd, J. E.: Mutation in the gene for bone

[124192]morphogenetic protein receptor II as a cause of primary pulmonary

[124193]hypertension in a large kindred. New Eng. J. Med. 345: 319–324,

[124194]2001. Note: Erratum. New Eng. J. Med. 345: 1506 only, 2001.

[124195]22081.

[124196]Rosenzweig, B. L.; Imamura, T.; Okadome, T.; Cox, G. N.; Yamashita,

[124197]H.; ten Dijke, P.; Heldin, C.–H.; Miyazono, K.: Cloning and characterization

[124198]of a human type II receptor for bone morphogenetic proteins. Proc.

[124199]Nat. Acad. Sci. 92: 7632–7636, 1995.

[124200]22082.

- [124201]Di Paola, R.; Frittitta, L.; Miscio, G.; Bozzali, M.; Baratta, R.; Centra, M.; Spampinato, D.; Santagati, M. G.; Ercolino, T.; Cisternino, C.; Soccio, T. Mastroianno, S.; Tassi, V.; Almgren, P.; Pizuti, A.; Vigneri, R.; Trischitta, V.: A variation in 3-prime UTR of hPTP1B increases specific gene expression and associates with insulin resistance. Am. J. Hum. Genet. 70: 806–812, 2002.
- [124202]22083.
- [124203]Bradley, D. G.; Farrar, G. J.; Sharp, E. M.; Kenna, P.; Humphries, M. M.; McConnell, D. J.; Daiger, S. P.; McWilliam, P.; Humphries, P.: Autosomal dominant retinitis pigmentosa: exclusion of the gene from the short arm of chromosome 1 including the region surrounding the Rhesus locus. Am. J. Hum. Genet. 44: 570–576, 1989.
- [124204]22084.

- [124214]Mutt, V.; Jorpes, J. E.; Magnusson, S.: Structure of porcine secretin:
- [124215]the amino acid sequence. *Europ. J. Biochem.* 15: 513–519, 1970.
- [124216]22085.
- [124217]Narducci, M. G.; Fiorenza, M. T.; Kang, S.–M.; Bevilacqua, A.;
- [124218]Di Giacomo, M.; Remotti, D.; Picchio, M. C.; Fidanza, V.; Cooper,
- [124219]M. D.; Croce, C. M.; Mangia, F.; Russo, G.: TCL1 participates in
- [124220]early embryonic development and is overexpressed in human seminomas. *Proc.*
- [124221]*Nat. Acad. Sci.* 99: 11712–11717, 2002.
- [124222]22086.
- [124223]Casey, G.; Yamanaka, Y.; Freiss, H.; Kobrin, M. S.; Lopez, M.
- [124224]E.; Buchler, M.; Beger, H. G.; Korc, M.: p53 mutations are common
- [124225]in pancreatic cancer and are absent in chronic pancreatitis. *Cancer*
- [124226]*Lett.* 69: 151–160, 1993.

[124227]22087.

[124228]Castedo, M.; Ferri, K. F.; Blanco, J.; Roumier, T.; Larochette,

[124229]N.; Barretina, J.; Amendola, A.; Nardacci, R.; Metivier, D.; Este,

[124230]J. A.; Piacentini, M.; Kroemer, G.: Human immunodeficiency virus

[124231]1 envelope glycoprotein complex-induced apoptosis involves mammalian

[124232]target of rapamycin/FKBP12-rapamycin-associated protein-mediated p53

[124233]phosphorylation. J. Exp. Med. 194: 1097-1110, 2001.

[124234]22088.

[124235]Chen, L.-C.; Neubauer, A.; Kurisu, W.; Waldman, F. M.; Ljung,

[124236]B.-M.; Goodson, W., III; Goldman, E. S.; Moore, D., II; Balazs, M.;

[124237]Liu, E.; Mayall, B. H.; Smith, H. S.: Loss of heterozygosity on the

[124238]short arm of chromosome 17 is associated with high proliferative capacity

[124239]and DNA aneuploidy in primary human breast cancer.

Proc. Nat. Acad.

[124240]Sci. 88: 3847–3851, 1991.

[124241]22089.

[124242]Chen, P.-L.; Chen, Y.; Bookstein, R.; Lee, W.-H.: Genetic mechanisms

[124243]of tumor suppression by the human p53 gene. Science
250: 1576–1580,

[124244]1990.

[124245]22090.

[124246]Chiba, I.; Takahashi, T.; Nau, M. M.; D'Amico, D.; Curiel, D.

[124247]T.; Mitsudomi, T.; Buchhagen, D. L.; Carbone, D.; Pianta-
dosi, S.;

[124248]Koga, H.; Reissman, P.; Slamon, D. J.; Holmes, E. C.;
Minna, J. D.

[124249]: Mutations in the p53 gene are frequent in primary, re-
sected non-small

[124250]cell lung cancer. Oncogene 5: 1603–1610, 1990.

[124251]22091.

[124252]Cho, Y.; Gorina, S.; Jeffrey, P. D.; Pavletich, N. P.: Crystal

[124253]structure of a p53 tumor suppressor–DNA complex: un-

derstanding tumorigenic

[124254]mutations. Science 265: 346–355, 1994.

[124255]22092.

[124256]Chumakov, A. M.; Miller, C. W.; Chen, D. L.; Koeffler, H. P.:

[124257]Analysis of p53 transactivation through high-affinity
binding sites. Oncogene 8:

[124258]3005–3011, 1993.

[124259]22093.

[124260]Chung, R. Y.; Whaley, J. M.; Anderson, K. M.; Freiman, R.
N.;

[124261]Menon, A. G.; Seizinger, B. R.: p53 gene mutations in hu-
man glioblastomas

[124262]associated with early age onset and better survival.
(Abstract) Am.

[124263]J. Hum. Genet. 47 (suppl.): A4, 1990.

[124264]22094.

[124265]Conseiller, E.; Debussche, L.; Landais, D.; Venot, C.; Mara-
trat,

[124266]M.; Sierra, V.; Tocque, B.; Bracco, L.: CTS1: a p53-derived
chimeric

[124267]tumor suppressor gene with enhanced in vitro apoptotic

properties. J.

[124268]Clin. Invest. 101: 120–127, 1998.

[124269]22095.

[124270]Cooper, D. N.; Youssoufian, H.: The CpG dinucleotide and human

[124271]genetic disease. Hum. Genet. 78: 151–155, 1988.

[124272]22096.

[124273]Crawford, L.: Human p53 and human tumours. BioEssays 3: 117–120,

[124274]1985.

[124275]22097.

[124276]Crook, T.; Wrede, D.; Tidy, J. A.; Mason, W. P.; Evans, D. J.;

[124277]Vousden, K. H.: Clonal p53 mutation in primary cervical cancer: association

[124278]with human–papillomavirus–negative tumours. Lancet 339: 1070–1073,

[124279]1992.

[124280]22098.

[124281]Culotta, E.; Koshland, D. E., Jr.: p53 sweeps through cancer

[124282]research. Science 262: 1958–1959, 1993.

[124283]22099.

[124284]Denissenko, M. F.; Chen, J. X.; Tang, M.; Pfeifer, G. P.: Cytosine

[124285]methylation determines hot spots of DNA damage in the human P53 gene. Proc.

[124286]Nat. Acad. Sci. 94: 3893–3898, 1997.

[124287]22100.

[124288]Denissenko, M. F.; Pao, A.; Tang, M.; Pfeifer, G. P.: Preferential

[124289]formation of benzo(a)pyrene adducts at lung cancer mutational hot

[124290]spots in P53. Science 274: 430–434, 1996.

[124291]22101.

[124292]Derry, W. B.; Putzke, A. P.; Rothman, J. H.: Caenorhabditis elegans

[124293]p53: role in apoptosis, meiosis, and stress resistance. Science 294:

[124294]591–595, 2001.

[124295]22102.

[124296]DiGiammarino, E. L.; Lee, A. S.; Cadwell, C.; Zhang, W.;
Bothner,
[124297]B.; Ribeiro, R. C.; Zambetti, G.; Kriwacki, R. W.: A novel
mechanism
[124298]of tumorigenesis involving pH-dependent destabilization
of a mutant
[124299]p53 tetramer. Nature Struct. Biol. 9: 12–16, 2002.
[124300]22103.

[124301]Donehower, L. A.; Harvey, M.; Slagle, B. L.; McArthur, M. J.;
[124302]Montgomery, C. A., Jr.; Butel, J. S.; Bradley, A.: Mice defi-
cient
[124303]for p53 are developmentally normal but susceptible to
spontaneous
[124304]tumours. Nature 356: 215–221, 1992.
[124305]22104.

[124306]Dumaz, N.; Drougard, C.; Sarasin, A.; Daya-Grosjean, L.:
Specific
[124307]UV-induced mutation spectrum in the p53 gene of skin
tumors from DNA-repair-deficient
[124308]xeroderma pigmentosum patients. Proc. Nat. Acad. Sci.
90: 10529–10533,

[124309]1993.

[124310]22105.

[124311]El-Deiry, W. S.; Kern, S. E.; Pietenpol, J. A.; Kinzler, K. W.;

[124312]Vogelstein, B.: Definition of a consensus binding site for
p53. Nature

[124313]Genet. 1: 45-49, 1992.

[124314]22106.

[124315]El-Deiry, W. S.; Tokino, T.; Velculescu, V. E.; Levy, D. B.;
Parsons,

[124316]R.; Trent, J. M.; Lin, D.; Mercer, E.; Kinzler, K. W.; Vogel-
stein,

[124317]B.: WAF1, a potential mediator of p53 tumor suppression.
Cell 75:

[124318]817-825, 1993.

[124319]22107.

[124320]Moore, K. J.; D'Amore-Bruno, M. A.; Korfhagen, T. R.;
Glasser,

[124321]S. W.; Whitsett, J. A.; Jenkins, N. A.; Copeland, N. G.: Chro-
mosomal

[124322]localization of three pulmonary surfactant protein genes
in the mouse. Genomics 12:

[124323]388–393, 1992.

[124324]22108.

[124325]Nogee, L. M.; Dunbar, A. E., III; Wert, S. E.; Askin, F.; Ham-
vas,

[124326]A.; Whitsett, J. A.: A mutation in the surfactant protein C
gene

[124327]associated with familial interstitial lung disease. New Eng.
J. Med. 344:

[124328]573–579, 2001.

[124329]22109.

[124330]Warr, R. G.; Hawgood, S.; Buckley, D. I.; Crisp, T. M.;
Schilling,

[124331]J.; Benson, B. J.; Ballard, P. L.; Clements, J. A.; White, R. T.:

[124332]Low molecular weight human pulmonary surfactant pro-
tein (SP5): isolation,

[124333]characterization, and cDNA and amino acid sequences.
Proc. Nat. Acad.

[124334]Sci. 84: 7915–7919, 1987.

[124335]22110.

[124336]Wood, S.; Yaremko, M. L.; Schertzer, M.; Kelemen, P. R.;
Minna,

[124337]J.; Westbrook, C. A.: Mapping of the pulmonary surfactant
SP5 (SFTP2)

[124338]locus to 8p21 and characterization of a microsatellite re-
peat marker

[124339]that shows frequent loss of heterozygosity in human car-
cinomas. Genomics 24:

[124340]597–600, 1994.

[124341]22111.

[124342]Kolble, K.; Lu, J.; Mole, S. E.; Kaluz, S.; Reid, K. B. M.: As-
signment

[124343]of the human pulmonary surfactant protein D gene
(SFTP4) to 10q22–q23

[124344]close to the surfactant protein A gene cluster. Genomics
17: 294–298,

[124345]1993.

[124346]22112.

[124347]Botas, C.; Poulain, F.; Akiyama, J.; Brown, C.; Allen, L.; Go-

erke,

[124348]J.; Clements, J.; Carlson, E.; Gillespie, A. M.; Epstein, C.;
Hawgood,

[124349]S.: Altered surfactant homeostasis and alveolar type II cell
morphology

[124350]in mice lacking surfactant protein D. Proc. Nat. Acad. Sci.
95:

[124351]11869–11874, 1998.

[124352]22113.

[124353]Crouch, E.; Rust, K.; Veile, R.; Donis-Keller, H.; Grosso, L.:

[124354]Genomic organization of human surfactant protein D
(SP-D): SP-D is

[124355]encoded on chromosome 10q22.2–23.1. J. Biol. Chem.
268: 2976–2983,

[124356]1993.

[124357]22114.

[124358]Holmskov, U.; Lawson, P.; Teisner, B.; Tornøe, I.; Willis, A.
C.;

[124359]Morgan, C.; Koch, C.; Reid, K. B. M.: Isolation and charac-
terization

[124360]of a new member of the scavenger receptor superfamily,

glycoprotein-340

[124361](gp-340), as a lung surfactant protein-D binding molecule. J. Biol.

[124362]Chem. 272: 13743-13749, 1997.

[124363]22115.

[124364]Holmskov, U.; Mollenhauer, J.; Madsen, J.; Vitved, L.; Gronlund,

[124365]J.; Tornoe, I.; Kliem, A.; Reid, K. B. M.; Poustka, A.; Skjodt, K.

[124366]: Cloning of gp-340, a putative opsonin receptor for lung surfactant

[124367]protein D. Proc. Nat. Acad. Sci. 96: 10794-10799, 1999.

[124368]22116.

[124369]LeVine, A. M.; Whitsett, J. A.; Hartshorn, K. L.; Crouch, E. C.;

[124370]Korfhagen, T. R.: Surfactant protein D enhances clearance of influenza

[124371]A virus from the lung in vivo. J. Immun. 167: 5868-5873, 2001.

[124372]22117.

[124373]Lu, J.; Willis, A. C.; Reid, K. B. M.: Purification, characteri-

zation

[124374]and cDNA cloning of human lung surfactant protein D.

Biochem. J. 284:

[124375]795–802, 1992.

[124376]22118.

[124377]Rust, K.; Grosso, L.; Zhang, V.; Chang, D.; Persson, A.;

Longmore,

[124378]W.; Cai, G.-Z.; Crouch, E.: Human surfactant protein D:

SP-D contains

[124379]a C-type lectin carbohydrate recognition domain. Arch.

Biochem. Biophys. 290:

[124380]116–126, 1991.

[124381]22119.

[124382]Wert, S. E.; Yoshida, M.; LeVine, A. M.; Ikegami, M.; Jones,

T.;

[124383]Ross, G. F.; Fisher, J. H.; Korfhagen, T. R.; Whitsett, J. A.:

Increased

[124384]metalloproteinase activity, oxidant production, and em-

physema in surfactant

[124385]protein D gene-inactivated mice. Proc. Nat. Acad. Sci. 97:

5972–5977,

[124386]2000.

[124387]22120.

[124388]Bunge, S.; Wedemann, H.; David, D.; Terwilliger, D. J.; van
den

[124389]Born, L. I.; Aulehla-Scholz, C.; Samanns, C.; Horn, M.; Ott,
J.; Schwinger,

[124390]E.; Schinzel, A.; Denton, M. J.; Gal, A.: Molecular analysis
and

[124391]genetic mapping of the rhodopsin gene in families with
autosomal dominant

[124392]retinitis pigmentosa. Genomics 17: 230–233, 1993.

[124393]22121.

[124394]Chang, H.-Y.; Ready, D. F.: Rescue of photoreceptor de-
generation

[124395]in rhodopsin-null Drosophila mutants by activated Rac1.
Science 290:

[124396]1978–1980, 2000.

[124397]22122.

[124398]Dikshit, M.; Agarwal, R.: Mutation analysis of codons 345
and

[124399]347 of rhodopsin gene in Indian retinitis pigmentosa pa-
tients. J.

[124400]Genet. 80: 111–116, 2001.

[124401]22123.

[124402]Dryja, T. P.; Hahn, L. B.; Cowley, G. S.; McGee, T. L.;
Berson,

[124403]E. L.: Mutation spectrum of the rhodopsin gene among
patients with

[124404]autosomal dominant retinitis pigmentosa. Proc. Nat. Acad.
Sci. 88:

[124405]9370–9374, 1991.

[124406]22124.

[124407]Dryja, T. P.; McGee, T. L.; Hahn, L. B.; Cowley, G. S.; Ols-
son,

[124408]J. E.; Reichel, E.; Sandberg, M. A.; Berson, E. L.: Mutations
within

[124409]the rhodopsin gene in patients with autosomal dominant
retinitis pigmentosa. New

[124410]Eng. J. Med. 323: 1302–1307, 1990.

[124411]22125.

[124412]Donis-Keller, H.; Green, P.; Helms, C.; Cartinhour, S.;
Weiffenbach,

[124413]B.; Stephens, K.; Keith, T. P.; Bowden, D. W.; Smith, D. R.;

Lander,

[124414]E. S.; Botstein, D.; Akots, G.; Rediker, K. S.; Gravius, T.:

Brown,

[124415]V. A.; Rising, M. B.; Parker, C.; Powers, J. A.; Watt, D. E.;

Kauffman,

[124416]E. R.; Bricker, A.; Phipps, P.; Muller-Kahle, H.; Fulton, T. R.;

Ng,

[124417]S.; Schumm, J. W.; Braman, J. C.; Knowlton, R. G.; Barker,

D. F.;

[124418]Crooks, S. M.; Lincoln, S. E.; Daly, M. J. and Abrahamson,

J.: A genetic

[124419]linkage map of the human genome. Cell 51: 319-337,

1987.

[124420]22126.

[124421]Dryja, T. P.; McGee, T. L.; Hahn, L. B.; Cowley, G. S.; Ols-

son,

[124422]J. E.; Reichel, E.; Sandberg, M. A.; Berson, E. L.: Missense

mutations

[124423]within the rhodopsin gene in patients with autosomal

dominant retinitis

[124424]pigmentosa. (Abstract) Am. J. Hum. Genet. 47 (suppl.):

A54, 1990.

[124425]22127.

[124426]Dryja, T. P.; McGee, T. L.; Reichel, E.; Hahn, L. B.; Cowley,
[124427]G. S.; Yandell, D. W.; Sandberg, M. A.; Berson, E. L.: A
point mutation
[124428]of the rhodopsin gene in one form of retinitis pigmentosa.
Nature 343:
[124429]364–366, 1990.
[124430]22128.
[124431]Elliott, R. W.; Sparkes, R. S.; Mohandas, T.; Grant, S. G.;
McGinnis,
[124432]J. F.: Localization of the rhodopsin gene to the distal half
of mouse
[124433]chromosome 6. Genomics 6: 635–644, 1990.
[124434]22129.
[124435]Farrar, G. J.; Findlay, J. B. C.; Kumar–Singh, R.; Kenna, P.;
[124436]Humphries, M. M.; Sharpe, E.; Humphries, P.: Autosomal
dominant retinitis
[124437]pigmentosa: a novel mutation in the rhodopsin gene in
the original
[124438]3q linked family. Hum. Molec. Genet. 1: 769–771, 1992.
[124439]22130.

- [124440]Farrar, G. J.; Kenna, P.; Redmond, R.; McWilliam, P.;
Bradley,
[124441]D. G.; Humphries, M. M.; Sharp, E. M.; Inglehearn, C. F.;
Bashir,
[124442]R.; Jay, M.; Watty, A.; Ludwig, M.; Schinzel, A.; Samanns,
C.; Gal,
[124443]A.; Bhattacharya, S.; Humphries, P.: Autosomal dominant
retinitis
[124444]pigmentosa: absence of the rhodopsin proline-
to-histidine substitution
[124445](codon 23) in pedigrees from Europe. Am. J. Hum. Genet.
47: 941-945,
[124446]1990.
[124447]22131.
[124448]Farrar, G. J.; Kenna, P.; Redmond, R.; Shiels, D.; McWilliam,
[124449]P.; Humphries, M. M.; Sharp, E. M.; Jordan, S.; Kumar-
Singh, R.; Humphries,
[124450]P.: Autosomal dominant retinitis pigmentosa: a mutation
in codon
[124451]178 of the rhodopsin gene in two families of Celtic origin.
Genomics 11:
[124452]1170-1171, 1991.

[124453]22132.

[124454]Fishman, G. A.; Stone, E. M.; Gilbert, L. D.; Kenna, P.;
Sheffield,

[124455]V. C.: Ocular findings associated with a rhodopsin gene
codon 58

[124456]transversion mutation in autosomal dominant retinitis
pigmentosa. Arch.

[124457]Ophthalmol. 109: 1387-1393, 1991.

[124458]22133.

[124459]Fishman, G. A.; Stone, E. M.; Gilbert, L. D.; Sheffield, V. C.

[124460]: Ocular findings associated with a rhodopsin gene codon
106 mutation:

[124461]glycine-to-arginine change in autosomal dominant retini-
tis pigmentosa. Arch.

[124462]Ophthalmol. 110: 646-653, 1992.

[124463]22134.

[124464]Fishman, G. A.; Vandenburgh, K.; Stone, E. M.; Gilbert, L.
D.;

[124465]Alexander, K. R.; Sheffield, V. C.: Ocular findings associ-
ated with

[124466]rhodopsin gene codon 267 and codon 190 mutations in

dominant retinitis

[124467]pigmentosa. Arch. Ophthalm. 110: 1582–1588, 1992.

[124468]22135.

[124469]Franke, R. R.; Konig, B.; Sakmar, T. P.; Khorana, H. G.;
Hofmann,

[124470]K. P.: Rhodopsin mutants that bind but fail to activate
transducin. Science 250:

[124471]123–125, 1990.

[124472]22136.

[124473]Gal, A.; Artlich, A.; Ludwig, M.; Niemeyer, G.; Olek, K.;
Schwinger,

[124474]E.; Schinzel, A.: Pro347-to-arg mutation of the rhodopsin
gene in

[124475]autosomal dominant retinitis pigmentosa. Genomics 11:
468–470, 1991.

[124476]22137.

[124477]Gal, A.; Samanns, C.; Watty, A.; Jimenez, J.; Ludwig, M.;
Chand,

[124478]A.; Pongratz, J.; Colantuoni, V.; Gusseck, H.; Olsson, J.;
Neugebauer,

[124479]M.; Schinzel, A.; Denton, M.: A gene for autosomal domi-

nant retinitis

[124480]pigmentosa is closely linked to D3S20 on 3q. (Abstract)

Retinitis

[124481]Pigmentosa Meeting, Dublin July, 1990.

[124482]22138.

[124483]Hao, W.; Wenzel, A.; Obin, M. S.; Chen, C.-K.; Brill, E.;

Krasnoperova,

[124484]N. V.; Eversole-Cire, P.; Kleyner, Y.; Taylor, A.; Simon, M.

I.; Grimm,

[124485]C.; Reme, C. E.; Lem, J.: Evidence for two apoptotic path-

ways in

[124486]light-induced retinal degeneration. Nature Genet. 32:

254-260, 2002.

[124487]22139.

[124488]Heckenlively, J. R.; Rodriguez, J. A.; Daiger, S. P.: Autosomal

mal

[124489]dominant sectoral retinitis pigmentosa: two families with

transversion

[124490]mutation in codon 23 of rhodopsin. Arch. Ophthal. 109:

84-91, 1991.

[124491]22140.

[124492]Humphries, M. M.; Rancourt, D.; Farrar, G. J.; Kenna, P.;

Hazel,

[124493]M.; Bush, R. A.; Sieving, P. A.; Sheils, D. M.; McNally, N.;
Creighton,

[124494]P.; Erven, A.; Boros, A.; Gulya, K.; Capecchi, M. R.;
Humphries, P.

[124495]: Retinopathy induced in mice by targeted disruption of
the rhodopsin

[124496]gene. *Nature Genet.* 15: 216–219, 1997.

[124497]22141.

[124498]Humphries, P.: Personal Communication. Dublin, Ireland
2/1990.

[124499]22142.

[124500]Humphries, P.; Farrar, G. J.; Kenna, P.; McWilliam, P.: Re-
tinitis

[124501]pigmentosa: genetic mapping in X-linked and autosomal
forms of the

[124502]disease. *Clin. Genet.* 38: 1–13, 1990.

[124503]22143.

[124504]Humphries, P.; Kenna, P.; Farrar, G. J.: On the molecular
genetics

[124505]of retinitis pigmentosa. *Science* 256: 804–808, 1992.

[124506]22144.

[124507]Whitmore, T. E.; Holloway, J. L.; Lofton-Day, C. E.; Maurer, M.

[124508]F.; Chen, L.; Quinton, T. J.; Vincent, J. B.; Scherer, S. W.; Lok,

[124509]S.: Human secretin (SCT): gene structure, chromosome location, and

[124510]distribution of mRNA. Cytogenet. Cell Genet. 90: 47-52, 2000.

[124511]22145.

[124512]Domino, S. E.; Zhang, L.; Gillespie, P. J.; Saunders, T. L.; Lowe,

[124513]J. B.: Deficiency of reproductive tract alpha(1,2)fucosylated glycans

[124514]and normal fertility in mice with targeted deletions of the FUT1 or

[124515]FUT2 alpha(1,2)fucosyltransferase locus. Molec. Cell. Biol. 21:

[124516]8336-8345, 2001.

[124517]22146.

[124518]Gedde-Dahl, T., Jr.; Olaisen, B.; Teisberg, P.; Wilhelmy, M. C.;

[124519]Mevag, B.; Helland, R.: The locus for apolipoprotein E (apoE) is

[124520]close to the Lutheran (Lu) blood group locus on chromosome 19. Hum.

[124521]Genet. 67: 178–182, 1984.

[124522]22147.

[124523]Grubb, R.: Zur Genetik des Lewis-Systems. Naturwissenschaften 21:

[124524]560–561, 1953.

[124525]22148.

[124526]Le Pendu, J.; Lemieux, R. U.; Lambert, F.; Dalix, A.-M.; Oriol,

[124527]R.: Distribution of H type 1 and H type 2 antigenic determinants

[124528]in human sera and saliva. Am. J. Hum. Genet. 34: 402–415, 1982.

[124529]22149.

[124530]Oriol, R.; Danilovs, J.; Hawkins, B. R.: A new genetic model

[124531]proposing that the Se gene is a structural gene closely linked to

[124532]the H gene. Am. J. Hum. Genet. 33: 421–431, 1981.

[124533]22150.

[124534]Reguigne-Arnould, I.; Couillin, P.; Mollicone, R.; Faure, S.;

[124535]Fletcher, A.; Kelly, R. J.; Lowe, J. B.; Oriol, R.: Relative positions

[124536]of two clusters of human alpha-L-fucosyltransferases in 19q (FUT1-FUT2)

[124537]and 19p (FUT6-FUT3-FUT5) within the microsatellite genetic map of

[124538]chromosome 19. Cytogenet. Cell Genet. 71: 158-162, 1995.

[124539]22151.

[124540]Kolanus, W.; Nagel, W.; Schiller, B.; Zeitlmann, L.; Godar, S.;

[124541]Stockinger, H.; Seed, B.: Alpha-L-beta-2 integrin/LFA-1 binding to

[124542]ICAM-1 induced by cytohesin-1, a cytoplasmic regulatory molecule. Cell 86:

[124543]233-242, 1996.

[124544]22152.

[124545]Ogasawara, M.; Kim, S.-C.; Adamik, R.; Togawa, A.; Ferrans, V.

[124546]J.; Takeda, K.; Kirby, M.; Moss, J.; Vaughan, M.: Similarities
in
[124547]function and gene structure of cytohesin-4 and cyto-
hesin-1, guanine
[124548]nucleotide-exchange proteins for ADP-ribosylation fac-
tors. J. Biol.
[124549]Chem. 275: 3221-3230, 2000.
[124550]22153.
[124551]Findlay, D. M.; Fisher, L. W.; McQuillan, C. I.; Termine, J.
D.;
[124552]Young, M. F.: Isolation of the osteonectin gene: evidence
that a
[124553]variable region of the osteonectin molecule is encoded
within one
[124554]exon. Biochemistry 27: 1483-1489, 1988.
[124555]22154.
[124556]Gilmour, D. T.; Lyon, G. J.; Carlton, M. B. L.; Sanes, J. R.;
Cunningham,
[124557]J. M.; Anderson, J. R.; Hogan, B. L. M.; Evans, M. J.;
Colledge, W.
[124558]H.: Mice deficient for the secreted glycoprotein SPARC/
osteonectin/BM40

[124559]develop normally but show severe age-onset cataract formation and

[124560]disruption of the lens. EMBO J. 17: 1860–1870, 1998.

[124561]22155.

[124562]Sanyanusin, P.; McNoe, L. A.; Sullivan, M. J.; Weaver, R. G.;

[124563]Eccles, M. R.: Mutation of PAX2 in two siblings with renal-coloboma

[124564]syndrome. Hum. Molec. Genet. 4: 2183–2184, 1995.

[124565]22156.

[124566]Sanyanusin, P.; Norrish, J. H.; Ward, T. A.; Nebel, A.; McNoe,

[124567]L. A.; Eccles, M. R.: Genomic structure of the human PAX2 gene. Genomics 35:

[124568]258–261, 1996.

[124569]22157.

[124570]Sanyanusin, P.; Schimmenti, L. A.; McNoe, L. A.; Ward, T. A.;

[124571]Pierpont, M. E. M.; Sullivan, M. J.; Dobyns, W. B.; Eccles, M. R.

[124572]: Mutation of the PAX2 gene in a family with optic nerve colobomas,

[124573]renal anomalies and vesicoureteral reflux. Nature Genet.
9: 358–364,

[124574]1995.

[124575]22158.

[124576]Schimmenti, L. A.; Cunliffe, H. E.; McNoe, L. A.; Ward, T.
A.;

[124577]French, M. C.; Shim, H. H.; Zhang, Y.–H.; Proesmans, W.;
Leys, A.;

[124578]Byerly, K. A.; Braddock, S. R.; Masuno, M.; Imaizumi, K.;
Devriendt,

[124579]K.; Eccles, M. R.: Further delineation of renal–coloboma
syndrome

[124580]in patients with extreme variability of phenotype and
identical PAX2

[124581]mutations. Am. J. Hum. Genet. 60: 869–878, 1997.

[124582]22159.

[124583]Schimmenti, L. A.; Pierpont, M. E.; Carpenter, B. L. M.;
Kashtan,

[124584]C. E.; Johnson, M. R.; Dobyns, W. B.: Autosomal dominant
optic nerve

[124585]colobomas, vesicoureteral reflux, and renal anomalies.
Am. J. Med.

[124586]Genet. 59: 204–208, 1995.

[124587]22160.

[124588]Schimmenti, L. A.; Shim, H. H.; Wirtschafter, J. D.; Pan-
zarino,

[124589]V. A.; Kashtan, C. E.; Kirkpatrick, S. J.; Wargowski, D. S.;
France,

[124590]T. D.; Michel, E.; Dobyns, W. B.: Homonucleotide expan-
sion and contraction

[124591]mutations of PAX2 and inclusion of Chiari 1 malformation
as part of

[124592]renal–coloboma syndrome. Hum. Mutat. 14: 369–376,
1999.

[124593]22161.

[124594]Stapleton, P.; Weith, A.; Urbanek, P.; Kozmik, Z.; Bus-
slinger,

[124595]M.: Chromosomal localization of seven PAX genes and
cloning of a

[124596]novel family member, PAX–9. Nature Genet. 3: 292–298,
1993.

[124597]22162.

[124598]Tellier, A.–L.; Amiel, J.; Delezoide, A.–L.; Audollent, S.;
Auge,

[124599]J.; Esnault, D.; Encha-Razavi, F.; Munnich, A.; Lyonnet, S.; Vekemans, M.; Attie-Bitach, T.: Expression of the PAX2 gene in human embryos and exclusion in the CHARGE syndrome. Am. J. Med. Genet. 93: 85-88, 2000.

[124602]2000.

[124603]22163.

[124604]Tellier, A.-L.; Amiel, J.; Salomon, R.; Jolly, D.; Delezoide, A.-L.; Auge, J.; Gubler, M.-C.; Munnich, A.; Lyonnet, S.; Antignac, C.; Vekemans, M.; Broyer, M.; Attie-Bitach, T.: PAX2 expression during early human development and its mutations in renal hypoplasia with or without coloboma. (Abstract) Am. J. Hum. Genet. 63 (suppl.): A7 only, 1998.

[124609]A7 only, 1998.

[124610]22164.

[124611]Ward, T. A.; Nebel, A.; Reeve, A. E.; Eccles, M. R.: Alternative

[124612] messenger RNA forms and open reading frames within an additional conserved

[124613] region of the human PAX-2 gene. Cell Growth Differ. 5: 1015-1021,

[124614] 1994.

[124615] 22165.

[124616] Weaver, R. G.; Cashwell, L. F.; Lorentz, W.; Whiteman, D.; Geisinger,

[124617] K. R.; Ball, M.: Optic nerve coloboma associated with renal disease. Am.

[124618] J. Med. Genet. 29: 597-605, 1988.

[124619] 22166.

[124620] Barr, F. G.; Nauta, L. E.; Davis, R. J.; Schafer, B. W.; Nycum,

[124621] L. M.; Biegel, J. A.: In vivo amplification of the PAX3-FKHR and

[124622] PAX7-FKHR fusion genes in alveolar rhabdomyosarcoma. Hum. Molec.

[124623] Genet. 5: 15-21, 1996.

[124624] 22167.

[124625] Burri, M.; Tromvoukis, Y.; Bopp, D.; Frigerio, G.; Noll, M.: Conservation

[124626]of the paired domain in metazoans and its structure in
three isolated

[124627]human genes. EMBO J. 8: 1183–1190, 1989.

[124628]22168.

[124629]Gruss, P.; Walther, C.: Pax in development. Cell 69:
719–722,

[124630]1992.

[124631]22169.

[124632]Schafer, B. W.; Mattei, M. G.: The human paired domain
gene PAX7

[124633](Hup1) maps to chromosome 1p35–1p36.2. Genomics 17:
249–251, 1993.

[124634]22170.

[124635]Seale, P.; Sabourin, L. A.; Girgis–Gabardo, A.; Mansouri,
A.; Gruss,

[124636]P.; Rudnicki, M. A.: Pax7 is required for the specification
of myogenic

[124637]satellite cells. Cell 102: 777–786, 2000.

[124638]22171.

[124639]Shapiro, D. N.; Sublett, J. E.; Li, B.; Valentine, M. B.; Morris,

[124640]S. W.; Noll, M.: The gene for PAX7, a member of the
paired-box-containing
[124641]genes, is localized on human chromosome arm 1p36. Ge-
nomics 17: 767-769,
[124642]1993.
[124643]22172.
[124644]Helwig, U.; Imai, K.; Schmahl, W.; Thomas, B. E.; Varnum,
D. S.;
[124645]Nadeau, J. H.; Balling, R.: Interaction between undulated
and Patch
[124646]leads to an extreme form of spina bifida in double-mutant
mice. Nature
[124647]Genet. 11: 60-63, 1995.
[124648]22173.
[124649]Sun, J.; Rose, J. B.; Bird, P.: Gene structure, chromosomal
localization,
[124650]and expression of the murine homologue of human pro-
teinase inhibitor
[124651]6 (PI-6) suggests divergence of PI-6 from the ovalbumin
serpins. J.
[124652]Biol. Chem. 270: 16089-16096, 1995.

[124653]22174.

[124654]Cowley, C. M. E.; Simrak, D.; Spurr, N. K.; Arnemann, J.;
Buxton,

[124655]R. S.: The plakophilin 1 (PKP1) and plakoglobin (JUP) genes
map to

[124656]human chromosomes 1q and 17, respectively. Hum.
Genet. 100: 486–488,

[124657]1997.

[124658]22175.

[124659]Guenet, J.–L.; Simon–Chazottes, D.; Ringwald, M.; Kemler,
R.:

[124660]The genes coding for alpha and beta catenin (Catn1 and
Catn2) and

[124661]plakoglobin (Jup) map to mouse chromosomes 18, 9, and
11, respectively. Mammalian

[124662]Genome 6: 363–366, 1995.

[124663]22176.

[124664]Ruiz, P.; Brinkmann, V.; Ledermann, B.; Behrend, M.;
Grund, C.;

[124665]Thalhammer, C.; Vogel, F.; Birchmeier, C.; Gunthert, U.;
Franke, W.

[124666]W.; Birchmeier, W.: Targeted mutation of plakoglobin in mice reveals

[124667]essential functions of desmosomes in the embryonic heart. J. Cell

[124668]Biol. 135: 215–225, 1996.

[124669]22177.

[124670]Buckley, M. F.; Goding, J. W.: Plasma cell membrane glycoprotein

[124671]gene Pca-1 (alkaline phosphodiesterase I) is linked to the proto-oncogene

[124672]Myb on mouse chromosome 10. Immunogenetics 36: 199–201, 1992.

[124673]22178.

[124674]Buckley, M. F.; Loveland, K. A.; McKinstry, W. J.; Garson, O. M.;

[124675]Goding, J. W.: Plasma cell membrane glycoprotein PC-1: cDNA cloning

[124676]of the human molecule, amino acid sequence, and chromosomal location. J.

[124677]Biol. Chem. 265: 17506–17511, 1990.

[124678]22179.

[124679]Frittitta, L.; Baratta, R.; Spampinato, D.; Di Paola, R.; Piz-
zuti,

[124680]A.; Vigneri, R.; Trischitta, V.: The Q121 PC-1 variant and
obesity

[124681]have additive and independent effects in causing insulin
resistance. J.

[124682]Clin. Endocr. Metab. 86: 5888-5891, 2001.

[124683]22180.

[124684]Harahap, A. R.; Goding, J. W.: Distribution of the murine
plasma

[124685]cell antigen PC-1 in non-lymphoid tissues. J. Immun. 141:
2317-2320,

[124686]1988.

[124687]22181.

[124688]Huang, R.; Rosenbach, M.; Vaughn, R.; Provvedini, D.;
Rebbe, N.;

[124689]Hickman, S.; Goding, J.; Terkeltaub, R.: Expression of the
murine

[124690]plasma cell nucleotide pyrophosphohydrolase PC-1 is
shared by human

[124691]liver, bone, and cartilage cells: regulation of PC-1 expres-
sion in

[124692]osteosarcoma cells by transforming growth factor-beta. J.
Clin. Invest. 94:
[124693]560-567, 1994.

[124694]22182.

[124695]Kahn, C. R.: Causes of insulin resistance. Nature 373:
384-385,
[124696]1995.

[124697]22183.

[124698]Maddux, B. A.; Goldfine, I. D.: Membrane glycoprotein PC-
1 inhibition
[124699]of insulin receptor function occurs via direct interaction
with the
[124700]receptor alpha-subunit. Diabetes 49: 13-19, 2000.

[124701]22184.

[124702]Maddux, B. A.; Sbraccia, P.; Kumakura, S.; Sasson, S.;
Youngren,
[124703]J.; Fisher, A.; Spencer, S.; Grupe, A.; Henzel, W.; Stewart, T.
A.;

[124704]Reaven, G. M.; Goldfine, I. D.: Membrane glycoprotein PC-
1 and insulin
[124705]resistance in non-insulin-dependent diabetes mellitus.

Nature 373:

[124706]448–451, 1995.

[124707]22185.

[124708]Nakamura, I.; Ikegawa, S.; Okawa, A.; Okuda, S.;
Koshizuka, Y.;

[124709]Kawaguchi, H.; Nakamura, K.; Koyama, T.; Goto, S.;
Toguchida, J.;

[124710]Matsushita, M.; Ochi, T.; Takaoka, K.; Nakamura, Y.: Asso-
ciation

[124711]of the human NPPS gene with ossification of the posterior
longitudinal

[124712]ligament of the spine (OPLL). Hum. Genet. 104: 492–497,
1999.

[124713]22186.

[124714]Okawa, A.; Nakamura, I.; Goto, S.; Moriya, H.; Nakamura,
Y.; Ikegawa,

[124715]S.: Mutation in Npps in a mouse model of ossification of
the posterior

[124716]longitudinal ligament of the spine. Nature Genet. 19:
271–273, 1998.

[124717]22187.

[124718]Pizzuti, A.; Frittitta, L.; Argiolas, A.; Baratta, R.; Goldfine,

[124719]I. D.; Bozzali, M.; Ercolino, T.; Scarlato, G.; Iacoviello, L.;
Vigneri,

[124720]R.; Tassi, V.; Trischitta, V.: A polymorphism (K121Q) of
the human

[124721]glycoprotein PC-1 gene coding region is strongly associ-
ated with insulin

[124722]resistance. Diabetes 48: 1881-1884, 1999.

[124723]22188.

[124724]Rebbe, N. F.; Tong, B. D.; Finley, E. M.; Hickman, S.: Iden-
tification

[124725]of nucleotide pyrophosphatase/alkaline phosphodi-
esterase I activity

[124726]associated with the mouse plasma cell differentiation
antigen PC-1. Proc.

[124727]Nat. Acad. Sci. 88: 5192-5196, 1991.

[124728]22189.

[124729]Sakaguchi, A. Y.; Lalley, P. A.; Zabel, B. U.; Ellis, R. W.;
Scolnick,

[124730]E. M.; Naylor, S. L.: Chromosome assignments of four
mouse cellular

[124731]homologs of sarcoma and leukemia virus oncogenes.
Proc. Nat. Acad.

[124732]Sci. 81: 525–529, 1984.

[124733]22190.

[124734]Takahashi, T.; Old, L. J.; Boyse, E. A.: Surface alloantigens

[124735]of plasma cells. J. Exp. Med. 131: 1325–1341, 1970.

[124736]22191.

[124737]van Driel, I. R.; Goding, J. W.: Plasma cell membrane glycoprotein

[124738]PC-1: primary structure deduced from cDNA clones. J. Biol. Chem. 262:

[124739]4882–4887, 1987.

[124740]22192.

[124741]Frank, S. L.; Klisak, I.; Sparkes, R. S.; Lusk, A. J.: A gene

[124742]homologous to plasminogen located on human chromosome 2q11–p11. Genomics 4:

[124743]449–451, 1989.

[124744]22193.

[124745]Hughes, A. E.; Ralston, S. H.; Marken, J.; Bell, C.; MacPherson,

[124746]H.; Wallace, R. G. H.; van Hul, W.; Whyte, M. P.; Nakatsuka,

K.; Hovy,

[124747]L.; Anderson, D. M.: Mutations in TNFRSF11A, affecting
the signal

[124748]peptide of RANK, cause familial expansile osteolysis. Na-
ture Genet. 24:

[124749]45-48, 2000.

[124750]22194.

[124751]Hughes, A. E.; Shearman, A. M.; Weber, J. L.; Barr, R. J.;
Wallace,

[124752]R. G. H.; Osterberg, P. H.; Nevin, N. C.; Mollan, R. A. B.:
Genetic

[124753]linkage of familial expansile osteolysis to chromosome
18q. Hum.

[124754]Molec. Genet. 3: 359-361, 1994.

[124755]22195.

[124756]Brandriff, B. F.; Gordon, L. A.; Tynan, K. T.; Olsen, A. S.;
Mohrenweiser,

[124757]H. W.; Fertitta, A.; Carrano, A. V.; Trask, B. J.: Order and
genomic

[124758]distances among members of the carcinoembryonic anti-
gen (CEA) gene

[124759]family determined by fluorescence in situ hybridization.

Genomics 12:

[124760]773–779, 1992.

[124761]22196.

[124762]Teglund, S.; Olsen, A.; Khan, W. N.; Frangsmyr, L.; Hammarstrom,

[124763]S.: The pregnancy-specific glycoprotein (PSG) gene cluster on human

[124764]chromosome 19: fine structure of the 11 PSG genes and identification

[124765]of 6 new genes forming a third subgroup within the carcinoembryonic

[124766]antigen (CEA) family. Genomics 23: 669–684, 1994.

[124767]22197.

[124768]Thompson, J.; Koumari, R.; Wagner, K.; Barnert, S.; Schleussner,

[124769]C.; Schrewe, H.; Zimmermann, W.; Muller, G.; Schempp, W.; Zaninetta,

[124770]D.; Ammaturo, D.; Hardman, N.: The human pregnancy-specific glycoprotein

[124771]genes are tightly linked on the long arm of chromosome 19 and are

[124772]coordinately expressed. Biochem. Biophys. Res. Commun.

167: 848–859,
 [124773]1990.

[124774]22198.

[124775]Olsen, A.; Teglund, S.; Nelson, D.; Gordon, L.; Copeland,
 A.; Georgescu,
 [124776]A.; Carrano, A.; Hammarstrom, S.: Gene organization of
 the pregnancy-specific
 [124777]glycoprotein region on human chromosome 19: assembly
 and analysis
 [124778]of a 700-kb cosmid contig spanning the region. Genomics
 23: 659–668,
 [124779]1994.

[124780]22199.

[124781]Thompson, J.; Zimmermann, W.; Osthus–Bugat, P.;
 Schleussner, C.;

[124782]Eades–Perner, A.–M.; Barnert, S.; von Kleist, S.; Willcocks,
 T.; Craig,
 [124783]I.; Tynan, K.; Olsen, A.; Mohrenweiser, H.: Long-range
 chromosomal
 [124784]mapping of the carcinoembryonic antigen (CEA) gene
 family cluster. Genomics 12:
 [124785]761–772, 1992.

[124786]22200.

[124787]Tynan, K.; Olsen, A.; Trask, B.; de Jong, P.; Thompson, J.; Zimmermann,

[124788]W.; Carrano, A.; Mohrenweiser, H.: Assembly and analysis of cosmid

[124789]contigs in the CEA-gene family region of human chromosome 19. Nucleic

[124790]Acids Res. 20: 1629–1636, 1992.

[124791]22201.

[124792]Sheer, D.; Sheppard, D. M.; Le Beau, M.; Rowley, J. D.; San Roman,

[124793]C.; Solomon, E.: Localization of the oncogene c-erbA1 immediately

[124794]proximal to the acute promyelocytic leukaemia breakpoint on chromosome

[124795]17. Ann. Hum. Genet. 49: 167–171, 1985.

[124796]22202.

[124797]Yoshitake, H.; Rittling, S. R.; Denhardt, D. T.; Noda, M.: Osteopontin-deficient

[124798]mice are resistant to ovariectomy-induced bone resorption. Proc.

[124799]Nat. Acad. Sci. 96: 8156–8160, 1999.

[124800]22203.

[124801]Edelmann, W.; Zervas, M.; Costello, P.; Roback. L.; Fischer, I.;

[124802]Hammarback, J. A.; Cowan, N.; Davies, P.; Wainer, B.; Kucherlapati,

[124803]R.: Neuronal abnormalities in microtubule-associated protein 1B mutant

[124804]mice. Proc. Nat. Acad. Sci. 93: 1270–1275, 1996.

[124805]22204.

[124806]Hammarback, J. A.; Obar, R. A.; Hughes, S. M.; Vallee, R. B.:

[124807]MAP1B is encoded as a polyprotein that is processed to form a complex

[124808]N-terminal microtubule-binding domain. Neuron 7: 129–139, 1991.

[124809]22205.

[124810]Lien, L. L.; Boyce, F. M.; Kleyn, P.; Brzustowicz, L. M.; Menninger,

[124811]J.; Ward, D. C.; Gilliam, T. C.; Kunkel, L. M.: Mapping of human

[124812]microtubule-associated protein 1B in proximity to the
spinal muscular

[124813]atrophy locus at 5q13. Proc. Nat. Acad. Sci. 88:
7873–7876, 1991.

[124814]22206.

[124815]Lien, L. L.; Feener, C. A.; Fischbach, N.; Kunkel, L. M.:
Cloning

[124816]of human microtubule-associated protein 1B and the
identification

[124817]of a related gene on chromosome 15. Genomics 22:
273–280, 1994.

[124818]22207.

[124819]Wirth, B.; Voosen, B.; Rohrig, D.; Knapp, M.; Piechaczek,
B.; Rudnik-Schoneborn,

[124820]S.; Zerres, K.: Fine mapping and narrowing of the genetic
interval

[124821]of the spinal muscular atrophy region by linkage studies.
Genomics 15:

[124822]113–118, 1993.

[124823]22208.

[124824]Zhang, Y. Q.; Bailey, A. M.; Matthies, H. J. G.; Renden, R.
B.;

[124825]Smith, M. A.; Speese, S. D.; Rubin, G. M.; Broadie, K.:

Drosophila

[124826]fragile X-related gene regulates the MAP1B homolog

Futsch to control

[124827]synaptic structure and function. *Cell* 107: 591–603, 2001.

[124828]22209.

[124829]Garner, C. C.; Tucker, R. P.; Matus, A.: Selective localization

[124830]of messenger RNA for cytoskeletal protein MAP2 in dendrites. *Nature* 336:

[124831]674–677, 1988.

[124832]22210.

[124833]Kalcheva, N.; Albala, J.; O'Guin, K.; Rubino, H.; Garner, C.; Shafit-Zagardo,

[124834]B.: Genomic structure of human microtubule-associated protein 2 (MAP-2)

[124835]and characterization of additional MAP-2 isoforms. *Proc. Nat. Acad.*

[124836]*Sci.* 92: 10894–10898, 1995.

[124837]22211.

[124838]Kindler, S.; Garner, C. C.: Four repeat MAP2 isoforms in

human

[124839]and rat brain. Molec. Brain Res. 26: 218–224, 1994.

[124840]22212.

[124841]Marsden, K. M.; Doll, T.; Ferralli, J.; Botteri, F.; Matus, A.

[124842]: Transgenic expression of embryonic MAP2 in adult
mouse brain: implications

[124843]for neuronal polarization. J. Neurosci. 16: 3265–3273,
1996.

[124844]22213.

[124845]Neve, R. L.; Harris, P.; Kosik, K. S.; Kurnit, D. M.; Donlon,
T.

[124846]A.: Identification of cDNA clones for the human micro-
tubule-associated

[124847]protein tau and chromosomal localization of the genes for
tau and

[124848]microtubule-associated protein 2. Molec. Brain Res. 1:
271–280,

[124849]1986.

[124850]22214.

[124851]Chapin, S. J.; Bulinski, J. C.: Non-neuronal 210 x 10(3) M(r)

[124852]microtubule-associated protein (MAP4) contains a domain

homologous

[124853]to the microtubule-binding domains of neuronal MAP2
and tau. J. Cell

[124854]Sci. 98: 27-36, 1991.

[124855]22215.

[124856]Adriaansen, H. J.; Geurts Van Kessel, A. H. M.; Wijdenes-
De Bresser,

[124857]J. H. F. M.; Van Drunen-Schoenmaker, E.; Van Dongen, J. J.
M.: Expression

[124858]of the myeloid differentiation antigen CD33 depends on
the presence

[124859]of human chromosome 19 in human-mouse hybrids. Ann.
Hum. Genet. 54:

[124860]115-119, 1990.

[124861]22216.

[124862]Bonthron, D. T.; Dunlop, N.; Barr, D. G. D.; El Sanousi, A.
A.;

[124863]Al-Gazali, L. I.: Organisation of the human PAX4 gene and
its exclusion

[124864]as a candidate for the Wolcott-Rallison syndrome. J. Med.
Genet. 35:

[124865]288-292, 1998.

[124866]22217.

[124867]Inoue, H.; Nomiyama, J.; Nakai, K.; Matsutani, A.;
Tanizawa, Y.;

[124868]Oka, Y.: Isolation of full-length cDNA of mouse PAX4 gene
and identification

[124869]of its human homologue. Biochem. Biophys. Res. Com-
mun. 243: 628-633,

[124870]1998.

[124871]22218.

[124872]Mansouri, A.; St-Onge, L.; Gruss, P.: Role of Pax genes in
endoderm-derived

[124873]organs. Trends Endocr. Metab. 10: 164-167, 1999.

[124874]22219.

[124875]Matsushita, T.; Yamaoka, T.; Otsuka, S.; Moritani, M.; Mat-
sumoto,

[124876]T.; Itakura, M.: Molecular cloning of mouse paired-
box-containing

[124877]gene (Pax)-4 from an islet beta cell line and deduced se-
quence of

[124878]human Pax-4. Biochem. Biophys. Res. Commun. 242:
176-180, 1998.

[124879]22220.

[124880]Sosa-Pineda, B.; Chowdhury, K.; Torres, M.; Oliver, G.;
Gruss,

[124881]P.: The Pax4 gene is essential for differentiation of in-
sulin-producing

[124882]beta cells in the mammalian pancreas. Nature 386:
399-402, 1997.

[124883]22221.

[124884]St-Onge, L.; Sosa-Pineda, B.; Chowdhury, K.; Mansouri, A.;
Gruss,

[124885]P.: Pax6 is required for differentiation of glucagon-pro-
ducing alpha-cells

[124886]in mouse pancreas. Nature 387: 406-409, 1997.

[124887]22222.

[124888]Tamura, T.; Izumikawa, Y.; Kishino, T.; Soejima, H.; Jinno,
Y.;

[124889]Niikawa, N.: Assignment of the human PAX4 gene to chro-
mosome band

[124890]7q32 by fluorescence in situ hybridization. Cytogenet. Cell
Genet. 66:

[124891]132-134, 1994.

[124892]22223.

[124893]Adams, B.; Dorfler, P.; Aguzzi, A.; Kozmik, Z.; Urbanek, P.;
Maurer-Fogy,

[124894]I.; Busslinger, M.: Pax-5 encodes the transcription factor
BSAP and

[124895]is expressed in B lymphocytes, the developing CNS, and
adult testis. Genes

[124896]Dev. 6: 1589-1607, 1992.

[124897]22224.

[124898]Busslinger, M.; Klix, N.; Pfeffer, P.; Graninger, P. G.;
Kozmik,

[124899]Z.: Deregulation of PAX-5 by translocation of the E- μ
enhancer of

[124900]the IgH locus adjacent to two alternative PAX-5 promoters
in a diffuse

[124901]large-cell lymphoma. Proc. Nat. Acad. Sci. 93:
6129-6134, 1996.

[124902]22225.

[124903]Mikkola, I.; Heavey, B.; Horcher, M.; Busslinger, M.: Rever-
sion

[124904]of B cell commitment upon loss of Pax5 expression. Sci-
ence 297:

[124905]110–113, 2002.

[124906]22226.

[124907]Nutt, S. L.; Heavey, B.; Rolink, A. G.; Busslinger, M.: Com-
mitment

[124908]to the B-lymphoid lineage depends on the transcription
factor Pax5. Nature 401:

[124909]556–562, 1999.

[124910]22227.

[124911]Nutt, S. L.; Vambrie, S.; Steinlein, P.; Kozmik, Z.; Rolink,
A.;

[124912]Weith, A.; Busslinger, M.: Independent regulation of the
two Pax5

[124913]alleles during B-cell development. Nature Genet. 21:
390–395, 1999.

[124914]22228.

[124915]Ohno, H.; Furukawa, T.; Fukuhara, S.; Zong, S. Q.; Kame-
saki, H.;

[124916]Shows, T. B.; Le Beau, M. M.; McKeithan, T. W.; Kawakami,
T.; Honjo,

[124917]T.: Molecular analysis of a chromosomal translocation,
t(9;14)(p13;q32),

[124918]in a diffuse large-cell lymphoma cell line expressing the
Ki-1 antigen. Proc.

[124919]Nat. Acad. Sci. 87: 628-632, 1990.

[124920]22229.

[124921]Rolink, A. G.; Nutt, S. L.; Melchers, F.; Busslinger, M.:
Long-term

[124922]in vivo reconstitution of T-cell development by
Pax5-deficient B-cell

[124923]progenitors. Nature 401: 603-606, 1999.

[124924]22230.

[124925]Vorechovsky, I.; Koskinen, S.; Paganelli, R.; Smith, C. I. E.;

[124926]Busslinger, M.; Hammarstrom, L.: The PAX5 gene: a link-
age and mutation

[124927]analysis in candidate human primary immunodeficiencies.
Immunogenetics 42:

[124928]149-152, 1995.

[124929]22231.

[124930]Walther, C.; Guenet, J.-L.; Simon, D.; Deutsch, U.; Jostes,
B.;

[124931]Goulding, M. D.; Plachov, D.; Balling, R.; Gruss, P.: Pax: a
murine

[124932]multigene family of paired box containing genes. Genomics 11: 424-434,

[124933]1991.

[124934]22232.

[124935]Koseki, H.; Zachgo, J.; Mizutani, Y.; Simon-Chazottes, D.; Guenet,

[124936]J.-L.; Balling, R.; Gossler, A.: Fine genetic mapping of the proximal

[124937]part of mouse chromosome 2 excludes Pax-8 as a candidate gene for

[124938]Danforth's short tail (Sd). Mammalian Genome 4: 324-327, 1993.

[124939]22233.

[124940]Kroll, T. G.; Sarraf, P.; Pecciarini, L.; Chen, C.-J.; Mueller,

[124941]E.; Splegeman, B. M.; Fletcher, J. A.:

PAX8-PPAR-gamma-1 fusion

[124942]in oncogene human thyroid carcinoma. Science 289: 1357-1360, 2000.

[124943]22234.

[124944]Macchia, P. E.; Lapi, P.; Krude, H.; Pirro, M. T.; Missero, C.;

[124945]Chiovato, L.; Souabni, A.; Baserga, M.; Tassi, V.; Pinchera,

A.; Fenzi,

[124946]G.; Gruters, A.; Busslinger, M.; Di Lauro, R.: PAX8 mutations associated

[124947]with congenital hypothyroidism caused by thyroid dysgenesis. Nature

[124948]Genet. 19: 83–86, 1998.

[124949]22235.

[124950]Mansouri, A.; Chowdhury, K.; Gruss, P.: Follicular cells of the

[124951]thyroid gland require Pax8 gene function. Nature Genet. 19: 87–90,

[124952]1998.

[124953]22236.

[124954]Pasca di Magliano, M.; Di Lauro, R.; Zannini, M.: Pax8 has a key

[124955]role in thyroid cell differentiation. Proc. Nat. Acad. Sci. 97:

[124956]13144–13149, 2000.

[124957]22237.

[124958]Plachov, D.; Chowdhury, K.; Walther, C.; Simon, D.; Guenet, J.–L.;

[124959]Gruss, P.: Pax–8, a murine paired box gene expressed in

the developing

[124960]excretory system and thyroid gland. Development 110:
643–651, 1990.

[124961]22238.

[124962]Tell, G.; Pellizzari, L.; Esposito, G.; Pucillo, C.; Macchia,

[124963]P. E.; Di Lauro, R.; Damante, G.: Structural defects of a
Pax8 mutant

[124964]that give rise to congenital hypothyroidism. Biochem. J.
341: 89–93,

[124965]1999.

[124966]22239.

[124967]Bongarzone, I.; Vigano, E.; Alberti, L.; Borrello, M. G.;

Pasini,

[124968]B.; Greco, A.; Mondellini, P.; Smith, D. P.; Ponder, B. A. J.;

Romeo,

[124969]G.; Pierotti, M. A.: Full activation of MEN2B mutant RET by
an additional

[124970]MEN2A mutation or by ligand GDNF stimulation. Onco-
gene 16: 2295–2301,

[124971]1998.

[124972]22240.

[124973] Carlson, K. M.; Bracamontes, J.; Jackson, C. E.; Clark, R.;
Lacroix,
[124974] A.; Wells, S. A., Jr.; Goodfellow, P. J.: Parent-of-origin ef-
fects
[124975] in multiple endocrine neoplasia type 2B. Am. J. Hum.
Genet. 55:
[124976] 1076–1082, 1994.
[124977] 22241.
[124978] Carlson, K. M.; Dou, S.; Chi, D.; Scavarda, N.; Toshima, K.;
Jackson,
[124979] C. E.; Wells, S. A., Jr.; Goodfellow, P. J.; Donis-Keller, H.:
Single
[124980] missense mutation in the tyrosine kinase catalytic domain
of the RET
[124981] protooncogene is associated with multiple endocrine neo-
plasia type
[124982] 2B. Proc. Nat. Acad. Sci. 91: 1579–1583, 1994.
[124983] 22242.
[124984] Mulligan, L. M.; Eng, C.; Healey, C. S.; Clayton, D.; Kwok, J.
[124985] B. J.; Gardner, E.; Ponder, M. A.; Frilling, A.; Jackson, C. E.;
Lehnert,

[124986]H.; Neumann, H. P. H.; Thibodeau, S. N.; Ponder, B. A. J.:

Specific

[124987]mutations of the RET proto-oncogene are related to disease phenotype

[124988]in MEN 2A and FMTC. Nature Genet. 6: 70–74, 1994.

[124989]22243.

[124990]Tartaglia, M.; Mehler, E. L.; Goldberg, R.; Zampino, G.; Brunner,

[124991]H. G.; Kremer, H.; van der Burgt, I.; Crosby, A. H.; Ion, A.; Jeffery,

[124992]S.; Kalidas, K.; Patton, M. A.; Kucherlapati, R. S.; Gelb, B. D.:

[124993]Mutations in PTPN11, encoding the protein tyrosine phosphatase SHP-2,

[124994]cause Noonan syndrome. Nature Genet. 29: 465–468, 2001. Note: Errata:

[124995]Nature Genet. 29: 491 only, 2001; Nature Genet. 30: 123 only, 2002.

[124996]22244.

[124997]Bruss, M.; Kunz, J.; Lingen, B.; Bonisch, H.: Chromosomal mapping

[124998]of the human gene for the tricyclic antidepressant-sensi-

tive noradrenaline

[124999] transporter. Hum. Genet. 91: 278–280, 1993.

[125000] 22245.

[125001] Esler, M.; Jennings, G.; Lambert, G.; Meredith, I.; Horne, M.;

[125002] Eisenhofer, G.: Overflow of catecholamine neurotransmitters to the

[125003] circulation: source, fate, and functions. Physiol. Rev. 70: 963–985,

[125004] 1990.

[125005] 22246.

[125006] Fritz, J. D.; Jayanthi, L. D.; Thoreson, M. A.; Blakely, R. D.

[125007]: Cloning and chromosomal mapping of the murine norepinephrine transporter. J.

[125008] Neurochem. 70: 2241–2251, 1998.

[125009] 22247.

[125010] Gelernter, J.; Kruger, S.; Kidd, K. K.; Amara, S.: TaqI RFLP at

[125011] norepinephrine transporter protein (NET) locus. Hum. Molec. Genet. 2:

[125012] 820 only, 1993.

[125013]22248.

[125014]Gelernter, J.; Kruger, S.; Pakstis, A. J.; Pacholczyk, T.;
Sparkes,

[125015]R. S.; Kidd, K. K.; Amara, S.: Assignment of the nore-
pinephrine transporter

[125016]protein (NET1) locus to chromosome 16. Genomics 18:
690-692, 1993.

[125017]22249.

[125018]Kaye, W. H.; Jimerson, D. C.; Lake, C. R.; Ebert, M. H.: Al-
tered

[125019]norepinephrine metabolism following long-term weight
recovery in patients

[125020]with anorexia nervosa. Psychiat. Res. 14: 333-342, 1985.

[125021]22250.

[125022]Pacholczyk, T.; Blakely, R. D.; Amara, S. G.: Expression
cloning

[125023]of a cocaine- and antidepressant-sensitive human nora-
drenaline transporter. Nature 350:

[125024]350-354, 1991.

[125025]22251.

[125026]Paczkowski, F. A.; Bonisch, H.; Bryan-Lluka, L. J.: Pharma-

cological

[125027]properties of the naturally occurring ala457pro variant of the human

[125028]norepinephrine transporter. Pharmacogenetics 12: 165–173, 2002.

[125029]22252.

[125030]Pirke, K. M.; Kellner, M.; Philipp, E.; Laessle, R.; Krieg, J.

[125031]C.; Fichter, M. M.: Plasma norepinephrine after a standardized test

[125032]meal in acute and remitted patients with anorexia nervosa and in healthy

[125033]controls. Biol. Psychiat. 31: 1074–1077, 1992.

[125034]22253.

[125035]Porzgen, P.; Bonisch, H.; Bruss, M.: Molecular cloning and organization

[125036]of the coding region of the human norepinephrine transporter gene. Biochem.

[125037]Biophys. Res. Commun. 215: 1145–1150, 1995. Notes: Erratum: Biochem.

[125038]Biophys. Res. Commun. 227: 642–643, 1996.

[125039]22254.

[125040]Shannon, J. R.; Flattem, N. L.; Jordan, J.; Jacob, G.; Black,

[125041]B. K.; Biaggioni, I.; Blakely, R. D.; Robertson, D.: Ortho-
static

[125042]intolerance and tachycardia associated with nore-
pinephrine-transporter

[125043]deficiency. New Eng. J. Med. 342: 541-549, 2000.

[125044]22255.

[125045]Urwin, R. E.; Bennetts, B.; Wilcken, B.; Lampropoulos, B.;
Beumont,

[125046]P.; Clarke, S.; Russell, J.; Tanner, S.; Nunn, K. P.: Anorexia
nervosa

[125047](restrictive subtype) is associated with a polymorphism in
the novel

[125048]norepinephrine transporter gene promoter polymorphic
region. Molec.

[125049]Psychiat. 7: 652-657, 2002.

[125050]22256.

[125051]Tomilin, A.; Remenyi, A.; Lins, K.; Bak, H.; Leidel, S.;
Vriend,

[125052]G.; Wilmanns, M.; Scholer, H. R.: Synergism with the coac-
tivator

[125053]OBF-1 (OCA-B, BOB-1) is mediated by a specific POU dimer configuration. Cell 103:

[125054]853-864, 2000.

[125055]22257.

[125056]Ko, H.-S.; Fast, P.; McBride, W.; Staudt, L. M.: A human protein

[125057]specific for the immunoglobulin octamer DNA motif contains a functional

[125058]homeobox domain. Cell 55: 135-144, 1988.

[125059]22258.

[125060]Schubart, K.; Massa, S.; Schubart, D.; Corcoran, L. M.; Rolink,

[125061]A. G.; Matthias, P.: B cell development and immunoglobulin gene transcription

[125062]in the absence of Oct-2 and OBF-1. Nature Immun. 2: 69-74, 2001.

[125063]22259.

[125064]Staudt, L. M.; Clerc, R. G.; Singh, H.; LeBowitz, J. H.; Sharp,

[125065]P. A.; Baltimore, D.: Cloning of a lymphoid-specific cDNA encoding

[125066]a protein binding the regulatory octamer DNA motif. Sci-

ence 241:

[125067]577–580, 1988.

[125068]22260.

[125069]Parker, R. C.; Mardon, G.; Lebo, R. V.; Varmus, H. E.;
Bishop,

[125070]J. M.: Isolation of duplicated human c-src genes located
on chromosomes

[125071]1 and 20. Molec. Cell. Biol. 5: 831–838, 1985.

[125072]22261.

[125073]Tronick, S. R.; Popescu, N. C.; Cheah, M. S. C.; Swan, D. C.;
Amsbaugh,

[125074]S. C.; Lengel, C. R.; DiPaolo, J. A.; Robbins, K. C.: Isolation
and

[125075]chromosomal localization of the human fgr protoonco-
gene, a distinct

[125076]member of the tyrosine kinase gene family. Proc. Nat.
Acad. Sci. 82:

[125077]6595–6599, 1985.

[125078]22262.

[125079]Bork, P.: The modular architecture of a new family of
growth regulators

[125080]related to connective tissue growth factor. FEBS Lett. 327:
125-130,

[125081]1993.

[125082]22263.

[125083]Sugaya, K.; Fukagawa, T.; Matsumoto, K.; Mita, K.; Takahashi, E.;

[125084]Ando, A.; Inoko, H.; Ikemura, T.: Three genes in the human MHC class

[125085]III region near the junction with the class II: gene for receptor

[125086]of advanced glycosylation end products, PBX2 homeobox gene and a Notch

[125087]homolog, human counterpart of mouse mammary tumor gene int-3. Genomics 23:

[125088]408-419, 1994.

[125089]22264.

[125090]Joliot, V.; Martinerie, C.; Dambrine, G.; Plassiart, G.; Brisac,

[125091]M.; Crochet, J.; Perbal, B.: Proviral rearrangements and overexpression

[125092]of a new cellular gene (nov) in myeloblastosis-associated virus type

[125093]1-induced nephroblastomas. *Molec. Cell. Biol.* 12: 10-21, 1992.

[125094]22265.

[125095]Kim, H.-S.; Nagalla, S. R.; Oh, Y.; Wilson, E.; Roberts, C. T., Jr.; Rosenfeld, R. G.: Identification of a family of low-affinity

[125097]insulin-like growth factor binding proteins (IFGBPs): characterization

[125098]of connective tissue growth factor as a member of the IGFBP superfamily. *Proc.*

[125099]*Nat. Acad. Sci.* 94: 12981-12986, 1997.

[125100]22266.

[125101]Martinerie, C.; Perbal, B.: Expression of a gene encoding a novel

[125102]potential IGF binding protein in human tissues. *C. R. Acad. Sci.*

[125103](Paris) 313 (ser. 3): 345-351, 1991.

[125104]22267.

[125105]Snaith, M. R.; Natarajan, D.; Taylor, L. B.; Choi, C.-P.; Martinerie,

[125106]C.; Perbal, B.; Schofield, P. N.; Boulter, C. A.: Genomic

structure

[125107]and chromosomal mapping of the mouse nov gene. Genomics 38: 425–428,

[125108]1996.

[125109]22268.

[125110]Soret, J.; Dambrine, G.; Perbal, B.: Induction of nephroblastoma

[125111]by myeloblastosis-associated virus type 1: state of proviral DNAs

[125112]in tumor cells. J. Virol. 63: 1803–1807, 1989.

[125113]22269.

[125114]Amson, R.; Sigaux, F.; Przedborski, S.; Flandrin, G.; Givol, D.;

[125115]Telerman, A.: The human protooncogene product p33pim is expressed

[125116]during fetal hematopoiesis and in diverse leukemias. Proc. Nat. Acad.

[125117]Sci. 86: 8857–8861, 1989.

[125118]22270.

[125119]Ark, B.; Gummere, G.; Bennett, D.; Artzt, K.: Mapping of the Pim-1

[125120]oncogene in mouse t-haplotypes and its use to define the relative

[125121]map positions of the tcl loci t-0(t-6) and t-w12 and the marker tf

[125122](tufted). Genomics 10: 385-389, 1991.

[125123]22271.

[125124]Cuypers, H. T.; Selten, G.; Berns, A.; Geurts van Kessel, A. H.

[125125]M.: Assignment of the human homologue of Pim-1, a mouse gene implicated

[125126]in leukemogenesis, to the pter-q12 region of chromosome 6. Hum. Genet. 72:

[125127]262-265, 1986.

[125128]22272.

[125129]Domen, J.; von Lindern, M.; Hermans, A.; Breuer, M.; Grosveld,

[125130]G.; Berns, A.: Comparison of the human and mouse PIM-1 cDNAs: nucleotide

[125131]sequence and immunological identification of the in vitro synthesized

[125132]PIM-1 protein. Oncogene Res. 1: 103-112, 1987.

[125133]22273.

[125134]Dunger, D. B.; Davies, K. E.; Pembrey, M.; Lake, B.; Pearson,
son,

[125135]P.; Williams, D.; Whitfield, A.; Dillon, M. J. D.: Deletion on
the

[125136]X chromosome detected by direct DNA analysis in one of
two unrelated

[125137]boys with glycerol kinase deficiency, adrenal hypoplasia,
and Duchenne

[125138]muscular dystrophy. Lancet I: 585-587, 1986.

[125139]22274.

[125140]Eriksson, A.; Lindstedt, S.; Ransnas, L.; von Wendt, L.: De-
ficiency

[125141]of glycerol kinase (EC 2.7.1.30). Clin. Chem. 29: 718-722,
1983.

[125142]22275.

[125143]Francke, U.; Harper, J. F.; Darras, B. T.; Cowan, J. M.; Mc-
Cabe,

[125144]E. R. B.; Kohlschutter, A.; Seltzer, W. K.; Saito, F.; Goto, J.;
Harpey,

[125145]J.-P.; Wise, J. E.: Congenital adrenal hypoplasia, myopathy,
and

[125146]glycerol kinase deficiency: molecular genetic evidence for deletions. Am.

[125147]J. Hum. Genet. 40: 212–227, 1987.

[125148]22276.

[125149]Gaudet, D.; Arsenault, S.; Perusse, L.; Vohl, M.–C.; St.–Pierre,

[125150]J.; Bergeron, J.; Despres, J.–P.; Dewar, K.; Daly, M. J.; Hudson,

[125151]T.; Rioux, J. D.: Glycerol as a correlate of impaired glucose tolerance:

[125152]dissection of a complex system by use of a simple genetic trait. Am.

[125153]J. Hum. Genet. 66: 1558–1568, 2000.

[125154]22277.

[125155]Ginns, E. I.; Barranger, J. A.; McClean, S. W.; Sliva, C.; Young,

[125156]R.; Schaefer, I.; Goodman, S. I.; McCabe, E. R. B.: A juvenile form

[125157]of glycerol kinase deficiency with episodic vomiting, acidemia, and

[125158]stupor. J. Pediat. 104: 736–739, 1984.

[125159]22278.

[125160]Goonewardena, P.; Dahl, N.; Ritzen, M.; Johnsen, E.; Petersson,

[125161]U.: Deletion in Xp associated with glycerol kinase deficiency, adrenal

[125162]aplasia and hypogonadotropic hypogonadism. (Abstract) Cytogenet.

[125163]Cell Genet. 46: 621 only, 1987.

[125164]22279.

[125165]Goussault, Y.; Turpin, E.; Neel, D.; Dreux, C.; Chanu, B.; Bakir,

[125166]R.; Rouffy, J.: 'Pseudohypertriglyceridemia' caused by hyperglycerolemia

[125167]due to congenital enzyme deficiency. Clin. Chim. Acta 123: 269-274,

[125168]1982.

[125169]22280.

[125170]Grier, R. E.; Howell, R. R.; Wu, D.; McCabe, E. R. B.: Isolated

[125171]glycerol kinase deficiency: nutritional and molecular genetic studies.

[125172](Abstract) Am. J. Hum. Genet. 45 (suppl.): A6 only, 1989.

[125173]22281.

[125174]Guggenheim, M. A.; McCabe, E. R. B.; Roig, M.; Goodman, S. I.;

[125175]Lum, G. M.; Bullen, W. W.; Ringel, S. P.: Glycerol kinase deficiency

[125176]with neuromuscular, skeletal, and adrenal abnormalities. Ann. Neurol. 7:

[125177]441–449, 1980.

[125178]22282.

[125179]Guo, W.; Worley, K.; Adams, V.; Mason, J.; Sylvester–Jackson,

[125180]D.; Zhang, Y.–H.; Towbin, J. A.; Fogt, D. D.; Madu, S.; Wheeler, D.

[125181]A.; McCabe, E. R. B.: Genomic scanning for expressed sequences in

[125182]Xp21 identifies the glycerol kinase gene. Nature Genet. 4: 367–371,

[125183]1993.

[125184]22283.

[125185]Hammond, J.; Howard, N. J.; Brookwell, R.; Purvis–Smith, S.; Wilcken,

- [125186]B.; Hoogenraad, N.: Proposed assignment of loci for X-linked adrenal hypoplasia and glycerol kinase genes. (Letter) Lancet I: 54 only, 1985.
- [125189]22284.
- [125190]Love, D. R.; Bloomfield, J. F.; Kenwrick, S. J.; Yates, J. R. W.; Davies, K. E.: Physical mapping distal to the DMD locus. Genomics 8: 106–112, 1990.
- [125193]22285.
- [125194]Mahbubul Huq, A. H. M.; Lovell, R. S.; Ou, C.-N.; Beaudet, A. L.; Craigen, W. J.: X-linked glycerol kinase deficiency in the mouse leads to growth retardation, altered fat metabolism, autonomous glucocorticoid secretion and neonatal death. Hum. Molec. Genet. 6: 1803–1809, 1997.
- [125198]22286.
- [125199]Mahbubul Huq, A. H. M.; Lovell, R. S.; Sampson, M. J.;

Decker,

[125200]W. K.; Dinulos, M. B.; Disteché, C. M.; Craigen, W. J.: Isolation,

[125201]mapping, and functional expression of the mouse X chromosome glycerol

[125202]kinase gene. Genomics 36: 530–534, 1996.

[125203]22287.

[125204]Nishimura, J.; Murakami, Y.; Kinoshita, T.: Paroxysmal nocturnal

[125205]hemoglobinuria: an acquired genetic disease. Am. J. Hemat. 62: 175–182,

[125206]1999.

[125207]22288.

[125208]Kosho, T.; Muroya, K.; Nagai, T.; Fujimoto, M.; Yokoya, S.; Sakamoto,

[125209]H.; Hirano, T.; Terasaki, H.; Ohashi, H.; Nishimura, G.; Sato, S.;

[125210]Matsuo, N.; Ogata, T.: Skeletal features and growth patterns in 14

[125211]patients with haploinsufficiency of SHOX: implications for the development

[125212]of Turner syndrome. J. Clin. Endocr. Metab. 84:

4613–4621, 1999.

[125213]22289.

[125214]Kuznetzova, T.; Baranov, A.; Ivaschenko, T.; Savitsky, G.
A.;

[125215]Lanceva, O. E.; Wang, M. R.; Giollant, M.; Malet, P.;
Kascheeva, T.;

[125216]Vakharlovsky, V.; Baranov, V. S.: X;Y translocation in a girl
with

[125217]short stature and some features of Turner's syndrome: cy-
togenetic

[125218]and molecular studies. J. Med. Genet. 31: 649–651, 1994.

[125219]22290.

[125220]May, C. A.; Shone, A. C.; Kalaydjieva, L.; Sajantila, A.; Jef-
freys,

[125221]A. J.: Crossover clustering and rapid decay of linkage dis-
equilibrium

[125222]in the Xp/Yp pseudoautosomal gene SHOX. Nature Genet.
31: 272–275,

[125223]2002.

[125224]22291.

[125225]Ogata, T.; Goodfellow, P.; Petit, C.; Aya, M.; Matsuo, N.:

Short

[125226] stature in a girl with a terminal Xp deletion distal to

DXYS15: localisation

[125227] of a growth gene(s) in the pseudoautosomal region. J.

Med. Genet. 29:

[125228] 455–459, 1992.

[125229] 22292.

[125230] Ogata, T.; Kosho, T.; Wakui, K.; Fukushima, Y.; Yoshimoto,

M.;

[125231] Miharu, N.: Short stature homeobox-containing gene du-

plication on

[125232] the der(X) chromosome in a female with 45,X/46,X, der(X),

gonadal

[125233] dysgenesis, and tall stature. J. Clin. Endocr. Metab. 85:

2927–2930,

[125234] 2000.

[125235] 22293.

[125236] Ogata, T.; Matsuo, N.; Nishimura, G.: SHOX haploinsuffi-

ciency

[125237] and overdosage: impact of gonadal function status. J.

Med. Genet. 38:

[125238] 1–6, 2001.

[125239]22294.

[125240]Ogata, T.; Muroya, K.; Sasaki, G.; Nishimura, G.; Kitoh, H.;
Hattori,

[125241]T.: SHOX nullizygosity and haploinsufficiency in a
Japanese family:

[125242]implication for the development of Turner skeletal fea-
tures. J. Clin.

[125243]Endocr. Metab. 87: 1390–1394, 2002.

[125244]22295.

[125245]Ogata, T.; Petit, C.; Rappold, G.; Matsuo, N.; Matsumoto,
T.;

[125246]Goodfellow, P.: Chromosomal localisation of a pseudoau-
tosomal growth

[125247]gene(s). J. Med. Genet. 29: 624–628, 1992.

[125248]22296.

[125249]Ogata, T.; Yoshizawa, A.; Muroya, K.; Matsuo, N.;
Fukushima, Y.;

[125250]Rappold, G.; Yokoya, S.: Short stature in a girl with partial
monosomy

[125251]of the pseudoautosomal region distal to DXYS15: further
evidence for

[125252]the assignment of the critical region for a pseudoautosomal growth

[125253]gene(s). J. Med. Genet. 32: 831–834, 1995.

[125254]22297.

[125255]Pfeiffer, R. A.: Observations in a case of an X/Y translocation,

[125256]t(X;Y)(p22;q11), in a mother and son. Cytogenet. Cell Genet. 26:

[125257]150–157, 1980.

[125258]22298.

[125259]Rao, E.; Blaschke, R. J.; Marchini, A.; Niesler, B.; Burnett,

[125260]M.; Rappold, G. A.: The Leri-Weill and Turner syndrome homeobox gene

[125261]SHOX encodes a cell-type specific transcriptional activator. Hum.

[125262]Molec. Genet. 10: 3083–3091, 2001.

[125263]22299.

[125264]Rao, E.; Weiss, B.; Fukami, M.; Mertz, A.; Meder, J.; Ogata, T.;

[125265]Heinrich, U.; Garcia-Heras, J.; Schiebel, K.; Rappold, G. A.: FISH-deletion

[125266] mapping defines a 270-kb short stature critical interval in the pseudoautosomal

[125267] region PAR1 on human sex chromosomes. Hum. Genet. 100: 236–239,

[125268] 1997.

[125269] 22300.

[125270] Rao, E.; Weiss, B.; Fukami, M.; Rump, A.; Niesler, B.; Mertz, A.; Muroya, K.; Binder, G.; Kirsch, S.; Winkelmann, M.; Nordsiek,

[125272] G.; Heinrich, U.; Breuning, M. H.; Ranke, M. B.; Rosenthal, A.; Ogata,

[125273] T.; Rappold, G. A.: Pseudoautosomal deletions encompassing a novel homeobox gene cause growth failure in idiopathic short stature and

[125274] Turner syndrome. Nature Genet. 16: 54–63, 1997.

[125275] 22301.

[125276] Rappold, G. A.; Fukami, M.; Niesler, B.; Schiller, S.; Zumkeller,

[125278] W.; Bettendorf, M.; Heinrich, U.; Vlachopapadoupoulou, E.; Reinehr,

[125279]T.; Onigata, K.; Ogata, T.: Deletions of the homeobox gene SHOX (short stature homeobox) are an important cause of growth failure in children with short stature. J. Clin. Endocr. Metab. 87: 1402–1406, 2002.

[125282]22302.

[125283]Schiller, S.; Spranger, S.; Schechinger, B.; Fukami, M.; Merker, S.; Drop, S. L. S.; Troger, J.; Knoblauch, H.; Kunze, J.; Seidel, J.; Rappold, G. A.: Phenotypic variation and genetic heterogeneity in Leri–Weill syndrome. Europ. J. Hum. Genet. 8: 54–62, 2000.

[125287]22303.

[125288]Shears, D. J.; Guillen–Navarro, E.; Sempere–Miralles, M.; Domingo–Jimenez, R.; Scambler, P. J.; Winter, R. M.: Pseudodominant inheritance of Langer mesomelic dysplasia caused by a SHOX homeobox missense mutation. Am.

[125291]J. Med. Genet. 110: 153–157, 2002.

[125292]22304.

[125293]Zinn, A. R.; Wei, F.; Zhang, L.; Elder, F. F.; Scott, C. I., Jr.;

[125294]Marttila, P.; Ross, J. L.: Complete SHOX deficiency causes
Langer

[125295]mesomelic dysplasia. Am. J. Med. Genet. 110: 158–163,
2002.

[125296]22305.

[125297]Zuffardi, O.; Maraschio, P.; Lo Curto, F.; Muller, U.; Gia-
rola,

[125298]A.; Perotti, L.: The role of Yp in sex determination: new
evidence

[125299]from X/Y translocations. Am. J. Med. Genet. 12: 175–184,
1982.

[125300]22306.

[125301]Acker, K. J.; Roels, H.; Beelaerts, W.; Pasternack, A.; Val-
cke,

[125302]R.: The histologic lesions of the kidney in the oculo-
cerebro-renal

[125303]syndrome of Lowe. Nephron 4: 193–214, 1967.

[125304]22307.

[125305]Attree, O.; Olivos, I. M.; Okabe, I.; Bailey, L. C.; Nelson, D.

[125306]L.; Lewis, R. A.; McInnes, R. R.; Nussbaum, R. L.: The
Lowe's oculocerebrorenal

[125307]syndrome gene encodes a protein highly homologous to
inositol polyphosphate-5-phosphatase. Nature 358:

[125308]239-242, 1992.

[125309]22308.

[125310]Auricchio, S.; Frischknecht, W.; Shmerling, D. H.: Primare
Tubulopathien.

[125311]III. Ein Fall von oculo-cerebro-renalem Syndrom
(Lowe-Syndrome). Helv.

[125312]Paediat. Acta 16: 647-655, 1961.

[125313]22309.

[125314]Bailey, L. C., Jr.; Olivos, I. M.; Leahey, A. M.; Attree, O. F.;

[125315]Okabe, I.; Lewis, R. A.; MacInnes, R. R.; Spinner, N. B.; Nel-
son,

[125316]D. L.; Nussbaum, R. L.: Characterization of a candidate
gene for

[125317]OCRL. (Abstract) Am. J. Hum. Genet. 51 (suppl.): A4 only,
1992.

[125318]22310.

[125319]Charnas, L. R.; Bernardini, I.; Rader, D.; Hoeg, J. M.; Gahl, W.

[125320]A.: Clinical and laboratory findings in the oculocerebrorenal syndrome

[125321]of Lowe, with special reference to growth and renal function. New

[125322]Eng. J. Med. 324: 1318–1325, 1991.

[125323]22311.

[125324]Chutorian, A.; Rowland, L. P.: Lowe's syndrome. Neurology 16:

[125325]115–122, 1966.

[125326]22312.

[125327]Delleman, J. W.; Bleeker-Wagemakers, E. M.; van Veelen, A. W. C.

[125328]: Opacities of the lens indicating carrier status in the oculo-cerebro-renal

[125329](Lowe) syndrome. J. Pediat. Ophthal. 14: 205–212, 1977.

[125330]22313.

[125331]Duden, R.; Griffiths, G.; Frank, R.; Argos, P.; Kreis, T. E.:

[125332]Beta-COP, a 110 kd protein associated with non-clathrin-coated vesicles

[125333]and the Golgi complex, shows homology to beta-adaptin.

Cell 64:

[125334]649–665, 1991.

[125335]22314.

[125336]Gardner, R. J. M.; Brown, N.: Lowe's syndrome: identification

[125337]of carriers by lens examination. J. Med. Genet. 13:
449–464, 1976.

[125338]22315.

[125339]Gropman, A.; Levin, S.; Yao, L.; Lin, T.; Suchy, S.; Sabnis,
S.;

[125340]Hadley, D.; Nussbaum, R.: Unusual renal features of Lowe
syndrome

[125341]in a mildly affected boy. Am. J. Med. Genet. 95: 461–466,
2000.

[125342]22316.

[125343]Harris, L. S.; Gitter, K. A.; Galin, M. A.; Plechaty, G. P.:

[125344]Oculo–cerebro–renal syndrome: report of a case in a baby
girl. Brit.

[125345]J. Ophthalm. 54: 278–280, 1970.

[125346]22317.

[125347]Hittner, H. M.; Carroll, A. J.; Prchal, J. T.: Linkage studies
 [125348]in carriers of Lowe oculo-cerebro-renal syndrome. Am. J.
 Hum. Genet. 34:
 [125349]966-971, 1982.
 [125350]22318.

[125351]Hodgson, S. V.; Heckmatt, J. Z.; Hughes, E.; Crolla, J. A.;
 Dubowitz,
 [125352]V.; Bobrow, M.: A balanced de novo X/autosome translo-
 cation in a

[125353]girl with manifestations of Lowe syndrome. Am. J. Med.
Genet. 23:

[125354]837–847, 1986.

[125355]22319.

[125356]Kawano, T.; Indo, Y.; Nakazato, H.; Shimadzu, M.; Matsuda, I.

[125357]: Oculocerebrorenal syndrome of Lowe: three mutations in
the OCRL1

[125358]gene derived from three patients with different phenotypes. Am. J.

[125359]Med. Genet. 77: 348–355, 1998.

[125360]22320.

[125361]Kenworthy, L.; Charnas, L.: Evidence for a discrete behavioral

[125362]phenotype in the oculocerebrorenal syndrome of Lowe.
Am. J. Med.

[125363]Genet. 59: 283–290, 1995.

[125364]22321.

[125365]Kenworthy, L.; Park, T.; Charnas, L. R.: Cognitive and behavioral

[125366]profile of the oculocerebrorenal syndrome of Lowe. Am. J.

Med. Genet. 46:

[125367]297–303, 1993.

[125368]22322.

[125369]Leahey, A.–M.; Charnas, L. R.; Nussbaum, R. L.: Nonsense mutations

[125370]in the OCRL–1 gene in patients with the oculocerebrorenal syndrome

[125371]of Lowe. Hum. Molec. Genet. 2: 461–463, 1993.

[125372]22323.

[125373]Leahey, A. M.; Olivos, I. M.; Bailey, L. C.; Nussbaum, R. L.:

[125374]Characterization of the molecular defects in Lowe's syndrome. (Abstract) Am.

[125375]J. Hum. Genet. 51 (suppl.): A220 only, 1992.

[125376]22324.

[125377]Lin, T.; Lewis, R. A.; Nussbaum, R. L.: Molecular confirmation

[125378]of carriers for Lowe syndrome. Ophthalmology 106: 119–122, 1999.

[125379]22325.

[125380]Lin, T.; Orrison, B. M.; Leahey, A.–M.; Suchy, S. F.; Bernard,

[125381]D. J.; Lewis, R. A.; Nussbaum, R. L.: Spectrum of mutations
in the

[125382]OCRL1 gene in the Lowe oculocerebrorenal syndrome.
Am. J. Hum. Genet. 60:

[125383]1384–1388, 1997.

[125384]22326.

[125385]Kutsche, K.; Yntema, H.; Brandt, A.; Jantke, I.; Nothwang,
H. G.;

[125386]Orth, U.; Boavida, M. G.; David, D.; Chelly, J.; Fryns, J.-P.;
Moraine,

[125387]C.; Ropers, H.-H.; Hamel, B. C. J.; van Bokhoven, H.; Gal,
A.: Mutations

[125388]in ARHGEF6, encoding a guanine nucleotide exchange
factor for Rho

[125389]GTPases, in patients with X-linked mental retardation. Na-
ture Genet. 26:

[125390]247–250, 2000.

[125391]22327.

[125392]Lower, K. M.; Gecz, J.: Characterization of ARHGEF6, a
guanine

[125393]nucleotide exchange factor for Rho GTPases and a candi-
date gene for

[125394]X-linked mental retardation: mutation screening in Borjeson-Forssman-Lehmann

[125395]syndrome and MRX27. Am. J. Med. Genet. 100: 43-48, 2001.

[125396]22328.

[125397]Manser, E.; Loo, T.-H.; Koh, C.-G.; Zhao, Z.-S.; Chen, X.-Q.; Tan,

[125398]L.; Tan, I.; Leung, T.; Lim, L.: PAK kinases are directly coupled

[125399]to the PIX family of nucleotide exchange factors. Molec. Cell 1:

[125400]183-192, 1998.

[125401]22329.

[125402]Nomura, N.; Miyajima, N.; Sazuka, T.; Tanaka, A.; Kawarabayashi,

[125403]Y.; Sato, S.; Nagase, T.; Seki, N.; Ishikawa, K.; Tabata, S.: Prediction

[125404]of the coding sequences of unidentified human genes. I. The coding

[125405]sequences of 40 new genes (KIAA0001-KIAA0040) deduced by analysis

[125406]of randomly sampled cDNA clones from human immature

myeloid cell line,

[125407]KG-1. DNA Res. 1: 27-35, 1994.

[125408]22330.

[125409]von Ballestrem, C. L.; Boavida, M. G.; Zuther, C.; Carreiro, M.

[125410]H.; David, D.; Gal, A.; Schwinger, E.: Jumping translocation in a

[125411]phenotypically normal female. Clin. Genet. 49: 156-159, 1996.

[125412]22331.

[125413]Yntema, H. G.; Hamel, B. C.; Smits, A. P.; van Roosmalen, T.; van

[125414]den Helm, B.; Kremer, H.; Ropers, H. H.; Smeets, D. F.; van Bokhoven,

[125415]H.: Localisation of a gene for non-specific X linked mental retardation

[125416](MRX46) to Xq25-q26. J. Med. Genet. 35: 801-805, 1998.

[125417]22332.

[125418]Grozingier, C. M.; Hassig, C. A.; Schreiber, S. L.: Three proteins

[125419]define a class of human histone deacetylases related to

yeast Hda1p. Proc.

[125420]Nat. Acad. Sci. 96: 4868–4873, 1999.

[125421]22333.

[125422]Hubbert, C.; Guardiola, A.; Shao, R.; Kawaguchi, Y.; Ito, A.;
Nixon,

[125423]A.; Yoshida, M.; Wang, X.-F.; Yao, T.-P.: HDAC6 is a mi-
crotubule-associated

[125424]deacetylase. Nature 417: 455–458, 2002.

[125425]22334.

[125426]Mahlknecht, U.; Schnittger, S.; Landgraf, F.; Schoch, C.;
Ottmann,

[125427]O. G.; Hiddemann, W.; Hoelzer, D.: Assignment of the hu-
man histone

[125428]deacetylase 6 gene (HDAC6) to X chromosome p11.23 by
in situ hybridization. Cytogenet.

[125429]Cell Genet. 93: 135–136, 2001.

[125430]22335.

[125431]Nagase, T.; Isikawa, K.; Suyama, M.; Kikuno, R.; Hirose, A.;
M.;

[125432]Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara,
O.: Prediction

[125433]of the coding sequences of unidentified human genes. XII.

The complete

[125434]sequences of 100 new cDNA clones from brain which code
for large proteins

[125435]in vitro. DNA Res. 5: 355–364, 1998.

[125436]22336.

[125437]Ichida, F.; Tsubata, S.; Bowles, K. R.; Haneda, N.; Uese, K.;

[125438]Miyawaki, T.; Dreyer, W. J.; Messina, J.; Li, H.; Bowles, N.
E.; Towbin,

[125439]J. A.: Novel gene mutations in patients with left ventricular
noncompaction

[125440]or Barth syndrome. Circulation 103: 1256–1263, 2001.

[125441]22337.

[125442]Johnston, J.; Kelley, R. I.; Feigenbaum, A.; Cox, G. F.; Iyer,

[125443]G. S.; Funanage, V. L.; Proujansky, R.: Mutation characteri-
zation

[125444]and genotype–phenotype correlation in Barth syndrome.
Am. J. Hum.

[125445]Genet. 61: 1053–1058, 1997.

[125446]22338.

[125447]Kelley, R. I.; Cheatham, J. P.; Clark, B. J.; Nigro, M. A.;
Powell,
[125448]B. R.; Sherwood, G. W.; Sladky, J. T.; Swisher, W. P.: X-
linked dilated
[125449]cardiomyopathy with neutropenia, growth retardation, and
3-methylglutaconic
[125450]aciduria. J. Pediatr. 119: 738-747, 1991.
[125451]22339.
[125452]Sakamoto, O.; Kitoh, T.; Ohura, T.; Ohya, N.; Iinuma, K.:
Novel
[125453]missense mutation (R94S) in the TAZ (G4.5) gene in a
Japanese patient
[125454]with Barth syndrome. J. Hum. Genet. 47: 229-231, 2002.
[125455]22340.
[125456]Sakamoto, O.; Ohura, T.; Katsushima, Y.; Fujiwara, I.;
Ogawa,
[125457]E.; Miyabayashi, S.; Iinuma, K.: A novel intronic mutation
of the
[125458]TAZ (G4.5) gene in a patient with Barth syndrome: cre-
ation of a 5-prime
[125459]splice donor site with variant GC consensus and elonga-
tion of the

[125460]upstream exon. Hum. Genet. 109: 559–563, 2001.

[125461]22341.

[125462]Strasser, K.; Masuda, S.; Mason, P.; Pfannstiel, J.; Oppizzi, M.;

[125463]Rodriguez–Navarro, S.; Rondon, A. G.; Aguilera, A.; Struhl, K.; Reed,

[125464]R.; Hurt, E.: TREX is a conserved complex coupling transcription

[125465]with messenger RNA export. Nature 417: 304–308, 2002.

[125466]22342.

[125467]Fortenberry, Y.; Hwang, J.–R.; Apletalina, E. V.; Lindberg, I.

[125468]: Functional characterization of proSAAS: similarities and differences

[125469]with 7B2. J. Biol. Chem. 277: 5175–5186, 2002.

[125470]22343.

[125471]Fricker, L. D.; McKinzie, A. A.; Sun, J.; Curran, E.; Qian, Y.;

[125472]Yan, L.; Patterson, S. D.; Courchesne, P. L.; Richards, B.; Levin,

[125473]N.; Mzhavia, N.; Devi, L. A.; Douglass, J.: Identification and characterization

[125474]of proSAAS, a granin-like neuroendocrine peptide precursor that inhibits
[125475]prohormone processing. J. Neurosci. 20: 639-648, 2000.
[125476]22344.
[125477]Omori, Y.; Mesnil, M.; Yamasaki, H.: Connexin 32 mutations from
[125478]X-linked Charcot-Marie-Tooth disease patients: functional defects
[125479]and dominant negative effects. Molec. Biol. Cell 7:
907-916, 1996.
[125480]22345.
[125481]Tabaraud, F.; Lagrange, E.; Sindou, P.; Vandenberghe, A.;
Levy,
[125482]N.; Vallat, J. M.: Demyelinating X-linked Charcot-Marie-Tooth disease:
[125483]unusual electrophysiological findings. Muscle Nerve 22:
1442-1447,
[125484]1999.
[125485]22346.
[125486]Moser, A. B.; Moser, H. W.: The prenatal diagnosis of X-linked

[125487]adrenoleukodystrophy. Prenatal Diag. 19: 46–48, 1999.

[125488]22347.

[125489]Sugarman, G. I.; Rimoin, D. L.; Lachman, R. S.: The facial–digital–genital

[125490](Aarskog) syndrome. Am. J. Dis. Child. 126: 248–252, 1973.

[125491]22348.

[125492]Tsukahara, M.; Fernandez, G. I.: Umbilical findings in Aarskog

[125493]syndrome. Clin. Genet. 45: 260–265, 1994.

[125494]22349.

[125495]Tyrkus, M.; Bawle, E.; Lipman, S.; Bozimowski, D.; Woolley, P.

[125496]V., Jr.: Aarskog–Scott syndrome inherited as an X-linked dominant

[125497]with full male–female expression. (Abstract) Am. J. Hum. Genet. 32:

[125498]134A only, 1980.

[125499]22350.

[125500]van den Bergh, P.; Fryns, J. P.; Wilms, G.; Piot, R.; Dra–lands,

- [125501]G.; van den Bergh, R.: Anomalous cerebral venous drainage in Aarskog
- [125502]syndrome. Clin. Genet. 25: 288–294, 1984.
- [125503]22351.
- [125504]Zheng, Y.; Fischer, D. J.; Santos, M. F.; Tigyi, G.; Pasteris,
- [125505]N. G.; Gorski, J. L.; Xu, Y.: The faciogenital dysplasia gene product
- [125506]FGD1 functions as a Cdc42Hs–specific guanine–nucleotide exchange factor. J.
- [125507]Biol. Chem. 271: 33169–33172, 1996.
- [125508]22352.
- [125509]Inaba, H.; Fujimaki, M.; Kazazian, H. H., Jr.; Antonarakis, S.
- [125510]E.: MspI polymorphic site in intron 22 of the factor VIII gene in
- [125511]the Japanese population. Hum. Genet. 84: 214–215, 1990.
- [125512]22353.
- [125513]Levinson, B.; Bermingham, J. R., Jr.; Metzenberg, A.; Ken-
- wrick,
- [125514]S.; Chapman, V.; Gitschier, J.: Sequence of the human factor VIII–associated

[125515]gene is conserved in mouse. Genomics 13: 862–865,
1992.

[125516]22354.

[125517]Levinson, B.; Kenwrick, S.; Lakich, D.; Hammonds, G.;
Gitschier,

[125518]J.: A transcribed gene in an intron of the human factor VIII
gene. Genomics 7:

[125519]1–11, 1990.

[125520]22355.

[125521]Peters, M. F.; Ross, C. A.: Isolation of a 40–kDa hunt-
ington-associated

[125522]protein. J. Biol. Chem. 276: 3188–3194, 2001.

[125523]22356.

[125524]Berra, B.; Gornati, R.; Rapelli, S.; Gatti, R.; Mancini, G. M.

[125525]S.; Ciana, G.; Bembi, B.: Infantile sialic acid storage dis-
ease:

[125526]biochemical studies. Am. J. Med. Genet. 58: 24–31, 1995.

[125527]22357.

[125528]Cameron, P. D.; Dubowitz, V.; Besley, G. T. N.; Fensom, A.
H.:

[125529]Sialic acid storage disease. Arch. Dis. Child. 65: 314–315, 1990.

[125530]22358.

[125531]Haataja, L.; Schleutker, J.; Laine, A.–P.; Renlund, M.; Savontaus,

[125532]M.–L.; Dib, C.; Weissenbach, J.; Peltonen, L.; Aula, P.: The genetic

[125533]locus for free sialic acid storage disease maps to the long arm of

[125534]chromosome 6. Am. J. Hum. Genet. 54: 1042–1049, 1994.

[125535]22359.

[125536]Mules, E. H.; Hayflick, S.; Miller, C. S.; Reynolds, L. W.; Thomas,

[125537]G. H.: Six novel deleterious and three neutral mutations in the gene

[125538]encoding the alpha–subunit of hexosaminidase A in non–Jewish individuals. Am.

[125539]J. Hum. Genet. 50: 834–841, 1992.

[125540]22360.

[125541]Myerowitz, R.: Tay–Sachs disease–causing mutations and neutral

[125542]polymorphisms in the Hex A gene. Hum. Mutat. 9:
195–208, 1997.

[125543]22361.

[125544]Myerowitz, R.: Splice junction mutation in some Ashkenazi
Jews

[125545]with Tay–Sachs disease: evidence against a single defect
within this

[125546]ethnic group. Proc. Nat. Acad. Sci. 85: 3955–3959, 1988.

[125547]22362.

[125548]Myerowitz, R.; Costigan, F. C.: The major defect in Ashke-
nazi

[125549]Jews with Tay–Sachs disease is an insertion in the gene for
the alpha–chain

[125550]of beta–hexosaminidase. J. Biol. Chem. 263:
18587–18589, 1988.

[125551]22363.

[125552]Myerowitz, R.; Hogikyan, N. D.: Different mutations in
Ashkenazi

[125553]Jewish and non–Jewish French Canadians with Tay–Sachs
disease. Science 232:

[125554]1646–1648, 1986.

[125555]22364.

[125556]Myerowitz, R.; Hogikyan, N. D.: A deletion involving Alu sequences

[125557]in the beta-hexosaminidase alpha-chain gene of French Canadians with

[125558]Tay-Sachs disease. J. Biol. Chem. 262: 15396-15399, 1987.

[125559]22365.

[125560]Myerowitz, R.; Piekarz, R.; Neufeld, E. F.; Shows, T. B.; Suzuki,

[125561]K.: Human beta-hexosaminidase alpha chain: coding sequence and homology

[125562]with the beta chain. Proc. Nat. Acad. Sci. 82: 7830-7834, 1985.

[125563]22366.

[125564]Myerowitz, R.; Proia, R. L.: cDNA clone for the alpha-chain of

[125565]human beta-hexosaminidase: deficiency of alpha-chain mRNA in Ashkenazi

[125566]Tay-Sachs fibroblasts. Proc. Nat. Acad. Sci. 81: 5394-5398, 1984.

[125567]22367.

[125568]Nakai, H.; Byers, M. G.; Nowak, N. J.; Shows, T. B.: Assignment

[125569]of beta-hexosaminidase A alpha-subunit to human chromosomal region

[125570]15q23-q24. Cytogenet. Cell Genet. 56: 164, 1991.

[125571]22368.

[125572]Nakano, T.; Muscillo, M.; Ohno, K.; Hoffman, A. J.; Suzuki, K.

[125573]: A point mutation in the coding sequence of the beta-hexosaminidase

[125574]alpha gene results in defective processing of the enzyme protein in

[125575]an unusual GM2-gangliosidosis variant. J. Neurochem. 51: 984-987,

[125576]1988.

[125577]22369.

[125578]Nakano, T.; Nanba, E.; Tanaka, A.; Ohno, K.; Suzuki, Y.; Suzuki,

[125579]K.: A new point mutation within exon 5 of beta-hexosaminidase alpha

[125580]gene in a Japanese infant with Tay-Sachs disease. Ann. Neurol. 27:

[125581]465–473, 1990.

[125582]22370.

[125583]Navon, R.; Kolodny, E. H.; Mitsumoto, H.; Thomas, G. H.;
Proia,

[125584]R. L.: Ashkenazi–Jewish and non–Jewish adult G(M2) gan-
gliosidosis

[125585]patients share a common genetic defect. Am. J. Hum.
Genet. 46: 817–821,

[125586]1990.

[125587]22371.

[125588]Navon, R.; Proia, R. L.: The mutations in Ashkenazi Jews
with

[125589]adult G(M2) gangliosidosis, the adult form of Tay–Sachs
disease. Science 243:

[125590]1471–1474, 1989.

[125591]22372.

[125592]Navon, R.; Proia, R. L.: Tay–Sachs disease in Moroccan
Jews:

[125593]deletion of a phenylalanine in the alpha–subunit of beta–
hexosaminidase. Am.

[125594]J. Hum. Genet. 48: 412–419, 1991.

[125595]22373.

[125596]Nishimoto, J.; Tanaka, A.; Nanba, E.; Suzuki, K.: Expression

[125597]of the beta-hexosaminidase alpha subunit gene with the four-base insertion

[125598]of infantile Jewish Tay-Sachs disease. J. Biol. Chem. 266: 14306-14309,

[125599]1991.

[125600]22374.

[125601]Koch, M. C.; Ricker, K.; Otto, M.; Grimm, T.; Hoffman, E. P.;

[125602]Rudel, R.; Bender, K.; Zoll, B.; Harper, P. S.; Lehmann-Horn, F.:

[125603]Confirmation of linkage of hyperkalaemic periodic paralysis to chromosome

[125604]17. J. Med. Genet. 28: 583-586, 1991.

[125605]22375.

[125606]Lehmann-Horn, F.; Rudel, R.; Ricker, K.: Non-dystrophic myotonias

[125607]and periodic paralyses. Neuromusc. Disord. 3: 161-168, 1993.

[125608]22376.

[125609]McClatchey, A. I.: Personal Communication. Boston, Mass.
3/16/1993.

[125610]22377.

[125611]McClatchey, A. I.; McKenna-Yasek, D.; Cros, D.; Worthen,
H. G.;

[125612]Kuncl, R. W.; DeSilva, S. M.; Cornblath, D. R.; Gusella, J. F.;
Brown,

[125613]R. H., Jr.: Novel mutations in families with unusual and
variable

[125614]disorders of the skeletal muscle sodium channel. Nature
Genet. 2:

[125615]148–152, 1992.

[125616]22378.

[125617]McClatchey, A. I.; Trofatter, J.; McKenna-Yasek, D.;
Raskind,

[125618]W.; Bird, T.; Pericak-Vance, M.; Gilchrist, J.; Arahata, K.;
Radosavljevic,

[125619]D.; Worthen, H. G.; Van den Bergh, P.; Haines, J. L.;
Gusella, J.

[125620]F.; Brown, R. H., Jr.: Dinucleotide repeat polymorphisms at
the SCN4A

[125621]locus suggest allelic heterogeneity of hyperkalemic periodic paralysis

[125622]and paramyotonia congenita. Am. J. Hum. Genet. 50: 896–901, 1992.

[125623]22379.

[125624]McClatchey, A. I.; Van den Bergh, P.; Pericak–Vance, M. A.; Raskind,

[125625]W.; Verellen, C.; McKenna–Yasek, D.; Rao, K.; Haines, J. L.; Bird,

[125626]T.; Brown, R. H., Jr.; Gusella, J. F.: Temperature–sensitive mutations

[125627]in the III–IV cytoplasmic loop region of the skeletal muscle sodium

[125628]channel gene in paramyotonia congenita. Cell 68: 769–774, 1992.

[125629]22380.

[125630]Moslehi, R.; Langlois, S.; Yam, I.; Friedman, J. M.: Linkage

[125631]of malignant hyperthermia and hyperkalemic periodic paralysis to the

[125632]adult skeletal muscle sodium channel (SCN4A) gene in a large pedigree. Am.

[125633]J. Med. Genet. 76: 21–27, 1998.

[125634]22381.

[125635]Ptacek, L. J.; George, A. L., Jr.; Barchi, R. L.; Griggs, R. C.;

[125636]Riggs, J. E.; Robertson, M.; Leppert, M. F.: Mutations in an
S4 segment

[125637]of the adult skeletal muscle sodium channel cause
paramyotonia congenita. Neuron 8:

[125638]891–897, 1992.

[125639]22382.

[125640]Ptacek, L. J.; George, A. L., Jr.; Griggs, R. C.; Tawil, R.;

Kallen,

[125641]R. G.; Barchi, R. L.; Robertson, M.; Leppert, M. F.: Identifi-
cation

[125642]of a mutation in the gene causing hyperkalemic periodic
paralysis. Cell 67:

[125643]1021–1027, 1991.

[125644]22383.

[125645]Ptacek, L. J.; Gouw, L.; Kwieninski, H.; McManis, P.;

Mendell,

[125646]J. R.; Barohn, A. L., Jr.; Robertson, M.; Leppert, M. F.:
Sodium

[125647]channel mutations in paramyotonia congenita and hyper-

kalemic periodic

[125648]paralysis. Ann. Neurol. 33: 300–307, 1993.

[125649]22384.

[125650]Ptacek, L. J.; Tawil, R.; Griggs, R. C.; Meola, G.; McManis, P.;

[125651]Barohn, R. J.; Mendell, J. R.; Harris, C.; Spitzer, R.; Santiago,

[125652]F.; Leppert, M. F.: Sodium channel mutations in acetazolamide-responsive

[125653]myotonia congenita, paramyotonia congenita, and hyperkalemic periodic

[125654]paralysis. Neurology 44: 1500–1503, 1994.

[125655]22385.

[125656]Ptacek, L. J.; Tawil, R.; Griggs, R. C.; Storvick, D.; Leppert,

[125657]M.: Linkage of atypical myotonia congenita to a sodium channel locus. Neurology 42:

[125658]431–433, 1992.

[125659]22386.

[125660]Ptacek, L. J.; Tyler, F.; Trimmer, J. S.; Agnew, W. S.; Leppert,

[125661]M.: Analysis in a large hyperkalemic periodic paralysis

pedigree

[125662]supports tight linkage to a sodium channel locus. Am. J. Hum. Genet. 49:

[125663]378–382, 1991.

[125664]22387.

[125665]Ricker, K.; Moxley, R. T., III; Heine, R.; Lehmann–Horn, F.:

[125666]Myotonia fluctuans: a third type of muscle sodium channel disease. Arch.

[125667]Neurol. 51: 1095–1102, 1994.

[125668]22388.

[125669]Rudolph, J. A.; Spier, S. J.; Byrns, G.; Hoffman, E. P.: Linkage

[125670]of hyperkalaemic periodic paralysis in Quarter horses to the horse

[125671]adult skeletal muscle sodium channel gene. Anim. Genet. 23: 241–250,

[125672]1992.

[125673]22389.

[125674]Rudolph, J. A.; Spier, S. J.; Byrns, G.; Rojas, C. V.; Bernoco,

[125675]D.; Hoffman, E. P.: Periodic paralysis in Quarter horses: a

sodium

[125676]channel mutation disseminated by selective breeding. *Nature Genet.* 2:

[125677]144–147, 1992.

[125678]22390.

[125679]Trudell, R. G.; Kaiser, K. K.; Griggs, R. C.: Acetazolamide responsive

[125680]myotonia congenita. *Neurology* 37: 488–491, 1987.

[125681]22391.

[125682]Tyler, F. H.; Stephens, F. E.; Gunn, F. D.; Perkoff, G. T.: Studies

[125683]in disorders of muscle. VII. Clinical manifestations and inheritance

[125684]of a type of periodic paralysis without hypopotassemia. *J. Clin.*

[125685]Invest. 30: 492–502, 1951.

[125686]22392.

[125687]European Polycystic Kidney Disease Consortium: The polycystic

[125688]kidney disease 1 gene encodes a 14 kb transcript and lies within a

[125689] duplicated region on chromosome 16. Cell 77: 881–894, 1994.

[125690] 22393.

[125691] Gabow, P. A.: Autosomal dominant polycystic kidney disease. New

[125692] Eng. J. Med. 329: 332–342, 1993.

[125693] 22394.

[125694] Gal, A.; Wirth, B.; Kaariainen, H.; Lucotte, G.; Landais, P.;

[125695] Gillessen–Kaesbach, G.; Muller–Wiefel, D. E.; Zerres, K.: Childhood

[125696] manifestation of autosomal dominant polycystic kidney disease: no

[125697] evidence for genetic heterogeneity. Clin. Genet. 35: 13–19, 1989.

[125698] 22395.

[125699] Germino, G. G.; Barton, N. J.; Lamb, J.; Higgs, D. R.; Harris,

[125700] P.; Xiao, G. H.; Scherer, G.; Nakamura, Y.; Reeders, S. T.: Identification

[125701] of a locus which shows no genetic recombination with the autosomal

[125702] dominant polycystic kidney disease gene on chromosome

16. Am. J.

[125703]Hum. Genet. 46: 925–933, 1990.

[125704]22396.

[125705]Germino, G. G.; Weinstat–Saslow, D.; Himmelbauer, H.;
Gillespie,

[125706]G. A. J.; Somlo, S.; Wirth, B.; Barton, N.; Harris, K. L.;
Frischauf,

[125707]A.–M.; Reeders, S. T.: The gene for autosomal dominant
polycystic

[125708]kidney disease lies in a 750–kb CpG-rich region. Ge-
nomics 13: 144–151,

[125709]1992.

[125710]22397.

[125711]Gillespie, G. A. J.; Somlo, S.; Germino, G. G.; Weinstat–
Saslow,

[125712]D.; Reeders, S. T.: CpG island in the region of an autoso-
mal dominant

[125713]polycystic kidney disease locus defines the 5–prime end
of a gene

[125714]encoding a putative proton channel. Proc. Nat. Acad. Sci.
88: 4289–4293,

[125715]1991.

[125716]22398.

[125717]Harris, P. C.; Barton, N. J.; Higgs, D. R.; Reeders, S. T.;
Wilkie,

[125718]A. O. M.: A long-range restriction map between the alpha-
globin complex

[125719]and a marker closely linked to the polycystic kidney dis-
ease 1 (PKD1)

[125720]locus. Genomics 7: 195-206, 1990.

[125721]22399.

[125722]Harris, P. C.; Thomas, S.; Ratcliffe, P. J.; Breuning, M. H.;

[125723]Coto, E.; Lopez-Larrea, C.: Rapid genetic analysis of fami-
lies with

[125724]polycystic kidney disease 1 by means of a microsatellite
marker. Lancet 338:

[125725]1484-1487, 1991.

[125726]22400.

[125727]Himmelbauer, H.; Pohlschmidt, M.; Snarey, A.; Germino, G.
G.;

[125728]Weinstat-Saslow, D.; Somlo, S.; Reeders, S. T.; Frischauf,
A.-M.:

[125729]Human-mouse homologies in the region of the polycystic

kidney disease

[125730]gene (PKD1). Genomics 13: 35–38, 1992.

[125731]22401.

[125732]Himmelbauer, H.; Pohlschmidt, M.; Snarey, A.; Somlo, S.;
Weinstat-Saslow,

[125733]D.; Germino, G.; Reeders, S.; Frischauf, A.–M.: A syntenic
region

[125734]between human 16p13.3 and mouse chromosome 17.
(Abstract) Cytogenet.

[125735]Cell Genet. 58: 2136, 1991.

[125736]22402.

[125737]Hughes, J.; Ward, C. J.; Peral, B.; Aspinwall, R.; Clark, K.;

[125738]San Millan, J. L.; Gamble, V.; Harris, P. C.: The polycystic
kidney

[125739]disease 1 (PKD1) gene encodes a novel protein with multi-
ple cell recognition

[125740]domains. Nature Genet. 10: 151–160, 1995.

[125741]22403.

[125742]International Polycystic Kidney Disease Consortium: Poly-
cystic

[125743]kidney disease: the complete structure of the PKD1 gene

and its protein. Cell 81:

[125744]289–298, 1995.

[125745]22404.

[125746]Keith, T.; Reeders, S. T.; Brown, V. A.; Green, P.; Phipps, P.;

[125747]Bricker, A.; Barton, N. J.; Collier, P. S.; Knowlton, R.; Nelson,
son,

[125748]C.; Helms, C.; Donis-Keller, H.: Linkage studies of autosomal dominant

[125749]polycystic kidney disease (PKD1) and a genetic map of chromosome 16

[125750]based on 40 RFLP markers. (Abstract) Am. J. Hum. Genet.
41: A172,

[125751]1987.

[125752]22405.

[125753]Fujii, J.; Zarain-Herzberg, A.; Willard, H. F.; Tada, M.; MacLennan,

[125754]D. H.: Structure of the rabbit phospholamban gene, cloning of the

[125755]human cDNA, and assignment of the gene to human chromosome 6. J.

[125756]Biol. Chem. 266: 11669–11675, 1991.

[125757]22406.

[125758]McTiernan, C. F.; Frye, C. S.; Lemster, B. H.; Kinder, E. A.;
Ogletree-Hughes,

[125759]M. L.; Moravec, C. S.; Feldman, A. M.: The human phos-
pholamban gene:

[125760]structure and expression. J. Molec. Cell Cardiol. 31:
679-692, 1999.

[125761]22407.

[125762]Otsu, K.; Fujii, J.; Periasamy, M.; Difilippantonio, M.; Up-
pender,

[125763]M.; Ward, D. C.; MacLennan, D. H.: Chromosome mapping
of five human

[125764]cardiac and skeletal muscle sarcoplasmic reticulum pro-
tein genes. Genomics 17:

[125765]507-509, 1993.

[125766]22408.

[125767]Barhanin, J.; Lesage, F.; Guillemare, E.; Fink, M.; Lazdun-
ski,

[125768]M.; Romey, G.: K(v)LQT1 and Isk (mink) proteins associate
to form

[125769]the I(Ks) cardiac potassium current. Nature 384: 78-80,
1996.

[125770]22409.

[125771]Marx, S. O.; Kurokawa, J.; Reiken, S.; Motoike, H.;

D'Armiento,

[125772]J.; Marks, A. R.; Kass, R. S.: Requirement of a macromolecular signaling

[125773]complex for beta adrenergic receptor modulation of the KCNQ1-KCNE1

[125774]potassium channel. Science 295: 496-499, 2002.

[125775]22410.

[125776]Sanguinetti, M. C.; Curran, M. E.; Zou, A.; Shen, J.; Spector,

[125777]P. S.; Atkinson, D. L.; Keating, M. T.: Coassembly of K(v)LQT1 and

[125778]minK (IsK) proteins to form cardiac I(Ks) potassium channel. Nature 384:

[125779]80-83, 1996.

[125780]22411.

[125781]Coleman, R. A.; Smith, W. L.; Narumiya, S.: VIII. International

[125782]union of pharmacology classification of prostanoid receptors: properties,

[125783]distribution, and structure of the receptors and their sub-

types. Pharm.

[125784]Rev. 46: 205–229, 1994.

[125785]22412.

[125786]Duncan, A. M. V.; Anderson, L. L.; Funk, C. D.;
Abramovitz, M.;

[125787]Adam, M.: Chromosomal localization of the human
prostanoid receptor

[125788]gene family. Genomics 25: 740–742, 1995.

[125789]22413.

[125790]White, D. M.; Mikol, D. D.; Espinosa, R.; Weimer, B.; Le
Beau,

[125791]M. M.; Stefansson, K.: Structure and chromosomal local-
ization of

[125792]the human gene for a brain form of prostaglandin D-2
synthase. J.

[125793]Biol. Chem. 267: 23202–23208, 1992.

[125794]22414.

[125795]Sales, K. J.; Katz, A. A.; Davis, M.; Hinz, S.; Soeters, R. P.;

[125796]Hofmeyr, M. D.; Millar, R. P.; Jabbour, H. N.: Cyclooxyge-
nase-2 expression

[125797]and prostaglandin E2 synthesis are up-regulated in carci-

nomas of the

[125798]cervix: a possible autocrine/paracrine regulation of neoplastic cell

[125799]function via EP2/EP4 receptors. J. Clin. Endocr. Metab. 86: 2243–2249,

[125800]2001.

[125801]22415.

[125802]Taketo, M.; Rochelle, J. M.; Sugimoto, Y.; Namba, T.; Honda, A.;

[125803]Negishi, M.; Ichikawa, A.; Narumiya, S.; Seldin, M. F.: Mapping of

[125804]the genes encoding mouse thromboxane A2 receptor and prostaglandin

[125805]E receptor subtypes EP2 and EP3. Genomics 19: 585–588, 1994.

[125806]22416.

[125807]Catella–Lawson, F.; Reilly, M. P.; Kapoor, S. C.; Cucchiara, A.

[125808]J.; DeMarco, S.; Tournier, B.; Vyas, S. N.; FitzGerald, G. A.: Cyclooxygenase

[125809]inhibitors and the antiplatelet effects of aspirin. New Eng. J. Med. 345:

[125810]1809–1817, 2001.

[125811]22417.

[125812]Dube, J.-N.; Drouin, J.; Aminian, M.; Plant, M. H.; La-neuville,

[125813]O.: Characterization of a partial prostaglandin endoperox-
ide H synthase-1

[125814]deficiency in a patient with a bleeding disorder. Brit. J.
Haemat. 113:

[125815]878–885, 2001.

[125816]22418.

[125817]Funk, C. D.; Funk, L. B.; Kennedy, M. E.; Pong, A. S.;
Fitzgerald,

[125818]G. A.: Human platelet/erythroleukemia cell prostaglandin
G/H synthase:

[125819]cDNA cloning, expression, and gene chromosomal as-
signment. FASEB

[125820]J. 5: 2304–2312, 1991.

[125821]22419.

[125822]Gavett, S. H.; Madison, S. L.; Chulada, P. C.; Scarborough,
P.

[125823]E.; Qu, W.; Boyle, J. E.; Tiano, H. F.; Lee, C. A.; Langenbach,

R.;

[125824]Roggli, V. L.; Zeldin, D. C.: Allergic lung responses are increased

[125825]in prostaglandin H synthase-deficient mice. J. Clin. Invest. 104:

[125826]721-732, 1999.

[125827]22420.

[125828]Kirschenbaum, A.; Liotta, D. R.; Yao, S.; Liu, X.-H.; Klausner,

[125829]A. P.; Unger, P.; Shapiro, E.; Leav, I.; Levine, A. C.: Immunohistochemical

[125830]localization of cyclooxygenase-1 and cyclooxygenase-2 in the human

[125831]fetal and adult male reproductive tracts. J. Clin. Endocr. Metab. 85:

[125832]3436-3441, 2000.

[125833]22421.

[125834]Langenbach, R.; Morham, S. G.; Tiano, H. F.; Loftin, C. D.; Ghanayem,

[125835]B. I.; Chulada, P. C.; Mahler, J. F.; Lee, C. A.; Goulding, E. H.;

[125836]Kluckman, K. D.; Kim, H. S.; Smithies, O.: Prostaglandin

synthase

[125837]1 gene disruption in mice reduces arachidonic acid–
induced inflammation

[125838]and indomethacin–induced gastric ulceration. Cell 83:
483–492, 1995.

[125839]22422.

[125840]Malkowski, M. G.; Ginell, S. L.; Smith, W. L.; Garavito, R. M.

[125841]: The productive conformation of arachidonic acid bound
to prostaglandin

[125842]synthase. Science 289: 1933–1937, 2000.

[125843]22423.

[125844]Pareti, F. I.; Manucci, P. M.; D'Angelo, A.; Smith, J. B.;
Sautebin,

[125845]L.; Galli, G.: Congenital deficiency of thromboxane and
prostacyclin. Lancet I:

[125846]898–900, 1980.

[125847]22424.

[125848]Picot, D.; Loll, P. J.; Garavito, R. M.: The x–ray crystal
structure

[125849]of the membrane protein prostaglandin H–2 synthase–1.
Nature 367:

[125850]243–249, 1994.

[125851]22425.

[125852]Pieke, S. A.; Kimberling, W. J.; Kenyon, J. B.; Gabow, P.:

Genetic

[125853]heterogeneity of polycystic kidney disease: an estimate of
the proportion

[125854]of families unlinked to chromosome 16. (Abstract) Am. J.
Hum. Genet. 45

[125855](suppl.): A58 only, 1989.

[125856]22426.

[125857]Qian, F.; Germino, F. J.; Cai, Y.; Zhang, X.; Somlo, S.; Ger-
mino,

[125858]G. G.: PKD1 interacts with PKD2 through a probable
coiled–coil domain. Nature

[125859]Genet. 16: 179–183, 1997.

[125860]22427.

[125861]Ravine, D.; Walker, R. G.; Gibson, R. N.; Forrest, S. M.;
Richards,

[125862]R. I.; Friend, K.; Sheffield, L. J.; Kincaid–Smith, P.; Danks,
D.

[125863]M.: Phenotype and genotype heterogeneity in autosomal

dominant polycystic

[125864]kidney disease. Lancet 340: 1330–1333, 1992.

[125865]22428.

[125866]Reynolds, D. M.; Hayashi, T.; Cai, Y.; Veldhuisen, B.; Watnick,

[125867]T. J.; Lens, X. M.; Mochizuki, T.; Qian, F.; Maeda, Y.; Li, L.; Fossdal,

[125868]R.; Coto, E.; Wu, G.; Breuning, M. H.; Germino, G. G.; Peters, D.

[125869]J. M.; Somlo, S.: Aberrant splicing in the PKD2 gene as a cause of

[125870]polycystic kidney disease. J. Am. Soc. Nephrol. 10: 2342–2351, 1999.

[125871]22429.

[125872]Romeo, G.; Devoto, M.; Costa, G.; Roncuzzi, L.; Catizone, L.;

[125873]Zucchelli, P.; Germino, G.-G.; Keith, T.; Weatherall, D. J.; Reeders,

[125874]S. T.: A second genetic locus for autosomal dominant polycystic kidney

[125875]disease. Lancet II: 8–11, 1988.

[125876]22430.

[125877]San Millan, J. L.; Viribay, M.; Peral, B.; Martinez, I.; Weissenbach,
[125878]J.; Moreno, F.: Refining the localization of the PKD2 locus
on chromosome
[125879]4q by linkage analysis in Spanish families with autosomal
dominant
[125880]polycystic kidney disease type 2. Am. J. Hum. Genet. 56:
248-253,
[125881]1995.
[125882]22431.
[125883]Scheffers, M. S.; Le, H.; van der Bent, P.; Leonhard, W.;
Prins,
[125884]F.; Spruit, L.; Breuning, M. H.; de Heer, E.; Peters, D. J. M.:
Distinct
[125885]subcellular expression of endogenous polycystin-2 in the
plasma membrane
[125886]and Golgi apparatus of MDCK cells. Hum. Molec. Genet.
11: 59-67,
[125887]2002.
[125888]22432.
[125889]Schneider, M. C.; Rodriguez, A. M.; Nomura, H.; Zhou, J.;
Morton,

- [125890]C. C.; Reeders, S. T.; Weremowicz, S.: A gene similar to PKD1 maps to chromosome 4q22: a candidate gene for PKD2. *Genomics* 38: 1-4, 1996.
- [125891]22433.
- [125892]Torra, R.; Badenas, C.; San Millan, J. L.; Perez-Oller, L.; Estivill, X.; Darnell, A.: A loss-of-function model for cystogenesis in human autosomal dominant polycystic kidney disease type 2. *Am. J. Hum. Genet.* 65: 345-352, 1999.
- [125893]22434.
- [125894]Tsiokas, L.; Kim, E.; Arnould, T.; Sukhatme, V. P.; Walz, G.: Homo- and heterodimeric interactions between the gene products of PKD1 and PKD2. *Proc. Nat. Acad. Sci.* 94: 6965-6970, 1997.
- [125895]22435.
- [125896]Veldhuisen, B.; Saris, J. J.; de Haij, S.; Hayashi, T.;

Reynolds,

[125904]D. M.; Mochizuki, T.; Elles, R.; Fossdal, R.; Bogdanova, N.;
van Dijk,

[125905]M. A.; Coto, E.; Ravine, D.; Norby, S.; Verellen-Dumoulin,
C.; Breuning,

[125906]M. H.; Somlo, S.; Peters, D. J. M.: A spectrum of mutations
in the

[125907]second gene for autosomal dominant polycystic kidney
disease (PKD2). Am.

[125908]J. Hum. Genet. 61: 547-555, 1997.

[125909]22436.

[125910]Viribay, M.; Hayashi, T.; Telleria, D.; Mochizuki, T.;
Reynolds,

[125911]D. M.; Alonso, R.; Lens, X. M.; Moreno, F.; Harris, P. C.;
Somlo,

[125912]S.; San Millan, J. L.: Novel stop and frameshifting muta-
tions in

[125913]the autosomal dominant polycystic kidney disease 2
(PKD2) gene. Hum.

[125914]Genet. 101: 229-234, 1997.

[125915]22437.

[125916]Watnick, T.; He, N.; Wang, K.; Liang, Y.; Parfrey, P.; Heffer-

ton,

[125917]D.; St. George-Hyslop, P.; Germino, G.; Pei, Y.: Mutations
of PKD1

[125918]in ADPKD2 cysts suggest a pathogenic effect of trans-
heterozygous

[125919]mutations. Nature Genet. 25: 143–144, 2000.

[125920]22438.

[125921]Wu, G.; D'Agati, V.; Cai, Y.; Markowitz, G.; Park, J. H.;
Reynolds,

[125922]D. M.; Maeda, Y.; Le, T. C.; Hou, H., Jr.; Kucherlapati, R.;
Edelmann,

[125923]W.; Somlo, S.: Somatic inactivation of Pkd2 results in poly-
cystic

[125924]kidney disease. Cell 98: 177–188, 1998.

[125925]22439.

[125926]Wu, G.; Markowitz, G. S.; Li, L.; D'Agati, V. D.; Factor, S.
M.;

[125927]Geng, L.; Tibara, S.; Tuchman, J.; Cai, Y.; Park, J. H.; van
Adelsberg,

[125928]J.; Hou, H., Jr.; Kucherlapati, R.; Edelmann, W.; Somlo, S.:
Cardiac

[125929]defects and renal failure in mice with targeted mutations

in Pkd2. Nature

[125930]Genet. 24: 75–78, 2000.

[125931]22440.

[125932]Wu, G.; Mochizuki, T.; Le, T. C.; Cai, Y.; Hayashi, T.;
Reynolds,

[125933]D. M.; Somlo, S.: Molecular cloning, cDNA sequence analysis, and

[125934]chromosomal localization of mouse Pkd2. Genomics 45:
220–223, 1997.

[125935]22441.

[125936]Xenophontos, S.; Constantinides, R.; Hayashi, T.;
Mochizuki, T.;

[125937]Somlo, S.; Pierides, A.; Constantinou Deltas, C.: A transition frameshift

[125938]mutation induced by a cytosine insertion in the polycystic
kidney

[125939]disease 2 gene (PKD2). Hum. Molec. Genet. 6: 949–952,
1997.

[125940]22442.

[125941]Zerres, K.; Mucher, G.; Rudnik–Schoneborn, S.: Autosomal
recessive

[125942]polycystic kidney disease does not map to the second

gene locus for

[125943]autosomal dominant polycystic kidney disease on chromosome 4. Hum.

[125944]Genet. 93: 697–698, 1994.

[125945]22443.

[125946]Zerres, K.; Volpel, M. C.; Weiss, H.: Cystic kidneys: genetics,

[125947]pathologic anatomy, clinical picture, and prenatal diagnosis. Hum.

[125948]Genet. 68: 104–135, 1984.

[125949]22444.

[125950]Ala-Mello, S.; Koskimies, O.; Rapola, J.; Kaariainen, H.:
Nephronophthisis

[125951]in Finland: epidemiology and comparison of genetically
classified

[125952]subgroups. Europ. J. Hum. Genet. 7: 205–211, 1999.

[125953]22445.

[125954]Chamberlin, B. C.; Hagge, W. W.; Stickler, G. B.: Juvenile
nephronophthisis

[125955]and medullary cystic disease. Mayo Clin. Proc. 52:
485–491, 1977.

[125956]22446.

[125957]Hussels-Maumenee, I.; Pierce, E. R.; Bias, W. B.; Schleuter-mann,

[125958]D. A.: Linkage studies of typical retinitis pigmentosa and common

[125959]markers. Am. J. Hum. Genet. 27: 505-508, 1975.

[125960]22447.

[125961]Olsson, J. E.; Samanns, C.; Jimenez, J.; Pongratz, J.; Chand,

[125962]A.; Watty, A.; Seuchter, S. A.; Denton, M.; Gal, A.: Gene of type

[125963]II autosomal dominant retinitis pigmentosa maps on the long arm of

[125964]chromosome 3. Am. J. Med. Genet. 35: 595-599, 1990.

[125965]22448.

[125966]Pierce, E. A.; Quinn, T.; Meehan, T.; McGee, T. L.; Berson, E.

[125967]L.; Dryja, T. P.: Mutations in a gene encoding a new oxygen-regulated

[125968]photoreceptor protein cause dominant retinitis pigmentosa. Nature

[125969]Genet. 22: 248-254, 1999.

[125970]22449.

[125971]Sullivan, L. S.; Heckenlively, J. R.; Bowne, S. J.; Zuo, J.;
Hide,

[125972]W. A.; Gal, A.; Denton, M.; Inglehearn, C. F.; Blanton, S. H.;
Daiger,

[125973]S. P.: Mutations in a novel retina-specific gene cause au-
tosomal

[125974]dominant retinitis pigmentosa. Nature Genet. 22:
255-259, 1999.

[125975]22450.

[125976]Inglehearn, C.; Farrar, J.; Denton, M.; Gal, A.; Humphries,
P.;

[125977]Bhattacharya, S.: Evidence against a second autosomal
dominant retinitis

[125978]pigmentosa locus close to rhodopsin on chromosome 3q.
(Letter) Am.

[125979]J. Hum. Genet. 53: 536-537, 1993.

[125980]22451.

[125981]McInnes, R. R.; Bascom, R. A.: Retinal genetics: a nullifying

[125982]effect for rhodopsin. Nature Genet. 1: 155-157, 1992.

[125983]22452.

- [125984]McWilliam, P.; Farrar, G. J.; Kenna, P.; Bradley, D. G.;
Humphries,
[125985]M. M.; Sharp, E. M.; McConnell, D. J.; Lawler, M.; Sheils, D.;
Ryan,
[125986]C.; Stevens, K.; Daiger, S. P.; Humphries, P.: Autosomal
dominant
[125987]retinitis pigmentosa (ADRP): localization of an ADRP gene
to the long
[125988]arm of chromosome 3. Genomics 5: 619–622, 1989.
[125989]22453.
[125990]Chautard–Freire–Maia, E. A.: Probable assignment of the
serum
[125991]cholinesterase (E1) and transferrin (Tf) loci to chromo-
some 1 in man. Hum.
[125992]Hered. 27: 134–142, 1977.
[125993]22454.
[125994]Chautard–Freire–Maia, E. A.: Probable assignment of the
E1 and
[125995]Tf loci to chromosome 1 in man. Ciencia e Cultura (Brazil)
28 (suppl.):
[125996]309–310, 1976.

[125997]22455.

[125998]Niswender, K. D.; Morton, G. J.; Stearns, W. H.; Rhodes, C. J.;

[125999]Myers, M. G., Jr.; Schwartz, M. W.: Key enzyme in leptin-induced

[126000]anorexia. Nature 413: 794-795, 2001.

[126001]22456.

[126002]Ogawa, Y.; Masuzaki, H.; Isse, N.; Okazaki, T.; Mori, K.; Shigemoto,

[126003]M.; Satoh, N.; Tamura, N.; Hosoda, K.; Yoshimasa, Y.; Jingami, H.;

[126004]Kawada, T.; Nakao, K.: Molecular cloning of rat obese cDNA and augmented

[126005]gene expression in genetically obese Zucker fatty (fa/fa) rats. J.

[126006]Clin. Invest. 96: 1647-1652, 1995.

[126007]22457.

[126008]Pasco, J. A.; Henry, M. J.; Kotowicz, M. A.; Collier, G. R.; Ball,

[126009]M. J.; Ugoni, A. M.; Nicholson, G. C.: Serum leptin levels are associated

- [126010] with bone mass in nonobese women. J. Clin. Endocr. Metab. 86: 1884–1887, 2001.
- [126011] 22458.
- [126012] Pellemounter, M. A.; Cullen, M. J.; Baker, M. B.; Hecht, R.; Winters, D.; Boone, T.; Collins, F.: Effects of the obese gene product on body weight regulation in ob/ob mice. Science 269: 540–542, 1995.
- [126013] 22459.
- [126014] Rink, T. J.: In search of a satiety factor. Nature 372: 408–409, 1994.
- [126015] 22460.
- [126016] Rosenbaum, M.; Leibel, R. L.: The role of leptin in human physiology. (Editorial) New Eng. J. Med. 341: 913–915, 1999.
- [126017] 22461.
- [126018] Saad, M. F.; Damani, S.; Gingerich, R. L.; Riad-Gabriel, M. G.;

[126024]Khan, A.; Boyadjian, R.; Jinagouda, S. D.; El-Tawil, K.;
Rude, R.

[126025]K.; Kamdar, V.: Sexual dimorphism in plasma leptin concentration. J.

[126026]Clin. Endocr. Metab. 82: 579–584, 1997.

[126027]22462.

[126028]Schubring, C.; Kiess, W.; Englaro, P.; Rascher, W.; Dotsch, J.;

[126029]Hanitsch, S.; Attanasio, A.; Blum, W. F.: Levels of leptin in maternal

[126030]serum, amniotic fluid, and arterial and venous cord blood: relation

[126031]to neonatal and placental weight. J. Clin. Endocr. Metab. 82: 1480–1483,

[126032]1997.

[126033]22463.

[126034]Shanley, L. J.; Irving, A. J.; Rae, M. G.; Ashford, M. L. J.;

[126035]Harvey, J.: Leptin inhibits rat hippocampal neurons via activation

[126036]of large conductance calcium-activated K(+) channels. Nature Neurosci. 5:

[126037]299–300, 2002.

[126038]22464.

[126039]Shimomura, I.; Hammer, R. E.; Ikemoto, S.; Brown, M. S.;
Goldstein,

[126040]J. L.: Leptin reverses insulin resistance and diabetes mellitus in

[126041]mice with congenital lipodystrophy. *Nature* 401: 73–76,
1999.

[126042]22465.

[126043]Shimomura, I.; Matsuda, M.; Hammer, R. E.; Bashmakov,
Y.; Brown,

[126044]M. S.; Goldstein, J. L.: Decreased IRS–2 and increased
SREBP–1c lead

[126045]to mixed insulin resistance and sensitivity in livers of
lipodystrophic

[126046]and ob/ob mice. *Molec. Cell* 6: 77–86, 2000.

[126047]22466.

[126048]Sierra–Honigsmann, M. R.; Nath, A. K.; Murakami, C.; Gar-
cia–Cardena,

[126049]G.; Papapetropoulos, A.; Sessa, W. C.; Madge, L. A.;
Schechner, J.

[126050]S.; Schwabb, M. B.; Polverini, P. J.; Flores-Riveros, J. R.: Biological

[126051]action of leptin as an angiogenic factor. Science 281: 1583–1586,

[126052]1998.

[126053]22467.

[126054]Sooranna, S. R.; Ward, S.; Bajoria, R.: Fetal leptin influences

[126055]birth weight in twins with discordant growth. Pediat. Res. 49: 667–672,

[126056]2001.

[126057]22468.

[126058]Stattin, P.; Soderberg, S.; Hallmans, G.; Bylund, A.; Kaaks, R.;

[126059]Stenman, U.-H.; Bergh, A.; Olsson, T.: Leptin is associated with

[126060]increased prostate cancer risk: a nested case-referent study. J.

[126061]Clin. Endocr. Metab. 86: 1341–1345, 2001.

[126062]22469.

[126063]Strobel, A.; Issad, T.; Camoin, L.; Ozata, M.; Strosberg, A.

D.

[126064]: A leptin missense mutation associated with hypogonadism and morbid

[126065]obesity. (Letter) Nature Genet. 18: 213–215, 1998.

[126066]22470.

[126067]Szczypka, M. S.; Rainey, M. A.; Palmiter, R. D.: Dopamine is

[126068]required for hyperphagia in Lep(ob/ob) mice. Nature Genet. 25: 102–104,

[126069]2000.

[126070]22471.

[126071]Trayhurn, P.; Thurlby, P. L.; James, W. P. T.: Thermogenic defect

[126072]in pre-obese ob/ob mice. (Letter) Nature 266: 60–62, 1977.

[126073]22472.

[126074]Verploegen, S. A. B. W.; Plaetinck, G.; Devos, R.; Van der Heyden,

[126075]J.; Guisez, Y.: A human leptin mutant induces weight gain in normal

[126076]mice. FEBS Lett. 405: 237–240, 1997.

[126077]22473.

[126078]Weigle, D. S.; Ganter, S. L.; Kuijper, J. L.; Leonetti, D. L.;

[126079]Boyko, E. J.; Fujimoto, W. Y.: Effect of regional fat distribution

[126080]and Prader–Willi syndrome on plasma leptin levels. J. Clin. Endocr.

[126081]Metab. 82: 566–570, 1997.

[126082]22474.

[126083]Wellhoener, P.; Fruehwald–Schultes, B.; Kern, W.; Dantz, D.; Kerner,

[126084]W.; Born, J.; Fehm, H. L.; Peters, A.: Glucose metabolism rather

[126085]than insulin is a main determinant of leptin secretion in humans. J.

[126086]Clin. Endocr. Metab. 85: 1267–1271, 2000.

[126087]22475.

[126088]Zhang, Y.; Proenca, R.; Maffei, M.; Barone, M.; Leopold, L.; Friedman,

[126089]J. M.: Positional cloning of the mouse obese gene and its human homologue. Nature 372:

[126090]425–432, 1994.

[126091]22476.

[126092]Zhao, A. Z.; Huan, J.-N.; Gupta, S.; Pal, R.; Sahu, A.: A
phosphatidylinositol

[126093]3-kinase-phosphodiesterase 3B-cyclic AMP pathway in
hypothalamic action

[126094]of leptin on feeding. *Nature Neurosci.* 5: 727-728, 2002.

[126095]22477.

[126096]Siracusa, L. D.; Rosner, M. H.; Vigano, M. A.; Gilbert, D. J.;

[126097]Staudt, L. M.; Copeland, N. G.; Jenkins, N. A.: Chromoso-
mal location

[126098]of the octamer transcription factors, Otf-1, Otf-2, and
Otf-3, defines

[126099]multiple Otf-3-related sequences dispersed in the mouse
genome. *Genomics* 10:

[126100]313-326, 1991.

[126101]22478.

[126102]Sturm, R. A.; Eyre, H. J.; Baker, E.; Sutherland, G. R.: The
human

[126103]OTF1 locus which overlaps the CD3Z gene is located at
1q22-q23. *Cytogenet.*

[126104]*Cell Genet.* 68: 231-232, 1995.

[126105]22479.

[126106]Marshall, J. B.; Diaz-Arias, A. A.; Bochna, G. S.; Vogele, K.

[126107]A.: Achalasia due to diffuse esophageal leiomyomatosis
and inherited

[126108]as an autosomal dominant disorder. Gastroenterology 98:
1358-1365,

[126109]1990.

[126110]22480.

[126111]Mattei, M. G.; d'Auriol, L.; Andre, C.; Passage, E.; Mattei, J.

[126112]F.; Galibert, F.: Assignment of the human c-kit proto-
oncogene to

[126113]the q11-q12 region of chromosome 4, using in situ hy-
bridization. (Abstract) Cytogenet.

[126114]Cell Genet. 46: 657, 1987.

[126115]22481.

[126116]Nagata, H.; Worobec, A. S.; Oh, C. K.; Chowdhury, B. A.;
Tannenbaum,

[126117]S.; Suzuki, Y.; Metcalfe, D. D.: Identification of a point mu-
tation

[126118]in the catalytic domain of the protooncogene c-kit in pe-
ripheral blood

[126119]mononuclear cells of patients who have mastocytosis with
an associated

[126120]hematologic disorder. Proc. Nat. Acad. Sci. 92:
10560–10564, 1995.

[126121]22482.

[126122]Nishida, T.; Hirota, S.; Taniguchi, M.; Hashimoto, K.;
Isozaki,

[126123]K.; Nakamura, H.; Kanakura, Y.; Tanaka, T.; Takabayashi,
A.; Matsuda,

[126124]H.; Kitamura, Y.: Familial gastrointestinal stromal tumours
with

[126125]germline mutation of the KIT gene. (Letter) Nature Genet.
19: 323–324,

[126126]1998.

[126127]22483.

[126128]Nocka, K.; Tan, J. C.; Chiu, E.; Chu, T. Y.; Ray, P.; Trakt-
man,

[126129]P.; Besmer, P.: Molecular bases of dominant negative and
loss of

[126130]function mutations at the murine c-kit/white spotting lo-
cus: W-37,

[126131]W-v, W-41 and W. EMBO J. 9: 1805–1813, 1990.

[126132]22484.

[126133]Nomura, K.; Hatayama, I.; Narita, T.; Kaneko, T.; Shiraishi, M.

[126134]: A novel KIT gene missense mutation in a Japanese family with piebaldism.

[126135](Letter) J. Invest. Derm. 111: 337–338, 1998.

[126136]22485.

[126137]Pignon, J.–M.; Giraudier, S.; Duquesnoy, P.; Jouault, H.; Imbert,

[126138]M.; Vainchenker, W.; Vernant, J.–P.; Tulliez, M.: A new c-kit mutation

[126139]in a case of aggressive mast cell disease. Brit. J. Haemat. 96:

[126140]374–376, 1997.

[126141]22486.

[126142]Reinsch, N.; Thomsen, H.; Xu, N.; Brink, M.; Looft, C.; Kalm,

[126143]E.; Brockmann, G. A.; Grupe, S.; Kuhn, C.; Schwerin, M.; Leyhe, B.;

[126144]Hiendleder, S.; Erhardt, G.; Medjugorac, I.; Russ, I.; Forster, M.;

[126145]Reents, R.; Averdunk, G.: A QTL for the degree of spotting
in cattle

[126146]shows synteny with the KIT locus on chromosome 6. J.
Hered. 90:

[126147]629–634, 1999.

[126148]22487.

[126149]Selmanowitz, V. J.; Rabinowitz, A. D.; Orentreich, N.;
Wenk, E.

[126150]: Pigmentary correction of piebaldism by autografts. I.
Procedures

[126151]and clinical findings. J. Derm. Surg. Oncol. 3: 615–622,
1977.

[126152]22488.

[126153]Spritz, R. A.; Beighton, P.: Piebaldism with deafness:
molecular

[126154]evidence for an expanded syndrome. Am. J. Med. Genet.
75: 101–103,

[126155]1998.

[126156]22489.

[126157]Spritz, R. A.; Droetto, S.; Fukushima, Y.: Deletion of the
KIT

[126158]and PDGFRA genes in a patient with piebaldism. Am. J.

Med. Genet. 44:

[126159]492–495, 1992.

[126160]22490.

[126161]Spritz, R. A.; Giebel, L. B.: Mutation of the c-kit

(mast/stem

[126162]cell growth factor receptor) proto-oncogene in human
piebaldism. (Abstract) Am.

[126163]J. Hum. Genet. 49 (suppl.): 38, 1991.

[126164]22491.

[126165]Spritz, R. A.; Giebel, L. B.; Holmes, S. A.: Dominant nega-
tive

[126166]and loss of function mutations of the c-kit (mast/stem
cell growth

[126167]factor receptor) proto-oncogene in human piebaldism.
Am. J. Hum.

[126168]Genet. 50: 261–269, 1992.

[126169]22492.

[126170]Spritz, R. A.; Holmes, S. A.; Ramesar, R.; Greenberg, J.;
Curtis,

[126171]D.; Beighton, P.: Mutations of the KIT (mast/stem cell

growth factor

[126172]receptor) proto-oncogene account for a continuous range
of phenotypes

[126173]in human piebaldism. Am. J. Hum. Genet. 51: 1058–1065,
1992.

[126174]22493.

[126175]Syrris, P.; Malik, N. M.; Murday, V. A.; Patton, M. A.; Carter,

[126176]N. D.; Hughes, H. E.; Metcalfe, K.: Three novel mutations
of the

[126177]proto-oncogene KIT cause human piebaldism. (Letter) Am.
J. Med. Genet. 95:

[126178]79–81, 2000.

[126179]22494.

[126180]Tan, J. C.; Nocka, K.; Ray, P.; Traktman, P.; Besmer, P.: The

[126181]dominant W42 spotting phenotype results from a mis-
sense mutation in

[126182]the c-kit receptor kinase. Science 247: 209–212, 1990.

[126183]22495.

[126184]Taylor, M. L.; Dastych, J.; Sehgal, D.; Sundstrom, M.; Nils-
son,

[126185]G.; Akin, C.; Mage, R. G.; Metcalfe, D. D.: The Kit-

activating mutation

[126186]D816V enhances stem cell factor-dependent chemotaxis.

Blood 98:

[126187]1195–1199, 2001.

[126188]22496.

[126189]Thomsen, L.; Robinson, T. L.; Lee, J. C. F.; Farraway, L. A.;

[126190]Hughes, M. J. G.; Andrews, D. W.; Huizinga, J. D.: Interstitial cells

[126191]of Cajal generate a rhythmic pacemaker current. Nature
Med. 4: 848–851,

[126192]1998.

[126193]22497.

[126194]Tian, Q.; Frierson, H. F., Jr.; Krystal, G. W.; Moskaluk, C. A.

[126195]: Activating c-kit gene mutations in human germ cell tumors. Am.

[126196]J. Path. 154: 1643–1647, 1999.

[126197]22498.

[126198]Tsujimura, T.; Furitsu, T.; Morimoto, M.; Isozaki, K.; Nomura,

[126199]S.; Matsuzawa, Y.; Kitamura, Y.; Kanakura, Y.: Ligand-

independent

[126200]activation of c-kit receptor tyrosine kinase in a murine mastocytoma

[126201]cell line P-815 generated by a point mutation. Blood 83: 2619-2626,

[126202]1994.

[126203]22499.

[126204]Vandenbark, G. R.; deCastro, C. M.; Taylor, H.; Dew-Knight, S.;

[126205]Kaufman, R. E.: Cloning and structural analysis of the human c-kit

[126206]gene. Oncogene 7: 1259-1266, 1992.

[126207]22500.

[126208]Worobec, A. S.; Semere, T.; Nagata, H.; Metcalfe, D. D.: Clinical

[126209]correlates of the presence of the asp816val c-kit mutation in the

[126210]peripheral blood mononuclear cells of patients with mastocytosis. Cancer 83:

[126211]2120-2129, 1998.

[126212]22501.

[126213]Yarden, Y.; Kuang, W.-J.; Yang-Feng, T.; Coussens, L.;
Munemitsu,

[126214]S.; Dull, T. J.; Chen, E.; Schlessinger, J.; Francke, U.; Ull-
rich,

[126215]A.: Human proto-oncogene c-kit: a new cell surface re-
ceptor tyrosine

[126216]kinase for an unidentified ligand. EMBO J. 6: 3341-3351,
1987.

[126217]22502.

[126218]Cheah, M. S. C.; Ley, T. J.; Tronick, S. R.; Robbins, K. C.:
Fgr

[126219]proto-oncogene mRNA induced in B lymphocytes by Ep-
stein-Barr virus

[126220]infection. Nature 319: 238-240, 1986.

[126221]22503.

[126222]Dracopoli, N. C.; Stanger, B. Z.; Lager, M.; Housman, D. E.:
Localization

[126223]of the FGR protooncogene on the genetic linkage map of
human chromosome

[126224]1p. Genomics 3: 124-128, 1988.

[126225]22504.

[126226]Le Beau, M. M.; Westbrook, C. A.; Diaz, M. O.; Rowley, J.
D.:

[126227]Evidence for two distinct c-src loci on human chromo-
somes 1 and 20. Nature 312:

[126228]70-71, 1984.

[126229]22505.

[126230]Lebo, R. V.; Cheung, M.-C.; Bruce, B. D.: Rapid gene map-
ping by

[126231]dual laser chromosome sorting and spot blot DNA analy-
sis. (Abstract) Am.

[126232]J. Hum. Genet. 36: 101S, 1984.

[126233]22506.

[126234]Nishizawa, M.; Semba, K.; Yoshida, M. C.; Yamamoto, T.;
Sasaki,

[126235]M.; Toyoshima, K.: Structure, expression, and chromoso-
mal location

[126236]of the human c-fgr gene. Molec. Cell. Biol. 6: 511-517,
1986.

[126237]22507.

[126238]Kaiser, U. B.; Halvorsen, L. M.; Chen, M. T.: SP1, steroido-
genic

- [126239] factor 1 (SF-1), and early growth response protein 1 (Egr-1) binding
- [126240] sites form a tripartite gonadotropin-releasing hormone response element
- [126241] in the rat luteinizing hormone-beta gene promoter: an integral role
- [126242] for SF-1. *Molec. Endocr.* 14: 1235-1245, 2000.
- [126243] 22508.
- [126244] Liao, W. X.; Goh, H. H.; Roy, A. C.: Functional characterization
- [126245] of a natural variant of luteinizing hormone. *Hum. Genet.* 111: 219-224,
- [126246] 2002.
- [126247] 22509.
- [126248] Liao, W. X.; Roy, a. C.; Chan, C.; Arulkumaran, S.; Ratnam, S.
- [126249] S.: A new molecular variant of luteinizing hormone associated with
- [126250] female infertility. *Fertil. Steril.* 69: 102-106, 1998.
- [126251] 22510.
- [126252] Manna, P. R.; Joshi, L.; Reinhold, V. N.; Aubert, M. L.; Sug-

anuma,

[126253]N.; Pettersson, K.; Huhtaniemi, I. T.: Synthesis, purification
and

[126254]structural and functional characterization of recombinant
form of

[126255]a common genetic variant of human luteinizing hormone.
Hum. Molec.

[126256]Genet. 11: 301–315, 2002.

[126257]22511.

[126258]Mohrenweiser, H. W.; Tynan, K. M.; Branscomb, E. W.; de
Jong,

[126259]P. J.; Olsen, A.; Trask, B.; Carrano, A. V.: Development of
an integrated

[126260]genetic, functional and physical map of human chromo-
some 19. (Abstract) Cytogenet.

[126261]Cell Genet. 58: 2021 only, 1991.

[126262]22512.

[126263]Muyan, M.; Furuhashi, M.; Sugahara, T.; Boime, I.: The car-
boxy-terminal

[126264]region of the beta-subunits of luteinizing hormone and
chorionic gonadotropin

[126265]differentially influence secretion and assembly of the het-

erodimers. Molec.

[126266]Endocr. 10: 1678–1687, 1996.

[126267]22513.

[126268]Park, I. J.; Burnett, L. S.; Jones, H. W., Jr.; Migeon, C. J.;

[126269]Blizzard, R. M.: A case of male pseudohermaphroditism
associated

[126270]with elevated LH, normal FSH and low testosterone possibly due to

[126271]the secretion of an abnormal LH molecule. Acta Endocr.
83: 173–181,

[126272]1976.

[126273]22514.

[126274]Pettersson, K.; Ding, Y.-Q.; Huhtaniemi, I.: An immunologically

[126275]anomalous luteinizing hormone variant in a healthy
woman. J. Clin.

[126276]Endocr. Metab. 74: 164–171, 1992.

[126277]22515.

[126278]Quirk, C. C.; Lozada, K. L.; Keri, R. A.; Nilson, J. H.: A single

[126279]Pitx1 binding site is essential for activity of the LH-beta

promoter

[126280]in transgenic mice. *Molec. Endocr.* 15: 734–746, 2001.

[126281]22516.

[126282]Raivio, T.; Huhtaniemi, I.; Anttila, R.; Siimes, M. A.; Hage-
nas,

[126283]L.; Nilsson, C.; Pettersson, K.; Dunkel, L.: The role of
luteinizing

[126284]hormone-beta gene polymorphism in the onset and pro-
gression of puberty

[126285]in healthy boys. *J. Clin. Endocr. Metab.* 81: 3278–3282,
1996.

[126286]22517.

[126287]Ramanujam, L. N.; Liao, W. X.; Roy, A. C.; Loganath, A.;
Goh,

[126288]H. H.; Ng, S. C.: Association of molecular variants of
luteinizing

[126289]hormone with menstrual disorders. *Clin. Endocrin.* 51:
243–246, 1999.

[126290]22518.

[126291]Ramanujam, L. N.; Liao, W. X.; Roy, A. C.; Ng, S. C.: Asso-
ciation

[126292]of molecular variants of luteinizing hormone with male in-

fertility. Hum.

[126293]Reprod. 15: 925–928, 2000.

[126294]22519.

[126295]Roy, A. C.; Liao, W. X.; Chen, Y.; Arulkumaran, S.; Ratnam, S.

[126296]S.: Identification of 7 novel mutations in LH beta-subunit gene by

[126297]SSCP. Molec. Cell Biochem. 165: 151–153, 1996.

[126298]22520.

[126299]Shome, B.; Parlow, A. F.: The primary structure of the hormone-specific,

[126300]beta subunit of human pituitary luteinizing hormone (hLH). J. Clin.

[126301]Endocr. 36: 618–621, 1973.

[126302]22521.

[126303]Tapanainen, J. S.; Koivunen, R.; Fauser, B. C. J. M.; Taylor,

[126304]A. E.; Clayton, R. N.; Rajkova, M.; White, D.; Franks, S.; Anttila,

[126305]L.; Pettersson, K. S. I.; Huhtaniemi, I. T.: A new contributing factor

[126306]to polycystic ovary syndrome: the genetic variant of

luteinizing hormone. J.

[126307]Clin. Endocr. Metab. 84: 1711–1715, 1999.

[126308]22522.

[126309]van den Beld, A. W.; Huhtaniemi, I. T.; Pettersson, K. S. L.;

[126310]Pols, H. A. P.; Grobbee, D. E.; de Jong, F. H.; Lamberts, S.
W. J.

[126311]: Luteinizing hormone and different genetic variants, as
indicators

[126312]of frailty in healthy elderly men. J. Clin. Endocr. Metab. 84:
1334–1339,

[126313]1999.

[126314]22523.

[126315]Weck, J.; Anderson, A. C.; Jenkins, S.; Fallest, P. C.; Shup-
nik,

[126316]M. A.: Divergent, composite gonadotropin–releasing hor-
mone–responsive

[126317]elements in the rat luteinizing hormone subunit genes.
Molec. Endocr. 14:

[126318]472–485, 2000.

[126319]22524.

[126320]Weiss, J.; Axelrod, L.; Whitcomb, R. W.; Harris, P. E.; Crow-

ley,

[126321]W. F.; Jameson, J. L.: Hypogonadism caused by a single amino acid

[126322]substitution in the beta subunit of luteinizing hormone.
New Eng.

[126323]J. Med. 326: 179–183, 1992.

[126324]22525.

[126325]Atger, M.; Misrahi, M.; Sar, S.; Le Flem, L.; Dessen, P.; Milgrom,

[126326]E.: Structure of the human luteinizing hormone–
choriogonadotropin

[126327]receptor gene: unusual promoter and 5–prime non–
coding regions. Molec.

[126328]Cell. Endocr. 111: 113–123, 1995.

[126329]22526.

[126330]Bardin, C. W.; Bullock, I. P.; Sherins, R. J.; Mowszowicz, I.;

[126331]Blackburn, W. R.: Androgen metabolism and mechanism of
action in

[126332]male pseudohermaphroditism: a study of testicular feminization. Recent

[126333]Prog. Horm. Res. 29: 65–109, 1973.

[126334]22527.

[126335]Berthezene, F.; Forest, M. G.; Grimaud, J. A.; Claustrat, B.;
Mornex,

[126336]R.: Leydig-cell agenesis: a cause of male pseudo-
hermaphroditism. New

[126337]Eng. J. Med. 295: 969-972, 1976.

[126338]22528.

[126339]Brown, D. M.; Markland, C.; Dehner, L. P.: Leydig cell hy-
poplasia--a

[126340]cause of male pseudohermaphroditism. J. Clin. Endocr.
Metab. 46:

[126341]1-7, 1978.

[126342]22529.

[126343]Canto, P.; Soderlund, D.; Ramon, G.; Nishimura, E.;
Mendez, J.

[126344]P.: Mutational analysis of the luteinizing hormone receptor
gene

[126345]in two individuals with Leydig cell tumors. Am. J. Med.
Genet. 108:

[126346]148-152, 2002.

[126347]22530.

[126348]Eblen, A.; Bao, S.; Lei, Z. M.; Nakajima, S. T.; Rao, C. V.:

The

[126349]presence of functional luteinizing hormone/chorionic gonadotropin

[126350]receptors in human sperm. J. Clin. Endocr. Metab. 86:
2643–2648,

[126351]2001.

[126352]22531.

[126353]El-Awady, M. K.; Temtamy, S. A.; Salam, M. A.; Gad, Y. Z.:

Familial

[126354]Leydig cell hypoplasia as a cause of male pseudo-hermaphroditism. Hum.

[126355]Hered. 37: 36–40, 1987.

[126356]22532.

[126357]Evans, B. A. J.; Bowen, D. J.; Smith, P. J.; Clayton, P. E.;

Gregory,

[126358]J. W.: A new point mutation in the luteinising hormone receptor gene

[126359]in familial and sporadic male limited precocious puberty: genotype

[126360]does not always correlate with phenotype. J. Med. Genet. 33: 143–147,

[126361]1996.

[126362]22533.

[126363]Gospodarowicz, D.: Properties of the luteinizing hormone
receptor

[126364]of isolated bovine corpus luteum plasma membrane. J.
Biol. Chem. 248:

[126365]5042–5049, 1973.

[126366]22534.

[126367]Gromoll, J.; Eiholzer, U.; Nieschlag, E.; Simoni, M.: Male
hypogonadism

[126368]caused by homozygous deletion of exon 10 of the
luteinizing hormone

[126369](LH) receptor: differential action of human chorionic go-
nadotropin

[126370]and LH. J. Clin. Endocr. Metab. 85: 2281–2286, 2000.

[126371]22535.

- [126372]Robert, B.; Barton, P.; Minty, A.; Daubas, P.; Weydert, A.;
Bonhomme,
[126373]F.; Catalan, J.; Chazottes, D.; Guenet, J. L.; Gros, F.; Buck-
ingham,
[126374]M. E.: Investigation of genetic linkage between myosin and
actin
[126375]genes using an interspecific mouse back-cross. *Nature*
314: 181-183,
[126376]1985.
[126377]22536.
- [126378]Saez, L. J.; Gianola, K. M.; McNally, E. M.; Feghali, R.; Eddy,
[126379]R.; Shows, T. B.; Leinwand, L. A.: Human cardiac myosin
heavy chain
[126380]genes and their linkage in the genome. *Nucleic Acids Res.*
15: 5443-5459,
[126381]1987.
[126382]22537.
- [126383]Ye, K.; Aghdasi, B.; Luo, H. R.; Moriarity, J. L.; Wu, F. Y.;
Hong,
[126384]J. J.; Hurt, K. J.; Bae, S. S.; Suh, P.-G.; Snyder, S. H.: Phos-
pholipase

[126385]C-gamma-1 is a physiological guanine nucleotide exchange factor for

[126386]the nuclear GTPase PIKE. Nature 415: 541-544, 2002.

[126387]22538.

[126388]Lazarou, L. P.; Davies, F.; Sarfarazi, M.; Coles, G. A.; Harper,

[126389]P. S.: Adult polycystic kidney disease and linked RFLPs at the alpha-globin

[126390]locus: a genetic study in the South Wales population. J. Med. Genet. 24:

[126391]466-473, 1987.

[126392]22539.

[126393]Burn, J.; Chapman, P.; Delhanty, J.; Wood, C.; Laloo, F.; Cachon-Gonzalez,

[126394]M. B.; Tsioupra, K.; Church, W.; Rhodes, M.; Gunn, A.: The UK northern

[126395]region genetic register for familial adenomatous polyposis coli: use

[126396]of age of onset, congenital hypertrophy of the retinal pigment epithelium,

[126397]and DNA markers in risk calculations. J. Med. Genet. 28: 289-296,

[126398]1991.

[126399]22540.

[126400]Tsuji, M.; Kawano, S.; Tsuji, S.; Sawaoka, H.; Hori, M.;
DuBois,

[126401]R. N.: Cyclooxygenase regulates angiogenesis induced by
colon cancer

[126402]cells. Cell 93: 705–716, 1998.

[126403]22541.

[126404]Vane, J. R.; Mitchell, J. A.; Appleton, I.; Tomlinson, A.;
Bishop-Bailey,

[126405]D.; Croxtall, J.; Willoughby, D. A.: Inducible isoforms of
cyclooxygenase

[126406]and nitric-oxide synthase in inflammation. Proc. Nat.
Acad. Sci. 91:

[126407]2046–2050, 1994.

[126408]22542.

[126409]Yokoyama, C.; Tanabe, T.: Cloning of human gene encod-
ing prostaglandin

[126410]endoperoxide synthase and primary structure of the en-
zyme. Biochem.

[126411]Biophys. Res. Commun. 165: 888–894, 1989.

[126412]22543.

[126413]Schwarz, H. P.; Heeb, M. J.; Lottenberg, R.; Roberts, H.;
Griffin,

[126414]J. H.: Familial protein S deficiency with a variant protein S
molecule

[126415]in plasma and platelets. Blood 74: 213–221, 1989.

[126416]22544.

[126417]Stanislovitis, P.; Watkins, P. C.; Eddy, R.; Cohen, E. H.;
Dackowski,

[126418]W. R.; Wydro, R. M.; Shows, T. B.: Regional assignment of
the gene

[126419]for protein S to the human chromosome region
3p21–3q21. (Abstract) Am.

[126420]J. Hum. Genet. 41: A187 only, 1987.

[126421]22545.

[126422]Watkins, P. C.; Eddy, R.; Fukushima, Y.; Byers, M. G.; Co-
hen,

[126423]E. H.; Dackowski, W. R.; Wydro, R. M.; Shows, T. B.: The
gene for

[126424]protein S maps near the centromere of human chromo-
some 3. Blood 71:

[126425]238-241, 1988.

[126426]22546.

[126427]Watkins, P. C.; Eddy, R.; Fukushima, Y.; Byers, M. G.; Cohen,

[126428]E. H.; Dackowski, W. R.; Wydro, R. M.; Shows, T. B.: Regional assignment

[126429]of the gene for protein S (PROS) to the human chromosome region 3p11.1-q11.2.

[126430](Abstract) Cytogenet. Cell Genet. 46: 712 only, 1987.

[126431]22547.

[126432]Zoller, B.; Garcia de Frutos, P.; Dahlback, B.: Evaluation of

[126433]the relationship between protein S and C4b-binding protein isoforms

[126434]in hereditary protein S deficiency demonstrating type I and type III

[126435]deficiencies to be phenotypic variants of the same genetic disease. Blood 85:

[126436]3524-3531, 1995.

[126437]22548.

[126438]Kaplan, R.; Morse, B.; Huebner, K.; Croce, C.; Howk, R.; Ravera,

- [126439]M.; Ricca, G.; Jaye, M.; Schlessinger, J.: Cloning of three human
- [126440]tyrosine phosphatases reveals a multigene family of receptor-linked
- [126441]protein-tyrosine-phosphatases expressed in brain. Proc. Nat. Acad.
- [126442]Sci. 87: 7000-7004, 1990.
- [126443]22549.
- [126444]Jirik, F. R.; Anderson, L. L.; Duncan, A. M. V.: The human protein-tyrosine
- [126445]phosphatase PTP-alpha/LRP gene (PTPA) is assigned to chromosome 20p13. Cytogenet.
- [126446]Cell Genet. 60: 117-118, 1992.
- [126447]22550.
- [126448]Jirik, F. R.; Janzen, N. M.; Melhado, I. G.; Harder, K. W.: Cloning
- [126449]and chromosomal assignment of a widely expressed human receptor-like
- [126450]protein-tyrosine phosphatase. FEBS Lett. 273: 239-242, 1990.
- [126451]22551.

[126452]Matthews, R. J.; Cahir, E. D.; Thomas, M. L.: Identification
of

[126453]an additional member of the protein-tyro-
sine-phosphatase family: evidence

[126454]for alternative splicing in the tyrosine phosphatase do-
main. Proc.

[126455]Nat. Acad. Sci. 87: 4444-4448, 1990.

[126456]22552.

[126457]Rao, V. V. N. G.; Loffler, C.; Sap, J.; Schlessinger, J.; Hans-
mann,

[126458]I.: The gene for receptor-linked protein-tyro-
sine-phosphatase (PTPA)

[126459]is assigned to human chromosome 20p12-pter by in situ
hybridization

[126460](ISH and FISH). Genomics 13: 906-907, 1992.

[126461]22553.

[126462]Schnittger, S.; Rao, V. V. N. G.; Deutsch, U.; Gruss, P.;
Balling,

[126463]R.; Hansmann, I.: PAX1, a member of the paired box-
containing class

[126464]of developmental control genes, is mapped to human
chromosome 20p11.2

[126465]by in situ hybridization (ISH and FISH). Genomics 14:
740–744, 1992.

[126466]22554.

[126467]Ahmad, F.; Azevedo, J. L., Jr.; Cortright, R.; Dohm, G. L.;
Goldstein,

[126468]B. J.: Alternations in skeletal muscle protein–tyrosine
phosphatase

[126469]activity and expression in insulin–resistant human obesity
and diabetes. J.

[126470]Clin. Invest. 100: 449–458, 1997.

[126471]22555.

[126472]Brown–Shimer, S.; Johnson, K. A.; Lawrence, J. B.; Johnson,
C.;

[126473]Bruskin, A.; Green, N. R.; Hill, D. E.: Molecular cloning and
chromosome

[126474]mapping of the human gene encoding protein phosphoty-
rosyl phosphatase

[126475]1B. Proc. Nat. Acad. Sci. 87: 5148–5152, 1990.

[126476]22556.

[126477]Charbonneau, H.; Tonks, N. K.; Kumar, S.; Diltz, C. D.;
Harrylock,

- [126478]M.; Cool, D. E.; Krebs, E. G.; Fischer, E. H.; Walsh, K. A.:
Human
[126479]placenta protein-tyrosine-phosphatase: amino acid se-
quence and relationship
[126480]to a family of receptor-like proteins. Proc. Nat. Acad. Sci.
86:
[126481]5252-5256, 1989.
[126482]22557.
[126483]Chernoff, J.; Schievella, A. R.; Jost, C. A.; Erikson, R. L.;
Neel,
[126484]B. G.: Cloning of a cDNA for a major human protein-tyro-
sine-phosphatase. Proc.
[126485]Nat. Acad. Sci. 87: 2735-2739, 1990.
[126486]22558.
[126487]Kerr, N. C.; Wilroy, R. S., Jr.; Kaufman, R. A.: Type 3 Pfeif-
fer
[126488]syndrome with normal thumbs. Am. J. Med. Genet. 66:
138-143, 1996.
[126489]22559.
[126490]Lajeunie, E.; Cameron, R.; El Ghouzzi, V.; de Parseval, N.;
Journeau,

- [126491]P.; Gonzales, M.; Delezoide, A.-L.; Bonaventure, J.; Le
Merrer, M.;
- [126492]Renier, D.: Clinical variability in patients with Apert's syn-
drome. J.
- [126493]Neurosurg. 90: 443-447, 1999.
- [126494]22560.
- [126495]Lajeunie, E.; Ma, H. W.; Bonaventure, J.; Munnich, A.; Le
Merrer,
- [126496]M.; Renier, D.: FGFR2 mutations in Pfeiffer syndrome.
(Letter) Nature
- [126497]Genet. 9: 108, 1995.
- [126498]22561.
- [126499]Ma, H. W.; Lajeunie, E.; Le\Merrer, M.; de\Parseval, N.;
Serville,
- [126500]F.; Weissenbach, J.; Munnich, A.; Renier, D.: No evidence
of genetic
- [126501]heterogeneity in Crouzon craniofacial dysostosis. Hum.
Genet. 96:
- [126502]731-735, 1995.
- [126503]22562.
- [126504]Mattei, M.-G.; Moreau, A.; Gesnel, M.-C.; Houssaint, E.;

Breathnach,

[126505]R.: Assignment by in situ hybridization of a fibroblast growth factor

[126506]receptor gene to human chromosome band 10q26. Hum. Genet. 87: 84–86,

[126507]1991.

[126508]22563.

[126509]Meyers, G. A.; Day, D.; Goldberg, R.; Daentl, D. L.; Przylepa,

[126510]K. A.; Abrams, L. J.; Graham, J. M., Jr.; Feingold, M.; Moeschler,

[126511]J. B.; Rawnsley, E.; Scott, A. F.; Jabs, E. W.: FGFR2 exon IIIa and

[126512]IIIc mutations in Crouzon, Jackson–Weiss, and Pfeiffer syndromes:

[126513]evidence for missense changes, insertions, and a deletion due to alternative

[126514]RNA splicing. Am. J. Hum. Genet. 58: 491–498, 1996.

[126515]22564.

[126516]Miki, T.; Fleming, T. P.; Bottaro, D. P.; Rubin, J. S.; Ron, D.;

[126517]Aaronson, S. A.: Expression cDNA cloning of the KGF re–

ceptor by creation

[126518]of a transforming autocrine loop. Science 251: 72–75,
1991.

[126519]22565.

[126520]Mulvihill, J. J.: Craniofacial syndromes: no such thing as a

[126521]single gene disease. Nature Genet. 9: 101–103, 1995.

[126522]22566.

[126523]Munro, C. S.; Wilkie, A. O. M.: Epidermal mosaicism pro-
ducing

[126524]localised acne: somatic mutation in FGFR2. Lancet 352:
704–705,

[126525]1998.

[126526]22567.

[126527]Okajima, K.; Robinson, L. K.; Hart, M. A.; Abuelo, D. N.;
Cowan,

[126528]L. S.; Hasegawa, T.; Maumenee, I. H.; Jabs, E. W.: Ocular
anterior

[126529]chamber dysgenesis in craniosynostosis syndromes with a
fibroblast

[126530]growth factor receptor 2 mutation. Am. J. Med. Genet. 85:
160–170,

[126531]1999.

[126532]22568.

[126533]Oldridge, M.; Lunt, P. W.; Zackai, E. H.; McDonald-McGinn, D.

[126534]M.; Muenke, M.; Moloney, D. M.; Twigg, S. R. F.; Heath, J. K.; Howard,

[126535]T. D.; Hoganson, G.; Gagnon, D. M.; Jabs, E. W.; Wilkie, A. O. M.

[126536]: Genotype-phenotype correlation for nucleotide substitutions in the

[126537]IgII-IgIII linker of FGFR2. Hum. Molec. Genet. 6: 137-143, 1997.

[126538]22569.

[126539]Oldridge, M.; Zackai, E. H.; McDonald-McGinn, D. M.; Iseki, S.;

[126540]Morriss-Kay, G. M.; Twigg, S. R. F.; Johnson, D.; Wall, S. A.; Jiang,

[126541]W.; Theda, C.; Jabs, E. W.; Wilkie, A. O. M.: De novo Alu-element

[126542]insertions in FGFR2 identify a distinct pathological basis for Apert

[126543]syndrome. Am. J. Hum. Genet. 64: 446-461, 1999.

[126544]22570.

[126545]Park, W.-J.; Meyers, G. A.; Li, X.; Theda, C.; Day, D.; Orlow,

[126546]S. J.; Jones, M. C.; Jabs, E. W.: Novel FGFR2 mutations in
Crouzon

[126547]and Jackson-Weiss syndromes show allelic heterogeneity
and phenotypic

[126548]variability. Hum. Molec. Genet. 4: 1229-1233, 1995.

[126549]22571.

[126550]Passo-Buenos, M. R.; Richieri-Costa, A.; Sertie, A. L.;
Kneppers,

[126551]A.: Presence of the Apert canonical S252W FGFR2 muta-
tion in a patient

[126552]without severe syndactyly. J. Med. Genet. 35: 677-679,
1998.

[126553]22572.

[126554]Priolo, M.; Lerone, M.; Baffico, M.; Baldi, M.; Ravazzolo, R.;

[126555]Cama, A.; Capra, V.; Silengo, M.: Pfeiffer syndrome type 2
associated

[126556]with a single amino acid deletion in the FGFR2 gene.
(Letter) Clin.

[126557]Genet. 58: 81-83, 2000.

[126558]22573.

[126559]Rees, J.: FGFR2 mutations and acne. (Letter) Lancet 352:
668–669,

[126560]1998.

[126561]22574.

[126562]Rutland, P.; Pulleyn, L. J.; Reardon, W.; Baraitser, M.; Hay-
ward,

[126563]R.; Jones, B.; Malcolm, S.; Winter, R. M.; Oldridge, M.;
Slaney, S.

[126564]F.; Poole, M. D.; Wilkie, A. O. M.: Identical mutations in the
FGFR2

[126565]gene cause both Pfeiffer and Crouzon syndrome pheno-
types. Nature

[126566]Genet. 9: 173–176, 1995.

[126567]22575.

[126568]Schaefer, F.; Anderson, C.; Can, B.; Say, B.: Novel mutation

[126569]in the FGFR2 gene at the same codon as the Crouzon syn-
drome mutations

[126570]in a severe Pfeiffer syndrome type 2 case. Am. J. Med.
Genet. 75:

[126571]252–255, 1998.

[126572]22576.

[126573]Shuster, S.: Dermatology in Clinical Medicine. Oxford: Oxford
ford

[126574]Univ. Press , 1978.

[126575]22577.

[126576]Slaney, S. F.; Oldridge, M.; Hurst, J. A.; Morriss-Kay, G. M.;

[126577]Hall, C. M.; Poole, M. D.; Wilkie, A. O. M.: Differential ef-
fects

[126578]of FGFR2 mutations on syndactyly and cleft palate in Apert
syndrome. Am.

[126579]J. Hum. Genet. 58: 923-932, 1996.

[126580]22578.

[126581]Steinberger, D.; Collmann, H.; Schmalenberger, B.; Muller,
U.

[126582]: A novel mutation (A886G) in exon 5 of FGFR2 in mem-
bers of a family

[126583]with Crouzon phenotype and plagiocephaly. J. Med. Genet.
34: 420-422,

[126584]1997.

[126585]22579.

[126586]Steinberger, D.; Mulliken, J. B.; Muller, U.: Predisposition
[126587]for cysteine substitutions in the immunoglobulin-like
chain of FGFR2
[126588]in Crouzon syndrome. Hum. Genet. 96: 113–115, 1995.
[126589]22580.

[126590]Vaughan, K. T.; Weber, F. E.; Ried, T.; Ward, D. C.;
Reinach, F.

[126591]C.; Fischman, D. A.: Human myosin-binding protein H
(MyBP-H): complete

[126592]primary sequence, genomic organization, and chromoso-
mal localization. Genomics 16:
[126593]34–40, 1993.

[126594]22581.

[126595]Torbergesen, T.: A family with dominant hereditary myoto-
nia, muscular

[126596]hypertrophy, and increased muscular irritability, distinct
from myotonia

[126597]congenita Thomsen. Acta Neurol. Scand. 51: 225–232,
1975.

[126598]22582.

[126599]Harding, A. E.: The clinical features and classification of

the

[126600]late onset autosomal dominant cerebellar ataxias: a study
of 11 families,

[126601]including descendants of 'the Drew family of Walworth'.

Brain 105:

[126602]1–28, 1982.

[126603]22583.

[126604]Havener, W. H.: Cerebellar–macular abiotrophy. Arch.

Ophthal. 45:

[126605]40–43, 1951.

[126606]22584.

[126607]Holmberg, M.; Johansson, J.; Forsgren, L.; Heijbel, J.;

Sandgren,

[126608]O.; Holmgren, G.: Localization of autosomal dominant
cerebellar ataxia

[126609]associated with retinal degeneration and anticipation to
chromosome

[126610]3p12–p21.1. Hum. Molec. Genet. 4: 1441–1445, 1995.

[126611]22585.

[126612]Jampel, R. S.; Okazaki, H.; Bernstein, H.: Ophthalmoplegia
and

- [126613]retinal degeneration associated with spinocerebellar ataxia. Arch.
- [126614]Ophthalmol. 66: 247–259, 1961.
- [126615]22586.
- [126616]Jonasson, J.; Juvonen, V.; Sistonen, P.; Ignatius, J.; Johansson,
- [126617]D.; Bjorck, E. J.; Wahlstrom, J.; Melberg, A.; Holmgren, G.; Forsgren,
- [126618]L.; Holmberg, M.: Evidence for a common spinocerebellar ataxia type
- [126619]7 (SCA7) founder mutation in Scandinavia. Europ. J. Hum. Genet. 8:
- [126620]918–922, 2000.
- [126621]22587.
- [126622]Kaytor, M. D.; Duvick, L. A.; Skinner, P. J.; Koob, M. D.; Ranum,
- [126623]L. P. W.; Orr, H. T.: Nuclear localization of the spinocerebellar
- [126624]ataxia type 7 protein, ataxin-7. Hum. Molec. Genet. 8: 1657–1664,
- [126625]1999.

[126626]22588.

[126627]Koob, M. D.; Benzow, K. A.; Bird, T. D.; Day, J. W.; Moseley,

[126628]M. L.; Ranum, L. P. W.: Rapid cloning of expanded trinucleotide repeat

[126629]sequences from genomic DNA. *Nature Genet.* 18: 72–75, 1998.

[126630]22589.

[126631]Krols, L.; Martin, J.-J.; David, G.; Van Regemorter, N.; Benomar,

[126632]A.; Lofgren, A.; Stevanin, G.; Durr, A.; Brice, A.; Van Broeckhoven,

[126633]C.: Refinement of the locus for autosomal dominant cerebellar ataxia

[126634]type II to chromosome 3p21.1–14.1. *Hum. Genet.* 99: 225–232, 1997.

[126635]22590.

[126636]Lebre, A.-S.; Jamot, L.; Takahashi, J.; Spassky, N.; Lepage,

[126637]C.; Ravise, N.; Zander, C.; Fujigasaki, H.; Kussel-Andermann, P.;

[126638]Duyckaerts, C.; Camonis, J. H.; Brice, A.: Ataxin-7 interacts with

[126639]a Cbl-associated protein that it recruits into neuronal in-
tranuclear

[126640]inclusions. Hum. Molec. Genet. 10: 1201-1213, 2001.

[126641]22591.

[126642]Lindblad, K.; Savontaus, M.-L.; Stevanin, G.; Holmberg, M.;
Digre,

[126643]K.; Zander, C.; Ehrsson, H.; David, G.; Benomar, A.;
Nikoskelainen,

[126644]E.; Trottier, Y.; Holmgren, G.; Ptacek, L. J.; Anttinen, A.;
Brice,

[126645]A.; Schalling, M.: An expanded CAG repeat sequence in
spinocerebellar

[126646]ataxia type 7. Genome Res. 6: 965-971, 1996.

[126647]22592.

[126648]Matilla, A.; Gorbea, C.; Einum, D. D.; Townsend, J.; Micha-
lik,

[126649]A.; van Broeckhoven, C.; Jensen, C. C.; Murphy, K. J.;
Ptacek, L.

[126650]J.; Fu, Y.-H.: Association of ataxin-7 with the proteasome
subunit

[126651]S4 of the 19S regulatory complex. Hum. Molec. Genet. 10:
2821-2831,

[126652]2001.

[126653]22593.

[126654]Michalik, A.; Del-Favero, J.; Mauger, C.; Lofgren, A.; Van Broeckhoven,

[126655]C.: Genomic organisation of the spinocerebellar ataxia type 7 (SCA7)

[126656]gene responsible for autosomal dominant cerebellar ataxia with retinal

[126657]degeneration. Hum. Genet. 105: 410-417, 1999.

[126658]22594.

[126659]Stevanin, G.; Giunti, P.; David, G.; Belal, S.; Durr, A.; Ruberg,

[126660]M.; Wood, N.; Brice, A.: De novo expansion of intermediate alleles

[126661]in spinocerebellar ataxia 7. Hum. Molec. Genet. 7: 1809-1813, 1998.

[126662]22595.

[126663]Trottier, Y.; Lutz, Y.; Stevanin, G.; Imbert, G.; Devys, D.; Cancel,

[126664]G.; Saudou, F.; Weber, C.; David, G.; Tora, L.; Agid, Y.; Brice, A.;

- [126665]Mandel, J.-L.: Polyglutamine expansion as a pathological epitope
in Huntington's disease and four dominant cerebellar ataxias. *Nature* 378: 403–406, 1995.
- [126668]22596.
- [126669]Wadia, N. H.; Swami, R. K.: A new form of heredo-familial spinocerebellar degeneration with slow eye movements (nine families). *Brain* 94: 359–374, 1971.
- [126672]22597.
- [126673]Weiner, L. P.; Konigsmark, B. W.; Stoll, J., Jr.; Magladery, J. W.: Hereditary olivopontocerebellar atrophy with retinal degeneration: report of a family through six generations. *Arch. Neurol.* 16: 364–376, 1967.
- [126676]1967.
- [126677]22598.
- [126678]Woodworth, J. A.; Beckett, R. S.; Netsky, M. G.: A composite

[126679] of hereditary ataxias: a familial disorder with features of olivopontocerebellar

[126680] atrophy, Leber's optic atrophy and Friedreich's ataxia.
Arch. Intern.

[126681] Med. 104: 594–606, 1959.

[126682] 22599.

[126683] Yvert, G.; Lindenberg, K. S.; Picaud, S.; Landwehrmeyer, G.
B.;

[126684] Sahel, J. –A.; Mandel, J.–L.: Expanded polyglutamines induce neurodegeneration

[126685] and trans-neuronal alterations in cerebellum and retina of SCA7 transgenic

[126686] mice. Hum. Molec. Genet. 9: 2491–2506, 2000.

[126687] 22600.

[126688] Zander, C.; Takahashi, J.; El Hachimi, K. H.; Fujigasaki, H.;

[126689] Albanese, V.; Lebre, A. S.; Stevanin, G.; Duyckaerts, C.;
Brice, A.

[126690]: Similarities between spinocerebellar ataxia type 7 (SCA7)
cell models

[126691] and human brain: proteins recruited in inclusions and activation of

[126692]caspase-3. Hum. Molec. Genet. 10: 2569-2579, 2001.

[126693]22601.

[126694]Cazzaniga, G.; Tosi, S.; Aloisi, A.; Giudici, G.; Daniotti, M.;

[126695]Pioltelli, P.; Kearney, L.; Biondi, A.: The tyrosine kinase
Abl-related

[126696]gene ARG is fused to ETV6 in an AML-M4Eo patient with a
t(1;12)(q25;p13):

[126697]molecular cloning of both reciprocal transcripts. Blood 94:
4370-4373,

[126698]1999.

[126699]22602.

[126700]Iijima, Y.; Ito, T.; Oikawa, T.; Eguchi, M.; Eguchi-Ishimae,
M.;

[126701]Kamada, N.; Kishi, K.; Asano, S.; Sakaki, Y.; Sato, Y.: A new
ETV6/TEL

[126702]partner gene, ARG (ABL-related gene or ABL2), identified
in an AML-M3

[126703]cell line with a t(1;12)(q25;p13) translocation. Blood 95:
2126-2131,

[126704]2000.

[126705]22603.

[126706] Mastick, C. C.; Brady, M. J.; Saltiel, A. R.: Insulin stimulates
[126707] the tyrosine phosphorylation of caveolin. *J. Cell Biol.* 129:
1523–1531,
[126708] 1995.
[126709] 22604.
[126710] Petrelli, A.; Gilestro, G. F.; Lanzardo, S.; Comoglio, P. M.;
Migone,
[126711] N.; Giordano, S.: The endophilin–CIN85–Cbl complex me-
diates ligand–dependent
[126712] downregulation of c–Met. *Nature* 416: 187–190, 2002.
[126713] 22605.
[126714] Soubeyran, P.; Kowanetz, K.; Szymkiewicz, I.; Langdon, W.
Y.; Dikic,
[126715] J.: Cbl–CIN85–endophilin complex mediates ligand–in-
duced downregulation
[126716] of EGF receptors. *Nature* 416: 183–187, 2002.
[126717] 22606.
[126718] Cannizzaro, L. A.; Madaule, P.; Hecht, F.; Axel, R.; Croce,
C.
[126719] M.; Huebner, K.: Chromosome localization of human ARH
genes, a ras–related

[126720]gene family. Genomics 6: 197–203, 1990.

[126721]22607.

[126722]Chardin, P.; Madaule, P.; Tavitian, A.: Coding sequence of human

[126723]rho cDNAs clone 6 and clone 9. Nucleic Acid Res. 16: 2717 only,

[126724]1988.

[126725]22608.

[126726]Madaule, P.; Axel, R.: A novel ras-related gene family. Cell 41:

[126727]31–40, 1985.

[126728]22609.

[126729]Maekawa, M.; Ishizaki, T.; Boku, S.; Watanabe, N.; Fujita, A.;

[126730]Iwamatsu, A.; Obinata, T.; Ohashi, K.; Mizuno, K.; Narumiya, S.:

[126731]Signaling from Rho to the actin cytoskeleton through protein kinases

[126732]ROCK and LIM-kinase. Science 285: 895–898, 1999.

[126733]22610.

[126734]Ridley, A. J.; Hall, A.: The small GTP-binding protein rho regulates

[126735]the assembly of focal adhesions and actin stress fibers in response

[126736]to growth factors. Cell 70: 389–399, 1992.

[126737]22611.

[126738]Fagan, K. P.; Oliveira, L.; Pittler, S. J.: Sequence of rho small

[126739]GTP-binding protein cDNAs from human retina and identification of

[126740]novel 5-prime end cloning artifacts. Exp. Eye Res. 59: 235–237,

[126741]1994.

[126742]22612.

[126743]Morris, S. W.; Valentine, M. B.; Kirstein, M. N.; Huebner, K.:

[126744]Reassignment of the human ARH9 RAS-related gene to chromosome 1p13–p21. Genomics 15:

[126745]677–679, 1993.

[126746]22613.

[126747]Maesaki, R.; Ihara, K.; Shimizu, T.; Kuroda, S.; Kaibuchi, K.;

- [126748]Hakoshima, T.: The structural basis of Rho effector recognition revealed
by the crystal structure of human RhoA complexed with the effector domain of PKN/PRK1. Molec. Cell 4: 793–803, 1999.
22614.
- [126752]Nakamura, M.; Nagano, T.; Chikama, T.; Nishida, T.: Role of the
small GTP-binding protein Rho in epithelial cell migration in the
rabbit cornea. Invest. Ophthalmol. Vis. Sci. 42: 941–947, 2001.
22615.
- [126756]Rao, P. V.; Deng, P.-F.; Kumar, J.; Epstein, D. L.: Modulation
of aqueous humor outflow facility by the Rho kinase-specific inhibitor
Y-27632. Invest. Ophthalmol. Vis. Sci. 42: 1029–1037, 2001.
22616.
- [126760]Alexander, C.; Votruba, M.; Pesch, U. E. A.; Thiselton, D. L.;

[126761]Mayer, S.; Moore, A.; Rodriguez, M.; Kellner, U.; Leo-Kottler, B.;

[126762]Auburger, G.; Bhattacharya, S. S.; Wissinger, B.: OPA1, encoding

[126763]a dynamin-related GTPase, is mutated in autosomal dominant optic atrophy

[126764]linked to chromosome 3q28. Nature Genet. 26: 211–215, 2000.

[126765]22617.

[126766]Rosenfeld, P. J.; Cowley, G. S.; McGee, T. L.; Sandberg, M. A.;

[126767]Berson, E. L.; Dryja, T. P.: A null mutation in the rhodopsin gene

[126768]causes rod photoreceptor dysfunction and autosomal recessive retinitis

[126769]pigmentosa. Nature Genet. 1: 209–213, 1992.

[126770]22618.

[126771]Pulst, S.-M.; Nechiporuk, A.; Nechiporuk, T.; Gispert, S.; Chen,

[126772]X.-N.; Lopes-Cendes, I.; Pearlman, S.; Starkman, S.; Orozco-Diaz,

[126773]G.; Lunkes, A.; DeJong, P.; Rouleau, G. A.; Auburger, G.;

Korenberg,

[126774]J. R.; Figueroa, C.; Sahba, S.: Moderate expansion of a normally

[126775]biallelic trinucleotide repeat in spinocerebellar ataxia type

2. Nature

[126776]Genet. 14: 269–276, 1996.

[126777]22619.

[126778]Sanpei, K.; Takano, H.; Igarashi, S.; Sato, T.; Oyake, M.;

Sasaki,

[126779]H.; Wakisaka, A.; Tashiro, K.; Ishida, Y.; Ikeuchi, T.; Koide, R.;

[126780]Saito, M.; Sato, A.; Tanaka, T.; Hanyu, S.; Takiyama, Y.; Nishizawa,

[126781]M.; Shimizu, N.; Nomura, Y.; Segawa, M.; Iwabuchi, K.; Eguchi, I.;

[126782]Tanaka, H.; Takahashi, H.; Tsuji, S.: Identification of the spinocerebellar

[126783]ataxia type 2 gene using a direct identification of repeat expansion

[126784]and cloning technique, DIRECT. Nature Genet. 14: 277–284, 1996.

[126785]22620.

[126786]Chen, K.; Yang, W.; Grimsby, J.; Shih, J. C.: The human
5-HT2

[126787]receptor is encoded by a multiple intron-exon gene. Brain
Res. Molec.

[126788]Brain Res. 14: 20-26, 1992.

[126789]22621.

[126790]Holmes, C.; Arranz, M. J.; Powell, J. F.; Collier, D. A.; Love-
stone,

[126791]S.: 5-HT-2A and 5-HT-2C receptor polymorphisms and
psychopathology

[126792]in late onset Alzheimer's disease. Hum. Molec. Genet. 7:
1507-1509,

[126793]1998.

[126794]22622.

[126795]Hsieh, C.-L.; Bowcock, A. M.; Farrer, L. A.; Hebert, J. M.;
Huang,

[126796]K. N.; Cavalli-Sforza, L. L.; Julius, D.; Francke, U.: The
serotonin

[126797]receptor subtype 2 locus HTR2 is on human chromosome
13 near genes

[126798]for esterase D and retinoblastoma-1 and on mouse chro-
mosome 14. Somat.

[126799]Cell Molec. Genet. 16: 567–574, 1990.

[126800]22623.

[126801]Kato, M. V.; Shimizu, T.; Nagayoshi, M.; Kaneko, A.;
Sasaki, M.

[126802]S.; Ikawa, Y.: Genomic imprinting of the human serotonin-
receptor

[126803](HTR2) gene involved in development of retinoblastoma.
Am. J. Hum.

[126804]Genet. 59: 1084–1090, 1996.

[126805]22624.

[126806]Liu, J.; Chen, Y.; Kozak, C. A.; Yu, L.: The 5-HT(2) sero-
tonin

[126807]receptor gene Htr-2 is tightly linked to Es-10 on mouse
chromosome

[126808]14. Genomics 11: 231–234, 1991.

[126809]22625.

[126810]Peroutka, S. J.: 5-Hydroxytryptamine receptor subtypes.
Ann.

[126811]Rev. Neurosci. 11: 45–60, 1988.

[126812]22626.

[126813]Sparkes, R. S.; Lan, N.; Klisak, I.; Mohandas, T.; Diep, A.;
Kojis,
[126814]T.; Heinzmann, C.; Shih, J. C.: Assignment of a serotonin
5HT-2 receptor
[126815]gene (HTR2) to human chromosome 13q14-q21 and
mouse chromosome 14. Genomics 9:
[126816]461-465, 1991.
[126817]22627.

[126818]Williams, J.; McGuffin, P.; Nothen, M.; Owen, M. J.; EMAS
Collaborative
[126819]Group: Meta-analysis of association between the 5-HT
(2A) receptor
[126820]T102C polymorphism and schizophrenia. (Letter) Lancet
349: 1221
[126821]only, 1997.
[126822]22628.

[126823]Williams, J.; Spurlock, G.; McGuffin, P.; Mallet, J.; Nothen,
M.
[126824]M.; Gill, M.; Aschauer, H.; Nylander, P. O.; Macciardi, F.;
Owen,
[126825]M. J.: Association between schizophrenia and T102C poly-
morphism of

- [126826]the 5-hydroxytryptophan type 2a-receptor gene. Euro-
pean Multicentre
[126827]Association Study of Schizophrenia (EMASS) Group. Lancet
347: 1294-1296,
[126828]1996.
[126829]22629.
- [126830]Wulff, C.; Wilson, H.; Rudge, J. S.; Wiegand, S. J.; Lunn, S.
F.;
- [126831]Fraser, H. M.: Luteal angiogenesis: prevention and inter-
vention by
[126832]treatment with vascular endothelial growth factor trapA40.
J. Clin.
[126833]Endocr. Metab. 86: 3377-3386, 2001.
[126834]22630.
- [126835]Philipson, L. H.; Hice, R. E.; Schaefer, K.; LaMendola, J.;
Bell,
- [126836]G. I.; Neldon, D. J.; Steiner, D. F.: Sequence and functional
expression
[126837]in *Xenopus* oocytes of a human insulinoma and islet
potassium channel. Proc.
[126838]Nat. Acad. Sci. 88: 53-57, 1991.

[126839]22631.

[126840]McPherson, J. D.; Wasmuth, J. J.; Chandy, K. G.; Swanson, R.; Dethlefs,

[126841]B.; Chandy, G.; Wymore, R.; Ghanshani, S.: Chromosomal localization

[126842]of 7 potassium channel genes. (Abstract) Cytogenet. Cell Genet. 58:

[126843]1979 only, 1991.

[126844]22632.

[126845]Tamkun, M. M.; Knoth, K. M.; Walbridge, J. A.; Kroemer, H.; Roden,

[126846]D. M.; Glover, D. M.: Molecular cloning and characterization of two

[126847]voltage-gated K⁺ channel cDNAs from human ventricle. FASEB J. 5:

[126848]331-337, 1991.

[126849]22633.

[126850]Phromchotikul, T.; Browne, D. L.; Curran, M. E.; Keating, M. T.;

[126851]Litt, M.: Dinucleotide repeat polymorphism at the KCNA5 locus. Hum.

[126852]Molec. Genet. 2: 1512 only, 1993.

[126853]22634.

[126854]Bardien-Kruger, S.; Wulff, H.; Arieff, Z.; Brink, P.; Chandy, K.

[126855]G.; Corfield, V.: Characterisation of the human voltage-gated potassium

[126856]channel gene, KCNA7, a candidate gene for inherited cardiac disorders,

[126857]and its exclusion as cause of progressive familial heart block I (PFHBI). Europ.

[126858]J. Hum. Genet. 10: 36-43, 2002.

[126859]22635.

[126860]Barton, J. W.; Hart, I. M.; Patterson, D.: Mapping of a locus

[126861]correcting lack of phosphoribosylaminoimidazole carboxylase activity

[126862]in Chinese hamster ovary cell Ade(-)D mutants to human chromosome

[126863]4. Genomics 9: 314-321, 1991.

[126864]22636.

[126865]Brayton, K. A.; Chen, Z.; Zhou, G.; Nagy, P. L.; Gavalas, A.; Trent,

[126866]J. M.; Deaven, L. L.; Dixon, J. E.; Zalkin, H.: Two genes for
de

[126867]novo purine nucleotide synthesis on human chromosome
4 are closely

[126868]linked and divergently transcribed. J. Biol. Chem. 269:
5313–5321,

[126869]1994.

[126870]22637.

[126871]Chu, E. H. Y.; Sun, N. C.; Chang, C. C.: Induction of aux-
otrophic

[126872]mutants by treatment of Chinese hamster cells with
5–bromodeoxyuridine

[126873]and black light. Proc. Nat. Acad. Sci. 68: 3459–3463,
1972.

[126874]22638.

[126875]Feldman, R. I.; Taylor, M. W.: Purine mutants of mam-
malian cell

[126876]lines. II. Identification of a phosphoribosylpyrophosphate
amidotransferase–deficient

[126877]mutant of Chinese hamster lung cells. Biochem. Genet.
13: 227–234,

[126878]1975.

[126879]22639.

[126880]Cavalas, A.; Dixon, J. E.; Brayton, K. A.; Zalkin, H.: Coexpression

[126881]of two closely linked avian genes for purine nucleotide synthesis

[126882]from a bidirectional promoter. Molec. Cell. Biol. 13: 4784-4792,

[126883]1993.

[126884]22640.

[126885]Jones, C.; Morse, H. G.; Palmer, D. K.: Comparative gene mapping

[126886]of man and higher primates involving genes assigned to human chromosomes

[126887]3, 4, 11, 21. (Abstract) Cytogenet. Cell Genet. 40: 662-663, 1985.

[126888]22641.

[126889]Stanley, W.; Chu, E. H. Y.: Assignment of the gene for phosphoribosylpyrophosphate

[126890]amidotransferase to the pter-q21 region of human chromosome 4. Cytogenet.

[126891]Cell Genet. 22: 228-231, 1978.

[126892]22642.

[126893]Zalkin, H.; Dixon, J. E.: De novo purine nucleotide biosynthesis. Prog.

[126894]Nucleic Acids Res. Molec. Biol. 42: 259–287, 1992.

[126895]22643.

[126896]Parfrey, P. S.; Bear, J. C.; Morgan, J.; Cramer, B. C.; McManamon,

[126897]P. J.; Gault, M. H.; Churchill, D. N.; Singh, M.; Hewitt, R.; Somlo,

[126898]S.; Reeders, S. T.: The diagnosis and prognosis of autosomal dominant

[126899]polycystic kidney disease. New Eng. J. Med. 323: 1085–1090, 1990.

[126900]22644.

[126901]Peral, B.; Gamble, V.; San Millan, J. L.; Strong, C.; Sloane-Stanley,

[126902]J.; Moreno, F.; Harris, P. C.: Splicing mutations of the polycystic

[126903]kidney disease 1 (PKD1) gene induced by intronic deletion. Hum. Molec.

[126904]Genet. 4: 569–574, 1995.

[126905]22645.

[126906]Peral, B.; Ong, A. C. M.; San Millan, J. L.; Gamble, V.; Rees,

[126907]L.; Harris, P. C.: A stable, nonsense mutation associated
with a

[126908]case of infantile onset polycystic kidney disease 1 (PKD1).
Hum.

[126909]Molec. Genet. 5: 539–542, 1996.

[126910]22646.

[126911]Peral, B.; San Millan, J. L.; Hernandez, C.; Valero, A.; Lath-
rop,

[126912]G. M.; Beckmann, J. S.; Moreno, F.: Estimating locus het-
erogeneity

[126913]in autosomal dominant polycystic kidney disease (ADPKD)
in the Spanish

[126914]population. J. Med. Genet. 30: 910–913, 1993.

[126915]22647.

[126916]Peral, B.; San Millan, J. L.; Ong, A. C. M.; Gamble, V.; Ward,

[126917]C. J.; Strong, C.; Harris, P. C.: Screening the 3–prime re-
gion of

[126918]the polycystic kidney disease 1 (PKD1) gene reveals six
novel mutations. Am.

[126919]J. Hum. Genet. 58: 86–96, 1996.

[126920]22648.

[126921]Pound, S. E.; Carothers, A. D.; Pignatelli, P. M.; Macnicol, A.

[126922]M.; Watson, M. L.; Wright, A. F.: Evidence for linkage dise-
quilibrium

[126923]between D16S94 and the adult onset polycystic kidney
disease (PKD1)

[126924]gene. J. Med. Genet. 29: 247–248, 1992.

[126925]22649.

[126926]Qian, F.; Watnick, T. J.; Onuchic, L. F.; Germino, G. G.: The

[126927]molecular basis of focal cyst formation in human autoso-
mal dominant

[126928]polycystic kidney disease type I. Cell 87: 979–987, 1996.

[126929]22650.

[126930]Reeders, S. T.: Personal Communication. Oxford, England
1986.

[126931]22651.

[126932]Reeders, S. T.: Multilocus polycystic disease. Nature
Genet. 1:

[126933]235–237, 1992.

[126934]22652.

[126935]Reeders, S. T.: Personal Communication. Oxford, England
10/11/1985.

[126936]22653.

[126937]Reeders, S. T.; Breuning, M. H.; Davies, K. E.; Nicholls, R.
D.;

[126938]Jarman, A. P.; Higgs, D. R.; Pearson, P. L.; Weatherall, D. J.:
A

[126939]highly polymorphic DNA marker linked to adult polycystic
kidney disease

[126940]on chromosome 16. Nature 317: 542–544, 1985.

[126941]22654.

[126942]Reeders, S. T.; Breuning, M. H.; Ryynanen, M. A.; Wright,
A. F.;

[126943]Davies, K. E.; King, A. W.; Watson, M. L.; Weatherall, D. J.:
A study

[126944]of genetic linkage heterogeneity in adult polycystic kidney
disease. Hum.

[126945]Genet. 76: 348–351, 1987.

[126946]22655.

[126947]Reeders, S. T.; Keith, T.; Green, P.; Germino, G. G.; Barton,

[126948]N. J.; Lehmann, O. J.; Brown, V. A.; Phipps, P.; Morgan, J.;
Bear,

[126949]J. C.; Parfrey, P.: Regional localization of the autosomal
dominant

[126950]polycystic kidney disease locus. Genomics 3: 150–155,
1988.

[126951]22656.

[126952]Ryynanen, M.; Dolata, M. M.; Lampainen, E.; Reeders, S. T.:
Localisation

[126953]of a mutation producing autosomal dominant polycystic
kidney disease

[126954]without renal failure. J. Med. Genet. 24: 462–465, 1987.

[126955]22657.

[126956]Liu, S. C.; Palek, J.; Prchal, J. T.: Defective spectrin dimer-
dimer

[126957]association in hereditary elliptocytosis. Proc. Nat. Acad.
Sci. 79:

[126958]2072–2076, 1982.

[126959]22658.

[126960]Torban, E.; Pelletier, J.; Goodyer, P.: F329L polymorphism

in

[126961]the human PAX8 gene. Am. J. Med. Genet. 72: 186–187,
1997.

[126962]22659.

[126963]Gartner, J.; Kearns, W.; Pearson, P.; Valle, D.: Characteri-
zation

[126964]and localization of the human 70-kD peroxisomal mem-
brane protein (PMP70)

[126965]gene. (Abstract) Am. J. Hum. Genet. 51 (suppl.): A168,
1992.

[126966]22660.

[126967]Gartner, J.; Kearns, W.; Rosenberg, C.; Pearson, P.;
Copeland,

[126968]N. G.; Gilbert, D. J.; Jenkins, N. A.; Valle, D.: Localization of

[126969]the 70-kDa peroxisomal membrane protein to human
1p21-p22 and mouse 22661.

[126970]Genomics 15: 412–414, 1993.

[126971]

[126972]3. Gartner, J.; Moser, H.; Valle, D.: Mutations in the 70K
peroxisomal

[126973]membrane protein gene in Zellweger syndrome. Nature

Genet. 1: 16–23,

[126974]1992.

[126975]22662.

[126976]Vilain, C.; Rydlewski, C.; Duprez, L.; Heinrichs, C.;

Abramowicz,

[126977]M.; Malvaux, P.; Renneboog, B.; Parma, J.; Costagliola, S.;

Vassart,

[126978]G.: Autosomal dominant transmission of congenital thyroid hypoplasia

[126979]due to loss-of-function mutation of PAX8. J. Clin. Endocr. Metab. 86:

[126980]234–238, 2001.

[126981]22663.

[126982]Walther, C.; Guenet, J.–L.; Simon, D.; Deutsch, U.; Jostes, B.;

[126983]Goulding, M.; Plachov, D.; Balling, R.; Gruss, P.: Pax: a murine

[126984]multigene family of paired box-containing genes. Genomics 11: 424–434,

[126985]1991.

[126986]22664.

[126987] Grueneberg, D. A.; Natesan, S.; Alexandre, C.; Gilman, M.
Z.:

[126988] Human and Drosophila homeodomain proteins that en-
hance the DNA-binding

[126989] activity of serum response factor. Science 257:
1089-1095, 1992.

[126990] 22665.

[126991] Kern, M. J.; Argao, E. A.; Birkenmeier, E. H.; Rowe, L. B.;
Potter,

[126992] S. S.: Genomic organization and chromosome localization
of the murine

[126993] homeobox gene Pmx. Genomics 19: 334-340, 1994.

[126994] 22666.

[126995] Nakamura, T.; Yamazaki, Y.; Hatano, Y.; Miura, I.: NUP98
is fused

[126996] to PMX1 homeobox gene in human acute myelogenous
leukemia with chromosome

[126997] translocation t(1;11)(q23;p15). Blood 94: 741-747, 1999.

[126998] 22667.

[126999] Norris, R. A.; Scott, K. K.; Moore, C. S.; Stetten, G.; Brown,

[127000] C. R.; Jabs, E. W.; Wulfsberg, E. A.; Yu, J.; Kern, M. J.: Hu-

man

[127001]PRRX1 and PRRX2 genes: cloning, expression, genomic localization,

[127002]and exclusion as disease genes for Nager syndrome.

Mammalian Genome 11:

[127003]1000–1005, 2000.

[127004]22668.

[127005]Riazuddin, S.; Castelein, C. M.; Ahmed, Z. M.; Lalwani, A. K.;

[127006]Mastroianni, M. A.; Naz, S.; Smith, T. N.; Liburd, N. A.; Friedman,

[127007]T. B.; Griffith, A. J.; Riazuddin, S.; Wilcox, E. R.: Dominant modifier

[127008]DFNM1 suppresses recessive deafness DFNB26. Nature Genet. 26: 431–434,

[127009]2000.

[127010]22669.

[127011]Schaller, J.: Illness resembling lupus erythematosus in mothers

[127012]of boys with chronic granulomatous disease. Ann. Intern. Med. 76:

[127013]747–750, 1972.

[127014]22670.

[127015]Theofilopoulos, A. N.; Dixon, F. J.: Murine models of systemic
temic

[127016]lupus erythematosus. Adv. Immun. 37: 269–390, 1985.

[127017]22671.

[127018]Eng, C.; Peacocke, M.: PTEN and inherited hamartoma-cancer syndromes.

[127019](Letter) Nature Genet. 19: 223 only, 1998.

[127020]22672.

[127021]Friedl, W.; Uhlhaas, S.; Schulmann, K.; Stolte, M.; Loff, S.;
Back,

[127022]W.; Mangold, E.; Stern, M.; Knaebel, H. P.; Sutter, C.; Weber, R.

[127023]G.; Pistorius, S.; Burger, B.; Propping, P.: Juvenile polyposis:

[127024]massive gastric polyposis is more common in MADH4 mutation carriers

[127025]than in BMPR1A mutation carriers. Hum. Genet. 111:
108–111, 2002.

[127026]22673.

[127027]Howe, J. R.; Bair, J. L.; Sayed, M. G.; Anderson, M. E.;

Mitros,

[127028]F. A.; Petersen, G. M.; Velculescu, V. E.; Traverso, G.; Vogelstein,

[127029]B.: Germline mutations of the gene encoding bone morphogenetic protein

[127030]receptor 1A in juvenile polyposis. *Nature Genet.* 28: 184–187, 2001.

[127031]22674.

[127032]Houlston, R.; Bevan, S.; Williams, A.; Young, J.; Dunlop, M.; Rozen,

[127033]P.; Eng, C.; Markie, D.; Woodford–Richens, K.; Rodriguez–Bigas, M.

[127034]A.; Leggett, B.; Neale, K.; Phillips, R.; Sheridan, E.; Hodgson, S.;

[127035]Iwama, T.; Eccles, D.; Bodmer, W.; Tomlinson, I.: Mutations in DPC4

[127036](SMAD4) cause juvenile polyposis syndrome, but only account for a

[127037]minority of cases. *Hum. Molec. Genet.* 7: 1907–1912, 1998.

[127038]22675.

[127039]Howe, J. R.; Ringold, J. C.; Summers, R. W.; Mitros, F. A.;

Nishimura,

[127040]D. Y.; Stone, E. M.: A gene for familial juvenile polyposis
maps

[127041]to chromosome 18q21.1. Am. J. Hum. Genet. 62:
1129–1136, 1998.

[127042]22676.

[127043]Howe, J. R.; Roth, S.; Ringold, J. C.; Summers, R. W.; Jarvi-
nen,

[127044]H. J.; Sistonen, P.; Tomlinson, I. P. M.; Houlston, R. S.; Be-
van,

[127045]S.; Mitros, F. A.; Stone, E. M.; Aaltonen, L. A.: Mutations in
the

[127046]SMAD4/DPC4 gene in juvenile polyposis. Science 280:
1086–1088, 1998.

[127047]22677.

[127048]Kurose, K.; Araki, T.; Matsunaka, T.; Takada, Y.; Emi, M.:
Variant

[127049]manifestation of Cowden disease in Japan: hamartoma-
tous polyposis

[127050]of the digestive tract with mutation of the PTEN gene.
(Letter) Am.

[127051]J. Hum. Genet. 64: 308–310, 1999.

[127052]22678.

[127053]Nelen, M. R.; Padberg, G. W.; Peeters, E. A.; Lin, A. Y.; van

[127054]den Helm, B.; Frants, R. R.; Coulon, V.; Goldstein, A. M.;
van Reen,

[127055]M. M.; Easton, D. F.; Eeles, R. A.; Hodgson, S.; and 10 others:
ers:

[127056]Localization of the gene for Cowden disease to chromosome 10q22–23. Nature

[127057]Genet. 13: 114–116, 1996.

[127058]22679.

[127059]Olschwang, S.; Serova–Sinilnikova, O. M.; Lenoir, G. M.;
Thomas,

[127060]G.: PTEN germ–line mutations in juvenile polyposis coli.
(Letter) Nature

[127061]Genet. 18: 12–14, 1998.

[127062]22680.

[127063]Meulenbelt, I.; Williams, C. J.; Te Koppele, J. M.; Van De
Giessen,

[127064]G. C.; Slagboom, P. E.: Population haplotype analysis and
evolutionary

[127065]relations of the COI2A1 gene. Ann. Hum. Genet. 60:

189-199, 1996.

[127066]22681.

[127067]Berry, C.; Cree, J.; Mann, T.: Aarskog's syndrome. Arch.
Dis.

[127068]Child. 55: 706-710, 1980.

[127069]22682.

[127070]Escobar, V.; Weaver, D. D.: Aarskog syndrome: new find-
ings and

[127071]genetic analysis. J.A.M.A. 240: 2638-2641, 1978.

[127072]22683.

[127073]Hanley, W. B.; McKusick, V. A.; Barranco, F. T.: Osteochon-
dritis

[127074]dissecans and associated malformations in brothers: a re-
view of familial

[127075]aspects. J. Bone Joint Surg. 49A: 925-937, 1967.

[127076]22684.

[127077]O'Reilly, M. S.; Holmgren, L.; Shing, Y.; Chen, C.; Rosen-
thal,

[127078]R. A.; Moses, M.; Lane, W. S.; Cao, Y.; Sage, E. H.; Folkman,
J.:

[127079]Angiostatin: a novel angiogenesis inhibitor that mediates

the suppression

[127080]of metastases by a Lewis lung carcinoma. Cell 79:
315–328, 1994.

[127081]22685.

[127082]Waite, K. A.; Eng, C.: Protean PTEN: form and function.
Am.

[127083]J. Hum. Genet. 70: 829–844, 2002.

[127084]22686.

[127085]Armstrong, J. G.; Davies, D. R.; Guy, S. P.; Frayling, I. M.;
Evans,

[127086]D. G. R.: APC mutations in familial adenomatous polyposis
families

[127087]in the northwest of England. Hum. Mutat. 10: 376–380,
1997.

[127088]22687.

[127089]Ashton–Rickardt, P. G.; Dunlop, M. G.; Nakamura, Y.; Mor-
ris, R.

[127090]G.; Purdie, C. A.; Steel, C. M.; Evans, H. J.; Bird, C. C.; Wyl-
lie,

[127091]A. H.: High frequency of APC loss in sporadic colorectal
carcinoma

[127092]due to breaks clustered in 5q21–22. Oncogene 4:

1169-1174, 1989.

[127093]22688.

[127094]Asman, H. B.; Pierce, E. R.: Familial multiple polyposis: a statistical

[127095]study of a large Kentucky kindred. Cancer 25: 972-981, 1970.

[127096]22689.

[127097]Baker, R. H.; Heinemann, M.-H.; Miller, H. H.; DeCosse, J. J.:

[127098]Hyperpigmented lesions of the retinal pigment epithelium in familial

[127099]adenomatous polyposis. Am. J. Med. Genet. 31: 427-435, 1988.

[127100]22690.

[127101]Bapat, B.; Odze, R.; Mitri, A.; Berk, T.; Ward, M.; Gallinger,

[127102]S.: Identification of somatic APC gene mutations in peri-ampullary

[127103]adenomas in a patient with familial adenomatous polyposis (FAP). Hum.

[127104]Molec. Genet. 2: 1957-1959, 1993.

[127105]22691.

- [127106] Barber, J. C. K.; Ellis, K. H.; Bowles, L. V.; Delhanty, J. D. A.; Ede, R. F.; Male, B. M.; Eccles, D. M.: Adenomatous polyposis coli and a cytogenetic deletion of chromosome 5 resulting from a maternal intrachromosomal insertion. J. Med. Genet. 31: 312-316, 1994.
- [127109] 22692.
- [127110] Bell, B.; Mazzaferri, E. L.: Familial adenomatous polyposis (Gardner's syndrome) and thyroid carcinoma: a case report and review of the literature. Digest. Dis. Sci. 38: 185-190, 1993.
- [127111] 22693.
- [127112] Benecke, E.: Ueber Epitheliome auf Atheromen (Epidermoide) und Dermoidcysten der Haut. Frankf. Z. Pathol. 42: 502-515, 1931.
- [127113] 22694.
- [127114] Berk, T.; Cohen, Z.; Cullen, J. B.: Familial polyposis and the role of the preventive registry. Canad. Med. Assoc. J. 124:

1427-1428,

[127120]1981.

[127121]22695.

[127122]Beroud, C.; Soussi, T.: APC gene: database of germline
and somatic

[127123]mutations in human tumors and cell lines. Nucleic Acids
Res. 24:

[127124]121-124, 1996.

[127125]22696.

[127126]Bisgaard, M. L.; Fenger, K.; Bulow, S.; Niebuhr, E.; Mohr, J.

[127127]: Familial adenomatous polyposis (FAP): frequency, pene-
trance, and

[127128]mutation rate. Hum. Mutat. 3: 121-125, 1994.

[127129]22697.

[127130]Blair, N. P.; Trempe, C. L.: Hypertrophy of the retinal pig-
ment

[127131]epithelium associated with Gardner's syndrome. Am. J.
Ophthalm. 90:

[127132]661-667, 1980.

[127133]22698.

- [127134] Bodmer, W. F.; Bailey, C. J.; Bodmer, J.; Bussey, H. J. R.; Ellis,
lis,
[127135] A.; Gorman, P.; Lucibello, F. C.; Murday, V. A.; Rider, S. H.;
Scambler,
[127136] P.; Sheer, D.; Solomon, E.; Spurr, N. K.: Localization of the
gene
[127137] for familial adenomatous polyposis on chromosome 5.
Nature 328:
[127138] 614–616, 1987.
[127139] 22699.
[127140] Boland, C. R.; Sato, J.; Appelman, H. D.; Bresalier, R. S.;
Feinberg,
[127141] A. P.: Microallelotyping defines the sequence and tempo
of allelic
[127142] losses at tumour suppressor gene loci during colorectal
cancer progression. Nature
[127143] Med. 1: 902–909, 1995.
[127144] 22700.
[127145] Boolbol, S. K.; Dannenberg, A. J.; Chadburn, A.; Martucci,
C.;
[127146] Guo, X.-J.; Ramonetti, J. T.; Abreu-Goris, M.; Newmark, H.
L.; Lipkin,

[127147]M. L.; DeCosse, J. J.; Bertagnolli, M. M.: Cyclooxygenase-2 overexpression

[127148]and tumor formation are blocked by sulindac in a murine model of familial

[127149]adenomatous polyposis. Cancer Res. 56: 2556-2560, 1996.

[127150]22701.

[127151]Brensinger, J. D.; Laken, S. J.; Luce, M. C.; Powell, S. M.; Vance,

[127152]G. H.; Ahnen, D. J.; Petersen, G. M.; Hamilton, S. R.; Giardiello,

[127153]F. M.: Variable phenotype of familial adenomatous polyposis in pedigrees

[127154]with 3-prime mutation in the APC gene. Gut 43: 548-552, 1998.

[127155]22702.

[127156]Bull, M. J.; Ellis, F. D.; Sato, S.; Weaver, D. D.: Hypertrophy

[127157]of retinal pigment epithelium in Gardner syndrome.

(Abstract) Proc.

[127158]Greenwood Genet. Center 4: 136 only, 1985.

[127159]22703.

[127160]Bunyan, D. J.; Shea-Simonds, J.; Reck, A. C.; Finnis, D.; Eccles,

[127161]D. M.: Genotype-phenotype correlations of new causative APC gene

[127162]mutations in patients with familial adenomatous polyposis. J. Med.

[127163]Genet. 32: 728-731, 1995.

[127164]22704.

[127165]Awata, H.; Endo, F.; Matsuda, I.: Structure of the human 4-hydroxyphenylpyruvic

[127166]acid dioxygenase gene (HPD). Genomics 23: 534-539, 1994.

[127167]22705.

[127168]Nagase, T.; Seki, N.; Tanaka, A.; Ishikawa, K.; Nomura, N.: Prediction

[127169]of the coding sequences of unidentified human genes. IV. The coding

[127170]sequences of 40 new genes (KIAA0121-KIAA0160) deduced by analysis

[127171]of cDNA clones from human cell line KG-1. DNA Res. 2: 167-174, 1995.

[127172]22706.

[127173]Parrish, J. E.; Ciccodicola, A.; Wehnert, M.; Cox, G. F.;
Chen,
[127174]E.; Nelson, D. L.: A muscle-specific DNase I-like gene in
human Xq28. Hum.
[127175]Molec. Genet. 4: 1557-1564, 1995.
[127176]22707.
[127177]Pergolizzi, R.; Appierto, V.; Bosetti, A.; DeBellis, G. L.;
Rovida,
[127178]E.; Biunno, I.: Cloning of a gene encoding a DNase I-like
endonuclease
[127179]in the human Xq28 region. Gene 168: 267-270, 1996.
[127180]22708.
[127181]Klink, A.; Schiebel, K.; Winkelmann, M.; Rao, E.; Hors-
themke, B.;
[127182]Ludecke, H.-J.; Claussen, U.; Scherer, G.; Rappold, G.: The
human
[127183]protein kinase gene PKX1 on Xp22.3 displays Xp/Yp ho-
mology and is
[127184]a site of chromosomal instability. Hum. Molec. Genet. 4:
869-878,
[127185]1995.

[127186]22709.

[127187]Schiebel, K.; Mertz, A.; Winkelmann, B.; Glaser, B.;
Schempp, W.;

[127188]Rappold, G.: FISH localization of the human Y-homolog of
protein

[127189]kinase PRKX (PRKY) to Yp11.2 and two pseudogenes to
15q26 and Xq12-q13. Cytogenet.

[127190]Cell Genet. 76: 49-52, 1997.

[127191]22710.

[127192]Brown, C. J.; Carrel, L.; Willard, H. F.: Expression of genes

[127193]from the human active and inactive X chromosomes. Am.
J. Hum. Genet. 60:

[127194]1333-1343, 1997.

[127195]22711.

[127196]Dong, B.; Horowitz, D. S.; Kobayashi, R.; Krainer, A. R.:
Purification

[127197]and cDNA cloning of HeLa cell p54(nrb), a nuclear protein
with two

[127198]RNA recognition motifs and extensive homology to human
splicing factor

[127199]PSF and Drosophila NONA/BJ6. Nucleic Acids Res. 21:

4085–4092, 1993.

[127200]22712.

[127201]Nelson, D. L.; Ballabio, A.; Cremers, F.; Monaco, A. P.;
Schlessinger,

[127202]D.: Report of the sixth international workshop on X chromosome mapping

[127203]1995. Cytogenet. Cell Genet. 71: 308–342, 1995.

[127204]22713.

[127205]Janssens, R.; Boeynaems, J.-M.; Godart, M.; Communi, D.:
Cloning

[127206]of a human heptahelical receptor closely related to the
P2Y(5) receptor. Biochem.

[127207]Biophys. Res. Commun. 236: 106–112, 1997.

[127208]22714.

[127209]O'Dowd, B. F.; Nguyen, T.; Jung, B. P.; Marchese, A.;
Cheng, R.;

[127210]Heng, H. H. Q.; Kolakowski, L. F., Jr.; Lynch, K. R.; George,
S. R.

[127211]: Cloning and chromosomal mapping of four putative
novel human G-protein-coupled

[127212]receptor genes. Gene 187: 75–81, 1997.

[127213]22715.

[127214]Bienvenu, T.; Der-Sarkissian, H.; Billuart, P.; Tissot, M.;
Des

[127215]Portes, V.; Bruls, T.; Chabrolle, J. P.; Chauveau, P.; Cherry,
M.;

[127216]Kahn, A.; Cohen, D.; Beldjord, C.; Chelly, J.; Cherif, D.:
Mapping

[127217]of the X-breakpoint involved in a balanced X;12 translo-
cation in a

[127218]female with mild mental retardation. Europ. J. Hum. Genet.
5: 105-109,

[127219]1997.

[127220]22716.

[127221]Billuart, P.; Bienvenu, T.; Ronce, N.; des Portes, V.; Vinet,
M.

[127222]C.; Zemni, R.; Crollius, H. R.; Carrie, A.; Fauchereau, F.;
Cherry,

[127223]M.; Briault, S.; Hamel, B.; Fryns, J.-P.; Beldjord, C.; Kahn,
A.;

[127224]Moraine, C.; Chelly, J.: Oligophrenin-1 encodes a rhoGAP
protein

[127225]involved in X-linked mental retardation. Nature 392:

823–826, 1998.

[127226]22717.

[127227]Kozak, M.: Point mutations define a sequence flanking the
AUG

[127228]initiator codon that modulates translation by eukaryotic
ribosomes. Cell 44:

[127229]283–292, 1986.

[127230]22718.

[127231]Tentler, D.; Gustavsson, P.; Leisti, J.; Schueler, M.; Chelly,

[127232]J.; Timonen, E.; Anneren, G.; Willard, H. F.; Dahl, N.: Dele-
tion

[127233]including the oligophrenin-1 gene associated with en-
larged cerebral

[127234]ventricles, cerebellar hypoplasia, seizures and ataxia. Eu-
rop. J.

[127235]Hum. Genet. 7: 541–548, 1999.

[127236]22719.

[127237]Greenfield, A.; Carrel, L.; Pennisi, D.; Philippe, C.; Quaderi,

[127238]N.; Siggers, P.; Steiner, K.; Tam, P. P. L.; Monaco, A. P.;
Willard,

[127239]H. F.; Koopman, P.: The UTX gene escapes X inactivation

in mice and

[127240]humans. Hum. Molec. Genet. 7: 737–742, 1998.

[127241]22720.

[127242]Greenfield, A. J.; Scott, D.; Pennisi, D.; Ehrmann, I.; Ellis,

[127243]P.; Cooper, L.; Simpson, E.; Koopman, P.: An H-YDb epitope is encoded

[127244]by a novel mouse Y chromosome gene. Nature Genet. 14: 474–478, 1996.

[127245]22721.

[127246]Lin, C.-S.; Aebersold, R. H.; Leavitt, J.: Correction of the N-terminal

[127247]sequences of the human plastin isoforms by using anchored polymerase

[127248]chain reaction: identification of a potential calcium-binding domain. Molec.

[127249]Cell. Biol. 10: 1818–1821, 1990.

[127250]22722.

[127251]Fukuda, M. N.; Sato, T.; Nakayama, J.; Klier, G.; Mikami, M.; Aoki,

[127252]D.; Nozawa, S.: Trophinin and tastin, a novel cell adhesion molecule

[127253]complex with potential involvement in embryo implantation. Genes

[127254]Dev. 9: 1199–1210, 1995.

[127255]22723.

[127256]Pack, S. D.; Tanigami, A.; Ledbetter, D. H.; Sato, T.; Fukuda,

[127257]M. N.: Assignment of trophoblast/endometrial epithelium cell adhesion

[127258]molecule trophinin gene TRO to human chromosome bands Xp11.22–p11.21

[127259]by in situ hybridization. Cytogenet. Cell Genet. 79: 123–124, 1997.

[127260]22724.

[127261]Suzuki, N.; Nakayama, J.; Shih, I. M.; Aoki, D.; Nozawa, S.; Fukuda,

[127262]M. N.: Expression of trophinin, tastin, and bystin by trophoblast

[127263]and endometrial cells in human placenta. Biol. Reprod. 60: 621–627,

[127264]1999.

[127265]22725.

[127266]Yan, M.; Wang, L.-C.; Hymowitz, S. G.; Schilbach, S.; Lee, J.;

[127267]Goddard, A.; de Vos, A. M.; Gao, W.-Q.; Dixit, V. M.: Two-amino acid

[127268]molecular switch in an epithelial morphogen that regulates binding

[127269]to two distinct receptors. Science 290: 523–527, 2000.

[127270]22726.

[127271]Sana, T. R.; Debets, R.; Timans, J. C.; Bazan, J. F.; Kastelein,

[127272]R. A.: Computational identification, cloning, and characterization

[127273]of IL1R9, a novel interleukin-1 receptor-like gene encoded over an

[127274]unusually large interval of human chromosome Xq22.2–q22.3. Genomics 69:

[127275]252–262, 2000.

[127276]22727.

[127277]Bech-Hansen, N. T.; Naylor, M. J.; Maybaum, T. A.; Sparkes, R.

[127278]L.; Koop, B.; Birch, D. G.; Bergen, A. A. B.; Prinsen, C. F. M.; Polomeno,

[127279]R. C.; Gal, A.; Drack, A. V.; Musarella, M. A.; Jacobson, S. G.; Young,

[127280]R. S. L.; Weleber, R. G.: Mutations in NYX, encoding the leucine-rich

[127281]proteoglycan nyctalopin, cause X-linked complete congenital stationary

[127282]night blindness. Nature Genet. 26: 319–323, 2000.

[127283]22728.

[127284]Pusch, C. M.; Zeitz, C.; Brandau, O.; Pesch, K.; Achatz, H.; Feil,

[127285]S.; Scharfe, C.; Maurer, J.; Jacobi, F. K.; Pinckers, A.; Andreasson,

[127286]S.; Hardcastle, A.; Wissinger, B.; Berger, W.; Meindl, A.: The complete

[127287]form of X-linked congenital stationary night blindness is caused by

[127288]mutations in a gene encoding a leucine-rich repeat protein. Nature

[127289]Genet. 26: 324–327, 2000.

[127290]22729.

[127291]Isbrandt, D.; Leicher, T.; Waldschutz, R.; Zhu, X.; Luhmann, U.;

[127292]Michel, U.; Sauter, K.; Pongs, O.: Gene structures and expression

[127293]profiles of three human KCND (Kv4) potassium channels mediating A-type

[127294]currents I(to) and I(sa). Genomics 64: 144–154, 2000.

[127295]22730.

[127296]Cao, Z.; Henzel, W.; Gao, X.: IRAK: a kinase associated with the

[127297]interleukin-1 receptor. Science 271: 1128–1131, 1996.

[127298]22731.

[127299]Kanakaraj, P.; Ngo, K.; Wu, Y.; Angulo, A.; Ghazal, P.; Harris,

[127300]C.A.; Siekierka, J. J.; Peterson, P. A.; Fung-Leung, W. P.: Defective

[127301]interleukin (IL)-18-mediated natural killer and T helper cell type

[127302]1 responses in IL-1 receptor-associated kinase (IRAK)-deficient mice. J.

[127303]Exp. Med. 189: 1129–1138, 1999.

[127304]22732.

[127305]Thomas, J. A.; Allen, J. L.; Tsen, M.; Dubnicoff, T.; Danao,

J.;

[127306]Liao, X. C.; Cao, Z.; Wasserman, S. A.: Impaired cytokine signaling

[127307]in mice lacking the IL-1 receptor-associated kinase. J. Immun. 163:

[127308]978-984, 1999.

[127309]22733.

[127310]van Vliet, J.; Turner, J.; Crossley, M.: Human Kruppel-like factor

[127311]8: a CACCC-box binding protein that associates with CtBP and represses

[127312]transcription. Nucleic Acids Res. 28: 1955-1962, 2000.

[127313]22734.

[127314]Lurquin, C.; De Smet, C.; Brasseur, F.; Muscatelli, F.; Martelange,

[127315]V.; De Plaen, E.; Brasseur, R.; Monaco, A. P.; Boon, T.: Two members

[127316]of the human MAGEB gene family located in Xp21.3 are expressed in

[127317]tumors of various histological origins. Genomics 46: 397-408, 1997.

[127318]22735.

[127319]Aguilar, M. J.; O'Brien, J. S.; Taber, P.: The syndrome of familial

[127320]leukodystrophy, adrenal insufficiency and cutaneous melanosis. In:

[127321]Aronson, S. M.; Volk, B. W.: Inborn Disorders of Sphingolipid Metabolism.

[127322]Oxford: Pergamon Press (pub.) 1967. Pp. 149–166.

[127323]22736.

[127324]Alpern, M.; Sack, G. H., Jr.; Krantz, D. H.; Jenness, J.; Zhang,

[127325]H.; Moser, H. W.: Chromosomal rearrangement segregating with adrenoleukodystrophy:

[127326]associated changes in color vision. Proc. Nat. Acad. Sci. 90: 9494–9498,

[127327]1993.

[127328]22737.

[127329]ASHG statement: Professional disclosure of familial genetic

[127330]information: the American Society of Human Genetics Social Issues

[127331]Subcommittee on Familial Disclosure. Am. J. Hum. Genet. 62: 474–483,

[127332]1998.

[127333]22738.

[127334]Aubourg, P.; Adamsbaum, C.; Lavallard-Rousseau, M.-C.;
Rocchiccioli,

[127335]F.; Cartier, N.; Jambaque, I.; Jakobezak, C.; Lemaitre, A.;
Boureau,

[127336]F.; Wolf, C.; Bougneres, P.-F.: A two-year trial of oleic and
erucic

[127337]acids ('Lorenzo's oil') as treatment for adrenomyeloneu-
ropathy. New

[127338]Eng. J. Med. 329: 745-752, 1993.

[127339]22739.

[127340]Aubourg, P.; Blanche, S.; Jambaque, I.; Rocchiccioli, F.;
Kalifa,

[127341]G.; Naud-Saudreau, C.; Rolland, M.-O.; Debre, M.; Chaus-
sain, J.-L.;

[127342]Griscelli, C.; Fischer, A.; Bougneres, P.-F.: Reversal of early
neurologic

[127343]and neuroradiologic manifestations of X-linked
adrenoleukodystrophy

[127344]by bone marrow transplantation. New Eng. J. Med. 322:
1860-1866,

[127345]1990.

[127346]22740.

[127347]Aubourg, P.; Feil, R.; Guidoux, S.; Kaplan, J.-C.; Moser, H.;
Kahn,

[127348]A.; Mandel, J.-L.: The red-green visual pigment gene re-
gion in adrenoleukodystrophy. Am.

[127349]J. Hum. Genet. 46: 459-469, 1990.

[127350]22741.

[127351]Aubourg, P.; Mosser, J.; Douar, A. M.; Sarde, C. O.; Lopez,
J.;

[127352]Mandel, J. L.: Adrenoleukodystrophy gene: unexpected
homology to

[127353]a protein involved in peroxisome biogenesis. Biochimie
75: 293-302,

[127354]1993.

[127355]22742.

[127356]Aubourg, P. R.; Sack, G. H., Jr.; Meyers, D. A.; Lease, J. J.;

[127357]Moser, H. W.: Linkage of adrenoleukodystrophy to a poly-
morphic DNA

[127358]probe. Ann. Neurol. 21: 349-352, 1987.

[127359]22743.

[127360]Aubourg, P. R.; Sack, G. H., Jr.; Moser, H.: Frequent alterations

[127361]of visual pigment genes in adrenoleukodystrophy. Am. J. Hum. Genet. 42:

[127362]408-413, 1988.

[127363]22744.

[127364]Barcelo, A.; Giros, M.; Martinez-Bermejo, A.; Sarde, C.-O.; Coll,

[127365]M. J.; Moser, J.; Mandel, J. L.; Estivill, X.; Pampols, T.: A frame

[127366]shift mutation in the candidate gene of X-linked adrenoleukodystrophy. VI.

[127367]International Congress of Inborn Errors of Metabolism. Milano, Italy May-27-31,

[127368]1994. Pp. 228 only.

[127369]22745.

[127370]Barcelo, A.; Giros, M.; Sarde, C. O.; Martinez-Bermejo, A.; Mandel,

[127371]J. L.; Pampols, T.; Estivill, X.: Identification of a new frameshift

[127372]mutation (1801delAG) in the ALD gene. Hum. Molec.

Genet. 3: 1889–1890,

[127373]1994.

[127374]22746.

[127375]Berg, K. A.; Beaty, T. H.; Raven, M. B.; Moser, A. B.; Moser,

[127376]H. W.: X-linked adrenoleukodystrophy in a 362 member
kindred. (Abstract) Am.

[127377]J. Hum. Genet. 45 (suppl.): A39, 1989.

[127378]22747.

[127379]Berger, J.; Molzer, B.; Fae, I.; Bernheimer, H.: X-linked
adrenoleukodystrophy

[127380](ALD): a novel mutation of the ALD gene in 6 members of
a family presenting

[127381]with 5 different phenotypes. Biochem. Biophys. Res. Com-
mun. 205:

[127382]1638–1643, 1994.

[127383]22748.

[127384]Bezman, L.; Moser, A. B.; Raymond, G. V.; Rinaldo, P.;
Watkins,

[127385]P. A.; Smith, K. D.; Kass, N. E.; Moser, H. W.:
Adrenoleukodystrophy:

[127386] incidence, new mutation rate, and results of extended family screening. Ann.

[127387] Neurol. 49: 512–517, 2001.

[127388] 22749.

[127389] Bezman, L.; Moser, H. W.: Incidence of X-linked adrenoleukodystrophy

[127390] and the relative frequency of its phenotypes. (Editorial) Am. J.

[127391]Med. Genet. 76: 415–419, 1998.

[127392]22750.

[127393]Blaw, M. E.: Melanodermic type leucodystrophy
(adrenoleukodystrophy).In:

[127394]Vinken, P. J.; Bruyn, G. W. (eds.): Handbook of Clinical
Neurology.

[127395]Vol. 10. Amsterdam: North Holland 1970. Pp. 128–133.

[127396]22751.

[127397]Boue, J.; Oberle, I.; Heilig, R.; Mandel, J. L.; Moser, A.;
Moser,

[127398]H.; Larsen, J. W., Jr.; Dumez, Y.; Boue, A.: First trimester
prenatal

[127399]diagnosis of adrenoleukodystrophy by determination of
very long chain

[127400]fatty acid levels and by linkage analysis to a DNA probe.
Hum. Genet. 69:

[127401]272–274, 1985.

[127402]22752.

[127403]Brady, R. O.: Inherited metabolic diseases of the nervous
system. Science 193:

[127404]733–739, 1976.

[127405]22753.

[127406]Braun, A.; Ambach, H.; Kammerer, S.; Rolinski, B.; Stockler, S.;

[127407]Rabl, W.; Gartner, J.; Zierz, S.; Roscher, A. A.: Mutations in the

[127408]gene for X-linked adrenoleukodystrophy in patients with different

[127409]clinical phenotypes. Am. J. Hum. Genet. 56: 854-861, 1995.

[127410]22754.

[127411]Jonsson, J. J.; Renieri, A.; Gallagher, P. G.; Kashtan, C. E.;

[127412]Cherniske, E. M.; Bruttini, M.; Piccini, M.; Vitelli, F.; Bal-labio,

[127413]A.; Pober, B. R.: Alport syndrome, mental retardation, midface hypoplasia,

[127414]and elliptocytosis: a new X linked contiguous gene deletion syndrome? J.

[127415]Med. Genet. 35: 273-278, 1998.

[127416]22755.

[127417]Kang, M. J.; Fujino, T.; Sasano, H.; Minekura, H.; Yabuki, N.;

[127418]Nagura, H.; Iijima, H.; Yamamoto, T. T.: A novel arachido-
nate-preferring

[127419]acyl-CoA synthetase is present in steroidogenic cells of
the rat adrenal,

[127420]ovary, and testis. Proc. Nat. Acad. Sci. 94: 2880-2884,
1997.

[127421]22756.

[127422]Meloni, I.; Muscettola, M.; Raynaud, M.; Longo, I.; Bruttini,
M.;

[127423]Moizard, M.-P.; Gomot, M.; Chelly, J.; des Portes, V.;
Fryns, J.-P.;

[127424]Ropers, H.-H.; Magi, B.; Bellan, C.; Volpi, N.; Yntema, H.
G.; Lewis,

[127425]S. E.; Schaffer, J. E.; Renieri, A.: FAFL4, encoding fatty
acid-CoA

[127426]ligase 4, is mutated in nonspecific X-linked mental retar-
dation. Nature

[127427]Genet. 30: 436-440, 2002.

[127428]22757.

[127429]Minekura, H.; Kang, M.-J.; Inagaki, Y.; Cho, Y.-Y.; Suzuki,
H.;

[127430]Fujino, T.; Yamamoto, T. T.: Exon/intron organization and

transcription

[127431]units of the human acyl-CoA synthetase 4 gene. Biochem. Biophys.

[127432]Res. Commun. 286: 80-86, 2001.

[127433]22758.

[127434]Piccini, M.; Vitelli, F.; Bruttini, M.; Pober, B. R.; Jonsson,

[127435]J. J.; Villanova, M.; Zollo, M.; Borsani, G.; Ballabio, A.; Renieri,

[127436]A.: FACL4, a new gene encoding long-chain acyl-CoA synthetase 4,

[127437]is deleted in a family with Alport syndrome, elliptocytosis, and mental

[127438]retardation. Genomics 47: 350-358, 1998.

[127439]22759.

[127440]Vitelli, F.; Meloni, I.; Fineschi, S.; Favara, F.; Tiziana Storlazzi,

[127441]C.; Rocchi, M.; Renieri, A.: Identification and characterization

[127442]of mouse orthologs of the AMMECR1 and FACL4 genes deleted in AMME

[127443]syndrome: orthology of Xq22.3 and MmuXF1-F3. Cyto-

genet. Cell Genet. 88:

[127444]259–263, 2000.

[127445]22760.

[127446]Lahn, B. T.; Page, D. C.: Functional coherence of the human Y

[127447]chromosome. Science 278: 675–680, 1997.

[127448]22761.

[127449]Chung, J.; Lee, S.–G.; Song, K.: Identification of a human homolog

[127450]of a putative RNA helicase gene (mDEAD3) expressed in mouse erythroid

[127451]cells. Korean J. Biochem. 27: 193–197, 1995.

[127452]22762.

[127453]Park, S. H.; Lee, S.–G.; Kim, Y.; Song, K.: Assignment of a human

[127454]putative RNA helicase gene, DDX3, to human X chromosome bands p11.3–p11.23. Cytogenet.

[127455]Cell Genet. 81: 178–179, 1998.

[127456]22763.

[127457]Gasper, N. J.; Kinzy, T. G.; Scherer, B. J.; Humbelin, M.;

Hershey,

[127458]J. W. B.; Merrick, W. C.: Translation initiation factor eIF-2:
cloning

[127459]and expression of the human cDNA encoding the gamma-
subunit. J. Biol.

[127460]Chem. 269: 3415-3422, 1994.

[127461]22764.

[127462]Ehrmann, I. E.; Ellis, P. S.; Mazeyrat, S.; Duthie, S.; Brock-
dorff,

[127463]N.; Mattei, M. G.; Gavin, M. A.; Affara, N. A.; Brown, G. M.;
Simpson,

[127464]E.; Mitchell, M. J.; Scott, D. M.: Characterization of genes
encoding

[127465]translation initiation factor eIF-2-gamma in mouse and
human: sex

[127466]chromosome localization, escape from X-inactivation and
evolution. Hum.

[127467]Molec. Genet. 7: 1725-1737, 1998.

[127468]22765.

[127469]Lee, S. M. Y.; Tsui, S. K. W.; Chan, K. K.; Garcia-Barcelo,
M.;

[127470]Waye, M. M. Y.; Fung, K. P.; Liew, C. C.; Lee, C. Y.: Chro-

mosomal

[127471]mapping, tissue distribution and cDNA sequence of four-
and-a-half

[127472]LIM domain protein 1 (FHL1). Gene 216: 163–170, 1998.

[127473]22766.

[127474]Morgan, M. J.; Madgwick, A. J.; Charleston, B.; Pell, J. M.;
Loughna,

[127475]P. T.: The developmental regulation of a novel muscle
LIM-protein. Biochem.

[127476]Biophys. Res. Commun. 212: 840–846, 1995.

[127477]22767.

[127478]Morgan, M. J.; Madgwick, A. J. A.: Slim defines a novel
family

[127479]of LIM-proteins expressed in skeletal muscle. Biochem.
Biophys. Res.

[127480]Commun. 225: 632–638, 1996.

[127481]22768.

[127482]Blanchet, P.; Wellemeyer, M. L.; Burton, G. V.: Retinitis
pigmentosa

[127483]following cytotoxic chemotherapy in Usher's syndrome.
Am. J. Med.

[127484]Sci. 303: 319–320, 1992.

[127485]22769.

[127486]Connor, W. E.; Weleber, R. G.; DeFrancesco, C.; Lin, D. S.;
Wolf,

[127487]D. P.: Sperm abnormalities in retinitis pigmentosa. Invest.
Ophthalm.

[127488]Vis. Sci. 38: 2619–2628, 1997.

[127489]22770.

[127490]Barnard, P. J.; Derry, J. M. J.; Ryder–Cook, A. S.; Barnard, E.

[127491]A.: Localization of the GABA(A) receptor alpha–3 subunit
gene on

[127492]the mouse X chromosome. (Abstract) Cytogenet. Cell
Genet. 51: 958

[127493]only, 1989.

[127494]22771.

[127495]Bell, M. V.; Bloomfield, J.; McKinley, M.; Patterson, M. N.;
Darlison,

[127496]M. G.; Barnard, E. A.; Davies, K. E.: Physical linkage of a
GABA–A

[127497]receptor subunit gene to the DXS374 locus in human
Xq28. Am. J. Hum.

[127498]Genet. 45: 883–888, 1989.

[127499]22772.

[127500]Buckle, V. J.; Fujita, N.; Bateson, A. N.; Darlison, M. G.;
Barnard,

[127501]E. A.: Localization of the human GABA-A3 receptor sub-
unit gene to

[127502]Xq28: a candidate gene for X-linked manic depression.
(Abstract) Cytogenet.

[127503]Cell Genet. 51: 972 only, 1989.

[127504]22773.

[127505]Buckle, V. J.; Fujita, N.; Ryder-Cook, A. S.; Derry, J. M. J.;

[127506]Barnard, P. J.; Lebo, R. V.; Schofield, P. R.; Seeburg, P. H.;
Bateson,

[127507]A. N.; Darlison, M. G.; Barnard, E. A.: Chromosomal local-
ization

[127508]of GABA-A receptor subunit genes: relationship to human
genetic disease. Neuron 3:

[127509]647–654, 1989.

[127510]22774.

[127511]Derry, J. M. J.; Barnard, P. J.: Mapping of the glycine recep-
tor

[127512]alpha-2-subunit gene and the GABA-A alpha-3-subunit gene on the mouse

[127513]X chromosome. Genomics 10: 593-597, 1991.

[127514]22775.

[127515]Buckle, V. J.; Edwards, J. H.; Evans, E. P.; Jonasson, J. A.; Lyon,

[127516]M. F.; Peters, J.; Searle, A. G.: Comparative maps of human and mouse

[127517]X chromosomes. (Abstract) Cytogenet. Cell Genet. 40: 594-595, 1985.

[127518]22776.

[127519]Mansour, I.; Delague, V.; Cazeneuve, C.; Dode, C.; Chouery, E.;

[127520]Pecheux, C.; Medlej-Hashim, M.; Salem, N.; El Zein, L.; Levan-Petit,

[127521]J.; Lefranc, G.; Goossens, M.; Delpech, M.; Amselem, S.; Loiselet,

[127522]J.; Grateau, G.; Megarbane, A.; Naman, R.: Familial Mediterranean

[127523]fever in Lebanon: mutation spectrum, evidence for cases in Maronites,

[127524]Greek Orthodoxes, Greek Catholics, Syriacs and Chiites

and for an

[127525]association between amyloidosis and M694V and M694I mutations. Europ.

[127526]J. Hum. Genet. 9: 51–55, 2001.

[127527]22777.

[127528]Matzner, Y.; Brzezinski, A.: C5a-inhibitor deficiency in peritoneal

[127529]fluids from patients with familial Mediterranean fever. New Eng.

[127530]J. Med. 311: 287–290, 1984.

[127531]22778.

[127532]Matzner, Y.; Partridge, R. E. H.; Levy, M.; Babior, B. M.: Diminished

[127533]activity of a chemotactic inhibitor in synovial fluids from patients

[127534]with familial Mediterranean fever. Blood 63: 629–633, 1984.

[127535]22779.

[127536]Medlej-Hashim, M.; Petit, I.; Adib, S.; Chouery, E.; Salem, N.;

[127537]Delague, V.; Rawashdeh, M.; Mansour, I.; Lefranc, G.; Na-

man, R.; Loiselet,
 [127538]J.; Lecron, J.-C.; Serre, J.-L.; Megarbane, A.: Familial
 Mediterranean
 [127539]fever: association of elevated IgD plasma levels with spe-
 cific MEFV
 [127540]mutations. *Europ. J. Hum. Genet.* 9: 849–854, 2001.
 [127541]22780.
 [127542]Medlej-Hashim, M.; Salem, N.; Chouery, E.; Rawashdeh,
 M.; Delague,
 [127543]V.; Haffar, M.; Mansour, I.; Naman, R.; Lefranc, G.; Loise-
 let, J.;
 [127544]Megarbane, A.: Familial Mediterranean fever: the potential
 for misdiagnosis
 [127545]of E148V using the E148Q usual RFLP detection method.
 (Letter) *Clin.*
 [127546]*Genet.* 61: 71–73, 2002.
 [127547]22781.
 [127548]Meyerhoff, J.: Personal Communication. Salt Lake City,
 Utah
 [127549]and Baltimore, Md. 1978.
 [127550]22782.

[127551]Meyerhoff, J.: Familial Mediterranean fever: report of a large

[127552]family, review of the literature, and discussion of the frequency

[127553]of amyloidosis. *Medicine* 59: 66–77, 1980.

[127554]22783.

[127555]Mollaret, P.: La meningite endothelio-leucocytaire multi-recurrente

[127556]benign: syndrome nouveau du maladie nouvelle? *Bull. Soc. Med. Hop.*

[127557]Paris 60: 121–122, 1944.

[127558]22784.

[127559]Ozdemir, A. I.; Sokmen, C.: Familial Mediterranean fever among

[127560]the Turkish people. *Am. J. Gastroent.* 51: 311–316, 1969.

[127561]22785.

[127562]Ozyilkan, E.; Simsek, H.; Telatar, H.: Absence of asthma in patients

[127563]with familial Mediterranean fever. *Israel J. Med. Sci.* 30: 237–238,

[127564]1994.

[127565]22786.

[127566]Papin, S.; Duquesnoy, P.; Cazeneuve, C.; Pantel, J.;
Coppey-Moisan,

[127567]M.; Dargemont, C.; Amselem, S.: Alternative splicing at the
MEFV

[127568]locus involved in familial Mediterranean fever regulates
translocation

[127569]of the marenostriin/pyrin protein to the nucleus. Hum.
Molec. Genet. 9:

[127570]3001–3009, 2000.

[127571]22787.

[127572]Pras, E.; Aksentijevich, I.; Gruberg, L.; Balow, J. E., Jr.;
Prosen,

[127573]L.; Dean, M.; Steinberg, A. D.; Pras, M.; Kastner, D. L.:
Mapping

[127574]of a gene causing familial Mediterranean fever to the short
arm of

[127575]chromosome 16. New Eng. J. Med. 326: 1509–1513,
1992.

[127576]22788.

[127577]Pras, E.; Aksentijevich, I.; Gruberg, L.; Shen, Y.; Prosen, L.;

[127578]Balow, J. E., Jr.; Richards, R. I.; Pras, M.; Kastner, D. L.: Lo-
cus

[127579]homogeneity in familial Mediterranean fever. (Abstract)
Am. J. Hum.

[127580]Genet. 51 (suppl.): A371, 1992.

[127581]22789.

[127582]Pras, E.; Aksentijevich, I.; Levy, E.; Gruberg, L.; Prosen, L.;

[127583]Dean, M.; Pras, M.; Kastner, D. L.: The gene causing famil-
ial Mediterranean

[127584]fever maps to the short arm of chromosome 16 in Druze
and Moslem Arab

[127585]families. Hum. Genet. 94: 576-577, 1994.

[127586]22790.

[127587]Pras, E.; Livneh, A.; Balow, J. E., Jr.; Pras, E.; Kastner, D.

[127588]L.; Pras, M.; Langevitz, P.: Clinical differences between
North African

[127589]and Iraqi Jews with familial Mediterranean fever. Am. J.
Med. Genet. 75:

[127590]216-219, 1998.

[127591]22791.

[127592]Pras, M.; Bronshpigel, N.; Zemer, D.; Gafni, J.: Variable incidence
of amyloidosis in familial Mediterranean fever among different ethnic
groups. Johns Hopkins Med. J. 150: 22–26, 1982.
22792.

[127596]Ravid, M.; Robson, M.; Kedar, I.: Prolonged colchicine
treatment
in four patients with amyloidosis. Ann. Intern. Med. 87:
568–570,
1977.

[127599]22793.

[127600]Rawashdeh, M. O.; Majeed, H. A.: Familial Mediterranean
fever
in Arab children: the high prevalence and gene frequency.
Europ.
J. Pediat. 155: 540–544, 1996.

[127603]22794.

[127604]Reimann, H. A.; Moadie, J.; Semerdijian, S.; Sahyoun, P. F.:
Periodic peritonitis--heredity and pathology: report of
seventy-two

[127606]cases. J.A.M.A. 154: 1254–1259, 1954.

[127607]22795.

[127608]Rogers, D. B.; Shohat, M.; Petersen, G. M.; Bickal, J.; Congleton,

[127609]J.; Schwabe, A. D.; Rotter, J. I.: Familial Mediterranean fever in

[127610]Armenians: autosomal recessive inheritance with high gene frequency. Am.

[127611]J. Med. Genet. 34: 168–172, 1989.

[127612]22796.

[127613]Rubinger, D.; Friedlaender, M. M.; Popovtzer, M. M.: Amelioration

[127614]of familial Mediterranean fever during hemodialysis. New Eng. J.

[127615]Med. 301: 142–144, 1979.

[127616]22797.

[127617]Saatci, U.; Ozen, S.; Ozdemir, S.; Bakkaloglu, A.; Besbas, N.;

[127618]Topaloglu, R.; Arslan, S.: Familial Mediterranean fever in children:

[127619]report of a large series and discussion of the risk and

prognostic

[127620]factors of amyloidosis. *Europ. J. Pediat.* 156: 619–623, 1997.

[127621]22798.

[127622]Sack, G. H., Jr.; Talbot, C. C., Jr.; McCarthy, B. G.; Harris,

[127623]E. L.; Kastner, D.; Gruberg, L.; Pras, M.: Exclusion of linkage between

[127624]familial Mediterranean fever and the human serum amyloid A (SAA) gene

[127625]cluster. *Hum. Genet.* 87: 506–508, 1991.

[127626]22799.

[127627]Schaner, P.; Richards, N.; Wadhwa, A.; Aksentijevich, I.; Kastner,

[127628]D.; Tucker, P.; Gumucio, D.: Episodic evolution of pyrin in primates:

[127629]human mutations recapitulate ancestral amino acid states. *Nature*

[127630]*Genet.* 27: 318–321, 2001.

[127631]22800.

[127632]Schlesinger, M.; Ilfeld, D. N.; Zamir, R.; Brautbar, C.: Familial

- [127633]Mediterranean fever: no linkage with HLA. Tissue Antigens
24: 65-66,
[127634]1984.
[127635]22801.
- [127636]Schwabe, A. D.; Lehman, T. J. A.: C5a-inhibitor deficiency-
-a
[127637]role in familial Mediterranean fever? (Editorial) New Eng. J.
Med. 311:
[127638]325-326, 1984.
[127639]22802.
- [127640]Schwabe, A. D.; Monroe, J. B.: Meningitis in familial
Mediterranean
[127641]fever. Am. J. Med. 85: 715-717, 1988.
[127642]22803.
- [127643]Schwabe, A. D.; Nishizawa, A.: Recurrent polyserositis
(familial
[127644]Mediterranean fever) in a Japanese. Jpn. J. Med. 26:
370-372, 1987.
[127645]22804.
- [127646]Schwabe, A. D.; Peters, R. S.: Familial Mediterranean fever
in

[127647]Armenians: analysis of 100 cases. *Medicine* 53: 453-462, 1974.

[127648]22805.

[127649]Schwabe, A. D.; Terasaki, P. I.; Barnett, E. V.; Territo, M. C.;

[127650]Klinenberg, J. R.; Peters, R. S.: Familial Mediterranean fever--recent

[127651]advances in pathogenesis and management. *West. J. Med.* 127: 15-23,

[127652]1977.

[127653]22806.

[127654]Chern, C. J.; Beutler, E.: Biochemical and electrophoretic studies

[127655]of erythrocyte pyridoxine kinase in white and black Americans. *Am.*

[127656]*J. Hum. Genet.* 28: 9-17, 1976.

[127657]22807.

[127658]Hanna, M. C.; Turner, A. J.; Kirkness, E. F.: Human pyridoxal

[127659]kinase: cDNA cloning, expression, and modulation by ligands of the

[127660]benzodiazepine receptor. J. Biol. Chem. 272:
10756–10760, 1997.

[127661]22808.

[127662]Dougherty, K. M.; Brandriss, M. C.; Valle, D.: Cloning hu-
man pyrroline–5–carboxylate

[127663]reductase cDNA by complementation in *Saccharomyces*
cerevisiae. J.

[127664]Biol. Chem. 267: 871–875, 1992.

[127665]22809.

[127666]Merrill, M. J.; Yeh, G. C.; Phang, J. M.: Purified human ery-
throcyte

[127667]pyrroline–5–carboxylate reductase: preferential oxidation
of NADPH. J.

[127668]Biol. Chem. 264: 9352–9358, 1989.

[127669]22810.

[127670]Bigley, R. H.; Stenzel, P.; Jones, R. T.; Campos, J. O.; Koler,

[127671]R. D.: Tissue distribution of human pyruvate kinase
isozymes. Enzym.

[127672]Biol. Clin. 9: 10–20, 1968.

[127673]22811.

[127674]Bunn, H. F.: Personal Communication. Boston, Mass.
4/30/1981.

[127675]22812.

[127676]Chern, C. J.; Kennett, R.; Engel, L. E.; Mellman, W. J.;
Croce,

[127677]C. M.: Assignment of the structural genes for the alpha
subunit of

[127678]hexosaminidase A, mannosephosphate isomerase and
pyruvate kinase to

[127679]the region q22-qter of human chromosome 15. Somat.
Cell Genet. 3:

[127680]553-560, 1977.

[127681]22813.

[127682]Junien, C.; Robinson-Skala, H.; Dreyfus, J. C.; Ravise, N.;
Boue,

[127683]J.; Boue, A.; Kaplan, J. C.: PK3: a new chromosome enzyme
marker

[127684]for gene dosage studies in chromosome 15 imbalance.
Hum. Genet. 54:

[127685]191-196, 1980.

[127686]22814.

[127687]Kahn, A.; Marie, J.; Garreau, H.; Sprengers, E. D.: Subunit structure,
[127688]interrelations and kinetic characteristics of the pyruvate kinase
[127689]from erythrocytes and liver. Biochim. Biophys. Acta 523: 58–74,
[127690]1978.
[127691]22815.
[127692]Levine, M.; Muirhead, H.; Stammers, D. K.; Stuart, D. I.: Structure
[127693]of pyruvate kinase and similarities with other enzymes: possible implications
[127694]for protein taxonomy and evolution. Nature 271: 626–630, 1978.
[127695]22816.
[127696]Nakashima, K.; Miwa, S.; Fujii, H.; Shinohara, K.; Yamauchi, K.;
[127697]Tsuji, Y.; Yanai, M.: Characterization of pyruvate kinase, PK Nagasaki. J.
[127698]Lab. Clin. Med. 90: 1012–1020, 1977.
[127699]22817.

- [127700]Ritter, H.; Friedrichson, U.; Schmitt, J.: Genetic variation of
[127701]mannose phosphate isomerase in man. Humangenetik 22:
261 only, 1974.
[127702]22818.
- [127703]Sheffield, V. C.; Fishman, G. A.; Beck, J. S.; Kimura, A. E.;
[127704]Stone, E. M.: Identification of novel rhodopsin mutations
associated
[127705]with retinitis pigmentosa by GC-clamped denaturing gra-
dient gel electrophoresis. Am.
[127706]J. Hum. Genet. 49: 699-706, 1991.
[127707]22819.
- [127708]Sieving, P. A.; Richards, J. E.; Bingham, E. L.; Naarendorp,
F.
[127709]: Dominant congenital complete nyctalopia and
gly90-to-asp rhodopsin
[127710]mutation. (Abstract) Invest. Ophthalm. Vis. Sci. 33: 1397,
1992.
[127711]22820.
- [127712]Sieving, P. A.; Richards, J. E.; Naarendorp, F.; Bingham, E.
L.;
[127713]Scott, K.; Alpern, M.: Dark-light: model for nightblindness

from

[127714]the human rhodopsin gly90-to-asp mutation. Proc. Nat.
Acad. Sci. 92:

[127715]880-884, 1995.

[127716]22821.

[127717]Sohocki, M. M.; Daiger, S. P.; Bowne, S. J.; Rodriquez, J. A.;

[127718]Northrup, H.; Heckenlively, J. R.; Birch, D. G.; Mintz-Hit-
tner, H.;

[127719]Ruiz, R. S.; Lewis, R. A.; Saperstein, D. A.; Sullivan, L. S.:
Prevalence

[127720]of mutations causing retinitis pigmentosa and other in-
herited retinopathies. Hum.

[127721]Mutat. 17: 42-51, 2001.

[127722]22822.

[127723]Souied, E.; Soubrane, G.; Benlian, P.; Coscas, G. J.; Gerber,

[127724]S.; Munnich, A.; Kaplan, J.: Retinitis punctata albescens
associated

[127725]with the arg135-to-trp mutation in the rhodopsin gene.
Am. J. Ophthal. 121:

[127726]19-25, 1996.

[127727]22823.

[127728]Sullivan, L. J.; Makris, G. S.; Dickinson, P.; Mulhall, L. E.

[127729]M.; App, D.; Forrest, S.; Cotton, R. G. H.; Loughnan, M. S.:

A new

[127730]codon 15 rhodopsin gene mutation in autosomal dominant
retinitis pigmentosa

[127731]is associated with sectorial disease. Arch. Ophthalmol. 111:
1512-1517,

[127732]1993.

[127733]22824.

[127734]Sung, C.-H.; Davenport, C. M.; Hennessey, J. C.; Maumenee, I.

[127735]H.; Jacobson, S. G.; Heckenlively, J. R.; Nowakowski, R.;
Fishman,

[127736]G.; Gouras, P.; Nathans, J.: Rhodopsin mutations in autosomal dominant

[127737]retinitis pigmentosa. Proc. Nat. Acad. Sci. 88: 6481-6485,
1991.

[127738]22825.

[127739]Sung, C.-H.; Davenport, C. M.; Nathans, J.: Rhodopsin
mutations

[127740]responsible for autosomal dominant retinitis pigmentosa:
clustering

[127741]of functional classes along the polypeptide chain. J. Biol.
Chem. 268:

[127742]26645–26649, 1993.

[127743]22826.

[127744]Sung, C.–H.; Makino, C.; Baylor, D.; Nathans, J.: A
rhodopsin

[127745]gene mutation responsible for autosomal dominant retini-
tis pigmentosa

[127746]results in a protein that is defective in localization to the
photoreceptor

[127747]outer segment. J. Neurosci. 14: 5818–5833, 1994.

[127748]22827.

[127749]Sung, C.–H.; Schneider, B. G.; Agarwal, N.; Papermaster, D.
S.;

[127750]Nathans, J.: Functional heterogeneity of mutant
rhodopsins responsible

[127751]for autosomal dominant retinitis pigmentosa. Proc. Nat.
Acad. Sci. 88:

[127752]8840–8844, 1991.

[127753]22828.

[127754]Vaithinathan, R.; Berson, E. L.; Dryja, T. P.: Further screen-

ing

[127755] of the rhodopsin gene in patients with autosomal dominant retinitis

[127756] pigmentosa. Genomics 21: 461–463, 1994.

[127757] 22829.

[127758] Weleber, R. G.; Pillers, D. M.; Powell, B. R.; Hanna, C. E.; Magenis,

[127759] R. E.; Buist, N. R. M.: Aland Island disease (Forsius–Eriksson syndrome)

[127760] associated with contiguous gene syndrome at Xp21: similarity to incomplete

[127761] congenital stationary night blindness. Arch. Ophthal. 107: 1170–1179,

[127762] 1989.

[127763] 22830.

[127764] Boehm, T.; Foroni, L.; Kaneko, Y.; Perutz, M. F.; Rabbitts, T.

[127765] H.: The rhombotin family of cysteine-rich LIM-domain oncogenes: distinct

[127766] members are involved in T-cell translocations to human chromosomes

[127767] 11p15 and 11p13. Proc. Nat. Acad. Sci. 88: 4367–4371,

1991.

[127768]22831.

[127769]Royer-Pokora, B.; Loos, U.; Ludwig, W.-D.: TTG-2, a new gene encoding

[127770]a cysteine-rich protein with the LIM motif, is overexpressed in acute

[127771]T-cell leukaemia with the t(11;14)(p13;q11). Oncogene 6: 1887-1893,

[127772]1991.

[127773]22832.

[127774]Yamada, Y.; Pannell, R.; Forster, A.; Rabbitts, T. H.: The oncogenic

[127775]LIM-only transcription factor Lmo2 regulates angiogenesis but not

[127776]vasculogenesis in mice. Proc. Nat. Acad. Sci. 97: 320-324, 2000.

[127777]22833.

[127778]Yamada, Y.; Warren, A. J.; Dobson, C.; Forster, A.; Pannell, R.;

[127779]Rabbitts, T. H.: The T cell leukemia LIM protein Lmo2 is necessary

[127780]for adult mouse hematopoiesis. Proc. Nat. Acad. Sci. 95:

3890-3895,

[127781]1998.

[127782]22834.

[127783]Yang-Feng, T. L.; Thelander, L.; Lewis, W. H.; Srinivasan,
P. R.;

[127784]Francke, U.: Gene localization of the ribonucleotide reduc-
tase M2

[127785]subunit and of related and co-amplified sequences on hu-
man and mouse

[127786]chromosomes. (Abstract) Am. J. Hum. Genet. 39: A174
only, 1986.

[127787]22835.

[127788]Lindgren, V.; Bryke, C. R.; Ozcelik, T.; Yang-Feng, T. L.;
Francke,

[127789]U.: Phenotypic, cytogenetic, and molecular studies of
three patients

[127790]with constitutional deletions of chromosome 5 in the re-
gion of the

[127791]gene for familial adenomatous polyposis. Am. J. Hum.
Genet. 50:

[127792]988-997, 1992.

[127793]22836.

[127794]Barletta, C.; Druck, T.; LaForgia, S.; Calabretta, B.;
Drabkin,
[127795]H.; Patterson, D.; Croce, C. M.; Huebner, K.: Chromosome
locations
[127796]of the MYB related genes, AMYB and BMYB. Cancer Res.
51: 3821-3824,
[127797]1991.
[127798]22837.
[127799]Nomura, N.; Takahashi, M.; Matsui, M.; Ishii, S.; Date, T.;
Sasamoto,
[127800]S.; Ishizaki, R.: Isolation of human cDNA clones of MYB-
related genes,
[127801]A-MYB and B-MYB. Nucleic Acids Res. 16: 11075-11089,
1988.
[127802]22838.
[127803]Takahashi, T.; Nakagoshi, H.; Sarai, A.; Nomura, N.; Ya-
mamoto,
[127804]T.; Ishii, S.: Human A-myb gene encodes a transcriptional
activator
[127805]containing the negative regulatory domains. FEBS Lett.
358: 89-96,
[127806]1995.

[127807]22839.

[127808]Boylan, K. B.; Takahashi, N.; Diamond, M.; Hood, L. E.;
Prusiner,

[127809]S. B.: DNA length polymorphism located 5-prime to the
human myelin

[127810]basic protein gene. Am. J. Hum. Genet. 40: 387-400,
1987.

[127811]22840.

[127812]Dayhoff, M. O.: Atlas of Protein Sequence and Structure.
Myelin

[127813]membrane encephalitogenic protein. Washington: National
Biomedical

[127814]Research Foundation (pub.) 5: 1972. Pp. D324 only.

[127815]22841.

[127816]Eylar, E. H.; Brostoff, S.; Hashim, G.; Westall, F. C.: Basic

[127817]A1 protein of the myelin membrane: the complete amino
acid sequence. J.

[127818]Biol. Chem. 246: 5770-5784, 1971.

[127819]22842.

[127820]Gomez, C. M.; Muggleton-Harris, A. L.; Whittingham, D.
G.; Hood,

[127821]L. E.; Readhead, C.: Rapid preimplantation detection of mutant (shiverer) and normal alleles of the mouse myelin basic protein gene allowing selective implantation and birth of live young. Proc. Nat. Acad. Sci. 87: 4481–4484, 1990.

[127822]22843.

[127823]Kamholz, J.; de Ferra, F.; Puckett, C.; Lazzarini, R.: Identification of three forms of human myelin basic protein by cDNA cloning. Proc. Nat. Acad. Sci. 83: 4962–4966, 1986.

[127824]22844.

[127825]Kamholz, J.; Spielman, R.; Gogolin, K.; Modi, W.; O'Brien, S.; Lazzarini, R.: The human myelin–basic–protein gene: chromosomal localization and RFLP analysis. Am. J. Hum. Genet. 40: 365–373, 1987.

[127826]22845.

[127827]Lalley, P. A.; McKusick, V. A.: Report of the committee on

comparative

[127835]mapping (HGM8). Cytogenet. Cell Genet. 40: 536–566,
1985.

[127836]22846.

[127837]Martensen, R. E.: Myelin basic protein speciation. Prog.
Clin.

[127838]Biol. Res. 146: 511–521, 1984.

[127839]22847.

[127840]Marty, M. C.; Alliot, F.; Rutin, J.; Fritz, R.; Trisler, D.; Pes-
sac,

[127841]B.: The myelin basic protein gene is expressed in differen-
tiated

[127842]blood cell lineages and in hemopoietic progenitors. Proc.
Nat. Acad.

[127843]Sci. 99: 8856–8861, 2002.

[127844]22848.

[127845]Molineaux, S. M.; Engh, H.; de Ferra, F.; Hudson, L.; Laz-
zarini,

[127846]R. A.: Recombination within the myelin basic protein gene
created

[127847]the dysmyelinating shiverer mouse mutation. Proc. Nat.

Acad. Sci. 83:

[127848]7542–7546, 1986.

[127849]22849.

[127850]Popko, B.; Puckett, C.; Lai, E.; Shine, H. D.; Readhead, C.;
Takahashi,

[127851]N.; Hunt, S. W., III; Sidman, R. L.; Hood, L.: Myelin deficient
mice:

[127852]expression of myelin basic protein and generation of mice
with varying

[127853]levels of myelin. Cell 48: 713–721, 1987.

[127854]22850.

[127855]Readhead, C.; Popko, B.; Takahashi, N.; Shine, H. D.;
Saavedra,

[127856]R. A.; Sidman, R. L.; Hood, L.: Expression of a myelin basic
protein

[127857]gene in transgenic shiverer mice: correction of the dys-
myelinating

[127858]phenotype. Cell 48: 703–712, 1987.

[127859]22851.

[127860]Roach, A.; Boylan, K.; Horvath, S.; Prusiner, S. B.; Hood, L.

[127861]E.: Characterization of cloned cDNA representing rat

myelin basic

[127862]protein: absence of expression in shiverer mutant mice.

Cell 34:

[127863]799–806, 1983.

[127864]22852.

[127865]Roach, A.; Takahashi, N.; Pravtcheva, D.; Ruddle, F.; Hood, L.

[127866]: Chromosomal mapping of mouse myelin basic protein gene and structure

[127867]and transcription of the partially deleted gene in shiverer mutant

[127868]mice. Cell 42: 149–155, 1985.

[127869]22853.

[127870]Saxe, D. F.; Takahashi, N.; Hood, L.; Simon, M. I.: Localization

[127871]of the human myelin basic protein gene (MBP) to region 18q22–qter

[127872]by in situ hybridization. Cytogenet. Cell Genet. 39: 246–249, 1985.

[127873]22854.

[127874]Sheremata, W.; Cosgrove, J. B. R.; Hylar, E. H.: Cellular hy-

persensitivity

[127875]to basic myelin (A1) protein and clinical multiple sclerosis.

New

[127876]Eng. J. Med. 291: 14-17, 1974.

[127877]22855.

[127878]Sidman, R.: Personal Communication. Boston, Mass. 1983.

[127879]22856.

[127880]Sidman, R. L.; Conover, C. S.; Carson, J. H.: Shiverer gene
maps

[127881]near the distal end of chromosome 18 in the house
mouse. Cytogenet.

[127882]Cell Genet. 39: 241-245, 1985.

[127883]22857.

[127884]Sparkes, R. S.; Mohandas, T.; Heinzmann, C.; Roth, H. J.;
Klisak,

[127885]I.; Campagnoni, A. T.: Assignment of the myelin basic
protein gene

[127886]to human chromosome 18q22-qter. Hum. Genet. 75:
147-150, 1987.

[127887]22858.

[127888]Takahashi, N.; Roach, A.; Teplow, D. B.; Prusiner, S. B.;

Hood,

[127889]L.: Cloning and characterization of the myelin basic protein gene

[127890]from mouse: one gene can encode both 14 kd and 18.5 kd MBPs by alternate

[127891]use of exons. Cell 42: 139–148, 1985.

[127892]22859.

[127893]Eiberg, H.; Mohr, J.; Nielsen, L. S.; Simonsen, N.: Genetics

[127894]and linkage relationships of the C3 polymorphism: discovery of C3–Se

[127895]linkage and assignment of LES–C3–DM–Se–PEPD–Lu synteny to chromosome

[127896]19. Clin. Genet. 24: 159–170, 1983.

[127897]22860.

[127898]Barouch, L. A.; Harrison, R. W.; Skaf, M. W.; Rosas, G. O.; Cappola,

[127899]T. P.; Kobeissi, Z. A.; Hobai, I. A.; Lemmon, C. A.; Burnett, A. L.;

[127900]O'Rourke, B.; Rodriguez, E. R.; Huang, P. L.; Lima, J. A. C.; Berkowitz,

[127901]D. E.; Hare, J. M.: Nitric oxide regulates the heart by spa-

tial confinement

[127902]of nitric oxide synthase isoforms. Nature 416: 337–340,
2002.

[127903]22861.

[127904]Dimmeler, S.; Fleming, I.; Fisslthaler, B.; Hermann, C.;
Busse,

[127905]R.; Zeiher, A. M.: Activation of nitric oxide synthase in en-
dothelial

[127906]cells by Akt-dependent phosphorylation. Nature 399:
601–605, 1999.

[127907]22862.

[127908]Fulton, D.; Gratton, J.-P.; McCabe, T. J.; Fontana, J.; Fujio,

[127909]Y.; Walsh, K.; Franke, T. F.; Papapetropoulos, A.; Sessa, W.
C.:

[127910]Regulation of endothelium-derived nitric oxide production
by the protein

[127911]kinase Akt. Nature 399: 597–601, 1999.

[127912]22863.

[127913]Kruh, G. D.; King, C. R.; Kraus, M. H.; Popescu, N. C.;
Amsbaugh,

[127914]S. C.; McBride, W. O.; Aaronson, S. A.: A novel human gene

closely

[127915]related to the abl proto-oncogene. Science 234:
1545-1548, 1986.

[127916]22864.

[127917]Audrezet, M.-P.; Chen, J.-M.; Le Marechal, C.;
Ruszniewski, P.;

[127918]Robaszkiewicz, M.; Raguene, O.; Quere, I.; Scotet, V.;
Ferec, C.

[127919]: Determination of the relative contribution of three
genes--the cystic

[127920]fibrosis transmembrane conductance regulator gene, the
cationic trypsinogen

[127921]gene, and the pancreatic secretory trypsin inhibitor gene--
to the

[127922]etiology of idiopathic chronic pancreatitis. Europ. J. Hum.
Genet. 10:

[127923]100-106, 2002.

[127924]22865.

[127925]Arnaud, J.; Vavrusa, B.; Sevin, J.; Constans, J.: Human red-
cell

[127926]acid phosphatase (ACP1): a new mutant (ACP1*KUK) de-
tected by isoelectric

[127927] focusing, kinetics of thermostability and substrate activity.

Hum.

[127928] Hered. 39: 288–293, 1989.

[127929] 22866.

[127930] Arnaud, J.; Vavrusa, B.; Wiederanders, G.; Constans, J.:

Human

[127931] red-cell acid phosphatase (ACP1): kinetic and thermodynamic characterization

[127932] of the KUK variant. Hum. Hered. 42: 140–142, 1992.

[127933] 22867.

[127934] Beemer, F. A.; van der Heiden, C.; Van Hemel, J. O.;

Jansen, M.

[127935]: Letter to the editors. (Letter) Clin. Genet. 24: 151, 1983.

[127936] 22868.

[127937] Berg, K.: Close linkage between APOB and ACP1 excluded.

(Abstract) Cytogenet.

[127938] Cell Genet. 46: 580, 1987.

[127939] 22869.

[127940] Bottini, E.; Carapella, E.; Orzalesi, M.; Lucarelli, P.; Pas-

cone,

[127941]R.; Gloria-Bottini, F.; Coccia, M.: Is there a role of erythrocyte

[127942]acid phosphatase polymorphism in intrauterine development? (Letter) Am.

[127943]J. Hum. Genet. 32: 764-767, 1980.

[127944]22870.

[127945]Seldin, M. F.; Kruh, G. D.: Mapping of Abll within a conserved

[127946]linkage group on distal mouse chromosome 1 syntenic with human chromosome

[127947]1 using an interspecific cross. Genomics 4: 221-223, 1989.

[127948]22871.

[127949]Bello, M. J.; Salagnon, N.; Rey, J. A.; Guichaoua, M. R.; Berge-Lefranc,

[127950]J. L.; Jordan, B. R.; Luciani, J. M.: Precise in situ localization

[127951]of NCAM, ETS1, and D11S29 on human meiotic chromosomes. Cytogenet.

[127952]Cell Genet. 52: 7-10, 1989.

[127953]22872.

[127954]Concannon, P.; Malhotra, U.; Charmley, P.; Reynolds, J.;

Lange,

[127955]K.; Gatti, R. A.: The ataxia-telangiectasia gene (ATA) on chromosome

[127956]11 is distinct from the ETS-1 gene. Am. J. Hum. Genet. 46: 789-794,

[127957]1990.

[127958]22873.

[127959]Gold, D. P.; van Dongen, J. J. M.; Morton, C. C.; Bruns, G. A.

[127960]P.; van den Elsen, P.; Geurts van Kessel, A. H. M.; Terhorst, C.:

[127961]The gene encoding the epsilon subunit of the T3/T-cell receptor complex

[127962]maps to chromosome 11 in humans and to chromosome 9 in mice. Proc.

[127963]Nat. Acad. Sci. 84: 1664-1668, 1987.

[127964]22874.

[127965]Ohtani, N.; Zebedee, Z.; Huot, T. J. G.; Stinson, J. A.; Sugimoto,

[127966]M.; Ohashi, Y.; Sharrocks, A. D.; Peters, G.; Hara, E.: Opposing

[127967]effects of Ets and Id proteins on p16(INK4A) expression

during cellular

[127968]senescence. Nature 409: 1067–1070, 2001.

[127969]22875.

[127970]Sacchi, N.; Watson, D. K.; Geurts van Kessel, A. H. M.;
Hagemeijer,

[127971]A.; Kersey, J.; Drabkin, H. D.; Patterson, D.; Papas, T. S.:
Hu-ets-1

[127972]and Hu-ets-2 genes are transposed in acute leukemias
with (4;11) and

[127973](8;21) translocations. Science 231: 379–382, 1986.

[127974]22876.

[127975]Watson, D. K.; McWilliams-Smith, M. J.; Kozak, C.; Reeves,
R.;

[127976]Gearhart, J.; Nunn, M. F.; Nash, W.; Fowle, J. R., III; Dues-
berg,

[127977]P.; Papas, T. S.; O'Brien, S. J.: Conserved chromosomal po-
sitions

[127978]of dual domains of the ets protooncogene in cats, mice,
and humans. Proc.

[127979]Nat. Acad. Sci. 83: 1792–1796, 1986.

[127980]22877.

[127981]Watson, D. K.; McWilliams-Smith, M. J.; Nunn, M. F.; Duesberg,

[127982]P. H.; O'Brien, S. J.; Papas, T. S.: The ets sequence from the transforming

[127983]gene of avian erythroblastosis virus, E26, has unique domains on human

[127984]chromosomes 11 and 21: both loci are transcriptionally active. Proc.

[127985]Nat. Acad. Sci. 82: 7294-7298, 1985.

[127986]22878.

[127987]Bellacosa, A.; Franke, T. F.; Gonzalez-Portal, M. E.; Datta, K.;

[127988]Taguchi, T.; Gardner, J.; Cheng, J. Q.; Testa, J. R.; Tschlis, P.

[127989]N.: Structure, expression and chromosomal mapping of c-akt: relationship

[127990]to v-akt and its implications. Oncogene 8: 745-754, 1993.

[127991]22879.

[127992]Brunet, A.; Bonni, A.; Zigmund, M. J.; Lin, M. Z.; Juo, P.; Hu,

[127993]L. S.; Anderson, M. J.; Arden, K. C.; Blenis, J.; Greenberg,

M. E.

[127994]: Akt promotes cell survival by phosphorylating and inhibiting a Forkhead

[127995]transcription factor. Cell 96: 857–868, 1999.

[127996]22880.

[127997]Chen, W. S.; Xu, P.-Z.; Gottlob, K.; Chen, M.-L.; Sokol, K.; Shiyanova,

[127998]T.; Roninson, I.; Weng, W.; Suzuki, R.; Tobe, K.; Kadowaki, T.; Hay,

[127999]N.: Growth retardation and increased apoptosis in mice with homozygous

[128000]disruption of the akt1 gene. Genes Dev. 15: 2203–2208, 2001.

[128001]22881.

[128002]Condorelli, G.; Drusco, A.; Stassi, G.; Bellacosa, A.; Roncarati,

[128003]R.; Iaccarino, G.; Russo, M. A.; Gu, Y.; Dalton, N.; Chung, C.; Latronico,

[128004]M. V. G.; Napoli, C.; Sadoshima, J.; Croce, C. M.; Ross, J., Jr.:

[128005]Akt induces enhanced myocardial contractility and cell size in vivo

[128006]in transgenic mice. Proc. Nat. Acad. Sci. 99:
12333–12338, 2002.

[128007]22882.

[128008]Delettre, C.; Lenaers, G.; Griffoin, J.–M.; Gigarel, N.;
Lorenzo,
[128009]C.; Belenguer, P.; Pelloquin, L.; Grosgeorge, J.; Turc–Carel,
C.;

[128010]Perret, E.; Astarie–Dequeker, C.; Lasquelllec, L.; Arnaud, B.;
Ducommun,
[128011]B.; Kaplan, J.; Hamel, C. P.: Nuclear gene OPA1, encoding a
mitochondrial
[128012]dynammin–related protein, is mutated in dominant optic at–
rophy. Nature
[128013]Genet. 26: 207–210, 2000.

[128014]22883.

[128015]Curry, C. J. R.; Magenis, R. E.; Brown, M.; Lanman, J. T., Jr.;

[128016]Tsai, J.; O’Lague, P.; Goodfellow, P.; Mohandas, T.;

Bergner, E. A.;

[128017]Shapiro, L. J.: Inherited chondrodysplasia punctata due to
a deletion
[128018]of the terminal short arm of an X chromosome. New Eng.
J. Med. 311:

[128019]1010–1015, 1984.

[128020]22884.

[128021]Bick, D.; Curry, C. J. R.; McGill, J. R.; Schorderet, D. F.; Bux,

[128022]R. C.; Moore, C. M.: Male infant with ichthyosis, Kallmann syndrome,

[128023]chondrodysplasia punctata, and an Xp chromosome deletion. Am. J.

[128024]Med. Genet. 33: 100–107, 1989.

[128025]22885.

[128026]Marlhens, F.; Chelly, J.; Kaplan, J. C.; Lefrancois, D.; Harpey,

[128027]J. P.; Dutrillaux, B.: Familial deletion of Xp21.2 with glycerol

[128028]kinase deficiency and congenital adrenal hypoplasia. Hum. Genet. 77:

[128029]379–383, 1987.

[128030]22886.

[128031]Lin, T.; Orrison, B. M.; Suchy, S. F.; Lewis, R. A.; Nussbaum,

[128032]R. L.: Mutations are not uniformly distributed throughout the OCRL1

[128033]gene in Lowe syndrome patients. Molec. Genet. Metab. 64:
58–61,

[128034]1998.

[128035]22887.

[128036]Poustka, A.; Dietrich, A.; Langenstein, G.; Toniolo, D.;
Warren,

[128037]S. T.; Lehrach, H.: Physical map of human Xq27–qter: lo-
calizing the

[128038]region of the fragile X mutation. Proc. Nat. Acad. Sci. 88:
8302–8306,

[128039]1991.

[128040]22888.

[128041]Ma, K.; Sharkey, A.; Kirsch, S.; Vogt, P.; Keil, R.; Hargreave,

[128042]T. B.; McBeath, S.; Chandley, A. C.: Towards the molecular
localisation

[128043]of the AZF locus: mapping of microdeletions in azoosper-
mic men within

[128044]14 subintervals of interval 6 of the human Y chromosome.
Hum. Molec.

[128045]Genet. 1: 29–33, 1992.

[128046]22889.

[128047]Adam, A.; Tippett, P.; Gavin, J.; Noades, J.; Sanger, R.;
Race,

[128048]R. R.: The linkage relation of Xg to G6PD in Israelis: the
evidence

[128049]of a second series of families. Am. J. Hum. Genet. 30:
211–218,

[128050]1966.

[128051]22890.

[128052]Ahluwalia, A.; Corcoran, C. M.; Vulliamy, T. J.; Ishwad, C.
S.;

[128053]Naidu, J. M.; Argusti, A.; Stevens, D. J.; Mason, P. J.; Luz-
zatto,

[128054]L.: G6PD Kalyan and G6PD Kerala; two deficient variants in
India

[128055]caused by the same 317 glu-to-lys mutation. Hum. Molec.
Genet. 3:

[128056]209–210, 1992.

[128057]22891.

[128058]Aksoy, K.; Yuregir, G. T.; Dikmen, N.; Unlukurt, I.: Three
new

[128059]G6PD variants, G6PD Adana, G6PD Samandag, and G6PD
Balcali in Cukurova,

[128060]Turkey. Hum. Genet. 76: 199–201, 1987.

[128061]22892.

[128062]Alfinito, F.; Cimmino, A.; Ferraro, F.; Cubellis, M. V.;
Vitagliano,

[128063]L.; Francese, M.; Zagari, A.; Rotoli, B.; Filosa, S.; Martini, G.

[128064]: Molecular characterization of G6PD deficiency in South-
ern Italy:

[128065]heterogeneity, correlation genotype–phenotype and de-
scription of a

[128066]new variant (G6PD Neapolis). Brit. J. Haemat. 98: 41–46,
1997.

[128067]22893.

[128068]Alperin, J. B.; Mills, G. C.: New variants of glucose-
6-phosphate

[128069]dehydrogenase (G6PD). Clin. Res. 20: 76, 1972.

[128070]22894.

[128071]Azevedo, E.; Kirkman, H. N.; Morrow, A. C.; Motulsky, A.
G.: Variants

[128072]of red cell glucose–6-phosphate dehydrogenase among
Asiatic Indians. Ann.

[128073]Hum. Genet. 31: 373–379, 1968.

[128074]22895.

[128075]Azevedo, E. S.; Yoshida, A.: Brazilian variant of glucose-6-phosphate

[128076]dehydrogenase (GD Minas Gerais). Nature 222: 380-382, 1969.

[128077]22896.

[128078]Babalola, A. O. G.; Beetlestone, J. G.; Luzzatto, L.: Genetic

[128079]variants of human erythrocyte glucose-6-phosphate dehydrogenase: kinetic

[128080]and thermodynamic parameters of variants A, B, and A- in relation

[128081]to quaternary structure. J. Biol. Chem. 251: 2993-3002, 1976.

[128082]22897.

[128083]Balinsky, D.; Cayanis, E.; Carter, G.; Jenkins, T.; Bersohn, I.

[128084]: A new variant of human erythrocyte glucose-6-phosphate dehydrogenase:

[128085]G6PD Port Elizabeth. Int. J. Biochem. 4: 235-244, 1973.

[128086]22898.

[128087]Balinsky, D.; Gomperts, E.; Cayanis, E.; Jenkins, T.; Bryer, D.;

[128088]Bersohn, I.; Metz, J.: Glucose 6-phosphate dehydrogenase
Johannesburg:

[128089]a new variant with reduced activity in a patient with con-
genital non-spherocytic

[128090]haemolytic anaemia. Brit. J. Haemat. 25: 385-392, 1973.

[128091]22899.

[128092]Araten, D. J.; Nafa, K.; Pakdeesuwan, K.; Luzzatto, L.:
Clonal

[128093]populations of hematopoietic cells with paroxysmal noc-
turnal hemoglobinuria

[128094]genotype and phenotype are present in normal individu-
als. Proc. Nat.

[128095]Acad. Sci. 96: 5209-5214, 1999.

[128096]22900.

[128097]Bessler, M.; Mason, P.; Hillmen, P.; Luzzatto, L.: Somatic
mutations

[128098]and cellular selection in paroxysmal nocturnal
haemoglobinuria. Lancet 343:

[128099]951-953, 1994.

[128100]22901.

[128101]Bessler, M.; Mason, P. J.; Hillmen, P.; Miyata, T.; Yamada,

N.;

[128102]Takeda, J.; Luzzatto, L.; Kinoshita, T.: Paroxysmal nocturnal haemoglobinuria

[128103](PNH) is caused by somatic mutations in the PIG-A gene.

EMBO J. 13:

[128104]110–117, 1994.

[128105]22902.

[128106]Brodsky, R. A.; Vala, M. S.; Barber, J. P.; Medof, M. E.;

Jones,

[128107]R. J.: Resistance to apoptosis caused by PIG-A gene mutations in

[128108]paroxysmal nocturnal hemoglobinuria. Proc. Nat. Acad.

Sci. 94: 8756–8760,

[128109]1997.

[128110]22903.

[128111]Hillmen, P.; Lewis, S. M.; Bessler, M.; Luzzatto, L.; Dacie, J.

[128112]V.: Natural history of paroxysmal nocturnal hemoglobinuria. New

[128113]Eng. J. Med. 333: 1253–1258, 1995.

[128114]22904.

[128115]Hirose, S.; Mohny, R. P.; Mutka, S. C.; Ravi, L.; Singleton,

D.

[128116]R.; Perry, G.; Tartakoff, A. M.; Medof, M. E.: Derivation and characterization

[128117]of glycoinositol-phospholipid anchor-defective human K562 cell clones. J.

[128118]Biol. Chem. 267: 5272-5278, 1992.

[128119]22905.

[128120]Hirose, S.; Ravi, L.; Prince, G. M.; Rosenfeld, M. G.; Silber,

[128121]R.; Andresen, S. W.; Hazra, S. V.; Medof, M. E.: Synthesis of mannosylglucosaminylinositol

[128122]phospholipids in normal but not paroxysmal nocturnal hemoglobinuria

[128123]cells. Proc. Nat. Acad. Sci. 89: 6025-6029, 1992.

[128124]22906.

[128125]Horikawa, K.; Kawaguchi, T.; Ishihara, S.; Nagakura, S.; Hida,

[128126]M.; Kagimoto, T.; Mitsuya, H.; Nakakuma, H.: Frequent detection of

[128127]T cells with mutations of the hypoxanthine-guanine phosphoribosyltransferase

[128128]gene in patients with paroxysmal nocturnal hemoglobin-

uria. Blood 99:

[128129]24–29, 2002.

[128130]22907.

[128131]Iida, Y.; Takeda, J.; Miyata, T.; Inoue, N.; Nishimura, J.; Kitanishi, T.; Maeda, K.; Kinoshita, T.: Characterization of genomic

PIG-A gene:

[128133]a gene for glycosylphosphatidylinositol-anchor biosynthesis and paroxysmal

[128134]nocturnal hemoglobinuria. Blood 83: 3126–3131, 1994.

[128135]22908.

[128136]Kawagoe, K.; Takeda, J.; Endo, Y.; Kinoshita, T.: Molecular cloning

[128137]of murine Pig-a, a gene for GPI-anchor biosynthesis, and demonstration

[128138]of interspecies conservation of its structure, function, and genetic

[128139]locus. Genomics 23: 566–574, 1994.

[128140]22909.

[128141]Keller, P.; Tremml, G.; Rosti, V.; Bessler, M.: X inactivation

[128142]and somatic cell selection rescue female mice carrying a

Piga-null

[128143]mutation. Proc. Nat. Acad. Sci. 96: 7479–7483, 1999.

[128144]22910.

[128145]Luzzatto, L.; Bessler, M.: The dual pathogenesis of paroxysmal

[128146]nocturnal hemoglobinuria. Curr. Opin. Hemat. 3: 101–110, 1996.

[128147]22911.

[128148]Luzzatto, L.; Bessler, M.; Rotoli, B.: Somatic mutations in paroxysmal

[128149]nocturnal hemoglobinuria: a blessing in disguise? Cell 88: 1–4,

[128150]1997.

[128151]22912.

[128152]Maugard, C.; Margueritte, G.; Tuffery, S.; Rabesandratana, H.;

[128153]Demaille, J.; Claustres, M.: Recurrent PIG-A mutation (IVS5+1G-A)

[128154]in a paediatric case of paroxysmal nocturnal haemoglobinuria: detection

[128155]by the protein truncation test. Brit. J. Haemat. 98: 21–24,

1997.

[128156]22913.

[128157]Miyata, T.; Takeda, J.; Iida, Y.; Yamada, N.; Inoue, N.;
Takahashi,

[128158]M.; Maeda, K.; Kitani, T.; Kinoshita, T.: The cloning of PIG-
A, a

[128159]component in the early step of GPI-anchor biosynthesis.
Science 259:

[128160]1318-1320, 1993.

[128161]22914.

[128162]Miyata, T.; Yamada, N.; Iida, Y.; Nishimura, J.; Takeda, J.;
Kitani,

[128163]T.; Kinoshita, T.: Abnormalities of PIG-A transcripts in
granulocytes

[128164]from patients with paroxysmal nocturnal hemoglobinuria.
New Eng.

[128165]J. Med. 330: 249-255, 1994.

[128166]22915.

[128167]Nafa, K.; Bessler, M.; Castro-Malaspina, H.; Jhanwar, S.;
Luzzatto,

[128168]L.: The spectrum of somatic mutations in the PIG-A gene

in paroxysmal

[128169]nocturnal hemoglobinuria includes large deletions and small duplications. Blood

[128170]Cells Molec. Dis. 24: 370–384, 1998.

[128171]22916.

[128172]Nafa, K.; Bessler, M.; Deeg, H. J.; Luzzatto, L.: New somatic

[128173]mutation in the PIG-A gene emerges at relapse of paroxysmal nocturnal

[128174]hemoglobinuria. Blood 92: 3422–3427, 1998.

[128175]22917.

[128176]Nafa, K.; Mason, P. J.; Hillmen, P.; Luzzatto, L.; Bessler, M.

[128177]: Mutations in the PIG-A gene causing paroxysmal nocturnal hemoglobinuria

[128178]are mainly of the frameshift type. Blood 86: 4650–4655, 1995.

[128179]22918.

[128180]Nagarajan, S.; Brodsky, R. A.; Young, N. S.; Medof, M. E.: Genetic

[128181]defects underlying paroxysmal nocturnal hemoglobinuria that arises

[128182]out of aplastic anemia. Blood 86: 4656–4661, 1995.

[128183]22919.

[128184]Nagarajan, S.; Brown, C. J.; Medof, M. E.: Identification of

[128185]a PIG-A related processed gene on chromosome 12. Hum.
Genet. 95:

[128186]691-697, 1995.

[128187]22920.

[128188]McCabe, E. R. B.: Personal Communication. Denver, Colo.
10/9/1985.

[128189]22921.

[128190]McCabe, E. R. B.; Fennessey, P. V.; Guggenheim, M. A.;
Miles,

[128191]B. S.; Bullen, W. W.; Sceats, D. J.; Goodman, S. I.: Human
glycerol

[128192]kinase deficiency with hyperglycerolemia and glyceroluria.
Biochem.

[128193]Biophys. Res. Commun. 78: 1327-1333, 1977.

[128194]22922.

[128195]Renier, W. O.; Nabben, F. A. E.; Hustinx, T. W. J.;
Veerkamp,

[128196]J. H.; Otten, B. J.; ter Laak, H. J.; ter Haar, B. G. A.;
Gabreels,

- [128197]F. J. M.: Congenital adrenal hypoplasia, progressive muscular dystrophy
- [128198]and severe mental retardation, in association with glycerol kinase
- [128199]deficiency, in male sibs. Clin. Genet. 24: 449–457, 1983.
- [128200]22923.
- [128201]Rose, C. I.; Haines, D. S. M.: Familial hyperglycerolemia. J.
- [128202]Clin. Invest. 61: 163–170, 1978.
- [128203]22924.
- [128204]Saito, F.; Goto, J.; Kakinuma, H.; Nakamura, F.; Murayama, S.;
- [128205]Nakano, I.; Tonomura, A.: Inherited Xp21 deletion in a boy with complex
- [128206]glycerol kinase deficiency syndrome. (Letter) Clin. Genet. 29: 92–93,
- [128207]1986.
- [128208]22925.
- [128209]Sargent, C. A.; Affara, N. A.; Bentley, E.; Pelmear, A.; Bailey,
- [128210]D. M. D.; Davey, P.; Dow, D.; Leversha, M.; Aplin, H.; Besley, G.

[128211]T. N.; Ferguson-Smith, M. A.: Cloning of the X-linked
glycerol kinase

[128212]deficiency gene and its identification by sequence com-
parison to the

[128213]*Bacillus subtilis* homologue. *Hum. Molec. Genet.* 2:
97-106, 1993.

[128214]22926.

[128215]Sargent, C. A.; Kidd, A.; Moore, S.; Dean, J.; Besley, G. T.
N.;

[128216]Affara, N. A.: Five cases of isolated glycerol kinase defi-
ciency,

[128217]including two families: failure to find genotype:phenotype
correlation. *J.*

[128218]*Med. Genet.* 37: 434-441, 2000.

[128219]22927.

[128220]Sargent, C. A.; Young, C.; Marsh, S.; Ferguson-Smith, M.
A.; Affara,

[128221]N. A.: The glycerol kinase gene family: structure of the Xp
gene,

[128222]and related intronless retroposons. *Hum. Molec. Genet.* 3:
1317-1324,

[128223]1994.

[128224]22928.

[128225]Sjarif, D. R.; Ploos van Amstel, J. K.; Duran, M.; Beemer, F.

[128226]A.; Poll-The, B. T.: Isolated and contiguous glycerol kinase
gene

[128227]disorders: a review. J. Inherit. Metab. Dis. 23: 529-547,
2000.

[128228]22929.

[128229]Sjarif, D. R.; Sinke, R. J.; Duran, M.; Beemer, F. A.; Kleijer,

[128230]W. J.; Ploos van Amstel, J. K.; Poll-The, B. T.: Clinical het-
erogeneity

[128231]and novel mutations in the glycerol kinase gene in three
families

[128232]with isolated glycerol kinase deficiency. J. Med. Genet. 35:
650-656,

[128233]1998.

[128234]22930.

[128235]von Petrykowski, W.; Beckmann, R.; Bohm, N.; Ketelsen,
U.-P.;

[128236]Ropers, H. H.; Sauer, M.: Adrenal insufficiency, myopathic
hypotonia,

[128237]severe psychomotor retardation, failure to thrive, consti-

pation and

[128238]bladder ectasia in 2 brothers: adrenomyodystrophy. *Helv. Paediat.*

[128239]*Acta* 37: 387–400, 1982.

[128240]22931.

[128241]Walker, A. P.; Muscatelli, F.; Monaco, A. P.: Isolation of the

[128242]human Xp21 glycerol kinase gene by positional cloning. *Hum. Molec.*

[128243]*Genet.* 2: 107–114, 1993.

[128244]22932.

[128245]Walker, A. P.; Muscatelli, F.; Stafford, A. N.; Chelly, J.; Dahl,

[128246]N.; Blomquist, H. K.; Delanghe, J.; Willems, P. J.; Steinmann, B.;

[128247]Monaco, A. P.: Mutations and phenotype in isolated glycerol kinase

[128248]deficiency. *Am. J. Hum. Genet.* 58: 1205–1211, 1996.

[128249]22933.

[128250]Wieringa, B.; Hustinx, T.; Scheres, J.; Hofker, M.; Schepens,

[128251]J.; Ropers, H. H.; ter Haar, B.: Glycerol kinase deficiency

syndrome

[128252]explained as X-chromosomal deletion. (Abstract) Cytogenet. Cell Genet. 40:

[128253]777 only, 1985.

[128254]22934.

[128255]Wieringa, B.; Hustinx, T.; Scheres, J.; Renier, W.; ter Haar,

[128256]B.: Complex glycerol kinase deficiency syndrome explained as X-chromosomal

[128257]deletion. (Letter) Clin. Genet. 27: 522-523, 1985.

[128258]22935.

[128259]Willard, H. F.: Cloning of the X-linked glycerol kinase gene. Hum.

[128260]Molec. Genet. 2: 95-96, 1993.

[128261]22936.

[128262]Zhang, Y.-H.; Dipple, K. M.; Vilain, E.; Huang, B.-L.; Finlayson,

[128263]G.; Therrell, B. L.; Worley, K.; Deininger, P.; McCabe, E. R. B.:

[128264]AluY insertion (IVS4-52ins316alu) in the glycerol kinase gene from

[128265]an individual with benign glycerol kinase deficiency. Hum.

Mutat. 15:

[128266]316–323, 2000.

[128267]22937.

[128268]Balinsky, D.; Rootman, A. J.; Nurse, G. T.; Cayanis, E.;
Lane,

[128269]A.; Jenkins, T.; Bersohn, I.: G6PD Kuanyama: a new variant
of human

[128270]erythrocyte glucose 6–phosphate dehydrogenase showing
slower than

[128271]normal electrophoretic mobility. S. Afr. J. Med. Sci. 39:
5–13,

[128272]1974.

[128273]22938.

[128274]Barretto, O. C.; Nonoyama, K.: Gd(+) Cuiaba, a new rare
glucose–6–phosphate

[128275]dehydrogenase variant presenting normal activity. Hum.
Genet. 77:

[128276]201–202, 1987.

[128277]22939.

[128278]Kornak, U.; Kasper, D.; Bosl, M. R.; Kaiser, E.; Schweizer,
M.;

[128279]Schulz, A.; Friedrich, W.; Delling, G.; Jentsch, T. J.: Loss of the

[128280]ClC-7 chloride channel leads to osteopetrosis in mice and man. Cell 104:

[128281]205–215, 2001.

[128282]22940.

[128283]Kornak, U.; Schulz, A.; Friedrich, W.; Uhlhaas, S.; Kremens, B.;

[128284]Voit, T.; Hasan, C.; Bode, U.; Jentsch, T. J.; Kubisch, C.: Mutations

[128285]in the $\alpha 3$ subunit of the vacuolar H(+)-ATPase cause infantile malignant

[128286]osteopetrosis. Hum. Molec. Genet. 9: 2059–2063, 2000.

[128287]22941.

[128288]Lowe, C. U.: Oculo-cerebral-renal syndrome. Maandschr. Kindergeneesk. 28:

[128289]77–80, 1960.

[128290]22942.

[128291]Lowe, C. U.; Terrey, M.; MacLachlan, E. A.: Organic-aciduria,

[128292]decreased renal ammonia production, hydrophthalmos,

and mental retardation. Am.

[128293]J. Dis. Child. 83: 164–184, 1952.

[128294]22943.

[128295]Martin, V. A. F.; Carson, N. A. J.: Inborn metabolic disorders

[128296]with associated ocular lesions in Northern Ireland. Trans. Ophthal.

[128297]Soc. U.K. 87: 847–870, 1967.

[128298]22944.

[128299]Matsuda, I.; Sugai, M.; Kajii, T.: Ornithine loading test in

[128300]Lowe's syndrome. J. Pediat. 77: 127–129, 1970.

[128301]22945.

[128302]Matsuda, I.; Takeda, T.; Sugai, M.; Matsuura, N.: Oculo-cerebrorenal

[128303]syndrome. Am. J. Dis. Child. 117: 205–212, 1969.

[128304]22946.

[128305]McCance, R. A.; Matheson, W. J.; Gresham, G. A.; Elkinton, J.

[128306]R.: The cerebro-ocular-renal dystrophies: a new variant. Arch. Dis.

[128307]Child. 35: 240–249, 1960.

[128308]22947.

[128309]Monnier, N.; Satre, V.; Lerouge, E.; Berthoin, F.; Lunardi, J.

[128310]: OCRL1 mutation analysis in French Lowe syndrome patients: implications

[128311]for molecular diagnosis strategy and genetic counseling.

Hum. Mutat. 16:

[128312]157–165, 2000.

[128313]22948.

[128314]Mueller, O. T.; Hartsfield, J. K., Jr.; Gallardo, L. A.; Essig,

[128315]Y.-P.; Miller, K. L.; Papenhausen, P. R.; Tedesco, T. A.:

Lowe oculocerebrorenal

[128316]syndrome in a female with a balanced X;20 translocation: mapping of

[128317]the X chromosome breakpoint. Am. J. Hum. Genet. 49:

804–810, 1991.

[128318]22949.

[128319]Nelson, D. L.; Ballabio, A.; Victoria, M. F.; Pieretti, M.; Bies,

[128320]R. D.; Gibbs, R. A.; Maley, J. A.; Chinault, A. C.; Webster, T.

D.;

[128321]Caskey, C. T.: Alu-primed polymerase chain reaction for regional

[128322]assignment of 110 yeast artificial chromosome clones from the human

[128323]X chromosome: identification of clones associated with a disease locus. Proc.

[128324]Nat. Acad. Sci. 88: 6157-6161, 1991.

[128325]22950.

[128326]Oetliker, O.; Rossi, E.: The influence of extracellular fluid

[128327]volume on the renal bicarbonate threshold: a study of two children

[128328]with Lowe's syndrome. Pediat. Res. 3: 140-148, 1969.

[128329]22951.

[128330]Olivos-Glander, I. M.; Janne, P. A.; Nussbaum, R. L.: The oculocerebrorenal

[128331]syndrome gene product is a 105-kD protein localized to the Golgi complex. Am.

[128332]J. Hum. Genet. 57: 817-823, 1995.

[128333]22952.

[128334]Pallisgaard, G.; Goldschmidt, E.: The oculo-cerebro-renal syndrome

[128335] of Lowe in four generations of one family. *Acta Paediat.*

Scand. 60:

[128336] 146–148, 1971.

[128337] 22953.

[128338] Reilly, D. S.; Lewis, R. A.; Ledbetter, D. H.; Nussbaum, R. L.

[128339] Tightly linked flanking markers for the Lowe oculocerebrorenal syndrome,

[128340] with application to carrier assessment. *Am. J. Hum. Genet.* 42: 748–755,

[128341] 1988.

[128342] 22954.

[128343] Reilly, D. S.; Lewis, R. A.; Nussbaum, R. L.: Genetic and physical

[128344] mapping of Xq24–q26 markers flanking the Lowe oculocerebrorenal syndrome. *Genomics* 8:

[128345] 62–70, 1990.

[128346] 22955.

[128347] Richards, W.; Donnell, G. N.; Wilson, W. A.; Stowens, D.; Perry,

[128348] T.: The oculo-cerebro-renal syndrome of Lowe. *Am. J. Dis. Child.* 109:

[128349]185–203, 1965.

[128350]22956.

[128351]Roschinger, W.; Muntau, A. C.; Rudolph, G.; Roscher, A. A.;
Kammerer,

[128352]S.: Carrier assessment in families with Lowe oculocere-
brorenal syndrome:

[128353]novel mutations in the OCRL1 gene and correlation of di-
rect DNA diagnosis

[128354]with ocular examination. Molec. Genet. Metab. 69:
213–222, 2000.

[128355]22957.

[128356]Satre, V.; Monnier, N.; Berthoin, F.; Ayuso, C.; Joannard,
A.;

[128357]Jouk, P.-S.; Lopez-Pajares, I.; Megabarne, A.; Philippe, H.
J.; Plauchu,

[128358]H.; Torres, M. L.; Lunardi, J.: Characterization of a
germline mosaicism

[128359]in families with Lowe syndrome, and identification of
seven novel

[128360]mutations in the OCRL1 gene. Am. J. Hum. Genet. 65:
68–76, 1999.

[128361]22958.

- [128362]Silver, D. N.; Lewis, R. A.; Ledbetter, D. H.; Bobrow, M.;
Nussbaum,
[128363]R. L.: Identification of tightly-linked flanking markers for
Lowe
[128364]syndrome. (Abstract) Cytogenet. Cell Genet. 46: 692 only,
1987.
[128365]22959.
- [128366]Silver, D. N.; Lewis, R. A.; Nussbaum, R. L.: Mapping the
Lowe
[128367]oculocerebrorenal syndrome to Xq24-q26 by use of re-
striction fragment
[128368]length polymorphisms. J. Clin. Invest. 79: 282-285, 1987.
[128369]22960.
- [128370]Streiff, E. B.; Straub, W.; Golay, L.: Les manifestations ocul-
laires
[128371]du syndrome de Lowe. Ophthalmologica 135: 632-639,
1958.
[128372]22961.
- [128373]Suchy, S. F.; Lin, T.; Horwitz, J. A.; O'Brien, W. E.; Nuss-
baum,
[128374]R. L.: First report of prenatal biochemical diagnosis of
Lowe syndrome. Prenatal

[128375]Diag. 18: 1117–1121, 1998.

[128376]22962.

[128377]Svorc, J.; Masopust, J.; Komarkova, A.; Macek, M.; Hyanek, J.

[128378]: Oculocerebrorenal syndrome in a female child. Am. J. Dis. Child. 114:

[128379]186–190, 1967.

[128380]22963.

[128381]Tripathi, R.; Cibis, G. W.; Harris, D. J.; Tripathi, B.: Lowe's

[128382]syndrome. Birth Defects Orig. Art. Ser. 18(6): 629–644, 1982.

[128383]22964.

[128384]Wadelius, C.; Fagerholm, P.; Pettersson, U.; Anneren, G.: Lowe

[128385]oculocerebrorenal syndrome: DNA-based linkage of the gene to Xq24–q26,

[128386]using tightly linked flanking markers and the correlation to lens

[128387]examination in carrier diagnosis. Am. J. Hum. Genet. 44: 241–247,

[128388]1989.

[128389]22965.

[128390]Wilson, W. A.; Richards, W.; Donnell, G. N.: Oculo-cere-bral-renal

[128391]syndrome of Lowe: a review of eight cases noting the ge-netic inheritance. Arch.

[128392]Ophthal. 70: 5-11, 1963.

[128393]22966.

[128394]Witzleben, C. L.; Schoen, E. J.; Tu, W. H.; McDonald, L. W.:

[128395]Progressive morphologic renal changes in the oculo-cerebro-renal syndrome

[128396]of Lowe. Am. J. Med. 44: 319-324, 1968.

[128397]22967.

[128398]Zhang, X.; Jefferson, A. B.; Auethavekiat, V.; Majerus, P. W.

[128399]: The protein deficient in Lowe syndrome is a phos-phatidylinositol-4,5-bisphosphate

[128400]5-phosphatase. Proc. Nat. Acad. Sci. 92: 4853-4856, 1995.

[128401]22968.

[128402]Schram, A. W.; Goldfischer, S.; van Roermund, C. W. T.; Brouwer-Kelder,

[128403]E. M.; Collins, J.; Hashimoto, T.; Heymans, H. S. A.; van
den Bosch,

[128404]H.; Schutgens, R. B. H.; Tager, J. M.; Wanders, R. J. A.: Hu-
man peroxisomal

[128405]3-oxoacyl-coenzyme A thiolase deficiency. Proc. Nat.
Acad. Sci. 84:

[128406]2494-2496, 1987.

[128407]22969.

[128408]Suzuki, Y.; Shimozawa, N.; Yajima, S.; Tomatsu, S.; Kondo,
N.;

[128409]Nakada, Y.; Akaboshi, S.; Iai, M.; Tanabe, Y.; Hashimoto,
T.; Wanders,

[128410]R. J. A.; Schutgens, R. B. H.; Moser, H. W.; Orii, T.: Novel
subtype
[128411]of peroxisomal acyl-CoA oxidase deficiency and bifunc-
tional enzyme
[128412]deficiency with detectable enzyme protein: identification
by means
[128413]of complementation analysis. Am. J. Hum. Genet. 54:
36-43, 1994.
[128414]22970.

[128415]Rosenberg, C. L.; Wong, E.; Petty, E. M.; Bale, A. E.; Tsuji-
moto,
[128416]Y.; Harris, N. L.; Arnold, A.: PRAD1, a candidate BCL1
oncogene:
[128417]mapping and expression in centrocytic lymphoma. Proc.
Nat. Acad.
[128418]Sci. 88: 9638-9642, 1991.
[128419]22971.

[128420]Tsujimoto, Y.; Jaffe, E.; Cossman, J.; Gorham, J.; Nowell, P.
[128421]C.; Croce, C. M.: Clustering of breakpoints on chromo-
some 11 in human
[128422]B-cell neoplasms with the t(11;14) chromosome translo-
cation. Nature 315:

[128423]340–343, 1985.

[128424]22972.

[128425]Tsujiimoto, Y.; Yunis, J.; Onorato–Showe, L.; Erikson, J.;
Nowell,

[128426]P. C.; Croce, C. M.: Molecular cloning of the chromosomal
breakpoint

[128427]of B–cell lymphomas and leukemias with the t(11;14)
chromosome translocation. Science 224:

[128428]1403–1406, 1984.

[128429]22973.

[128430]Barbany, G.; Hoglund, M.; Simonsson, B.: Complete
molecular remission

[128431]in chronic myelogenous leukemia after imatinib therapy.
(Letter) New

[128432]Eng. J. Med. 347: 539–540, 2002.

[128433]22974.

[128434]Budarf, M.; Canaani, E.; Emanuel, B. S.: Linear order of the
four

[128435]BCR–related loci in 22q11. Genomics 3: 168–171, 1988.

[128436]22975.

[128437]Castellanos, A.; Pintado, B.; Weruaga, E.; Arevalo, R.;
Lopez,
[128438]A.; Orfao, A.; Sanchez-Garcia, I.: A BCR-ABL(p190) fusion
gene made
[128439]by homologous recombination causes B-cell acute lym-
phoblastic leukemias
[128440]in chimeric mice with independence of the endogenous
bcr product. Blood 90:
[128441]2168-2174, 1997. 1. Brunning, R. D.: Philadelphia chro-
mosome positive
[128442]leukemia. Hum. Path. 11: 307-309, 1980.
[128443]22976.
[128444]Chisoe, S. L.; Bodenteich, A.; Wang, Y.-F.; Wang, Y.-P.;
Burian,
[128445]D.; Clifton, S. W.; Crabtree, J.; Freeman, A.; Iyer, K.; Jian,
L.;
[128446]Ma, Y.; McLaury, H.-J.; Pan, H.-Q.; Sarhan, O. H.; Toth, S.;
Wang,
[128447]Z.; Zhang, G.; Heisterkamp, N.; Groffen, J.; Roe, B. A.: Se-
quence
[128448]and analysis of the human ABL gene, the BCR gene, and
regions involved

[128449]in the Philadelphia chromosomal translocation. Genomics
27: 67-82,

[128450]1995.

[128451]22977.

[128452]Court Brown, W. M.; Doll, R.: Mortality from cancer and
other

[128453]causes after radiotherapy for ankylosing spondylitis. Brit.
Med.

[128454]J. 2: 1327-1332, 1965.

[128455]22978.

[128456]Croce, C. M.; Huebner, K.; Isobe, M.; Fainstain, E.; Lifshitz,

[128457]B.; Shtivelman, E.; Canaani, E.: Mapping of four distinct
BCR-related

[128458]loci to chromosome region 22q11: order of BCR loci rela-
tive to chronic

[128459]myelogenous leukemia and acute lymphoblastic leukemia
breakpoints. Proc.

[128460]Nat. Acad. Sci. 84: 7174-7178, 1987.

[128461]22979.

[128462]Daley, G. Q.; Van Etten, R. A.; Baltimore, D.: Induction of
chronic

[128463]myelogenous leukemia in mice by the P210(bcr/abl) gene
of the Philadelphia

[128464]chromosome. Science 247: 824–830, 1990.

[128465]22980.

[128466]de Klein, A.; Geurts van Kessel, A.; Grosveld, G.; Bartram,
C.

[128467]R.; Hagemeijer, A.; Bootsma, D.; Spurr, N. K.; Heisterkamp,
N.; Groffen,

[128468]J.; Stephenson, J. R.: A cellular oncogene is translocated to
the

[128469]Philadelphia chromosome in chronic myelocytic
leukaemia. Nature 300:

[128470]765–767, 1982.

[128471]22981.

[128472]Diekmann, D.; Brill, S.; Garrett, M. D.; Totty, N.; Hsuan, J.;

[128473]Monfries, C.; Hall, C.; Lim, L.; Hall, A.: Bcr encodes a GT-
Pase-activating

[128474]protein for p21(rac). Nature 351: 400–402, 1991.

[128475]22982.

[128476]Druker, B. J.; Sawyers, C. L.; Kantarjian, H.; Resta, D. J.;
Reese,

[128477]S. F.; Ford, J. M.; Capdeville, R.; Talpaz, M.: Activity of a specific

[128478]inhibitor of the BCR-ABL tyrosine kinase in the blast crisis of chronic

[128479]myeloid leukemia and acute lymphoblastic leukemia with the Philadelphia

[128480]chromosome. New Eng. J. Med. 344: 1038-1042, 2001.

[128481]22983.

[128482]Druker, B. J.; Talpaz, M.; Resta, D. J.; Peng, B.; Buchdunger,

[128483]E.; Ford, J. M.; Lydon, N. B.; Kantarjian, H.; Capdeville, R.; Ohno-Jones,

[128484]S.; Sawyers, C. L.: Efficacy and safety of a specific inhibitor of

[128485]the BCR-ABL tyrosine kinase in chronic myeloid leukemia. New Eng.

[128486]J. Med. 344: 1031-1037, 2001.

[128487]22984.

[128488]Emanuel, B. S.; Selden, J. R.; Wang, E.; Nowell, P. C.; Croce,

[128489]C. M.: In situ hybridization and translocation breakpoint mapping.

[128490]I. Non-identical 22q11 breakpoints for the t(9;22) of CML
and the

[128491]t(8;22) of Burkitt lymphoma. Cytogenet. Cell Genet. 38:
127-131,

[128492]1984.

[128493]22985.

[128494]Misrahi, M.; Meduri, G.; Pissard, S.; Bouvattier, C.; Beau, I.;

[128495]Loosfelt, H.; Jolivet, A.; Rappaport, R.; Milgrom, E.;
Bougnères,

[128496]P.: Comparison of immunocytochemical and molecular
features with

[128497]the phenotype in a case of incomplete male pseudo-
hermaphroditism associated

[128498]with a mutation of the luteinizing hormone receptor. J.
Clin. Endocr.

[128499]Metab. 82: 2159-2165, 1997.

[128500]22986.

[128501]Perez-Palacios, G.; Scaglia, H. E.; Kofman, S.; Saavedra, D.;

[128502]Ochoa, S.; Laraza, O.; Perez, A. E.: Inherited deficiency of
gonadotropin

[128503]receptor in Leydig cells: a new form of male pseudo-

hermaphroditism.

[128504](Abstract) Am. J. Hum. Genet. 27: 71A, 1975.

[128505]22987.

[128506]Perez-Palacios, G.; Scaglia, H. E.; Kofman-Alfaro, S.;
Saavedra,

[128507]D.; Ochoa, S.; Larraza, O.; Perez, A. E.: Inherited male
pseudohermaphroditism

[128508]due to gonadotrophin unresponsiveness. Acta Endocr. 98:
148-155,

[128509]1981.

[128510]22988.

[128511]Rodien, P.; Cetani, F.; Costagliola, S.; Tonacchera, M.;
Duprez,

[128512]L.; Minegishi, T.; Govaerts, C.; Vassart, G.: Evidences for
an allelic

[128513]variant of the human LC/CG receptor rather than a gene
duplication:

[128514]functional comparison of wild-type and variant receptors.
J. Clin.

[128515]Endocr. Metab. 83: 4431-4434, 1998.

[128516]22989.

[128517]Rousseau-Merck, M. F.; Misrahi, M.; Atger, M.; Loosfelt, H.; Milgrom,

[128518]E.; Berger, R.: Localization of the human luteinizing hormone/choriogonadotropin

[128519]receptor gene (LHCGR) to chromosome 2p21. Cytogenet. Cell Genet. 54:

[128520]77-79, 1990.

[128521]22990.

[128522]Saldanha, P. H.; Arnhold, I. J. P.; Mendonca, B. B.; Bloise, W.;

[128523]Toledo, S. P. A.: A clinico-genetic investigation of Leydig cell

[128524]hypoplasia. Am. J. Med. Genet. 26: 337-344, 1987.

[128525]22991.

[128526]Schwartz, M.; Imperato-McGinley, J.; Peterson, R. E.; Cooper,

[128527]G.; Morris, P. L.; MacGillivray, M.; Hensle, T.: Male pseudo-hermaphroditism

[128528]secondary to an abnormality in Leydig cell differentiation. J. Clin.

[128529]Endocr. Metab. 53: 123-127, 1981.

[128530]22992.

[128531]Segaloff, D. L.; Ascoli, M.: The lutropin/chori-
gonadotropin

[128532]receptor: 4 years later. Endocr. Rev. 14: 324-347, 1993.

[128533]22993.

[128534]Shenker, A.; Laue, L.; Kosugi, S.; Merendino, J. J., Jr.;
Minegishi,

[128535]T.; Cutler, G. B., Jr.: A constitutively activating mutation of
the

[128536]lutinizing hormone receptor in familial male precocious
puberty. Nature 365:

[128537]652-654, 1993.

[128538]22994.

[128539]Rosenthal, I. M.; Refetoff, S.; Rich, B.; Barnes, R. B.; Sun-
thornthepvarakul,

[128540]T.; Parma, J.; Rosenfield, R. L.: Response to challenge with
gonadotropin-releasing

[128541]hormone agonist in a mother and her two sons with a
constitutively

[128542]activating mutation of the luteinizing hormone receptor-
-a clinical

[128543]research center study. J. Clin. Endocr. Metab. 81:
3802–3806, 1996.

[128544]22995.

[128545]Stavrou, S. S.; Zhu, Y.-S.; Cai, L.-Q.; Katz, M. D.; Herrera,
[128546]C.; Defillo-Ricart, M.; Imperato-McGinley, J.: A novel mu-
tation of

[128547]the human luteinizing hormone receptor in 46XY and
46XX sisters. J.

[128548]Clin. Endocr. Metab. 83: 2091–2098, 1998.

[128549]22996.

[128550]Toledo, S. P. A.; Brunner, H. G.; Kraaij, R.; Post, M.; Dahia,
[128551]P. L. M.; Hayashida, C. Y.; Kremer, H.; Themmen, A. P. N.:
An inactivating

[128552]mutation of the luteinizing hormone receptor causes
amenorrhea in

[128553]a 46,XX female. J. Clin. Endocr. Metab. 81: 3850–3854,
1996.

[128554]22997.

[128555]Tsai-Morris, C-H.; Geng, Y.; Buczko, E.; Dufau, M. L.: A
novel

[128556]human luteinizing hormone receptor gene. J. Clin. Endocr.

Metab. 83:

[128557]288–291, 1998.

[128558]22998.

[128559]Wu, S.–M.; Chan, W.–Y.: Male pseudohermaphroditism due to inactivating

[128560]luteinizing hormone receptor mutations. Arch. Med. Res. 30: 495–500,

[128561]1999.

[128562]22999.

[128563]Wu, S.–M.; Hallermeier, K. M.; Laue, L.; Brain, C.; Berry, A.

[128564]C.; Grant, D. B.; Griffin, J. E.; Wilson, J. D.; Cutler, G. B., Jr.;

[128565]Chan, W.–Y.: Inactivation of the luteinizing hormone/chorionic gonadotropin

[128566]receptor by an insertional mutation in Leydig cell hypoplasia. Molec.

[128567]Endocr. 12: 1651–1660, 1998.

[128568]23000.

[128569]Yano, K.; Hidaka, A.; Saji, M.; Polymeropoulos, M. H.; Okuno,

[128570]A.; Kohn, L. D.; Cutler, G. B., Jr.: A sporadic case of male–

limited

[128571]precocious puberty has the same constitutively activating point mutation

[128572]in luteinizing hormone/choriogonadotropin receptor gene as familial

[128573]cases. J. Clin. Endocr. Metab. 79: 1818–1823, 1994.

[128574]23001.

[128575]Yano, K.; Kohn, L. D.; Saji, M.; Okuno, A.; Cutler, G. B., Jr.

[128576]: Phe576 plays an important role in the secondary structure and intracellular

[128577]signaling of the human luteinizing hormone/chorionic gonadotropin

[128578]receptor. J. Clin. Endocr. Metab. 82: 2586–2591, 1997.

[128579]23002.

[128580]Yano, K.; Saji, M.; Hidaka, A.; Moriya, N.; Okuno, A.; Kohn, L.

[128581]D.; Cutler, G. B.: A new constitutively activating point mutation

[128582]in the luteinizing hormone/choriogonadotropin receptor gene in cases

[128583]of male-limited precocious puberty. J. Clin. Endocr. 80:

1162–1168,

[128584]1995.

[128585]23003.

[128586]Zenteno, J. C.; Canto, P.; Kofman–Alfaro, S.; Mendez, J. P.:

[128587]Evidence for genetic heterogeneity in male pseudo–
hermaphroditism due

[128588]to Leydig cell hypoplasia. J. Clin. Endocr. Metab. 84:
3803–3806,

[128589]1999.

[128590]23004.

[128591]Grandori, C.; Mac, J.; Siebelt, F.; Ayer, D. E.; Eisenman, R.
N.

[128592]: Myc–Max heterodimers activate a DEAD box gene and
interact with

[128593]multiple E box–related sites in vivo. EMBO J. 15:
4344–4357, 1996.

[128594]23005.

[128595]Deltas, C. C.: Mutations of the human polycystic kidney
disease

[128596]2 (PKD2) gene. Hum. Mutat. 18: 13–24, 2001.

[128597]23006.

- [128598]Elles, R. G.; Read, A. P.; Hodgkinson, K. A.; Watters, A.; Harris,
R.: Recombination or heterogeneity: is there a second locus for adult polycystic kidney disease? J. Med. Genet. 27: 413–417, 1990.
- [128601]23007.
- [128602]Fossdal, R.; Boovarsson, M.; Asmundsson, P.; Ragnarsson, J.; Peters, D.; Breuning, M. H.; Jensson, O.: Icelandic families with autosomal dominant polycystic kidney disease: families unlinked to chromosome 16p13.3 revealed by linkage analysis. Hum. Genet. 91: 609–613, 1993.
- [128606]23008.
- [128607]Gallagher, A. R.; Cedzich, A.; Gretz, N.; Somlo, S.; Witzgall, R.: The polycystic kidney disease protein PKD2 interacts with Hax-1, a protein associated with the actin cytoskeleton. Proc. Nat. Acad.
- [128610]Sci. 97: 4017–4022, 2000.

[128611]23009.

[128612]Gonzalez-Perrett, S.; Kim, K.; Ibarra, C.; Damiano, A. E.;
Zotta,

[128613]E.; Batelli, M.; Harris, P. C.; Reisin, I. L.; Arnaout, M. A.;
Cantiello,

[128614]H. F.: Polycystin-2, the protein mutated in autosomal
dominant polycystic

[128615]kidney disease (ADPKD), is a Ca^{2+} -permeable nonselective
cation

[128616]channel. Proc. Nat. Acad. Sci. 98: 1182-1187, 2001.

[128617]23010.

[128618]Grantham, J. J.; Calvet, J. P.: Polycystic kidney disease: in

[128619]danger of being X-rated? Proc. Nat. Acad. Sci. 98:
790-792, 2001.

[128620]23011.

[128621]Hanaoka, K.; Qian, F.; Boletta, A.; Bhunia, A. K.; Piontek,
K.;

[128622]Tsiokas, L.; Sukhatme, V. P.; Guggino, W. B.; Germino, G.
G.: Co-assembly

[128623]of polycystin-1 and -2 produces unique cation-permeable
currents. Nature 408:

[128624]990–994, 2000.

[128625]23012.

[128626]Hateboer, N.; van Dijk, M. A.; Bogdanova, N.; Coto, E.;
Saggar–Malik,

[128627]A. K.; San Millan, J. L.; Torra, R.; Breuning, M.; Ravine, D.:
Comparison

[128628]of phenotypes of polycystic kidney disease types 1 and 2.
Lancet 353:

[128629]103–107, 1999.

[128630]23013.

[128631]Hayashi, T.; Mochizuki, T.; Reynolds, D. M.; Wu, G.; Cai,
Y.;

[128632]Somlo, S.: Characterization of the exon structure of the
polycystic

[128633]kidney disease 2 gene (PKD2). Genomics 44: 131–136,
1997.

[128634]23014.

[128635]Jeffery, S.; Saggar–Malik, A. K.; Morgan, S.; MacGregor, G.
A.

[128636]: A family with autosomal dominant polycystic kidney dis-
ease not linked

[128637]to chromosome 16p13.3. Clin. Genet. 44: 173–176, 1993.

[128638]23015.

[128639]Kimberling, W. J.; Fain, P. R.; Kenyon, J. B.; Goldgar, D.;
Sujansky,

[128640]E.; Gabow, P. A.: Linkage heterogeneity of autosomal
dominant polycystic

[128641]kidney disease. New Eng. J. Med. 319: 913–918, 1988.

[128642]23016.

[128643]Kimberling, W. J.; Kumar, S.; Gabow, P. A.; Kenyon, J. B.;
Connolly,

[128644]C. J.; Somlo, S.: Autosomal dominant polycystic kidney
disease: localization

[128645]of the second gene to chromosome 4q13–q23. Genomics
18: 467–472,

[128646]1993.

[128647]23017.

[128648]Koptides, M.; Hadjimichael, C.; Koupepidou, P.; Pierides,
A.;

[128649]Deltas, C. C.: Germinal and somatic mutations in the PKD2
gene of

[128650]renal cysts in autosomal dominant polycystic kidney dis–

ease. Hum.

[128651]Molec. Genet. 8: 509–513, 1999.

[128652]23018.

[128653]Koptides, M.; Mean, R.; Demetriou, K.; Pierides, A.; Deltas, C.

[128654]C.: Genetic evidence for a trans–heterozygous model for cystogenesis

[128655]in autosomal dominant polycystic kidney disease. Hum. Molec. Genet. 9:

[128656]447–452, 2000.

[128657]23019.

[128658]Koulou, P.; Cai, Y.; Geng, L.; Maeda, Y.; Nishimura, S.; Witzgall,

[128659]R.; Ehrlich, B. E.; Somlo, S.: Polycystin–2 is an intracellular calcium

[128660]release channel. Nature Cell Biol. 4: 191–197, 2002.

[128661]23020.

[128662]Kumar, S.; Kimberling, W. J.; Gabow, P. A.; Kenyon, J. B.: Genetic

[128663]linkage studies of autosomal dominant polycystic kidney disease: search

[128664]for the second gene in a large Sicilian family. Hum. Genet.

87:

[128665]129–133, 1991.

[128666]23021.

[128667]Kumar, S.; Kimberling, W. J.; Gabow, P. A.; Shugart, Y. Y.;

Pieke–Dahl,

[128668]S.: Exclusion of autosomal dominant polycystic kidney
disease type

[128669]II (ADPKD2) from 160 cM of chromosome 1. J. Med. Genet.

27: 697–700,

[128670]1990.

[128671]23022.

[128672]Mochizuki, T.; Wu, G.; Hayashi, T.; Xenophontos, S. L.;

Veldhuisen,

[128673]B.; Saris, J. J.; Reynolds, D. M.; Cai, Y.; Gabow, P. A.;

Pierides,

[128674]A.; Kimberling, W. J.; Breuning, M. H.; Constantinou

Deltas, C.; Peters,

[128675]D. J. M.; Somlo, S.: PKD2, a gene for polycystic kidney dis-
ease that

[128676]encodes an integral membrane protein. Science 272:

1339–1342, 1996.

[128677]23023.

[128678]Newby, L. J.; Streets, A. J.; Zhao, Y.; Harris, P. C.; Ward, C.

[128679]J.; Ong, A. C.: Identification, characterization, and localization

[128680]of a novel kidney polycystin-1-polycystin-2 complex. J. Biol. Chem. 277:

[128681]20763-20773, 2002.

[128682]23024.

[128683]Norby, S.; Schwartz, M.: Possible locus for polycystic kidney

[128684]disease on chromosome 2. (Letter) Lancet 336: 323-324, 1990.

[128685]23025.

[128686]Parfrey, P. S. N.; Bear, J. C.; Morgan, J.; Cramer, B. C.; McManamon,

[128687]P. J.; Gault, M. H.; Churchill, D.; Singh, M.; Hewitt, R.; Somlo,

[128688]S.; Reeders, S.: The diagnosis and prognosis of autosomal dominant

[128689]polycystic kidney disease. New Eng. J. Med. 323: 1085-1090, 1990.

[128690]23026.

[128691]Pei, Y.; He, N.; Wang, K.; Kasenda, M.; Paterson, A. D.;
Chan,

[128692]G.; Liang, Y.; Roscoe, J.; Brissenden, J.; Hefferton, D.; Par-
frey,

[128693]P.; Somlo, S.; St. George-Hyslop, P.: A spectrum of muta-
tions in

[128694]the polycystic kidney disease-2 (PKD2) gene from eight
Canadian kindreds. J.

[128695]Am. Soc. Nephrol. 9: 1853-1860, 1998.

[128696]23027.

[128697]Pei, Y.; Paterson, A. D.; Wang, K. R.; He, N.; Hefferton, D.;

[128698]Watnick, T.; Germino, G. G.; Parfrey, P.; Somlo, S.; St.
George-Hyslop,

[128699]P.: Bilineal disease and trans-heterozygotes in autosomal
dominant

[128700]polycystic kidney disease. Am. J. Hum. Genet. 68:
355-363, 2001.

[128701]23028.

[128702]Pei, Y.; Wang, K.; Kasenda, M.; Paterson, A. D.; Liang, Y.;
Huang,

[128703]E.; Lian, J.; Rogovea, E.; Somlo, S.; St. George-Hyslop, P.: A novel

[128704]frameshift mutation induced by an adenosine insertion in the polycystic

[128705]kidney disease 2 (PKD2) gene. *Kidney Int.* 53: 1127–1132, 1998.

[128706]23029.

[128707]Cleutjens, K. B. J. M.; van Eekelen, C. C. E. M.; van der Ko-rput,

[128708]H. A. G. M.; Brinkmann, A. O.; Trapman, J.: Two androgen response

[128709]regions cooperate in steroid hormone regulated activity of the prostate-specific

[128710]antigen promoter. *J. Biol. Chem.* 271: 6379–6388, 1996.

[128711]23030.

[128712]Evans, B. A.; Drinkwater, C. C.; Richards, R. I.: Mouse glandular

[128713]kallikrein genes: structure and partial sequence analysis of the kallikrein

[128714]gene locus. *J. Biol. Chem.* 262: 8027–8034, 1987.

[128715]23031.

- [128716]Lundwall, A.; Lilja, H.: Molecular cloning of human prostate specific antigen cDNA. FEBS Lett. 214: 317–322, 1987.
- [128718]23032.
- [128719]Melegos, D. N.; Yu, H.; Ashok, M.; Wang, C.; Stanczyk, F.; Diamandis, E. P.: Prostate-specific antigen in female serum, a potential new marker of androgen excess. J. Clin. Endocr. Metab. 82: 777–780, 1997.
- [128722]1997.
- [128723]23033.
- [128724]Prendergast, G. C.; Lawe, D.; Ziff, E. B.: Association of Myn, the murine homolog of Max, with c-Myc stimulates methylation-sensitive DNA binding and Ras cotransformation. Cell 65: 395–408, 1991.
- [128727]23034.
- [128728]Wagner, A. J.; Le Beau, M. M.; Diaz, M. O.; Hay, N.: Expression,

[128729]regulation, and chromosomal localization of the Max gene. Proc. Nat.

[128730]Acad. Sci. 89: 3111–3115, 1992.

[128731]23035.

[128732]Zervos, A. S.; Faccio, L.; Gatto, J. P.; Kyriakis, J. M.; Brent,

[128733]R.: Mxi2, a mitogen-activated protein kinase that recognizes and

[128734]phosphorylates Max protein. Proc. Nat. Acad. Sci. 92: 10531–10534,

[128735]1995.

[128736]23036.

[128737]Emi, M.; Katagiri, T.; Harada, Y.; Saito, H.; Inazawa, J.; Ito,

[128738]I.; Kasumi, F.; Nakamura, Y.: A novel metalloprotease/disintegrin-like

[128739]gene at 17q21.3 is somatically rearranged in two primary breast cancers. Nature

[128740]Genet. 5: 151–157, 1993.

[128741]23037.

[128742]Katagiri, T.; Harada, Y.; Emi, M.; Nakamura, Y.: Human metalloprotease/disintegrin-like

[128743](MDC) gene: exon-intron organization and alternative splicing. Cytogenet.

[128744]Cell Genet. 68: 39-44, 1995.

[128745]23038.

[128746]Begley, C. G.; Lipkowitz, S.; Gobel, V.; Mahon, K. A.; Bertness,

[128747]V.; Green, A. R.; Gough, N. M.; Kirsch, I. R.: Molecular characterization

[128748]of NSCL, a gene encoding a helix-loop-helix protein expressed in the

[128749]developing nervous system. Proc. Nat. Acad. Sci. 89: 38-42, 1992.

[128750]23039.

[128751]Brown, L.; Espinosa, R., III; Le Beau, M. M.; Siciliano, M. J.;

[128752]Baer, R.: HEN1 and HEN2: a subgroup of basic helix-loop-helix genes

[128753]that are coexpressed in a human neuroblastoma. Proc. Nat. Acad. Sci. 89:

[128754]8492-8496, 1992.

[128755]23040.

[128756]Cogliati, T.; Good, D. J.; Haigney, M.; Delgado-Romero, P.;

Eckhaus,

[128757]M. A.; Koch, W. J.; Kirsch, I. R.: Predisposition to arrhythmia and

[128758]autonomic dysfunction in Nhlh1-deficient mice. Molec. Cell. Biol. 22:

[128759]4977-4983, 2002.

[128760]23041.

[128761]Lipkowitz, S.; Gobel, V.; Varterasian, M. L.; Nakahara, K.; Tchorz,

[128762]K.; Kirsch, I. R.: A comparative structural characterization of the

[128763]human NSCL-1 and NSCL-2 genes: two basic helix-loop-helix genes expressed

[128764]in the developing nervous system. J. Biol. Chem. 267: 21065-21071,

[128765]1992.

[128766]23042.

[128767]Mullick, A.; Groulx, N.; Trasler, D.; Gros, P.: Nhlh1, a basic
[128768]helix-loop-helix transcription factor, is very tightly linked
to the

[128769]mouse looptail (Lp) mutation. Mammalian Genome 6:

700–704, 1995.

[128770]23043.

[128771]Meeker, T. C.; Nagarajan, L.; ar-Rushdi, A.; Rovera, G.; Huebner,

[128772]K.; Croce, C. M.: Characterization of the human PIM-1 gene: a putative

[128773]proto-oncogene coding for a tissue specific member of the protein

[128774]kinase family. *Oncogene Res.* 1: 87–101, 1987.

[128775]23044.

[128776]Pasqualucci, L.; Neumeister, P.; Goossens, T.; Nanjangud, G.; Chaganti,

[128777]R. S. K.; Kuppers, R.; Dalla-Favera, R.: Hypermutation of multiple

[128778]proto-oncogenes in B-cell diffuse large-cell lymphomas. *Nature* 412:

[128779]341–346, 2001.

[128780]23045.

[128781]Ragoussis, J.; Senger, G.; Mockridge, I.; Sanseau, P.; Ruddy, S.;

[128782]Dudley, K.; Sheer, D.; Trowsdale, J.: A testis-expressed Zn

finger

[128783] gene (ZNF76) in human 6p21.3 centromeric to the MHC is closely linked

[128784] to the human homolog of the t-complex gene tcp-11.

Genomics 14:

[128785] 673-679, 1992.

[128786] 23046.

[128787] Saris, C. J. M.; Domen, J.; Berns, A.: The pim-1 oncogene encodes

[128788] two related protein-serine/threonine kinases by alternative initiation

[128789] at AUG and CUG. EMBO J. 10: 655-664, 1991.

[128790] 23047.

[128791] Selten, G.; Cuypers, H. T.; Boelens, W.; Robanus-Maandag, E.;

[128792] Verbeek, J.; Domen, J.; van Beveren, C.; Berns, A.: The primary structure

[128793] of the putative oncogene pim-1 shows extensive homology with protein

[128794] kinases. Cell 46: 603-611, 1986.

[128795] 23048.

[128796]Zoghbi, H. Y.; Ballantyne, C. M.; O'Brien, W. E.; McCall, A. E.;

[128797]Kwiatkowski, T. J., Jr.; Ledbetter, S. A.; Beaudet, A. L.:
Deletion

[128798]and linkage mapping of eight markers from the proximal
short arm of

[128799]chromosome 6. Genomics 6: 352–357, 1990.

[128800]23049.

[128801]Knebelmann, B.; Forestier, L.; Drouot, L.; Quinones, S.;
Chuet,

[128802]C.; Benessy, F.; Saus, J.; Antignac, C.: Splice-mediated in-
sertion

[128803]of an Alu sequence in the COL4A3 mRNA causing autosomal
recessive

[128804]Alport syndrome. Hum. Molec. Genet. 4: 675–679, 1995.

[128805]23050.

[128806]Gromoll, J.; Partsch, C.-J.; Simoni, M.; Nordhoff, V.; Sippell,

[128807]W. G.; Nieschlag, E.; Saxena, B. B.: A mutation in the first
transmembrane

[128808]domain of the lutropin receptor causes male precocious
puberty. J.

[128809]Clin. Endocr. Metab. 83: 476–480, 1998.

[128810]23051.

[128811]Kawate, N.; Kletter, G. B.; Wilson, B. E.; Netzloff, M. L.;
Menon,

[128812]K. M. J.: Identification of constitutively activating mutation
of

[128813]the luteinising hormone receptor in a family with male
limited gonadotrophin

[128814]independent precocious puberty (testotoxicosis). J. Med.
Genet. 32:

[128815]553–554, 1995.

[128816]23052.

[128817]Kosugi, S.; Van Dop, C.; Geffner, M. E.; Rabl, W.; Carel,
J.–C.;

[128818]Chaussain, J.–L.; Mori, T.; Merendino, J. J., Jr.; Shenker, A.:
Characterization

[128819]of heterogeneous mutations causing constitutive activa-
tion of the

[128820]luteinizing hormone receptor in familial male precocious
puberty. Hum.

[128821]Molec. Genet. 4: 183–188, 1995.

[128822]23053.

[128823]Kremer, H.; Kraaij, R.; Toledo, S. P. A.; Post, M.; Fridman, J.

[128824]B.; Hayashida, C. Y.; van Reen, M.; Milgrom, E.; Ropers,
H.-H.; Mariman,

[128825]E.; Themmen, A. P. N.; Brunner, H. G.: Male pseudo-
hermaphroditism

[128826]due to a homozygous missense mutation of the luteiniz-
ing hormone receptor

[128827]gene. *Nature Genet.* 9: 160-164, 1995.

[128828]23054.

[128829]Kremer, H.; Mariman, E.; Otten, B. J.; Moll, G. W., Jr.;
Stoelinga,

[128830]G. B. A.; Wit, J. M.; Jansen, M.; Drop, S. L.; Faas, B.; Ropers,
H.-H.;

[128831]Brunner, H. G.: Cosegregation of missense mutations of
the luteinizing

[128832]hormone receptor gene with familial male-limited preco-
cious puberty. *Hum.*

[128833]*Molec. Genet.* 2: 1779-1783, 1993.

[128834]23055.

[128835]Kremer, H.; Martens, J. W. M.; van Reen, M.; Verhoef-Post,

M.;

[128836]Wit, J. M.; Otten, B. J.; Drop, S. L. S.; Delemarre-van de
Waal, H.

[128837]A.; Pombo-Arias, M.; De Luca, F.; Potau, N.; Buckler, J. M.
H.; and

[128838]9 others: A limited repertoire of mutations of the luteiniz-
ing hormone

[128839](LH) receptor gene in familial and sporadic patients with
male LH-independent

[128840]precocious puberty. J. Clin. Endocr. Metab. 84:
1136-1140, 1999.

[128841]23056.

[128842]Latronico, A. C.; Abell, A. N.; Arnhold, I. J.; Liu, X.; Lins,

[128843]T. S. S.; Brito, V. N.; Billerbeck, A. E.; Segaloff, D. L.; Men-
donca,

[128844]B. B.: A unique constitutively activating mutation in third
transmembrane

[128845]helix of luteinizing hormone receptor causes sporadic
male gonadotropin-independent

[128846]precocious puberty. J. Clin. Endocr. Metab. 83:
2435-2440, 1998.

[128847]23057.

[128848]Latronico, A. C.; Anasti, J.; Arnhold, I. J. P.; Mendonca, B.

[128849]B.; Domenice, S.; Albano, M. C.; Zachman, K.; Wajchenberg, B. L.;

[128850]Tsigos, C.: A novel mutation of the luteinizing hormone receptor

[128851]gene causing male gonadotropin-independent precocious puberty. J.

[128852]Clin. Endocr. Metab. 80: 2490–2494, 1995.

[128853]23058.

[128854]Latronico, A. C.; Anasti, J.; Arnhold, I. J. P.; Rapaport, R.;

[128855]Mendonca, B. B.; Bloise, W.; Castro, M.; Tsigos, C.; Chrousos, G.

[128856]P.: Testicular and ovarian resistance to luteinizing hormone caused

[128857]by inactivating mutations of the luteinizing hormone-receptor gene. New

[128858]Eng. J. Med. 334: 507–512, 1996.

[128859]23059.

[128860]Laue, L.; Chan, W.-Y.; Hsueh, A. J. W.; Kudo, M.; Hsu, S. Y.;

[128861]Wu, S.-M.; Blomberg, L.; Cutler, G. B., Jr.: Genetic heterogeneity

[128862]of constitutively activating mutations of the human
luteinizing hormone
[128863]receptor in familial male-limited precocious puberty. Proc.
Nat.
[128864]Acad. Sci. 92: 1906-1910, 1995.
[128865]23060.
[128866]Laue, L.; Wu, S.-M.; Kudo, M.; Hsueh, A. J. W.; Cutler, G.
B.,
[128867]Jr.; Griffin, J. E.; Wilson, J. D.; Brain, C.; Berry, A. C.; Grant,
[128868]D. B.; Chan, W.-Y.: A nonsense mutation of the human
luteinizing
[128869]hormone receptor gene in Leydig cell hypoplasia. Hum.
Molec. Genet. 4:
[128870]1429-1433, 1995.
[128871]23061.
[128872]Lee, C. Y.; Ryan, R. J.: The uptake of human luteinizing
hormone
[128873](NLH) by slices of luteinized rat ovaries. Endocrinology 89:
1515-1523,
[128874]1971.
[128875]23062.

[128876]Latronico, A. C.; Chai, Y.; Arnhold, I. J. P.; Liu, X.; Mendonca,
[128877]B. B.; Segaloff, D. L.: A homozygous microdeletion in helix
7 of
[128878]the luteinizing hormone receptor associated with familial
testicular
[128879]and ovarian resistance is due to both decreased cell surface expression
[128880]and impaired effector activation by the cell surface receptor. Molec.
[128881]Endocr. 12: 442-450, 1998.
[128882]23063.
[128883]Latronico, A. C.; Shinozaki, H.; Guerra, G., Jr.; Pereira, M.
[128884]A. A.; Helena, S.; Lemos Marini, V.; Baptista, M. T. M.; Arnhold,
[128885]I. J. P.; Fanelli, F.; Mendonca, B. B.; Segaloff, D. L.: Gonadotropin-independent
[128886]precocious puberty due to luteinizing hormone receptor mutations in
[128887]Brazilian boys: a novel constitutively activating mutation in the
[128888]first transmembrane helix. J. Clin. Endocr. Metab. 85:

4799–4805,

[128889]2000.

[128890]23064.

[128891]Lee, C. Y.; Ryan, R. J.: Luteinizing hormone receptors: specific

[128892]binding of human luteinizing hormone to homogenates of
luteinized

[128893]rat ovaries. Proc. Nat. Acad. Sci. 69: 3520–3523, 1972.

[128894]23065.

[128895]Liu, G.; Duranteau, L.; Carel, J.-C.; Monroe, J.; Doyle, D. A.;

[128896]Shenker, A.: Leydig-cell tumors caused by an activating
mutation

[128897]of the gene encoding the luteinizing hormone receptor.
New Eng. J.

[128898]Med. 341: 1731–1736, 1999.

[128899]23066.

[128900]Loosfelt, H.; Misrahi, M.; Atger, M.; Salesse, R.; Vu Hai-Luu

[128901]Thi, M. T.; Jolivet, A.; Guiochon-Mantel, A.; Sar, S.; Jallal,
B.;

[128902]Garnier, J.; Milgrom, E.: Cloning and sequencing of

porcine LH-hCG

[128903]receptor cDNA: variants lacking transmembrane domain.

Science 245:

[128904]525–528, 1989.

[128905]23067.

[128906]Martens, J. W. M.; Verhoef-Post, M.; Abelin, N.; Erzabella, M.;

[128907]Toledo, S. P. A.; Brunner, H. G.; Themmen, A. P. N.: A homozygous

[128908]mutation in the luteinizing hormone receptor causes partial Leydig

[128909]cell hypoplasia: correlation between receptor activity and phenotype. Molec.

[128910]Endocr. 12: 775–784, 1998.

[128911]23068.

[128912]McFarland, K. C.; Sprengel, R.; Phillips, H. S.; Kohler, M.; Rosemblit,

[128913]N.; Nikolics, K.; Segaloff, D. L.; Seeburg, P. H.: Lutropin-choriogonadotropin

[128914]receptor: an unusual member of the G protein-coupled receptor family. Science 245:

[128915]494–499, 1989.

[128916]23069.

[128917]Min, L.; Ascoli, M.: Effect of activating and inactivating mutations

[128918]on the phosphorylation and trafficking of the human lutropin/choriogonadotropin

[128919]receptor. *Molec. Endocr.* 14: 1797–1810, 2000.

[128920]23070.

[128921]Minegishi, T.; Nakamura, K.; Takakura, Y.; Miyamoto, K.; Hasegawa,

[128922]Y.; Ibuki, Y.; Igarashi, M.: Cloning and sequencing of human LH/hCG

[128923]receptor cDNA. *Biochem. Biophys. Res. Commun.* 172: 1049–1054, 1990.

[128924]23071.

[128925]Chapin, S. J.; Lue, C.-M.; Yu, M. T.; Bulinski, J. C.: Differential

[128926]expression of alternatively spliced forms of MAP4: a repertoire of

[128927]structurally different microtubule-binding domains. *Biochemistry* 34:

[128928]2289–2301, 1995.

[128929]23072.

[128930]Laird, P. W.; van der Lugt, N. M. T.; Clarke, A.; Domen, J.;
Linders,

[128931]K.; McWhir, J.; Berns, A.; Hooper, M.: In vivo analysis of
Pim-1

[128932]deficiency. Nucleic Acids Res. 21: 4750-4755, 1993.

[128933]23073.

[128934]Gartner, J.; Obie, C.; Watkins, P.; Valle, D.: Restoration of

[128935]peroxisome biogenesis in a peroxisome-deficient mam-
malian cell line

[128936]by expression of either the 35 kDa or the 70 kDa peroxi-
somal membrane

[128937]proteins. J. Inherit. Metab. Dis. 17: 327-329, 1994.

[128938]23074.

[128939]Coto, E.; Sanz de Castro, S.; Aguado, S.; Alvarez, J.; Arias,
M.;

[128940]Menendez, M. J.; Lopez-Larrea, C.: DNA microsatellite
analysis of

[128941]families with autosomal dominant polycystic kidney dis-
ease types 1

[128942]and 2: evaluation of clinical heterogeneity between both

forms of

[128943]the disease. J. Med. Genet. 32: 442–445, 1995.

[128944]23075.

[128945]Maric, S. C.; Crozat, A.; Janne, O. A.: Structure and organization

[128946]of the human S-adenosylmethionine decarboxylase gene. J. Biol. Chem. 267:

[128947]18915–18923, 1992.

[128948]23076.

[128949]Maric, S. C.; Crozat, A.; Louhimo, J.; Knuutila, S.; Janne, O.

[128950]A.: The human S-adenosylmethionine decarboxylase gene: nucleotide

[128951]sequence of a pseudogene and chromosomal localization of the active

[128952]gene (AMD1) and the pseudogene (AMD2). Cytogenet. Cell Genet. 70:

[128953]195–199, 1995.

[128954]23077.

[128955]Mangan, M. E.; Olmsted, J. B.: The gene for microtubule-associated

[128956]protein 4 (Mtap4) maps to the distal region of mouse

chromosome 9. Mammalian

[128957]Genome 7: 918–925, 1996.

[128958]23078.

[128959]West, R. R.; Tenbarger, K. M.; Olmsted, J. B.: A model for
microtubule-associated

[128960]protein 4 structure: domains defined by comparisons of
human, mouse,

[128961]and bovine sequences. J. Biol. Chem. 266: 21886–21896,
1991.

[128962]23079.

[128963]Abel, K. J.; Boehnke, M.; Prahalad, M.; Ho, P.; Flejter, W. L.;

[128964]Watkins, M.; VanderStoep, J.; Chandrasekharappa, S. C.;
Collins, F.

[128965]S.; Glover, T. W.; Weber, B. L.: A radiation hybrid map of
the BRCA1

[128966]region of chromosome 17q12–q21. Genomics 17:
632–641, 1993.

[128967]23080.

[128968]Alonso, A. D. C.; Grundke-Iqbal, I.; Iqbal, K.: Alzheimer's
disease

[128969]hyperphosphorylated tau sequesters normal tau into tan–

gles of filaments

[128970]and disassembles microtubules. Nature Med. 2: 783–787,
1996.

[128971]23081.

[128972]Arima, K.; Kowalska, A.; Hasegawa, M.; Mukoyama, M.;
Watanabe,

[128973]R.; Kawai, M.; Takahashi, K.; Iwatsubo, T.; Tabira, T.;
Sunohara,

[128974]N.: Two brothers with frontotemporal dementia and
parkinsonism with

[128975]an N279K mutation of the tau gene. Neurology 54:
1787–1795, 2000.

[128976]23082.

[128977]Baker, M.; Litvan, I.; Houlden, H.; Adamson, J.; Dickson,
D.; Perez–Tur,

[128978]J.; Hardy, J.; Lynch, T.; Bigio, E.; Hutton, M.: Association of
an

[128979]extended haplotype in the tau gene with progressive
supranuclear palsy. Hum.

[128980]Molec. Genet. 8: 711–715, 1999.

[128981]23083.

[128982]Brown, J.; Lantos, P. L.; Roques, P.; Fidani, L.; Rossor, M.

N.

[128983]: Familial dementia with swollen achromatic neurons and corticobasal

[128984]inclusion bodies: a clinical and pathological study. J. Neurol. Sci. 135:

[128985]21–30, 1996.

[128986]23084.

[128987]Clark, L. N.; Poorkaj, P.; Wszolek, Z.; Geschwind, D. H.; Nasreddine,

[128988]Z. S.; Miller, B.; Li, D.; Payami, H.; Awert, F.; Markopoulou, K.;

[128989]Andreadis, A.; D'Souza, I.; Lee, V. M.-Y.; Reed, L.; Trojanowski,

[128990]J. Q.; Zhukareva, V.; Bird, T.; Schellenberg, G.; Wilhelmsen, K. C.

[128991]: Pathogenic implications of mutations in the tau gene in pallido-ponto-nigral

[128992]degeneration and related neurodegenerative disorders linked to chromosome

[128993]17. Proc. Nat. Acad. Sci. 95: 13103–13107, 1998.

[128994]23085.

[128995]Connell, J. W.; Gibb, G. M.; Betts, J. C.; Blackstock, W. P.;

Gallo,

[128996]J.-M.; Lovestone, S.; Hutton, M.; Anderton, B. H.: Effects of
FTDP-17

[128997]mutations on the in vitro phosphorylation of tau by glyco-
gen synthase

[128998]kinase 3-beta identified by mass spectrometry demon-
strate certain

[128999]mutations exert long-range conformational changes. FEBS
Lett. 493:

[129000]40-44, 2001.

[129001]23086.

[129002]Conrad, C.; Andreadis, A.; Trojanowski, J. Q.; Dickson, D.
W.; Kang,

[129003]D.; Chen, X.; Weiderholt, W.; Hansen, L.; Masliah, E.; Thal,
L. J.;

[129004]Katzman, R.; Xia, Y.; Saitoh, T.: Genetic evidence for the
involvement

[129005]of tau in progressive supranuclear palsy. Ann. Neurol. 41:
277-281,

[129006]1997.

[129007]23087.

[129008]Conrad, C.; Vianna, C.; Freeman, M.; Davies, P.: A poly-

morphic

[129009] gene nested within an intron of the tau gene: implications
for Alzheimer's

[129010] disease. Proc. Nat. Acad. Sci. 99: 7751–7756, 2002.

[129011] 23088.

[129012] Delisle, M. B.; Murrell, J. R.; Richardson, R.; Trofatter, J.

[129013] A.; Rascol, O.; Soulages, X.; Mohr, M.; Calvas, P.; Ghetti,
B.: A

[129014] mutation at codon 279 (N279K) in exon 10 of the tau
gene causes a

[129015] tauopathy with dementia and supranuclear palsy. Acta
Neuropath. 98:

[129016] 62–77, 1999.

[129017] 23089.

[129018] Donlon, T. A.; Harris, P.; Neve, R. L.: Localization of mi-
crotubule-associated

[129019] protein tau (MTBT1) to chromosome 17q21. (Abstract) Cy-
togenet. Cell

[129020] Genet. 46: 607, 1987.

[129021] 23090.

[129022] Goedert, M.; Crowther, R. A.; Spillantini, M. G.: Tau muta-

tions

[129023]cause frontotemporal dementias. Neuron 21: 955–958, 1998.

[129024]23091.

[129025]Goedert, M.; Spillantini, M. G.; Potier, M. C.; Ulrich, J.; Crowther,

[129026]R. A.: Cloning and sequencing of the cDNA encoding an isoform of

[129027]microtubule-associated protein tau containing four tandem repeats:

[129028]differential expression of tau protein mRNAs in human brain. EMBO

[129029]J. 8: 393–399, 1989.

[129030]23092.

[129031]Goedert, M.; Wischik, C. M.; Crowther, R. A.; Walker, J. E.; Klug,

[129032]A.: Cloning and sequencing of the cDNA encoding a core protein of

[129033]the paired helical filament of Alzheimer disease: identification as

[129034]the microtubule-associated protein tau. Proc. Nat. Acad. Sci. 85:

[129035]4051–4055, 1988.

[129036]23093.

[129037]Goode, B. L.; Chau, M.; Denis, P. E.; Feinstein, S. C.: Structural

[129038]and functional differences between 3-repeat and 4-repeat tau isoforms:

[129039]implications for normal tau function and the onset of neurodegenerative

[129040]disease. J. Biol. Chem. 275: 38182–38189, 2000.

[129041]23094.

[129042]Heutink, P.: Untangling tau-related dementia. Hum. Molec. Genet. 9:

[129043]979–986, 2000.

[129044]23095.

[129045]Heutink, P.; Stevens, M.; Rizzu, P.; Bakker, E.; Kros, J. M.;

[129046]Tibben, A.; Niermeijer, M. F.; van Duijn, C. M.; Oostra, B. A.; van

[129047]Swieten, J. C.: Hereditary frontotemporal dementia is linked to chromosome

[129048]17q21–q22: a genetic and clinicopathological study of three Dutch

[129049]families. Ann. Neurol. 41: 150–159, 1997.

[129050]23096.

[129051]Hiesberger, T.; Trommsdorff, M.; Howell, B. W.; Goffinet, A.;

[129052]Mumby, M. C.; Cooper, J. A.; Herz, J.: Direct binding of reelin to

[129053]VLDL receptor and apoE receptor 2 induces tyrosine phosphorylation

[129054]of disabled-1 and modulates tau phosphorylation. Neuron 24: 481–489,

[129055]1999.

[129056]23097.

[129057]Hong, M.; Zhukareva, V.; Vogelsberg–Ragaglia, V.; Ws-zolek, Z.;

[129058]Reed, L.; Miller, B. I.; Geschwind, D. H.; Bird, T. D.; McKeel, D.;

[129059]Goate, A.; Morris, J. C.; Wilhelmsen, K. C.; Schellenberg, G. D.;

[129060]Trojanowski, J. Q.; Lee, V. M.–Y.: Mutation-specific functional impairments

[129061]in distinct tau isoforms of hereditary FTDP-17. Science 282: 1914–1917,

[129062]1998.

[129063]23098.

[129064]Hutton, M.: Missense and splice site mutations in tau associated

[129065]with FTDP-17: multiple pathogenic mechanisms. Neurology 56 (suppl.

[129066]4): S21-S25, 2001.

[129067]23099.

[129068]Hutton, M.; Lendon, C. L.; Rizzu, P.; Baker, M.; Froelich, S.;

[129069]Houlden, H.; Pickering-Brown, S.; Chakraverty, S.; Isaacs, A.; Grover,

[129070]A.; Hackett, J.; Adamson, J.; and 39 others: Association of missense

[129071]and 5-prime-splice-site mutations in tau with the inherited dementia

[129072]FTDP-17. Nature 393: 702-705, 1998.

[129073]23100.

[129074]Iijima, M.; Tabira, T.; Poorkaj, P.; Schellenberg, G. D.; Trojanowski,

[129075]J. Q.; Lee, V. M.; Schmidt, M. L.; Takahashi, K.; Nabika, T.; Matsumoto,

[129076]T.; Yamashita, Y.; Yoshioka, S.; Ishino, H.: A distinct familial
 [129077]presenile dementia with a novel missense mutation in the
 tau gene. Neuroreport 10:
 [129078]497–501, 1999.
 [129079]23101.

[129080]Ishihara, T.; Hong, M.; Zhang, B.; Nakagawa, Y.; Lee, M. K.;
 Trojanowski,
 [129081]J. Q.; Lee, V. M.-Y.: Age-dependent emergence and progression of
 [129082]a tauopathy in transgenic mice overexpressing the shortest human tau
 [129083]isoform. Neuron 24: 751–762, 1999.
 [129084]23102.

[129085]Janssen, J. C.; Warrington, E. K.; Morris, H. R.; Lantos, P.;
 [129086]Brown, J.; Revesz, T.; Wood, N.; Khan, M. N.; Cipolotti, L.;
 Fox,
 [129087]N. C.; Rossor, M. N.: Clinical features of frontotemporal
 dementia
 [129088]due to the intronic tau 10 +16 mutation. Neurology 58:
 1161–1168,

[129089]2002.

[129090]23103.

[129091]Braun, T.; Grzeschik, K.-H.; Bober, E.; Arnold, H.-H.: The
MYF

[129092]genes, a group of human muscle determining factors, are
localized

[129093]on different human chromosomes. (Abstract) Cytogenet.
Cell Genet. 51:

[129094]969 only, 1989.

[129095]23104.

[129096]Henry, I.; Puech, A.; Antignac, C.; Couillin, P.; Jeanpierre,
M.;

[129097]Ahnine, L.; Barichard, F.; Boehm, T.; Augereau, P.; Scrable,
H.; Rabbitts,

[129098]T. H.; Rochefort, H.; Cavenee, W.; Junien, C.: Subregional
mapping

[129099]of BWS, CTSD, MYOD1, and a T-ALL breakpoint in 11p15.
(Abstract) Cytogenet.

[129100]Cell Genet. 51: 1013 only, 1989.

[129101]23105.

[129102]Kim, Y.-J.; Noguchi, S.; Hayashi, Y. K.; Tsukahara, T.;

Shimizu,

[129103]T.; Arahata, K.: The product of an oculopharyngeal muscular dystrophy

[129104]gene, poly(A)-binding protein 2, interacts with SKIP and stimulates

[129105]muscle-specific gene expression. Hum. Molec. Genet. 10: 1129-1139,

[129106]2001.

[129107]23106.

[129108]Olson, E. N.: MyoD family: a paradigm for development. Genes

[129109]Dev. 4: 1454-1461, 1990.

[129110]23107.

[129111]Troger, J.; Neyer, S.; Heufler, C.; Huemer, H.; Schmid, E.; Griesser,

[129112]U.; Kralinger, M.; Kremser, B.; Baldissera, I.; Kieselbach, G.: Substance

[129113]P and vasoactive intestinal polypeptide in the streptozotocin-induced

[129114]diabetic rat retina. Invest. Ophthalm. Vis. Sci. 42: 1045-1050, 2001.

[129115]23108.

[129116]Castagliuolo, I.; Riegler, M.; Pasha, A.; Nikulasson, S.; Lu, B.;

[129117]Gerard, C.; Gerard, N. P.; Pothoulakis, C.: Neurokinin-1 (NK-1) receptor

[129118]is required in Clostridium difficile-induced enteritis. J. Clin.

[129119]Invest. 101: 1547-1550, 1998.

[129120]23109.

[129121]De Felipe, C.; Herrero, J. F.; O'Brien, J. A.; Palmer, J. A.; Doyle,

[129122]C. A.; Smith, A. J. H.; Laird, J. M. A.; Belmonte, C.; Cervero, F.;

[129123]Hunt, S. P.: Altered nociception, analgesia and aggression in mice

[129124]lacking the receptor for substance P. Nature 392: 394-397, 1998.

[129125]23110.

[129126]Gerard, N. P.; Garraway, L. A.; Eddy, R. L., Jr.; Shows, T. B.;

[129127]Iijima, H.; Paquet, J. L.; Gerard, C.: Human substance P receptor

[129128](NK-1): organization of the gene, chromosome localization, and functional

[129129]expression of cDNA clones. Biochemistry 30:
10640–10646, 1991.

[129130]23111.

[129131]Hopkins, B.; Powell, S. J.; Danks, P.; Briggs, I.; Graham, A.:

[129132]Isolation and characterisation of the human lung NK-1 re-
ceptor cDNA. Biochem.

[129133]Biophys. Res. Commun. 180: 1110–1117, 1991.

[129134]23112.

[129135]Murtra, P.; Sheasby, A. M.; Hunt, S. P.; De Felipe, C.: Re-
warding

[129136]effects of opiates are absent in mice lacking the receptor
for substance

[129137]P. Nature 45: 180–183, 2000.

[129138]23113.

[129139]Page, N. M.; Woods, R. J.; Gardiner, S. M.; Lomthaisong, K.;
Gladwell,

[129140]R. T.; Butlin, D. J.; Manyonda, I. T.; Lowry, P. J.: Excessive
placental

[129141]secretion of neurokinin B during the third trimester causes
pre-eclampsia. Nature 405:

[129142]797–800, 2000.

[129143]23114.

[129144]Ishikawa, T.; Umesono, K.; Mangelsdorf, D. J.; Aburatani, H.; Stanger,

[129145]B. Z.; Shibasaki, Y.; Imawari, M.; Evans, R. M.; Takaku, F.:
A functional

[129146]retinoic acid receptor encoded by the gene on human
chromosome 12.

[129147]Molec. Endocr. 4: 837-844, 1990.

[129148]23115.

[129149]Krust, A.; Kastner, P.; Petkovich, M.; Zelent, A.; Chambon,
P.

[129150]: A third human retinoic acid receptor, hRAR-gamma.
Proc. Nat. Acad.

[129151]Sci. 86: 5310-5314, 1989.

[129152]23116.

[129153]Mattei, M.-G.; Riviere, M.; Krust, A.; Ingvarsson, S.;
Vennstrom,

[129154]B.; Islam, M. Q.; Levan, G.; Kautner, P.; Zelent, A.; Cham-
bon, P.;

[129155]Szpirer, J.; Szpirer, C.: Chromosomal assignment of
retinoic acid

[129156]receptor (RAR) genes in the human, mouse, and rat
genomes. Genomics 10:

[129157]1061–1069, 1991.

[129158]23117.

[129159]Ryseck, R.-P.; Macdonald-Bravo, H.; Mattei, M. G.;
Siegfried, R.

[129160]L.; Bravo, R.: Structure, mapping and expression of a
growth factor

[129161]inducible gene encoding a putative nuclear hormonal
binding receptor.

[129162]EMBO J. 8: 3327–3335, 1989.

[129163]23118.

[129164]Abramson, D. H.; Ellsworth, R. M.; Zimmerman, L. E.:
Monocular

[129165]cancer in retinoblastoma survivors. Trans. Am. Acad.
Ophthalmol. Otolaryng. 81:

[129166]454–457, 1976.

[129167]23119.

[129168]Aherne, G. E. S.; Roberts, D. F.: Retinoblastoma--a clinical
survey

[129169]and its genetic implications. Clin. Genet. 8: 275–290,

1975.

[129170]23120.

[129171]Alonso, J.; Garcia-Miguel, P.; Abelairas, J.; Mendiola, M.;
Sarret,

[129172]E.; Vendrell, M. T.; Navajas, A.; Pestana, A.: Spectrum of
germline

[129173]RB1 gene mutations in Spanish retinoblastoma patients:
phenotypic

[129174]and molecular epidemiological implications. Hum. Mutat.
17: 412-422,

[129175]2001.

[129176]23121.

[129177]Amoaku, W. M. K.; Willshaw, H. E.; Parkes, S. E.; Shah, K. J.;

[129178]Mann, J. R.: Trilateral retinoblastoma: a report of five pa-
tients. Cancer 78:

[129179]858-863, 1996.

[129180]23122.

[129181]Bader, J. L.; Meadows, A. T.; Zimmerman, L. E.; Rorke, L.
B.; Voute,

[129182]P. A.; Champion, L. A. A.; Miller, R. W.: Bilateral
retinoblastoma

- [129183]with ectopic intracranial retinoblastoma: trilateral retinoblastoma. Cancer
- [129184]Genet. Cytogenet. 5: 203–213, 1982.
- [129185]23123.
- [129186]Balaban–Malenbaum, G.; Gilbert, F.; Nichols, W. W.; Hill, R.; Shields, J.; Meadows, A. T.: A deleted chromosome no. 13 in human retinoblastoma cells: relevance to tumorigenesis. Cancer Genet. Cytogenet. 3: 243–250, 1981.
- [129189]1981.
- [129190]23124.
- [129191]Bandara, L. R.; Adamczewski, J. P.; Hunt, T.; La Thangue, N. B.
- [129192]: Cyclin A and the retinoblastoma gene product complex with a common transcription factor. Nature 352: 249–251, 1991.
- [129194]23125.
- [129195]Benedict, W. F.; Murphree, A. L.; Banerjee, A.; Spina, C. A.; Sparkes, M. C.; Sparkes, R. S.: Patient with 13 chromosome dele-

tion: evidence

[129197] that the retinoblastoma gene is a recessive cancer gene.

Science 219:

[129198] 973–975, 1983.

[129199] 23126.

[129200] Benedict, W. F.; Xu, H.-J.; Hu, S.-X.; Takahashi, R.: Role of

[129201] the retinoblastoma gene in the initiation and progression
of human

[129202] cancer. J. Clin. Invest. 85: 988–993, 1990.

[129203] 23127.

[129204] Bia, B.; Cowell, J. K.: Independent constitutional germline
mutations

[129205] occurring in the RB1 gene in cousins with bilateral
retinoblastoma. Oncogene 11:

[129206] 977–979, 1995.

[129207] 23128.

[129208] Blanquet, V.; Turleau, C.; de Grouchy, J.; Creau-Goldberg,
N.

[129209]: Physical map around the retinoblastoma gene: possible
genomic imprinting

[129210] suggested by Nrul digestion. Genomics 10: 350–355,

1991.

[129211]23129.

[129212]Blanquet, V.; Turleau, C.; Gross-Morand, M. S.; Sena-
maud-Beaufort,

[129213]C.; Doz, F.; Besmond, C.: Spectrum of germline mutations
in the RB1

[129214]gene: a study of 232 patients with hereditary and non
hereditary retinoblastoma. Hum.

[129215]Molec. Genet. 4: 383-388, 1995.

[129216]23130.

[129217]Jiang, Z.; Cote, J.; Kwon, J. M.; Goate, A. M.; Wu, J. Y.:
Aberrant

[129218]splicing of tau pre-mRNA caused by intronic mutations
associated with

[129219]the inherited dementia frontotemporal dementia with
Parkinsonism linked

[129220]to chromosome 17. Molec. Cell. Biol. 20: 4036-4048,
2000.

[129221]23131.

[129222]Magenis, R. E.; Smith, L.; Nadeau, J. H.; Johnson, K. R.;
Mountjoy,

[129223]K. G.; Cone, R. D.: Mapping of the ACTH, MSH, and neural

(MC3 and

[129224]MC4) melanocortin receptors in the mouse and human.

Mammalian Genome 5:

[129225]503–508, 1994.

[129226]23132.

[129227]Futterweit, W.; Ritch, R.; Teekhasaene, C.; Nelson, E. S.:

Coexistence

[129228]of Prader–Willi syndrome, congenital ectropion uveae with
glaucoma,

[129229]and factor XI deficiency. J.A.M.A. 255: 3280–3282, 1986.

[129230]23133.

[129231]Chow, D.; He, X.; Snow, A. L.; Rose–John, S.; Garcia, K. C.:

Structure

[129232]of an extracellular gp130 cytokine receptor signaling
complex. Science 291:

[129233]2150–2155, 2001.

[129234]23134.

[129235]Funatsu, H.; Yamashita, H.; Noma, H.; Mimura, T.; Ya–
mashita, T.;

[129236]Hori, S.: Increased levels of vascular endothelial growth
factor

[129237]and interleukin-6 in the aqueous humor of diabetics with
macular edema. Am.

[129238]J. Ophthalm. 133: 70-77, 2002.

[129239]23135.

[129240]Kelsell, D. P.; Risk, J. M.; Leigh, I. M.; Stevens, H. P.; Ellis,

[129241]A.; Hennies, H. C.; Reis, A.; Weissenbach, J.; Bishop, D. T.;
Spurr,

[129242]N. K.; Field, J. K.: Close mapping of the focal non-
epidermolytic

[129243]palmoplantar keratoderma (PPK) locus associated with oe-
sophageal cancer

[129244](TOC). Hum. Molec. Genet. 5: 857-860, 1996.

[129245]23136.

[129246]McLean, W. H. I.: Personal Communication. Philadelphia,
Pa.

[129247]2/26/1997.

[129248]23137.

[129249]McLean, W. H. I.; Rugg, E. L.; Lunny, D. P.; Morley, S. M.;
Lane,

[129250]E. B.; Swensson, O.; Dopping-Hepenstal, P. J. C.; Griffiths,
W. A.

- [129251]D.; Eady, R. A. J.; Higgins, C.; Navsaria, H. A.; Leigh, I. M.;
Strachan,
- [129252]T.; Kunkeler, L.; Munro, C. S.: Keratin 16 and keratin 17
mutations
- [129253]cause pachyonychia congenita. *Nature Genet.* 9: 273–278,
1995.
- [129254]23138.
- [129255]Shamsheer, M. K.; Navsaria, H. A.; Stevens, H. P.; Ratnavel,
R.
- [129256]C.; Purkis, P. E.; McLean, W. H.; Cook, L. J.; Griffiths, W. A.
D.;
- [129257]Geschmeissner, S.; Spurr, N.; Leigh, I. M.: Novel mutations
in keratin
- [129258]16 gene underly focal nonepidermolytic palmoplantar
keratoderma (NEPPK)
- [129259]in two families. *Hum. Mol. Genet.* 4: 1875–1881, 1995.
- [129260]23139.
- [129261]Smith, F. J. D.; Del Monaco, M.; Steijlen, P. M.; Munro, C.
S.;
- [129262]Morvay, M.; Coleman, C. M.; Rietveld, F. J. R.; Uitto, J.;
McLean,
- [129263]W. H. I.: Novel proline substitution mutations in keratin 16

in two

[129264]cases of pachyonychia congenita type 1. Brit. J. Derm.

141: 1010–1016,

[129265]1999.

[129266]23140.

[129267]Smith, F. J. D.; McKusick, V. A.; Nielsen, K.; Pfendner, E.;

Uitto,

[129268]J.; McLean, W. H. I.: Cloning of multiple keratin 16 genes

facilitates

[129269]prenatal diagnosis of pachyonychia congenita type 1. Pre-

natal Diag. 19:

[129270]941–946, 1999.

[129271]23141.

[129272]Stevens, H. P.; Kelsell, D. P.; Spurr, N. K.; Bishop, D. T.;

Purkis,

[129273]P. E.; Griffiths, W. A. D.; Rustin, M. H. A.; Leigh, I. M.: Ker-

atin

[129274]staining and linkage of non-epidermolytic focal palmo-

plantar keratodermas

[129275](PPK) to 17q. (Abstract) Brit. J. Derm. 131: 425, 1994.

[129276]23142.

- [129277] Terrinoni, A.; Puddu, P.; Didona, B.; De Laurenzi, V.;
Candi, E.;
- [129278] Smith, F. J. D.; McLean, W. H. I.; Melino, G.: A mutation in
the
- [129279] V1 domain of K16 is responsible for unilateral palmoplan-
tar verrucous
- [129280] nevus. J. Invest. Derm. 114: 1136–1140, 2000.
- [129281] 23143.
- [129282] Lewis, J.; McGowan, E.; Rockwood, J.; Melrose, H.;
Nacharaju,
- [129283] P.; Van Slegtenhorst, M.; Gwinn-Hardy, K.; Murphy, M. P.;
Baker, M.;
- [129284] Yu, X.; Duff, K.; Hardy, J.; Corral, A.; Lin, W.-L.; Yen, S.-H.;
Dickson,
- [129285] D. W.; Davies, P.; Hutton, M.: Neurofibrillary tangles, amy-
otrophy
- [129286] and progressive motor disturbance in mice expressing
mutant (P301L)
- [129287] tau protein. Nature Genet. 25: 402–405, 2000. Note: Erra-
tum: Nature
- [129288] Genet. 26: 127 only, 2000.
- [129289] 23144.

[129290]Lippa, C. F.; Zhukareva, V.; Kawarai, T.; Uryu, K.; Shafiq, M.;

[129291]Nee, L. E.; Grafman, J.; Liang, Y.; St George-Hyslop, P. H.; Trojanowski,

[129292]J. Q.; Lee, V. M.-Y.: Frontotemporal dementia with novel tau pathology

[129293]and a glu342val tau mutation. *Ann. Neurol.* 48: 850–858, 2000.

[129294]23145.

[129295]Litvan, I.; Baker, M.; Hutton, M.: Tau genotype: no effect on

[129296]onset, symptom severity, or survival in progressive supranuclear palsy. *Neurology* 57:

[129297]138–140, 2001.

[129298]23146.

[129299]Martin, E. R.; Scott, W. K.; Nance, M. A.; Watts, R. L.; Hubble,

[129300]J. P.; Koller, W. C.; Lyons, K.; Pahwa, R.; Stern, M. B.; Colcher,

[129301]A.; Hiner, B. C.; Jankovic, J.; and 20 others: Association of single-nucleotide

[129302]polymorphisms of the Tau gene with late-onset Parkinson

disease. J.A.M.A. 286:

[129303]2245–2250, 2001.

[129304]23147.

[129305]Moser, A. B.; Rasmussen, M.; Naidu, S.; Watkins, P. A.;
McGuinness,

[129306]M.; Hajra, A. K.; Chen, G.; Raymond, G.; Liu, A.; Gordon,
D.; Garnaas,

[129307]K.; Walton, D. S.; Skjedal, O. H.; Guggenheim, M. A.; Jack-
son, L.

[129308]G.; Elias, E. R.; Moser, H. W.: Phenotype of patients with
peroxisomal

[129309]disorders subdivided into sixteen complementation
groups. J. Pediat. 127:

[129310]13–22, 1995.

[129311]23148.

[129312]Paton, B. C.; Heron, S. E.; Nelson, P. V.; Morris, C. P.; Pou-
los,

[129313]A.: Absence of mutations raises doubts about the role of
the 70-kD

[129314]peroxisomal membrane protein in Zellweger syndrome.
(Letter) Am.

[129315]J. Hum. Genet. 60: 1535–1539, 1997.

[129316]23149.

[129317]Aljada, A.; Ghanim, H.; Friedman, J.; Garg, R.; Mohanty, P.;
Dandona,

[129318]P.: Troglitazone reduces the expression of PPAR-gamma
while stimulating

[129319]that of PPAR-alpha in mononuclear cells in obese sub-
jects. J. Clin.

[129320]Endocr. Metab. 86: 3130-3133, 2001.

[129321]23150.

[129322]Costet, P.; Legendre, C.; More, J.; Edgar, A.; Galtier, P.;
Pineau,

[129323]T.: Peroxisome proliferator-activated receptor alpha-
isoform deficiency

[129324]leads to progressive dyslipidemia with sexually dimorphic
obesity

[129325]and steatosis. J. Biol. Chem. 273: 29577-29585, 1998.

[129326]23151.

[129327]Djouadi, F.; Weinheimer, C. J.; Saffitz, J. E.; Pitchford, C.;

[129328]Bastin, J.; Gonzalez, F. J.; Kelly, D. P.: A gender-related
defect

[129329]in lipid metabolism and glucose homeostasis in peroxi-

some proliferator-activated

[129330]receptor alpha-deficient mice. J. Clin. Invest. 102:

1083-1091,

[129331]1998.

[129332]23152.

[129333]Kersten, S.; Desvergne, B.; Wahli, W.: Roles of PPARs in
health

[129334]and disease. Nature 405: 421-424, 2000.

[129335]23153.

[129336]Kersten, S.; Seydoux, J.; Peters, J. M.; Gonzalez, F. J.;
Desvergne,

[129337]B.; Wahli, W.: Peroxisome proliferator-activated receptor
alpha mediates

[129338]the adaptive response to fasting. J. Clin. Invest. 103:
1489-1498,

[129339]1999.

[129340]23154.

[129341]Lee, S. S.-T.; Pineau, T.; Drago, J.; Lee, E. J.; Owens, J. W.;

[129342]Kroetz, D. L.; Fernandez-Salguero, P. M.; Westphal, H.;

Gonzalez,

[129343]F. J.: Targeted disruption of the alpha isoform of the per-

oxisome

[129344]proliferator-activated receptor gene in mice results in
abolishment

[129345]of the pleiotropic effects of peroxisome proliferators.

Molec. Cell.

[129346]Biol. 15: 3012-3022, 1995.

[129347]23155.

[129348]Lee, Y.; Yu, X.; Gonzales, F.; Mangelsdorf, D. J.; Wang,
M.-Y.;

[129349]Richardson, C.; Witters, L. A.; Unger, R. H.: PPAR-alpha is
necessary

[129350]for the lipopenic action of hyperleptinemia on white adi-
pose and liver

[129351]tissue. Proc. Nat. Acad. Sci. 99: 11848-11853, 2002.

[129352]23156.

[129353]Leone, T. C.; Weinheimer, C. J.; Kelly, D. P.: A critical role

[129354]for the peroxisome proliferator-activated receptor alpha
(PPAR-alpha)

[129355]in the cellular fasting response: the PPAR-alpha-null
mouse as a model

[129356]of fatty acid oxidation disorders. Proc. Nat. Acad. Sci. 96:

7473–7478,

[129357]1999.

[129358]23157.

[129359]Michalik, L.; Desvergne, B.; Tan, N. S.; Basu–Modak, S.; Escher,

[129360]P.; Rieusset, J.; Peters, J. M.; Kaya, G.; Gonzalez, F. J.; Zakany,

[129361]J.; Metzger, D.; Chambon, P.; Duboule, D.; Wahli, W.: Impaired skin

[129362]wound healing in peroxisome proliferator–activated receptor (PPAR)–alpha

[129363]and PPAR–beta mutant mice. J. Cell Biol. 154: 799–814, 2001.

[129364]23158.

[129365]Sher, T.; Yi, H.–F.; McBride, O. W.; Gonzalez, F. J.: cDNA cloning,

[129366]chromosomal mapping, and functional characterization of the human

[129367]peroxisome proliferator activated receptor. Biochemistry 32: 5598–5604,

[129368]1993.

[129369]23159.

[129370] Xu, H. E.; Stanley, T. B.; Montana, V. G.; Lambert, M. H.;
Shearer,

[129371] B. G.; Cobb, J. E.; McKee, D. D.; Galardi, C. M.; Plunket, K.
D.;

[129372] Nolte, R. T.; Parks, D. J.; Moore, J. T.; Kliewer, S. A.; Will-
son,

[129373] T. M.; Stimmel, J. B.: Structural basis for antagonist-mediated recruitment

[129374] of nuclear co-repressors by PPAR-alpha. Nature 415:
813-817, 2002.

[129375] 23160.

[129376] Cameron, P.; Limjuco, G.; Rodkey, J.; Bennett, C.; Schmidt,
J.

[129377] A.: Amino acid sequence analysis of human interleukin 1
(IL-1): evidence

[129378] for biochemically distinct forms of IL-1. J. Exp. Med. 162:
790-801,

[129379] 1985.

[129380] 23161.

[129381] D'Eustachio, P.; Jadidi, S.; Fuhlbrigge, R. C.; Gray, P. W.;
Chaplin,

[129382] D. D.: Interleukin-1 alpha and beta genes: linkage on

chromosome

[129383]2 in the mouse. Immunogenetics 26: 339-343, 1987.

[129384]23162.

[129385]Dinarello, C. A.: An update on human interleukin-1: from molecular

[129386]biology to clinical relevance. J. Clin. Immun. 5: 287-297, 1985.

[129387]23163.

[129388]El-Omar, E. M.; Carrington, M.; Chow, W.-H.; McColl, K. E. L.;

[129389]Bream, J. H.; Young, H. A.; Herrera, J.; Lissowska, J.; Yuan, C.-C.;

[129390]Rothman, N.; Lanyon, G.; Martin, M.; Fraumeni, J. F., Jr.; Rabkin,

[129391]C. S.: Interleukin-1 polymorphisms associated with increased risk

[129392]of gastric cancer. Nature 404: 398-402, 2000. Note: Erratum: Nature

[129393]412: 99 only, 2001.

[129394]23164.

[129395]Furutani, Y.; Notake, M.; Yamayoshi, M.; Yamagishi, J.;

Nomura,

[129396]H.; Ohue, M.; Furuta, R.; Fukui, T.; Yamada, M.; Nakamura,
S.: Cloning

[129397]and characterization of the cDNAs for human and rabbit
interleukin-1

[129398]precursor. Nucleic Acids Res. 13: 5869-5882, 1985.

[129399]23165.

[129400]Hamajima, N.; Matsuo, K.; Saito, T.; Tajima, K.; Okuma, K.;
Yamao,

[129401]K.; Tominaga, S.: Interleukin 1 polymorphisms, lifestyle
factors,

[129402]and helicobacter pylori infection. Jpn. J. Cancer Res. 92:
383-389,

[129403]2001.

[129404]23166.

[129405]Le Beau, M. M.; Rowley, J. D.: Personal Communication.
Chicago,

[129406]Ill. 6/18/1986.

[129407]23167.

[129408]March, C. J.; Mosley, B.; Larsen, A.; Cerretti, D. P.; Braedt,

[129409]G.; Price, V.; Gillis, S.; Henney, C. S.; Kronheim, S. R.;

Grabstein,

[129410]K.; Conlon, P. J.; Hopp, T. P.; Cosman, D.: Cloning, sequence and

[129411]expression of two distinct human interleukin-1 complementary DNAs. *Nature* 315:

[129412]641-647, 1985.

[129413]23168.

[129414]Nicklin, M. J. H.; Weith, A.; Duff, G. W.: A physical map of

[129415]the region encompassing the human interleukin-1-alpha, interleukin-1-beta,

[129416]and interleukin-1 receptor antagonist genes. *Genomics* 19: 382-384,

[129417]1994.

[129418]23169.

[129419]Samad, T. A.; Moore, K. A.; Sapirstein, A.; Billet, S.; Allchorne,

[129420]A.; Poole, S.; Bonventre, J. V.; Woolf, C. J.: Interleukin-1-beta-mediated

[129421]induction of Cox-2 in the CNS contributes to inflammatory pain hypersensitivity. *Nature* 410:

[129422]471-475, 2001.

[129423]23170.

[129424]Vidal-Vanaclocha, F.; Fantuzzi, G.; Mendoza, L.; Fuentes,
A. M.;

[129425]Anasagasti, M. J.; Martin, J.; Carrascal, T.; Walsh, P.;
Reznikov,

[129426]L. L.; Kim, S.-H.; Novick, D.; Rubinstein, M.; Dinarello, C.
A.:

[129427]IL-18 regulates IL-1-beta-dependent hepatic melanoma
metastasis via

[129428]vascular cell adhesion molecule-1. Proc. Nat. Acad. Sci.
97: 734-739,

[129429]2000.

[129430]23171.

[129431]Webb, A.; Collins, K.; Auron, P.; Eddy, R.; Nakai, H.; Byers,

[129432]M.; Shows, T. B.: Genetics of acute phase response: a gene
for interleukin-1

[129433]is on chromosome 2. (Abstract) Am. J. Hum. Genet. 37:
A142 only,

[129434]1985.

[129435]23172.

[129436]Webb, A. C.; Collins, K. L.; Auron, P. E.; Eddy, R. L.; Nakai,

[129437]H.; Byers, M.; Shows, T. B.: The gene for interleukin-1 (IL1)
is

[129438]on human chromosome 2. (Abstract) Cytogenet. Cell
Genet. 40: 774

[129439]only, 1985.

[129440]23173.

[129441]Webb, A. C.; Collins, K. L.; Auron, P. E.; Eddy, R. L.; Nakai,

[129442]H.; Byers, M. G.; Haley, L. L.; Henry, W. M.; Shows, T. B.:
Interleukin-1

[129443]gene (IL1) assigned to long arm of human chromosome 2.

Lymphokine

[129444]Res. 5: 77–85, 1986.

[129445]23174.

[129446]Ansano, M.; Toda, M.; Sakaguchi, N.; Sakaguchi, S.: Autoimmune

[129447]disease as a consequence of developmental abnormality of a T cell

[129448]subpopulation. J. Exp. Med. 184: 387–396, 1996.

[129449]23175.

[129450]Cosman, D.; Cerretti, D. P.; Larsen, A.; Park, L.; March, C.; Dower,

[129451]S.; Gillis, S.; Urdal, D.: Cloning, sequence and expression of human

[129452]interleukin–2 receptor. Nature 312: 768–771, 1984.

[129453]23176.

[129454]Ferrari, S.; Cannizzaro, L. A.; Battini, R.; Huebner, K.; Baserga,

[129455]R.: The gene encoding human vimentin is located on the short arm

[129456]of chromosome 10. Am. J. Hum. Genet. 41: 616–626, 1987.

[129457]23177.

[129458]Greene, W. C.; Leonard, W. J.; Depper, J. M.; Nelson, D. L.;
Waldmann,

[129459]T. A.: The human interleukin-2 receptor: normal and ab-
normal expression

[129460]in T cells and in leukemias induced by the human T-
lymphotropic retroviruses. Ann.

[129461]Intern. Med. 105: 560-572, 1986.

[129462]23178.

[129463]Hatakeyama, M.; Minamoto, S.; Taniguchi, T.: Intracyto-
plasmic

[129464]phosphorylation sites of Tac antigen (p55) are not essen-
tial for the

[129465]conformation, function, and regulation of the human in-
terleukin 2

[129466]receptor. Proc. Nat. Acad. Sci. 83: 9650-9654, 1986.

[129467]23179.

[129468]Hatakeyama, M.; Minamoto, S.; Uchiyama, T.; Hardy, R. R.;
Yamada,

[129469]G.; Taniguchi, T.: Reconstitution of functional receptor for
human

[129470]interleukin-2 in mouse cells. Nature 318: 467-470, 1985.

[129471]23180.

[129472]Ihle, J. N.; Kerr, I. M.: Jaks and Stats in signaling by the cytokine

[129473]receptor superfamily. Trends Genet. 11: 69-74, 1995.

[129474]23181.

[129475]Ishida, N.; Kanamori, H.; Noma, T.; Nikaido, T.; Sabe, H.; Suzuki,

[129476]N.; Shimizu, A.; Honjo, T.: Molecular cloning and structure of the

[129477]human interleukin 2 receptor gene. Nucleic Acids Res. 13: 7579-7589,

[129478]1985.

[129479]23182.

[129480]Kondo, S.; Shimizu, A.; Maeda, M.; Tagaya, Y.; Yodoi, J.; Honjo,

[129481]T.: Expression of functional human interleukin-2 receptor in mouse

[129482]T cells by cDNA transfection. Nature 320: 75-77, 1986.

[129483]23183.

[129484]Leonard, W. J.; Depper, J. M.; Crabtree, G. R.; Rudikoff, S.;

[129485]Pumphrey, J.; Robb, R. J.; Kronke, M.; Svetlik, P. B.; Pepper,
N.

[129486]J.; Waldmann, T. A.; Greene, W. C.: Molecular cloning and
expression

[129487]of cDNAs for the human interleukin-2 receptor. Nature
311: 626-631,

[129488]1984.

[129489]23184.

[129490]Lekanne Deprez, R. H.; Riegman, P. H. J.; Groen, N. A.;
Warringa,

[129491]U. L.; van Biezen, N. A.; Molijn, A. C.; Bootsma, D.; de
Jong, P.

[129492]J.; Menon, A. G.; Kley, N. A.; Seizinger, B. R.; Zwarthoff, E.
C.

[129493]: Cloning and characterization of MN1, a gene from chro-
mosome 22q11,

[129494]which is disrupted by a balanced translocation in a
meningioma. Oncogene 10:

[129495]1521-1528, 1995.

[129496]23185.

[129497]Dahia, P. L. M.; Aguiar, R. C. T.; Alberta, J.; Kum, J. B.;
 Caron,
 [129498]S.; Sill, H.; Marsh, D. J.; Ritz, J.; Freedman, A.; Stiles, C.;
 Eng,
 [129499]C.: PTEN is inversely correlated with the cell survival factor
 Akt/PKB
 [129500]and is inactivated via multiple mechanisms in haemato-
 logical malignancies. Hum.
 [129501]Molec. Genet. 8: 185-193, 1999.
 [129502]23186.
 [129503]Bryson, G. L. M.; Massa, H.; Trask, B. J.; Van Etten, R. L.:
 Gene
 [129504]structure, sequence, and chromosomal localization of the
 human red
 [129505]cell-type low molecular weight acid phosphotyrosyl phos-
 phatase gene,
 [129506]ACP1. Genomics 30: 133-140, 1995.
 [129507]23187.
 [129508]Chen, T.-R.; McMorris, F. A.; Creagan, R.; Ricciuti, F. C.;
 Tischfield,
 [129509]J.; Ruddle, F. H.: Assignment of the genes for malate oxi-
 doreductase

[129510]decarboxylating to chromosome 6 and peptidase B and lactate dehydrogenase

[129511]B to chromosome 12 in man. Am. J. Hum. Genet. 25: 200–207, 1973.

[129512]23188.

[129513]Hamerton, J. L.; Mohandas, T.; McAlpine, P. J.; Douglas, G. R.

[129514]; Localization of human gene loci using spontaneous chromosome rearrangements

[129515]in human–Chinese hamster somatic cell hybrids. Am. J. Hum. Genet. 27:

[129516]595–608, 1975.

[129517]23189.

[129518]Herbschleb–Voogt, E.; Meera Khan, P.: Defining the locus of origin

[129519]of a genetically determined electrophoretic variant of a multilocus

[129520]enzyme system; the Calcutta–1 of human LDH system is a B–locus variant. Hum.

[129521]Genet. 57: 290–295, 1981.

[129522]23190.

- [129523] Houki, N.; Matsushima, Y.; Kitamura, M.; Tukada, T.;
Nishina, T.;
- [129524] Nakayama, T.: A case of deficiency of lactate dehydroge-
nase H-subunit. Jpn.
- [129525] J. Clin. Chem. 15: 85-90, 1986.
- [129526] 23191.
- [129527] Kitamura, M.; Iijima, N.; Hashimoto, F.; Hiratsuka, A.:
Hereditary
- [129528] deficiency of subunit H of lactate dehydrogenase. Clin.
Chim. Acta 34:
- [129529] 419-423, 1971.
- [129530] 23192.
- [129531] Maekawa, M.; Sudo, K.; Kitajima, M.; Matsuura, Y.; Li, S.
S.-L.;
- [129532] Kanno, T.: Analysis of a genetic mutation in an elec-
trophoretic variant
- [129533] of the human lactate dehydrogenase-B(H) subunit. Hum.
Genet. 91:
- [129534] 423-426, 1993.
- [129535] 23193.
- [129536] Maekawa, M.; Sudo, K.; Nagura, K.; Li, S. S.-L.; Kanno, T.:

Population

- [129537]screening of lactate dehydrogenase deficiencies in
Fukuoka Prefecture
- [129538]in Japan and molecular characterization of three independent mutations
- [129539]in the lactate dehydrogenase-B(H) gene. Hum. Genet. 93:
74-76, 1994.
- [129540]23194.
- [129541]Malpuech, G.; Kaplan, J. C.; Rethore, M. O.; Junien, C.;
Geneix,
- [129542]A.: Une observation de deletion partielle du bras court du
chromosome
- [129543]12: localisation du gene de la lacticodeshydrogenase B.
Lyon Med. 233:
- [129544]275-279, 1975.
- [129545]23195.
- [129546]Mayeda, K.; Weiss, L.; Lindahl, R.; Dully, M.: Localization of
- [129547]the human lactate dehydrogenase B gene on the short arm
of chromosome
- [129548]12. Am. J. Hum. Genet. 26: 59-64, 1974.
- [129549]23196.

- [129550]Miwa, S.; Nishima, T.; Kanehashi, Y.; Kitamura, M.; Hiratsuka,
A.; Shizume, K.: Studies on erythrocyte metabolism in a
case with
hereditary deficiency of H-subunit of lactate dehydroge-
nase. Acta
Haemat. Jpn. 34: 228-232, 1971.
23197.
- [129555]Bell, D. W.; Varley, J. M.; Szydlo, T. E.; Kang, D. H.; Wahrer,
D. C. R.; Shannon, K. E.; Lubratovich, M.; Verselis, S. J.; Is-
selbacher,
K. J.; Fraumeni, J. F.; Birch, J. M.; Li, F. P.; Garber, J. E.;
Haber,
D. A.: Heterozygous germ line hCHK2 mutations in Li-
Fraumeni syndrome. Science 286:
2528-2531, 1999.
23198.
- [129561]Birch, J. M.; Hartley, A. L.; Tricker, K. J.; Prosser, J.; Condie,
A.; Kelsey, A. M.; Harris, M.; Jones, P. H. M.; Binchy, A.;
Crowther,
D.; Craft, A. W.; Eden, O. B.; Evans, D. G. R.; Thompson, E.;

Mann,

[129564]J. R.; Martin, J.; Mitchell, E. L. D.; Santibanez-Koref, M. F.:

Prevalence

[129565]and diversity of constitutional mutations in the p53 gene
among 21

[129566]Li-Fraumeni families. Cancer Res. 54: 1298-1304, 1994.

[129567]23199.

[129568]Goy, A.; Gilles, F.; Remache, Y.; Zelenetz, A. D.: Physical
linkage

[129569]of the lysyl oxidase-like (LOXL1) gene to the PML gene on
human chromosome

[129570]15q22. Cytogenet. Cell Genet. 88: 22-24, 2000.

[129571]23200.

[129572]Kenyon, K.; Modi, W. S.; Contente, S.; Friedman, R. M.: A
novel

[129573]human cDNA with a predicted protein similar to lysyl ox-
idase maps

[129574]to chromosome 15q24-q25. J. Biol. Chem. 268:
18435-18437, 1993.

[129575]23201.

[129576]Kim, Y.; Boyd, C. D.; Csiszar, K.: A new gene with se-

quence and

[129577] structural similarity to the gene encoding human lysyl oxidase. J.

[129578] Biol. Chem. 270: 7176–7182, 1995.

[129579] 23202.

[129580] Wydner, K. S.; Kim, Y.; Csiszar, K.; Boyd, C. D.; Passmore, H.

[129581] C.: An intron capture strategy used to identify and map a lysyl oxidase-like

[129582] gene on chromosome 9 in the mouse. Genomics 40: 342–345, 1997.

[129583] 23203.

[129584] Schmidt, B.; Servidei, S.; Gabbai, A. A.; Silva, A. C.; de Sousa

[129585] Bulle de Oliveira, A.; DiMauro, S.: McArdle's disease in two generations:

[129586] autosomal recessive transmission with manifesting heterozygote. Neurology 37:

[129587] 1558–1561, 1987.

[129588] 23204.

[129589] Arch, E. M.; Goodman, B. K.; Van Wesep, R. A.; Liaw, D.;

Clarke,

[129590]K.; Parsons, R.; McKusick, V. A.; Geraghty, M. T.: Deletion
of PTEN

[129591]in a patient with Bannayan–Riley–Ruvalcaba syndrome
suggests allelism

[129592]with Cowden disease. Am. J. Med. Genet. 71: 489–493,
1997.

[129593]23205.

[129594]Celebi, J. T.; Tsou, H. C.; Chen, F. F.; Zhang, H.; Ping, X. L.;

[129595]Lebwohl, M. G.; Kezis, J.; Peacocke, M.: Phenotypic find-
ings of Cowden

[129596]syndrome and Bannayan–Zonana syndrome in a family as-
sociated with

[129597]a single germline mutation in PTEN. J. Med. Genet. 36:
360–364,

[129598]1999.

[129599]23206.

[129600]Dudek, H.; Datta, S. R.; Franke, T. F.; Birnbaum, M. J.; Yao,
R.;

[129601]Cooper, G. M.; Segal, R. A.; Kaplan, D. R.; Greenberg, M.
E.: Regulation

[129602]of neuronal survival by the serine–threonine protein ki-

nase Akt. Science 275:

[129603]661–663, 1997.

[129604]23207.

[129605]Franke, T. F.; Kaplan, D. R.; Cantley, L. C.; Toker, A.: Direct

[129606]regulation of the Akt proto-oncogene product by phosphatidylinositol-3,4-bisphosphate. Science 275:

[129607]665–667, 1997.

[129608]23208.

[129609]Franke, T. F.; Yang, S.-I.; Chan, T. O.; Datta, K.; Kazanietz, A.; Morrison, D. K.; Kaplan, D. R.; Tsichlis, P. N.: The protein

[129610]kinase encoded by the Akt proto-oncogene is a target of the PDGF-activated

[129611]phosphatidylinositol 3-kinase. Cell 81: 727–736, 1995.

[129612]23209.

[129613]Furnari, F. B.; Huang, H. J.; Cavenee, W. K.: The phospho-

[129614]inositol phosphatase activity of PTEN mediates a serum-sensitive G1 growth

[129615]arrest in glioma cells. Cancer Res. 58: 5002–5008, 1998.

[129616]

[129617]23210.

[129618]Hajdуч, E.; Litherland, G. J.; Hundal, H. S.: Protein kinase

[129619]B (PKB/Akt)--a key regulator of glucose transport? FEBS
Lett. 492:

[129620]199-203, 2001.

[129621]23211.

[129622]Hemmings, B. A.: Akt signaling: linking membrane events
to life

[129623]and death decisions. Science 275: 628-630, 1997.

[129624]23212.

[129625]Jones, P. F.; Jakubowicz, T.; Pitossi, F. J.; Maurer, F.; Hem-
mings,

[129626]B. A.: Molecular cloning and identification of a serine/
threonine

[129627]protein kinase of the second-messenger subfamily. Proc.
Nat. Acad.

[129628]Sci. 88: 4171-4175, 1991.

[129629]23213.

[129630]Lee, M.-J.; Thangada, S.; Paik, J.-H.; Sapkota, G. P.; An-
cellin,

- [129631]N.; Chae, S.-S.; Wu, M.; Morales-Ruiz, M.; Sessa, W. C.; Alessi, D.
- [129632]R.; Hla, T.: Akt-mediated phosphorylation of the G protein-coupled
- [129633]receptor EDG-1 is required for endothelial cell chemotaxis. *Molec.*
- [129634]*Cell* 8: 693-704, 2001.
- [129635]23214.
- [129636]Maira, S.-M.; Galetic, I.; Brazil, D. P.; Kaeck, S.; Ingley, E.;
- [129637]Thelen, M.; Hemmings, B. A.: Carboxyl-terminal modulator protein
- [129638](CTMP), a negative regulator of PKB/Akt and v-Akt at the plasma membrane. *Scien*
- [129639]*ce* 294: 374-380, 2001.
- [129640]23215.
- [129641]Powell, D. W.; Rane, M. J.; Chen, Q.; Singh, S.; McLeish, K. R.
- [129642]: Identification of 14-3-3-zeta as a protein kinase B/Akt substrate. *J.*
- [129643]*Biol. Chem.* 277: 21639-21642, 2002.
- [129644]23216.

[129645]Staal, S. P.: Molecular cloning of the akt oncogene and its human

[129646]homologues AKT1 and AKT2: amplification of AKT1 in a primary human

[129647]gastric adenocarcinoma. Proc. Nat. Acad. Sci. 84: 5034–5037, 1987.

[129648]23217.

[129649]Staal, S. P.; Huebner, K.; Croce, C. M.; Parsa, N. Z.; Testa,

[129650]J. R.: The AKT1 proto-oncogene maps to human chromosome 14, band

[129651]q32. Genomics 2: 96–98, 1988.

[129652]23218.

[129653]Vanhaesebroeck, B.; Alessi, D. R.: The PI3K–PDK1 connection:

[129654]more than just a road to PKB. Biochem. J. 346: 561–576, 2000.

[129655]23219.

[129656]Weng, L.–P.; Brown, J. L.; Eng, C.: PTEN induces apoptosis and

[129657]cell cycle arrest through phosphoinositol–3–kinase/Akt–dependent and

[129658]—independent pathways. Hum. Molec. Genet. 10:
237–242, 2001.

[129659]23220.

[129660]Yang, J.; Cron, P.; Thompson, V.; Good, V. M.; Hess, D.;
Hemmings,

[129661]B. A.; Barford, D.: Molecular mechanism for the regulation
of protein

[129662]kinase B/Akt by hydrophobic motif phosphorylation.
Molec. Cell 9:

[129663]1227–1240, 2002.

[129664]23221.

[129665]Le Beau, M. M.; Rowley, J. D.; Sacchi, N.; Watson, D. K.; Pa-
pas,

[129666]T. S.; Diaz, M. O.: Hu-ets-2 is translocated to chromo-
some 8 in the

[129667]t(8;21) in acute myelogenous leukemia. Cancer Genet. Cy-
togenet. 23:

[129668]269–274, 1986.

[129669]23222.

[129670]Mavrothalassitis, G. J.; Watson, D. K.; Papas, T. S.: Molecu-
lar

- [129671]and functional characterization of the promoter of ETS2,
the human
- [129672]c-ets-2 gene. Proc. Nat. Acad. Sci. 87: 1047-1051, 1990.
- [129673]23223.
- [129674]Sacchi, N.; Cheng, S. V.; Tanzi, R. E.; Gusella, J. F.;
Drabkin,
- [129675]H. A.; Patterson, D.; Haines, J. H.; Papas, T. S.: The ETS
genes
- [129676]on chromosome 21 are distal to the breakpoint of the
acute myelogenous
- [129677]leukemia translocation (8;21). Genomics 3: 110-116,
1988.
- [129678]23224.
- [129679]Sumarsono, S. H.; Wilson, T. J.; Tymms, M. J.; Venter, D. J.;
Corrick,
- [129680]C. M.; Kola, R.; Lahoud, M. H.; Papas, T. S.; Seth, A.; Kola,
I.:
- [129681]Down's syndrome-like skeletal abnormalities in Ets2
transgenic mice. Nature 379:
- [129682]534-540, 1996.
- [129683]23225.

- [129684]Chu, E. H. Y.; Chang, C. C.; Sun, N. C.: Synteny of the human
man
- [129685]genes for gal-1-PT, ACP-1, MDH-1, and gal-
plus-activator and assignment
- [129686]to chromosome 2. Birth Defects Orig. Art. Ser. XI(3):
103-106, 1975.
- [129687]Note: Alternate: Cytogenet. Cell Genet. 14: 273-276,
1975...
- [129688]23226.
- [129689]Rasmussen, S. K.; Lautier, C.; Hansen, L.; Echwald, S. M.;
Hansen,
- [129690]T.; Ekstrom, C. T.; Urhammer, S. A.; Borch-Johnsen, K.;
Grigorescu,
- [129691]F.; Smith, R. J.; Pedersen, O.: Studies of the variability of
the
- [129692]genes encoding the insulin-like growth factor I receptor
and its ligand
- [129693]in relation to type 2 diabetes mellitus. J. Clin. Endocr.
Metab. 85:
- [129694]1606-1610, 2000.
- [129695]23227.
- [129696]Dissing, J.; Johnsen, A. H.: Human red cell acid phos-

phatase (ACP1):

[129697]the primary structure of the two pairs of isozymes encoded by the

[129698]ACP1*A and ACP1*C alleles. Biochim. Biophys. Acta 1121: 261-268,

[129699]1992.

[129700]23228.

[129701]Dissing, J.; Johnsen, A. H.; Sensabaugh, G. F.: Human red cell

[129702]acid phosphatase (ACP1): the amino acid sequence of the two isozymes

[129703]Bf and Bs encoded by the ACP1*B allele. J. Biol. Chem. 266: 20619-20625,

[129704]1991.

[129705]23229.

[129706]Emanuel, B. S.; Zackai, E. H.; Van Dyke, D. C.; Swallow, D. M.;

[129707]Allen, F. H.; Mellman, W. J.: Deletion mapping: further evidence

[129708]for the location of acid phosphatase (ACP-1) within 2p23. Am. J.

[129709]Med. Genet. 4: 167-172, 1979.

[129710]23230.

[129711]Ferguson-Smith, M. A.; Newman, B. F.; Ellis, P. M.; Thomson, D.

[129712]M. G.; Riley, I. D.: Assignment by deletion of human red cell acid

[129713]phosphatase gene locus to the short arm of chromosome 2. Nature 243:

[129714]271-273, 1973.

[129715]23231.

[129716]Fisher, R. A.; Harris, H.: Studies on the separate isoenzymes

[129717]of red cell acid phosphatase phenotypes A and B: chromatographic separation

[129718]of the isoenzymes. Ann. Hum. Genet. 34: 431-438, 1971.

[129719]23232.

[129720]Fuhrmann, W.; Lichte, K. H.: Human red cell acid phosphatase

[129721]polymorphism: a study on gene frequency and forensic use of the system

[129722]in cases of disputed paternity. Humangenetik 3: 121-126, 1966.

[129723]23233.

[129724]Giblett, E. R.; Scott, N. M.: Red cell acid phosphatase:
racial

[129725]distribution and report of a new phenotype. Am. J. Hum.
Genet. 17:

[129726]425-432, 1965.

[129727]23234.

[129728]Herbich, J.; Fisher, R. A.; Hopkinson, D. A.: Atypical segre-
gation

[129729]of human red cell acid phosphatase phenotypes: evidence
for a rare

[129730]'silent' allele P(O). Ann. Hum. Genet. 34: 145-152, 1970.

[129731]23235.

[129732]Herbich, J.; Meinhardt, K.: The rare 'silent' allele P(O) or P(V)

[129733](P Vienna) of human red cell acid phosphatase, typed in a
second family. Humangenetik 15:

[129734]345-348, 1972.

[129735]23236.

[129736]Hopkinson, D. A.; Spencer, N.; Harris, H.: Red cell acid
phosphatase

[129737]variants: a new human polymorphism. Nature 199:
969-971, 1963.

[129738]23237.

[129739]Hulten, M.; Lindsten, J.; Pen-Ming, L. M.; Fraccaro, M.;
Mannini,

[129740]A.; Tiepolo, L.; Robson, E. B.; Heiken, A.; Tillinger, K. G.:
Possible

[129741]localization of the genes for the Kidd blood group on an
autosome

[129742]involved in a reciprocal translocation. Nature 211:
1067-1068, 1966.

[129743]23238.

[129744]Junien, C.; Kaplan, J.-C.; Bernheim, A.; Berger, R.: Regional

[129745]assignment of red cell acid phosphatase locus to band
2p25. Hum.

[129746]Genet. 48: 17-21, 1979.

[129747]23239.

[129748]Karp, G. W., Jr.; Sutton, H. E.: Some new phenotypes of
human

[129749]red cell acid phosphatase. Am. J. Hum. Genet. 19: 54-62,
1967.

[129750]23240.

[129751]Lothe, R. A.; Gedde-Dahl, T.; Olaisen, B.; Bakker, E.; Pearson,
son,

[129752]P.: Very close linkage between D2S1 and ACP1 on chromosome 2p. Ann.

[129753]Hum. Genet. 50: 361-367, 1986.

[129754]23241.

[129755]Mace, M. A.; Cook, P. J. L.; Robson, E. B.: Linkage data on red

[129756]cell acid phosphatase from family studies. Ann. Hum. Genet. 38:

[129757]471-477, 1975.

[129758]23242.

[129759]Mace, M. A.; Robson, E. B.: Linkage data on ACP-1 and MNSs. Cytogenet.

[129760]Cell Genet. 13: 123-125, 1974.

[129761]23243.

[129762]Magenis, R. E.; Koler, R. D.; Lovrien, E. W.; Bigley, R. H.; Duval,

[129763]M. C.; Overton, K. M.: Gene dosage: evidence for assignment of erythrocyte

[129764]acid phosphatase locus to chromosome 2. Proc. Nat.
Acad. Sci. 72:

[129765]4526–4530, 1975.

[129766]23244.

[129767]Mayr, W. R.: No close linkage between MNSs and red cell
acid

[129768]phosphatase. Hum. Hered. 26: 1–3, 1976.

[129769]23245.

[129770]Miller, S. A.; Nelson, M. S.; Dykes, D. D.; Polesky, H. F.:
Comparison

[129771]of acid phosphatase ACP1 variants by isoelectric focusing
and conventional

[129772]electrophoresis: identification of three new alleles,
ACP1*N, ACP1*P

[129773]and ACP1*S. Hum. Hered. 37: 371–375, 1987.

[129774]23246.

[129775]Mohrenweiser, H. W.; Novotny, J. E.: ACP-1-GUA-1: a low-
activity

[129776]variant of human erythrocyte acid phosphatase: associa-
tion with increased

[129777]glutathione reductase activity. Am. J. Hum. Genet. 34:

425-433,

[129778]1982.

[129779]23247.

[129780]Nezbeda, P.: Occurrence of the ACP-1 null allele in
Czechoslovakia. Hum.

[129781]Genet. 46: 227-229, 1979.

[129782]23248.

[129783]Palmarino, R.; Agostino, R.; Gloria, F.; Lucarelli, P.; Bus-
inco,

[129784]L.; Antognoni, G.; Maggioni, G.; Workman, P. L.; Bottini, E.:
Red

[129785]cell acid phosphatase: another polymorphism correlated
with malaria? Am.

[129786]J. Phys. Anthropol. 43: 177-186, 1975.

[129787]23249.

[129788]Povey, S.; Swallow, D. M.; Bobrow, M.; Craig, I.; Van
Heyningen,

[129789]V.: Probable assignment of the locus determining human
red cell acid

[129790]phosphatase ACP(1) to chromosome 2 using somatic cell
hybrids. Ann.

[129791]Hum. Genet. 38: 1-5, 1974.

[129792]23250.

[129793]Renwick, J. H.: Assignment and map-positioning of human loci

[129794]using chromosomal variation. Ann. Hum. Genet. 35: 79-97, 1971.

[129795]23251.

[129796]Sensabaugh, G. F.; Golden, V. L.: Phenotype dependence in the

[129797]inhibition of red cell acid phosphatase (ACP) by folates. Am. J.

[129798]Hum. Genet. 30: 553-560, 1978.

[129799]23252.

[129800]Swallow, D. M.; Povey, S.; Harris, H.: Activity of the 'red cell'

[129801]acid phosphatase locus in other tissues. Ann. Hum. Genet. 37: 31-38,

[129802]1973.

[129803]23253.

[129804]Wu, R. M.; Cheng, C. W.; Chen, K. H.; Lu, S. L.; Shan, D. E.;

- [129805]Ho, Y. F.; Chern, H. D.: The COMT L allele modifies the association
society
- [129806]between MAOB polymorphism and PD in Taiwanese. Neurology 56: 375–382,
2001.
- [129807]23254.
- [129808]Leonard, W. J.; Depper, J. M.; Kanehisa, M.; Kronke, M.;
Peffer,
N. J.; Svetlik, P. B.; Sullivan, M.; Greene, W. C.: Structure of
the human interleukin–2 receptor gene. Science 230:
633–639, 1985.
- [129812]23255.
- [129813]Leonard, W. J.; Depper, J. M.; Robb, R. J.; Waldmann, T. A.;
Greene,
W. C.: Characterization of the human receptor for T-cell
growth factor. Proc.
Nat. Acad. Sci. 80: 6957–6961, 1983.
- [129816]23256.
- [129817]Marx, J. L.: The interleukin–2 receptor gene is cloned. Science 226:
1064–1065, 1985.

[129819]23257.

[129820]Nikaido, T.; Shimizu, A.; Ishida, N.; Sabe, H.; Teshigawara, K.;

[129821]Maeda, M.; Uchiyama, T.; Yodoi, J.; Honjo, T.: Molecular cloning

[129822]of a cDNA encoding human interleukin-2 receptor. Nature 311: 631-635,

[129823]1984.

[129824]23258.

[129825]Robb, R. J.; Rusk, C. M.; Neeper, M. P.: Structure-function relationships

[129826]for the interleukin 2 receptor: location of ligand and anti-body binding

[129827]sites on the Tac receptor chain by mutational analysis. Proc. Nat.

[129828]Acad. Sci. 85: 5654-5658, 1988.

[129829]23259.

[129830]Sharfe, N.; Dadi, H. K.; Shahar, M.; Roifman, C. M.: Human immune

[129831]disorder arising from mutation of the alpha chain of the interleukin-2

[129832]receptor. Proc. Nat. Acad. Sci. 94: 3168–3171, 1997.

[129833]23260.

[129834]Shevach, E. M.: Certified professionals: CD4(+)CD25(+) suppressor

[129835]T cells. J. Exp. Med. 193: F41–F45, 2001.

[129836]23261.

[129837]Tsudo, M.; Kozak, R. W.; Goldman, C. K.; Waldmann, T. A.: Demonstration

[129838]of a non-Tac peptide that binds interleukin 2: a potential participant

[129839]in a multichain interleukin 2 receptor complex. Proc. Nat. Acad.

[129840]Sci. 83: 9694–9698, 1986.

[129841]23262.

[129842]Urdal, D. L.; March, C. J.; Gillis, S.; Larsen, A.; Dower, S.

[129843]K.: Purification and chemical characterization of the receptor for

[129844]interleukin 2 from activated human T lymphocytes and from a human

[129845]T-cell lymphoma cell line. Proc. Nat. Acad. Sci. 81: 6481–6485,

[129846]1984.

[129847]23263.

[129848]Mohrenweiser, H. W.; Neel, J. V.: Frequency of thermostability

[129849]variants: estimation of total 'rare' variant frequency in human populations. Proc.

[129850]Nat. Acad. Sci. 78: 5729-5733, 1981.

[129851]23264.

[129852]Farrer, L. A.; Conneally, P. M.: A genetic model for age at onset

[129853]in Huntington disease. Am. J. Hum. Genet. 37: 350-357, 1985.

[129854]23265.

[129855]Farrer, L. A.; Conneally, P. M.; Yu, P.: The natural history

[129856]of Huntington disease: possible role of 'aging genes.'. Am. J. Med.

[129857]Genet. 18: 115-123, 1984.

[129858]23266.

[129859]Farrer, L. A.; Cupples, L. A.; Wiater, P.; Conneally, P. M.; Gusella,

[129860]J. F.; Myers, R. H.: The normal Huntington disease (HD) allele, or
[129861]a closely linked gene, influences age at onset of HD. Am.
J. Hum.
[129862]Genet. 53: 125–130, 1993.
[129863]23267.
[129864]Ferrante, R. J.; Kowall, N. W.; Beal, M. F.; Richardson, E. P.,
[129865]Jr.; Bird, E. D.; Martin, J. B.: Selective sparing of a class of
[129866]striatal neurons in Huntington's disease. Science 230:
561–563,
[129867]1985.
[129868]23268.
[129869]Folstein, S.; Abbott, M.; Moser, R.; Parhad, I.; Clark, A.;
Folstein,
[129870]M.: A phenocopy of Huntington's disease: lacunar infarcts
of the
[129871]corpus striatum. Johns Hopkins Med. J. 148: 104–108,
1981.
[129872]23269.
[129873]Folstein, S. E.; Abbott, M. H.; Franz, M. L.; Huang, S.;
Chase,

- [129874]G. A.; Folstein, M. F.: Phenotypic heterogeneity in Huntington disease. J. Neurogenet. 1: 175–184, 1984.
- [129875]23270.
- [129876]23270.
- [129877]Folstein, S. E.; Chase, G. A.; Wahl, W. E.; McDonnell, A. M.;
- [129878]Folstein, M. F.: Huntington disease in Maryland: clinical aspects of racial variation. Am. J. Hum. Genet. 41: 168–179, 1987.
- [129880]23271.
- [129881]Folstein, S. E.; Phillips, J. A., III; Meyers, D. A.; Chase, G. A.; Abbott, M. H.; Franz, M. L.; Waber, P. G.; Kazazian, H. H., Jr.;
- [129882]Conneally, P. M.; Hobbs, W.; Tanzi, R.; Faryniarz, A.; Gibbons, K.;
- [129883]Gusella, J.: Huntington's disease: two families with differing clinical features show linkage to the G8 probe. Science 229: 776–779, 1985.
- [129884]23272.
- [129885]Freeman, T. B.; Cicchetti, F.; Hauser, R. A.; Deacon, T. W.; Li,

- [129888]X.-J.; Hersch, S. M.; Nauert, G. M.; Sanberg, P. R.; Kordower, J.
- [129889]H.; Saporta, S.; Isacson, O.: Transplanted fetal striatum in Huntington's
- [129890]disease: phenotypic development and lack of pathology. Proc. Nat.
- [129891]Acad. Sci. 97: 13877-13882, 2000.
- [129892]23273.
- [129893]Froster-Iskenius, U. G.; Hayden, M. R.; Wang, H. S.; Kalousek,
- [129894]D. K.; Horsman, D.; Pfeiffer, R. A.; Schottky, A.; Schwinger, E.:
- [129895]A family with Huntington disease and reciprocal translocation 4;5. Am.
- [129896]J. Hum. Genet. 38: 759-767, 1986.
- [129897]23274.
- [129898]Gellera, C.; Meoni, C.; Castellotti, B.; Zappacosta, B.; Girotti,
- [129899]F.; Taroni, F.; DiDonato, S.: Errors in Huntington disease diagnostic
- [129900]test caused by trinucleotide deletion in the IT15 gene. (Letter) Am.

[129901]J. Hum. Genet. 59: 475–477, 1996.

[129902]23275.

[129903]Gervais, F. G.; Singaraja, R.; Xanthoudakis, S.; Gutekunst, C.-A.;

[129904]Leavitt, B. R.; Metzler, M.; Hackam, A. S.; Tam, J.; Vaillancourt,

[129905]J. P.; Houtzager, V.; Rasper, D. M.; Roy, S.; Hayden, M. R.; Nicholson,

[129906]D. W.: Recruitment and activation of caspase-8 by the huntingtin-interacting

[129907]protein Hip-1 and a novel partner Hippi. Nature Cell Biol. 4: 95–105,

[129908]2002.

[129909]23276.

[129910]Gilliam, T. C.; Bucan, M.; MacDonald, M. E.; Zimmer, M.; Haines,

[129911]J. L.; Cheng, S. V.; Pohl, T. M.; Meyers, R. H.; Whaley, W. L.; Allitto,

[129912]B. A.; Faryniarz, A.; Wasmuth, J. J.; Frischauf, A.-M.; Conneally,

[129913]P. M.; Lehrach, H.; Gusella, J. F.: A DNA segment encoding two genes

[129914]very tightly linked to Huntington's disease. Science 238:
950-952,

[129915]1987.

[129916]23277.

[129917]Gilliam, T. C.; Tanzi, R. E.; Haines, J. L.; Bonner, T. I.; Fary-
niarz,

[129918]A. G.; Hobbs, W. J.; MacDonald, M. E.; Cheng, S. V.; Fol-
stein, S.

[129919]E.; Conneally, P. M.; Wexler, N. S.; Gusella, J. F.: Localiza-
tion

[129920]of the Huntington's disease gene to a small segment of
chromosome

[129921]4 flanked by D4S10 and the telomere. Cell 50: 565-571,
1987.

[129922]23278.

[129923]Giordani, B.; Berent, S.; Boivin, M. J.; Penney, J. B.; Lehti-
nen,

[129924]S.; Markel, D. S.; Hollingsworth, Z.; Butterbaugh, G.;
Hichwa, R.

[129925]D.; Gusella, J. F.; Young, A. B.: Longitudinal neuropsychol-
ogical

[129926]and genetic linkage analysis of persons at risk for Hunt-

ington's disease. Arch.

[129927]Neurol. 52: 59-64, 1995.

[129928]23279.

[129929]Goetz, I.; Roberts, E.; Comings, D. E.: Fibroblasts in Huntington's

[129930]disease. New Eng. J. Med. 293: 1225-1227, 1975.

[129931]23280.

[129932]Goldberg, Y. P.; Andrew, S. E.; Theilmann, J.; Kremer, B.; Squitieri,

[129933]F.; Telenius, H.; Brown, J. D.; Hayden, M. R.: Familial predisposition

[129934]to recurrent mutations causing Huntington's disease: genetic risk

[129935]to sibs of sporadic cases. J. Med. Genet. 30: 987-990, 1993.

[129936]23281.

[129937]Goldberg, Y. P.; Kalchman, M. A.; Metzler, M.; Nasir, J.; Zeisler,

[129938]J.; Graham, R.; Koide, H. B.; O'Kusky, J.; Sharp, A. H.; Ross, C.

[129939]A.; Jirik, F.; Hayden, M. R.: Absence of disease phenotype

and intergenerational

[129940]stability of the CAG repeat in transgenic mice expressing the human

[129941]Huntington disease transcript. Hum. Molec. Genet. 5: 177-185, 1996.

[129942]23282.

[129943]Goldberg, Y. P.; Nicholson, D. W.; Rasper, D. M.; Kalchman, M.

[129944]A.; Koide, H. B.; Graham, R. K.; Bromm, M.; Kazemi-Esfarjani, P.;

[129945]Thornberry, N. A.; Vaillancourt, J. P.; Hayden, M. R.: Cleavage of

[129946]huntingtin by apopain, a proapoptotic cysteine protease, is modulated

[129947]by the polyglutamine tract. Nature Genet. 13: 442-449, 1996.

[129948]23283.

[129949]Goldberg, Y. P.; Rommens, J. M.; Andrew, S. E.; Hutchinson, G.

[129950]B.; Lin, B.; Theilmann, J.; Graham, R.; Graves, M. L.; Starr, E.;

[129951]McDonald, H.; Nasir, J.; Schappert, K.; Kalchman, M. A.;

Clarke, L.

[129952]A.; Hayden, M. R.: Identification of an Alu retrotransposi-
tion event

[129953]in close proximity to a strong candidate gene for Hunting-
ton's disease. Nature 362:

[129954]370-373, 1993.

[129955]23284.

[129956]Green, H.: Human genetic diseases due to codon reitera-
tion: relationship

[129957]to an evolutionary mechanism. Cell 74: 955-956, 1993.

[129958]23285.

[129959]Greenberg, L. J.; Martell, R. W.; Theilman, J.; Hayden, M.
R.;

[129960]Joubert, J.: Genetic linkage between Huntington disease
and the D4S10

[129961]locus in South African families: further evidence against
non-allelic

[129962]heterogeneity. Hum. Genet. 87: 701-708, 1991.

[129963]23286.

[129964]Gu, M.; Gash, M. T.; Mann, V. M.; Javoy-Agid, F.; Cooper, J.
M.;

[129965]Schapira, A. H. V.: Mitochondrial defect in Huntington's disease

[129966]caudate nucleus. Ann. Neurol. 39: 385–389, 1996.

[129967]23287.

[129968]Gusella, J.; Tanzi, R. E.; Bader, P. I.; Phelan, M. C.; Stevenson,

[129969]R.; Hayden, M. R.; Hofman, K. J.; Faryniarz, A. G.; Gibbons, K.:

[129970]Deletion of Huntington's disease linked G8 (D4S10) locus in Wolf–Hirschhorn

[129971]syndrome. Nature 318: 75–78, 1985.

[129972]23288.

[129973]Gusella, J. F.; Gibbons, K.; Hobbs, W.; Heft, R.; Anderson, M.;

[129974]Rashtchian, R.; Folstein, S.; Wallace, P.; Conneally, P. M.; Tanzi,

[129975]R.: The G8 locus linked to Huntington's disease. (Abstract) Am.

[129976]J. Hum. Genet. 36: 139S, 1984.

[129977]23289.

[129978]Gusella, J. F.; McNeil, S.; Persichetti, F.; Srinidhi, J.; Novel–

letto,

[129979]A.; Bird, E.; Faber, P.; Vonsattel, J.-P.; Myers, R. H.; Mac-Donald,

[129980]M. E.: Huntington's disease. Cold Spring Harbor Symp. Quant. Biol. 61:

[129981]615-626, 1996.

[129982]23290.

[129983]Dey, B. R.; Furlanetto, R. W.; Nissley, S. P.: Cloning of human

[129984]p55-gamma, a regulatory subunit of phosphatidylinositol 3-kinase,

[129985]by a yeast two-hybrid library screen with the insulin-like growth

[129986]factor-I receptor. Gene 209: 175-183, 1998.

[129987]23291.

[129988]Francke, U.; Yang-Feng, T. L.; Brissenden, J. E.; Ullrich, A.:

[129989]Chromosomal mapping of genes involved in growth control. Cold Spring

[129990]Harbor Symp. Quant. Biol. 51: 855-866, 1986.

[129991]23292.

[129992]Maor, S. B.; Abramovitch, S.; Erdos, M. R.; Brody, L. C.;

Werner,

[129993]H.: BRCA1 suppresses insulin-like growth factor-I receptor promoter

[129994]activity: potential interaction between BRCA1 and Sp1.
Molec. Genet.

[129995]Metab. 69: 130-136, 2000.

[129996]23293.

[129997]Poduslo, S. E.; Dean, M.; Kolch, U.; O'Brien, S. J.: Detecting
[129998]high-resolution polymorphisms in human coding loci by
combining PCR

[129999]and single-strand conformation polymorphism (SSCP)
analysis. Am.

[130000]J. Hum. Genet. 49: 106-111, 1991.

[130001]23294.

[130002]Okumura, N.; Terasawa, F.; Ueno, I.; Oki, K.; Yamauchi, K.;
Hidaka,

[130003]H.; Tozuka, M.; Okura, M.; Katsuyama, T.: Genetic analyses in homozygous

[130004]and heterozygous variants of lactate dehydrogenase-B (H)
subunit:

[130005]LD-B Matsumoto I and II (LD-B W323R). Clin. Chim. Acta

287: 163-171,

[130006]1999.

[130007]23295.

[130008]Rethore, M.-O.; Junien, C.; Malpuech, G.; Baccichetti, C.;
Tenconi,

[130009]R.; Kaplan, J. C.; de Romeuf, J.; Lejeune, J.: Localisation du
gene

[130010]de la glyceraldehyde 3-phosphate dehydrogenase (G3PD)
sur le segment

[130011]distal du bras court du chromosome 12. Ann. Genet. 19:
140-142,

[130012]1976.

[130013]23296.

[130014]Rethore, M.-O.; Kaplan, J.-C.; Junien, C.; Cruveiller, J.;
Dutrillaux,

[130015]B.; Aurias, A.; Carpentier, S.; Lafourcade, J.; Lejeune, J.:
Augmentation

[130016]de l'activite de la LDH-B chez un garçon trisomique 12p
par malsegregation

[130017]d'une translocation maternelle t(12;14)(q12;p11). Ann.
Genet. 18:

[130018]81-87, 1975.

[130019]23297.

[130020]Sakai, I.; Sharief, F. S.; Pan, Y.-C. E.; Li, S. S.-L.: The cDNA

[130021]and protein sequences of human lactate dehydrogenase B.

Biochem.

[130022]J. 248: 933-936, 1987.

[130023]23298.

[130024]Santachiara, A. S.; Nabholz, M.; Miggiano, V.; Darlington,

A.

[130025]J.; Bodmer, W. F.: Linkage between human lactate dehy-

drogenase B

[130026]and peptidase B genes. Nature 227: 248-251, 1970.

[130027]23299.

[130028]Steinbach, P.; Rehder, H.: Tetrasomy for the short arm of

chromosome

[130029]12 with accessory isochromosome (+i[12p]) and a marked

LDH-B gene

[130030]dosage effect. Clin. Genet. 32: 1-4, 1987.

[130031]23300.

[130032]Sudo, K.: Personal Communication. Komae City, Japan

7/12/1993.

[130033]23301.

[130034]Sudo, K.; Maekawa, M.; Ikawa, S.; Machida, K.; Kitamura,
M.; Li,

[130035]S. S.-L.: A missense mutation found in human lactate de-
hydrogenase-B

[130036](H) variant gene. Biochem. Biophys. Res. Commun. 168:
672-676, 1990.

[130037]23302.

[130038]Sudo, K.; Maekawa, M.; Luedemann, M. M.; Deaven, L. L.;
Li, S.

[130039]S.-L.: Human lactate dehydrogenase-B processed pseudo-
gene: nucleotide

[130040]sequence analysis and assignment to the X-chromosome.
Biochem. Biophys.

[130041]Res. Commun. 171: 67-74, 1990.

[130042]23303.

[130043]Sudo, K.; Maekawa, M.; Tomonaga, A.; Tsukada, T.;
Nakayama, T.;

[130044]Kitamura, M.; Li, S. S.-L.; Kanno, T.; Toriumi, J.: Molecular
characterization

[130045]of genetic mutations in human lactate dehydrogenase
(LDH)B (H) variant. Hum.

[130046]Genet. 89: 158-162, 1992.

[130047]23304.

[130048]Van Someren, H.; Meera Khan, P.; Westerveld, A.; Bootsma,
D.:

[130049]Human genetics---two new linkage groups carrying differ-
ent loci for

[130050]LDH and glutamic-pyruvic transaminase found. Nature
240: 221-222,

[130051]1972.

[130052]23305.

[130053]Weiss, L.; Mayeda, K.; Lindahl, R.; Dully, M.: Localization of

[130054]human LDH-B gene of the short arm of chromosome 12.
(Abstract) Am.

[130055]J. Hum. Genet. 25: 85A only, 1973.

[130056]23306.

[130057]Askew, D. S.; Bartholomew, C.; Buchberg, A. M.; Valentine,
M. B.;

[130058]Jenkins, N. A.; Copeland, N. G.; Ihle, J. N.: His-1 and His-
2: identification

[130059]and chromosomal mapping of two commonly rearranged
sites of viral

[130060]integration in a myeloid leukemia. Oncogene 6:
2041–2047, 1991.

[130061]23307.

[130062]Zhong, S.; Delva, L.; Rachez, C.; Cenciarelli, C.; Gandini,
D.;

[130063]Zhang, H.; Kalantry, S.; Freedman, L. P.; Pandolfi, P. P.: A
RA-dependent,

[130064]tumour–growth suppressive transcription complex is the
target of the

[130065]PML–RAR–alpha and T18 oncoproteins. Nature Genet. 23:
287–295, 1999.

[130066]23308.

[130067]Kamb, A.; Gruis, N. A.; Weaver–Feldhaus, J.; Liu, Q.;
Harshman,

[130068]K.; Tavtigian, S. V.; Stockert, E.; Day, R. S., III; Johnson, B.
E.;

[130069]Skolnick, M. H.: A cell cycle regulator potentially involved
in genesis

[130070]of many tumor types. Science 264: 436–440, 1994.

[130071]23309.

[130072]Bonner, T.; O'Brien, S. J.; Nash, W. G.; Rapp, U. R.; Morton,
C.

[130073]C.; Leder, P.: The human homologs of the raf (mil) oncogene are located
[130074]on human chromosomes 3 and 4. Science 223: 71-74,
1984.
[130075]23310.

[130076]Bonner, T. I.; Kerby, S. B.; Suttrave, P.; Gunnell, M. A.; Mark,
[130077]G.; Rapp, U. R.: Structure and biological activity of human
homologs
[130078]of the raf/mil oncogene. Molec. Cell. Biol. 5: 1400-1407,
1985.
[130079]23311.

[130080]Bonner, T. I.; Oppermann, H.; Seeburg, P.; Kerby, S. B.;
Gunnell,
[130081]M. A.; Young, A. C.; Rapp, U. R.: The complete coding se-
quence of
[130082]the human raf oncogene and the corresponding structure
of the c-raf-1
[130083]gene. Nucleic Acids Res. 14: 1009-1015, 1986.
[130084]23312.

[130085]Bonner, T. I.; Rapp, U. R.; Nash, W. G.; O'Brien, S. J.: Two
human

[130086]homologues to a new retroviral oncogene raf-1 and raf-2
are assigned

[130087]to human chromosomes 3 and 4, respectively. (Abstract)
Cytogenet.

[130088]Cell Genet. 37: 424, 1984.

[130089]23313.

[130090]Fukui, M.; Yamamoto, T.; Kawai, S.; Maruo, K.; Toyoshima,
K.:

[130091]Detection of a raf-related and two other transforming
DNA sequences

[130092]in human tumors maintained in nude mice. Proc. Nat.
Acad. Sci. 82:

[130093]5954-5958, 1985.

[130094]23314.

[130095]Hiroshige, S.; Carlock, L.; Smith, M.: Regional assignment
of

[130096]the RAF2 gene to the region 4pter-4p15. (Abstract) Am. J.
Hum. Genet. 39:

[130097]A157, 1986.

[130098]23315.

[130099]Hoppyan, S.; Gokgoz, N.; Poon, R.; Gensure, R. C.; Yu, C.;

Cole,

[130100]W. G.; Bell, R. S.; Juppner, H.; Andrulis, I. L.; Wunder, J. S.;

Alman,

[130101]B. A.: A mutant PTH/PTHrP type I receptor in enchondro-
matosis. Nature

[130102]Genet. 30: 306–310, 2002.

[130103]23316.

[130104]Sakamoto, O.; Iwama, A.; Amitani, R.; Takehara, T.; Yam-
aguchi,

[130105]N.; Yamamoto, T.; Masuyama, K.; Yamanaka, T.; Ando, M.;
Suda, T.:

[130106]Role of macrophage–stimulating protein and its receptor,
RON tyrosine

[130107]kinase, in ciliary motility. J. Clin. Invest. 99: 701–709,
1997.

[130108]23317.

[130109]Fukuyama, R.; Ichijoh, Y.; Minoshima, S.; Kitamura, N.;
Shimizu,

[130110]N.: Assignment of hepatocyte growth factor (HGF) gene to
chromosome

[130111]7q21.1. (Abstract) Cytogenet. Cell Genet. 58: 1921 only,
1991.

[130112]23318.

[130113]Fukuyama, R.; Ichijoh, Y.; Minoshima, S.; Kitamura, N.;
Shimizu,

[130114]N.: Regional localization of the hepatocyte growth factor
(HGF) gene

[130115]to human chromosome 7 band q21.1. Genomics 11:
410-415, 1991.

[130116]23319.

[130117]Gherardi, E.; Stoker, M.: Hepatocytes and scatter factor.
Nature 346:

[130118]228 only, 1990.

[130119]23320.

[130120]Gohda, E.; Tsubouchi, H.; Nakayama, H.; Hirono, S.;
Sakiyama, O.;

[130121]Takahashi, K.; Miyazaki, H.; Hashimoto, S.; Daikuhara, Y.:
Purification

[130122]and partial characterization of hepatocyte growth factor
from plasma

[130123]of a patient with fulminant hepatic failure. J. Clin. Invest.
81:

[130124]414-419, 1988.

[130125]23321.

[130126]Kilby, M. D.; Afford, S.; Li, X. F.; Strain, A. J.; Ahmed, A.;

[130127]Whittle, M. J.: Localisation of hepatocyte growth factor and
its

[130128]receptor (c-met) protein and mRNA in human term pla-
centa. Growth

[130129]Factors 13: 133-139, 1996.

[130130]23322.

[130131]Lai, L.; Goldschneider, I.: Cutting edge: identification of a

[130132]hybrid cytokine consisting of IL-7 and the beta-chain of
the hepatocyte

[130133]growth factor/scatter factor. J. Immun. 167: 3550-3554,
2001.

[130134]23323.

[130135]Maina, F.; Casagrande, F.; Audero, E.; Simeone, A.; Co-
moglio, P.

[130136]M.; Klein, R.; Ponzetto, C.: Uncoupling of Grb2 from the
Met receptor

[130137]in vivo reveals complex roles in muscle development. Cell
87: 531-542,

[130138]1996.

[130139]23324.

[130140]Miyazawa, K.; Tsubouchi, H.; Naka, D.; Takahashi, K.; Okigaki,

[130141]M.; Arakaki, N.; Nakayama, H.; Hirono, S.; Sakiyama, O.; Takahashi,

[130142]K.; Gohda, E.; Daikuhara, Y.; Kitamura, N.: Molecular cloning and

[130143]sequence analysis of cDNA for human hepatocyte growth factor. Biochem.

[130144]Biophys. Res. Commun. 163: 967–973, 1989.

[130145]23325.

[130146]Nakamura, T.; Nishizawa, T.; Hagiya, M.; Seki, T.; Shimonishi,

[130147]M.; Sugimura, A.; Tashiro, K.; Shimizu, S.: Molecular cloning and

[130148]expression of human hepatocyte growth factor. Nature 342: 440–443,

[130149]1989.

[130150]23326.

[130151]Noonan, F. P.; Recio, J. A.; Takayama, H.; Duray, P.; Anver, M.

[130152]R.; Rush, W. L.; De Fabo, E. C.; Merlino, G.: Neonatal sunburn and

[130153]melanoma in mice: severe sunburn in newborn, but not adult, mice is

[130154]linked with melanoma in later life. Nature 413: 271–272, 2001.

[130155]23327.

[130156]Powell, E. M.; Mars, W. M.; Levitt, P.: Hepatocyte growth factor/scatter

[130157]factor is a motogen for interneurons migrating from the ventral to

[130158]dorsal telencephalon. Neuron 30: 79–89, 2001.

[130159]23328.

[130160]Rubin, J. S.; Chan, A. M.–L.; Bottaro, D. P.; Burgess, W. H.;

[130161]Taylor, W. G.; Cech, A. C.; Hirschfield, D. W.; Wong, J.; Miki, T.;

[130162]Finch, P. W.; Aaronson, S. A.: A broad-spectrum human lung fibroblast-derived

[130163]mitogen is a variant of hepatocyte growth factor. Proc. Nat. Acad.

[130164]Sci. 88: 415–419, 1991.

[130165]23329.

[130166]Saccone, S.; Narsimhan, R. P.; Gaudino, G.; Dalpra, L.; Co-
moglio,

[130167]P. M.; Della Valle, G.: Regional mapping of the human
hepatocyte

[130168]growth factor (HGF)-scatter factor gene to chromosome
7q21.1. Genomics 13:

[130169]912-914, 1992.

[130170]23330.

[130171]Schmidt, C.; Blatt, F.; Goedecke, S.; Brinkmann, V.; Zschi-
esche,

[130172]W.; Sharpe, M.; Gherardi, E.; Birchmeier, C.: Scatter factor/
hepatocyte

[130173]growth factor is essential for liver development. Nature
373: 699-702,

[130174]1995.

[130175]23331.

[130176]Uehara, Y.; Minowa, O.; Mori, C.; Shiota, K.; Kuno, J.;
Noda,

[130177]T.; Kitamura, N.: Placental defect and embryonic lethality
in mice

[130178]lacking hepatocyte growth factor/scatter factor. Nature
373: 702–705,

[130179]1995.

[130180]23332.

[130181]Weidner, K. M.; Arakaki, N.; Hartmann, G.; Vandekerck-
hove, J.;

[130182]Weingart, S.; Rieder, H.; Fonatsch, C.; Tsubouchi, H.;
Hishida, T.;

[130183]Daikuhara, Y.; Birchmeier, W.: Evidence for the identity of
human

[130184]scatter factor and human hepatocyte growth factor. Proc.
Nat. Acad.

[130185]Sci. 88: 7001–7005, 1991.

[130186]23333.

[130187]Zarnegar, R.; Petersen, B.; DeFrances, M. C.; Michalopou-
los, G.

[130188]: Localization of hepatocyte growth factor (HGF) gene on
human chromosome

[130189]7. Genomics 12: 147–150, 1992.

[130190]23334.

[130191]Gusella, J. F.; Tanzi, R.; Anderson, M. A.; Ottina, K.; Wal-

lace,

[130192]M.; Conneally, P. M.: Linkage analysis of Huntington's disease using

[130193]RFLPs. (Abstract) Cytogenet. Cell Genet. 37: 484–485, 1984.

[130194]23335.

[130195]Gusella, J. F.; Tanzi, R. E.; Anderson, M. A.; Hobbs, W.; Gibbons,

[130196]K.; Raschtchian, R.; Gilliam, T. C.; Wallace, M. R.; Wexler, N. S.;

[130197]Conneally, P. M.: DNA markers for nervous system diseases. Science 225:

[130198]1320–1326, 1984.

[130199]23336.

[130200]Gusella, J. F.; Wexler, N. S.; Conneally, P. M.; Naylor, S. L.;

[130201]Anderson, M. A.; Tanzi, R. E.; Watkins, P. C.; Ottina, K.; Wallace,

[130202]M. R.; Sakaguchi, A. Y.; Young, A. B.; Shoulson, I.; Bonilla, E.;

[130203]Martin, J. B.: A polymorphic DNA marker genetically linked to Huntington's

[130204]disease. Nature 306: 234–238, 1983.

[130205]23337.

[130206]Gutekunst, C.-A.; Levey, A. I.; Heilman, C. J.; Whaley, W. L.;

[130207]Yi, H.; Nash, N. R.; Rees, H. D.; Madden, J. J.; Hersch, S. M.: Identification

[130208]and localization of huntingtin in brain and human lymphoblastoid cell

[130209]lines with anti-fusion protein antibodies. Proc. Nat. Acad. Sci. 92:

[130210]8710-8714, 1995.

[130211]

[130212]100. Haines, J.; Tanzi, R.; Wexler, N.; Harper, P.; Folstein, S.;

[130213]Cassiman, J.; Meyers, R.; Young, A.; Hayden, M.; Falek, A.; Tolosa,

[130214]E.; Crespi, S.; Campanella, G.; Holmgren, G.; Anvret, M.; Kanazawa,

[130215]I.; Gusella, J.; Conneally, M.: No evidence of linkage heterogeneity

[130216]between Huntington disease (HD) and G8 (D4S10). (Abstract) Am. J.

[130217]Hum. Genet. 39: A156, 1986.

[130218]

[130219]101. Hansson, O.; Petersen, A.; Leist, M.; Nicotera, P.;
Castilho,

[130220]R. F.; Brundin, P.: Transgenic mice expressing a Hunting-
ton's disease

[130221]mutation are resistant to quinolinic acid-induced striatal
excitotoxicity. Proc.

[130222]Nat. Acad. Sci. 96: 8727-8732, 1999.

[130223]

[130224]102. Harding, A. E.: Genetic aspects of autosomal domi-
nant late onset

[130225]cerebellar ataxia. J. Med. Genet. 18: 436-441, 1981.

[130226]

[130227]103. Harper, P. S.: The epidemiology of Huntington's dis-
ease. Hum.

[130228]Genet. 89: 365-376, 1992.

[130229]

[130230]104. Harper, P. S.: Localization of the gene for Hunting-
ton's chorea. Trends

[130231]Neurosci. 7: 1-2, 1984.

[130232]

[130233]105. Harper, P. S.; Lim, C.; Craufurd, D.: Ten years of
presymptomatic

[130234]testing for Huntington's disease: the experience of the UK
Huntington's

[130235]Disease Prediction Consortium. J. Med. Genet. 37:
567-571, 2000.

[130236]

[130237]106. Harper, P. S.; Sarfarazi, M.: Genetic prediction and
family

[130238]structure in Huntington's chorea. Brit. Med. J. 290:
1929-1931,

[130239]1985.

[130240]

[130241]107. Harper, P. S.; Tyler, A.; Walker, D. A.; Newcombe, R.
G.; Davies,

[130242]K.: Huntington's chorea: the basis for long-term preven-
tion. Lancet II:

[130243]346-349, 1979.

[130244]

[130245]108. Harper, P. S.; Youngman, S.; Anderson, M. A.; Sar-

farazi, M.;

[130246]Quarrell, O.; Tanzi, R.; Shaw, D.; Wallace, P.; Conneally, P.
M.;

[130247]Gusella, J. F.: Genetic linkage between Huntington's dis-
ease and

[130248]the DNA polymorphism G8 in South Wales families. J. Med.
Genet. 22:

[130249]447-450, 1985.

[130250]

[130251]109. Hayden, M. R.: Huntington's Chorea. Berlin and New
York: Springer-Verlag

[130252](pub.) 1981.

[130253]

[130254]110. Hayden, M. R.; Beighton, P.: Genetic aspects of Hunt-
ington's

[130255]chorea: results of a national survey. Am. J. Med. Genet.
11: 135-141,

[130256]1982.

[130257]

[130258]111. Hayden, M. R.; Hewitt, J.; Wasmuth, J. J.; Kastelein, J.
J.;

[130259]Langlois, S.; Conneally, M.; Haines, J.; Smith, B.; Hilbert, C.; Allard,

[130260]D.: A polymorphic DNA marker that represents a conserved expressed

[130261]sequence in the region of the Huntington disease gene. Am. J. Hum.

[130262]Genet. 42: 125–131, 1988.

[130263]

[130264]112. Hayden, M. R.; Martin, W. R. W.; Stoessl, A. J.; Clark, C.; Hollenberg,

[130265]S.; Adam, M. J.; Ammann, W.; Harrop, R.; Rogers, J.; Ruth, T.; Sayre,

[130266]C.; Pate, B. D.: Positron emission tomography in the early diagnosis

[130267]of Huntington's disease. Neurology 36: 888–894, 1986.

[130268]

[130269]113. Hayes, C. V.: Genetic testing for Huntington's disease—a family

[130270]issue. (Editorial) New Eng. J. Med. 327: 1449–1451, 1992.

[130271]

[130272]114. Heiser, V.; Scherzinger, E.; Boeddrich, A.; Nordhoff,

E.; Lurz,

[130273]R.; Schugardt, N.; Lehrach, H.; Wanker, E. E.: Inhibition of huntingtin

[130274]fibrillogenesis by specific antibodies and small molecules: implications

[130275]for Huntington's disease therapy. Proc. Nat. Acad. Sci. 97: 6739–6744,

[130276]2000.

[130277]

[130278]115. Hilditch–Maguire, P.; Trettel, F.; Passani, L. A.; Auerbach,

[130279]A.; Persichetti, F.; MacDonald, M. E.: Huntingtin: an iron-regulated

[130280]protein essential for normal nuclear and perinuclear organelles. Hum.

[130281]Molec. Genet. 9: 2789–2797, 2000.

[130282]

[130283]116. Hodge, S. E.; Spence, M. A.; Crandall, B. F.; Sparkes, R. S.;

[130284]Sparkes, M. C.; Crist, M.; Tideman, S.: Huntington disease: linkage

[130285]analysis with age-of-onset corrections. Am. J. Med. Genet.

5: 247–254,

[130286]1980.

[130287]

[130288]117. Hodgson, J. G.; Agopyan, N.; Gutekunst, C.–A.; Leavitt, B. R.;

[130289]LePiane, F.; Singaraja, R.; Smith, D. J.; Bissada, N.; McCutcheon,

[130290]K.; Nasir, J.; Jamot, L.; Li, X.–J.; Stevens, M. E.; Rosemond, E.;

[130291]Roder, J. C.; Phillips, A. G.; Rubin, E. M.; Hersch, S. M.; Hayden,

[130292]M. R.: A YAC mouse model for Huntington's disease with full-length

[130293]mutant huntingtin, cytoplasmic toxicity, and selective striatal neurodegeneration. *Neuron* 23:

[130294]181–192, 1999.

[130295]

[130296]118. Hodgson, J. G.; Smith, D. J.; McCutcheon, K.; Koide, H. B.; Nishiyama,

[130297]K.; Dinulos, M. B.; Stevens, M. E.; Bissada, N.; Nasir, J.; Kanazawa,

[130298]J.; Disteche, C. M.; Rubin, E. M.; Hayden, M. R.: Human

huntingtin

[130299] derived from YAC transgenes compensates for loss of
murine huntingtin

[130300] by rescue of the embryonic lethal phenotype. Hum. Molec.
Genet. 5:

[130301] 1875–1885, 1996.

[130302]

[130303] 119. Holmgren, G.; Winnberg Almqvist, E.; Anvret, M.;
Conneally, M.;

[130304] Hobbs, W.; Mattsson, B.; Wahlstrom, J.; Winblad, B.;
Gusella, J. F.

[130305]: Linkage of G8 (D4S10) in two Swedish families with
Huntington's

[130306] disease. Clin. Genet. 32: 289–294, 1987.

[130307]

[130308] 120. Hoogeveen, A. T.; Willemsen, R.; Meyer, N.; de Rooij,
K. E.;

[130309] Roos, R. A. C.; van Ommen, G.–J. B.; Galjaard, H.: Charac-
terization

[130310] and localization of the Huntington disease gene product.
Hum. Molec.

[130311] Genet. 2: 2069–2073, 1993.

[130312]

[130313]121. Horton, T. M.; Graham, B. H.; Corral-Debrinski, M.;
Shoffner,

[130314]J. M.; Kaufman, A. E.; Beal, M. F.; Wallace, D. C.: Marked
increase

[130315]in mitochondrial DNA deletion levels in the cerebral cortex
of Huntington's

[130316]disease patients. *Neurology* 45: 1879–1883, 1995.

[130317]

[130318]122. Huntington's Disease Collaborative Research Group:
A novel

[130319]gene containing a trinucleotide repeat that is expanded
and unstable

[130320]on Huntington's disease chromosomes. *Cell* 72: 971–983,
1993.

[130321]

[130322]123. Huntington, G.: On chorea. *Med. Surg. Reporter* 26:
317–321,

[130323]1872.

[130324]

[130325]124. Ikeda, H.; Yamaguchi, M.; Sugai, S.; Aze, Y.; Naru-

miya, S.; Kakizuka,

[130326]A.: Expanded polyglutamine in the Machado–Joseph disease protein

[130327]induces cell death in vitro and in vivo. *Nature Genet.* 13: 196–202,

[130328]1996.

[130329]

[130330]125. Ikonen, E.; Ignatius, J.; Norio, R.; Palo, J.; Peltonen, L.:

[130331]Huntington disease in Finland: a molecular and genealogical study. *Hum.*

[130332]*Genet.* 89: 275–280, 1992.

[130333]

[130334]126. Illarioshkin, S. N.; Igarashi, S.; Onodera, O.; Markova, E. D.;

[130335]Nikolskaya, N. N.; Tanaka, H.; Chabrashwili, T. Z.; Insarova, N. G.;

[130336]Endo, K.; Ivanova–Smolenskaya, I. A.; Tsuji, S.: Trinucleotide repeat

[130337]length and rate of progression of Huntington's disease. *Ann. Neurol.* 36:

[130338]630–635, 1994.

[130339]

[130340]127. Jana, N. R.; Zemskov, E. A.; Wang, G.; Nukina, N.: Altered proteasomal

[130341]function due to the expression of polyglutamine-expanded truncated

[130342]N-terminal huntingtin induces apoptosis by caspase activation through

[130343]mitochondrial cytochrome c release. Hum. Molec. Genet. 10: 1049-1059,

[130344]2001.

[130345]

[130346]128. Kahlem, P.; Green, H.; Djian, P.: Transglutaminase action imitates

[130347]Huntington's disease: selective polymerization of huntingtin containing

[130348]expanded polyglutamine. Molec. Cell 1: 595-601, 1998.

[130349]

[130350]129. Kanazawa, I.; Kondo, I.; Ikeda, J.-E.; Ikeda, T.; Shizu, Y.;

[130351]Yoshida, M.; Narabayashi, H.; Kuroda, S.; Tsunoda, H.; Mizuta, E.;

- [130352] Okuno, Y.; Sugawara, K.; Murata, M.; Takahashi, M.;
Gusella, J. F.
- [130353] Studies on DNA markers (D4S10 and D4S43/S127) ge-
netically linked
to Huntington's disease in Japanese families. Hum. Genet.
85: 257-260,
1990.
- [130355]
- [130356]
- [130357] 130. Karpuj, M. V.; Becher, M. W.; Springer, J. E.; Chabas,
D.; Youssef,
S.; Pedotti, R.; Mitchell, D.; Steinman, L.: Prolonged sur-
vival and
decreased abnormal movements in transgenic model of
Huntington disease,
with administration of the transglutaminase inhibitor cys-
tamine. Nature
Med. 8: 143-149, 2002. Note: Erratum: Nature Med. 8:
303 only, 2002.
- [130362]
- [130363] 131. Kazantsev, A.; Walker, H. A.; Slepko, N.; Bear, J. E.;
Preisinger,
E.; Steffan, J. S.; Zhu, Y.-Z.; Gertler, F. B.; Housman, D. E.;

Marsh,

[130365]J. L.; Thompson, L. M.: A bivalent Huntingtin binding peptide suppresses

[130366]polyglutamine aggregation and pathogenesis in *Drosophila*. *Nature*

[130367]*Genet.* 30: 367–376, 2002.

[130368]

[130369]132. Kazemi-Esfarjani, P.; Benzer, S.: Genetic suppression of polyglutamine

[130370]toxicity in *Drosophila*. *Science* 287: 1837–1840, 2000.

[130371]

[130372]133. Keene, C. D.; Rodrigues, C. M. P.; Eich, T.; Chhabra, M. S.;

[130373]Steer, C. J.; Low, W. C.: Tauroursodeoxycholic acid, a bile acid,

[130374]is neuroprotective in a transgenic animal model of Huntington's disease. *Proc.*

[130375]*Nat. Acad. Sci.* 99: 10671–10676, 2002.

[130376]

[130377]134. Keene, C. D.; Rodrigues, C. M. P.; Eich, T.; Linehan-Stieers,

[130378]C.; Abt, A.; Kren, B. T.; Steer, C. J.; Low, W. C.: A bile acid protects

[130379]against motor and cognitive deficits and reduces striatal degeneration

[130380]in the 3-nitropropionic acid model of Huntington's disease. Exp.

[130381]Neurol. 171: 351-360, 2001.

[130382]

[130383]135. Kehoe, P.; Krawczak, M.; Harper, P. S.; Owen, M. J.; Jones, A.

[130384]L.: Age of onset in Huntington disease: sex specific influence of

[130385]apolipoprotein E genotype and normal CAG repeat length. J. Med. Genet. 36:

[130386]108-111, 1999.

[130387]

[130388]136. Kennedy, L.; Shelbourne, P. F.: Dramatic mutation instability

[130389]in HD mouse striatum: does polyglutamine load contribute to cell-specific

[130390]vulnerability in Huntington's disease? Hum. Molec. Genet. 9: 2539-2544,

[130391]2000.

[130392]

[130393]137. Kerbeshian, J.; Burd, L.; Leech, C.; Rorabaugh, A.:
Huntington

[130394]disease and childhood-onset Tourette syndrome. Am. J.
Med. Genet. 39:

[130395]1-3, 1991.

[130396]

[130397]138. Khoshnan, A.; Ko, J.; Patterson, P. H.: Effects of intra-
cellular

[130398]expression of anti-huntingtin antibodies of various speci-
ficities

[130399]on mutant huntingtin aggregation and toxicity. Proc. Nat.
Acad. Sci. 99:

[130400]1002-1007, 2002.

[130401]

[130402]139. Klawans, H. L., Jr.; Paulson, G. W.; Ringel, S. P.; Bar-
beau,

[130403]A.: L-dopa in the detection of presymptomatic Hunting-
ton's chorea. New

[130404]Eng. J. Med. 286: 1332-1334, 1972.

[130405]

[130406]140. Ko, J.; Ou, S.; Patterson, P. H.: New anti-huntingtin
monoclonal

[130407]antibodies: implications for huntingtin conformation and
its binding

[130408]proteins. Brain Res. Bull. 56: 319–329, 2001.

[130409]

[130410]141. Kovtun, I. V.; McMurray, C. T.: Trinucleotide expansion in haploid

[130411]germ cells by gap repair. Nature Genet. 27: 407–411,
2001.

[130412]

[130413]142. Kovtun, I. V.; Therneau, T. M.; McMurray, C. T.: Gender of the

[130414]embryo contributes to CAG instability in transgenic mice
containing

[130415]a Huntington's disease gene. Hum. Molec. Genet. 9:
2767–2775, 2000.

[130416]

[130417]143. Kremer, B.; Goldberg, P.; Andrew, S. E.; Theilmann, J.;
Telenius,

[130418]H.; Zeisler, J.; Squitieri, F.; Lin, B.; Bassett, A.; Almqvist, E.;

[130419]Bird, T. D.; Hayden, M. R.: A worldwide study of the Huntington's

[130420]disease mutation: the sensitivity and specificity of measuring CAG

[130421]repeats. New Eng. J. Med. 330: 1401–1406, 1994.

[130422]

[130423]144. Laccone, F.; Christian, W.: A recurrent expansion of a maternal

[130424]allele with 36 CAG repeats causes Huntington disease in two sisters. Am.

[130425]J. Hum. Genet. 66: 1145–1148, 2000.

[130426]

[130427]145. Landegent, J. E.; Jansen in de Wal, N.; Fisser–Groen, Y. M.;

[130428]Bakker, E.; van der Ploeg, M.; Pearson, P. L.: Fine mapping of the

[130429]Huntington disease linked D4S10 locus by non-radioactive in situ hybridization. Hum.

[130430]Genet. 73: 354–357, 1986.

[130431]

[130432]146. Lanska, D. J.; Lavine, L.; Lanska, M. J.; Schoenberg, B. S.:

[130433]Huntington's disease mortality in the United States. Neu-
rology 38:

[130434]769-772, 1988.

[130435]

[130436]147. Lazzarini, A.; McCormack, M. K.; Lepore, F.: Maternal
transmission

[130437]of juvenile Huntington's disease in U.S. black families.
(Abstract) Am.

[130438]J. Hum. Genet. 36: 62S, 1984.

[130439]

[130440]148. Leavitt, B. R.; Guttman, J. A.; Hodgson, J. G.; Kimel,
G. H.;

[130441]Singaraja, R.; Vogl, A. W.; Hayden, M. R.: Wild-type hunt-
ingtin reduces

[130442]the cellular toxicity of mutant huntingtin in vivo. Am. J.
Hum. Genet. 68:

[130443]313-324, 2001.

[130444]

[130445]149. Leeflang, E. P.; Tavare, S.; Marjoram, P.; Neal, C. O.

S.; Srinidhi,

[130446]J.; MacDonald, M. E.; de Young, M.; Wexler, N. S.; Gusella,
J. F.;

[130447]Arnhem, N.: Analysis of germline mutation spectra at the
Huntington's

[130448]disease locus supports a mitotic mutation mechanism.

Hum. Molec.

[130449]Genet. 8: 173–183, 1999.

[130450]

[130451]150. Leeflang, E. P.; Zhang, L.; Tavaré, S.; Hubert, R.;

Srinidhi,

[130452]J.; MacDonald, M. E.; Myers, R. H.; de Young, M.; Wexler,

N. S.; Gusella,

[130453]J. F.; Arnheim, N.: Single sperm analysis of the trinucleotide repeats

in the Huntington's disease gene: quantification of the

[130454]mutation frequency

spectrum. Hum. Molec. Genet. 4: 1519–1526, 1995.

[130455]

[130456]

[130457]151. Leung, C. M.; Chan, Y. W.; Chang, C. M.; Yu, Y. L.;

Chen, C.

[130458]N.: Huntington's disease in Chinese: a hypothesis of its

origin. J.

[130459]Neurol. Neurosurg. Psychiat. 55: 681–684, 1992.

[130460]

[130461]152. Li, H.; Li, S.-H.; Johnston, H.; Shelbourne, P. F.; Li,

X.-J.

[130462]: Amino-terminal fragments of mutant huntingtin show selective accumulation

[130463]in striatal neurons and synaptic toxicity. Nature Genet. 25: 385-389,

[130464]2000.

[130465]

[130466]153. Li, S.-H.; Lam, S.; Cheng, A. L.; Li, X.-J.: Intracellular huntingtin

[130467]increases the expression of caspase-1 and induces apoptosis. Hum.

[130468]Molec. Genet. 9: 2859-2867, 2000.

[130469]

[130470]154. Li, X.-J.; Li, S.-H.; Sharp, A. H.; Nucifora, F. C., Jr.; Schilling,

[130471]G.; Lanahan, A.; Worley, P.; Snyder, S. H.; Ross, C. A.: A huntingtin-associated

[130472]protein enriched in brain and implications for pathology. Nature 378:

[130473]398-402, 1995.

[130474]

- [130475]155. Lin, B.; Nasir, J.; Kalchman, M. A.; McDonald, H.; Zeisler, J.;
- [130476]Goldberg, Y. P.; Hayden, M. R.: Structural analysis of the 5-prime
- [130477]region of mouse and human Huntington disease genes reveals conservation
- [130478]of putative promoter region and di- and trinucleotide polymorphisms. *Genomics* 25:
- [130479]707-715, 1995.
- [130480]
- [130481]156. Lin, C.-H.; Tallaksen-Greene, S.; Chien, W.-M.; Cearley, J. A.;
- [130482]Jackson, W. S.; Crouse, A. B.; Ren, S.; Li, X.-J.; Albin, R. L.; Detloff,
- [130483]P. J.: Neurological abnormalities in a knock-in mouse model of Huntington's
- [130484]disease. *Hum. Molec. Genet.* 10: 137-144, 2001.
- [130485]
- [130486]157. Lindblad, A. N.: To test or not to test: an ethical conflict
- [130487]with presymptomatic testing of individuals at 25% risk for Huntington's

[130488]disorder. Clin. Genet. 60: 442–446, 2001.

[130489]

[130490]158. Lovestone, S.; Hodgson, S.; Sham, P.; Differ, A.–M.;
Levy, R.

[130491]: Familial psychiatric presentation of Huntington's disease.
J. Med.

[130492]Genet. 33: 128–131, 1996.

[130493]

[130494]159. Lunkes, A.; Lindenberg, K. S.; Ben–Haiem, L.; Weber,
C.; Devys,

[130495]D.; Landwehrmeyer, G. B.; Mandel, J.–L.; Trottier, Y.: Pro-
teases

[130496]acting on mutant huntingtin generate cleaved products
that differentially

[130497]build up cytoplasmic and nuclear inclusions. Molec. Cell
10: 259–269,

[130498]2002.

[130499]

[130500]160. Lunkes, A.; Mandel, J.–L.: A cellular model that reca-
pitulates

[130501]major pathogenic steps of Huntington's disease. Hum.

Molec. Genet. 7:

[130502]1355–1361, 1998.

[130503]

[130504]161. Luthi–Carter, R.; Strand, A.; Peters, N. L.; Solano, S.
M.; Hollingsworth,

[130505]Z. R.; Menon, A. S.; Frey, A. S.; Spektor, B. S.; Penney, E. B.;
Schilling,

[130506]G.; Ross, C. A.; Borchelt, D. R.; Tapscott, S. J.; Young, A.
B.; Cha,

[130507]J.–H. J.; Olson, J. M.: Decreased expression of striatal sig-
naling

[130508]genes in a mouse model of Huntington's disease. Hum.
Molec. Genet. 9:

[130509]1259–1271, 2000.

[130510]

[130511]162. Lyon, R. L.: Huntington's chorea in the Moray Firth
area. Brit.

[130512]Med. J. 1: 1301–1306, 1962.

[130513]

[130514]163. MacDonald, M. E.; Barnes, G.; Srinidhi, J.; Duyao, M.
P.; Ambrose,

- [130515]C. M.; Myers, R. H.; Gray, J.; Conneally, P. M.; Young, A.; Penney,
- [130516]J.; Shoulson, I.; Hollingsworth, Z.; Koroshetz, W.; Bird, E.; Vonsattel,
- [130517]J. P.; Bonilla, E.; Moscovitz, C.; Penchaszadeh, G.; Brzus-towicz,
- [130518]L.; Alvir, J.; Bickham Conde, J.; Cha, J.-H.; Dure, L.; Gomez, F.;
- [130519]Ramos-Arroyo, M.; Sanchez-Ramos, J.; Snodgrass, S. R.; de Young, M.;
- [130520]Wexler, N. S.; MacFarlane, H.; Anderson, M. A.; Jenkins, B.; Gusella,
- [130521]J. F.: Gametic but not somatic instability of CAG repeat length in
- [130522]Huntington's disease. *J. Med. Genet.* 30: 982-986, 1993.
- [130523]
- [130524]164. MacDonald, M. E.; Cheng, S. V.; Zimmer, M.; Haines, J. L.; Poustka,
- [130525]A.; Allitto, B.; Smith, B.; Whaley, W. L.; Romano, D. M.; Jagadeesh,
- [130526]J.; Myers, R. H.; Lehrach, H.; Wasmuth, J. J.; Frischauf, A.-M.; Gusella,

[130527]J. F.: Clustering of multiallele DNA markers near the Huntington's

[130528]disease gene. J. Clin. Invest. 84: 1013–1016, 1989.

[130529]

[130530]165. MacDonald, M. E.; Duyao, M.; Calzonetti, T.; Auerbach, A.; Ryan,

[130531]A.; Barnes, G.; White, J. K.; Auerbach, W.; Vonsattel, J.-P.; Gusella,

[130532]J. F.; Joyner, A. L.: Targeted inactivation of the mouse Huntington's

[130533]disease gene homolog Hdh. Cold Spring Harbor Symp. Quant. Biol. 61:

[130534]627–638, 1996.

[130535]

[130536]166. MacDonald, M. E.; Haines, J. L.; Zimmer, M.; Cheng, S. V.; Youngman,

[130537]S.; Whaley, W. L.; Wexler, N.; Bucan, M.; Allitto, B. A.; Smith, B.;

[130538]Leavitt, J.; Poustka, A.; Harper, P.; Lehrach, H.; Wasmuth, J. J.;

[130539]Frischauf, A.-M.; Gusella, J. F.: Recombination events suggest potential

[130540]sites for the Huntington's disease gene. Neuron 3:
183–190, 1989.

[130541]

[130542]167. MacDonald, M. E.; Novelletto, A.; Lin, C.; Tagle, D.;
Barnes,

[130543]G.; Bates, G.; Taylor, S.; Allitto, B.; Altherr, M.; Myers, R.;
Lehrach,

[130544]H.; Collins, F. S.; Wasmuth, J. J.; Frontali, M.; Gusella, J. F.:

[130545]The Huntington's disease candidate region exhibits many
different

[130546]haplotypes. Nature Genet. 1: 99–103, 1992.

[130547]

[130548]168. MacDonald, M. E.; Vonsattel, J. P.; Shrinidhi, J.;
Couropmitree,

[130549]N. N.; Cupples, L. A.; Bird, E. D.; Gusella, J. F.; Myers, R.
H.:

[130550]Evidence for the GluR6 gene associated with younger on-
set age of Huntington's

[130551]disease. Neurology 53: 1330–1332, 1999.

[130552]

[130553]169. Magenis, E.; Gusella, J.; Weliky, K.; Haight, G.;

Sheehy, B.

[130554] Huntington disease-linked (HD) restriction fragment polymorphism

[130555] localized to band p16 of chromosome 4 by in situ hybridization. (Abstract) Cytogenet.

[130556] Cell Genet. 40: 685, 1985.

[130557]

[130558] 170. Magenis, R. E.; Gusella, J.; Weliky, K.; Olson, S.; Haight, G.;

[130559] Toth-Fejel, S.; Sheehy, R.: Huntington disease-linked restriction

[130560] fragment length polymorphism localized within band p16.1 of chromosome

[130561] 4 by in situ hybridization. Am. J. Hum. Genet. 39: 383-391, 1986.

[130562]

[130563] 171. Maltzberger, J. T.: Even unto the twelfth generation-Huntington's

[130564] chorea. J. Hist. Med. Allied Sci. 16: 1-17, 1961.

[130565]

[130566] 172. Mangiarini, L.; Sathasivam, K.; Mahal, A.; Mott, R.;

Seller,

[130567]M.; Bates, G. P.: Instability of highly expanded CAG repeats in mice

[130568]transgenic for the Huntington's disease mutation. Nature Genet. 15:

[130569]197–200, 1997.

[130570]

[130571]173. Mangiarini, L.; Sathasivam, K.; Seller, M.; Cozens, B.; Harper,

[130572]A.; Hetherington, C.; Lawton, M.; Trottier, Y.; Lehrach, H.; Davies,

[130573]S. W.; Bates, G. P.: Exon 1 of the HD gene with an expanded CAG repeat

[130574]is sufficient to cause a progressive neurological phenotype in transgenic

[130575]mice. Cell 87: 493–506, 1996.

[130576]

[130577]174. Martin, J. B.; Gusella, J. F.: Huntington's disease: pathogenesis

[130578]and management. New Eng. J. Med. 315: 1267–1276, 1986.

[130579]

[130580]175. Marx, J. L.: A parent's sex may affect gene expression. Science 239:

[130581]352–353, 1988.

[130582]

[130583]176. Mastromauro, C. A.; Meissen, G. J.; Cupples, L. A.; Kiely, D.

[130584]K.; Berkman, B.; Myers, R. H.: Estimation of fertility and fitness

[130585]in Huntington disease in New England. Am. J. Med. Genet. 33: 248–254,

[130586]1989.

[130587]

[130588]177. Masuda, N.; Goto, J.; Murayama, N.; Watanabe, M.; Kondo, I.;

[130589]Kanazawa, I.: Analysis of triplet repeats in the huntingtin gene

[130590]in Japanese families affected with Huntington's disease. J. Med.

[130591]Genet. 32: 701–705, 1995.

[130592]

[130593]178. Mazziotta, J. C.; Phelps, M. E.; Pahl, J. J.; Huang,

S.-C.; Baxter,

[130594]L. R.; Riege, W. H.; Hoffman, J. M.; Kuhl, D. E.; Lanto, A. B.;

Wapenski,

[130595]J. A.; Markham, C. H.: Reduced cerebral glucose

metabolism in asymptomatic

[130596]subjects at risk for Huntington's disease. New Eng. J. Med.

316:

[130597]357-362, 1987.

[130598]

[130599]179. McKeown, C.; Read, A. P.; Dodge, A.; Stecko, O.;

Mercer, A.;

[130600]Harris, R.: Wolf-Hirschhorn locus is distal to D4S10 on

short arm

[130601]of chromosome 4. J. Med. Genet. 24: 410-412, 1987.

[130602]

[130603]180. McKusick, V. A.: Personal Communication. Baltimore,

Md. 1977.

[130604]

[130605]181. McLaughlin, B. A.; Spencer, C.; Eberwine, J.: CAG

trinucleotide

[130606]RNA repeats interact with RNA-binding proteins. Am. J.

Hum. Genet. 59:

[130607]561–569, 1996.

[130608]

[130609]182. Meissen, G. J.; Myers, R. H.; Mastromauro, C. A.; Koro-

roshetz,

[130610]W. J.; Klinger, K. W.; Farrer, L. A.; Watkins, P. A.; Gusella, J.

[130611]F.; Bird, E. D.; Martin, J. B.: Predictive testing for Hunting-

ton's

[130612]disease with use of a linked DNA marker. New Eng. J. Med.

318: 535–542,

[130613]1988.

[130614]

[130615]183. Metzler, M.; Helgason, C. D.; Dragatsis, I.; Zhang, T.;

Gan,

[130616]L.; Pineault, N.; Zeitlin, S. O.; Humphries, R. K.; Hayden, M.

R.

[130617]: Huntingtin is required for normal hematopoiesis. Hum.

Molec. Genet. 9:

[130618]387–394, 2000.

[130619]

[130620]184. Millan, F. A.; Curits, A.; Mennie, M.; Holloway, S.;

Boxer, M.;

[130621]Faed, M. J. W.; Crawford, J. W.; Liston, W. A.; Brock, D. J.
H.:

[130622]Prenatal exclusion testing for Huntington's disease: a
problem of

[130623]too much information. J. Med. Genet. 26: 83–85, 1989.

[130624]

[130625]185. Mochizuki, H.; Kamakura, K.; Kumada, M.; Goto, J.;
Kanazawa,

[130626]I.; Motoyoshi, K.: A patient with Huntington's disease pre-
senting

[130627]with laryngeal chorea. Europ. Neurol. 41: 119–120, 1999.

[130628]

[130629]186. Morris, M. J.; Tyler, A.; Lazarou, L.; Meredith, L.;
Harper,

[130630]P. S.: Problems in genetic prediction for Huntington's dis-
ease. Lancet II:

[130631]601–603, 1989.

[130632]

[130633]187. Morrison, P. J.; Johnston, W. P.; Nevin, N. C.: The epi-
demiology

[130634]of Huntington's disease in Northern Ireland. J. Med. Genet.

32:

[130635]524–530, 1995.

[130636]

[130637]188. Mouchiroud, D.; D'Onofrio, G.; Aissani, B.; Macaya, G.; Gautier,

[130638]C.; Bernardi, G.: The distribution of genes in the human genome. Gene 100:

[130639]181–187, 1991.

[130640]

[130641]189. Muchowski, P. J.; Ning, K.; D'Souza-Schorey, C.; Fields, S.:

[130642]Requirement of an intact microtubule cytoskeleton for aggregation

[130643]and inclusion body formation by a mutant huntingtin fragment. Proc.

[130644]Nat. Acad. Sci. 99: 727–732, 2002.

[130645]

[130646]190. Myers, R. H.; Cupples, L. A.; Schoenfeld, M.; D'Agostino, R.

[130647]B.; Terrin, N. C.; Goldmakher, N.; Wolf, P. A.: Maternal fac-

tors

[130648]in onset of Huntington disease. Am. J. Hum. Genet. 37:
511–523,

[130649]1985.

[130650]

[130651]191. Myers, R. H.; Goldman, D.; Bird, E. D.; Sax, D. S.;
Merril, C.

[130652]R.; Schoenfeld, M.; Wolf, P. A.: Maternal transmission in
Huntington's

[130653]disease. Lancet I: 208–210, 1983.

[130654]

[130655]192. Myers, R. H.; Leavitt, J.; Farrer, L. A.; Jagadeesh, J.;
McFarlane,

[130656]H.; Mastromauro, C. A.; Mark, R. J.; Gusella, J. F.: Homozy-
gote for

[130657]Huntington disease. Am. J. Hum. Genet. 45: 615–618,
1989.

[130658]

[130659]193. Myers, R. H.; Madden, J. J.; Teague, J. L.; Falek, A.:
Factors

[130660]related to onset age of Huntington disease. Am. J. Hum.

Genet. 34:

[130661]481–488, 1982.

[130662]

[130663]194. Myrianthopoulos, N. C.: Huntington's chorea. J. Med.

Genet. 3:

[130664]298–314, 1966.

[130665]

[130666]195. Narain, Y.; Wyttenbach, A.; Rankin, J.; Furlong, R. A.;

Rubinsztein,

[130667]D. C.: A molecular investigation of true dominance in

Huntington's

[130668]disease. J. Med. Genet. 36: 739–746, 1999.

[130669]

[130670]196. Nasir, J.; Floresco, S. B.; O'Kusky, J. R.; Diewert, V. M.;

Richman,

[130671]J. M.; Zeisler, J.; Borowski, A.; Marth, J. D.; Phillips, A. G.;

Hayden,

[130672]M. R.: Targeted disruption of the Huntington's disease

gene results

[130673]in embryonic lethality and behavioral and morphological

changes in

[130674]heterozygotes. Cell 81: 811–823, 1995.

[130675]

[130676]197. Nasir, J.; Lin, B.; Bucan, M.; Koizumi, T.; Nadeau, J. H.; Hayden,

[130677]M. R.: The murine homologues of the Huntington disease gene (Hdh)

[130678]and the alpha-adducin gene (Add1) map to mouse chromosome 5 within

[130679]a region of conserved synteny with human chromosome 4p16.3. Genomics 22:

[130680]198–201, 1994.

[130681]

[130682]198. Navarrete, C.; Martinez, I.; Salamanca, F.: Paternal line of

[130683]transmission in chorea of Huntington with very early onset. Genet.

[130684]Counsel. 5: 175–178, 1994.

[130685]

[130686]199. Ona, V. O.; Li, M.; Vonsattel, J. P. G.; Andrews, L. J.; Khan,

[130687]S. Q.; Chung, W. M.; Frey, A. S.; Menon, A. S.; Li, X.-J.;

Stieg,

[130688]P. E.; Yuan, J.; Penney, J. B.; Young, A. B.; Cha, J.-H. J.;

Friedlander,

[130689]R. M.: Inhibition of caspase-1 slows disease progression
in a mouse

[130690]model of Huntington's disease. Nature 399: 263-267,
1999.

[130691]

[130692]200. Ordway, J. M.; Tallaksen-Greene, S.; Gutekunst,
C.-A.; Bernstein,

[130693]E. M.; Cearley, J. A.; Wiener, H. W.; Dure, L. S., IV; Lindsey,
R.;

[130694]Hersch, S. M.; Jope, R. S.; Albin, R. L.; Detloff, P. J.: Ectopi-
cally

[130695]expressed CAG repeats cause intranuclear inclusions and
a progressive

[130696]late onset neurological phenotype in the mouse. Cell 91:
753-763,

[130697]1997.

[130698]

[130699]201. Orr, H. T.; Zoghbi, H. Y.: Reversing neurodegenera-
tion: a promise

[130700]unfolds. Cell 101: 1–4, 2000.

[130701]

[130702]202. Osler, W.: Remarks on the varieties of chronic chorea, and a

[130703]report upon two families of the hereditary form, with one autopsy. J.

[130704]Nerv. Ment. Dis. 18: 97–111, 1893.

[130705]

[130706]203. Palo, J.; Somer, H.; Ikonen, E.; Karila, L.; Peltonen, L.: Low

[130707]prevalence of Huntington's disease in Finland. Lancet I: 805–806,

[130708]1987.

[130709]

[130710]204. Panov, A. V.; Gutekunst, C.–A.; Leavitt, B. R.; Hayden, M. R.;

[130711]Burke, J. R.; Strittmatter, W. J.; Greenamyre, J. T.: Early mitochondrial

[130712]calcium defects in Huntington's disease are a direct effect of polyglutamines. Nature

[130713]Neurosci. 5: 731–736, 2002.

[130714]

[130715]205. Parrish, J. E.; Nelson, D. L.: Methods for finding
genes: a

[130716]major rate-limiting step in positional cloning. Genet. Anal.
Tech.

[130717]Appl. 10: 29-41, 1993.

[130718]

[130719]206. Paulson, H. L.; Bonini, N. M.; Roth, K. A.: Polyglu-
tamine disease

[130720]and neuronal cell death. Proc. Nat. Acad. Sci. 97:
12957-12958,

[130721]2000.

[130722]

[130723]207. Peel, A. L.; Rao, R. V.; Cottrell, B. A.; Hayden, M. R.;
Ellerby,

[130724]L. M.; Bredesen, D. E.: Double-stranded RNA-dependent
protein kinase,

[130725]PKR, binds preferentially to Huntington's disease (HD)
transcripts

[130726]and is activated in HD tissue. Hum. Molec. Genet. 10:
1531-1538,

[130727]2001.

[130728]

[130729]208. Pericak-Vance, M. A.; Conneally, P. M.; Merritt, A. D.;
Roos,
[130730]R.; Norton, J. A., Jr.; Vance, J. M.: Genetic linkage studies
in
[130731]Huntington disease. Cytogenet. Cell Genet. 22: 640-645,
1978.

[130732]

[130733]209. Perry, T. L.; Hansen, S.; Kloster, M.: Huntington's
chorea:
[130734]deficiency of gamma-aminobutyric acid in brain. New Eng.
J. Med. 288:
[130735]337-342, 1973.

[130736]

[130737]210. Petersen, A.; Larsen, K. E.; Behr, G. G.; Romero, N.;
Przedborski,
[130738]S.; Brundin, P.; Sulzer, D.: Expanded CAG repeats in exon
1 of the
[130739]Huntington's disease gene stimulate dopamine-mediated
striatal neuron

[130740]autophagy and degeneration. Hum. Molec. Genet. 10:
1243–1254, 2001.

[130741]

[130742]211. Peyser, C. E.; Folstein, M.; Chase, G. A.; Starkstein, S.;
Brandt,

[130743]J.; Cockrell, J. R.; Bylsma, F.; Coyle, J. T.; McHugh, P. R.;
Folstein,

[130744]S. E.: Trial of d-alpha-tocopherol in Huntington's disease.
Am.

[130745]J. Psychiat. 152: 1771–1775, 1995.

[130746]

[130747]212. Portera-Cailliau, C.; Hedreen, J. C.; Price, D. L.; Koli-
atsos,

[130748]V. E.: Evidence for apoptotic cell death in Huntington dis-
ease and

[130749]excitotoxic animal models. J. Neurosci. 15: 3775–3787,
1995.

[130750]

[130751]213. Pridmore, S. A.: The large Huntington's disease fam-
ily of Tasmania. Med.

[130752]J. Aust. 153: 593–595, 1990.

[130753]

[130754]214. Quarrell, O. W. J.; Meredith, A. L.; Tyler, A.; Youngman, S.;

[130755]Upadhyaya, M.; Harper, P. S.: Exclusion testing for Huntington's

[130756]disease in pregnancy with a closely linked DNA marker.

Lancet I:

[130757]1281-1283, 1987.

[130758]

[130759]215. Quarrell, O. W. J.; Tyler, A.; Cole, G.; Harper, P. S.:
The

[130760]problem of isolated cases of Huntington's disease in South
Wales 1974-1984. Clin.

[130761]Genet. 30: 433-439, 1986.

[130762]

[130763]216. Quarrell, O. W. J.; Tyler, A.; Jones, M. P.; Nordin, M.;
Harper,

[130764]P. S.: Population studies of Huntington's disease in Wales.
Clin.

[130765]Genet. 33: 189-195, 1988.

[130766]

[130767]217. Ranen, N. G.; Stine, O. C.; Abbott, M. H.; Sherr, M.;
Codori,

[130768]A.-M.; Franz, M. L.; Chao, N. I.; Chung, A. S.; Pleasant, N.;
Callahan,

[130769]C.; Kasch, L. M.; Ghaffari, M.; Chase, G. A.; Kazazian, H.
H.; Brandt,

[130770]J.; Folstein, S. E.; Ross, C. A.: Anticipation and instability of
[130771]IT-15 (CAG) n repeats in parent-offspring pairs with Hunt-
ington disease. Am.

[130772]J. Hum. Genet. 57: 593-602, 1995.

[130773]

[130774]218. Read, A. P.: Huntington's disease: testing the test.
Nature

[130775]Genet. 4: 329-330, 1993.

[130776]

[130777]219. Reddy, P. H.; Williams, M.; Charles, V.; Garrett, L.;
Pike-Buchanan,

[130778]L.; Whetsell, W. O., Jr.; Miller, G.; Tagle, D. A.: Behavioral
abnormalities

[130779]and selective neuronal loss in HD transgenic mice ex-
pressing mutated

[130780]full-length HD cDNA. Nature Genet. 20: 198–202, 1998.

[130781]

[130782]220. Reed, T. E.; Chandler, J. H.: Huntington's chorea in Michigan.

[130783]I. Demography and genetics. Am. J. Hum. Genet. 10: 203–225, 1958.

[130784]

[130785]221. Reed, T. E.; Neel, J. V.: Huntington's chorea in Michigan. II.

[130786]Selection and mutation. Am. J. Hum. Genet. 11: 107–136, 1959.

[130787]

[130788]222. Reik, W.: Genomic imprinting: a possible mechanism for the parental

[130789]origin effect in Huntington's chorea. J. Med. Genet. 25: 805–808,

[130790]1988.

[130791]

[130792]223. Reik, W.: Genomic imprinting and genetic disorders in man. Trends

[130793]Genet. 5: 331–336, 1989.

[130794]

[130795]224. Reik, W.; Collick, A.; Norris, M. L.; Barton, S. C.;
Surani,

[130796]M. A.: Genomic imprinting determines methylation of
parental alleles

[130797]in transgenic mice. *Nature* 328: 248–251, 1987.

[130798]

[130799]225. Reiner, A.; Albin, R. L.; Anderson, K. D.; D'Amato, C.
J.; Penney,

[130800]J. B.; Young, A. B.: Differential loss of striatal projection
neurons

[130801]in Huntington disease. *Proc. Nat. Acad. Sci.* 85:
5733–5737, 1988.

[130802]

[130803]226. Richfield, E. K.; Herkenham, M.: Selective vulnerabil-
ity in

[130804]Huntington's disease: preferential loss of cannabinoid re-
ceptors in

[130805]lateral globus pallidus. *Ann. Neurol.* 36: 577–584, 1994.

[130806]

[130807]227. Ridley, R. M.; Frith, C. D.; Crow, T. J.; Conneally, P.

M.:

[130808]Anticipation in Huntington's disease is inherited through the male

[130809]line but may originate in the female. J. Med. Genet. 25: 589-595,

[130810]1988.

[130811]

[130812]228. Ridley, R. M.; Frith, C. D.; Farrer, L. A.; Conneally, P. M.

[130813]: Patterns of inheritance of the symptoms of Huntington's disease

[130814]suggestive of an effect of genomic imprinting. J. Med. Genet. 28:

[130815]224-231, 1991.

[130816]

[130817]229. Robbins, C.; Theilmann, J.; Youngman, S.; Haines, J.; Altherr,

[130818]M. J.; Harper, P. S.; Payne, C.; Junker, A.; Wasmuth, J.; Hayden,

[130819]M. R.: Evidence from family studies that the gene causing Huntington

[130820]disease is telomeric to D4S95 and D4S90. Am. J. Hum.

Genet. 44:

[130821]422-425, 1989.

[130822]

[130823]230. Rosenberg, N. K.; Sorensen, S. A.; Christensen, A.-L.:
Neuropsychological

[130824]characteristics of Huntington's disease carriers: a double
blind study. J.

[130825]Med. Genet. 32: 600-604, 1995.

[130826]

[130827]231. Rosenblatt, A.; Brinkman, R. R.; Liang, K. Y.;
Almqvist, E. W.;

[130828]Margolis, R. L.; Huang, C. Y.; Sherr, M.; Franz, M. L.; Ab-
bott, M.

[130829]H.; Hayden, M. R.; Ross, C. A.: Familial influence on age of
onset

[130830]among siblings with Huntington disease. Am. J. Med.
Genet. 105:

[130831]399-403, 2001.

[130832]

[130833]232. Roses, A. D.: From genes to mechanisms to thera-
pies: lessons

[130834]to be learned from neurological disorders. Nature Med. 2:
267–269,
[130835]1996.

[130836]

[130837]233. Rubinsztein, D. C.; Amos, W.; Leggo, J.; Goodburn,
S.; Ramesar,

[130838]R. S.; Old, J.; Bontrop, R.; McMahon, R.; Barton, D. E.; Fer-
guson-Smith,

[130839]M. A.: Mutational bias provides a model for the evolution
of Huntington's

[130840]disease and predicts a general increase in disease preva-
lence. Nature

[130841]Genet. 7: 525–530, 1994.

[130842]

[130843]234. Rubinsztein, D. C.; Leggo, J.; Coles, R.; et al.: Pheno-
typic

[130844]characterization of individuals with 30–40 CAG repeats in
the Huntington

[130845]disease (HD) gene reveals HD cases with 36 repeats and
apparently

[130846]normal elderly individuals with 36–39 repeats. Am. J.
Hum. Genet. 59:

[130847]16–22, 1996.

[130848]

[130849]235. Sabl, J. F.; Laird, C. D.: Epigene conversion: a proposal with

[130850]implications for gene mapping in humans. *Am. J. Hum. Genet.* 50:

[130851]1171–1177, 1992.

[130852]

[130853]236. Saccone, S.; De Sario, A.; Della Valle, G.; Bernardi, G.:
The

[130854]highest gene concentrations in the human genome are in
telomeric bands

[130855]of metaphase chromosomes. *Proc. Nat. Acad. Sci.* 89:
4913–4917, 1992.

[130856]

[130857]237. Sapienza, C.; Peterson, A. C.; Rossant, J.; Balling, R.:
Degree

[130858]of methylation of transgenes is independent of gamete of
origin. *Nature* 328:

[130859]251–254, 1987.

[130860]

- [130861]238. Sapp, E.; Ge, P.; Aizawa, H.; Bird, E.; Penney, J.;
Young, A.
[130862]B.; Vonsattel, J.-P.; DiFiglia, M.: Evidence for a preferential
loss
[130863]of enkephalin immunoreactivity in the external globus
pallidus in
[130864]low grade Huntington's disease using high resolution im-
age analysis. Neuroscience 64:
[130865]397-404, 1995.
[130866]
[130867]239. Sathasivam, K.; Hobbs, C.; Turmaine, M.; Mangiarini,
L.; Mahal,
[130868]A.; Bertaux, F.; Wanker, E. E.; Doherty, P.; Davies, S. W.;
Bates,
[130869]G. P.: Formation of polyglutamine inclusions in non-CNS
tissue. Hum.
[130870]Molec. Genet. 8: 813-822, 1999.
[130871]
[130872]240. Sathasivam, K.; Woodman, B.; Mahal, A.; Bertaux, F.;
Wanker,
[130873]E. E.; Shima, D. T.; Bates, G. P.: Centrosome disorganiza-
tion in

[130874] fibroblast cultures derived from R6/2 Huntington's disease (HD) transgenic

[130875] mice and HD patients. Hum. Molec. Genet. 10: 2425–2435, 2001.

[130876]

[130877] 241. Saudou, F.; Finkbeiner, S.; Devys, D.; Greenberg, M. E.: Huntingtin

[130878] acts in the nucleus to induce apoptosis but death does not correlate

[130879] with the formation of intranuclear inclusions. Cell 95: 55–66, 1998.

[130880]

[130881] 242. Sax, D. S.; Bird, E. D.; Gusella, J. F.; Myers, R. H.: Phenotypic

[130882] variation in 2 Huntington's disease families with linkage to chromosome

[130883] 4. Neurology 39: 1332–1336, 1989.

[130884]

[130885] 243. Scherzinger, E.; Lurz, R.; Turmaine, M.; Mangiarini, L.; Hollenbach,

[130886] B.; Hasenbank, R.; Bates, G. P.; Davies, S. W.; Lehrach, H.; Wanker,

[130887]E. E.: Huntingtin-encoded polyglutamine expansions form amyloid-like

[130888]protein aggregates in vitro and in vivo. Cell 90: 549–558, 1997.

[130889]

[130890]244. Scherzinger, E.; Sittler, A.; Schweiger, K.; Heiser, V.; Lurz,

[130891]R.; Hasenbank, R.; Bates, G. P.; Lehrach, H.; Wanker, E. E.: Self-assembly

[130892]of polyglutamine-containing huntingtin fragments into amyloid-like

[130893]fibrils: implications for Huntington's disease pathology. Proc. Nat.

[130894]Acad. Sci. 96: 4604–4609, 1999.

[130895]

[130896]245. Schilling, G.; Becher, M. W.; Sharp, A. H.; Jinnah, H. A.; Duan,

[130897]K.; Kotzuk, J. A.; Slunt, H. H.; Ratovitski, T.; Cooper, J. K.; Jenkins,

[130898]N. A.; Copeland, N. G.; Price, D. L.; Ross, C. A.; Borchelt, D. R.

[130899]: Intracellular inclusions and neuritic aggregates in trans-

genic mice

[130900]expressing a mutant N-terminal fragment of huntingtin.

Hum. Molec.

[130901]Genet. 8: 397-407, 1999.

[130902]

[130903]246. Schwarcz, R.; Okuno, E.; White, R. J.; Bird, E. D.;

Whetsell,

[130904]W. O., Jr.: 3-Hydroxyanthranilate oxygenase activity is increased

[130905]in the brains of Huntington disease victims. Proc. Nat.

Acad. Sci. 85:

[130906]4079-4081, 1988.

[130907]

[130908]247. Scrimgeour, E. M.: Possible introduction of Huntington's chorea

[130909]into Pacific Islands by New England whalers. Am. J. Med.

Genet. 15:

[130910]607-613, 1983.

[130911]

[130912]248. Scrimgeour, E. M.; Samman, Y.; Brock, D. J. H.: Huntington's

[130913]disease in a Sudanese family from Khartoum. Hum. Genet.
96: 624–625,
[130914]1995.

[130915]

[130916]249. Shelbourne, P. F.; Killeen, N.; Hevner, R. F.; Johnston,
H. M.;

[130917]Tecott, L.; Lewandoski, M.; Ennis, M.; Ramirez, L.; Li, Z.;
Iannicola,

[130918]C.; Littman, D. R.; Myers, R. M.: A Huntington's disease
CAG expansion

[130919]at the murine Hdh locus is unstable and associated with
behavioural

[130920]abnormalities in mice. Hum. Molec. Genet. 8: 763–774,
1999.

[130921]

[130922]250. Shiwach, R.: Psychopathology in Huntington's disease
patients. Acta

[130923]Psychiat. Scand. 90: 241–246, 1994.

[130924]

[130925]251. Shiwach, R. S.; Norbury, C. G.: A controlled psychi-
atric study

[130926] of individuals at risk for Huntington's disease. Brit. J. Psychiatry. 165:

[130927] 500–505, 1994.

[130928]

[130929] 252. Silber, E.; Kromberg, J.; Temlett, J. A.; Krause, A.; Saffer,

[130930] D.: Huntington's disease confirmed by genetic testing in five African

[130931] families. Mov. Disord. 13: 726–730, 1998.

[130932]

[130933] 253. Simpson, S. A.; Johnston, A. W.: The prevalence and patterns

[130934] of care of Huntington's chorea in Grampian. Brit. J. Psychiatry. 155:

[130935] 799–804, 1989.

[130936]

[130937] 254. Sisodia, S. S.: Nuclear inclusions in glutamine repeat disorders:

[130938] are they pernicious, coincidental, or beneficial? Cell 95: 1–4,

[130939] 1998.

[130940]

[130941]255. Sittler, A.; Lurz, R.; Lueder, G.; Priller, J.; Hayer-Hartl,

[130942]M. K.; Hartl, F. U.; Lehrach, H.; Wanker, E. E.: Gel-

danamycin activates

[130943]a heat shock response and inhibits huntingtin aggregation

in a cell

[130944]culture model of Huntington's disease. Hum. Molec.

Genet. 10: 1307-1315,

[130945]2001.

[130946]

[130947]256. Skraastad, M. I.; Van de Vosse, E.; Belfroid, R.; Hold,

K.; Vegter-van

[130948]der Vlis, M.; Sandkuijl, L. A.; Bakker, E.; van Ommen, G. J.

B.:

[130949]Significant linkage disequilibrium between the Huntington

disease

[130950]gene and the loci D4S10 and D4S95 in the Dutch popula-

tion. Am. J.

[130951]Hum. Genet. 51: 730-735, 1992.

[130952]

[130953]257. Snell, R. G.; MacMillan, J. C.; Cheadle, J. P.; Fenton, I.;

Lazarou,

[130954]L. P.; Davies, P.; MacDonald, M. E.; Gusella, J. F.; Harper, P. S.;

[130955]Shaw, D. J.: Relationship between trinucleotide repeat expansion

[130956]and phenotypic variation in Huntington's disease. Nature Genet. 4:

[130957]393–397, 1993.

[130958]

[130959]258. Snell, R. G.; Youngman, S.; Lehrach, H.; Sarfarazi, M.; Harper,

[130960]P. S.; Shaw, D. J.: A new probe (2R3) in the region of Huntington's

[130961]disease. (Abstract) Cytogenet. Cell Genet. 51: 1083, 1989.

[130962]

[130963]259. Stebbins, C. E.; Russo, A. A.; Schneider, C.; Rosen, N.; Hartl,

[130964]F. U.; Pavletich, N. P.: Crystal structure of an Hsp90–geldanamycin

[130965]complex: targeting of a protein chaperone by an antitumor agent. Cell 89:

[130966]239–250, 1997.

[130967]

[130968]260. Steffan, J. S.; Bodal, L.; Pallos, J.; Poelman, M.; Mc-Campbell,

[130969]A.; Apostol, B. L.; Kazantsev, A.; Schmidt, E.; Zhu, Y.-Z.; Greenwald,

[130970]M.; Kurokawa, R.; Housman, D. E.; Jackson, G. R.; Marsh, J. L.; Thompson,

[130971]L. M.: Histone deacetylase inhibitors arrest polyglutamine-dependent

[130972]neurodegeneration in *Drosophila*. *Nature* 413: 739-743, 2001.

[130973]

[130974]261. Steffan, J. S.; Kazantsev, A.; Spasic-Boskovic, O.; Greenwald,

[130975]M.; Zhu, Y.-Z.; Gohler, H.; Wanker, E. E.; Bates, G. P.; Housman,

[130976]D. E.; Thompson, L. M.: The Huntington's disease protein interacts

[130977]with p53 and CREB-binding protein and represses transcription. *Proc.*

[130978]*Nat. Acad. Sci.* 97: 6763-6768, 2000.

[130979]

[130980]262. Stine, O. C.; Smith, K. D.: The estimation of selection coefficients

[130981]in Afrikaners: Huntington disease, porphyria variegata, and lipoid

[130982]proteinosis. Am. J. Hum. Genet. 46: 452–458, 1990.

[130983]

[130984]263. Strobel, S. A.; Doucette–Stamm, L. A.; Riba, L.; Housman, D.

[130985]E.; Dervan, P. B.: Site–specific cleavage of human chromosome 4 mediated

[130986]by triple–helix formation. Science 254: 1639–1642, 1991.

[130987]

[130988]264. Tolmie, J. L.; Davidson, H. R.; May, H. M.; McIntosh, K.; Paterson,

[130989]J. S.; Smith, B.: The prenatal exclusion test for Huntington's disease:

[130990]experience in the West of Scotland, 1986–1993. J. Med. Genet. 32:

[130991]97–101, 1995.

[130992]

[130993]265. Tranebjaerg, L.; Petersen, A.; Hove, K.; Rehder, H.;

Mikkelsen,

[130994]M.: Clinical and cytogenetic studies in a large (4;8)
translocation

[130995]family with pre- and postnatal Wolf syndrome. Ann.
Genet. 27: 224-229,

[130996]1984.

[130997]

[130998]266. Trettel, F.; Rigamonti, D.; Hilditch-Maguire, P.;
Wheeler, V.

[130999]C.; Sharp, A. H.; Persichetti, F.; Cattaneo, E.; MacDonald,
M. E.

[131000]: Dominant phenotypes produced by the HD mutation in
STHdh(Q111) striatal

[131001]cells. Hum. Molec. Genet. 9: 2799-2809, 2000.

[131002]

[131003]267. Trottier, Y.; Biancalana, V.; Mandel, J.-L.: Instability
of

[131004]CAG repeats in Huntington's disease: relation to parental
transmission

[131005]and age of onset. J. Med. Genet. 31: 377-382, 1994.

[131006]

- [131007]268. Trottier, Y.; Devys, D.; Imbert, G.; Saudou, F.; An, I.;
Lutz,
[131008]Y.; Weber, C.; Agid, Y.; Hirsch, E. C.; Mandel, J.-L.: Cellular
localization
[131009]of the Huntington's disease protein and discrimination of
the normal
[131010]and mutated form. *Nature Genet.* 10: 104–110, 1995.
[131011]
[131012]269. Tyler, A.; Quarrell, O. W. J.; Lazarou, L. P.; Meredith,
A. L.;
[131013]Harper, P. S.: Exclusion testing in pregnancy for Hunting-
ton's disease. *J.*
[131014]*Med. Genet.* 27: 488–495, 1990.
[131015]
[131016]270. van Dellen, A.; Blakemore, C.; Deacon, R.; York, D.;
Hannan,
[131017]A. J.: Delaying the onset of Huntington's in mice: this un-
remitting
[131018]disease develops later in animals stimulated by their envi-
ronment.
[131019](Letter) *Nature* 404: 721–722, 2000.

[131020]

[131021]271. van der Weiden, R. M. F.: George Huntington and
George Sumner

[131022]Huntington: a tale of two doctors. (Letter) Hist. Phil. Life
Sci. 11:

[131023]297–304, 1989.

[131024]

[131025]272. van Dijk, J. G.; van der Velde, E. A.; Roos, R. A. C.;
Bruyn,

[131026]G. W.: Juvenile Huntington disease. Hum. Genet. 73:
235–239, 1986.

[131027]

[131028]273. Vessie, P. R.: Original article on the transmission of
Huntington's

[131029]chorea for 300 years--the Bures family group. J. Nerv.
Ment. Dis. 76:

[131030]553–573, 1932.

[131031]

[131032]274. Volkers, W. S.; Went, L. N.; Vegter–van der Vlis, M.;
Harper,

[131033]P. S.; Caro, A.: Genetic linkage studies in Huntington's

chorea. Ann.

[131034]Hum. Genet. 44: 75–79, 1980.

[131035]

[131036]275. Walker, D. A.; Harper, P. S.; Wells, C. E. C.; Tyler, A.;
Davies,

[131037]K.; Newcombe, R. G.: Huntington's chorea in South Wales:
a genetic

[131038]and epidemiological study. Clin. Genet. 19: 213–221,
1981.

[131039]

[131040]276. Wallace, D. C.; Hall, A. C.: Evidence of genetic het-
erogeneity

[131041]in Huntington's chorea. J. Neurol. Neurosurg. Psychiat. 35:
789–800,

[131042]1972.

[131043]

[131044]277. Wang, H. S.; Greenberg, C. R.; Hewitt, J.; Kalousek,
D.; Hayden,

[131045]M. R.: Subregional assignment of the linked marker G8
(D4S10) for

[131046]Huntington disease to chromosome 4p16.1–16.3. Am. J.

Hum. Genet. 39:

[131047]392-396, 1986.

[131048]

[131049]278. Wang, H. S.; Greenberg, C. R.; Kalousek, D.; Gusella,
J.; Horsman,

[131050]D.; Hayden, M. R.: Subregional assignment of the linked
marker D4S10

[131051](G8) for Huntington disease by in situ hybridization.
(Abstract) Cytogenet.

[131052]Cell Genet. 40: 772, 1985.

[131053]

[131054]279. Warner, J.; Barron, L.; St Clair, D.; Brock, D.: Reliabil-
ity

[131055]of clinical diagnosis of Huntington's disease. (Letter) J.
Neurol.

[131056]Neurosurg. Psychiat. 63: 1277, 1994.

[131057]

[131058]280. Wasmuth, J. J.; Hewitt, J.; Smith, B.; Allard, D.; Haines,
J.

[131059]L.; Skarecky, D.; Partlow, E.; Hayden, M. R.: A highly poly-
morphic

[131060] locus very tightly linked to the Huntington's disease gene.

Nature 332:

[131061] 734–736, 1988.

[131062]

[131063] 281. Went, L. N.; Vegter–van der Vlis, M.; Bruyn, G. W.:

Parental

[131064] transmission in Huntington's disease. Lancet I:

1100–1102, 1984.

[131065]

[131066] 282. Wexler, N. S.: The Tiresias complex: Huntington's disease as

[131067] a paradigm of testing for late–onset disorders. FASEB J. 6: 2820–2825,

[131068] 1992.

[131069]

[131070] 283. Wexler, N. S.; Bonilla, E.; Young, A. B.; Shoulson, I.; Gomez,

[131071] F.; Starosta, S.; Travers, H.; Villalobas, M.; de Quiroz, I.; Erbe,

[131072] R.; Penney, J. B.; Uzzell, R. S.; Burnham, F. A.; Daugherty, L.; Jones,

- [131073]B.; Mapstone, C.; Rivas, M.; Messer, E.; Wexler, A.; Snodgrass, R.;
- [131074]Rosenzweig, G.; Esteves, J.; Marsol, N.; Bailey, S.; Brinley, F. J.;
- [131075]Goldstein, E.; Greene, A. E.; Kidd, J. R.; Kidd, K. K.; Gusella, J.
- [131076]F.; Conneally, P. M.; Moreno, H.: Huntington's disease in Venezuela
- [131077]and gene linkage. (Abstract) Cytogenet. Cell Genet. 37: 605, 1984.
- [131078]
- [131079]284. Wexler, N. S.; Young, A.; Tanzi, R.; Starosta, S.; Gomez, F.;
- [131080]Travers, H.; Snodgrass, S. R.; Moreno, H.; Shoulson, I.; Penney, J.;
- [131081]Conneally, P. M.; Gusella, J.: Huntington's disease homozygotes identified.
- [131082](Abstract) Am. J. Hum. Genet. 37: A82, 1985.
- [131083]
- [131084]285. Wexler, N. S.; Young, A. B.; Tanzi, R. E.; Travers, H.; Starosta-Rubinstein,
- [131085]S.; Penney, J. B.; Snodgrass, S. R.; Shoulson, I.; Gomez, F.;

Ramos

[131086]Arroyo, M. A.; Penchaszadeh, G. K.; Moreno, H.; Gibbons, K.; Faryniarz,

[131087]A.; Hobbs, W.; Anderson, M. A.; Bonilla, E.; Conneally, P. M.; Gusella,

[131088]J. F.: Homozygotes for Huntington's disease. *Nature* 326: 194–197,

[131089]1987.

[131090]

[131091]286. Wheeler, V. C.; Gutekunst, C.–A.; Vrbanc, V.; Lebel, L.–A.;

[131092]Schilling, G.; Hersch, S.; Friedlander, R. M.; Gusella, J. F.; Vonsattel,

[131093]J.–P.; Borchelt, D. R.; MacDonald, M. E.: Early phenotypes that presage

[131094]late-onset neurodegenerative disease allow testing of modifiers in

[131095]Hdh CAG knock-in mice. *Hum. Molec. Genet.* 11: 633–640, 2002.

[131096]

[131097]287. Wheeler, V. C.; White, J. K.; Gutekunst, C.–A.; Vrbanc, V.;

- [131098]Weaver, M.; Li, X. J.; Li, S.-H.; Yi, H.; Vonsattel, J.-P.;
Gusella,
[131099]J. F.; Hersch, S.; Auerbach, W.; Joyner, A. L.; MacDonald,
M. E.:
[131100]Long glutamine tracts cause nuclear localization of a novel
form of
[131101]huntingtin in medium spiny striatal neurons in Hdh-Q92
and Hdh-Q111
[131102]knock-in mice. Hum. Molec. Genet. 9: 503-513, 2000.
[131103]
[131104]288. Wiggins, S.; Whyte, P.; Huggins, M.; Adam, S.; Theil-
mann, J.;
[131105]Bloch, M.; Sheps, S. B.; Schechter, M. T.; Hayden, M. R.:
The psychological
[131106]consequences of predictive testing for Huntington's dis-
ease. New
[131107]Eng. J. Med. 327: 1401-1405, 1992.
[131108]
[131109]289. Wolff, G.; Deuschl, G.; Wienker, T. F.; Hummel, K.;
Bender, K.;
[131110]Lucking, C. H.; Schumacher, M.; Hammer, J.; Oepen, G.:
New mutation

[131111]to Huntington's disease. J. Med. Genet. 26: 18-27, 1989.

[131112]

[131113]290. Wright, H. H.; Still, C. N.; Abramson, R. K.: Hunting-
ton's disease

[131114]in black kindreds in South Carolina. Arch. Neurol. 38:
412-414,

[131115]1981.

[131116]

[131117]291. Xiang, F.; Almqvist, E. W.; Huq, M.; Lundin, A.; Hay-
den, M. R.;

[131118]Edstrom, L.; Anvret, M.; Zhang, Z.: A Huntington disease-
like neurodegenerative

[131119]disorder maps to chromosome 20p. Am. J. Hum. Genet.
63: 1431-1438,

[131120]1998.

[131121]

[131122]292. Yamamoto, A.; Lucas, J. J.; Hen, R.: Reversal of neu-
ropathology

[131123]and motor dysfunction in a conditional model of Hunting-
ton's disease. Cell 101:

[131124]57-66, 2000.

[131125]

[131126]293. Youngman, S.; Sarfarazi, M.; Quarrell, O. W. J.; Conneally, P.

[131127]M.; Gibbons, K.; Harper, P. S.; Shaw, D. J.; Tanzi, R. E.; Wallace,

[131128]M. R.; Gusella, J. F.: Studies of a DNA marker (G8) genetically linked

[131129]to Huntington disease in British families. Hum. Genet. 73: 333-339,

[131130]1986.

[131131]

[131132]294. Zabel, B. U.; Naylor, S. L.; Sakaguchi, A. Y.; Gusella, J. F.

[131133]: Regional localization of a DNA polymorphism (D4S10) linked to Huntington's

[131134]disease at 4p16-p15. (Abstract) Cytogenet. Cell Genet. 40: 787,

[131135]1985.

[131136]

[131137]295. Zabel, B. U.; Naylor, S. L.; Sakaguchi, A. Y.; Gusella, J. F.

[131138]: Mapping of the DNA locus D4S10 and the linked Huntington's disease

[131139] gene to 4p16-p15. Cytogenet. Cell Genet. 42: 187-190, 1986.

[131140]

[131141] 296. Zeitlin, S.; Liu, J.-P.; Chapman, D. L.; Papaioannou, V. E.;

[131142] Efstratiadis, A.: Increased apoptosis and early embryonic lethality

[131143] in mice nullizygous for the Huntington's disease gene homologue. Nature

[131144] Genet. 11: 155-163, 1995.

[131145]

[131146] 297. Zlotogora, J.: Dominance and homozygosity. Am. J. Med. Genet. 68:

[131147] 412-416, 1997.

[131148]

[131149] 298. Zuccato, C.; Ciammola, A.; Rigamonti, D.; Leavitt, B. R.; Goffredo,

[131150] D.; Conti, L.; MacDonald, M. E.; Friedlander, R. M.; Silani, V.; Hayden,

[131151]M. R.; Timmusk, T.; Sipione, S.; Cattaneo, E.: Loss of huntingtin-mediated

[131152]BDNF gene transcription in Huntington's disease. Science 293: 493–498,

[131153]2001.

[131154]

[131155]299. Zuhlke, C.; Riess, O.; Bockel, B.; Lange, H.; Thies, U.: Mitotic

[131156]stability and meiotic variability of the (CAG)_n repeat in the Huntington

[131157]disease gene. Hum. Molec. Genet. 2: 2063–2067, 1993.

[131158]

[131159]300. Zuo, J.; Robbins, C.; Taillon-Miller, P.; Cox, D. R.; Myers,

[131160]R. M.: Cloning of the Huntington disease region in yeast artificial

[131161]chromosomes. Hum. Molec. Genet. 1: 149–159, 1992.

[131162]23338.

[131163]Calandro, L. M.; Baer, D. M.; Sensabaugh, G. F.: Characterization

[131164]of a recombinant that locates the hereditary hemochro-

matosis gene

[131165]telomeric to HLA-F. Hum. Genet. 96: 339-342, 1995.

[131166]23339.

[131167]Campbell, K.; Flavin, N.; Ivens, A.; Robert, B.; Buckingham, M.;

[131168]Williamson, R.: The human homeobox gene HOX7 maps to 4p16.1 and is

[131169]deleted in Wolf-Hirschhorn syndrome patients. (Abstract) Am. J. Hum.

[131170]Genet. 45 (suppl.): A179 only, 1989.

[131171]23340.

[131172]Gehring, W. J.; Qian, Y. Q.; Billeter, M.; Furukubo-Tokunaga, K.;

[131173]Schier, A. F.; Resendez-Perez, D.; Affolter, M.; Otting, G.; Wuthrich,

[131174]K.: Homeodomain-DNA recognition. Cell 78: 211-223, 1994.

[131175]23341.

[131176]Greenberg, F.; Elder, F. F. B.; Ledbetter, D. H.; Altherr, M.;

[131177]Wasmuth, J. J.: Molecular confirmation of Wolf-Hirschhorn syndrome

[131178]with apparently normal chromosomes. (Abstract) Am. J.
Hum. Genet. 45

[131179](suppl.): A76 only, 1989.

[131180]23342.

[131181]Hewitt, J. E.; Clark, L. N.; Ivens, A.; Williamson, R.: Struc-
ture

[131182]and sequence of the human homeobox gene HOX7. Ge-
nomics 11: 670-678,

[131183]1991.

[131184]23343.

[131185]Hirschhorn, K.; Cooper, H. L.; Firschein, I. L.: Deletion of
short

[131186]arms of chromosome 4-5 in a child with defects of mid-
line fusion. Humangenetik 1:

[131187]479-482, 1965.

[131188]23344.

[131189]Hu, G.; Vastardis, H.; Bendall, A. J.; Wang, Z.; Logan, M.;
Zhang,

[131190]H.; Nelson, C.; Stein, S.; Greenfield, N.; Seidman, C. E.;
Seidman,

[131191]J. G.; Abate-Shen, C.: Haploinsufficiency of MSX1: a mech-

anism for

[131192]selective tooth agenesis. *Molec. Cell. Biol.* 18: 6044–6051,
1998.

[131193]23345.

[131194]Hwang, S.-J.; Beaty, T. H.; McIntosh, I.; Hefferon, T.;
Panny,

[131195]S. R.: Association between homeobox-containing gene
MSX1 and the

[131196]occurrence of limb deficiency. *Am. J. Med. Genet.* 75:
419–423, 1998.

[131197]23346.

[131198]Ivens, A.; Flavin, N.; Williamson, R.; Dixon, M.; Bates, G.;
Buckingham,

[131199]M.; Robert, B.: The human homeobox gene HOX7 maps to
chromosome 4p16.1

[131200]and may be implicated in Wolf–Hirschhorn syndrome.
Hum. Genet. 84:

[131201]473–476, 1990.

[131202]23347.

[131203]Jumlongras, D.; Bei, M.; Stimson, J. M.; Wang, W.-F.; De-
Palma,

[131204]S. R.; Seidman, C. E.; Felbor, U.; Maas, R.; Seidman, J. G.;

Olsen,

[131205]B. R.: A nonsense mutation in MSX1 causes Witkop syndrome. Am. J.

[131206]Hum. Genet. 69: 67–74, 2001.

[131207]23348.

[131208]Odelberg, S. J.; Kollhoff, A.; Keating, M. T.: Dedifferentiation

[131209]of mammalian myotubes induced by msx1. Cell 103: 1099–1109, 2000.

[131210]23349.

[131211]Robert, B.; Sassoon, D.; Jacq, B.; Gehring, W.; Buckingham, M.

[131212]: Hox-7, a mouse homeobox gene with a novel pattern of expression

[131213]during embryogenesis. EMBO J. 8: 91–100, 1989.

[131214]23350.

[131215]Satokata, I.; Maas, R.: Msx1 deficient mice exhibited cleft palate

[131216]and abnormalities of craniofacial and tooth development. Nature Genet. 6:

[131217]348–355, 1994.

[131218]23351.

[131219]Scarel, R. M.; Trevilatto, P. C.; Di Hipolito, O., Jr.; Ca-
margo,

[131220]L. E. A.; Line, S. R. P.: Absence of mutations in the home-
odomain

[131221]of the MSX1 gene in patients with hypodontia. Am. J. Med.
Genet. 92:

[131222]346–349, 2000.

[131223]23352.

[131224]Stimson, J. M.; Sivers, J. E.; Hlava, G. L.: Features of
oligodontia

[131225]in three generations. J. Clin. Pediat. Dent. 21: 269–275,
1997.

[131226]23353.

[131227]van den Boogaard, M.–J. H.; Dorland, M.; Beemer, F. A.;
Ploos

[131228]van Amstel, H. K.: MSX1 mutation is associated with oro-
facial clefting

[131229]and tooth agenesis in humans. (Letter) Nature Genet. 24:
342–343,

[131230]2000. Note: Erratum: Nature Genet. 25: 125 only, 2000.

[131231]23354.

[131232]Vastardis, H.; Karimbux, N.; Guthua, S. W.; Seidman, J. G.;
Seidman,

[131233]C. E.: A human MSX1 homeodomain missense mutation
causes evidence

[131234]tooth agenesis. Nature Genet. 13: 417–421, 1996.

[131235]23355.

[131236]Wolf, U.; Reinwein, H.; Porsch, R.; Schroter, R.; Baitsch, H.

[131237]: Defizienz an den kurzen Armen eines Chromosoms Nr.
4. Humangenetik 1:

[131238]397–413, 1965.

[131239]23356.

[131240]Del Campo, M.; Jones, M. C.; Veraksa, A. N.; Curry, C. J.;
Jones,

[131241]K. L.; Mascarello, J. T.; Ali-Kahn-Catts, Z.; Drumheller, T.;
McGinnis,

[131242]W.: Monodactylous limbs and abnormal genitalia are asso-
ciated with

[131243]hemizygosity for the human 2q31 region that includes the
HOXD cluster. Am.

[131244]J. Hum. Genet. 65: 104–110, 1999.

[131245]23357.

[131246]Zakany, J.; Kmita, M.; Alarcon, P.; de la Pompa, J.-L.;
Duboule,

[131247]D.: Localized and transient transcription of Hox genes
suggests a

[131248]link between patterning and the segmentation clock. Cell
106: 207–217,

[131249]2001.

[131250]23358.

[131251]Lafuse, W. P.; Zwilling, B. S.: Localization of the inhibin
beta-B

[131252]gene on mouse chromosome 1. Mammalian Genome 4:
399–400, 1993.

[131253]23359.

[131254]Salmenkivi, K.; Arola, J.; Voutilainen, R.; Ilvesmaki, V.;
Haglund,

[131255]C.; Kahri, A. I.; Heikkila, P.; Liu, J.: Inhibin/activin beta-
B-subunit

[131256]expression in pheochromocytomas favors benign diagno-
sis. J. Clin.

[131257]Endocr. Metab. 86: 2231–2235, 2001.

[131258]23360.

[131259]Schrewe, H.; Gendron-Maguire, M.; Harbison, M. L.; Gridley, T.

[131260]: Mice homozygous for a null mutation of activin beta B are viable

[131261]and fertile. Mech. Dev. 47: 43-51, 1994.

[131262]23361.

[131263]Kotanides, H.; Reich, N. C.: Interleukin-4-induced STAT6 recognizes

[131264]and activates a target site in the promoter of the interleukin-4 receptor

[131265]gene. J. Biol. Chem. 271: 25555-25561, 1996.

[131266]23362.

[131267]Caggana, M.; Walker, K.; Reilly, A. A.; Conroy, J. M.; Duva, S.;

[131268]Walsh, A. C.: Population-based studies reveal differences in the

[131269]allelic frequencies of two functionally significant human interleukin-4

[131270]receptor polymorphisms in several ethnic groups. Genet. Med. 1:

[131271]267-271, 1999.

[131272]23363.

[131273]Deichmann, K. A.; Heinzmann, A.; Forster, J.; Dischinger, S.; Mehl,

[131274]C.; Brueggemolte, E.; Hildebrandt, F.; Moseler, M.; Kuehr, J.: Linkage

[131275]and allelic association of atopy and markers flanking the IL4-receptor

[131276]gene. Clin. Exp. Allergy 28: 151-155, 1998.

[131277]23364.

[131278]MacLeod, J. N.; Lee, A. K.; Liebhaber, S. A.; Cooke, N. E.: Developmental

[131279]control and alternative splicing of the placentally expressed transcripts

[131280]from the human growth hormone gene cluster. J. Biol. Chem. 267:

[131281]14219-14226, 1992.

[131282]23365.

[131283]Wurzel, J. M.; Parks, J. S.; Herd, J. E.; Nielsen, P. V.: A gene

[131284]deletion is responsible for absence of human chorionic somatomammotropin. DNA 1:

[131285]251-257, 1982.

[131286]23366.

[131287]Gallin, J. I.: Personal Communication. Bethesda, Md.
10/12/1990.

[131288]23367.

[131289]McCombs, J. L.; Teng, C. T.; Pentecost, B. T.; Magnuson, V.
L.;

[131290]Moore, C. M.; McGill, J. R.: Chromosomal localization of
human lactotransferrin

[131291]gene (LTF) by in situ hybridization. Cytogenet. Cell Genet.
47:

[131292]16-17, 1988.

[131293]23368.

[131294]Moriuchi, M.; Moriuchi, H.: A milk protein lactoferrin en-
hances

[131295]human T cell leukemia virus type I and suppresses HIV-1
infection. J.

[131296]Immun. 166: 4231-4236, 2001.

[131297]23369.

[131298]Naylor, S. L.; Marshall, A.; Solomon, A.; McGill, J. R.; Mc-
Combs,

[131299]J.; Magnuson, V. L.; Moore, C. M.; Lalley, P. A.; Pentecost, B. T.;

[131300]Teng, C.: Lactoferrin maps to human chromosome 3(q21-q23) and mouse

[131301]chromosome 9. (Abstract) Cytogenet. Cell Genet. 46: 669 only, 1987.

[131302]23370.

[131303]Powell, M. J.; Ogden, J. E.: Nucleotide sequence of human lactoferrin

[131304]cDNA. Nucleic Acids Res. 18: 4013 only, 1990.

[131305]23371.

[131306]Qiu, J.; Hendrixson, D. R.; Baker, E. N.; Murphy, T. F.; St. Geme,

[131307]J. W., III; Plaut, A. G.: Human milk lactoferrin inactivates two

[131308]putative colonization factors expressed by Haemophilus influenzae. Proc.

[131309]Nat. Acad. Sci. 95: 12641-12646, 1998.

[131310]23372.

[131311]Rado, T. A.; Wei, X.; Benz, E. J., Jr.: Isolation of lactoferrin

[131312]cDNA from a human myeloid library and expression of

mRNA during normal

[131313]and leukemic myelopoiesis. Blood 70: 989–993, 1987.

[131314]23373.

[131315]Singh, P. K.; Parsek, M. R.; Greenberg, E. P.; Welsh, M. J.: A

[131316]component of innate immunity prevents bacterial biofilm
development. Nature 417:

[131317]552–555, 2002.

[131318]23374.

[131319]Teng, C. T.; Pentecost, B. T.; Marshall, A.; Solomon, A.;
Bowman,

[131320]B. H.; Lalley, P. A.; Naylor, S. L.: Assignment of the lacto-
transferrin

[131321]gene to human chromosome 3 and to mouse chromosome
9. Somat. Cell

[131322]Molec. Genet. 13: 689–693, 1987.

[131323]23375.

[131324]Yang, F.; Lum, J.; Baldwin, W. D.; Brune, J. L.; van Bragt, P.;

[131325]Bowman, B. H.: Genetic analysis of human iron binding
glycoproteins.

[131326](Abstract) Am. J. Hum. Genet. 35: 184A only, 1983.

[131327]23376.

[131328]Tanaka, A. R.; Ikeda, Y.; Abe-Dohmae, S.; Arakawa, R.;
Sadanami,

[131329]K.; Kidera, A.; Nakagawa, S.; Nagase, T.; Aoki, R.; Kioka,
N.; Amachi,

[131330]T.; Yokoyama, S.; Ueda, K.: Human ABCA1 contains a large
amino terminal

[131331]extracellular domain homologous to an epitope of Sjo-
gren's syndrome. Biochem.

[131332]Biophys. Res. Commun. 283: 1019-1025, 2001.

[131333]23377.

[131334]Aoki, N.; Suzuki, H.; Ito, K.; Ito, M.: A novel point mutation

[131335]of the FALDH gene in a Japanese family with Sjogren-
Larsson syndrome. J.

[131336]Invest. Derm. 114: 1065-1066, 2000.

[131337]23378.

[131338]Blumel, J.; Watkins, M.; Eggers, G. W. N.: Spastic
quadriplegia

[131339]combined with congenital ichthyosiform erythroderma
and oligophrenia. Am.

[131340]J. Dis. Child. 96: 724-726, 1958.

[131341]23379.

[131342]Chang, C.; Yoshida, A.: Human fatty aldehyde dehydrogenase gene

[131343](ALDH10): organization and tissue-dependent expression. Genomics 40:

[131344]80–85, 1997.

[131345]23380.

[131346]De Laurenzi, V.; Rogers, G. R.; Hamrock, D. J.; Marekov, L. N.;

[131347]Steinert, P. M.; Compton, J. G.; Markova, N.; Rizzo, W. B.: Sjogren–Larsson

[131348]syndrome is caused by mutations in the fatty aldehyde dehydrogenase

[131349]gene. Nature Genet. 12: 52–57, 1996.

[131350]23381.

[131351]De Laurenzi, V.; Rogers, G. R.; Tarcsa, E.; Carney, G.; Marekov,

[131352]L.; Bale, S. J.; Compton, J. G.; Markova, N.; Steinert, P. M.; Rizzo,

[131353]W. B.: Sjogren–Larsson syndrome is caused by a common mutation in

[131354]northern European and Swedish patients. J. Invest. Derm.
109: 79–83,

[131355]1997.

[131356]23382.

[131357]Gedde–Dahl, T., Jr.; Rajka, G.; Larsen, T. E.; Jellum, E.: Au-
tosomal

[131358]recessive ichthyosis in Norway: II. Sjogren–Larsson–like
ichthyosis

[131359]without CNS or eye involvement. Clin. Genet. 26:
242–244, 1984.

[131360]23383.

[131361]Gilbert, W. R., Jr.; Smith, J. L.; Nyhan, W. L.: The Sjogren–
Larsson

[131362]syndrome. Arch. Ophthal. 80: 308–316, 1968.

[131363]23384.

[131364]Goldsmith, L. A.; Baden, H. P.; Canty, T. G.: Sjogren–Lars-
son

[131365]syndrome. Acta Derm. Venerol. 51: 374–378, 1971.

[131366]23385.

[131367]Gustavson, K. H.; Jagell, S.: Dermatoglyphic patterns in the
Sjogren–Larsson

[131368]syndrome. Clin. Genet. 17: 120–124, 1980.

[131369]23386.

[131370]Heijer, A.; Reed, W. B.: Sjogren–Larsson syndrome: congenital

[131371]ichthyosis, spastic paralysis, and oligophrenia. Arch. Derm. 92:

[131372]545–552, 1965.

[131373]23387.

[131374]Jl, L.; Oostheim, W.; van Werkhoven, M.; Willemsen, M. A. A.

[131375]P.; Wanders, R. J. A.: Molecular basis of Sjogren–Larsson syndrome:

[131376]frequency of the 1297–1298 del GA and 943C–T mutation in 29 patients. J.

[131377]Inherit. Metab. Dis. 22: 319–321, 1999.

[131378]23388.

[131379]Jagell, S.; Gustavson, K.–H.; Holmgren, G.: Sjogren–Larsson syndrome

[131380]in Sweden: a clinical, genetic and epidemiological study. Clin. Genet. 19:

[131381]233–256, 1981.

[131382]23389.

[131383]Jagell, S.; Linden, S.: Ichthyosis in the Sjogren-Larsson syndrome. Clin.

[131384]Genet. 21: 243-252, 1982.

[131385]23390.

[131386]Jagell, S.; Polland, W.; Sandgren, O.: Specific changes in the

[131387]fundus typical for the Sjogren-Larsson syndrome: an ophthalmological

[131388]study of 35 patients. Acta Ophthal. 58: 321-330, 1980.

[131389]23391.

[131390]Shohat, M.; Bu, X.; Shohat, T.; Fischel-Ghodsian, N.; Margal, N.;

[131391]Nakamura, Y.; Schwabe, A. D.; Schlezinger, M.; Danon, Y.; Rotter,

[131392]J. I.: The gene for familial Mediterranean fever in both Armenians

[131393]and non-Ashkenazi Jews is linked to the alpha-globin complex on 16p:

[131394]evidence for locus homogeneity. Am. J. Hum. Genet. 51: 1349-1354,

[131395]1992.

[131396]23392.

[131397]Shohat, M.; Korenberg, J. R.; Schlesinger, M.; Shohat, T.;
Petersen,

[131398]G. M.; Schwabe, A.; Rotter, J. I.: The gene for serum amy-
loid P component

[131399](APCS) is not a genetic marker for amyloidosis in familial
Mediterranean

[131400]fever. (Abstract) Cytogenet. Cell Genet. 51: 1079, 1989.

[131401]23393.

[131402]Shohat, M.; Korenberg, J. R.; Schwabe, A. D.; Rotter, J. I.:

[131403]Hypothesis: familial Mediterranean fever--a genetic disorder of the

[131404]lipocortin family? Am. J. Med. Genet. 34: 163-167, 1989.

[131405]23394.

[131406]Shohat, M.; Korenberg, J. R.; Shohat, T.; Sack, G.; Schwabe,
A.;

[131407]Rotter, J. I.: Serum amyloid A (SAA) locus is not associated
with

[131408]familial Mediterranean fever (FMF) or FMF-amyloidosis.
(Abstract) Cytogenet.

[131409]Cell Genet. 51: 1079, 1989.

[131410]23395.

[131411]Shohat, M.; Livneh, A.; Zemer, D.; Pras, M.; Sohar, E.: Twin

[131412]studies in familial Mediterranean fever. Am. J. Med. Genet.
44:

[131413]179–182, 1992.

[131414]23396.

[131415]Shohat, M.; Shohat, T.; Rotter, J. I.; Schlesinger, M.; Pe-
tersen,

[131416]G. M.; Pribyl, T.; Sack, G.; Schwabe, A. D.; Korenberg, J. R.:
Serum

[131417]amyloid A and P protein genes in familial Mediterranean
fever. Genomics 8:

[131418]83–89, 1990.

[131419]

[131420]100. Shohat, T.; Shohat, M.; Petersen, G. M.; Sparkes, R.
S.; Langfield,

[131421]D.; Bickal, J.; Korenberg, J. R.; Schwabe, A. D.; Rotter, J. I.:

[131422]Genetic marker family studies in familial Mediterranean
fever (FMF)

[131423]in Armenians. Clin. Genet. 38: 332–339, 1990.

[131424]

[131425]101. Shohat, T.; Shohat, M.; Tyan, D. B.; Wang, S. J.;
Sparkes, R.

[131426]S.; Schwabe, A. D.; Rotter, J. I.: Familial Mediterranean
fever--linkage

[131427]studies with genetic markers on chromosome 6. Tissue
Antigens 36:

[131428]103–107, 1990.

[131429]

[131430]102. Siegal, S.: Benign paroxysmal peritonitis. Ann. Intern.
Med. 23:

[131431]1–22, 1945.

[131432]

[131433]103. Siegal, S.: Familial paroxysmal polyserositis: analysis
of fifty

[131434]cases. Am. J. Med. 36: 893–918, 1964.

[131435]

[131436]104. Sohar, E.; Gafni, J.; Pras, M.; Heller, H.: Familial
Mediterranean

[131437]fever: a survey of 470 cases and review of literature. Am.
J. Med. 43:

[131438]227-253, 1967.

[131439]

[131440]105. Sokmen, C.: Comment. Trans. Assoc. Am. Phys. 72:
120-121,

[131441]1959.

[131442]

[131443]106. Sood, R.; Blake, T.; Aksentijevich, I.; Wood, G.; Chen,
X.; Gardner,

[131444]D.; Shelton, D. A.; Mangelsdorf, M.; Orsborn, A.; Pras, E.;
Balow,

[131445]J. E. Jr.; Centola, M. and 16 others: Construction of a 1-Mb
restriction-mapped

[131446]cosmid contig containing the candidate region for the fa-
milial Mediterranean

[131447]fever locus (MEFV) on chromosome 16p13.3. Genomics
42: 83-95, 1997.

[131448]

[131449]107. Tamir, N.; Langevitz, P.; Zemer, D.; Pras, E.; Shinar,
Y.; Padeh,

[131450]S.; Zaks, N.; Pras, M.; Livneh, A.: Late-onset familial
Mediterranean

[131451]fever (FMF): a subset with distinct clinical, demographic,
and molecular

[131452]genetic characteristics. Am. J. Med. Genet. 87: 30–35,
1999.

[131453]

[131454]108. Touitou, I.: The spectrum of familial Mediterranean
fever (FMF)

[131455]mutations. Europ. J. Hum. Genet. 9: 473–483, 2001.

[131456]

[131457]109. Tutar, H. E.; Imamoglu, A.; Atalay, S.: Recurrent peri-
carditis

[131458]in familial Mediterranean fever. (Letter) Acta Pediat. 88:
1045–1046,

[131459]1999.

[131460]

[131461]110. Wolff, S. M.; Dinarello, C. A.; Dale, D. C.; Goldfinger,
S. E.;

[131462]Alling, D. W.: Colchicine therapy of familial Mediterranean
fever.

[131463](Abstract) Clin. Res. 22: 567A, 1974.

[131464]

[131465]111. Yazigi, A.; Abou-Charaf, L. C.: Colchicine for recurrent pericarditis

[131466]in children. Acta Paediat. 87: 603-604, 1998.

[131467]

[131468]112. Yilmaz, E.; Ozen, S.; Balci, B.; Duzova, A.; Topaloglu, R.; Besbas,

[131469]N.; Saatci, U.; Bakkaloglu, A.; Ozguc, M.: Mutation frequency of

[131470]familial Mediterranean fever and evidence for a high carrier rate

[131471]in the Turkish population. *Europ. J. Hum. Genet.* 9: 553–555, 2001.

[131472]

[131473]113. Yuval, Y.; Hemo-Zisser, M.; Zemer, D.; Sohar, E.; Pras, M.:

[131474]Dominant inheritance in two families with familial Mediterranean fever

[131475](FMF). *Am. J. Med. Genet.* 57: 455–457, 1995.

[131476]

[131477]114. Zemer, D.; Livneh, A.; Pras, M.; Sohar, E.: Familial Mediterranean

[131478]fever in the colchicine era: the fate of one family. *Am. J. Med.*

[131479]*Genet.* 45: 340–344, 1993.

[131480]

[131481]115. Zemer, D.; Pras, M.; Sohar, E.; Modan, M.; Cabili, S.;
Gafni,

[131482]J.: Colchicine in the prevention and treatment of the amy-
loidosis

[131483]of familial Mediterranean fever. New Eng. J. Med. 314:
1001-1005,

[131484]1986.

[131485]

[131486]116. Zemer, D.; Revach, M.; Pras, M.; Modan, B.; Schor, S.;
Sohar,

[131487]E.; Gafni, J.: A controlled trial of colchicine in preventing
attacks

[131488]of familial Mediterranean fever. New Eng. J. Med. 291:
932-934,

[131489]1974.

[131490]23397.

[131491]Labay, V.; Raz, T.; Baron, D.; Mandel, H.; Williams, H.; Bar-
rett,

[131492]T.; Szargel, R.; McDonald, L.; Shalata, A.; Nosaka, K.; Gre-
gory, S.;

[131493]Cohen, N.: Mutations in SLC19A2 cause thiamine-responsive megaloblastic

[131494]anaemia associated with diabetes mellitus and deafness. Nature Genet. 22:

[131495]300–304, 1999.

[131496]23398.

[131497]Millat, G.; Marcais, C.; Rafi, M. A.; Yamamoto, T.; Morris, J.

[131498]A.; Pentchev, P. G.; Ohno, K.; Wenger, D. A.; Vanier, M. T.: Niemann–Pick

[131499]C1 disease: the I1061T substitution is a frequent mutant allele in

[131500]patients of western European descent and correlates with a classic

[131501]juvenile phenotype. Am. J. Hum. Genet. 65: 1321–1329, 1999.

[131502]23399.

[131503]Agostoni, C.; Massetto, N.; Biasucci, G.; Rottoli, A.; Bonvissuto,

[131504]M.; Bruzzese, M.; Giovannini, M.; Riva, E.: Effects of long-chain

[131505]polyunsaturated fatty acid supplementation on fatty acid status and

- [131506]visual function in treated children with hyperphenylalaninemia. J.
- [131507]Pediat. 137: 504–509, 2000.
- [131508]23400.
- [131509]Eisensmith, R. C.; Okano, Y.; Dasovich, M.; Wang, T.; Guttler,
- [131510]F.; Lou, H.; Guldberg, P.; Lichter-Konecki, U.; Konecki, D. S.; Svensson,
- [131511]E.; Hagenfeldt, L.; Rey, F.; Munnich, A.; Lyonnet, S.; Cockburn, F.;
- [131512]Connor, J. M.; Pembrey, M. E.; Smith, I.; Gitzelmann, R.; Steinmann,
- [131513]B.; Apold, J.; Eiken, H. G.; Giovannini, M.; Riva, E.; Longhi, R.;
- [131514]Romano, C.; Cerone, R.; Naughten, E. R.; Mullins, C.; Cahalane, S.;
- [131515]Ozalp, I.; Fekete, G.; Schuler, D.; Berencsi, G. Y.; Nasz, I.; Brdicka,
- [131516]R.; Kamaryt, J.; Pijackova, A.; Cabalska, B.; Boszkowa, K.; Schwartz,
- [131517]E.; Kalinin, V. N.; Jin, L.; Chakraborty, R.; Woo, S. L. C.: Multiple

[131518]origins for phenylketonuria in Europe. Am. J. Hum. Genet.
51: 1355–1365,
[131519]1992.
[131520]23401.
[131521]Kirk, E. P. E.; Fletcher, J. M.; Sharp, P.; Carey, B.; Poulos,
[131522]A.: X-linked adrenoleukodystrophy: the Australasian ex-
perience. Am.
[131523]J. Med. Genet. 76: 420–423, 1998.
[131524]23402.
[131525]Kobayashi, T.; Noda, S.; Umezaki, H.; Goto, I.; Suzuki, S.;
Kitaguchi,
[131526]T.; Kuroiwa, Y.: Familial spinocerebellar degeneration as
an expression
[131527]of adrenoleukodystrophy. J. Neurol. Neurosurg. Psychiat.
49: 1438–1440,
[131528]1986.
[131529]23403.
[131530]Kobayashi, T.; Yamada, T.; Yasutake, T.; Shinnoh, N.;
Goto, I.;
[131531]Iwaki, T.: Adrenoleukodystrophy gene encodes an 80 kDa
membrane protein. Biochem.

[131532]Biophys. Res. Commun. 201: 1029–1034, 1994.

[131533]23404.

[131534]Kok, F.; Neumann, S.; Sarde, C.–O.; Zheng, S.; Wu, K.–H.;
Wei,

[131535]H.–M.; Bergin, J.; Watkins, P. A.; Gould, S.; Sack, G.; Moser,
H.;

[131536]Mandel, J.–L.; Smith, K. D.: Mutational analysis of patients
with

[131537]X-linked adrenoleukodystrophy. Hum. Mutat. 6: 104–115,
1995.

[131538]23405.

[131539]Kolodny, E. H.: The adrenoleukodystrophy–adreno–
myeloneuropathy

[131540]complex: is it treatable? (Editorial) Ann. Neurol. 21:
230–231,

[131541]1987.

[131542]23406.

[131543]Korenke, G. C.; Fuchs, S.; Krasemann, E.; et al.: Cerebral
adrenoleukodystrophy

[131544](ALD) in only one of monozygous twins with an identical
ALD genotype. Ann.

[131545]Neurol. 40: 254–257, 1996.

[131546]23407.

[131547]Krasemann, E. W.; Meier, V.; Korenke, G. C.; Hunneman, D.
H.;

[131548]Hanefeld, F.: Identification of mutations in the ALD-gene
of 20 families

[131549]with adrenoleukodystrophy/adrenomyeloneuropathy.
Hum. Genet. 97:

[131550]194–197, 1996.

[131551]23408.

[131552]Kruse, B.; Barker, P. B.; van Zijl, P. C. M.; Duyn, J. H.; Moo-
nen,

[131553]C. T. W.; Moser, H. W.: Multislice proton magnetic reso-
nance spectroscopic

[131554]imaging in X-linked adrenoleukodystrophy. Ann. Neurol.
36: 595–608,

[131555]1994.

[131556]23409.

[131557]Lachtermacher, M. B. R.; Seuanez, H. N.; Moser, A. B.;
Moser,

[131558]H. W.; Smith, K. D.: Determination of 30 X-linked

adrenoleukodystrophy

[131559] mutations, including 15 not previously described. Hum.

Mutat. 15:

[131560] 348–353, 2000.

[131561] 23410.

[131562] Laureti, S.; Casucci, G.; Santeusano, F.; Angeletti, G.;

Aubourg,

[131563] P.; Brunetti, P.: X-linked adrenoleukodystrophy is a fre-

quent cause

[131564] of idiopathic Addison's disease in young adult male pa-

tients. J.

[131565] Clin. Endocr. Metab. 81: 470–474, 1996.

[131566] 23411.

[131567] Lazo, O.; Contreras, M.; Hashmi, M.; Stanley, W.; Irazu, C.;

Singh,

[131568] J.: Peroxisomal lignoceroyl-CoA ligase deficiency in child-

hood adrenoleukodystrophy

[131569] and adrenomyeloneuropathy. Proc. Nat. Acad. Sci. 85:

7647–7651,

[131570] 1988.

[131571] 23412.

[131572]Ligtenberg, M. J. L.; Kemp, S.; Sarde, C.-O.; van Geel, B.
M.;

[131573]Kleijer, W. J.; Barth, P. G.; Mandel, J.-L.; van Oost, B. A.;
Bolhuis,

[131574]P. A.: Spectrum of mutations in the gene encoding the
adrenoleukodystrophy

[131575]protein. Am. J. Hum. Genet. 56: 44-50, 1995.

[131576]23413.

[131577]Luciani, G. B.; Pessotto, R.; Mazzucco, A.:
Adrenoleukodystrophy

[131578]presenting as postperfusion syndrome. (Letter) New Eng.
J. Med. 336:

[131579]731-732, 1997.

[131580]23414.

[131581]Maestri, N. E.; Beaty, T. H.: Predictions of a 2-locus model

[131582]for disease heterogeneity: application to adrenoleukodys-
trophy. Am.

[131583]J. Med. Genet. 44: 576-582, 1992.

[131584]23415.

[131585]Malm, G.; Ringden, O.; Anvret, M.; von Döbeln, U.; Hagen-
feldt,

[131586]L.; Isberg, B.; Knuutila, S.; Nennesmo, I.; Winiarski, J.; Marcus,
cus,

[131587]C.: Treatment of adrenoleukodystrophy with bone marrow
transplantation. Acta

[131588]Paediat. 86: 484-492, 1997.

[131589]23416.

[131590]Manz, H. J.; Schuelein, M.; McCullough, D. C.; Kishimoto,
Y.;

[131591]Eiben, R. M.: New phenotypic variant of adrenoleukodys-
trophy: pathologic,

[131592]ultrastructural, and biochemical study in two brothers. J.
Neurol.

[131593]Sci. 45: 245-260, 1980.

[131594]23417.

[131595]Martin, J. J.; Dompas, B.; Ceuterick, C.; Jacobs, K.: Adreno-
myeloneuropathy

[131596]and adrenoleukodystrophy in two brothers. Europ. Neurol.
19: 281-287,

[131597]1980.

[131598]23418.

[131599]Menkes, J. H.; Corbo, L. M.: Adrenoleukodystrophy: accu-

mutation

[131600]of cholesterol esters with very long chain fatty acids. Neu-
rology 27:

[131601]928-932, 1977.

[131602]23419.

[131603]Migeon, B. R.; Moser, H. W.; Moser, A. B.; Axelman, J.; Sil-
lence,

[131604]D.; Norum, R. A.: Adrenoleukodystrophy: evidence for X
linkage inactivation,

[131605]and selection favoring the mutant allele in heterozygous
cells. Proc.

[131606]Nat. Acad. Sci. 78: 5066-5070, 1981.

[131607]23420.

[131608]Millat, G.; Marcais, C.; Tomasetto, C.; Chikh, K.; Fensom,
A.

[131609]H.; Harzer, K.; Wenger, D. A.; Ohno, K.; Vanier, M. T.: Nie-
mann-Pick

[131610]C1 disease: correlations between NPC1 mutations, levels
of NPC1 protein,

[131611]and phenotypes emphasize the functional significance of
the putative

[131612]sterol-sensing domain and of the cysteine-rich luminal

loop. Am.

[131613]J. Hum. Genet. 68: 1373–1385, 2001.

[131614]23421.

[131615]Morris, J. A.; Zhang, D.; Coleman, K. G.; Nagle, J.;

Pentchev,

[131616]P. G.; Carstea, E. D.: The genomic organization and poly-

morphism

[131617]analysis of the human Niemann–Pick C1 gene. Biochem.

Biophys. Res.

[131618]Commun. 261: 493–498, 1999.

[131619]23422.

[131620]Omura, K.; Suzuki, Y.; Norose, N.; Sato, M.; Maruyama, K.;

Koeda,

[131621]T.: Type C Niemann–Pick disease: clinical and biochemical

studies

[131622]on 6 cases. Brain Dev. 11: 57–61, 1989.

[131623]23423.

[131624]Palmer, M.; Green, W. R.; Maumenee, I. H.; Valle, D. L.;

Singer,

[131625]H. S.; Morton, S. J.; Moser, H. W.: Niemann–Pick disease–

–type C:

[131626]ocular histopathologic and electron microscopic studies.

Arch. Ophthalmol. 103:

[131627]817–822, 1985.

[131628]23424.

[131629]Patel, S. C.; Suresh, S.; Kumar, U.; Hu, C. Y.; Cooney, A.;

Blanchette–Mackie,

[131630]E. J.; Neufeld, E. B.; Patel, R. C.; Brady, R. O.; Patel, Y. C.;

Pentchev,

[131631]P. G.; Ong, W.–Y.: Localization of Niemann–Pick C1 protein

in astrocytes:

[131632]implications for neuronal degeneration in Niemann–Pick

type C disease. Proc.

[131633]Nat. Acad. Sci. 96: 1657–1662, 1999.

[131634]23425.

[131635]Pentchev, P. G.; Boothe, A. D.; Kruth, H. S.; Weintraub, H.;

Stivers,

[131636]J.; Brady, R. O.: A genetic storage disorder in BALB/C mice

with

[131637]a metabolic block in esterification of exogenous chole-

sterol. J.

[131638]Biol. Chem. 259: 5784–5791, 1984.

[131639]23426.

[131640]Sun, X.-H.: Constitutive expression of the Id1 gene impairs mouse

[131641]B cell development. Cell 79: 893–900, 1994.

[131642]23427.

[131643]Nowell, P. C.; Hungerford, D. A.: Chromosome studies on normal

[131644]and leukemic human leukocytes. J. Nat. Cancer Inst. 25: 85–109,

[131645]1960.

[131646]23428.

[131647]Dowbenko, D. J.; Diep, A.; Taylor, B. A.; Lysis, A. J.; Lasky,

[131648]L. A.: Characterization of the murine homing receptor gene reveals

[131649]correspondence between protein domains and coding exons. Genomics 9:

[131650]270–277, 1991.

[131651]23429.

[131652]Aarskog, N. K.; Vedeler, C. A.: Real-time quantitative polymerase

[131653]chain reaction: a new method that detects both the peripheral myelin

[131654]protein 22 duplication in Charcot-Marie-Tooth type 1A disease and

[131655]the peripheral myelin protein 22 deletion in hereditary neuropathy

[131656]with liability to pressure palsies. Hum. Genet. 107: 494-498, 2000.

[131657]23430.

[131658]Chance, P. F.; Alderson, M. K.; Leppig, K. A.; Lensch, M. W.; Matsunami,

[131659]N.; Smith, B.; Swanson, P. D.; Odelberg, S. J.; Distèche, C. M.; Bird,

[131660]T. D.: DNA deletion associated with hereditary neuropathy with liability

[131661]to pressure palsies. Cell 72: 143-151, 1993.

[131662]23431.

[131663]Lehmann, A. R.; Bootsma, D.; Clarkson, S. G.; Cleaver, J. E.;

[131664]McAlpine, P. J.; Tanaka, K.; Thompson, L. H.; Wood, R. D.: Nomenclature

[131665]of human DNA repair genes. Mutat. Res. 315: 41-42,

1994.

[131666]23432.

[131667]Abruzzo, L. V.; Jaffe, E. S.; Cotelingam, J. D.; Whang-Peng, J.;

[131668]Del Duca, V., Jr.; Medeiros, L. J.: T-cell lymphoblastic lymphoma

[131669]with eosinophilia associated with subsequent myeloid malignancy. Am.

[131670]J. Surg. Path. 16: 236-245, 1992.

[131671]23433.

[131672]Chesi, M.; Nardini, E.; Brents, L. A.; Schrock, E.; Ried, T.; Kuehl,

[131673]W. M.; Bergsagel, P. L.: Frequent translocation t(4;14)(p16.3;q32.3)

[131674]in multiple myeloma is associated with increased expression and activating

[131675]mutations of fibroblast growth factor receptor 3. Nature Genet. 16:

[131676]260-264, 1997.

[131677]23434.

[131678]Demiroglu, A.; Steer, E. J.; Heath, C.; Taylor, K.; Bentley,

M.;

[131679] Allen, S. L.; Koduru, P.; Brody, J. P.; Hawson, G.; Rodwell, R.; Doody,

[131680] M.-L.; Carnicero, F.; Reiter, A.; Goldman, J. M.; Melo, J. V.; Cross,

[131681] N. C. P.: The t(8;22) in chronic myeloid leukemia fuses BCR to FGFR1:

[131682] transforming activity and specific inhibition of FGFR1 fusion proteins. Blood 98:

[131683] 3778-3783, 2001.

[131684] 23435.

[131685] Guasch, G.; Mack, G. J.; Popovici, C.; Dastugue, N.; Birnbaum,

[131686] D.; Rattner, J. B.; Pebusque, M.-J.: FGFR1 is fused to the centrosome-associated

[131687] protein CEP110 in the 8p12 stem cell myeloproliferative disorder with

[131688] t(8;9)(p12;q33). Blood 95: 1788-1796, 2000.

[131689] 23436.

[131690] Inhorn, R. C.; Aster, J. C.; Roach, S. A.; Slapak, C. A.; Soifer,

[131691] R.; Tantravahi, R.; Stone, R. M.: A syndrome of lym-

phoblastic lymphoma,
[131692]eosinophilia, and myeloid hyperplasia/malignancy associ-
ated with t(8;13)(p11;q11):
[131693]description of a distinctive clinicopathologic entity. Blood
85:
[131694]1881-1887, 1995.
[131695]23437.
[131696]Lee, P. L.; Johnson, D. E.; Cousens, L. S.; Fried, V. A.;
Williams,
[131697]L. T.: Purification and complementary DNA cloning of a
receptor for
[131698]basic fibroblast growth factor. Science 245: 57-60, 1989.
[131699]23438.
[131700]Mavrothalassitis, G.; Tzimagiorgis, G.; Mitsialis, A.; Zannis,
[131701]V.; Plaitakis, A.; Papamatheakis, J.; Moschonas, N.: Isola-
tion and
[131702]characterization of cDNA clones encoding human liver
glutamate dehydrogenase:
[131703]evidence for a small gene family. Proc. Nat. Acad. Sci. 85:
3494-3498,
[131704]1988.

[131705]23439.

[131706]Michaelidis, T. M.; Tzimagiorgis, G.; Moschonas, N. K.; Papamatheakis,

[131707]J.: The human glutamate dehydrogenase gene family:
gene organization

[131708]and structural characterization. Genomics 16: 150–160,
1993.

[131709]23440.

[131710]Miki, Y.; Taki, T.; Ohura, T.; Kato, H.; Yanagisawa, M.;
Hayashi,

[131711]Y.: Novel missense mutations in the glutamate dehydro-
genase gene

[131712]in the congenital hyperinsulinism–hyperammonemia syn-
drome. J. Pediat. 136:

[131713]69–72, 2000.

[131714]23441.

[131715]Nakatani, Y.; Schneider, M.; Banner, C.; Freese, E.: Com-
plete

[131716]nucleotide sequence of human glutamate dehydrogenase
cDNA. Nucleic

[131717]Acids Res. 16: 6237 only, 1988.

[131718]23442.

[131719]Nelson, R. L.; Povey, M. S.; Hopkinson, D. A.; Harris, H.:
Electrophoresis

[131720]of human L-glutamate dehydrogenase: tissue distribution
and preliminary

[131721]population survey. Biochem. Genet. 15: 87-91, 1977.

[131722]23443.

[131723]Plaitakis, A.; Berl, S.; Yahr, M. D.: Abnormal glutamate
metabolism

[131724]in an adult-onset degenerative neurological disorder. Sci-
ence 216:

[131725]193-196, 1982.

[131726]23444.

[131727]Plaitakis, A.; Berl, S.; Yahr, M. D.: Neurological disorders

[131728]associated with deficiency of glutamate dehydrogenase.
Ann. Neurol. 15:

[131729]144-153, 1984.

[131730]23445.

[131731]Plaitakis, A.; Nicklas, W. J.; Desnick, R. J.: Glutamate dehy-
drogenase

[131732]deficiency in three patients with spinocerebellar syndrome. Ann.

[131733]Neurol. 7: 297–303, 1980.

[131734]23446.

[131735]Santer, R.; Kinner, M.; Passarge, M.; Superti-Furga, A.; Mayatepek,

[131736]E.; Meissner, T.; Schneppenheim, R.; Schaub, J.: Novel missense mutations

[131737]outside the allosteric domain of glutamate dehydrogenase are prevalent

[131738]in European patients with the congenital hyperinsulinism–hyperammonemia

[131739]syndrome. Hum. Genet. 108: 66–71, 2001.

[131740]23447.

[131741]Shaughnessy, J., Jr.; Mock, B.; Duncan, R.; Potter, M.; Banner,

[131742]C.: A restriction fragment length polymorphism at murine Glud locus

[131743]cosegregates with Rib-1, Es-10, and Tcra on chromosome 14. Nucleic

[131744]Acids Res. 17: 2881 only, 1989.

[131745]23448.

[131746]Smith, T. J.; Peterson, P. E.; Schmidt, T.; Fang, J.; Stanley,

[131747]C. A.: Structures of bovine glutamate dehydrogenase
complexes elucidate

[131748]the mechanism of purine regulation. J. Molec. Biol. 307:
707–720,

[131749]2001.

[131750]23449.

[131751]Sorbi, S.; Tonini, S.; Giannini, E.; Piacentini, S.; Marini, P.;

[131752]Amaducci, L.: Abnormal platelet glutamate dehydrogenase
activity

[131753]and activation in dominant and nondominant olivoponto-
cerebellar atrophy. Ann.

[131754]Neurol. 19: 239–245, 1986.

[131755]23450.

[131756]Stanley, C. A.; Lieu, Y.; Hsu, B.; Poncz, M.: Hypoglycemia
in

[131757]infants with hyperinsulinism and hyperammonemia: gain
of function

[131758]mutations in the pathway of leucine-mediated insulin se-
cretion. (Abstract) Diabetes 46

[131759](suppl. 1): 217A only, 1997.

[131760]23451.

[131761]Stanley, C. A.; Lieu, Y. K.; Hsu, B. Y. L.; Burlina, A. B.;
Greenberg,

[131762]C. R.; Hopwood, N. J.; Perlman, K.; Rich, B. H.; Zammarchi,
E.; Poncz,

[131763]M.: Hyperinsulinism and hyperammonemia in infants with
regulatory

[131764]mutations of the glutamate dehydrogenase gene. New
Eng. J. Med. 338:

[131765]1352–1357, 1998.

[131766]23452.

[131767]Tzimagiorgis, G.; Adamson, M. C.; Kozak, C. A.;
Moschonas, N.

[131768]K.: Chromosomal mapping of glutamate dehydrogenase
gene sequences

[131769]to mouse chromosomes 7 and 14. Genomics 10: 83–88,
1991.

[131770]23453.

[131771]Tzimagiorgis, G.; Leversha, M. A.; Chroniary, K.; Gouliel-
mos,

[131772]G.; Sargent, C. A.; Ferguson-Smith, M.; Moschonas, N. K.:

Structure

[131773]and expression analysis of a member of the human glutamate dehydrogenase

[131774](GLUD) gene family mapped to chromosome 10p11.2.

Hum. Genet. 91:

[131775]433-438, 1993.

[131776]23454.

[131777]Yamaguchi, T.; Hayashi, K.; Murakami, H.; Ota, K.;

Maruyama, S.

[131778]: Glutamate dehydrogenase deficiency in spinocerebellar degeneration. Neurochem.

[131779]Res. 7: 627-636, 1982.

[131780]23455.

[131781]Agus, D. B.; Gambhir, S. S.; Pardridge, W. M.; Spielholz, C.;

Baselga,

[131782]J.; Vera, J. C.; Golde, D. W.: Vitamin C crosses the blood-brain

[131783]barrier in the oxidized form through the glucose transporters. J.

[131784]Clin. Invest. 100: 2842-2848, 1997.

[131785]23456.

[131786]Ardinger, R. H., Jr.; Buetow, K. H.; Weiss, M. J.; Nemer, M.;
DeHaven,

[131787]C. R.; Murray, J. C.: Multipoint linkage relationships of 6
loci

[131788]on 1p (ALPL, GLUT, PGD, PGM1, PND, RH). (Abstract) Am.
J. Hum. Genet. 41:

[131789]A154 only, 1987.

[131790]23457.

[131791]Baroni, M. G.; Oelbaum, R. S.; Pozzilli, P.; Stocks, J.; Li,
S.-R.;

[131792]Fiore, V.; Galton, D. J.: Polymorphisms at the GLUT1
(HepG2) and

[131793]GLUT4 (muscle/adipocyte) glucose transporter genes and
non-insulin-dependent

[131794]diabetes mellitus (NIDDM). Hum. Genet. 88: 557-561,
1992.

[131795]23458.

[131796]Brockmann, K.; Wang, D.; Korenke, C. G.; von Moers, A.;
Ho, Y.-Y.;

[131797]Pascual, J. M.; Kuang, K.; Yang, H.; Ma, L.; Kranz-Eble, P.;
Fischbarg,

[131798]J.; Hanefeld, F.; De Vivo, D. C.: Autosomal dominant Glut-1 deficiency

[131799]syndrome and familial epilepsy. Ann. Neurol. 50: 476-485, 2001.

[131800]23459.

[131801]Klepper, J.; Willemsen, M.; Verrips, A.; Guertsen, E.; Hermann,

[131802]R.; Kutzick, C.; Florcken, A.; Voit, T.: Autosomal dominant transmission

[131803]of GLUT1 deficiency. Hum. Molec. Genet. 10: 63-68, 2001.

[131804]23460.

[131805]Lazar, V.; Bidart, J.-M.; Caillou, B.; Mahe, C.; Lacroix, L.; Filetti,

[131806]S.; Schlumberger, M.: Expression of the Na(+)/I(-) symporter gene

[131807]in human thyroid tumors: a comparison study with other thyroid-specific

[131808]genes. J. Clin. Endocr. Metab. 84: 3228-3234, 1999.

[131809]23461.

[131810]Li, S. R.; Baroni, M. G.; Oelbaum, R. S.; Stock, J.; Galton, D.

[131811]J.: Association of genetic variant of the glucose transporter with

[131812]non-insulin-dependent diabetes mellitus. Lancet II: 368-370, 1988.

[131813]23462.

[131814]Mueckler, M.; Caruso, C.; Baldwin, S. A.; Panico, M.; Blench, I.;

[131815]Morris, H. R.; Allard, W. J.; Lienhard, G. E.; Lodish, H. F.: Sequence

[131816]and structure of a human glucose transporter. Science 229: 941-945,

[131817]1985.

[131818]23463.

[131819]Sarkar, H. K.; Thorens, B.; Lodish, H. F.; Kaback, H. R.: Expression

[131820]of the human erythrocyte glucose transporter in Escherichia coli. Proc.

[131821]Nat. Acad. Sci. 85: 5463-5467, 1988.

[131822]23464.

[131823]Seidner, G.; Alvarez, M. G.; Yeh, J.-I.; O'Driscoll, K. R.; Klepper,

[131824]J.; Stump, T. S.; Wang, D.; Spinner, N. B.; Birnbaum, M. J.;

De Vivo,

[131825]D. C.: GLUT-1 deficiency syndrome caused by haploinsufficiency of

[131826]the blood-brain barrier hexose carrier. *Nature Genet.* 18: 188-191,

[131827]1998.

[131828]23465.

[131829]Shepherd, P. R.; Kahn, B. B.: Glucose transporters and insulin

[131830]action: implications for insulin resistance and diabetes mellitus. *New*

[131831]*Eng. J. Med.* 341: 248-257, 1999.

[131832]23466.

[131833]Shows, T. B.; Eddy, R. L.; Byers, M. G.; Fukushima, Y.; Dehaven,

[131834]C. R.; Murray, J. C.; Bell, G. I.: Polymorphic human glucose transporter

[131835]gene (GLUT) is on chromosome 1p31.3-p35. *Diabetes* 36: 546-549, 1987.

[131836]23467.

- [131837]Kelley, M. J.; Pech, M.; Seuanez, H. N.; Rubin, J. S.; O'Brien,
[131838]S. J.; Aaronson, S. A.: Emergence of the keratinocyte
growth factor
[131839]multigene family during the great ape radiation. Proc. Nat.
Acad.
[131840]Sci. 89: 9287–9291, 1992.
[131841]23468.
- [131842]Mattei, M.–G.; deLapeyriere, O.; Bresnick, J.; Dickson, C.;
Birnbaum,
[131843]D.; Mason, I.: Mouse Fgf7 (fibroblast growth factor 7) and
Fgf8 (fibroblast
[131844]growth factor 8) genes map to chromosomes 2 and 19 re-
spectively. Mammalian
[131845]Genome 6: 196–197, 1995.
[131846]23469.
- [131847]Rubin, J. S.; Osada, H.; Finch, P. W.; Taylor, W. G.;
Rudikoff,
[131848]S.; Aaronson, S. A.: Purification and characterization of a
newly
[131849]identified growth factor specific for epithelial cells. Proc.
Nat.

[131850]Acad. Sci. 86: 802–806, 1989.

[131851]23470.

[131852]Werner, S.; Smola, H.; Liao, X.; Longaker, M. T.; Krieg, T.;
Hofschneider,

[131853]P. H.; Williams, L. T.: The function of KGF in morphogene-
sis of epithelium

[131854]and reepithelialization of wounds. Science 266: 819–822,
1994.

[131855]23471.

[131856]Zimonjic, D. B.; Kelley, M. J.; Rubin, J. S.; Aaronson, S. A.;

[131857]Popescu, N. C.: Fluorescence in situ hybridization analysis
of keratinocyte

[131858]growth factor gene amplification and dispersion in evolu-
tion of great

[131859]apes and humans. Proc. Nat. Acad. Sci. 94: 11461–11465,
1997.

[131860]23472.

[131861]Hoeck, H. C.; Vestergaard, P.; Jakobsen, P. E.; Falhof, J.;

Laurberg,

[131862]P.: Diagnosis of growth hormone (GH) deficiency in adults
with hypothalamic–pituitary

[131863] disorders: comparison of test results using pyridostigmine plus GH-releasing

[131864] hormone (GHRH), clonidine plus GHRH, and insulin-induced hypoglycemia

[131865] as GH secretagogues. J. Clin. Endocr. Metab. 85: 1467-1472, 2000.

[131866] 23473.

[131867] Igarashi, Y.; Ogawa, M.; Kamijo, T.; Iwatani, N.; Nishi, Y.; Kohno,

[131868] H.; Masumura, T.; Koga, J.: A new mutation causing inherited growth

[131869] hormone deficiency: a compound heterozygote of a 6.7 kb deletion and

[131870] a two base deletion in the third exon of the GH-1 gene. Hum. Molec.

[131871] Genet. 2: 1073-1074, 1993.

[131872] 23474.

[131873] Johannsson, G.; Marin, P.; Lonn, L.; Ottosson, M.; Stenlof, K.;

[131874] Bjorntorp, P.; Sjostrom, L.; Bengtsson, B.-A.: Growth hormone treatment

[131875] of abdominally obese men reduces abdominal fat mass,

improves glucose

[131876]and lipoprotein metabolism, and reduces diastolic blood pressure. J.

[131877]Clin. Endocr. Metab. 82: 727–734, 1997.

[131878]23475.

[131879]Kamijo, T.; Hayashi, Y.; Shimatsu, A.; Kinoshita, E.; Yoshimoto,

[131880]M.; Ogawa, M.; Seo, H.: Mutations in intron 3 of GH-1 gene associated

[131881]with isolated GH deficiency type II in three Japanese families. Clin.

[131882]Endocr. 51: 355–360, 1999.

[131883]23476.

[131884]Kowarski, A. A.; Schneider J.; Ben-Galim, E.; Weldon, V. V.; Daughaday,

[131885]W. H.: Growth failure with normal serum RIA-GH and low somatomedin

[131886]activity: somatomedin restoration and growth acceleration after exogenous

[131887]GH. J. Clin. Endocr. Metab. 47: 461–464, 1978.

[131888]23477.

[131889]Lebo, R. V.: Personal Communication. San Francisco, Calif.

[131890]7/31/1980.

[131891]23478.

[131892]Leiberman, E.; Pesler, D.; Parvari, R.; Elbedour, K.; Abdul-Latif,

[131893]H.; Brown, M. R.; Parks, J. S.; Carmi, R.: Short stature in carriers

[131894]of recessive mutation causing familial isolated growth hormone deficiency. Am.

[131895]J. Med. Genet. 90: 188-192, 2000.

[131896]23479.

[131897]Maghnie, M.; Strigazzi, C.; Tinelli, C.; Autelli, M.; Cisternino,

[131898]M.; Loche, S.; Severi, F.: Growth hormone (GH) deficiency (GHD) of

[131899]childhood onset: reassessment of GH status and evaluation of the predictive

[131900]criteria for permanent GHD in young adults. J. Clin. Endocr. Metab. 84:

[131901]1324-1328, 1999.

[131902]23480.

[131903]Mahajan, T.; Lightman, S. L.: A simple test for growth hormone

[131904]deficiency in adults. J. Clin. Endocr. Metab. 85:
1473–1476, 2000.

[131905]23481.

[131906]Martial, J. A.; Hallewell, R. A.; Baxter, J. D.; Goodman, H.
M.

[131907]: Human growth hormone: complementary DNA cloning
and expression in

[131908]bacteria. Science 205: 602–607, 1979.

[131909]23482.

[131910]Martinez, A. S.; Domene, H. M.; Ropelato, M. G.; Jasper, H.
G.;

[131911]Pennisi, P. A.; Escobar, M. E.; Heinrich, J. J.: Estrogen
priming

[131912]effect on growth hormone (GH) provocative test: a useful
tool for

[131913]the diagnosis of GH deficiency. J. Clin. Endocr. Metab. 85:
4168–4172,

[131914]2000.

[131915]23483.

[131916]Masuda, N.; Watahiki, M.; Tanaka, M.; Yamakawa, M.; Shimizu, K.;

[131917]Nagai, J.; Nakashima, K.: Molecular cloning of cDNA encoding 20 kDa

[131918]variant human growth hormone and the alternative splicing mechanism. Biochim.

[131919]Biophys. Acta 949: 125–131, 1988.

[131920]23484.

[131921]McCarthy, E. M. S.; Phillips, J. A., III: Characterization of

[131922]an intron splice enhancer that regulates alternative splicing of human

[131923]GH pre-mRNA. Hum. Molec. Genet. 7: 1491–1496, 1998.

[131924]23485.

[131925]Mendlewicz, J.; Linkowski, P.; Kerkhofs, M.; Leproult, R.; Copinschi,

[131926]G.; Van Cauter, E.: Genetic control of 24-hour growth hormone secretion

[131927]in man: a twin study. J. Clin. Endocr. Metab. 84: 856–862, 1999.

[131928]23486.

[131929]Mericq, V.; Cassorla, F.; Salazar, T.; Avila, A.; Iniguez, G.;

- [131930] Bowers, C. Y.; Merriam, G. R.: Effects of eight months treatment
- [131931] with graded doses of a growth hormone (GH)-releasing peptide in GH-deficient
- [131932] children. J. Clin. Endocr. Metab. 83: 2355–2360, 1998.
- [131933] 23487.
- [131934] Morgan, J. R.; Barrandon, Y.; Green, H.; Mulligan, R. C.: Expression
- [131935] of an exogenous growth hormone gene by transplantable human epidermal
- [131936] cells. Science 237: 1476–1479, 1987.
- [131937] 23488.
- [131938] Moseley, C. T.; Mullis, P. E.; Prince, M. A.; Phillips, J. A.,
- [131939] III: An exon splice enhancer mutation causes autosomal dominant GH
- [131940] deficiency. J. Clin. Endocr. Metab. 87: 847–852, 2002.
- [131941] 23489.
- [131942] Mullis, P. E.; Akinci, A.; Kanaka, C.; Eble, A.; Brook, C. G.
- [131943] D.: Prevalence of human growth hormone-1 gene deletions among patients

[131944]with isolated growth hormone deficiency from different populations. *Pediat.*

[131945]*Res.* 31: 532–534, 1992.

[131946]23490.

[131947]Niall, H. D.; Hogan, M. L.; Sauer, R.; Rosenblum, I. Y.; Greenwood,

[131948]F. C.: Sequence of pituitary and placental lactogenic and growth

[131949]hormones: evolution from a primordial peptide by gene reduplication. *Proc.*

[131950]*Nat. Acad. Sci.* 68: 866–869, 1971.

[131951]23491.

[131952]Nishi, Y.; Tanaka, T.; Takano, K.; Fujieda, K.; Igarashi, Y.;

[131953]Hanew, K.; Hirano, T.; Yokoya, S; Tachibana, K.; Saito, T.; Watanabe,

[131954]S.: Recent status in the occurrence of leukemia in growth hormone-treated

[131955]patients in Japan. *J. Clin. Endocr. Metab.* 84: 1961–1965, 1999.

[131956]23492.

[131957]Paladini, A. C.; Pena, C.; Retegui, L. A.: The intriguing na-

ture

[131958]of the multiple actions of growth hormone. Trends
Biochem. Sci. 4:

[131959]256–260, 1979.

[131960]23493.

[131961]Pena-Almazan, S.; Buchlis, J.; Miller, S.; Shine, B.;
Macgillivray,

[131962]M.: Linear growth characteristics of congenitally GH-
deficient infants

[131963]from birth to one year of age. J. Clin. Endocr. Metab. 86:
5691–5694,

[131964]2001.

[131965]23494.

[131966]Pfeifer, M.; Verhovec, R.; Zizek, B.; Prezelj, J.; Poredos, P.;

[131967]Clayton, R. N.: Growth hormone (GH) treatment reverses
early atherosclerotic

[131968]changes in GH-deficient adults. J. Clin. Endocr. Metab. 84:
453–457,

[131969]1999.

[131970]23495.

[131971]Phillips, J. A., III: Inherited defects in growth hormone syn–

thesis

[131972]and action.In: Scriver, C. R.; Beaudet, A. L.; Sly, W. S.;
Valle,

[131973]D. (eds.): The Metabolic and Molecular Bases of Inherited
Disease.

[131974]Vol. II. New York: McGraw-Hill (7th ed.): 1995. Pp.
3023-3044.

[131975]23496.

[131976]Phillips, J. A., III: Personal Communication. Baltimore, Md.

[131977]1/17/1983.

[131978]23497.

[131979]Phillips, J. A., III; Cogan, J. D.: Genetic basis of endocrine

[131980]disease 6: molecular basis of familial human growth hor-
mone deficiency. J.

[131981]Clin. Endocr. Metab. 78: 11-16, 1994.

[131982]23498.

[131983]Phillips, J. A., III; Hjelle, B. L.; Seeburg, P. H.; Plotnick,

[131984]L. P.; Migeon, C. J.; Zachmann, M.: Heterogeneity in the
molecular

[131985]basis of familial growth hormone deficiency (IGHD).

(Abstract) Am.

[131986]J. Hum. Genet. 33: 52A, 1981.

[131987]23499.

[131988]Rappaport, R.; Mugnier, E.; Limoni, C.; Crosnier, H.; Czernichow,

[131989]P.; Leger, J.; Limal, J.-M.; Rochiccioli, P.; Soskin, S.;
Brauner,

[131990]R.; Pinto, G.; Pawels, C.; and 27 others: A 5-year prospective
study of growth hormone (GH)-deficient children treated

[131991]with GH before
the age of 3 years. J. Clin. Endocr. Metab. 82: 452-456,

[131992]1997.

[131993]23500.

[131994]Buckingham, M.; Alonso, S.; Barton, P.; Cohen, A.; Daubas,
P.;

[131995]Garner, I.; Robert, B.; Weydert, A.: Actin and myosin multi-
gene families:

[131996]their expression during the formation and maturation of
striated muscle. Am.

[131997]J. Med. Genet. 25: 623-634, 1986.

[131998]23501.

- [131999]Edwards, Y. H.; Parkar, M.; Povey, S.; West, L. F.; Parrington,
ton,
- [132000]J. M.; Solomon, E.: Human myosin heavy chain genes assigned to chromosome
signed to chromosome
- [132001]17 using a human cDNA clone as probe. Ann. Hum. Genet.
49: 101-109,
- [132002]1985.
- [132003]23502.
- [132004]Epp, T. A.; Dixon, I. M. C.; Wang, H.-Y.; Sole, M. J.; Liew,
C.-C.
- [132005]: Structural organization of the human cardiac alpha-
myosin heavy
- [132006]chain gene (MYH6). Genomics 18: 505-509, 1993.
- [132007]23503.
- [132008]Geisterfer-Lowrance, A. A. T.; Kass, S.; Tanigawa, G.; Vos-
berg,
- [132009]H.-P.; McKenna, W.; Seidman, C. E.; Seidman, J. G.: A
molecular basis
- [132010]for familial hypertrophic cardiomyopathy: a beta cardiac
myosin heavy
- [132011]chain gene missense mutation. Cell 62: 999-1006, 1990.

[132012]23504.

[132013]Jaenicke, T.; Diederich, K. W.; Haas, W.; Schleich, J.;
Lichter,

[132014]P.; Pfordt, M.; Bach, A.; Vosberg, H.-P.: The complete se-
quence of

[132015]the human beta-myosin heavy chain gene and a compara-
tive analysis

[132016]of its product. Genomics 8: 194-206, 1990.

[132017]23505.

[132018]Kurabayashi, M.; Tsuchimochi, H.; Komuro, I.; Takaku, F.;
Yazaki,

[132019]Y.: Molecular cloning and characterization of human car-
diac alpha-

[132020]and beta-form myosin heavy chain complementary DNA
clones: regulation

[132021]of expression during development and pressure overload
in human atrium. J.

[132022]Clin. Invest. 82: 524-531, 1988.

[132023]23506.

[132024]Lichter, P.; Umeda, P. K.; Levin, J. E.; Vosberg, H.-P.: Par-
tial

[132025] characterization of the human beta-myosin heavy-chain
gene which is
[132026] expressed in heart and skeletal muscle. *Europ. J. Biochem.*
160:
[132027] 419-426, 1986.
[132028] 23507.
[132029] Mahdavi, V.; Chambers, A. P.; Nadal-Ginard, B.: Cardiac
alpha-
[132030] and beta-myosin heavy chain genes are organized in tan-
dem. *Proc.*
[132031] *Nat. Acad. Sci.* 81: 2626-2630, 1984.
[132032] 23508.
[132033] Mahdavi, V.; Periasamy, M.; Nadal-Ginard, B.: Molecular
characterization
[132034] of two myosin heavy chain genes expressed in the adult
heart. *Nature* 297:
[132035] 659-664, 1982.
[132036] 23509.
[132037] Matsuoka, R.; Chambers, A.; Kimura, M.; Kanda, N.; Bruns,
G.;
[132038] Yoshida, M.; Takao, A.: Molecular cloning and chromoso-

mal localization

[132039] of a gene coding for human cardiac myosin heavy-chain.

Am. J. Med.

[132040] Genet. 29: 369–376, 1988.

[132041] 23510.

[132042] Matsuoka, R.; Yoshida, M. C.; Kanda, N.; Kimura, M.;

Ozasa, H.;

[132043] Takao, A.: Human cardiac myosin heavy chain gene

mapped within chromosome

[132044] region 14q11.2–q13. Am. J. Med. Genet. 32: 279–284,

1989.

[132045] 23511.

[132046] Nadal-Ginard, B.; Mahdavi, V.: Molecular basis of cardiac

performance:

[132047] plasticity of the myocardium generated through protein

isoform switches. J.

[132048] Clin. Invest. 84: 1693–1700, 1989.

[132049] 23512.

[132050] Niimura, H.; Patton, K. K.; McKenna, W. J.; Soultis, J.;

Maron,

[132051] B. J.; Seidman, J. G.; Seidman, C. E.: Sarcomere protein

gene mutations

[132052]in hypertrophic cardiomyopathy of the elderly. Circulation
105:

[132053]446-451, 2002.

[132054]23513.

[132055]Rappold, G. A.; Vosberg, H.-P.: Chromosomal localization
of a

[132056]human myosin heavy-chain gene by in situ hybridization.
Hum. Genet. 65:

[132057]195-197, 1983.

[132058]23514.

[132059]Adams, J. G., III; Morrison, W. T.; Steinberg, M. H.: A dou-
ble

[132060]crossover within a single human gene: Hb Parchman
(NH-2-delta-beta-delta-COOH).

[132061](Abstract) Am. J. Hum. Genet. 33: 34A only, 1981.

[132062]23515.

[132063]Adams, J. G., III; Morrison, W. T.; Steinberg, M. H.:
Hemoglobin

[132064]Parchman: double crossover within a single human gene.
Science 218:

[132065]291-293, 1982.

[132066]23516.

[132067]Ahern, E. J.; Ahern, V. N.; Aarons, G. H.; Jones, R. T.;
Brimhall,

[132068]B.: Hemoglobin Lepore Washington in two Jamaican families: interaction

[132069]with beta chain variants. Blood 40: 246-256, 1972.

[132070]23517.

[132071]Baglioni, C.: The fusion of two peptide chains in
hemoglobin Lepore

[132072]and its interpretation as a genetic deletion. Proc. Nat.
Acad. Sci. 48:

[132073]1880-1886, 1962.

[132074]23518.

[132075]Baglioni, C.; Ventruto, V.: Human abnormal hemoglobins.
II. A

[132076]chemical study of hemoglobin Lepore from a homozygote
individual. Europ.

[132077]J. Biochem. 5: 29-32, 1968.

[132078]23519.

[132079]McLean, R. H.; Hoefnagel, D.: Partial lipodystrophy and familial
milial

[132080]C3 deficiency. Hum. Hered. 30: 149–154, 1980.

[132081]23520.

[132082]Reaven, G. M.: Role of insulin resistance in human disease.
Diabetes 37:

[132083]1595–1607, 1988.

[132084]23521.

[132085]Sparkes, R. S.; Zollman, S.; Klisak, I.; Kirchgessner, T. G.;

[132086]Komaromy, M. C.; Mohandas, T.; Schotz, M. C.; Lusic, A. J.:
Human

[132087]genes involved in lipolysis of plasma lipoproteins: mapping of loci

[132088]for lipoprotein lipase to 8p22 and hepatic lipase to 15q21.
Genomics 1:

[132089]138–144, 1987.

[132090]23522.

[132091]Nurden, A. T.; Didry, D.; Kieffer, N.; McEver, R. P.: Residual

[132092]amounts of glycoproteins IIb and IIIa may be present in
the platelets

[132093] of most patients with Glanzmann's thrombasthenia. Blood
65: 1021-1024,

[132094] 1985.

[132095] 23523.

[132096] Nurden, A. T.; Rosa, J.-P.; Fournier, D.; Legrand, C.; Didry,

[132097] D.; Parquet, A.; Pidard, D.: A variant of Glanzmann's
thrombasthenia

[132098] with abnormal glycoprotein IIb-IIIa complexes in the
platelet membrane. J.

[132099] Clin. Invest. 79: 962-969, 1987.

[132100] 23524.

[132101] Budka, H.; Sluga, E.; Heiss, W.-D.: Spastic paraplegia as-
sociated

[132102] with Addison's disease: adult variant of adreno-
leukodystrophy. J.

[132103] Neurol. 213: 237-250, 1976.

[132104] 23525.

[132105] Cremers, F. P. M.; Pfeiffer, R. A.; van de Pol, T. J. R.;
Hofker,

[132106] M. H.; Kruse, T. A.; Wieringa, B.; Ropers, H. H.: An intersti-
tial

[132107]duplication of the X chromosome in a male allows physical
fine mapping

[132108]of probes from the Xq13-q22 region. Hum. Genet. 77:
23-27, 1987.

[132109]23526.

[132110]Bauer, P.; Knoblich, R.; Bauer, C.; Finckh, U.; Hufen, A.;
Kropp,

[132111]J.; Braun, S.; Kustermann-Kuhn, B.; Schmidt, D.; Harzer,
K.; Rolfs,

[132112]A.: NPC1: complete genomic sequence, mutation analysis,
and characterization

[132113]of haplotypes. Hum. Mutat. 19: 30-38, 2002.

[132114]23527.

[132115]Besley, G. T. N.; Hoogeboom, A. J. M.; Hoogeveen, A.;
Kleijer,

[132116]W. J.; Galjaard, H.: Somatic cell hybridisation studies
showing different

[132117]gene mutations in Niemann-Pick variants. Hum. Genet.
54: 409-412,

[132118]1980.

[132119]23528.

[132120]Blanchette-Mackie, E. J.; Dwyer, N. K.; Amende, L. M.;
Kruth, H.

[132121]S.; Butler, J. D.; Sokol, J.; Comly, M. E.; Vanier, M. T.; Au-
gust,

[132122]J. T.; Brady, R. O.; Pentchev, P. G.: Type-C Niemann-Pick
disease:

[132123]low density lipoprotein uptake is associated with prema-
ture cholesterol

[132124]accumulation in the Golgi complex and excessive choles-
terol storage

[132125]in lysosomes. Proc. Nat. Acad. Sci. 85: 8022-8026, 1988.

[132126]23529.

[132127]Boustany, R.-M. N.; Kaye, E.; Alroy, J.: Ultrastructural find-
ings

[132128]in skin from patients with Niemann-Pick disease, type C.
Pediat.

[132129]Neurol. 6: 177-183, 1990.

[132130]23530.

[132131]Brady, R. O.: Personal Communication. Bethesda, Md.
3/7/1986.

[132132]23531.

- [132133]Brady, R. O.: Sphingomyelin lipidoses: Niemann–Pick disease.In:
- [132134]Stanbury, J. B.; Wyngaarden, J. B.; Fredrickson, D. S.; Goldstein,
- [132135]J. L.; Brown, M. S.: The Metabolic Basis of Inherited Disease.
- [132136]New York: McGraw–Hill (pub.) (5th ed.): 1983. Pp. 731 only.
- [132137]23532.
- [132138]Brown, D. E.; Thrall, M. A.; Walkley, S. U.; Wenger, D. A.; Mitchell,
- [132139]T. W.; Smith, M. O.; Royals, K. L.; March, P. A.; Allison, R. W.:
- [132140]Feline Niemann–Pick disease type C. Am. J. Path. 144: 1412–1415,
- [132141]1994.
- [132142]23533.
- [132143]Carstea, E. D.; Morris, J. A.; Coleman, K. G.; Loftus, S. K.;
- [132144]Zhang, D.; Cummings, C.; Gu, J.; Rosenfeld, M. A.; Pavan, W. J.; Krizman,
- [132145]D. B.; Nagle, J.; Polymeropoulos, M. H.; and 26 others: Niemann–Pick

[132146]C1 disease gene: homology to mediators of cholesterol homeostasis. Science 277:

[132147]228–231, 1997.

[132148]23534.

[132149]Carstea, E. D.; Parker, C. C.; Fandino, L. B.; Vanier, M. T.;

[132150]Overhauser, J.; Weissenbach, J.; Pentchev, P. G.; Brady, R. O.; Polymeropoulos,

[132151]M. H.: Localizing the human Niemann–Pick C gene to 18q11–12. (Abstract) Am.

[132152]J. Hum. Genet. 55 (suppl.): A182, 1994.

[132153]23535.

[132154]Carstea, E. D.; Polymeropoulos, M. H.; Parker, C. C.; Detera–Wadleigh,

[132155]S. D.; O'Neill, R. R.; Patterson, M. C.; Goldin, E.; Xiao, H.; Straub,

[132156]R. E.; Vanier, M. T.; Brady, R. O.; Pentchev, P. G.: Linkage of Niemann–Pick

[132157]disease type C to human chromosome 18. Proc. Nat. Acad. Sci. 90:

[132158]2002–2004, 1993.

[132159]23536.

[132160]Ceuterick, C.; Martin, J. J.: Nerve biopsy findings in Niemann-Pick

[132161]type II (NPC). (Letter) Acta Neuropath. 88: 602-603, 1994.

[132162]23537.

[132163]Christomanou, H.: Niemann-Pick disease, type C: evidence for

[132164]the deficiency of an activating factor stimulating sphingomyelin and

[132165]glucocerebroside degradation. Hoppe Seylers Z. Physiol. Chem. 361:

[132166]1489-1502, 1980.

[132167]23538.

[132168]Davies, J. P.; Chen, F. W.; Ioannou, Y. A.: Transmembrane molecular

[132169]pump activity of Niemann-Pick C1 protein. Science 290: 2295-2298,

[132170]2000.

[132171]23539.

[132172]Erickson, R. P.; Garver, W. S.; Camargo, F.; Hossain, G. S.; Heidenreich,

[132173]R. A.: Pharmacological and genetic modifications of so-

matic cholesterol

[132174]do not substantially alter the course of CNS disease in
Niemann–Pick

[132175]C mice. J. Inherit. Metab. Dis. 23: 54–62, 2000.

[132176]23540.

[132177]Fink, J. K.; Filling–Katz, M. R.; Sokol, J.; Cogan, D. G.;
Pikus,

[132178]A.; Sonies, B.; Soong, B.; Pentchev, P. G.; Comly, M. E.;
Brady, R.

[132179]O.; Barton, N. W.: Clinical spectrum of Niemann–Pick dis-
ease type

[132180]C. Neurology 39: 1040–1049, 1989.

[132181]23541.

[132182]Gilbert, E. F.; Callahan, J.; Viseskul, C.; Opitz, J. M.: Nie-
mann–Pick

[132183]disease type C: pathological, histochemical, ultrastructural
and biochemical

[132184]studies. Europ. J. Pediat. 136: 263–274, 1981.

[132185]23542.

[132186]Greer, W. L.; Dobson, M. J.; Girouard, G. S.; Byers, D. M.;
Riddell,

- [132187]D. C.; Neumann, P. E.: Mutations in NPC1 highlight a conserved NPC1-specific
cysteine-rich domain. *Am. J. Hum. Genet.* 65: 1252–1260,
1999.
- [132189]23543.
- [132190]Greer, W. L.; Riddell, D. C.; Byers, D. M.; Welch, J. P.;
Girouard,
[132191]G. S.; Sparrow, S. M.; Gillan, T. L.; Neumann, P. E.: Linkage
of
[132192]Niemann–Pick disease type D to the same region of human chromosome
18 as Niemann–Pick disease type C. *Am. J. Hum. Genet.*
61: 139–142,
[132194]1997.
- [132195]23544.
- [132196]Greer, W. L.; Riddell, D. C.; Gillan, T. L.; Girouard, G. S.;
[132197]Sparrow, S. M.; Byers, D. M.; Dobson, M. J.; Neumann, P.
E.: The
[132198]Nova Scotia (type D) form of Niemann–Pick disease is
caused by a 3097G–T
[132199]transversion in NPC1. *Am. J. Hum. Genet.* 63: 52–54,
1998.

[132200]23545.

[132201]Imrie, J.; Wraith, J. E.: Niemann–Pick disease type C. Arch.

[132202]Dis. Child. 84: 427–429, 2001.

[132203]23546.

[132204]Ioannou, Y. A.: The structure and function of the Niemann–Pick

[132205]C1 protein. Molec. Genet. Metab. 71: 175–181, 2000.

[132206]23547.

[132207]Kruth, H. S.; Comly, M. E.; Butler, J. D.; Vanier, M. T.; Fink,

[132208]J. K.; Wenger, D. A.; Patel, S.; Pentchev, P. G.: Type C Niemann–Pick

[132209]disease: abnormal metabolism of low density lipoprotein in homozygous

[132210]and heterozygous fibroblasts. J. Biol. Chem. 261: 16769–16774, 1986.

[132211]23548.

[132212]Liscum, L.; Faust, J. R.: Low density lipoprotein (LDL)–mediated

[132213]suppression of cholesterol synthesis and LDL uptake is defective in

[132214]Niemann–Pick type C fibroblasts. J. Biol. Chem. 262:
17002–17008,

[132215]1987.

[132216]23549.

[132217]Liu, Y.; Wu, Y.–P.; Wada, R.; Neufeld, E. B.; Mullin, K. A.;
Howard,

[132218]A. C.; Pentchev, P. G.; Vanier, M. T.; Suzuki, K.; Proia, R. L.:

[132219]Alleviation of neuronal ganglioside storage does not im-
prove the clinical

[132220]course of the Niemann–Pick C disease mouse. Hum.
Molec. Genet. 9:

[132221]1087–1092, 2000.

[132222]23550.

[132223]Loftus, S. K.; Morris, J. A.; Carstea, E. D.; Gu, J. Z.; Cum-
mings,

[132224]C.; Brown, A.; Ellison, J.; Ohno, K.; Rosenfeld, M. A.; Tagle,
D.

[132225]A.; Pentchev, P. G.; Pavan, W. J.: Murine model of Nie-
mann–Pick C

[132226]disease: mutation in a cholesterol homeostasis gene. Sci-
ence 277:

[132227]232–235, 1997.

[132228]23551.

[132229]Lowenthal, A. C.; Cummings, J. F.; Wenger, D. A.; Thrall, M. A.;

[132230]Wood, P. A.; de Lahunta, A.: Feline sphingolipidosis re-
sembling Niemann–Pick

[132231]disease type C. Acta Neuropath. 81: 189–197, 1990.

[132232]23552.

[132233]Meiner, V.; Shpitzen, S.; Mandel, H.; Klar, A.; Ben–Neria, Z.;

[132234]Zlotogora, J.; Sagi, M.; Lossos, A.; Bargal, R.; Sury, V.; Carmi,

[132235]R.; Leitersdorf, E.; Zeigler, M.: Clinical–biochemical corre-
lation

[132236]in molecularly characterized patients with Niemann–Pick
type C. Genet.

[132237]Med. 3: 343–348, 2001.

[132238]23553.

[132239]Cappa, M.; Bertini, E.; del Balzo, P.; Cambiaso, P.; Di Biase,

[132240]A.; Salvati, S.: High dose immunoglobulin IV treatment in
adrenoleukodystrophy. J.

[132241]Neurol. Neurosurg. Psychiat. 57 (suppl.): 69–70, 1994.

[132242]23554.

[132243]Cartier, N.; Sarde, C.–O.; Douar, A.–M.; Mosser, J.; Mandel, J.–L.;

[132244]Aubourg, P.: Abnormal messenger RNA expression and a missense mutation

[132245]in patients with X-linked adrenoleukodystrophy. Hum. Molec. Genet. 2:

[132246]1949–1951, 1993.

[132247]23555.

[132248]Cotrufo, R.; Melone, M. A. B.; Monsurro, M. R.; Di Iorio, G.;

[132249]Carella, C.; Moser, H. W.: Phenotype heterogeneity among hemizygotes

[132250]in a family biochemically screened for adrenoleukodystrophy. Am.

[132251]J. Med. Genet. 26: 833–838, 1987.

[132252]23556.

[132253]Davis, L. E.; Snyder, R. D.; Orth, D. N.; Nicholson, W. E.; Kornfeld,

[132254]M.; Seelinger, D. F.: Adrenoleukodystrophy and adrenomyeloneuropathy

[132255]associated with partial adrenal insufficiency in three generations

[132256]of a kindred. Am. J. Med. 66: 342–347, 1979.

[132257]23557.

[132258]Di Chiro, G.; Eiben, R. M.; Manz, H. J.; Jacobs, I. B.; Schellinger,

[132259]D.: A new CT pattern in adrenoleukodystrophy. Radiology 137: 687–692,

[132260]1980.

[132261]23558.

[132262]Di Rocco, M.; Doria-Lamba, L.; Caruso, U.: Monozygotic twins

[132263]with X-linked adrenoleukodystrophy and different phenotypes. Ann.

[132264]Neurol. 50: 424 only, 2001.

[132265]23559.

[132266]Dodd, A.; Rowland, S. A.; Hawkes, S. L. J.; Kennedy, M. A.; Love,

[132267]D. R.: Mutations in the adrenoleukodystrophy gene. Hum. Mutat. 9:

[132268]500–511, 1997.

[132269]23560.

[132270]Dvorakova, L.; Storkanova, G.; Unterrainer, G.; Hujova, J.;
Kmoch,

[132271]S.; Zeman, J.; Hrebicek, M.; Berger, J.: Eight novel ABCD1
gene mutations

[132272]and three polymorphisms in patients with X-linked
adrenoleukodystrophy:

[132273]the first polymorphism causing an amino acid exchange.
Hum. Mutat. 18:

[132274]52–60, 2001.

[132275]23561.

[132276]Eichler, E. E.; Budarf, M. L.; Rocchi, M.; Deaven, L. L.;
Doggett,

[132277]N. A.; Baldini, A.; Nelson, D. L.; Mohrenweiser, H. W.: In-
terchromosomal

[132278]duplications of the adrenoleukodystrophy locus: a phe-
nomenon of pericentromeric

[132279]plasticity. Hum. Molec. Genet. 6: 991–1002, 1997.

[132280]23562.

[132281]El-Deiry, S. S.; Naidu, S.; Blevins, L. S.; Ladenson, P. W.:

[132282]Assessment of adrenal function in women heterozygous

for adrenoleukodystrophy. J.

[132283]Clin. Endocr. Metab. 82: 856–860, 1997.

[132284]23563.

[132285]Fanconi, A.; Prader, A.; Isler, W.; Luthy, F.; Siebenmann, R.

[132286]E.: Morbus Addison mit Hirnsklerose im Kindesalter. Ein
hereditaeres

[132287]Syndrom mit X-chromosomaler Vererbung? Helv. Paediat.
Acta 18: 480–501,

[132288]1963.

[132289]23564.

[132290]Fanen, P.; Guidoux, S.; Sarde, C.-O.; Mandel, J.-L.;
Goossens,

[132291]M.; Aubourg, P.: Identification of mutations in the putative
ATP-binding

[132292]domain of the adrenoleukodystrophy gene. J. Clin. Invest.
94: 516–520,

[132293]1994.

[132294]23565.

[132295]Federico, A.; Dotti, M. T.; Annunziata, P.; Bonuccelli, U.;
Fenzi,

[132296]G.; Ciacci, G.; Malandrini, A.; Meucci, G.; Guazzi, G. C.:

Adrenomyeloneurodystrophy

[132297] with late cerebral involvement and evidence of a multiple autoimmune

[132298] disorder. J. Inherit. Metab. Dis. 11 (suppl. 2): 169–172, 1988.

[132299] 23566.

[132300] Feil, R.; Aubourg, P.; Mosser, J.; Douar, A.–M.; Le Paslier, D.;

[132301] Philippe, C.; Mandel, J.–L.: Adrenoleukodystrophy: a complex chromosomal

[132302] rearrangement in the Xq28 red/green–color–pigment gene region indicates

[132303] two possible gene locations. Am. J. Hum. Genet. 49: 1361–1371, 1991.

[132304] 23567.

[132305] Forss–Petter, S.; Werner, H.; Berger, J.; Lassmann, H.; Molzer,

[132306] B.; Schwab, M. H.; Bernheimer, H.; Zimmermann, F.; Nave, K.–A.: Targeted

[132307] inactivation of the X–linked adrenoleukodystrophy gene in mice. J.

[132308] Neurosci. Res. 50: 829–843, 1997.

[132309]23568.

[132310]Fuchs, S.; Sarde, C. O.; Wedemann, H.; Schwinger, E.;
Mandel,

[132311]J. L.; Gal, A.: Missense mutations are frequent in the gene
for X-chromosomal

[132312]adrenoleukodystrophy (ALD). Hum. Molec. Genet. 3:
1903-1905, 1994.

[132313]23569.

[132314]Griffin, J. W.; Goren, E.; Schaumburg, H. H.; Engel, W. K.;
Loriaux,

[132315]L.: Adrenomyeloneuropathy: a probable variant of
adrenoleukodystrophy.

[132316]J. Clinical and endocrinologic aspects. Neurology 27:
1107-1113,

[132317]1977.

[132318]23570.

[132319]Guimaraes, C. P.; Lemos, M.; Menezes, I.; Coelho, T.; Sa-
Miranda,

[132320]C.; Azevedo, J. E.: Characterisation of two mutations in the
ABCD1

[132321]gene leading to low levels of normal ALDP. Hum. Genet.
109: 616-622,

[132322]2001.

[132323]23571.

[132324]Gumbinas, M.; Liu, H. M.; Dawson, G.; Larsen, M.; Green,
O.:

[132325]Progressive spastic paraparesis and adrenal insufficiency.
Arch.

[132326]Neurol. 33: 678-680, 1976.

[132327]23572.

[132328]Haberfeld, W.; Spieler, F.: Zur diffusen Hirn-
Ruckenmarksklerose

[132329]im Kindesalter. Dt. Z. Nervheilk 40: 436-463, 1910.

[132330]23573.

[132331]Harris-Jones, J. N.; Nixon, P. G. F.: Familial Addison's dis-
ease

[132332]with spastic paraplegia. J. Clin. Endocr. 15: 739-744,
1955.

[132333]23574.

[132334]Hashmi, M.; Stanley, W.; Singh, I.: Lignoceroyl-CoASH lig-
ase:

[132335]enzyme defect in fatty acid beta-oxidation system in X-
linked childhood

[132336]adrenoleukodystrophy. FEBS Lett. 196: 247–250, 1986.

[132337]23575.

[132338]Hecaen, H.; De Ajuriaguerra, J.: Balint's syndrome (psychic paralysis

[132339]of visual fixation) and its minor forms. Brain 77: 373–400, 1954.

[132340]23576.

[132341]Heffungs, W.; Hameister, H.; Ropers, H. H.: Addison disease and

[132342]cerebral sclerosis in an apparently heterozygous girl: evidence for

[132343]inactivation of the adrenoleukodystrophy locus. Clin. Genet. 18:

[132344]184–188, 1980.

[132345]23577.

[132346]Ho, J. K.; Moser, H.; Kishimoto, Y.; Hamilton, J. A.: Interactions

[132347]of a very long chain fatty acid with model membranes and serum albumin:

[132348]implications for the pathogenesis of adrenoleukodystrophy. J. Clin.

[132349]Invest. 96: 1455–1463, 1995.

[132350]23578.

[132351]Hoefnagel, D.; Brun, A.; Ingbar, S. H.; Goldman, H.: Addison's

[132352]disease and diffuse cerebral sclerosis. J. Neurol. Neurosurg. Psychiat. 30:

[132353]56–60, 1967.

[132354]23579.

[132355]Hoefnagel, D.; Van den Noort, S.; Ingbar, S. H.: Diffuse cerebral

[132356]sclerosis with endocrine abnormalities in young males. Brain 85:

[132357]553–568, 1962.

[132358]23580.

[132359]Holmberg, B. H.; Hagg, E.; Hagenfeldt, L.: Adrenomyeloneuropathy--report

[132360]on a family. J. Intern. Med. 230: 535–538, 1991.

[132361]23581.

[132362]Igarashi, M.; Schaumburg, H. H.; Powers, J.; Kishimoto, Y.; Kolodny,

[132363]E. H.; Suzuki, K.: Fatty acid abnormality in
adrenoleukodystrophy. J.

[132364]Neurochem. 26: 851–860, 1976.

[132365]23582.

[132366]Kano, S.; Watanabe, M.; Kanai, M.; Koike, R.; Onodera, O.;
Tsuji,

[132367]S.; Okamoto, K.; Shoji, M.: A Japanese family with
adrenoleukodystrophy

[132368]with a codon 291 deletion: a clinical, biochemical, patho-
logical,

[132369]and genetic report. J. Neurol. Sci. 158: 187–192, 1998.

[132370]23583.

[132371]Kemp, S.; Ligtenberg, M. J. L.; van Geel, B. M.; Barth, P. G.;

[132372]Sarde, C.-O.; van Oost, B. A.; Bolhuis, P. A.: Two intronic
mutations

[132373]in the adrenoleukodystrophy gene. Hum. Mutat. 6:
272–273, 1995.

[132374]23584.

[132375]Kemp, S.; Pujol, A.; Waterham, H. R.; van Geel, B. M.;
Boehm,

[132376]C. D.; Raymond, G. V.; Cutting, G. R.; Wanders, R. J. A.;

Moser, H.

[132377]W.: ABCD1 mutations and the X-linked adrenoleukodys-
trophy mutation

[132378]database: role in diagnosis and clinical correlations. Hum.
Mutat. 18:

[132379]499–515, 2001.

[132380]23585.

[132381]Ito, M.; Yuan, C.-X.; Malik, S.; Gu, W.; Fondell, J. D.; Yama-
mura,

[132382]S.; Fu, Z.-Y.; Zhang, X.; Qin, J.; Roeder, R. G.: Identity be-
tween

[132383]TRAP and SMCC complexes indicates novel pathways for
the function

[132384]of nuclear receptors and diverse mammalian activators.
Molec. Cell 3:

[132385]361–370, 1999.

[132386]23586.

[132387]Nagase, T.; Seki, N.; Ishikawa, K.; Tanaka, A.; Nomura, N.:
Prediction

[132388]of the coding sequences of unidentified human genes. V.
The coding

[132389]sequences of 40 new genes (KIAA0161–KIAA0200) de-

duced by analysis

[132390]of cDNA clones from human cell line KG-1. DNA Res. 3:
17-24, 1996.

[132391]23587.

[132392]Maeda, K.; Matsushashi, S.; Hori, K.; Xin, Z.; Mukai, T.;
Tabuchi,

[132393]K.; Egashira, M.; Niikawa, N.: Cloning and characterization
of a

[132394]novel human gene, TM4SF6, encoding a protein belonging
to the transmembrane

[132395]4 superfamily, and mapped to Xq22. Genomics 52:
240-242, 1998.

[132396]23588.

[132397]Todd, S. C.; Doctor, V. S.; Levy, S.: Sequences and expres-
sion

[132398]of six new members of the tetraspanin/TM4SF family.
Biochim. Biophys.

[132399]Acta 1399: 101-104, 1998.

[132400]23589.

[132401]Vaccari, T.; Beltrame, M.; Ferrari, S.; Bianchi, M. E.: Hmg4,

[132402]a new member of the Hmg1/2 gene family. Genomics 49:

247–252, 1998.

[132403]23590.

[132404]Wilke, K.; Wiemann, S.; Gaul, R.; Gong, W.; Poustka, A.:

Isolation

[132405]of human and mouse HMG2a cDNAs: evidence for an
HMG2a-specific 3-prime

[132406]untranslated region. *Gene* 198: 269–274, 1997.

[132407]23591.

[132408]Bird, A. P.: HTF islands as gene markers in the vertebrate
nucleus. *Trends*

[132409]*Genet.* 3: 342–347, 1987.

[132410]23592.

[132411]Anson, D. S.; Blake, D. J.; Winship, P. R.; Birnbaum, D.;
Brownlee,

[132412]G. G.: Nullisomic deletion of the mcf.2 transforming gene
in two

[132413]haemophilia B patients. *EMBO J.* 7: 2795–2799, 1988.

[132414]23593.

[132415]Gong, Y.; Slee, R. B.; Fukai, N.; Rawadi, G.; Roman–Roman,
S.;

[132416]Reginato, A. M.; Wang, H.; Cundy, T.; Glorieux, F. H.; Lev,

D.; Zacharin,
 [132417]M.; Oexle, K.; and 50 others: LDL receptor-related protein
 5 (LRP5)
 [132418]affects bone accrual and eye development. Cell 107:
 513–523, 2001.
 [132419]23594.

[132420]Abadie, V.; Jaruzelska, J.; Lyonnet, S.; Millasseau, P.;
 Berthelon,
 [132421]M.; Rey, F.; Munnich, A.; Rey, J.: Illegitimate transcription
 of
 [132422]the phenylalanine hydroxylase gene in lymphocytes for
 identification
 [132423]of mutations in phenylketonuria. Hum. Molec. Genet. 2:
 31–34, 1993.
 [132424]23595.

[132425]Abadie, V.; Lyonnet, S.; Maurin, N.; Berthelon, M.; Caillaud,
 C.;

[132426]Giraud, F.; Mattei, J.-F.; Rey, J.; Rey, F.; Munnich, A.: CpG
 dinucleotides
 [132427]are mutation hot spots in phenylketonuria. Genomics 5:
 936–939,
 [132428]1989.

[132429]23596.

[132430]Abadie, V.; Lyonnet, S.; Maurin, N.; Berthelon, M.; Caillaud, C.;

[132431]Giraud, F.; Mattei, J.-F.; Rey, J.; Rey, F.; Munnich, A.: CpG dinucleotides

[132432]are mutation hot spots in phenylketonuria. (Abstract) Am. J. Hum.

[132433]Genet. 45 (suppl.): A169 only, 1989.

[132434]23597.

[132435]Weiss, P.; Tietze, F.; Gahl, W. A.; Seppala, R.; Ashwell, G.:

[132436]Identification of the metabolic defect in sialuria. J. Biol. Chem. 264:

[132437]17635-17636, 1989.

[132438]23598.

[132439]Kennedy, M. A.; Rowland, S. A.; Miller, A. L.; Morris, C. M.;

[132440]Neville, L. A.; Dodd, A.; Fifield, W. J.; Love, D. R.: Structure

[132441]and location of the murine adrenoleukodystrophy gene. Genomics 32:

[132442]395-400, 1996.

[132443]23599.

- [132444] Bassi, M. T.; Ramesar, R. S.; Caciotti, B.; Winship, I. M.; De
[132445] Grandi, A.; Riboni, M.; Townes, P. L.; Beighton, P.; Ballabio,
A.;
[132446] Borsani, G.: X-linked late-onset sensorineural deafness
caused by
[132447] a deletion involving OA1 and a novel gene containing WD-
40 repeats. Am.
[132448] J. Hum. Genet. 64: 1604-1616, 1999.
[132449] 23600.
[132450] Converse, P. J.: Personal Communication. Baltimore, Md.
8/24/2001.
[132451] 23601.
[132452] Disteche, C. M.; Dinulos, M. B.; Bassi, M. T.; Elliott, R. W.;
[132453] Rugarli, E. I.: Mapping of the murine Tbl1 gene reveals a
new rearrangement
[132454] between mouse and human X chromosomes. Mammalian
Genome 9: 1062-1064,
[132455] 1998.
[132456] 23602.
[132457] Dong, X.; Tsuda, L.; Zavitz, K. H.; Lin, M.; Li, S.; Carthew,
R.

[132458]W.; Zipursky, S. L.: ebi regulates epidermal growth factor receptor

[132459]signaling pathways in Drosophila. Genes Dev. 13: 954–965, 1999.

[132460]23603.

[132461]Matsuzawa, S.; Reed, J. C.: Siah–1, SIP, and Ebi collaborate in

[132462]a novel pathway for beta–catenin degradation linked to p53 responses. Molec.

[132463]Cell 7: 915–926, 2001.

[132464]23604.

[132465]Zhang, J.; Kalkum, M.; Chait, B. T.; Roeder, R. G.: The N–CoR–HDAC3

[132466]nuclear receptor corepressor complex inhibits the JNK pathway through

[132467]the integral subunit GPS2. Molec. Cell 9: 611–623, 2002.

[132468]23605.

[132469]Mu, J.; Skurat, A. V.; Roach, P. J.: Glycogenin–2, a novel self–glucosylating

[132470]protein involved in liver glycogen biosynthesis. J. Biol. Chem. 272:

[132471]27589–27597, 1997.

[132472]23606.

[132473]Bartley, J. A.; Miller, D. K.; Hayford, J. T.; McCabe, E. R. B.

[132474]: Concordance of X-linked glycerol kinase deficiency with
X-linked

[132475]congenital adrenal hypoplasia. *Lancet* II: 733–736, 1982.

[132476]23607.

[132477]Jansen, G. A.; Ofman, R.; Ferdinandusse, S.; IJlst, L.; Mui-
jsers,

[132478]A. O.; Skjeldal, O. H.; Stokke, O.; Jakobs, C.; Besley, G. T.
N.;

[132479]Wraith, J. E.; Wanders, R. J. A.: Refsum disease is caused
by mutations

[132480]in the phytanoyl–CoA hydroxylase gene. *Nature Genet.* 17:
190–193,

[132481]1997.

[132482]23608.

[132483]Mihalik, S. J.; Morrell, J. C.; Kim, D.; Sacksteder, K. A.;
Watkins,

[132484]P. A.; Gould, S. J.: Identification of PAHX, a Refsum disease
gene. *Nature*

[132485]Genet. 17: 185–189, 1997.

[132486]23609.

[132487]Wilcken, B.; Don, N.; Greenaway, R.; Hammond, J.; Sosula, L.:

[132488]Sialuria: a second case. J. Inherit. Metab. Dis. 10: 97–102, 1987.

[132489]23610.

[132490]Bickel, H.; Gerrard, J.; Hickmans, E. M.: Influence of phenylalanine

[132491]intake on phenylketonuria. Lancet 2: 812–813, 1953.

[132492]23611.

[132493]Bowden, J. A.; McArthur, C. L., III: Possible biochemical model

[132494]for phenylketonuria. Nature 235: 230, 1972.

[132495]23612.

[132496]Brenton, D. P.; Lilburn, M.: Maternal phenylketonuria: a study

[132497]from the United Kingdom. Europ. J. Pediat. 155 (suppl. 1): 177–180,

[132498]1996.

[132499]23613.

[132500]Burgard, P.; Rupp, A.; Konecki, D. S.; Trefz, F. K.; Schmidt,
[132501]H.; Lichter-Konecki, U.: Phenylalanine hydroxylase geno-
types, predicted
[132502]residual enzyme activity and phenotypic parameters of di-
agnosis and
[132503]treatment of phenylketonuria. *Europ. J. Pediat.* 155
(suppl.): S11-S15,
[132504]1996.
[132505]23614.

[132506]Byck, S.; Tyfield, L.; Carter, K.; Scriver, C. R.: Prediction
[132507]of multiple hypermutable codons in the human PAH gene:
codon 280 contains
[132508]recurrent mutations in Quebec and other populations.
Hum. Mutat. 9:
[132509]316-321, 1997.
[132510]23615.

[132511]Cabalska, B.: Personal Communication. Warsaw, Poland
1980.
[132512]23616.

[132513]Caillaud, C.; Lyonnet, S.; Rey, F.; Melle, D.; Frebourg, T.;
Berthelon,

[132514]M.; Vilarinho, L.; Vaz Osorio, R.; Rey, J.; Munnich, A.: A
3-base

[132515]pair in-frame deletion of the phenylalanine hydroxylase
gene results

[132516]in a kinetic variant of phenylketonuria. J. Biol. Chem. 266:
9351-9354,

[132517]1991.

[132518]23617.

[132519]Carter, C. O.; Woolf, L. I.: The birthplaces of parents and
grandparents

[132520]of a series of patients with phenylketonuria in southeast
England. Ann.

[132521]Hum. Genet. 25: 57-64, 1961.

[132522]23618.

[132523]Carter, K. C.; Byck, S.; Waters, P. J.; Richards, B.; Nowacki,

[132524]P. M.; Laframboise, R.; Lambert, M.; Treacy, E.; Sriver, C.
R.:

[132525]Mutation at the phenylalanine hydroxylase gene (PAH) and
its use to

[132526]document population genetic variation: the Quebec expe-
rience. Europ.

[132527]J. Hum. Genet. 6: 61–70, 1998.

[132528]23619.

[132529]Centerwall, W. R.; Neff, C. A.: Phenylketonuria: a case report

[132530]of children of Jewish ancestry. Arch. Pediat. 78: 379–384, 1961.

[132531]23620.

[132532]Chao, H.-K.; Hsiao, K.-J.; Su, T.-S.: A silent mutation induces

[132533]exon skipping in the phenylalanine hydroxylase gene in phenylketonuria. Hum.

[132534]Genet. 108: 14–19, 2001.

[132535]23621.

[132536]Chen, K.-J.; Chao, H.-K.; Hsiao, K.-J.; Su, T.-S.: Identification

[132537]and characterization of a novel liver-specific enhancer of the human

[132538]phenylalanine hydroxylase gene. Hum. Genet. 110: 235–243, 2002.

[132539]23622.

[132540]Chen, S.-H.; Hsiao, K.-J.; Lin, L.-H.; Liu, T.-T.; Tang,

R.-B.;

[132541]Su, T.-S.: Study of restriction fragment length polymorphisms at

[132542]the human phenylalanine hydroxylase locus and evaluation of its potential

[132543]application in prenatal diagnosis of phenylketonuria in Chinese. Hum.

[132544]Genet. 81: 226-230, 1989.

[132545]23623.

[132546]Cipic-Schmidt, S.; Trefz, F. K.; Funders, B.; Seidlitz, G.; Ullrich,

[132547]K.: German maternal phenylketonuria study. Europ. J. Pediat. 155

[132548](suppl. 1): 173-176, 1996.

[132549]23624.

[132550]Cohen, B. E.; Bodonyi, E.; Szeinberg, A.: Phenylketonuria in

[132551]Jews. Lancet I: 344-345, 1961.

[132552]23625.

[132553]Corsello, G.; Bosco, P.; Cali, F.; Greco, D.; Cammarata, M.; Ciaccio,

[132554]M.; Piccione, M.; Romano, V.: Maternal phenylketonuria in

two Sicilian

[132555]families identified by maternal blood phenylalanine level
screening

[132556]and identification of a new phenylalanine hydroxylase
gene mutation

[132557](P407L). (Letter) Europ. J. Pediat. 158: 83–88, 1999.

[132558]23626.

[132559]Coskun, T.; Ozalp, I.; Kale, G.; Gogus, S.: Scleroderma-like

[132560]skin lesions in two patients with phenylketonuria. Europ. J.
Pediat. 150:

[132561]109–110, 1990.

[132562]23627.

[132563]Cotton, R. G. H.: Heterogeneity of phenylketonuria at the
clinical,

[132564]protein and DNA levels. J. Inherit. Metab. Dis. 13:
739–750, 1990.

[132565]23628.

[132566]Cunningham, G. C.; Day, R. W.; Berman, J. L.; Hsia, D.
Y.–Y.:

[132567]Phenylalanine tolerance tests in families with phenylke-
tonuria and

- [132568]hyperphenylalaninemia. Am. J. Dis. Child. 117: 626-635, 1969.
- [132569]23629.
- [132570]Daiger, S. P.; Chakraborty, R.; Reed, L.; Fekete, G.; Schuler, D.; Berenssi, G.; Nasz, I.; Brdicka, R.; Kamaryt, J.; Pijackova, A.;
- [132572]Moore, S.; Sullivan, S.; Woo, S. L. C.: Polymorphic DNA haplotypes
- [132573]at the phenylalanine hydroxylase (PAH) locus in European families
- [132574]with phenylketonuria (PKU). Am. J. Hum. Genet. 45: 310-318, 1989.
- [132575]23630.
- [132576]Gaedigk, R.; Duncan, A. M. V.; Miyazaki, I.; Robinson, B. H.; Dosch, H.-M.: ICA1 encoding p69, a protein linked to the development of
- [132578]type 1 diabetes, maps to human chromosome 7p22. Cytogenet. Cell Genet. 66:
- [132579]274-276, 1994.
- [132580]23631.

[132581]Gaedigk, R.; Karges, W.; Hui, M. F.; Scherer, S. W.; Dosch, H.-M.

[132582]Genomic organization and transcript analysis of ICAp69, a target

[132583]antigen in diabetic autoimmunity. Genomics 38: 382–391, 1996.

[132584]23632.

[132585]Pietropaolo, M.; Castano, L.; Babu, S.; Buelow, R.; Kuo, Y.-L.

[132586]S.; Martin, S.; Martin, A.; Powers, A. C.; Prochazka, M.; Naggert,

[132587]J.; Leiter, E. H.; Eisenbarth, G. S.: Islet cell autoantigen 69 kD

[132588](ICA69): molecular cloning and characterization of a novel diabetes-associated

[132589]autoantigen. J. Clin. Invest. 92: 359–371, 1993.

[132590]23633.

[132591]Diaz, M. O.; Le Beau, M. M.; Pitha, P.; Rowley, J. D.: Interferon

[132592]and c-ets-1 genes in the translocation (9;11)(p22;q23) in human acute

[132593]monocytic leukemia. Science 231: 265–267, 1986.

[132594]23634.

[132595]Ohlsson, M.; Feder, J.; Cavalli-Sforza, L. L.; von Gabain, A.

[132596]: Close linkage of alpha and beta interferons and infrequent duplication

[132597]of beta interferon in humans. Proc. Nat. Acad. Sci. 82: 4473-4476,

[132598]1985.

[132599]23635.

[132600]Owerbach, D.; Rutter, W. J.; Shows, T. B.; Gray, P.; Goeddel,

[132601]D. V.; Lawn, R. M.: Leukocyte and fibroblast interferon genes are

[132602]located on human chromosome 9. Proc. Nat. Acad. Sci. 78: 3123-3127,

[132603]1981.

[132604]23636.

[132605]Siegal, F. P.; Kadowaki, N.; Shodell, M.; Fitzgerald-Bocarsly,

[132606]P. A.; Shah, K.; Ho, S.; Antonenko, S.; Liu, Y.-J.: The nature of

[132607]the principal type 1 interferon-producing cells in human

blood. Science 284:

[132608]1835–1837, 1999.

[132609]23637.

[132610]Heath, P.; Elvin, P.; Jenner, D.; Gammack, A.; Morten, J.;
Markham,

[132611]A.: Localisation of a cDNA clone for human cytokeratin 18
to chromosome

[132612]17p11–p12 by in situ hybridisation. Hum. Genet. 85:
669–670, 1990.

[132613]23638.

[132614]Ku, N.–O.; Michie, S.; Oshima, R. G.; Omary, M. B.: Chronic
hepatitis,

[132615]hepatocyte fragility, and increased soluble phosphogly-
cokeratins in

[132616]transgenic mice expressing a keratin 18 conserved argi-
nine mutant. J.

[132617]Cell Biol. 131: 1303–1314, 1995.

[132618]23639.

[132619]Ku, N.–O.; Michie, S. A.; Soetikno, R. M.; Resurreccion, E.
Z.;

[132620]Broome, R. L.; Oshima, R. G.; Omary, M. B.: Susceptibility

to hepatotoxicity

[132621]in transgenic mice that express a dominant-negative human keratin

[132622]18 mutant. J. Clin. Invest. 98: 1034–1046, 1996.

[132623]23640.

[132624]Kulesh, D. A.; Oshima, R. G.: Complete structure of the gene for

[132625]human keratin 18. Genomics 4: 339–347, 1989.

[132626]23641.

[132627]Lendahl, U.; Zimmerman, L. B.; McKay, R. D. G.: CNS stem cells

[132628]express a new class of intermediate filament protein. Cell 60: 585–595,

[132629]1990.

[132630]23642.

[132631]Steinert, P. M.; Roop, D. R.: Molecular and cellular biology of

[132632]intermediate filaments. Ann. Rev. Biochem. 57: 593–625, 1988.

[132633]23643.

[132634]Waseem, A.; Gough, A. C.; Spurr, N. K.; Lane, E. B.: Local–

ization

[132635]of the gene for human simple epithelial keratin 18 to
chromosome 12

[132636]using polymerase chain reaction. Genomics 7: 188–194,
1990.

[132637]23644.

[132638]Erikson, J.; Griffin, C.; ar-Rushdi, A.; Valtieri, M.; Hoxie,

[132639]J.; Finan, J.; Emanuel, B. S.; Rovera, G.; Nowell, P. C.;
Croce, C.

[132640]M.: Heterogeneity of chromosome 22 breakpoint in
Philadelphia-positive

[132641](Ph+) acute lymphocytic leukemia. Proc. Nat. Acad. Sci.
83: 1807–1811,

[132642]1986.

[132643]23645.

[132644]Fioretos, T.; Heisterkamp, N.; Groffen, J.: No evidence for
genomic

[132645]imprinting of the human BCR gene. Blood 83: 3441–3444,
1994.

[132646]23646.

[132647]Fitzgerald, P. H.: Evidence that chromosome band 22q12

is concerned

[132648]with cell proliferation in chronic myeloid leukemia. Hum.

Genet. 33:

[132649]269–274, 1976.

[132650]23647.

[132651]Ganesan, T. S.; Rassool, F.; Guo, A.-P.; Th'ng, K. H.;

Dowding,

[132652]C.; Hibbin, J. A.; Young, B. D.; White, H.; Kumaran, T. O.;

Galton,

[132653]D. A. G.; Goldman, J. M.: Rearrangement of the bcr gene in

Philadelphia

[132654]chromosome-negative chronic myeloid leukemia. Blood

68: 957–960,

[132655]1986.

[132656]23648.

[132657]Goldman, J. M.; Melo, J. V.: Targeting the BCR-ABL tyro-

sine kinase

[132658]in chronic myeloid leukemia. (Editorial) New Eng. J. Med.

344: 1084–1086,

[132659]2001.

[132660]23649.

[132661]Gorre, M. E.; Mohammed, M.; Ellwood, K.; Hsu, N.; Paquette, R.;

[132662]Rao, P. N.; Sawyers, C. L.: Clinical resistance to STI-571 cancer

[132663]therapy caused by BCR-ABL gene mutation or amplification. Science 293:

[132664]876-883, 2001.

[132665]23650.

[132666]Groffen, J.; Stephenson, J. R.; Heisterkamp, N.; de Klein, A.;

[132667]Bartram, C. R.; Grosveld, G.: Philadelphia chromosomal breakpoints

[132668]are clustered within a limited region, bcr, on chromosome 22. Cell 36:

[132669]93-99, 1984.

[132670]23651.

[132671]Grossman, A.; Silver, R. T.; Arlin, Z.; Coleman, M.; Camposano,

[132672]E.; Gascon, P.; Benn, P. A.: Fine mapping of chromosome 22 breakpoints

[132673]within the breakpoint cluster region (bcr) implies a role for bcr

[132674]lexon 3 in determining disease duration in chronic myeloid leukemia. Am.

[132675]J. Hum. Genet. 45: 729–738, 1989.

[132676]23652.

[132677]Grosveld, G.; Verwoerd, T.; van Agthoven, T.; de Klein, A.; Ramachandran,

[132678]K. L.; Heisterkamp, N.; Stam, K.; Groffen, J.: The chronic myelocytic

[132679]cell line K562 contains a breakpoint in bcr and produces a chimeric

[132680]bcr/c-abl transcript. Molec. Cell. Biol. 6: 607–616, 1986.

[132681]23653.

[132682]Haas, O. A.: Are ABL and BCR imprinted? No definitive answers,

[132683]but more questions. Leukemia 9: 740–745, 1995.

[132684]23654.

[132685]Haas, O. A.; Argyriou-Tirita, A.; Lion, T.: Parental origin of

[132686]chromosomes involved in the translocation t(9;22). Nature 359: 414–416,

[132687]1992.

[132688]23655.

[132689]Hariharan, I. K.; Adams, J. M.: cDNA sequence for human
bcr,

[132690]the gene that translocates to the abl oncogene in chronic
myeloid

[132691]leukaemia. EMBO J. 6: 115–119, 1987.

[132692]23656.

[132693]Heisterkamp, N.; Jenster, G.; ten Hoeve, J.; Zovich, D.; Pat-
tengale,

[132694]P. K.; Groffen, J.: Acute leukaemia in bcr/abl transgenic
mice. Nature 344:

[132695]251–253, 1990.

[132696]23657.

[132697]Heisterkamp, N.; Stam, K.; Groffen, J.; de Klein, A.;
Grosveld,

[132698]G.: Structural organization of the bcr gene and its role in
the Ph-1

[132699]translocation. Nature 315: 758–761, 1985.

[132700]23658.

[132701]Hermans, A.; Heisterkamp, N.; von Lindern, M.; van Baal,
S.; Meijer,

- [132702]D.; van der Plas, D.; Wiedemann, L. M.; Groffen, J.;
Bootsma, D.;
- [132703]Grosveld, G.: Unique fusion of bcr and c-abl genes in
Philadelphia
- [132704]chromosome positive acute lymphoblastic leukemia. Cell
51: 33-40,
- [132705]1987.
- [132706]23659.
- [132707]Huettner, C. S.; Zhang, P.; Van Etten, R. A.; Tenen, D. G.:
Reversibility
- [132708]of acute B-cell leukaemia induced by BCR-ABL1. Nature
Genet. 24:
- [132709]57-60, 2000.
- [132710]23660.
- [132711]Jacobs, A.: Benzene and leukemia. Brit. J. Haemat. 72:
119-121,
- [132712]1989.
- [132713]23661.
- [132714]Klein, G.: Specific chromosomal translocations and the
genesis
- [132715]of the B-cell-derived tumors in mice and men. Cell 32:

311-315,

[132716]1983.

[132717]23662.

[132718]Koeffler, H. P.; Golde, D. W.: Chronic myelogenous leukemia--new

[132719]concepts. New Eng. J. Med. 304: 1201-1209 and 1269-1274, 1981.

[132720]23663.

[132721]Kohno, S.-I.; Sandberg, A. A.: Chromosomes and causation of human

[132722]cancer and leukemia. XXXIX. Usual and unusual findings in Ph(1)-positive

[132723]CML. Cancer 46: 2227-2237, 1980.

[132724]23664.

[132725]Laurent, E.; Talpaz, M.; Kantarjian, H.; Kurzrock, R.: The BCR

[132726]gene and Philadelphia chromosome-positive leukemogenesis. Cancer

[132727]Res. 61: 2343-2355, 2001.

[132728]23665.

[132729]Lillicrap, D. A.; Sterndale, H.: Familial chronic myeloid

leukaemia.

[132730](Letter) Lancet II: 699, 1984.

[132731]23666.

[132732]Lim, Y.-M.; Wong, S.; Lau, G.; Witte, O. N.; Colicelli, J.:

BCR/ABL

[132733]inhibition by an escort/phosphatase fusion protein. Proc.

Nat. Acad.

[132734]Sci. 97: 12233-12238, 2000.

[132735]23667.

[132736]Litz, C. E.; Copenhaver, C. M.: Paternal origin of the rear-

ranged

[132737]major breakpoint cluster region in chronic myeloid

leukemia. Blood 83:

[132738]3445-3448, 1994.

[132739]23668.

[132740]Maru, Y.; Witte, O. N.: The BCR gene encodes a novel ser-

ine/threonine

[132741]kinase activity within a single exon. Cell 67: 459-468,

1991.

[132742]23669.

[132743]Melo, J. V.; Yan, X.-H.; Diamond, J.; Goldman, J. M.: Lack

of

[132744]imprinting of the ABL gene. (Letter) Nature Genet. 8:
318–319, 1994.

[132745]23670.

[132746]Melo, J. V.; Yan, X.-H.; Diamond, J.; Goldman, J. M.: Bal-
anced

[132747]parental contribution to the ABL component of the BCR-
ABL gene in

[132748]chronic myeloid leukemia. Leukemia 9: 734–745, 1995.

[132749]23671.

[132750]Mes-Masson, A.-M.; McLaughlin, J.; Daley, G. Q.; Paskind,
M.;

[132751]Witte, O. N.: Overlapping cDNA clones define the complete
coding

[132752]region for the P210(c-abl) gene product associated with
chronic myelogeneous

[132753]leukemia cells containing the Philadelphia chromosome.
Proc. Nat.

[132754]Acad. Sci. 83: 9768–9772, 1986.

[132755]23672.

[132756]Mills, K. I.; MacKenzie, E. D.; Birnie, G. D.: The site of the

[132757]breakpoint within the bcr is a prognostic factor in
Philadelphia-positive

[132758]CML patients. Blood 72: 1237-1241, 1988.

[132759]23673.

[132760]Mittelman, F.; Levan, G.: Clustering of aberrations to specific
cific

[132761]chromosomes in human neoplasms. III. Incidence and geographic
ographic distribution

[132762]of chromosome aberrations in 856 cases. Hereditas 89:
207-232, 1978.

[132763]23674.

[132764]Nowell, P. C.; Hungerford, D. A.: A minute chromosome in
human

[132765]chronic granulocytic leukemia. Science 132: 1497, 1960.

[132766]23675.

[132767]Kusari, J.; Takata, Y.; Hatada, E.; Freidenberg, G.; Kolter-
man,

[132768]O.; Olefsky, J. M.: Insulin resistance and diabetes due to
different

[132769]mutations in the tyrosine kinase domain of both insulin
receptor gene

[132770]alleles. J. Biol. Chem. 266: 5260–5267, 1991.

[132771]23676.

[132772]Lander, E. S.; Botstein, D.: Homozygosity mapping: a way
to map

[132773]human recessive traits with the DNA of inbred children.
Science 236:

[132774]1567–1570, 1987.

[132775]23677.

[132776]Sagane, K.; Ohya, Y.; Hasegawa, Y.; Tanaka, I.: Metallo-
proteinase-like,

[132777]disintegrin-like, cysteine-rich proteins MDC2 and MDC3:
novel human

[132778]cellular disintegrins highly expressed in the brain.
Biochem. J. 334:

[132779]93–98, 1998.

[132780]23678.

[132781]Splendore, A.; Passos-Bueno, M. R.; Jabs, E. W.; Van
Maldergem,

[132782]L.; Wulfsberg, E. A.: TCOF1 mutations excluded from a
role in other

[132783]first and second branchial arch-related disorders. (Letter)

Am. J.

[132784]Med. Genet. 111: 324–327, 2002.

[132785]23679.

[132786]Alsobrook, J. P., II; Zohar, A. H.; Leboyer, M.; Chabane, N.;
Ebstein,

[132787]R. P.; Pauls, D. L.: Association between the COMT locus
and obsessive–compulsive

[132788]disorder in females but not males. Am. J. Med. Genet.
(Neuropsychiat.

[132789]Genet.) 114: 116–120, 2002.

[132790]23680.

[132791]Karayiorgou, M.; Sobin, C.; Blundell, M. L.; Galke, B. L.;
Malinova,

[132792]L.; Goldberg, P.; Ott, J.; Gogos, J. A.: Family–based associ-
ation

[132793]studies support a sexually dimorphic effect of COMT and
MAOA on genetic

[132794]susceptibility to obsessive–compulsive disorder. Biol. Psy-
chiat. 45:

[132795]1178–1189, 1999.

[132796]23681.

[132797]Taira, M.; Yoshida, T.; Miyagawa, K.; Sakamoto, H.; Terada, M.;

[132798]Sugimura, T.: cDNA sequence of human transforming gene hst and identification

[132799]of the coding sequence required for transforming activity. Proc.

[132800]Nat. Acad. Sci. 84: 2980–2984, 1987.

[132801]23682.

[132802]Allikmets, R.; Shroyer, N. F.; Singh, N.; Seddon, J. M.; Lewis,

[132803]R. A.; Bernstein, P. S.; Peiffer, A.; Zabriskie, N. A.; Hutchinson,

[132804]A.; Dean, M.; Lupski, J. R.; Leppert, M.: Mutation of the Stargardt

[132805]disease gene (ABCR) in age-related macular degeneration. Science 277:

[132806]1805–1807, 1997.

[132807]23683.

[132808]De La Paz, M. A.; Guy, V. K.; Abou-Donia, S.; Heinis, R.; Bracken,

[132809]B.; Vance, J. M.; Gilbert, J. R.; Gass, J. D. M.; Haines, J. L.; Pericak-Vance,

- [132810]M. A.: Analysis of the Stargardt disease gene (ABCR) in age-related
macular degeneration. Ophthalmology 106: 1531–1536, 1999.
- [132811]23684.
- [132812]Farndon, J. R.; Leight, G. S.; Dilley, W. G.; Baylin, S. B.; Smallridge,
R. C.; Harrison, T. S.; Wells, S. A., Jr.: Familial medullary thyroid
carcinoma without associated endocrinopathies: a distinct clinical
entity. Brit. J. Surg. 73: 278–281, 1986.
- [132813]23685.
- [132814]Taylor, M. D.; Liu, L.; Raffel, C.; Hui, C.; Mainprize, T. G.; Zhang, X.; Agatep, R.; Chiappa, S.; Gao, L.; Lowrance, A.; Hao, A.; Goldstein, A. M.; Stavrou, T.; Scherer, S. W.; Dura, W. T.; Wainwright,
B.; Squire, J. A.; Rutka, J. T.; Hogg, D.: Mutations in SUFU predispose
to medulloblastoma. Nature Genet. 31: 306–310, 2002.

[132823]23686.

[132824]Ball, E. W.; Meynell, M. J.; Beale, D.; Kynoch, P.; Lehmann, H.;

[132825]Strelton, A. O. W.: Haemoglobin alpha(2) prime: alpha 2 gamma 2 (16

[132826]glycine to arginine). Nature 209: 1217–1218, 1966.

[132827]23687.

[132828]Barnabas, J.; Muller, C. J.: Haemoglobin Lepore (Hollandia). Nature 194:

[132829]931–932, 1962.

[132830]23688.

[132831]Beaven, G. H.; Gratzner, W. B.; Stevens, B. L.; Shooter, E. M.;

[132832]Ellis, M. J.; White, J. C.; Gillespie, J. E. O.: An abnormal haemoglobin

[132833](Lepore–Cyprus) resembling haemoglobin Lepore and its interaction

[132834]with thalassaemia. Brit. J. Haemat. 10: 159–170, 1964.

[132835]23689.

[132836]Brennan, S. O.; Williamson, D.; Smith, M. B.; Cauchi, M. N.; Macphee,

[132837]A.; Carrell, R. W.: HbA(2) Victoria, delta24 (B6) gly-to-asp:
a new

[132838]delta chain variant occurring with beta-thalassemia.
Hemoglobin 8:

[132839]163-168, 1984.

[132840]23690.

[132841]Camaschella, C.; Alfarano, A.; Gottardi, E.; Travi, M.; Prim-
ignani,

[132842]P.; Cappio, F. C.; Saglio, G.: Prenatal diagnosis of fetal
hemoglobin

[132843]Lepore-Boston disease on maternal peripheral blood.
Blood 75: 2102-2106,

[132844]1990.

[132845]23691.

[132846]Codrington, J. F.; Kutlar, F.; Harris, H. F.; Wilson, J. B.;
Stoming,

[132847]T. A.; Huisman, T. H. J.: Hb A(2)-Wrens or
delta98(FG5)val-to-met,

[132848]an unstable delta-chain variant identified by sequence
analysis of

[132849]amplified DNA. Biochim. Biophys. Acta 1009: 87-89,
1989.

[132850]23692.

[132851]Curtain, C. C.: A structural study of abnormal
haemoglobins occurring

[132852]in New Guinea. Aust. J. Exp. Biol. Med. Sci. 42: 89–97,
1964.

[132853]23693.

[132854]De Angioletti, M.; Di Girgenti, C.; Messineo, R.; Capra, M.;
Carestia,

[132855]C.: Hb A2–Monreale (delta–146 (HC3) his-to-arg), a novel
delta chain

[132856]variant detected in west Sicily. Hemoglobin 26: 1–5, 2002.

[132857]23694.

[132858]De Jong, W. W.; Went, L. N.: Hemoglobin A(2)–NYU in the
Netherlands:

[132859]incidence of delta–chain variants in human populations.
Hum. Hered. 24:

[132860]32–39, 1974.

[132861]23695.

[132862]De Jong, W. W. W.; Bernini, L. F.: Haemoglobin Babinga
(delta

[132863]136 glycine-aspartic acid): a new delta chain variant. Na–

ture 219:

[132864]1360–1362, 1968.

[132865]23696.

[132866]Efremov, G. D.: Hemoglobins Lepore and anti-Lepore.

Hemoglobin 2:

[132867]197–233, 1978.

[132868]23697.

[132869]McMichael, A.; Makgoba, W.: Complexity in human histocompatibility

[132870]loci. Nature 293: 701–702, 1981.

[132871]23698.

[132872]Moen, T.; Albrechtsen, D.; Flatmark, A.; Jakobsen, A.; Jervell,

[132873]J.; Halvorsen, S.; Solheim, B. G.; Thorsby, E.: Importance of HLA-DR

[132874]matching in cadaveric renal transplantation: a prospective one-center

[132875]study of 170 transplants. New Eng. J. Med. 303: 850–854, 1980.

[132876]23699.

[132877]Olerup, O.; Troye-Blomberg, M.; Schreuder, G. M. T.; Ri-

ley, E.

[132878]M.: HLA-DR and -DQ gene polymorphism in West Africans
is twice as

[132879]extensive as in North European Caucasians: evolutionary
implications. Proc.

[132880]Nat. Acad. Sci. 88: 8480-8484, 1991.

[132881]23700.

[132882]Park, M. S.; Terasaki, P. I.; Bernoco, D.: Relationship be-
tween

[132883]MT and DR antigens.In: Terasaki, P. I.: Histocompatibility
Testing

[132884]1980. Los Angeles: UCLA Press (pub.) 1980.

[132885]23701.

[132886]Park, M. S.; Terasaki, P. I.; Bernoco, D.; Iwaki, Y.: Evidence

[132887]for a second B-cell locus separate from the DR locus.
Transplant.

[132888]Proc. 10: 823-828, 1978.

[132889]23702.

[132890]Pisella, P.-J.; Brignole, F.; Debbasch, C.; Lozato, P.-A.;
Creuzot-Garcher,

[132891]C.; Bara, J.; Saiag, P.; Warnet, J.-M.; Baudouin, C.: Flow cy-

tometric

[132892]analysis of conjunctival epithelium in ocular rosacea and
keratoconjunctivitis

[132893]sicca. Ophthalmology 107: 1841-1849, 2000.

[132894]23703.

[132895]Rollini, P.; Mach, B.; Gorski, J.: Linkage map of three HLA-
DR

[132896]beta-chain genes: evidence for a recent duplication event.
Proc.

[132897]Nat. Acad. Sci. 82: 7197-7201, 1985.

[132898]23704.

[132899]Sachs, J. A.; Jaraquemada, D.; Festenstein, H.: Intra HLA-D
region

[132900]recombinant maps HLA-DR between HLA-B and HLA-D.
Tissue Antigens 17:

[132901]43-56, 1981.

[132902]23705.

[132903]Schwartz, B. D.: Personal Communication. St. Louis, Mo.
2/22/1983.

[132904]23706.

[132905]Shackelford, D. A.; Mann, D. L.; van Rood, J. J.; Ferrara, G.

[132906]B.; Strominger, J. L.: Human B-cell alloantigens DC1, MT1,
and LB12

[132907]are identical to each other but distinct from the HLA-DR
antigen. Proc.

[132908]Nat. Acad. Sci. 78: 4566-4570, 1981.

[132909]23707.

[132910]Strominger, J. L.: Biology of the human histocompatibility
leukocyte

[132911]antigen (HLA) system and a hypothesis regarding the gen-
eration of

[132912]autoimmune diseases. J. Clin. Invest. 77: 1411-1415,
1986.

[132913]23708.

[132914]Suciu-Foca, N.; Weiner, J.; Rohowsky, C.; McKiernan, P.;
Susinno,

[132915]E.; Rubinstein, P.: Indications that Dw determinants are
controlled

[132916]by distinct (but closely linked) genes. Transplant. Proc. 10:
799-804,

[132917]1978.

[132918]23709.

[132919]Thursz, M. R.; Thomas, H. C.; Greenwood, B. M.; Hill, A. V.
S.

[132920]Heterozygote advantage for HLA class-II type in hepatitis
B virus

[132921]infection. (Letter) Nature Genet. 17: 11-12, 1997.

[132922]23710.

[132923]Tiercy, J.-M.; Gorski, J.; Jeannet, M.; Mach, B.: Identifica-
tion

[132924]and distribution of three serologically undetected alleles
of HLA-DR

[132925]by oligonucleotide-DNA typing analysis. Proc. Nat. Acad.
Sci. 85:

[132926]198-202, 1988.

[132927]23711.

[132928]Tiwari, J. L.; Terasaki, P. I.: HLA and Disease Associations.

[132929]New York: Springer-Verlag (pub.) 1985.

[132930]23712.

[132931]Trowsdale, J.: Genetics and polymorphism: class II anti-
gens. Brit.

[132932]Med. Bull. 43: 15-36, 1987.

[132933]23713.

[132934]Tsubota, K.; Fukagawa, K.; Fujihara, T.; Shimmura, S.;
Saito,

[132935]I.; Saito, K.; Takeuchi, T.: Regulation of human leukocyte
antigen

[132936]expression in human conjunctival epithelium. Invest.
Ophthalm. Vis.

[132937]Sci. 40: 28–34, 1999.

[132938]23714.

[132939]Walker, L. E.; Hewick, R.; Hunkapiller, M. W.; Hood, L. E.;
Dreyer,

[132940]W. J.; Reisfeld, R. A.: N-terminal amino acid sequences of
the alpha

[132941]and beta chains of HLA-DR1 and HLA-DR2 antigens. Bio-
chemistry 23:

[132942]185–188, 1983.

[132943]23715.

[132944]Spelbrink, J. N.; Li, F.-Y.; Tiranti, V.; Nikali, K.; Yuan,
Q.-P.;

[132945]Tariq, M.; Wanrooij, S.; Garrido, N.; Comi, G.; Morandi, L.;
Santoro,

[132946]L.; Toscano, A.; and 9 others: Human mitochondrial DNA deletions

[132947]associated with mutations in the gene encoding twinkle, a phage T7

[132948]gene 4-like protein localized in mitochondria. Nature Genet. 28:

[132949]223-231, 2001.

[132950]23716.

[132951]Farfel, Z.; Iiri, T.; Shapira, H.; Roitman, A.; Mouallem, M.;

[132952]Bourne, H. R.: Pseudohypoparathyroidism, a novel mutation in the

[132953]beta/gamma- contact region of Gs-alpha impairs receptor stimulation. J.

[132954]Biol. Chem. 271: 19653-19655, 1996.

[132955]23717.

[132956]Heisler, L. K.; Cowley, M. A.; Tecott, L. H.; Fan, W.; Low, M.

[132957]J.; Smart, J. L.; Rubinstein, M.; Tatro, J. B.; Marcus, J. N.; Holstege,

[132958]H.; Lee, C. E.; Cone, R. D.; Elmquist, J. K.: Activation of central

[132959]melanocortin pathways by fenfluramine. Science 297:

609–611, 2002.

[132960]23718.

[132961]Folster–Holst, R.; Moises, H. W.; Yang, L.; Fritsch, W.;
Weissenbach,

[132962]J.; Christophers, E.: Linkage between atopy and the IgE
high–affinity

[132963]receptor gene at 11q13 in atopic dermatitis families.
Hum. Genet. 102:

[132964]236–239, 1998.

[132965]23719.

[132966]Hill, M. R.; Cookson, W. O. C. M.: A new variant of the beta
subunit

[132967]of the high–affinity receptor for immunoglobulin E
(Fc–epsilon–RI–beta

[132968]E237G): associations with measures of atopy and
bronchial hyper–responsiveness. Hum.

[132969]Molec. Genet. 5: 959–962, 1996.

[132970]23720.

[132971]Hupp, K.; Siwarski, D.; Mock, B. A.; Kinet, J.–P.: Gene map–
ping

[132972]of the three subunits of the high affinity FcR for IgE to

mouse chromosomes

[132973]1 and 19. J. Immun. 143: 3787–3791, 1989.

[132974]23721.

[132975]Kuster, H.; Zhang, L.; Brini, A. T.; MacGlashan, D. W. J.;
Kinet,

[132976]J.-P.: The gene and cDNA for the human high affinity im-
munoglobulin

[132977]E receptor beta chain and expression of the complete hu-
man receptor. J.

[132978]Biol. Chem. 267: 12782–12787, 1992.

[132979]23722.

[132980]Nagata, H.; Mutoh, H.; Kumahara, K.; Arimoto, Y.;
Tomemori, T.;

[132981]Sakurai, D.; Arase, K.; Ohno, K.; Yamakoshi, T.; Nakano,
K.; Okawa,

[132982]T.; Numata, T.; Konno, A.: Association between nasal al-
lergy and

[132983]a coding variant of the Fc-epsilon-RI-beta gene
Glu237Gly in a Japanese

[132984]population. Hum. Genet. 109: 262–266, 2001.

[132985]23723.

[132986] Shirakawa, T.; Li, A.; Dubowitz, M.; Dekker, J. W.; Shaw, A. E.;

[132987] Faux, J. A.; Ra, C.; Cookson, W. O. C. M.; Hopkin, J. M.: Association

[132988] between atopy and variants of the beta subunit of the high-affinity

[132989] immunoglobulin E receptor. *Nature Genet.* 7: 125–130, 1994.

[132990] 23724.

[132991] Shirakawa, T.; Mao, X.-Q.; Sasaki, S.; Enomoto, T.; Kawai, M.;

[132992] Morimoto, K.; Hopkin, J.: Association between atopic asthma and a

[132993] coding variant of Fc-epsilon-RI-beta in a Japanese population. *Hum.*

[132994] *Molec. Genet.* 5: 1129–1130, 1996.

[132995] 23725.

[132996] Szepietowski, P.; Gaudray, P.: FCER1B, a candidate gene for atopy,

[132997] is located in 11q13 between CD20 and TCN1. *Genomics* 19: 399–400,

[132998] 1994.

[132999]23726.

[133000]Allen, G.; Fantes, K. H.: A family of structural genes for
human

[133001]lymphoblastoid (leucocyte-type) interferon. Nature 287:
408–411,

[133002]1980.

[133003]23727.

[133004]Diaz, M. O.; Bohlander, S.; Allen, G.: Nomenclature of the
human

[133005]interferon genes. J. Interferon Cytokine Res. 16: 179–180,
1996.

[133006]23728.

[133007]Diaz, M. O.; Pomykala, H. M.; Bohlander, S. K.; Maltepe, E.;
Malik,

[133008]K.; Brownstein, B.; Olopade, O. I.: Structure of the human
type-I

[133009]interferon gene cluster determined from a YAC clone con-
tig. Genomics 22:

[133010]540–552, 1994.

[133011]23729.

[133012]Douglas, R. M.; Moore, B. W.; Miles, H. B.; Davies, L. M.;

Graham,

[133013]N. M. H.; Ryan, P.; Worswick, D. A.; Albrecht, J. K.: Prophylactic

[133014]efficacy of intranasal alpha-2-interferon against rhinovirus infections

[133015]in the family setting. New Eng. J. Med. 314: 65-70, 1986.

[133016]23730.

[133017]Edge, M. D.; Green, A. R.; Heathcliffe, G. R.; Meacock, P. A.;

[133018]Schuch, W.; Scanlon, D. B.; Atkinson, T. C.; Newton, C. R.; Markham,

[133019]A. F.: Total synthesis of a human leukocyte interferon gene. Nature 292:

[133020]756-762, 1981.

[133021]23731.

[133022]Fountain, J. W.; Karayiorgou, M.; Taruscio, D.; Graw, S. L.; Buckler,

[133023]A. J.; Ward, D. C.; Dracopoli, N. C.; Housman, D. E.: Genetic and

[133024]physical map of the interferon region on chromosome 9p. Genomics 14:

[133025]105-112, 1992.

[133026]23732.

[133027]Gillespie, D.; Carter, W.: Concerted evolution of human interferon

[133028]alpha genes. J. Interferon Res. 3: 83-88, 1983.

[133029]23733.

[133030]Hayden, F. G.; Albrecht, J. K.; Kaiser, D. L.; Gwaltney, J. M.,

[133031]Jr.: Prevention of natural colds by contact prophylaxis with intranasal

[133032]alpha-2-interferon. New Eng. J. Med. 314: 71-75, 1986.

[133033]23734.

[133034]Hitzeman, R. A.; Hagie, F. E.; Levine, H. L.; Goeddel, D. V.;

[133035]Ammerer, G.; Hall, B. D.: Expression of a human gene for interferon

[133036]in yeast. Nature 293: 717-722, 1981.

[133037]23735.

[133038]Imai, M.; Sano, T.; Yanase, Y.; Miyamoto, K.; Yonehara, S.; Mori,

[133039]H.; Honda, T.; Fukuda, S.; Nakamura, T.; Miyakawa, Y.; Mayumi, M.

[133040]: Demonstration of two subtypes of human leukocyte in-

terferon (IFN- α)

[133041]by monoclonal antibodies. J. Immun. 128: 2824-2825,
1982.

[133042]23736.

[133043]Isaacs, D.; Clarke, J. R.; Tyrrell, D. A. J.; Webster, A. D. B.;

[133044]Valman, H. B.: Deficient production of leucocyte interferon
(interferon- α)

[133045]in vitro and in vivo in children with recurrent respiratory
tract

[133046]infections. Lancet II: 950-952, 1981.

[133047]23737.

[133048]Lawn, R. M.; Adelman, J.; Dull, T. J.; Gross, M.; Goeddel,
D.;

[133049]Ullrich, A.: DNA sequence of two closely linked human
leukocyte interferon

[133050]genes. Science 212: 1159-1162, 1981.

[133051]23738.

[133052]Lawn, R. M.; Goeddel, D. V.; Ullrich, A.: The human inter-
feron

[133053]gene family.(Abstract) Sixth Int. Cong. Hum. Genet.,
Jerusalem 55

[133054]only, 1981.

[133055]23739.

[133056]Miyata, T.; Hayashida, H.: Recent divergence from a common ancestor

[133057]of human IFN- α genes. Nature 295: 165-168, 1982.

[133058]23740.

[133059]Mory, Y.; Chernajovsky, Y.; Feinstein, S. I.; Chen, L.; Weissenbach,

[133060]J.; Revel, M.: Expression of the cloned human interferon beta-1 gene

[133061]in E. coli.(Abstract) Sixth Int. Cong. Hum. Genet., Jerusalem 56

[133062]only, 1981.

[133063]23741.

[133064]Reis, A.: Personal Communication. Berlin, Germany 3/28/1994.

[133065]23742.

[133066]Blanchet-Bardon, C.; Nazzaro, V.; Chevrant-Breton, J.; Espie, M.;

[133067]Kerbrat, P.; Le Marec, B.: Hereditary epidermolytic palmo-plantar

[133068]keratoderma associated with breast and ovarian cancer in
a large kindred. Brit.

[133069]J. Derm. 117: 363-370, 1987.

[133070]23743.

[133071]Appel, S.; Filter, M.; Reis, A.; Hennies, H. C.; Bergheim, A.;

[133072]Ogilvie, E.; Arndt, S.; Simmons, A.; Lovett, M.; Hide, W.;
Ramsay,

[133073]M.; Reichwald, K.; Zimmermann, W.; Rosenthal, A.: Physi-
cal and transcriptional

[133074]map of the critical region for keratolytic winter erythema
(KWE) on

[133075]chromosome 8p22-p23 between D8S550 and D8S1759.
Europ. J. Hum. Genet. 10:

[133076]17-25, 2002.

[133077]23744.

[133078]Clarke, C. A.; McConnell, R. B.: Six cases of carcinoma of
the

[133079]oesophagus occurring in one family. Brit. Med. J. 2:
1137-1138,

[133080]1954.

[133081]23745.

[133082]Ellis, A.; Field, J. K.; Field, E. A.; Friedmann, P. S.; Fryer,
[133083]A.; Howard, P.; Leigh, I. M.; Risk, J.; Shaw, J. M.; Whittaker,
J.
[133084]: Tylosis associated with carcinoma of the oesophagus
and oral leukoplakia
[133085]in a large Liverpool family--a review of six generations.
Europ.
[133086]J. Cancer 30B: 102-112, 1994.
[133087]23746.
[133088]Harper, P. S.; Harper, R. M. J.; Howel-Evans, A. W.: Carci-
noma
[133089]of the oesophagus with tylosis. Quart. J. Med. 39:
317-333, 1970.
[133090]23747.
[133091]Olavarria, E.; Craddock, C.; Dazzi, F.; Marin, D.; Marktel,
S.;
[133092]Apperley, J. F.; Goldman, J. M.: Imatinib mesylate (STI571)
in the
[133093]treatment of relapse of chronic myeloid leukemia after al-
logeneic
[133094]stem cell transplantation. Blood 99: 3861-3862, 2002.

[133095]23748.

[133096]Pegoraro, L.; Matera, L.; Ritz, J.; Levis, A.; Palumbo, A.; Biagini,

[133097]G.: Establishment of a Ph(1)-positive human cell line (BV173). J.

[133098]Nat. Cancer Inst. 70: 447-451, 1983.

[133099]23749.

[133100]Perrotti, D.; Cesi, V.; Trotta, R.; Guerzoni, C.; Santilli, G.;

[133101]Campbell, K.; Iervolino, A.; Condorelli, F.; Gambacorti-Passerini,

[133102]C.; Caligiuri, M. A.; Calabretta, B.: BCR-ABL suppresses C/EBP-alpha

[133103]expression through inhibitory action of hnRNP E2. Nature Genet. 30:

[133104]48-58, 2002.

[133105]23750.

[133106]Prakash, O.; Yunis, J. J.: High resolution chromosomes of the

[133107]t(9;22) positive leukemias. Cancer Genet. Cytogenet. 11: 361-367,

[133108]1984.

[133109]23751.

[133110]Priest, J. R.; Robison, L. L.; McKenna, R. W.; Lindquist, L.
L.;

[133111]Warkentin, P. I.; LeBien, T. W.; Woods, W. G.; Kersey, J. H.;
Coccia,

[133112]P. F.; Nesbit, M. E., Jr.: Philadelphia chromosome positive
childhood

[133113]acute lymphoblastic leukemia. Blood 56: 15-22, 1980.

[133114]23752.

[133115]Rowley, J. D.: A new consistent chromosomal abnormality
in chronic

[133116]myelogenous leukemia identified by quinacrine fluores-
cence and Giemsa

[133117]staining. Nature 243: 290-293, 1973.

[133118]23753.

[133119]Rubin, C. M.; Carrino, J. J.; Dickler, M. N.; Leibowitz, D.;
Smith,

[133120]S. D.; Westbrook, C. A.: Heterogeneity of genomic fusion
of BCR and

[133121]ABL in Philadelphia chromosome-positive acute lym-
phoblastic leukemia. Proc.

[133122]Nat. Acad. Sci. 85: 2795–2799, 1988.

[133123]23754.

[133124]Saglio, G.; Storlazzi, C. T.; Giugliano, E.; Surace, C.; Anelli,

[133125]L.; Rege-Cambrin, G.; Zagaria, A.; Velasco, A. J.; Heiniger, A.; Scaravaglio,

[133126]P.; Gomez, A. T.; Gomez, J. R.; Archidiacono, N.; Banfi, S.; Rocchi,

[133127]M.: A 76-kb duplicon maps close to the BCR gene on chromosome 22

[133128]and the ABL gene on chromosome 9: possible involvement in the genesis

[133129]of the Philadelphia chromosome translocation. Proc. Nat. Acad. Sci. 99:

[133130]9882–9887, 2002.

[133131]23755.

[133132]Savage, D. G.; Antman, K. H.: Imatinib mesylate—a new oral targeted

[133133]therapy. New Eng. J. Med. 346: 683–693, 2002.

[133134]23756.

[133135]Sawyers, C. L.: Chronic myeloid leukemia. New Eng. J. Med. 340:

[133136]1330–1340, 1999.

[133137]23757.

[133138]Schaefer–Rego, K.; Dudek, H.; Popenoe, D.; Arlin, Z.;
Mears, J.

[133139]G.; Bank, A.; Leibowitz, D.: CML patients in blast crisis
have breakpoints

[133140]localized to a specific region of the BCR. Blood 70:
448–455, 1987.

[133141]23758.

[133142]Shtivelman, E.; Gale, R. P.; Dreazen, O.; Berrebi, A.; Zaizov,

[133143]R.; Kubonishi, I.; Miyoshi, I.; Canaani, E.: bcr–abl RNA in
patients

[133144]with chronic myelogenous leukemia. Blood 69: 971–973,
1987.

[133145]23759.

[133146]Shtivelman, E.; Lifshitz, B.; Gale, R. P.; Canaani, E.: Fused

[133147]transcript of abl and bcr genes in chronic myelogenous
leukaemia. Nature 315:

[133148]550–554, 1985.

[133149]23760.

[133150]Skorski, T.; Nieborowska-Skorska, M.; Nicolaides, N. C.;
 Szczalik,
 [133151]C.; Iversen, P.; Iozzo, R. V.; Zon, G.; Calabretta, B.: Sup-
 pression
 [133152]of Philadelphia-1 leukemia cell growth in mice by BCR-
 ABL antisense
 [133153]oligodeoxynucleotide. Proc. Nat. Acad. Sci. 91:
 4504-4508, 1994.
 [133154]23761.

[133155]Stam, K.; Heisterkamp, N.; Grosveld, G.; de Klein, A.;
 Verma,
 [133156]R. S.; Coleman, M.; Dosik, H.; Groffen, J.: Evidence of a
 new chimeric
 [133157]bcr/c-abl mRNA in patients with chronic myelocytic
 leukemia and the
 [133158]Philadelphia chromosome. New Eng. J. Med. 313:
 1429-1433, 1985.
 [133159]23762.

[133160]Stam, K.; Heisterkamp, N.; Reynolds, F. H., Jr.; Groffen, J.:
 [133161]Evidence that the phl gene encodes a 160,000-dalton
 phosphoprotein
 [133162]with associated kinase activity. Molec. Cell. Biol. 7:

1955–1960,

[133163]1987.

[133164]23763.

[133165]Swan, D. C.; McBride, O. W.; Robbins, K. C.; Keithley, D. A.;

[133166]Reddy, E. P.; Aaronson, S. A.: Chromosomal mapping of
the simian

[133167]sarcoma virus onc gene analogue in human cells. Proc.
Nat. Acad.

[133168]Sci. 79: 4691–4695, 1982.

[133169]23764.

[133170]Tanabe, T.; Kuwabara, T.; Warashina, M.; Tani, K.; Taira,
K.;

[133171]Asano, S.: Oncogene inactivation in a mouse model: tissue
invasion

[133172]by leukaemic cells is stalled by loading them with a de-
signer ribozyme. Nature 406:

[133173]473–474, 2000.

[133174]23765.

[133175]Teyssier, J. R.; Bartram, C. R.; Deville, J.; Potron, G.; Pi-
geon,

[133176]F.: C-abl oncogene and chromosome 22 'bcr' juxtaposi-

tion in chronic

[133177]myelogenous leukemia. New Eng. J. Med. 312:

1393–1394, 1985.

[133178]23766.

[133179]Tkachuk, D. C.; Westbrook, C. A.; Andreeff, M.; Donlon, T.

A.;

[133180]Cleary, M. L.; Suryanarayan, K.; Homge, M.; Redner, A.;

Gray, J.;

[133181]Pinkel, D.: Detection of bcr–abl fusion in chronic myeloge-
neous leukemia

[133182]by in situ hybridization. Science 250: 559–562, 1990.

[133183]23767.

[133184]Verhest, A.; Monsieur, R.: Philadelphia chromosome–posi-
tive thrombocythemia

[133185]with leukemic transformation. (Letter) New Eng. J. Med.

308: 1603,

[133186]1983.

[133187]23768.

[133188]Verma, R. S.; Dosik, H.: Heteromorphisms of the Philadel-
phia

[133189](Ph–1) chromosome in patients with chronic myelogenous

leukaemia (CML).

[133190]l. Classification and clinical significance. Brit. J. Haemat.

45:

[133191]215-222, 1980.

[133192]23769.

[133193]Aisenberg, A. C.; Wilkes, B. M.; Jacobson, J. O.: The bcl-2
gene

[133194]is rearranged in many diffuse B-cell lymphomas. Blood
71: 969-972,

[133195]1988.

[133196]23770.

[133197]Bakhshi, A.; Jensen, J. P.; Goldman, P.; Wright, J. J.;
McBride,

[133198]O. W.; Epstein, A. L.; Korsmeyer, S. J.: Cloning the chro-
mosomal

[133199]breakpoint of t(14;18) human lymphomas: clustering
around J(H) on

[133200]chromosome 14 and near a transcriptional unit on 18. Cell
41: 899-906,

[133201]1985.

[133202]23771.

[133203]Bakhshi, A.; Wright, J. J.; Graninger, W.; Seto, M.; Owens, J.;

[133204]Cossman, J.; Jensen, J. P.; Goldman, P.; Korsmeyer, S. J.:
Mechanism

[133205]of the t(14;18) chromosomal translocation: structural
analysis of

[133206]both derivative 14 and 18 reciprocal partners. Proc. Nat.
Acad. Sci. 84:

[133207]2396–2400, 1987.

[133208]23772.

[133209]Chen, J.; Flannery, J. G.; LaVail, M. M.; Steinberg, R. H.; Xu,

[133210]J.; Simon, M. I.: Bcl-2 overexpression reduces apoptotic
photoreceptor

[133211]cell death in three different retinal degenerations. Proc.
Nat. Acad.

[133212]Sci. 93: 7042–7047, 1996.

[133213]23773.

[133214]Cleary, M. L.; Sklar, J.: Nucleotide sequence of a t(14;18)
chromosomal

[133215]breakpoint in follicular lymphoma and demonstration of a
breakpoint–cluster

[133216]region near a transcriptionally active locus on chromosome 18. Proc.

[133217]Nat. Acad. Sci. 82: 7439-7443, 1985.

[133218]23774.

[133219]Cleary, M. L.; Smith, S. D.; Sklar, J.: Cloning and structural

[133220]analysis of cDNAs for bcl-2 and a hybrid bcl-2/immunoglobulin transcript

[133221]resulting from the t(14;18) translocation. Cell 47: 19-28, 1986.

[133222]23775.

[133223]Cockerham, G. C.; Hidayat, A. A.; Bijwaard, K. E.; Sheng, Z.-M.

[133224]: Re-evaluation of 'reactive lymphoid hyperplasia of the uvea': an

[133225]immunohistochemical and molecular analysis of 10 cases. Ophthalmology 107:

[133226]151-158, 2000.

[133227]23776.

[133228]Croce, C. M.: Personal Communication. Philadelphia, Pa. 1989.

[133229]23777.

[133230]Efremov, G. D.; Rudivic, R.; Niazi, G. A.; Hunter, E., Jr.;
Huisman,

[133231]T. H. J.; Schroeder, W. A.: An individual with Hb-Lep-
ore-Baltimore-delta

[133232]beta-thalassaemia in a Yugoslavian family. Scand. J.
Haemat. 16:

[133233]81-89, 1976.

[133234]23778.

[133235]Fessas, P.; Stamatoyannopoulos, G.; Karaklis, A.:
Hemoglobin

[133236]'Pylos': study of a hemoglobinopathy resembling tha-
lassemia in the

[133237]heterozygous, homozygous and double heterozygous
state. Blood 19:

[133238]1-22, 1962.

[133239]23779.

[133240]Fioretti, G.; De Angioletti, M.; Masciangelo, F.; Lacerra, G.;

[133241]Scarallo, A.; de Bonis, C.; Pagano, L.; Guarino, E.; De Rosa,
L.;

[133242]Salvati, F.; Carestia, C.: Origin heterogeneity of Hb Lep-
ore-Boston

[133243]gene in Italy. Am. J. Hum. Genet. 50: 781–786, 1992.

[133244]23780.

[133245]Fujita, S.; Ohta, Y.; Saito, S.; Kobayashi, Y.; Naritomi, Y.;

[133246]Kawaguchi, T.; Imamura, T.; Wada, Y.; Hayashi, A.:

Hemoglobin A(2)

[133247]Honai (delta90(F6) glu-to-val): a new delta chain variant.

Hemoglobin 9:

[133248]597–607, 1985.

[133249]23781.

[133250]Galanello, R.; Gasperini, D.; Perseu, L.; Barella, S.; Ideo, A.;

[133251]Cao, A.: Hb A(2)–Sant' Antioco (delta93(F9) cys-to-gly): a
new delta

[133252]chain variant identified by sequencing of amplified DNA.

Hemoglobin 18:

[133253]437–439, 1994.

[133254]23782.

[133255]Gasperini, D.; Perseu, L.; Cossu, P.; Podda, R.; Cao, A.;

Galanello,

[133256]R.: A novel delta-0-thalassemia mutation: TCG-to-TAG
(trp-to-stop)

[133257]at codon 37. Hum. Mutat. 3: 71–72, 1994.

[133258]23783.

[133259]Gerald, P. S.; Diamond, L. K.: A new hereditary
hemoglobinopathy

[133260](the Lepore trait) and its interaction with thalassemia trait.
Blood 12:

[133261]835–844, 1958.

[133262]23784.

[133263]Harano, T.; Harano, K.; Kushida, Y.; Ueda, S.; Kawakami,
H.:

[133264]Hb A(2)–Niigata [δ –1(NA1)val–to–ala]: a new delta
chain variant

[133265]found in the Japanese population. Hemoglobin 15:
335–339, 1991.

[133266]23785.

[133267]Horton, B.; Payne, R. A.; Bridges, M. T.; Huisman, T. H. J.:

[133268]Studies on an abnormal minor hemoglobin component
Hb–beta(2). Clin.

[133269]Chim. Acta 6: 246–253, 1961.

[133270]23786.

[133271] Huehns, E. R.: A third haemoglobin abnormality in two individuals

[133272] with Hb-H disease. In: Lehmann, H.; Betke, K.:
Haemoglobin-Colloquium.

[133273] Stuttgart: Georg Thieme Verlag (pub.) 1962. Pp. 76 only.
[133274] 23787.

[133275] Huehns, E. R.; Dance, N.; Beaven, G. H.; Stevens, B. L.:
Further

[133276] investigations in haemoglobin H disease. (Abstract) Proc.
9th Cong.

[133277] Int. Soc. Hemat., Mexico City 7-9, 1962.

[133278] 23788.

[133279] Huisman, T. H. J.; Sydenstricker, V. P.: Haemoglobin: difference

[133280] in gross structure of two electrophoretically identical minor haemoglobin

[133281] components. Nature 193: 489-491, 1962.

[133282] 23789.

[133283] Jeffreys, A. J.: DNA sequence variants in the G-gamma, A-gamma,

[133284] delta and beta globin genes of man. Cell 18: 1-10, 1979.

[133285]23790.

[133286]Jones, R. T.; Brimhall, B.; Huehns, E. R.; Barnicot, N. A.:
Hemoglobin

[133287]Sphakia: a delta-chain variant of hemoglobin A2 from
Crete. Science 151:

[133288]1406–1408, 1966.

[133289]23791.

[133290]Jones, R. T.; Brimhall, B.; Huisman, T. H. J.: Structural
characterization

[133291]of two delta chain variants: hemoglobin A-prime-2 (B2)
and hemoglobin

[133292]Flatbush. J. Biol. Chem. 242: 5141–5145, 1967.

[133293]23792.

[133294]Juricic, D.; Crepinko, I.; Efremov, G. D.; Lam, H.; Webber,
B.

[133295]B.; Headlee, M. G.; Huisman, T. H. J.: Hb A(2)–Zagreb or
delta125(H3)gln-to-glu,

[133296]a new delta chain variant in association with delta-
beta-thalassemia. Hemoglobin 7:

[133297]443–448, 1983.

[133298]23793.

[133299]Klein, H. L.; Petes, T. D.: Intrachromosomal gene conversion

[133300]in yeast. Nature 289: 144–148, 1981.

[133301]23794.

[133302]Labie, D.; Schroeder, W. A.; Huisman, T. H. J.: The amino acid

[133303]sequence of the delta–beta chains of haemoglobin Lepore Augusta =

[133304]Lepore Washington. Biochim. Biophys. Acta 127: 428–437, 1966.

[133305]23795.

[133306]Lanclos, K. D.; Patterson, J.; Efremov, G. D.; Wong, S. C.; Villegas,

[133307]A.; Ojwang, P. J.; Wilson, J. B.; Kutlar, F.; Huisman, T. H. J.:

[133308]Characterization of chromosomes with hybrid genes for Hb Lepore–Washington,

[133309]Hb Lepore–Baltimore, Hb P–Nilotic, and Hb Kenya. Hum. Genet. 77:

[133310]40–45, 1987.

[133311]23796.

[133312]Lee, R. C.; Huisman, T. H. J.: A variant of hemoglobin A–2

found

[133313]in a Negro family. Blood 24: 495–501, 1964.

[133314]23797.

[133315]Lehmann, H.; Jenkins, T.; Plowman, D.; Nurse, G. T.: Homozygosity

[133316]for the delta-chain variant haemoglobin A(2)-prime (HbB-2) (delta16

[133317]gly-to-arg). Hemoglobin 9: 363–372, 1985.

[133318]23798.

[133319]Leung, H.; Gilbert, A. T.; Fleming, P. J.; Wong, J.; Hughes, W.

[133320]G.; Hussein, S.; Nash, A. R.: Hb A(2)-Parkville or delta47(CD6) asp-to-val,

[133321]a new delta chain variant. Hemoglobin 15: 407–416, 1991.

[133322]23799.

[133323]Lie-Injo, L. E.; Pribadi, W.; Westendorp-Boerma, F.; Efremov,

[133324]G. D.; Wilson, J. B.; Reynolds, C. A.; Huisman, T. H. J.: Hemoglobin

[133325]A(2)-Indonesia or alpha(2) beta(2) 69(E13)gly to arg.

Biochim. Biophys.

[133326]Acta 229: 335–342, 1971.

[133327]23800.

[133328]Losekoot, M.; Fodde, R.; Giordano, P. C.; Bernini, L. F.: A novel

[133329]delta-0-thalassemia arising from a frameshift insertion, detected

[133330]by direct sequencing of enzymatically amplified DNA. Hum. Genet. 83:

[133331]75–78, 1989.

[133332]23801.

[133333]Loudianos, G.; Cao, A.; Pirastu, M.; Vassilopoulos, G.; Kol-
lia,

[133334]P.; Loukopoulos, D.: Molecular basis of the delta tha-
lassemia in

[133335]cis to hemoglobin Knossos variant. (Letter) Blood 77:
2087–2088,

[133336]1991.

[133337]23802.

[133338]Loudianos, G.; Murru, S.; Kanavakis, E.; Metaxotou-Mavro-
mati,

[133339]A.; Theodoropoulou, D.; Kattamis, C.; Cao, A.; Pirastu, M.:

A new

[133340]delta chain variant hemoglobin A(2)-Corfu or al-

pha(2)delta(2) 116

[133341]arg-to-cys (G18), detected by delta-globin gene analysis

in a Greek

[133342]family. Hum. Genet. 87: 237-238, 1991.

[133343]23803.

[133344]Loudianos, G.; Murru, S.; Ristaldi, M. S.; Cossu, P.; Pilia,

G.;

[133345]Porcu, S.; Sciarratta, G. V.; Parodi, M. I.; Cao, A.; Pirastu,

M.

[133346]: A novel delta-thalassemia mutation: a G-to-C substitu-

tion at codon

[133347]30 of the delta-globin gene in a person of southern Italian

origin. Hum.

[133348]Mutat. 1: 169-171, 1992.

[133349]23804.

[133350]Loudianos, G.; Porcu, S.; Cossu, P.; Tannoia, N.; Vitucci,

A.;

[133351]Campanale, D.; Cao, A.; Pirastu, M.: A new delta-chain

variant hemoglobin

[133352]A(2)-Puglia or delta26 glu-to-asp (B8), detected by DNA analysis in

[133353]a family of southern Italian extraction. Hum. Mutat. 2: 327-329,

[133354]1993.

[133355]23805.

[133356]Martin, S. L.; Zimmer, E. A.; Kan, Y. W.; Wilson, A. C.: Silent

[133357]delta-globin gene in Old World monkeys. Proc. Nat. Acad. Sci. 77:

[133358]3563-3566, 1980.

[133359]23806.

[133360]Matsuda, M.; Sakamoto, N.; Fukumaki, Y.: Delta-thalassemia caused

[133361]by disruption of the site for an erythroid-specific transcription

[133362]factor, GATA-1, in the delta-globin gene promoter. Blood 80: 1347-1351,

[133363]1992.

[133364]23807.

[133365]Moi, P.; Loudianos, G.; Lavinha, J.; Murru, S.; Cossu, P.; Casu,

[133366]R.; Oggiano, L.; Longinotti, M.; Cao, A.; Pirastu, M.: Delta-thalassemia

[133367]due to a mutation in an erythroid-specific binding protein sequence

[133368]3-prime to the delta-globin gene. Blood 79: 512-516, 1992.

[133369]23808.

[133370]Moi, P.; Paglietti, E.; Sanna, A.; Brancati, C.; Tagarelli, A.;

[133371]Galanello, R.; Cao, A.; Pirastu, M.: Delineation of the molecular

[133372]basis of delta- and normal HbA2 beta-thalassemia. Blood 72: 530-533,

[133373]1988.

[133374]23809.

[133375]Guida, P. M.; Todd, J. E.; Moore, S. W.; Beal, J. M.: Zollinger-Ellison

[133376]syndrome with interesting variations: report of twelve cases including

[133377]one of carcinoid of the duodenum. Am. J. Surg. 112: 807-817, 1966.

[133378]23810.

- [133379]Guo, S.S.; Sawicki, M. P.: Molecular and genetic mechanisms of
tumorigenesis in multiple endocrine neoplasia type-1.
Molec. Endocr. 15:
1653-1664, 2001.
- [133382]23811.
- [133383]Guru, S. C.; Agarwal, S. K.; Manickam, P.; Olufemi, S.-E.;
Crabtree,
J. S.; Weisemann, J. M.; Kester, M. B.; Kim, Y. S.; Wang, Y.;
Emmert-Buck,
M. R.; Liotta, L. A.; Spiegel, A. M.; Boguski, M. S.; Roe, B.
A.;
- [133386]Collins, F. S.; Marx, S. J.; Burns, L.; Chandrasekharappa, S.
C.:
- [133387]A transcript map for the 2.8-Mb region containing the
multiple endocrine
neoplasia type 1 locus. Genome Res. 7: 725-735, 1997.
- [133389]23812.
- [133390]Guru, S. C.; Crabtree, J. S.; Brown, K. D.; Dunn, K. J.; Man-
ickam,
P.; Prasad, N. B.; Wangsa, D.; Burns, A. L.; Spiegel, A. M.;
Marx,

[133392]S. J.; Pavan, W. J.; Collins, F. S.; Chandrasekharappa, S. C.:
Isolation,
[133393]genomic organization, and expression analysis of Men1,
the murine
[133394]homolog of the MEN1 gene. Mammalian Genome 10:
592–596, 1999.
[133395]23813.
[133396]Guru, S. C.; Goldsmith, P. K.; Burns, A. L.; Marx, S. J.;
Spiegel,
[133397]A. M.; Collins, F. S.; Chandrasekharappa, S. C.: Menin, the
product
[133398]of the MEN1 gene, is a nuclear protein. Proc. Nat. Acad.
Sci. 95:
[133399]1630–1634, 1998.
[133400]23814.
[133401]Heppner, C.; Kester, M. B.; Agarwal, S. K.; Debelenko, L.
V.;
[133402]Emmert–Buck, M. R.; Guru, S. C.; Manickam, P.; Olufemi,
S.–E.; Skarulis,
[133403]M. C.; Doppman, J. L.; Alexander, R. H.; Kim, Y. S.; Sagggar,
S. K.;
[133404]Lubensky, I. A.; Zhuang, Z.; Liotta, L. A.; Chan–

drasekharappa, S.

[133405]C.; Collins, F. S.; Spiegel, A. M.; Burns, A. L.; Marx, S. J.:

Somatic

[133406]mutation of the MEN1 gene in parathyroid tumours. Na-

ture Genet. 16:

[133407]375-378, 1997.

[133408]23815.

[133409]Heppner, C.; Reincke, M.; Agarwal, S. K.; Mora, P.; Allolio,

B.;

[133410]Burns, A. L.; Spiegel, A. M.; Marx, S. J.: MEN1 gene analysis

in

[133411]sporadic adrenocortical neoplasms. J. Clin. Endocr. Metab.

84: 216-219,

[133412]1999.

[133413]23816.

[133414]Hershon, K. S.; Kelly, W. A.; Shaw, C. M.; Schwartz, R.;

Bierman,

[133415]E. L.: Prolactinomas as part of the multiple endocrine neo-

plastic

[133416]syndrome type 1. Am. J. Med. 74: 713-720, 1983.

[133417]23817.

[133418]Hessman, O.; Skogseid, B.; Westin, G.; Akerstrom, G.: Multiple

[133419]allelic deletions and intratumoral genetic heterogeneity in MEN1 pancreatic

[133420]tumors. J. Clin. Endocr. Metab. 86: 1355–1361, 2001.

[133421]23818.

[133422]Johnson, G. J.; Summerskill, W. H. J.; Anderson, V. E.; Keating,

[133423]F. R.: Clinical and genetic investigation of a large kindred with

[133424]multiple endocrine adenomatosis. New Eng. J. Med. 277: 1379–1385,

[133425]1967.

[133426]23819.

[133427]Jones, B. S.; O'Hagan, J. J.; Phear, D. N.; Sheville, E.: A case

[133428]of the Zollinger–Ellison syndrome associated with hyperplasia of salivary

[133429]and Brunner's glands. Gut 11: 837–839, 1970.

[133430]23820.

[133431]Kaji, H.; Canaff, L.; Lebrun, J.-J.; Goltzman, D.; Hendy, G. N.

- [133432]: Inactivation of menin, a Smad3-interacting protein,
blocks transforming
- [133433]growth factor type beta signaling. Proc. Nat. Acad. Sci. 98:
3837–3842,
- [133434]2001.
- [133435]23821.
- [133436]Kjellman, M.; Roshani, L.; Teh, B. T.; Kallioniemi, O.-P.;
Hoog,
- [133437]A.; Gray, S.; Farnebo, L.-O.; Holst, M.; Backdahl, M.; Lars-
son, C.
- [133438]: Genotyping of adrenocortical tumors: very frequent
deletions of
- [133439]the MEN1 locus in 11q13 and of a 1-centimorgan region
in 2p16. J.
- [133440]Clin. Endocr. Metab. 84: 730–735, 1999.
- [133441]23822.
- [133442]Lamers, C. B. H. W.; Froeling, P. G. A. M.: Clinical signifi-
cance
- [133443]of hyperparathyroidism in familial multiple endocrine ade-
nomatosis. Am.
- [133444]J. Med. 66: 422–424, 1979.

[133445]23823.

[133446]Lamers, C. B. H. W.; Stadil, F.; van Tongeren, J. H.: Prevalence

[133447]of endocrine abnormalities in patients with the Zollinger–Ellison

[133448]syndrome in their families. Am. J. Med. 64: 607–612, 1978.

[133449]23824.

[133450]Larsson, C.; Shepherd, J.; Nakamura, Y.; Blomberg, C.; Weber,

[133451]G.; Werelius, B.; Hayward, N.; Teh, B.; Tokino, T.; Seizinger, B.;

[133452]Skogseid, B.; Oberg, K.; Nordenskjold, M.: Predictive testing for

[133453]multiple endocrine neoplasia type 1 using DNA polymorphisms. J. Clin.

[133454]Invest. 89: 1344–1349, 1992.

[133455]23825.

[133456]Larsson, C.; Skogseid, B.; Oberg, K.; Nakamura, Y.; Nordenskjold,

[133457]M.: Multiple endocrine neoplasia type 1 gene maps to chromosome 11

[133458]and is lost in insulinoma. Nature 332: 85–87, 1988.

[133459]23826.

[133460]Larsson, C.; Skogseid, B.; Oberg, K.; Nordenskjold, M.:

Linkage

[133461]of multiple endocrine neoplasia type 1 (MEN1) to markers
at 11q13.

[133462](Abstract) Cytogenet. Cell Genet. 51: 1029, 1989.

[133463]23827.

[133464]Lemmens, I.; Van de Ven, W. J. M.; Kas, K.; Zhang, C. X.;

Giraud,

[133465]S.; Wautot, V.; Buisson, N.; De Witte, K.; Salandre, J.;

Lenoir, G.;

[133466]Pugeat, M.; Calender, A.; and 19 others: Identification of
the

[133467]multiple endocrine neoplasia type 1 (MEN1) gene. Hum.
Molec. Genet. 6:

[133468]1177–1183, 1997.

[133469]23828.

[133470]Long, T. T.; Barton, T. K.; Draffin, R.; Reeves, W. J.; Mc-

Carty,

[133471]K. S., Jr.: Conservative management of the Zollinger-Elli-

son syndrome:

[133472] ectopic gastrin production by an ovarian cystadenoma.

J.A.M.A. 243:

[133473] 1837–1839, 1980.

[133474] 23829.

[133475] Lulu, D. J.; Corcoran, T. E.; Andre, M.: Familial endocrine
adenomatosis

[133476] with associated Zollinger–Ellison syndrome: Wermer's
syndrome. Am.

[133477] J. Surg. 115: 695–701, 1968.

[133478] 23830.

[133479] Mallette, L. E.; Bilezikian, J. P.; Ketcham, A. S.; Aurbach, G.

[133480] D.: Parathyroid carcinoma in familial hyperparathyroidism.
Am. J.

[133481] Med. 57: 642–648, 1974.

[133482] 23831.

[133483] Maton, P. N.; Gardner, J. D.; Jensen, R. T.: Cushing's syn-
drome

[133484] in patients with the Zollinger–Ellison syndrome. New Eng.
J. Med. 315:

[133485] 1–5, 1986.

[133486]23832.

[133487]McCarthy, D. M.: Zollinger–Ellison syndrome. Ann. Rev.
Med. 33:

[133488]197–215, 1982.

[133489]23833.

[133490]McCarthy, D. M.; Olinger, E. J.; May, R. J.; Long, B. W.;
Gardner,

[133491]J. D.: H₂–histamine receptor blocking agents in the
Zollinger–Ellison

[133492]syndrome: experience in seven cases and implications for
long-term

[133493]therapy. Ann. Intern. Med. 87: 668–675, 1977.

[133494]23834.

[133495]Mee, A. S.; Ismail, S.; Bornman, P. C.; Marks, I. N.: Chang-
ing

[133496]concepts in the presentation, diagnosis and management
of the Zollinger–Ellison

[133497]syndrome. Quart. J. Med. 206: 256–267, 1983.

[133498]23835.

[133499]Nakamura, Y.; Larsson, C.; Julier, C.; Bystrom, C.; Skog-
seid,

[133500]B.; Wells, S.; Oberg, K.; Carlson, M.; Taggart, T.; O'Connell, P.;

[133501]Leppert, M.; Lalouel, J.-M.; Nordenskjold, M.; White, R.:
Localization

[133502]of the genetic defect in multiple endocrine neoplasia type
I within

[133503]a small region of chromosome 11. Am. J. Hum. Genet. 44:
751-755,

[133504]1989.

[133505]23836.

[133506]Nordenskjold, M.; Jones, C. A.; Glaser, T.; Werelius, B.;
Larsson,

[133507]C.; Jansson, M.: Mapping of chromosome 11q probes on
somatic cell

[133508]hybrids. (Abstract) Cytogenet. Cell Genet. 51: 1054, 1989.

[133509]23837.

[133510]Basson, C. T.; Bachinsky, D. R.; Lin, R. C.; Levi, T.; Elkins,

[133511]J. A.; Soultz, J.; Grayzel, D.; Kroumpouzou, E.; Traill, T. A.;
Leblanc-Straceski,

[133512]J.; Renault, B.; Kucherlapati, R.; Seidman, J. G.; Seidman, C.
E.

[133513]: Mutations in human cause limb and cardiac malforma-
tion in Holt-Oram

[133514]syndrome. Nature Genet. 15: 30-35, 1997.

[133515]23838.

[133516]Basson, C. T.; Huang, T.; Lin, R. C.; Bachinsky, D. R.; Wer-
emowicz,

[133517]S.; Vaglio, A.; Bruzzone, R.; Quadrelli, R.; Lerone, M.;
Romeo, G.;

[133518]Silengo, M.; Pereira, A.; Krieger, J.; Mesquita, S. F.; Kamis-

ago,

[133519]M.; Morton, C. C.; Pierpont, M. E. M.; Muller, C. W.; Seidman, J.

[133520]G.; Seidman, C. E.: Different TBX5 interactions in heart and limb

[133521]defined by Holt–Oram syndrome mutations. Proc. Nat. Acad. Sci. 96:

[133522]2919–2924, 1999.

[133523]23839.

[133524]Glenn, C. L.; Wang, W. Y. S.; Benjafield, A. V.; Morris, B. J.

[133525]: Linkage and association of tumor necrosis factor receptor 2 locus

[133526]with hypertension, hypercholesterolemia and plasma shed receptor. Hum.

[133527]Molec. Genet. 9: 1943–1949, 2000.

[133528]23840.

[133529]Jones, C.: Synteny between the Pro(plus) marker and human glutamate

[133530]oxaloacetate transaminase. Somat. Cell Genet. 1: 345–354, 1975.

[133531]23841.

[133532]Lasky, L. A.; Singer, M. S.; Dowbenko, D.; Imai, Y.; Henzel, W.

[133533]J.; Grimley, C.; Fennie, C.; Gillett, N.; Watson, S. R.; Rosen, S.

[133534]D.: An endothelial ligand for L-selectin is a novel mucin-like molecule. Cell 69:

[133535]927-938, 1992.

[133536]23842.

[133537]Lasky, L. A.; Singer, M. S.; Yednock, T. A.; Dowbenko, D.; Fennie,

[133538]C.; Rodriguez, H.; Nguyen, T.; Stachel, S.; Rosen, S. D.: Cloning

[133539]of a lymphocyte homing receptor reveals a lectin domain. Cell 56:

[133540]1045-1055, 1989.

[133541]23843.

[133542]Ord, D. C.; Ernst, T. J.; Zhou, L.-J.; Rambaldi, A.; Spertini,

[133543]O.; Griffin, J.; Tedder, T. F.: Structure of the gene encoding the

[133544]human leukocyte adhesion molecule-1 (TQ1, Leu-8) of lymphocytes and

[133545]neutrophils. J. Biol. Chem. 265: 7760–7767, 1990.

[133546]23844.

[133547]Siegelman, M. H.; Weissman, I. L.: Human homologue of mouse lymph

[133548]node homing receptor: evolutionary conservation at tandem cell interaction

[133549]domains. Proc. Nat. Acad. Sci. 86: 5562–5566, 1989.

[133550]23845.

[133551]Sitrin, R. G.; Pan, P. M.; Blackwood, R. A.; Huang, J.; Petty,

[133552]H. R.: Cutting edge: evidence for a signaling partnership between

[133553]urokinase receptors (CD87) and L-selectin (CD62L) in human polymorphonuclear

[133554]neutrophils. J. Immun. 166: 4822–4825, 2001.

[133555]23846.

[133556]Tedder, T. F.; Isaacs, C. M.; Ernst, T. J.; Demetri, G. D.; Adler,

[133557]D. A.; Disteche, C. M.: Isolation and chromosomal localization of

[133558]cDNAs encoding a novel human lymphocyte cell surface molecule, LAM-1:

[133559]homology with the mouse lymphocyte homing receptor
and other human

[133560]adhesion proteins. J. Exp. Med. 170: 123–133, 1989.

[133561]23847.

[133562]Aizawa, S.; Nakano, H.; Ishida, T.; Horie, R.; Nagai, M.; Ito,

[133563]K.; Yagita, H.; Okumura, K.; Inoue, J.; Watanabe, T.: Tumor
necrosis

[133564]factor receptor-associated factor (TRAF) 5 and TRAF2 are
involved

[133565]in CD30-mediated NF-kappa-B activation. J. Biol. Chem.
272: 2042–2045,

[133566]1997.

[133567]23848.

[133568]Durkop, H.; Latza, U.; Hummel, M.; Eitelbach, F.; Seed, B.;
Stein,

[133569]H.: Molecular cloning and expression of a new member of
the nerve

[133570]growth factor receptor family that is characteristic for
Hodgkin's

[133571]disease. Cell 68: 421–427, 1992.

[133572]23849.

[133573]Kurts, C.; Carbone, F. R.; Krummel, M. F.; Koch, K. M.;
Miller,

[133574]J. F. A. P.; Heath, W. R.: Signalling through CD30 protects
against

[133575]autoimmune diabetes mediated by CD8 T cells. Nature
398: 341–344,

[133576]1999.

[133577]23850.

[133578]McClive, P. J.; Morahan, G.: Assignment of the mouse ho-
mologues

[133579]of 6 loci from HSA1p to chromosomes 3 and 4. Genomics
23: 243–246,

[133580]1994.

[133581]23851.

[133582]Stein, H.; Gerdes, J.; Schwab, U.; Lemke, H.; Mason, D. Y.;
Ziegler,

[133583]A.; Schienle, W.; Diehl, V.: Identification of Hodgkin and
Sternberg–Reed

[133584]cells as a unique cell type derived from a newly detected
small cell

[133585]population. Int. J. Cancer 30: 445–449, 1982.

[133586]23852.

[133587]de Lau, W.; Clevers, H.: LEF1 turns over a new leaf. *Nature Genet.* 28:

[133588]3–5, 2001.

[133589]23853.

[133590]Hovanes, K.; Li, T. W. H.; Munguia, J. E.; Truong, T.; Milovanovic,

[133591]T.; Marsh, J. L.; Holcombe, R. F.; Waterman, M. L.: Beta-catenin-sensitive

[133592]isoforms of lymphoid enhancer factor-1 are selectively expressed in

[133593]colon cancer. *Nature Genet.* 28: 53–57, 2001.

[133594]23854.

[133595]Love, J. J.; Li, X.; Case, D. A.; Giese, K.; Grosschedl, R.; Wright,

[133596]P. E.: Structural basis for DNA bending by the architectural transcription

[133597]factor LEF-1. *Nature* 376: 791–795, 1995.

[133598]23855.

[133599]Milatovich, A.; Travis, A.; Grosschedl, R.; Francke, U.: LEF1,

[133600]the gene for lymphoid enhancer-binding factor 1, mapped
to human chromosome
[133601]4 (q23-q25) and distal mouse chromosome 3 by Southern
blot analysis
[133602]and fluorescence in situ hybridization. (Abstract) Cyto-
genet. Cell
[133603]Genet. 58: 1888 only, 1991.
[133604]23856.
[133605]Milatovich, A.; Travis, A.; Grosschedl, R.; Francke, U.: Gene
[133606]for lymphoid enhancer-binding factor 1 (LEF1) mapped to
human chromosome
[133607]4 (q23-q25) and mouse chromosome 3 near Egf. Ge-
nomics 11: 1040-1048,
[133608]1991.
[133609]23857.
[133610]van Genderen, C.; Okamura, R. M.; Farinas, I.; Quo, R. G.;
Parslow,
[133611]T. G.; Bruhn, L.; Grosschedl, R.: Development of several
organs that
[133612]require inductive epithelial-mesenchymal interactions is
impaired

[133613]in LEF-1-deficient mice. Genes Dev. 8: 2691-2703, 1994.

[133614]23858.

[133615]Waterman, M. L.; Fischer, W. H.; Jones, K. A.: A thymus-specific

[133616]member of the HMG protein family regulates the human T cell receptor

[133617]C alpha enhancer. Genes Dev. 5: 656-669, 1991.

[133618]23859.

[133619]Zhou, P.; Byrne, C.; Jacobs, J.; Fuchs, E.: Lymphoid enhancer

[133620]factor 1 directs hair follicle patterning and epithelial cell fate. Genes

[133621]Dev. 9: 700-713, 1995.

[133622]23860.

[133623]White, P. S.; Jensen, S. J.; Rajalingam, V.; Stairs, D.; Sulman,

[133624]E. P.; Maris, J. M.; Biegel, J. A.; Wooster, R.; Brodeur, G. M.:

[133625]Physical mapping of the CA6, ENO1, and SLC2A5 (GLUT5) genes and reassignment

[133626]of SLC2A5 to 1p36.2. Cytogenet. Cell Genet. 81: 60-64, 1998.

[133627]23861.

[133628]Puranam, R. S.; Eubanks, J. H.; Heinemann, S. F.; McNamara, J.

[133629]O.: Chromosomal localization of gene for human glutamate receptor

[133630]subunit-7. Somat. Cell Molec. Genet. 19: 581-588, 1993.

[133631]23862.

[133632]Paschen, W.; Hedreen, J. C.; Ross, C. A.: RNA editing of the glutamate

[133633]receptor subunits GluR2 and GluR6 in human brain tissue. J. Neurochem. 63:

[133634]1596-1602, 1994.

[133635]23863.

[133636]Gregor, P.; Reeves, R. H.; Jabs, E. W.; Yang, X.; Dackowski, W.;

[133637]Rochelle, J. M.; Brown, R. H., Jr.; Haines, J. L.; O'Hara, B. F.;

[133638]Uhl, G. R.; Seldin, M. F.: Chromosomal localization of glutamate

[133639]receptor genes: relationship to familial amyotrophic lateral sclerosis

[133640]and other neurological disorders of mice and humans.

Proc. Nat. Acad.

[133641]Sci. 90: 3053–3057, 1993.

[133642]23864.

[133643]McNamara, J. O.; Eubanks, J. H.; McPherson, J. D.; Wasmuth, J.

[133644]J.; Evans, G. A.; Heinemann, S. F.: Chromosomal localization of human

[133645]glutamate receptor genes. J. Neurosci. 12: 2555–2562, 1992.

[133646]23865.

[133647]Armstrong, N.; Gouaux, E.: Mechanisms for activation and antagonism

[133648]of an AMPA-sensitive glutamate receptor: crystal structures of the

[133649]GluR2 ligand binding core. Neuron 28: 165–181, 2000.

[133650]23866.

[133651]Brusa, R.; Zimmermann, F.; Koh, D.-S.; Feldmeyer, D.; Gass, P.;

[133652]Seeburg, P. H.; Sprengel, R.: Early-onset epilepsy and postnatal

[133653]lethality associated with an editing-deficient GluR-B allele

in mice. Science 270:

[133654]1677–1680, 1995.

[133655]23867.

[133656]Hollmann, M.; Hartley, M.; Heinemann, S.: Ca(2+) permeability

[133657]of KA–AMPA–gated glutamate receptor channels depends on subunit composition. Science 252:

[133658]851–853, 1991.

[133659]23868.

[133660]Kohler, M.; Kornau, H. C.; Seeburg, P. H.: The organization of

[133661]the gene for the functionally dominant alpha-amino-3-hydroxy-5-methylisoxazole-4-propionic

[133662]acid receptor subunit Glu4-B. J. Biol. Chem. 269: 17367–17370, 1994.

[133663]23869.

[133664]Sun, W.; Ferrer-Montiel, A. V.; Schinder, A. F.; McPherson, J.

[133665]P.; Evans, G. A.; Montal, M.: Molecular cloning, chromosomal mapping,

[133666]and functional expression of human brain glutamate re-

ceptors. Proc.

[133667]Nat. Acad. Sci. 89: 1443-1447, 1992.

[133668]23870.

[133669]Sun, Y.; Olson, R.; Horning, M.; Armstrong, N.; Mayer, M.;
Gouaux,

[133670]E.: Mechanism of glutamate receptor desensitization. Na-
ture 417:

[133671]245-253, 2002.

[133672]23871.

[133673]Shows, T. B.: Linkage of loci for human pyruvate kinase
and mannosephosphate

[133674]isomerase in somatic cell hybrids. (Abstract) Am. J. Hum.
Genet. 24:

[133675]13A only, 1972.

[133676]23872.

[133677]Shows, T. B.: Somatic cell genetics of enzyme markers as-
sociated

[133678]with three human linkage groups.In: Davidson, R. L.: Proc.
Conf.

[133679]Somatic Cell Hybridization, Orlando, Fla. (pub.) 1973.

[133680]23873.

- [133681]Champion, M. J.; Brown, J. A.; Shows, T. B.: Assignment of cytoplasmic
- [133682]alpha-mannosidase (MAN-A) and confirmation of the mitochondrial isocitrate
- [133683]dehydrogenase (IDH-M) genes to the q11--qter region of chromosome
- [133684]15 in man. Cytogenet. Cell Genet. 22: 498-502, 1978.
- [133685]23874.
- [133686]Ferguson-Smith, M. A.; Westerveld, A.: Report of the committee
- [133687]on the genetic constitution of chromosomes 13, 14, 15, 16, 17, 18,
- [133688]19, 20, 21, and 22 (HGM5). Cytogenet. Cell Genet. 25: 59-73, 1979.
- [133689]23875.
- [133690]Neri, G.; Ricci, R.; Pelino, A.; Bova, R.; Tedeschi, B.; Serra, A.: A boy with ring chromosome 15 derived from a
- t(15q;15q) Robertsonian
- [133692]translocation in the mother: cytogenetic and biochemical findings. Am.
- [133693]J. Med. Genet. 14: 307-314, 1983.

[133694]23876.

[133695]Chui, D.; Oh-Eda, M.; Liao, Y.-F.; Panneerselvam, K.; Lai, A.;

[133696]Marek, K. W.; Freeze, H. H.; Moremen, K. W.; Fukuda, M. N.; Marth,

[133697]J. D.: Alpha-mannosidase-II deficiency results in dyserythropoiesis

[133698]and unveils an alternate pathway in oligosaccharide biosynthesis. Cell 90:

[133699]157-167, 1997.

[133700]23877.

[133701]Chui, D.; Sellakumar, G.; Green, R. S.; Sutton-Smith, M.; McQuistan,

[133702]T.; Marek, K. W.; Morris, H. R.; Dell, A.; Marth, J. D.: Genetic

[133703]remodeling of protein glycosylation in vivo induces autoimmune disease. Proc.

[133704]Nat. Acad. Sci. 98: 1142-1147, 2001.

[133705]23878.

[133706]Gasparini, P.; del Giudice, E. M.; Delaunay, J.; Totaro, A.; Granatiero,

- [133707]M.; Melchionda, S.; Zelante, L.; Iolascon, A.: Localization of the
- [133708]congenital dyserythropoietic anemia II locus to chromosome 20q11.2
- [133709]by genomewide search. *Am. J. Hum. Genet.* 61: 1112–1116, 1997.
- [133710]23879.
- [133711]Misago, M.; Liao, Y.-F.; Kudo, S.; Eto, S.; Mattei, M.-G.; Moremen,
- [133712]K. W.; Fukuda, M. N.: Molecular cloning and expression of cDNAs encoding
- [133713]human alpha-mannosidase II and a previously unrecognized alpha-mannosidase
- [133714]II(X) isozyme. *Proc. Nat. Acad. Sci.* 92: 11766–11770, 1995.
- [133715]23880.
- [133716]Moremen, K. W.; Robbins, P. W.: Isolation, characterization, and
- [133717]expression of cDNAs encoding murine alpha-mannosidase II, a Golgi
- [133718]enzyme that controls conversion of high mannose to complex N-glycans. *J.*

[133719]Cell Biol. 115: 1521–1534, 1991.

[133720]23881.

[133721]Wada, A.; Sakamoto, H.; Katoh, O.; Yoshida, T.; Yokota, J.;
Little,

[133722]P. F. R.; Sugimura, T.; Terada, M.: Two homologous onco-
genes, HST1

[133723]and INT2, are closely located in human genome. Biochem.
Biophys.

[133724]Res. Commun. 157: 828–835, 1988.

[133725]23882.

[133726]Yoshida, T.; Tsutsumi, M.; Sakamoto, H.; Miyagawa, K.;
Teshima,

[133727]S.; Sugimura, T.; Terada, M.: Expression of the HST1
oncogene in

[133728]human germ cell tumors. Biochem. Biophys. Res. Com-
mun. 155: 1324–1329,

[133729]1988.

[133730]23883.

[133731]Birchmeier, C.; Birnbaum, D.; Waitches, G.; Fasano, O.;
Wigler,

[133732]M.: Characterization of an activated human ROS gene.

Molec. Cell.

[133733]Biol. 6: 3109–3116, 1986.

[133734]23884.

[133735]Birchmeier, C.; Sharma, S.; Wigler, M.: Expression and re-
arrangement

[133736]of the ROS1 gene in human glioblastoma cells. Proc. Nat.
Acad. Sci. 84:

[133737]9270–9274, 1987.

[133738]23885.

[133739]Nagarajan, L.; Louie, E.; Tsujimoto, Y.; Balduzzi, P. C.;
Huebner,

[133740]K.; Croce, C. M.: The human c-ros gene (ROS) is located at
chromosome

[133741]region 6q16–6q22. Proc. Nat. Acad. Sci. 83: 6568–6572,
1986.

[133742]23886.

[133743]Rabin, M.; Birnbaum, D.; Wigler, M.; Ruddle, F. H.: MCF3
oncogene

[133744]mapped to region on chromosome 6 associated with ma-
lignant transformation.

[133745](Abstract) Am. J. Hum. Genet. 37 only, 1985.

[133746]23887.

[133747]Rabin, M.; Birnbaum, D.; Young, D.; Birchmeier, C.; Wigler, M.;

[133748]Ruddle, F. H.: Human ROS1 and MAS1 oncogenes located in regions of

[133749]chromosome 6 associated with tumor-specific rearrangements. Oncogene

[133750]Res. 1: 169-178, 1987.

[133751]23888.

[133752]Satoh, H.; Yoshida, M. C.; Matsushime, H.; Shibuya, M.; Sasaki,

[133753]M.: Regional localization of the human c-ros-1 on 6q22 and flt on

[133754]13q12. Jpn. J. Cancer Res. 78: 772-775, 1987.

[133755]23889.

[133756]Yoshida, M. C.; Satoh, H.; Matsushime, H.; Shibuya, M.; Sasaki,

[133757]M.: Two ros-related protooncogenes, c-ros-1 and flt, are regionally

[133758]mapped on human chromosomes 6q22 and 13q12, respectively. (Abstract) Cytogenet.

[133759]Cell Genet. 46: 724 only, 1987.

[133760]23890.

[133761]Fong, G.-H.; Rossant, J.; Breitman, M. L.: Role of the Flt-1
receptor

[133762]tyrosine kinase in regulating the assembly of vascular en-
dothelium. Nature 376:

[133763]65-69, 1995.

[133764]23891.

[133765]He, Y.; Smith, S. K.; Day, K. A.; Clark, D. E.; Licence, D. R.;

[133766]Charnock-Jones, D. S.: Alternative splicing of vascular en-
dothelial

[133767]growth factor (VEGF)-R1 (FLT-1) pre-mRNA is important
for the regulation

[133768]of VEGF activity. Molec. Endocr. 13: 537-545, 1999.

[133769]23892.

[133770]Imbert, A.; Rosnet, O.; Marchetto, S.; Ollendorff, V.; Birn-
baum,

[133771]D.; Pebusque, M.-J.: Characterization of a yeast artificial
chromosome

[133772]from human chromosome band 13q12 containing the
FLT1 and FLT3 receptor-type

[133773] tyrosine kinase genes. Cytogenet. Cell Genet. 67:
175–177, 1994.

[133774] 23893.

[133775] Kendall, R. L.; Wang, G.; Thomas, K. A.: Identification of a
natural

[133776] soluble form of the vascular endothelial growth factor re-
ceptor, FLT-1,

[133777] and its heterodimerization with KDR. Biochem. Biophys.
Res. Commun. 226:

[133778] 324–328, 1996.

[133779] 23894.

[133780] Shalaby, F.; Rossant, J.; Yamaguchi, T. P.; Gertsenstein, M.;
Wu,

[133781] X.-F.; Breitman, M. L.; Schuh, A. C.: Failure of blood-is-
land formation

[133782] and vasculogenesis in Flk-1-deficient mice. Nature 376:
62–65, 1995.

[133783] 23895.

[133784] Wiesmann, C.; Fuh, G.; Christinger, H. W.; Eigenbrot, C.;
Wells,

[133785] J. A.; de Vos, A. M.: Crystal structure at 1.7 Angstrom res-
olution

[133786] of VEGF in complex with domain 2 of the Flt-1 receptor.

Cell 91:

[133787] 695–704, 1997. Cell 91:695–704, 1997

[133788] 23896.

[133789] Dabiri, G. A.; Young, C. L.; Rosenbloom, J.; Southwick, F.

S.:

[133790] Molecular cloning of human macrophage capping protein

cDNA: a unique

[133791] member of the gelsolin/villin family expressed primarily in
macrophages. J.

[133792] Biol. Chem. 267: 16545–16552, 1992.

[133793] 23897.

[133794] Mishra, V. S.; Henske, E. P.; Kwiatkowski, D. J.; Southwick,

F.

[133795] S.: The human actin–regulatory protein cap G: gene struc-
ture and

[133796] chromosome location. Genomics 23: 560–565, 1994.

[133797] 23898.

[133798] Southwick, F. S.; DiNubile, M. J.: Rabbit alveolar

macrophages

[133799] contain a Ca(2+)-sensitive, 41,000–Dalton protein which

reversibly

[133800]blocks the 'barbed' ends of actin filaments but does not sever them. J.

[133801]Biol. Chem. 261: 14191-14195, 1986.

[133802]23899.

[133803]Fukai, K.; Oh, J.; Frenk, E.; Almodovar, C.; Spritz, R. A.: Linkage

[133804]disequilibrium mapping of the gene for Hermansky-Pudlak syndrome to

[133805]chromosome 10q23.1-q23.3. Hum. Molec. Genet. 4: 1665-1669, 1995.

[133806]23900.

[133807]Le Beau, M. M.; Diaz, M. O.; Plowman, G. D.; Brown, J. P.; Rowley,

[133808]J. D.: Chromosomal sublocalization of the human p97 melanoma antigen. Hum.

[133809]Genet. 72: 294-296, 1986.

[133810]23901.

[133811]Plowman, G. D.; Brown, J. P.; Enns, C. A.; Schroder, J.; Nikinmaa,

[133812]B.; Sussman, H. H.; Hellstrom, K. E.; Hellstrom, I.: Assign-

ment of

[133813]the gene for human melanoma-associated antigen p97 to
chromosome 3. Nature 303:

[133814]70–72, 1983.

[133815]23902.

[133816]Rose, T. M.; Plowman, G. D.; Teplow, D. B.; Dreyer, W. J.;
Hellstrom,

[133817]K. E.; Brown, J. P.: Primary structure of the human
melanoma-associated

[133818]antigen p97 (melanotransferrin) deduced from the mRNA
sequence. Proc.

[133819]Nat. Acad. Sci. 83: 1261–1265, 1986.

[133820]23903.

[133821]Seligman, P. A.; Butler, C. D.; Massey, E. J.; Kaur, J. A.;
Brown,

[133822]J. P.; Plowman, G. D.; Miller, Y.; Jones, C.: The p97 antigen
is

[133823]mapped to the q24–qter region of chromosome 3, the
same region as

[133824]the transferrin receptor. Am. J. Hum. Genet. 38: 540–548,
1986.

[133825]23904.

[133826]Yang, F.; Lum, J. B.; McGill, J. R.; Moore, C. M.; Naylor, S. L.;

[133827]van Bragt, P. H.; Baldwin, W. D.; Bowman, B. H.: Human transferrin:

[133828]cDNA characterization and chromosomal localization. Proc. Nat. Acad.

[133829]Sci. 81: 2752–2756, 1984.

[133830]23905.

[133831]Randerson–Moor, J. A.; Harland, M.; Williams, S.; Cuthbert–Heavens,

[133832]D.; Sheridan, E.; Aveyard, J.; Sibley, K.; Whitaker, L.; Knowles,

[133833]M.; Bishop, J. N.; Bishop, D. T.: A germline deletion of p14(ARF)

[133834]but not CDKN2A in a melanoma–neural system tumour syndrome family. Hum.

[133835]Molec. Genet. 10: 55–62, 2001.

[133836]23906.

[133837]Fong, L. Y. Y.; Fidanza, V.; Zanesi, N.; Lock, L. F.; Siracusa,

[133838]L. D.; Mancini, R.; Siprashvili, Z.; Ottey, M.; Martin, S. E.; Druck,

[133839]T.; McCue, P. A.; Croce, C. M.; Huebner, K.: Muir-Torre-like syndrome
 [133840]in Fhit-deficient mice. Proc. Nat. Acad. Sci. 97:
 4742-4747, 2000.
 [133841]23907.

[133842]Chemelli, R. M.; Willie, J. T.; Sinton, C. M.; Elmquist, J. K.;
 [133843]Scammell, T.; Lee, C.; Richardson, J. A.; Williams, S. C.;
 Xiong,
 [133844]Y.; Kisanuki, Y.; Fitch, T. E.; Nakazato, M.; Hammer, R. E.;
 Saper,
 [133845]C. B.; Yanagisawa, M.: Narcolepsy in orexin knockout
 mice: molecular
 [133846]genetics of sleep regulation. Cell 437-451, 1999.
 [133847]23908.

[133848]Adkins, S.; Gan, K. N.; Mody, M.; La Du, B. N.: Molecular
 basis
 [133849]for the polymorphic forms of human serum paraoxonase/
 arylesterase:
 [133850]glutamine or arginine at position 191, for the respective A
 and B
 [133851]allozymes. Am. J. Hum. Genet. 52: 598-608, 1993.

[133852]23909.

[133853]Antikainen, M.; Murtomaki, S.; Syvanne, M.; Pahlman, R.;
Tahvanainen,

[133854]E.; Jauhiainen, M.; Frick, M. H.; Ehnholm, C.: The gln-
arg191 polymorphism

[133855]of the human paraoxonase gene (HUMPONA) is not asso-
ciated with the

[133856]risk of coronary artery disease in Finns. J. Clin. Invest. 98:
883-885,

[133857]1996.

[133858]23910.

[133859]Augustinsson, K.-B.; Henricson, B.: A genetically con-
trolled esterase

[133860]in rat plasma. Biochim. Biophys. Acta 124: 323-331,
1966.

[133861]23911.

[133862]Barbieri, M.; Bonafe, M.; Marfella, R.; Ragno, E.; Giugliano,
D.;

[133863]Franceschi, C.; Paolisso, G.: LL-paraoxonase genotype is
associated

[133864]with a more severe degree of homeostasis model assess-
ment IR in healthy

[133865]subjects. J. Clin. Endocr. Metab. 87: 222–225, 2002.

[133866]23912.

[133867]Brophy, V. H.; Jampsa, R. L.; Clendenning, J. B.; McKinstry, L.

[133868]A.; Jarvik, G. P.; Furlong, C. E.: Effects of 5–prime regulatory–region

[133869]polymorphisms on paraoxonase–gene (PON1) expression. Am. J. Hum.

[133870]Genet. 68: 1428–1436, 2001.

[133871]23913.

[133872]Cherry, N.; Mackness, M.; Durrington, P.; Povey, A.; Dippnall,

[133873]M.; Smith, T.; Mackness, B.: Paraoxonase (PON1) polymorphisms in

[133874]farmers attributing ill health to sheep dip. Lancet 359: 763–764,

[133875]2002.

[133876]23914.

[133877]Clendenning, J. B.; Humbert, R.; Green, E. D.; Wood, C.; Traver,

[133878]D.; Furlong, C. E.: Structural organization of the human

PON1 gene. Genomics 35:

[133879]586–589, 1996.

[133880]23915.

[133881]Coates, P. M.; Mestriner, M. A.; Hopkinson, D. A.: A preliminary

[133882]interpretation of the esterase isozymes of human tissues.
Ann. Hum.

[133883]Genet. 39: 1–20, 1975.

[133884]23916.

[133885]Davies, H. G.; Richter, R. J.; Keifer, M.; Broomfield, C. A.;
Sowalla,

[133886]J.; Furlong, C. E.: The effect of the human serum paraoxonase polymorphism

[133887]is reversed with diazoxon, soman and sarin. Nature Genet.
14: 334–336,

[133888]1996.

[133889]23917.

[133890]Deakin, S.; Leviev, I.; Nicaud, V.; Meynet, M.–C. B.; Tired,
L.;

[133891]James, R. W.: Paraoxonase–1 L55M polymorphism is associated with

[133892]an abnormal oral glucose tolerance test and differentiates
high risk

[133893]coronary disease families. J. Clin. Endocr. Metab. 87:
1268–1273,

[133894]2002.

[133895]23918.

[133896]Eckerson, H. W.; Romson, J.; Wyte, C.; La Du, B. N.: The
human

[133897]serum paraoxonase polymorphism: identification of phe-
notypes by their

[133898]response to salts. Am. J. Hum. Genet. 35: 214–227, 1983.

[133899]23919.

[133900]Eckerson, H. W.; Wyte, C. M.; La Du, B. N.: The human
serum paraoxonase/arylesterase

[133901]polymorphism. Am. J. Hum. Genet. 35: 1126–1136, 1983.

[133902]23920.

[133903]Eiberg, H.; Mohr, J.: Linkage relations of the paraoxonase
polymorphism

[133904]with 43 marker systems. (Abstract) Cytogenet. Cell Genet.
25: 150,

[133905]1979.

[133906]23921.

[133907]Rey-Campos, J.; Chouard, T.; Yaniv, M.; Cereghini, S.:
vHNF1

[133908]is a homeoprotein that activates transcription and forms
heterodimers

[133909]with HNF1. EMBO J. 10: 1445-1457, 1991.

[133910]23922.

[133911]Shih, D. Q.; Bussen, M.; Sehayek, E.; Ananthanarayanan,
M.; Shneider,

[133912]B. L.; Suchy, F. J.; Shefer, S.; Bollileni, J. S.; Gonzalez, F. J.;

[133913]Breslow, J. L.; Stoffel, M.: Hepatocyte nuclear factor-
1-alpha is

[133914]an essential regulator of bile acid and plasma cholesterol
metabolism. Nature

[133915]Genet. 27: 375-382, 2001.

[133916]23923.

[133917]Urhammer, S. A.; Hansen, T.; Ekstrom, C. T.; Eiberg, H.;
Pederson,

[133918]O.: The Ala/Val98 polymorphism of the hepatocyte nu-
clear factor-1-alpha

[133919]gene contributes to the interindividual variation in serum

C-peptide

[133920]response during an oral glucose tolerance test: evidence from studies

[133921]of 231 glucose-tolerant first degree relatives of type 2 diabetic

[133922]probands. J. Clin. Endocr. Metab. 83: 4506-4509, 1998.

[133923]23924.

[133924]Urhammer, S. A.; Moller, A. M.; Nyholm, B.; Ekstrom, C. T.; Eiberg,

[133925]H.; Clausen, J. O.; Hansen, T.; Pedersen, O.; Schmitz, O.: The effect

[133926]of two frequent amino acid variants of the hepatocyte nuclear factor-1-alpha

[133927]gene on estimates of the pancreatic beta-cell function in Caucasian

[133928]glucose-tolerant first-degree relatives of type 2 diabetic patients. J.

[133929]Clin. Endocr. Metab. 83: 3992-3995, 1998.

[133930]23925.

[133931]Urhammer, S. A.; Rasmussen, S. K.; Kaisaki, P. J.; Oda, N.; Yamagata,

[133932]K.; Moller, A. M.; Fridberg, M.; Hansen, L.; Hansen, T.; Bell,

G.

[133933]J.; Pedersen, O.: Genetic variation in the hepatocyte nuclear factor-1-alpha

[133934]gene in Danish Caucasians with late-onset NIDDM. Diabetologia 40:

[133935]473-475, 1997.

[133936]23926.

[133937]Vaxillaire, M.; Rouard, M.; Yamagata, K.; Oda, N.; Kaisaki, P.

[133938]J.; Boriraj, V. V.; Chevre, J.-C.; Boccio, V.; Cox, R. D.; Lathrop,

[133939]G. M.; Dussoix, P.; Philippe, J.; Timsit, J.; Charpentier, G.; Velho,

[133940]G.; Bell, G. I.; Froguel, P.: Identification of nine novel mutations

[133941]in the hepatocyte nuclear factor 1 alpha gene associated with maturity-onset

[133942]diabetes of the young (MODY3). Hum. Molec. Genet. 6: 583-586, 1997.

[133943]23927.

[133944]Yamada, S.; Nishigori, H.; Onda, H.; Utsugi, T.; Yanagawa, T.;

[133945]Maruyama, T.; Onigata, K.; Nagashima, K.; Nagai, R.;
Morikawa, A.;

[133946]Takeuchi, T.; Takeda, J.: Identification of mutations in the
hepatocyte

[133947]nuclear factor (HNF)-1-alpha gene in Japanese subjects
with IDDM. Diabetes 46:

[133948]1643-1647, 1997.

[133949]23928.

[133950]Yamagata, K.; Oda, N.; Kalsaki, P. J.; Menzel, S.; Furuta, H.;

[133951]Vaxillaire, M.; Southam, L.; Cox, R. D.; Lathrop, G. M.;
Borhaj, V.

[133952]V.; Chen, X.; Cox, N. J.; Oda, Y.; Yano, H.; Le Beau, M. M.;
Yamada,

[133953]S.; Nishigori, H.; Takeda, J.; Fajans, S. S.; Hattersley, A. T.;
Iwasaki,

[133954]N.; Hansen, T.; Pedersen, O.; Polonsky, K. S.; Turner, R. C.;
Velho,

[133955]G.; Chevre, J.-C.; Froguel, P.; Bell, G. I.: Mutations in the
hepatocyte

[133956]nuclear factor-1-alpha gene in maturity-onset diabetes of
the young

[133957](MODY3). Nature 384: 455-457, 1996.

[133958]23929.

[133959]Yoshiuchi, I.; Yamagata, K.; Yoshimoto, M.; Zhu, Q.; Yang, Q.;

[133960]Nammo, T.; Uenaka, R.; Kinoshita, E.; Hanafusa, T.; Miyagawa, J.;

[133961]Matsuzawa, Y.: Analysis of a non-functional HNF-1-alpha (TCF1) mutation

[133962]in Japanese subjects with familial type 1 diabetes. Hum. Mutat. 18:

[133963]345-351, 2001.

[133964]23930.

[133965]Dhanasekaran, S. M.; Barrette, T. R.; Ghosh, D.; Shah, R.; Varambally,

[133966]S.; Kurachi, K.; Pienta, K. J.; Rubin, M. A.; Chinnaiyan, A. M.:

[133967]Delineation of prognostic biomarkers in prostate cancer. Nature 412:

[133968]822-826, 2001.

[133969]23931.

[133970]Leytus, S. P.; Loeb, K. R.; Hagen, F. S.; Kurachi, K.; Davie, E.

[133971]W.: A novel trypsin-like serine protease (hepsin) with a putative

[133972]transmembrane domain expressed by human liver and hepatoma cells. *Biochemistry* 27:

[133973]1067-1074, 1988.

[133974]23932.

[133975]Tsuji, A.; Torres-Rosado, A.; Arai, T.; Le Beau, M. M.; Lemons,

[133976]R. S.; Chou, S.-H.; Kurachi, K.: Hepsin, a cell membrane-associated

[133977]protease: characterization, tissue distribution, and gene localization. *J.*

[133978]*Biol. Chem.* 266: 16948-16953, 1991.

[133979]23933.

[133980]Wu, Q.; Yu, D.; Post, J.; Halks-Miller, M.; Sadler, J. E.; Morser,

[133981]J.: Generation and characterization of mice deficient in hepsin,

[133982]a hepatic transmembrane serine protease. *J. Clin. Invest.* 101: 321-326,

[133983]1998.

[133984]23934.

[133985]Cannella, B.; Hoban, C. J.; Gao, Y.-L.; Garcia-Arenas, R.;
Lawson,

[133986]D.; Marchionni, M.; Gwynne, D.; Raine, C. S.: The neureg-
ulin, glial

[133987]growth factor 2, diminishes autoimmune demyelination
and enhances

[133988]remyelination in a chronic relapsing model for multiple
sclerosis. Proc.

[133989]Nat. Acad. Sci. 95: 10100-10105, 1998.

[133990]23935.

[133991]Falls, D. L.; Rosen, K. M.; Corfas, G.; Lane, W. S.; Fis-
chbach,

[133992]G. D.: ARIA, a protein that stimulates acetylcholine recep-
tor synthesis,

[133993]is a member of the Neu ligand family. Cell 72: 801-815,
1993.

[133994]23936.

[133995]Fernandez, P.-A.; Tang, D. G.; Cheng, L.; Prochiantz, A.;
Mudge,

[133996]A. W.; Raff, M. C.: Evidence that axon-derived neuregulin
promotes

[133997]oligodendrocyte survival in the developing rat optic nerve.

Neuron 28:

[133998]81–90, 2000.

[133999]23937.

[134000]Holmes, W. E.; Sliwkowski, M. X.; Akita, R. W.; Henzel, W.

J.;

[134001]Lee, J.; Park, J. W.; Yansura, D.; Abadi, N.; Raab, H.; Lewis,

G.

[134002]D.; Shepard, H. M.; Kuang, W.-J.; Wood, W. I.; Goeddel, D.

V.; Vandlen,

[134003]R. L.: Identification of heregulin, a specific activator of
p185(erbB2). Science 256:

[134004]1205–1210, 1992.

[134005]23938.

[134006]Huang, Y. Z.; Won, S.; Ali, D. W.; Wang, Q.; Tanowitz, M.;

Du,

[134007]Q. S.; Pelkey, K. A.; Yang, D. J.; Xiong, W. C.; Salter, M. W.;

Mei,

[134008]L.: Regulation of neuregulin signaling by PSD-95 interact-
ing with

[134009]ErbB4 at CNS synapses. Neuron 26: 443–455, 2000.

[134010]23939.

[134011]Lee, J.; Wood, W. I.: Assignment of heregulin (HGL) to human chromosome

[134012]8p22-p11 by PCR analysis of somatic cell hybrid DNA. Genomics 16:

[134013]790-791, 1993.

[134014]23940.

[134015]Martinou, J.-C.; Falls, D. L.; Fischbach, G. D.; Merlie, J. P.

[134016]: Acetylcholine receptor-inducing activity stimulates expression of

[134017]the epsilon-subunit gene of the muscle acetylcholine receptor. Proc.

[134018]Nat. Acad. Sci. 88: 7669-7673, 1991.

[134019]23941.

[134020]Ni, C.-Y.; Murphy, M. P.; Golde, T. E.; Carpenter, G.: Gamma-secretase

[134021]cleavage and nuclear localization of ErbB-4 receptor tyrosine kinase. Science 294:

[134022]2179-2181, 2001.

[134023]23942.

- [134024]Orr-Urtreger, A.; Trakhtenbrot, L.; Ben-Levy, R.; Wen, D.;
Rechavi,
[134025]G.; Lonai, P.; Yarden, Y.: Neural expression and chromo-
somal mapping
[134026]of NEU differentiation factor to 8p12-p21. Proc. Nat.
Acad. Sci. 90:
[134027]1867-1871, 1993.
[134028]23943.
[134029]Boyd, D.; Jain, S. K.; Crampton, J.; Barrett, K. J.; Drysdale,
[134030]J.: Isolation and characterization of a cDNA clone for hu-
man ferritin
[134031]heavy chain. Proc. Nat. Acad. Sci. 81: 4751-4755, 1984.
[134032]23944.
[134033]Caskey, J. H.; Jones, C.; Miller, Y. E.; Seligman, P. A.: Hu-
man
[134034]ferritin gene is assigned to chromosome 19. Proc. Nat.
Acad. Sci. 80:
[134035]482-486, 1983.
[134036]23945.
[134037]Costanzo, F.; Colombo, M.; Staempfli, S.; Santoro, C.;
Marone,

[134038]M.; Frank, R.; Delius, H.; Cortese, R.: Structure of gene and pseudogenes

[134039]of human apoferritin H. Nucleic Acids Res. 14: 721–736, 1986.

[134040]23946.

[134041]Cragg, S. J.; Drysdale, J.; Worwood, M.: Genes for the 'H' subunit

[134042]of human ferritin are present on a number of human chromosomes. Hum.

[134043]Genet. 71: 108–112, 1985.

[134044]23947.

[134045]Eisenstein, R. S.: Iron regulatory proteins and the molecular

[134046]control of mammalian iron metabolism. Annu. Rev. Nutr. 20: 627–662,

[134047]2000.

[134048]23948.

[134049]Ferreira, C.; Bucchini, D.; Martin, M.–E.; Levi, S.; Arosio, P.;

[134050]Grandchamp, B.; Beaumont, C.: Early embryonic lethality of H ferritin

[134051]gene deletion in mice. J. Biol. Chem. 275: 3021–3024,

2000.

[134052]23949.

[134053]Ferreira, C.; Santambrogio, P.; Martin, M.-E.; Andrieu, V.;
Feldmann,

[134054]G.; Henin, D.; Beaumont, C.: H ferritin knockout mice: a
model of

[134055]hyperferritinemia in the absence of iron overload. Blood
98: 525-532,

[134056]2001.

[134057]23950.

[134058]Gailani, M. R.; Petty, E. M.; Horsthemke, B.; Arnold, A.;
Marx,

[134059]S. J.; Bale, A. E.: Physical mapping of chromosome
11q12-13 by pulsed

[134060]field gel electrophoresis (PFGE). (Abstract) Cytogenet. Cell
Genet. 58:

[134061]1959 only, 1991.

[134062]23951.

[134063]Gatti, R. A.; Shaked, R.; Mohandas, T. K.; Salser, W.: Hu-
man

[134064]ferritin genes: chromosomal assignments and polymor-

phisms. Am. J.

[134065]Hum. Genet. 41: 654–667, 1987.

[134066]23952.

[134067]Harrison, P. M.; Arosio, P.: The ferritins: molecular properties,
ties,

[134068]iron storage function and cellular regulation. Biochim.
Biophys.

[134069]Acta 1275: 161–203, 1996.

[134070]23953.

[134071]Hentze, M. W.; Keim, S.; Papadopoulos, P.; O'Brien, S.;
Modi,

[134072]W.; Drysdale, J.; Leonard, W. J.; Harford, J. B.; Klausner, R.
D.

[134073]: Cloning, characterization, expression, and chromosomal
localization

[134074]of a human ferritin heavy-chain gene. Proc. Nat. Acad. Sci.
83:

[134075]7226–7230, 1986.

[134076]23954.

[134077]Kato, J.; Fujikawa, K.; Kanda, M.; Fukuda, N.; Sasaki, K.;
Takayama,

[134078]T.; Kobune, M.; Takada, K.; Takimoto, R.; Hamada, H.; Ikeda, T.; Niitsu, Y.: A mutation, in the iron-responsive element of H ferritin mRNA, causing autosomal dominant iron overload. *Am. J. Hum. Genet.* 69: 191–197, 2001.

[134082]23955.

[134083]Leibold, E. A.; Munro, H. N.: Cytoplasmic protein binds in vitro to a highly conserved sequence in the 5-prime untranslated region of ferritin heavy- and light-subunit mRNAs. *Proc. Nat. Acad. Sci.* 85: 2171–2175, 1988.

[134087]23956.

[134088]Lauren, P.: The two histological main types of gastric carcinoma: diffuse and so-called intestinal-type carcinoma. An attempt at a histo-clinical classification. *Acta Path. Microbiol. Scand.* 64: 31–49, 1965.

[134091]23957.

[134092]La Vecchia, C.; Negri, E.; Franceschi, S.; Gentile, A.: Family

[134093]history and the risk of stomach and colorectal cancer.

Cancer 70:

[134094]50–55, 1992.

[134095]23958.

[134096]Zanghieri, G.; Di Gregorio, C.; Sacchetti, C.; Fante, R.; Sas-
satelli,

[134097]R.; Cannizzo, G.; Carriero, A.; Ponz de Leon, M.: Familial
occurrence

[134098]of gastric cancer in the 2-year experience of a popula-
tion-based registry. Cancer 66:

[134099]2047–2051, 1990.

[134100]23959.

[134101]Canfield, V. A.; Okamoto, C. T.; Chow, D.; Dorfman, J.;
Gros, P.;

[134102]Forte, J. G.; Levenson, R.: Cloning of the H,K-ATPase beta
subunit:

[134103]tissue-specific expression, chromosomal assignment, and
relationship

[134104]to Na,K-ATPase beta subunits. J. Biol. Chem. 265:

19878–19884, 1990.

[134105]23960.

[134106]Maeda, M.; Oshiman, K.-I.; Tamura, S.; Futai, M.: Human gastric

[134107](H(+) + K(+))-ATPase gene: similarity to (Na(+) + K(+))-ATPase genes

[134108]in exon/intron organization but difference in control region. J.

[134109]Biol. Chem. 265: 9027–9032, 1990.

[134110]23961.

[134111]Song, I.; Yamada, T.; Trent, J. M.: Mapping of the gene encoding

[134112]the alpha-subunit of the human H(+),K(+)-ATPase to chromosome 19q13.1

[134113]by fluorescent in situ hybridization. Genomics 14: 547–548, 1992.

[134114]23962.

[134115]Anderson, L. A.; Friedman, L.; Osborne-Lawrence, S.; Lynch, E.;

[134116]Weissenbach, J.; Bowcock, A.; King, M.-C.: High-density genetic map

[134117]of the BRCA1 region of chromosome 17q12–q21. Ge-

nomics 17: 618–623,

[134118]1993.

[134119]23963.

[134120]Cheung, A. T.; Dayanandan, B.; Lewis, J. T.; Korbitt, G. S.;
Rajotte,

[134121]R. V.; Bryer–Ash, M.; Boylan, M. O.; Wolfe, M. M.; Kieffer,
T. J.

[134122]: Glucose–dependent insulin release from genetically en-
gineered K

[134123]cells. Science 290: 1959–1962, 2000.

[134124]23964.

[134125]Fukushima, Y.; Byers, M. G.; Eddy, R. L.; Haley, L. L.;
Henry,

[134126]W. M.; Inagaki, N.; Seino, Y.; Takeda, J.; Yano, H.; Yamada,
Y.; Imura,

[134127]H.; Bell, G. I.; Shows, T. B.: Localization of the human gas-
tric

[134128]inhibitory polypeptide gene (GIP) to chromosome
17q21.3–q22. (Abstract) Cytogenet.

[134129]Cell Genet. 51: 1001 only, 1989.

[134130]23965.

[134131]Inagaki, N.; Seino, Y.; Takeda, J.; Yano, H.; Yamada, Y.;
Bell,

[134132]G. I.; Eddy, R. L.; Fukushima, Y.; Byers, M. G.; Shows, T. B.;
Imura,

[134133]H.: Gastric inhibitory polypeptide: structure and chromo-
somal localization

[134134]of the human gene. Molec. Endocr. 3: 1014-1021, 1989.

[134135]23966.

[134136]Lacroix, A.; Bolte, E.; Tremblay, J.; Dupre, J.; Poitras, P.;
Fournier,

[134137]H.; Garon, J.; Garrel, D.; Bayard, F.; Taillefer, R.; Flanagan,
R.

[134138]J.; Hamet, P.: Gastric inhibitory polypeptide-dependent
cortisol

[134139]hypersecretion--a new cause of Cushing's syndrome. New
Eng. J. Med. 327:

[134140]974-980, 1992.

[134141]23967.

[134142]Lewis, T. B.; O'Connell, P.; Leach, R. J.: Localization of glu-
cose-dependent

[134143]insulinotropic polypeptide (GIP) to a gene cluster on chro-
mosome 17q. Genomics 19:

[134144]589–591, 1994.

[134145]23968.

[134146]Mohlke, K. L.; Nichols, W. C.; Westrick, R. J.; Novak, E. K.;
Cooney,

[134147]K. A.; Swank, R. T.; Ginsburg, D.: A novel modifier gene
for plasma

[134148]von Willebrand factor level maps to distal mouse chromo-
some 11. Proc.

[134149]Nat. Acad. Sci. 93: 15352–15357, 1996.

[134150]23969.

[134151]Reznik, Y.; Allali-Zerah, V.; Chayvialle, J. A.; Leroyer, R.;
Leymarie,

[134152]P.; Travert, G.; Lebrethon, M.-C.; Budi, I.; Balliere, A.-M.;
Mahoudeau,

[134153]J.: Food-dependent Cushing's syndrome mediated by
aberrant adrenal

[134154]sensitivity to gastric inhibitory polypeptide. New Eng. J.
Med. 327:

[134155]981–986, 1992.

[134156]23970.

[134157]Ambrose, C. M.; Duyao, M. P.; Barnes, G.; Bates, G. P.; Lin,

C.

[134158]S.; Srinidhi, J.; Baxendale, S.; Hummerich, H.; Lehrach, H.;
Altherr,

[134159]M.; Wasmuth, J.; Buckler, A.; Church, D.; Housman, D.;
Berks, M.;

[134160]Micklem, G.; Durbin, R.; Dodge, A.; Read, A.; Gusella, J.;
MacDonald,

[134161]M. E.: Structure and expression of the Huntington's dis-
ease gene:

[134162]evidence against simple inactivation due to an expanded
CAG repeat. Somat.

[134163]Cell Molec. Genet. 20: 27-38, 1994.

[134164]23971.

[134165]Andrew, S. E.; Goldberg, Y. P.; Kremer, B.; Squitieri, F.;
Theilmann,

[134166]J.; Zeisler, J.; Telenius, H.; Adam, S.; Almquist, E.; Anvret,
M.;

[134167]Lucotte, G.; Stoessl, A. J.; Campanella, G.; Hayden, M. R.:
Huntington

[134168]disease without CAG expansion: phenocopies or errors in
assignment? Am.

[134169]J. Hum. Genet. 54: 852-863, 1994.

[134170]23972.

[134171]Andrew, S. E.; Goldberg, Y. P.; Kremer, B.; Telenius, H.;
Theilmann,

[134172]J.; Adam, S.; Starr, E.; Squitieri, F.; Lin, B.; Kalchman, M. A.;

[134173]Graham, R. K.; Hayden, M. R.: The relationship between
trinucleotide

[134174](CAG) repeat length and clinical features of Huntington's
disease. Nature

[134175]Genet. 4: 398–403, 1993.

[134176]23973.

[134177]Aronin, N.; Chase, K.; Young, C.; Sapp, E.; Schwarz, C.;
Matta,

[134178]N.; Kornreich, R.; Landwehrmeyer, B.; Bird, E.; Beal, M. F.;
Vonsattel,

[134179]J.–P.; Smith, T.; Carraway, R.; Boyce, F. M.; Young, A. B.;
Penney,

[134180]J. B.; DiFiglia, M.: CAG expansion affects the expression of
mutant

[134181]huntingtin in Huntington's disease brain. Neuron 15:
1193–1201,

[134182]1995.

[134183]23974.

[134184]Auerbach, W.; Hurlbert, M. S.; Hilditch-Maguire, P.;
Wadghiri,

[134185]Y. Z.; Wheeler, V. C.; Cohen, S. I.; Joyner, A. L.; MacDonald,
M.

[134186]E.; Turbull, D. H.: The HD mutation causes progressive
lethal neurological

[134187]disease in mice expressing reduced levels of huntingtin.
Hum. Molec.

[134188]Genet. 10: 2515–2523, 2001.

[134189]23975.

[134190]Bamford, K. A.; Caine, E. D.; Kido, D. K.; Cox, C.; Shoul-
son,

[134191]J.: A prospective evaluation of cognitive decline in early
Huntington's

[134192]disease: functional and radiographic correlates. Neurology
45: 1867–1873,

[134193]1995.

[134194]23976.

[134195]Barbeau, A.: Parental ascent in the juvenile form of Hunt-
ington's

[134196]chorea. (Letter) Lancet II: 937 only, 1970.

[134197]23977.

[134198]Barkley, D. S.; Hardiwidjaja, S.; Menkes, J. H.: Abnormalities
ties

[134199]in growth of skin fibroblasts of patients with Huntington's
disease. Ann.

[134200]Neurol. 1: 426-430, 1977.

[134201]23978.

[134202]Barnes, G. T.; Duyao, M. P.; Ambrose, C. M.; McNeil, S.;
Perischetti,

[134203]F.; Srinidhi, J.; Gusella, J. F.; MacDonald, M. E.: Mouse
Huntington's

[134204]disease gene homolog (Hdh). Somat. Cell Molec. Genet.
20: 87-97,

[134205]1994.

[134206]23979.

[134207]Bates, G. P.; Mangiarini, L.; Mahal, A.; Davies, S. W.: Trans-
genic

[134208]models of Huntington's disease. Hum. Molec. Genet. 6:
1633-1637,

[134209]1997.

[134210]23980.

[134211]Bates, G. P.; Valdes, J.; Hummerich, H.; Baxendale, S.; Le
Paslier,

[134212]D. L.; Monaco, A. P.; Tagle, D.; MacDonald, M. E.; Altherr,
M.; Ross,

[134213]M.; Brownstein, B. H.; Bentley, D.; Wasmuth, J. J.; Gusella,
J. F.;

[134214]Cohen, D.; Collins, F.; Lehrach, H.: Characterization of a
yeast

[134215]artificial chromosome contig spanning the Huntington's
disease gene

[134216]candidate region. Nature Genet. 1: 180–187, 1992.

[134217]23981.

[134218]Baxendale, S.; Abdulla, S.; Elgar, G.; Buck, D.; Berks, M.;
Micklem,

[134219]G.; Durbin, R.; Bates, G.; Brenner, S.; Beck, S.; Lehrach, H.:
Comparative

[134220]sequence analysis of the human and pufferfish Hunting-
ton's disease

[134221]genes. Nature Genet. 10: 67–76, 1995.

[134222]23982.

- [134223]Behan, P. O.; Bone, I.: Hereditary chorea without dementia.
J.
- [134224]Neurol. Neurosurg. Psychiat. 40: 687–691, 1977.
- [134225]23983.
- [134226]Bernardi, G.: The isochore organization of the human
genome. Annu.
- [134227]Rev. Genet. 23: 637–661, 1989.
- [134228]23984.
- [134229]Bibb, J. A.; Yan, Z.; Svenningsson, P.; Snyder, G. L.; Pieri-
bone,
- [134230]V. A.; Horiuchi, A.; Nairn, A. C.; Messer, A.; Greengard, P.:
Severe
- [134231]deficiencies in dopamine signaling in presymptomatic
Huntington's
- [134232]disease mice. Proc. Nat. Acad. Sci. 97: 6809–6814, 2000.
- [134233]23985.
- [134234]Bird, E. D.; Caro, A. J.; Pilling, J. B.: A sex related factor
- [134235]in the inheritance of Huntington's chorea. Ann. Hum.
Genet. 37:
- [134236]255–260, 1974.

[134237]23986.

[134238]Bloch, M.; Hayden, M. R.: Predictive testing for Huntington disease

[134239]in childhood: challenges and implications. Am. J. Hum. Genet. 46:

[134240]1–4, 1990.

[134241]23987.

[134242]Bloch, M.; Hayden, M. R.: Preclinical testing in Huntington disease.

[134243](Letter) Am. J. Med. Genet. 27: 733–734, 1987.

[134244]23988.

[134245]Boehnke, M.; Conneally, P. M.; Lange, K.: Two models for a maternal

[134246]factor in the inheritance of Huntington disease. Am. J. Hum. Genet. 35:

[134247]845–860, 1983.

[134248]23989.

[134249]Brackenridge, C. J.: The relation of type of initial symptoms

[134250]and line of transmission to ages at onset and death in Huntington's

[134251]disease. Clin. Genet. 2: 287–297, 1971.

[134252]23990.

[134253]Brackenridge, C. J.: Familial correlations for age at onset
and

[134254]age at death in Huntington's disease. J. Med. Genet. 9:
23–32, 1972.

[134255]23991.

[134256]Brackenridge, C. J.: Relationship of parental age of rigidity
[134257]in Huntington's disease. J. Med. Genet. 11: 136–140,
1974.

[134258]23992.

[134259]Brackenridge, C. J.; Case, J.; Chiu, E.; Probert, D. N.;
Teltscher,

[134260]B.; Wallace, D. C.: A linkage study of the loci for Hunting-
ton's

[134261]disease and some common polymorphic markers. Ann.
Hum. Genet. 42:

[134262]201–211, 1978.

[134263]23993.

[134264]Brinkman, R. R.; Mezei, M. M.; Theilmann, J.; Almqvist, E.;
Hayden,

[134265]M. R.: The likelihood of being affected with Huntington disease by

[134266]a particular age, for a specific CAG size. Am. J. Hum. Genet. 60:

[134267]1202–1210, 1997.

[134268]23994.

[134269]Brothers, C. R. D.: The history and incidence of Huntington's

[134270]chorea in Tasmania. Proc. Roy. Aust. Coll. Physicians 4: 48–50,

[134271]1949.

[134272]23995.

[134273]Buetow, K. H.; Shiang, R.; Yang, P.; Nakamura, Y.; Lathrop, G.

[134274]M.; White, R.; Wasmuth, J. J.; Wood, S.; Berdahl, L. D.; Lysens,

[134275]N. J.; Ritty, T. M.; Wise, M. E.; Murray, J. C.: A detailed multipoint

[134276]map of human chromosome 4 provides evidence for linkage heterogeneity

[134277]and position-specific recombination rates. Am. J. Hum. Genet. 48:

[134278]911-925, 1991.

[134279]23996.

[134280]Bundey, S.: New mutations in Huntington's chorea. (Letter)
J.

[134281]Med. Genet. 20: 76-77, 1983.

[134282]23997.

[134283]Byers, R. K.; Dodge, J. A.: Huntington's chorea in children:

[134284]report of four cases. Neurology 17: 587-596, 1967.

[134285]23998.

[134286]Campbell, A. M. G.; Corner, B. D.; Norman, R. M.; Urich,
H.:

[134287]The rigid form of Huntington's disease. J. Neurol. Neuro-
surg. Psychiat. 24:

[134288]71-77, 1961.

[134289]23999.

[134290]Cariello, L.; de Cristofaro, T.; Zanetti, L.; Cuomo, T.; Di
Maio,

[134291]L.; Campanella, G.; Rinaldi, S.; Zanetti, P.; Di Lauro, R.;
Varrone,

[134292]S.: Transglutaminase activity is related to CAG repeat

length in

[134293]patients with Huntington's disease. Hum. Genet. 98:
633–635, 1996.

[134294]24000.

[134295]Marshall, D.: Ectodermal dysplasia: report of kindred with
ocular

[134296]abnormalities and hearing defect. Am. J. Ophthalm. 45:
143–156, 1958.

[134297]24001.

[134298]Korkko, J.; Ritvaniemi, P.; Haataja, L.; Kaariainen, H.;
Kivirikko,

[134299]K. I.; Prockop, D. J.; Ala-Kokko, L.: Mutation in type II pro-
collagen

[134300](COL2A1) that substitutes aspartate for glycine alpha-I-67
and that

[134301]causes cataracts and retinal detachment: evidence for
molecular heterogeneity

[134302]in the Wagner syndrome and the Stickler syndrome
(arthro-ophthalmopathy). Am.

[134303]J. Hum. Genet. 53: 55–61, 1993.

[134304]24002.

[134305]Fryns, J. P.; Kleczkowska, A.; Moerman, P.; Vandenberghe,

K.:

[134306]Hereditary hydronephrosis and the short arm of chromosome 6. (Letter) Hum.

[134307]Genet. 91: 514–515, 1993.

[134308]24003.

[134309]Groenen, P. M. A.; Garcia, E.; Debeer, P.; Devriendt, K.; Fryns,

[134310]J. P.; Van de Ven, W. J. M.: Structure, sequence, and chromosome

[134311]19 localization of human USF2 and its rearrangement in a patient with

[134312]multicystic renal dysplasia. Genomics 38: 141–148, 1996.

[134313]24004.

[134314]Groenen, P. M. A.; Vanderlinden, G.; Devriendt, K.; Fryns, J.-P;

[134315]Van de Ven, W. J. M.: Rearrangement of the human CDC5L gene by a

[134316]t(6;19)(p21;q13.1) in a patient with multicystic renal dysplasia. Genomics 49:

[134317]218–229, 1998.

[134318]24005.

[134319]Sepehrnia, B.; Kamboh, M. I.; Adams–Campbell, L. L.;
Nwankwo,

[134320]M.; Ferrell, R. E.: Genetic studies of human apolipoproteins. VII.

[134321]Population distribution of polymorphisms of apolipoproteins A–I, A–II,

[134322]A–IV, C–II, E, and H in Nigeria. Am. J. Hum. Genet. 43:
847–853,

[134323]1988.

[134324]24006.

[134325]Bost, F.; Bourguignon, J.; Martin, J.–P.; Sesboue, R.;
Thiberville,

[134326]L.; Diarra–Mehrpour, M.: Isolation and characterization of
the human

[134327]inter–alpha–trypsin inhibitor heavy–chain H1 gene. Europ.
J. Biochem. 218:

[134328]283–291, 1993.

[134329]24007.

[134330]Galizzi, J.–P.; Zuber, C. E.; Harada, N.; Gorman, D. M.;
Kastelein,

[134331]R.; Banchereau, J.; Howard, M.; Miyajima, A.: Molecular
cloning of

[134332]a cDNA encoding the human interleukin 4 receptor. Int. Immun. 2:

[134333]669–675, 1990.

[134334]24008.

[134335]Grimbacher, B.; Holland, S. M.; Puck, J. M.: The interleukin-4

[134336]receptor variant Q576R in hyper-IgE syndrome. (Letter) New Eng. J.

[134337]Med. 338: 1073–1074, 1998.

[134338]24009.

[134339]Idzerda, R. L.; March, C. J.; Mosley, B.; Lyman, S. D.; VandenBos,

[134340]T.; Gimpel, S. D.; Din, W. S.; Grabstein, K. H.; Widmer, M. B.; Park,

[134341]L. S.; Cosman, D.; Beckmann, M. P.: Human interleukin-4 receptor

[134342]confers biological responsiveness and defines a novel receptor superfamily. J.

[134343]Exp. Med. 171: 861–873, 1990.

[134344]24010.

[134345]Khurana Hershey, G. K.; Friedrich, M. F.; Esswein, L. A.;

Thomas,

[134346]M. L.; Chatila, T. A.: The association of atopy with a gain-of-function

[134347]mutation in the alpha subunit of the interleukin-4 receptor. New

[134348]Eng. J. Med. 337: 1720-1725, 1997.

[134349]24011.

[134350]Kruse, S.; Japha, T.; Tedner, M.; Sparholt, S. H.; Forster, J.;

[134351]Kuehr, J.; Deichmann, K. A.; Abello, A.: The polymorphisms S503P

[134352]and Q576R in the interleukin-4 receptor alpha gene are associated

[134353]with atopy and influence the signal transduction. Immunology 96:

[134354]365-371, 1999.

[134355]24012.

[134356]Mitsuyasu, H.; Izuhara, K.; Mao, X.-Q.; Gao, P.-S.; Arinobu, Y.;

[134357]Enomoto, T.; Kawai, M.; Sasaki, S.; Dake, Y.; Hamasaki, N.; Shirakawa,

[134358]T.; Hopkin, J. M.: Ile50val variant of IL4R-alpha upregu-

lates IgE

[134359]synthesis and associates with atopic asthma. (Letter) Nature Genet. 19:

[134360]119–120, 1998.

[134361]24013.

[134362]Corcoran, A. E.; Riddell, A.; Krooshoop, D.; Venkitaraman, A. R.

[134363]: Impaired immunoglobulin gene rearrangement in mice lacking the IL-7

[134364]receptor. Nature 391: 904–907, 1998.

[134365]24014.

[134366]Ober, C.; Leavitt, S. A.; Tsalenko, A.; Howard, T. D.; Hoki, D.

[134367]M.; Daniel, R.; Newman, D. L.; Wu, X.; Parry, R.; Lester, L. A.; Solway,

[134368]J.; Blumenthal, M.; King, R. A.; Xu, J.; Meyers, D. A.; Bleecker,

[134369]E. R.; Cox, N. J.: Variation in the interleukin 4–receptor alpha

[134370]gene confers susceptibility to asthma and atopy in ethnically diverse

[134371]populations. Am. J. Hum. Genet. 66: 517–526, 2000.

[134372]24015.

[134373]Ober, C.; Tsalenko, A.; Willadsen, S.; Newman, D.; Daniel, R.;

[134374]Wu, X.; Andal, J.; Hoki, D.; Schneider, D.; True, K.; Schou, C.; Parry,

[134375]R.; Cox, N.: Genome-wide screen for atopy susceptibility alleles

[134376]in the Hutterites. Clin. Exp. Allergy 29 (suppl. 4): 11–15, 1999.

[134377]24016.

[134378]Patuzzo, C.; Trabetti, E.; Malerba, G.; Martinati, L. C.; Boner,

[134379]A. L.; Pescolliderungg, L.; Zanoni, G.; Pignatti, P. F.: No linkage

[134380]or association of the IL-4R-alpha gene Q576R mutation with atopic

[134381]asthma in Italian families. (Letter) J. Med. Genet. 37: 382–384,

[134382]2000.

[134383]24017.

[134384]Pritchard, M. A.; Baker, E.; Whitmore, S. A.; Sutherland, G. R.;

[134385] Idzerda, R. L.; Park, L. S.; Cosman, D.; Jenkins, N. A.;

Gilbert,

[134386] D. J.; Copeland, N. G.; Beckmann, M. P.: The interleukin-4
receptor

[134387] gene (IL4R) maps to 16p11.2-16p12.1 in human and to
the distal region

[134388] of mouse chromosome 7. Genomics 10: 801-806, 1991.

[134389] 24018.

[134390] Suzuki, H.; Chung, F.; Palmer, E.; Sasaki, T.; Ohara, N.;
Taylor,

[134391] B. A.; Ohara, J.-I.: Gene mapping of mouse IL-4 receptor:
the loci

[134392] of interleukin 4 (IL-4) receptor gene and lymphocyte func-
tion associated

[134393] antigen 1 (LFA-1) gene are closely linked on chromosome
7. Immunogenetics 34:

[134394] 252-256, 1991.

[134395] 24019.

[134396] Zurawski, S. M.; Vega, F., Jr.; Huyghe, B.; Zurawski, G.: Re-
ceptors

[134397] for interleukin-13 and interleukin-4 are complex and
share a novel

[134398] component that functions in signal transduction. EMBO J.
12: 2663–2670,

[134399] 1993.

[134400] 24020.

[134401] Koshland, M. E.: The coming of age of the immunoglobulin J chain. Annu.

[134402] Rev. Immun. 3: 425–453, 1985.

[134403] 24021.

[134404] Max, E. E.; McBride, O. W.; Morton, C. C.; Robinson, M. A.:
Human

[134405] J chain gene: chromosomal localization and associated restriction

[134406] fragment length polymorphisms. Proc. Nat. Acad. Sci. 83:
5592–5596,

[134407] 1986.

[134408] 24022.

[134409] Diarra-Mehrpour, M.; Bourguignon, J.; Sesboue, R.; Mattei,
M.-G.;

[134410] Passage, E.; Salier, J.-P.; Martin, J.-P.: Human plasma inter-alpha-trypsin

[134411] inhibitor is encoded by four genes on three chromosomes.

Europ. J.

[134412]Biochem. 179: 147–154, 1989.

[134413]24023.

[134414]Enghild, J. J.; Thogersen, I. B.; Pizzo, S. V.; Salvesen, G.:

[134415]Analysis of inter- α -trypsin inhibitor and a novel
trypsin inhibitor,

[134416]pre- α -trypsin inhibitor, from human plasma:
polypeptide chain

[134417]stoichiometry and assembly by glycan. J. Biol. Chem. 264:
15975–15981,

[134418]1989.

[134419]24024.

[134420]Salier, J. P.; Diarra-Mehrpour, M.; Sesboue, R.; Bour-
guignon, J.;

[134421]Benarous, R.; Ohkubo, I.; Kurachi, S.; Kurachi, K.; Martin, J.
P.

[134422]: Isolation and characterization of cDNAs encoding the
heavy chain

[134423]of human inter- α -trypsin inhibitor (I- α -TI): un-
ambiguous evidence

[134424]for multipolypeptide chain structure of I- α -TI. Proc.

Nat. Acad.

[134425]Sci. 84: 8272–8276, 1987.

[134426]24025.

[134427]Salier, J. P.; Simon, D.; Rouet, P.; Raguenes, G.; Muscatelli,

[134428]F.; Gebhard, W.; Guenet, J. L.; Mattei, M. G.: Homologous
chromosomal

[134429]locations of the four genes for inter-alpha-inhibitor and
pre-alpha-inhibitor

[134430]family in human and mouse: assignment of the ancestral
gene for the

[134431]lipocalin superfamily. Genomics 14: 83–88, 1992.

[134432]24026.

[134433]Diarra-Mehrpour, M.; Sarafan, N.; Bourguignon, J.; Bonnet,
F.;

[134434]Bost, F.; Martin, J.-P.: Human inter-alpha-trypsin inhibitor
heavy

[134435]chain H3 gene: genomic organization, promoter analysis,
and gene linkage. J.

[134436]Biol. Chem. 273: 26809–26819, 1998.

[134437]24027.

[134438]Goodwin, R. G.; Friend, D.; Ziegler, S. F.; Jerzy, R.; Falk, B.

[134439]A.; Gimpel, S.; Cosman, D.; Dower, S. K.; March, C. J.; Narmen, A.

[134440]E.; Park, L. S.: Cloning of the human and murine interleukin-7 receptors:

[134441]demonstration of a soluble form and homology to a new receptor superfamily. Cell 60:

[134442]941-951, 1990.

[134443]24028.

[134444]Kondo, M.; Takeshita, T.; Higuchi, M.; Nakamura, M.; Sudo, T.;

[134445]Nishikawa, S.-I.; Sugamura, K.: Functional participation of the IL-2

[134446]receptor gamma chain in IL-7 receptor complexes. Science 263: 1453-1454,

[134447]1994.

[134448]24029.

[134449]Lai, S. Y.; Molden, J.; Goldsmith, M. A.: Shared gamma(c) subunit

[134450]within the human interleukin-7 receptor complex: a molecular basis

[134451]for the pathogenesis of X-linked severe combined immunodeficiency. J.

[134452]Clin. Invest. 99: 169–177, 1997.

[134453]24030.

[134454]Lynch, M.; Baker, E.; Park, L. S.; Sutherland, G. R.; Goodwin,

[134455]R. G.: The interleukin–7 receptor gene is at 5p13. Hum. Genet. 89:

[134456]566–568, 1992.

[134457]24031.

[134458]Maki, K.; Sunaga, S.; Komagata, Y.; Kodaira, Y.; Mabuchi, A.; Karasuyama,

[134459]H.; Yokomuro, K.; Miyazaki, J.; Ikuta, K.: Interleukin 7 receptor–deficient

[134460]mice lack gamma/delta T cells. Proc. Nat. Acad. Sci. 93: 7172–7177,

[134461]1996.

[134462]24032.

[134463]Noguchi, M.; Nakamura, Y.; Russell, S. M.; Ziegler, S. F.; Tsang,

[134464]M.; Cao, X.; Leonard, W. J.: Interleukin–2 receptor gamma chain:

[134465]a functional component of the interleukin–7 receptor. Sci-

ence 262:

[134466]1877–1880, 1993.

[134467]24033.

[134468]Peschon, J. J.; Morrissey, P. J.; Grabstein, K. H. et al.: Early

[134469]lymphocyte expansion is severely impaired in interleukin
7 receptor-deficient

[134470]mice. *J. Exp. Med.* 180: 1955–1960, 1994.

[134471]24034.

[134472]Puel, A.; Ziegler, S. F.; Buckley, R. H.; Leonard, W. J.: De-
fective

[134473]IL7R expression in T-B+NK+ severe combined immunod-
efficiency. *Nature*

[134474]*Genet.* 20: 394–397, 1998.

[134475]24035.

[134476]Mezei, M. M.; Mankodi, A.; Brais, B.; Marineau, C.; Thorn-
ton, C.

[134477]A.; Rouleau, G. A.; Karpati, G.: Minimal expansion of the
GCG repeat

[134478]in the PABP2 gene does not predispose to sporadic inclu-
sion body myositis. *Neurology* 52:

[134479]669–670, 1999.

[134480]24036.

[134481]Aleman, A.; Verhaar, H. J. J.; De Haan, E. H. F.; De Vries, W.

[134482]R.; Samson, M. M.; Drent, M. L.; Van Der Veen, E. A.;

Koppeschaar,

[134483]H. P. F.: Insulin-like growth factor-I and cognitive function
in

[134484]healthy older men. *J. Clin. Endocr. Metab.* 84: 471-475,
1999.

[134485]24037.

[134486]Baker, J.; Liu, J.-P.; Robertson, E. J.; Efstratiadis, A.: Role

[134487]of insulin-like growth factors in embryonic and postnatal
growth. *Cell* 75:

[134488]73-82, 1993.

[134489]24038.

[134490]Bowcock, A.; Sartorelli, V.: Polymorphism and mapping of
the IGF1

[134491]gene, and absence of association with stature among
African Pygmies. *Hum.*

[134492]*Genet.* 85: 349-354, 1990.

[134493]24039.

- [134494]Camacho–Hubner, C.; Woods, K. A.; Miraki–Moud, F.; Hindmarsh, P.
- [134495]C.; Clark, A. J.; Hansson, Y.; Johnston, A.; Baxter, R. C.; Savage,
- [134496]M. O.: Effects of recombinant human insulin–like growth factor I
- [134497](IGF–I) therapy on the growth hormone–IGF system of a patient with
- [134498]a partial IGF–I gene deletion. J. Clin. Endocr. Metab. 84: 1611–1616,
- [134499]1999.
- [134500]24040.
- [134501]Goddard, A. D.; Covello, R.; Luoh, S.–M.; Clackson, T.; Attie,
- [134502]K. M.; Gesundheit, N.; Rundle, A. C.; Wells, J. A.; Carlsson, L. M.
- [134503]S.: Mutations of the growth hormone receptor in children with idiopathic
- [134504]short stature. New Eng. J. Med. 333: 1093–1098, 1995.
- [134505]24041.
- [134506]Guler, H.–P.; Binz, K.; Eigenmann, E.; Jaggi, S.; Zimmermann, D.;

- [134507]Zapf, J.; Froesch, E. R.: Small stature and insulin-like growth factors:
- [134508]prolonged treatment of mini-poodles with recombinant human insulin-like
- [134509]growth factor I. Acta Endocr. 121: 456–464, 1989.
- [134510]24042.
- [134511]Schmale, H.; Ivell, R.; Breindl, M.; Darmer, D.; Richter, D.:
- [134512]The mutant vasopressin gene from diabetes insipidus (Brattleboro)
- [134513]rats is transcribed but the message is not efficiently translated. EMBO
- [134514]J. 3: 3289–3293, 1984.
- [134515]24043.
- [134516]Willcutts, M. D.; Felner, E.; White, P. C.: Autosomal recessive
- [134517]familial neurohypophyseal diabetes insipidus with continued secretion
- [134518]of mutant weakly active vasopressin. Hum. Molec. Genet. 8: 1303–1307,
- [134519]1999.
- [134520]24044.

[134521]Bichet, D. G.; Hendy, G. N.; Lonergan, M.; Arthus, M.-F.;
Ligier,

[134522]S.; Pausova, Z.; Kluge, R.; Zingg, H.; Saenger, P.; Oppen-
heimer, E.;

[134523]Hirsch, D. J.; Gilgenkrantz, S.; Salles, J.-P.; Oberle, I.; Man-
del,

[134524]J.-L.; Gregory, M. C.; Fujiwara, T. M.; Morgan, K.; Scriver, C. R.

[134525]: X-linked nephrogenic diabetes insipidus: from the ship Hopewell

[134526]to RFLP studies. Am. J. Hum. Genet. 51: 1089–1102, 1992.

[134527]24045.

[134528]Ten Bessel, R. W.; Peters, E. R.: Progressive hydronephrosis,

[134529]hydroureter, and dilatation of the bladder in siblings with congenital

[134530]nephrogenic diabetes insipidus. J. Pediatr. 77: 439–443, 1970.

[134531]24046.

[134532]Fajans, S. S.: Maturity-onset diabetes of the young (MODY). Diabetes

[134533]Metab. Rev. 5: 579–606, 1989.

[134534]24047.

[134535]Harrison, G. S.; Drabkin, H. A.; Kao, F.-T.; Hartz, J.; Hart, I.

[134536]M.; Chu, E. H. Y.; Wu, B. J.; Morimoto, R. I.: Chromosomal location

[134537] of human genes encoding major heat-shock protein
HSP70. Somat. Cell

[134538] Molec. Genet. 13: 119–130, 1987.

[134539] 24048.

[134540] Imai, Y.; Soda, M.; Hatakeyama, S.; Akagi, T.; Hashikawa,
T.; Nakayama,

[134541] K.; Takahashi, R.: CHIP is associated with Parkin, a gene
responsible

[134542] for familial Parkinson's disease, and enhances its ubiquitin
ligase

[134543] activity. Molec. Cell 10: 55–67, 2002.

[134544] 24049.

[134545] Laroia, G.; Cuesta, R.; Brewer, G.; Schneider, R. J.: Control

[134546] of mRNA decay by heat shock-ubiquitin-proteasome
pathway. Science 284:

[134547] 499–502, 1999.

[134548] 24050.

[134549] Milner, C. M.; Campbell, R. D.: Polymorphic analysis of the
three

[134550] MHC-linked HSP70 genes. Immunogenetics 36: 357–362,
1992.

[134551]24051.

[134552]Milner, C. M.; Campbell, R. D.: Structure and expression of
the

[134553]three MHC-linked HSP70 genes. Immunogenetics 32:
242-251, 1990.

[134554]24052.

[134555]Sargent, C. A.; Dunham, I.; Trowsdale, J.; Campbell, R. D.:
Human

[134556]major histocompatibility complex contains genes for the
major heat

[134557]shock protein HSP70. Proc. Nat. Acad. Sci. 86:
1968-1972, 1989.

[134558]24053.

[134559]Bonnycastle, L. L. C.; Yu, C.-E.; Hunt, C. R.; Trask, B. J.;
Clancy,

[134560]K. P.; Weber, J. L.; Patterson, D.; Schellenberg, G. D.:
Cloning,

[134561]sequencing, and mapping of the human chromosome 14
heat shock protein

[134562]gene (HSPA2). Genomics 23: 85-93, 1994.

[134563]24054.

[134564]Dix, D. J.; Allen, J. W.; Collins, B. W.; Mori, C.; Nakamura, N.;

[134565]Poorman–Allen, P.; Goulding, E. H.; Eddy, E. M.: Targeted gene disruption

[134566]of Hsp70–2 results in failed meiosis, germ cell apoptosis, and male

[134567]infertility. Proc. Nat. Acad. Sci. 93: 3264–3268, 1996.

[134568]24055.

[134569]Hunt, C. R.; Gasser, D. L.; Chaplin, D. D.; Pierce, J. C.; Kozak,

[134570]C. A.: Chromosomal localization of five murine HSP70 gene family

[134571]members: Hsp70–1, Hsp70–2, Hsp70–3, Hsc70t and Grp78. Genomics 16:

[134572]193–198, 1993.

[134573]24056.

[134574]Roux, A.–F.; Nguyen, V. T. T.; Squire, J. A.; Cox, D. W.: A heat

[134575]shock gene at 14q22: mapping and expression. Hum. Molec. Genet. 3:

[134576]1819–1822, 1994.

[134577]24057.

[134578]Wolpowitz, D.; Mason, T. B. A.; Dietrich, P.; Mendelsohn, M.;

[134579]Talmage, D. A.; Role, L. W.: Cysteine-rich domain isoforms of the

[134580]neuregulin-1 gene are required for maintenance of peripheral synapses. Neuron 25:

[134581]79-91, 2000.

[134582]24058.

[134583]Bolton, P.; Powell, J.; Rutter, M.; Buckle, V.; Yates, J. R. W.;

[134584]Ishikawa-Brush, Y.; Monaco, A. P.: Autism, mental retardation, multiple

[134585]exostoses and short stature in a female with 46,X,t(X;8)(p22.13;q22.1). Psychiat.

[134586]Genet. 5: 51-55, 1995.

[134587]24059.

[134588]David, G.; van der Schueren, B.; Marynen, P.; Cassiman, J.-J.;

[134589]van den Berghe, H.: Molecular cloning of amphiglycan, a novel integral

[134590]membrane heparan sulfate proteoglycan expressed by ep-

ithelial and

[134591]fibroblastic cells. J. Cell Biol. 118: 961–969, 1992.

[134592]24060.

[134593]Ishikawa–Brush, Y.; Powell, J. F.; Bolton, P.; Miller, A. P.;
Francis,

[134594]F.; Willard, H. F.; Lehrach, H.; Monaco, A. P.: Autism and
multiple

[134595]exostoses associated with an X;8 translocation occurring
within the

[134596]GRPR gene and 3–prime to the SDC2 gene. Hum. Molec.
Genet. 6: 1241–1250,

[134597]1997.

[134598]24061.

[134599]Lories, V.; De Broeck, H.; David, G.; Cassiman, J. J.; Van
den

[134600]Berghe, H.: Heparan sulfate proteoglycans of human lung
fibroblasts:

[134601]structural heterogeneity of the core proteins of the hy-
drophobic cell-associated

[134602]forms. J. Biol. Chem. 262: 854–859, 1987.

[134603]24062.

[134604] Marynen, P.; Zhang, J.; Cassiman, J.-J.; David, G.: The gene
for
[134605] the 48kDa and the 90kDa core proteins of cell surface-
associated heparan
[134606] sulphate proteoglycan of human lung fibroblast maps to
chromosome
[134607] 8q2. (Abstract) Cytogenet. Cell Genet. 51: 1040, 1989.
[134608] 24063.

[134609] Marynen, P.; Zhang, J.; Cassiman, J.-J.; Van den Berghe,
H.; David,
[134610] G.: Partial primary structure of the 48- and 90-kilodalton
core proteins
[134611] of cell surface-associated heparan sulfate proteoglycans
of lung fibroblasts:
[134612] prediction of an integral membrane domain and evidence
for multiple
[134613] distinct core proteins at the cell surface of human lung fi-
broblasts. J.
[134614] Biol. Chem. 264: 7017-7024, 1989.
[134615] 24064.

[134616] Spring, J.; Goldberger, O. A.; Jenkins, N. A.; Gilbert, D. J.;

- [134617] Copeland, N. G.; Bernfield, M.: Mapping of the syndecan genes in
the mouse: linkage with members of the Myc gene family.
Genomics 21:
597–601, 1994.
- [134620] 24065.
- [134621] Glaser, T.; Matthews, K. E.; Hudson, J. W.; Seth, P.; Hous-
man,
D. E.; Crerar, M. M.: Localization of the muscle, liver and
brain
glycogen phosphorylase genes on linkage maps of mouse
chromosomes
19, 12 and 2, respectively. Genomics 5: 510–521, 1989.
- [134625] 24066.
- [134626] Newgard, C. B.; Littman, D. R.; van Genderen, C.; Smith,
M.; Fletterick,
R. J.: Human brain glycogen phosphorylase: cloning, se-
quence analysis,
chromosomal mapping, tissue expression, and compari-
son with the human
liver and muscle isozymes. J. Biol. Chem. 263:
3850–3857, 1988.

[134630]24067.

[134631]Yasuda, K.; Espinosa, R., III; Davis, E. M.; Le Beau, M. M.;
Bell,

[134632]G. I.: Human somatostatin receptor genes: localization of
SSTR5 to

[134633]human chromosome 20p11.2. Genomics 17: 785-786,
1993.

[134634]24068.

[134635]Barclay, A. N.: Personal Communication. Oxford, England
1/9/1989.

[134636]24069.

[134637]Douglas, J.; Albertson, D. G.; Barclay, A. N.; Davis, M.;
Rabbitts,

[134638]P. H.: RFLP and mapping of human MOX-1 gene on chro-
mosome 3. Nucleic

[134639]Acids Res. 16: 9067 only, 1988.

[134640]24070.

[134641]Hoek, R. M.; Ruuls, S. R.; Murphy, C. A.; Wright, G. J.; God-
dard,

[134642]R.; Zurawski, S. M.; Blom, B.; Homola, M. E.; Streit, W. J.;
Brown,

[134643]M. H.; Barclay, A. N.; Sedgwick, J. D.: Down-regulation of the macrophage

[134644]lineage through interaction with OX2 (CD200). Science 290: 1768-1771,

[134645]2000.

[134646]24071.

[134647]McCaughan, G. W.; Clark, M. J.; Barclay, A. N.: Characterization

[134648]of the human homolog of the rat MRC OX-2 membrane glycoprotein. Immunogenetics 25:

[134649]329-335, 1987.

[134650]24072.

[134651]McCaughan, G. W.; Clark, M. J.; Hurst, J.; Grosveld, F.; Barclay,

[134652]A. N.: The gene for MRC OX-2 membrane glycoprotein is localized on

[134653]human chromosome 3. Immunogenetics 25: 133-135, 1987.

[134654]24073.

[134655]Wright, G. J.; Jones, M.; Puklavec, M. J.; Brown, M. H.; Barclay,

[134656]A. N.: The unusual distribution of the neuronal/lymphoid cell surface

[134657]CD200 (OX2) glycoprotein is conserved in humans. Immunology 102:

[134658]173–179, 2001.

[134659]24074.

[134660]Buijs, A.; Sherr, S.; van Baal, S.; van Bezouw, S.; van der Plas,

[134661]D.; Geurts van Kessel, A.; Riegman, P.; Lekanne Deprez, R.; Zwarthoff,

[134662]E.; Hagemeijer, A.; Grosveld, G.: Translocation (12;22)(p13;q11)

[134663]in myeloproliferative disorders results in fusion of the ETS-like

[134664]TEL gene on 12q13 to the MN1 gene on 22q11. Oncogene 10: 1511–1519,

[134665]1995.

[134666]24075.

[134667]Lekanne Deprez, R. H.; Groen, N. A.; van Biezen, N. A.; Hagemeijer,

[134668]A.; van Drunen, E.; Koper, J. W.; Avezaat, C. J. J.; Bootsma, D.;

[134669]Zwarthoff, E. C.: A t(4;22) in a meningioma points to the
localization
[134670]of a putative tumor-suppressor gene. Am. J. Hum. Genet.
48: 783-790,
[134671]1991.
[134672]24076.
[134673]Reche, P. A.; Soumelis, V.; Gorman, D. M.; Clifford, T.; Liu,
[134674]M.; Travis, M.; Zurawski, S. M.; Johnston, J.; Liu, Y.-J.;
Spits,
[134675]H.; de Waal Malefyt, R.; Kastelein, R. A.; Bazan, J. F.: Hu-
man thymic
[134676]stromal lymphopoietin preferentially stimulates myeloid
cells. J.
[134677]Immun. 167: 336-343, 2001.
[134678]24077.
[134679]Chrousos, G. P.: Personal Communication. Bethesda, Md.
10/21/2002.
[134680]24078.
[134681]Chrousos, G. P.; MacLusky, N. J.; Brandon, D. D.; Tomita,
M.; Renquist,
[134682]D. M.; Loriaux, D. L.; Lipsett, M. B.: Progesterone resis-

tance.In:

[134683]Chrousos, G. P.; Loriaux, D. L.; Lipsett, M. B.: Steroid Hormone

[134684]Resistance: Mechanisms and Clinical Aspects. New York: Plenum Press

[134685](pub.) 1986.

[134686]24079.

[134687]Keller, D. W.; Wiest, W. G.; Askin, F. B.; Johnson, L. W.; Strickler,

[134688]R. C.: Pseudocorpus luteum insufficiency: a local defect of progesterone

[134689]action on endometrial stroma. J. Clin. Endocr. Metab. 48: 127-132,

[134690]1979.

[134691]24080.

[134692]Daiger, S. P.; Reed, L.; Huang, S.-S.; Zeng, Y.-T.; Wang, T.;

[134693]Lo, W. H. Y.; Okano, Y.; Hase, Y.; Fukuda, Y.; Oura, T.; Tada, K.;

[134694]Woo, S. L. C.: Polymorphic DNA haplotypes at the phenylalanine hydroxylase

[134695](PAH) locus in Asian families with phenylketonuria (PKU).

Am. J.

[134696]Hum. Genet. 45: 319–324, 1989.

[134697]24081.

[134698]Desviat, L. R.; Perez, B.; De Lucca, M.; Cornejo, V.;

Schmidt,

[134699]B.; Ugarte, M.: Evidence in Latin America of recurrence of

V388M,

[134700]a phenylketonuria mutation with high in vitro residual ac-

tivity. Am.

[134701]J. Hum. Genet. 57: 337–342, 1995.

[134702]24082.

[134703]Desviat, L. R.; Perez, B.; Ugarte, M.: A new PKU mutation

associated

[134704]with haplotype 12. Hum. Molec. Genet. 1: 765–766, 1992.

[134705]24083.

[134706]Desviat, L. R.; Perez, B.; Ugarte, M.: Phenylketonuria in

Spanish

[134707]Gypsies: prevalence of the IVS10nt546 mutation on haplo-

type 34. Hum.

[134708]Mutat. 9: 66–68, 1997.

[134709]24084.

[134710]Dianzani, I.; Camaschella, C.; Saglio, G.; Ferrero, G. B.; Ramus,

[134711]S.; Ponzzone, A.; Cotton, R. G. H.: Molecular analysis of contiguous

[134712]exons of the phenylalanine hydroxylase gene: identification of a new

[134713]PKU mutation. J. Med. Genet. 30: 228–231, 1993.

[134714]24085.

[134715]Dianzani, I.; Devoto, M.; Camaschella, C.; Saglio, G.; Ferrero,

[134716]G. B.; Cerone, R.; Romano, C.; Romeo, G.; Giovannini, M.; Riva, E.;

[134717]Angeneydt, F.; Trefz, F. K.; Okano, Y.; Woo, S. L. C.: Haplotype

[134718]distribution and molecular defects at the phenylalanine hydroxylase

[134719]locus in Italy. Hum. Genet. 86: 69–72, 1990.

[134720]24086.

[134721]Dianzani, I.; Forrest, S. M.; Camaschella, C.; Saglio, G.; Ponzzone,

[134722]A.; Cotton, R. G. H.: Screening for mutations in the phenylalanine

[134723]hydroxylase gene from Italian patients with phenylketonuria by using
[134724]the chemical cleavage method: a new splice mutation. Am. J. Hum.
[134725]Genet. 48: 631–635, 1991.
[134726]24087.
[134727]DiLella, A. G.; Huang, W.-M.; Woo, S. L. C.: Screening for phenylketonuria
[134728]mutations by DNA amplification with the polymerase chain reaction. Lancet I:
[134729]497–499, 1988.
[134730]24088.
[134731]DiLella, A. G.; Kwok, S. C. M.; Ledley, F. D.; Marvit, J.; Woo, S. L. C.: Molecular structure and polymorphic map of the human phenylalanine
[134732]hydroxylase gene. Biochemistry 25: 743–749, 1986.
[134733]24089.
[134734]DiLella, A. G.; Marvit, J.; Brayton, K.; Woo, S. L. C.: An amino-acid
[134735]substitution involved in phenylketonuria is in linkage disequilibrium

[134737]with DNA haplotype 2. Nature 327: 333–336, 1987.

[134738]24090.

[134739]DiLella, A. G.; Marvit, J.; Guttler, F.; Woo, S. L. C.: Molecular
lar

[134740]genetics of phenylketonuria. (Abstract) 7th Int. Cong.
Hum. Genet.,

[134741]Berlin 665–666, 1986.

[134742]24091.

[134743]DiLella, A. G.; Marvit, J.; Lidsky, A. S.; Guttler, F.; Woo, S.

[134744]L. C.: Tight linkage between a splicing mutation and a
specific DNA

[134745]haplotype in phenylketonuria. Nature 322: 799–803,
1986.

[134746]24092.

[134747]Drogari, E.; Smith, I.; Beasley, M.; Lloyd, J. K.: Timing of

[134748]strict diet in relation to fetal damage in maternal
phenylketonuria:

[134749]an international collaborative study by the MRC/DHSS
phenylketonuria

[134750]register. Lancet II: 927–930, 1987.

[134751]24093.

[134752]Dworniczak, B.; Aulehla-Scholz, C.; Horst, J.: Phenylalanine

[134753]hydroxylase gene: silent mutation uncovers evolutionary origin of

[134754]different alleles. Clin. Genet. 38: 270-273, 1990.

[134755]24094.

[134756]Dworniczak, B.; Aulehla-Scholz, C.; Horst, J.: Phenylketonuria:

[134757]detection of a frequent haplotype 4 allele mutation. Hum. Genet. 84:

[134758]95-96, 1989.

[134759]24095.

[134760]Dworniczak, B.; Aulehla-Scholz, C.; Kalaydjieva, L.; Bartholome,

[134761]K.; Grudza, K.; Horst, J.: Aberrant splicing of phenylalanine hydroxylase

[134762]mRNA: the major cause for phenylketonuria in parts of southern Europe. Genomics 11:

[134763]242-246, 1991.

[134764]24096.

[134765]Dworniczak, B.; Grudda, K.; Stumper, J.; Bartholome, K.;
Aulehla-Scholz,

[134766]C.; Horst, J.: Phenylalanine hydroxylase gene: novel mis-
sense mutation

[134767]in exon 7 causing severe phenylketonuria. Genomics 9:
193-199, 1991.

[134768]24097.

[134769]Dworniczak, B.; Kalaydjieva, L.; Aulehla-Scholz, C.; Ullrich,

[134770]K.; Kremensky, I.; Radeva, B.; Horst, J.: Recurrent non-
sense mutation

[134771]in exon 7 of the phenylalanine hydroxylase gene. Hum.
Genet. 87:

[134772]731-733, 1991.

[134773]24098.

[134774]Dworniczak, B.; Kalaydjieva, L.; Pankoke, S.; Aulehla-
Scholz,

[134775]C.; Allen, G.; Horst, J.: Analysis of exon 7 of the human
phenylalanine

[134776]hydroxylase gene: a mutation hot spot? Hum. Mutat. 1:
138-146, 1992.

[134777]24099.

[134778]Economou-Petersen, E.; Henriksen, K. F.; Guldberg, P.;
Guttler,

[134779]F.: Molecular basis for nonphenylketonuria hyperphenyl-
alaninemia. Genomics 14:

[134780]1–5, 1992.

[134781]24100.

[134782]Eigel, A.; Dworniczak, B.; Kalaydjieva, L.; Horst, J.: A
frameshift

[134783]mutation in exon 2 of the phenylalanine hydroxylase gene
linked to

[134784]RFLP haplotype 1. Hum. Genet. 87: 739–741, 1991.

[134785]24101.

[134786]Eiken, H. G.; Knappskog, P. M.; Apold, J.; Flatmark, T.: PKU

[134787]mutation G46S is associated with increased aggregation
and degradation

[134788]of the phenylalanine hydroxylase enzyme. Hum. Mutat. 7:
228–238,

[134789]1996.

[134790]24102.

[134791]Eiken, H. G.; Knappskog, P. M.; Apold, J.; Skjelkvale, L.;
Boman,

- [134792]H.: A de novo phenylketonuria mutation: ATG (met) to ATA (ile) in the start codon of the phenylalanine hydroxylase gene. Hum. Mutat. 1: 388–391, 1992.
- [134793]24103.
- [134794]Eiken, H. G.; Knappskog, P. M.; Boman, H.; Thune, K. S.; Kaada, G.; Motzfeldt, K.; Apold, J.: Relative frequency, heterogeneity and geographic clustering of PKU mutations in Norway. Europ. J. Hum. Genet. 4: 205–213, 1996.
- [134795]24104.
- [134796]Eiken, H. G.; Knappskog, P. M.; Guldberg, P.; Apold, J.: DGGE analysis as supplement to SSCP analysis of the phenylalanine hydroxylase gene: detection of eight (one de novo, seven inherited) of nine remaining Norwegian PKU mutations. Hum. Mutat. 8: 19–22, 1996.

[134805]24105.

[134806]Eiken, H. G.; Stangeland, K.; Skjelkvale, L.; Knappskog, P.
M.;

[134807]Boman, H.; Apol, J.: PKU mutations R408Q and F299C in
Norway: haplotype

[134808]associations, geographic distributions and phenotype
characteristics. Hum.

[134809]Genet. 88: 608–612, 1992.

[134810]24106.

[134811]Dautigny, A.; Bernier, I.; Colombani, J.; Jolles, P.: Human
platelets

[134812]as a source of HL-A antigens: a study of various solubi-
lization techniques. Biochimie 57:

[134813]1197–1201, 1975.

[134814]24107.

[134815]Moser, A. B.; Kreiter, N.; Bezman, L.; Lu, S.; Raymond, G.
V.;

[134816]Naidu, S.; Moser, H. W.: Plasma very long chain fatty acids
in 3,000

[134817]peroxisome disease patients and 29,000 controls. Ann.
Neurol. 45:

[134818]100–110, 1999.

[134819]24108.

[134820]Hoskins, J. A.; Jack, G.; Wade, H. E.; Peiris, R. J. D.; Wright,

[134821]E. C.; Starr, D. J. T.; Stern, J.: Enzymatic control of phenyl-
alanine

[134822]intake in phenylketonuria. *Lancet* I: 392–394, 1980.

[134823]

[134824]100. Howell, R. R.; Stevenson, R. E.: The offspring of
phenylketonuric

[134825]women. *Soc. Biol.* 18 (suppl.): S19–S29, 1971.

[134826]

[134827]101. Hsia, D. Y.–Y.: Phenylketonuria and its variants. *Prog.*
Med.

[134828]*Genet.* 7: 29–68, 1970.

[134829]

[134830]102. Huang, S.; Ren, Z.; Zeng, Y.; Woo, S. L. C.: PAH 399
GTA(val)–to–GTT(val),

[134831]a new silent mutation found in the Chinese. *Hum. Genet.*
86: 305–306,

[134832]1991.

[134833]

[134834]103. Huang, S.-Z.; Zhou, X.-D.; Ren, Z.-R.; Zeng, Y.-T.;
Woo, S. L.

[134835]C.: Prenatal detection of an arg-to-ter mutation at codon
111 of

[134836]the PAH gene using DNA amplification. Prenatal Diag. 10:
289-293,

[134837]1990.

[134838]

[134839]104. Huntley, C. C.; Stevenson, R. E.: Maternal phenylke-
tonuria:

[134840]course of two pregnancies. Obstet. Gynec. 34: 694-700,
1969.

[134841]

[134842]105. Hutchesson, A. C. J.; Hall, S. K.; Preece, M. A.; Green,
A.:

[134843]Screening for tyrosinaemia type I. Arch. Child. Dis. 74:
F191-F194,

[134844]1996.

[134845]

[134846]106. Ivaschenko, T.; Baranov, V. S.: Rapid and efficient

PCR/Styl

[134847]test for identification of common mutation R408W in
phenylketonuria

[134848]patients. J. Med. Genet. 30: 153–154, 1993.

[134849]

[134850]107. Jaruzelska, J.; Henriksen, K. F.; Guttler, F.; Riess, O.;
Borski,

[134851]K.; Blin, N.; Slomski, R.: The codon 408 mutation associ-
ated with

[134852]haplotype 2 is predominant in Polish families with
phenylketonuria. Hum.

[134853]Genet. 86: 247–250, 1991.

[134854]

[134855]108. Jaruzelska, J.; Melle, D.; Matuszak, R.; Borski, K.;
Munnich,

[134856]A.: A new 15 bp deletion in exon 11 of the phenylalanine
hydroxylase

[134857]gene in phenylketonuria. Hum. Molec. Genet. 1: 763–764,
1992.

[134858]

[134859]109. Jervis, G. A.: Phenylpyruvic oligophrenia deficiency of

phenylalanine-oxidizing

[134860]system. Proc. Soc. Exper. Biol. Med. 82: 514-515, 1953.

[134861]

[134862]110. Jervis, G. A.: Studies on phenylpyruvic oligophrenia:
the position

[134863]of the metabolic error. J. Biol. Chem. 169: 651-656, 1947.

[134864]

[134865]111. John, S. W. M.; Rozen, R.; Laframboise, R.; Laberge,
C.; Sriver,

[134866]C. R.: Five mutations at the PAH locus account for almost
90% of

[134867]PKU mutations in French-Canadians from eastern Quebec.
Hum. Mutat. 1:

[134868]72-74, 1992.

[134869]

[134870]112. John, S. W. M.; Rozen, R.; Laframboise, R.; Laberge,
C.; Sriver,

[134871]C. R.: Novel PKU mutation on haplotype 2 in French-
Canadians. Am.

[134872]J. Hum. Genet. 45: 905-909, 1989.

[134873]

- [134874]113. John, S. W. M.; Rozen, R.; Scriver, C. R.; Laframboise, R.; Laberge, C.: Recurrent mutation, gene conversion, or recombination at the human phenylalanine hydroxylase locus: evidence in French-Canadians and a catalog of mutations. Am. J. Hum. Genet. 46: 970-974, 1990.
- [134878]
- [134879]114. John, S. W. M.; Scriver, C. R.; Laframboise, R.; Rozen, R.: In vitro and in vivo correlations for I65T and M1V mutations at the phenylalanine hydroxylase locus. Hum. Mutat. 1: 147-153, 1992.
- [134882]
- [134883]115. Justice, M. J.; Siracusa, L. D.; Gilbert, D. J.; Heisterkamp, N.; Groffen, J.; Chada, K.; Silan, C. M.; Copeland, N. G.; Jenkins, N. A.: A genetic linkage map of mouse chromosome 10: localization

[134886] of eighteen molecular markers using a single interspecific backcross. Genetics 125:

[134887] 855–866, 1990.

[134888]

[134889] 116. Kalanin, J.; Takarada, Y.; Kagawa, S.; Yamashita, K.; Ohtsuka,

[134890] N.; Matsuoka, A.: Gypsy phenylketonuria: a point mutation of the

[134891] phenylalanine hydroxylase gene in gypsy families from Slovakia. Am.

[134892] J. Med. Genet. 49: 235–239, 1994.

[134893]

[134894] 117. Kalaydjieva, L.; Dworniczak, B.; Aulehla-Scholz, C.; Devoto,

[134895] M.; Romeo, G.; Sturhmann, M.; Kucinskas, V.; Yurgelyavicius, V.; Horst,

[134896] J.: Silent mutations in the phenylalanine hydroxylase gene as an

[134897] aid to the diagnosis of phenylketonuria. J. Med. Genet. 28: 686–690,

[134898] 1991.

[134899]

[134900]118. Kalaydjieva, L.; Dworniczak, B.; Aulehla-Scholz, C.;
Kremensky,

[134901]I.; Bronzova, J.; Eigel, A.; Horst, J.: Classical phenylke-
tonuria

[134902]in Bulgaria: RFLP haplotypes and frequency of the major
mutations. J.

[134903]Med. Genet. 27: 742-745, 1990.

[134904]

[134905]119. Kalaydjieva, L.; Dworniczak, B.; Kucinskas, V.;
Yurgeliavicius,

[134906]V.; Kunert, E.; Horst, J.: Geographical distribution gradi-
ents of

[134907]the major PKU mutations and the linked haplotypes. Hum.
Genet. 86:

[134908]411-413, 1991.

[134909]

[134910]120. Kamaryt, J.; Mrskos, A.; Podhradská, O.; Kolcova, V.;
Cabalska,

[134911]B.; Duzynska, N.; Borzymowska, J.: PKU locus: genetic
linkage with

[134912]human amylase (AMY) loci and assignment to linkage group 1. Hum.

[134913]Genet. 43: 205–210, 1978.

[134914]

[134915]121. Kaufman, S.: Phenylketonuria: biochemical mechanisms. Adv.

[134916]Neurochem. 2: 1–132, 1976.

[134917]

[134918]122. Kaufman, S.: A model of human phenylalanine metabolism in normal

[134919]subjects and in phenylketonuric patients. Proc. Nat. Acad. Sci. 96:

[134920]3160–3164, 1999.

[134921]

[134922]123. Kaufman, S.: Differential diagnosis of variant forms of hyperphenylalaninemia. Pediatrics 65:

[134923]840–842, 1980.

[134924]

[134925]124. Kaufman, S.; Fisher, D. B.: Purification and some physical properties

[134926]of phenylalanine hydroxylase from rat liver. J. Biol. Chem.
245:

[134927]4745–4750, 1970.

[134928]

[134929]125. Kaufman, S.; Max, E. E.; Kang, E. S.: Phenylalanine
hydroxylase

[134930]activity in liver biopsies from hyperphenylalaninemia het-
erozygotes:

[134931]deviation from proportionality with gene dosage. Pediat.
Res. 9:

[134932]632–634, 1975.

[134933]

[134934]126. Kawashima, H.; Kawano, M.; Masaki, A.; Sato, T.:
Three cases

[134935]of untreated classical PKU: a report on cataracts and brain
calcification. Am.

[134936]J. Med. Genet. 29: 89–93, 1988.

[134937]

[134938]127. Kerr, G. R.; Chamove, A. S.; Harlow, H. F.; Waisman,
H. A.:

[134939]'Fetal PKU': the effect of maternal hyperphenylalaninemia

during pregnancy

[134940]in the rhesus monkey (*Macaca mulatta*). *Pediatrics* 42:
27–36, 1968.

[134941]

[134942]128. Kidd, K. K.: Phenylketonuria: population genetics of a
disease. *Nature* 327:

[134943]282–283, 1987.

[134944]

[134945]129. Kleiman, S.; Vanagaite, L.; Bernstein, J.; Schwartz, G.;
Brand,

[134946]N.; Elitzur, A.; Woo, S. L. C.; Shiloh, Y.: Phenylketonuria:
variable

[134947]phenotypic outcomes of the R261Q mutation and mater-
nal PKU in the

[134948]offspring of a healthy homozygote. *J. Med. Genet.* 30:
284–288, 1993.

[134949]

[134950]130. Knapp, A.; Tintschewa, R.; Scheibe, E.; Schiebe, E.;
Jager, B.;

[134951]Biebler, K. E.: The genetic linkage between the PKU locus
and the

[134952]loci for amylase–1, amylase–2, Fy, PGM–1, and Rh and the

question

[134953] of assignment of the PKU locus to chromosome no. 1.

Hum. Genet. 60:

[134954] 122–125, 1982.

[134955]

[134956] 131. Knappskog, P. M.; Eiken, H. G.; Martinez, A.; Flatmark, T.; Apold,

[134957] J.: The PKU mutation S349P causes complete loss of catalytic activity

[134958] in the recombinant phenylalanine hydroxylase enzyme.

Hum. Genet. 95:

[134959] 171–173, 1995.

[134960]

[134961] 132. Koch, R.; Moats, R.; Guttler, F.; Guldberg, P.; Nelson, M., Jr.

[134962] Blood–brain phenylalanine relationships in persons with phenylketonuria. Pediatrics 106:

[134963] 1093–1096, 2000.

[134964]

[134965] 133. Komrower, G. M.; Sardharwalla, I. B.; Coutts, J. M. J.; Ingham,

[134966]D.: Management of maternal phenylketonuria: an emerging clinical

[134967]problem. Brit. Med. J. I: 1383–1387, 1979.

[134968]

[134969]134. Konecki, D. S.; Lichter-Konecki, U.: The phenylketonuria locus:

[134970]current knowledge about alleles and mutations of the phenylalanine

[134971]hydroxylase gene in various populations. Hum. Genet. 87: 377–388,

[134972]1991.

[134973]

[134974]135. Konecki, D. S.; Schlotter, M.; Trefz, F. K.; Lichter-Konecki,

[134975]U.: The identification of two mis-sense mutations at the PAH gene

[134976]locus in a Turkish patient with phenylketonuria. Hum. Genet. 87:

[134977]389–393, 1991.

[134978]

[134979]136. Kwok, S. C. M.; Ledley, F. D.; DiLella, A. G.; Robson,

K. J.

[134980]H.; Woo, S. L. C.: Nucleotide sequence of a full-length complementary

[134981]DNA clone and amino acid sequence of human phenylalanine hydroxylase. *Biochemistry* 24:

[134982]556–561, 1985.

[134983]

[134984]137. Labrune, P.; Melle, D.; Rey, F.; Berthelon, M.; Cailaud, C.;

[134985]Rey, J.; Munnich, A.; Lyonnet, S.: Single-strand conformation polymorphism

[134986]for detection of mutations and base substitutions in phenylketonuria. *Am.*

[134987]*J. Hum. Genet.* 48: 1115–1120, 1991.

[134988]

[134989]138. Lasala, J. M.; Coscia, C. J.: Accumulation of a tetrahydroisoquinoline

[134990]in phenylketonuria. *Science* 203: 283–284, 1979.

[134991]

[134992]139. Leandro, P.; Rivera, I.; Ribeiro, V.; de Almeida, I. T.; da Silva,

[134993]C.; Lechner, M. C.: Mutation analysis of phenylketonuria in
South

[134994]and Central Portugal: prevalence of V388M mutation.
Hum. Mutat. 6:

[134995]192–194, 1995.

[134996]

[134997]140. Ledbetter, S. A.; Ledbetter, D. H.; Ledley, F. D.; Woo,
S.:

[134998]Localization of phenylalanine hydroxylase (PAH) and al-
pha-1 antitrypsin

[134999](AAT) loci in mouse genome by synteny and in situ hy-
bridization. (Abstract) Am.

[135000]J. Hum. Genet. 41: A173 only, 1987.

[135001]

[135002]141. Ledley, F. D.; DiLella, A. G.; Woo, S. L. C.: Molecular
biology

[135003]of phenylalanine hydroxylase and phenylketonuria. Trends
Genet. 1:

[135004]309–313, 1985.

[135005]

[135006]142. Ledley, F. D.; Grenett, H. E.; DiLella, A. G.; Kwok, S. C.

M.;

[135007]Woo, S. L. C.: Gene transfer and expression of human
phenylalanine

[135008]hydroxylase. Science 228: 77-79, 1985.

[135009]

[135010]143. Ledley, F. D.; Grenett, H. E.; McGinnis-Shelnutt, M.;
Woo, S.

[135011]L. C.: Retroviral-mediated gene transfer of human pheny-
lalanine hydroxylase

[135012]into NIH 3T3 and hepatoma cells. Proc. Nat. Acad. Sci. 83:
409-413,

[135013]1986.

[135014]

[135015]144. Ledley, F. D.; Koch, R.; Jew, K.; Beaudet, A.; O'Brien,
W. E.;

[135016]Bartos, D. P.; Woo, S. L. C.: Phenylalanine hydroxylase ex-
pression

[135017]in liver of a fetus with phenylketonuria. J. Pediat. 113:
463-468,

[135018]1988.

[135019]

[135020]145. Ledley, F. D.; Levy, H. L.; Woo, S. L. C.: Molecular analysis

[135021]of the inheritance of phenylketonuria and mild hyperphenylalaninemia

[135022]in families with both disorders. New Eng. J. Med. 314: 1276–1280,

[135023]1986.

[135024]

[135025]146. Lenke, R. R.; Levy, H. L.: Maternal phenylketonuria and hyperphenylalaninemia:

[135026]an international survey of the outcome of untreated and treated pregnancies. New

[135027]Eng. J. Med. 303: 1202–1208, 1980.

[135028]

[135029]147. Leuzzi, V.; Bianchi, M. C.; Tosetti, M.; Carducci, C.; Carducci,

[135030]C.; Antonozzi, I.: Clinical significance of brain phenylalanine concentration

[135031]assessed by in vivo proton magnetic resonance spectroscopy in phenylketonuria. J.

[135032]Inherit. Metab. Dis. 23: 563–570, 2000.

[135033]

[135034]148. Levy, H. L.: Molecular genetics of phenylketonuria and its implications.

[135035](Editorial) Am. J. Hum. Genet. 45: 667–670, 1989.

[135036]

[135037]149. Levy, H. L.; Guldberg, P.; Guttler, F.; Hanley, W. B.; Matalon,

[135038]R.; Rouse, B. M.; Trefz, F.; Azen, C.; Allred, E. N.; de la Cruz,

[135039]F.; Koch, R.: Congenital heart disease in maternal phenylketonuria:

[135040]report from the Maternal PKU Collaborative Study. Pediat. Res. 49:

[135041]636–642, 2001.

[135042]

[135043]150. Levy, H. L.; Karolkewicz, V.; Houghton, S. A.; Mac-Cready, R.

[135044]A.: Screening the 'normal' population in Massachusetts for phenylketonuria. New

[135045]Eng. J. Med. 282: 1455–1458, 1970.

[135046]

[135047]151. Levy, H. L.; Lobbregt, D.; Barnes, P. D.; Poussaint, T.
Y.:

[135048]Maternal phenylketonuria: magnetic resonance imaging of
the brain

[135049]in offspring. J. Pediat. 128: 770–775, 1996.

[135050]

[135051]152. Levy, H. L.; Waisbren, S. E.: Effects of untreated ma-
ternal

[135052]phenylketonuria and hyperphenylalaninemia on the fetus.
New Eng.

[135053]J. Med. 309: 1269–1274, 1983.

[135054]

[135055]153. Li, J.; Eisensmith, R. C.; Wang, T.; Lo, W. H. Y.;
Huang, S.-Z.;

[135056]Zeng, Y.-T.; Yuan, L.-F.; Liu, S.-R.; Woo, S. L. C.: Identifi-
cation

[135057]of three novel missense PKU mutations among Chinese.
Genomics 13:

[135058]894–895, 1992.

[135059]

[135060]154. Lichter-Konecki, U.; Konecki, D. S.; DiLella, A. G.;

Brayton,

[135061]K.; Marvit, J.; Hahn, T. M.; Trefz, F. K.; Woo, S. L. C.:

Phenylalanine

[135062]hydroxylase deficiency caused by a single base substitution in an

[135063]exon of the human phenylalanine hydroxylase gene. Biochemistry 27:

[135064]2881-2885, 1988.

[135065]

[135066]155. Lidsky, A. S.; Guttler, F.; Woo, S. L. C.: Prenatal diagnosis

[135067]of classic phenylketonuria by DNA analysis. Lancet I: 549-551, 1985.

[135068]

[135069]156. Lidsky, A. S.; Law, M. L.; Morse, H. G.; Kao, F.-T.; Rabin, M.;

[135070]Ruddle, F. H.; Woo, S. L. C.: Regional mapping of the phenylalanine

[135071]hydroxylase gene and the phenylketonuria locus in the human genome. Proc.

[135072]Nat. Acad. Sci. 82: 6221-6225, 1985.

[135073]

[135074]157. Lidsky, A. S.; Ledley, F. D.; DiLella, A. G.; Kwok, S. C. M.;

[135075]Daiger, S. P.; Robson, K. J. H.; Woo, S. L. C.: Extensive restriction

[135076]site polymorphism at the human phenylalanine hydroxylase locus and

[135077]application in prenatal diagnosis of phenylketonuria. Am. J. Hum.

[135078]Genet. 37: 619–634, 1985.

[135079]

[135080]158. Lidsky, A. S.; Robson, K. J. H.; Thirumalachary, C.; Barker,

[135081]P. E.; Ruddle, F. H.; Woo, S. L. C.: The PKU locus in man is on chromosome

[135082]12. Am. J. Hum. Genet. 36: 527–533, 1984.

[135083]

[135084]159. Lin, C.-H.; Hsiao, K.-J.; Tsai, T.-F.; Chao, H.-K.; Su, T.-S.

[135085]: Identification of a missense phenylketonuria mutation at codon 408

[135086]in Chinese. Hum. Genet. 89: 593–596, 1992.

[135087]

[135088]160. Liu, S. R.; Zuo, Q. H.: Newborn screening for
phenylketonuria

[135089]in eleven districts. Chinese Med. J. 99: 113–118, 1986.

[135090]

[135091]161. Lyonnet, S.; Caillaud, C.; Rey, F.; Berthelon, M.;
Frezal, J.;

[135092]Rey, J.; Munnich, A.: Molecular genetics of phenylke-
tonuria in Mediterranean

[135093]countries: a mutation associated with partial phenylala-
nine hydroxylase

[135094]deficiency. Am. J. Hum. Genet. 44: 511–517, 1989.

[135095]

[135096]162. Lyonnet, S.; Melle, D.; de Braekeleer, M.; Lafram-
boise, R.; Rey,

[135097]F.; John, S. W. M.; Berthelon, M.; Berthelot, J.; Journal, H.;
Le

[135098]Marec, B.; Parent, P.; de Parscau, L.; Saudubray, J.-M.;
Rozen, R.;

[135099]Rey, J.; Munnich, A.; Scriver, C. R.: Time and space clusters
of

[135100]the French-Canadian M1V phenylketonuria mutation in
France. Am. J.

[135101]Hum. Genet. 51: 191-196, 1992.

[135102]

[135103]163. Marvit, J.; DiLella, A. G.; Brayton, K.; Ledley, F. D.;
Robson,

[135104]K. J. H.; Woo, S. L. C.: GT to AT transition at a splice donor
site

[135105]causes skipping of the preceding exon in phenylketonuria.
Nucleic

[135106]Acids Res. 15: 5613-5628, 1987.

[135107]

[135108]164. Matalon, R.; Justice, P.; Deanching, M. N.: Phenylala-
nine hydroxylase

[135109]in human placenta: novel system for study of phenylke-
tonuria. (Letter) Lancet I:

[135110]853-854, 1977.

[135111]

[135112]165. McDonald, J. D.; Bode, V. C.; Dove, W. F.; Shedlovsky,
A.: Pah(hph-5):

[135113]a mouse mutant deficient in phenylalanine hydroxylase.

Proc. Nat.

[135114] Acad. Sci. 87: 1965–1967, 1990.

[135115]

[135116] 166. McDonald, J. D.; Charlton, C. K.: Characterization of mutations

[135117] at the mouse phenylalanine hydroxylase locus. Genomics 39: 402–405,

[135118] 1997.

[135119]

[135120] 167. Melle, D.; Verelst, P.; Rey, F.; Berthelon, M.; Francois, B.;

[135121] Munnich, A.; Lyonnet, S.: Two distinct mutations at a single BamHI

[135122] site in phenylketonuria. J. Med. Genet. 28: 38–40, 1991.

[135123]

[135124] 168. Menkes, J. H.; Aeberhard, E.: Maternal phenylketonuria. J.

[135125] Pediat. 74: 924–931, 1969.

[135126]

[135127] 169. Murphey, R. M.: Phenylketonuria (PKU) and the single

gene: an

[135128]old story retold. Behav. Genet. 13: 141–157, 1983.

[135129]

[135130]170. Nowacki, P.; Byck, S.; Prevost, L.; Scriver, C. R.: The
PAH

[135131]mutation analysis consortium database: update 1996. Nu-
cleic Acids

[135132]Res. 25: 139–142, 1997.

[135133]

[135134]171. Nyhan, W. L.: Fifty years ago: Asbjorn Folling and
phenylketonuria. Trends

[135135]Biochem. Sci. 9: 71–72, 1984.

[135136]

[135137]172. O'Connell, P.; Leppert, M.; Hoff, M.; Kumlin, E.;
Thomas, W.;

[135138]Cai, G.; Law, M.; White, R.: A linkage map for human chro-
mosome 12.

[135139](Abstract) Am. J. Hum. Genet. 37: A169 only, 1985.

[135140]

[135141]173. O'Flynn, M. E.; Tillman, P.; Hsia, D. Y.-Y.: Hyper-

phenylalaninemia

[135142]without phenylketonuria. Am. J. Dis. Child. 113: 22-30,
1967.

[135143]

[135144]174. Okano, Y.; Asada, M.; Kang, Y.; Nishi, Y.; Hase, Y.;
Oura, T.;

[135145]Isshiki, G.: Molecular characterization of phenylketonuria
in Japanese

[135146]patients. Hum. Genet. 103: 613-618, 1998.

[135147]

[135148]175. Okano, Y.; Eisensmith, R. C.; Dasovich, M.; Wang, T.;
Guttler,

[135149]F.; Woo, S. L. C.: A prevalent missense mutation in North-
ern Europe

[135150]associated with hyperphenylalaninaemia. Europ. J. Pediat.
150: 347-352,

[135151]1991.

[135152]

[135153]176. Okano, Y.; Wang, T.; Eisensmith, R. C.; Guttler, F.;
Woo, S.

[135154]L. C.: Recurrent mutation in the human phenylalanine hy-

droxylase

[135155]gene. Am. J. Hum. Genet. 46: 919–924, 1990.

[135156]

[135157]177. Okano, Y.; Wang, T.; Eisensmith, R. C.; Longhi, R.;
Riva, E.;

[135158]Giovannini, M.; Cerone, R.; Romano, C.; Woo, S. L. C.:
Phenylketonuria

[135159]missense mutations in the Mediterranean. Genomics 9:
96–103, 1991.

[135160]

[135161]178. Okano, Y.; Wang, T.; Eisensmith, R. C.; Steinmann, B.;
Gitzelmann,

[135162]R.; Woo, S. L. C.: Missense mutations associated with RFLP
haplotypes

[135163]1 and 4 of the human phenylalanine hydroxylase gene.
Am. J. Hum.

[135164]Genet. 46: 18–25, 1990.

[135165]

[135166]179. Okano, Y.; Wang, T.; Eisensmith, R. C.; Woo, S. L. C.:
PKU mutations

[135167]among Caucasians. (Abstract) Am. J. Hum. Genet. 45:

A211 only, 1989.

[135168]

[135169]180. Paine, R. S.: The variability in manifestations of untreated

[135170]patients with phenylketonuria (phenylpyruvic aciduria).
Pediatrics 20:

[135171]290-302, 1957.

[135172]

[135173]181. Paul, T. D.; Brandt, I. K.; Elsas, L. J.; Jackson, C. E.;
Mamunes,

[135174]P.; Nance, C. S.; Nance, W. E.: Phenylketonuria heterozygote detection

[135175]in families with affected children. Am. J. Hum. Genet. 30:
293-301,

[135176]1978.

[135177]

[135178]182. Paul, T. D.; Brandt, I. K.; Elsas, L. J.; Jackson, C. E.;
Nance,

[135179]C. S.; Nance, W. E.: Linkage analysis using heterozygote detection

[135180]in phenylketonuria. Clin. Genet. 16: 217-232, 1979.

[135181]

[135182]183. Perez, B.; Desviat, L. R.; Die, M.; Cornejo, V.;
Chamoles, N.

[135183]A.; Nicolini, H.; Ugarte, M.: Presence of the Mediterranean
PKU mutation

[135184]IVS10 in Latin America. Hum. Molec. Genet. 2:
1289–1290, 1993.

[135185]

[135186]184. Perez, B.; Desviat, L. R.; Die, M.; Ugarte, M.: Mutation
analysis

[135187]of phenylketonuria in Spain: prevalence of two Mediter-
ranean mutations. Hum.

[135188]Genet. 89: 341–342, 1992.

[135189]

[135190]185. Perry, T. L.; Hansen, S.; Tischler, B.; Bunting, R.; Dia-
mond,

[135191]S.: Glutamine depletion in phenylketonuria: possible cause
of the

[135192]mental defect. New Eng. J. Med. 282: 761–766, 1970.

[135193]

[135194]186. Pitt, D. B.; O'Day, J.: Phenylketonuria does not cause

cataracts. Europ.

[135195]J. Pediat. 150: 661–664, 1991.

[135196]

[135197]187. Ramus, S. J.; Forrest, S. M.; Cotton, R. G. H.: Illegitimate

[135198]transcription of phenylalanine hydroxylase for detection of mutations

[135199]in patients with phenylketonuria. Hum. Mutat. 1: 154–158, 1992.

[135200]

[135201]188. Ramus, S. J.; Forrest, S. M.; Saleeba, J. A.; Cotton, R. G. H.

[135202]: CpG hotspot causes second mutation in codon 408 of the phenylalanine

[135203]hydroxylase gene. Hum. Genet. 90: 147–148, 1992.

[135204]

[135205]189. Rao, D. C.; Keats, B. J.; Lalouel, J. M.; Morton, N. E.; Yee,

[135206]S.: A maximum likelihood map of chromosome 1. Am. J. Hum. Genet. 31:

[135207]680–696, 1979.

[135208]

[135209]190. Rey, F.; Berthelon, M.; Caillaud, C.; Lyonnet, S.;
Abadie, V.;

[135210]Blandin-Savoja, F.; Feingold, J.; Saudubray, J.-M.; Frezal,
J.; Munnich,

[135211]A.; Rey, J.: Clinical and molecular heterogeneity of phenyl-
alanine

[135212]hydroxylase deficiencies in France. Am. J. Hum. Genet. 43:
914-921,

[135213]1988.

[135214]

[135215]191. Riess, O.; Michel, A.; Speer, A.; Cobet, G.; Coutelle,
C.: Introduction

[135216]of genomic diagnosis of classical phenylketonuria to the
health care

[135217]system of the German Democratic Republic. Clin. Genet.
32: 209-215,

[135218]1987.

[135219]

[135220]192. Rohr, F. J.; Doherty, L. B.; Waisbren, S. E.; Bailey, I. V.;

[135221]Ampola, M. G.; Benacerraf, B.; Levy, H. L.: New England

maternal

[135222]PKU project: prospective study of untreated and treated pregnancies

[135223]and their outcomes. J. Pediat. 110: 391–398, 1987.

[135224]

[135225]193. Romeo, G.; Menozzi, P.; Ferlini, A.; Prosperi, L.; Cerone, R.;

[135226]Scalisi, S.; Romano, C.; Antonozzi, I.; Riva, E.; Piceni Sereni, L.;

[135227]Zammarchi, E.; Lenzi, G.; Sartorio, R.; Andria, G.; Cioni, M.; Fois,

[135228]A.; Burrioni, M.; Burlina, A. B.; Carnevale, F.: Incidence of classic

[135229]PKU in Italy estimated from consanguineous marriages and from neonatal

[135230]screening. Clin. Genet. 24: 339–345, 1983.

[135231]

[135232]194. Rosenblatt, D. S.; Scriver, C. R.: Heterogeneity in genetic

[135233]control of phenylalanine metabolism in man. Nature 218: 677–678,

[135234]1968.

[135235]

[135236]195. Rouse, B.; Azen, C.; Koch, R.; Matalon, R.; Hanley, W.;
de la

[135237]Cruz, F.; Trefz, F.; Friedman, E.; Shifrin, H.: Maternal
phenylketonuria

[135238]collaborative study (MPKUCS) offspring: facial anomalies,
malformations,

[135239]and early neurological sequelae. Am. J. Med. Genet. 69:
89–95, 1997.

[135240]

[135241]196. Rouse, B.; Matalon, R.; Koch, R.; Azen, C.; Levy, H.;
Hanley,

[135242]W.; Trefz, F.; de la Cruz, F.: Maternal phenylketonuria syn-
drome:

[135243]congenital heart defects, microcephaly, and developmen-
tal outcomes. J.

[135244]Pediat. 136: 57–61, 2000.

[135245]

[135246]197. Sarkissian, C. N.; Shao, Z.; Blain, F.; Peevers, R.; Su,
H.;

[135247]Heft, R.; Chang, T. M. S.; Scriver, C. R.: A different ap-
proach to

[135248] treatment of phenylketonuria: phenylalanine degradation
with recombinant

[135249] phenylalanine ammonia lyase. Proc. Nat. Acad. Sci. 96:
2339–2344,

[135250] 1999.

[135251]

[135252] 198. Saugstad, L. F.: Frequency of phenylketonuria in Nor-
way. Clin.

[135253] Genet. 7: 40–51, 1975.

[135254]

[135255] 199. Saugstad, L. F.: Anthropological significance of
phenylketonuria. Clin.

[135256] Genet. 7: 52–61, 1975.

[135257]

[135258] 200. Scott, T. M.; Fyfe, W. M.; Hart, D. M.: Maternal
phenylketonuria:

[135259] abnormal baby despite low phenylalanine diet during
pregnancy. Arch.

[135260] Dis. Child. 55: 634–649, 1980.

[135261]

[135262]201. Scriver, C. R.: Personal Communication. Montreal,
Quebec, Canada

[135263]1974.

[135264]

[135265]202. Scriver, C. R.; Clow, C. L.: Phenylketonuria and other
phenylalanine

[135266]hydroxylation mutants in man. Ann. Rev. Genet. 14:
179–202, 1980.

[135267]

[135268]203. Scriver, C. R.; Clow, C. L.: Phenylketonuria: epitome
of human

[135269]biochemical genetics. New Eng. J. Med. 303: 1336–1342
and 1394–1400,

[135270]1980.

[135271]

[135272]204. Scriver, C. R.; Eisensmith, R. C.; Woo, S. L. C.; Kauf-
man, S.

[135273]: The hyperphenylalaninemias of man and mouse. Annu.
Rev. Genet. 28:

[135274]141–165, 1994.

[135275]

[135276]205. Scriver, C. R.; Gregory, D. M.; Sovetts, D.; Clow, C. L.;
Tissenbaum,

[135277]G.: Normal plasma free amino acid values in adults: the
influence

[135278]of some common physiological variables. Metabolism 34:
868–873,

[135279]1985.

[135280]

[135281]206. Scriver, C. R.; Kaufman, S.; Woo, S. L. C.: The hyper-
phenylalaninemias.In:

[135282]Scriver, C. R.; Beaudet, A. L.; Sly, W. S.; Valle, D.: The
Metabolic

[135283]Basis of Inherited Disease. New York: McGraw–Hill (pub.)
(6th

[135284]ed.) I: 1989. Pp. 495–546.

[135285]

[135286]207. Shimizu, A.; Sakai, Y.; Ohno, K.; Masaki, S.; Kuwano,
R.; Takahashi,

[135287]Y.; Miyashita, N.; Watanabe, T.: A molecular genetic link-
age map

[135288]of mouse chromosome 10, including the Myb, S100b, Pah,
Sl, and Ifg

[135289]genes. Biochem. Genet. 30: 529–535, 1992.

[135290]

[135291]208. Smith, C. B.; Kang, J.: Cerebral protein synthesis in a genetic

[135292]mouse model of phenylketonuria. Proc. Nat. Acad. Sci. 97: 11014–11019,

[135293]2000.

[135294]

[135295]209. Smith, I.; Macartney, F. J.; Erdohazi, M.; Pincott, J. R.; Wolff,

[135296]O. H.; Brenton, D. P.; Biddle, S. A.; Fairweather, D. V. I.; Dobbing,

[135297]J.: Fetal damage despite low-phenylalanine diet after conception

[135298]in a phenylketonuric woman. Lancet I: 17–19, 1979.

[135299]

[135300]210. Steffens, C.: No difference in dermatoglyphics of fingers and

[135301]palms between phenylketonuria patients and controls. (Letter) Hum.

[135302]Genet. 69: 195 only, 1985.

[135303]

[135304]211. Stegink, L. D.; Filer, L. J., Jr.; Baker, G. L.; Bell, E. F.;

[135305]Ziegler, E. E.; Brummel, M. C.; Krause, W. L.: Repeated in-
gestion

[135306]of aspartame-sweetened beverage: effect on plasma
amino acid concentrations

[135307]in individuals heterozygous for phenylketonuria.

Metabolism 38:

[135308]78-84, 1989.

[135309]

[135310]212. Stuhmann, M.; Riess, O.; Monch, E.; Kurdoglu, G.:

Haplotype

[135311]analysis of the phenylalanine hydroxylase gene in Turkish
phenylketonuria

[135312]families. Clin. Genet. 36: 117-121, 1989.

[135313]

[135314]213. Superti-Furga, A.; Steinmann, B.; Duc, G.; Gitzel-
mann, R.: Maternal

[135315]phenylketonuria syndrome in cousins caused by mild, un-
recognized phenylketonuria

[135316]in their mothers homozygous for the phenylalanine hy-

droxylase arg261-to-gln

[135317]mutation. *Europ. J. Pediat.* 150: 493–497, 1991.

[135318]

[135319]214. Svensson, E.; Andersson, B.; Hagenfeldt, L.: Two mutations within

[135320]the coding sequence of the phenylalanine hydroxylase gene. *Hum. Genet.* 85:

[135321]300–304, 1990.

[135322]

[135323]215. Svensson, E.; Eisensmith, R. C.; Dworniczak, B.; von Dobeln,

[135324]U.; Hagenfeldt, L.; Horst, J.; Woo, S. L. C.: Two missense mutations

[135325]causing mild hyperphenylalaninemia associated with DNA haplotype 12. *Hum.*

[135326]*Mutat.* 1: 129–137, 1992.

[135327]

[135328]216. Takahashi, K.; Kure, S.; Matsubara, Y.; Narisawa, K.: Novel

[135329]phenylketonuria mutation detected by analysis of ectopically transcribed

[135330]phenylalanine hydroxylase mRNA from lymphoblast.

(Letter) Lancet 340:

[135331]1473 only, 1992.

[135332]

[135333]217. Teebi, A. S.; Al-Awadi, S. A.; Farag, T. I.; Naguib, K.
K.; El-Khalifa,

[135334]M. Y.: Phenylketonuria in Kuwait and Arab countries. Eu-
rop. J. Pediat. 146:

[135335]59-60, 1987.

[135336]

[135337]218. Thompson, A. J.; Tillotson, S.; Smith, I.; Kendall, B.;
Moore,

[135338]S. G.; Brenton, D. P.: Brain MRI changes in phenylke-
tonuria: associations

[135339]with dietary status. Brain 116: 811-821, 1993.

[135340]

[135341]219. Tourian, A. Y.; Sidbury, J. B., Jr.: Phenylketonuria.In:
Stanbury,

[135342]J. B.; Wyngaarden, J. B.; Fredrickson, D. S.: Metabolic Basis
of

[135343]Inherited Disease. New York: McGraw-Hill (pub.) (4th ed.):

1978.

[135344]Pp. 240–255.

[135345]

[135346]220. Tsai, T.–F.; Hsiao, K.–J.; Su, T.–S.: Phenylketonuria
mutation

[135347]in Chinese haplotype 44 identical with haplotype 2 muta-
tion in northern–European

[135348]Caucasians. Hum. Genet. 84: 409–411, 1990.

[135349]

[135350]221. Tyfield, L. A.; Meredith, A. L.; Osborn, M. J.; Harper,
P. S.

[135351]: Identification of the haplotype pattern associated with
the mutant

[135352]PKU allele in the Gypsy population of Wales. J. Med. Genet.
26:

[135353]499–503, 1989.

[135354]

[135355]222. Tyfield, L. A.; Stephenson, A.; Cockburn, F.; Harvie,
A.; Bidwell,

[135356]J. L.; Wood, N. A. P.; Pilz, D. T.; Harper, P.; Smith, I.: Se-
quence

[135357]variation at the phenylalanine hydroxylase gene in the
British Isles. Am.

[135358]J. Hum. Genet. 60: 388–396, 1997.

[135359]

[135360]223. Ullrich, K.; Weglage, J.; Schuierer, G.; Funders, B.;
Pietsch,

[135361]M.; Koch, H. G.; Hahn-Ullrich, H.: Cranial MRI in PKU: eval-
uation

[135362]of a critical threshold for blood phenylalanine. (Letter)
Neuropediatrics 25:

[135363]278–279, 1994.

[135364]

[135365]224. Usha, R.; Uma, R.; Farag, T. I.; Girish, Y.; Al-Ghanim,
M. M.

[135366]A.; Al-Najdi, K.; Al-Awadi, S. A.; El-Badramany, M. H.: Late
diagnosis

[135367]of phenylketonuria in a Bedouin mother. Am. J. Med.
Genet. 44: 713–715,

[135368]1992.

[135369]

[135370]225. Vajro, P.; Strisciuglio, P.; Houssin, D.; Huault, G.;

Laurent,

[135371]J.; Alvarez, F.; Bernard, O.: Correction of phenylketonuria after

[135372]liver transplantation in a child with cirrhosis. (Letter) New Eng.

[135373]J. Med. 329: 363 only, 1993.

[135374]

[135375]226. van Spronsen, F. J.; Reijngoud, D.-J.; Smit, G. P. A.; Nagel,

[135376]G. T.; Stellaard, F.; Berger, R.; Heymans, H. S. A.: Phenylketonuria:

[135377]the in vivo hydroxylation rate of phenylalanine into tyrosine is decreased. J.

[135378]Clin. Invest. 101: 2875-2880, 1998.

[135379]

[135380]227. Waisbren, S. E.; Hanley, W.; Levy, H. L.; Shifrin, H.; Allred,

[135381]E.; Azen, C.; Chang, P.-N.; Cipic-Schmidt, S.; de la Cruz, F.; Hall,

[135382]R.; Matalon, R.; Nanson, J.; Rouse, B.; Trefz, F.; Koch, R.: Outcome

[135383]at age 4 years in offspring of women with maternal

phenylketonuria:

[135384]the maternal PKU collaborative study. J.A.M.A. 283:
756-762, 2000.

[135385]

[135386]228. Wang, T.; Okano, Y.; Eisensmith, R.; Huang, S.-Z.;
Zeng, Y.-T.;

[135387]Lo, W. H. Y.; Woo, S. L. C.: Molecular genetics of phenylke-
tonuria

[135388]in Orientals: linkage disequilibrium between a termination
mutation

[135389]and haplotype 4 of the phenylalanine hydroxylase gene.
Am. J. Hum.

[135390]Genet. 45: 675-680, 1989.

[135391]

[135392]229. Wang, T.; Okano, Y.; Eisensmith, R. C.; Fekete, G.;
Schuler,

[135393]D.; Berencsi, G.; Nasz, I.; Woo, S. L. C.: Molecular genetics
of

[135394]PKU in Eastern Europe: a nonsense mutation associated
with haplotype

[135395]4 of the phenylalanine hydroxylase gene. Somat. Cell
Molec. Genet. 16:

[135396]85–89, 1990.

[135397]

[135398]230. Wang, T.; Okano, Y.; Eisensmith, R. C.; Harvey, M. L.;
Lo, W.

[135399]H. Y.; Huang, S.-Z.; Zeng, Y.-T.; Yuan, L.-F.; Furuyama, J.;
Oura,

[135400]T.; Sommer, S. S.; Woo, S. L. C.: Founder effect of a preva-
lent phenylketonuria

[135401]mutation in the Oriental population. Proc. Nat. Acad. Sci.
88: 2146–2150,

[135402]1991.

[135403]

[135404]231. Wang, T.; Okano, Y.; Eisensmith, R. C.; Lo, W. H. Y.;
Huang,

[135405]S.-Z.; Zeng, Y.-T.; Woo, S. L. C.: Identification of a novel
phenylketonuria

[135406](PKU) mutation in the Chinese: further evidence for multi-
ple origins

[135407]of PKU in Asia. Am. J. Hum. Genet. 48: 628–630, 1991.

[135408]

[135409]232. Wang, T.; Okano, Y.; Eisensmith, R. C.; Lo, W. H. Y.;

Huang,

[135410]S.-Z.; Zeng, Y.-T.; Yuan, L.-F.; Liu, S.-R.; Woo, S. L. C.:

Identification

[135411]of three novel PKU mutations among Chinese: evidence

for recombination

[135412]or recurrent mutation at the PAH locus. Genomics 13:

230-231, 1992.

[135413]

[135414]233. Wang, T.; Okano, Y.; Eisensmith, R. C.; Lo, W. H. Y.;

Huang,

[135415]S.-Z.; Zeng, Y.-T.; Yuan, L.-F.; Liu, S.-R.; Woo, S. L. C.:

Missense

[135416]mutations prevalent in Orientals with phenylketonuria:

molecular characterization

[135417]and clinical implications. Genomics 10: 449-456, 1991.

[135418]

[135419]234. Wang, T.; Woo, S. L. C.: Personal Communication.

Houston, Tex.

[135420]1/20/1990.

[135421]

[135422]235. Waters, P. J.; Parniak, M. A.; Akerman, B. R.; Scriver,

C. R.

[135423] Characterization of phenylketonuria missense substitutions, distant

[135424] from the phenylalanine hydroxylase active site, illustrates a paradigm

[135425] for mechanism and potential modulation of phenotype. Molec. Genet.

[135426] Metab. 69: 101–110, 2000.

[135427]

[135428] 236. Weglage, J.; Grenzebach, M.; Pietsch, M.; Feldmann, R.; Linnenbank,

[135429] R.; Denecke, J.; Koch, H. G.: Behavioural and emotional problems

[135430] in early-treated adolescents with phenylketonuria in comparison with

[135431] diabetic patients and healthy controls. J. Inherit. Metab. Dis. 23:

[135432] 487–496, 2000.

[135433]

[135434] 237. Weglage, J.; Pietsch, M.; Denecke, J.; Sprinz, A.; Feldmann,

[135435] R.; Grenzebach, M.; Ullrich, K.: Regression of neuropsych-

chological

[135436]deficits in early-treated phenylketonurics during adolescence. J.

[135437]Inherit. Metab. Dis. 22: 693–705, 1999.

[135438]

[135439]238. Weinstein, M.; Eisensmith, R. C.; Abadie, V.; Avigad, S.; Lyonnet,

[135440]S.; Schwartz, G.; Munnich, A.; Woo, S. L. C.; Shiloh, Y.: A missense

[135441]mutation, S349P, completely inactivates phenylalanine hydroxylase

[135442]in North African Jews with phenylketonuria. Hum. Genet. 90: 645–649,

[135443]1993.

[135444]

[135445]239. Woo, S. L. C.: Personal Communication. Houston, Tex. 1/11/1983.

[135446]

[135447]240. Woo, S. L. C.: Collation of RFLP haplotypes at the human phenylalanine

[135448]hydroxylase (PAH) locus. (Letter) Am. J. Hum. Genet. 43:

781-783,

[135449]1988.

[135450]

[135451]241. Woo, S. L. C.: Molecular basis and population genetics of phenylketonuria. Biochemistry 28:

[135452]1-7, 1989.

[135453]

[135454]242. Woo, S. L. C.; Chandra, T.; Stackhouse, R.; Robson, K. J. H.

[135455]: Analysis of phenylketonuria (PKU) by cloning and characterization

[135456]of the human phenylalanine hydroxylase gene. (Abstract) Am. J. Hum.

[135457]Genet. 34: 168A only, 1982.

[135458]

[135459]243. Woo, S. L. C.; Lidsky, A.; Law, M.; Kao, F. T.: Regional mapping

[135460]of the human phenylalanine hydroxylase gene and PKU locus to 12q21-qter.

[135461](Abstract) Am. J. Hum. Genet. 36: 210S only, 1984.

[135462]

[135463]244. Woo, S. L. C.; Lidsky, A. S.; Guttler, F.; Chandra, T.;
Robson,

[135464]K. J. H.: Cloned human phenylalanine hydroxylase gene
permits prenatal

[135465]diagnosis and carrier detection of classical phenylke-
tonuria. Nature 306:

[135466]151-155, 1983.

[135467]

[135468]245. Woo, S. L. C.; Lidsky, A. S.; Guttler, F.; Thiru-
malachary, C.;

[135469]Robson, K. J. H.: Prenatal diagnosis of classical phenylke-
tonuria

[135470]by gene mapping. J.A.M.A. 251: 1998-2002, 1984.

[135471]

[135472]246. Woolf, L. I.: The heterozygote advantage in
phenylketonuria.

[135473](Letter) Am. J. Hum. Genet. 38: 773-775, 1986.

[135474]

[135475]247. Woolf, L. I.; Cranston, W. I.; Goodwin, B. L.: Genetics
of phenylketonuria.

[135476]I. Heterozygosity for phenylketonuria. II. Third allele at the

phenylalanine

[135477]hydroxylase locus in man. Nature 213: 882–885, 1967.

[135478]

[135479]248. Woolf, L. I.; Goodwin, B. L.; Cranston, W. I.; Wade, D. N.; Woolf,

[135480]F.; Hudson, F. P.; McBean, M. S.: A third allele at the phenylalanine–hydroxylase

[135481]locus in mild phenylketonuria (hyperphenylalaninaemia). Lancet I:

[135482]114–117, 1968.

[135483]

[135484]249. Woolf, L. I.; McBean, M. S.; Woolf, F. M.; Cahalane, S. F.:

[135485]Phenylketonuria as a balanced polymorphism: the nature of the heterozygote

[135486]advantage. Ann. Hum. Genet. 38: 461–469, 1975.

[135487]

[135488]250. Yu, J. S.; O'Halloran, M. T.: Atypical phenylketonuria in a

[135489]family with a phenylketonuric mother. Pediatrics 46: 707–711, 1970.

[135490]

[135491]251. Zschocke, J.; Graham, C. A.; Stewart, F. J.; Carson, D. J.; Nevin,

[135492]N. C.: Non-phenylketonuria hyperphenylalaninaemia in Northern Ireland:

[135493]frequent mutation allows screening and early diagnosis. Hum. Mutat. 4:

[135494]114–118, 1994.

[135495]

[135496]252. Zschocke, J.; Mallory, J. P.; Eiken, H. G.; Nevin, N. C.: Phenylketonuria

[135497]and the peoples of Northern Ireland. Hum. Genet. 100: 189–194, 1997.

[135498]

[135499]253. Zschocke, J.; Quak, E.; Knauer, A.; Fritz, B.; Aslan, M.; Hoffmann,

[135500]G. F.: Large heterozygous deletion masquerading as homozygous missense

[135501]mutation: a pitfall in diagnostic mutation analysis. J. Inherit.

[135502]Metab. Dis. 22: 687–692, 1999.

[135503]

[135504]254. Zygulska, M.; Eigel, A.; Aulehla-Scholz, C.; Pietrzyk, J. J.;

[135505]Horst, J.: Molecular analysis of PKU haplotypes in the population

[135506]of southern Poland. Hum. Genet. 86: 292-294, 1991.

[135507]

[135508]255. Zygulska, M.; Eigel, A.; Dworniczak, B.; Sutkowska, A.; Pietrzyk,

[135509]J. J.; Horst, J.: Phenylketonuria in Poland: 66% of PKU alleles are

[135510]caused by three mutations. Hum. Genet. 88: 91-94, 1991.

[135511]24109.

[135512]Bessant, D. A. R.; Payne, A. M.; Plant, C.; Bird, A. C.; Bhat-tacharya,

[135513]S. S.: Further refinement of the Usher 2A locus at 1q41. J. Med.

[135514]Genet. 35: 773-774, 1998.

[135515]24110.

[135516]Konrad, M.; Saunier, S.; Heidet, L.; Silbermann, F.; Bennisy,

[135517]F.; Calado, J.; Le Paslier, D.; Broyer, M.; Gubler, M.-C.;
Antignac,

[135518]C.: Large homozygous deletions of the 2q13 region are a
major cause

[135519]of juvenile nephronophthisis. Hum. Molec. Genet. 5:
367-371, 1996.

[135520]24111.

[135521]Degos, L.; Dautigny, A.; Brouet, J. C.; Colombani, M.; Ar-
daillou,

[135522]N.; Caen, J. P.; Colombani, J.: A molecular defect in throm-
basthenic

[135523]platelets. J. Clin. Invest. 56: 236-240, 1975.

[135524]24112.

[135525]Djaffar, I.; Vilette, D.; Pidard, D.; Wautier, J.-L.; Rosa, J.-P.

[135526]: Human platelet antigen 3 (HPA-3): localization of the de-
terminant

[135527]of the alloantibody Lek(a)(HPA-3a) to the C-terminus of
platelet glycoprotein

[135528]IIb heavy chain and contribution of O-linked carbohy-
drates. Thromb.

[135529]Haemost. 69: 485-489, 1993.

[135530]24113.

[135531]Friedman, L. L.; Bowie, E. J. W.; Thompson, J. H., Jr.;
Brown,

[135532]A. L., Jr.; Owen, C. A., Jr.: Familial Glanzmann's thrombas-
thenia. Mayo

[135533]Clin. Proc. 39: 908-918, 1964.

[135534]24114.

[135535]Giltay, J. C.; Leeksa, O. C.; Breederveld, C.; van Mourik, J.

[135536]A.: Normal synthesis and expression of endothelial IIb/IIIa
in Glanzmann's

[135537]thrombasthenia. Blood 69: 809-812, 1987.

[135538]24115.

[135539]Grimaldi, C. M.; Chen, F.; Wu, C.; Weiss, H. J.; Coller, B. S.;

[135540]French, D. L.: Glycoprotein IIb leu214pro mutation pro-
duces Glanzmann

[135541]thrombasthenia with both quantitative and qualitative ab-
normalities

[135542]in GPIIb/IIIa. Blood 91: 1562-1571, 1998.

[135543]24116.

[135544]Gross, R.; Gerok, W.; Lohr, G. W.; Vogell, W.; Waller, H. D.;

[135545]Theopold, W.: Ueber die Natur der Thrombasthenie.

Thrombopathie Glanzmann-Naegeli. Klin.

[135546]Wschr. 38: 193-206, 1960.

[135547]24117.

[135548]Hathaway, W. E.: Bleeding disorders due to platelet dys-

function. Am.

[135549]J. Dis. Child. 121: 127-134, 1971.

[135550]24118.

[135551]Herrmann, F. H.; Meyer, M.; Gogstad, G. O.; Solum, N. O.:

Glycoprotein

[135552]IIb-IIIa complex in platelets of patients and heterozygotes

of Glanzmann's

[135553]thrombasthenia. Thromb. Res. 32: 615-622, 1983.

[135554]24119.

[135555]Herrmann, F. H.; Meyer, M.; Ihle, E.: Protein and glycopro-

tein

[135556]abnormalities in an unusual subtype of Glanzmann's

thrombasthenia. Haemostasis 12:

[135557]337-344, 1982.

[135558]24120.

[135559]Kanska, B.; Niewiarowski, S.; Ostrowski, L.; Poplawski, A.;
Prokopowicz,

[135560]J.: Macrothrombocytic thrombopathia. Clinical, coagula-
tion and hereditary

[135561]aspects. Thromb. Diath. Haemorrh. 10: 88-100, 1963.

[135562]24121.

[135563]Karpatkin, S.; Weiss, H. J.: Deficiency of glutathione per-
oxidase

[135564]associated with high levels of reduced glutathione in
Glanzmann's

[135565]thrombasthenia. New Eng. J. Med. 287: 1062-1066, 1972.

[135566]24122.

[135567]Kato, A.; Yamamoto, K.; Miyazaki, S.; Jung, S. M.; Moroi,
M.;

[135568]Aoki, N.: Molecular basis for Glanzmann's thrombasthenia
(GT) in

[135569]a compound heterozygote with glycoprotein IIb gene: a
proposal for

[135570]the classification of GT based on the biosynthetic pathway

of glycoprotein

[135571]Iib-IIIa complex. Blood 79: 3212-3218, 1992.

[135572]24123.

[135573]Khanduri, U.; Pulimood, R.; Sudarsanam, A.; Carman, R.
H.; Jadhav,

[135574]M.; Pereira, S.: Glanzmann's thrombasthenia: a review and
report

[135575]of 42 cases from South India. Thromb. Haemost. 46:
717-721, 1981.

[135576]24124.

[135577]Kunicki, T. J.; Aster, R. H.: Deletion of the platelet-specific

[135578]alloantigen PI(A1) from platelets in Glanzmann's throm-
basthenia. J.

[135579]Clin. Invest. 61: 1225-1231, 1978.

[135580]24125.

[135581]Lelong, J. C.: La thrombopathie de Glanzmann-Naegeli.
Paris:

[135582]R. Foulon et Cie. (pub.) 1960.

[135583]24126.

[135584]Levy, J. M.; Mayer, G.; Sacrez, R.; Ruff, R.; Francfort, J. J.;

[135585]Rodier, L.: Thrombasthenie de Glanzmann-Naegeli: etude
d'un groupe

[135586]ethnique a forte endogamie. Ann. Pediat. 18: 129-137,
1971.

[135587]24127.

[135588]Luo, X. Y.; Bray, P. F.; Magenis, R. E.: Precise localization

[135589]of the gene for platelet membrane glycoprotein IIb to
17q21.32 using

[135590]structural rearrangements of chromosome 17. (Abstract)
Cytogenet.

[135591]Cell Genet. 51: 1036 only, 1989.

[135592]24128.

[135593]Marx, R.; Jean, G.: Studien zur Pathogenese der Throm-
basthenie

[135594]Glanzmann-Naegeli. Klin. Wschr. 40: 942-953, 1962.

[135595]24129.

[135596]McEver, R. P.; Baenziger, J. U.; Majerus, P. W.: Isolation and

[135597]structural characterization of the polypeptide subunits of
membrane

[135598]glycoprotein IIb-IIIa from human platelets. Blood 59:
80-85, 1982.

[135599]24130.

[135600]McEver, R. P.; Baenziger, N. L.; Majerus, P. W.: Isolation
and

[135601]quantitation of the platelet membrane glycoprotein defi-
cient in thrombasthenia

[135602]using a monoclonal hybridoma antibody. J. Clin. Invest.
66: 1311-1318,

[135603]1980.

[135604]24131.

[135605]Meyer, M.; Herrmann, F. H.: Diversity of glycoprotein defi-
ciencies

[135606]in Glanzmann's thrombasthenia. Thromb. Haemost. 54:
626-629, 1985.

[135607]24132.

[135608]Moser, K.; Lechner, K.; Vinazzer, H.: A hitherto not de-
scribed

[135609]enzyme defect in thrombasthenia: glutathione reductase
deficiency. Thromb.

[135610]Diath. Haemorrh. 19: 46-52, 1968.

[135611]24133.

[135612]Nachman, R. L.: Thrombasthenia: immunologic evidence

of a platelet

[135613]protein abnormality. J. Lab. Clin. Med. 67: 411-419, 1966.

[135614]24134.

[135615]Newman, P. J.: Platelet GPIIb-IIIa: molecular variations and

[135616]alloantigens. Thromb. Haemost. 66: 111-118, 1991.

[135617]24135.

[135618]Nurden, A. T.; Caen, J. P.: An abnormal platelet glycoprotein

[135619]pattern in three cases of Glanzmann's thrombasthenia.
Brit. J. Haemat. 28:

[135620]253-260, 1974.

[135621]24136.

[135622]Eisensmith, R. C.; Woo, S. L. C.: Molecular basis of
phenylketonuria

[135623]and related hyperphenylalaninemias: mutations and polymorphisms in

[135624]the human phenylalanine hydroxylase gene. Hum. Mutat.
1: 13-23,

[135625]1992.

[135626]24137.

[135627]Eisensmith, R. C.; Woo, S. L. C.: Updated listing of haplo-
types

[135628]at the human phenylalanine hydroxylase (PAH) locus.
(Letter) Am.

[135629]J. Hum. Genet. 51: 1445-1448, 1992.

[135630]24138.

[135631]Eisensmith, R. C.; Woo, S. L. C.: Gene therapy for
phenylketonuria. Europ.

[135632]J. Pediat. 155 (suppl. 1): S16-S19, 1996.

[135633]24139.

[135634]Fisch, R. O.; Tagatz, G.; Stassart, J. P.: Gestational carrier-
-a

[135635]reproductive haven for offspring of mothers with
phenylketonuria (PKU):

[135636]an alternative therapy for maternal PKU. J. Inherit. Metab.
Dis. 16:

[135637]957-961, 1993.

[135638]24140.

[135639]Flatz, G.; Oelbe, M.; Herrmann, H.: Ethnic distribution of
phenylketonuria

[135640]in the north German population. Hum. Genet. 65:

396–399, 1984.

[135641]24141.

[135642]Folling, A.: Ueber Ausscheidung von Phenylbrenztrauben-
saeure

[135643]in den Harn als Stoffwechselanomalie in Verbindung mit
Imbezillitaet. Ztschr.

[135644]Physiol. Chem. 227: 169–176, 1934.

[135645]24142.

[135646]Forrest, S. M.; Dahl, H. H.; Howells, D. W.; Dianzani, I.;
Cotton,

[135647]R. G. H.: Mutation detection in phenylketonuria by using
chemical

[135648]cleavage of mismatch: importance of using probes from
both normal

[135649]and patient samples. Am. J. Hum. Genet. 49: 175–183,
1991.

[135650]24143.

[135651]Frankenburg, W. K.; Duncan, B. R.; Coffelt, R. W.; Koch, R.;
Coldwell,

[135652]J. G.; Son, C. D.: Maternal phenylketonuria: implications
for growth

[135653]and development. J. Pediat. 73: 560–570, 1968.

[135654]24144.

[135655]Friedman, P. A.; Fisher, D. B.; Kang, E. S.; Kaufman, S.: De-
tection

[135656]of hepatic phenylalanine 4-hydroxylase in classical
phenylketonuria. Proc.

[135657]Nat. Acad. Sci. 70: 552-556, 1973.

[135658]24145.

[135659]Gjetting, T.; Petersen, M.; Guldberg, P.; Guttler, F.: Mis-
sense

[135660]mutations in the N-terminal domain of human phenylala-
nine hydroxylase

[135661]interfere with binding of regulatory phenylalanine. Am. J.
Hum. Genet. 68:

[135662]1353-1360, 2001.

[135663]24146.

[135664]Greeves, L. G.; Patterson, C. C.; Carson, D. J.; Thom, R.;
Wolfenden,

[135665]M. C.; Zshock, J.; Graham, C. A.; Nevin, N. C.; Trimble, E.
R.:

[135666]Effect of genotype on changes in intelligence quotient af-
ter dietary

- [135667]relaxation in phenylketonuria and hyperphenylalani-
naemia. Arch. Dis.
- [135668]Child. 82: 216–221, 2000.
- [135669]24147.
- [135670]Gregory, D. M.; Sovetts, D.; Clow, C. L.; Sriver, C. R.:
Plasma
- [135671]free amino acid values in normal children and adoles-
cents. Metabolism 35:
- [135672]967–969, 1986.
- [135673]24148.
- [135674]Griffiths, P. V.; Demellweek, C.; Fay, N.; Robinson, P. H.;
Davidson,
- [135675]D. C.: Wechsler subscale IQ and subtest profile in early
treated
- [135676]phenylketonuria. Arch. Dis. Child. 82: 209–215, 2000.
- [135677]24149.
- [135678]Guldborg, P.; Henriksen, K. F.; Sipila, I.; Guttler, F.; de la
- [135679]Chapelle, A.: Phenylketonuria in a low incidence popula-
tion: molecular
- [135680]characterization of mutations in Finland. J. Med. Genet.
32: 976–978,

[135681]1995.

[135682]24150.

[135683]Guldberg, P.; Henriksen, K. F.; Thony, B.; Blau, N.; Guttler,

[135684]F.: Molecular heterogeneity of nonphenylketonuria hyper-
phenylalaninemia

[135685]in 25 Danish patients. Genomics 21: 453–455, 1994.

[135686]24151.

[135687]Guldberg, P.; Lou, H. C.; Henriksen, K. F.; Mikkelsen, I.;
Olsen,

[135688]B.; Holck, B.; Guttler, F.: A novel missense mutation in the
phenylalanine

[135689]hydroxylase gene of a homozygous Pakistani patient with
non-PKU hyperphenylalaninemia. Hum.

[135690]Molec. Genet. 2: 1061–1062, 1993.

[135691]24152.

[135692]Guldberg, P.; Zschocke, J.; Dagbjartsson, A.; Henriksen, K.
F.;

[135693]Guttler, F.: A molecular survey of phenylketonuria in Ice-
land: identification

[135694]of a founding mutation and evidence of predominant
Norse settlement. Europ.

[135695]J. Hum. Genet. 5: 376–381, 1997.

[135696]24153.

[135697]Guthrie, R.: The introduction of newborn screening for phenylketonuria:

[135698]a personal history. Europ. J. Pediat. 155 (suppl. 1): 4–5, 1996.

[135699]24154.

[135700]Guthrie, R.; Susi, A.: A simple phenylalanine method for detecting

[135701]phenylketonuria in large populations of newborn infants. Pediatrics 32:

[135702]338–343, 1963.

[135703]24155.

[135704]Guldborg, P.; Rey, F.; Zschocke, J.; Romano, V.; Francois, B.;

[135705]Michiels, L.; Ullrich, K.; Hoffmann, G. F.; Burgard, P.; Schmidt,

[135706]H.; Meli, C.; Riva, E.; Dianzani, I.; Ponzone, A.; Rey, J.; Gütler,

[135707]F.: A European multicenter study of phenylalanine hydroxylase deficiency:

- [135708]classification of 105 mutations and a general system for
genotype-based
- [135709]prediction of metabolic phenotype. Am. J. Hum. Genet. 63:
71-79,
- [135710]1998.
- [135711]24156.
- [135712]Guttler, F.: Hyperphenylalaninemia: diagnosis and classifi-
cation
- [135713]of the various types of phenylalanine hydroxylase defi-
ciency in childhood. Acta
- [135714]Paediat. Scand. 280 (suppl.): 1-80, 1980.
- [135715]24157.
- [135716]Guttler, F.; Azen, C.; Guldberg, P.; Romstad, A.; Hanley, W.
B.;
- [135717]Levy, H. L.; Matalon, R.; Rouse, B. M.; Trefz, F.; de la Cruz,
F.;
- [135718]Koch, R.: Relationship among genotype, biochemical phe-
notype, and
- [135719]cognitive performance in females with phenylalanine hy-
droxylase deficiency:
- [135720]report from the maternal phenylketonuria collaborative
study. Pediatrics 104:

[135721]258–262, 1999.

[135722]24158.

[135723]Guttler, F.; Hansen, G.: Heterozygote detection in phenylketonuria. Clin.

[135724]Genet. 11: 137–146, 1977.

[135725]24159.

[135726]Guttler, F.; Ledley, F. D.; Lidsky, A. S.; DiLella, A. G.; Sullivan,

[135727]S. E.; Woo, S. L. C.: Correlation between polymorphic DNA haplotypes

[135728]at phenylalanine hydroxylase locus and clinical phenotypes of phenylketonuria. J.

[135729]Pediat. 110: 68–71, 1987.

[135730]24160.

[135731]Guttler, F.; Woo, S. L. C.: Molecular genetics of PKU. J. Inherit.

[135732]Metab. Dis. 9 (suppl. 1): 58–68, 1986.

[135733]24161.

[135734]Hanley, W. B.; Clarke, J. T. R.; Schoonheydt, W.: Maternal phenylketonuria

[135735](PKU)--a review. Clin. Biochem. 20: 149-156, 1987.

[135736]24162.

[135737]Hanley, W. B.; Koch, R.; Levy, H. L.; Matalon, R.; Rouse, B.;

[135738]Azen, C.; de la Cruz, F.: The North American Maternal
Phenylketonuria

[135739]Collaborative Study, developmental assessment of the off-
spring: preliminary

[135740]report. Europ. J. Pediat. 155 (suppl. 1): 169-172, 1996.

[135741]24163.

[135742]Hanley, W. B.; Lee, A. W.; Hanley, A. J. G.; Lehotay, D. C.;
Austin,

[135743]V. J.; Schoonheydt, W. E.; Platt, B. A.; Clarke, J. T. R.: 'Hypo-
tyrosinemia'

[135744]in phenylketonuria. Molec. Genet. Metab. 69: 286-294,
2000.

[135745]24164.

[135746]Hertzberg, M.; Jahromi, K.; Ferguson, V.; Dahl, H. H. M.;
Mercer,

[135747]J.; Mickleson, K. N. P.; Trent, R. J.: Phenylalanine hydroxy-
lase

[135748]gene haplotypes in Polynesians: evolutionary origins and

absence of

[135749]alleles associated with severe phenylketonuria. Am. J.

Hum. Genet. 44:

[135750]382–387, 1989.

[135751]24165.

[135752]Hoang, L.; Byck, S.; Prevost, L.; Scriver, C. R.: PAH Muta-

tion

[135753]Analysis Consortium Database: a database for disease-

producing and

[135754]other allelic variation at the human PAH locus. Nucleic

Acids Res. 24:

[135755]127–131, 1996.

[135756]24166.

[135757]Hofman, K. J.; Steel, G.; Kazazian, H. H.; Valle, D.:

Phenylketonuria

[135758]in U.S. blacks: molecular analysis of the phenylalanine hy-

droxylase

[135759]gene. Am. J. Hum. Genet. 48: 791–798, 1991.

[135760]24167.

[135761]Garland, D.; Rao, P. V.; Del Corso, A.; Mura, U.; Zigler, J.

S.,

[135762]Jr.: Zeta-crystallin is a major protein in the lens of
Camelus dromedarius.

[135763]Arch. Biochem. Biophys. 285: 134-136, 1991.

[135764]24168.

[135765]Gonzalez, P.; Rao, P. V.; Zigler, J. S., Jr.: Organization of
[135766]the human zeta-crystallin/quinone reductase gene
(CRYZ). Genomics 21:
[135767]317-324, 1994.

[135768]24169.

[135769]Heinzmann, C.; Kojis, T. L.; Gonzalez, P.; Rao, P. V.; Zigler,
[135770]J. S., Jr.; Polymeropoulos, M. H.; Klisak, I.; Sparkes, R. S.;
Mohandas,
[135771]T.; Bateman, J. B.: Assignment of the zeta-crystallin gene
(CRYZ)
[135772]to human chromosome 1p22-p31 and identification of
restriction fragment
[135773]length polymorphisms. Genomics 23: 403-407, 1994.

[135774]24170.

[135775]Huang, Q.-L.; Russell, P.; Stone, S. H.; Zigler, J. S., Jr.:
Zeta-crystallin,

[135776]a novel lens protein from the guinea pig. Curr. Eye Res. 6:
725-732,

[135777]1987.

[135778]24171.

[135779]Rodriguez, I. R.; Gonzalez, P.; Zigler, J. S., Jr.; Borrás, T.

[135780]; A guinea-pig hereditary cataract contains a splice site
deletion

[135781]in a crystallin gene. Biochim. Biophys. Acta 1180: 44-52,
1992.

[135782]24172.

[135783]Morrison, N. A.; Yeoman, R.; Kelly, P. J.; Eisman, J. A.:
Contribution

[135784]of trans-acting factor alleles to normal physiological vari-
ability:

[135785]vitamin D receptor gene polymorphisms and circulating
osteocalcin. Proc.

[135786]Nat. Acad. Sci. 89: 6665-6669, 1992.

[135787]24173.

[135788]Pan, L. C.; Price, P. A.: The propeptide of rat bone
gamma-carboxyglutamic

[135789]acid protein shares homology with other vitamin K-

dependent protein

[135790]precursors. Proc. Nat. Acad. Sci. 82: 6109–6113, 1985.

[135791]24174.

[135792]Puchacz, E.; Lian, J. B.; Stein, G. S.; Wozney, J.; Huebner, K.;

[135793]Croce, C.: Chromosomal localization of the human osteocalcin gene. Endocrinology 124:

[135794]2648–2650, 1989.

[135795]24175.

[135796]Tabas, J. A.; Zasloff, M.; Wasmuth, J. J.; Emanuel, B. S.; Altherr,

[135797]M. R.; McPherson, J. D.; Wozney, J. M.; Kaplan, F. S.: Bone morphogenetic

[135798]protein: chromosomal localization of human genes for BMP1, BMP2A,

[135799]and BMP3. Genomics 9: 283–289, 1991.

[135800]24176.

[135801]Wozney, J. M.; Rosen, V.; Celeste, A. J.; Mitsock, L. M.; Whitters,

[135802]M. J.; Kriz, R. W.; Hewick, R. M.; Wang, E. A.: Novel regulators

[135803]of bone formation: molecular clones and activities. Science 242:

[135804]1528–1534, 1988.

[135805]24177.

[135806]Connor, J. M.: Fibrodysplasia ossificans progressiva: lessons

[135807]from rare maladies. (Editorial) New Eng. J. Med. 335: 591–593, 1996.

[135808]24178.

[135809]Dooley, C. A.; Attia, G. R.; Rainey, W. E.; Moore, D. R.; Carr,

[135810]B. R.: Bone morphogenetic protein inhibits ovarian androgen production. J.

[135811]Clin. Endocr. Metab. 85: 3331–3337, 2000.

[135812]24179.

[135813]McAlpine, P. J.: Personal Communication. Winnipeg, Manitoba,

[135814]Canada 7/15/1992.

[135815]24180.

[135816]Monsoro–Burq, A.–H.; le Douarin, N. M.: BMP4 plays a key role

[135817]in left-right patterning in chick embryos by maintaining
Sonic Hedgehog

[135818]asymmetry. *Molec. Cell* 7: 789-799, 2001.

[135819]24181.

[135820]Shafritz, A. B.; Shore, E. M.; Gannon, F. H.; Zasloff, M. A.;
Taub,

[135821]R.; Muenke, M.; Kaplan, F. S.: Overexpression of an os-
teogenic morphogen

[135822]in fibrodysplasia ossificans progressiva. *New Eng. J. Med.*
335:

[135823]555-561, 1996.

[135824]24182.

[135825]Tabas, J. A.; Hahn, G. V.; Cohen, R. B.; Seaunez, H. N.;
Modi,

[135826]W. S.; Wozney, J. M.; Zasloff, M.; Kaplan, F. S.: Chromoso-
mal assignment

[135827]of the human gene for bone morphogenetic protein 4.
Clin. Orthop.

[135828]*Rel. Res.* 293: 310-316, 1993.

[135829]24183.

[135830]Tucker, A. S.; Matthews, K. L.; Sharpe, P. T.: Transforma-

tion

[135831] of tooth type induced by inhibition of BMP signaling. *Science* 282:

[135832] 1136–1138, 1998.

[135833] 24184.

[135834] van den Wijngaard, A.; Olde Weghuis, D.; Boersma, C. J. C.; van

[135835] Zoelen, E. J. J.; Geurts van Kessel, A.; Olijve, W.: Fine mapping

[135836] of the human bone morphogenetic protein–4 gene (BMP4) to chromosome

[135837] 14q22–q23 by in situ hybridization. *Genomics* 27: 559–560, 1995.

[135838] 24185.

[135839] van den Wijngaard, A.; van Kraay, M.; van Zoelen, E. J. J.; Olijve,

[135840] W.; Boersma, C. J. C.: Genomic organization of the human bone morphogenetic

[135841] protein–4 gene: molecular basis for multiple transcripts. *Biochem.*

[135842] *Biophys. Res. Commun.* 219: 789–794, 1996.

[135843] 24186.

- [135844]Amano, S.; Scott, I. C.; Takahara, K.; Koch, M.; Champli-
aud, M.-F.;
- [135845]Gerecke, D. R.; Keene, D. R.; Hudson, D. L.; Nishiyama, T.;
Lee, S.;
- [135846]Greenspan, D. S.; Burgeson, R. E.: Bone morphogenetic
protein 1 is
- [135847]an extracellular processing enzyme of the laminin 5
gamma-2 chain. J.
- [135848]Biol. Chem. 275: 22728-22735, 2000.
- [135849]24187.
- [135850]Bond, J. S.; Beynon, R. J.: The astacin family of metalloen-
dopeptidases. Protein
- [135851]Sci. 4: 1247-1261, 1995.
- [135852]24188.
- [135853]Ceci, J. D.; Kingsley, D. M.; Silan, C. M.; Copeland, N. G.;
- Jenkins,
- [135854]N. A.: An interspecific backcross linkage map of the prox-
imal half
- [135855]of mouse chromosome 14. Genomics 6: 673-678, 1990.
- [135856]24189.
- [135857]Kessler, E.; Takahara, K.; Biniaminov, L.; Brusel, M.;

Greenspan,

[135858]D.: Bone morphogenic protein-1: the type I procollagen
C-proteinase. Science 271:

[135859]360-362, 1996.

[135860]24190.

[135861]Kremmidiotis, G.; Baker, E.; Crawford, J.; Eyre, H. J.; Nah-
mias,

[135862]J.; Callen, D. F.: Localization of human cadherin genes to
chromosome

[135863]regions exhibiting cancer-related loss of heterozygosity.
Genomics 49:

[135864]467-471, 1998.

[135865]24191.

[135866]Herrenknecht, K.; Ozawa, M.; Eckerskorn, C.; Lottspeich,
F.; Lenter,

[135867]M.; Kemler, R.: The uvomorulin-anchorage protein alpha-
catenin is

[135868]a vinculin homologue. Proc. Nat. Acad. Sci. 88:
9156-9160, 1991.

[135869]24192.

[135870]Blomback, M.; Blomback, B.; Mammen, E. F.; Prasad, A. S.:

Fibrinogen

[135871]Detroit--a molecular defect in the N-terminal disulphide knot of human

[135872]fibrinogen? Nature 218: 134-137, 1968.

[135873]24193.

[135874]Magnuson, V. L.; McCombs, J. L.; Lee, C.-C.; Yang, F.; Bowman,

[135875]B. H.; McGill, J. R.: Human alpha-2-HS-glycoprotein localized to

[135876]3q27-q29 by in situ hybridization. Cytogenet. Cell Genet. 47: 72-74,

[135877]1988.

[135878]24194.

[135879]Hromas, R.; Collins, S.; Raskind, W.; Deaven, L.; Kaushansky, K.

[135880]: Hem-1, a potential membrane protein, with expression restricted

[135881]to blood cells. Biochim. Biophys. Acta 1090: 241-244, 1991.

[135882]24195.

[135883]Chen, Y.-H.; Lin, S.-J.; Lin, M.-W.; Tsai, H.-L.; Kuo, S.-S.;

Chen,

[135884]J.-W.; Charng, M.-J.; Wu, T.-C.; Chen, L.-C.; Ding, P. Y.-A.;

Pan,

[135885]W.-H.; Jou, Y.-S.; Chau, L.-Y.: Microsatellite polymorphism

in promoter

[135886]of heme oxygenase-1 gene is associated with susceptibil-

ity to coronary

[135887]artery disease in type 2 diabetic patients. Hum. Genet.

111: 1-8,

[135888]2002.

[135889]24196.

[135890]Fujita, T.; Toda, K.; Karimova, A.; Yan, S.-F.; Naka, Y.; Yet,

[135891]S.-F.; Pinsky, D. J.: Paradoxical rescue from ischemic lung

injury

[135892]by inhaled carbon monoxide driven by derepression of

fibrinolysis. Nature

[135893]Med. 7: 598-604, 2001.

[135894]24197.

[135895]Kutty, R. K.; Kutty, G.; Rodriguez, I. R.; Chader, G. J.; Wig-

gart,

[135896]B.: Chromosomal localization of the human oxygenase

genes: heme oxygenase-1

[135897](HMOX1) maps to chromosome 22q12 and heme oxygenase-2 (HMOX2) maps

[135898]to chromosome 16p13.3. Genomics 20: 513-516, 1994.

[135899]24198.

[135900]Poss, K. D.; Tonegawa, S.: Reduced stress defense in heme oxygenase

[135901]1-deficient cells. Proc. Nat. Acad. Sci. 94: 10925-10930, 1997.

[135902]24199.

[135903]Saito-Ohara, F.; Ikeuchi, T.; Matsumoto, M.; Kurata, S.:
Assignment

[135904]of the mouse heme oxygenase genes: heme oxygenase-1 (Hmox1) to chromosome

[135905]10 band C1 and heme oxygenase-2 (Hmox2) to chromosome 16 band B1. Cytogenet.

[135906]Cell Genet. 77: 180-181, 1997.

[135907]24200.

[135908]Seroussi, E.; Kedra, D.; Kost-Alimova, M.; Sandberg-Nordqvist,

[135909]A.-C.; Fransson, I.; Jacobs, J. F. M.; Fu, Y.; Pan, H.-Q.; Roe,

B.

- [135910]A.; Imreh, S.; Dumanski, J. P.: TOM1 genes map to human chromosome 22q13.1 and mouse chromosome 8C1 and encode proteins similar to the endosomal proteins HGS and STAM. Genomics 57: 380–388, 1999.
- [135913]24201.
- [135914]Yachie, A.; Niida, Y.; Wada, T.; Igarashi, N.; Kaneda, H.; Toma, T.; Ohta, K.; Kasahara, Y.; Koizumi, S.: Oxidative stress causes enhanced endothelial cell injury in human heme oxygenase–1 deficiency. J. Clin. Invest. 103: 129–135, 1999.
- [135918]24202.
- [135919]Yamada, N.; Yamaya, M.; Okinaga, S.; Nakayama, K.; Sekizawa, K.; Shibahara, S.; Sasaki, H.: Microsatellite polymorphism in the heme oxygenase–1 gene promoter is associated with susceptibility to emphysema. Am.

[135922]J. Hum. Genet. 66: 187–195, 2000. Note: Erratum: Am. J.

Hum. Genet.

[135923]68: 1542 only, 2001.

[135924]24203.

[135925]Yoshida, T.; Biro, P.; Cohen, T.; Muller, R. M.; Shibahara, S.

[135926]: Human heme oxygenase cDNA and induction of its
mRNA by hemin. Europ.

[135927]J. Biochem. 171: 457–461, 1988.

[135928]24204.

[135929]Stevenson, R. E.; Schroer, R. J.; Taylor, H. A.; Compton, J.
D.;

[135930]Livingston, R. E., III: Dislocated lens, dolichostenomelia,
and joint

[135931]stiffness. Proc. Greenwood Genet. Center 1: 16–22, 1982.

[135932]24205.

[135933]Celli, J.; Duijf, P.; Hamel, B. C. J.; Bamshad, M.; Kramer, B.;

[135934]Smits, A. P. T.; Newbury-Ecob, R.; Hennekam, R. C. M.;

Van Buggenhout,

[135935]G.; van Haeringen, A.; Woods, C. G.; van Essen, A. J.; de
Waal, R.;

- [135936]Vriend, G.; Haber, D. A.; Yang, A.; McKeon, F.; Brunner, H. G.; van Bokhoven, H.: Heterozygous germline mutations in the p53 homolog p63 are the cause of EEC syndrome. Cell 99: 143–153, 1999.
- [135939]24206.
- [135940]O'Farrell, P. H.: High resolution two-dimensional electrophoresis of proteins. J. Biol. Chem. 250: 4007–4021, 1975.
- [135942]24207.
- [135943]Osawa, M.; Umetsu, K.; Ohki, T.; Nagasawa, T.; Suzuki, T.; Takeichi, S.: Molecular evidence for human alpha2-HS glycoprotein (AHSG) polymorphism. Hum. Genet. 99: 18–21, 1997.
- [135946]24208.
- [135947]Pedersen, K. O.: Fetuin, a new globulin isolated from serum.
- [135948](Letter) Nature 154: 575, 1944.
- [135949]24209.

- [135950]Rizzu, P.; Baldini, A.: Three members of the human cystatin gene
- [135951]superfamily, AHSG, HRG, and KNG, map within one megabase of genomic
- [135952]DNA at 3q27. Cytogenet. Cell Genet. 70: 26–28, 1995.
- [135953]24210.
- [135954]Terkelsen, O. B. F.; Jahnen–Dechent, W.; Nielsen, H.; Moos, T.;
- [135955]Fink, E.; Nawratil, P.; Muller–Esterl, W.; Mollgard, K.: Rat fetuin:
- [135956]distribution of protein and mRNA in embryonic and neonatal rat tissues. Anat.
- [135957]Embryol. 197: 125–133, 1998.
- [135958]24211.
- [135959]Umetsu, K.; Kashimura, S.; Ikeda, N.; Suzuki, T.: A new alpha-2-HS-glycoprotein
- [135960]typing by isoelectric focusing. Hum. Genet. 67: 70–71, 1984.
- [135961]24212.
- [135962]Umetsu, K.; Kashimura, S.; Ikeda, N.; Suzuki, T.: A new alpha-2-HS-glycoprotein

[135963]allele (AHS*5*) in two Japanese families. Hum. Genet. 68:
264–265,

[135964]1984.

[135965]24213.

[135966]Umetsu, K.; Yuasa, I.; Ikuta, M.; Suzuki, T.: Alpha-
2-HS-glycoprotein

[135967](A2HS) polymorphism in a Japanese population: existence
of two new

[135968]variants. Jpn. J. Hum. Genet. 32: 263–266, 1987.

[135969]24214.

[135970]Umetsu, K.; Yuasa, I.; Suzuki, T.: The polymorphism of
desialyzed

[135971]alpha-2-HS-glycoprotein (AHS): isoelectric focusing in
2.5M urea as

[135972]a method for identification of genetic variants. Hum.
Genet. 73:

[135973]372–373, 1986.

[135974]24215.

[135975]Yoshioka, Y.; Gejyo, F.; Marti, T.; Rickli, E. E.; Burgi, W.;

[135976]Offner, G. D.; Troxler, R. F.; Schmid, K.: The complete
amino acid

[135977]sequence of the A-chain of human plasma alpha-
2HS-glycoprotein. J.

[135978]Biol. Chem. 261: 1665-1676, 1986.

[135979]24216.

[135980]Zelinski, T.; Kaita, H.; Lewis, M.; Coghlan, G.; Craig, D.:
The

[135981]sequence of chromosome 3 loci AHS:TF:CHE1. Hum.
Hered. 37: 1-6,

[135982]1987.

[135983]24217.

[135984]Cleve, H.: Genetic studies on the deficiency of beta-
2-glycoprotein

[135985]I of human serum. Humangenetik 5: 294-304, 1968.

[135986]24218.

[135987]Cleve, H.; Rittner, C.: Further family studies on the genetic

[135988]control of beta-2-glycoprotein I concentration in human
serum. Humangenetik 7:

[135989]93-97, 1969.

[135990]24219.

[135991]Eiberg, H.; Mohr, J.; Nielsen, L. S.: The beta-

2-glycoprotein

[135992]I (BG): allele frequencies and linkage relationships.

(Abstract) Cytogenet.

[135993]Cell Genet. 37: 462 only, 1984.

[135994]24220.

[135995]Eiberg, H.; Nielsen, L. S.; Mohr, J.: Exclusion mapping of apolipoprotein

[135996]H (APOH) and relationship between electrophoretic and quantitative

[135997]polymorphism. (Abstract) Cytogenet. Cell Genet. 51: 994 only, 1989.

[135998]24221.

[135999]Haagerup, A.; Kristensen, T.; Kruse, T. A.: Polymorphism and genetic

[136000]mapping of the gene encoding human beta-2-glycoprotein I to chromosome

[136001]17. (Abstract) Cytogenet. Cell Genet. 58: 2005 only, 1991.

[136002]24222.

[136003]Haupt, H.; Schwick, H. G.; Storiko, K.: Ueber einen erblichen

[136004]beta-2-Glykoprotein I-Mangel. Humangenetik 5:

291-293, 1968.

[136005]24223.

[136006]Hoeg, J. M.; Segal, P.; Gregg, R. E.; Chang, Y. S.; Lindgren, F.

[136007]T.; Adamson, G. L.; Frank, M.; Brickman, C.; Brewer, H. B., Jr.:

[136008]Characterization of plasma lipids and lipoproteins in patients with

[136009]beta-2-glycoprotein I (apolipoprotein H) deficiency. Atherosclerosis 55:

[136010]25-34, 1985.

[136011]24224.

[136012]Kamboh, M. I.; Ferrell, R. E.; Sepehrnia, B.: Genetic studies of human apolipoproteins. IV. Structural heterogeneity of apolipoprotein

[136014]H (beta-2-glycoprotein I). Am. J. Hum. Genet. 42: 452-457, 1988.

[136015]24225.

[136016]Koppe, A. L.; Walter, H.; Chopra, V. P.; Bajatzadeh, M.: Investigations

[136017]on the genetics and population genetics of the beta-

2-glycoprotein

[136018] polymorphism. Humangenetik 9: 164–171, 1970.

[136019] 24226.

[136020] Lozier, J.; Takahashi, N.; Putnam, F. W.: Complete amino acid

[136021] sequence of human plasma beta-2-glycoprotein I. Proc. Nat. Acad.

[136022] Sci. 81: 3640–3644, 1984.

[136023] 24227.

[136024] Mehdi, H.; Nunn, M.; Steel, D. M.; Whitehead, A. S.; Perez, M.;

[136025] Walker, L.; Peeples, M. E.: Nucleotide sequence and expression of

[136026] the human gene encoding apolipoprotein H (beta-2-glycoprotein I). Gene 108:

[136027] 293–298, 1991.

[136028] 24228.

[136029] Nakaya, Y.; Schaefer, E. J.; Brewer, H. B.: Activation of human

[136030] post-heparin lipase by apolipoprotein H (beta-2-glycoprotein I). Biochem.

[136031]Biophys. Res. Commun. 95: 1168–1172, 1980.

[136032]24229.

[136033]Nonaka, M.; Matsuda, Y.; Shiroishi, T.; Moriwaki, K.; Nonaka,

[136034]M.; Natsuume–Sakai, S.: Molecular cloning of mouse beta-2-glycoprotein

[136035]and mapping of the gene to chromosome 11. Genomics 13: 1082–1087,

[136036]1992.

[136037]24230.

[136038]Rahimi, A. G.; Goedde, H. W.; Flatz, G.; Kaifie, S.; Benkmann,

[136039]H.–G.; Delbruck, H.: Serum protein polymorphisms in four populations

[136040]of Afghanistan. Am. J. Hum. Genet. 29: 356–360, 1977.

[136041]24231.

[136042]Richter, A.; Cleve, H.: Genetic variations of human serum beta-2-glycoprotein

[136043]demonstrated by isoelectric focusing. Electrophoresis 9: 317–322,

[136044]1988.

[136045]24232.

[136046]Sanghera, D. K.; Nestlerode, C. S.; Ferrell, R. E.; Kamboh, M.

[136047]I.: Chimpanzee apolipoprotein H (beta-2-glycoprotein I): report on

[136048]the gene structure, a common polymorphism, and a high prevalence of

[136049]antiphospholipid antibodies. Hum. Genet. 109: 63-72, 2001.

[136050]24233.

[136051]Sanghera, D. K.; Wagenknecht, D. R.; McIntyre, J. A.; Kamboh,

[136052]M. I.: Identification of structural mutations in the fifth domain

[136053]of apolipoprotein H (beta-2-glycoprotein I) which affect phospholipid

[136054]binding. Hum. Molec. Genet. 6: 311-316, 1997.

[136055]24234.

[136056]Sepehrnia, B.; Kamboh, M. I.; Adams-Campbell, L. L.; Bunker, C.

[136057]H.; Nwankwo, M.; Majumder, P. P.; Ferrell, R. E.: Genetic studies

[136058] of human apolipoproteins. VIII. Role of the apolipoprotein
H polymorphism

[136059] in relation to serum lipoprotein concentrations. Hum.
Genet. 82:

[136060] 118–122, 1989.

[136061] 24235.

[136062] Kost, G. J.; Verity, A.: A new variant of late-onset
myophosphorylase

[136063] deficiency. Muscle Nerve 3: 195–201, 1980.

[136064] 24236.

[136065] Rosenmann, A.; Scriver, C. R.; Clow, C. L.; Levy, H. L.: His-
tidinaemia.

[136066] Part II: Impact; a retrospective study. J. Inherit. Metab. Dis.
6:

[136067] 54–57, 1983.

[136068] 24237.

[136069] Johnson, J. L.; Coyne, K. E.; Rajagopalan, K. V.; Van Hove,
J.

[136070] L. K.; Mackay, M.; Pitt, J.; Boneh, A.: Molybdopterin syn-
thase mutations

[136071] in a mild case of molybdenum cofactor deficiency. Am. J.

Med. Genet. 104:

[136072]169–173, 2001.

[136073]24238.

[136074]Johnson, J. L.; Wuebbens, M. M.; Mandell, R.; Shih, V. E.:

Molybdenum

[136075]cofactor biosynthesis in humans: identification of two
complementation

[136076]groups of cofactor-deficient patients and preliminary
characterization

[136077]of a diffusible molybdopterin precursor. J. Clin. Invest. 83:
897–903,

[136078]1989.

[136079]24239.

[136080]Reiss, J.; Christensen, E.; Dorche, C.: Molybdenum cofac-
tor deficiency:

[136081]first prenatal genetic analysis. Prenatal Diag. 19:
386–388, 1999.

[136082]24240.

[136083]Reiss, J.; Christensen, E.; Kurlemann, G.; Zabet, M.-T.;
Dorche,

[136084]C.: Genomic structure and mutational spectrum of the bi-

cistronic

[136085]MOCS1 gene defective in molybdenum cofactor deficiency
type A. Hum.

[136086]Genet. 103: 639–644, 1998.

[136087]24241.

[136088]Reiss, J.; Cohen, N.; Dorche, C.; Mandel, H.; Mendel, R. R.;
Stallmeyer,

[136089]B.; Zobot, M.–T.; Dierks, T.: Mutations in a polycistronic
nuclear

[136090]gene associated with molybdenum cofactor deficiency.
Nature Genet. 20:

[136091]51–53, 1998.

[136092]24242.

[136093]Reiss, J.; Dorche, B.; Stallmeyer, B.; Mendel, R. R.; Cohen,
N.;

[136094]Zobot, M. T.: Human molybdopterin synthase gene: ge-
nomic structure

[136095]and mutations in molybdenum cofactor deficiency type B.
Am. J. Hum.

[136096]Genet. 64: 706–711, 1999.

[136097]24243.

- [136098]Inui, K.; Emmett, M.; Wenger, D. A.: Immunological evidence for
deficiency in an activator protein for sulfatide sulfatase in
a variant
form of metachromatic leukodystrophy. Proc. Nat. Acad.
Sci. 80:
3074–3077, 1983.
- [136102]24244.
- [136103]Austin, J.; Armstrong, D.; Fouch, S.; Mitchell, C.; Stumpf,
D.
A.; Shearer, L.; Briner, O.: Metachromatic leukodystrophy
(MLD).
VIII. MLD in adults: diagnosis and pathogenesis. Arch.
Neurol. 18:
225–240, 1968.
- [136107]24245.
- [136108]Austin, J.; McAfee, D.; Armstrong, D.; O'Rourke, M.;
Shearer, L.;
Bachhawat, B. K.: Abnormal sulphatase activities in two
human diseases
(metachromatic leukodystrophy and gargoylism).
Biochem. J. 93: 15C–17C,

[136111]1964.

[136112]24246.

[136113]Austin, J.; McAfee, D.; Shearer, L.: Metachromatic form of diffuse

[136114]cerebral sclerosis. IV. Low sulfatase activity in the urine of nine

[136115]living patients with metachromatic leukodystrophy (MLD). Arch. Neurol. 12:

[136116]447-455, 1965.

[136117]24247.

[136118]Austin, J. H.: Some recent findings in leukodystrophies and in

[136119]gargoylism.In: Aronson, S. M.; Volk, B. W.: Inborn Disorders of Sphingolipid

[136120]Metabolism. Oxford: Pergamon Press (pub.) 1967. Pp. 359-387.

[136121]24248.

[136122]Austin, J. H.: Metachromatic form of diffuse cerebral sclerosis.

[136123]III. Significance of sulfatide and other lipid abnormalities in white

[136124]matter and kidney. Neurology 10: 470–483, 1960.

[136125]24249.

[136126]Baldinger, S.; Pierpont, M. E.; Wenger, D. A.: Pseudodeficiency

[136127]of arylsulfatase A: a counseling dilemma. Clin. Genet. 31: 70–76,

[136128]1987.

[136129]24250.

[136130]Barth, M. L.; Fensom, A.; Harris, A.: Missense mutations in the

[136131]aryl sulphatase A genes of metachromatic leukodystrophy patients. Hum.

[136132]Molec. Gen. 2: 2117–2121, 1993.

[136133]24251.

[136134]Barth, M. L.; Fensom, A.; Harris, A.: Identification of seven

[136135]novel mutations associated with metachromatic leukodystrophy. Hum.

[136136]Mutat. 6: 170–176, 1995.

[136137]24252.

[136138]Lebo, R. V.; Anderson, L. A.; DiMauro, S.; Lynch, E.;

Hwang, P.;

[136139]Fletcher, R.: Rare McArdle disease locus polymorphic site on 11q13

[136140]contains CpG sequence. Hum. Genet. 86: 17–24, 1990.

[136141]24253.

[136142]Lebo, R. V.; Gorin, F.; Fletcher, R. J.; Kao, F.-T.; Cheung,

[136143]M.-C.; Bruce, B. D.; Kan, Y. W.: High-resolution chromosome sorting

[136144]and DNA spot-blot analysis assign McArdle's syndrome to chromosome

[136145]11. Science 225: 57–59, 1984.

[136146]24254.

[136147]Lehoczky, T.; Halasy, M.; Simon, G.; Harnos, G.: Glycogenic myopathy:

[136148]a case of skeletal muscle-glycogenosis in twins. J. Neurol. Sci. 2:

[136149]366–384, 1965.

[136150]24255.

[136151]Mantle, D.; Lauffart, B.; Atack, J.; Lane, R. J. M.: Absence

[136152]of biochemical heterogeneity in McArdle's disease: a high

resolution

[136153]SDS–polyacrylamide gel electrophoresis study. J. Neurol. Sci. 78:

[136154]63–70, 1987.

[136155]24256.

[136156]Martin, M. A.; Rubio, J. C.; Garcia, A.; Fernandez, M. A.; Campos,

[136157]Y.; Krawczak, M.; Cooper, D. N.; Arenas, J.: Resolution of a mispaired

[136158]secondary structure intermediate could account for a novel micro–insertion/deletion

[136159](387 insA/del 8 bp) in the PYGM gene causing McArdle's disease. Clin.

[136160]Genet. 59: 48–51, 2001.

[136161]24257.

[136162]Martinuzzi, A.; Vergani, L.; Carrozzo, R.; Fanin, M.; Bartoloni,

[136163]L.; Angelini, C.; Askanas, V.; Engel, W. K.: Expression of muscle–type

[136164]phosphorylase in innervated and aneural cultured muscle of patients

[136165]with myophosphorylase deficiency. J. Clin. Invest. 92:

1774–1780,

[136166]1993.

[136167]24258.

[136168]McArdle, B.: Myopathy due to a defect in muscle glycogen breakdown. Clin.

[136169]Sci. 10: 13–33, 1951.

[136170]24259.

[136171]Mellick, R. S.; Mahler, R. F.; Hughes, B. P.: McArdle's syndrome.

[136172]Phosphorylase-deficient myopathy. Lancet I: 1045–1048, 1962.

[136173]24260.

[136174]Miranda, A. F.; Nette, E. G.; Hartlage, P. L.; DiMauro, S.: Phosphorylase

[136175]isoenzymes in normal and myophosphorylase-deficient human heart. Neurology 29:

[136176]1538–1541, 1979.

[136177]24261.

[136178]Papadimitriou, A.; Manta, P.; Divari, R.; Karabetsos, A.; Papadimitriou,

[136179]E.; Bresolin, N.: McArdle's disease: two clinical expressions

in

[136180]the same pedigree. J. Neurol. 237: 267–270, 1990.

[136181]24262.

[136182]Pearson, C. M.; Rimer, D. G.; Mommaerts, W. F. H. M.: A
metabolic

[136183]myopathy due to absence of muscle phosphorylase. Am. J.
Med. 30:

[136184]502–517, 1961.

[136185]24263.

[136186]Ross, B. D.; Radda, G. K.; Gadian, D. G.; Rocker, G.; Esiri,
M.;

[136187]Falconer–Smith, J.: Examination of a case of suspected
McArdle's

[136188]syndrome by (31)P nuclear magnetic resonance. New Eng.
J. Med. 304:

[136189]1338–1342, 1981.

[136190]24264.

[136191]Rowland, L. P.; Lovelace, R. E.; Schotland, D. L.; Araki, S.;

[136192]Carmel, P.: The clinical diagnosis of McArdle's disease:
identification

[136193]of another family with deficiency of muscle phosphory–

lase. Neurology 16:

[136194]93–100, 1966.

[136195]24265.

[136196]Sarova–Pinhas, I.; Sadeh, M.: McArdle disease in a Druze family. Israel

[136197]J. Med. Sci. 25: 64–68, 1989.

[136198]24266.

[136199]Schmid, R.; Hammaker, L.: Hereditary absence of muscle phosphorylase

[136200](McArdle's syndrome). New Eng. J. Med. 264: 223–225, 1961.

[136201]24267.

[136202]Schmid, R.; Mahler, R.: Chronic progressive myopathy with myoglobinuria:

[136203]demonstration of a glycogenolytic defect in the muscle. J. Clin.

[136204]Invest. 38: 2044–2058, 1959.

[136205]24268.

[136206]Servidei, S.; Shanske, S.; Zeviani, M.; Lebo, R.; Fletterick,

[136207]R.; DiMauro, S.: McArdle's disease: biochemical and

molecular genetic

[136208] studies. Ann. Neurol. 24: 774-781, 1988.

[136209] 24269.

[136210] Slonim, A. E.; Goans, P. J.: Myopathy in McArdle's syndrome:

[136211] improvement with a high-protein diet. New Eng. J. Med. 312: 355-359,

[136212] 1985.

[136213] 24270.

[136214] Tabata, T.; Kikunami, K.; Matsushita, Y.; Inoue, T.; Inoue, T.;

[136215] Okamoto, T.; Kono, N.; Takahashi, M.; Tarui, S.; Morii, H.: Acute

[136216] renal failure in McArdle's disease. Nephron 44: 371-374, 1986.

[136217] 24271.

[136218] Tsujino, S.; Rubin, L. A.; Shanske, S.; DiMauro, S.: An A-to-C

[136219] substitution involving the translation initiation codon in a patient

[136220] with myophosphorylase deficiency (McArdle's disease).

Hum. Mutat. 4:

[136221]73–75, 1994.

[136222]24272.

[136223]Tsuchino, S.; Shanske, S.; DiMauro, S.: Molecular genetic heterogeneity

[136224]of myophosphorylase deficiency (McArdle's disease). New Eng. J. Med. 329:

[136225]241–245, 1993.

[136226]24273.

[136227]Tsuchino, S.; Shanske, S.; Martinuzzi, A.; Heiman–Patterson, T.;

[136228]DiMauro, S.: Two novel missense mutations (E654K, L396P) in Caucasian

[136229]patients with myophosphorylase deficiency (McArdle's disease). Hum.

[136230]Mutat. 6: 276–277, 1995.

[136231]24274.

[136232]Tsuchino, S.; Shanske, S.; Nonaka, I.; Eto, Y.; Mendell, J. R.;

[136233]Fenichel, G. M.; DiMauro, S.: Three new mutations in patients with

[136234]myophosphorylase deficiency (McArdle disease). Am. J.

Hum. Genet. 54:

[136235]44–52, 1994.

[136236]24275.

[136237]Vorgerd, M.; Kubisch, C.; Burwinkel, B.; Reichmann, H.;
Mortier,

[136238]W.; Tettenborn, B.; Pongratz, D.; Lindemuth, R.; Tegen-
thoff, M.; Malin,

[136239]J.-P.; Kilimann, M. W.: Mutation analysis in myophospho-
rylase deficiency

[136240](McArdle's disease). Ann. Neurol. 43: 326–331, 1998.

[136241]24276.

[136242]Hers, H. G.: Etudes enzymatiques sur fragments hepa-
tiques: application

[136243]a la classification des glycogenoses. Rev. Int. Hepat. 9:
35–55,

[136244]1959.

[136245]24277.

[136246]Scriver, C. R.; Levy, H. L.: Histidinaemia. Part I: Reconciling

[136247]retrospective and prospective findings. J. Inherit. Metab.
Dis. 6:

[136248]51–53, 1983.

[136249]24278.

[136250]Suchi, M.; Harada, N.; Wada, Y.; Takagi, Y.: Molecular cloning

[136251]of a cDNA encoding human histidase. Biochim. Biophys. Acta 1216:

[136252]293–295, 1993.

[136253]24279.

[136254]Suchi, M.; Sano, H.; Mizuno, H.; Wada, Y.: Molecular cloning

[136255]and structural characterization of the human histidase gene (HAL). Genomics 29:

[136256]98–104, 1995.

[136257]24280.

[136258]Taylor, R. G.; Garcia-Heras, J.; Sadler, S. J.; Lafreniere, R.

[136259]G.; Willard, H. F.; Ledbetter, D. H.; McInnes, R. R.: Localization

[136260]of histidase to human chromosome region 12q22–q24.1 and mouse chromosome

[136261]region 10C2–D1. Cytogenet. Cell Genet. 56: 178–181, 1991.

[136262]24281.

[136263]Taylor, R. G.; Garcia-Heras, J.; Sexsmith, E.; Lambert, M.;
Ledbetter,
[136264]D.; McInnes, R. R.: Molecular characterization of histidase
cDNAs,
[136265]and sublocalization of the human histidase gene to
12q22-23, and the
[136266]mouse gene to 10C2-D1.(Abstract) Am. J. Hum. Genet. 45
(suppl.):
[136267]A164, 1989.
[136268]24282.

[136269]Taylor, R. G.; Grieco, D.; Clarke, G. A.; McInnes, R. R.; Tay-
lor,
[136270]B. A.: Identification of the mutation in murine histidinemia
(his)
[136271]and genetic mapping of the murine histidase locus (Hal)
on chromosome
[136272]10. Genomics 16: 231-240, 1993.
[136273]24283.

[136274]Wadman, S. K.; Van Sprang, F. J.; Van Stekelenburg, G. J.;
de
[136275]Bree, P. K.: Three new cases of histidinemia: clinical and
biochemical

[136276]data. Acta Paediat. Scand. 56: 485–492, 1967.

[136277]24284.

[136278]Woody, N. C.; Snyder, C. H.; Harris, J. A.: Histidinemia.
Am.

[136279]J. Dis. Child. 110: 606–613, 1965.

[136280]24285.

[136281]Martin–Subero, J. I.; Gesk, S.; Harder, L.; Sonoki, T.;
Tucker,

[136282]P. W.; Schlegelberger, B.; Grote, W.; Novo, F. J.; Calasanz,
M. J.;

[136283]Hansmann, M. L.; Dyer, M. J. S.; Siebert, R.: Recurrent in-
volvement

[136284]of the REL and BCL11A loci in classical Hodgkin lym-
phoma. Blood 99:

[136285]1474–1477, 2002.

[136286]24286.

[136287]Hankinson, S. E.; Willett, W. C.; Colditz, G. A.; Hunter, D.
J.;

[136288]Michaud, D. S.; Deroo, B.; Rosner, B.; Speizer, F. E.; Pollak,
M.

[136289]: Circulating concentrations of insulin–like growth factor–I

and risk

[136290]of breast cancer. Lancet 351: 1393–1396, 1998.

[136291]24287.

[136292]Esumi, N.; Budarf, M.; Ciccarelli, L.; Sellinger, B.; Kozak, C.

[136293]A.; Wistow, G.: Conserved gene structure and genomic
linkage for

[136294]D-dopachrome tautomerase (DDT) and MIF. Mammalian
Genome 9: 753–757,

[136295]1998.

[136296]24288.

[136297]Roger, T.; David, J.; Glauser, M. P.; Calandra, T.: MIF regu-
lates

[136298]innate immune response through modulation of Toll-like
receptor 4. Nature 414:

[136299]920–924, 2001.

[136300]24289.

[136301]Emi, M.; Asaoka, H.; Matsumoto, A.; Itakura, H.; Kurihara,
Y.;

[136302]Wada, Y.; Kanamori, H.; Yazaki, Y.; Takahashi, E.; Lepert,
M.; Lalouel,

[136303]J.–M.; Kodama, T.; Mukai, T.: Structure, organization, and

chromosomal

[136304]mapping of the human macrophage scavenger receptor
gene. J. Biol.

[136305]Chem. 268: 2120–2125, 1993.

[136306]24290.

[136307]Latil, A.; Lidereau, R.: Genetic aspects of prostate cancer.
Virchows

[136308]Arch. 432: 389–406, 1998.

[136309]24291.

[136310]Matsumoto, A.; Naito, M.; Itakura, H.; Ikemoto, S.; Asaoka,
H.;

[136311]Hayakawa, I.; Kanamori, H.; Aburatani, H.; Takaku, F.;

Suzuki, H.;

[136312]Kobari, Y.; Miyai, T.; Takahashi, K.; Cohen, E. H.; Wydro,
R.; Housman,

[136313]D. E.; Kodama, T.: Human macrophage scavenger recep-
tors: primary

[136314]structure, expression, and localization in atherosclerotic
lesions. Proc.

[136315]Nat. Acad. Sci. 87: 9133–9137, 1990.

[136316]24292.

[136317] Xu, J.; Zheng, S. L.; Hawkins, G. A.; Faith, D. A.; Kelly, B.;

[136318] Isaacs, S. D.; Wiley, K. E.; Chang, B.; Ewing, C. M.; Bu-
jnovszky,

[136319] P.; Carpten, J. D.; Bleecker, E. R.; Walsh, P. C.; Trent, J. M.;
Meyers,

[136320] D. A.; Isaacs, W. B.: Linkage and association studies of
prostate

[136321] cancer susceptibility: evidence for linkage at 8p22-23.
Am. J. Hum.

[136322] Genet. 69: 341-350, 2001.

[136323] 24293.

[136324] Xu, J.; Zheng, S. L.; Komiya, A.; Mychaleckyj, J. C.; Isaacs,
S.

[136325] D.; Hu, J. J.; Sterling, D.; Lange, E. M.; Hawkins, G. A.;
Turner,

[136326] A.; Ewing, C. M.; Faith, D. A.; and 19 others: Germline mu-
tations

[136327] and sequence variants of the macrophage scavenger re-
ceptor 1 gene

[136328] are associated with prostate cancer risk. Nature Genet. 32:
321-325,

[136329] 2002.

[136330]24294.

[136331]Holness, C. L.; Simmons, D. L.: Molecular cloning of CD68,
a human

[136332]macrophage marker related to lysosomal glycoproteins.
Blood 81:

[136333]1607-1613, 1993.

[136334]24295.

[136335]Jones, E.; Quinn, C. M.; See, C. G.; Montgomery, D. S.;
Ford, M.

[136336]J., Kolble, K.; Gordon, S.; Greaves, D. R.: The linked human
elongation

[136337]initiation factor 4A1 (EIF4A1) and CD68 genes map to
chromosome 17p13. Genomics 53:

[136338]248-250, 1998.

[136339]24296.

[136340]Chavrier, P.; Janssen-Timmen, U.; Mattei, M.-G.; Zerial,
M.; Bravo,

[136341]R.; Charnay, P.: Structure, chromosome location, and ex-
pression of

[136342]the mouse zinc finger gene Krox-20: multiple gene prod-
ucts and coregulation

[136343]with the proto-oncogene c-fos. Molec. Cell. Biol. 9:
787-797, 1989.

[136344]24297.

[136345]Le Beau, M. M.; Espinosa, R., III; Neuman, W. L.; Stock, W.;
Roulston,

[136346]D.; Larson, R. A.; Keinanen, M.; Westbrook, C. A.: Cytoge-
netic and

[136347]molecular delineation of the smallest commonly deleted
region of chromosome

[136348]5 in malignant myeloid diseases. Proc. Nat. Acad. Sci. 90:
5484-5488,

[136349]1993.

[136350]24298.

[136351]Barrow, L. L.; Simin, K.; Jones, J. M.; Lee, D. C.; Meisler, M.

[136352]H.: Conserved linkage of early growth response 4, annexin
4, and

[136353]transforming growth factor alpha on mouse chromosome
6. Genomics 19:

[136354]388-390, 1994.

[136355]24299.

[136356]Crosby, S. D.; Veile, R. A.; Donis-Keller, H.; Baraban, J. M.;

[136357]Bhat, R. V.; Simburger, K. S.; Milbrandt, J.: Neural-specific expression,
[136358]genomic structure, and chromosomal localization of the gene encoding
[136359]the zinc-finger transcription factor NGFI-C. Proc. Nat. Acad. Sci. 89:
[136360]4739-4743, 1992.
[136361]24300.

[136362]Boerkoel, C. F.; Takashima, H.; Bacino, C. A.; Daentl, D.; Lupski,
[136363]J. R.: EGR2 mutation R359W causes a spectrum of Dejerine-Sottas neuropathy. Neurogenetics 3:
[136364]153-157, 2001.
[136365]24301.

[136366]Willcutt, E. G.; Pennington, B. F.; Smith, S. D.; Cardon, L. R.;

[136367]Gayan, J.; Knopik, V. S.; Olson, R. K.; DeFries, J. C.: Quantitative
[136368]trait locus for reading disability on chromosome 6p is pleiotropic
[136369]for attention-deficit/hyperactivity disorder. Am. J. Med. Genet. 114:

[136370]260–268, 2002.

[136371]24302.

[136372]Koiwai, O.; Nishizawa, M.; Hasada, K.; Aono, S.; Adachi, Y.;
Mamiya,

[136373]N.; Sato, H.: Gilbert's syndrome is caused by a heterozy-
gous missense

[136374]mutation in the gene for bilirubin UDP-
glucuronosyltransferase. Hum.

[136375]Molec. Genet. 4: 1183–1186, 1995.

[136376]24303.

[136377]Maruo, Y.; Wada, S.; Yamamoto, K.; Sato, H.; Yamano, T.;
Shimada,

[136378]M.: A case of anorexia nervosa with hyperbilirubinaemia in
a patient

[136379]homozygous for a mutation in the bilirubin UDP-
glucuronosyltransferase

[136380]gene. Europ. J. Pediat. 158: 547–549, 1999.

[136381]24304.

[136382]Andres, D. A.; Milatovich, A.; Ozcelik, T.; Wenzlau, J. M.;
Brown,

[136383]M. S.; Goldstein, J. L.; Francke, U.: cDNA cloning of the two

subunits

[136384]of human CAAX farnesyltransferase and chromosomal
mapping of FNTA

[136385]and FNTB loci and related sequences. Genomics 18:
105–112, 1993.

[136386]24305.

[136387]Long, S. B.; Casey, P. J.; Beese, L. S.: Reaction path of pro-
tein

[136388]farnesyltransferase at atomic resolution. Nature 419:
645–650, 2002.

[136389]24306.

[136390]Roifman, C. M.; Zhang, J.; Chitayat, D.; Sharfe, N.: A partial

[136391]deficiency of interleukin-7R-alpha is sufficient to abro-
gate T-cell

[136392]development and cause severe combined immunodeficiency. Blood 96:

[136393]2803–2807, 2000.

[136394]24307.

[136395]Bowne, S. J.; Sullivan, L. S.; Blanton, S. H.; Cepko, C. L.;
Blackshaw,

[136396]S.; Birch, D. G.; Highbanks-Wheaton, D.; Heckenlively, J.

R.; Daiger,

[136397]S. P.: Mutations in the inosine monophosphate dehydrogenase 1 gene

[136398](IMPDH1) cause the RP10 form of autosomal dominant retinitis pigmentosa. Hum.

[136399]Molec. Genet. 11: 559-568, 2002.

[136400]24308.

[136401]Collart, F. R.; Huberman, E.: Cloning and sequence analysis of

[136402]the human and Chinese hamster inosine-5-prime-monophosphate dehydrogenase

[136403]cDNAs. J. Biol. Chem. 263: 15769-15772, 1988.

[136404]24309.

[136405]Doggett, N. A.; Callen, D. F.; Chen, Z. L.; Moore, S.; Tesmer,

[136406]J. G.; Duesing, L. A.; Stallings, R. L.: Identification and regional

[136407]localization of a human IMP dehydrogenase-like locus (IMPDHL1) at

[136408]16p13.13. Genomics 18: 687-689, 1993.

[136409]24310.

[136410]Gu, J. J.; Kaiser-Rogers, K.; Rao, K.; Mitchell, B. S.: Assignment
[136411]of the human type I IMP dehydrogenase gene (IMPDH1) to
[136412]7q31.3-q32. Genomics 24: 179-181, 1994.
[136413]24311.
[136414]Natsumeda, Y.; Ohno, S.; Kawasaki, H.; Konno, Y.; Weber,
G.; Suzuki,
[136415]K.: Two distinct cDNAs for human IMP dehydrogenase. J.
Biol. Chem. 265:
[136416]5292-5295, 1990.
[136417]24312.
[136418]Kennan, A.; Aherne, A.; Palfi, A.; Humphries, M.; McKee,
A.; Stitt,
[136419]A.; Simpson, D. A. C.; Demtroder, K.; Orntoft, T.; Ayuso,
C.; Kenna,
[136420]P. F.; Farrar, G. J.; Humphries, P.: Identification of an IM-
PDH1
[136421]mutation in autosomal dominant retinitis pigmentosa
(RP10) revealed
[136422]following comparative microarray analysis of transcripts
derived from

- [136423]retinas of wild-type and Rho-/- mice. Hum. Molec. Genet.
11: 547-558,
[136424]2002.
[136425]24313.
[136426]Crompton, M. R.; Moss, S. E.; Crompton, M. J.: Diversity in
the
[136427]lipocortin/calpactin family. Cell 55: 1-3, 1988.
[136428]24314.
[136429]Horlick, K. R.; Cheng, I. C.; Wong, W. T.; Wakeland, E. K.;
Nick,
[136430]H. S.: Mouse lipocortin I gene structure and chromosomal
assignment:
[136431]gene duplication and the origins of a gene family. Ge-
nomics 10:
[136432]365-374, 1991.
[136433]24315.
[136434]Huebner, K.; Cannizzaro, L. A.; Croce, C. M.; Frey, A. Z.;
Wallner,
[136435]B. P.; Hecht, B. K.; Hecht, F.: Chromosome localization of
the human
[136436]genes for lipocortin I and the lipocortin II family.

(Abstract) Cytogenet.

[136437]Cell Genet. 46: 631 only, 1987.

[136438]24316.

[136439]Huebner, K.; Cannizzaro, L. A.; Frey, A. Z.; Hecht, B. K.;
Hecht,

[136440]F.; Croce, C. M.; Wallner, B. P.: Chromosomal localization
of the

[136441]human genes for lipocortin I and lipocortin II. Oncogene
Res. 2:

[136442]299-310, 1988.

[136443]24317.

[136444]Pepinsky, R. B.; Tizard, R.; Mattaliano, R. J.; Sinclair, L. K.;

[136445]Miller, G. T.; Browning, J. L.; Chow, E. P.; Burne, C.; Huang,
K.-S.;

[136446]Pratt, D.; Wachter, L.; Hession, C.; Frey, A. Z.; Wallner, B.
P.:

[136447]Five distinct calcium and phospholipid binding proteins
share homology

[136448]with lipocortin I. J. Biol. Chem. 263: 10799-10811, 1988.

[136449]24318.

[136450]Shohat, M.; Korenberg, J. R.; Schwabe, A. D.; Rotter, J. I.:

Hypothesis:

[136451]familial Mediterranean fever--a genetic disorder of the lipocortin

[136452]family? Am. J. Med. Genet. 34: 163-167, 1989.

[136453]24319.

[136454]Wallner, B. P.; Mattaliano, R. J.; Hession, C.; Cate, R. L.; Tizard,

[136455]R.; Sinclair, L. K.; Foeller, C.; Chow, E. P.; Browning, J. L.; Ramachandran,

[136456]K. L.; Pepinsky, R. B.: Cloning and expression of human lipocortin,

[136457]a phospholipase A2 inhibitor with potential anti-inflammatory activity. Nature 320:

[136458]77-81, 1986.

[136459]24320.

[136460]Harman, S. M.; Metter, E. J.; Blackman, M. R.; Landis, P. K.;

[136461]Carter, H. B.: Serum levels of insulin-like growth factor I (IGF-I),

[136462]IGF-II, IGF-binding protein-3, and prostate-specific antigen as predictors

[136463]of clinical prostate cancer. J. Clin. Endocr. Metab. 85:

4258–4265,

[136464]2000.

[136465]24321.

[136466]Holly, J.: Insulin-like growth factor-I and new opportunities

[136467]for cancer prevention. *Lancet* 351: 1373–1375, 1998.

[136468]24322.

[136469]Hoppener, J. W. M.; de Pagter-Holthuisen, P.; Geurts van Kessel,

[136470]A. H. M.; Jansen, M.; Kittur, S. D.; Antonarakis, S. E.; Lips, C.

[136471]J. M.; Sussenbach, J. S.: The human gene encoding insulin-like growth

[136472]factor I is located on chromosome 12. *Hum. Genet.* 69: 157–160, 1985.

[136473]24323.

[136474]Le Bouc, Y.; Dreyer, D.; Jaeger, F.; Binoux, M.; Sondermeyer,

[136475]P.: Complete characterization of the human IGF-I nucleotide sequence

[136476]isolated from a newly constructed adult liver cDNA library.

FEBS

[136477]Lett. 196: 108–112, 1986.

[136478]24324.

[136479]Hellstrom, A.; Perruzzi, C.; Ju, M.; Engstrom, E.; Hard,
A.-L.;

[136480]Liu, J.-L.; Albertsson-Wikland, K.; Carlsson, B.; Niklasson,
A.; Sjodell,

[136481]L.; LeRoith, D.; Senger, D. R.; Smith, L. E. H.: Low IGF-I
suppresses

[136482]VEGF–survival signaling in retinal endothelial cells: direct
correlation

[136483]with clinical retinopathy of prematurity. Proc. Nat. Acad.
Sci. 98:

[136484]5804–5808, 2001.

[136485]24325.

[136486]Lembo, G.; Rockman, H. A.; Hunter, J. J.; Steinmetz, H.;
Koch,

[136487]W. J.; Ma, L.; Printz, M. P.; Ross, J., Jr.; Chien, K. R.; Powell–
Braxton,

[136488]L.: Elevated blood pressure and enhanced myocardial con-
tractility

[136489]in mice with severe IGF–1 deficiency. J. Clin. Invest. 98:

2648–2655,

[136490]1996.

[136491]24326.

[136492]Li, C. H.; Yamashiro, D.; Gospodarowicz, D.; Kaplan, S. L.;

Van

[136493]Vliet, G.: Total synthesis of insulin-like growth factor I

(somatomedin

[136494]C). Proc. Nat. Acad. Sci. 80: 2216–2220, 1983.

[136495]24327.

[136496]Liu, J.-P.; Baker, J.; Perkins, A. S.; Robertson, E. J.; Efstra-

tiadis,

[136497]A.: Mice carrying null mutations of the genes encoding in-

sulin-like

[136498]growth factor I (Igf-1) and type 1 IGF receptor (Igf1r). Cell

75:

[136499]59–72, 1993.

[136500]24328.

[136501]Mathews, L. S.; Norstedt, G.; Palmiter, R. D.: Regulation of

[136502]insulin-like growth factor I gene expression by growth

hormone. Proc.

[136503]Nat. Acad. Sci. 83: 9343–9347, 1986.

[136504]24329.

[136505]Morton, C.; Rall, L.; Bell, G.; Shows, T.: Human insulin-like

[136506]growth factor-1 (IGF1) is encoded at 12q22-q24.1, and
insulin-like

[136507]growth factor-2 (IGF2) is at 11p15. (Abstract) Cytogenet.
Cell Genet. 40:

[136508]703 only, 1985.

[136509]24330.

[136510]Musaro, A.; McCullagh, K.; Paul, A.; Houghton, L.; Do-
browolny,

[136511]G.; Molinaro, M.; Barton, E. R.; Sweeney, H. L.; Rosenthal,
N.: Localized

[136512]Igf-1 transgene expression sustains hypertrophy and re-
generation in

[136513]senescent skeletal muscle. Nature Genet. 27: 195-200,
2001.

[136514]24331.

[136515]Musaro, A.; McCullagh, K. J. A.; Naya, F. J.; Olson, E. N.;
Rosenthal,

[136516]N.: IGF-1 induces skeletal myocyte hypertrophy through
calcineurin

[136517]in association with GATA-2 and NF-ATc1. Nature 400:
581-585, 1999.

[136518]24332.

[136519]Playford, M. P.; Bicknell, D.; Bodmer, W. F.; Macaulay, V. M.

[136520]; Insulin-like growth factor 1 regulates the location, sta-
bility,

[136521]and transcriptional activity of beta-catenin. Proc. Nat.
Acad. Sci. 97:

[136522]12103-12108, 2000.

[136523]24333.

[136524]Powell-Braxton, L.; Hollingshead, P.; Warburton, C.; Dowd,
M.;

[136525]Pitts-Meek, S.; Dalton, D.; Gillett, N.; Stewart, T. A.: IGF-I
is

[136526]required for normal embryonic growth in mice. Genes
Dev. 7: 2609-2617,

[136527]1993.

[136528]24334.

[136529]Rapp, R.; Deger, A.; Blum, W.; Koch, R.; Weber, U.: Charac-
terization

[136530]of the protein which binds insulin-like growth factor in

human serum. Europ.

[136531]J. Biochem. 172: 421–425, 1988.

[136532]24335.

[136533]Rotwein, P.: Two insulin-like growth factor I messenger
RNAs

[136534]are expressed in human liver. Proc. Nat. Acad. Sci. 83:
77–81, 1986.

[136535]24336.

[136536]Schoenle, E. J.; Zenobi, P. D.; Torresani, T.; Werder, E. A.;

[136537]Zachmann, M.; Froesch, E. R.: Recombinant human in-
sulin-like growth

[136538]factor I (rhIGF I) reduces hyperglycaemia in patients with
extreme

[136539]insulin resistance. Diabetologia 34: 675–679, 1991.

[136540]24337.

[136541]Semsarian, C.; Wu, M.-J.; Ju, Y.-K.; Marciniak, T.; Yeoh, T.;

[136542]Allen, D. G.; Harvey, R. P.; Graham, R. M.: Skeletal muscle
hypertrophy

[136543]is mediated by a Ca^{2+} -dependent calcineurin signalling
pathway. Nature 400:

[136544]576–581, 1999.

[136545]24338.

[136546]Svoboda, M. E.; Van Wyk, J. J.; Klapper, D. G.; Fellows, R. E.;

[136547]Grissom, F. E.; Schleuter, R. J.: Purification of somatomedin-C from

[136548]human plasma: chemical and biological properties, partial sequence

[136549]analysis, and relationship to other somatomedins. Biochemistry 19:

[136550]790-797, 1980.

[136551]24339.

[136552]Taylor, B. A.; Grieco, D.: Localization of the gene encoding

[136553]insulin-like growth factor I on mouse chromosome 10. Cytogenet. Cell

[136554]Genet. 56: 57-58, 1991.

[136555]24340.

[136556]Tricoli, J. V.; Rall, L. B.; Scott, J.; Bell, G. I.; Shows, T.

[136557]B.: Insulin-like growth factor genes: chromosome organization and

[136558]association with disease. (Abstract) Am. J. Hum. Genet. 36: 121S

[136559]only, 1984.

[136560]24341.

[136561]Tricoli, J. V.; Rall, L. B.; Scott, J.; Bell, G. I.; Shows, T.

[136562]B.: Localization of insulin-like growth factor genes to human chromosomes

[136563]11 and 12. Nature 310: 784–786, 1984.

[136564]24342.

[136565]Ullrich, A.; Berman, C. H.; Dull, T. J.; Gray, A.; Lee, J. M.

[136566]: Isolation of the human insulin-like growth factor I gene using a

[136567]single synthetic DNA probe. EMBO J. 3: 361–364, 1984.

[136568]24343.

[136569]Gough, N. M.; Rakar, S.; Harpur, A.; Wilks, A. F.: Localization

[136570]of genes for two members of the JAK family of protein tyrosine kinases

[136571]to murine chromosomes 4 and 19. Mammalian Genome 6: 247–248, 1995.

[136572]24344.

[136573]Ihle, J. N.: Cytokine receptor signalling. Nature 377: 591–594,

[136574]1995.

[136575]24345.

- [136576]Pritchard, M. A.; Baker, E.; Callen, D. F.; Sutherland, G. R.;
[136577]Wilks, A. F.: Two members of the JAK family of protein ty-
rosine kinases
[136578]map to chromosomes 1p31.3 and 9p24. Mammalian
Genome 3: 36-38, 1992.
[136579]24346.
- [136580]Campbell, G. S.; Argetsinger, L. S.; Ihle, J. N.; Kelly, P. A.;
[136581]Rillema, J. A.; Carter-Su, C.: Activation of JAK2 tyrosine ki-
nase
[136582]by prolactin receptors in Nb2 cells and mouse mammary
gland explants. Proc.
[136583]Nat. Acad. Sci. 91: 5232-5236, 1994.
[136584]24347.
- [136585]Huang, L. J.; Constantinescu, S. N.; Lodish, H. F.: The N-
terminal
[136586]domain of Janus kinase 2 is required for Golgi processing
and cell
[136587]surface expression of erythropoietin receptor. Molec. Cell
8: 1327-1338,
[136588]2001.
[136589]24348.

[136590]Neubauer, H.; Cumano, A.; Muller, M.; Wu, H.; Huffstadt, U.; Pfeffer,

[136591]K.: Jak2 deficiency defines an essential developmental checkpoint

[136592]in definitive hematopoiesis. Cell 93: 397–409, 1998.

[136593]24349.

[136594]Parganas, E.; Wang, D.; Stravopodis, D.; Topham, D. J.; Marine,

[136595]J.–C.; Teglund, S.; Vanin, E. F.; Bodner, S.; Colamonici, O. R.; van

[136596]Deursen, J. M.; Grosveld, G.; Ihle, J. N.: Jak2 is essential for

[136597]signaling through a variety of cytokine receptors. Cell 93: 385–395,

[136598]1998.

[136599]24350.

[136600]Saltzman, A.; Stone, M.; Franks, C.; Searfoss, G.; Munro, R.; Jaye,

[136601]M.; Ivashchenko, Y.: Cloning and characterization of human Jak–2

[136602]kinase: high mRNA expression in immune cells and muscle tissue. Biochem.

[136603]Biophys. Res. Commun. 246: 627–633, 1998.

[136604]24351.

[136605]Schwaller, J.; Parganas, E.; Wang, D.; Cain, D.; Aster, J. C.;

[136606]Williams, I. R.; Lee, C.-K.; Gerthner, R.; Kitamura, T.;

Frantsve,

[136607]J.; Anastasiadou, E.; Loh, M. L.; Levy, D. E.; Ihle, J. N.;

Gilliland,

[136608]D. G.: Stat5 is essential for the myelo- and lymphoprolif-
erative

[136609]disease induced by TEL/JAK2. Molec. Cell 6: 693–704,
2000.

[136610]24352.

[136611]Watling, D.; Guschin, D.; Muller, M.; Silvennoinen, O.; Wit-
thuhn,

[136612]B. A.; Quelle, F. W.; Rogers, N. C.; Schindler, C.; Stark, G.
R.;

[136613]Ihle, J. N.; Kerr, I. M.: Complementation by the protein ty-
rosine

[136614]kinase JAK2 of a mutant cell line defective in the inter-
feron-gamma

[136615]signal transduction pathway. Nature 366: 166–170, 1993.

[136616]24353.

[136617]Cihakova, D.; Trebusak, K.; Heino, M.; Fadeyev, V.; Tiul-
pakov,

[136618]A.; Battelino, T.; Tar, A.; Halasz, Z.; Blumel, P.; Tawfik, S.;
Krohn,

[136619]K.; Lebl, J.; Peterson, P.; MEWPE-APECED Study Group:
Novel AIRE

[136620]mutations and P450 cytochrome autoantibodies in Central
and Eastern

[136621]European patients with APECED. Hum. Mutat. 18:
225–232, 2001.

[136622]24354.

[136623]Clemente, M. G.; Obermayer–Straub, P.; Meloni, A.; Strass-
burg,

[136624]C. P.; Arangino, V.; Tukey, R. H.; de Virgiliis, S.; Manns, M.
P.

[136625]: Cytochrome P450 1A2 is a hepatic autoantigen in au-
toimmune polyglandular

[136626]syndrome type 1. J. Clin. Endocr. Metab. 82: 1353–1361,
1997.

[136627]24355.

[136628]Craig, J. M.; Schiff, L. H.; Boone, J. E.: Chronic moniliasis

[136629]associated with Addison's disease. Am. J. Dis. Child. 89:
669–684,
[136630]1955.
[136631]24356.
[136632]Ekwall, O.; Hedstrand, H.; Haavik, J.; Perheentupa, J.; Bet-
terle,
[136633]C.; Gustafsson, J.; Husebye, E.; Rorsman, F.; Kampe, O.:
Pteridin-dependent
[136634]hydroxylases as autoantigens in autoimmune polyen-
docrine syndrome
[136635]type I. J. Clin. Endocr. Metab. 85: 2944–2950, 2000.
[136636]24357.
[136637]Cetani, F.; Barbesino, G.; Borsari, S.; Pardi, E.; Cianferotti,
[136638]L.; Pinchera, A.; Marcocci, C.: A novel mutation of the au-
toimmune
[136639]regulator gene in an Italian kindred with autoimmune
polyendocrinopathy–candidiasis–ectodermal
[136640]dystrophy, acting in a dominant fashion and strongly
cosegregating
[136641]with hypothyroid autoimmune thyroiditis. J. Clin. Endocr.
Metab. 86:

[136642]4747–4752, 2001.

[136643]24358.

[136644]Finnish–German APECED Consortium: An autoimmune disease, APECED,

[136645]caused by mutations in a novel gene featuring two PHD-type zinc-finger

[136646]domains. *Nature Genet.* 17: 399–403, 1997.

[136647]24359.

[136648]Foz, M.; Mirada, A.; Guardia, J.: Endocrine disorders in a family.

[136649](Letter) *Lancet* II: 269 only, 1970.

[136650]24360.

[136651]Gass, J. D. M.: The syndrome of keratoconjunctivitis, superficial

[136652]moniliiasis, idiopathic hypoparathyroidism and Addison's disease. *Am.*

[136653]*J. Ophthalm.* 54: 660–674, 1962.

[136654]24361.

[136655]Gylling, M.; Tuomi, T.; Bjorses, P.; Kontiainen, S.; Partanen,

[136656]J.; Christie, M. R.; Knip, M.; Perheentupa, J.; Miettinen, A.:

Beta-cell

[136657]autoantibodies, human leukocyte antigen II alleles, and
type 1 diabetes

[136658]in autoimmune polyendocrinopathy–candida–
sis–ectodermal dystrophy. J.

[136659]Clin. Endocr. Metab. 85: 4434–4440, 2000.

[136660]24362.

[136661]Hedstrand, H.; Ekwall, O.; Haavik, J.; Landgren, E.; Betterle,

[136662]C.; Perheentupa, J.; Gustafsson, J.; Husebye, E.; Rorsman,
F.; Kampe,

[136663]O.: Identification of tyrosine hydroxylase as an autoanti-
gen in autoimmune

[136664]polyendocrine syndrome type I. Biochem. Biophys. Res.
Commun. 267:

[136665]456–461, 2000.

[136666]24363.

[136667]Heino, M.; Peterson, P.; Kudoh, J.; Shimizu, N.; An-
tonarakis,

[136668]S. E.; Scott, H. S.; Krohn, K.: APECED mutations in the au-
toimmune

[136669]regulator (AIRE) gene. Hum. Mutat. 18: 205–211, 2001.

[136670]24364.

[136671]Heino, M.; Scott, H. S.; Chen, Q.; Peterson, P.; Maenpaa, U.;

[136672]Papasavvas, M.–P.; Mittaz, L.; Barras, C.; Rossier, C.; Chrousos,

[136673]G. P.; Stratakis, C. A.; Nagamine, K.; Kudoh, J.; Shimizu, N.; Maclaren,

[136674]N.; Antonarakis, S. E.; Krohn, K.: Mutation analyses of North American

[136675]APS–1 patients. Hum. Mutat. 13: 69–74, 1999.

[136676]24365.

[136677]Hendrix, T. R.: Personal Communication. Baltimore, Md. 1/14/1985.

[136678]24366.

[136679]Hiekkala, H.: Idiopathic hypoparathyroidism, adrenal insufficiency

[136680]and moniliasis in children. Ann. Paediat. Fenn. 10: 213–222, 1964.

[136681]24367.

[136682]Hung, W.; Migeon, C. J.; Parrott, R. H.: A possible autoim–

mune

[136683]basis for Addison's disease in three siblings, one with idiopathic

[136684]hypoparathyroidism, pernicious anemia and superficial moniliasis. New

[136685]Eng. J. Med. 269: 658–663, 1963.

[136686]24368.

[136687]Husebye, E. S.; Gebre-Medhin, G.; Tuomi, T.; Perheentupa, J.;

[136688]Landin-Olsson, M.; Gustafsson, J.; Rorsman, F.; Kampe, O.: Autoantibodies

[136689]against aromatic L-amino acid decarboxylase in autoimmune polyendocrine

[136690]syndrome type I. J. Clin. Endocr. Metab. 82: 147–150, 1997.

[136691]24369.

[136692]Ishii, T.; Suzuki, Y.; Ando, N.; Matsuo, N.; Ogata, T.: Novel

[136693]mutations of the autoimmune regulator gene in two siblings with autoimmune

[136694]polyendocrinopathy–candidiasis–ectodermal dystrophy. J. Clin. Endocr.

[136695]Metab. 85: 2922–2926, 2000.

[136696]24370.

[136697]Kenny, F. M.; Holliday, M. D.: Hypoparathyroidism, moniliasis,

[136698]Addison's and Hashimoto's disease. Hypercalcemia treated with intravenously

[136699]administered sodium sulfate. New Eng. J. Med. 271: 708-713, 1964.

[136700]24371.

[136701]Krohn, K.; Uibo, R.; Aavik, E.; Peterson, P.; Savilahti, K.:

[136702]Identification by molecular cloning of an autoantigen associated with

[136703]Addison's disease as steroid 17-alpha-hydroxylase. Lancet 339: 770-773,

[136704]1992.

[136705]24372.

[136706]Kunin, A. S.; MacKay, B. R.; Burns, S. L.; Halberstam, M. J.:

[136707]The syndrome of hypoparathyroidism and adrenocortical insufficiency,

[136708]a possible sequel of hepatitis: case report and review of the literature. Am.

[136709]J. Med. 34: 856-866, 1963.

[136710]24373.

[136711]Louria, D. B.; Shannon, D. C.; Johnson, G.; Caroline, L.;
Okas,

[136712]A.; Taschdjian, C.: The susceptibility to moniliasis in chil-
dren

[136713]with endocrine hypofunction. Trans. Assoc. Am. Phys. 80:
236-249,

[136714]1967.

[136715]24374.

[136716]Maghnie, M.; Cosi, G.; Genovese, E.; Manca-Bitti, M. L.;
Cohen,

[136717]A.; Zecca, S.; Tinelli, C.; Gallucci, M.; Bernasconi, S.;
Boscherini,

[136718]B.; Severi, F.; Arico, M.: Central diabetes insipidus in chil-
dren

[136719]and young adults. New Eng. J. Med. 343: 998-1007, 2000.

[136720]24375.

[136721]Marieb, N. J.; Melby, J. C.; Lyall, S. S.: Isolated hypoaldos-
teronism

[136722]associated with idiopathic hypoparathyroidism. Arch. In-
tern. Med. 134:

[136723]424–429, 1974.

[136724]24376.

[136725]McKusick, V. A.: Personal Communication. Baltimore, Md.
1985.

[136726]24377.

[136727]Meloni, A.; Perniola, R.; Faa, V.; Corvaglia, E.; Cao, A.;
Rosatelli,

[136728]M. C.: Delineation of the molecular defects in the AIRE
gene in autoimmune

[136729]polyendocrinopathy–candidiasis–ectodermal dystrophy
patients from

[136730]Southern Italy. J. Clin. Endocr. Metab. 87: 841–846, 2002.

[136731]24378.

[136732]Nagamine, K.; Peterson, P.; Scott, H. S.; Kudoh, J.; Mi-
noshima,

[136733]S.; Heino, M.; Krohn, K. J. E.; Lalioti, M. D.; Mullis, P. E.;
Antonarakis,

[136734]S. E.; Kawasaki, K.; Asakawa, S.; Ito, F.; Shimizu, N.: Posi-
tional

[136735]cloning of the APECED gene. Nature Genet. 17: 393–398,
1997.

[136736]24379.

[136737]Neufeld, M.; Maclaren, N.; Blizzard, R.: Autoimmune
polyglandular

[136738]syndrome. *Pediat. Ann.* 9: 154–162, 1980.

[136739]24380.

[136740]Neufeld, M.; Maclaren, N. K.; Blizzard, R. M.: Two types of
autoimmune

[136741]Addison's disease associated with different polyglandular
autoimmune

[136742](PGA) syndromes. *Medicine* 60: 355–362, 1981.

[136743]24381.

[136744]Nithiyananthan, R.; Heward, J. M.; Allahabadia, A.; Barnett,
A.

[136745]H.; Franklyn, J. A.; Gough, S. C. L.: A heterozygous dele-
tion of

[136746]the autoimmune regulator (AIRE1) gene, autoimmune thy-
roid disease,

[136747]and type 1 diabetes: no evidence for association. *J. Clin.*
Endocr.

[136748]*Metab.* 85: 1320–1322, 2000.

[136749]24382.

- [136750]Okano, M.; Bell, D. W.; Haber, D. A.; Li, E.: DNA methyltransferases
- [136751]Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian
- [136752]development. Cell 99: 247–257, 1999.
- [136753]24383.
- [136754]Wijmenga, C.; van den Heuvel, L. P. W. J.; Strengman, E.; Luyten,
- [136755]J. A. F. M.; van der Burgt, I. J. A. M.; de Groot, R.; Smeets, D.
- [136756]F. C. M.; Draaisma, J. M. T.; van Dongen, J. J.; De Abreu, R. A.;
- [136757]Pearson, P. L.; Sandkuijl, L. A.; Weemaes, C. M. R.: Localization
- [136758]of the ICF syndrome to chromosome 20 by homozygosity mapping. Am.
- [136759]J. Hum. Genet. 63: 803–809, 1998.
- [136760]24384.
- [136761]Xu, G.-L.; Bestor, T. H.; Bourc'his, D.; Hsieh, C.-L.; Tommerup,
- [136762]N.; Bugge, M.; Hulten, M.; Qu, X.; Russo, J. J.; Viegas-Pequignot,

[136763]E.: Chromosome instability and immunodeficiency syndrome caused by

[136764]mutations in a DNA methyltransferase gene. Nature 402: 187-191,

[136765]1999.

[136766]24385.

[136767]Yamamoto, T.; Ninomiya, H.; Matsumoto, M.; Ohta, Y.; Nanba, E.;

[136768]Tsutsumi, Y.; Yamakawa, K.; Millat, G.; Vanier, M. T.; Pentchev, P.

[136769]G.; Ohno, K.: Genotype-phenotype relationship of Niemann-Pick disease

[136770]type C: a possible correlation between clinical onset and levels of

[136771]NPC1 protein in isolated skin fibroblasts. J. Med. Genet. 37: 707-711,

[136772]2000.

[136773]24386.

[136774]Alagille, D.; Josso, F.; Binet, J. L.; Blin, M. L.: La dystrophie

[136775]thrombocytaire hemorrhagique: discussion nosologique. Nouv. Rev.

[136776]Franc. Hemat. 4: 755–790, 1964.

[136777]24387.

[136778]Awidi, A. S.: Increased incidence of Glanzmann's throm-
basthenia

[136779]in Jordan as compared with Scandinavia. Scand. J. Haemat.
30: 218–222,

[136780]1983.

[136781]24388.

[136782]Basani, R .B.; Vilaire, G.; Shattil, S. J.; Kolodziej, M. A.;
Bennett,

[136783]J. S.; Poncz, M.: Glanzmann thrombasthenia due to a two
amino acid

[136784]deletion in the fourth calcium-binding domain of alphaIIb:
demonstration

[136785]of the importance of calcium-binding domains in the con-
formation of

[136786]alphaIIb/beta3. Blood 88: 167–173, 1996.

[136787]24389.

[136788]Bellucci, S.; Devergie, A.; Gluckman, E.; Tobelem, G.;
Lethielleux,

[136789]P.; Benbunan, M.; Schaison, G.; Boiron, M.: Complete cor-

rection of

[136790]Glanzmann's thrombasthenia by allogeneic bone-marrow transplantation. Brit.

[136791]J. Haemat. 59: 635-641, 1985.

[136792]24390.

[136793]Beutler, E.: Glanzmann's thrombasthenia and reduced glutathione. New

[136794]Eng. J. Med. 287: 1094-1095, 1972.

[136795]24391.

[136796]Bowie, E. J. W.; Owen, C. A., Jr.: Thrombopathy. Seminars Hemat. 5:

[136797]73-82, 1968.

[136798]24392.

[136799]Bowie, E. J. W.; Thompson, J. H., Jr.; Owen, C. A., Jr.: A new

[136800]abnormality of platelet function. Thromb. Diath. Haemorrh. 11: 195-203,

[136801]1964.

[136802]24393.

[136803]Bray, P. F.: Inherited diseases of platelet glycoproteins: considerations

[136804]for rapid molecular characterization. Thromb. Haemost.
72: 492–502,

[136805]1994.

[136806]24394.

[136807]Bray, P. F.; Barsh, G.; Rosa, J.-P.; Luo, X. Y.; Magenis, E.;

[136808]Shuman, M. A.: Physical linkage of the genes for platelet
membrane

[136809]glycoproteins IIb and IIIa. Proc. Nat. Acad. Sci. 85:
8683–8687,

[136810]1988.

[136811]24395.

[136812]Bray, P. F.; Rosa, J.-P.; Johnston, G. I.; Shiu, D. T.; Cook,

[136813]R. G.; Lau, C.; Kan, Y. W.; McEver, R. P.; Shuman, M. A.:
Platelet

[136814]glycoprotein IIb: chromosomal localization and tissue ex-
pression. J.

[136815]Clin. Invest. 80: 1812–1817, 1987.

[136816]24396.

[136817]Bray, P. F.; Rosa, J.-P.; Lingappa, V. R.; Kan, Y. W.; McEver,

[136818]R. P.; Shuman, M. A.: Biogenesis of the platelet receptor

for fibrinogen:

[136819]evidence for separate precursors for glycoproteins IIb and IIIa. Proc.

[136820]Nat. Acad. Sci. 83: 1480-1484, 1986.

[136821]24397.

[136822]Burk, C. D.; Newman, P. J.; Lyman, S.; Gill, J.; Collier, B. S.;

[136823]Poncz, M.: A deletion in the gene for glycoprotein IIb associated

[136824]with Glanzmann's thrombasthenia. J. Clin. Invest. 87: 270-276, 1991.

[136825]24398.

[136826]Collier, B. S.; Seligsohn, U.; Little, P. A.: Type I Glanzmann

[136827]thrombasthenia patients from the Iraqi-Jewish and Arab populations

[136828]in Israel can be differentiated by platelet glycoprotein IIIa immunoblot

[136829]analysis. Blood 69: 1696-1703, 1987.

[136830]24399.

[136831]Corby, D. G.; Zirbel, C. L.; Lindley, A.; Schulman, I.: Thrombasthenia. Am.

[136832]J. Dis. Child. 121: 140-144, 1971.

[136833]24400.

[136834]Cronberg, S.; Nilsson, I. M.; Zetterqvist, E.: Investigation
[136835]of a family with members with both severe and mild de-
gree of thrombasthenia. Acta

[136836]Paediat. Scand. 56: 189–197, 1967.

[136837]24401.

[136838]Chuang, D. T.; Shih, V. E.: Disorders of branched chain
amino

[136839]acid and keto acid metabolism.In: Scriver, C. R.; Beaudet,
A. L.;

[136840]Sly, W. S.; Valle, D.: Metabolic and Molecular Bases of In-
herited

[136841]Disease. New York: McGraw–Hill (pub.) (7th ed.): 1995. Pp.
1239–1277.

[136842]24402.

[136843]Barth, M. L.; Fensom, A.; Harris, A.: Missense mutations in
the

[136844]arylsulphatase A genes of metachromatic leukodystrophy
patients. Hum.

[136845]Molec. Genet. 2: 2117–2121, 1993.

[136846]24403.

[136847]Barth, M. L.; Fensom, A.; Harris, A.: Prevalence of common mutations

[136848]in the arylsulphatase A gene in metachromatic leukodys- trophy patients

[136849]diagnosed in Britain. Hum. Genet. 91: 73–77, 1993.

[136850]24404.

[136851]Lake, B. D.; Smith, V. V.; Judge, M. R.; Harper, J. I.; Besley,

[136852]G. T. N.: Hexanol dehydrogenase activity shown by en- zyme histochemistry

[136853]on skin biopsies allows differentiation of Sjogren–Larsson syndrome

[136854]from other ichthyoses. J. Inherit. Metab. Dis. 14: 338–340, 1991.

[136855]24405.

[136856]Link, J. K.; Roldan, E. C.: Mental deficiency, spasticity, and

[136857]congenital ichthyosis: report of a case. J. Pediat. 52: 712–714,

[136858]1958.

[136859]24406.

[136860]Pigg, M.; Jagell, S.; Sillen, A.; Weissenbach, J.; Gustavson,

[136861]K.-H.; Wadelius, C.: The Sjogren-Larsson syndrome gene is close to

[136862]D17S805 as determined by linkage analysis and allelic association. Nature

[136863]Genet. 8: 361-364, 1994.

[136864]24407.

[136865]Rayner, A.; Lampert, R. P.; Rennert, O. M.: Familial ichthyosis,

[136866]dwarfism, mental retardation, and renal disease. J. Pediat. 92:

[136867]766-768, 1978.

[136868]24408.

[136869]Richards, B. W.: Congenital ichthyosis, spastic diplegia and

[136870]mental deficiency. (Letter) Brit. Med. J. 2: 714 only, 1960.

[136871]24409.

[136872]Rizzo, W. B.; Carney, G.; Lin, Z.: The molecular basis of Sjogren-Larsson

[136873]syndrome: mutation analysis of the fatty aldehyde dehydrogenase gene. Am.

[136874]J. Hum. Genet. 65: 1547-1560, 1999.

[136875]24410.

[136876]Rizzo, W. B.; Craft, D. A.: Sjogren-Larsson syndrome: deficient

[136877]activity of the fatty aldehyde dehydrogenase component of fatty alcohol:NAD+

[136878]oxidoreductase in cultured fibroblasts. J. Clin. Invest. 88: 1643-1648,

[136879]1991.

[136880]24411.

[136881]Rizzo, W. B.; Dammann, A. L.; Craft, D.; Black, S.; Henderson

[136882]Tilton, A.; Africk, D.; Chaves-Carballo, E.: Sjogren-Larsson syndrome:

[136883]deficient fatty alcohol:NAD+ oxidoreductase (FAO) activity in mixed

[136884]leukocytes. (Abstract) Am. J. Hum. Genet. 41: A16 only, 1987.

[136885]24412.

[136886]Rizzo, W. B.; Dammann, A. L.; Craft, D. A.: Sjogren-Larsson syndrome:

[136887]impaired fatty alcohol oxidation in cultured fibroblasts due to deficient

[136888]fatty alcohol:nicotinamide adenine dinucleotide oxidore-
ductase activity. J.

[136889]Clin. Invest. 81: 738-744, 1988.

[136890]24413.

[136891]Rizzo, W. B.; Dammann, A. L.; Craft, D. A.; Black, S. H.;
Henderson

[136892]Tilton, A.; Africk, D.; Chaves-Carballo, E.; Holmgren, G.;
Jagell,

[136893]S.: Sjogren-Larsson syndrome: inherited defect in the fatty
alcohol

[136894]cycle. J. Pediat. 115: 228-234, 1989.

[136895]24414.

[136896]Rogers, G. R.; Markova, N. G.; De Laurenzi, V.; Rizzo, W.
B.;

[136897]Compton, J. G.: Genomic organization and expression of
the human

[136898]fatty aldehyde dehydrogenase gene (FALDH). Genomics
39: 127-135,

[136899]1997.

[136900]24415.

[136901]Rogers, G. R.; Rizzo, W. B.; Zlotogorski, A.; Hashem, N.;

Lee,

[136902]M.; Bale, S. J.; Compton, J. G.: Sjogren-Larsson syndrome (SLS) and

[136903]fatty aldehyde dehydrogenase (FALDH) genes map to chromosome 17p.

[136904](Abstract) Am. J. Hum. Genet. 57: A202 only, 1995.

[136905]24416.

[136906]Rogers, G. R.; Rizzo, W. B.; Zlotogorski, A.; Hashem, N.; Lee,

[136907]M.; Compton, J. G.; Bale, S. J.: Genetic homogeneity in Sjogren-Larsson

[136908]syndrome: linkage to chromosome 17p in families of different non-Swedish

[136909]ethnic origins. Am. J. Hum. Genet. 57: 1123-1129, 1995.

[136910]24417.

[136911]Selmanowitz, V. J.; Porter, M. J.: The Sjogren-Larsson syndrome. Am.

[136912]J. Med. 42: 412-422, 1967.

[136913]24418.

[136914]Sillen, A.; Anton-Lamprecht, I.; Braun-Quentin, C.; Kraus, C.

[136915]S.; Sayli, B. S.; Ayuso, C.; Jagell, S.; Kuster, W.; Wadelius, C.

[136916]: Spectrum of mutations and sequence variants in the
FALDH gene in

[136917]patients with Sjogren–Larsson syndrome. Hum. Mutat. 12:
377–384,

[136918]1998.

[136919]24419.

[136920]Sillen, A.; Holmgren, G.; Wadelius, C.: First prenatal diag-
nosis

[136921]by mutation analysis in a family with Sjogren–Larsson
syndrome. Prenat.

[136922]Diag. 17: 1147–1149, 1997.

[136923]24420.

[136924]Sillen, A.; Jagell, S.; Wadelius, C.: A missense mutation in

[136925]the FALDH gene identified in Sjogren–Larsson syndrome
patients originating

[136926]from the northern part of Sweden. Hum. Genet. 100:
201–203, 1997.

[136927]24421.

[136928]Sjogren, T.: Oligophrenia combined with congenital
ichthyosiform

[136929]erythrodermia, spastic syndrome and macular retinal de-
generation:

[136930]a clinical and genetic study. Acta Genet. Statist. Med. 6:
80-91,

[136931]1956.

[136932]24422.

[136933]Sjogren, T.; Larsson, T.: Oligophrenia in combination with
congenital

[136934]ichthyosis and spastic disorders: a clinical and genetic
study. Acta

[136935]Psychiat. Neurol. Scand. 32 (suppl. 113): 1-112, 1957.

[136936]24423.

[136937]Tsukamoto, N.; Chang, C.; Yoshida, A.: Mutations associ-
ated with

[136938]Sjogren-Larsson syndrome. Ann. Hum. Genet. 61:
235-242, 1997.

[136939]24424.

[136940]Willemsen, M. A. A. P.; Cruysberg, J. R. M.; Rotteveel, J. J.;

[136941]Aandekerck, A. L.; Van Domburg, P. H. M. F.; Deutman, A.
F.: Juvenile

[136942]macular dystrophy associated with deficient activity of

fatty aldehyde

[136943]dehydrogenase in Sjogren–Larsson syndrome. Am. J. Ophthalmol. 130:

[136944]782–789, 2000.

[136945]24425.

[136946]Zaleski, W. A.: Congenital ichthyosis, mental retardation and

[136947]spasticity (Sjogren–Larsson syndrome). Canad. Med. Assoc. J. 86:

[136948]951–954, 1962.

[136949]24426.

[136950]Zhang, M.; Dwyer, N. K.; Love, D. C.; Cooney, A.; Comly, M.; Neufeld,

[136951]E.; Pentchev, P. G.; Blanchette–Mackie, E. J.; Hanover, J. A.: Cessation

[136952]of rapid late endosomal tubulovesicular trafficking in Niemann–Pick

[136953]type C1 disease. Proc. Nat. Acad. Sci. 98: 4466–4471, 2001.

[136954]24427.

[136955]Loeffen, J.; Elpeleg, O.; Smeitink, J.; Smeets, R.; Stockler–

Ipsiroglu,

[136956]S.; Mandel, H.; Sengers, R.; Trijbels, F.; van den Heuvel, L.:

Mutations

[136957]in the complex I NDUF52 gene of patients with cardiomyopathy and encephalomyopathy. Ann.

[136958]Neurol. 49: 195–201, 2001.

[136959]24428.

[136960]Smeitink, J.; van den Heuvel, L.: Human mitochondrial complex

[136961]I in health and disease. Am. J. Hum. Genet. 64: 1505–1510, 1999.

[136962]24429.

[136963]Baumgartner, E. R.; Suormala, T.; Wick, H.; Bausch, J.; Bonjour,

[136964]J.–P.: Biotinidase deficiency: factors responsible for the increased

[136965]biotin requirement. J. Inherit. Metab. Dis. 8 (suppl. 1): 59–64,

[136966]1985.

[136967]24430.

[136968]Burlina, A. B.; Sherwood, W. G.; Marchioro, M. V.; Dalla

Bernardina,

[136969]B.; Gaburro, D.: Neonatal screening for biotinidase deficiency in

[136970]north eastern Italy. *Europ. J. Pediat.* 147: 317–318, 1988.

[136971]24431.

[136972]Cole, H.; Reynolds, T. R.; Lockyer, J. M.; Buck, G. A.; Den-
son,

[136973]T.; Spence, J. E.; Hymes, J.; Wolf, B.: Human serum bio-
tinidase:

[136974]cDNA cloning, sequence, and characterization. *J. Biol.*
Chem. 269:

[136975]6566–6570, 1994.

[136976]24432.

[136977]Cole, H.; Weremowicz, S.; Morton, C. C.; Wolf, B.: Localiza-
tion

[136978]of serum biotinidase (BTD) to human chromosome 3 in
band p25. *Genomics* 22:

[136979]662–663, 1994.

[136980]24433.

[136981]Dionisi-Vici, C.; Bachmann, C.; Graziani, M. C.; Sabetta,
G.:

- [136982]Laryngeal stridor as a leading symptom in a biotinidase-deficient
patient. J. Inherit. Metab. Dis. 11: 312-313, 1988.
- [136983]24434.
- [136984]Dunkel, G.; Scriver, C. R.; Clow, C. L.; Melancon, S.;
Lemieux,
B.; Grenier, A.; Laberge, C.: Prospective ascertainment of
complete
and partial serum biotinidase deficiency in the newborn. J.
Inherit.
Metab. Dis. 12: 131-138, 1989.
- [136985]24435.
- [136986]Fischer, A.; Munnich, A.; Saudubray, J. M.; Mamas, S.;
Coude, F.
X.; Charpentier, C.; Dray, F.; Frezal, J.; Griscelli, C.: Biotin-
responsive
immunoregulatory dysfunction in multiple carboxylase
deficiency. J.
Clin. Immun. 2: 35-38, 1982.
- [136987]24436.
- [136988]Gaudry, M.; Munnich, A.; Saudubray, J. M.; Ogier, H.;

Mitchell,

[136996]G.; Marsac, C.; Causse, M.; Marquet, A.; Frezal, J.: Deficient liver

[136997]biotinidase activity in multiple carboxylase deficiency.

(Letter) Lancet II:

[136998]397, 1983.

[136999]24437.

[137000]Giardini, O.; D'Eufemia, P.; Cantani, A.; Ruberto, U.; Sabetta,

[137001]G.; Castro, M.; Cardi, E.: Congenital lactic acidosis associated

[137002]with laryngeal stridor: report of 2 cases. Riv. Ital. Pediat. 7:

[137003]139-142, 1981.

[137004]24438.

[137005]Gordon, A.: Biotinidase mutational 'hotspot'. (Letter) Nature

[137006]Genet. 13: 144-145, 1996.

[137007]24439.

[137008]Greter, J.; Holme, E.; Lindstedt, S.; Koivikko, M.: Biotin-responsive

[137009]3-methylcrotonylglycinuria with biotinidase deficiency. J.

Inherit.

[137010]Metab. Dis. 8: 103-104, 1985.

[137011]24440.

[137012]Hart, P. S.; Hymes, J.; Wolf, B.: Biochemical and immunological

[137013]characterization of serum biotinidase in profound biotinidase deficiency. Am.

[137014]J. Hum. Genet. 50: 126-136, 1992.

[137015]24441.

[137016]Heard, G. S.; Wolf, B.; Jefferson, L. G.; Weissbecker, K. A.;

[137017]Nance, W. E.; Secor McVoy, J. R.; Napolitano, A.; Mitchell, P. L.;

[137018]Lambert, F. W.; Linyear, A. S.: Neonatal screening for biotinidase

[137019]deficiency: results of a 1-year pilot study. J. Pediat. 108: 40-46,

[137020]1986.

[137021]24442.

[137022]Hymes, J.; Stanley, C. M.; Wolf, B.: Mutations in BTBD caus-

ing

[137023]biotinidase deficiency. Hum. Mutat. 18: 375–381, 2001.

[137024]24443.

[137025]Kalayci, O.; Coskun, T.; Tokatli, A.; Demir, E.; Erdem, G.;
Gungor,

[137026]C.; Yukselen, A.; Ozalp, I.: Infantile spasms as the initial
symptom

[137027]of biotinidase deficiency. J. Pediat. 124: 103–104, 1994.

[137028]24444.

[137029]Kennedy, R.; Girdwood, R. W. A.; King, M. D.: Neonatal
screening

[137030]for biotinidase deficiency: a pilot study in Scotland. J. In-
herit.

[137031]Metab. Dis. 12: 344–345, 1989.

[137032]24445.

[137033]Minns, R. A.; Kirk, J.: Biotinidase deficiency in Scotland.
(Letter) Dev.

[137034]Med. Child Neurol. 36: 748–749, 1994.

[137035]24446.

[137036]Muhl, A.; Moslinger, D.; Item, C. B.; Stockler-Ipsiroglu, S.:

[137037]Molecular characterisation of 34 patients with biotinidase

deficiency

[137038]ascertained by newborn screening and family investigation. *Europ.*

[137039]*J. Hum. Genet.* 9: 237-243, 2001.

[137040]24447.

[137041]Munnich, A.; Saudubray, J. M.; Carre, G.; Coude, F. X.; Ogier,

[137042]H.; Charpentier, C.; Frezal, J.: Defective biotin absorption in multiple

[137043]carboxylase deficiency. (Letter) *Lancet* II: 263 only, 1981.

[137044]24448.

[137045]Norrsgard, K. J.; Pomponio, R. J.; Hymes, J.; Wolf, B.: Mutations

[137046]causing profound biotinidase deficiency in children ascertained by

[137047]newborn screening in the United States occur at different frequencies

[137048]than in symptomatic children. *Pediat. Res.* 46: 20-27, 1999.

[137049]24449.

[137050]Norrsgard, K. J.; Swango, K. L.; Wolf, B.: Mutation (D444H)

in

[137051]the biotinidase gene causes approximately 50% loss of
enzyme activity

[137052]and is common in the general population. (Abstract) Am.
J. Hum. Genet.

[137053](Suppl.) 59: A275 only, 1996.

[137054]24450.

[137055]Pomponio, R. J.; Coskun, T.; Demirkol, M.; Tokatli, A.;
Ozalp,

[137056]I.; Huner, G.; Baykal, T.; Wolf, B.: Novel mutations cause
biotinidase

[137057]deficiency in Turkish children. J. Inherit. Metab. Dis. 23:
120-128,

[137058]2000.

[137059]24451.

[137060]Pomponio, R. J.; Hymes, J.; Reynolds, T. R.; Meyers, G. A.;
Fleischhauer,

[137061]K.; Buck, G. A.; Wolf, B.: Mutations in the human bio-
tinidase gene

[137062]that cause profound biotinidase deficiency in symptomatic
children:

[137063]molecular, biochemical, and clinical analysis. Pediat. Res.

42:

[137064]840–848, 1997.

[137065]24452.

[137066]Pomponio, R. J.; Narasimahan, V.; Reynolds, T. R.; Buck, G. A.;

[137067]Povirk, L. F.; Wolf, B.: Deletion/insertion mutation that causes

[137068]biotinidase deficiency may result from the formation of a quasipalindromic

[137069]structure. Hum. Molec. Genet. 5: 1657–1661, 1996.

[137070]24453.

[137071]Pomponio, R. J.; Norrgard, K. J.; Hymes, J.; Reynolds, T. R.;

[137072]Buck, G. A.; Baumgartner, R.; Suormala, T.; Wolf, B.: Arg538 to cys

[137073]mutation in a CpG dinucleotide of the human biotinidase gene is the

[137074]second most common cause of profound biotinidase deficiency in symptomatic

[137075]children. Hum. Genet. 99: 506–512, 1997.

[137076]24454.

[137077]Pomponio, R. J.; Reynolds, T. R.; Cole, H.; Buck, G. A.;

Wolf,

[137078]B.: Mutational hotspot in the human biotinidase gene
causes profound

[137079]biotinidase deficiency. Nature Genet. 11: 96–98, 1995.

[137080]24455.

[137081]Pomponio, R. J.; Reynolds, T. R.; Mandel, H.; Admoni, O.;
Melone,

[137082]P. D.; Buck, G. A.; Wolf, B.: Profound biotinidase deficiency
caused

[137083]by a point mutation that creates a downstream cryptic
3–prime splice

[137084]acceptor site within an exon of the human biotinidase
gene. Hum.

[137085]Molec. Genet. 6: 739–745, 1997.

[137086]24456.

[137087]Packman, S.; Sweetman, L.; Baker, H.; Wall, S.: The neona-
tal

[137088]form of biotin–responsive multiple carboxylase deficiency.
J. Pediat. 99:

[137089]418–420, 1981.

[137090]24457.

[137091]Packman, S.; Sweetman, L.; Yoshino, M.; Baker, H.; Cowan, M.:

[137092]Biotin-responsive multiple carboxylase deficiency of infantile onset. J.

[137093]Pediat. 99: 421-423, 1981.

[137094]24458.

[137095]Daiger, S. P.; Lidsky, A. S.; Chakraborty, R.; Koch, R.; Gut-
tler,

[137096]F.; Woo, S. L. C.: Polymorphic DNA haplotypes at the
phenylalanine

[137097]hydroxylase locus in prenatal diagnosis of phenylke-
tonuria. Lancet I:

[137098]229-231, 1986.

[137099]24459.

[137100]Pizarro, T. T.; Michie, M. H.; Bentz, M.; Woraratanadharm,
J.;

[137101]Smith, M. F., Jr.; Foley, E.; Moskaluk, C. A.; Bickston, S. J.;
Cominelli,

[137102]F.: IL-18, a novel immunoregulatory cytokine, is up-
regulated in

[137103]Crohn's disease: expression and localization in intestinal
mucosal

[137104]cells. J. Immun. 162: 6829–6835, 1999.

[137105]24460.

[137106]Oggiano, L.; Pirastu, M.; Moi, P.; Longinotti, M.; Perseu, L.;

[137107]Cao, A.: Molecular characterization of a normal HbA-2
beta-thalassemia

[137108]determinant in a Sardinian family. Brit. J. Haemat. 67:
225–229,

[137109]1987.

[137110]24461.

[137111]Ohba, Y.; Igarashi, M.; Tsukahara, M.; Nakashima, M.;
Sanada,

[137112]C.; Ami, M.; Arai, Y.; Miyaji, T.: Hb A(2) Yokoshima,
delta25(B7)

[137113]gly-to-asp, a new delta chain variant found in a Japanese
family. Hemoglobin 9:

[137114]613–615, 1985.

[137115]24462.

[137116]Ostertag, W.; Smith, E. W.: Hemoglobin–Lepore–Baltimore,
a third

[137117]type of a delta, beta crossover (delta 50, beta 86). Europ.
J. Biochem. 10:

[137118]371–376, 1969.

[137119]24463.

[137120]Papadakis, M.; Drakoulakou, O.; Papapanagiotou, E.;
Pessini, D.;

[137121]Loutradi–Anagnostou, A.: Hb A(2)–Agrinio
(delta-43(CD2)glu-to-gly

[137122](GAG-to-GGG)): a new delta-chain variant detected in a
Greek family. Hemoglobin 19:

[137123]295–299, 1995.

[137124]24464.

[137125]Petes, T. D.: Evidence that structural variants within the
human

[137126]delta-globin protein may reflect genetic interactions be-
tween the

[137127]delta- and beta-globin genes. (Letter) Am. J. Hum. Genet.
34: 820–823,

[137128]1982.

[137129]24465.

[137130]Ramirez, F.; Mears, J. G.; Nudel, U.; Bank, A.; Luzzatto, L.;

[137131]Di Prisco, G.; D'Avino, R.; Pepe, G.; Gambino, R.; Cimino,
R.; Quattrin,

[137132]N.: Defects in DNA and globin messenger RNA in homozygotes for hemoglobin

[137133]Lepore. J. Clin. Invest. 63: 736–742, 1979.

[137134]24466.

[137135]Ranney, H. M.; Jacobs, A. S.: Simultaneous occurrence of haemoglobins

[137136]C and Lepore in an Afro–American. Nature 204: 163–166, 1964.

[137137]24467.

[137138]Ranney, H. M.; Jacobs, A. S.; Ramot, B.; Bradley, T. B., Jr.:

[137139]Hemoglobin NYU, a delta chain variant, alpha 2 delta 2(12 lys). J.

[137140]Clin. Invest. 48: 2057–2062, 1969.

[137141]24468.

[137142]Renda, M.; Piazza, T.; Ciaccio, C.; Maggio, A.: Delta(+)-27 homozygosis

[137143]in a Sicilian family. Haematologica 77: 82–83, 1992.

[137144]24469.

[137145]Rieder, R. F.; Clegg, J. B.; Weiss, H. J.; Christy, N. P.; Rabinowitz,

[137146]R.: Hemoglobin A-2-Roosevelt: delta 20 val-to-glu.

Biochim. Biophys.

[137147]Acta 439: 501-504, 1976.

[137148]24470.

[137149]Romero Garcia, C.; Navarro, J. L.; Lam, H.; Webber, B. B.;

Headlee,

[137150]M. G.; Wilson, J. B.; Huisman, T. H. J.: Hb A(2)-Manzanares

or delta

[137151]121(GH4) glu-to-val, an unstable delta chain variant observed in a

[137152]Spanish family. Hemoglobin 7: 435-442, 1983.

[137153]24471.

[137154]Salkie, M. L.; Gordon, P. A.; Rigal, W. M.; Lam, H.; Wilson,

J.

[137155]B.; Headlee, M. E.; Huisman, T. H. J.: Hb A2-Canada or

delta 99(G1)

[137156]asp-to-asn, a newly discovered delta chain variant with increased

[137157]oxygen affinity occurring in cis to beta-thalassemia.

Hemoglobin 6:

[137158]223-231, 1982.

[137159]24472.

[137160]Schiliro, G.; Russo-Mancuso, G.; Dibenedetto, S. P.; Samperi,

[137161]P.; Di Cataldo, A.; Ragusa, R.; Testa, R.: Six rare hemoglobin variants

[137162]found in Sicily. Hemoglobin 15: 431-437, 1991.

[137163]24473.

[137164]Sharma, R. S.; Harding, D. L.; Wong, S. D.; Wilson, J. B.; Gravely,

[137165]M. E.; Huisman, T. H. J.: A new delta chain variant, hemoglobin-A2

[137166]Melbourne, or delta 43 glu-to-lys (CD2). Biochim. Biophys. Acta 359:

[137167]233-235, 1974.

[137168]24474.

[137169]Sharma, R. S.; Williams, L.; Wilson, J. B.; Huisman, T. H. J.

[137170]: Hemoglobin A(2) Coburg or alpha-2-delta-2 116 arg-to-his (G18). Biochim.

[137171]Biophys. Acta 393: 379-382, 1975.

[137172]24475.

[137173]Spritz, R. A.; DeRiel, J. K.; Forget, B. G.; Weissman, S. M.:

[137174]Complete nucleotide sequence of the human delta-globin gene. Cell 21:

[137175]639-646, 1980.

[137176]24476.

[137177]Trifillis, P.; Ioannou, P.; Schwartz, E.; Surrey, S.: Identification

[137178]of four novel delta-globin gene mutations in Greek Cypriots using

[137179]polymerase chain reaction and automated fluorescence-based DNA sequence

[137180]analysis. Blood 78: 3298-3305, 1991.

[137181]24477.

[137182]Vella, F.; Graham, B.: A variant of hemoglobin A(2) in Alberta

[137183]Indians. Clin. Biochem. 2: 455-460, 1969.

[137184]24478.

[137185]Williamson, D.; Brennan, S. O.; Strosberg, H.; Whitty, J.; Carell,

[137186]R. W.: Hemoglobin A(2) Fitzroy delta142 ala-to-asg: a new delta-chain

[137187]variant. Hemoglobin 8: 325–332, 1984.

[137188]24479.

[137189]Wilson, J. T.; Wilson, L. B.; Ohta, Y.: A case of homozygous

[137190]delta-thalassemia not due to a deletion of the delta-
globin structural

[137191]gene. Biochem. Biophys. Res. Commun. 99: 1035–1039,
1981.

[137192]24480.

[137193]Le Coniat, M.; Kinet, J. P.; Berger, R.: The human genes for
the

[137194]alpha and gamma subunits of the mast cell receptor for
immunoglobulin

[137195]E are located on human chromosome band 1q23. Im-
munogenetics 32:

[137196]183–186, 1990.

[137197]24481.

[137198]Pestka, S.: The human interferons--from protein purifica-
tion

[137199]and sequence to cloning and expression in bacteria: be-
fore, between,

[137200]and beyond. Arch. Biochem. Biophys. 221: 1–37, 1983.

[137201]24482.

[137202]Sehgal, P. B.; Sagar, A. D.; Braude, I. A.: Further heterogeneity

[137203]of human alpha-interferon mRNA species. Science 214: 803–805, 1981.

[137204]24483.

[137205]Shows, T. B.; Sakaguchi, A. Y.; Naylor, S. L.; Goeddel, D. V.;

[137206]Lawn, R. M.: Clustering of leukocyte and fibroblast interferon genes

[137207]on human chromosome 9. Science 218: 373–374, 1982.

[137208]24484.

[137209]Slate, D. L.; D'Eustachio, P.; Pravtcheva, D.; Cunningham, A.

[137210]C.; Nagata, S.; Weissmann, C.; Ruddle, F. H.: Chromosomal location

[137211]of a human alpha interferon gene family. J. Exp. Med. 155: 1019–1024,

[137212]1982.

[137213]24485.

[137214]Trent, J. M.; Olson, S.; Lawn, R. M.: Chromosomal localization

tion

[137215]of human leukocyte, fibroblast and immune interferon
genes by means

[137216]of in situ hybridization. Proc. Nat. Acad. Sci. 79:
7809-7813, 1982.

[137217]24486.

[137218]Ullrich, A.; Gray, A.; Goeddel, D. V.; Dull, T. J.: Nucleotide

[137219]sequence of a portion of human chromosome 9 contain-
ing a leukocyte

[137220]interferon gene cluster. J. Molec. Biol. 156: 467-486,
1982.

[137221]24487.

[137222]Virelizier, J. L.; Griscelli, C.: Defaut selectif de secretion

[137223]d'interferon associe a un deficit d'activite cytotoxique na-
turelle. Arch.

[137224]Franc. Pediat. 38: 77-81, 1981.

[137225]24488.

[137226]Virelizier, J. L.; Lenoir, G.; Griscelli, C.: Persistent Epstein-
Barr

[137227]virus infection in a child with hypergammaglobulinaemia
and immunoblastic

- [137228] proliferation associated with a selective defect in interferon secretion. Lancet II: 231–234, 1978.
- [137229] 24489.
- [137230] It Hart, L. M.; Stolk, R. P.; Heine, R. J.; Grobbee, D. E.; van der Does, F. E. E.; Maassen, J. A.: Association of the insulin-receptor variant met-985 with hyperglycemia and non-insulin-dependent diabetes mellitus in the Netherlands: a population-based study. Am. J. Hum. Genet. 59: 1119–1125, 1996.
- [137231] 24490.
- [137232] Accili, D.; Drago, J.; Lee, E. J.; Johnson, M. D.; Cool, M. H.; Salvatore, P.; Asico, L. D.; Jose, P. A.; Taylor, S. I.; Westphal, H.: Early neonatal death in mice homozygous for a null allele of the insulin receptor gene. Nature Genet. 12: 106–109, 1996.
- [137233] 24491.

[137242]Accili, D.; Frapier, C.; Mosthaf, L.; McKeon, C.; Elbein, S.
C.;

[137243]Permutt, M. A.; Ramos, E.; Lander, E.; Ullrich, A.; Taylor, S.
I.

[137244]: A mutation in the insulin receptor gene that impairs
transport of

[137245]the receptor to the plasma membrane and causes insulin-
resistant diabetes. EMBO

[137246]J. 8: 2509–2517, 1989.

[137247]24492.

[137248]Al-Gazali, L. I.; Khalil, M.; Devadas, K.: A syndrome of in-
sulin

[137249]resistance resembling leprechaunism in five sibs of con-
sanguineous

[137250]parents. J. Med. Genet. 30: 470–475, 1993.

[137251]24493.

[137252]Bar, R. S.; Muggeo, M.; Roth, J.; Kahn, C. R.; Havrankova,
J.;

[137253]Imperato-McGinley, J.: Insulin resistance, acanthosis ni-
gricans,

[137254]and normal insulin receptors in a young woman: evidence
for a postreceptor

[137255]defect. J. Clin. Endocr. Metab. 47: 620–625, 1978.

[137256]24494.

[137257]Barbetti, F.; Gejman, P. V.; Taylor, S. I.; Raben, N.; Cama, A.;

[137258]Bonora, E.; Pizzo, P.; Moghetti, P.; Muggeo, M.; Roth, J.:
Detection

[137259]of mutations in insulin receptor gene by denaturing gradient gel electrophoresis. Diabetes 41:

[137260]408–415, 1992.

[137261]24495.

[137262]Barnes, N. D.; Palumbo, P. J.; Hayles, A. B.; Folgar, H.: Insulin

[137263]resistance, skin changes, and virilization: a recessively inherited

[137264]syndrome possibly due to pineal gland dysfunction. Diabetologia 10:

[137265]285–289, 1974.

[137266]24496.

[137267]Belke, D. D.; Betuing, S.; Tuttle, M. J.; Graveleau, C.;
Young,

[137268]M. E.; Pham, M.; Zhang, D.; Cooksey, R. C.; McClain, D. A.;

Litwin,

[137269]S. E.; Taegtmeier, H.; Severson, D.; Kahn, C. R.; Abel, E.
D.: Insulin

[137270]signaling coordinately regulates cardiac size, metabolism,
and contractile

[137271]protein isoform expression. J. Clin. Invest. 109: 629–639,
2002.

[137272]24497.

[137273]Benecke, H.; Flier, J. S.; Moller, D. E.: Alternatively spliced

[137274]variants of the insulin receptor protein: expression in nor-
mal and

[137275]diabetic human tissues. J. Clin. Invest. 89: 2066–2070,
1992.

[137276]24498.

[137277]Bruning, J. C.; Gautam, D.; Burks, D. J.; Gillette, J.; Schu-
bert,

[137278]M.; Orban, P. C.; Klein, R.; Krone, W.; Muller-Wieland, D.;
Kahn,

[137279]C. R.: Role of brain insulin receptor in control of body
weight and

[137280]reproduction. Science 289: 2122–2125, 2000.

[137281]24499.

[137282]Bruning, J. C.; Michael, M. D.; Winnay, J. N.; Hayashi, T.;
Horsch,

[137283]D.; Accili, D.; Goodyear, L. J.; Kahn, C. R.: A muscle-spe-
cific insulin

[137284]receptor knockout exhibits features of the metabolic syn-
drome of NIDDM

[137285]without altering glucose tolerance. Molec. Cell 2:
559-569, 1998.

[137286]24500.

[137287]Cama, A.; de la Luz Sierra, M.; Ottini, L.; Kadowaki, T.;
Gorden,

[137288]P.; Imperato-McGinley, J.; Taylor, S. I.: A mutation in the
tyrosine

[137289]kinase domain of the insulin receptor associated with in-
sulin resistance

[137290]in an obese woman. J. Clin. Endocr. Metab. 73: 894-901,
1991.

[137291]24501.

[137292]Li, Q. Y.; Newbury-Ecob, R. A.; Terrett, J. A.; Wilson, D. I.;

[137293]Curtis, A. R. J.; Yi, C. H.; Gebuhr, T.; Bullen, P. J.; Robson,
S.

[137294]C.; Strachan, T.; Bonnet, D.; Lyonnet, S.; Young, I. D.; Rae-

burn,

[137295]J. A.; Buckler, A. J.; Law, D. J.; Brook, J. D.: Holt-Oram syndrome

[137296]is caused by mutations in TBX5, a member of the
Brachyury (T) gene

[137297]family. Nature Genet. 15: 21-29, 1997.

[137298]24502.

[137299]Terrett, J. A.; Newbury-Ecob, R.; Cross, G. S.; Fenton, I.;
Raeburn,

[137300]J. A.; Young, I. D.; Brook, J. D.: Holt-Oram syndrome is a
genetically

[137301]heterogeneous disease with one locus mapping to human
chromosome 12q. Nature

[137302]Genet. 6: 401-404, 1994.

[137303]24503.

[137304]Yang, J.; Hu, D.; Xia, J.; Yang, Y.; Ying, B.; Hu, J.; Zhou, X.

[137305]: Three novel TBX5 mutations in Chinese patients with
Holt-Oram syndrome. Am.

[137306]J. Med. Genet. 92: 237-240, 2000.

[137307]24504.

[137308]Henry, I.; Humphries, S. E.; Tata, F.; Barichard, F.; Holm,

M.;

[137309]Williamson, R.; Junien, C.: The gene for HMG CoA reductase (HMGCR)

[137310]is on human chromosome 5. (Abstract) Cytogenet. Cell Genet. 40:

[137311]649-650, 1985.

[137312]24505.

[137313]Humphries, S. E.; Tata, F.; Henry, I.; Barichard, F.; Holm, M.;

[137314]Junien, C.; Williamson, R.: The isolation, characterisation, and

[137315]chromosomal assignment of the gene for human 3-hydroxy-3-methylglutaryl

[137316]coenzyme A reductase, (HMG-CoA reductase). Hum. Genet. 71: 254-258,

[137317]1985.

[137318]24506.

[137319]Rutherford, S.; Johnson, M. P.; Curtain, R. P.; Griffiths, L.

[137320]R.: Chromosome 17 and the inducible nitric oxide synthase gene in

[137321]human essential hypertension. Hum. Genet. 109:

408-415, 2001.

[137322]24507.

[137323]Walter, J. W.; North, P. E.; Waner, M.; Mizeracki, A.; Blei, F.;

[137324]Walker, J. W. T.; Reinisch, J. F.; Marchuk, D. A.: Somatic
mutation

[137325]of vascular endothelial growth factor receptors in juvenile
hemangioma. Genes

[137326]Chromosomes Cancer 33: 295-303, 2002.

[137327]24508.

[137328]Ragoussis, J.; Senger, G.; Trowsdale, J.; Campbell, I. G.:
Genomic

[137329]organization of the human folate receptor genes on chro-
mosome 11q13. Genomics 14:

[137330]423-430, 1992.

[137331]24509.

[137332]Barber, R. C.; Shaw, G. M.; Lammer, E. J.; Greer, K. A.;
Biela,

[137333]T. A.; Lacey, S. W.; Wasserman, C. R.; Finnell, R. H.: Lack of
association

[137334]between mutations in the folate receptor-alpha gene and
spina bifida. Am.

[137335]J. Med. Genet. 76: 310–317, 1998.

[137336]24510.

[137337]Bowcock, A. M.; Lacey, S.; Saltman, D.; Mohandas, T. K.;
Kamen,

[137338]B. A.; Taggart, R. T.: Localization of the folate receptor
gene to

[137339]chromosome 11q13. (Abstract) Cytogenet. Cell Genet. 58:
1955 only,

[137340]1991.

[137341]24511.

[137342]Campbell, I. G.; Jones, T. A.; Foulkes, W. D.; Trowsdale, J.:

[137343]Folate-binding protein is a marker for ovarian cancer.
Cancer Res. 51:

[137344]5329–5338, 1991.

[137345]24512.

[137346]Kranes, A.; Balogh, K., Jr.: Liver disease in a patient with

[137347]von Hippel–Lindau disease. New Eng. J. Med. 275:
950–959, 1966.

[137348]24513.

[137349]Lamiell, J. M.: Personal Communication. San Francisco,

Calif.

[137350]9/1987.

[137351]24514.

[137352]Lamiell, J. M.; Salazar, F. G.; Hsia, Y. E.: Von Hippel–Lindau

[137353]disease affecting 43 members of a single kindred.

Medicine 68: 1–29,

[137354]1989.

[137355]24515.

[137356]Latif, F.; Tory, K.; Gnarr, J.; Yao, M.; Duh, F.–M.; Orcutt,

[137357]M. L.; Stackhouse, T.; Kuzmin, I.; Modi, W.; Geil, L.;

Schmidt, L.;

[137358]Zhou, F.; Li, H.; Wei, M. H.; Chen, F.; Glenn, G.; Choyke, P.;

Walther,

[137359]M. M.; Weng, Y.; Duan, D.–S. R.; Dean, M.; Glavac, D.;

Richards, F.

[137360]M.; Crossey, P. A.; Ferguson–Smith, M. A.; Le Paslier, D.;

Chumakov,

[137361]J.; Cohen, D.; Chinault, A. C.; Maher, E. R.; Linehan, W. M.;

Zbar,

[137362]B.; Lerman, M. I.: Identification of the von Hippel–Lindau

disease

[137363]tumor suppressor gene. Science 260: 1317–1320, 1993.

[137364]24516.

[137365]Lee, S.; Chen, D. Y. T.; Humphrey, J. S.; Gnarr, J. R.; Linehan,

[137366]W. M.; Klausner, R. D.: Nuclear/cytoplasmic localization of the von

[137367]Hippel–Lindau tumor suppressor gene product is determined by cell

[137368]density. Proc. Nat. Acad. Sci. 93: 1770–1775, 1996.

[137369]24517.

[137370]Lenz, T.; Thiede, H. M.; Nussberger, J.; Atlas, S. A.; Distler,

[137371]A.; Schulte, K. L.: Hyperreninemia and secondary hyperaldosteronism

[137372]in a patient with pheochromocytoma and von Hippel–Lindau disease. Nephron 62:

[137373]345–350, 1992.

[137374]24518.

[137375]Lindau, A.: Zur Frage der Angiomatosis Retinae und Ihrer Hirncomplication. Acta

[137376]Ophthal. 4: 193–226, 1927.

[137377]24519.

[137378]Loeb, D. B.; Pericak-Vance, M. A.; Stajich, J. M.; Vance, J. M.

[137379]: A novel mutation in the von Hippel-Lindau gene. Hum. Molec. Genet. 3:

[137380]1423-1424, 1994.

[137381]24520.

[137382]Lui, W. O.; Chen, J.; Glasker, S.; Bender, B. U.; Madura, C.;

[137383]Khoo, S. K.; Kort, E.; Larsson, C.; Neumann, H. P. H.; Teh, B. T.

[137384]: Selective loss of chromosome 11 in pheochromocytomas associated

[137385]with the VHL syndrome. Oncogene 21: 1117-1122, 2002.

[137386]24521.

[137387]Maddock, I. R.; Moran, A.; Maher, E. R.; Teare, M. D.; Norman,

[137388]A.; Payne, S. J.; Whitehouse, R.; Dodd, C.; Lavin, M.; Hartley, N.;

[137389]Super, M.; Evans, D. G. R.: A genetic register for von Hippel-Lindau

[137390]disease. J. Med. Genet. 33: 120-127, 1996.

[137391]24522.

[137392]Maher, E. R.; Bentley, E.; Yates, J. R. W.; Barton, D.; Jennings,

[137393]A.; Fellows, I. W.; Ponder, M. A.; Ponder, B. A. J.; Benjamin,

[137394]Harris, R.; Ferguson-Smith, M. A.: Mapping of von Hippel-Lindau disease

[137395]to chromosome 3p confirmed by genetic linkage analysis. J. Neurol.

[137396]Sci. 100: 27-30, 1990.

[137397]24523.

[137398]Maher, E. R.; Bentley, E.; Yates, J. R. W.; Latif, F.; Lerman,

[137399]M.; Zbar, B.; Affara, N. A.; Ferguson-Smith, M. A.: Mapping of the

[137400]von Hippel-Lindau disease locus to a small region of chromosome 3p

[137401]by genetic linkage analysis. Genomics 10: 957-960, 1991.

[137402]24524.

[137403]Maher, E. R.; Iselius, L.; Yates, J. R. W.; Littler, M.; Benjamin,

[137404]C.; Harris, R.; Sampson, J.; Williams, A.; Ferguson-Smith,

M. A.;

[137405]Morton, N.: Von Hippel–Lindau disease: a genetic study. J. Med.

[137406]Genet. 28: 443–447, 1991.

[137407]24525.

[137408]Maher, E. R.; Yates, J. R. W.; Ferguson–Smith, M. A.: Statistical

[137409]analysis of the two stage mutation model in von Hippel–Lindau disease,

[137410]and in sporadic cerebellar haemangioblastoma and renal cell carcinoma. J.

[137411]Med. Genet. 27: 311–314, 1990.

[137412]24526.

[137413]Mahon, P. C.; Hirota, K.; Semenza, G. L.: FIH–1: a novel protein

[137414]that interacts with HIF–1–alpha and VHL to mediate repression of HIF–1

[137415]transcriptional activity. Genes Dev. 15: 2675–2686, 2001.

[137416]

[137417]100. Manski, T. J.; Heffner, D. K.; Glenn, G. M.; Patronas, N. J.;

[137418]Pikus, A. T.; Katz, D.; Lebovics, R.; Sledjeski, K.; Choyke, P. L.;

[137419]Zbar, B.; Linehan, W. M.; Oldfield, E. H.: Endolymphatic sac tumors:

[137420]a source of morbid hearing loss in von Hippel–Lindau disease. J.A.M.A. 277:

[137421]1461–1466, 1997.

[137422]

[137423]101. Maxwell, P. H.; Wiesener, M. S.; Chang, G.–W.; Clifford, S. C.;

[137424]Vaux, E. C.; Cockman, M. E.; Wykoff, C. C.; Pugh, C. W.; Maher, E.

[137425]R.; Ratcliffe, P. J.: The tumour suppressor protein VHL targets hypoxia-inducible

[137426]factors for oxygen-dependent proteolysis. Nature 399: 271–275, 1999.

[137427]

[137428]102. McCabe, C. M.; Flynn, H. W., Jr.; Shields, C. L.; Shields, J.

[137429]A.; Regillo, C. D.; McDonald, H. R.; Berrocal, M. H.; Gass, J. D.

[137430]M.; Mieler, W. F.: Juxtapapillary capillary hemangiomas:

clinical

[137431] features and visual acuity outcomes. Ophthalmology 107:
2240–2249,

[137432] 2000.

[137433]

[137434] 103. McKusick, V. A.; Abbey, H.; Bettersby, E. J.; Borhani,
N. O.;

[137435] Boyer, S. H., IV; Cohen, B. H.; Ferguson–Smith, M. A.;
Franke, F.

[137436] R.; Gordon, H.; Handmaker, S. D.; Harris, W. S.; Hawkins,
M. R.;

[137437] and 13 others: Medical genetics 1960. J. Chronic Dis. 14:
1–198,

[137438] 1961. Fig. 71.

[137439]

[137440] 104. Melmon, K. L.; Rosen, S. W.: Lindau's disease: review
of the

[137441] literature and study of a large kindred. Am. J. Med. 36:
595–617,

[137442] 1964.

[137443]

- [137444]105. Min, J.-H.; Yang, H.; Ivan, M.; Gertler, F.; Kaelin, W. G., Jr.;
[137445]Pavletich, N. P.: Structure of an HIF-1- α -pVHL complex: hydroxyproline
[137446]recognition in signaling. Science 296: 1886-1889, 2002.
[137447]
[137448]106. Mukhopadhyay, D.; Knebelmann, B.; Cohen, H. T.;
Ananth, S.; Sukhatme,
[137449]V. P.: The von Hippel-Lindau tumor suppressor gene
product interacts
[137450]with Sp1 to repress vascular endothelial growth factor
promoter activity. Molec
[137451]. Cell. Biol. 17: 5629-5639, 1997.
[137452]
[137453]107. Neumann, H. P. H.; Berger, D. P.; Sigmund, G.; Blum,
U.; Schmidt,
[137454]D.; Parmer, R. J.; Volk, B.; Kirste, G.: Pheochromocytomas,
multiple
[137455]endocrine neoplasia type 2, and von Hippel-Lindau disease. New Eng.
[137456]J. Med. 329: 1531-1538, 1993.

[137457]

- [137458]108. Neumann, H. P. H.; Eng, C.; Mulligan, L. M.; Glavac, D.; Zauner, J.; Ponder, B. A. J.; Crossey, P. A.; Maher, E. R.; Brauch, H.:
Consequences
[137460]of direct genetic testing for germline mutations in the
clinical management
[137461]of families with multiple endocrine neoplasia, type II.
J.A.M.A. 274:
[137462]1149–1151, 1995.

[137463]

- [137464]109. Neumann, H. P. H.; Wiestler, O. D.: Clustering of features of
[137465]von Hippel–Lindau syndrome: evidence for a complex genetic locus. Lancet 337:
[137466]1052–1054, 1991.

[137467]

- [137468]110. Nibbelink, D. W.; Peters, B. H.; McCormick, W. F.: On the association
[137469]of pheochromocytoma and cerebellar hemangioblastoma. Neurology 19:

[137470]455–460, 1969.

[137471]

[137472]111. Oberstrass, J.; Reifenberger, G.; Reifenberger, J.;
Wechsler,

[137473]W.; Collins, V. P.: Mutation of the von Hippel–Lindau tu-
mour suppressor

[137474]gene in capillary haemangioblastomas of the central ner-
vous system. J.

[137475]Path. 179: 151–156, 1996.

[137476]

[137477]112. Ohh, M.; Yauch, R. L.; Lonergan, K. M.; Whaley, J. M.;
Stemmer–Rachamimov,

[137478]A. O.; Louis, D. N.; Gavin, B. J.; Kley, N.; Kaelin, W. G., Jr.;
Iliopoulos,

[137479]O.: The von Hippel–Lindau tumor suppressor protein is
required for

[137480]proper assembly of an extracellular fibronectin matrix.
Molec. Cell 1:

[137481]959–968, 1998.

[137482]

[137483]113. Olschwang, S.; Richard, S.; Boisson, C.; Giraud, S.;

Laurent-Puig,

[137484]P.; Resche, F.; Thomas, G.: Germline mutation profile of the VHL

[137485]gene in von Hippel-Lindau disease and in sporadic hemangioblastoma. Hum.

[137486]Mutat. 12: 424-430, 1998.

[137487]

[137488]114. Otenasek, F. J.; Silver, M. L.: Spinal hemangioma (hemangioblastoma)

[137489]in Lindau's disease: report of six cases in a single family. J. Neurosurg. 18:

[137490]295-300, 1961.

[137491]

[137492]115. Pack, S. D.; Zbar, B.; Pak, E.; Ault, D. O.; Humphrey, J. S.;

[137493]Pham, T.; Hurley, K.; Weil, R. J.; Park, W.-S.; Kuzmin, I.; Stolle,

[137494]C.; Glenn, G.; Liotta, L. A.; Lerman, M. I.; Klausner, R. D.; Linehan,

[137495]W. M.; Zhuang, Z.: Constitutional von Hippel-Lindau (VHL) gene deletions

[137496]detected in VHL families by fluorescence in situ hybridiza-

tion. Cancer

[137497]Res. 59: 5560–5564, 1999.

[137498]

[137499]116. Pause, A.; Lee, S.; Lonergan, K. M.; Klausner, R. D.:

The von

[137500]Hippel–Lindau tumor suppressor gene is required for cell
cycle exit

[137501]upon serum withdrawal. Proc. Nat. Acad. Sci. 95:
993–998, 1998.

[137502]

[137503]117. Price, E. B., Jr.: Papillary cystadenoma of the epi-
didymis:

[137504]a clinicopathologic analysis of 20 cases. Arch. Path. 91:
456–470,

[137505]1971.

[137506]

[137507]118. Probst, A.; Lotz, M.; Heitz, P.: Von Hippel–Lindau's
disease,

[137508]syringomyelia and multiple endocrine tumors: a complex
neuroendocrinopathy. Virchows

[137509]Arch. Path. Anat. Histol. 378: 265–272, 1978.

[137510]

[137511]119. Prowse, A. H.; Webster, A. R.; Richards, F. M.;
Richard, S.;

[137512]Olschwang, S.; Resche, F.; Affara, N. A.; Maher, E. R.: So-
matic inactivation

[137513]of the VHL gene in Von Hippel–Lindau disease tumors.
Am. J. Hum.

[137514]Genet. 60: 765–771, 1997.

[137515]

[137516]120. Rho, Y. M.: Von Hippel–Lindau's disease: a report of
5 cases. Canad.

[137517]Med. Assoc. J. 101: 135–142, 1969.

[137518]

[137519]121. Richard, S.; Croisille, L.; Yvart, J.; Casadeval, N.; Es-
chwege,

[137520]P.; Aghakhani, N.; David, P.; Gaudric, A.; Scigalla, P.; Her-
mine,

[137521]O.: Paradoxical secondary polycythemia in von Hippel–
Lindau patients

[137522]treated with anti–vascular endothelial growth factor re-
ceptor therapy. Blood 99:

[137523]3851–3853, 2002.

[137524]

[137525]122. Richards, F. M.; Crossey, P. A.; Phipps, M. E.; Foster, K.; Latif,

[137526]F.; Evans, G.; Sampson, J.; Lerman, M. I.; Zbar, B.; Affara, N. A.;

[137527]Ferguson-Smith, M. A.; Maher, E. R.: Detailed mapping of germline

[137528]deletions of the von Hippel-Lindau disease tumour suppressor gene. Hum.

[137529]Molec. Genet. 3: 595–598, 1994.

[137530]

[137531]123. Richards, F. M.; Maher, E. R.; Latif, F.; Phipps, M. E.; Tory,

[137532]K.; Lush, M.; Crossey, P. A.; Oostra, B.; Gustavson, K. H.; Green,

[137533]J.; Turner, G.; Yates, J. R. W.; Linehan, W. M.; Affara, N. A.; Lerman,

[137534]M.; Zbar, B.; Ferguson-Smith, M. A.: Detailed genetic mapping of

[137535]the von Hippel-Lindau disease tumour suppressor gene. J. Med. Genet. 30:

[137536]104–107, 1993.

[137537]

[137538]124. Richards, F. M.; Phipps, M. E.; Latif, F.; Yao, M.;
Crossey,

[137539]P. A.; Foster, K.; Linehan, W. M.; Affara, N. A.; Lerman, M.
I.; Zbar,

[137540]B.; Ferguson-Smith, M. A.; Maher, E. R.: Mapping the von
Hippel–Lindau

[137541]disease tumour suppressor gene: identification of
germline deletions

[137542]by pulsed field gel electrophoresis. Hum. Molec. Genet. 2:
879–882,

[137543]1993.

[137544]

[137545]125. Richards, F. M.; Schofield, P. N.; Fleming, S.; Maher,
E. R.

[137546]: Expression of the von Hippel–Lindau disease tumour
suppressor gene

[137547]during human embryogenesis. Hum. Molec. Genet. 5:
639–644, 1996.

[137548]

[137549]126. Rubenstein, J. L.; Yaari, H.: von Hippel–Lindau and
the genetics

[137550]of astrocytoma. J. Nat. Cancer Inst. 86: 142–143, 1994.

[137551]

[137552]127. Sander, S.; Normann, T.; Mathisen, W.: Pheochromo-
cytoma associated

[137553]with von–Hippel–Lindau's disease in a family. Scand. J.
Urol. Nephrol. 4:

[137554]259–263, 1970.

[137555]

[137556]128. Schechterman, L.: Lindau's disease: report of an un-
usual case

[137557]and two additional cases in a Negro family. Med. Ann.
D.C. 30: 64–76,

[137558]1961.

[137559]

[137560]129. Schimke, R. N.; Collins, D. L.; Rothberg, P. G.: Func-
tioning

[137561]carotid paraganglioma in the von Hippel–Lindau syn-
drome. (Letter) Am.

[137562]J. Med. Genet. 80: 533–534, 1998.

[137563]

[137564]130. Schoenfeld, A.; Davidowitz, E. J.; Burk, R. D.: A second major

[137565]native von Hippel-Lindau gene product, initiated from an internal

[137566]translation start site, functions as a tumor suppressor.

Proc. Nat.

[137567]Acad. Sci. 95: 8817-8822, 1998.

[137568]

[137569]131. Seizinger, B. R.; Farmer, G.; Haines, J.; Anderson, K.; Whaley,

[137570]J.; Hettlich, C.; Decker, J.; Rouleau, G.; Smith, D.; Drabkin, H.;

[137571]Filling-Katz, M.; Neumann, H.; Collins, D.; Hsia, E.; Green, J.; Waziri,

[137572]M.; Gusella, J.; Li, F.: Isolating the gene(s) for von Hippel-Lindau

[137573]disease and renal cell carcinoma. (Abstract) Am. J. Hum. Genet. 45

[137574](suppl.): A32, 1989.

[137575]

[137576]132. Seizinger, B. R.; Rouleau, G. A.; Ozelius, L. J.; Lane,
[137577]A. H.; Farmer, G. E.; Lamiell, J. M.; Haines, J.; Yuen, J.
[137578]W.; Collins, D.; Majoor-Krakauer, D.; et al.: Von Hippel-
Lindau
[137579]disease maps to the region of chromosome 3 associated
with renal cell
[137580]carcinoma. Nature 332: 268-269, 1988.

[137581]

[137582]133. Seizinger, B. R.; Smith, D. I.; Filling-Katz, M. R.; Neumann,

[137583]H.; Green, J. S.; Choyke, P. L.; Anderson, K. M.; Freiman, R. N.;

[137584]Klauck, S. M.; Whaley, J.; Decker, H.-J. H.; Hsia, Y. E.; Collins,

[137585]D.; Halperin, J.; Lamiell, J. M.; Oostra, B.; Waziri, M. H.; Gorin,

[137586]M. B.; Scherer, G.; Drabkin, H. A.; Aronin, N.; Schinzel, A.; Martuza,

[137587]R. L.; Gusella, J. F.; Haines, J. L.: Genetic flanking markers refine

[137588]diagnostic criteria and provide insights into the genetics of Von

[137589]Hippel Lindau disease. Proc. Nat. Acad. Sci. 88: 2864-2868, 1991.

[137590]

[137591]134. Sgambati, M. T.; Stolle, C.; Choyke, P. L.; Walther, M. M.; Zbar,

[137592]B.; Linehan, W. M.; Glenn, G. M.: Mosaicism in von Hippel-Lindau

[137593]disease: lessons from kindreds with germline mutations
identified

[137594]in offspring with mosaic parents. Am. J. Hum. Genet. 66:
84–91,

[137595]2000.

[137596]

[137597]135. Sharp, W. V.; Platt, R. L.: Familial pheochromocytoma:
association

[137598]with von-Hippel-Lindau's disease. Angiology 22:
141–146, 1971.

[137599]

[137600]136. Shokeir, M. H. K.: Von Hippel–Lindau syndrome: a
report on three

[137601]kindreds. J. Med. Genet. 7: 155–157, 1970.

[137602]

[137603]137. Silver, M. L.: Hereditary vascular tumors of the ner-
vous system. J.A.M.A.

[137604]156: 1053–1056, 1954.

[137605]

[137606]138. Stolle, C.; Glenn, G.; Zbar, B.; Humphrey, J. S.;
Choyke, P.;

[137607]Walther, M.; Pack, S.; Hurley, K.; Audrey, C.; Klausner, R.;
Linehan,

[137608]W. M.: Improved detection of germline mutations in the
von Hippel–Lindau

[137609]disease tumor suppressor gene. Hum. Mutat. 12:
417–423, 1998.

[137610]

[137611]139. Thomas, M.; Burnside, R. M.: Von Hippel–Lindau dis-
ease. Am.

[137612]J. Ophthalm. 51: 140–146, 1961.

[137613]

[137614]140. Tisherman, S. E.; Gregg, F. J.; Danowski, T. S.: Famil-
ial pheochromocytoma

[137615]. J.A.M.A. 182: 152–156, 1962.

[137616]

[137617]141. Tisherman, S. E.; Tisherman, B. G.; Tisherman, S. A.;
Dunmire,

[137618]S.; Levey, G. S.; Mulvihill, J. J.: Three-decade investigation
of

[137619]familial pheochromocytoma: an allele of von Hippel–Lin-
dau disease? Arch.

[137620]Int. Med. 153: 2550–2556, 1993.

[137621]

[137622]142. Tory, K.; Brauch, H.; Linehan, M.; Barba, D.; Oldfield, E.; Filling-Katz,

[137623]M.; Seizinger, B.; Nakamura, Y.; White, R.; Marshall, F. F.; Lerman,

[137624]M. I.; Zbar, B.: Specific genetic change in tumors associated with

[137625]von Hippel–Lindau disease. J. Nat. Cancer Inst. 81: 1097–1101, 1989.

[137626]

[137627]143. Tsuda, H.; Fukushima, S.; Takahashi, M.; Hikosaka, Y.; Hayashi,

[137628]K.: Familial bilateral papillary cystadenoma of the epididymis: report

[137629]of three cases in siblings. Cancer 37: 1831–1839, 1976.

[137630]

[137631]144. Tyers, M.; Willems, A. R.: One ring to rule a super-family of

[137632]E3 ubiquitin ligases. Science 284: 602–604, 1999.

[137633]

- [137634]145. Vance, J. M.; Small, K. W.; Jones, M. A.; Stajich, J. M.; Yamaoka, L. H.; Roses, A. D.; Hung, W.-Y.; Pericak-Vance, M. A.: Confirmation of linkage in von Hippel-Lindau disease. *Genomics* 6: 565-567, 1990.
- [137637]
- [137638]146. van der Harst, E.; de Krijger, R. R.; Dinjens, W. N. M.; Weeks, L. E.; Bonjer, H. J.; Bruining, H. A.; Lamberts, S. W. J.; Koper, J. W.: Germline mutations in the *vhl* gene in patients presenting with pheochromocytomas. *Int. J. Cancer*. 77: 337-340, 1998.
- [137642]
- [137643]147. Versteeg, R.: Aberrant methylation in cancer. (Editorial) *Am. J. Hum. Genet.* 60: 751-754, 1997.
- [137645]
- [137646]148. Vogelstein, B.: Personal Communication. Baltimore, Md. 1/6/1995.

[137647]

[137648]149. von Hippel, E.: Ueber eine sehr seltene Erkrankung
der Netzhaut. Albrecht

[137649]von Graefes Arch. Ophthal. 59: 83–106, 1904.

[137650]

[137651]150. Webster, A. R.; Maher, E. R.; Bird, A. C.; Moore, A. T.:
Risk

[137652]of multisystem disease in isolated ocular angioma
(haemangioblastoma). J.

[137653]Med. Genet. 37: 62–63, 2000.

[137654]

[137655]151. Webster, A. R.; Richards, F. M.; MacDonald, F. E.;
Moore, A.

[137656]T.; Maher, E. R.: An analysis of phenotypic variation in the
familial

[137657]cancer syndrome von Hippel–Lindau disease: evidence for
modifier effects. Am.

[137658]J. Hum. Genet. 63: 1025–1035, 1998.

[137659]

[137660]152. Wells, R. A.; Reenders, S. T.; Green, J.; Johnson, G. J.;
Robson,

- [137661]K. J. H.: Linkage studies of von Hippel–Lindau syndrome using multiple–locus hypervariable DNA probes. (Abstract) Cytogenet. Cell Genet. 46: 714, 1987.
- [137664]
- [137665]153. Wesolowski, D. P.; Ellwood, R. A.; Schwab, R. E.; Farah, J.: Hippel–Lindau syndrome in identical twins. Brit. J. Radiol. 54: 982–986, 1981.
- [137668]
- [137669]154. Wiesener, M. S.; Seyfarth, M.; Warnecke, C.; Jurgensen, J. S.; Rosenberger, C.; Morgan, N. V.; Maher, E. R.; Frei, U.; Eckardt, K.–U. Paraneoplastic erythrocytosis associated with an inactivating point mutation of the von Hippel–Lindau gene in a renal cell carcinoma. Blood 99: 3562–3565, 2002.

[137674]

[137675]155. Wise, K. S.; Gibson, J. A.: Von Hippel–Lindau's disease and

[137676]phaeochromocytoma. Brit. Med. J. 1: 441, 1971.

[137677]

[137678]156. Wyburn–Mason, R.: Arteriovenous aneurysm of mid-brain and retina,

[137679]facial naevi and mental changes. Brain 66: 163–203, 1943.

[137680]

[137681]157. Zatyka, M.; da Silva, N. F.; Clifford, S. C.; Morris, M. R.;

[137682]Wiesener, M. S.; Eckardt, K.–U.; Houlston, R. S.; Richards, F. M.;

[137683]Latif, F.; Maher, E. R.: Identification of cyclin D1 and other novel

[137684]targets for the von Hippel–Lindau tumor suppressor gene by expression

[137685]array analysis and investigation of cyclin D1 genotype as a modifier

[137686]in von Hippel–Lindau disease. Cancer Res. 62: 3803–3811, 2002.

[137687]

[137688]158. Zbar, B.; Kishida, T.; Chen, F.; Schmidt, L.; Maher, E. R.; Richards,

[137689]F. M.; Crossey, P. A.; Webster, A. R.; Affara, N. A.; Ferguson-Smith,

[137690]M. A.; Brauch, H.; Glavac, D.; and 14 others: Germline mutations

[137691]in the Von Hippel-Lindau disease (VHL) gene in families from North

[137692]America, Europe, and Japan. Hum. Mutat. 8: 348-357, 1996.

[137693]

[137694]159. Zhuang, Z.; Emmert-Buck, M. R.; Roth, M. J.; Gnarr, J.; Linehan,

[137695]W. M.; Liotta, L. A.; Lubensky, I. A.: Von Hippel-Lindau disease

[137696]gene deletion detected in microdissected sporadic human colon carcinoma

[137697]specimens. Hum. Path. 27: 152-156, 1996.

[137698]24527.

[137699]Devlin, R. H.; Deeb, S.; Brunsell, J.; Hayden, M. R.: Partial

[137700] gene duplication involving exon–Alu interchange results
in lipoprotein

[137701] lipase deficiency. *Am. J. Hum. Genet.* 46: 112–119, 1990.

[137702] 24528.

[137703] Dichek, H. L.; Fojo, S. S.; Beg, O. U.; Skarlatos, S. I.; Brun-
zell,

[137704] J. D.; Cutler, G. B., Jr.; Brewer, H. B., Jr.: Identification of

[137705] 2 separate allelic mutations in the lipoprotein lipase gene
of a patient

[137706] with the familial hyperchylomicronemia syndrome. *J. Biol.*
Chem. 266:

[137707] 473–477, 1991.

[137708] 24529.

[137709] Eckel, R. H.: Lipoprotein lipase: a multifunctional enzyme
relevant

[137710] to common metabolic diseases. *New Eng. J. Med.* 320:
1060–1068, 1989.

[137711] 24530.

[137712] Emi, M.; Hata, A.; Robertson, M.; Iverius, P.–H.; Hegele, R.;

[137713] Lalouel, J.–M.: Lipoprotein lipase deficiency resulting from
a nonsense

[137714]mutation in exon 3 of the lipoprotein lipase gene. Am. J. Hum. Genet. 47:

[137715]107–111, 1990.

[137716]24531.

[137717]Emi, M.; Wilson, D. E.; Iverius, P.–H.; Wu, L.; Hata, A.; Hegele,

[137718]R.; Williams, R. R.; Lalouel, J.–M.: Missense mutation (gly-to-glu188)

[137719]of human lipoprotein lipase imparting functional deficiency. J. Biol.

[137720]Chem. 265: 5910–5916, 1990.

[137721]24532.

[137722]Faustinella, F.; Chang, A.; Van Biervliet, J. P.; Rosseneu, M.;

[137723]Vinaimont, N.; Smith, L. C.; Chen, S.–H.; Chan, L.: Catalytic triad

[137724]residue mutation (asp156-to-gly) causing familial lipoprotein lipase

[137725]deficiency: co-inheritance with a nonsense mutation (ser447-to-ter)

[137726]in a Turkish family. J. Biol. Chem. 266: 14418–14424, 1991.

[137727]24533.

[137728]Feoli-Fonseca, J. C.; Levy, E.; Godard, M.; Lambert, M.: Familial

[137729]lipoprotein lipase deficiency in infancy: clinical, biochemical, and

[137730]molecular study. J. Pediat. 133: 417-423, 1998.

[137731]24534.

[137732]Fisher, R. M.; Humphries, S. E.; Talmud, P. J.: Common variation

[137733]in the lipoprotein lipase gene: effects on plasma lipids and risk

[137734]of atherosclerosis. Atherosclerosis 135: 145-159, 1997.

[137735]24535.

[137736]Franklin, S. M.: Splenomegaly with lipaemia. Proc. Roy. Soc.

[137737]Med. 30: 711, 1937.

[137738]24536.

[137739]Funke, H.; Klug, J.; Assmann, G.: Hind III RFLP in the lipoprotein

[137740]lipase gene, (LPL). Nucleic Acids Res. 15: 9102, 1987.

[137741]24537.

[137742]Funke, H.; Reckwerth, A.; Stapenhorst, D.; Schulze Beier-
ing, M.;

[137743]Jansen, M.; Assmann, G.: Bst NI (Eco RI) RFLP in the
lipoprotein

[137744]lipase gene (LPL). Nucleic Acids Res. 16: 2741, 1988.

[137745]24538.

[137746]Gagne, C.; Brun, L. D.; Julien, P.; Moorjani, S.; Lupien, P. J.

[137747]: Primary lipoprotein lipase activity deficiency: clinical in-
vestigation

[137748]of a French Canadian population. Canad. Med. Assoc. J.
140: 405-411,

[137749]1989.

[137750]24539.

[137751]Gilbert, B.; Rouis, M.; Griglio, S.; de Lumley, L.; Laplaud,
P.-M.

[137752]: Lipoprotein lipase (LPL) deficiency: a new patient ho-
mozygote for

[137753]the preponderant mutation gly88-to-glu in the human
LPL gene and review

[137754]of reported mutations: 75% are clustered in exons 5 and

6. Ann. Genet. 44:

[137755]25–32, 2001.

[137756]24540.

[137757]Ginzinger, D. G.; Lewis, M. E. S.; Ma, Y.; Jones, B. R.; Liu,

[137758]G.; Jones, S. D.; Hayden, M. R.: A mutation in the lipoprotein lipase

[137759]gene is the molecular basis of chylomicronemia in a colony of domestic

[137760]cats. J. Clin. Invest. 97: 1257–1266, 1996.

[137761]24541.

[137762]Gotoda, T.; Yamada, N.; Kawamura, M.; Kozaki, K.; Mori, N.; Ishibashi,

[137763]S.; Shimano, H.; Takaku, F.; Yazaki, Y.; Furuichi, Y.; Murase, T.

[137764]: Heterogeneous mutations in the human lipoprotein lipase gene in

[137765]patients with familial lipoprotein lipase deficiency. J. Clin. Invest. 88:

[137766]1856–1864, 1991.

[137767]24542.

[137768]Gotoda, T.; Yamada, N.; Murase, T.; Miyake, S.; Murakami,

R.;

[137769]Kawamura, M.; Kozaki, K.; Mori, N.; Shimano, H.; Shimada, M.; Yazaki,

[137770]Y.: A newly identified null allelic mutation in the human lipoprotein

[137771]lipase (LPL) gene of a compound heterozygote with familial LPL deficiency. *Biochim.*

[137772]*Biophys. Acta* 1138: 353-356, 1992.

[137773]24543.

[137774]Hata, A.; Emi, M.; Luc, G.; Basdevant, A.; Gambert, P.; Iverius,

[137775]P.-H.; Lalouel, J.-M.: Compound heterozygote for lipoprotein lipase

[137776]deficiency: ser-to-thr(244) and transition in 3-prime splice site

[137777]of intron 2 (AG-to-AA) in the lipoprotein lipase gene. *Am. J. Hum.*

[137778]*Genet.* 47: 721-726, 1990.

[137779]24544.

[137780]Hata, A.; Ridinger, D. N.; Sutherland, S. D.; Emi, M.; Kwong,

[137781]L. K.; Shuhua, J.; Lubbers, A.; Guy-Grand, B.; Basdevant,

A.; Iverius,

[137782]P. H.; Wilson, D. E.; Lalouel, J.-M.: Missense mutations in
exon

[137783]5 of the human lipoprotein lipase gene: inactivation cor-
relates with

[137784]loss of dimerization. J. Biol. Chem. 267: 20132-20139,
1992.

[137785]24545.

[137786]Haubenwallner, S.; Horl, G.; Shachter, N. S.; Presta, E.;
Fried,

[137787]S. K.; Hofler, G.; Kostner, G. M.; Breslow, J. L.; Zechner, R.:
A

[137788]novel missense mutation in the gene for lipoprotein lipase
resulting

[137789]in a highly conservative amino acid substitution
(asp180-to-glu) causes

[137790]familial chylomicronemia (type I hyperlipoproteinemia).
Genomics 18:

[137791]392-396, 1993.

[137792]24546.

[137793]Havel, R. J.; Gordon, R. S.: Idiopathic hyperlipemia:
metabolic

[137794]studies in an affected family. J. Clin. Invest. 39:
1777-1790, 1960.

[137795]24547.

[137796]Heaney, A. P.; Sharer, N.; Rameh, B.; Braganza, J. M.; Dur-
rington,

[137797]P. N.: Prevention of recurrent pancreatitis in familial
lipoprotein

[137798]lipase deficiency with high-dose antioxidant therapy. J.
Clin. Endocr.

[137799]Metab. 84: 1203-1205, 1999.

[137800]24548.

[137801]Heinzmann, C.; Ladas, J.; Antonarakis, S.; Kirchgessner,
T.;

[137802]Schotz, M.; Lasis, A. J.: RFLP for the human lipoprotein li-
pase (LPL)

[137803]gene: HindIII. Nucleic Acids Res. 15: 6763, 1987.

[137804]24549.

[137805]Heizmann, C.; Kirchgessner, T.; Kwiterovich, P. O.; Ladas,
J.

[137806]A.; Derby, C.; Antonarakis, S. E.; Lasis, A. J.: DNA poly-
morphism

[137807]haplotypes of the human lipoprotein lipase gene: possible association

[137808]with high density lipoprotein levels. Hum. Genet. 86: 578–584, 1991.

[137809]24550.

[137810]Henderson, H.; Ma, Y.; Kastelein, J.; Roederer, G.; Julien, P.;

[137811]Brunzell, J.; Hayden, M. R.: Identification of the molecular defects

[137812]underlying chylomicronemia in the majority of 75 separate probands

[137813]with LPL deficiency.(Abstract) Clin. Res. 39: 336A, 1991.

[137814]24551.

[137815]Henderson, H. E.; Bijvoet, S. M.; Mannens, M. A. M. M.; Bruin,

[137816]T.; Erkelens, D. W.; Hayden, M. R.; Kastelein, J. J. P.: Ile225-to-thr

[137817]loop mutation in the lipoprotein lipase (LPL) gene is a de novo event. Am.

[137818]J. Med. Genet. 78: 313–316, 1998.

[137819]24552.

[137820]Sjlander, A.; Birgander, R.; Rannug, A.; Alexandrie, A.-K.;
Tornling,

[137821]G.; Beckman, G.: Association between the p21 codon
31A1 (arg) allele

[137822]and lung cancer. Hum. Hered. 46: 221-225, 1996.

[137823]24553.

[137824]Fryns, J. P.; van den Berghe, K.; van Assche, A.; van den
Berghe,

[137825]H.: Prenatal diagnosis of campomelic dwarfism. Clin.
Genet. 19:

[137826]199-201, 1981.

[137827]24554.

[137828]Mellows, H. J.; Pryse-Davies, J.; Bennett, M. J.; Carter, C. O.

[137829]: The campomelic syndrome in two female siblings. Clin.
Genet. 18:

[137830]137-141, 1980.

[137831]24555.

[137832]Bjursell, C.; Wahlstrom, J.; Berg, K.; Stibler, H.; Kristians-
son,

[137833]B.; Matthijs, G.; Martinsson, T.: Detailed mapping of the
phosphomannomutase

[137834]2 (PMM2) gene and mutation detection enable improved analysis for

[137835]Scandinavian CDG type I families. *Europ. J. Hum. Genet.* 6: 603–611,

[137836]1998.

[137837]24556.

[137838]Satoyoshi, E.; Kowa, H.: A myopathy due to glycolytic abnormality. *Arch. Neurol.* 17: 248–256, 1967.

[137840]24557.

[137841]Servidei, S.; Bonilla, E.; Diedrich, R. G.; Kornfeld, M.; Oates,

[137842]J. D.; Davidson, M.; Vora, S.; DiMauro, S.: Fatal infantile form

[137843]of muscle phosphofructokinase deficiency. *Neurology* 36: 1465–1470,

[137844]1986.

[137845]24558.

[137846]Tani, K.; Fujii, H.; Takegawa, S.; Miwa, S.; Koyama, W.; Kanayama,

[137847]M.; Imanaka, A.; Imanaka, F.; Kuramoto, A.: Two cases of

phosphofructokinase

[137848]deficiency associated with congenital hemolytic anemia
found in Japan. Am.

[137849]J. Hemat. 14: 165-174, 1983.

[137850]24559.

[137851]Tarui, S.; Okuno, G.; Ikura, Y.; Tanaka, T.; Suda, M.;
Nishikawa,

[137852]M.: Phosphofructokinase deficiency in skeletal muscle: a
new type

[137853]of glycogenosis. Biochem. Biophys. Res. Commun. 19:
517-523, 1965.

[137854]24560.

[137855]Tsuji, S.; Servidei, S.; Tonin, P.; Shanske, S.; Azan, G.;
DiMauro,

[137856]S.: Identification of three novel mutations in non-
Ashkenazi Italian

[137857]patients with muscle phosphofructokinase deficiency. Am.
J. Hum.

[137858]Genet. 54: 812-819, 1994.

[137859]24561.

[137860]Matsumoto, K.; Yoshitomi, H.; Rossant, J.; Zaret, K. S.:

Liver

[137861]organogenesis promoted by endothelial cells prior to vascular function. Science 294:

[137862]559–563, 2001.

[137863]24562.

[137864]Matthews, W.; Jordan, C. T.; Gavin, M.; Jenkins, N. A.; Copeland,

[137865]N. G.; Lemischka, I. R.: A receptor tyrosine kinase cDNA isolated

[137866]from a population of enriched primitive hematopoietic cells and exhibiting

[137867]close genetic linkage to c-kit. Proc. Nat. Acad. Sci. 88: 9026–9030,

[137868]1991.

[137869]24563.

[137870]Millauer, B.; Shawver, L. K.; Plate, K. H.; Risau, W.; Ullrich,

[137871]A.: Glioblastoma growth inhibited in vivo by a dominant-negative

[137872]Flk-1 mutant. Nature 367: 576–579, 1994.

[137873]24564.

[137874]Millauer, B.; Witzigmann-Voos, S.; Schnurch, H.; Martinez,

R.;

[137875]Moller, N. P. H.; Risau, W.; Ullrich, A.: High affinity VEGF binding

[137876]and developmental expression suggest Flk-1 as a major regulator of

[137877]vasculogenesis and angiogenesis. Cell 72: 835-846, 1993.

[137878]24565.

[137879]Plate, K. H.; Breier, G.; Millauer, B.; Ullrich, A.; Risau, W.

[137880]: Up-regulation of vascular endothelial growth factor and its cognate

[137881]receptors in a rat glioma model of tumor angiogenesis. Cancer Res. 53:

[137882]5822-5827, 1993.

[137883]24566.

[137884]Quinn, T. P.; Peters, K. G.; De Vries, C.; Ferrara, N.; Williams,

[137885]L. T.: Fetal liver kinase 1 is a receptor for vascular endothelial

[137886]growth factor and is selectively expressed in vascular endothelium. Proc.

[137887]Nat. Acad. Sci. 90: 7533-7537, 1993.

[137888]24567.

[137889]Sait, S. N. J.; Dougher-Vermazen, M.; Shows, T. B.; Ter-
man, B.

[137890]J.: The kinase insert domain receptor gene (KDR) has been
relocated

[137891]to chromosome 4q11-q12. Cytogenet. Cell Genet. 70:
145-146, 1995.

[137892]24568.

[137893]Shalaby, F.; Rossant, J.; Yamaguchi, T. P.; Gertsenstein, M.;

[137894]Wu, X.-F.; Breitman, M. L.; Schuh, A. C.: Failure of blood-
island

[137895]formation and vasculogenesis in Flk-1-deficient mice. Na-
ture 376:

[137896]62-65, 1995.

[137897]24569.

[137898]Shay-Salit, A.; Shushy, M.; Wolfowitz, E.; Yahav, H.; Bre-
viario,

[137899]F.; Dejana, E.; Resnick, N.: VEGF receptor 2 and the ad-
herens junction

[137900]as a mechanical transducer in vascular endothelial cells.
Proc. Nat.

[137901]Acad. Sci. 99: 9462–9467, 2002.

[137902]24570.

[137903]Spritz, R. A.; Strunk, K. M.; Lee, S.-T.; Lu-Kuo, J. M.; Ward,

[137904]D. C.; Le Paslier, D.; Altherr, M. R.; Dorman, T. E.; Moir, D.
T.

[137905]; A YAC contig spanning a cluster of human type III recep-
tor protein

[137906]tyrosine kinase genes (PDGFRA-KIT-KDR) in chromosome
segment 4q12. Genomics 22:

[137907]431–436, 1994.

[137908]24571.

[137909]Terman, B.; Carrion, M.; Eddy, R. L.; Shows, T. B.: Tyrosine

[137910]kinase growth factor receptor (KDR) localized to chromo-
some 4 (Abstract) Cytogenet.

[137911]Cell Genet. 58: 1890 only, 1991.

[137912]24572.

[137913]Terman, B. I.; Carrion, M. E.; Kovacs, E.; Rasmussen, B. A.;
Eddy,

[137914]R. L.; Shows, T. B.: Identification of a new endothelial cell
growth

[137915]factor receptor tyrosine kinase. Oncogene 6: 1677–1683, 1991.

[137916]24573.

[137917]Terman, B. I.; Jani-Sait, S.; Carrion, M. E.; Shows, T. B.: The

[137918]KDR gene maps to human chromosome 4q31.2–q32, a locus which is distinct

[137919]from locations for other type III growth factor receptor tyrosine

[137920]kinases. Cytogenet. Cell Genet. 60: 214–215, 1992.

[137921]24574.

[137922]Yamashita, J.; Itoh, H.; Hirashima, M.; Ogawa, M.; Nishikawa,

[137923]S.; Yurugi, T.; Naito, M.; Nakao, K.; Nishikawa, S.–I.: Flk1–positive

[137924]cells derived from embryonic stem cells serve as vascular progenitors. Nature 408:

[137925]92–96, 2000.

[137926]24575.

[137927]Ziegler, B. L.; Valtieri, M.; Porada, G. A.; De Maria, R.; Muller,

[137928]R.; Masella, B.; Gabbianelli, M.; Casella, I.; Pelosi, E.; Bock,

T.;

[137929]Zanjani, E. D.; Peschle, C.: KDR receptor: a key marker
defining

[137930]hematopoietic stem cells. Science 285: 1553–1558, 1999.

[137931]24576.

[137932]Ala-Mello, S.; Sankila, E.-M.; Koskimies, O.; de la
Chapelle, A.;

[137933]Kaariainen, H.: Molecular studies in Finnish patients with
familial

[137934]juvenile nephronophthisis exclude a founder effect and
support a common

[137935]mutation causing mechanism. J. Med. Genet. 35:
279–283, 1998.

[137936]24577.

[137937]Alexander, F.; Campbell, S.: Familial uremic medullary
cystic

[137938]disease. Pediatrics 45: 1024–1028, 1970.

[137939]24578.

[137940]Antignac, C.; Arduy, C. H.; Beckmann, J. S.; Benessy, F.;

Gros,

[137941]F.; Medhioub, M.; Hildebrandt, F.; Dufier, J.-L.;

Kleinknecht, C.;

[137942]Broyer, M.; Weissenbach, J.; Habib, R.; Cohen, D.: A gene
for familial

[137943]juvenile nephronophthisis (recessive medullary cystic kidney disease)

[137944]maps to chromosome 2p. Nature Genet. 3: 342-345,
1993.

[137945]24579.

[137946]Benzing, T.; Gerke, P.; Hopker, K.; Hildebrandt, F.; Kim, E.;
Walz,

[137947]G.: Nephrocystin interacts with Pyk2, p130(Cas), and
tensin and triggers

[137948]phosphorylation of Pyk2. Proc. Nat. Acad. Sci. 98:
9784-9789, 2001.

[137949]24580.

[137950]Betz, R.; Rensing, C.; Otto, E.; Mincheva, A.; Zehnder, D.;
Lichter,

[137951]P.; Hildebrandt, F.: Children with ocular motor apraxia
type Cogan

[137952]carry deletions in the gene (NPHP1) for juvenile
nephronophthisis. J.

[137953]Pediat. 136: 828-831, 2000.

[137954]24581.

[137955]Broberger, O.; Winberg, J.; Zetterstrom, R.: Juvenile nephronophthisis.

[137956]I. A genetically determined nephropathy with hypotonic polyuria and

[137957]azotaemia. Acta Paediat. 49: 470–479, 1960.

[137958]24582.

[137959]Caridi, G.; Murer, L.; Bellantuono, R.; Sorino, P.; Caringella,

[137960]D. A.; Gusmano, R.; Ghiggeri, G. M.: Renal–retinal syndromes: association

[137961]of retinal anomalies and recessive nephronophthisis in patients with

[137962]homozygous deletion of the NPH1 locus. Am. J. Kidney Dis. 32: 1059–1062,

[137963]1998.

[137964]24583.

[137965]Cohen, A. H.; Hoyer, J. R.: Nephronophthisis: a primary tubular

[137966]basement membrane defect. Lab. Invest. 55: 564–572, 1986.

[137967]24584.

[137968]Collan, Y.; Sipponen, P.; Haapanen, E.; Lindahl, J.; Jokinen,
 [137969]E. J.; Hjelt, L.: Hereditary nephronophthisis with a life span
 of
 [137970]3 decades--light and electron microscopical, immunohis-
 tochemical,
 [137971]clinical and family studies. Wilhelm Roux. Arch. Develop.
 Biol. 376:
 [137972]195-208, 1977.
 [137973]24585.

[137974]Donaldson, J. C.; Dempsey, P. J.; Reddy, S.; Bouton, A. H.;
 Coffey,
 [137975]R. J.; Hanks, S. K.: Crk-associated substrate p130(Cas) in-
 teracts
 [137976]with nephrocystin and both proteins localize to cell-cell
 contacts
 [137977]of polarized epithelial cells. Exp. Cell Res. 256: 168-175,
 2000.
 [137978]24586.

[137979]Fanconi, G.; Hanhart, E.; Von Albertini, A.; Uehlinger, E.;
 Dolivo,
 [137980]G.; Prader, A.: Die familiaere juvenile Nephronophthise.
 (Die idiopathische

[137981]Parenchymatose). *Helv. Paediat. Acta* 6: 1–49, 1951.

[137982]24587.

[137983]Gibson, A. A. M.; Arneil, G. C.: Nephronophthisis: report of

[137984]8 cases in Britain. *Arch. Dis. Child.* 47: 84–89, 1972.

[137985]24588.

[137986]Giselson, N.; Heinegard, D.; Holmberg, C. G.; Lindberg, L.
G.;

[137987]Lindstedt, E.; Lindstedt, G.; Schersten, B.: Renal medullary
cystic

[137988]disease or familial juvenile nephronophthisis: a renal
tubular disease. *Am.*

[137989]*J. Med.* 48: 174–184, 1970.

[137990]24589.

[137991]Hackzell, G.; Lundmark, C.: Familial juvenile nephronoph-
thisis. *Acta*

[137992]*Paediat.* 47: 428–440, 1958.

[137993]24590.

[137994]Herdman, R. C.; Good, R. A.; Vernier, R. L.: Medullary cys-
tic

[137995]disease in two siblings. *Am. J. Med.* 43: 335–344, 1967.

[137996]24591.

[137997]Hildebrandt, F.: Personal Communication. Freiburg, Germany

[137998]10/1/1997.

[137999]24592.

[138000]Hildebrandt, F.; Otto, E.; Rensing, C.; Nothwang, H. G.; Vollmer,

[138001]M.; Adolphs, J.; Hanusch, H.; Brandis, M.: A novel gene encoding

[138002]an SH3 domain protein is mutated in nephronophthisis type 1. Nature

[138003]Genet. 17: 149–153, 1997.

[138004]24593.

[138005]Hildebrandt, F.; Singh–Sawhney, I.; Schnieders, B.; Papenfuss,

[138006]T.; Brandis, M.; Members of the APN Study Group: Refined genetic

[138007]mapping of a gene for familial juvenile nephronophthisis (NPH1) and

[138008]physical mapping of linked markers. Genomics 25: 360–364, 1995.

[138009]24594.

[138010]Khoury, Z.; Brezis, M.; Mogle, P.: Familial medullary
sponge

[138011]kidney in association with congenital absence of teeth
(anodontia). Nephron 48:

[138012]231–233, 1988.

[138013]24595.

[138014]Cho, K. R.; Oliner, J. D.; Simons, J. W.; Hedrick, L.; Fearon,

[138015]E. R.; Preisinger, A. C.; Hedge, P.; Silverman, G. A.; Vogel-
stein,

[138016]B.: The DCC gene: structural analysis and mutations in
colorectal

[138017]carcinomas. Genomics 19: 525–531, 1994.

[138018]24596.

[138019]Fazeli, A.; Dickinson, S. L.; Hermiston, M. L.; Tighe, R. V.;
Steen,

[138020]R. G.; Small, C. G.; Stoeckli, E. T.; Keino-Masu, K.; Masu,
M.; Rayburn,

[138021]H.; Simons, J.; Bronson, R. T.; Gordon, J. I.; Tessier-Lavi-
gne, M.;

[138022]Weinberg, R. A.: Phenotype of mice lacking functional

Deleted in

[138023]colorectal cancer (Dcc) gene. Nature 386: 796–804, 1997.

[138024]24597.

[138025]Fearon, E. R.; Cho, K. R.; Nigro, J. M.; Kern, S. E.; Simons, J.

[138026]W.; Ruppert, J. M.; Hamilton, S. R.; Preisinger, A. C.;

Thomas, G.;

[138027]Kinzler, K. W.; Vogelstein, B.: Identification of a chromo-
some 18q

[138028]gene that is altered in colorectal cancers. Science 247:
49–56,

[138029]1990.

[138030]24598.

[138031]Gotley, D. C.; Reeder, J. A.; Fawcett, J.; Walsh, M. D.; Bates,

[138032]P.; Simmons, D. L.; Antalis, T. M.: The deleted in colon
cancer (DCC)

[138033]gene is consistently expressed in colorectal cancers and
metastases. Oncogene 13:

[138034]787–795, 1996.

[138035]24599.

[138036]Hohne, M. W.; Halatsch, M.–E.; Kahl, G. F.; Weinel, R. J.:

Frequent

- [138037]loss of expression of the potential tumor suppressor gene
DCC in ductal
[138038]pancreatic adenocarcinoma. Cancer Res. 52: 2616–2619,
1992.
[138039]24600.
- [138040]Jen, J.; Kim, H.; Piantadosi, S.; Liu, Z.-F.; Levitt, R. C.; Sis-
tonen,
[138041]P.; Kinzler, K. W.; Vogelstein, B.; Hamilton, S. R.: Allelic
loss
[138042]of chromosome 18q and prognosis in colorectal cancer.
New Eng. J.
[138043]Med. 331: 213–221, 1994.
[138044]24601.
- [138045]Keino-Masu, K.; Masu, M.; Hinck, L.; Leonardo, E. D.;
Chan, S.
[138046]S.-Y.; Culotti, J. G.; Tessier-Lavigne, M.: Deleted in col-
orectal
[138047]cancer (DCC) encodes a netrin receptor. Cell 87: 175–185,
1996.
[138048]24602.
- [138049]Keynes, R.; Cook, G. M. W.: Axon guidance molecules. Cell

[138050]161–169, 1995.

[138051]24603.

[138052]Kolodziej, P. A.; Timpe, L. C.; Mitchell, K. J.; Fried, S. R.;

[138053]Goodman, C. S.; Jan, L. Y.; Jan, Y. N.: Frazzled encodes a
Drosophila

[138054]member of the DCC immunoglobulin subfamily and is re-
quired for CNS

[138055]and motor axon guidance. Cell 87: 197–204, 1996.

[138056]24604.

[138057]Maesawa, C.; Tamura, G.; Ogasawara, S.; Suzuki, Y.;
Sakata, K.;

[138058]Sugimura, J.; Nishizuka, S.; Sato, N.; Ishida, K.; Saito, K.;
Satodate,

[138059]R.: Loss of heterozygosity at the DCC gene locus is not
crucial for

[138060]the acquisition of metastatic potential in oesophageal
squamous cell

[138061]carcinoma. Europ. J. Cancer. 32A: 896–898, 1996.

[138062]24605.

[138063]Mehlen, P.; Rabizadeh, S.; Snipas, S. J.; Assa-Munt, N.;

Salvesen,

[138064]G. S.; Bredesen, D. E.: The DCC gene product induces apoptosis by

[138065]a mechanism requiring receptor proteolysis. Nature 395: 801–804,

[138066]1998.

[138067]24606.

[138068]Miyake, S.; Nagai, K.; Yoshino, K.; Oto, M.; Endo, M.; Yuasa,

[138069]Y.: Point mutations and allelic deletion of tumor suppressor gene

[138070]DCC in human esophageal squamous cell carcinomas and their relation

[138071]to metastasis. Cancer Res. 54: 3007–3010, 1994.

[138072]24607.

[138073]Nigro, J. M.; Cho, K. R.; Fearon, E. R.; Kern, S. E.; Ruppert,

[138074]J. M.; Oliner, J. D.; Kinzler, K. W.; Vogelstein, B.: Scrambled exons. Cell 64:

[138075]607–613, 1991.

[138076]24608.

[138077]Shibata, D.; Reale, M. A.; Lavin, P.; Silverman, M.; Fearon,

E.

[138078]R.; Steele, G., Jr.; Jessup, J. M.; Loda, M.; Summerhayes, I.

C.:

[138079]The DCC protein and prognosis in colorectal cancer. New

Eng. J. Med. 335:

[138080]1727-1732, 1996.

[138081]24609.

[138082]Stein, E.; Tessier-Lavigne, M.: Hierarchical organization of

[138083]guidance receptors: silencing of netrin attraction by Slit
through

[138084]a Robo/DCC receptor complex. Science 291: 1928-1938,
2001.

[138085]24610.

[138086]Tanaka, K.; Oshimura, M.; Kikuchi, R.; Seki, M.; Hayashi,
T.;

[138087]Miyaki, M.: Suppression of tumorigenicity in human colon
carcinoma

[138088]cells by introduction of normal chromosome 5 or 18. Na-
ture 349:

[138089]340-342, 1991.

[138090]24611.

[138091]Uchino, S.; Tsuda, H.; Noguchi, M.; Yokota, J.; Terada, M.;
Saito,
[138092]T.; Kobayashi, M.; Sugimura, T.; Hirohashi, S.: Frequent
loss of
[138093]heterozygosity at the DCC locus in gastric cancer. Cancer
Res. 52:
[138094]3099–3102, 1992.
[138095]24612.
[138096]Vogelstein, B.: Personal Communication. Baltimore, Md.
11/30/1995.
[138097]24613.
[138098]Zetter, B. R.: Adhesion molecules in tumor metastasis.
Semin.
[138099]Cancer Biol. 4: 219–229, 1993.
[138100]24614.
[138101]Devor, E. J.; Grandy, D. K.; Civelli, O.; Litt, M.; Burgess, A.
[138102]K.; Isenberg, K. E.; van de Wetering, B. J. M.; Oostra, B.:
Genetic
[138103]linkage is excluded for the D2–dopamine receptor
lambda–HD2G1 and
[138104]flanking loci on chromosome 11q22–q23 in Tourette syn–

drome. Hum.

[138105]Hered. 40: 105–108, 1990.

[138106]24615.

[138107]Voutsinas, S.; Wynne–Davies, R.: The infrequency of malignant

[138108]disease in diaphyseal aclasis and neurofibromatosis. J. Med. Genet. 20:

[138109]345–349, 1983.

[138110]24616.

[138111]Freemantle, S. J.; Taylor, S. M.; Krystal, G.; Moran, R. G.: Upstream

[138112]organization of and multiple transcripts from the human folylpoly–gamma–glutamate

[138113]synthetase gene. J. Biol. Chem. 270: 9579–9584, 1995.

[138114]24617.

[138115]Garrow, T. A.; Admon, A.; Shane, B.: Expression cloning of a human

[138116]cDNA encoding folylpoly(gamma–glutamate) synthetase and determination

[138117]of its primary structure. Proc. Nat. Acad. Sci. 89: 9151–9155, 1992.

[138118]24618.

[138119]Jones, C.; Kao, F.-T.: Regional mapping of the folylpolyglutamate

[138120]synthetase gene (FPGS) to 9cen-q34. (Abstract) Cytogenet. Cell Genet. 37:

[138121]499-500, 1984.

[138122]24619.

[138123]Jones, C.; Kao, F.-T.; Taylor, R. T.: Chromosomal assignment of

[138124]the gene for folylpolyglutamate synthetase to human chromosome 9. Cytogenet.

[138125]Cell Genet. 28: 181-194, 1980.

[138126]24620.

[138127]Jones, C.; Kao, F. T.: Assignment of the human gene complementing

[138128]the auxotrophic marker GAT-minus to chromosome 9. (Abstract) Cytogenet.

[138129]Cell Genet. 25: 168, 1979.

[138130]24621.

[138131]Kao, F. T.; Puck, T. T.: Genetics of somatic mammalian cells.

[138132]VII. Induction and isolation of nutritional mutants in Chinese hamster

[138133]cells. Proc. Nat. Acad. Sci. 60: 1275–1281, 1968.

[138134]24622.

[138135]Sussman, D. J.; Milman, G.; Shane, B.: Characterization of human

[138136]folylpolyglutamate synthetase expressed in Chinese hamster ovary cells. Somat.

[138137]Cell Molec. Genet. 12: 531–540, 1986.

[138138]24623.

[138139]Taylor, R. T.; Hanna, M. L.: Folate-dependent enzymes in cultured

[138140]Chinese hamster cells: folylpolyglutamate synthetase and its absence

[138141]in mutants auxotrophic for glycine, adenosine and thymidine. Arch.

[138142]Biochem. Biophys. 181: 331–344, 1977.

[138143]24624.

[138144]Taylor, S. M.; Freemantle, S. J.; Moran, R. G.: Structural organization

[138145]of the human folylpoly- γ -glutamate synthetase

gene: evidence for

[138146]a single genomic locus. Cancer Res. 55: 6030–6034,
1995.

[138147]24625.

[138148]Buters, J. T. M.; Tang, B.-K.; Pineau, T.; Gelboin, H. V.;
Kimura,

[138149]S.; Gonzalez, F. J.: Role of CYP1A2 in caffeine pharmacoki-
netics

[138150]and metabolism: studies using mice deficient in CYP1A2.
Pharmacogenetics 6:

[138151]291–296, 1996.

[138152]24626.

[138153]Butler, M. A.; Iwasaki, M.; Guengerich, F. P.; Kadlubar, F. F.

[138154]: Human cytochrome P-450(PA) (P-450IA2), the
phenacetin O-deethylase,

[138155]is primarily responsible for the hepatic 3-demethylation
of caffeine

[138156]and N-oxidation of carcinogenic arylamines. Proc. Nat.
Acad. Sci. 86:

[138157]7696–7700, 1989.

[138158]24627.

[138159]Christiansen, L.; Bygum, A.; Jensen, A.; Thomsen, K.;
Brandrup,
[138160]F.; Horder, M.; Petersen, N. E.: Association between
CYP1A2 polymorphism
[138161]and susceptibility to porphyria cutanea tarda. Hum. Genet.
107:
[138162]612–614, 2000.
[138163]24628.

[138164]Devonshire, H. W.; Kong, I.; Cooper, M.; Sloan, T. P.; Idle,
J.
[138165]R.; Smith, R. L.: The contribution of genetically determined
oxidation
[138166]status to inter-individual variation in phenacetin disposi-
tion. Brit.
[138167]J. Clin. Pharm. 16: 157–166, 1983.
[138168]24629.

[138169]Ikeya, K.; Jaiswal, A. K.; Owens, R. A.; Jones, J. E.; Nebert,
[138170]D. W.; Kimura, S.: Human CYP1A2: sequence, gene struc-
ture, comparison
[138171]with the mouse and rat orthologous gene, and differences
in liver

[138172]1A2 mRNA expression. Molec. Endocr. 3: 1399–1408,
1989.

[138173]24630.

[138174]Jaiswal, A. K.; Nebert, D. W.; McBride, O. W.; Gonzalez, F. J.

[138175]: Human P(3)450: cDNA and complete protein sequence,
repetitive Alu

[138176]sequences in the 3–prime nontranslated region, and lo-
calization of

[138177]gene to chromosome 15. J. Exp. Path. 3: 1–17, 1987.

[138178]24631.

[138179]Liang, H.–C. L.; Li, H.; McKinnon, R. A.; Duffy, J. J.; Potter,

[138180]S. S.; Puga, A.; Nebert, D. W.: Cyp1a2(–/–) null mutant
mice develop

[138181]normally but show deficient drug metabolism. Proc. Nat.
Acad. Sci. 93:

[138182]1671–1676, 1996.

[138183]24632.

[138184]Nebert, D. W.: Personal Communication. Bethesda, Md.
2/3/1988.

[138185]24633.

[138186]Sesardic, D.; Boobis, A. R.; Edwards, R. J.; Davies, D. S.: A

[138187]form of cytochrome P450 in man, orthologous to form d
in the rat,

[138188]catalyses the O-deethylation of phenacetin and is in-
ducible by cigarette

[138189]smoking. Brit. J. Clin. Pharm. 26: 363–372, 1988.

[138190]24634.

[138191]Tantcheva-Poor, I.; Zaigler, M.; Rietbrock, S.; Fuhr, U.: Es-
timation

[138192]of cytochrome P-450 CYP1A2 activity in 863 healthy Cau-
casians using

[138193]a saliva-based caffeine test. Pharmacogenetics 9:
131–144, 1999.

[138194]24635.

[138195]Wooding, S. P.; Watkins, W. S.; Bamshad, M. J.; Dunn, D.
M.; Weiss,

[138196]R. B.; Jorde, L. B.: DNA sequence variation in a 3.7-kb
noncoding

[138197]sequence 5-prime of the CYP1A2 gene: implications for
human population

[138198]history and natural selection. Am. J. Hum. Genet. 71:
528–542, 2002.

[138199]24636.

[138200]Bale, A. E.; Mitchell, A. L.; Gonzalez, F. J.; McBride, O. W.:

[138201]Localization of CYP2F1 by multipoint linkage analysis and
pulsed-field

[138202]gel electrophoresis. Genomics 10: 284-286, 1991.

[138203]24637.

[138204]Mitchell, A. L.; Bale, A. E.; Gonzalez, F.; McBride, O. W.:
Mapping

[138205]of cytochrome p450 IIA, IIB, and IIF subfamilies on chro-
mosome 19

[138206]by linkage analysis. (Abstract) Cytogenet. Cell Genet. 51:
1045,

[138207]1989.

[138208]24638.

[138209]Nhamburo, P. T.; Kimura, S.; McBride, O. W.; Kozak, C. A.;
Gelboin,

[138210]H. V.; Gonzalez, F. J.: The human CYP2F gene subfamily:
identification

[138211]of a cDNA encoding a new cytochrome P450, cDNA-di-
rected expression,

[138212]and chromosome mapping. Biochemistry 29: 5491-5499,

1990.

[138213]24639.

[138214]Thum, T.; Borlak, J.: Gene expression in distinct regions of
the

[138215]heart. Lancet 355: 979–983, 2000.

[138216]24640.

[138217]Wells, D. E.; Hill, A.; Lin, X.; Ahn, J.; Brown, N.; Wagner, M.

[138218]J.: Identification of novel mutations in the human EXT1 tu-
mor suppressor

[138219]gene. Hum. Genet. 99: 612–615, 1997.

[138220]24641.

[138221]Wicklund, C. L.; Pauli, R. M.; Johnston, D.; Hecht, J. T.:

Natural

[138222]history study of hereditary multiple exostoses. Am. J. Med.
Genet. 55:

[138223]43–46, 1995.

[138224]24642.

[138225]Wuyts, W.; Van Hul, W.: Molecular basis of multiple exos-
toses:

[138226]mutations in the EXT1 and EXT2 genes. Hum. Mutat. 15:

220–227, 2000.

[138227]24643.

[138228]Wuyts, W.; Van Hul, W.; De Boulle, K.; Hendrickx, J.;
Bakker,

[138229]E.; Vanhoenacker, F.; Mollica, F.; Ludecke, H.-J.; Sayli, B.
S.; Pazzaglia,

[138230]U. E.; Mortier, G.; Hamel, B.; Conrad, E. U.; Matsushita, M.;
Raskind,

[138231]W. H.; Willems, P. J.: Mutations in the EXT1 and EXT2
genes in hereditary

[138232]multiple exostoses. Am. J. Hum. Genet. 62: 346–354,
1998.

[138233]24644.

[138234]Xu, L.; Xia, J.; Jiang, H.; Zhou, J.; Li, H.; Wang, D.; Pan, Q.;

[138235]Long, Z.; Fan, C.; Deng, H.-X.: Mutation analysis of heredit-
itary multiple

[138236]exostoses in the Chinese. Hum. Genet. 105: 45–50, 1999.

[138237]24645.

[138238]Yoshiura, K.; Inazawa, J.; Koyama, K.; Nakamura, Y.; Ni-
ikawa,

[138239]N.: Mapping of the 8q23 translocation breakpoint of

t(8;13) observed

[138240]in a patient with multiple exostoses. Genes Chromosomes
Cancer 9:

[138241]57–61, 1994.

[138242]24646.

[138243]Bellaïche, Y.; The, I.; Perrimon, N.: Tout-velu is a
Drosophila

[138244]homologue of the putative tumour suppressor EXT-1 and
is needed for

[138245]Hh diffusion. Nature 394: 85–88, 1998.

[138246]24647.

[138247]Clines, G. A.; Ashley, J. A.; Shah, S.; Lovett, M.: The struc-
ture

[138248]of the human multiple exostoses 2 gene and characteriza-
tion of homologs

[138249]in mouse and Caenorhabditis elegans. Genome Res. 7:
359–367, 1997.

[138250]24648.

[138251]Francannet, C.; Cohen-Tanugi, A.; Le Merrer, M.; Munnich,
A.; Bonaventure,

[138252]J.; Legeai-Mallet, L.: Genotype–phenotype correlation in

hereditary

[138253]multiple exostoses. J. Med. Genet. 38: 430–434, 2001.

[138254]24649.

[138255]Ligon, A. H.; Potocki, L.; Shaffer, L. G.; Stickens, D.; Evans,

[138256]G. A.: Gene for multiple exostoses (EXT2) maps to

11(p11.2p12) and

[138257]is deleted in patients with a contiguous gene syndrome.

(Letter) Am.

[138258]J. Med. Genet. 75: 538–540, 1998.

[138259]24650.

[138260]Lind, T.; Tufaro, F.; McCormick, C.; Lindahl, U.; Lidholt, K.

[138261]: The putative tumor suppressors EXT1 and EXT2 are gly-

cosyltransferases

[138262]required for the biosynthesis of heparan sulfate. J. Biol.

Chem. 273:

[138263]26265–26268, 1998.

[138264]24651.

[138265]McGaughan, J. M.; Ward, H. B.; Evans, D. G. R.: WAGR

syndrome

[138266]and multiple exostoses in a patient with

del(11)(p11.2p14.2). J.

[138267]Med. Genet. 32: 823–824, 1995.

[138268]24652.

[138269]Potocki, L.; Greenberg, F.; Shaffer, L. G.: Interstitial deletion

[138270]of 11(p12p11.2): a rare chromosomal syndrome with mental retardation,

[138271]parietal foramina, and multiple exostoses. (Abstract) Am. J. Hum.

[138272]Genet. 57: A123, 1995.

[138273]24653.

[138274]Potocki, L.; Shaffer, L. G.: Interstitial deletion of 11(p11.2p12):

[138275]a newly described contiguous gene deletion syndrome involving the

[138276]gene for hereditary multiple exostoses (EXT2). Am. J. Med. Genet. 62:

[138277]319–325, 1996.

[138278]24654.

[138279]Quirini, G. E.; Meyer, J. R.; Herman, M.; Russell, E. J.: Osteochondroma

[138280]of the thoracic spine: an unusual cause of spinal cord compression. Am.

[138281]J. Neuroradiol. 17: 961-964, 1996.

[138282]24655.

[138283]Simmons, A. D.; Musy, M. M.; Lopes, C. S.; Hwang, L.-Y.; Yang,

[138284]Y.-P.; Lovett, M.: A direct interaction between EXT proteins and

[138285]glycosyltransferases is defective in hereditary multiple exostoses. Hum.

[138286]Molec. Genet. 8: 2155-2164, 1999.

[138287]24656.

[138288]Stickens, D.; Clines, G.; Burbee, D.; Ramos, P.; Thomas, S.; Hogue,

[138289]D.; Hecht, J. T.; Lovett, M.; Evans, G. A.: The EXT2 multiple exostoses

[138290]gene defines a family of putative tumour suppressor genes. Nature

[138291]Genet. 14: 25-32, 1996.

[138292]24657.

[138293]Stickens, D.; Evans, G. A.: Isolation and characterization of

- [138294] the murine homolog of the human EXT2 multiple exostoses gene. *Biochem. Molec. Med.* 61: 16–21, 1997.
- [138295] 24658.
- [138296] Wu, Y.-Q.; Heutink, P.; de Vries, B. B. A.; Sandkuijl, L. A.; van den Ouweland, A. M. W.; Niermeijer, M. F.; Galjaard, H.; Reyniers, E.; Willems, P. J.; Halley, D. J. J.: Assignment of a second locus for multiple exostoses to the pericentromeric region of chromosome 11. *Hum. Molec. Genet.* 3: 167–171, 1994.
- [138297] 24659.
- [138298] Joutel, A.; Favrole, P.; Labauge, P.; Chabriat, H.; Lescoat, C.; Andreux, F.; Domenga, V.; Cecillon, M.; Vahedi, K.; Ducros, A.; Cave-Riant, F.; Bousser, M. G.; Tournier-Lasserre, E.: Skin biopsy immunostaining with a Notch3 monoclonal antibody for CADASIL diagnosis. *Lancet* 358:

[138307]2049–2051, 2001.

[138308]24660.

[138309]Aoki, M.; Abe, K.; Kameya, T.; Watanabe, M.; Itoyama, Y.:
Maternal

[138310]anticipation of DRPLA. Hum. Molec. Genet. 3: 1197–1198,
1994.

[138311]24661.

[138312]Barinaga, M.: An intriguing new lead on Huntington's dis-
ease. Science 271:

[138313]1233–1234, 1996.

[138314]24662.

[138315]Burke, J. R.; Enghild, J. J.; Martin, M. E.; Jou, Y.–S.; Myers,

[138316]R. M.; Roses, A. D.; Vance, J. M.; Strittmatter, W. J.: Hunt-
ingtin

[138317]and DRPLA proteins selectively interact with the enzyme
GAPDH. Nature

[138318]Med. 2: 347–350, 1996.

[138319]24663.

[138320]Burke, J. R.; Wingfield, M. S.; Lewis, K. E.; Roses, A. D.;
Lee,

[138321]J. E.; Hulette, C.; Pericak-Vance, M. A.; Vance, J. M.: The
Haw River

[138322]syndrome: dentatorubropallidoluysian atrophy (DRPLA) in
an African-American

[138323]family. Nature Genet. 7: 521-524, 1994.

[138324]24664.

[138325]Cancel, G.; Durr, A.; Stevanin, G.; Chneiweiss, H.; Duyck-
aerts,

[138326]C.; Serdaru, M.; de Toffol, B.; Agid, Y.; Brice, A.: Is DRPLA
also

[138327]linked to 14q? (Letter) Nature Genet. 6: 8, 1994.

[138328]24665.

[138329]Connarty, M.; Dennis, N. R.; Patch, C.; Macpherson, J. N.;
Harvey,

[138330]J. F.: Molecular re-investigation of patients with Hunting-
ton's disease

[138331]in Wessex reveals a family with dentatorubral and palli-
doluysian atrophy. Hum.

[138332]Genet. 97: 76-78, 1996.

[138333]24666.

[138334]Dubourg, O.; Durr, A.; Chneiweiss, H.; Stevanin, G.; Can-

cel, G.;

[138335]Penet, C.; Agid, Y.; Brice, A.: La forme ataxo-choreique de
DRPLA

[138336]existe-t-elle en Europe? Recherche de la mutation dans
120 familles. Rev.

[138337]Neurol. (Paris) 151: 657-660, 1995.

[138338]24667.

[138339]Hayashi, Y.; Kakita, A.; Yamada, M.; Koide, R.; Igarashi, S.;
Takano,

[138340]H.; Ikeuchi, T.; Wakabayashi, K.; Egawa, S.; Tsuji, S.; Taka-
hashi,

[138341]H.: Hereditary dentatorubral-pallidoluysian atrophy: de-
tection of

[138342]widespread ubiquitinated neuronal and glial intranuclear
inclusions

[138343]in the brain. Acta Neuropath. 96: 547-552, 1998.

[138344]24668.

[138345]Hirayama, K.; Iizuka, R.; Maehara, K.; Watanabe, T.: Clini-
copathological

[138346]study of dentatorubro-pallidoluysian atrophy. Part 1. Its
clinical

[138347]form and analysis of symptomatology. Shinkei Kenkyu no

Shinpo 25:

[138348]725–736, 1981.

[138349]24669.

[138350]Hirayama, K.; Takayanagi, T.; Nakamura, R.; Yanagisawa, N.; Hattori,

[138351]T.; Kita, K.; Yanagimoto, S.; Fujita, M.; Nagaoka, M.; Sato-mura, Y.;

[138352]Sobue, I.; Iizuka, R.; Toyokura, Y.; Satoyoshi, E.:
Spinocerebellar

[138353]degenerations in Japan: a nationwide epidemiological and
clinical

[138354]study. *Acta Neurol. Scand.* 89 (suppl. 153): 1–22, 1994.

[138355]24670.

[138356]Iizuka, R.; Hirayama, K.; Maehara, K.: Dentatorubro-palli-doluysian

[138357]atrophy: a clinico-pathological study. *J. Neurol. Neuro-surg. Psychiat.* 47:

[138358]1288–1298, 1984.

[138359]24671.

[138360]Ikeuchi, T.; Igarashi, S.; Takiyama, Y.; Onodera, O.; Oyake, M.;

[138361]Takano, H.; Koide, R.; Tanaka, H.; Tsuji, S.: Non-mendelian transmission

[138362]in dentatorubral-pallidoluysian atrophy and Machado-Joseph disease:

[138363]the mutant allele is preferentially transmitted in male meiosis. Am.

[138364]J. Hum. Genet. 58: 730-733, 1996.

[138365]24672.

[138366]Iwabuchi, K.; Amano, N.; Yokoi, S.; Nakano, T.; Yagishita, S.

[138367]: Two familial cases of dentato-rubro-pallidoluysian atrophy with

[138368]pseudo-Huntington's chorea (in Japanese). Clin. Neurol.

25: 1052-1060,

[138369]1985.

[138370]24673.

[138371]Eubanks, J. H.; Djabali, M.; Selleri, L.; Grandy, D. K.; Civelli,

[138372]O.; McElligott, D. L.; Evans, G. A.: Structure and linkage of the

[138373]D2 dopamine receptor and neural cell adhesion molecule genes on human

[138374]chromosome 11q23. Genomics 14: 1010-1018, 1992.

[138375]24674.

[138376]Gejman, P. V.; Ram, A.; Gelernter, J.; Friedman, E.; Cao, Q.;

[138377]Pickar, D.; Blum, K.; Noble, E. P.; Kranzler, H. R.; O'Malley, S.;

[138378]Hamer, D. H.; Whitsitt, F.; Rao, P.; DeLisi, L. E.; Virkkunen, M.;

[138379]Linnoila, M.; Goldman, D.; Gershon, E. S.: No structural mutation

[138380]in the dopamine D2 receptor gene in alcoholism or schizophrenia: analysis

[138381]using denaturing gradient gel electrophoresis. J.A.M.A. 271: 204-208,

[138382]1994.

[138383]24675.

[138384]Gelernter, J.; Grandy, D. K.; Bunzow, J.; Pakstis, A. J.; Civelli,

[138385]O.; Retief, A. E.; Litt, M.; Kidd, K. K.: D(2) dopamine receptor

[138386]locus (probe hD2G1) maps close to D11S29 (probe L7) and is also linked

[138387]to PBGD (probe PBGD0.0) and D11S84 (probe p-2-7-ID6)
on 11q. (Abstract) Cytogenet.

[138388]Cell Genet. 51: 1002 only, 1989.

[138389]24676.

[138390]Gelernter, J.; Pakstis, A. J.; Grandy, D.; Litt, M.; Retief, A.

[138391]E.; Kennedy, J. L.; Hing-Loh, A.; Schoolfield, G.; Civelli, O.;
Kidd,

[138392]K. K.: Linkage map of eight human chromosome 11q
markers, including

[138393]DRD2, spanning 60 cM. Cytogenet. Cell Genet. 60: 26-28,
1992.

[138394]24677.

[138395]Gelernter, J.; Pakstis, A. J.; Pauls, D. L.; Kurlan, R.;
Gancher,

[138396]S. T.; Civelli, O.; Grandy, D.; Kidd, K. K.: Gilles de la
Tourette

[138397]syndrome is not linked to D2-dopamine receptor. Arch.
Gen. Psychiat. 47:

[138398]1073-1077, 1990.

[138399]24678.

[138400]Giros, B.; Sokoloff, P.; Martres, M.-P.; Riou, J.-F.; Emorine,

[138401]L. J.; Schwartz, J.-C.: Alternative splicing directs the expression

[138402]of two D(2) dopamine receptor isoforms. *Nature* 342: 923–926, 1989.

[138403]24679.

[138404]Grandy, D. K.; Litt, M.; Allen, L.; Bunzow, J. R.; Marchionni,

[138405]M.; Makam, H.; Reed, L.; Magenis, R. E.; Civelli, O.: The human dopamine

[138406]D(2) receptor gene is located on chromosome 11 at q22–q23 and identifies

[138407]a TaqI RFLP. *Am. J. Hum. Genet.* 45: 778–785, 1989.

[138408]24680.

[138409]Grandy, D. K.; Marchionni, M. A.; Makam, H.; Stofko, R. E.; Alfano,

[138410]M.; Frothingham, L.; Fischer, J. B.; Burke–Howie, K. J.; Bunzow, J.

[138411]R.; Server, A. C.; Civelli, O.: Cloning of the cDNA and gene for

[138412]a human D2 dopamine receptor. *Proc. Nat. Acad. Sci.* 86: 9762–9766,

[138413]1989.

[138414]24681.

[138415]Klein, C.; Brin, M. F.; Kramer, P.; Sena-Esteves, M.; de Leon,

[138416]D.; Doheny, D.; Bressman, S.; Fahn, S.; Breakefield, X. O.; Ozelius,

[138417]L. J.: Association of a missense change in the D2 dopamine receptor

[138418]with myoclonus dystonia. Proc. Nat. Acad. Sci. 96: 5173–5176, 1999.

[138419]24682.

[138420]Maldonado, R.; Salardi, A.; Valverde, O.; Samad, T. A.; Roques,

[138421]B. P.; Borrelli, E.: Absence of opiate rewarding effects in mice

[138422]lacking dopamine D2 receptors. Nature 388: 586–589, 1997.

[138423]24683.

[138424]Milligan, G.: Receptors as kissing cousins. Science 288: 65–67,

[138425]2000.

[138426]24684.

[138427]Monsma, F. J., Jr.; McVittie, L. D.; Gerfen, C. R.; Mahan, L.
[138428]C.; Sibley, D. R.: Multiple D(2) dopamine receptors produced by alternative
[138429]RNA splicing. *Nature* 342: 926–929, 1989.
[138430]24685.
[138431]Oliveri, R. L.; Annesi, G.; Zappia, M.; Civitelli, D.; Montesanti,
[138432]R.; Branca, D.; Nicoletti, G.; Spadafora, P.; Pasqua, A. A.;
Cittadella,
[138433]R.; Andreoli, V.; Gambardella, A.; Aguglia, U.; Quattrone,
A.: Dopamine
[138434]D2 receptor gene polymorphism and the risk of levodopa-induced dyskinesias
[138435]in PD. *Neurology* 53: 1425–1430, 1999.
[138436]24686.
[138437]Rocheville, M.; Lange, D. C.; Kumar, U.; Patel, S. C.; Patel,
[138438]R. C.; Patel, Y. C.: Receptors for dopamine and somatostatin: formation
[138439]of hetero-oligomers with enhanced functional activity.
Science 288:
[138440]154–157, 2000.

[138441]24687.

[138442]Smith, M.; Wasmuth, J.; McPherson, J. D.; Wagner, C.;
Grandy,

[138443]D.; Civelli, O.; Potkin, S.; Litt, M.: Cosegregation of an
11q22.3-9p22

[138444]translocation with affective disorder: proximity of the
dopamine D2

[138445]receptor gene relative to the translocation breakpoint.
(Abstract) Am.

[138446]J. Hum. Genet. 45 (suppl.): A220 only, 1989.

[138447]24688.

[138448]St Clair, D.; Blackwood, D.; Muir, W.; Carothers, A.; Walker,

[138449]M.; Spowart, G.; Gosden, C.; Evans, H. J.: Association
within a family

[138450]of a balanced autosomal translocation with major mental
illness. Lancet 336:

[138451]13-16, 1990.

[138452]24689.

[138453]Suarez, B. K.; Parsian, A.; Hampe, C. L.; Todd, R. D.; Reich,

[138454]T.; Cloninger, C. R.: Linkage disequilibria at the D2
dopamine receptor

[138455]locus (DRD2) in alcoholics and controls. Genomics 19:
12–20, 1994.

[138456]24690.

[138457]Usiello, A.; Baik, J.–H.; Rouge–Pont, F.; Picetti, R.; Dierich,

[138458]A.; LeMeur, M.; Piazza, P. V.; Borrelli, E.: Distinct functions
of

[138459]the two isoforms of dopamine D2 receptors. Nature 408:
199–203,

[138460]2000.

[138461]24691.

[138462]Wang, J.; Liu, Z.–L.; Chen, B.: Association study of
dopamine

[138463]D2, D3 receptor gene polymorphisms with motor fluctua-
tions in PD. Neurology 56:

[138464]1757–1759, 2001.

[138465]24692.

[138466]Accili, D.; Fishburn, C. S.; Drago, J.; Steiner, H.; Lachowicz,

[138467]J. E.; Park, B.–H.; Gauda, E. B.; Lee, E. J.; Cool, M. H.; Sib-
ley,

[138468]D. R.; Gerfen, C. R.; Westphal, H.; Fuchs, S.: A targeted
mutation

[138469]of the D3 dopamine receptor gene is associated with hyperactivity

[138470]in mice. Proc. Nat. Acad. Sci. 93: 1945–1949, 1996.

[138471]24693.

[138472]Asico, L. D.; Ladines, C.; Fuchs, S.; Accili, D.; Carey, R. M.;

[138473]Semeraro, C.; Pocchiari, F.; Felder, R. A.; Eisner, G. M.; Jose, P.

[138474]A.: Disruption of the dopamine D3 receptor gene produces renin-dependent

[138475]hypertension. J. Clin. Invest. 102: 493–498, 1998.

[138476]24694.

[138477]Crocq, M.–A.; Mant, R.; Asherson, P.; Williams, J.; Hode, Y.; Mayerova,

[138478]A.; Collier, D.; Lannfelt, L.; Sokoloff, P.; Schwartz, J.–C.; Gill,

[138479]M.; Macher, J.–P.; McGuffin, P.; Owen, M. J.: Association between

[138480]schizophrenia and homozygosity at the dopamine D3 receptor gene. J.

[138481]Med. Genet. 29: 858–860, 1992.

[138482]24695.

[138483]Guillin, O.; Diaz, J.; Carroll, P.; Griffon, N.; Schwartz, J.-C.;

[138484]Sokoloff, P.: BDNF controls dopamine D3 receptor expression and triggers

[138485]behavioural sensitization. *Nature* 411: 86–89, 2001.

[138486]24696.

[138487]Ilani, T.; Ben-Shachar, D.; Strous, R. D.; Mazor, M.; Sheinkman,

[138488]A.; Kotler, M.; Fuchs, S.: A peripheral marker for schizophrenia:

[138489]increased levels of D3 dopamine receptor mRNA in blood lymphocytes. *Proc.*

[138490]*Nat. Acad. Sci.* 98: 625–628, 2001.

[138491]24697.

[138492]Le Coniat, M.; Sokoloff, P.; Hillion, J.; Martres, M.-P.; Giros,

[138493]B.; Pilon, C.; Schwartz, J.-C.; Berger, R.: Chromosomal localization

[138494]of the human D-3 dopamine receptor gene. *Hum. Genet.* 87: 618–620,

[138495]1991.

[138496]24698.

- [138497]Nanko, S.; Sasaki, T.; Fukuda, R.; Hattori, M.; Dai, X. Y.; Kazamatsuri, H.; Kuwata, S.; Juji, T.; Gill, M.: A study of the association between schizophrenia and the dopamine D3 receptor gene. Hum. Genet. 92: 336–338, 1993.
- [138501]24699.
- [138502]Nothen, M. M.; Cichon, S.; Propping, P.; Fimmers, R.; Schwab, S. G.; Wildenauer, D. B.: Excess of homozygosity at the dopamine D3 receptor gene in schizophrenia not confirmed. (Letter) J. Med. Genet. 30: 708–712, 1993.
- [138506]24700.
- [138507]Bevilacqua, M.; Butcher, E.; Furie, B.; Furie, B.; Gallatin, M.; Gimbrone, M.; Harlan, J.; Kishimoto, K.; Lasky, L.; McEver, R.; Paulson, J.; Rosen, S.; Seed, B.; Siegelman, M.; Springer, T.; Stoolman, L.;

- [138510] Tedder, T.; Varki, A.; Wagner, D.; Weissman, I.; Zimmerman, G.: Selectins:
[138511] a family of adhesion receptors. (Letter) Cell 67: 233 only,
1991.
[138512] 24701.
- [138513] Bevilacqua, M. P.; Stengelin, S.; Gimbrone, M. A., Jr.; Seed,
B.
[138514]: Endothelial leukocyte adhesion molecule 1: an inducible
receptor
[138515] for neutrophils related to complement regulatory proteins
and lectins. Science 243:
[138516] 1160–1165, 1989.
[138517] 24702.
- [138518] Collins, T.; Williams, A.; Johnston, G. I.; Kim, J.; Eddy, R.;
[138519] Shows, T.; Gimbrone, M. A., Jr.; Bevilacqua, M. P.: Struc-
ture and
[138520] chromosomal location of the gene for endothelial-leuko-
cyte adhesion
[138521] molecule 1. J. Biol. Chem. 266: 2466–2473, 1991.
[138522] 24703.
- [138523] DeLisser, H. M.; Christofidou-Solomidou, M.; Sun, J.;

Nakada, M.

[138524]T.; Sullivan, K. E.: Loss of endothelial surface expression
of E-selectin

[138525]in a patient with recurrent infections. Blood 94: 884–894,
1999.

[138526]24704.

[138527]Wang, N.; Chintala, S. K.; Fini, M. E.; Schuman, J. S.: Acti-
vation

[138528]of a tissue-specific stress response in the aqueous out-
flow pathway

[138529]of the eye defines the glaucoma disease phenotype. Na-
ture Med. 7:

[138530]304–309, 2001.

[138531]24705.

[138532]Watson, M. L.; Kingsmore, S. F.; Johnston, G. I.; Siegelman,
M.

[138533]H.; Le Beau, M. M.; Lemons, R. S.; Bora, N. S.; Howard, T.
A.; Weissman,

[138534]I. L.; McEver, R. P.; Seldin, M. F.: Genomic organization of
the

[138535]selectin family of leukocyte adhesion molecules on human
and mouse

[138536]chromosome 1. J. Exp. Med. 172: 263–272, 1990.

[138537]24706.

[138538]Wenzel, K.; Felix, S.; Kleber, F. X.; Brachold, R.; Menke, T.;

[138539]Schattke, S.; Schulte, K. L.; Glaser, C.; Rohde, K.; Baumann,
G.;

[138540]Speer, A.: E-selectin polymorphism and atherosclerosis:
an association

[138541]study. Hum. Molec. Genet. 3: 1935–1937, 1994.

[138542]24707.

[138543]Zheng, F.; Chevalier, J. A.; Zhang, L. Q.; Virgil, D.; Ye, S.
Q.;

[138544]Kwiterovich, P. O.: An HphI polymorphism in the E-
selectin gene is

[138545]associated with premature coronary artery disease. Clin.
Genet. 59:

[138546]58–64, 2001.

[138547]24708.

[138548]Burgess, W. H.; Mehlman, T.; Marshak, D. R.; Fraser, B. A.;
Maciag,

[138549]T.: Structural evidence that endothelial cell growth factor
beta

[138550] is the precursor of both endothelial cell growth factor al-
pha and

[138551] acidic fibroblast growth factor. Proc. Nat. Acad. Sci. 83:
7216-7220,

[138552] 1986.

[138553] 24709.

[138554] Eckenstein, F. P.: Fibroblast growth factors in the nervous
system. J.

[138555] Neurobiol. 25: 1467-1480, 1994.

[138556] 24710.

[138557] Gautschi-Sova, P.; Muller, T.; Bohlen, P.: Amino acid se-
quence

[138558] of human acidic fibroblast growth factor. Biochem. Bio-
phys. Res.

[138559] Commun. 140: 874-880, 1986.

[138560] 24711.

[138561] Jaye, M.; Howk, R.; Burgess, W.; Ricca, G. A.; Chiu, I.-M.;
Ravera,

[138562] M. W.; O'Brien, S. J.; Modi, W. S.; Maciag, T.; Drohan, W. N.:
Human

[138563] endothelial cell growth factor: cloning, nucleotide se-

quence, and

[138564]chromosome localization. Science 233: 541–545, 1986.

[138565]24712.

[138566]Jung, J.; Zheng, M.; Goldfarb, M.; Zaret, K. S.: Initiation of

[138567]mammalian liver development from endoderm by fibroblast growth factors. Science 284:

[138568]1998–2003, 1999.

[138569]24713.

[138570]Mergia, A.; Eddy, R.; Abraham, J. A.; Fiddes, J. C.; Shows, T.

[138571]B.: The genes for basic and acidic fibroblast growth factors are

[138572]on different human chromosomes. Biochem. Biophys. Res. Commun. 138:

[138573]644–651, 1986.

[138574]24714.

[138575]Pellegrini, L.; Burke, D. F.; von Delft, F.; Mulloy, B.; Blundell,

[138576]T. L.: Crystal structure of fibroblast growth factor receptor ectodomain

[138577]bound to ligand and heparin. Nature 407: 1029–1034,

2000.

[138578]24715.

[138579]Plotnikov, A. N.; Hubbard, S. R.; Schlessinger, J.; Moham-
madi,

[138580]M.: Crystal structures of two FGF-FGFR complexes reveal
the determinants

[138581]of ligand-receptor specificity. Cell 101: 413-424, 2000.

[138582]24716.

[138583]Wang, W.-P.; Lehtoma, K.; Varban, M. L.; Krishnan, I.;
Chiu, I.-M.

[138584]: Cloning of the gene coding for human class 1 heparin-
binding growth

[138585]factor and its expression in fetal tissues. Molec. Cell. Biol.
9:

[138586]2387-2395, 1989.

[138587]24717.

[138588]Kayes-Wandover, K.; White, P. C.: Steroidogenic enzyme
gene expression

[138589]in the human heart. J. Clin. Endocr. Metab. 85:
2519-2525, 2000.

[138590]24718.

[138591]Carrero-Valenzuela, R. D.; Quan, F.; Lightowlers, R.; Ken-
naway,

[138592]N. G.; Litt, M.; Forte, M.: Human cytochrome c oxidase
subunit VIb:

[138593]characterization and mapping of a multigene family. Gene
102: 229-236,

[138594]1991.

[138595]24719.

[138596]Taanman, J.-W.; Schrage, C.; Ponne, N.; Bolhuis, P.; de
Vries,

[138597]H.; Agsteribbe, E.: Nucleotide sequence of cDNA encoding
subunit

[138598]VIb of human cytochrome c oxidase. Nucleic Acids Res.
17: 1766

[138599]only, 1989.

[138600]24720.

[138601]Taanman, J.-W.; van der Veen, A. Y.; Schrage, C.; de Vries, H.;

[138602]Buys, C. H. C. M.: Assignment of the gene coding for human cytochrome

[138603]c oxidase subunit VIb to chromosome 19, band q13.1, by fluorescence

[138604]in situ hybridisation. Hum. Genet. 87: 325–327, 1991.

[138605]24721.

[138606]Freddi, S.; Savarirayan, R.; Bateman, J. F.: Molecular diagnosis

[138607]of Stickler syndrome: A COL2A1 stop codon mutation screening strategy

[138608]that is not compromised by mutant mRNA instability. Am. J. Med. Genet. 90:

[138609]398–406, 2000.

[138610]24722.

[138611]Griffith, A. J.; Sprunger, L. K.; Sirko-Osadsa, D. A.; Tiller,

[138612]G. E.; Meisler, M. H.; Warman, M. L.: Marshall syndrome associated

[138613]with a splicing defect at the COL11A1 gene. Am. J. Hum.

Genet. 62:

[138614]816-823, 1998.

[138615]24723.

[138616]Cox, D. R.; Epstein, L. B.; Epstein, C. J.: Genes coding for sensitivity

[138617]to interferon (IfRec) and soluble superoxide dismutase (SOD-1) are

[138618]linked in mouse and man and map to mouse chromosome 16. Proc. Nat.

[138619]Acad. Sci. 77: 2168-2172, 1980.

[138620]24724.

[138621]De Clercq, E.; Edy, V. G.; Cassiman, J.-J.: Chromosome 21 does

[138622]not code for an interferon receptor. Nature 264: 249-251, 1976.

[138623]24725.

[138624]Epstein, L. B.; Epstein, C. J.: Localization of the gene AVG for

[138625]the antiviral expression of immune and classical interferon to the

[138626]distal portion of the long arm of chromosome 21. J. Infect.

[138627](suppl.): A56-A62, 1976.

[138628]24726.

[138629]Faltynek, C. R.; Branca, A. A.; McCandless, S.; Baglioni, C.:

[138630]Characterization of an interferon receptor on human lymphoblastoid

[138631]cells. Proc. Nat. Acad. Sci. 80: 3269-3273, 1983.

[138632]24727.

[138633]Fournier, A.; Zhang, Z. Q.; Tan, Y. H.: Human beta:alpha but not

[138634]gamma interferon binding site is a product of the chromosome 21 interferon

[138635]action gene. Somat. Cell Molec. Genet. 11: 291-295, 1985.

[138636]24728.

[138637]Lutfalla, G.; Gardiner, K.; Proudhon, D.; Vielh, E.; Uze, G.:

[138638]The structure of the human interferon alpha/beta receptor gene. J.

[138639]Biol. Chem. 267: 2802-2809, 1992.

[138640]24729.

[138641]Lutfalla, G.; Roeckel, N.; Mogensen, K. E.; Mattei, M. G.;
Uze,

[138642]G.: Assignment of the human interferon- α receptor
gene to chromosome

[138643]21q22.1 by in situ hybridization. J. Interferon Res. 10:
515-517,

[138644]1990.

[138645]24730.

[138646]Maroun, L. E.: Interferon action and chromosome 21 tri-
somy. (Letter) J.

[138647]Theor. Biol. 86: 603-606, 1980.

[138648]24731.

[138649]Novick, D.; Cohen, B.; Rubinstein, M.: The human inter-
feron α/β

[138650]receptor: characterization and molecular cloning. Cell 77:
391-400,

[138651]1994.

[138652]24732.

[138653]Raziuddin, A.; Sarkar, F. H.; Dutkowski, R.; Shulman, L.;
Ruddle,

[138654]F. H.; Gupta, S. L.: Receptors for human α and β in-

terferon

[138655]but not for gamma interferon are specified by human chromosome 21. Proc.

[138656]Nat. Acad. Sci. 81: 5504-5508, 1984.

[138657]24733.

[138658]Revel, M.; Bash, D.; Ruddle, F. H.: Antibodies to a cell-surface

[138659]component coded by human chromosome 21 inhibit action of interferon. Nature 260:

[138660]139-141, 1976.

[138661]24734.

[138662]Calabrese, G.; Crescenzi, C.; Morizio, E.; Palka, G.; Guerra, E.;

[138663]Alberti, S.: Assignment of TACSTD1 (alias TROP1, M4S1) to human chromosome

[138664]2p21 and refinement of mapping of TACSTD2 (alias TROP2, M1S1) to human

[138665]chromosome 1p32 by in situ hybridization. Cytogenet. Cell Genet. 92:

[138666]164-165, 2001.

[138667]24735.

[138668]Fornaro, M.; Dell'Arciprete, R.; Stella, M.; Bucci, C.; Nutini,
[138669]M.; Capri, M. G.; Alberti, S.: Cloning of the gene encoding
Trop-2,
[138670]a cell-surface glycoprotein expressed by human carcino-
mas. Int. J.
[138671]Cancer 62: 610-618, 1995.
[138672]24736.
[138673]Linnenbach, A. J.; Seng, B. A.; Wu, S.; Robbins, S.; Scollon,
M.;
[138674]Pyrç, J. J.; Druck, T.; Huebner, K.: Retroposition in a family
of
[138675]carcinoma-associated antigen genes. Molec. Cell Biol. 13:
1507-1515,
[138676]1993.
[138677]24737.
[138678]Linnenbach, A. J.; Wojcierowski, J.; Wu, S.; Pyrç, J. J.; Ross,
[138679]A. H.; Dietzschold, B.; Speicher, D.; Koprowski, H.: Se-
quence investigation
[138680]of the major gastrointestinal tumor-associated antigen
gene family,
[138681]GA733. Proc. Nat. Acad. Sci. 86: 27-31, 1989.

[138682]24738.

[138683]Ren, Z.; Lin, P.-Y.; Klintworth, G. K.; Iwata, F.; Munier, F.
L.;

[138684]Schorderet, D. F.; El Matri, L.; Theendakara, V.; Basti, S.;
Reddy,

[138685]M.; Hejtmancik, J. F.: Allelic and locus heterogeneity in au-
tosomal

[138686]recessive gelatinous drop-like corneal dystrophy. Hum.
Genet. 110:

[138687]568-577, 2002.

[138688]24739.

[138689]TsujiKawa, M.; Kurahashi, H.; Tanaka, T.; Nishida, K.; Shi-
momura,

[138690]Y.; Tano, Y.; Nakamura, Y.: Identification of the gene re-
sponsible

[138691]for gelatinous drop-like corneal dystrophy. Nature Genet.
21: 420-423,

[138692]1999.

[138693]24740.

[138694]Briegel, K.; Lim, K.-C.; Plank, C.; Beug, H.; Engel, J. D.;
Zenke,

- [138695]M.: Ectopic expression of a conditional GATA-2/estrogen receptor
- [138696]chimera arrests erythroid differentiation in a hormone-dependent manner. *Genes Dev.* 7: 1097-1109, 1993.
- [138698]24741.
- [138699]Ciciotte, S. L.; Tsai, F.-Y.; Peters, L. L.: Gata2 maps to mouse chromosome 6. *Mammalian Genome* 8: 69-70, 1997.
- [138701]24742.
- [138702]Dasen, J. S.; O'Connell, S. M.; Flynn, S. E.; Treier, M.; Gleiberman, A. S.; Szeto, D. P.; Hooshmand, F.; Aggarwal, A. K.; Rosenfeld, M.
- [138704]G.: Reciprocal interactions of Pit1 and GATA2 mediate signaling gradient-induced determination of pituitary cell types. *Cell* 97: 587-598, 1999.
- [138706]24743.
- [138707]Dorfman, D. M.; Wilson, D. B.; Bruns, G. A.; Orkin, S. H.: Human

[138708]transcription factor GATA-2: evidence for regulation of
preproendothelin-1

[138709]gene expression in endothelial cells. J. Biol. Chem. 267:
1279-1285,

[138710]1992.

[138711]24744.

[138712]Lee, M.-E.; Temizer, D. H.; Clifford, J. A.; Quertermous, T.:

[138713]Cloning of the GATA-binding protein that regulates en-
dothelin-1 gene

[138714]expression in endothelial cells. J. Biol. Chem. 266:
16188-16192,

[138715]1991.

[138716]24745.

[138717]Tsai, F.-Y.; Keller, G.; Kuo, F. C.; Weiss, M.; Chen, J.;
Rosenblatt,

[138718]M.; Alt, F. W.; Orkin, S. H.: An early haematopoietic defect
in mice

[138719]lacking the transcription factor GATA2. Nature 371:
221-226, 1994.

[138720]24746.

[138721]Zhou, Y.; Lim, K.-C.; Onodera, K.; Takahashi, S.; Ohta, J.;

Minegishi,

[138722]N.; Tsai, F.-Y.; Orkin, S. H.; Yamamoto, M.; Engel, J. D.:

Rescue

[138723]of the embryonic lethal hematopoietic defect reveals a
critical role

[138724]for GATA-2 in urogenital development. EMBO J. 17:
6689-6700, 1998.

[138725]24747.

[138726]Becker-Follmann, J.; Gaa, A.; Bausch, E.; Natt, E.; Scherer,
G.;

[138727]von Deimling, O.: High-resolution mapping of a linkage
group on mouse

[138728]chromosome 8 conserved on human chromosome 16Q.
Mammalian Genome 8:

[138729]172-177, 1997.

[138730]24748.

[138731]Brooks, P. C.; Silletti, S.; von Schalscha, T. L.; Friedlander,

[138732]M.; Cheresh, D. A.: Disruption of angiogenesis by PEX, a
noncatalytic

[138733]metalloproteinase fragment with integrin binding activity.
Cell 92:

[138734]391-400, 1998.

[138735]24749.

[138736]Collier, I. E.; Bruns, G. A. P.; Goldberg, G. I.; Gerhard, D. S.

[138737]: On the structure and chromosome location of the 72-
and 92-kDa human

[138738]type IV collagenase genes. *Genomics* 9: 429-434, 1991.

[138739]24750.

[138740]Corry, D. B.; Rishi, K.; Kanellis, J.; Kiss, A.; Song, L.; Xu,

[138741]J.; Feng, L.; Werb, Z.; Kheradmand, F.: Decreased allergic
lung inflammatory

[138742]cell egression and increased susceptibility to asphyxiation
in MMP2-deficiency. *Nature*

[138743]*Immun.* 3: 347-353, 2002.

[138744]24751.

[138745]Devarajan, P.; Johnston, J. J.; Ginsberg, S. S.; Van Wart, H.
E.;

[138746]Berliner, N.: Structure and expression of neutrophil gelati-
nase cDNA:

[138747]identity with type IV collagenase from HT1080 cells. *J.*
Biol. Chem. 267:

[138748]25228-25232, 1992.

[138749]24752.

[138750]Fan, Y.-S.; Eddy, R. L.; Huhtala, P.; Byers, M. G.; Haley, L. L.;

[138751]Henry, W. M.; Tryggvason, K.; Shows, T. B.: Collagenase type IV (CLG4)

[138752]is mapped to human chromosome 16q21. (Abstract) Cytogenet. Cell Genet. 51:

[138753]996, 1989.

[138754]24753.

[138755]Huhtala, P.; Chow, L. T.; Tryggvason, K.: Structure of the human

[138756]type IV collagenase gene. J. Biol. Chem. 265: 11077-11082, 1990.

[138757]24754.

[138758]Huhtala, P.; Eddy, R. L.; Fan, Y. S.; Byers, M. G.; Shows, T.

[138759]B.; Tryggvason, K.: Completion of the primary structure of the human

[138760]type IV collagenase preproenzyme and assignment of the gene (CLG4)

[138761]to the q21 region of chromosome 16. Genomics 6: 554-559, 1990.

[138762]24755.

[138763]Irwin, J. C.; Kirk, D.; Gwatkin, R. B. L.; Navre, M.; Cannon,

[138764]P.; Giudice, L. C.: Human endometrial matrix metallopro-
teinase-2,

[138765]a putative menstrual proteinase: hormonal regulation in
cultured stromal

[138766]cells and messenger RNA expression during the menstrual
cycle. J.

[138767]Clin. Invest. 97: 438-447, 1996.

[138768]24756.

[138769]Itoh, T.; Tanioka, M.; Yoshida, H.; Yoshioka, T.; Nishimoto,
H.;

[138770]Itohara, S.: Reduced angiogenesis and tumor progression
in gelatinase

[138771]A-deficient mice. Cancer Res. 58: 1048-1051, 1998.

[138772]24757.

[138773]Kato, T.; Kure, T.; Chang, J.-H.; Gabison, E. E.; Itoh, T.;
Itohara,

[138774]S.; Azar, D. T.: Diminished corneal angiogenesis in gelati-
nase A-deficient

[138775]mice. FEBS Lett. 508: 187-190, 2001.

[138776]24758.

[138777]Morgunova, E.; Tuuttila, A.; Bergmann, U.; Isupov, M.;
Lindqvist,

[138778]Y.; Schneider, G.; Tryggvason, K.: Structure of human pro-
matrix

[138779]metalloproteinase-2: activation mechanism revealed. Sci-
ence 284:

[138780]1667-1670, 1999.

[138781]24759.

[138782]Vu, T. H.: Don't mess with the matrix. Nature Genet. 28:
202-204,

[138783]2001.

[138784]24760.

[138785]Willecke, K.; Jungbluth, S.; Dahl, E.; Hennemann, H.;
Heynkes,

[138786]R.; Grzeschik, K.-H.: Six genes of the human connexin
gene family

[138787]coding for gap junctional proteins are assigned to four
different

[138788]human chromosomes. Europ. J. Cell Biol. 53: 275-280,
1990.

[138789]24761.

[138790]Reed, K. E.; Westphale, E. M.; Larson, D. M.; Wang, H.-Z.;
Veenstra,

[138791]R. D.; Beyer, E. C.: Molecular cloning and functional ex-
pression

[138792]of human connexin37, an endothelial cell gap junction
protein. J.

[138793]Clin. Invest. 91: 997-1004, 1993.

[138794]24762.

[138795]Van Camp, G.; Coucke, P.; Speleman, F.; Van Roy, N.;
Beyer, E.

[138796]C.; Oostra, B. A.; Willems, P. J.: The gene for human gap
junction

[138797]protein connexin37 (GJA4) maps to chromosome 1p35.1,
in the vicinity

[138798]of D1S195. Genomics 30: 402-403, 1995.

[138799]24763.

[138800]Gelb, B. D.; Zhang, J.; Cotter, P. D.; Gershin, I. F.; Desnick,

[138801]R. J.: Physical mapping of the human connexin 40 (GJA5)
flavin-containing

[138802]monooxygenase 5, and natriuretic peptide receptor A

genes on 1q21. Genomics 39:

[138803]409–411, 1997.

[138804]24764.

[138805]Kanter, H. L.; Saffitz, J. E.; Beyer, E. C.: Cardiac myocytes

[138806]express multiple gap junction proteins. Circ. Res. 70:

438–444,

[138807]1992.

[138808]24765.

[138809]Kanter, H. L.; Saffitz, J. E.; Beyer, E. C.: Molecular cloning

[138810]of two human cardiac gap junction proteins, connexin40

and connexin45. J.

[138811]Molec. Cell. Cardiol. 26: 861–868, 1994.

[138812]24766.

[138813]Oviedo–Orta, E.; Hoy, T.; Evans, W. H.: Intercellular com-

munication

[138814]in the immune system: differential expression of con-

nexin40 and 43,

[138815]and perturbation of gap junction channel functions in pe-

ripheral blood

[138816]and tonsil human lymphocyte subpopulations. Immunol–

ogy 99: 578–590,

[138817]2000.

[138818]24767.

[138819]Seul, K. H.; Tadros, P. N.; Beyer, E. C.: Mouse connexin40:
gene

[138820]structure and promoter analysis. Genomics 46: 120–126,
1997.

[138821]24768.

[138822]Britz–Cunningham, S. H.; Shah, M. M.; Zuppan, C. W.;
Fletcher,

[138823]W. H.: Mutations of the connexin43 gap–junction gene in
patients

[138824]with heart malformations and defects of laterality. New
Eng. J. Med. 332:

[138825]1323–1329, 1995.

[138826]24769.

[138827]Brueckner, M.; D'Eustachio, P.; Horwich, A. L.: Linkage
mapping

[138828]of a mouse gene, iv, that controls left–right asymmetry of
the heart

[138829]and viscera. Proc. Nat. Acad. Sci. 86: 5035–5038, 1989.

[138830]24770.

[138831]Burdine, R. D.; Schier, A. F.: Conserved and divergent mechanisms

[138832]in left-right axis formation. Genes Dev. 14: 763-776, 2000.

[138833]24771.

[138834]Casey, B.; Ballabio, A.: Connexin43 mutations in sporadic and

[138835]familial defects of laterality. (Letter) New Eng. J. Med. 333: 941,

[138836]1995.

[138837]24772.

[138838]Corcos, I. A.; Meese, E. U.; Loch-Caruso, R.: Human connexin 43

[138839]gene locus, GJA1, sublocalized to band 6q21-q23.2. Cytogenet. Cell

[138840]Genet. 64: 31-32, 1993.

[138841]24773.

[138842]Debrus, S.; Tuffery, S.; Matsuoka, R.; Galal, O.; Sarda, P.; Sauer,

[138843]U.; Bozio, A.; Tanman, B.; Toutain, A.; Claustres, M.; Le

Paslier,

[138844]D.; Bouvagnet, P.: Lack of evidence for connexin 43 gene mutations

[138845]in human autosomal recessive lateralization defects. J. Molec. Cell.

[138846]Cardiol. 29: 1423–1431, 1997.

[138847]24774.

[138848]Przylepa, K. A.; Paznekas, W.; Zhang, M.; Golabi, M.; Bias, W.;

[138849]Bamshad, M. J.; Carey, J. C.; Hall, B. D.; Stevenson, R.; Orlow, S.

[138850]J.; Cohen, M. M., Jr.; Jabs, E. W.: Fibroblast growth factor receptor

[138851]2 mutations in Beare–Stevenson cutis gyrata syndrome. Nature Genet. 13:

[138852]492–494, 1996.

[138853]24775.

[138854]Hai, T.; Liu, F.; Coukos, W. J.; Green, M. R.: Transcription factor

[138855]ATF cDNA clones: an extensive family of leucine zipper proteins able

[138856]to selectively form DNA-binding heterodimers. Genes Dev.

3: 2083–2090,

[138857]1989.

[138858]24776.

[138859]Herzig, S.; Long, F.; Jhala, U. S.; Hedrick, S.; Quinn, R.;
Bauer,

[138860]A.; Rudolph, D.; Schutz, G.; Yoon, C.; Puigserver, P.;
Spiegelman,

[138861]B.; Montminy, M.: CREB regulates hepatic gluconeogenesis
through

[138862]the coactivator PGC-1. *Nature* 413: 179–183, 2001.

[138863]24777.

[138864]Benson, A. M.; Hunkeler, M. J.; Talalay, P.: Increase of
NAD(P)H:quinone

[138865]reductase by dietary antioxidants: possible role in protec-
tion against

[138866]carcinogenesis and toxicity. *Proc. Nat. Acad. Sci.* 77:
5216–5220,

[138867]1980.

[138868]24778.

[138869]Chen, L. Z.; Harris, P. C.; Apostolou, S.; Baker, E.; Holman,
K.;

[138870] Lane, S. A.; Nancarrow, J. K.; Whitmore, S. A.; Stallings, R. L.;

[138871] Hildebrand, C. E.; Richards, R. I.; Sutherland, G. R.; Callen, D.

[138872] F.: A refined physical map of the long arm of human chromosome 16. Genomics 10:

[138873] 308–312, 1991.

[138874] 24779.

[138875] Edwards, Y. H.; Hopkinson, D. A.; Carritt, B.: A genetic characterization

[138876] of the human diaphorase–4 deficiency. Ann. Hum. Genet. 47: 97–105,

[138877] 1983.

[138878] 24780.

[138879] Grzeschik, K.–H.: Assignment of a structural gene for a fourth

[138880] human diaphorase (DIA–4) to chromosome 16 in man–mouse somatic cell

[138881] hybrids. Hum. Genet. 53: 189–193, 1980.

[138882] 24781.

[138883] Jaiswal, A. K.; McBride, O. W.; Adesnik, M.; Nebert, D. W.:

Human

[138884]dioxin-inducible cytosolic NAD(P)H:menadione oxidoreductase: cDNA

[138885]sequence and localization of gene to chromosome 16. J. Biol. Chem. 263:

[138886]13572-13578, 1988.

[138887]24782.

[138888]Kelsey, K. T.; Ross, D.; Traver, R. D.; Christiani, D. C.; Zuo,

[138889]Z. F.; Spitz, M. R.; Wang, M.; Xu, X.; Lee, B. K.; Schwartz, B. S.;

[138890]Wiencke, J. K.: Ethnic variation in the prevalence of a common NAD(P)H

[138891]quinone oxidoreductase polymorphism and its implications for anti-cancer

[138892]chemotherapy. Brit. J. Cancer 76: 852-854, 1997.

[138893]24783.

[138894]Lavinha, J.; Morrison, N.; Glasgow, L.; Ferguson-Smith, M. A.:

[138895]Further evidence for the regional localization of human APRT and DIA4

[138896]on chromosome 16. (Abstract) Cytogenet. Cell Genet. 37:

517 only,

[138897]1984.

[138898]24784.

[138899]Moran, J. L.; Siegel, D.; Ross, D.: A potential mechanism
underlying

[138900]the increased susceptibility of individuals with a polymor-
phism in

[138901]NAD(P)H:quinone oxidoreductase 1 (NQO1) to benzene
toxicity. Proc.

[138902]Nat. Acad. Sci. 96: 8150–8155, 1999.

[138903]24785.

[138904]Povey, S.; Wilson, D.; Edwards, Y. H.: Assignment of a hu-
man diaphorase

[138905](DIA-4) to chromosome 16. Ann. Hum. Genet. 43:
349–353, 1980.

[138906]24786.

[138907]Radjendirane, V.; Joseph, P.; Lee, Y.-H.; Kimura, S.; Klein-
Szanto,

[138908]A. J. P.; Gonzalez, F. J.; Jaiswal, A. K.: Disruption of the DT
diaphorase

[138909](NQO1) gene in mice leads to increased menadione toxic-

ity. J. Biol.

[138910]Chem. 273: 7382–7389, 1998.

[138911]24787.

[138912]Rothman, N.; Smith, M. T.; Hayes, R. B.; Traver, R. D.;
Hoener,

[138913]B.; Campleman, S.; Li, G. L.; Dosemeci, M.; Linet, M.;
Zhang, L.;

[138914]Xi, L.; Wacholder, S.; Lu, W.; Meyer, K. B.; Titenko–Holland,
N.;

[138915]Stewart, J. T.; Yin, S.; Ross, D.: Benzene poisoning, a risk
factor

[138916]for hematological malignancy, is associated with the
NQO1 609C–T mutation

[138917]and rapid fractional excretion of chlorzoxazone. Cancer
Res. 57:

[138918]2839–2842, 1997.

[138919]24788.

[138920]Smith, M. T.: Benzene, NQO1, and genetic susceptibility to
cancer.

[138921](Commentary) Proc. Nat. Acad. Sci. 96: 7624–7626, 1999.

[138922]24789.

[138923]Traver, R. D.; Horikoshi, T.; Danenberg, K. D.; Stadlbauer, T.

[138924]H.; Danenberg, P. V.; Ross, D.; Gibson, N. W.:

NAD(P)H:quinone oxidoreductase

[138925]gene expression in human colon carcinoma cells: characterization of

[138926]a mutation which modulates DT-diaphorase activity and mitomycin sensitivity. Cancer

[138927]Res. 52: 797-802, 1992.

[138928]24790.

[138929]Traver, R. D.; Siegel, D.; Beall, H. D.; Phillips, R. M.; Gibson,

[138930]N. W.; Franklin, W. A.; Ross, D.: Characterization of a polymorphism

[138931]in NAD(P)H: quinone oxidoreductase (DT-diaphorase). Brit. J. Cancer 75:

[138932]69-75, 1997.

[138933]24791.

[138934]Zhang, L.; Rothman, N.; Wang, Y.; Hayes, R. B.; Li, G.; Dosemeci,

[138935]M.; Yin, S.; Kolachana, P.; Titenko-Holland, N.; Smith, M. T.: Increased

[138936]aneusomy and long arm deletion of chromosomes 5 and 7
in the lymphocytes

[138937]of Chinese workers exposed to benzene. Carcinogenesis
19: 1955–1961,

[138938]1998.

[138939]24792.

[138940]Zhang, L.; Wang, Y.; Shang, N.; Smith, M. T.: Benzene
metabolites

[138941]induce the loss of long arm deletion of chromosomes 5
and 7 in human

[138942]lymphocytes. Leukemia Res. 22: 105–113, 1998.

[138943]24793.

[138944]Roychoudhury, A. K.; Nei, M.: Human Polymorphic Genes:
World Distribution.

[138945]New York: Oxford Univ. Press (pub.) 1988.

[138946]24794.

[138947]Hark, A. T.; Schoenherr, C. J.; Katz, D. J.; Ingram, R. S.;
Levrose,

[138948]J. M.; Tilghman, S. M.: CTCF mediates methylation–sensi-
tive enhancer–blocking

[138949]activity at the H19/Igf2 locus. Nature 405: 486–489,

2000.

[138950]24795.

[138951]Hark, A. T.; Tilghman, S. M.: Chromatin conformation of
the H19

[138952]epigenetic mark. Hum. Molec. Genet. 7: 1979–1985,
1998.

[138953]24796.

[138954]Jones, B. K.; Levorse, J.; Tilghman, S. M.: Deletion of a nu-
clease-sensitive

[138955]region between the Igf2 and H19 genes leads to Igf2 mis-
regulation

[138956]and increased adiposity. Hum. Molec. Genet. 10:
807–814, 2001.

[138957]24797.

[138958]Jones, J. M.; Meisler, M. H.; Seldin, M. F.; Lee, B. K.; Eicher,

[138959]E. M.: Localization of insulin-2 (Ins-2) and the obesity
mutant tubby

[138960](tub) to distinct regions of mouse chromosome 7. Ge-
nomics 14: 197–199,

[138961]1992.

[138962]24798.

[138963]Rainier, S.; Johnson, L. A.; Dobry, C. J.; Ping, A. J.; Grundy,
[138964]P. E.; Feinberg, A. P.: Relaxation of imprinted genes in hu-
man cancer. Nature 362:
[138965]747–749, 1993.
[138966]24799.
[138967]Srivastava, M.; Hsieh, S.; Grinberg, A.; Williams–Simons, L.;
[138968]Huang, S.–P.; Pfeifer, K.: H19 and Igf2 monoallelic expres-
sion is
[138969]regulated in two distinct ways by a shared cis acting regu-
latory region
[138970]upstream of H19. Genes Dev. 14: 1186–1195, 2000.
[138971]24800.
[138972]Thorvaldsen, J. L.; Duran, K. L.; Bartolomei, M. S.: Deletion
[138973]of the H19 differentially methylated domain results in loss
of imprinted
[138974]expression of H19 and Igf2. Genes Dev. 12: 3693–3702,
1998.
[138975]24801.
[138976]Webber, A.; Ingram, R. S.; Levorse, J. M.; Tilghman, S. M.:
Location

[138977]of enhancers is essential for imprinting of H19 and Igf2
genes. Nature 391:

[138978]711–715, 1998.

[138979]24802.

[138980]Zhang, Y.; Tycko, B.: Monoallelic expression of the human
H19

[138981]gene. Nature Genet. 1: 40–44, 1992.

[138982]24803.

[138983]Campanelli, J. T.; Hoch, W.; Rupp, F.; Kreiner, T.; Scheller,
R.

[138984]H.: Agrin mediates cell contact–induced acetylcholine re-
ceptor clustering. Cell 67:

[138985]909–916, 1991.

[138986]24804.

[138987]DeChiara, T. M.; Bowen, D. C.; Valenzuela, D. M.; Sim-
mons, M. V.;

[138988]Poueymirou, W. T.; Thomas, S.; Kinetz, E.; Compton, D. L.;
Rojas,

[138989]E.; Park, J. S.; Smith, C.; DiStefano, P. S.; Glass, D. J.; Bur-
den,

[138990]S. J.; Yancopoulos, G. D.: The receptor tyrosine kinase

MuSK is required

[138991]for neuromuscular junction formation in vivo. Cell 85:
501–512,

[138992]1996.

[138993]24805.

[138994]Chan, S. Y.; Empig, C. J.; Welte, F. J.; Speck, R. F.;
Schmaljohn,

[138995]A.; Kreisberg, J. F.; Goldsmith, M. A.: Folate receptor- α is
pha is

[138996]a cofactor for cellular entry by Marburg and Ebola viruses.
Cell 106:

[138997]117–126, 2001.

[138998]24806.

[138999]Czeizel, A. E.; Dudas, I.: Prevention of the first occurrence

[139000]of neural-tube defects by periconceptional vitamin sup-
plementation. New

[139001]Eng. J. Med. 327: 1832–1835, 1992.

[139002]24807.

[139003]De Marco, P.; Moroni, A.; Merello, E.; de Franchis, R.; An-
dreussi,

[139004]L.; Finnell, R. H.; Barber, R. C.; Cama, A.; Capra, V.: Folate

pathway

[139005]gene alterations in patients with neural tube defects. Am. J. Med.

[139006]Genet. 95: 216–223, 2000.

[139007]24808.

[139008]Elwood, P. C.: Molecular cloning and characterization of the human

[139009]folate-binding protein cDNA from placenta and malignant tissue culture

[139010](KB) cells. J. Biol. Chem. 264: 14893–14901, 1989.

[139011]24809.

[139012]Elwood, P. C.; Nachmanoff, K.; Saikawa, Y.; Page, S. T.; Pacheco,

[139013]P.; Roberts, S.; Chung, K.–N.: The divergent 5–prime termini of the

[139014]alpha human folate receptor (hFR) mRNAs originate from two tissue-specific

[139015]promoters and alternative splicing: characterization of the alpha

[139016]hFR gene structure. Biochemistry 36: 1467–1478, 1997.

[139017]24810.

- [139018]Boman, B. M.; Wildrick, D. M.; Alfaro, S. R.: Chromosome
18 allele
- [139019]Loss at the D18S6 locus in human colorectal carcinomas.
Biochem.
- [139020]Biophys. Res. Commun. 155: 463–469, 1988.
- [139021]24811.
- [139022]Chan, S. S.–Y.; Zheng, H.; Su, M.–W.; Wilk, R.; Killeen, M.
T.;
- [139023]Hedgecock, E. M.; Culotti, J. G.: UNC–40, a *C. elegans* ho–
molog of
- [139024]the DCC (deleted in colorectal cancer), is required in
motile cells
- [139025]responding to UNC–6 netrin cues. Cell 87: 187–195,
1996.
- [139026]24812.
- [139027]Sarmanova, J.; Benesova, K.; Gut, I.; Nedelcheva–Kris–
tensen, V.;
- [139028]Tynkova, L.; Soucek, P.: Genetic polymorphisms of bio–
transformation
- [139029]enzymes in patients with Hodgkin's and non–Hodgkin's
lymphomas. Hum.
- [139030]Molec. Genet. 10: 1265–1273, 2001.

[139031]24813.

[139032]Albrecht, F. E.; Drago, J.; Felder, R. A.; Printz, M. P.; Eisner,

[139033]G. M.; Robillard, J. E.; Sibley, D. R.; Westphal, H. J.; Jose, P.

[139034]A.: Role of the D-1A dopamine receptor in the pathogenesis of genetic

[139035]hypertension. J. Clin. Invest. 97: 2283-2288, 1996.

[139036]24814.

[139037]Bermak, J. C.; Li, M.; Bullock, C.; Zhou, Q.-Y.: Regulation of

[139038]transport of the dopamine D1 receptor by a new membrane-associated

[139039]ER protein. Nature Cell Biol. 3: 492-498, 2001.

[139040]24815.

[139041]Boulton, J.; Lewis, M. S.; Wainscoat, J. S.: Physical linkage

[139042]of glucocorticoid receptor (GRL) and dopamine D1 receptor (DRD1) on

[139043]the long arm of chromosome 5.(Abstract) Cytogenet. Cell Genet. 58:

[139044]1894 only, 1991.

[139045]24816.

[139046]Castner, S. A.; Williams, G. V.; Goldman-Rakic, P. S.: Re-
versal

[139047]of antipsychotic-induced working memory deficits by
short-term dopamine

[139048]D1 receptor stimulation. Science 287: 2020-2022, 2000.

[139049]24817.

[139050]Dearry, A.; Gingrich, J. A.; Falardeau, P.; Fremeau, R. T.,
Jr.;

[139051]Bates, M. D.; Caron, M. G.: Molecular cloning and expres-
sion of the

[139052]gene for a human D(1) dopamine receptor. Nature 347:
72-76, 1990.

[139053]24818.

[139054]Grandy, D. K.; Zhou, Q.-Y.; Allen, L.; Litt, R.; Magenis, R.
E.;

[139055]Civelli, O.; Litt, M.: A human D(1) dopamine receptor gene
is located

[139056]on chromosome 5 at q35.1 and identifies an EcoRI RFLP.
Am. J. Hum.

[139057]Genet. 47: 828-834, 1990.

[139058]24819.

[139059]Krushkal, J.; Xiong, M.; Ferrell, R.; Sing, C. F.; Turner, S. T.;

[139060]Boerwinkle, E.: Linkage and association of adrenergic and
dopamine

[139061]receptor genes in the distal portion of the long arm of
chromosome

[139062]5 with systolic blood pressure variation. Hum. Molec.
Genet. 7:

[139063]1379–1383, 1998.

[139064]24820.

[139065]Mayerhofer, A.; Hemmings, H. C., Jr.; Snyder, G. L.; Green-
gard,

[139066]P.; Boddien, S.; Berg, U.; Brucker, C.: Functional
dopamine-1 receptors

[139067]and DARPP-32 are expressed in human ovary and granu-
losa luteal cells

[139068]in vitro. J. Clin. Endocr. Metab. 84: 257–264, 1999.

[139069]24821.

[139070]Sunahara, R. K.; Niznik, H. B.; Weiner, D. M.; Stormann, T.
M.;

[139071]Brann, M. R.; Kennedy, J. L.; Gelernter, J. E.; Rozmahel, R.;

Yang,

[139072]Y.; Israel, Y.; Seeman, P.; O'Dowd, B. F.: Human dopamine
D1 receptor

[139073]encoded by an intronless gene on chromosome 5. Nature
347: 80–83,

[139074]1990.

[139075]24822.

[139076]Wilkie, T. M.; Chen, Y.; Gilbert, D. J.; Moore, K. J.; Yu, L.;

[139077]Simon, M. I.; Copeland, N. G.; Jenkins, N. A.: Identification,
chromosomal

[139078]location, and genome organization of mammalian G-
protein-coupled receptors. Genomics 18:

[139079]175–184, 1993.

[139080]24823.

[139081]Xu, M.; Moratalla, R.; Gold, L. H.; Hiroi, N.; Koob, G. F.;
Graybiel,

[139082]A. M.; Tonegawa, S.: Dopamine D1 receptor mutant mice
are deficient

[139083]in striatal expression of dynorphin and in dopamine-mediated behavioral

[139084]responses. Cell 79: 729–742, 1994.

[139085]24824.

[139086]Zhou, Q.-Y.; Grandy, D. K.; Thambi, L.; Kushner, J. A.; Van Tol,

[139087]H. H. M.; Cone, R.; Pribnow, D.; Salon, J.; Bunzow, J. R.; Civelli,

[139088]O.: Cloning and expression of human and rat D(1) dopamine receptors. Nature 347:

[139089]76-80, 1990.

[139090]24825.

[139091]Balk, J.-H.; Picetti, R.; Salardi, A.; Thirlet, G.; Dierich, A.;

[139092]Depaulls, A.; Le Meur, M.; Borrelli, E.: Parkinsonian-like locomotor

[139093]impairment in mice lacking dopamine D2 receptors. Nature 377: 424-428,

[139094]1995.

[139095]24826.

[139096]Basu, S.; Nagy, J. A.; Pal, S.; Vasile, E.; Eckelhoefer, I. A.;

[139097]Bliss, V. S.; Manseau, E. J.; Dasgupta, P. S.; Dvorak, H. F.; Mukhopadhyay,

[139098]D.: The neurotransmitter dopamine inhibits angiogenesis induced by

[139099]vascular permeability factor/vascular endothelial growth factor. Nature

[139100]Med. 7: 569–574, 2001.

[139101]24827.

[139102]Blum, K.; Noble, E. P.; Sheridan, P. J.; Montgomery, A.; Ritchie,

[139103]T.; Jagadeeswaran, P.; Nogami, H.; Briggs, A. H.; Cohn, J. B.: Allelic

[139104]association of human dopamine D(2) receptor gene in alcoholism. J.A.M.A. 263:

[139105]2055–2060, 1990.

[139106]24828.

[139107]Bolos, A. M.; Dean, M.; Lucas–Derse, S.; Ramsburg, M.; Brown, G.

[139108]L.; Goldman, D.: Population and pedigree studies reveal a lack of

[139109]association between the dopamine D(2) receptor gene and alcoholism. J.A.M.A. 264:

[139110]3156–3160, 1990.

[139111]24829.

[139112]Bunzow, J. R.; Van Tol, H. H.; Grandy, D. K.; Albert, P.; Sa–

lon,

[139113]J.; Christie, M.; Machida, C. A.; Neve, K. A.; Civelli, O.:

Cloning

[139114]and expression of a rat D2 dopamine receptor cDNA. Nature 336: 783–787,

[139115]1988.

[139116]24830.

[139117]Dal Toso, R.; Sommer, B.; Ewert, M.; Herb, A.; Pritchett, D. B.;

[139118]Bach, A.; Shivers, B. D.; Seeburg, P. H.: The dopamine D(2) receptor:

[139119]two molecular forms generated by alternative splicing.

EMBO J. 8:

[139120]4025–4034, 1989.

[139121]24831.

[139122]Matsutani, A.; Hing, A.; Steinbrueck, T.; Janssen, R.; Weber,

[139123]J.; Permutt, M. A.; Donis-Keller, H.: Mapping the human liver/islet

[139124]glucose transporter (GLUT2) gene within a genetic linkage map of chromosome

[139125]3q using a (CA)_n dinucleotide repeat polymorphism and

characterization

[139126]of the polymorphism in three racial groups. Genomics 13:
495–501,

[139127]1992.

[139128]24832.

[139129]Meera Khan, P.: Personal Communication. Leiden, The
Netherlands

[139130]1977.

[139131]24833.

[139132]Garman, S. C.; Kinet, J.-P.; Jardetzky, T. S.: Crystal struc-
ture

[139133]of the human high-affinity IgE receptor. Cell 95: 951–961,
1998.

[139134]24834.

[139135]Garman, S. C.; Wurzburg, B. A.; Tarchevskaya, S. S.; Kinet,
J.-P.;

[139136]Jardetzky, T. S.: Structure of the Fc fragment of human IgE
bound

[139137]to its high-affinity receptor Fc-epsilon-RI-alpha. Nature
406: 259–266,

[139138]2000.

[139139]24835.

[139140]Liu, F.-T.; Albrandt, K.; Robertson, M. W.: cDNA hetero-
geneity

[139141]suggests structural variants related to the high-affinity IgE
receptor. Proc.

[139142]Nat. Acad. Sci. 85: 5639-5643, 1988.

[139143]24836.

[139144]Shimizu, A.; Tepler, I.; Benfey, P. N.; Berenstein, E. H.; Sir-
aganian,

[139145]R. P.; Leder, P.: Human and rat mast cell high-affinity im-
munoglobulin

[139146]E receptors: characterization of putative alpha-chain gene
products. Proc.

[139147]Nat. Acad. Sci. 85: 1907-1911, 1988.

[139148]24837.

[139149]Miyajima, I.; Dombrowicz, D.; Martin, T. R.; Ravetch, J. V.;
Kinet,

[139150]J.-P.; Galli, S. J.: Systemic anaphylaxis in the mouse can be
mediated

[139151]largely through IgG1 and Fc-gamma-RIII: assessment of
the cardiopulmonary

[139152]changes, mast cell degranulation, and death associated with active

[139153]or IgE- or IgG1-dependent passive anaphylaxis. J. Clin. Invest. 99:

[139154]901-914, 1997.

[139155]24838.

[139156]Tepler, I.; Morton, C. C.; Shimizu, A.; Holcombe, R. F.; Eddy,

[139157]R.; Shows, T. B.; Leder, P.: The gene for the human mast cell high-affinity

[139158]IgE receptor alpha chain: chromosomal localization to 1q21-q23 and

[139159]RFLP analysis. Am. J. Hum. Genet. 45: 761-765, 1989.

[139160]24839.

[139161]Zhu, D.; Kepley, C. L.; Zhang, M.; Zhang, K.; Saxon, A.: A novel

[139162]human immunoglobulin Fc-gamma-Fc-epsilon bifunctional fusion protein

[139163]inhibits Fc-epsilon-RI-mediated degranulation. Nature Med. 8: 518-521,

[139164]2002.

[139165]24840.

[139166]Bain, G.; Robanus Maandag, E. C.; Izon, D. J.; Amsen, D.;
Kruisbeek,

[139167]A. M.; Weintraub, B. C.; Krop, I.; Schlissel, M. S.; Feeney, A.
J.;

[139168]van Roon, M.; van der Valk, M.; te Riele, H. P. J.; Berns, A.;
Murre,

[139169]C.: E2A proteins are required for proper B cell develop-
ment and initiation

[139170]of immunoglobulin gene rearrangements. Cell 79:
885–892, 1994.

[139171]24841.

[139172]El Ghouzzi, V.; Legeai-Mallet, L.; Aresta, S.; Benoist, C.;
Munnich,

[139173]A.; de Gunzburg, J.; Bonaventure, J.: Saethre–Chotzen mu-
tations cause

[139174]TWIST protein degradation or impaired nuclear location.
Hum. Molec.

[139175]Genet. 9: 813–819, 2000.

[139176]24842.

[139177]Henthorn, P.; Kiledjian, M.; Kadesch, T.: Two distinct tran-
scription

[139178]factors that bind the immunoglobulin enhancer mu-
E5/kappa-E2 motif. Science 247:

[139179]467-470, 1990.

[139180]24843.

[139181]Kamps, M. P.; Murre, C.; Sun, X.; Baltimore, D.: A new
homeobox

[139182]gene contributes the DNA binding domain of the t(1;19)
translocation

[139183]protein in pre-B ALL. Cell 60: 547-555, 1990.

[139184]24844.

[139185]Mellentin, J. D.; Murre, C.; Donlon, T. A.; McCaw, P. S.;
Smith,

[139186]S. D.; Carroll, A. J.; McDonald, M. E.; Baltimore, D.; Cleary,
M.

[139187]L.: The gene for enhancer binding proteins E12/E47 lies at
the t(1;19)

[139188]breakpoint in acute leukemias. Science 246: 379-382,
1989.

[139189]24845.

[139190]Murre, C.; McCaw, P. S.; Baltimore, D. :Cell 56: 777-783,
1989.

[139191]24846.

[139192]Nourse, J.; Mellentin, J. D.; Galili, N.; Wilkinson, J.; Stan-
bridge,

[139193]E.; Smith, S. D.; Cleary, M. L.: Chromosomal translocation
t(1;19)

[139194]results in synthesis of a homeobox fusion mRNA that
codes for a potential

[139195]chimeric transcription factor. Cell 60: 535–545, 1990.

[139196]24847.

[139197]Matsutani, A.; Koranyi, L.; Cox, N.; Permutt, M. A.: Poly-
morphisms

[139198]of GLUT2 and GLUT4 genes: use in evaluation of genetic
susceptibility

[139199]to NIDDM in Blacks. Diabetes 39: 1534–1542, 1990.

[139200]24848.

[139201]Moller, A. M.; Jensen, N. M.; Pildal, J.; Drivsholm, T.;
Borch-Johnsen,

[139202]K.; Urhammer, S. A.; Hansen, T.; Pedersen, O.: Studies of
genetic

[139203]variability of the glucose transporter 2 promoter in pa-
tients with

- [139204]type 2 diabetes mellitus. J. Clin. Endocr. Metab. 86:
2181–2186,
[139205]2001.
[139206]24849.
[139207]Mueckler, M.; Caruso, C.; Baldwin, S. A.; Panico, M.;
Blench,
[139208]J.; Morris, H. R.; Allard, W. J.; Lienhard, G. E.; Lodish, H. F.:
[139209]Sequence and structure of a human glucose transporter.
Science 229:
[139210]941–945, 1985.
[139211]24850.
[139212]Mueckler, M.; Kruse, M.; Strube, M.; Riggs, A. C.; Chiu, K.
C.;
[139213]Permutt, M. A.: A mutation in the Glut2 glucose trans-
porter gene
[139214]of a diabetic patient abolishes transport activity. J. Biol.
Chem. 269:
[139215]17765–17767, 1994.
[139216]24851.
[139217]Muller, D.; Santer, R.; Krawinkel, M.; Christiansen, B.;
Schaub,

- [139218]J.: Fanconi-Bickel syndrome presenting in neonatal screening for galactosemia. J. Inherit. Metab. Dis. 20: 607-608, 1997.
- [139220]24852.
- [139221]Oka, Y.; Asano, T.; Shibaskai, Y.; Lin, J.-L.; Tsukuda, K.; Katagiri, H.; Akanuma, Y.; Takaku, F.: C-terminal truncated glucose transporter is locked into an inward-facing form without transport activity. Nature 345: 550-553, 1990.
- [139225]24853.
- [139226]Orci, L.; Thorens, B.; Ravazzola, M.; Lodish, H. F.: Localization of the pancreatic beta cell glucose transporter to specific plasma membrane domains. Science 245: 295-297, 1989.
- [139229]24854.
- [139230]Permutt, M. A.; Koranyi, L.; Keller, K.; Lacy, P. E.; Scharp, D. W.; Mueckler, M.: Cloning and functional expression of a human

[139232]pancreatic islet glucose-transporter cDNA. Proc. Nat.
Acad. Sci. 86:

[139233]8688-8692, 1989.

[139234]24855.

[139235]Sakamoto, O.; Ogawa, E.; Ohura, T.; Igarashi, Y.; Matsub-
ara, Y.;

[139236]Narisawa, K.; Iinuma, K.: Mutation analysis of the GLUT2
gene in

[139237]patients with Fanconi-Bickel syndrome. Pediat. Res. 48:
586-589,

[139238]2000.

[139239]24856.

[139240]Sanjad, S. A.; Kaddoura, R. E.; Nazer, H. M.; Akhtar, M.;
Sakati,

[139241]N. A.: Fanconi's syndrome with hepatorenal glycogenosis
associated

[139242]with phosphorylase b kinase deficiency. Am. J. Dis. Child.
147:

[139243]957-959, 1993.

[139244]24857.

[139245]Santer, R.; Groth, S.; Kinner, M.; Dombrowski, A.; Berry, G.

T.;

[139246]Brodehl, J.; Leonard, J. V.; Moses, S.; Norgren, S.; Skovby, F.; Schneppenheim,

[139247]R.; Steinmann, B.; Schaub, J.: The mutation spectrum of the facilitative

[139248]glucose transporter gene SLC2A2 (GLUT2) in patients with Fanconi-Bickel

[139249]syndrome. Hum. Genet. 110: 21-29, 2002.

[139250]24858.

[139251]Santer, R.; Schneppenheim, R.; Dombrowski, A.; Gotze, H.; Steinmann,

[139252]B.; Schaub, J.: Fanconi-Bickel syndrome--a congenital defect of the

[139253]liver-type facilitative glucose transporter. J. Inherit. Metab. Dis. 21:

[139254]191-194, 1998.

[139255]24859.

[139256]Santer, R.; Schneppenheim, R.; Dombrowski, A.; Gotze, H.; Steinmann,

[139257]B.; Schaub, J.: Mutations in GLUT2, the gene for the liver-type glucose

[139258]transporter, in patients with Fanconi-Bickel syndrome.

Nature Genet. 17:

[139259]324–326, 1997. Note: Erratum: Nature Genet. 18: 298 only, 1998.

[139260]24860.

[139261]Steinmann, B.; Zeller, L.: Personal Communication. Zurich and

[139262]Santa Maria Calanca, Switzerland 8/2/1997.

[139263]24861.

[139264]Takeda, J.; Kayano, T.; Fukumoto, H.; Bell, G. I.: Organization

[139265]of the human GLUT2 (pancreatic beta-cell and hepatocyte) glucose transporter

[139266]gene. Diabetes 42: 773–777, 1993.

[139267]24862.

[139268]Tanizawa, Y.; Riggs, A. C.; Chiu, K. C.; Janssen, R. C.; Bell,

[139269]D. S. H.; Go, R. P. C.; Roseman, J. M.; Acton, R. T.; Permutt, M.

[139270]A.: Variability of the pancreatic islet beta cell/liver (GLUT 2)

[139271]glucose transporter gene in NIDDM patients. Diabetologia 37: 420–427,

[139272]1994.

[139273]24863.

[139274]Thorens, B.; Deriaz, N.; Bosco, D.; DeVos, A.; Pipeleers, D.;

[139275]Schuit, F.; Meda, P.; Porret, A.: Protein kinase A-
dependent phosphorylation

[139276]of GLUT2 in pancreatic beta cells. J. Biol. Chem. 271:
8075-8081,

[139277]1996.

[139278]24864.

[139279]Fan, Y.-S.; Eddy, R. L.; Byers, M. G.; Haley, L. L.; Henry, W.

[139280]M.; Kayano, T.; Shows, T. B.; Bell, G. I.: Assignment of
genes encoding

[139281]three human glucose transporter/transporter-like pro-
teins (GLUT4,

[139282]GLUT5 and GLUT6) to chromosomes 17, 1 and 5, respec-
tively. (Abstract) Cytogenet.

[139283]Cell Genet. 51: 997 only, 1989.

[139284]24865.

[139285]Gould, G. W.; Holman, G. D.: The glucose transporter fam-
ily: structure,

[139286]function and tissue-specific expression. Biochem. J. 295:
329-341,

[139287]1993.

[139288]24866.

[139289]Hauguel-de Mouzon, S.; Challier, J. C.; Kacemi, A.;
Cauzac, M.;

[139290]Malek, A.; Girard, J.: The GLUT3 glucose transporter iso-
form is differentially

[139291]expressed within human placental cell types. J. Clin. En-
docr. Metab. 82:

[139292]2689-2694, 1997.

[139293]24867.

[139294]Kayano, T.; Burant, C. F.; Fukumoto, H.; Gould, G. W.; Fan,
Y.

[139295]S.; Eddy, R. L.; Byers, M. G.; Shows, T. B.; Seino, S.; Bell, G.
I.

[139296]: Human facilitative glucose transporters: isolation, func-
tional characterization,

[139297]and gene localization of cDNAs encoding an isoform
(GLUT5) expressed

[139298]in small intestine, kidney, muscle, and adipose tissue and
an unusual

- [139299]glucose transporter pseudogene-like sequence (GLUT6). J. Biol. Chem. 265: 13276-13282, 1990.
- [139300]24868.
- [139301]Kayano, T.; Fukumoto, H.; Eddy, R. L.; Fan, Y.-S.; Byers, M. G.;
- [139302]Shows, T. B.; Bell, G. I.: Evidence for a family of human glucose transporter-like proteins: sequence and gene localization of a protein expressed in fetal skeletal muscle and other tissues. J. Biol. Chem. 263: 15245-15248, 1988.
- [139303]24869.
- [139304]Gitelman, B. J.; Tomkins, D. J.; Partington, M. W.; Roberts, M.;
- [139305]H.; Simpson, N. E.: Gene dosage studies of glutamic oxaloacetic transaminase (GOT) and hexokinase (HK) in two patients with possible partial trisomy 10q. (Abstract) Am. J. Hum. Genet. 32: 41A only, 1980.

[139312]24870.

[139313]Junien, C.; Despoisse, S.; Turleau, C.; de Grouchy, J.;
Bucher,

[139314]T.; Fundele, R.: Assignment of phosphoglycerate mutase
(PGAMA) to

[139315]human chromosome 10: regional mapping of GOT1 and
PGAMA to subbands

[139316]10q26.1 (or q25.3). Ann. Genet. 25: 25-27, 1982.

[139317]24871.

[139318]O'Brien, S. J.; Bonner, T. I.; Cohen, M.; O'Connell, C.; Nash,

[139319]W. G.: Mapping of an endogenous retroviral sequence to
human chromosome

[139320]18. Nature 303: 74-77, 1983.

[139321]24872.

[139322]Overhauser, J.; Mewar, R.; Rojas, K.; Lia, K.; Kline, A. D.;
Silverman,

[139323]G. A.: STS map of genes and anonymous DNA fragments
on human chromosome

[139324]18 using a panel of somatic cell hybrids. Genomics 15:
387-391,

[139325]1993.

[139326]24873.

[139327]Yunis, J. J.; Oken, M. M.; Kaplan, M. E.; Ensrud, K. M.;
Howe,

[139328]R. R.; Theologides, A.: Distinctive chromosomal abnormalities in

[139329]histologic subtypes of non-Hodgkin's lymphoma. New
Eng. J. Med. 307:

[139330]1231-1236, 1982.

[139331]24874.

[139332]Abdalla, S. A.; Pece-Barbara, N.; Vera, S.; Tapia, E.; Paez,
E.;

[139333]Bernabeu, C.; Letarte, M.: Analysis of ALK-1 and endoglin
in newborns

[139334]from families with hereditary hemorrhagic telangiectasia
type 2. Hum.

[139335]Molec. Genet. 9: 1227-1237, 2000.

[139336]24875.

[139337]Bellon, T.; Corbi, A.; Lastres, P.; Cales, C.; Cebrian, M.;
Vera,

[139338]S.; Cheifetz, S.; Massague, J.; Letarte, M.; Bernabeu, C.:
Identification

- [139339]and expression of two forms of the human transforming growth factor- β -binding
- [139340]protein endoglin with distinct cytoplasmic regions. *Europ. J. Immun.* 23:
- [139341]2340–2345, 1993.
- [139342]24876.
- [139343]Bourdeau, A.; Dumont, D. J.; Letarte, M.: A murine model of hereditary
- [139344]hemorrhagic telangiectasia. *J. Clin. Invest.* 104: 1343–1351, 1999.
- [139345]24877.
- [139346]Fernandez-Ruiz, E.; St. Jacques, S.; Bellon, T.; Letarte, M.; Bernabeu,
- [139347]C.: Assignment of the human endoglin gene (END) to 9q34-qter. *Cytogenet. Cell Genet.* 64: 204–207, 1993.
- [139348]24878.
- [139349]24878.
- [139350]Gallione, C. J.; Klaus, D. J.; Yeh, E. Y.; Stenzel, T. T.; Xue,
- [139351]Y.; Anthony, K. B.; McAllister, K. A.; Baldwin, M. A.; Berg, J. N.;
- [139352]Lux, A.; Smith, J. D.; Vary, C. P. H.; Craigen, W. J.; Wester-

mann,

[139353]C. J. J.; Warner, M. L.; Miller, Y. E.; Jackson, C. E.;

Guttmacher,

[139354]A. E.; Marchuk, D. A.: Mutation and expression analysis of
the endoglin

[139355]gene in hereditary hemorrhagic telangiectasia reveals null
alleles. Hum.

[139356]Mutat. 11: 286–294, 1998.

[139357]24879.

[139358]Gallione, C. J.; Scheessele, E. A.; Reinhardt, D.; Duits, A. J.;

[139359]Berg, J. N.; Westermann, C. J. J.; Marchuk, D. A.: Two com-
mon endoglin

[139360]mutations in families with hereditary hemorrhagic telang-
iectasia in

[139361]the Netherlands Antilles: evidence for a founder effect.
Hum. Genet. 107:

[139362]40–44, 2000.

[139363]24880.

[139364]Gougos, A.; Letarte, M.: Primary structure of endoglin, an
RGD-containing

[139365]glycoprotein of human endothelial cells. J. Biol. Chem.

265: 8361–8364,
[139366]1990.

[139367]24881.

[139368]Kozak, M.: Context effects and inefficient initiation at
non-AUG
[139369]codons in eucaryotic cell-free translation systems. Molec.
Cell.
[139370]Biol. 9: 5073–5080, 1989.

[139371]24882.

[139372]Li, D. Y.; Sorensen, L. K.; Brooke, B. S.; Urness, L. D.;
Davis,
[139373]E. C.; Taylor, D. G.; Boak, B. B.; Wendel, D. P.: Defective
angiogenesis
[139374]in mice lacking endoglin. Science 284: 1534–1537, 1999.

[139375]24883.

[139376]Lux, A.; Gallione, C. J.; Marchuk, D. A.: Expression analysis
[139377]of endoglin missense and truncation mutations: insights
into protein
[139378]structure and disease mechanisms. Hum. Molec. Genet. 9:
745–755,
[139379]2000.

[139380]24884.

[139381]McAllister, K. A.; Grogg, K. M.; Johnson, D. W.; Gallione, C.

[139382]J.; Baldwin, M. A.; Jackson, C. E.; Helmbold, E. A.; Markel,
D. S.;

[139383]McKinnon, W. C.; Murrell, J.; McCormick, M. K.; Pericak-
Vance, M.

[139384]A.; Heutink, P.; Oostra, B. A.; Haitjema, T.; Westerman, C.
J. J.;

[139385]Porteous, M. E.; Guttmacher, A. E.; Letarte, M.; Marchuk,
D. A.:

[139386]Endoglin, a TGF- β binding protein of endothelial cells,
is the

[139387]gene for hereditary haemorrhagic telangiectasia type 1.
Nature Genet. 8:

[139388]345-351, 1994.

[139389]24885.

[139390]Paquet, M.-E.; Pece-Barbara, N.; Vera, S.; Cymerman, U.;
Karabegovic,

[139391]A.; Shovlin, C.; Letarte, M.: Analysis of several endoglin
mutants

[139392]reveals no endogenous mature or secreted protein capa-
ble of interfering

[139393]with normal endoglin function. Hum. Molec. Genet. 10:
1347–1357,

[139394]2001.

[139395]24886.

[139396]Pece, N.; Vera, S.; Cymerman, U.; White, R. I., Jr.; Wrana, J.

[139397]L.; Letarte, M.: Mutant endoglin in hereditary hemorrhagic
telangiectasia

[139398]type 1 is transiently expressed intracellularly and is not a
dominant

[139399]negative. J. Clin. Invest. 100: 2568–2579, 1997.

[139400]24887.

[139401]Pece-Barbara, N.; Cymerman, U.; Vera, S.; Marchuk, D. A.;
Letarte,

[139402]M.: Expression analysis of four endoglin missense muta-
tions suggests

[139403]that haploinsufficiency is the predominant mechanism for
hereditary

[139404]hemorrhagic telangiectasia type 1. Hum. Molec. Genet. 8:
2171–2181,

[139405]1999.

[139406]24888.

[139407]Pilz, A.; Woodward, K.; Peters, J.; Povey, S.; Abbott, C.:

Comparative

[139408]mapping of 38 human chromosome 9 loci in the laboratory mouse. (Abstract) Ann.

[139409]Hum. Genet. 58: 231-232, 1994.

[139410]24889.

[139411]Qureshi, S. T.; Gros, P.; Letarte, M.; Malo, D.: The murine endoglin

[139412]gene (Eng) maps to chromosome 2. Genomics 26: 165-166, 1995.

[139413]24890.

[139414]Abbott, C.; Jackson, I. J.; Carritt, B.; Povey, S.: The human

[139415]homolog of the mouse brown gene maps to the short arm of chromosome

[139416]9 and extends the known region of homology with mouse chromosome 4. Genomics 11:

[139417]471-473, 1991.

[139418]24891.

[139419]Box, N. F.; Wyeth, J. R.; Mayne, C. J.; O'Gorman, L. E.; Martin,

[139420]N. G.; Sturm, R. A.: Complete sequence and polymorphism

study of

[139421]the human TYRP1 gene encoding tyrosinase-related protein 1. Mammalian

[139422]Genome 9: 50-53, 1998.

[139423]24892.

[139424]Chintamaneni, C. D.; Ramsay, M.; Colman, M.-A.; Fox, M. F.; Pickard,

[139425]R. T.; Kwon, B. S.: Mapping the human CAS2 gene, the homologue of

[139426]the mouse brown (b) locus, to human chromosome 9p22-pter. Biochem.

[139427]Biophys. Res. Commun. 178: 227-235, 1991.

[139428]24893.

[139429]Cohen, T.; Muller, R. M.; Tomita, Y.; Shibahara, S.: Nucleotide

[139430]sequence of the cDNA encoding human tyrosinase-related protein. Nucleic

[139431]Acids Res. 18: 2807-2808, 1990.

[139432]24894.

[139433]Halaban, R.; Moellmann, G.: Murine and human b locus pigmentation

- [139434]genes encode a glycoprotein (gp75) with catalase activity.
Proc.
- [139435]Nat. Acad. Sci. 87: 4809–4813, 1990.
- [139436]24895.
- [139437]Bode–Lesniewska, B.; Dours–Zimmermann, M. T.; Oder–
matt, B. F.;
- [139438]Briner, J.; Heitz, P. U.; Zimmermann, D. R.: Distribution of
the
- [139439]large aggregating proteoglycan versican in adult human
tissues. J.
- [139440]Histochem. Cytochem. 44: 303–312, 1996.
- [139441]24896.
- [139442]Dours–Zimmermann, M. T.; Zimmermann, D. R.: A novel
glycosaminoglycan
- [139443]attachment domain identified in two alternative splice
variants of
- [139444]human versican. J. Biol. Chem. 269: 32992–32998, 1994.
- [139445]24897.
- [139446]Iozzo, R. V.; Naso, M. F.; Cannizzaro, L. A.; Wasmuth, J. J.;
McPherson,
- [139447]J. D.: Mapping of the versican proteoglycan gene (CSPG2)

to the long

[139448]arm of human chromosome 5 (5q12–5q14). Genomics 14:
845–851, 1992.

[139449]24898.

[139450]Kjellen, L.; Lindahl, U.: Proteoglycans: structures and in-
teractions. Annu.

[139451]Rev. Biochem. 60: 443–475, 1991.

[139452]24899.

[139453]Naso, M. F.; Morgan, J. L.; Buchberg, A. M.; Siracusa, L. D.;
Iozzo,

[139454]R. V.: Expression pattern and mapping of the murine ver-
sican gene

[139455](Cspg2) to chromosome 13. Genomics 29: 297–300,
1995.

[139456]24900.

[139457]Naso, M. F.; Zimmermann, D. R.; Iozzo, R. V.: Characteri-
zation

[139458]of the complete genomic structure of the human versican
gene and functional

[139459]analysis of its promoter. J. Biol. Chem. 269:
32999–33008, 1994.

[139460]24901.

[139461]Zako, M.; Shinomura, T.; Ujita, M.; Ito, K.; Kimata, K.: Ex-
pression
[139462]of PG-M(V3), an alternatively spliced form of PG-M with-
out a chondroitin
[139463]sulfate attachment region in mouse and human tissues. J.
Biol. Chem. 270:
[139464]3914-3918, 1995.
[139465]24902.

[139466]Zimmermann, D. R.; Ruoslahti, E.: Multiple domains of the
large
[139467]fibroblast proteoglycan, versican. EMBO J. 8: 2975-2981,
1989.
[139468]24903.

[139469]Grunder, S.; Geisler, H.-S.; Rainer, S.; Fink, J. K.: Acid-
sensing
[139470]ion channel (ASIC) 4 gene: physical mapping, genomic or-
ganisation,
[139471]and evaluation as a candidate for paroxysmal dystonia.
Europ. J.
[139472]Hum. Genet. 9: 672-676, 2001.
[139473]24904.

[139474] de Vries, W. N.; de Wet, W. J.: The molecular defect in an autosomal

[139475] dominant form of osteogenesis imperfecta: synthesis of type I procollagen

[139476] containing cysteine in the triple-helical domain of pro-alpha-1(I)

[139477] chains. J. Biol. Chem. 261: 9056-9064, 1986.

[139478] 24905.

[139479] de Vries, W. N.; de Wet, W. J.: Development and diseases of cartilage

[139480] and bone matrix. In: Sen, A.; Thornhill, T.: UCLA Symposia on Molecular

[139481] and Cellular Biology--New Series. New York: Alan R. Liss (pub.)

[139482] 46: 1987.

[139483] 24906.

[139484] Deak, S. B.; Scholz, P. M.; Amenta, P. S.; Constantinou, C. D.;

[139485] Levi-Minzi, S. A.; Gonzalez-Lavin, L.; Mackenzie, J. W.: The substitution

[139486] of arginine for glycine 85 of the alpha-1(I) procollagen chain results

[139487]in mild osteogenesis imperfecta: the mutation provides direct evidence

[139488]for three discrete domains of cooperative melting of intact type I

[139489]collagen. J. Biol. Chem. 266: 21827–21832, 1991.

[139490]24907.

[139491]Diesel, A. J.; Schumacher, A. M.; Flavell, R. A.: A Hind III

[139492]restriction site polymorphism in the human collagen alpha-1(I)-like

[139493]gene on chromosome no. 7. Hum. Genet. 62: 175–176, 1982.

[139494]24908.

[139495]Forlino, A.; Zolezzi, F.; Valli, M.; Pignatti, P. F.; Cetta, G.;

[139496]Brunelli, P. C.; Mottes, M.: Severe (type III) osteogenesis imperfecta

[139497]due to glycine substitutions in the central domain of the collagen

[139498]triple helix. Hum. Molec. Genet. 3: 2201–2206, 1994.

[139499]24909.

[139500]Huerre, C.; Junien, C.; Weil, D.; Chu, M.-L.; Morabito, M.;
Van

- [139501] Cong, N.; Myers, J. C.; Foubert, C.; Gross, M.-S.; Prockop, D. J.;
- [139502] Boue, A.; Kaplan, J.-C.; de la Chapelle, A.; Ramirez, F.: Human type
- [139503] procollagen genes are located on different chromosomes. Proc. Nat.
- [139504] Acad. Sci. 79: 6627-6630, 1982.
- [139505] 24910.
- [139506] Izant, J. G.; Weintraub, H.: Inhibition of thymidine kinase gene
- [139507] expression by anti-sense RNA: a molecular approach to genetic analysis. Cell 36:
- [139508] 1007-1015, 1984.
- [139509] 24911.
- [139510] Khillan, J. S.; Li, S.-W.; Prockop, D. J.: Partial rescue of
- [139511] a lethal phenotype of fragile bones in transgenic mice with a chimeric
- [139512] antisense gene directed against a mutated collagen gene. Proc. Nat.
- [139513] Acad. Sci. 91: 6298-6302, 1994.
- [139514] 24912.

[139515]Korkko, J.; Ala-Kokko, L.; De Paepe, A.; Nuytinck, L.; Earley,

[139516]J.; Prockop, D. J.: Analysis of the COL1A1 and COL1A2 genes by PCR

[139517]amplification and scanning by conformation-sensitive gel electrophoresis

[139518]identifies only COL1A1 mutations in 15 patients with osteogenesis

[139519]imperfecta type I: identification of common sequences of null-allele

[139520]mutations. Am. J. Hum. Genet. 62: 98-110, 1998.

[139521]24913.

[139522]Korkko, J.; Kuivaniemi, H.; Paassilta, P.; Zhuang, J.; Tromp,

[139523]G.; DePaepe, A.; Prockop, D. J.; Ala-Kokko, L.: Two new recurrent

[139524]nucleotide mutations in the COL1A1 gene in four patients with osteogenesis

[139525]imperfecta: about one-fifth are recurrent. Hum. Mutat. 9: 148-156,

[139526]1997.

[139527]24914.

- [139528]Lazarides, E.; Lukens, L. N.: Collagen synthesis on polysomes
- [139529]in vivo and in vitro. Nature N.B. 232: 37–40, 1971.
- [139530]24915.
- [139531]Lund, A. M.; Skovby, F.; Schwartz, M.: (G586V) substitutions
- [139532]in the alpha-1 and alpha-2 chains of collagen I: effect of alpha-chain
- [139533]stoichiometry on the phenotype of osteogenesis imperfecta? Hum. Mutat. 9:
- [139534]431–436, 1997.
- [139535]24916.
- [139536]Mackay, K.; Lund, A. M.; Raghunath, M.; Steinmann, B.; Dagleish,
- [139537]R.: SSCP detection of a gly565-to-val substitution in the pro-alpha-1(I)
- [139538]collagen chain resulting in osteogenesis imperfecta type II. Hum.
- [139539]Genet. 91: 439–444, 1993.
- [139540]24917.
- [139541]Marini, J. C.; Grange, D. K.; Gottesman, G. S.; Lewis, M. B.;

[139542]Koeplin, D. A.: Osteogenesis imperfecta type IV: detection of a point

[139543]mutation in one alpha-1(I) collagen allele (COL1A1) by RNA/RNA hybrid

[139544]analysis. J. Biol. Chem. 264: 11893-11900, 1989.

[139545]24918.

[139546]Marini, J. C.; Lewis, M. B.; Chen, K.: Moderately severe osteogenesis

[139547]imperfecta associated with substitutions of serine for glycine in

[139548]the alpha-1(I) chain of type I collagen. Am. J. Med. Genet. 45:

[139549]241-245, 1993.

[139550]24919.

[139551]Mayer, S. A.; Rubin, B. S.; Starman, B. J.; Byers, P. H.: Spontaneous

[139552]multivessel cervical artery dissection in a patient with a substitution

[139553]of alanine for glycine (G13A) in the alpha-1(I) chain of type I collagen. Neurology 47:

[139554]552-556, 1996.

[139555]24920.

[139556]Mizuno, T.; Chou, M.; Inouye, M.: A unique mechanism
regulating

[139557]gene expression: translational inhibition by a complemen-
tary RNA transcript

[139558](micRNA). Proc. Nat. Acad. Sci. 81: 1966–1970, 1984.

[139559]24921.

[139560]Mottes, M.; Gomez Lira, M. M.; Valli, M.; Scarano, G.;
Lonardo,

[139561]F.; Forlino, A.; Cetta, G.; Pignatti, P. F.: Paternal mosaicism
for

[139562]a COL1A1 dominant mutation (alpha-1 ser-415) causes
recurrent osteogenesis

[139563]imperfecta. Hum. Mutat. 2: 196–204, 1993.

[139564]24922.

[139565]Mottes, M.; Sangalli, A.; Valli, M.; Gomez Lira, M.; Tenni,
R.;

[139566]Buttitta, P.; Pignatti, P. F.; Cetta, G.: Mild dominant osteo-
genesis

[139567]imperfecta with intrafamilial variability: the cause is a ser-
ine for

[139568]glycine alpha-1(I) 901 substitution in a type-I collagen gene. Hum.

[139569]Genet. 89: 480-484, 1992.

[139570]24923.

[139571]Munke, M.; Harbers, K.; Jaenisch, R.; Francke, U.: Chromosomal

[139572]mapping of four different integration sites of Moloney murine leukemia

[139573]virus including the locus for alpha-1(I) collagen in mouse. Cytogenet.

[139574]Cell Genet. 43: 140-149, 1986.

[139575]24924.

[139576]Namikawa, C.; Suzumori, K.; Fukushima, Y.; Sasaki, M.; Hata, A.

[139577]: Recurrence of osteogenesis imperfecta because of paternal mosaicism:

[139578]gly862-to-ser substitution in a type I collagen gene (COL1A1). Hum.

[139579]Genet. 95: 666-670, 1995.

[139580]24925.

[139581]Nicholls, A. C.; Oliver, J.; Renouf, D. V.; Keston, M.; Pope,

[139582]F. M.: Substitution of cysteine for glycine at residue 415 of
one

[139583]allele of the alpha-1(I) chain of type I procollagen in type
III/IV

[139584]osteogenesis imperfecta. J. Med. Genet. 28: 757-764,
1991.

[139585]24926.

[139586]Steinmann, B.; Rao, V. H.; Vogel, A.; Bruckner, P.; Gitzel-
mann,

[139587]R.; Byers, P. H.: Cysteine in the triple-helical domain of
one allelic

[139588]product of the alpha-1(I) gene of type I collagen produces
a lethal

[139589]form of osteogenesis imperfecta. J. Biol. Chem. 250:
11129-11139,

[139590]1984.

[139591]24927.

[139592]Albert, L. J.; Inman, R. D.: Molecular mimicry and autoim-
munity. New

[139593]Eng. J. Med. 341: 2068-2074, 1999.

[139594]24928.

- [139595]Asada, H.; Kawamura, Y.; Maruyama, K.; Kume, H.; Ding, R.-G.; Ji, F. Y.; Kanbara, N.; Kuzume, H.; Sanbo, M.; Yagi, T.; Obata, K.: Mice lacking the 65 kDa isoform of glutamic acid decarboxylase (GAD65) maintain normal levels of GAD67 and GABA in their brains but are susceptible to seizures. *Biochem. Biophys. Res. Commun.* 229: 891–895, 1996.
- [139600]24929.
- [139601]Baekkeskov, S.; Aanstoot, H.-J.; Christgau, S.; Reetz, A.; Solimena, M.; Cascalho, M.; Folli, F.; Richter-Olesen, H.; Camilli, P.-D.: Identification of the 64K autoantigen in insulin-dependent diabetes as the GABA-synthesizing enzyme glutamic acid decarboxylase. *Nature* 347: 151–156, 1990.
- [139605]151–156, 1990.
- [139606]24930.
- [139607]Bu, D.-F.; Erlander, M. G.; Hitz, B. C.; Tillakaratne, N. J. K.;

- [139608]Kaufman, D. L.; Wagner-McPherson, C. B.; Evans, G. A.;
Tobin, A. J.
- [139609]Two human glutamate decarboxylases, 65-kDa GAD and
67-kDa GAD, are
- [139610]each encoded by a single gene. Proc. Nat. Acad. Sci. 89:
2115–2119,
- [139611]1992.
- [139612]24931.
- [139613]Cram, D. S.; Barnett, L. D.; Joseph, J. L.; Harrison, L. C.:
Cloning
- [139614]and partial nucleotide sequence of human glutamic acid
decarboxylase
- [139615]cDNA from brain and pancreatic islets. Biochem. Biophys.
Res. Commun. 176:
- [139616]1239–1244, 1991.
- [139617]24932.
- [139618]De Aizpurua, H. J.; Wilson, Y. M.; Harrison, L. C.: Glutamic
acid

- [139619]decarboxylase autoantibodies in preclinical insulin-dependent diabetes. Proc.
- [139620]Nat. Acad. Sci. 89: 9841-9845, 1992.
- [139621]24933.
- [139622]Edelhoff, S.; Grubin, C. E.; Karlsen, A. E.; Adler, D. A.; Foster,
- [139623]D.; Disteché, C. M.; Lernmark, A.: Mapping of glutamic acid decarboxylase
- [139624](GAD) genes. Genomics 17: 93-97, 1993.
- [139625]24934.
- [139626]Jones, D. B.; Armstrong, N. W.: Cocksackie virus and diabetes revisited.
- [139627](Letter) Nature Med. 1: 284 only, 1995.
- [139628]24935.
- [139629]Karlsen, A. E.; Hagopian, W. A.; Grubin, C. E.; Dube, S.; Disteché,
- [139630]C. M.; Adler, D. A.; Barmeier, H.; Mathewes, S.; Grant, F. J.; Foster,
- [139631]D.; Lernmark, A.: Cloning and primary structure of a human islet
- [139632]isoform of glutamic acid decarboxylase from chromosome

10. Proc.

[139633]Nat. Acad. Sci. 88: 8337–8341, 1991.

[139634]24936.

[139635]Kash, S. F.; Johnson, R. S.; Tecott, L. H.; Noebels, J. L.;
Mayfield,

[139636]R. D.; Hanahan, D.; Baekkeskov, S.: Epilepsy in mice deficient in

[139637]the 65-kDa isoform of glutamic acid decarboxylase. Proc.
Nat. Acad.

[139638]Sci. 94: 14060–14065, 1997.

[139639]24937.

[139640]Kaufman, D. L.; Erlander, M. G.; Clare-Salzler, M.; Atkinson,
son,

[139641]M. A.; Maclaren, N. K.; Tobin, A. J.: Autoimmunity to two
forms of

[139642]glutamate decarboxylase in insulin-dependent diabetes
mellitus. J.

[139643]Clin. Invest. 89: 283–292, 1992.

[139644]24938.

[139645]Lohmann, T.; Hawa, M.; Leslie, R. D. G.; Lane, R.; Picard, J.;

[139646]Londei, M.: Immune reactivity to glutamic acid decarboxy-

lase 65 in

[139647]stiff-man syndrome and type 1 diabetes mellitus. Lancet
356: 31-35,

[139648]2000.

[139649]24939.

[139650]Richter, W.; Endl, J.; Eiermann, T. H.; Brandt, M.; Kientsch-En-
gel,

[139651]R.; Thivolet, C.; Jungfer, H.; Scherbaum, W. A.: Human
monoclonal

[139652]islet cell antibodies from a patient with insulin-dependent
diabetes

[139653]mellitus reveal glutamate decarboxylase as the target
antigen. Proc.

[139654]Nat. Acad. Sci. 89: 8467-8471, 1992.

[139655]24940.

[139656]Tian, J.; Lehmann, P. V.; Kaufman, D. L.: T cell cross-
reactivity

[139657]between coxsackievirus and glutamate decarboxylase is
associated with

[139658]a murine diabetes susceptibility allele. J. Exp. Med. 180:
1979-1984,

[139659]1994.

[139660]24941.

[139661]Chiang, S.-H.; Baumann, C. A.; Kanzaki, M.; Thurmond, D. C.; Watson,

[139662]R. T.; Neudauer, C. L.; Macara, I. G.; Pessin, J. E.; Saltiel, A.

[139663]R.: Insulin-stimulated GLUT4 translocation requires the CAP-dependent

[139664]activation of TC10. *Nature* 410: 944-948, 2001.

[139665]24942.

[139666]Ribon, V.; Printen, J. A.; Hoffman, N. G.; Kay, B. K.; Saltiel,

[139667]A. R.: A novel, multifunctional c-Cbl binding protein in insulin

[139668]receptor signaling in 3T3-L1 adipocytes. *Molec. Cell. Biol.* 18:

[139669]872-879, 1998.

[139670]24943.

[139671]Astrin, K. H.; Arredondo-Vega, F. X.; Desnick, R. J.; Smith, M.

[139672]: Assignment of the gene for cytosolic alanine amino-transferase (AAT1)

[139673]to human chromosome 8. *Ann. Hum. Genet.* 46: 125-133, 1982.

[139674]24944.

[139675]Chen, S.-H.; Giblett, E. R.: Polymorphism of soluble glutamic-pyruvic

[139676]transaminase: a new genetic marker in man. Science 173: 148-149,

[139677]1971.

[139678]24945.

[139679]Cook, P. J. L.; Jeremiah, S. J.; Buckton, K. E.: Exclusion mapping

[139680]of GPT. (Abstract) Cytogenet. Cell Genet. 32: 258 only, 1982.

[139681]24946.

[139682]Mombaerts, P.; Wang, F.; Dulac, C.; Vassar, R.; Chao, S. K.; Nemes,

[139683]A.; Mendelsohn, M.; Edmondson, J.; Axel, R.: The molecular biology

[139684]of olfactory perception. Cold Spring Harbor Symp. Quant. Biol. 61:

[139685]135-145, 1996.

[139686]24947.

[139687]Reed, R. R.: Genetic approaches to mammalian olfaction.

Cold

[139688]Spring Harbor Symp. Quant. Biol. 61: 165–172, 1996.

[139689]24948.

[139690]Ronnett, G. V.; Snyder, S. H.: Molecular messengers of olfaction. Trends

[139691]Neurosci. 15: 508–513, 1992.

[139692]24949.

[139693]Schwab, S. G.; Hallmayer, J.; Lerer, B.; Albus, M.; Bornmann, M.;

[139694]Honig, S.; Strauss, M.; Segman, R.; Lichtermann, D.; Knapp, M.; Trixler,

[139695]M.; Maier, W.; Wildenauer, D. B.: Support for a chromosome 18p locus

[139696]conferring susceptibility to functional psychoses in families with

[139697]schizophrenia, by association and linkage analysis. Am. J. Hum. Genet. 63:

[139698]1139–1152, 1998.

[139699]24950.

[139700]Davignon, I.; Barnard, M.; Gavrilova, O.; Sweet, K.; Wilkie, T.

[139701]M.: Gene structure of murine Gna11 and Gna15: tandemly duplicated

[139702]Gq class G protein alpha subunit genes. Genomics 31: 359–366, 1996.

[139703]24951.

[139704]Jiang, M.; Pandey, S.; Tran, V. T.; Fong, H. K.: Guanine nucleotide-binding

[139705]regulatory proteins in retinal pigment epithelial cells. Proc. Nat.

[139706]Acad. Sci. 88: 3907–3911, 1991.

[139707]24952.

[139708]Offermanns, S.; Zhao, L.-P.; Gohla, A.; Sarosi, I.; Simon, M. I.;

[139709]Wilkie, T. M.: Embryonic cardiomyocyte hypoplasia and craniofacial

[139710]defects in G-alpha-q/G-alpha-11-mutant mice. EMBO J. 17: 4304–4312,

[139711]1998.

[139712]24953.

[139713]Strathmann, M. P.; Simon, M. I.: G-alpha-12 and G-alpha-13 subunits

[139714]define a fourth class of G protein alpha subunits. Proc.
Nat. Acad.

[139715]Sci. 88: 5582–5586, 1991.

[139716]24954.

[139717]Ahmed, S. F.; Barr, D. G. D.; Bonthron, D. T.: GNAS1 muta-
tions

[139718]and progressive osseous heteroplasia. (Letter) New Eng. J.
Med. 346:

[139719]1669–1670, 2002.

[139720]24955.

[139721]Ahmed, S. F.; Dixon, P. H.; Bonthron, D. T.; Stirling, H. F.;
Barr,

[139722]D. G. D.; Kelnar, C. J. H.; Thakker, R. V.: GNAS1 mutational
analysis

[139723]in pseudohypoparathyroidism. Clin. Endocr. 49: 525–531,
1998.

[139724]24956.

[139725]Ahrens, W.; Hiort, O.; Staedt, P.; Kirschner, T.; Marschke,
C.;

[139726]Kruse, K.: Analysis of the GNAS1 gene in Albright's hered-
itary osteodystrophy. J.

[139727]Clin. Endocr. Metab. 86: 4630–4634, 2001.

[139728]24957.

[139729]Aldred, M. A.; Trembath, R. C.: Activating and inactivating mutations

[139730]in the human GNAS1 gene. Hum. Mutat. 16: 183–189, 2000.

[139731]24958.

[139732]Ashley, P. L.; Ellison, J.; Sullivan, K. A.; Bourne, H. R.; Cox,

[139733]D. R.: Chromosomal assignment of the murine Gi and Gs genes. (Abstract) Am.

[139734]J. Hum. Genet. 41: A155 only, 1987.

[139735]24959.

[139736]Ballare, E.; Mantovani, S.; Lania, A.; Di Blasio, A. M.; Vallar,

[139737]L.; Spada, A.: Activating mutations of the GS- α gene are associated

[139738]with low levels of GS- α protein in growth hormone-secreting tumors. J.

[139739]Clin. Endocr. Metab. 83: 4386–4390, 1998.

[139740]24960.

[139741]Bastepe, M.; Pincus, J. E.; Sugimoto, T.; Tojo, K. Kanatani,

M.;

[139742]Azuma, Y.; Kruse, K.; Rosenbloom, A. L.; Koshiyama, H.; Juppner, H.

[139743]; Positional dissociation between the genetic mutation responsible

[139744]for pseudohypoparathyroidism type Ib and the associated methylation

[139745]defect at exon A/B: evidence for a long-range regulatory element within

[139746]the imprinted GNAS1 locus. Hum. Molec. Genet. 10: 1231-1241, 2001.

[139747]24961.

[139748]Bastepe, M.; Juppner, H.: GNAS1 mutations and progressive osseous

[139749]heteroplasia. (Letter) New Eng. J. Med. 346: 1671 only, 2002.

[139750]24962.

[139751]Bastepe, M.; Lane, A. H.; Juppner, H.: Parental uniparental isodisomy

[139752]of chromosome 20q--and the resulting changes in GNAS1 methylation--as

[139753]a plausible cause of pseudohypoparathyroidism. Am. J.

Hum. Genet. 68:

[139754]1283–1289, 2001.

[139755]24963.

[139756]Bianco, P.; Riminucci, M.; Majolagbe, A.; Kuznetsov, S. A.;

Collins,

[139757]M. T.; Mankani, M. H.; Corsi, A.; Bone, H. G.; Wientroub,

S.; Spiegel,

[139758]A. M.; Fisher, L. W.; Robey, P. G.: Mutations of the GNAS1

gene,

[139759]stromal cell dysfunction, and osteomalacic changes in

non-McCune-Albright

[139760]fibrous dysplasia of bone. J. Bone Miner. Res. 15:

120–128, 2000.

[139761]24964.

[139762]Billestrup, N.; Swanson, L. W.; Vale, W.: Growth hormone–

releasing

[139763]factor stimulates proliferation of somatotrophs in vitro.

Proc. Nat.

[139764]Acad. Sci. 83: 6854–6857, 1986.

[139765]24965.

[139766]Candelieri, G. A.; Glorieux, F. H.; Prud'Homme, J.;

St.-Arnaud,

[139767]R.: Increased expression of the c-fos proto-oncogene in bone from

[139768]patients with fibrous dysplasia. New Eng. J. Med. 332: 1546-1551,

[139769]1995.

[139770]24966.

[139771]Candelieri, G. A.; Roughley, P. J.; Glorieux, F. H.: Polymerase

[139772]chain reaction-based technique for the selective enrichment and analysis

[139773]of mosaic arg201 mutations in G alpha s from patients with fibrous

[139774]dysplasia of bone. Bone 21: 201-206, 1997.

[139775]24967.

[139776]Carel, J. C.; Le Stunff, C.; Condamine, L.; Mallet, E.; Chaus-

[139777]J. L.; Adnot, P.; Garabedian, M.; Bougneres, P.: Resistance to the

[139778]lipolytic action of epinephrine: a new feature of protein GS deficiency. J.

[139779]Clin. Endocr. Metab. 84: 4127-4131, 1999.

[139780]24968.

[139781]Carter, A.; Bardin, C.; Collins, R.; Simons, C.; Bray, P.;
Spiegel,

[139782]A.: Reduced expression of multiple forms of the alpha
subunit of

[139783]the stimulatory GTP-binding protein in pseudohy-
poparathyroidism type

[139784]1a. Proc. Nat. Acad. Sci. 84: 7266-7269, 1987.

[139785]24969.

[139786]Cattanach, B. M.; Kirk, M.: Differential activity of mater-
nally

[139787]and paternally derived chromosome regions in mice. Na-
ture 315: 496-498,

[139788]1985.

[139789]24970.

[139790]Coutant, R.; Lumbroso, S.; Rey, R.; Lahlou, N.; Venara, M.;
Rouleau,

[139791]S.; Sultan, C.; Limal, J.-M.: Macroorchidism due to au-
tonomous hyperfunction

[139792]of Sertoli cells and GS-alpha gene mutation: an unusual
expression

[139793]of McCune–Albright syndrome in a prepubertal boy. J.
Clin. Endocr.

[139794]Metab. 86: 1778–1781, 2001.

[139795]24971.

[139796]Kim, H.–S.; Nagalla, S. R.; Oh, Y.; Wilson, E.; Roberts, C. T.,
[139797]Jr.; Rosenfeld, R. G.: Identification of a family of low–
affinity

[139798]insulin–like growth factor binding proteins (IGFBPs): char–
acterization

[139799]of connective tissue growth factor as a member of the
IGFBP superfamily. Proc.

[139800]Nat. Acad. Sci. 94: 12981–12986, 1997.

[139801]24972.

[139802]Martinerie, C.; Viegas–Pequignot, E.; Guenard, I.; Dutril–
laux,

[139803]B.; Nguyen, V. C.; Bernheim, A.; Perbal, B.: Physical map–
ping of

[139804]human loci homologous to the chicken nov proto–
oncogene. Oncogene 7:

[139805]2529–2534, 1992.

[139806]24973.

- [139807]Nakanishi, T.; Yamaai, T.; Asano, M.; Nawachi, K.; Suzuki, M.;
- [139808]Sugimoto, T.; Takigawa, M.: Overexpression of connective tissue growth
- [139809]factor/hypertrophic chondrocyte-specific gene product 24 decreases
- [139810]bone density in adult mice and induces dwarfism. Biochem. Biophys. Res. Commun. 281: 678-681, 2001.
- [139812]24974.
- [139813]O'Donovan, N.; Galvin, M.; Morgan, J. G.: Physical mapping of
- [139814]the CXC chemokine locus on human chromosome 4. Cytogenet. Cell Genet. 84:
- [139815]39-42, 1999.
- [139816]24975.
- [139817]McGill, J. R.; Boyd, D.; Barrett, K. J.; Drysdale, J. W.; Moore, C. M.: Localization of human ferritin H (heavy) and L (light) subunits
- [139819]by in situ hybridization. (Abstract) Am. J. Hum. Genet. 36: 146S

[139820]only, 1984.

[139821]24976.

[139822]Murray, M. T.; White, K.; Munro, H. N.: Conservation of
ferritin

[139823]heavy subunit gene structure: implications for the regula-
tion of ferritin

[139824]gene expression. Proc. Nat. Acad. Sci. 84: 7438-7442,
1987.

[139825]24977.

[139826]Papadopoulos, P.; Bhavsar, D.; Zappone, E.; David, V.;
Jones,

[139827]C.; Worwood, M.; Drysdale, J.: A second human ferritin H
locus on

[139828]chromosome 11. Cytogenet. Cell Genet. 61: 107-108,
1992.

[139829]24978.

[139830]Richard, C. W.; Withers, D. A.; Meeker, T. C.; Myers, R. M.:

[139831]A radiation hybrid map of the proximal long arm of hu-
man chromosome

[139832]11 containing the MEN-1 and bcl-1 disease locus.
(Abstract) Cytogenet.

[139833]Cell Genet. 58: 1970 only, 1991.

[139834]24979.

[139835]Worwood, M.; Brook, J. D.; Cragg, S. J.; Hellkuhl, B.; Jones,

[139836]B. M.; Perera, P.; Roberts, S. H.; Shaw, D. J.: Assignment of
human

[139837]ferritin genes to chromosomes 11 and 19q13.3-19qter.

Hum. Genet. 69:

[139838]371-374, 1985.

[139839]24980.

[139840]Wu, K.-J.; Polack, A.; Dalla-Favera, R.: Coordinated regu-
lation

[139841]of iron-controlling genes, H-ferritin and IRP2, by c-MYC.
Science 283:

[139842]676-679, 1999.

[139843]24981.

[139844]Yachou, A.; Mattei, M. G.; Roeckel, N.; Grandchamp, B.;
Beaumont,

[139845]C.: Mouse ferritin H sequences map to chromosomes 3, 6,
and 19. Genomics 9:

[139846]204-206, 1991.

[139847]24982.

[139848]Yachou, A.-K.; Renaudie, F.; Guenet, J.-L.; Simon-
Chazottes, D.;

[139849]Jones, R.; Grandchamp, B.; Beaumont, C.: Mouse ferritin H
multigene

[139850]family is polymorphic and contains a single multiallelic
functional

[139851]gene located on chromosome 19. Genomics 10: 531-538,
1991.

[139852]24983.

[139853]Youssofian, H.; Chance, P.; Tuck-Muller, C. M.; Jabs, E.
W.:

[139854]Association of a new chromosomal deletion
[del(1)(q32q42)] with diaphragmatic

[139855]hernia: assignment of a human ferritin gene. Hum. Genet.
78: 267-270,

[139856]1988.

[139857]24984.

[139858]Takeda, J.; Seino, Y.; Tanaka, K.-I.; Fukumoto, H.; Kayano,
T.;

[139859]Takahashi, H.; Mitani, T.; Kurono, M.; Suzuki, T.; Tobe, T.;
Imura,

- [139860]H.: Sequence of an intestinal cDNA encoding human gastric inhibitory polypeptide precursor. Proc. Nat. Acad. Sci. 84: 7005–7008, 1987.
- [139862]24985.
- [139863]Xu, W.; Gorman, P. A.; Rider, S. H.; Hedge, P. J.; Moore, G.; Prichard, C.; Sheer, D.; Solomon, E.: Construction of a genetic map of human chromosome 17 by use of chromosome-mediated gene transfer. Proc. Nat. Acad. Sci. 85: 8563–8567, 1988.
- [139867]24986.
- [139868]Fishman, G. I.; Eddy, R. L.; Shows, T. B.; Rosenthal, L.; Leinwand, L. A.: The human connexin gene family of gap junction proteins: distinct chromosomal locations but similar structures. Genomics 10: 250–256, 1991.
- [139872]24987.
- [139873]Campbell, H. D.; Webb, G. C.; Kono, T.; Taniguchi, T.;

Ford, J.

[139874]H.; Young, I. G.: Assignment of the interleukin-2 receptor
beta chain

[139875]gene (IL-2rb) to band E on mouse chromosome 15. Ge-
nomics 12: 179-180,

[139876]1992.

[139877]24988.

[139878]Burgeson, R. E.; Chiquet, M.; Deutzmann, R.; Ekblom, P.;
Engel,

[139879]J.; Kleinman, H.; Martin, G. R.; Meneguzzi, G.; Paulsson,
M.; Sanes,

[139880]J.; Timpl, R.; Tryggvason, K.; Yamada, Y.; Yurchenco, P. D.:
A new

[139881]nomenclature for the laminins. Matrix Biol. 14: 209-211,
1994.

[139882]24989.

[139883]Gremlich, S.; Porret, A.; Hani, E. H.; Cherif, D.; Vionnet, N.;

[139884]Froguel, P.; Thorens, B.: Cloning, functional expression,
and chromosomal

[139885]localization of the human pancreatic islet glucose-de-
pendent insulinotropic

[139886]polypeptide receptor. Diabetes 44: 1202-1208, 1995.

[139887]24990.

[139888]Miyawaki, K.; Yamada, Y.; Ban, N.; Ihara, Y.; Tsukiyama, K.;
Zhou,

[139889]H.; Fujimoto, S.; Oku, A.; Tsuda, K.; Toyokuni, S.; Hiai, H.;
Mizunoya,

[139890]W.; and 9 others: Inhibition of gastric inhibitory polypep-
tide

[139891]signaling prevents obesity. Nature Med. 8: 738-742,
2002.

[139892]24991.

[139893]Stoffel, M.; Fernald, A. A.; Le Beau, M. M.; Bell, G. I.: As-
signment

[139894]of the gastric inhibitory polypeptide receptor gene (GIPR)
to chromosome

[139895]bands 19q13.2-q13.3 by fluorescence in situ hybridiza-
tion. Genomics 28:

[139896]607-609, 1995.

[139897]24992.

[139898]Usdin, T. B.; Mezey, E.; Button, D. C.; Brownstein, M. J.;
Bonner,

[139899]T. I.: Gastric inhibitory polypeptide receptor, a member of
the secretin-vasoactive

[139900]intestinal peptide receptor family, is widely distributed in
peripheral

[139901]organs and the brain. *Endocrinology* 133: 2861–2870,
1993.

[139902]24993.

[139903]Volz, A.; Groke, R.; Lankat-Buttgereit, B.; Fehmann, H.-C.;
Bode,

[139904]H. P.; Goke, B.: Molecular cloning, functional expression,
and signal

[139905]transduction of the GIP-receptor cloned from a human in-
sulinoma. *FEBS*

[139906]*Lett.* 373: 23–29, 1995.

[139907]24994.

[139908]Yamada, Y.; Hayami, T.; Nakamura, K.; Kaisaki, P. J.;
Someya, Y.;

[139909]Wang, C.-Z.; Seino, S.; Seino, Y.: Human gastric inhibitory
polypeptide

[139910]receptor: cloning of the gene (GIPR) and cDNA. *Genomics*
29: 773–776,

[139911]1995.

[139912]24995.

- [139913] Miyawaki, K.; Yamada, Y.; Yano, H.; Niwa, H.; Ban, N.; Ihara, Y.;
- [139914] Kubota, A.; Fujimoto, S.; Kajikawa, M.; Kuroe, A.; Tsuda, K.; Hashimoto,
- [139915] H.; Yamashita, T.; Jomori, T.; Tashiro, F.; Miyazaki, J.; Seino, Y.
- [139916] Glucose intolerance caused by a defect in the entero-insular axis:
- [139917] a study in gastric inhibitory polypeptide receptor knock-out mice. *Proc.*
- [139918] *Nat. Acad. Sci.* 96: 14843–14847, 1999.
- [139919] 24996.
- [139920] Bentley, P. H.; Kenner, G. W.; Sheppard, R. C.: Structure of human
- [139921] gastrins I and II. *Nature* 209: 583–585, 1966.
- [139922] 24997.
- [139923] Boel, E.; Vuust, J.; Norris, F.; Norris, K.; Wind, A.; Rehfeld,
- [139924] J. F.; Marcker, K. A.: Molecular cloning of human gastrin cDNA: evidence
- [139925] for evolution of gastrin by gene duplication. *Proc. Nat. Acad. Sci.* 80:

[139926]2866–2869, 1983.

[139927]24998.

[139928]Flejter, W. L.; Barcroft, C. L.; Guo, S.–W.; Lynch, E. D.;
Boehnke,

[139929]M.; Chandrasekharappa, S.; Hayes, S.; Collins, F. S.; Weber,
B. L.;

[139930]Glover, T. W.: Multicolor FISH mapping with Alu–
PCR–amplified YAC

[139931]clone DNA determines the order of markers in the BRCA1
region on chromosome

[139932]17q12–q21. Genomics 17: 624–631, 1993.

[139933]24999.

[139934]Flejter, W. L.; Kukowska–Latallo, J. F.; Kioussis, S.; Chan–
drasekharappa,

[139935]S. C.; King, S. E.; Chamberlain, J. S.: Tetranucleotide repeat
polymorphism

[139936]at D17S846 maps within 40 kb of GAS at 17q12–q22.
Hum. Molec. Genet. 2:

[139937]1080 only, 1993.

[139938]25000.

[139939]Friis–Hansen, L.; Rourke, I. J.; Bundgaard, J. R.; Rehfeld, J.

- [139940]F.; Samuelson, L. C.: Molecular structure and genetic mapping of
[139941]the mouse gastrin gene. FEBS Lett. 386: 128–132, 1996.
[139942]25001.
- [139943]Fukushige, S.; Murotsu, T.; Matsubara, K.: Chromosomal assignment
[139944]of human genes for gastrin, thyrotropin (TSH)–beta sub-unit and C-erb-2
[139945]by chromosome sorting combined with velocity sedimentation and southern
[139946]hybridization. Biochem. Biophys. Res. Commun. 134: 477–483, 1986.
[139947]25002.
- [139948]Gregory, R. A.; Tracy, H. J.; Agarwal, K. L.; Grossman, M. I.:
[139949]Amino acid constitution of two gastrins isolated from Zollinger–Ellison
[139950]tumor tissue. Gut 10: 603–608, 1969.
[139951]25003.
- [139952]Ito, R.; Sato, K.; Helmer, T.; Jay, G.; Agarwal, K.: Structural
[139953]analysis of the gene encoding human gastrin: the large intron contains

[139954]an Alu sequence. Proc. Nat. Acad. Sci. 81: 4662–4666,
1984.

[139955]25004.

[139956]Lund, T.; Geurts van Kessel, A. H. M.; Huan, S.; Dixon, J. E.

[139957]: The genes for human gastrin and cholecystokinin are located on different

[139958]chromosomes. Hum. Genet. 73: 77–80, 1986.

[139959]25005.

[139960]Lund, T.; Geurts van Kessel, A. H. M.; Westerveld, A.: The human

[139961]gastrin gene is located at human chromosome 17.
(Abstract) Cytogenet.

[139962]Cell Genet. 40: 683 only, 1985.

[139963]25006.

[139964]Wiborg, O.; Berglund, L.; Boel, E.; Norris, F.; Norris, K.; Rehfeld,

[139965]J. F.; Marcker, K. A.; Vuust, J.: Structure of a human gastrin gene. Proc.

[139966]Nat. Acad. Sci. 81: 1067–1069, 1984.

[139967]25007.

- [139968]Justice, M. J.; Gilbert, D. J.; Kinzler, K. W.; Vogelstein, B.;
[139969]Buchberg, A. M.; Ceci, J. D.; Matsuda, Y.; Chapman, V. M.;
Patriotis,
[139970]C.; Makris, A.; Tsiichlis, P. N.; Jenkins, N. A.; Copeland, N.
G.:
[139971]A molecular genetic linkage map of mouse chromosome
18 reveals extensive
[139972]linkage conservation with human chromosomes 5 and 18.
Genomics 13:
[139973]1281–1288, 1992.
[139974]25008.
[139975]Cox, D. W.; Francke, U.: Direct assignment of orosomu-
coid to human
[139976]chromosome 9 and alpha-2-HS-glycoprotein to chromo-
some 3 using human
[139977]fetal liver x rat hepatoma hybrids. Hum. Genet. 70:
109–115, 1985.
[139978]25009.
[139979]Umetsu, K.; Yuasa, I.; Nishimura, H.; Sasaki, H.; Suzuki, T.:
[139980]Genetic polymorphisms of orosomucoid and alpha-
2-HS-glycoprotein in

[139981]a Philippine population. Hum. Hered. 38: 287–290, 1988.

[139982]25010.

[139983]Triantafilou, K.; Triantafilou, M.; Dedrick, R. L.: A
CD14-independent

[139984]LPS receptor cluster. Nature Immun. 2: 338–345, 2001.

[139985]25011.

[139986]Klinger, H. P.: Suppression of tumorigenicity in somatic
cell

[139987]hybrids. I. Suppression and reexpression of tumorigenicity
in diploid

[139988]human x D98/AH2 hybrids and independent segregation
of tumorigenicity

[139989]from other cell phenotypes. Cytogenet. Cell Genet. 27:
254–266,

[139990]1980.

[139991]25012.

[139992]Lichy, J. H.; Modi, W. S.; Seuanetz, H. N.; Howley, P. M.:
Identification

[139993]of a human chromosome 11 gene which is differentially
regulated in

[139994]tumorigenic and nontumorigenic somatic cell hybrids of

HeLa cells. Cell

[139995]Growth Differ. 3: 541–548, 1992.

[139996]25013.

[139997]Stanbridge, E. J.: Suppression of malignancy in human cells. Nature 260:

[139998]17–20, 1976.

[139999]25014.

[140000]Corrigall, V. M.; Arastu, M.; Khan, S.; Shah, C.; Fife, M.; Smeets,

[140001]T.; Tak, P.–P.; Panayi, G. S.: Functional IL–2 receptor beta (CD122)

[140002]and gamma (CD132) chains are expressed by fibroblast-like synoviocytes:

[140003]activation by IL–2 stimulates monocyte chemoattractant protein–1 production. J.

[140004]Immun. 166: 4141–4147, 2001.

[140005]25015.

[140006]Gnarra, J. R.; Otani, H.; Wang, M. G.; McBride, O. W.; Sharon,

[140007]M.; Leonard, W. J.: Human interleukin 2 receptor beta-chain gene:

[140008]chromosomal localization and identification of 5-prime regulatory

[140009]sequences. Proc. Nat. Acad. Sci. 87: 3440–3444, 1990.

[140010]25016.

[140011]Hatakeyama, M.; Tsudo, M.; Minamoto, S.; Kono, T.; Doi, T.; Miyata,

[140012]T.; Miyasaka, M.; Taniguchi, T.: Interleukin-2 receptor beta chain

[140013]gene: generation of three receptor forms by cloned human alpha and

[140014]beta chain cDNA's. Science 244: 551–556, 1989.

[140015]25017.

[140016]Lamaze, C.; Dujeancourt, A.; Baba, T.; Lo, C. G.; Benmerah, A.;

[140017]Dautry-Varsat, A.: Interleukin 2 receptors and detergent-resistant

[140018]membrane domains define a clathrin-independent endocytic pathway. Molec.

[140019]Cell 7: 661–671, 2001.

[140020]25018.

[140021]Shibuya, H.; Yoneyama, M.; Nakamura, Y.; Harada, H.;

Hatakeyama,

[140022]M.; Minamoto, S.; Kono, T.; Doi, T.; White, R.; Taniguchi, T.: The

[140023]human interleukin-2 receptor beta-chain gene: genomic organization,

[140024]promoter analysis and chromosomal assignment. Nucleic Acids Res. 18:

[140025]3697-3703, 1990.

[140026]25019.

[140027]Suzuki, H.; Kundig, T. M.; Furlonger, C.; Wakeham, A.; Timms, E.;

[140028]Matsuyama, T.; Schmits, R.; Simard, J. J. L.; Ohashi, P. S.; Griesser,

[140029]H.; Taniguchi, T.; Paige, C. J.; Mak, T. W.: Deregulated T cell activation

[140030]and autoimmunity in mice lacking interleukin-2 receptor beta. Science 268:

[140031]1472-1476, 1995.

[140032]25020.

[140033]Tsuda, M.; Kitamura, F.; Miyasaka, M.: Characterization of the

[140034]interleukin 2 receptor beta chain using three distinct

monoclonal

[140035]antibodies. Proc. Nat. Acad. Sci. 86: 1982–1986, 1989.

[140036]25021.

[140037]Ballard, F. J.; Baxter, R. C.; Binoux, M.; Clemmons, D. R.;
Drop,

[140038]S. L. S.; Hall, K.; Hintz, R. L.; Rechler, M. M.; Rutanen, E.
M.;

[140039]Schwander, J. C.: Report on the nomenclature of the IGF
binding proteins. J.

[140040]Clin. Endocr. Metab. 70: 817–818, 1990.

[140041]25022.

[140042]Popovici, R. M.; Lu, M.; Bhatia, S.; Faessen, G. H.; Giaccia,
A.

[140043]J.; Giudice, L. C.: Hypoxia regulates insulin-like growth
factor-binding

[140044]protein 1 in human fetal hepatocytes in primary culture:
suggestive

[140045]molecular mechanisms for in utero fetal growth restriction
caused

[140046]by uteroplacental insufficiency. J. Clin. Endocr. Metab. 86:
2653–2659,

[140047]2001.

[140048]25023.

[140049]Allander, S. V.; Larsson, C.; Ehrenborg, E.; Suwanichkul,
A.; Weber,

[140050]G.; Morris, S. L.; Bajalica, S.; Kiefer, M. C.; Luthman, H.;
Powell,

[140051]D. R.: Characterization of the chromosomal gene and pro-
moter for

[140052]human insulin-like growth factor binding protein-5. J.
Biol. Chem. 269:

[140053]10891-10898, 1994.

[140054]25024.

[140055]Bauer, M.; Dieterich, W.; Ehnis, T.; Schuppan, D.: Complete
primary

[140056]structure of human collagen type XIV (undulin). Biochim.
Biophys.

[140057]Acta 1354: 183-188, 1997.

[140058]25025.

[140059]Dublet, B.; van der Rest, M.: Type XIV collagen, a new ho-
motrimeric

[140060]molecule extracted from fetal bovine skin and tendon,
with a triple

[140061]helical disulfide-bonded domain homologous to type IX
and type XII

[140062]collagens. J. Biol. Chem. 266: 6853-6858, 1991.

[140063]25026.

[140064]Just, M.; Herbst, H.; Hummel, M.; Durkop, H.; Tripier, D.;
Stein,

[140065]H.; Schuppan, D.: Undulin is a novel member of the fi-
bronectin-tenascin

[140066]family of extracellular matrix glycoproteins. J. Biol. Chem.
266:

[140067]17326-17332, 1991.

[140068]25027.

[140069]Schnittger, S.; Herbst, H.; Schuppan, D.; Dannenberg, C.;
Bauer,

[140070]M.; Fonatsch, C.: Localization of the undulin gene (UND) to
human

[140071]chromosome band 8q23. Cytogenet. Cell Genet. 68:
233-234, 1995.

[140072]25028.

[140073]Schuppan, D.; Cantaluppi, M. C.; Becker, J.; Veit, A.; Bunte,
T.;

- [140074]Troyer, D.; Schuppan, F.; Schmid, M.; Ackermann, R.;
Hahn, E. G.:
- [140075]Undulin, an extracellular matrix glycoprotein associated
with collagen
- [140076]fibrils. J. Biol. Chem. 265: 8823–8832, 1990.
- [140077]25029.
- [140078]Eklund, L.; Piuholo, J.; Komulainen, J.; Sormunen, R.; Ong-
varrasopone,
- [140079]C.; Fassler, R.; Muona, A.; Ilves, M.; Ruskoaho, H.; Takala,
T. E.
- [140080]S.; Pihlajaniemi, T.: Lack of type XV collagen causes a
skeletal
- [140081]myopathy and cardiovascular defects in mice. Proc. Nat.
Acad. Sci. 98:
- [140082]1194–1199, 2001.
- [140083]25030.
- [140084]Hagg, P. M.; Hagg, P. O.; Peltonen, S.; Autio-Harminen,
H.; Pihlajaniemi,
- [140085]T.: Location of type XV collagen in human tissues and its
accumulation
- [140086]in the interstitial matrix of the fibrotic kidney. Am. J. Path.
150:

[140087]2075–2086, 1997.

[140088]25031.

[140089]Hagg, P. M.; Horelli-Kuitunen, N.; Eklund, L.; Palotie, A.;
Pihlajaniemi,

[140090]T.: Cloning of mouse type XV collagen sequences and
mapping of the

[140091]corresponding gene to 4B1–3: comparison of mouse and
human alpha-1(XV)

[140092]collagen sequences indicates divergence in the number of
small collagenous

[140093]domains. Genomics 45: 31–41, 1997.

[140094]25032.

[140095]Hagg, P. M.; Muona, A.; Lietard, J.; Kivirikko, S.; Pihla-
janiemi,

[140096]T.: Complete exon–intron organization of the human gene
for the alpha-1

[140097]chain of type XV collagen (COL15A1) and comparison with
the homologous

[140098]Col18a1 gene. J. Biol. Chem. 273: 17824–17831, 1998.

[140099]25033.

[140100]Huebner, K.; Cannizzaro, L. A.; Jabs, E. W.; Kivirikko, S.;

Manzone,

[140101]H.; Pihlajaniemi, T.; Myers, J. C.: Chromosomal assignment
of a gene

[140102]encoding a new collagen type (COL15A1) to 9q21-q22.

Genomics 14:

[140103]220-224, 1992.

[140104]25034.

[140105]Kivirikko, S.; Heinamaki, P.; Rehn, M.; Honkanen, N.; My-
ers, J.

[140106]C.; Pihlajaniemi, T.: Primary structure of the alpha-1 chain
of human

[140107]type XV collagen and exon-intron organization in the
3-prime region

[140108]of the corresponding gene. J. Biol. Chem. 269:
4773-4779, 1994.

[140109]25035.

[140110]Muragaki, Y.; Abe, N.; Ninomiya, Y.; Olsen, B. R.; Ooshima,
A.

[140111]: The human alpha-1(XV) collagen chain contains a large
amino-terminal

[140112]non-triple helical domain with a tandem repeat structure
and homology

[140113]to alpha-1(XVIII) collagen. J. Biol. Chem. 269: 4042-4046, 1994.

[140114]25036.

[140115]Myers, J. C.; Dion, A. S.; Abraham, V.; Amenta, P. S.: Type XV

[140116]collagen exhibits a widespread distribution in human tissues but a

[140117]distinct localization in basement membrane zones. Cell Tissue Res. 286:

[140118]493-505, 1996.

[140119]25037.

[140120]Myers, J. C.; Kivirikko, S.; Gordon, M. K.; Dion, A. S.; Pihlajaniemi,

[140121]T.: Identification of a previously unknown human collagen chain,

[140122]alpha-1(XV), characterized by extensive interruptions in the triple-helical

[140123]region. Proc. Nat. Acad. Sci. 89: 10144-10148, 1992.

[140124]25038.

[140125]Ramchandran, R.; Dhanabal, M.; Volk, R.; Waterman, M. J. F.; Segal,

[140126]M.; Lu, H.; Knebelmann, B.; Sukhatme, V. P.: Antiangiogenic activity

[140127]of restin, NC10 domain of human collagen XV: comparison to endostatin. Biochem.

[140128]Biophys. Res. Commun. 255: 735–739, 1999.

[140129]25039.

[140130]Rehn, M.; Hintikka, E.; Pihlajaniemi, T.: Primary structure of

[140131]the alpha 1 chain of mouse type XVIII collagen, partial structure

[140132]of the corresponding gene, and comparison of the alpha 1(XVIII) chain

[140133]with its homologue, the alpha 1(XV) collagen chain. J. Biol. Chem. 269:

[140134]13929–13935, 1994.

[140135]25040.

[140136]Sasaki, T.; Larsson, H.; Tisi, D.; Claesson-Welsh, L.; Hohenester,

[140137]E.; Timpl, R.: Endostatins derived from collagens XV and XVIII differ

[140138]in structural and binding properties, tissue distribution and anti-angiogenic

[140139]activity. J. Molec. Biol. 301: 1179–1190, 2000.

[140140]25041.

[140141]Pan, T.–C.; Zhang, R.–Z.; Mattei, M.–G.; Timpl, R.; Chu, M.–L.

[140142]: Cloning and chromosomal location of human alpha-1(XVI) collagen.

[140143]Proc. Nat. Acad. Sci. 89: 6565–6569, 1992.

[140144]25042.

[140145]Yamaguchi, N.; Kimura, S.; McBride, O. W.; Hori, H.; Yamada, Y.;

[140146]Kanamori, T.; Yamakoshi, H.; Nagai, Y.: Molecular cloning and partial

[140147]characterization of a novel collagen chain, alpha-1(XVI), consisting

[140148]of repetitive collagenous domains and cysteine-containing non-collagenous

[140149]segments. J. Biochem. 112: 856–863, 1992.

[140150]25043.

[140151]O'Reilly, M. S.; Boehm, T.; Shing, Y.; Fukai, N.; Vasios, G.; Lane,

[140152]W. S.; Flynn, E.; Birkhead, J. R.; Olsen, B. R.; Folkman, J.:

Endostatin:

- [140153]an endogenous inhibitor of angiogenesis and tumor growth. Cell 88: 277–285, 1997.
- [140155]25044.
- [140156]Sakai, L. Y.; Keene, D. R.; Glanville, R. W.; Bachinger, H. P.
- [140157]: Purification and partial characterization of fibrillin, a cysteine-rich structural component of connective tissue microfibrils. J. Biol. Chem. 266: 14763–14770, 1991.
- [140160]25045.
- [140161]Eiberg, H.; Bisgaard, M. L.; Mohr, J.: Linkage between alpha-1-B-glycoprotein (A1BG) and Lutheran (LU) red blood group system: assignment to chromosome 19: new genetic variants of A1BG. Clin. Genet. 36: 415–418, 1989.
- [140164]25046.
- [140165]Eiberg, H.; Nielsen, L. S.; Gahne, B.; Juneja, R. K.; Mohr, J.
- [140166]: Exclusion data for the alpha-1-B glycoprotein (A1BG)

polymorphism.

[140167](Abstract) Cytogenet. Cell Genet. 51: 994 only, 1989.

[140168]25047.

[140169]Gahne, B.; Juneja, R. K.; Stratil, A.: Genetic polymorphism
of

[140170]human plasma alpha-1-B-glycoprotein: phenotyping by
immunoblotting

[140171]or by a simple method of 2-D electrophoresis. Hum.
Genet. 76: 111-115,

[140172]1987.

[140173]25048.

[140174]Ishioka, N.; Takahashi, N.; Putnam, F. W.: Amino acid se-
quence

[140175]of human plasma alpha-1B-glycoprotein: homology to the
immunoglobulin

[140176]supergene family. Proc. Nat. Acad. Sci. 83: 2363-2367,
1986.

[140177]25049.

[140178]Juneja, R. K.; Weitkamp, L. R.; Stratil, A.; Gahne, B.; Gut-
tormsen,

[140179]S. A.: Further studies of the plasma alpha-

1-B-glycoprotein polymorphism:

[140180]two new alleles and allele frequencies in Caucasians and
in American

[140181]blacks. Hum. Hered. 38: 267-272, 1988.

[140182]25050.

[140183]Anderson, L.; Anderson, N. G.: High resolution two-
dimensional

[140184]electrophoresis of human plasma proteins. Proc. Nat.
Acad. Sci. 74:

[140185]5421-5425, 1977.

[140186]25051.

[140187]Boutin, B.; Feng, S. H.; Arnaud, P.: The genetic polymor-
phism

[140188]of alpha(2)-HS glycoprotein: study by ultrathin-layer iso-
electric

[140189]focusing and immunoblot. Am. J. Hum. Genet. 37:
1098-1105, 1985.

[140190]25052.

[140191]Cox, D. W.; Andrews, B. J.: Silver stain immunofixation for
alpha-2-HS-glycoprotein:

[140192]a new method for detection of protein heterogeneity. In:

Stathakos,

[140193]D.: Electrophoresis '82. Berlin: Walter de Gruyter (pub.)
1983.

[140194]Pp. 243–247.

[140195]25053.

[140196]Cox, D. W.; Andrews, B. J.; Wills, D. E.: Genetic polymor-
phism

[140197]of alpha-2-HS-glycoprotein. Am. J. Hum. Genet. 38:
699–706, 1986.

[140198]25054.

[140199]Cox, D. W.; Francke, U.; Allderdice, P. W.; McAlpine, P. J.:
Gene

[140200]mapping of human serum proteins using hepatoma hy-
brids and human chromosome

[140201]deletions and duplications. Genetics 107 (suppl.):
s22–s23, 1984.

[140202]25055.

[140203]Eiberg, H.; Mohr, J.; Nielsen, L. S.: A2HS: new methods of
phenotyping

[140204]and analysis of linkage relations: assignment to chromo-
some 3. (Abstract) Cytogenet.

[140205]Cell Genet. 37: 461 only, 1984.

[140206]25056.

[140207]Gejyo, F.; Chang, J.-L.; Burgi, W.; Schmid, K.; Offner, G. D.;

[140208]Troxler, R. F.; Van Halbeek, H.; Dorland, L.; Gerwig, G. J.;
Vliegthart,

[140209]J. F. G.: Characterization of the B-chain of human plasma
alpha(2)HS-glycoprotein:

[140210]the complete amino acid sequence and primary structure
of its heteroglycan. J.

[140211]Biol. Chem. 258: 4966-4971, 1983.

[140212]25057.

[140213]Jahnen-Dechent, W.: Personal Communication. Mainz,
Germany

[140214]7/22/1998.

[140215]25058.

[140216]Jahnen-Dechent, W.; Schinke, T.; Trindl, A.; Muller-Esterl,
W.;

[140217]Sablitzky, F.; Kaiser, S.; Blessing, M.: Cloning and targeted
deletion

[140218]of the mouse fetuin gene. J. Biol. Chem. 272:
31496-31503, 1997.

[140219]25059.

- [140220]Lee, C.-C.; Bowman, B. H.; Yang, F.: Human alpha-2-HS-glycoprotein: the A and B chains with a connecting sequence are encoded by a single mRNA transcript. Proc. Nat. Acad. Sci. 84: 4403-4407, 1987.
- [140221]25060.
- [140222]Fajans, S. S.; Bell, G. I.; Polonsky, K. S.: Molecular mechanisms and clinical pathophysiology of maturity-onset diabetes of the young. New Eng. J. Med. 345: 971-980, 2001.
- [140223]25061.
- [140224]Lindner, T.; Gragnoli, C.; Furuta, H.; Cockburn, B. N.; Petzold, C.; Rietzsch, H.; Weiss, U.; Schulze, J.; Bell, G. I.: Hepatic function in a family with a nonsense mutation (R154X) in the hepatocyte nuclear factor-4-alpha/MODY1 gene. J. Clin. Invest. 100: 1400-1405, 1997.
- [140225]25062.

- [140233]Moller, A. M.; Dalgaard, L. T.; Ambye, L.; Hansen, L.; Schmitz, O.; Hansen, T.; Pedersen, O.: A novel Phe75fsdelT mutation in the hepatocyte nuclear factor-4-alpha gene in a Danish pedigree with maturity-onset diabetes of the young. J. Clin. Endocr. Metab. 84: 367-369, 1999.
- [140237]25063.
- [140238]Thomas, H.; Jaschkowitz, K.; Bulman, M.; Frayling, T. M.; Mitchell, S. M. S.; Roosen, S.; Lingott-Frieg, A.; Tack, C. J.; Ellard, S.; Ryffel, G. U.; Hattersley, A. T.: A distant upstream promoter of the HNF-4-alpha gene connects the transcription factors involved in maturity-onset diabetes of the young. Hum. Molec. Genet. 10: 2089-2097, 2001.
- [140243]2001.
- [140244]25064.
- [140245]Froguel, P.; Vaxillaire, M.; Sun, F.; Velho, G.; Zouali, H.; Butel,

- [140246]M. O.; Lesage, S.; Vionnet, N.; Clement, K.; Fougerousse, F.; Tanizawa,
- [140247]Y.; Weissenbach, J.; Beckmann, J. S.; Lathrop, G. M.; Passa, P.; Permutt,
- [140248]M. A.; Cohen, D.: Close linkage of glucokinase locus on chromosome
- [140249]7p to early-onset non-insulin-dependent diabetes mellitus. Nature 356:
- [140250]162-164, 1992.
- [140251]25065.
- [140252]Vionnet, N.; Stoffel, M.; Takeda, J.; Yasuda, K.; Bell, G. I.;
- [140253]Zouali, H.; Lesage, S.; Velho, G.; Iris, F.; Passa, P.; Froguel, P.;
- [140254]Cohen, D.: Nonsense mutation in the glucokinase gene causes early-onset
- [140255]non-insulin-dependent diabetes mellitus. Nature 356: 721-722, 1992.
- [140256]25066.
- [140257]Dunger, D. B.; Ong, K. K. L.; Huxtable, S. J.; Sherriff, A.; Woods,
- [140258]K. A.; Ahmed, M. L.; Golding, J.; Pembrey, M. E.; Ring, S.;
- ALSPAC

[140259]Study Team; Bennett, S. T.; Todd, J. A.: Association of the
INS

[140260]VNTR with size at birth. Nature Genet. 19: 98–100, 1998.

[140261]25067.

[140262]Joseph, L. J.; Le Beau, M. M.; Jamieson, G. A., Jr.; Acharya,
S.;

[140263]Shows, T. B.; Rowley, J. D.; Sukhatme, V. P.: Molecular
cloning,

[140264]sequencing, and mapping of EGR2, a human early growth
response gene

[140265]encoding a protein with 'zinc-binding finger' structure.
Proc. Nat.

[140266]Acad. Sci. 85: 7164–7168, 1988.

[140267]25068.

[140268]Nagarajan, R.; Svaren, J.; Le, N.; Araki, T.; Watson, M.; Mil-
brandt,

[140269]J.: EGR2 mutations in inherited neuropathies dominant-
negatively

[140270]inhibit myelin gene expression. Neuron 30: 355–368,
2001.

[140271]25069.

[140272] Timmerman, V.; De Jonghe, P.; Ceuterick, C.; De Vriendt, E.; Lofgren, A.; Nelis, E.; Warner, L. E.; Lupski, J. R.; Martin, J.-J.; Van Broeckhoven, C.: Novel missense mutation in the early growth response 2 gene associated with Dejerine-Sottas syndrome phenotype. *Neurology* 52: 1827-1832, 1999.

[140276] 25070.

[140277] Warner, L. E.; Mancias, P.; Butler, I.; Lupski, J. R.: Mutation in the early growth response 2 (EGR2) transcription factor associated with recessive congenital hypomyelinating neuropathy (CHN). (Abstract) *Am. J. Hum. Genet.* 61 (suppl.): A350 only, 1997.

[140281] 25071.

[140282] Warner, L. E.; Mancias, P.; Butler, I. J.; McDonald, C. M.; Keppen, L.; Koob, K. G.; Lupski, J. R.: Mutations in the early growth response

- [140285]2 (EGR2) gene are associated with hereditary myelinopathies. Nature
- [140286]Genet. 18: 382–384, 1998.
- [140287]25072.
- [140288]Warner, L. E.; Svaren, J.; Milbrandt, J.; Lupski, J. R.: Functional
- [140289]consequences of mutations in the early growth response 2 gene (EGR2)
- [140290]correlate with severity of human myelinopathies. Hum. Molec. Genet. 8:
- [140291]1245–1251, 1999.
- [140292]25073.
- [140293]Wu, J.; Joseph, L.; Sukhatme, V. P.; Kidd, K. K.: A HindIII polymorphism
- [140294]identified by the human early growth response gene 2 (EGR2) on chromosome 25074.
- [140295]Nucleic Acids Res. 16: 11855 only, 1988.
- [140296]25075.
- [140297]Kaneda, Y.; Hayes, H.; Uchida, T.; Yoshida, M. C.; Okada, Y.:
- [140298]Regional assignment of five genes on human chromosome

19. Chromosoma 95:

[140299]8-12, 1987.

[140300]25076.

[140301]Boyle, J. M.; Hey, Y.; Fox, M.: Human ecto-5-prime nucleotidase

[140302]maps to chromosome 6q14-q21. (Abstract) Cytogenet. Cell Genet. 51:

[140303]968 only, 1989.

[140304]25077.

[140305]Boyle, J. M.; Hey, Y.; Geurts van Kessel, A.; Fox, M.: Assignment

[140306]of ecto-5-prime-nucleotidase to human chromosome 6. Hum. Genet. 81:

[140307]88-92, 1988.

[140308]25078.

[140309]Misumi, Y.; Ogata, S.; Ohkubo, K.; Hirose, S.; Ikehara, Y.: Primary

[140310]structure of human placental 5-prime-nucleotidase and identification

[140311]of the glycolipid anchor in the mature form. Europ. J. Biochem. 191:

[140312]563–569, 1990.

[140313]25079.

[140314]Monreal, A. W.; Ferguson, B. M.; Headon, D. J.; Street, S. L.;

[140315]Overbeek, P. A.; Zonana, J.: Mutations in the human homologue of

[140316]mouse dl cause autosomal recessive and dominant hypohidrotic ectodermal

[140317]dysplasia. *Nature Genet.* 22: 366–369, 1999.

[140318]25080.

[140319]Schrijver, I.; Liu, W.; Brenn, T.; Furthmayr, H.; Francke, U.

[140320]: Cysteine substitutions in epidermal growth factor-like domains of

[140321]fibrillin-1: distinct effects on biochemical and clinical phenotypes. *Am.*

[140322]*J. Hum. Genet.* 65: 1007–1020, 1999.

[140323]25081.

[140324]Schrijver, I.; Liu, W.; Francke, U.: The pathogenicity of the

[140325]Pro1148-to-ala substitution in the FBN1 gene: causing or predisposing

[140326]to Marfan syndrome and aortic aneurysm, or clinically innocent? Hum.

[140327]Genet. 99: 607–611, 1997.

[140328]25082.

[140329]Schrijver, I.; Liu, W.; Odom, R.; Brenn, T.; Oefner, P.; Furthmayr,

[140330]H.; Francke, U.: Premature termination mutations in FBN1: distinct

[140331]effects on differential allelic expression and on protein and clinical

[140332]phenotypes. Am. J. Hum. Genet. 71: 223–237, 2002.

[140333]25083.

[140334]Siracusa, L. D.; McGrath, R.; Ma, Q.; Moskow, J. J.; Manne, J.;

[140335]Christner, P. J.; Buchberg, A. M.; Jimenez, S. A.: A tandem duplication

[140336]within the fibrillin 1 gene is associated with the mouse tight skin

[140337]mutation. Genome Res. 6: 300–313, 1996.

[140338]25084.

[140339]Sood, S.; Eldadah, Z. A.; Krause, W. L.; McIntosh, I.; Dietz,

- [140340]H. C.: Mutation in fibrillin-1 and the Marfanoid-craniosynostosis
- [140341](Shprintzen-Goldberg) syndrome. Nature Genet. 12: 209-211, 1996.
- [140342]25085.
- [140343]Stahl-Hallengren, C.; Ukkonen, T.; Kainulainen, K.; Kristofersson,
- [140344]U.; Saxne, T.; Tornqvist, K.; Peltonen, L.: An extra cysteine in
- [140345]one of the non-calcium-binding epidermal growth factor-like motifs
- [140346]of the FBN1 polypeptide is connected to a novel variant of Marfan
- [140347]syndrome. J. Clin. Invest. 94: 709-713, 1994.
- [140348]25086.
- [140349]Tiecke, F.; Katzke, S.; Booms, P.; Robinson, P. N.; Neumann, L.;
- [140350]Godfrey, M.; Mathews, K. R.; Scheuner, M.; Hinkel, G. K.; Brenner,
- [140351]R. E.; Hovels-Gurich, H. H.; Hagemeier, C.; Fuchs, J.; Skovby, F.;
- [140352]Rosenberg, T.: Classic, atypically severe and neonatal

Marfan syndrome:

- [140353] twelve mutations and genotype-phenotype correlations in
FBN1 exons
[140354] 24-40. *Europ. J. Hum. Genet.* 9: 13-21, 2001.
- [140355] 25087.
- [140356] Trask, T. M.; Ritty, T. M.; Broekelmann, T.; Tisdale, C.;
Mecham,
[140357] R. P.: N-terminal domains of fibrillin 1 and fibrillin 2 di-
rect the
[140358] formation of homodimers: a possible first step in mi-
crofibril assembly. *Biochem.*
[140359] *J.* 340: 693-701, 1999.
- [140360] 25088.
- [140361] Tynan, K.; Comeau, K.; Pearson, M.; Wilgenbus, P.; Levitt,
D.;
- [140362] Gasner, C.; Berg, M. A.; Miller, D. C.; Francke, U.: Mutation
screening
[140363] of complete fibrillin-1 coding sequence: report of five new
mutations,
[140364] including two in 8-cysteine domains. *Hum. Molec. Genet.*
2: 1813-1821,
[140365] 1993.

[140366]25089.

[140367]Velinov, M.; Sarfarazi, M.; Young, K.; Hodes, M. E.; Con-
neally,

[140368]P. M.; Jackson, C. E.; Tsipouras, P.: Limb-girdle muscular
dystrophy

[140369]is closely linked to the fibrillin locus on chromosome 15.
Connect.

[140370]Tissue Res. 29: 13-21, 1993.

[140371]25090.

[140372]Wang, M.; Mathews, K. R.; Imaizumi, K.; Beiraghi, S.;
Blumberg,

[140373]B.; Scheuner, M.; Graham, J. M., Jr.; Godfrey, M.: P1148A in
fibrillin-1

[140374]is not a mutation anymore. (Letter) Nature Genet. 15: 12
only, 1997.

[140375]25091.

[140376]Wang, M.; Wang, J.-Y.; Cisler, J.; Imaizumi, K.; Burton, B.
K.;

[140377]Jones, M. C.; Lamberti, J. J.; Godfrey, M.: Three novel fib-
rillin

[140378]mutations in exons 25 and 27: classic versus neonatal
Marfan syndrome. Hum.

[140379]Mutat. 9: 359–362, 1997.

[140380]25092.

[140381]Watanabe, Y.; Yano, S.; Koga, Y.; Yukizane, S.; Nishiyori, A.;

[140382]Yoshino, M.; Kato, H.; Ogata, T.; Adachi, M.: P1148A in fibrillin-1

[140383]is not a mutation leading to Shprintzen-Goldberg syndrome. (Letter) Hum.

[140384]Mutat. 10: 326–327, 1997.

[140385]25093.

[140386]Whiteman, P.; Smallridge, R. S.; Knott, V.; Cordle, J. J.; Downing,

[140387]A. K.; Handford, P. A.: A G1127S change in calcium-binding epidermal

[140388]growth factor-like domain 13 of human fibrillin-1 causes short range

[140389]conformational effects. J. Biol. Chem. 276: 17156–17162, 2001.

[140390]25094.

[140391]Alving, B. M.; Henschen, A. H.: Fibrinogen Giessen I: a congenital

[140392]homozygously expressed dysfibrinogenemia with A-alpha
arg-to-his substitution. Am.

[140393]J. Hemat. 25: 479-482, 1987.

[140394]25095.

[140395]Arocha-Pinango, C. L.; Rodriguez, S.; Nagy, H.; Perez Re-
quejo,

[140396]J. L.: Fibrinogen Lima: a new dysfibrinogenaemia with a
high-molecular

[140397]weight alpha-chain and effective polymerization. Blood
Coagul. Fibrinolysis 1:

[140398]561-565, 1990.

[140399]25096.

[140400]Aschbacher, A.; Buetow, K.; Chung, D.; Walsh, S.; Murray,
J.:

[140401]Linkage disequilibrium of RFLP's associated with alpha,
beta, and

[140402]gamma fibrinogen predict gene order on chromosome 4.
(Abstract) Am.

[140403]J. Hum. Genet. 37: A186 only, 1985.

[140404]25097.

[140405]Asl, L. H.; Liepnieks, J. J.; Uemichi, T.; Rebibou, J.-M.; Jus-

trabo,

[140406]E.; Droz, D.; Mousson, C.; Chalopin, J.-M.; Benson, M. D.;
Delpech,

[140407]M.; Grateau, G.: Renal amyloidosis with a frame shift mu-
tation in

[140408]fibrinogen A(alpha)-chain gene producing a novel amyloid
protein. Blood 90:

[140409]4799-4805, 1997.

[140410]25098.

[140411]Asselta, R.; Duga, S.; Spena, S.; Santagostino, E.; Peyvandi,
F.;

[140412]Piseddu, G.; Targhetta, R.; Malcovati, M.; Mannucci, P. M.;
Tenchini,

[140413]M. L.: Congenital afibrinogenemia: mutations leading to
premature

[140414]termination codons in fibrinogen A-alpha-chain gene are
not associated

[140415]with the decay of the mutant mRNAs. Blood 98:
3685-3692, 2001.

[140416]25099.

[140417]Aznar, J.; Fernandez-Pavon, A.; Reganon, E.; Vila, V.; Orel-
lana,

[140418]F.: Fibrinogen Valencia--a new case of congenital dysfibrinogenemia. Thromb.

[140419]Diath. Haemorrh. 32: 564-577, 1974.

[140420]25100.

[140421]Barthels, M.; Sandvoss, G.: Fibrinogen Hannover, another abnormal

[140422]fibrinogen. Blut 34: 99-106, 1977.

[140423]25101.

[140424]Beck, E. A.; Charache, P.; Jackson, D. P.: A new inherited coagulation

[140425]disorder caused by an abnormal fibrinogen ('fibrinogen Baltimore'). Nature 208:

[140426]143-145, 1965.

[140427]25102.

[140428]Beck, E. A.; Shainoff, J. R.; Vogel, A.; Jackson, D. P.: Functional

[140429]evaluation of an inherited abnormal fibrinogen: fibrinogen 'Baltimore.'. J.

[140430]Clin. Invest. 50: 1874-1884, 1971.

[140431]25103.

[140432]Benson, M. D.; Liepnieks, J.; Uemichi, T.; Wheeler, G.; Cor-
rea,

[140433]R.: Hereditary renal amyloidosis associated with a mutant
fibrinogen

[140434]alpha-chain. Nature Genet. 3: 252-255, 1993.

[140435]25104.

[140436]Blomback, B.; Blomback, M.: Molecular defects and vari-
ants of

[140437]fibrinogen. Nouv. Rev. Franc. Hemat. 10: 671-678, 1970.

[140438]25105.

[140439]Blomback, B.; Hessel, B.; Fields, R.; Procyk, R.: Fibrinogen

[140440]Aarhus: an abnormal fibrinogen with A alpha 19 arg-
to-gly substitution.In:

[140441]Mosesson, M. W.; Amrani, D. L.; Siebenlist, K. R.; Di Orio,
J. P.

[140442]: Fibrinogen 3: Biochemistry, Biological Functions, Gene
Regulation

[140443]and Expression.. Amsterdam: Elsevier (pub.) 1988. Pp.
263-266.

[140444]25106.

[140445]Ballo, R.; Beighton, P. H.; Ramesar, R. S.: Stickler-like syn-

drome

[140446]due to a dominant negative mutation in the COL2A1 gene.

Am. J. Med.

[140447]Genet. 80: 6–11, 1998.

[140448]25107.

[140449]Chan, E. F.; Gat, U.; McNiff, J. M.; Fuchs, E.: A common human

[140450]skin tumour is caused by activating mutations in beta-catenin. Nature

[140451]Genet. 21: 410–413, 1999.

[140452]25108.

[140453]Oh, S. P.; Kamagata, Y.; Muragaki, Y.; Timmons, S.; Ooshima, A.;

[140454]Olsen, B. R.: Isolation and sequencing of cDNAs for proteins with

[140455]multiple domains of Gly–X–Y repeats identify a novel family of collagenous

[140456]proteins. Proc. Nat. Acad. Sci. 91: 4229–4233, 1994.

[140457]25109.

[140458]Oh, S. P.; Warman, M. L.; Seldin, M. F.; Cheng, S.–D.; Knoll, J.

[140459]H. M.; Timmons, S.; Olsen, B. R.: Cloning of cDNA and genomic DNA
[140460]encoding human type XVIII collagen and localization of the
alpha-1(XVIII)
[140461]collagen gene to mouse chromosome 10 and human
chromosome 21. Genomics 19:
[140462]494-499, 1994.
[140463]25110.
[140464]Rehn, M.; Hintikka, E.; Pihlajaniemi, T.: Characterization of
[140465]the mouse gene for the alpha-1 chain of type XVIII collagen (Col18a1)
[140466]reveals that the three variant N-terminal polypeptide
forms are transcribed
[140467]from two widely separated promoters. Genomics 32:
436-446, 1996.
[140468]25111.
[140469]Rehn, M.; Pihlajaniemi, T.: Alpha-1(XVIII), a collagen chain
with
[140470]frequent interruptions in the collagenous sequence, a distinct tissue
[140471]distribution, and homology with type XV collagen. Proc.
Nat. Acad.

[140472]Sci. 91: 4234–4238, 1994.

[140473]25112.

[140474]Saarela, J.; Ylikarppa, R.; Rehn, M.; Purmonen, S.; Pihlajaniemi,

[140475]T.: Complete primary structure of two variant forms of human type

[140476]XVIII collagen and tissue-specific differences in the expression of

[140477]the corresponding transcripts. Matrix Biol. 16: 319–328, 1998.

[140478]25113.

[140479]Horelli-Kuitunen, N.; Kvist, A.-P.; Helaakoski, T.; Kivirikko,

[140480]K.; Pihlajaniemi, T.; Palotie, A.: The order and transcriptional

[140481]orientation of the human COL13A1 and P4HA genes on chromosome 10 long

[140482]arm determined by high-resolution FISH. Genomics 46: 299–302, 1997.

[140483]25114.

[140484]Pajunen, L.; Tamminen, M.; Solomon, E.; Pihlajaniemi, T.: Assignment

[140485] of the gene coding for the alpha 1 chain of collagen type
XIII (COL13A1)

[140486] to human chromosome region 10q11-qter. Cytogenet.
Cell Genet. 52:

[140487] 190-193, 1989.

[140488] 25115.

[140489] Shows, T. B.; Tikka, L.; Byers, M. G.; Eddy, R. L.; Haley, L.
L.;

[140490] Henry, W. M.; Prockop, D. J.; Tryggvason, K.: Assignment
of the human

[140491] collagen alpha-1(XIII) chain gene (COL13A1) to the q22
region of chromosome

[140492] 10. Genomics 5: 128-133, 1989.

[140493] 25116.

[140494] Tikka, L.; Pihlajaniemi, T.; Henttu, P.; Prockop, D. J.; Tryg-
gvason,

[140495] K.: Gene structure for the alpha-1 chain of a human
short-chain collagen

[140496] (type XIII) with alternatively spliced transcripts and trans-
lation

[140497] termination codon at the 5-prime end of the last exon.
Proc. Nat.

[140498]Acad. Sci. 85: 7491–7495, 1988.

[140499]25117.

[140500]Cunningham, B. A.; Hemperly, J. J.; Murray, B. A.; Prediger, E.

[140501]A.; Brackenbury, R.; Edelman, G. M.: Neural cell adhesion molecule:

[140502]structure, immunoglobulin-like domains, cell surface modulation, and

[140503]alternative RNA splicing. Science 236: 799–806, 1987.

[140504]25118.

[140505]D'Eustachio, P.; Davisson, M. T.: Resolution of the staggerer

[140506](sg) mutation from the neural cell adhesion molecule locus (Ncam)

[140507]on mouse chromosome 9. Mammalian Genome 4: 278–280, 1993.

[140508]25119.

[140509]D'Eustachio, P.; Owens, G. C.; Edelman, G. M.; Cunningham, B. A.

[140510]: Chromosomal location of the gene encoding the neural cell adhesion

[140511]molecule (N-CAM) in the mouse. Proc. Nat. Acad. Sci. 82:
7631-7635,

[140512]1985.

[140513]25120.

[140514]Lin, D. M.; Fetter, R. D.; Kopczynski, C.; Grenningloh, G.;
Goodman,

[140515]C. S.: Genetic analysis of fasciclin II in Drosophila: defasci-
cultation,

[140516]refasciculation, and altered fasciculation. Neuron 13:
1055-1069,

[140517]1994.

[140518]25121.

[140519]Altare, F.; Jouanguy, E.; Lamhamedi-Cherradi, S.; Fon-
daneche, M.-C.;

[140520]Fizame, C.; Ribierre, F.; Merlin, G.; Dembic, Z.; Schreiber,
R.; Lisowska-Groszpiere,

[140521]B.; Fischer, A.; Seboun, E.; Casanova, J.-L.: A causative re-
lationship

[140522]between mutant IFNGR1 alleles and impaired cellular re-
sponse to IFN-gamma

[140523]in a compound heterozygous child. (Letter) Am. J. Hum.
Genet. 62:

[140524]723–726, 1998.

[140525]25122.

[140526]Dalton, D. K.; Pitts–Meek, S.; Keshav, S.; Figari, I. S.;
Bradley,

[140527]A.; Stewart, T. A.: Multiple defects of immune cell function
in mice

[140528]with disrupted interferon–gamma genes. Science 259:
1739–1742, 1993.

[140529]25123.

[140530]Dessein, A. J.; Hillaire, D.; Elwali, N. E. M. A.; Marquet, S.;

[140531]Mohamed–Ali, Q.; Mirghani, A.; Henri, S.; Abdelhameed, A.
A.; Saeed,

[140532]O. K.; Magzoub, M. M. A.; Abel, L.: Severe hepatic fibrosis
in Schistosoma

[140533]mansoni infection is controlled by a major locus that is
closely linked

[140534]to the interferon–gamma receptor gene. Am. J. Hum.
Genet. 65: 709–721,

[140535]1999.

[140536]25124.

[140537]Fellous, M.: Personal Communication. Paris, France

10/24/1986.

[140538]25125.

[140539]Fellous, M.; Couillin, P.; Rosa, F.; Metezeau, P.; Foubert, C.;

[140540]Gross, M. S.; Frezal, J.; Van Cong, N.: Receptor for human gamma

[140541]interferon is specified by human chromosome 18.
(Abstract) Cytogenet.

[140542]Cell Genet. 40: 627-628, 1985.

[140543]25126.

[140544]Jouanguy, E.; Altare, F.; Lamhamedi, S.; Revy, P.; Emile, J.-F.;

[140545]Newport, M.; Levin, M.; Blanche, S.; Seboun, E.; Fischer, A.; Casanova,

[140546]J.-L.: Interferon-gamma-receptor deficiency in an infant with fatal

[140547]bacille Calmette-Guerin infection. New Eng. J. Med. 335: 1956-1961,

[140548]1996.

[140549]25127.

[140550]Jouanguy, E.; Dupuis, S.; Pallier, A.; Doffinger, R.; Fon-

daneche,

[140551]M.-C.; Fieschi, C.; Lamhamedi-Cherradi, S.; Altare, F.;
Emile, J.-F.;

[140552]Lutz, P.; Bordigoni, P.; Cokugras, H.; Akcakaya, N.; Land-
man-Parker,

[140553]J.; Donnadieu, J.; Camcioglu, Y.; Casanova, J.-L.: In a novel
form

[140554]of IFN-gamma receptor 1 deficiency, cell surface recep-
tors fail to

[140555]bind IFN-gamma. J. Clin. Invest. 105: 1429-1436, 2000.

[140556]25128.

[140557]Jouanguy, E.; Lamhamedi-Cherradi, S.; Lammas, D.; Dor-
man, S. E.;

[140558]Fondaneche, M.-C.; Dupuis, S.; Doffinger, R.; Altare, F.;
Girdlestone,

[140559]J.; Emile, J.-F.; Ducoulombier, H.; Edgar, D.; and 10 others:
A

[140560]human IFNGR1 small deletion hotspot associated with
dominant susceptibility

[140561]to mycobacterial infection. Nature Genet. 21: 370-378,
1999.

[140562]25129.

[140563]Kaplan, D. H.; Shankaran, V.; Dighe, A. S.; Stockert, E.;
 Aguet,
 [140564]M.; Old, L. J.; Schreiber, R. D.: Demonstration of an inter-
 feron
 [140565]gamma-dependent tumor surveillance system in immuno-
 competent mice. Proc.
 [140566]Nat. Acad. Sci. 95: 7556-7561, 1998.
 [140567]25130.
 [140568]Le Coniat, M.; Alcaide-Loridan, C.; Fellous, M.; Berger, R.:
 [140569]Human interferon gamma receptor 1 (IFNGR1) gene maps
 to chromosome
 [140570]region 6q23-6q24. Hum. Genet. 84: 92-94, 1989.
 [140571]25131.
 [140572]Lekstrom-Himes, J. A.; Gallin, J. I.: Immunodeficiency dis-
 eases
 [140573]caused by defects in phagocytes. New Eng. J. Med. 343:
 1703-1714,
 [140574]2000.
 [140575]25132.
 [140576]Levin, M.; Newport, M. J.; D'Souza, S.; Kalabalikis, P.;
 Brown,

- [140577]J. N.; Lenicker, H. M.; Agius, P. V.; Davies, E. G.; Thrasher, A.;
- [140578]Klein, N.; Blackwell, J. M.: Familial disseminated atypical mycobacterial
- [140579]infection in childhood: a human mycobacterial susceptibility gene? Lancet 345:
- [140580]79–83, 1995.
- [140581]25133.
- [140582]Mariano, T. M.; Kozak, C. A.; Langer, J. A.; Pestka, S.: The
- [140583]mouse immune interferon receptor gene is located on chromosome 10. J.
- [140584]Biol. Chem. 262: 5812–5814, 1987.
- [140585]25134.
- [140586]Deloukas, P.; Schuler, G. D.; Gyapay, G.; Beasley, E. M.; Soderlund,
- [140587]C.; Rodriguez–Tome, P.; Hui, L.; Matise, T. C.; McKusick, K. B.; Beckmann,
- [140588]J. S.; Bentolila, S.; Bihoreau, M.–T.; and 53 others: A physical
- [140589]map of 30,000 human genes. Science 282: 744–746, 1998.

[140590]25135.

[140591]Manser, E.; Leung, T.; Salihuddin, H.; Tan, L.; Lim, L.: A
non-receptor

[140592]tyrosine kinase that inhibits the GTPase activity of
p21cdc42. Nature 363:

[140593]364–367, 1993.

[140594]25136.

[140595]Wu, W. J.; Erickson, J. W.; Lin, R.; Cerione, R. A.: The
gamma-subunit

[140596]of the coatomer complex binds Cdc42 to mediate trans-
formation. Nature 405:

[140597]800–804, 2000.

[140598]25137.

[140599]Basu, S.; Binder, R. J.; Ramalingam, T.; Srivastava, P. K.:
CD91

[140600]is a common receptor for heat shock proteins gp96,
hsp90, hsp70, and

[140601]calreticulin. Immunity 14: 303–313, 2001.

[140602]25138.

[140603]Beisiegel, U.; Weber, W.; Ihrke, G.; Herz, J.; Stanley, K. K.:

[140604]The LDL-receptor-related protein, LRP, is an apolipoprotein E-binding

[140605]protein. *Nature* 341: 162-164, 1989.

[140606]25139.

[140607]Binder, R. J.; Han, D. K.; Srivastava, P. K.: CD91: a receptor

[140608]for heat shock protein gp96. *Nature Immun.* 1: 151-155, 2000.

[140609]25140.

[140610]Forus, A.; Maelandsmo, G. M.; Fodstad, Y.; Myklebost, O.: The

[140611]genes for the alpha-2-macroglobulin receptor/LDL receptor-related

[140612]protein and GLI are located within a chromosomal segment of about

[140613]300 kilobases and are coamplified in a rhabdomyosarcoma cell line.

[140614](Abstract) *Cytogenet. Cell Genet.* 58: 1977 only, 1991.

[140615]25141.

[140616]Forus, A.; Myklebost, O.: A physical map of a 1.3-Mb region on

[140617]the long arm of chromosome 12, spanning the GLI and

LRP loci. Genomics 14:

[140618]117–120, 1992.

[140619]25142.

[140620]Hilliker, C.; Van Leuven, F.; Van Den Berghe, H.: Assignment of

[140621]the gene coding for the alpha(2)-macroglobulin receptor to mouse chromosome

[140622]15 and to human chromosome 12q13–q14 by isotopic and nonisotopic in

[140623]situ hybridization. Genomics 13: 472–474, 1992.

[140624]25143.

[140625]Kristensen, T.; Moestrup, S. K.; Gliemann, J.; Bendtsen, L.; Sand,

[140626]O.; Sottrup-Jensen, L.: Evidence that the newly cloned low-density-lipoprotein

[140627]receptor related protein (LRP) is the alpha-2-macroglobulin receptor. FEBS

[140628]Lett. 276: 151–155, 1990.

[140629]25144.

[140630]Lendon, C. L.; Talbot, C. J.; Craddock, N. J.; Han, S. W.; Wragg,

- [140631] M.; Morris, J. C.; Goate, A. M.: Genetic association studies between
- [140632] dementia of the Alzheimer's type and three receptors for apolipoprotein
- [140633] E in a Caucasian population. *Neurosci. Lett.* 222: 187-190, 1997.
- [140634] 25145.
- [140635] Myklebost, O.; Arheden, K.; Rogne, S.; Geurts van Kessel, A.;
- [140636] Mandahl, N.; Herz, J.; Stanley, K.; Heim, S.; Mitelman, F.: The gene
- [140637] for the human putative apoE receptor is on chromosome 12 in the segment

[140638]q13–14. Genomics 5: 65–69, 1989.

[140639]25146.

[140640]Pericak–Vance, M. A.; Bass, M. P.; Yamaoka, L. H.; Gaskell, P.

[140641]C.; Scott, W. K.; Terwedow, H. A.; Menold, M. M.; Conneally, P. M.;

[140642]Small, G. W.; Vance, J. M.; Saunders, A. M.; Roses, A. D.; Haines,

[140643]J. L.: Complete genomic screen in late-onset familial Alzheimer disease:

[140644]evidence for a new locus on chromosome 12. J.A.M.A. 278: 1237–1241,

[140645]1997.

[140646]25147.

[140647]Scott, W. K.; Yamaoka, L. H.; Bass, M. P.; Gaskell, P. C.; Conneally,

[140648]P. M.; Small, G. W.; Farrer, L. A.; Auerbach, S. A.; Saunders, A.

[140649]M.; Roses, A. D.; Haines, J. L.; Pericak–Vance, M. A.: No genetic

[140650]association between the LRP receptor and sporadic or late-onset familial

[140651]Alzheimer disease. Neurogenetics 1: 179–183, 1998.

[140652]25148.

[140653]Strickland, D. K.; Ashcom, J. D.; Williams, S.; Burgess, W.
H.;

[140654]Migliorini, M.; Argraves, W. S.: Sequence identity between
the alpha-2-macroglobulin

[140655]receptor and low density lipoprotein receptor-related pro-
tein suggests

[140656]that this molecule is a multifunctional receptor. J. Biol.
Chem. 265:

[140657]17401–17404, 1990.

[140658]25149.

[140659]Yochem, J.; Greenwald, I.: A gene for a low density
lipoprotein

[140660]receptor-related protein in the nematode *Caenorhabditis*
elegans. Proc.

[140661]Nat. Acad. Sci. 90: 4572–4576, 1993.

[140662]25150.

[140663]Ladías, J. A. A.; Karathanasis, S. K.: Regulation of the
apolipoprotein

[140664]AI gene by ARP-1, a novel member of the steroid receptor

superfamily. Science 251:

[140665]561–565, 1991.

[140666]25151.

[140667]Modi, W. S.; Seuanes, H.; Mietus-Snyder, M.; O'Brien, S. J.;
Karathanasis,

[140668]S. K.: Chromosomal localization of the ARP-1 gene to
15q26. (Abstract) Cytogenet.

[140669]Cell Genet. 58: 1995 only, 1991.

[140670]25152.

[140671]Pereira, F. A.; Qiu, Y.; Zhou, G.; Tsai, M.-J.; Tsai, S. Y.: The
[140672]orphan nuclear receptor COUP-TFII is required for angio-
genesis and

[140673]heart development. Genes Dev. 13: 1037–1049, 1999.

[140674]25153.

[140675]Widom, R. L.; Ladas, J. A. A.; Kouidou, S.; Karathanasis, S.
K.

[140676]: Synergistic interactions between transcription factors
control expression

[140677]of the apolipoprotein AI gene in liver cells. Molec. Cell.
Biol. 11:

[140678]677–687, 1991.

[140679]25154.

[140680]Agre, P.; Smith, B. L.; Baumgarten, R.; Preston, G. M.;
Pressman,

[140681]E.; Wilson, P.; Illum, N.; Anstee, D. J.; Lande, M. B.; Zeidel,
M.

[140682]L.: Human red cell aquaporin CHIP. II. Expression during
normal fetal

[140683]development and in a novel form of congenital dysery-
thropoietic anemia. J.

[140684]Clin. Invest. 94: 1050-1058, 1994.

[140685]25155.

[140686]Ikeda, H.; Yamaguchi, M.; Sugai, S.; Aze, Y.; Narumiya, S.;
Kakizuka,

[140687]A.: Expanded polyglutamine in the Machado-Joseph dis-
ease protein

[140688]induces cell death in vitro and in vivo. Nature Genet. 13:
196-202,

[140689]1996.

[140690]25156.

[140691]Santisteban, I.; Arredondo-Vega, F. X.; Kelly, S.; Mary, A.;
Fischer,

[140692]A.; Hummell, D. S.; Lawton, A.; Sorensen, R. U.; Stiehm, E.
R.; Uribe,
[140693]L.; Weinberg, K.; Hershfield, M. S.: Novel splicing, mis-
sense, and
[140694]deletion mutations in seven adenosine deaminase-defi-
cient patients
[140695]with late/delayed onset of combined immunodeficiency
disease: contribution
[140696]of genotype to phenotype. J. Clin. Invest. 92: 2291-2302,
1993.
[140697]
[140698]100. Schmalstieg, F. C.; Mills, G. C.; Tsuda, H.; Goldman,
A. S.:
[140699]Severe combined immunodeficiency in a child with a
healthy adenosine
[140700]deaminase deficient mother. Pediat. Res. 17: 935-940,
1983.
[140701]
[140702]101. Schrader, W. P.; Pollara, B.; Meuwissen, H. J.: Charac-
terization
[140703]of the residual adenosine deaminating activity in the
spleen of a

[140704]patient with combined immunodeficiency disease and adenosine deaminase

[140705]deficiency. Proc. Nat. Acad. Sci. 75: 446–450, 1978.

[140706]

[140707]102. Scott, C. R.; Chen, S.–H.; Giblett, E. R.: Deletion of the carrier

[140708]state in combined immunodeficiency disease associated with deaminase

[140709]deficiency. J. Clin. Invest. 53: 1194–1196, 1974.

[140710]

[140711]103. Shovlin, C. L.; Hughes, J. M. B.; Simmonds, H. A.; Fairbanks,

[140712]L.; Deacock, S.; Lechler, R.; Roberts, I.; Webster, A. D. B.: Adult

[140713]presentation of adenosine deaminase deficiency. (Letter) Lancet 341:

[140714]1471, 1993.

[140715]

[140716]104. Shovlin, C. L.; Simmonds, H. A.; Fairbanks, L. D.; Deacock, S.

[140717]J.; Hughes, J. M. B.; Lechler, R. I.; Webster, A. D. B.; Sun,

X.-M.;

[140718]Webb, J. C.; Soutar, A. K.: Adult onset immunodeficiency
caused by
[140719]inherited adenosine deaminase deficiency. J. Immun. 153:
2331-2339,
[140720]1994.

[140721]

[140722]105. Spencer, N.; Hopkinson, D. A.; Harris, H.: Adenosine
deaminase
[140723]polymorphism in man. Ann. Hum. Genet. 32: 9-14, 1968.

[140724]

[140725]106. Stephan, V.; Wahn, V.; Le Deist, F.; Dirksen, U.; Bro-
ker, B.;
[140726]Muller-Fleckenstein, I.; Horneff, G.; Schroten, H.; Fischer,
A.; de

[140727]Saint Basile, G.: Atypical X-linked severe combined im-
munodeficiency
[140728]due to possible spontaneous reversion of the genetic de-
fect in T cells. New
[140729]Eng. J. Med. 335: 1563-1567, 1996.

[140730]

[140731]107. Tariverdian, G.; Ritter, H.: Adenosine deaminase polymorphism

[140732](EC 3.5.4.4): formal genetics and linkage relations. Humangenetik 7:

[140733]176–178, 1969.

[140734]

[140735]108. Tischfield, J. A.; Creagan, R. P.; Nichols, E. A.; Ruddle, F.

[140736]H.: Assignment of a gene for adenosine deaminase to human chromosome

[140737]20. Hum. Hered. 24: 1–11, 1974.

[140738]

[140739]109. Tzall, S.; Ellenbogen, A.; Eng, F.; Hirschhorn, R.: Identification

[140740]and characterization of nine RFLPs at the adenosine deaminase (ADA)

[140741]locus. Am. J. Hum. Genet. 44: 864–875, 1989.

[140742]

[140743]110. Umetsu, D. T.; Schlossman, C. M.; Ochs, H. D.; Herschfield, M.

[140744]S.: Heterogeneity of phenotype in two siblings with

adenosine deaminase

[140745]deficiency. J. Allergy Clin. Immun. 93: 543-550, 1994.

[140746]

[140747]111. Valerio, D.; Dekker, B. M. M.; Duyvesteyn, M. G. C.;
van der

[140748]Voorn, L.; Berkvens, T. M.; van Ormondt, H.; van der Eb, A.
J.: One

[140749]adenosine deaminase allele in a patient with severe com-
bined immunodeficiency

[140750]contains a point mutation abolishing enzyme activity.
EMBO J. 5:

[140751]113-119, 1986.

[140752]

[140753]112. Valerio, D.; Duyvesteyn, M. G. C.; Dekker, B. M. M.;
Weeda, G.;

[140754]Berkvens, T. M.; van der Voorn, L.; van Ormondt, H.; van
der Eb, A.

[140755]J.: Adenosine deaminase: characterization and expression
of a gene

[140756]with a remarkable promoter. EMBO J. 4: 437-443, 1985.

[140757]

- [140758]113. Valerio, D.; Duyvesteyn, M. G. C.; Meera Khan, P.; Pearson, P.
- [140759]L.; Geurts van Kessel, A.; van Ormondt, H.: Direct assignment of
- [140760]ADA gene to chromosome 20. (Abstract) Cytogenet. Cell Genet. 37:
- [140761]599, 1984.
- [140762]
- [140763]114. Valerio, D.; Duyvesteyn, M. G. C.; van Ormondt, H.; Meera Khan,
- [140764]P.; van der Eb, A. J.: Adenosine deaminase (ADA) deficiency in cells
- [140765]derived from humans with severe combined immunodeficiency is due to
- [140766]an aberration of the ADA protein. Nucleic Acids Res. 12: 1015-1024,
- [140767]1984.
- [140768]
- [140769]115. Valerio, D.; McIvor, R. S.; Williams, S. R.; Duyvesteyn, M. G.
- [140770]C.; van Ormondt, H.; van der Eb, A. J.; Martin, D. W., Jr.: Cloning

[140771] of human adenosine deaminase cDNA and expression in mouse cells. Gene 31:

[140772] 147–153, 1984.

[140773]

[140774] 116. Van der Weyden, M. B.; Kelley, W. N.: Adenosine deaminase deficiency

[140775] in severe combined immunodeficiency: evidence for a posttranslational

[140776] defect. (Abstract) J. Clin. Invest. 53: 81A–82A, 1974.

[140777]

[140778] 117. Wakamiya, M.; Blackburn, M. R.; Jurecic, R.; McArthur, M. J.;

[140779] Geske, R. S.; Cartwright, J., Jr.; Mitani, K.; Vaishnav, S.; Belmont,

[140780] J. W.; Kellems, R. E.; Finegold, M. J.; Montgomery, Jr., C. A.; Bradley,

[140781] A.; Caskey, C. T.: Disruption of the adenosine deaminase gene causes

[140782] hepatocellular impairment and perinatal lethality in mice. Proc.

[140783] Nat. Acad. Sci. 92: 3673–3677, 1995.

[140784]

[140785]118. Weitkamp, L. R.: Further data on the genetic linkage relations

[140786]of the adenosine deaminase locus. Hum. Hered. 21: 351-356, 1971.

[140787]

[140788]119. Weitkamp, L. R.: Genetic linkage relationships of the ADA and

[140789]6-PGD loci in 'Humangenetik.' (Letter) Humangenetik 15: 359-360,

[140790]1972.

[140791]

[140792]120. Wiginton, D. A.; Adrian, G. S.; Friedman, R. L.; Suttle, D. P.;

[140793]Hutton, J. J.: Cloning of cDNA sequences of human adenosine deaminase. Proc.

[140794]Nat. Acad. Sci. 80: 7481-7485, 1983.

[140795]

[140796]121. Wiginton, D. A.; Adrian, G. S.; Hutton, J. J.: Sequence of human

[140797]adenosine deaminase cDNA including the coding region

and a small intron. Nucleic

[140798]Acids Res. 12: 2439–2446, 1984.

[140799]

[140800]122. Wiginton, D. A.; Hutton, J. J.: Immunoreactive protein
in adenosine

[140801]deaminase deficient human lymphoblast cell lines. J. Biol.
Chem. 257:

[140802]3211–3217, 1982.

[140803]

[140804]123. Wiginton, D. A.; Kaplan, D. J.; States, J. C.; Akeson, A.
L.;

[140805]Perme, C. M.; Bilyk, I. J.; Vaughn, A. J.; Lattier, D. L.; Hut-
ton,

[140806]J. J.: Complete sequence and structure of the gene for hu-
man adenosine

[140807]deaminase. Biochemistry 25: 8234–8244, 1986.

[140808]

[140809]124. Yokoyama, S.; Hayashi, T.; Yoshimura, Y.; Irimada,
K.; Saito,

[140810]T.; Akiba, T.; Tsuchiya, S.: Severe combined immunodeficiency disease

[140811]with adenosine deaminase deficiency. Tohoku J. Exp. Med.
129: 197–202,

[140812]1979.

[140813]

[140814]125. Yount, J.; Nichols, P.; Ochs, H. D.; Hammar, S. P.;
Scott, C.

[140815]R.; Chen, S.–H.; Giblett, E. R.; Wedgwood, R. J.: Absence of
erythrocyte

[140816]adenosine deaminase associated with severe combined
immunodeficiency. J.

[140817]Pediat. 84: 173–177, 1974.

[140818]

[140819]126. Ziegler, J. B.; Lee, C. H.; Van Der Weyden, M. B.; Bag-
nara, A.

[140820]S.; Beveridge, J.: Severe combined immunodeficiency and
adenosine

[140821]deaminase deficiency: failure of enzyme replacement
therapy. Arch.

[140822]Dis. Child. 55: 452–457, 1980.

[140823]

[140824]127. Ziegler, J. B.; Van Der Weyden, M. B.; Lee, C. H.;

Daniel, A.

[140825] Prenatal diagnosis for adenosine deaminase deficiency. J. Med.

[140826] Genet. 18: 154–156, 1981.

[140827] 25157.

[140828] Dooley, T. P.: Personal Communication. San Antonio, Tex. 7/21/1992.

[140829] 25158.

[140830] Dooley, T. P.; Weiland, K. L.; Simon, M.: cDNA sequence of human

[140831] p11 calpactin I light chain. Genomics 13: 866–868, 1992.

[140832] 25159.

[140833] Harder, T.; Kube, E.; Gerke, V.: Cloning and characterization

[140834] of the human gene encoding p11: structural similarity to other members

[140835] of the S-100 gene family. Gene 113: 269–274, 1992.

[140836] 25160.

[140837] Kube, E.; Weber, K.; Gerke, V.: Primary structure of human, chicken,

[140838] and *Xenopus laevis* p11, a cellular ligand of the Src-kinase

substrate,

[140839]annexin II. Gene 102: 255–259, 1991.

[140840]25161.

[140841]Okuse, K.; Malik–Hall, M.; Baker, M. D.; Poon, W.–Y. L.;
Kong,

[140842]H.; Chao, M. V.; Wood, J. N.: Annexin II light chain regu-
lates sensory

[140843]neuron–specific sodium channel expression. Nature 417:
653–656,

[140844]2002.

[140845]25162.

[140846]Saris, C. J. M.; Kristensen, T.; D'Eustachio, P.; Hicks, L. J.;

[140847]Noonan, D. J.; Hunter, T.; Tack, B. F.: cDNA sequence and
tissue

[140848]distribution of the mRNA for bovine and murine p11, the
S100–related

[140849]light chain of the protein–tyrosine kinase substrate p36
(calpactin

[140850]l). J. Biol. Chem. 262: 10663–10671, 1987.

[140851]25163.

[140852]Eikenboom, J. C. J.; Vink, T.; Briet, E.; Sixma, J. J.; Reitsma,

[140853]P. H.: Multiple substitutions in the von Willebrand factor
gene that
[140854]mimic the pseudogene sequence. Proc. Nat. Acad. Sci. 91:
2221–2224,
[140855]1994.
[140856]25164.
[140857]Gralnick, H. R.; Coller, B. S.: Molecular defects in
haemophilia
[140858]A and von Willebrand's disease. Lancet I: 837–838, 1976.
[140859]25165.
[140860]Watanabe, H.; Orii, K. E.; Fukao, T.; Song, X.–Q.; Aoyama,
T.;
[140861]Jlst, L.; Ruiter, J.; Wanders, R. J. A.; Kondo, N.: Molecular
basis
[140862]of very long chain acyl–CoA dehydrogenase deficiency in
three Israeli
[140863]patients: identification of a complex mutant allele with
P65L and
[140864]K247Q mutations, the former being an exonic mutation
causing exon
[140865]3 skipping. Hum. Mutat. 15: 430–438, 2000.

[140866]25166.

[140867]Gavish, H.; dos Santos, C. C.; Buchwald, M.: Generation of a non-functional

[140868]Fanconi anemia group C protein (FACC) by site-directed in vitro mutagenesis.

[140869](Abstract) Am. J. Hum. Genet. 51 (suppl.): A128 only, 1992.

[140870]25167.

[140871]Gavish, H.; dos Santos, C. C.; Buchwald, M.: A leu554-to-pro substitution

[140872]completely abolishes the functional complementing activity of the

[140873]Fanconi anemia (FACC) protein. Hum. Molec. Genet. 2: 123-126, 1993.

[140874]25168.

[140875]Ahmad, W.; Haque, M. F.; Brancolini, V.; Tsou, H. C.; Haque, S.;

[140876]Lam, H.; Aita, V. M.; Owen, J.; deBlaquiere, M.; Frank, J.; Cserhalmi-Friedman,

[140877]P. B.; Leask, A.; McGrath, J. A.; Peacocke, M.; Ahmad, M.; Ott, J.;

[140878]Christiano, A. M.: Alopecia universalis associated with a

mutation

[140879]in the human hairless gene. Science 279: 720–724, 1998.

[140880]25169.

[140881]Matthijs, G.; Schollen, E.; Heykants, L.; Grunewald, S.:

Phosphomannomutase

[140882]deficiency: the molecular basis of the classical Jaeken syndrome (CDGS

[140883]type Ia). Molec. Genet. Metab. 68: 220–226, 1999.

[140884]25170.

[140885]Matthijs, G.; Schollen, E.; Pardon, E.; Veiga-Da-Cunha, M.;

Jaeken,

[140886]J.; Cassiman, J.-J.; Van Schaftingen, E.: Mutations in PMM2, a phosphomannomutase

[140887]gene on chromosome 16p13, in carbohydrate-deficient glycoprotein type

[140888]I syndrome (Jaeken syndrome). Nature Genet. 16: 88–92, 1997.

[140889]25171.

[140890]Matthijs, G.; Schollen, E.; Pirard, M.; Budarf, M. L.; Van Schaftingen,

[140891]E.; Cassiman, J.-J.: PMM (PMM1), the human homologue of

SEC53 or

[140892] yeast phosphomannomutase, is localized on chromosome
22q13. Genomics 40:

[140893] 41–47, 1997.

[140894] 25172.

[140895] Broeks, A.; de Klein, A.; Floore, A. N.; Muijtjens, M.; Klei-
jer,

[140896] W. J.; Jaspers, N. G.; van't Veer, L. J.: ATM germline muta-
tions

[140897] in classical ataxia–telangiectasia patients in the Dutch
population. Hum.

[140898] Mutat. 12: 330–337, 1998.

[140899] 25173.

[140900] Munoz, F.; Lestringant, G.; Sybert, V.; Frydman, M.; Al-
swaini,

[140901] A.; Frossard, P. M.; Jorgenson, R.; Zonana, J.: Definitive
evidence

[140902] for an autosomal recessive form of hypohidrotic ectoder-
mal dysplasia

[140903] clinically indistinguishable from the more common X-
linked disorder. Am.

[140904] J. Hum. Genet. 61: 94–100, 1997.

[140905]25174.

[140906]Van Schaftingen, E.; Jaeken, J.: Phosphomannomutase deficiency

[140907]is a cause of carbohydrate-deficient glycoprotein syndrome type I. FEBS

[140908]Lett. 377: 318-320, 1995.

[140909]25175.

[140910]Carritt, B.: Somatic cell genetic evidence for the presence of

[140911]a gene for citrullinemia on human chromosome 9. Cytogenet. Cell Genet. 19:

[140912]44-48, 1977.

[140913]25176.

[140914]Carritt, B.; Goldfarb, P. S. G.; Hooper, M. L.; Slack, C.: Chromosome

[140915]assignment of a human gene for argininosuccinate synthetase expression

[140916]in Chinese hamster-human somatic cell hybrids. Exp. Cell Res. 106:

[140917]71-78, 1977.

[140918]25177.

[140919]Daiger, S. P.; Hoffman, N. S.; Wildin, R. S.; Su, T.-S.: Multi-
ple,
[140920]independent restriction site polymorphisms in human
DNA detected with
[140921]a cDNA probe to argininosuccinate synthetase (AS). Am. J.
Hum. Genet. 36:
[140922]736-749, 1984.
[140923]25178.
[140924]Daiger, S. P.; Wildin, R. S.; Su, T.-S.: Polymorphic variants
[140925]of restriction fragments of human DNA detected with a
probe to argininosuccinate
[140926]synthetase. (Abstract) Am. J. Hum. Genet. 33: 136A only,
1981.
[140927]25179.
[140928]Gripp, K. W.; Stolle, C. A.; Celle, L.; McDonald-McGinn, D.
M.;
[140929]Whitaker, L. A.; Zackai, E. H.: TWIST gene mutation in a
patient
[140930]with radial aplasia and craniosynostosis: further evidence
for heterogeneity
[140931]of Baller-Gerold syndrome. Am. J. Med. Genet. 82:
170-176, 1999.

[140932]25180.

[140933]Seto, M. L.; Lee, S. J.; Sze, R. W.; Cunningham, M. L.: Another

[140934]TWIST on Baller-Gerold syndrome. Am. J. Med. Genet.
104: 323-330,

[140935]2001.

[140936]25181.

[140937]Karathanasis, S. K.: Apolipoprotein multigene family: tandem organization

[140938]of human apolipoprotein AI, CIII, and AIV genes. Proc. Nat.
Acad.

[140939]Sci. 82: 6374-6378, 1985.

[140940]25182.

[140941]Bailey, R.; Miller, D. A.: Mapping the human apolipoprotein
genes

[140942]CII and E to band 19q13.1 by in situ hybridization.
(Abstract) Am.

[140943]J. Hum. Genet. 41: A156 only, 1987.

[140944]25183.

[140945]Bartres-Faz, D.; Clemente, I. C.; Junque, C.; Valveny, N.;
Lopez-Alomar,

[140946]A.; Sanchez-Aldeguer, J.; Lopez-Guillen, A.; Moral, P.:
APOE and
[140947]APOC1 genetic polymorphisms in age-associated memory
impairment. Neurogenetics 3:
[140948]215–219, 2001.
[140949]25184.
[140950]Crook, T. H.; Bartus, R. T.; Ferris, S. H.; Whitehouse, P.;
Cohen,
[140951]G. D.; Gershon, S.: Age-associated memory impairment:
proposed diagnostic
[140952]criteria and measures of clinical change. Report of a Na-
tional Institute
[140953]of Mental Health work group. Dev. Neuropsychol. 2:
261–276, 1986.
[140954]25185.
[140955]Davison, P. J.; Norton, P.; Wallis, S. C.; Gill, L.; Cook, M.;
[140956]Williamson, R.; Humphries, S. E.: There are two gene se-
quences for
[140957]human apolipoprotein CI (APO CI) on chromosome 19,
one of which is
[140958]4 kb from the gene for APO E. Biochem. Biophys. Res.
Commun. 136:

[140959]876–884, 1986.

[140960]25186.

[140961]Gautier, T.; Masson, D.; Jong, M. C.; Duverneuil, L.; Le
Guern,

[140962]N.; Deckert, V.; Pais de Barros, J.–P.; Dumont, L.; Bataille,
A.;

[140963]Zak, Z.; Jiang, X.–C.; Tall, A. R.; Havekes, L. M.; Lagrost, L.:

[140964]Apolipoprotein CI deficiency markedly augments plasma
lipoprotein

[140965]changes mediated by human cholesteryl ester transfer
protein (CETP)

[140966]in CETP transgenic/ApoCI–knocked out mice. J. Biol.
Chem. 277: 31354–31363,

[140967]2002.

[140968]25187.

[140969]Lauer, S. J.; Walker, D.; Elshourbagy, N. A.; Reardon, C. A.;
Levy–Wilson,

[140970]B.; Taylor, J. M.: Two copies of the human apolipoprotein
C–I gene

[140971]are linked closely to the apolipoprotein E gene. J. Biol.
Chem. 263:

[140972]7277-7286, 1988.

[140973]25188.

[140974]Myklebost, O.; Rogne, S.: The gene for human apolipoprotein CII

[140975]is located 4.3 kilobases away from the apolipoprotein E gene on chromosome

[140976]19. Hum. Genet. 73: 286-289, 1986.

[140977]25189.

[140978]Myklebost, O.; Rogne, S.: A physical map of the apolipoprotein

[140979]gene cluster on human chromosome 19. Hum. Genet. 78: 244-247, 1988.

[140980]25190.

[140981]Richards, A. J.; Baguley, D. M.; Yates, J. R. W.; Lane, C.; Nicol,

[140982]M.; Harper, P. S.; Scott, J. D.; Snead, M. P.: Variation in the vitreous

[140983]phenotype of Stickler syndrome can be caused by different amino acid

[140984]substitutions in the X position of the type II collagen Gly-X-Y triple

[140985]helix. Am. J. Hum. Genet. 67: 1083–1094, 2000.

[140986]25191.

[140987]Stickler, G. B.; Belau, P. G.; Farrell, F. J.; Jones, J. D.; Pugh,

[140988]D. G.; Steinberg, A. G.; Ward, L. E.: Hereditary progressive
arthro–ophthalmopathy. Mayo

[140989]Clin. Proc. 40: 433–455, 1965.

[140990]25192.

[140991]Wilkin, D. J.; Liberfarb, R.; Davis, J.; Levy, H. P.; Cole, W.

[140992]G.; Francomano, C. A.; Cohn, D. H.: Rapid determination
of COL2A1

[140993]mutations in individuals with Stickler syndrome: analysis
of potential

[140994]premature termination codons. Am. J. Med. Genet. 94:
141–148, 2000.

[140995]25193.

[140996]Williams, C. J.; Ganguly, A.; Considine, E.; McCarron, S.;
Prockop,

[140997]D. J.; Walsh–Vockley, C.; Michels, V. V.: A(–2)–to–G transi-
tion at

[140998]the 3–prime acceptor splice site of IVS17 characterizes the
COL2A1

[140999]gene mutation in the original Stickler syndrome kindred.
Am. J. Med.

[141000]Genet. 63: 461–467, 1996.

[141001]25194.

[141002]Zlotogora, J.; Sagi, M.; Schuper, A.; Leiba, H.; Merin, S.:
Variability

[141003]of Stickler syndrome. Am. J. Med. Genet. 42: 337–339,
1992.

[141004]25195.

[141005]Brown, S.; Wiebel, F. J.; Gelboin, H. V.; Minna, J. D.: As-
signment

[141006]of a locus required for flavoprotein–linked monooxyge-
nase expression

[141007]to human chromosome 2. Proc. Nat. Acad. Sci. 73:
4628–4632, 1976.

[141008]25196.

[141009]Chen, Y. T.; Tukey, R. H.; Swan, D. C.; Negishi, N.; Nebert,
D.

[141010]W.: Characterization of the human P1–450 genomic gene.
(Abstract) Clin.

[141011]Res. 31: 456A, 1983.

[141012]25197.

[141013]Corchero, J.; Pimprale, S.; Kimura, S.; Gonzalez, F. J.: Organization

[141014]of the CYP1A cluster on human chromosome 15: implications for gene

[141015]regulation. Pharmacogenetics 11: 1–6, 2001.

[141016]25198.

[141017]Gorman, C.; Padmanabhan, R.; Howard, B. H.: High efficiency DNA-mediated

[141018]transformation of primate cells. Science 221: 551–553, 1983.

[141019]25199.

[141020]Gorman, C. M.: CAT: an easy assay for gene expression (citation

[141021]classic). Current Contents (Life Sciences) 36(22): 8, 1993.

[141022]25200.

[141023]Hildebrand, C. E.; Gonzalez, F. J.; Kozak, C. A.; Nebert, D. W.

[141024]: Regional linkage analysis of the dioxin-inducible P-450 gene family

[141025]on mouse chromosome 9. Biochem. Biophys. Res. Com-

mun. 130: 396–406,

[141026]1985.

[141027]25201.

[141028]Lee, S.; Wu, X.; Reid, M. E.; Zelinski, T.; Redman, C. M.:

Molecular

[141029]basis of the Kell (K1) phenotype. Blood 85: 912–916,

1995.

[141030]25202.

[141031]Lee, S.; Zambas, E. D.; Marsh, W. L.; Redman, C. M.: The
human

[141032]Kell blood group gene maps to chromosome 7q33 and its
expression is

[141033]restricted to erythroid cells. Blood 81: 2804–2809, 1993.

[141034]25203.

[141035]Lee, S.; Zambas, E. D.; Marsh, W. L.; Redman, C. M.:

Molecular

[141036]cloning and primary structure of Kell blood group protein.
Proc.

[141037]Nat. Acad. Sci. 88: 6353–6357, 1991.

[141038]25204.

[141039]Marsh, W. L.: Molecular biology of blood groups: cloning

of the

[141040]Kell gene. Transfusion 32: 98–101, 1992.

[141041]25205.

[141042]Morton, N. E.; Krieger, H.; Steinberg, A. G.; Rosenfield, R.
E.

[141043]: Genetic evidence confirming the localization of Sutter in
the Kell

[141044]blood-group system. Vox Sang. 10: 608–613, 1965.

[141045]25206.

[141046]Murphy, M. T.; Morrison, N.; Miles, J. S.; Fraser, R. H.;
Spurr,

[141047]N. K.; Boyd, E.: Regional chromosomal assignment of the
Kell blood

[141048]group locus (KEL) to chromosome 7q33–q35 by fluores-
cence in situ hybridization:

[141049]evidence for the polypeptide nature of antigenic variation.
Hum.

[141050]Genet. 91: 585–588, 1993.

[141051]25207.

[141052]Parke, J. T.; Riccardi, V. M.; Lewis, R. A.; Ferrell, R. E.:

[141053]A syndrome of microcephaly and retinal pigmentary ab-

normalities without

[141054]mental retardation in a family with coincidental autosomal dominant

[141055]hyperreflexia. Am. J. Med. Genet. 17: 585-594, 1984.

[141056]25208.

[141057]Purohit, K. R.; Weber, J. L.; Ward, L. J.; Keats, B. J. B.: The

[141058]Kell blood group locus is close to the cystic fibrosis locus on chromosome

[141059]7. Hum. Genet. 89: 457-458, 1992.

[141060]25209.

[141061]Russo, D.; Wu, X.; Redman, C. M.; Lee, S.: Expression of Kell

[141062]blood group protein in nonerythroid tissues Blood 96: 340-346, 2000.

[141063]25210.

[141064]Stroup, M.; MacIlroy, M.; Walker, R.; Aydelotte, J. V.: Evidence

[141065]that Sutter belongs to the Kell blood group system. Transfusion 5:

[141066]309-314, 1965.

[141067]25211.

[141068]Wagner, T.; Berer, A.; Lanzer, G.; Geissler, K.: Kell is not
[141069]restricted to the erythropoietic lineage but is also ex-
pressed on
[141070]myeloid progenitor cells. Brit. J. Haemat. 110: 409–411,
2000.
[141071]25212.

[141072]Weitkamp, L. R.; Townes, P. L.; Johnston, E.: Linkage data
on
[141073]urinary pepsinogen and the Kell blood group. Birth De-
fects Orig.
[141074]Art. Ser. 11(3): 281–282, 1975. Note: Alternate: Cyto-
genet. Cell
[141075]Genet. 14: 451–452, 1975.
[141076]25213.

[141077]Wimer, B. M.; Marsh, W. L.; Taswell, H. F.: Clinical charac-
teristics
[141078]of the McLeod blood group phenotype. (Abstract) 19th
Annual Meeting
[141079]of the Am. Soc. Hemat., Boston , 1976.
[141080]25214.

[141081]Yu, L.-C.; Twu, Y.-C.; Chang, C.-Y.; Lin, M.: Molecular ba-

sis

[141082]of the Kell-null phenotype: a mutation at the splice site of human

[141083]KEL gene abolishes the expression of Kell blood group antigens. J.

[141084]Biol. Chem. 276: 10247-10252, 2001.

[141085]25215.

[141086]Zelinski, T. A.; Coghlan, G. E.; Myal, Y.; White, L. J.; Philipps,

[141087]S. E.: Assignment of the Kell blood group locus to chromosome 7q.

[141088](Abstract) Cytogenet. Cell Genet. 58: 1927 only, 1991.

[141089]25216.

[141090]Field, L. L.; Marazita, M. L.; Spence, M. A.; Crandall, B. F.;

[141091]Sparkes, R. S.: Is JK linked to IGK on chromosome 2? (Abstract) Cytogenet.

[141092]Cell Genet. 40: 628-629, 1985.

[141093]25217.

[141094]Gedde-Dahl, T.: Personal Communication. Oslo, Norway 9/26/1986.

[141095]25218.

[141096]Geitvik, G. A.; Hoyheim, B.; Gedde-Dahl, T.; Grzeschik, K.
H.;

[141097]Lothe, R.; Tomter, H.; Olaisen, B.: The Kidd (JK) blood
group locus

[141098]assigned to chromosome 18 by close linkage to a DNA-
RFLP. Hum. Genet. 77:

[141099]205-209, 1987.

[141100]25219.

[141101]Hulten, M.; Lindsten, J.; Pen-Ming, L. M.; Fraccaro, M.;
Mannini,

[141102]A.; Trepolo, L.; Robson, E. B.; Heiken, A.; Tellingén, K. G.:
Possible

[141103]localization of the genes for the Kidd blood group on an
autosome

[141104]involved in a reciprocal translocation. Nature 211:
1067-1068, 1968.

[141105]25220.

[141106]Leppert, M.; Ferrell, R.; Kamboh, M. I.; Beasley, J.;
O'Connell,

[141107]P.; Lathrop, M.; Lalouel, J.-M.; White, R.: Linkage of the
polymorphic

[141108]protein markers F13B, C1S, C1R, and blood group antigen

Kidd in CEPH

[141109]reference families. (Abstract) Cytogenet. Cell Genet. 46:
647, 1987.

[141110]25221.

[141111]Lucien, N.; Chiaroni, J.; Cartron, J.-P.; Bailly, P.: Partial

[141112]deletion in the JK locus causing a Jk(null) phenotype.

Blood 99:

[141113]1079–1081, 2002.

[141114]25222.

[141115]Lucien, N.; Sidoux-Walter, F.; Olives, B.; Moulds, J.; Le

Pennec,

[141116]P.-Y.; Cartron, J.-P.; Bailly, P.: Characterization of the gene

encoding

[141117]the human Kidd blood group/urea transporter protein:

evidence for

[141118]splice site mutations in Jk-null individuals. J. Biol. Chem.

273:

[141119]12973–12980, 1998.

[141120]25223.

[141121]Olives, B.; Mattei, M.-G.; Huet, M.; Neau, P.; Martial, S.;

Cartron,

- [141122]J.-P.; Bailly, P.: Kidd blood group and urea transport function of
- [141123]human erythrocytes are carried by the same protein. J. Biol. Chem. 270:
- [141124]15607–15610, 1995.
- [141125]25224.
- [141126]Albertsen, H.; Plaetke, R.; Ballard, L.; Fujimoto, E.; Connolly,
- [141127]J.; Lawrence, E.; Rodriguez, P.; Robertson, M.; Bradley, P.; Milner,
- [141128]B.; Fuhrman, D.; Marks, A.; Sargent, R.; Cartwright, P.; Matsunami,
- [141129]N.; White, R.: Genetic mapping of the BRCA1 region on chromosome
- [141130]17q21. Am. J. Hum. Genet. 54: 516–525, 1994.
- [141131]25225.
- [141132]Andersen, T. I.; Borresen, A.-L.; Moller, P.: A common BRCA1 mutation
- [141133]in Norwegian breast and ovarian cancer families? Am. J. Hum. Genet. 59:
- [141134]486–487, 1996.

[141135]25226.

[141136]Anderson, S. F.; Schlegel, B. P.; Nakajima, T.; Wolpin, E. S.;

[141137]Parvin, J. D.: BRCA1 protein is linked to the RNA polymerase II holoenzyme

[141138]complex via RNA helicase A. *Nature Genet.* 19: 254–256, 1998.

[141139]25227.

[141140]Arason, A.; Barkardottir, R. B.; Egilsson, V.: Linkage analysis

[141141]of chromosome 17q markers and breast–ovarian cancer in Icelandic families,

[141142]and possible relationship to prostatic cancer. *Am. J. Hum. Genet.* 52:

[141143]711–717, 1993.

[141144]25228.

[141145]Womack, J. E.: Personal Communication. College Station, Texas

[141146]2/26/1990.

[141147]25229.

[141148]Hiraoka, L. R.; Hsu, L.; Hsieh, C.–L.: Assignment of ALDH3 to

[141149]human chromosome 17p11.2 and ALDH5 to human chromosome 9p13. Genomics 25:

[141150]323–325, 1995.

[141151]25230.

[141152]Vasiliou, V.; Bairoch, A.; Tipton, K. F.; Nebert, D. W.: Eukaryotic

[141153]aldehyde dehydrogenase (ALDH) genes: human polymorphisms, and recommended

[141154]nomenclature based on divergent evolution and chromosomal mapping. Pharmacogenetics 9:

[141155]421–434, 1999.

[141156]25231.

[141157]Harada, S.; Agarwal, D. P.; Goedde, H. W.: Electrophoretic and

[141158]biochemical studies of human aldehyde dehydrogenase isozymes in various

[141159]tissues. Life Sci. 26: 1773–1780, 1980.

[141160]25232.

[141161]Hsu, L. C.; Chang, W.-C.; Yoshida, A.: Cloning of a new human

[141162]aldehyde dehydrogenase gene and comparison with liver

cytosolic ALDH1

[141163]and mitochondrial ALDH2 genes. (Abstract) Am. J. Hum. Genet. 45

[141164](suppl.): A196 only, 1989.

[141165]25233.

[141166]Hsu, L. C.; Chang, W. C.: Cloning and characterization of a new

[141167]functional human aldehyde dehydrogenase gene. J. Biol. Chem. 266:

[141168]12257-12265, 1991.

[141169]25234.

[141170]Ashworth, A.: Two acetyl-CoA acetyltransferase genes located in

[141171]the t-complex region of mouse chromosome 17 partially overlap the

[141172]Tcp-1 and Tcp-1x genes. Genomics 18: 195-198, 1993.

[141173]25235.

[141174]Bennett, M. J.; Hosking, G. P.; Smith, M. F.; Gray, R. G. F.; Middleton,

[141175]B.: Biochemical investigations on a patient with a defect in cytosolic

[141176]acetoacetyl-CoA thiolase, associated with mental retardation. J.

[141177]Inherit. Metab. Dis. 7: 125-128, 1984.

[141178]25236.

[141179]Wunderle, V. M.; Critcher, R.; Hastie, N.; Goodfellow, P. N.;

[141180]Schedl, A.: Deletion of long-range regulatory elements upstream of

[141181]SOX9 causes campomelic dysplasia. Proc. Nat. Acad. Sci. 95: 10649-10654,

[141182]1998.

[141183]25237.

[141184]Young, I. D.; Zuccollo, J. M.; Maltby, E. L.; Broderick, N. J.

[141185]: Campomelic dysplasia associated with a de novo 2q;17q reciprocal

[141186]translocation. J. Med. Genet. 29: 251-252, 1992.

[141187]25238.

[141188]Lynch, H. T.; Schuelke, G. S.; Kimberling, W. J.; Albano, W. A.;

[141189]Lynch, J. F.; Biscione, K. A.; Lipkin, M. L.; Deschner, E. E.; Mikol,

- [141190]Y. B.; Sandberg, A. A.; Elston, R. C.; Bailey-Wilson, J. E.;
Danes,
[141191]B. S.: Hereditary nonpolyposis colorectal cancer (Lynch
syndromes
[141192]I and II). II. Biomarker studies. Cancer 56: 939-951, 1985.
[141193]25239.
[141194]Bourn, D.; Carter, S. A.; Mason, S.; Evans, D. G. R.; Stra-
chan,
[141195]T.: Germline mutations in the neurofibromatosis type 2
tumor suppressor
[141196]gene. Hum. Molec. Genet. 3: 813-816, 1994.
[141197]25240.
[141198]Bouzas, E. A.; Freidlin, V.; Parry, D. M.; Eldridge, R.;
Kaiser-Kupfer,
[141199]M. I.: Lens opacities in neurofibromatosis 2: further sig-
nificant
[141200]correlations. Brit. J. Ophthal. 77: 354-357, 1993.
[141201]25241.
[141202]Bouzas, E. A.; Parry, D. M.; Eldridge, R.; Kaiser-Kupfer, M.
I.
[141203]: Visual impairment in patients with neurofibromatosis 2.

Neurology 43:

[141204]622–623, 1993.

[141205]25242.

[141206]Bruder, C. E. G.; Hirvela, C.; Tapia-Paez, I.; Fransson, I.;
Segraves,

[141207]R.; Hamilton, G.; Zhang, X. X.; Evans, D. G.; Wallace, A. J.;
Baser,

[141208]M. E.; Zucman-Rossi, J.; Hergersberg, M.; and 22 others:
High resolution

[141209]deletion analysis of constitutional DNA from neurofibro-
matosis type

[141210]2 (NF2) patients using microarray-CGH. Hum. Molec.
Genet. 10: 271–282,

[141211]2001.

[141212]25243.

[141213]Claudio, J. O.; Malo, D.; Rouleau, G. A.: The mouse neu-
rofibromatosis

[141214]type 2 gene maps to chromosome 11. Genomics 21:
437–439, 1994.

[141215]25244.

[141216]Egan, R. A.; Thompson, C. R.; MacCollin, M.; Lessell, S.:

Monocular

[141217]levator paresis in neurofibromatosis type 2. Neurology
56: 1222–1224,

[141218]2001.

[141219]25245.

[141220]Eldridge, R.: Central neurofibromatosis with bilateral
acoustic

[141221]neuroma. Adv. Neurol. 29: 57–65, 1981.

[141222]25246.

[141223]Eldridge, R.; Parry, D. M.; Kaiser–Kupfer, M. I.: Neurofibro-
matosis

[141224]2 (NF2): clinical heterogeneity and natural history based
on 39 individuals

[141225]in 9 families and 16 sporadic cases. (Abstract) Am. J. Hum.
Genet. 49

[141226](suppl.): 133, 1991.

[141227]25247.

[141228]Evans, D. G. R.; Birch, J. M.; Ramsden, R. T.: Paediatric
presentation

[141229]of type 2 neurofibromatosis. Arch. Dis. Child. 81:
496–499, 1999.

[141230]25248.

[141231]Evans, D. G. R.; Bourn, D.; Wallace, A.; Ramsden, R. T.;
Mitchell,

[141232]J. D.; Strachan, T.: Diagnostic issues in a family with late
onset

[141233]type 2 neurofibromatosis. J. Med. Genet. 32: 470-474,
1995.

[141234]25249.

[141235]Evans, D. G. R.; Huson, S. M.; Donnai, D.; Neary, W.; Blair,
V.;

[141236]Newton, V.; Strachan, T.; Harris, R.: A genetic study of
type 2 neurofibromatosis

[141237]in the United Kingdom. II. Guidelines for genetic coun-
selling. J.

[141238]Med. Genet. 29: 847-852, 1992.

[141239]25250.

[141240]Evans, D. G. R.; Huson, S. M.; Donnai, D.; Neary, W.; Blair,
V.;

[141241]Teare, D.; Newton, V.; Strachan, T.; Ramsden, R.; Harris,
R.: A genetic

[141242]study of type 2 neurofibromatosis in the United Kingdom.
I. Prevalence,

[141243]mutation rate, fitness, and confirmation of maternal transmission

[141244]effect on severity. J. Med. Genet. 29: 841–846, 1992.

[141245]25251.

[141246]Evans, D. G. R.; Trueman, L.; Wallace, A.; Collins, S.; Strachan,

[141247]T.: Genotype/phenotype correlations in type 2 neurofibromatosis (NF2):

[141248]evidence for more severe disease associated with truncating mutations. J.

[141249]Med. Genet. 35: 450–455, 1998.

[141250]25252.

[141251]Evans, D. G. R.; Wallace, A. J.; Wu, C. L.; Trueman, L.; Ramsden,

[141252]R. T.; Strachan, T.: Somatic mosaicism: a common cause of classic

[141253]disease in tumor-prone syndromes? Lessons from type 2 neurofibromatosis. Am.

[141254]J. Hum. Genet. 63: 727–736, 1998.

[141255]25253.

[141256]Fabricant, R. N.; Todaro, G. J.; Eldridge, R.: Increased lev-

els

[141257]of a nerve-growth factor cross-reacting protein in 'central'
neurofibromatosis. Lancet I:

[141258]4-7, 1979.

[141259]25254.

[141260]Feiling, A.; Ward, E.: A familial form of acoustic tumour.
Brit.

[141261]Med. J. 1: 496-497, 1920.

[141262]25255.

[141263]Fernandez-Valle, C.; Tang, Y.; Ricard, J.; Rodenas-Ruano,
A.;

[141264]Taylor, A.; Hackler, E.; Biggerstaff, J.; Iacovelli, J.: Paxillin

[141265]binds schwannomin and regulates its density-dependent
localization

[141266]and effect on cell morphology. Nature Genet. 31:
354-362, 2002.

[141267]25256.

[141268]Gardner, W. J.; Frazier, C. H.: Hereditary bilateral acoustic

[141269]tumors. J. Hered. 22: 7-8, 1933.

[141270]25257.

[141271]Gardner, W. J.; Frazier, C. H.: Bilateral acoustic neurofibromas.

[141272]A clinical study and survey of a family of five generations with bilateral

[141273]deafness in 38 members. Arch. Neurol. Psychiat. 23: 266–302, 1930.

[141274]25258.

[141275]Gardner, W. J.; Turner, O.: Bilateral acoustic neurofibromas:

[141276]further clinical and pathologic data on hereditary deafness and Recklinghausen's

[141277]disease. Arch. Neurol. Psychiat. 44: 76–99, 1940.

[141278]25259.

[141279]Giovannini, M.; Robanus-Maandag, E.; van der Valk, M.; Niwa-Kawakita,

[141280]M.; Abramowski, V.; Goutebroze, L.; Woodruff, J. M.; Berns, A.; Thomas,

[141281]G.: Conditional biallelic Nf2 mutation in the mouse promotes manifestations

[141282]of human neurofibromatosis type 2. Genes Dev. 14: 1617–1630, 2000.

[141283]25260.

[141284]Gutmann, D. H.: The neurofibromatoses: when less is more. Hum.

[141285]Molec. Genet. 10: 747-755, 2001.

[141286]25261.

[141287]Gutmann, D. H.; Haipek, C. A.; Burke, S. P.; Sun, C.-X.;
Scoles,

[141288]D. R.; Pulst, S. M.: The NF2 interactor, hepatocyte growth
factor-regulated

[141289]tyrosine kinase substrate (HRS), associates with merlin in
the 'open'

[141290]conformation and suppresses cell growth and motility.
Hum. Molec.

[141291]Genet. 10: 825-834, 2001.

[141292]25262.

[141293]Gutmann, D. H.; Hirbe, A. C.; Haipek, C. A.: Functional
analysis

[141294]of neurofibromatosis 2 (NF2) missense mutations. Hum.
Molec. Genet. 10:

[141295]1519-1529, 2001.

[141296]25263.

[141297]Gutmann, D. H.; Sherman, L.; Seftor, L.; Haipek, C.; Lu, K.

H.;

[141298]Hendrix, M.: Increased expression of the NF2 tumor suppressor gene

[141299]product, merlin, impairs cell motility, adhesion and spreading. Hum.

[141300]Molec. Genet. 8: 267-275, 1999.

[141301]25264.

[141302]Jacoby, L. B.; Jones, D.; Davis, K.; Kronn, D.; Short, M. P.;

[141303]Gusella, J.; MacCollin, M.: Molecular analysis of the NF2 tumor-suppressor

[141304]gene in schwannomatosis. Am. J. Hum. Genet. 61: 1293-1302, 1997.

[141305]25265.

[141306]Kaiser-Kupfer, M. I.; Freidlin, V.; Datiles, M. B.; Edwards, P.

[141307]A.; Sherman, J. L.; Parry, D.; McCain, L. M.; Eldridge, R.:

The association

[141308]of posterior capsular lens opacities with bilateral acoustic neuromas

[141309]in patients with neurofibromatosis type 2. Arch. Ophthalmol. 107: 541-544,

[141310]1989.

[141311]25266.

[141312]Kanter, W. R.; Eldridge, R.; Fabricant, R.; Allen, J. C.; Koerber,

[141313]T.: Central neurofibromatosis with bilateral acoustic neuroma: genetic,

[141314]clinical and biochemical distinctions from peripheral neurofibromatosis. Neurology 30:

[141315]851–859, 1980.

[141316]25267.

[141317]Kehrer-Sawatzki, H.; Udart, M.; Krone, W.; Baden, R.; Fahsold,

[141318]R.; Thomas, G.; Schmucker, B.; Assum, G.: Mutational analysis and

[141319]expression studies of the neurofibromatosis type 2 (NF2) gene in a

[141320]patient with a ring chromosome 22 and NF2. Hum. Genet. 100: 67–74,

[141321]1997.

[141322]25268.

[141323]Kedes, L.; Ng, S.-Y.; Lin, C.-S.; Gunning, P.; Eddy, R.; Shows,

[141324]T.; Leavitt, J.: The human beta-actin multigene family.
Trans. Assoc.

[141325]Am. Phys. 98: 42-46, 1985.

[141326]25269.

[141327]Leavitt, J.; Bushar, G.; Kakunaga, T.; Hamada, H.; Hirakawa, T.;

[141328]Goldman, D.; Merrill, C.: Variations in expression of mutant beta-actin

[141329]accompanying incremental increases in human fibroblast tumorigenicity. Cell 28:

[141330]259-268, 1982.

[141331]25270.

[141332]Nakajima-Iijima, S.; Hamada, H.; Reddy, P.; Kakunaga, T.:
Molecular

[141333]structure of the human cytoplasmic beta-actin gene; interspecies homology

[141334]of sequences in the introns. Proc. Nat. Acad. Sci. 82:
6133-6137,

[141335]1985.

[141336]25271.

[141337]Ng, S.-Y.; Gunning, P.; Eddy, R.; Ponte, P.; Leavitt, J.;

Kedes,

[141338]L.; Shows, T.: Chromosome 7 assignment of the human
beta-actin functional

[141339]gene (ACTB) and the chromosomal dispersion of pseudo-
genes. (Abstract) Cytogenet.

[141340]Cell Genet. 40: 712 only, 1985.

[141341]25272.

[141342]Ng, S.-Y.; Gunning, P.; Eddy, R.; Ponte, P.; Leavitt, J.;
Shows,

[141343]T.; Kedes, L.: Evolution of the functional human beta-actin
gene

[141344]and its multi-pseudogene family: conservation of the
noncoding regions

[141345]and chromosomal dispersion of pseudogenes. Molec. Cell.
Biol. 5:

[141346]2720-2732, 1985.

[141347]25273.

[141348]Toyama, S.; Toyama, S.: A variant form of beta-actin in a
mutant

[141349]of KB cells resistant to cytochalasin B. Cell 37: 609-614,
1984.

[141350]25274.

[141351] Vandekerckhove, J.; Weber, K.: Mammalian cytoplasmic
actins are
[141352] the products of at least two genes and differ in primary
structure
[141353] in at least 25 identified positions from skeletal muscle
actins. Proc.
[141354] Nat. Acad. Sci. 75: 1106–1110, 1978.
[141355] 25275.
[141356] Alderdice, P. W.; Kaita, H.; Lewis, M.; McAlpine, P. J.;
Wong,
[141357] P.; Anderson, J.; Giblett, E. R.: Segregation of marker loci
in families
[141358] with an inherited paracentric insertion of chromosome 9.
Am. J. Hum.
[141359] Genet. 39: 612–617, 1986.
[141360] 25276.
[141361] Beutler, E.; Carson, D.; Dannawi, H.; Forman, L.; Kuhl, W.;
West,
[141362] C.; Westwood, B.: Metabolic compensation for profound
erythrocyte
[141363] adenylate kinase deficiency: a hereditary enzyme defect
without hemolytic

[141364]anemia. J. Clin. Invest. 72: 648–655, 1983.

[141365]25277.

[141366]Beutler, E.; Carson, D. A.; Dannawi, H.; Forman, L.; Kuhl, W.;

[141367]West, C.; Westwood, B.: Red cell adenylate kinase deficiency: another

[141368]non-disease? (Abstract) Blood 60: 33A only, 1982.

[141369]25278.

[141370]Bianchi, P.; Zappa, M.; Bredi, E.; Vercellati, C.; Pelissero, G.;

[141371]Barraco, F.; Zanella, A.: A case of complete adenylate kinase deficiency

[141372]due to a nonsense mutation in AK-1 gene (arg107-to-stop, CGA-to-TGA)

[141373]associated with chronic haemolytic anaemia. Brit. J. Haemat. 105:

[141374]75–79, 1999.

[141375]25279.

[141376]Boivin, P.; Galand, C.; Hakim, J.; Simony, D.; Seligman, M.: Une

[141377]nouvelle erythroenzymopathie: anemie hemolytique con-

genitale non spherocytaire

[141378]et deficit hereditaire en adenylate-kinase erythrocytaire.

Presse

[141379]Med. 79: 215-218, 1971.

[141380]25280.

[141381]Boivin, P.; Galand, C.; Hakim, J.; Simony, D.; Seligman, M.:

Deficit

[141382]congenital en adenylate-kinase erythrocytaire. (Letter)

Presse Med. 78:

[141383]1443 only, 1970.

[141384]25281.

[141385]Bowman, J. E.; Frischer, H.; Ajmar, F.; Carson, P. E.; Gower,

M.

[141386]K.: Population, family and biochemical investigation of
human adenylate

[141387]kinase polymorphism. Nature 214: 1156-1158, 1967.

[141388]25282.

[141389]Brock, D. J. H.: Evidence against a common subunit in
adenylate

[141390]kinase and pyruvate kinase. Humangenetik 10: 30-34,
1970.

[141391]25283.

[141392]Cavalli-Sforza, L. L.; King, M. C.; Go, R. C. P.; Namboodiri, K.

[141393]K.; Lynch, H. T.; Wong, L.; Kaplan, E. B.; Elston, R. C.: Possible

[141394]linkage between transcobalamin II (TC II) and adenylate kinase (AK).

[141395](Abstract) Cytogenet. Cell Genet. 25: 140-141, 1979.

[141396]25284.

[141397]Cook, P. J. L.; Robson, E. B.; Buckton, K. E.; Slaughter, C. A.;

[141398]Gray, J. E.; Blank, C. E.; James, F. E.; Ridler, M. A. C.; Insley,

[141399]J.; Hulten, M.: Segregation of ABO, AK(1) and ACONs in families with

[141400]abnormalities of chromosome 9. Ann. Hum. Genet. 41: 365-377, 1978.

[141401]25285.

[141402]Fenger, K.; Sorensen, S. A.: Evaluation of a possible sex difference

[141403]in recombination for the ABO-AK linkage. Am. J. Hum. Genet. 27:

[141404]784–788, 1975.

[141405]25286.

[141406]Ferguson-Smith, M. A.; Aitken, D. A.; Turleau, C.; de
Grouchy,

[141407]J.: Localisation of the human ABO: Np-1: AK-1 linkage
group by regional

[141408]assignment of AK-1 to 9q34. Hum. Genet. 34: 35–43,
1976.

[141409]25287.

[141410]Fildes, R. A.; Harris, H.: Genetically determined variation
of

[141411]adenylate kinase in man. Nature 209: 261–262, 1966.

[141412]25288.

[141413]Geurts van Kessel, A. H. M.; Hagemeijer, A.; Westerveld,
A.; Meera

[141414]Khan, P.; de Groot, P. G.; Pearson, P. L.: Characterization
of chromosomal

[141415]abnormalities in chronic myeloid leukemia using somatic
cell hybrids.

[141416](Abstract) Cytogenet. Cell Genet. 32: 280 only, 1982.

[141417]25289.

- [141418]Lachant, N. A.; Zerez, C. R.; Barredo, J.; Lee, D. W.; Savely,
[141419]S. M.; Tanaka, K. R.: Hereditary erythrocyte adenylate ki-
nase deficiency:
[141420]a defect of multiple phosphotransferases? Blood 77:
2774-2784, 1991.
[141421]25290.
- [141422]Matsuura, S.; Igarashi, M.; Tanizawa, Y.; Yamada, M.;
Kishi, F.;
[141423]Kajii, T.; Fujii, H.; Miwa, S.; Sakurai, M.; Nakazawa, A.: Hu-
man
[141424]adenylate kinase deficiency associated with hemolytic
anemia: a single
[141425]base substitution affecting solubility and catalytic activity
of the
[141426]cytosolic adenylate kinase. J. Biol. Chem. 264:
10148-10155, 1989.
[141427]25291.
- [141428]Miwa, S.; Fujii, H.; Tani, K.; Takahashi, K.; Takizawa, T.;
Igarashi,
[141429]T.: Red cell adenylate kinase deficiency associated with
hereditary
[141430]nonspherocytic hemolytic anemia: clinical and biochemical

studies. Am.

[141431]J. Hemat. 14: 325-333, 1983.

[141432]25292.

[141433]Mohandas, T.; Sparkes, R. S.; Sparkes, M. C.; Shulkin, J. D.;

[141434]Toomey, K. E.; Funderburk, S. J.: Regional localization of
human

[141435]gene loci on chromosome 9: studies of somatic cell hy-
brids containing

[141436]human translocations. Am. J. Hum. Genet. 31: 586-600,
1979.

[141437]25293.

[141438]Povey, S.; Slaughter, C. A.; Wilson, D. E.; Gormley, I. P.;
Buckton,

[141439]K. E.; Perry, P.; Bobrow, M.: Evidence for the assignment of
loci

[141440]AK 1, AK 3 and ACON to chromosome 9 in man. Ann.
Hum. Genet. 39:

[141441]413-422, 1976.

[141442]25294.

[141443]Qualtieri, A.; Pedace, V.; Bisconte, M. G.; Bria, M.; Gulino,

[141444]B.; Andreoli, V.; Brancati, C.: Severe erythrocyte adenylate

kinase

[141445]deficiency due to homozygous A-to-G substitution at
codon 164 of human

[141446]AK1 gene associated with chronic haemolytic anaemia.
Brit. J. Haemat. 99:

[141447]770-776, 1997.

[141448]25295.

[141449]Rapley, S.; Robson, E. B.; Harris, H.; Smith, S. M.: Data on
[141450]the incidence, segregation and linkage relations of the
adenylate

[141451]kinase (AK) polymorphism. Ann. Hum. Genet. 31:
237-242, 1967.

[141452]25296.

[141453]Ohno, S.: Original domain for the serum albumin family
arose

[141454]from repeated sequences. Proc. Nat. Acad. Sci. 78:
7657-7661, 1981.

[141455]25297.

[141456]Ott, H.: Analbuminemia. In: Linneweh, F.: Erbliche Stoff-
wechselkrankheiten.

[141457]Munich: Urban und Schwarzenberg (pub.) 1962. Pp. 44.

[141458]25298.

[141459]Peach, R. J.; Brennan, S. O.: Structural characterization of

[141460]a glycoprotein variant of human serum albumin: albumin

Casebrook (494

[141461]asp-to-asn). Biochim. Biophys. Acta 1097: 49-54, 1991.

[141462]25299.

[141463]Peach, R. J.; Fellowes, A. P.; Brennan, S. O.; George, P. M.:

[141464]Albumin Rugby Park: a truncated albumin variant caused

by a G-to-C

[141465]splice-site mutation in intron 13. Biochim. Biophys. Acta

1180:

[141466]107-110, 1992.

[141467]25300.

[141468]Petersen, C. E.; Scottolini, A. G.; Cody, L. R.; Mandel, M.;

Reimer,

[141469]N.; Bhagavan, N. V.: A point mutation in the human serum

albumin

[141470]gene results in familial dysalbuminaemic hyperthyroxi-

naemia. J. Med.

[141471]Genet. 31: 355-359, 1994.

[141472]25301.

[141473]Pinkert, C. A.; Ornitz, D. M.; Brinster, R. L.; Palmiter, R. D.

[141474]: An albumin enhancer located 10 kb upstream functions
along with

[141475]its promoter to direct efficient, liver-specific expression in
transgenic

[141476]mice. Genes Dev. 1: 268-276, 1987.

[141477]25302.

[141478]Pohlenz, J.; Sadow, P. M.; Koffler, T.; Schonberger, W.;
Weiss,

[141479]R. E.; Refetoff, S.: Congenital hypothyroidism in a child
with unsuspected

[141480]familial dysalbuminemic hyperthyroxinemia caused by a
mutation (R218H)

[141481]in the human albumin gene. J. Pediat. 139: 887-891,
2001.

[141482]25303.

[141483]Prager, E. M.; Wilson, A. C.; Lowenstein, J. M.; Sarich, V. M.

[141484]: Mammoth albumin. Science 209: 287-289, 1980.

[141485]25304.

[141486]Premachandra, B. N.; Wolfe, B.; Williams, I. K.: Coexistence
[141487]of familial dysalbuminemic hyperthyroxinemia with famil-
ial hypercholesterolemia
[141488]and multiple lipoprotein type hyperlipidemia. Am. J. Med.
84: 345-351,
[141489]1988.
[141490]25305.
[141491]Putnam, F. W.: Personal Communication. Bloomington,
Ind. 8/4/1993.
[141492]25306.
[141493]Rajatanavin, R.; Fournier, L.; DeCosimo, D.; Abreau, C.;
Braverman,
[141494]L. E.: Elevated serum free thyroxine by thyroxine analog
radioimmunoassays
[141495]in euthyroid patients with familial dysalbuminemic hyper-
thyroxinemia. Ann.
[141496]Intern. Med. 97: 865-866, 1982.
[141497]25307.
[141498]Rajatanavin, R.; Young, R. A.; Braverman, L. E.: Effect of
chloride
[141499]on serum thyroxine binding in familial dysalbuminemic

hyperthyroxinemia. J.

[141500]Clin. Endocr. Metab. 58: 388–391, 1984.

[141501]25308.

[141502]Rochu, D.; Fine, J. M.: New method for identifying genetic variants

[141503]of human proalbumin. Clin. Chem. 32: 2063–2065, 1986.

[141504]25309.

[141505]Ruffner, D. E.; Dugaiczky, A.: Splicing mutation in human hereditary

[141506]analbuminemia. Proc. Nat. Acad. Sci. 85: 2125–2129, 1988.

[141507]25310.

[141508]Ruiz, M.; Rajatanavin, R.; Young, R. A.; Taylor, C.; Brown, R.;

[141509]Braverman, L. E.; Ingbar, S. H.: Familial dysalbuminemic hyperthyroxinemia:

[141510]a syndrome that can be confused with thyrotoxicosis. New Eng. J.

[141511]Med. 306: 635–639, 1982.

[141512]

[141513]100. Sakamoto, Y.; Davis, E.; Madison, J.; Watkins, S.;

McLaughlin,

[141514]H.; Leahy, D. T.; Putnam, F. W.: Purification and structural study

[141515]of two albumin variants in an Irish population. Clin. Chim. Acta 204:

[141516]179–188, 1991.

[141517]

[141518]101. Sanders, G. T. B.; Tarnoky, A. L.: Albumin Amsterdam: a new

[141519]European albumin variant. IRCS Med. Sci. 7: 581 only, 1979.

[141520]

[141521]102. Sarcione, E. J.; Aungst, C. W.: Studies in bisalbuminemia: binding

[141522]properties of the two albumins. Blood 20: 156–164, 1962.

[141523]

[141524]103. Sargent, T. D.; Wu, J.-R.; Sala-Trepat, J. M.; Wallace, R. B.;

[141525]Reyes, A. A.; Bonner, J.: The rat serum albumin gene: analysis of

[141526]cloned sequences. Proc. Nat. Acad. Sci. 76: 3256–3260,

1979.

[141527]

[141528]104. Sarich, V. M.: Generation time and albumin evolution.
Biochem.

[141529]Genet. 7: 205–212, 1972.

[141530]

[141531]105. Schell, L. M.; Agarwal, S. S.; Blumberg, B. S.; Levy, H.;
Bennett,

[141532]H.; Laughlin, W. S.; Martin, J. P.: Distribution of albumin
variants

[141533]Naskapi and Mexico among Aleuts, Frobisher Bay Eski-
mos, and Micmac,

[141534]Naskapi, Mohawk, Omaha and Apache Indians. Am. J.
Phys. Anthrop. 49:

[141535]111–118, 1978.

[141536]

[141537]106. Schell, L. M.; Blumberg, B. S.: The genetics of human
serum

[141538]albumin.In: Rosenoer, V. M.; Oratz, M.; Rothschild, M. A.:
Albumin

[141539]Structure, Function and Uses. Oxford: Pergamon Press

(pub.) 1977.

[141540]Pp. 113–141.

[141541]

[141542]107. Shalaby, F.; Shafritz, D. A.: Exon skipping during
splicing

[141543]of albumin mRNA precursors in Nagase analbuminemic
rats. Proc. Nat.

[141544]Acad. Sci. 87: 2652–2656, 1990.

[141545]

[141546]108. Shashaty, G.; Atamer, M.: Acquired bisalbuminemia
with hyperamylasemia. Digest.

[141547]Dis. 17: 59–67, 1972.

[141548]

[141549]109. Shibata, T.; Abe, T.: Linkage between the loci for
serum albumin

[141550]and vitamin D binding protein (GC) in the Japanese quail.
Animal

[141551]Genet. 27: 195–197, 1996.

[141552]

[141553]110. Silverberg, J. D. H.; Premachandra, B. N.: Familial hy–

perthyroxinemia

[141554] due to abnormal thyroid hormone binding. Ann. Intern. Med. 96: 183–186,

[141555] 1982.

[141556]

[141557] 111. Sunthornthepvarakul, T.; Likitmaskul, S.; Ngown-garmratana, S.;

[141558] Angsusingha, K.; Kitvitayasak, S.; Scherberg, N. H.; Refetoff, S.

[141559]: Familial dysalbuminemic hypertriiodothyroninemia: a new, dominantly

[141560] inherited albumin defect. J. Clin. Endocr. Metab. 83: 1448–1454,

[141561] 1998.

[141562]

[141563] 112. Swain, B. K.; Talukder, G.; Sharma, A.: Bisalbuminaemia: reports

[141564] from Calcutta. Biomedicine 33: 172–173, 1980.

[141565]

[141566] 113. Takahashi, N.; Takahashi, Y.; Blumberg, B. S.; Putnam, F. W.

[141567]: Amino acid substitutions in genetic variants of human serum albumin

[141568]and in sequences inferred from molecular cloning. Proc. Nat. Acad.

[141569]Sci. 84: 4413–4417, 1987.

[141570]

[141571]114. Takahashi, N.; Takahashi, Y.; Isobe, T.; Putnam, F. W.; Fujita,

[141572]M.; Satoh, C.; Neel, J. V.: Amino acid substitutions in inherited

[141573]albumin variants from Amerindian and Japanese populations. Proc.

[141574]Nat. Acad. Sci. 84: 8001–8005, 1987.

[141575]

[141576]115. Takahashi, N.; Takahashi, Y.; Putnam, F. W.: Structural changes

[141577]and metal binding by proalbumins and other amino-terminal genetic

[141578]variants of human serum albumin. Proc. Nat. Acad. Sci. 84: 7403–7407,

[141579]1987.

[141580]

[141581]116. Tarnoky, A. L.; Lestas, A. N.: A new type of bisalbuminaemia. Clin.

[141582]Chim. Acta 9: 551-558, 1964.

[141583]

[141584]117. Urano, Y.; Sakai, M.; Watanabe, K.; Tamaoki, T.: Tandem arrangement

[141585]of the albumin and alpha-fetoprotein genes in the human genome. Gene 32:

[141586]255-261, 1984.

[141587]

[141588]118. Vanzetti, G.; Porta, F.; Prencipe, L.; Scherini, A.; Fraccaro,

[141589]M.: A homozygote for a serum albumin variant of the fast type. Hum.

[141590]Genet. 46: 5-9, 1979.

[141591]

[141592]119. Vaysse, J.; Pilardeau, P.; Garnier, M.: Trisalbuminemia. (Letter) New

[141593]Eng. J. Med. 305: 833-834, 1981.

[141594]

[141595]120. Wada, N.; Chiba, H.; Shimizu, C.; Kijima, H.; Kubo, M.; Koike

[141596]T.: A novel missense mutation in codon 218 of the albumin gene in

[141597]a distinct phenotype of familial dysalbuminemic hyperthyroxinemia

[141598]in a Japanese kindred. J. Clin. Endocr. Metab. 82: 3246–3250, 1997.

[141599]

[141600]121. Watkins, S.; Madison, J.; Galliano, M.; Minchiotti, L.; Putnam,

[141601]F. W.: A nucleotide insertion and frameshift cause analbuminemia

[141602]in an Italian family. Proc. Nat. Acad. Sci. 91: 2275–2279, 1994.

[141603]

[141604]122. Weitkamp, L. R.: Comparative gene mapping: linkage between the

[141605]albumin and Gc loci in the horse. (Abstract) Am. J. Hum. Genet. 30:

[141606]128A only, 1978.

[141607]

[141608]123. Weitkamp, L. R.; Buck, A. A.: Phenotype frequencies
for four

[141609]serum proteins in Afghanistan: two 'new' albumin variants.
Humangenetik 15:

[141610]335–340, 1972.

[141611]

[141612]124. Weitkamp, L. R.; Chagnon, N. A.: Albumin Maku: a
new variant

[141613]of human serum albumin. Nature 217: 759–760, 1968.

[141614]

[141615]125. Weitkamp, L. R.; Franglen, G.; Rokala, D. A.; Polesky,
H. F.;

[141616]Simpson, N. E.; Sunderman, F. W., Jr.; Bell, H. E.; Saave, J.;
Lisker,

[141617]R.; Bohls, S. W.: An electrophoretic comparison of human
serum albumin

[141618]variants: eight distinguishable types. Hum. Hered. 19:
159–169,

[141619]1969.

[141620]

- [141621]126. Weitkamp, L. R.; Renwick, J. H.; Berger, J. P.; Shreffler, D.
C.; Drachmann, O.; Wuhrmann, F.; Braend, M.; Franglen, G.: Additional
data and summary for albumin-GC linkage in man. Hum. Hered. 20:
1-7, 1970.
- [141625]
- [141626]127. Weitkamp, L. R.; Robson, E. B.; Shreffler, D. C.; Corney, G.
An unusual human serum albumin variant: further data on genetic
linkage between loci for human serum albumin and group-specific component
(GC). Am. J. Hum. Genet. 20: 392-397, 1968.
- [141630]
- [141631]128. Weitkamp, L. R.; Rucknagel, D. L.; Gershowitz, H.: Genetic linkage
between structural loci for albumin and group specific component (GC). Am.
J. Hum. Genet. 18: 559-571, 1966.

[141634]

[141635]129. Weitkamp, L. R.; Salzano, F. M.; Neel, J. V.; Porta, F.;
Geerdink,

[141636]R. A.; Tarnoky, A. L.: Human serum albumin: twenty-three
genetic

[141637]variants and their population distribution. Ann. Hum.
Genet. 36:

[141638]381–392, 1973.

[141639]

[141640]130. Weitkamp, L. R.; Shreffler, D. C.; Robbins, J. L.;
Drachmann,

[141641]O.; Adner, P. L.; Weime, R. J.; Simon, N. M.; Cooke, K. B.;
Sandor,

[141642]G.; Wuhrmann, F.; Braend, M.; Tarnoky, A. L.: An elec-
trophoretic

[141643]comparison of serum albumin variants from nineteen un-
related families. Acta

[141644]Genet. Statist. Med. 17: 399–405, 1967.

[141645]

[141646]131. Wieme, R. J.: On the presence of two albumins in
certain normal

[141647]human sera and its genetic determination. Clin. Chim.

Acta 5: 443–445,

[141648]1960.

[141649]

[141650]132. Williams, D. I.; Martin, N. H.: Bisalbuminemia with
curious

[141651]acrocyanotic skin changes (two cases). Proc. Roy. Soc.
Med. 53:

[141652]566–568, 1960.

[141653]

[141654]133. Yabu, Y.; Amir, S. M.; Ruiz, M.; Braverman, L. E.; Ing-
bar, S.

[141655]H.: Heterogeneity of thyroxine binding by serum albumins
in normal

[141656]subjects and patients with familial dysalbuminemic hyper-
thyroxinemia. J.

[141657]Clin. Endocr. Metab. 60: 451–459, 1985.

[141658]

[141659]134. Yabu, Y.; Miyai, K.; Kobayashi, A.; Miki, K.; Doi, K.;
Takamatsu,

[141660]J.; Mozai, T.; Matsuzuka, F.; Kuma, K.: A new type of albu-
min with

[141661]predominantly increased binding affinity for
3,3–prime,5–triiodothyronine

[141662]in a patient with Graves' disease. J. Endocr. Invest. 10:
163–169,

[141663]1987.

[141664]

[141665]135. Yeo, P. P. B.; Yabu, Y.; Etzkorn, J. R.; Rajatanavin, R.;
Braverman,

[141666]L. E.; Ingbar, S. H.: A four generation study of dysalbu-
minemic hyperthyroxinemia:

[141667]diagnosis in the presence of an acquired excess of thy-
roxine–binding

[141668]globulin. J. Endocr. Invest. 10: 33–38, 1987.

[141669]

[141670]136. Ying, Q.; Liang, Z.; Wu, H.; Wang, L.: The gene fre–

quency of

[141671]serum albumin variants in Chinese and the electrophoretic
characterization

[141672]of several serum albumin variants. Scientia Sinica 24:
1597–1602,

[141673]1981.

[141674]25311.

[141675]Osier, M. V.; Pakstis, A. J.; Soodyall, H.; Comas, D.; Gold-
man,

[141676]D.; Odunsi, A.; Okonofua, F.; Parnas, J.; Schulz, L. O.;
Bertranpetit,

[141677]J.; Bonne–Tamir, B.; Lu, R.–B.; Kidd, J. R.; Kidd, K. K.: A
global

[141678]perspective on genetic variation at the ADH genes reveals
unusual

[141679]patterns of linkage disequilibrium and diversity. Am. J.
Hum. Genet. 71:

[141680]84–99, 2002.

[141681]25312.

[141682]Comings, D. E.; Gade–Andavolu, R.; Gonzalez, N.; Blake,
H.; Wu,

[141683]S.; MacMurray, J. P.: Additive effect of three noradrenergic

genes

[141684](ADRA2A, ADRA2C, DBH) on attention-deficit hyperactivity disorder

[141685]and learning disabilities in Tourette syndrome subjects.

Clin. Genet. 55:

[141686]160–172, 1999.

[141687]25313.

[141688]Halperin, J. M.; Newcorn, J. H.; Koda, V. H.; Pick, L.; McKay,

[141689]K. E.; Knott, P.: Noradrenergic mechanisms in ADHD children with

[141690]and without reading disabilities: a replication and extension. J.

[141691]Am. Acad. Child Adolesc. Psychiat. 36: 1688–1697, 1997.

[141692]25314.

[141693]Hein, L.; Altman, J. D.; Kobilka, B. K.: Two functionally distinct

[141694]alpha-2-adrenergic receptors regulate sympathetic neurotransmission. Nature 402:

[141695]181–184, 1999.

[141696]25315.

[141697]Hoehe, M.; Berrettini, W.; Leppert, M.; Lalouel, J.-M.; Byer-

ley,

[141698]W.; Gershon, E.; White, R.: Genetic mapping of adrenergic receptor

[141699]genes. (Abstract) Am. J. Hum. Genet. 45 (suppl.): A143 only, 1989.

[141700]25316.

[141701]Hoehe, M. R.; Berrettini, W. H.; Lentes, K.-U.: Dra I identifies

[141702]a two allele DNA polymorphism in the human alpha-2-adrenergic receptor

[141703]gene (ADRAR), using a 5.5 kb probe (p ADRAR). Nucleic Acids Res. 16:

[141704]9070 only, 1988.

[141705]25317.

[141706]Kobilka, B. K.; Matsui, H.; Kobilka, T. S.; Yang-Feng, T. L.; Francke,

[141707]U.; Caron, M. G.; Lefkowitz, R. J.; Regan, J. W.: Cloning, sequencing,

[141708]and expression of the gene coding for the human platelet alpha-2-adrenergic

[141709]receptor. Science 238: 650-656, 1987.

[141710]25318.

[141711]Oakey, R. J.; Caron, M. G.; Lefkowitz, R. J.; Seldin, M. F.:
Genomic

[141712]organization of adrenergic and serotonin receptors in the
mouse: linkage

[141713]mapping of sequence-related genes provides a method
for examining

[141714]mammalian chromosome evolution. Genomics 10:
338-344, 1991.

[141715]25319.

[141716]Philipp, M.; Brede, M. E.; Hadamek, K.; Gessler, M.; Lohse,
M.

[141717]J.; Hein, L.: Placental alpha-2-adrenoceptors control vas-
cular development

[141718]at the interface between mother and embryo. Nature
Genet. 31: 311-315,

[141719]2002.

[141720]25320.

[141721]Surprenant, A.; Horstman, D. A.; Akbarali, H.; Limbird, L.
E.:

[141722]A point mutation of the alpha-2-adrenoceptor that blocks
coupling

[141723]to potassium but not calcium currents. Science 257:

977-980, 1992.

[141724]25321.

[141725]Yang-Feng, T. L.; Kobilka, B. K.; Caron, M. G.; Lefkowitz, R.

[141726]J.; Francke, U.: Chromosomal assignment of genes for an alpha-adrenergic

[141727]receptor (ADRAR) and for another member of this receptor family coupled

[141728]to guanine nucleotide regulatory proteins (RG21). (Abstract) Cytogenet.

[141729]Cell Genet. 46: 722-723, 1987.

[141730]25322.

[141731]Bruno, J. F.; Whittaker, J.; Song, J.; Berelowitz, M.: Molecular

[141732]cloning and sequencing of a cDNA encoding a human alpha-1A adrenergic

[141733]receptor. Biochem. Biophys. Res. Commun. 179: 1485-1490, 1991.

[141734]25323.

[141735]Esbenshade, T. A.; Hirasawa, A.; Tsujimoto, G.; Tanaka, T.; Yano,

[141736]J.; Minneman, K. P.; Murphy, T. J.: Cloning of the human

alpha-1D-adrenergic

[141737]receptor and inducible expression of three subtypes in
SK-N-MC cells. Molec.

[141738]Pharm. 47: 977-985, 1995.

[141739]25324.

[141740]Forray, C.; Bard, J. A.; Wetzel, J. M.; Chiu, G.; Shapiro, E.;

[141741]Tang, R.; Lepor, H.; Hartig, P. R.; Weinshank, R. L.;
Branchek, T.

[141742]A.; Gluchowski, C.: The alpha-1-adrenergic receptor that
mediates

[141743]smooth muscle contraction in human prostate has the
pharmacological

[141744]properties of the cloned human alpha-1C subtype. Molec.
Pharm. 45:

[141745]703-708, 1994.

[141746]25325.

[141747]Loftus, S. K.; Shiang, R.; Warrington, J. A.; Bengtsson, U.;
McPherson,

[141748]J. D.; Wasmuth, J. J.: Genes encoding adrenergic receptors
are not

[141749]clustered on the long arm of human chromosome 5. Cyto-

genet. Cell

[141750]Genet. 67: 69–74, 1994.

[141751]25326.

[141752]Lomasney, J. W.; Cotecchia, S.; Lorenz, W.; Leung, W.-Y.;
Schwinn,

[141753]D. A.; Yang-Feng, T. L.; Brownstein, M.; Lefkowitz, R. J.;
Caron,

[141754]M. G.: Molecular cloning and expression of the cDNA for
the alpha-1A-adrenergic

[141755]receptor: the gene for which is located on human chromo-
some 5. J.

[141756]Biol. Chem. 266: 6365–6369, 1991.

[141757]25327.

[141758]Schwinn, D. A.; Johnston, G. I.; Page, S. O.; Mosley, M. J.;
Wilson,

[141759]K. H.; Worman, N. P.; Campbell, S.; Fidock, M. D.; Furness,
L. M.;

[141760]Parry-Smith, D. J.; Peter, B.; Bailey, D. S.: Cloning and
pharmacological

[141761]characterization of human alpha-1 adrenergic receptors:
sequence corrections

[141762]and direct comparison with other species homologues. J.

Pharm. Exp.

[141763]Ther. 272: 134–142, 1995.

[141764]25328.

[141765]Schwinn, D. A.; Lomasney, J. W.: Pharmacologic characterization

[141766]of cloned alpha-1-adrenoceptor subtypes: selective antagonists suggest

[141767]the existence of a fourth subtype. Europ. J. Pharm. 227: 433–436,

[141768]1992.

[141769]25329.

[141770]Weinberg, D. H.; Trivedi, P.; Tan, C. P.; Mitra, S.; Perkins-Barrow,

[141771]A.; Borkowski, D.; Strader, C. D.; Bayne, M.: Cloning, expression

[141772]and characterization of human alpha adrenergic receptors alpha-1A,

[141773]alpha-1B, and alpha-1C. Biochem. Biophys. Res. Commun. 201: 1296–1304,

[141774]1994.

[141775]25330.

[141776]Yang-Feng, T. L.; Han, H.; Lomasney, J. W.; Caron, M. G.:
Localization
[141777]of the cDNA for an alpha-1-adrenergic receptor subtype
(ADRA1D) to
[141778]chromosome band 20p13. Cytogenet. Cell Genet. 66:
170-171, 1994.
[141779]25331.

[141780]Markert, M. L.; Hershfield, M. S.; Wiginton, D. A.; States, J.
[141781]C.; Ward, F. E.; Bigner, S. H.; Buckley, R. H.; Kaufman, R.
E.; Hutton,
[141782]J. J.: Identification of a deletion in the adenosine deami-
nase gene
[141783]in a child with severe combined immunodeficiency. J. Im-
mun. 138:
[141784]3203-3206, 1987.
[141785]25332.

[141786]Markert, M. L.; Hutton, J. J.; Wiginton, D. A.; States, J. C.;
[141787]Kaufman, R. E.: Adenosine deaminase (ADA) deficiency
due to deletion
[141788]of the ADA gene promoter and first exon by homologous
recombination

[141789]between two Alu elements. J. Clin. Invest. 81: 1323–1327, 1988.

[141790]25333.

[141791]Markert, M. L.; Norby–Slycord, C.; Ward, F. E.: A high proportion

[141792]of ADA point mutations associated with a specific alanine-to-valine

[141793]substitution. Am. J. Hum. Genet. 45: 354–361, 1989.

[141794]25334.

[141795]Meuwissen, H. J.; Pollara, B.; Pickering, R. J.: Combined immunodeficiency

[141796]disease associated with adenosine deaminase deficiency (report on

[141797]a workshop held in Albany, New York, October 1, 1973). J. Pediat. 86:

[141798]169–181, 1975.

[141799]25335.

[141800]Migchielsen, A. A. J.; Breuer, M. L.; van Roon, M. A.; te Riele,

[141801]H.; Zurcher, C.; Ossendorp, F.; Toutain, S.; Hershfield, M. S.; Berns,

[141802]A.; Valerio, D.: Adenosine-deaminase-deficient mice die perinatally

[141803]and exhibit liver-cell degeneration, atelectasis and small intestinal

[141804]cell death. Nature Genet. 10: 279-287, 1995.

[141805]25336.

[141806]Mitchell, B. S.; Mejias, E.; Daddona, P. E.; Kelley, W. N.: Purinogenic

[141807]immunodeficiency disease: selective toxicity of deoxyribonucleosides

[141808]for T-cells. Proc. Nat. Acad. Sci. 75: 5011-5014, 1978.

[141809]25337.

[141810]Moen, R. C.; Horowitz, S. D.; Sondel, P. M.; Borchering, W. R.;

[141811]Trigg, M. E.; Billing, R.; Hong, R.: Immunologic reconstitution after

[141812]haploidentical bone marrow transplantation for immune deficiency disorders:

[141813]treatment of bone marrow cells with monoclonal antibody CT-2 and complement. Blood 70:

[141814]664-669, 1987.

[141815]25338.

[141816]Mohandas, T.; Sparkes, R. S.; Suh, E. J.; Hershfield, M. S.:

[141817]Regional localization of the human genes for S-adenosylhomocysteine

[141818]hydrolase (cen-q131) and adenosine deaminase (q131-qter) on chromosome

[141819]20. Hum. Genet. 66: 292-295, 1984.

[141820]25339.

[141821]Nielsen, K. B.; Tommerup, N.; Jespersen, B.; Nygaard, P.; Kleif,

[141822]L.: Segregation of a t(3;20) translocation through three generations

[141823]resulting in unbalanced karyotypes in six persons. J. Med. Genet. 23:

[141824]446-451, 1986.

[141825]25340.

[141826]Orkin, S. H.; Daddona, P. E.; Shewach, D. S.; Markham, A. F.;

[141827]Bruns, G. A.; Goff, S. C.; Kelley, W. N.: Molecular cloning of human

[141828]adenosine deaminase gene sequences. J. Biol. Chem. 258:

12753–12756,

[141829]1983.

[141830]25341.

[141831]Ozsahin, H.; Arredondo–Vega, F. X.; Santisteban, I.;

Fuhrer, H.;

[141832]Tuchs Schmid, P.; Jochum, W.; Aguzzi, A.; Lederman, H. M.;

Fleischman,

[141833]A.; Winkelstein, J. A.; Seger, R. A.; Hershfield, M. S.:

Adenosine

[141834]deaminase deficiency in adults. Blood 89: 2849–2855,

1997.

[141835]25342.

[141836]Onodera, M.; Ariga, T.; Kawamura, N.; Kobayashi, I.;

Ohtsu, M.;

[141837]Yamada, M.; Tame, A.; Furuta, H.; Okano, M.; Matsumoto,

S.; Kotani,

[141838]H.; McGarrity, G. J.; Blaese, R. M.; Sakiyama, Y.: Successful

peripheral

[141839]T–lymphocyte–directed gene transfer for a patient with

severe combined

[141840]immune deficiency caused by adenosine deaminase defi–

ciency. Blood 91:

[141841]30–36, 1998.

[141842]25343.

[141843]Palmer, T. D.; Hock, R. A.; Osborne, W. R. A.; Miller, A. D.:

[141844]Efficient retrovirus-mediated transfer and expression of a
human adenosine

[141845]deaminase gene in diploid skin fibroblasts from an
adenosine deaminase-deficient

[141846]human. Proc. Nat. Acad. Sci. 84: 1055–1059, 1987.

[141847]25344.

[141848]Parkman, R.; Gelfand, E. W.; Rosen, F. S.; Sanderson, A.;
Hirschhorn,

[141849]R.: Severe combined immunodeficiency and adenosine
deaminase deficiency. New

[141850]Eng. J. Med. 292: 714–719, 1975.

[141851]25345.

[141852]Petersen, M. B.; Tranebjaerg, L.; Tommerup, N.; Nygaard,
P.; Edwards,

[141853]H.: New assignment of the adenosine deaminase gene lo-
cus to chromosome

[141854]20q13.11 by study of a patient with interstitial deletion
20q. J.

[141855]Med. Genet. 24: 93–96, 1987.

[141856]25346.

[141857]Polmar, S. H.; Stern, R. C.; Schwartz, A. L.; Wetzler, E. M.;

[141858]Chase, P. A.; Hirschhorn, R.: Enzyme replacement therapy
for adenosine

[141859]deaminase deficiency and severe combined immunodeficiency. New Eng.

[141860]J. Med. 295: 1337–1343, 1976.

[141861]25347.

[141862]Ratech, H.; Greco, M. A.; Gallo, G.; Rimoin, D. L.; Kamino,
H.;

[141863]Hirschhorn, R.: Pathologic findings in adenosine–deami-
nase–deficient

[141864]severe combined immunodeficiency. I. Kidney, adrenal,
and chondro–osseous

[141865]tissue alterations. Am. J. Path. 120: 157–169, 1985.

[141866]25348.

[141867]Ritter, H.; Wendt, G. G.; Tariverdian, G.; Zelch, J.; Rube, M.;

[141868]Kirchberg, G.: Genetics and linkage analysis of adenosine
deaminase. Humangenetik 14:

[141869]69–71, 1971.

[141870]25349.

[141871]Rogers, M. H.; Lwin, R.; Fairbanks, L.; Gerritsen, B.; Gaspar,

[141872]H. B.: Cognitive and behavioral abnormalities in adenosine deaminase

[141873]deficient severe combined immunodeficiency. *J. Pediat.* 139: 44–50,

[141874]2001.

[141875]25350.

[141876]Rothschild, C. B.; Akots, G.; Hayworth, R.; Pettenati, M. J.;

[141877]Rao, P. N.; Wood, P.; Stolz, F.–M.; Hansmann, I.; Serino, K.; Keith,

[141878]T. P.; Fajans, S. S.; Bowden, D. W.: A genetic map of chromosome

[141879]20q12–q13.1: multiple highly polymorphic microsatellite and RFLP markers

[141880]linked to the maturity-onset diabetes of the young (MODY) locus. *Am.*

[141881]*J. Hum. Genet.* 52: 110–123, 1993.

[141882]25351.

[141883]Rubinstein, A.; Hirschhorn, R.; Sicklick, M.; Murphy, R. A.:

[141884]In vivo and in vitro effects of thymosin and adenosine
deaminase on

[141885]adenosine-deaminase-deficient lymphocytes. New Eng. J.
Med. 300:

[141886]387-392, 1979.

[141887]25352.

[141888]Rudd, N. L.; Bain, H. W.; Giblett, E.; Chen, S.-H.; Worton, R.

[141889]G.: Partial trisomy 20 confirmed by gene dosage studies.
Am. J.

[141890]Med. Genet. 4: 357-364, 1979.

[141891]25353.

[141892]Koch, G.; Shows, T. B.: Somatic cell genetics of adenosine
deaminase

[141893]expression and severe combined immune deficiency dis-
ease in man. Proc.

[141894]Nat. Acad. Sci. 77: 4211-4215, 1980.

[141895]25354.

[141896]Gautam, M.; Noakes, P. G.; Moscoso, L.; Rupp, F.; Scheller,
R.

[141897]H.; Merlie, J. P.; Sanes, J. R.: Defective neuromuscular synaptogenesis

[141898]in agrin-deficient mutant mice. Cell 85: 525–535, 1996.

[141899]25355.

[141900]Glass, D. J.; Bowen, D. C.; Stitt, T. N.; Radziejewski, C.; Bruno,

[141901]J.; Ryan, T. E.; Gies, D. R.; Shah, S.; Mattsson, K.; Burden, S. J.;

[141902]DiStefano, P. S.; Valenzuela, D. M.; DeChiara, T. M.; Yancopoulos,

[141903]G. D.: Agrin acts via a MuSK receptor complex. Cell 85: 513–523,

[141904]1996.

[141905]25356.

[141906]Khan, A. A.; Bose, C.; Yam, L. S.; Soloski, M. J.; Rupp, F.: Physiological

[141907]regulation of the immunological synapse by agrin. Science 292: 1681–1686,

[141908]2001.

[141909]25357.

[141910]Lin, W.; Burgess, R. W.; Dominguez, B.; Pfaff, S. L.; Sanes,

J.

[141911]R.; Lee, K.-F.: Distinct roles of nerve and muscle in post-synaptic

[141912]differentiation of the neuromuscular synapse. Nature 410: 1057-1064,

[141913]2001.

[141914]25358.

[141915]McMahan, U. J.: The agrin hypothesis Cold Spring Harb. Symp.

[141916]Quant. Biol. 50: 407-418, 1990.

[141917]25359.

[141918]Rupp, F.; Ozcelik, T.; Linial, M.; Peterson, K.; Francke, U.; Scheller,

[141919]R.: Structure and chromosomal localization of the mammalian agrin

[141920]gene. J. Neurosci. 12: 3535-3544, 1992.

[141921]25360.

[141922]Rupp, F.; Payan, D. G.; Magill-Solc, C.; Cowan, D. M.; Scheller,

[141923]R. H.: Structure and expression of a rat agrin. Neuron 6: 811-823,

[141924]1991.

[141925]25361.

[141926]D'Alfonso, S.; Richiardi, P. M.: A polymorphic variation in a
[141927]putative regulation box of the TNFA promoter region. Im-
munogenetics 39:

[141928]150–154, 1994.

[141929]25362.

[141930]Mietus-Snyder, M.; Charmley, P.; Korf, B.; Ladas, J. A. A.:
Gatti,

[141931]R. A. and Karathanasis, S. K.: Genetic linkage of the hu-
man apolipoprotein

[141932]AI-CIII-AIV gene cluster and the neural cell adhesion
molecule (NCAM)

[141933]gene. Genomics 7: 633–637, 1990.

[141934]25363.

[141935]Mietus-Snyder, M.; Korf, B.; Ladas, J. A.; Karathanasis, S.
K.

[141936]: Linkage of the human apolipoproteins A1, C3, A4 and
the neural cell

[141937]adhesion molecule (NCAM) genes. (Abstract) Cytogenet.
Cell Genet. 51:

[141938]1044 only, 1989.

[141939]25364.

[141940]Nguyen, C.; Mattei, M. G.; Goridis, C.; Mattei, J. F.; Jordan,

[141941]B. R.: Localization of the human N-CAM gene to chromosome 11 by in

[141942]situ hybridization with a murine N-CAM cDNA probe.
(Abstract) Cytogenet.

[141943]Cell Genet. 40: 713 only, 1985.

[141944]25365.

[141945]Nguyen, C.; Mattei, M. G.; Mattei, J.-F.; Santoni, M.-J.;
Goridis,

[141946]C.; Jordan, B. R.: Localization of the human NCAM gene to
band q23

[141947]of chromosome 11: the third gene coding for a cell interaction molecule

[141948]mapped to the distal portion of the long arm of chromosome 11. J.

[141949]Cell Biol. 102: 711-715, 1986.

[141950]25366.

[141951]Rabinowitz, J. E.; Rutishauser, U.; Magnuson, T.: Targeted
mutation

[141952]of Ncam to produce a secreted molecule results in a dominant embryonic

[141953]lethality. Proc. Nat. Acad. Sci. 93: 6421–6424, 1996.

[141954]25367.

[141955]Rutishauser, U.; Acheson, A.; Hall, A. K.; Mann, D. M.; Sunshine,

[141956]J.: The neural cell adhesion molecule (NCAM) as a regulator of cell-cell

[141957]interactions. Science 240: 53–57, 1988.

[141958]25368.

[141959]Rutishauser, U.; Goridis, C.: NCAM: the molecule and its genetics. Trends

[141960]Genet. 2: 72–76, 1986.

[141961]25369.

[141962]Telatar, M.; Lange, E.; Uhrhammer, N.; Gatti, R. A.: New localization

[141963]of NCAM, proximal to DRD2 at chromosome 11q23. Mammalian Genome 6:

[141964]59–60, 1995.

[141965]25370.

- [141966]Durbin, H.; Novelli, M.; Bodmer, W.: Detection of a 4-bp insertion
- [141967](CACA) functional polymorphism at nucleotide 241 of the cellular adhesion
- [141968]regulatory molecule CMAR (formerly CAR). Genomics 19: 181–182, 1994.
- [141969]25371.
- [141970]Koyama, K.; Emi, M.; Nakamura, Y.: The cell adhesion regulator
- [141971](CAR) gene, TaqI and insertion/deletion polymorphisms, and regional
- [141972]assignment to the peritelomeric region of 16q by linkage analysis. Genomics 16:
- [141973]264–265, 1993.
- [141974]25372.
- [141975]Pullman, W. E.; Bodmer, W. F.: Cloning and characterization of
- [141976]a gene that regulates cell adhesion. Nature 356: 529–532, 1992.
- [141977]25373.
- [141978]Agellon, L. B.; Drover, V. A. B.; Cheema, S. K.; Gbaguidi, G. F.;

[141979]Walsh, A.: Dietary cholesterol fails to stimulate the human cholesterol

[141980]7-alpha-hydroxylase gene (CYP7A1) in transgenic mice. J. Biol. Chem. 277:

[141981]20131-20134, 2002.

[141982]25374.

[141983]Angelin, B.; Einarsson, K.; Hellstrom, K.; Leijid, B.: Bile acid

[141984]kinetics in relation to endogenous triglyceride metabolism in various

[141985]types of hyperlipoproteinemia. J. Lipid Res. 19: 1004-1016, 1978.

[141986]25375.

[141987]Angelin, B.; Hershon, K. S.; Brunzell, J. D.: Bile acid metabolism

[141988]in hereditary forms of hypertriglyceridemia: evidence for an increased

[141989]synthesis rate in monogenic familial hypertriglyceridemia. Proc.

[141990]Nat. Acad. Sci. 84: 5434-5438, 1987.

[141991]25376.

[141992]Cohen, J. C.; Cali, J. J.; Jelinek, D. F.; Mehrabian, M.;

Sparkes,

[141993]R. S.; Lusi, A. J.; Russell, D. W.; Hobbs, H. H.: Cloning of
the

[141994]human cholesterol 7- α -hydroxylase gene (CYP7) and
localization

[141995]to chromosome 8q11-q12. Genomics 14: 153-161, 1992.

[141996]25377.

[141997]Drover, V. A. B.; Wong, N. C. W.; Agellon, L. B.: A distinct
thyroid

[141998]hormone response element mediates repression of the
human cholesterol

[141999]7- α -hydroxylase (CYP7A1) gene promoter. Molec. En-
docr. 16: 14-23,

[142000]2002.

[142001]25378.

[142002]Goodwin, B.; Jones, S. A.; Price, R. R.; Watson, M. A.; Mc-
Kee,

[142003]D. D.; Moore, L. B.; Galardi, C.; Wilson, J. G.; Lewis, M. C.;
Roth,

[142004]M. E.; Maloney, P. R.; Willson, T. M.; Kliewer, S. A.: A regu-
latory

[142005]cascade of the nuclear receptors FXR, SHP-1, and LXR-1

represses bile

[142006]acid biosynthesis. Molec. Cell 6: 517–526, 2000.

[142007]25379.

[142008]Molowa, D. T.; Chen, W. S.; Cimis, G. M.; Tan, C. P.: Transcriptional

[142009]regulation of the human cholesterol 7 alpha-hydroxylase gene. Biochemistry 31:

[142010]2539–2544, 1992.

[142011]25380.

[142012]Nitta, M.; Ku, S.; Brown, C.; Okamoto, A. Y.; Shan, B.: CPF: an

[142013]orphan nuclear receptor that regulates liver-specific expression of

[142014]the human cholesterol 7-alpha-hydroxylase gene. Proc. Nat. Acad.

[142015]Sci. 96: 6660–6665, 1999.

[142016]25381.

[142017]Noshiro, M.; Okuda, K.: Molecular cloning and sequence analysis

[142018]of cDNA encoding human cholesterol 7 alpha-hydroxylase. FEBS Lett. 268:

[142019]137–140, 1990.

[142020]25382.

[142021]Paumgartner, G.; Sauerbruch, T.: Gallstones: pathogenesis. Lancet 338:

[142022]1117–1121, 1991.

[142023]25383.

[142024]Wang, J.; Freeman, D. J.; Grundy, S. M.; Levine, D. M.;
Guerra,

[142025]R.; Cohen, J. C.: Linkage between cholesterol
7- α -hydroxylase

[142026]and high plasma low-density lipoprotein cholesterol concentrations. J.

[142027]Clin. Invest. 101: 1283–1291, 1998.

[142028]25384.

[142029]Gress, C. J.; Jacenko, O.: Growth plate compressions and
altered

[142030]hematopoiesis in collagen X null mice. J. Cell Biol. 149:
983–993,

[142031]2000.

[142032]25385.

[142033]Hasegawa, T.; Kozlowski, K.; Nishimura, G.; Hara, H.;
Hasegawa,

[142034]Y.; Aso, T.; Koto, S.; Nagai, T.; Tsuchiya, Y.: Japanese type
of

[142035]spondylo-metaphyseal dysplasia. *Pediat. Radiol.* 24:
194-197, 1994.

[142036]25386.

[142037]Ikegawa, S.; Nakamura, K.; Nagano, A.; Haga, N.; Naka-
mura, Y.:

[142038]Mutations in the N-terminal globular domain of the type X
collagen

[142039]gene (COL10A1) in patients with Schmid metaphyseal
chondrodysplasia. *Hum.*

[142040]*Mutat.* 9: 131-135, 1997.

[142041]25387.

[142042]Ikegawa, S.; Nishimura, G.; Nagai, T.; Hasegawa, T.;
Ohashi, H.;

[142043]Nakamura, Y.: Mutation of the type X collagen gene
(COL10A1) causes

[142044]spondylometaphyseal dysplasia. *Am. J. Hum. Genet.* 63:
1659-1662,

[142045]1998.

[142046]25388.

[142047]Kirsch, T.; von der Mark, K.: Isolation of human type X collagen

[142048]and immunolocalization in fetal human cartilage. *Europ. J. Biochem.* 196:

[142049]575–580, 1991.

[142050]25389.

[142051]Kuivaniemi, H.; Tromp, G.; Prockop, D. J.: Mutations in fibrillar

[142052]collagens (types I, II, III, and XI), fibril-associated collagen (type

[142053]IX), and network-forming collagen (type X) cause a spectrum of diseases

[142054]of bone, cartilage, and blood vessels. *Hum. Mutat.* 9: 300–315, 1997.

[142055]25390.

[142056]McIntosh, I.; Abbot, M. H.; Francomano, C. A.: Concentration

[142057]of mutations causing Schmid metaphyseal chondrodysplasia in the C-terminal

[142058]noncollagenous domain of type X collagen. *Hum. Mut.* 5: 121–125,

[142059]1995.

[142060]25391.

[142061]McIntosh, I.; Abbot, M. H.; Warman, M. L.; Olsen, B. R.;
Francomano,

[142062]C. A.: Additional mutations of type X collagen confirm
COL10A1 as

[142063]the Schmid metaphyseal chondrodysplasia locus. Hum.
Molec. Genet. 3:

[142064]303–307, 1994.

[142065]25392.

[142066]Nielsen, V. H.; Bendixen, C.; Arnbjerg, J.; Sorensen, C. M.;
Jensen,

[142067]H. E.; Shukri, N. M.; Thomsen, B.: Abnormal growth plate
function

[142068]in pigs carrying a dominant mutation in type X collagen.
Mammalian

[142069]Genome 11: 1087–1092, 2000.

[142070]25393.

[142071]Pokharel, R. K.; Alimsardjono, H.; Uno, K.; Fujii, S.; Shiba,

[142072]R.; Matsuo, M.: A novel mutation substituting tryptophan
with arginine

[142073]in the carboxyl-terminal, noncollagenous domain of collagen X in a
[142074]case of Schmid metaphyseal chondrodysplasia. Biochem. Biophys. Res. Commun. 217: 1157-1162, 1995.
[142075]25394.
[142076]Rosati, R.; Horan, G. S. B.; Pinero, G. J.; Garofalo, S.; Keene, D. R.; Horton, W. A.; Vuorio, E.; de Crombrughe, B.; Behringer, R. R.: Normal long bone growth and development in type X collagen-null mice. Nature Genet. 8: 129-135, 1994.
[142080]25395.
[142082]Sawai, H.; Ida, A.; Nakata, Y.; Koyama, K.: Novel missense mutation resulting in the substitution of tyrosine by cysteine at codon 597 of the type X collagen gene associated with Schmid metaphyseal chondrodysplasia. J. Hum. Genet. 43: 259-261, 1998.

[142086]25396.

[142087]Schmid, T. M.; Linsenmayer, T. F.: Immunohistochemical
localization

[142088]of short chain cartilage collagen (type X) in avian tissues.
J. Cell

[142089]Biol. 100: 598-605, 1985.

[142090]25397.

[142091]Stratakis, C. A.; Orban, Z.; Burns, A. L.; Vottero, A.; Mitsi-
ades,

[142092]C. S.; Marx, S. J.; Abbassi, V.; Chrousos, G. P.: Dideoxyfin-
gerprinting

[142093](ddF) analysis of the type X collagen gene (COL10A1) and
identification

[142094]of a novel mutation (S671P) in a kindred with Schmid
metaphyseal chondrodysplasia. Biochem.

[142095]Molec. Med. 59: 112-117, 1996.

[142096]25398.

[142097]Sweetman, W. A.; Rash, B.; Sykes, B.; Beighton, P.; Hecht, J.

[142098]T.; Zabel, B.; Thomas, J. T.; Boot-Handford, R.; Grant, M.
E.; Wallis,

[142099]G. A.: SSCP and segregation analysis of the human type X

collagen

[142100]gene (COL10A1) in heritable forms of chondrodysplasia.

Am. J. Hum.

[142101]Genet. 51: 841–849, 1992.

[142102]25399.

[142103]Thomas, J. T.; Cresswell, C. J.; Rash, B.; Hoyland, J.;

Freemont,

[142104]A. J.; Grant, M. E.; Boot-Handford, R. P.: The human colla-

gen X gene:

[142105]complete primary sequence and reexpression in os-

teoarthritis. Biochem.

[142106]Soc. Trans. 19: 804–808, 1991.

[142107]25400.

[142108]Thomas, J. T.; Cresswell, C. J.; Rash, B.; Nicolai, H.; Jones,

[142109]T.; Solomon, E.; Grant, M. E.; Boot-Handford, R. P.: The

human collagen

[142110]X gene: complete primary translated sequence and chro-

mosomal localization. Biochem.

[142111]J. 280: 617–623, 1991.

[142112]25401.

[142113]Wallis, G. A.; Rash, B.; Sweetman, W. A.; Thomas, J. T.; Su-

per,

[142114]M.; Evans, G.; Grant, M. E.; Boot-Handford, R. P.: Amino acid substitutions

[142115]of conserved residues in the carboxyl-terminal domain of the alpha-I(X)

[142116]chain of type X collagen occur in two unrelated families with metaphyseal

[142117]chondrodysplasia type Schmid. Am. J. Hum. Genet. 54: 169-178, 1994.

[142118]25402.

[142119]Haeffliger, J.-A.; Bruzzone, R.; Jenkins, N. A.; Gilbert, D. J.;

[142120]Copeland, N. G.; Paul, D. L.: Four novel members of the connexin

[142121]family of gap junction proteins: molecular cloning, expression, and

[142122]chromosome mapping. J. Biol. Chem. 267: 2057-2064, 1992.

[142123]25403.

[142124]Hsieh, C.-L.; Kumar, N. M.; Gilula, N. B.; Francke, U.: Distribution

[142125]of genes for gap junction membrane channel proteins on human and mouse

[142126]chromosomes. Somat. Cell Molec. Genet. 17: 191–200,
1991.

[142127]25404.

[142128]Hattersley, A. T.; Beards, F.; Ballantyne, E.; Appleton, M.;
Harvey,

[142129]R.; Ellard, S.: Mutations in the glucokinase gene of the fe-
tus result

[142130]in reduced birth weight. Nature Genet. 19: 268–270,
1998.

[142131]25405.

[142132]McCarthy, M.: Weighing in on diabetes risk. Nature Genet.
19:

[142133]209–210, 1998.

[142134]25406.

[142135]Neel, J. V.: Diabetes mellitus: a 'thrifty' genotype rendered

[142136]detrimental by 'progress'? Am. J. Hum. Genet. 14:
353–362, 1962.

[142137]25407.

[142138]Spielman, R. S.; McGinnis, R. E.; Ewens, W. J.: Transmission

[142139]test for linkage disequilibrium: the insulin gene region
and insulin-dependent

[142140]diabetes mellitus (IDDM). Am. J. Hum. Genet. 52:
506–516, 1993.

[142141]25408.

[142142]Altshuler, D.; Hirschhorn, J. N.; Klannemark, M.; Lindgren,
C.

[142143]M.; Vohl, M.–C.; Nemesh, J.; Lane, C. R.; Schaffner, S. F.;
Bolk,

[142144]S.; Brewer, C.; Tuomi, T.; Gaudet, D.; Hudson, T. J.; Daly,
M.; Groop,

[142145]L.; Lander, E. S.: The common PPAR–gamma pro12ala
polymorphism is

[142146]associated with decreased risk of type 2 diabetes. Nature
Genet. 76–80,

[142147]2000.

[142148]25409.

[142149]DeStefano, A. L.; Baldwin, C. T.; Burzstyn, M.; Gavras, I.;
Handy,

[142150]D. E.; Joost, O.; Martel, T.; Nicolaou, M.; Schwartz, F.;
Streeten,

[142151]D. H. P.; Farrer, L. A.; Gavras, H.: Autosomal dominant or–
thostatic

[142152]hypotensive disorder maps to chromosome 18q. Am. J.

Hum. Genet. 63:

[142153]1425–1430, 1998.

[142154]25410.

[142155]Zneimer, S. M.; Womack, J. E.: Regional localization of the
fibronectin

[142156]and gamma crystallin genes to mouse chromosome 1 by
in situ hybridization. Cytogenet.

[142157]Cell Genet. 48: 238–241, 1988.

[142158]25411.

[142159]den Dunnen, J. T.; Moormann, R. J. M.; Cremers, F. P. M.;
Schoenmakers,

[142160]J. G. G.: Two human gamma–crystallin genes are linked
and riddled

[142161]with Alu–repeats. Gene 38: 197–204, 1985.

[142162]25412.

[142163]Daniels, G.; King, M.–J.; Avent, N. D.; Khalid, G.; Reid, M.;

[142164]Mallinson, G.; Symthe, J.; Cedergren, B.: A point mutation
in the

[142165]GYPC gene results in the expression of the blood group
An(a) antigen

[142166]on glycophorin D but not on glycophorin C: further evi–

dence that glycophorin

[142167]D is a product of the GYPC gene. Blood 82: 3198–3203, 1993.

[142168]25413.

[142169]El-Maliki, B.; Blanchard, D.; Dahr, W.; Beyreuther, K.; Cartron,

[142170]J.-P.: Structural homology between glycophorins C and D of human

[142171]erythrocytes. Europ. J. Biochem. 183: 639–643, 1989.

[142172]25414.

[142173]Jorgensen, J.; Drachmann, O.; Gavin, J.: Duch, Dh(a), a low frequency

[142174]red cell antigen. Hum. Hered. 32: 73–75, 1982.

[142175]25415.

[142176]King, M. J.; Avent, N. D.; Mallinson, G.; Reid, M. E.: Point

[142177]mutation in the glycophorin C gene results in the expression of the

[142178]blood group antigen Dh(a). Vox Sang. 63: 56–58, 1992.

[142179]25416.

[142180]Le Van Kim, C.; Colin, Y.; Blanchard, D.; Dahr, W.; London,

J.;

[142181]Cartron, J.-P.: Gerbich blood group deficiency of the
Ge:-1,-2,-3

[142182]and Ge:-1,-2,3 types: immunochemical study and ge-
nomic analysis with

[142183]cDNA probes. *Europ. J. Biochem.* 165: 571-579, 1987.

[142184]25417.

[142185]Mattei, M. G.; Colin, Y.; Le Van Kim, C.; Mattei, J. F.;
Cartron,

[142186]J. P.: Localization of the gene for human erythrocyte gly-
cophorin

[142187]C to chromosome 2, q14-q21. *Hum. Genet.* 74: 420-422,
1986.

[142188]25418.

[142189]Pasvol, G.; Anstee, D. J.; Tanner, M. J. A.: Glycophorin C
and

[142190]the invasion of red cells by *Plasmodium falciparum*. *Lancet*
I: 907-908,

[142191]1984.

[142192]25419.

[142193]Race, R. R.; Sanger, R.: *Blood Groups in Man*. Oxford:

Blackwell

[142194]Sci. Publ. (pub.) (6th ed.): 1975. Pp. 416–421.

[142195]25420.

[142196]Reid, M. E.: The Gerbich blood group antigens: a review.
Med.

[142197]Lab. Sci. 43: 177–182, 1972.

[142198]25421.

[142199]Reid, M. E.; Sullivan, C.; Taylor, M.; Anstee, D. J.: Inheri-
tance

[142200]of human-erythrocyte Gerbich blood group antigens. Am.
J. Hum. Genet. 41:

[142201]1117–1123, 1987.

[142202]25422.

[142203]Simmons, R. T.; Albrey, J. A.: A 'new' blood group antigen
Webb

[142204](Wb) of low frequency found in two Australian families.
Med. J. Aust. I:

[142205]8–10, 1963.

[142206]25423.

[142207]Sondag, D.; Alloisio, N.; Blanchard, D.; Ducluzeau, M.–T.;

Colonna,

[142208]P.; Bachir, D.; Bloy, C.; Cartron, J.-P.; Delaunay, J.: Gerbich reactivity

[142209]in 4.1(-) hereditary elliptocytosis and protein 4.1 level in blood

[142210]group Gerbich deficiency. Brit. J. Haemat. 65: 43-50, 1987.

[142211]25424.

[142212]Spring, F. A.: Immunochemical characterisation of the low-incidence

[142213]antigen, Dh(a). Vox Sang. 61: 65-68, 1991.

[142214]25425.

[142215]Telen, M. J.; Le Van Kim, C.; Guizzo, M. L.; Cartron, J.-P.; Colin,

[142216]Y.: Erythrocyte Webb-type glycophorin C variant lacks N-glycosylation

[142217]due to an asparagine to serine substitution. Am. J. Hemat. 37: 51-52,

[142218]1991.

[142219]25426.

[142220]Winardi, R.; Reid, M.; Conboy, J.; Mohandas, N.: Molecular

analysis

[142221]of glycophorin C deficiency in human erythrocytes. Blood
81: 2799-2803,

[142222]1993.

[142223]25427.

[142224]Bierhuizen, M. F. A.; Mattei, M.-G.; Fukuda, M.: Expression
of

[142225]the developmental I antigen by a cloned human cDNA en-
coding a member

[142226]of a beta-1,6-N-acetylglucosaminyltransferase gene fam-
ily. Genes

[142227]Dev. 7: 468-478, 1993.

[142228]25428.

[142229]Lin-Chu, M.; Broadberry, R. E.; Okubo, Y.; Tanaka, M.: The
i phenotype

[142230]and congenital cataracts among Chinese in Taiwan (Letter)
Transfusion 31:

[142231]676-677, 1991.

[142232]25429.

[142233]Ogata, H.; Okubo, Y.; Akabane, T.: Phenotype i associated
with

[142234]congenital cataract in Japanese. Transfusion 19: 166–168,
1979.

[142235]25430.

[142236]Yeh, J.-C.; Ong, E.; Fukuda, M.: Molecular cloning and ex-
pression

[142237]of a novel beta-1,6-N-acetylglucosaminyltransferase that
forms core

[142238]2, core 4, and I branches. J. Biol. Chem. 274: 3215–3221,
1999.

[142239]25431.

[142240]Yu, L.-C.; Twu, Y.-C.; Chang, C.-Y.; Lin, M.: Molecular ba-
sis

[142241]of the adult i phenotype and the gene responsible for the
expression

[142242]of the human blood group I antigen. Blood 98:
3840–3845, 2001.

[142243]25432.

[142244]Allen, F. H., Jr.; Krabbe, S. M.; Corcoran, P. A.: A new phe-
notype

[142245](McLeod) in the Kell blood-group system. Vox Sang. 6:
555–560, 1961.

[142246]25433.

[142247]Avent, N. D.; Martin, P. G.: Kell typing by allele-specific
PCR

[142248](ASP). Brit. J. Haemat. 93: 728–730, 1996.

[142249]25434.

[142250]Chown, B.; Lewis, M.; Kaita, K.: A 'new' Kell blood-group
phenotype. Nature 180:

[142251]711 only, 1957.

[142252]25435.

[142253]Conneally, P. M.; Nance, W. E.; Huntzinger, R. S.: Linkage
analysis

[142254]of Kell–Sutter and PTC loci. (Abstract) Am. J. Hum. Genet.
26: 22A

[142255]only, 1974.

[142256]25436.

[142257]Abdelhak, S.; Kalatzis, V.; Heilig, R.; Compain, S.; Samson,
D.;

[142258]Vincent, C.; Weil, D.; Cruaud, C.; Sahly, I.; Leibovici, M.;
Bitner-Grindzicz,

[142259]M.; Francis, M.; Lacombe, D.; Vigneron, J.; Charachon, R.;
Boven,

[142260]K.; Bedbeder, P.; Van Regemorter, N.; Weissenbach, J.; Pe-

tit, C.:

[142261]A human homologue of the *Drosophila* eyes absent gene underlies branchio-oto-renal

[142262](BOR) syndrome and identifies a novel gene family. *Nature Genet.* 15:

[142263]157–164, 1997.

[142264]25437.

[142265]Tommerup, N.; Schempp, W.; Meinecke, P.; Pedersen, S.; Bolund,

[142266]L.; Brandt, C.; Goodpasture, C.; Guldberg, P.; Held, K.; Reinwein,

[142267]H.; Saugstad, O. D.; Scherer, G.; Skjeldal, O.; Toder, R.; Westvik,

[142268]J.; van der Hagen, C. B.; Wolf, U.: Assignment of an autosomal sex

[142269]reversal locus (SRA1) and campomelic dysplasia (CMPD1) to 17q24.3–q25.1. *Nature*

[142270]*Genet.* 4: 170–174, 1993.

[142271]25438.

[142272]Phillips, D. J.; Albertsson-Wikland, K.; Eriksson, K.; Wide, L.

[142273]: Changes in the isoforms of luteinizing hormone and fol-

lile-stimulating

[142274]hormone during puberty in normal children. J. Clin. Endocr. Metab. 82:

[142275]3103-3106, 1997.

[142276]25439.

[142277]Anderson, M. J.; Shelton, G. D.; Cavenee, W. K.; Arden, K. C.:

[142278]Embryonic expression of the tumor-associated PAX3-FKHR fusion protein

[142279]interferes with the developmental functions of Pax3. Proc. Nat. Acad.

[142280]Sci. 98: 1589-1594, 2001.

[142281]25440.

[142282]Anderson, M. J.; Viars, C. S.; Czekay, S.; Cavenee, W. K.; Arden,

[142283]K. C.: Cloning and characterization of three human fork-head genes

[142284]that comprise an FKHR-like gene subfamily. Genomics 47: 187-199,

[142285]1998.

[142286]25441.

[142287] Barr, F. G.; Galili, N.; Holick, J.; Biegel, J. A.; Rovera, G.; Emanuel, B. S.: Rearrangement of the PAX3 paired box gene in the paediatric solid tumor alveolar rhabdomyosarcoma. Nature Genet. 3: 113-117, 1993.

[142290] 25442.

[142291] Davis, R. J.; Barr, F. G.: Fusion genes resulting from alternative chromosomal translocations are overexpressed by gene-specific mechanisms in alveolar rhabdomyosarcoma. Proc. Nat. Acad. Sci. 94: 8047-8051, 1997.

[142295] 25443.

[142296] Fredericks, W. J.; Galili, N.; Mukhopadhyay, S.; Rovera, G.; Bennicelli, J.; Barr, F. G.; Rauscher, F. J., III: The PAX3-FKHR fusion protein created by the t(2;13) translocation in alveolar rhabdomyosarcomas

[142300] is a more potent transcriptional activator than PAX3.

Molec. Cell.

[142301] Biol. 15: 1522–1535, 1995.

[142302] 25444.

[142303] Galili, N.; Davis, R. J.; Fredericks, W. J.; Mukhopadhyay, S.;

[142304] Rauscher, F. J., III; Emanuel, B. S.; Rovera, G.; Barr, F. G.:

Fusion

[142305] of a fork head domain gene to PAX3 in the solid tumour
alveolar rhabdomyosarcoma. Nature

[142306] Genet. 5: 230–235, 1993.

[142307] 25445.

[142308] Khan, J.; Bittner, M. L.; Saal, L. H.; Teichmann, U.; Azorsa,
D.

[142309] O.; Gooden, G. C.; Pavan, W. J.; Trent, J. M.; Meltzer, P. S.:
cDNA

[142310] microarrays detect activation of a myogenic transcription
program

[142311] by the PAX3–FKHR fusion oncogene. Proc. Nat. Acad. Sci.
96: 13264–13269,

[142312] 1999.

[142313] 25446.

[142314]Lahat, H.; Pras, E.; Olender, T.; Avidan, N.; Ben-Asher, E.;
Man,

[142315]O.; Levy-Nissenbaum, E.; Khoury, A.; Lorber, A.; Goldman,
B.; Lancet,

[142316]D.; Eldar, M.: A missense mutation in a highly conserved
region of

[142317]CASQ2 is associated with autosomal recessive cate-
cholamine-induced

[142318]polymorphic ventricular tachycardia in Bedouin families
from Israel. Am.

[142319]J. Hum. Genet. 69: 1378-1384, 2001.

[142320]25447.

[142321]Viskin, S.; Belhassen, B.: Polymorphic ventricular tach-
yarrhythmias

[142322]in the absence of organic heart disease: classification, dif-
ferential

[142323]diagnosis, and implication for therapy. Prog. Cardiovas.
Dis. 41:

[142324]17-34, 1998.

[142325]25448.

[142326]Angle, C. R.: Congenital bowing and angulation of the
long bones. Pediatrics 13:

[142327]257–268, 1954.

[142328]25449.

[142329]Bain, A. D.; Barrett, H. S.: Congenital bowing of the long bones:

[142330]report of a case. Arch. Dis. Child. 34: 516–524, 1959.

[142331]25450.

[142332]Bell, D. M.; Leung, K. K. H.; Wheatley, S. C.; Ng, L. J.; Zhou,

[142333]S.; Ling, K. W.; Sham, M. H.; Koopman, P.; Tam, P. P. L.; Cheah, K.

[142334]S. E.: SOX9 directly regulates the type-II collagen gene. Nature

[142335]Genet. 16: 174–178, 1997.

[142336]25451.

[142337]Bi, W.; Huang, W.; Whitworth, D. J.; Deng, J. M.; Zhang, Z.; Behringer,

[142338]R. R.; de Crombrughe, B.: Haploinsufficiency of Sox9 results in

[142339]defective cartilage primordia and premature skeletal mineralization. Proc.

[142340]Nat. Acad. Sci. 98: 6698–6703, 2001.

[142341]25452.

[142342]Bishop, C. E.; Whitworth, D. J.; Qin, Y.; Agoulnik, A. I.;
Agoulnik,

[142343]J. U.; Harrison, W. R.; Behringer, R. R.; Overbeek, P. A.: A
transgenic

[142344]insertion upstream of Sox9 is associated with dominant
XX sex reversal

[142345]in the mouse. Nature Genet. 26: 490–494, 2000.

[142346]25453.

[142347]Caffey, J. P.: Prenatal bowing and thickening of tubular
bones,

[142348]with multiple cutaneous dimples in arms and legs: a con-
genital syndrome

[142349]of mechanical origin. Am. J. Dis. Child. 74: 543–562,
1947.

[142350]25454.

[142351]Cameron, F. J.; Hageman, R. M.; Cooke–Yarborough, C.;
Kwok, C.;

[142352]Goodwin, L. L.; Sillence, D. O.; Sinclair, A. H.: A novel germ
line

[142353]mutation in SOX9 causes familial campomelic dysplasia
and sex reversal. Hum.

[142354]Molec. Genet. 5: 1625–1630, 1996.

[142355]25455.

[142356]Cameron, F. J.; Sinclair, A. H.: Mutations in SRY and SOX9:
testis-determining

[142357]genes. Hum. Mutat. 9: 388–395, 1997.

[142358]25456.

[142359]Cooke, C. T.; Mulcahy, M. T.; Cullity, G. J.; Watson, M.;
Sprague,

[142360]P.: Campomelic dysplasia with sex reversal: morphological
and cytogenetic

[142361]studies of a case. Pathology 17: 526–529, 1985.

[142362]25457.

[142363]Cremin, B. J.; Orsmond, G.; Beighton, P.: Autosomal reces-
sive

[142364]inheritance in campomelic dwarfism.(Letter) Lancet I:
488–489,

[142365]1973.

[142366]25458.

[142367]Dagna Bricarelli, F.; Fraccaro, M.; Lindsten, J.; Muller, U.;

[142368]Baggio, P.; Carbone, L. D. L.; Hjerpe, A.; Lindgren, F.;

Mayerova,

[142369]A.; Ringertz, H.; Ritzen, E. M.; Rovetta, D. C.; Sicchero, C.;
Wolf,

[142370]U.: Sex-reversed XY females with campomelic dysplasia
are H-Y negative. Hum.

[142371]Genet. 57: 15-22, 1981.

[142372]25459.

[142373]Ebensperger, C.; Jager, R. J.; Lattermann, U.; Dagna
Bricarelli,

[142374]F.; Keutel, J.; Lindsten, J.; Rehder, H.; Muller, U.; Wolf, U.:
No

[142375]evidence of mutations in four candidate genes for male
sex determination/differentiation

[142376]in sex-reversed XY females with campomelic dysplasia.
Ann. Genet. 34:

[142377]233-238, 1991.

[142378]25460.

[142379]Fontaine, G.; Walbaum, R.; Farriaux, J. P.; Tilmont, P.;
Peuzin,

[142380]F.; Delecour, M.: Le conseil genetique dans la dysplasie
campomelique

[142381](a propos de deux observations). J. Genet. Hum. 28:

267–279, 1980.

[142382]25461.

[142383]Foster, J. W.; Dominguez–Steglich, M. A.; Guioli, S.; Kwok, C.;

[142384]Weller, P. A.; Stevanovic, M.; Weissenbach, J.; Mansour, S.; Young,

[142385]J. D.; Goodfellow, P. N.; Brook, J. D.; Schafer, A. J.: Campomelic

[142386]dysplasia and autosomal sex reversal caused by mutations in an SRY–related

[142387]gene. *Nature* 372: 525–530, 1994.

[142388]25462.

[142389]Friedrich, U.; Schaefer, E.; Meinecke, P.: Campomelic dysplasia

[142390]without overt campomelia. *Clin. Dysmorph.* 1: 172–178, 1992.

[142391]25463.

[142392]Gasca, S.; Canizares, J.; de Santa Barbara, P.; Mejean, C.; Poulat,

[142393]F.; Berta, P.; Boizet–Bonhoure, B.: A nuclear export signal within

[142394]the high mobility group domain regulates the nucleocyto–

plasmic translocation

[142395]of SOX9 during sexual determination. Proc. Nat. Acad. Sci.
99: 11199–11204,

[142396]2002.

[142397]25464.

[142398]Glass, R. B. J.; Rosenbaum, K. N.: Acampomelic cam-
pomelic dysplasia:

[142399]further radiographic variations. Am. J. Med. Genet. 69:
29–32, 1997.

[142400]25465.

[142401]Hall, B.; Spranger, J. W.: Campomelic dysplasia: further
elucidation

[142402]of a distinct entity. Am. J. Dis. Child. 134: 285–289, 1980.

[142403]25466.

[142404]Hoefnagel, D.; Wurster-Hill, D. H.; Dupree, W. B.;
Benirschke,

[142405]K.; Fuld, G. L.: Camptomelic dwarfism associated with XY-
gonadal

[142406]dysgenesis and chromosome anomalies. Clin. Genet. 13:
489–499, 1978.

[142407]25467.

[142408]Houston, C. S.; Opitz, J. M.; Spranger, J. W.; Macpherson, R.

[142409]J.; Reed, M. H.; Gilbert, E. F.; Herrmann, J.; Schinzel, A.:
The

[142410]campomelic syndrome: review, report of 17 cases, and
follow-up on

[142411]the currently 17-year-old boy first reported by Maroteaux
et al in

[142412]1971. Am. J. Med. Genet. 15: 3–28, 1983.

[142413]25468.

[142414]Hovmoller, M. L.; Osuna, A.; Eklof, O.; Fredga, K.; Hjerpe, A.;

[142415]Lindsten, J.; Ritzen, M.; Stanescu, V.; Svenningsen, N.:
Camptomelic

[142416]dwarfism. A genetically determined mesenchymal disorder
combined with

[142417]sex reversal. Hereditas 86: 51–62, 1977.

[142418]25469.

[142419]Huang, W.; Chung, U.; Kronenberg, H. M.; de Crombrughe, B.:

[142420]The chondrogenic transcription factor Sox9 is a target of
signaling

[142421]by the parathyroid hormone-related peptide in the growth plate of

[142422]endochondral bones. Proc. Nat. Acad. Sci. 98: 160–165, 2001.

[142423]25470.

[142424]Kanai, Y.; Koopman, P.: Structural and functional characterization

[142425]of the mouse Sox9 promoter: implications for campomelic dysplasia. Hum.

[142426]Molec. Genet. 8: 691–696, 1999.

[142427]25471.

[142428]Medema, R. H.; Kops, G. J. P. L.; Bos, J. L.; Burgering, B. M.

[142429]T.: AFX-like forkhead transcription factors mediate cell-cycle regulation

[142430]by Ras and PKB through p27(kip1). Nature 404: 782–787, 2000.

[142431]25472.

[142432]Eicher, E. M.; Womack, J. E.: Chromosomal location of soluble

[142433]glutamic-pyruvic transaminase-1 (Gpt-1) in the mouse. Biochem. Genet. 15:

[142434]1-8, 1977.

[142435]25473.

[142436]Falk, C. T.; Huss, J.: Linkage data on the chromosome 16 markers

[142437]HP and PGP: no additional support for or against the mapping of GPT.

[142438](Abstract) Cytogenet. Cell Genet. 40: 626 only, 1985.

[142439]25474.

[142440]Ferrell, R. E.: Personal Communication. Pittsburgh, Pa. 1/22/1988.

[142441]25475.

[142442]Jeremiah, S. J.; Kielty, C.; Povey, S.; McLellan, T.: Immunological

[142443]characterization of human GPT in hybrids supports assignment to chromosome 25476.

[142444](Abstract) Cytogenet. Cell Genet. 37: 498 only, 1984.

[142445]

[142446]8. Kalimanovska, V.; Majkic-Singh, N.; Stojanov, M.; Grozdanic, V.;

[142447]Vucetic, G.; Andelic, M.; Gligorovic, V.; Tomasevic, R.: Human red

[142448]cell glutamic-pyruvic transaminase polymorphism in Serbia, Yugoslavia. Hum.

[142449]Hered. 33: 319-321, 1983.

[142450]25477.

[142451]Kielty, C. M.; Povey, S.; Hopkinson, D. A.: Regulation of expression

[142452]of liver-specific enzymes: II. Activation and chromosomal localization

[142453]of soluble glutamate-pyruvate transaminase. Ann. Hum. Genet. 46:

[142454]135-143, 1982.

[142455]25478.

[142456]King, M.-C.; Go, R. C. P.; Elston, R. C.; Lynch, H. T.; Petrakis,

[142457]N. L.: Allele increasing susceptibility to human breast cancer may

[142458]be linked to the glutamate-pyruvate transaminase locus. Science 208:

[142459]406-408, 1980.

[142460]25479.

[142461]Kompf, J.: Population genetics of soluble glutamic-pyruvic

transaminase

[142462](EC:2.6.1.2): gene frequencies in southwestern Germany.

Humangenetik 14:

[142463]76–77, 1971.

[142464]25480.

[142465]Lahav, M.; Szeinberg, A.: A red-cell glutamic-pyruvic

transaminase

[142466]polymorphism in several population groups in Israel.

Hum. Hered. 22:

[142467]533–538, 1972.

[142468]25481.

[142469]Marazita, M. L.; Spence, M. A.; Sparkes, R. S.; Field, L. L.;

[142470]Crandall, B. F.; Sparkes, M. C.; Crist, M.: Linkage relations

of

[142471]GPT (glutamic-pyruvate transaminase). (Abstract) Cyto-

genet. Cell

[142472]Genet. 40: 690 only, 1985.

[142473]25482.

[142474]McLellan, T.: Two previously undetected variants of glu-

tamic-pyruvic

[142475]transaminase found by acidic polyacrylamide gel elec-

trophoresis. Am.

[142476]J. Hum. Genet. 34: 623-628, 1982.

[142477]25483.

[142478]McLellan, T.; Cannon, L. A.; Bishop, D. T.; Skolnick, M. H.:

[142479]The cumulative lod score between a breast cancer susceptibility locus

[142480]and GPT is -3.86. (Abstract) Cytogenet. Cell Genet. 37: 536-537,

[142481]1984.

[142482]25484.

[142483]Mithal, Y.; Lane, A. B.; Jenkins, T.: Absence of red cell glutamic-pyruvate

[142484]transaminase: discovery of a 'silent' allele homozygote. Am. J. Hum.

[142485]Genet. 32: 42-46, 1980.

[142486]25485.

[142487]O'Connell, P.; Nakamura, Y.; Lathrop, G. M.; Leppert, M.; Cartwright,

[142488]P.; Lalouel, J.-M.; White, R.: Three genetic linkage groups on chromosome

[142489]8. (Abstract) Cytogenet. Cell Genet. 46: 672 only, 1987.

[142490]25486.

[142491]Olaisen, B.: Genetics of the GPT system: family, mother-child

[142492]and association studies. Clin. Genet. 7: 245-254, 1975.

[142493]25487.

[142494]Pelzer, C. F.; Norum, R. A.: Identification of human red cell
[142495]glutamate-pyruvate transaminase (GPT) phenotypes by
isoelectric focusing. Am.

[142496]J. Hum. Genet. 37: 147-152, 1985.

[142497]25488.

[142498]Rocha, J.; Amorim, A.; Almeida, V. M.; Oliveira, J. P.; Leao,
[142499]M.; Tavares, M. C.; Pereira, M. S.; Vidal-Pinheiro, L.: Gene
dosage

[142500]evidence for the regional assignment of GPT
(glutamate-pyruvate transaminase;

[142501]E.C. 2.6.1.2) locus to 8q24.2-8qter. Hum. Genet. 80:
299-300, 1988.

[142502]25489.

[142503]Sanders, M. F.; King, M. C.; Lattanzio, D.; Crandall, J.; Le-
ung,

[142504]R.: Absence of linkage between HP and GPT. (Abstract)
Cytogenet.

[142505]Cell Genet. 37: 536–537, 1984.

[142506]25490.

[142507]Santachiara Benerecetti, A. S.; Beretta, M.; Pampiglione, S.:

[142508]Red cell glutamic–pyruvic transaminase polymorphism in
a sample of

[142509]the Italian population: a new variant allele: GPT(8). Hum.
Hered. 25:

[142510]276–278, 1975.

[142511]25491.

[142512]Sohocki, M. M.; Sullivan, L. S.; Harrison, W. R.; Sodergren,
E.

[142513]J.; Elder, F. F. B.; Weinstock, G.; Tanase, S.; Daiger, S. P.:
Human

[142514]glutamate pyruvate transaminase (GPT): localization to
8q24.3, cDNA

[142515]and genomic sequences, and polymorphic sites. Genomics
40: 247–252,

[142516]1997.

[142517]25492.

- [142518]Sparkes, M. C.; Crist, M.; Sparkes, R. S.: Glutamate pyruvate
transaminase null allele in seven new families. Hum.
Genet. 65:
147–148, 1983.
- [142521]25493.
- [142522]Wijnen, L. M. M.; Meera Khan, P.: Assignment of GPT to
human
chromosome 16. (Abstract) Cytogenet. Cell Genet. 32: 327
only, 1982.
- [142524]25494.
- [142525]Cremonini, N.; Graziano, E.; Chiarini, V.; Sforza, A.;
Zampa,
G. A.: Atypical McCune–Albright syndrome associated with
growth hormone–prolactin
pituitary adenoma: natural history, long–term follow–up,
and SMS 201–995--bromocriptine
combined treatment results. J. Clin. Endocr. Metab. 75:
1166–1169,
1992.
- [142530]25495.

[142531]DeChiara, T.; Robertson, E. J.; Efstratiadis, A.: Parental imprinting
[142532]of the mouse insulin-like growth factor II gene. Cell 64:
849–859,
[142533]1991.
[142534]25496.
[142535]Eddy, M. C.; Jan de Beur, S. M.; Yandow, S. M.; McAlister, W.
[142536]H.; Shore, E. M.; Kaplan, F. S.; Whyte, M. P.; Levine, M. A.:
Deficiency
[142537]of the alpha-subunit of the stimulatory G protein and severe extraskelatal
[142538]ossification. J. Bone Miner. Res. 15: 2074–2083, 2000.
[142539]25497.
[142540]Falconer, M. A.; Cope, C. L.; Robb-Smith, A. H. T.: Fibrous dysplasia
[142541]of bone with endocrine disorders and cutaneous pigmentation (Albright's
[142542]disease). Quart. J. Med. 11: 121–154, 1942.
[142543]25498.
[142544]Jackson, I. J.: A cDNA encoding tyrosinase-related protein

maps

[142545]to the brown locus in mouse. Proc. Nat. Acad. Sci. 85:
4392–4396,

[142546]1988. Note: Erratum: Proc. Nat. Acad. Sci. 86: 997 only,
1989.

[142547]25499.

[142548]Albertsen, H. M.; Smith, S. A.; Mazoyer, S.; Fujimoto, E.;
Stevens,

[142549]J.; Williams, B.; Rodriguez, P.; Cropp, C. S.; Slijepcevic, P.;
Carlson,

[142550]M.; Robertson, M.; Bradley, P.; Lawrence, E.; Harrington,
T.; Mei

[142551]Sheng, Z.; Hoopes, R.; Sternberg, N.; Brothman, A.; Calla-
han, R.;

[142552]Ponder, B. A. J.; White, R.: A physical map and candidate
genes in

[142553]the BRCA1 region on chromosome 17q12–21. Nature
Genet. 7: 472–479,

[142554]1994.

[142555]25500.

[142556]Ninomiya, Y.; Olsen, B. R.: Synthesis and characterization
of

- [142557]cDNA encoding a cartilage-specific short collagen. Proc.
Nat. Acad.
- [142558]Sci. 81: 3014-3018, 1984.
- [142559]25501.
- [142560]Khatib, Z. A.; Inaba, T.; Valentine, M.; Look, A. T.: Chro-
mosomal
- [142561]localization and cDNA cloning of the human DBP and TEF
genes. Genomics 23:
- [142562]344-351, 1994.
- [142563]25502.
- [142564]Pilla, M.; Perachon, S.; Sautel, F.; Garrido, F.; Mann, A.;
Wermuth,
- [142565]C. G.; Schwartz, J.-C.; Everitt, B. J.; Sokoloff, P.: Selective
inhibition
- [142566]of cocaine-seeking behaviour by a partial dopamine D3
receptor agonist. Nature 400:
- [142567]371-375, 1999.
- [142568]25503.
- [142569]Kaplan, R.; Jaye, M.; Burgess, W. H.; Schlaepfer, D. D.;
Haigler,
- [142570]H. T.: Cloning and expression of cDNA for human en-

donexin II, a Ca^{2+}

[142571]and phospholipid binding protein. J. Biol. Chem. 263:
8037–8043,

[142572]1988.

[142573]25504.

[142574]Arinami, T.; Ishikawa, M.; Inoue, A.; Yanagisawa, M.;
Masaki, T.;

[142575]Yoshida, M. C.; Hamaguchi, H.: Chromosomal assign-
ments of the human

[142576]endothelin family genes: the endothelin-1 gene (EDN1) to
6p23–p24,

[142577]the endothelin-2 gene (EDN2) to 1p34, and the endothe-
lin-3 gene (EDN3)

[142578]to 20q13.2–q13.3. Am. J. Hum. Genet. 48: 990–996,
1991.

[142579]25505.

[142580]Bourgeois, C.; Robert, B.; Rebourcet, R.; Mondon, F.;
Mignot, T.–M.;

[142581]Duc–Goiran, P.; Ferre, F.: Endothelin-1 and ET(A) receptor
expression

[142582]in vascular smooth muscle cells from human placenta: a
new ET(A) receptor

[142583] messenger ribonucleic acid is generated by alternative splicing of

[142584] exon 3. J. Clin. Endocr. Metab. 82: 3116–3123, 1997.

[142585] 25506.

[142586] Inoue, A.; Yanagisawa, M.; Kimura, S.; Kasuya, Y.;

Miyauchi, T.;

[142587] Goto, K.; Masaki, T.: The human endothelin family: three structurally

[142588] and pharmacologically distinct iso-peptides predicted by three separate

[142589] genes. Proc. Nat. Acad. Sci. 86: 2863–2867, 1989.

[142590] 25507.

[142591] Muller, G.; Ruppert, S.; Schmid, E.; Schutz, G.: Functional analysis

[142592] of alternatively spliced tyrosinase gene transcripts. EMBO J. 7:

[142593] 2723–2730, 1988.

[142594] 25508.

[142595] Murty, V. V. V. S.; Bouchard, B.; Mathew, S.; Vijayasaradhi, S.;

[142596] Houghton, A. N.: Assignment of the human TYRP (brown)

locus to chromosome

[142597]region 9p23 by nonradioactive in situ hybridization. Genomics 13:

[142598]227-229, 1992.

[142599]25509.

[142600]Ramsay, M.; Colman, M. A.; Jenkins, T.; Fox, M.; Chintamani,

[142601]C.; Pickard, R.; Kwon, B.: The human CAS2 locus (homologous to the

[142602]mouse b locus) maps to 9p22-pter. (Abstract) Cytogenet. Cell Genet. 58:

[142603]1943 only, 1991.

[142604]25510.

[142605]Shibahara, S.; Tomita, Y.; Yoshizawa, M.; Shibata, K.; Tagami,

[142606]H.: Identification of mutation in the pigment cell-specific gene

[142607]located at the brown locus in mouse. Pigment Cell Res. Suppl. 2:

[142608]90-95, 1992.

[142609]25511.

- [142610]Shaikh, S.; Ball, D.; Craddock, N.; Castle, D.; Hunt, N.;
Mant,
[142611]R.; Owen, M.; Collier, D.; Gill, M.: The dopamine D3 recep-
tor gene:
[142612]no association with bipolar affective disorder. J. Med.
Genet. 30:
[142613]308-309, 1993.
[142614]25512.
- [142615]Sokoloff, P.; Giros, B.; Martres, M.-P.; Bouthenet, M.-L.;
Schwartz,
[142616]J.-C.: Molecular cloning and characterization of a novel
dopamine
[142617]receptor (D-3) as a target for neuroleptics. Nature 347:
146-151,
[142618]1990.
[142619]25513.
- [142620]Spurlock, G.; Williams, J.; McGuffin, P.; Aschauer, H. N.;
Lenzinger,
[142621]E.; Fuchs, K.; Sieghart, W. C.; Meszaros, K.; Fathi, N.; Lau-
rent,
[142622]C.; Mallet, J.; Macciardi, F.; Pedrini, S.; Gill, M.; Hawi, Z.;
Gibson,

- [142623]S.; Jazin, E. E.; Yang, H.-T.; Adolfsson, R.; Pato, C. N.;
Dourado,
- [142624]A. M.; Owen, M. J.: European Multicentre Association Study
of Schizophrenia:
- [142625]a study of the DRD2 ser311-to-cys and DRD3 ser9-to-gly
polymorphisms. Am.
- [142626]J. Med. Genet. 81: 24–28, 1998.
- [142627]25514.
- [142628]Barsh, G. S.; Seeburg, P. H.; Gelinas, R. E.: The human
growth
- [142629]hormone gene family:structure and evolution of the chro-
mosomal locus. Nucleic
- [142630]Acids Res. 11: 3939–3958, 1983.
- [142631]25515.
- [142632]Halford, S.; Freedman, M. S.; Bellingham, J.; Inglis, S. L.;
Poopalasundaram,
- [142633]S.; Soni, B. G.; Foster, R. G.; Hunt, D. M.: Characterization
of
- [142634]a novel human opsin gene with wide tissue expression
and identification
- [142635]of embedded and flanking genes on chromosome 1q43.
Genomics 72:

[142636]203–208, 2001.

[142637]25516.

[142638]Haneda, M.; Chan, S. J.; Kwok, S. C. M.; Rubenstein, A. H.;
Steiner,

[142639]D. F.: Studies on mutant human insulin genes: identifica-
tion and

[142640]sequence analysis of a gene encoding (Ser-B24) insulin.
Proc. Nat.

[142641]Acad. Sci. 80: 6366–6370, 1983.

[142642]25517.

[142643]Hart, A. W.; Baeza, N.; Apelqvist, A.; Edlund, H.: Attenua-
tion

[142644]of FGF signalling in mouse beta-cells leads to diabetes.
Nature 408:

[142645]864–868, 2000.

[142646]25518.

[142647]Mammarella, S.; Romano, F.; Di Valerio, A.; Creati, B.; Es-
posito,

[142648]D. L.; Palmirotta, R.; Capani, F.; Vitullo, P.; Volpe, G.; Bat-
tista,

[142649]P.; Della Loggia, F.; Mariani–Costantini, R.; Cama, A.: In–

teraction

[142650]between the G1057D variant of IRS-2 and overweight in
the pathogenesis

[142651]of type 2 diabetes. Hum. Molec. Genet. 9: 2517-2521,
2000.

[142652]25519.

[142653]Triggs-Raine, B. L.; Kirkpatrick, R. D.; Kelly, S. L.; Norquay,

[142654]L. D.; Cattini, P. A.; Yamagata, K.; Hanley, A. J. G.; Zinman,
B.;

[142655]Harris, S. B.; Barrett, P. H.; Hegele, R. A.: HNF1-alpha
G319S, a

[142656]transactivation-deficient mutant, is associated with al-
tered dynamics

[142657]of diabetes onset in an Oji-Cree community. Proc. Nat.
Acad. Sci. 99:

[142658]4614-4619, 2002.

[142659]25520.

[142660]Tuomi, T.; Carlsson, A.; Li, H.; Isomaa, B.; Miettinen, A.;
Nilsson,

[142661]A.; Nissen, M.; Ehrnstrom, B.-O.; Forsen, B.; Snickars, B.;
Lahti,

[142662]K.; Forsblom, C.; Saloranta, C.; Taskinen, M.-R.; Groop, L.

C.: Clinical

[142663]and genetic characteristics of type 2 diabetes with and without GAD

[142664]antibodies. Diabetes 48: 150–157, 1999.

[142665]25521.

[142666]Yuan, M.; Konstantopoulos, N.; Lee, J.; Hansen, L.; Li, Z.-W.;

[142667]Karin, M.; Shoelson, S. E.: Reversal of obesity- and diet-induced

[142668]insulin resistance with salicylates or targeted disruption of Ikk-beta. Science 293:

[142669]1673–1677, 2001.

[142670]25522.

[142671]Nuytinck, L.; Dalgleish, R.; Spotila, L.; Renard, J.-P.; Van Regemorter,

[142672]N.; De Paepe, A.: Substitution of glycine-661 by serine in the alpha-1(I)

[142673]and alpha-2(I) chains of type I collagen results in different clinical

[142674]and biochemical phenotypes. Hum. Genet. 97: 324–329, 1996.

[142675]25523.

[142676]Nuytinck, L.; Freund, M.; Lagae, L.; Pierard, G. E.; Hermanns-Le, T.; De Paepe, A.: Classical Ehlers–Danlos syndrome caused by a mutation in type I collagen. *Am. J. Hum. Genet.* 66: 1398–1402, 2000.

[142679]25524.

[142680]Oliver, J. E.; Thompson, E. M.; Pope, F. M.; Nicholls, A. C.: Mutation in the carboxy-terminal propeptide of the pro- $\alpha 1(I)$ chain of type I collagen in a child with severe osteogenesis imperfecta (OI type III): possible implications for protein folding. *Hum. Mutat.* 7: 318–326, 1996.

[142684]318–326, 1996.

[142685]25525.

[142686]Pace, J. M.; Atkinson, M.; Willing, M. C.; Wallis, G.; Byers, P. H.: Deletions and duplications of Gly–Xaa–Yaa triplet repeats in the triple helical domains of type I collagen chains disrupt helix

[142689]formation and result in several types of osteogenesis im-
perfecta. Hum.

[142690]Mutat. 18: 319-326, 2001.

[142691]25526.

[142692]Pack, M.; Constantinou, C. D.; Kalia, K.; Nielsen, K. B.;
Prockop,

[142693]D. J.: Substitution of serine for alpha-1(I)-glycine 844 in a
severe

[142694]variant of osteogenesis imperfecta minimally destabilizes
the triple

[142695]helix of type I procollagen: the effects of glycine substitu-
tions

[142696]on thermal stability are either position or amino acid spe-
cific. J.

[142697]Biol. Chem. 264: 19694-19699, 1989.

[142698]25527.

[142699]Pereira, R.; Halford, K.; Sokolov, B. P.; Khillan, J. S.;
Prockop,

[142700]D. J.: Phenotypic variability and incomplete penetrance of
spontaneous

[142701]fractures in an inbred strain of transgenic mice expressing
a mutated

[142702]collagen gene (COL1A1). J. Clin. Invest. 93: 1765–1769,
1994.

[142703]25528.

[142704]Pereira, R.; Khillan, J. S.; Helminen, H. J.; Hume, E. L.;
Prockop,

[142705]D. J.: Transgenic mice expressing a partially deleted gene
for type

[142706]I procollagen (COL1A1): a breeding line with a phenotype
of spontaneous

[142707]fractures and decreased bone collagen and mineral. J.
Clin. Invest. 91:

[142708]709–716, 1993.

[142709]25529.

[142710]Pereira, R. F.; O'Hara, M. D.; Laptev, A. V.; Halford, K. W.;

[142711]Pollard, M. D.; Class, R.; Simon, D.; Livezey, K.; Prockop,
D. J.

[142712]: Marrow stromal cells as a source of progenitor cells for
nonhematopoietic

[142713]tissues in transgenic mice with a phenotype of osteogene-
sis imperfecta. Proc.

[142714]Nat. Acad. Sci. 95: 1142–1147, 1998.

[142715]25530.

[142716]Pestka, S.; Daugherty, B. L.; Jung, V.; Hotta, K.; Pestka, R.

[142717]K.: Anti-mRNA: specific inhibition of translation of single
mRNA

[142718]molecules. Proc. Nat. Acad. Sci. 81: 7525-7528, 1984.

[142719]25531.

[142720]Raghunath, M.; Steinmann, B.; Delozier-Blanchet, C.; Ex-
termann,

[142721]P.; Superti-Furga, A.: Prenatal diagnosis of collagen disor-
ders by

[142722]direct biochemical analysis of chorionic villus biopsies.
Pediat.

[142723]Res. 36: 441-448, 1994.

[142724]25532.

[142725]Redford-Badwal, D. A.; Stover, M. L.; Valli, M.; McKinstry,
M.

[142726]B.; Rowe, D. W.: Nuclear retention of COL1A1 messenger
RNA identifies

[142727]null alleles causing mild osteogenesis imperfecta. J. Clin.
Invest. 97:

[142728]1035-1040, 1996.

[142729]25533.

[142730]Retief, E.; Parker, M. I.; Retief, A. E.: Regional chromosome

[142731]mapping of human collagen genes alpha 2(I) and alpha
1(I) (COLIA2

[142732]and COLIA1). Hum. Genet. 69: 304–308, 1985.

[142733]25534.

[142734]Sainz, J.; Van Tornout, J. M.; Sayre, J.; Kaufman, F.;
Gilsanz,

[142735]V.: Association of collagen type 1 alpha-1 gene polymor-
phism with

[142736]bone density in early childhood. J. Clin. Endocr. Metab. 84:
853–855,

[142737]1999.

[142738]25535.

[142739]Schnieke, A.; Harbers, K.; Jaenisch, R.: Embryonic lethal
mutation

[142740]in mice induced by retrovirus insertion into the alpha-1 (I)
collagen

[142741]gene. Nature 304: 315–320, 1983.

[142742]25536.

[142743]Schwarze, U.; Starman, B. J.; Byers, P. H.: Redefinition of
exon
[142744]7 in the COL1A1 gene of type I collagen by an intron 8
splice-donor-site
[142745]mutation in a form of osteogenesis imperfecta: influence
of intron
[142746]splice order on outcome of splice-site mutation. Am. J.
Hum. Genet. 65:
[142747]336-344, 1999.
[142748]25537.
[142749]Sippola-Thiele, M.; Tromp, G. C.; Prockop, D. J.; Ramirez,
F.
[142750]: Assessment of small polymorphisms in defined human
collagen gene
[142751]segments. Hum. Genet. 72: 245-247, 1986.
[142752]25538.
[142753]Solomon, E.; Hiorns, L.; Sheer, D.; Rowe, D.: Confirmation
that
[142754]the type I collagen gene on chromosome 17 is
COL1A1(alpha-1(I)), using
[142755]a human genomic probe. Ann. Hum. Genet. 48: 39-42,
1984.

[142756]25539.

[142757]Solomon, E.; Hiorns, L. R.; Cheah, K. S. E.; Parkar, M.;
Weiss,

[142758]E.; Flavell, R. A.: Assignment of a human alpha-1(I)-like
collagen

[142759]gene to chromosome 12, by molecular hybridiza-
tion.(Abstract) Cytogenet.

[142760]Cell Genet. 37: 588-589, 1984.

[142761]25540.

[142762]Solomon, E.; Sykes, B.: Assignment of alpha-1 (I), alpha-2,
and

[142763]possibly alpha-1 (III), chains of human collagen to chro-
mosome 7.(Abstract) Cytogenet.

[142764]Cell Genet. 25: 205, 1979.

[142765]25541.

[142766]Steinmann, B.: Personal Communication. Zurich, Switzer-
land

[142767]11/28/1994.

[142768]25542.

[142769]Barker, P. E.; Shipp, M. A.; D'Adamio, L.; Masteller, E. L.;
Reinherz,

[142770]E. L.: The common acute lymphoblastic leukemia antigen gene maps

[142771]to chromosomal region 3(q21-q27). J. Immun. 142: 283-287, 1989.

[142772]25543.

[142773]D'Adamio, L.; Shipp, M. A.; Masteller, E. L.; Reinherz, E. L.:

[142774]Organization of the gene encoding common acute lymphoblastic leukemia

[142775]antigen (neutral endopeptidase 24.11): multiple miniexons and separate

[142776]5'-prime untranslated regions. Proc. Nat. Acad. Sci. 86: 7103-7107,

[142777]1989.

[142778]25544.

[142779]Debiec, H.; Guigonis, V.; Mougenot, B.; Decobert, F.; Haymann,

[142780]J.-P.; Bensman, A.; Deschenes, G.; Ronco, P. M.: Antenatal membranous

[142781]glomerulonephritis due to anti-neutral endopeptidase antibodies. New

[142782]Eng. J. Med. 346: 2053-2060, 2002.

[142783]25545.

[142784]Letarte, M.; Vera, S.; Tran, R.; Addis, J. B. L.; Onizuka, R. J.;

[142785]Quackenbush, E. J.; Jongeneel, C. V.; McInnes, R. R.: Com-
mon acute

[142786]lymphocytic leukemia antigen is identical to neutral en-
dopeptidase. J.

[142787]Exp. Med. 168: 1247-1253, 1988.

[142788]25546.

[142789]Shipp, M. A.; Vijayaraghavan, J.; Schmidt, E. V.; Masteller,
E.

[142790]L.; D'Adamio, L.; Hersh, L. B.; Reinherz, E. L.: Common
acute lymphoblastic

[142791]leukemia antigen (CALLA) is active neutral endopeptidase
24.11 ('enkephalinase'):

[142792]direct evidence by cDNA transfection analysis. Proc. Nat.
Acad. Sci. 86:

[142793]297-301, 1989.

[142794]25547.

[142795]Tran-Paterson, R.; Willard, H. F.; Letarte, M.: The common
acute

[142796]lymphoblastic leukemia antigen (neutral endopeptidase-

–3.4.24.11)

[142797]gene is located on human chromosome 3. Cancer Genet.

Cytogenet. 42:

[142798]129–134, 1989.

[142799]25548.

[142800]Boyd, Y.; Cockburn, D.; Holt, S.; Munro, E.; van Ommen, G.

J.;

[142801]Gillard, B.; Affara, N.; Ferguson-Smith, M.; Craig, I.: Map-

ping of

[142802]12 translocation breakpoints in the Xp21 region with re-

spect to the

[142803]locus for Duchenne muscular dystrophy. Cytogenet. Cell

Genet. 48:

[142804]28–34, 1988.

[142805]25549.

[142806]Sellar, G. C.; Blake, D. J.; Reid, K. B.: Characterization and

[142807]organization of the genes encoding the A-, B-, and C-

chains of human

[142808]complement subcomponent C1q: the complete derived

amino acid sequence

[142809]of human C1q. Biochem. J. 274: 481–490, 1991.

[142810]25550.

[142811]Topaloglu, R.; Bakkaloglu, A.; Slingsby, J. H.; Mihatsch, M. J.;

[142812]Pascual, M.; Norsworthy, P.; Morley, B. J.; Saatci, U.; Schif-ferli,

[142813]J. A.; Walport, M. J.: Molecular basis of hereditary C1q de-ficiency

[142814]associated with SLE and IgA nephropathy in a Turkish family. Kidney

[142815]Int. 50: 635-642, 1996.

[142816]25551.

[142817]Skok, J.; Solomon, E.; Reid, K. B. M.; Thompson, R. A.: Dis-tinct

[142818]genes for fibroblast and serum C1q. Nature 292: 549-551, 1981.

[142819]25552.

[142820]Hannema, A. J.; Kluin-Nelemans, J. C.; Hack, C. E.; Eeren-berg-Belmer,

[142821]A. J. M.; Mallee, C.; van Helden, H. P. T.: SLE like syndrome and

[142822]functional deficiency of C1q in members of a large family. Clin.

[142823]Exp. Immun. 55: 106–114, 1984.

[142824]25553.

[142825]McAdam, R. A.; Goundis, D.; Reid, K. B. M.: A homozygous point

[142826]mutation results in a stop codon in the C1q deficient individual. Immunogenetics 27:

[142827]259–264, 1988.

[142828]25554.

[142829]Sellar, G. C.; Cockburn, D.; Reid, K. B. M.: Localization of the

[142830]gene cluster encoding the A, B, and C chains of human C1q to 1p34.1–1p36.3. Immunogenetics 35:

[142831]214–216, 1992.

[142832]25555.

[142833]Solomon, E.; Skok, J.; Griffin, J.; Reid, K. B. M.: Human C1q

[142834]B chain (C1QB) is on chromosome 1p. (Abstract) Cytogenet. Cell Genet. 40:

[142835]749 only, 1985.

[142836]25556.

[142837]Thompson, R. A.; Haeney, M.; Reid, K. B. M.; Davis, J. G.;

White,

[142838]R. H.; Cameron, A. H.: A genetic defect of the C1q sub-component of

[142839]complement associated with childhood (immune complex) nephritis. New

[142840]Eng. J. Med. 303: 22-24, 1980.

[142841]25557.

[142842]Lahn, B. T.; Page, D. C.: Four evolutionary strata on the human

[142843]X chromosome. Science 286: 964-967, 1999.

[142844]25558.

[142845]Lee, F. A.; Issacs, H.; Strauss, J.: The 'campptomelic' syndrome.

[142846]Short life-span dwarfism with respiratory distress, hypotonia, peculiar

[142847]facies, and multiple skeletal and cartilaginous deformities. Am.

[142848]J. Dis. Child. 124: 485-496, 1972.

[142849]25559.

[142850]Lynch, S. A.; Gaunt, M. L.; Minford, A. M. B.: Campomelic dysplasia:

[142851]evidence of autosomal dominant inheritance. J. Med.
Genet. 30: 683–686,

[142852]1993.

[142853]25560.

[142854]Macpherson, R. I.; Skinner, S. A.; Donnenfeld, A. E.: Acam-
pomelic

[142855]campomelic dysplasia. Pediat. Radiol. 20: 90–93, 1989.

[142856]25561.

[142857]Mansour, S.; Hall, C. M.; Pembrey, M. E.; Young, I. D.: A
clinical

[142858]and genetic study of campomelic dysplasia. J. Med. Genet.
32: 415–420,

[142859]1995.

[142860]25562.

[142861]Maraia, R.; Saal, H. M.; Wangsa, D.: A chromosome 17q de
novo

[142862]paracentric inversion in a patient with campomelic dyspla-
sia; case

[142863]report and etiologic hypothesis. Clin. Genet. 39: 401–408,
1991.

[142864]25563.

- [142865]Maroteaux, P.; Spranger, J. W.; Opitz, J. M.; Kucera, J.;
Lowry,
[142866]R. B.; Schimke, R. N.; Kagan, S. M.: Le syndrome cam-
pomelique. Presse
[142867]Med. 22: 1157-1162, 1971.
[142868]25564.
[142869]Meyer, J.; Sudbeck, P.; Held, M.; Wagner, T.; Schmitz, M.
L.;
[142870]Bricarelli, F. D.; Eggermont, E.; Friedrich, U.; Haas, O. A.;
Kobelt,
[142871]A.; Leroy, J. G.; van Maldergem, L.; Michel, E.; Mitulla, B.;
Pfeiffer,
[142872]R. A.; Schinzel, A.; Schmidt, H.; Scherer, G.: Mutational
analysis
[142873]of the SOX9 gene in campomelic dysplasia and autosomal
sex reversal:
[142874]lack of genotype/phenotype correlations. Hum. Molec.
Genet. 6: 91-98,
[142875]1997.
[142876]25565.
[142877]Moedjono, S. J.; Crandall, B. F.; Sparkes, R. S.; Feldman, G.

[142878]M.; Austin, G. E.; Perry, S.: The campomelic syndrome in a
singleton

[142879]and monozygotic twins. Clin. Genet. 18: 397–401, 1980.

[142880]25566.

[142881]Moog, U.; Jansen, N. J. G.; Scherer, G.; Schrander-Stumpel,
C.

[142882]T. R. M.: Acampomelic campomelic syndrome. Am. J. Med.
Genet. 104:

[142883]239–245, 2001.

[142884]25567.

[142885]Morais da Silva, S.; Hacker, A.; Harley, V.; Goodfellow, P.;
Swain,

[142886]A.; Lovell-Badge, R.: Sox9 expression during gonadal de-
velopment

[142887]implies a conserved role for the gene in testis differentia-
tion in

[142888]mammals and birds. Nature Genet. 14: 62–68, 1996.

[142889]25568.

[142890]Murakami, S.; Kan, M.; McKeehan, W. L.; de Crombrughe,
B.: Up-regulation

[142891]of the chondrogenic Sox9 gene by fibroblast growth fac-

tors is mediated

[142892]by the mitogen-activated protein kinase pathway. Proc.
Nat. Acad.

[142893]Sci. 97: 1113-1118, 2000.

[142894]25569.

[142895]Ninomiya, S.; Isomura, M.; Narahara, K.; Seino, Y.; Naka-
mura,

[142896]Y.: Isolation of a testis-specific cDNA on chromosome 17q
from a

[142897]region adjacent to the breakpoint of t(12;17) observed in
a patient

[142898]with acampomelic campomelic dysplasia and sex reversal.
Hum. Molec.

[142899]Genet. 5: 69-72, 1996.

[142900]25570.

[142901]Ninomiya, S.; Yokoyama, Y.; Teraoka, M.; Mori, R.; Inoue,
C.;

[142902]Yamashita, S.; Tamai, H.; Funato, M.; Seino, Y.: A novel
mutation

[142903](296 del G) of the SOX9 gene in a patient with campomelic
syndrome

[142904]and sex reversal. Clin. Genet. 58: 224-227, 2000.

[142905]25571.

[142906]Olney, P. N.; Kean, L. S.; Graham, D.; Elsas, L. J.; May, K.
M.

[142907]: Campomelic syndrome and deletion of SOX9. Am. J. Med.
Genet. 84:

[142908]20–24, 1999.

[142909]25572.

[142910]Ozkilic, A.; Seven, M.; Yuksel, A.: A case of acampomelic
campomelic

[142911]dysplasia. Genet. Counsel. 13: 23–28, 2002.

[142912]25573.

[142913]Patel, M.; Dorman, K. S.; Zhang, Y.–H.; Huang, B.–L.;
Arnold,

[142914]A. P.; Sinsheimer, J. S.; Vilain, E.; McCabe, E. R. B.: Primate
DAX1,

[142915]SRY, and SOX9: evolutionary stratification of sex–
determination pathway. Am.

[142916]J. Hum. Genet. 68: 275–280, 2001.

[142917]25574.

[142918]Pfeifer, D.; Kist, R.; Dewar, K.; Devon, K.; Lander, E. S.; Bir–
ren,

- [142919]B.; Korniszewski, L.; Back, E.; Scherer, G.: Campomelic dysplasia translocation breakpoints are scattered over 1 Mb proximal to SOX9: evidence for an extended control region. Am. J. Hum. Genet. 65: 111–124, 1999.
- [142922]25575.
- [142923]Puck, S. M.; Haseltine, F. P.; Francke, U.: Absence of H-Y antigen in an XY female with campomelic dysplasia. Hum. Genet. 57: 23–27, 1981.
- [142926]25576.
- [142927]Rimoin, D. L.: Personal Communication. Torrance, Calif. 8/12/1976.
- [142929]25577.
- [142930]Rodriguez, J. I.: Vascular anomalies in campomelic syndrome. Am. J. Med. Genet. 46: 185–192, 1993.
- [142932]25578.

- [142933]Savarirayan, R.; Bankier, A.: Acampomelic campomelic dysplasia
with de novo 5q;17q reciprocal translocation and severe phenotype. J.
Med. Genet. 35: 597–599, 1998.
25579.
- [142937]Schimke, R. N.: XY sex–reversed campomelia--possibly an X–linked
disorder? (Letter) Clin. Genet. 16: 62–63, 1979.
25580.
- [142940]Shafai, T.; Schwartz, L.: Camptomelic syndrome in siblings. J.
Pediat. 89: 512–513, 1976.
25581.
- [142943]Spranger, J.: Advances in bone dysplasias. (Abstract) Sixth
Int. Cong. Hum. Genet. Jerusalem , 1981.
25582.
- [142946]Stuve, A.; Wiedemann, H.–R.: Congenital bowing of the long bones
in two sisters. (Letter) Lancet I: 495, 1971.

[142948]25583.

[142949]Sudbeck, P.; Schmitz, M. L.; Baeuerle, P. A.; Scherer, G.:
Sex

[142950]reversal by loss of the C-terminal transactivation domain
of human

[142951]SOX9. Nature Genet. 13: 230–232, 1996.

[142952]25584.

[142953]Thong, M.–K.; Scherer, G.; Kozlowski, K.; Haan, E.; Morris,
L.

[142954]: Acampomelic campomelic dysplasia with SOX9 mutation.
Am. J. Med.

[142955]Genet. 93: 421–425, 2000.

[142956]25585.

[142957]Thurmon, T. F.; De Fraites, E. B.; Anderson, E. E.: Familial

[142958]campomelic dwarfism. J. Pediat. 83: 841–843, 1973.

[142959]25586.

[142960]Vidal, V. P. I.; Chaboissier, M.–C.; de Rooij, D. G.; Schedl,

[142961]A.: Sox9 induces testis development in XX transgenic
mice. Nature

[142962]Genet. 28: 216–217, 2001.

[142963]25587.

[142964]Wagner, T.; Wirth, J.; Meyer, J.; Zabel, B.; Held, M.; Zimmer,

[142965]J.; Pasantes, J.; Dagna Bricarelli, F.; Keutel, J.; Hustert, E.; Wolf,

[142966]U.; Tommerup, N.; Schempp, W.; Scherer, G.: Autosomal sex reversal

[142967]and campomelic dysplasia are caused by mutations in and around the

[142968]SRY-related gene SOX9. Cell 79: 1111-1120, 1994.

[142969]25588.

[142970]Weller, S. D. V.: Hypophosphatasia with congenital dimples. Proc.

[142971]Roy. Soc. Med. 52: 637, 1959.

[142972]25589.

[142973]Wirth, J.; Wagner, T.; Meyer, J.; Pfeiffer, R. A.; Tietze, H.-U.;

[142974]Schempp, W.; Scherer, G.: Translocation breakpoints in three patients

[142975]with campomelic dysplasia and autosomal sex reversal map more than

[142976]130 kb from SOX9. Hum. Genet. 97: 186–193, 1996.

[142977]25590.

[142978]Wright, E.; Hargrave, M. R.; Christiansen, J.; Cooper, L.;
Kun,

[142979]J.; Evans, T.; Gangadharan, U.; Greenfield, A.; Koopman,
P.: The

[142980]Sry-related gene Sox9 is expressed during chondrogenesis in mouse

[142981]embryos. Nature Genet. 9: 15–20, 1995.

[142982]25591.

[142983]Albritton, L. M.; Bowcock, A. M.; Eddy, R. L.; Morton, C. C.;
Tseng,

[142984]L.; Farrer, L. A.; Cavalli-Sforza, L. L.; Shows, T. B.; Cunningham,

[142985]J. M.: The human cationic amino acid transporter (ATRC1):
physical

[142986]and genetic mapping to 13q12–q14. Genomics 12:
430–434, 1992.

[142987]25592.

[142988]Bowcock, A. M.; Gerken, S. C.; Barnes, R. I.; Shiang, R.;
Jabs,

- [142989]E. W.; Warren, A. C.; Antonarakis, S.; Retief, A. E.; Vergnaud, G.;
- [142990]Leppert, M.; Lalouel, J.-M.; White, R. L.; Cavalli-Sforza, L. L.;
- [142991]The CEPH consortium linkage map of human chromosome 13. *Genomics* 16: 486–496, 1993.
- [142992]25593.
- [142993]Kim, J. W.; Closs, E. I.; Albritton, L. M.; Cunningham, J. M.:
- [142995]Transport of cationic amino acids by the mouse ecotropic retrovirus receptor. *Nature* 352: 725–728, 1991.
- [142996]25594.
- [142998]Kozak, C. A.; Albritton, L. M.; Cunningham, J. M.: Genetic mapping of a cloned sequence responsible for susceptibility to leukemia viruses. *J. Virol.* 64: 3119–3121, 1990.
- [143001]25595.
- [143002]Oie, H. K.; Gazdar, A. F.; Lalley, P. A.; Russell, E. K.; Minna,

- [143003]J. D.; DeLarco, J.; Todaro, G. J.; Francke, U.: Mouse chromosome
- [143004]5 codes for ecotropic murine leukaemia virus cell-surface receptor. *Nature* 274:
- [143005]60-62, 1978.
- [143006]25596.
- [143007]Ruddle, N. H.; Conta, B. S.; Leinwand, L.; Kozak, C.; Ruddle, F.;
- [143008]Besmer, P.; Baltimore, D.: Assignment of the receptor for ecotropic
- [143009]murine leukemia virus to mouse chromosome 5. *J. Exp. Med.* 148: 451-465,
- [143010]1978.
- [143011]25597.
- [143012]Cook, P. W.; Piepkorn, M.; Clegg, C. H.; Plowman, G. D.; DeMay,
- [143013]J. M.; Brown, J. R.; Pittelkow, M. R.: Transgenic expression of the
- [143014]human amphiregulin gene induces a psoriasis-like phenotype. *J. Clin.*
- [143015]*Invest.* 100: 2286-2294, 1997.

[143016]25598.

[143017]Disteche, C. M.; Plowman, G. D.; Gronwald, R. G. K.; Kelly, J.;

[143018]Bowen-Pope, D.; Adler, D. A.; Murray, J. C.: Mapping of the amphiregulin

[143019]and the platelet-growth factor receptor alpha genes to the proximal

[143020]long arm of chromosome 4. (Abstract) Cytogenet. Cell Genet. 51:

[143021]990 only, 1989.

[143022]25599.

[143023]Kimura, H.; Fischer, W. H.; Schubert, D.: Structure, expression

[143024]and function of a schwannoma-derived growth factor. Nature 348:

[143025]257-260, 1990.

[143026]25600.

[143027]Gillessen-Kaesbach, G.; Demuth, S.; Thiele, H.; Theile, U.; Lich,

[143028]C.; Horsthemke, B.: A previously unrecognised phenotype characterised

- [143029]by obesity, muscular hypotonia, and ability to speak in patients with
- [143030]Angelman syndrome caused by an imprinting defect. *Eur. J. Hum. Genet.* 7: 638–644, 1999.
- [143031]25601.
- [143032]25601.
- [143033]Acosta, J.; Hettinga, J.; Fluckiger, R.; Krumrei, N.; Goldfine, A.; Angarita, L.; Halperin, J.: Molecular basis for a link between
- [143034]complement and the vascular complications of diabetes. *Proc. Nat. Acad. Sci.* 97: 5450–5455, 2000.
- [143035]25602.
- [143036]25602.
- [143037]Bickmore, W. A.; Longbottom, D.; Oghene, K.; Fletcher, J. M.; van
- [143038]Heyningen, V.: Colocalization of the human CD59 gene to 11p13 with
- [143039]the MIC11 cell surface antigen. *Genomics* 17: 129–135, 1993.
- [143040]25603.
- [143041]25603.
- [143042]Davies, A.; Simmons, D. L.; Hale, G.; Harrison, R. A.; Tighe,

H.;

[143043]Lachmann, P. J.; Waldmann, H.: CD59, an LY-6-like protein expressed

[143044]in human lymphoid cells, regulates the action of the complement membrane

[143045]attack complex on homologous cells. J. Exp. Med. 170: 637-654, 1989.

[143046]25604.

[143047]Forsberg, U. H.; Bazil, V.; Stefanova, I.; Schroder, J.: Gene

[143048]for human CD59 (likely Ly-6 homologue) is located on the short arm

[143049]of chromosome 11. Immunogenetics 30: 188-193, 1989.

[143050]25605.

[143051]Harada, R.; Okada, N.; Fujita, T.; Okada, H.: Purification of

[143052]1F5 antigen that prevents complement attack on homologous cell membranes. J.

[143053]Immun. 144: 1823-1828, 1990.

[143054]25606.

[143055]Heckl-Ostreicher, B.; Ragg, S.; Drechsler, M.; Scherthan, H.; Royer-Pokora,

[143056]B.: Localization of the human CD59 gene by fluorescence

in situ hybridization

[143057]and pulsed-field gel electrophoresis. Cytogenet. Cell
Genet. 63:

[143058]144–146, 1993.

[143059]25607.

[143060]Holt, D. S.; Botto, M.; Bygrave, A. E.; Hanna, S. M.; Walport,

[143061]M. J.; Morgan, B. P.: Targeted deletion of the CD59 gene
causes spontaneous

[143062]intravascular hemolysis and hemoglobinuria. Blood 98:
442–449, 2001.

[143063]25608.

[143064]Holt, D. S.; Powell, M. B.; Rushmere, N. K.; Morgan, B. P.:
Genomic

[143065]structure and chromosome location of the gene encoding
mouse CD59. Cytogenet.

[143066]Cell Genet. 89: 264–267, 2000.

[143067]25609.

[143068]Huppi, K.; Duncan, R.; Potter, M.: Myc-1 is centromeric to
the

[143069]linkage group Ly-6-Sis-Gdc-1 on mouse chromosome 15.
Immunogenetics 27:

[143070]215–219, 1988.

[143071]25610.

[143072]Kamiura, S.; Nolan, C. M.; Meruelo, D.: Long-range physical map

[143073]of the Ly-6 complex: mapping the Ly-6 multigene family by field-inversion

[143074]and two-dimensional gel electrophoresis. Genomics 12: 89–105, 1992.

[143075]25611.

[143076]Low, M. G.; Saltiel, A. R.: Structural and functional roles of

[143077]glycosyl-phosphatidylinositol in membranes. Science 239: 268–275,

[143078]1988.

[143079]25612.

[143080]Mahoney, J. F.; Urakaze, M.; Hall, S.; DeGasperi, R.; Chang, H.-M.;

[143081]Sugiyama, E.; Warren, C. D.; Borowitz, M.; Nicholson-Weller, A.; Rosse,

[143082]W. F.; Yeh, E. T. H.: Defective glycosylphosphatidylinositol anchor

[143083]synthesis in paroxysmal nocturnal hemoglobinuria granu-

locytes. Blood 79:

[143084]1400-1403, 1992.

[143085]25613.

[143086]Mao, M.; Yu, M.; Tong, J.-H.; Ye, J.; Zhu, J.; Huang, Q.-H.;

Fu,

[143087]G.; Yu, L.; Zhao, S.-Y.; Waxman, S.; Lanotte, M.; Wang,

Z.-Y.; Tan,

[143088]J.-Z.; Chan, S.-J.; Chen, Z.: RIG-E, a human homolog of

the murine

[143089]Ly-6 family, is induced by retinoic acid during the differ-

entiation

[143090]of acute promyelocytic leukemia cell. Proc. Nat. Acad. Sci.

93:

[143091]5910-5914, 1996.

[143092]25614.

[143093]Meri, S.; Morgan, B. P.; Davies, A.; Daniels, R. H.;

Olavesen,

[143094]M. G.; Waldmann, H.; Lachmann, P. J.: Human protectin

(CD59), an

[143095]18,000-20,000 MW complement lysis restricting factor,

inhibits C5b-8

[143096]catalysed insertion of C9 into lipid bilayers. Immunology

71: 1-9,

[143097]1990.

[143098]25615.

[143099]Meri, S.; Morgan, B. P.; Wing, M.; Jones, J.; Davies, A.; Po-
dack,

[143100]E.; Lachmann, P. J.: Human protectin (CD59), an 18-20-kD
homologous

[143101]complement restriction factor, does not restrict perforin-
mediated

[143102]lysis. J. Exp. Med. 172: 367-370, 1990.

[143103]25616.

[143104]Motoyama, N.; Okada, N.; Yamashina, M.; Okada, H.:
Paroxysmal

[143105]nocturnal hemoglobinuria due to hereditary nucleotide
deletion in

[143106]the HRF20 (CD59) gene. Europ. J. Immun. 22: 2669-2673,
1992.

[143107]25617.

[143108]Okada, N.; Harada, R.; Fujiita, T.; Okada, H.: A novel
membrane

[143109]glycoprotein capable of inhibiting membrane attack by

homologous complement. Int.

[143110]Immun. 1: 205–208, 1989.

[143111]25618.

[143112]Stubbs, L.; Carver, E.; Ashworth, L.; Lopez–Molina, L.: Location

[143113]of the DBP transcription factor gene in human and mouse.
Mammalian

[143114]Genome 7: 65–67, 1996.

[143115]25619.

[143116]Szpirer, C.; Riviere, M.; Cortese, R.; Nakamura, T.; Islam, M.

[143117]Q.; Levan, G.; Szpirer, J.: Chromosomal localization in man and rat

[143118]of the genes encoding the liver–enriched transcription factors C/EBP,

[143119]DBP, and HNF1/LFB–1 (CEBP, DBP, and transcription factor 1, TCF1,

[143120]respectively) and of the hepatocyte growth factor/scatter factor gene

[143121](HGF). Genomics 13: 293–300, 1992.

[143122]25620.

- [143123]Gersten, K. M.; Natsuka, S.; Trinchera, M.; Petryniak, B.;
Kelly,
- [143124]R. J.; Hiraiwa, N.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N.
G.;
- [143125]Lowe, J. B.: Molecular cloning, expression, chromosomal
assignment,
- [143126]and tissue-specific expression of a murine alpha-
(1,3)-fucosyltransferase
- [143127]locus corresponding to the human ELAM-1 ligand fucosyl
transferase. J.
- [143128]Biol. Chem. 270: 25047-25056, 1995.
- [143129]25621.
- [143130]Angeletti, R. H.: Chromogranins and neuroendocrine se-
cretion.
- [143131](Editorial) Lab. Invest. 55: 387-390, 1986.
- [143132]25622.
- [143133]Bhargava, G.; Russell, J.; Sherwood, L. M.: Phosphorylation
of
- [143134]parathyroid secretory protein. Proc. Nat. Acad. Sci. 80:
878-881,
- [143135]1983.

[143136]25623.

[143137]Cetin, Y.; Aunis, D.; Bader, M.-F.; Galindo, E.; Jorns, A.;
Bargsten,

[143138]G.; Grube, D.: Chromostatin, a chromogranin A-derived
bioactive peptide,

[143139]is present in human pancreatic insulin (beta) cells. Proc.
Nat. Acad.

[143140]Sci. 90: 2360-2364, 1993.

[143141]25624.

[143142]Janssen, J. W. G.; Schulz, A. S.; Steenvoorden, A. C. M.;
Schmidberger,

[143143]M.; Strehl, S.; Ambros, P. F.; Bartram, C. R.: A novel puta-
tive tyrosine

[143144]kinase receptor with oncogenic potential. Oncogene 6:
2113-2120,

[143145]1991.

[143146]25625.

[143147]Liu, E.; Hjelle, B.; Bishop, J. M.: Transforming genes in
chronic

[143148]myelogenous leukemia. Proc. Nat. Acad. Sci. 85:
1952-1956, 1988.

[143149]25626.

[143150]Lu, Q.; Lemke, G.: Homeostatic regulation of the immune system

[143151]by receptor tyrosine kinases of the Tyro 3 family. Science 293:

[143152]306–311, 2001.

[143153]25627.

[143154]Manfioletti, G.; Brancolini, C.; Avanzi, G.; Schneider, C.: The

[143155]protein encoded by a growth arrest–specific gene (gas6) is a new member

[143156]of the vitamin K–dependent proteins related to protein S, a negative

[143157]coregulator in the blood coagulation cascade. Molec. Cell. Biol. 13:

[143158]4976–4985, 1993.

[143159]25628.

[143160]O'Bryan, J. P.; Frye, R. A.; Cogswell, P. C.; Neubauer, A.; Kitch,

[143161]B.; Prokop, C.; Espinosa, R., III; Le Beau, M. M.; Earp, H. S.; Liu,

- [143162]E. T.: Axl, a transforming gene isolated from primary human myeloid leukemia cells, encodes a novel receptor tyrosine kinase. Molec. Cell. Biol. 11: 5016–5031, 1991.
- [143163]25629.
- [143164]Varnum, B. C.; Young, C.; Elliott, G.; Garcia, A.; Bartley, T. D.; Fridell, Y.-W.; Hunt, R. W.; Trail, G.; Clogston, C.; Toso, R.
- [143165]J.; Yanagihara, D.; Bennett, L.; Sylber, M.; Merewether, L. A.; Tseng, A.; Escobar, E.; Liu, E. T.; Yamane, H. K.: Axl receptor tyrosine kinase stimulated by the vitamin K-dependent protein encoded by growth-arrest-specific gene 6. Nature 373: 623–626, 1995.
- [143166]25630.
- [143167]Galbraith, G. M. P.; Pandey, J. P.: Tumor necrosis factor alpha (TNF-alpha) gene polymorphism in alopecia areata. Hum. Genet. 96:

[143175]433–436, 1995.

[143176]25631.

[143177]Tazi–Ahnini, R.; di Giovine, F. S.; McDonagh, A. J. G.; Messenger,

[143178]A. G.; Amadou, C.; Cox, A.; Duff, G. W.; Cork, M. J.: Structure and

[143179]polymorphism of the human gene for the interferon–induced p78 protein

[143180](MX1): evidence of association with alopecia areata in the Down syndrome

[143181]region. Hum. Genet. 106: 639–645, 2000.

[143182]25632.

[143183]Wilson, A. G.; di Giovine, F. S.; Blakemore, A. I. F.; Duff, G.

[143184]W.: Single base polymorphism in the human tumour necrosis factor

[143185]alpha (TNF–alpha) gene detectable by NcoI restriction of PCR product. Hum.

[143186]Molec. Genet. 1: 353 only, 1992.

[143187]25633.

[143188]Belanger, L.; Roy, S.; Allard, D.: New albumin gene 3–prime adjacent

[143189]to the alpha-1-fetoprotein locus. J. Biol. Chem. 269:

5481-5484,

[143190]1994.

[143191]25634.

[143192]Lichenstein, H. S.; Lyons, D. E.; Wurfel, M. M.; Johnson, D.

A.;

[143193]McGinley, M. D.; Leidli, J. C.; Trollinger, D. B.; Mayer, J. P.;

Wright,

[143194]S. D.; Zukowski, M. M.: Afamin is a new member of the al-

bumin, alpha-fetoprotein,

[143195]and vitamin D-binding protein gene family. J. Biol. Chem.

269: 18149-18154,

[143196]1994.

[143197]25635.

[143198]Nishio, H.; Dugaiczky, A.: Complete structure of the hu-

man alpha-albumin

[143199]gene, a new member of the serum albumin multigene

family. Proc. Nat.

[143200]Acad. Sci. 93: 7557-7561, 1996.

[143201]25636.

[143202]Song, Y.-H.; Naumova, A. K.; Liebhaber, S. A.; Cooke, N.

E.: Physical

[143203]and meiotic mapping of the region of human chromosome
4q11-q13 encompassing

[143204]the vitamin D binding protein DBP/Gc-globulin and albu-
min multigene

[143205]cluster. Genome Res. 9: 581-587, 1999.

[143206]25637.

[143207]Ettinger, S.; Weksler, M. E.; Zhou, X.; Blass, J.; Szabo, P.:

[143208]Chromosomal fragility associated with familial Alzheimer's
disease. Ann.

[143209]Neurol. 36: 190-199, 1994.

[143210]25638.

[143211]Fox, N. C.; Kennedy, A. M.; Harvey, R. J.; Lantos, P. L.;
Roques,

[143212]P. K.; Collinge, J.; Hardy, J.; Hutton, M.; Stevens, J. M.;
Warrington,

[143213]E. K.; Rossor, M. N.: Clinicopathological features of famil-
ial Alzheimer's

[143214]disease associated with the M139V mutation in the prese-
nilin 1 gene:

[143215]pedigree but not mutation specific age at onset provides

evidence

[143216]for a further genetic factor. Brain 120: 491–501, 1997.

[143217]25639.

[143218]Seri, M.; Celli, I.; Betsos, N.; Claudiani, F.; Camera, G.;

Romeo,

[143219]G.: A cys634gly substitution of the RET proto-oncogene in
a family

[143220]with recurrence of multiple endocrine neoplasia type 2A
and cutaneous

[143221]lichen amyloidosis. Clin. Genet. 51: 86–90, 1997.

[143222]25640.

[143223]Georgakopoulos, A.; Marambaud, P.; Efthimiopoulos, S.;

Shioi,

[143224]J.; Cui, W.; Li, H.-C.; Schutte, M.; Gordon, R.; Holstein, G.

R.;

[143225]Martinelli, G.; Mehta, P.; Friedrich, V. L., Jr.; Robakis, N. K.:

[143226]Presenilin-1 forms complexes with the cadherin/catenin
cell-cell adhesion

[143227]system and is recruited to intercellular and synaptic con-
tacts. Molec.

[143228]Cell 4: 893–902, 1999.

[143229]25641.

[143230]Grilli, M.; Diodato, E.; Lozza, G.; Brusa, R.; Casarini, M.;
Uberti,

[143231]D.; Rozmahel, R.; Westaway, D.; St George-Hyslop, P.;
Memo, M.; Ongini,

[143232]E.: Presenilin-1 regulates the neuronal threshold to exci-
totoxicity

[143233]both physiologically and pathologically. Proc. Nat. Acad.
Sci. 97:

[143234]12822-12827, 2000.

[143235]25642.

[143236]Gustafson, L.; Brun, A.; Englund, E.; Hagnell, O.; Nilsson,
K.;

[143237]Stensmyr, M.; Ohlin, A.-K.; Abrahamson, M.: A 50-year
perspective

[143238]of a family with chromosome-14-linked Alzheimer's dis-
ease. Hum. Genet. 102:

[143239]253-257, 1998.

[143240]25643.

[143241]Handler, M.; Yang, X.; Shen, J.: Presenilin-1 regulates neu-
ronal

[143242]differentiation during neurogenesis. Development 127:
2593–2606,

[143243]2000.

[143244]25644.

[143245]Hartmann, D.: From Alzheimer's disease to skin tumors:
the catenin

[143246]connection. Proc. Nat. Acad. Sci. 98: 10522–10523, 2001.

[143247]25645.

[143248]Harvey, R. J.; Ellison, D.; Hardy, J.; Hutton, M.; Roques, P.

[143249]K.; Collinge, J.; Fox, N. C.; Rossor, M. N.: Chromosome 14
familial

[143250]Alzheimer's disease: the clinical and neuropathological
characteristics

[143251]of a family with a leucine-to-serine (L250S) substitution at
codon

[143252]250 of the presenilin 1 gene. J. Neurol. Neurosurg. Psy-
chiat. 64:

[143253]44–49, 1998.

[143254]25646.

[143255]Hsiao, K.; Chapman, P.; Nilsen, S.; Eckman, C.; Harigaya,
Y.;

[143256]Younkin, S.; Yang, F.; Cole, G.: Correlative memory deficits, A- β elevation, and amyloid plaques in transgenic mice. Science 274: 99–103, 1996.

[143259]25647.

[143260]Hull, M.; Fiebich, B. L.; Dykieriek, P.; Schmidtke, K.; Nitzsche, E.; Orszagh, M.; Deuschl, G.; Moser, E.; Schumacher, M.; Lucking, C.; Berger, M.; Bauer, J.: Early-onset Alzheimer's disease due to mutations of the presenilin-1 gene on chromosome 14: a 7-year follow-up of a patient with a mutation at codon 139. Europ. Arch. Psychiat. Clin. Neurosci. 248: 123–129, 1998.

[143266]25648.

[143267]Jarrett, J. T.; Berger, E. P.; Lansbury, P. T.: The carboxy terminus of the β amyloid protein is critical for the seeding of amyloid

[143269]formation: implications for the pathogenesis of
Alzheimer's disease. Biochemistry 32:

[143270]4693–4697, 1993.

[143271]25649.

[143272]Johnson, K. A.; Lopera, F.; Jones, K.; Becker, A.; Sperling,
R.;

[143273]Hilson, J.; Londono, J.; Siegert, I.; Arcos, M.; Moreno, S.;
Madrigal,

[143274]L.; Ossa, J.; Pineda, N.; Ardila, A.; Roselli, M.; Albert, M. S.;

[143275]Kosik, K. S.; Rios, A.: Presenilin-1-associated abnormali-
ties in

[143276]regional cerebral perfusion. Neurology 56: 1545–1551,
2001.

[143277]25650.

[143278]Kamal, A.; Almenar-Queralt, A.; LeBlanc, J. F.; Roberts, E.
A.;

[143279]Goldstein, L. S. B.: Kinesin-mediated axonal transport of a
membrane

[143280]compartment containing beta-secretase and presenilin-1
requires APP. Nature 414:

[143281]643–648, 2001.

[143282]25651.

[143283]Kamal, A.; Stokin, G. B.; Yang, Z.; Xia, C.; Goldstein, L. S.

[143284]: Axonal transport of amyloid precursor protein is mediated by direct

[143285]binding to the kinesin light chain subunit of kinesin-I.
Neuron 28:

[143286]449–459, 2000.

[143287]25652.

[143288]Kang, D. E.; Soriano, S.; Xia, X.; Eberhart, C. G.; De
Strooper,

[143289]B.; Zheng, H.; Koo, E. H.: Presenilin couples the paired
phosphorylation

[143290]of beta-catenin independent of Axin: implications for
beta-catenin

[143291]activation in tumorigenesis. Cell 110: 751–762, 2002.

[143292]25653.

[143293]Kopan, R.; Goate, A.: A common enzyme connects Notch
signaling

[143294]and Alzheimer's disease. Genes Dev. 14: 2799–2806,
2000.

[143295]25654.

[143296]Kovacs, D. M.; Fausett, H. J.; Page, K. J.; Kim, T.-W.; Moir,
 [143297]R. D.; Merriam, D. E.; Hollister, R. D.; Hallmark, O. G.;
 Mancini,
 [143298]R.; Felsenstein, K. M.; Hyman, B. T.; Tanzi, R. E.; Wasco,
 W.: Alzheimer-associated
 [143299]presenilins 1 and 2: neuronal expression in brain and lo-
 calization
 [143300]to intracellular membranes in mammalian cells. Nature
 Med. 2: 224-229,
 [143301]1996.
 [143302]25655.
 [143303]Kwok, J. B. J.; Taddei, K.; Hallupp, M.; Fisher, C.; Brooks,
 W.
 [143304]S.; Broe, G. A.; Hardy, J.; Fulham, M. J.; Nicholson, G. A.;
 Stell,
 [143305]R.; St. George Hyslop, P. H.; Fraser, P. E.; and 6 others:
 Two
 [143306]novel (M233T and R278T) presenilin-1 mutations in early-
 onset Alzheimer's
 [143307]disease pedigrees and preliminary evidence for associa-
 tion of presenilin-1
 [143308]mutations with a novel phenotype. Neuroreport 8:

1537–1542, 1997.

[143309]25656.

[143310]Lambert, J.-C.; Mann, D. M. A.; Harris, J. M.; Chartier-Harlin,

[143311]M.-C.; Cumming, A.; Coates, J.; Lemmon, H.; StClair, D.; Iwatsubo,

[143312]T.; Lendon, C.: The -48 C/T polymorphism in the presenilin 1 promoter

[143313]is associated with an increased risk of developing Alzheimer's disease

[143314]and an increased A-beta load in brain. J. Med. Genet. 38: 353–355,

[143315]2001.

[143316]25657.

[143317]Leissring, M. A.; Akbari, Y.; Fanger, C. M.; Cahalin, M. D.; Mattson,

[143318]M. P.; LaFerla, F. M.: Capacitative calcium entry deficits and elevated

[143319]luminal calcium content in mutant presenilin-1 knockin mice. J. Cell

[143320]Biol. 149: 793–797, 2000.

[143321]25658.

- [143322]Lemere, C. A.; Lopera, F.; Kosik, K. S.; Lendon, C. L.; Ossa,
[143323]J.; Saido, T. C.; Yamaguchi, H.; Ruiz, A.; Martinez, A.;
Madrigal,
[143324]L.; Hincapie, L.; Arango, J. C.; Anthony, D. C.; Koo, E. H.;
Goate,
[143325]A. M.; Selkoe, D. J.; Arango, J. C.: The E280A presenilin 1
Alzheimer
[143326]mutation produces increased A-beta-42 deposition and
severe cerebellar
[143327]pathology. Nature Med. 2: 1146-1150, 1996.
[143328]25659.
[143329]Lewis, P. A.; Perez-Tur, J.; Golde, T. E.; Hardy, J.: The pre-
senilin
[143330]1 C92S mutation increases A-beta-42 production.
Biochem. Biophys.
[143331]Res. Commun. 277: 261-263, 2000.
[143332]25660.
[143333]Li, J.; Xu, M.; Zhou, H.; Ma, J.; Potter, H.: Alzheimer prese-
nilins
[143334]in the nuclear membrane, interphase kinetochores, and
centrosomes

[143335] suggest a role in chromosome segregation. Cell 90:
917–927, 1997.

[143336] 25661.

[143337] Li, Y.–M.; Lai, M.–T.; Xu, M.; Huang, Q.; DiMuzio–Mower,
J.; Sardana,

[143338] M. K.; Shi, X.–P.; Yin, K.–C.; Shafer, J. A.; Gardell, S. J.: Pre–
senilin

[143339] 1 is linked with gamma–secretase activity in the detergent
solubilized

[143340] state. Proc. Nat. Acad. Sci. 97: 6138–6143, 2000.

[143341] 25662.

[143342] Matsubara–Tsutsui, M.; Yasuda, M.; Yamagata, H.; No–
mura, T.; Taguchi,

[143343] K.; Kohara, K.; Miyoshi, K.; Miki, T.: Molecular evidence of
presenilin

[143344] 1 mutation in familial early onset dementia. Am. J. Med.
Genet. 114:

[143345] 292–298, 2002.

[143346] 25663.

[143347] Hofstra, R. M. W.; Sijmons, R. H.; Stelwagen, T.; Stulp, R.
P.;

[143348]Kousseff, B. G.; Lips, C. J. M.; Steijlen, P. M.; Van Voorst
Vader,

[143349]P. C.; Buys, C. H. C. M.: RET mutation screening in familial
cutaneous

[143350]lichen amyloidosis and in skin amyloidosis associated with
multiple

[143351]endocrine neoplasia. J. Invest. Derm. 107: 215–218, 1996.

[143352]25664.

[143353]Bordet, T.; Lesbordes, J.-C.; Rouhani, S.; Castelnau-
Ptakhine,

[143354]L.; Schmalbruch, H.; Haase, G.; Kahn, A.: Protective effects
of cardiotrophin-1

[143355]adenoviral gene transfer on neuromuscular degeneration
in transgenic

[143356]ALS mice. Hum. Molec. Genet. 10: 1925–1933, 2001.

[143357]25665.

[143358]Narahara, K.; Kikkawa, K.; Kimira, S.; Kimoto, H.; Ogata,
M.;

[143359]Kasai, R.; Hamawaki, M.; Matsuoka, K.: Regional mapping
of catalase

[143360]and Wilms tumor--aniridia, genitourinary abnormalities,
and mental

- [143361]retardation triad loci to the chromosome segment
11p1305–p1306. Hum.
- [143362]Genet. 66: 181–185, 1984.
- [143363]25666.
- [143364]Turleau, C.; de Grouchy, J.; Nihoul–Fekete, C.; Dufier, J. L.;
- [143365]Chavin–Colin, F.; Junien, C.: Del11p13/nephroblastoma
without aniridia. Hum.
- [143366]Genet. 67: 455–456, 1984.
- [143367]25667.
- [143368]Newport, M. J.; Huxley, C. M.; Huston, S.; Hawrylowicz, C.
M.;
- [143369]Oostra, B. A.; Williamson, R.; Levin, M.: A mutation in the
interferon–gamma–receptor
- [143370]gene and susceptibility to mycobacterial infection. New
Eng. J. Med. 335:
- [143371]1941–1949, 1996.
- [143372]25668.
- [143373]Cowan, C. A.; Yokoyama, N.; Bianchi, L. M.; Henkemeyer,
M.; Frittsch,
- [143374]B.: EphB2 guides axons at the midline and is necessary for
normal

[143375]vestibular function. Neuron 26: 417–430, 2000.

[143376]25669.

[143377]Novick, D.; Orchansky, P.; Revel, M.; Rubinstein, M.: The human

[143378]interferon–gamma receptor: purification, characterization, and preparation

[143379]of antibodies. J. Biol. Chem. 262: 8483–8487, 1987.

[143380]25670.

[143381]Orchansky, P.; Rubinstein, M.; Fischer, D. G.: The interferon–gamma

[143382]receptor in human monocytes is different from the one in nonhematopoietic

[143383]cells. J. Immun. 136: 169–173, 1986.

[143384]25671.

[143385]Papanicolaou, G. J.; Parsa, N. Z.; Meltzer, P. S.; Trent, J. M.

[143386]: Assignment of interferon gamma receptor (IFNGR1) to human chromosome

[143387]bands 6q24.1–q24.2 by in situ hybridization. Cytogenet. Cell Genet. 76:

[143388]181–182, 1997. Note: Erratum: Cytogenet. Cell Genet. 78: 132 only,

[143389]1997.

[143390]25672.

[143391]Pierre–Audigier, C.; Jouanguy, E.; Lamhamedi, S.; Altare, F.;

[143392]Rauzier, J.; Vincent, V.; Canioni, D.; Emile, J. F.; Fischer, A.;

[143393]Blanche, S.; Gaillard, J. L.; Casanova, J. L.: Fatal disseminated

[143394]Mycobacterium smegmatis infection in a child with inherited interferon

[143395]gamma receptor deficiency. Clin. Infect. Dis. 24: 982–984, 1997.

[143396]25673.

[143397]Rashidbaigi, A.; Langer, J. A.; Jung, V.; Jones, C.; Morse, H.

[143398]G.; Tischfield, J. A.; Trill, J. J.; Kung, H.–F.; Pestka, S.: The

[143399]gene for the human immune interferon receptor is located on chromosome

[143400]6. Proc. Nat. Acad. Sci. 83: 384–388, 1986.

[143401]25674.

[143402]Rettig, W. J.; Grzeschik, K.–H.; Yenamandra, A. K.; Garcia, E.;

[143403]Old, L. J.: Definition of selectable cell surface markers for human

[143404]chromosomes and chromosome segments in rodent-human hybrids. Somat.

[143405]Cell Molec. Genet. 14: 223-231, 1988.

[143406]25675.

[143407]Shankaran, V.; Ikeda, H.; Bruce, A. T.; White, J. M.; Swanson,

[143408]P. E.; Old, L. J.; Schreiber, R. D.: IFN-gamma and lymphocytes prevent

[143409]primary tumour development and shape tumour immunogenicity. Nature 410:

[143410]1107-1111, 2001.

[143411]25676.

[143412]Bauer, R.; Imhof, A.; Pscherer, A.; Kopp, H.; Moser, M.; Seegers,

[143413]S.; Kerscher, M.; Tainsky, M. A.; Hofstaedter, F.; Buettner, R.:

[143414]The genomic structure of the human AP-2 transcription factor. Nucleic

[143415]Acids Res. 22: 1413-1420, 1994.

[143416]25677.

[143417]Buettner, R.; Kannan, P.; Imhof, A.; Bauer, R.; Yim, S. O.;
Glockshuber,

[143418]R.; Van Dyke, M. W.; Tainsky, M. A.: An alternatively
spliced mRNA

[143419]from the AP-2 gene encodes a negative regulator of tran-
scriptional

[143420]activation by AP-2. Molec. Cell. Biol. 13: 4174-4185,
1993.

[143421]25678.

[143422]Ahmad, I.: Mash-1 is expressed during ROD photorecep-
tor differentiation

[143423]and binds an E-box, E(opsin-1), in the rat opsin gene.
Brain Res.

[143424]Dev. Brain Res. 90: 184-189, 1995.

[143425]25679.

[143426]Ball, D. W.; Azzoli, C. G.; Baylin, S. B.; Chi, D.; Dou, S.; Do-
nis-Keller,

[143427]H.; Cumaraswamy, A.; Borges, M.; Nelkin, B. D.: Identifica-
tion of

[143428]a human achaete-scute homolog highly expressed in neu-
roendocrine tumors. Proc.

[143429]Nat. Acad. Sci. 90: 5648–5652, 1993.

[143430]25680.

[143431]Guillemot, F.; Lo, L.-C.; Johnson, J. E.; Auerbach, A.; Anderson,

[143432]D. J.; Joyner, A. L.: Mammalian achaete–scute homolog 1 is required

[143433]for the early development of olfactory and autonomic neurons. Cell 75:

[143434]463–476, 1993.

[143435]25681.

[143436]Renault, B.; Lieman, J.; Ward, D.; Krauter, K.; Kucherlapati, R.

[143437]: Localization of the human achaete–scute homolog gene (ASCL1) distal

[143438]to phenylalanine hydroxylase (PAH) and proximal to tumor rejection

[143439]antigen (TRA1) on chromosome 12q22–q23. Genomics 30: 81–83, 1995.

[143440]25682.

[143441]Deen, P. M. T.; Weghuis, D. O.; Geurts van Kessel, A.; Wieringa,

[143442]B.; van Os, C. H.: The human gene for water channel
aquaporin 1 (AQP1)

[143443]is localized on chromosome 7p15–p14. Cytogenet. Cell
Genet. 65:

[143444]243–246, 1994.

[143445]25683.

[143446]de Groot, B. L.; Grubmuller, H.: Water permeation across
biological

[143447]membranes: mechanism and dynamics of aquaporin–1
and Glpf. Science 294:

[143448]2353–2357, 2001.

[143449]25684.

[143450]de la Chapelle, A.; Vuopio, P.; Sanger, R.; Teesdale, P.:
Monosomy–7

[143451]and the Colton blood–groups. (Letter) Lancet II: 817 only,
1975.

[143452]25685.

[143453]Denker, B. M.; Smith, B. L.; Kuhajda, F. P.; Agre, P.: Identi-
fication,

[143454]purification, and partial characterization of a novel M(r)
28,000

[143455]integral membrane protein from erythrocytes and renal tubules. J.

[143456]Biol. Chem. 263: 15634–15642, 1988.

[143457]25686.

[143458]Fushimi, K.; Uchida, S.; Hara, Y.; Hirata, Y.; Marumo, F.; Sasaki,

[143459]S.: Cloning and expression of apical membrane water channel of rat

[143460]kidney collecting tubule. Nature 361: 549–552, 1993.

[143461]25687.

[143462]Hoffman, J. F.: Aquaporin: a wee burn runs through it. (Editorial) J.

[143463]Clin. Invest. 92: 1604–1605, 1993.

[143464]25688.

[143465]Keen, T. J.; Inglehearn, C. F.; Patel, R. J.; Green, E. D.; Peluso,

[143466]D. C.; Bhattacharya, S. S.: Localization of the aquaporin 1 (AQP1)

[143467]gene within a YAC contig containing the polymorphic markers D7S632

[143468]and D7S526. Genomics 25: 599–600, 1995.

[143469]25689.

[143470]King, L. S.; Choi, M.; Fernandez, P. C.; Cartron, J.-P.; Agre,

[143471]P.: Defective urinary concentrating ability due to a complete deficiency

[143472]of aquaporin-1. New Eng. J. Med. 345: 175-179, 2001.

[143473]25690.

[143474]King, L. S.; Nielsen, S.; Agre, P.; Brown, R. H.: Decreased pulmonary

[143475]vascular permeability in aquaporin-1-null humans. Proc. Nat. Acad.

[143476]Sci. 99: 1059-1063, 2002.

[143477]25691.

[143478]Knepper, M. A.: The aquaporin family of molecular water channels. Proc.

[143479]Nat. Acad. Sci. 91: 6255-6258, 1994.

[143480]25692.

[143481]Ma, T.; Yang, B.; Gillespie, A.; Carlson, E. J.; Epstein, C. J.;

[143482]Verkman, A. S.: Severely impaired urinary concentrating ability in

[143483]transgenic mice lacking aquaporin-1 water channels. J.

Biol. Chem. 273:

[143484]4296–4299, 1998.

[143485]25693.

[143486]Maurel, C.; Reizer, J.; Schroeder, J. I.; Chrispeels, M.: The

[143487]vacuolar membrane protein gamma-TIP creates water
specific channels

[143488]in *Xenopus* oocytes. EMBO J. 12: 2241–2247, 1993.

[143489]25694.

[143490]Moon, C.; Preston, G. M.; Griffin, C. A.; Jabs, E. W.; Agre, P.

[143491]: The human aquaporin-CHIP gene: structure, organiza-
tion, and chromosomal

[143492]localization. J. Biol. Chem. 268: 15772–15778, 1993.

[143493]25695.

[143494]Moon, C.; Williams, J. B.; Preston, G. M.; Copeland, N. G.;
Gilbert,

[143495]D. J.; Nathans, D.; Jenkins, N. A.; Agre, P.: The mouse
Aquaporin-1

[143496]gene. Genomics 30: 354–357, 1995.

[143497]25696.

[143498]Murata, K.; Mitsuoka, K.; Hirai, T.; Walz, T.; Agre, P.; Heymann,

[143499]J. B.; Engel, A.; Fujiyoshi, Y.: Structural determinants of water

[143500]permeation through aquaporin-1. Nature 407: 599-605, 2000.

[143501]25697.

[143502]Parsons, S. F.; Jones, J.; Anstee, D. J.; Judson, P. A.; Gardner,

[143503]B.; Wiener, E.; Poole, J.; Illum, N.; Wickramasinghe, S. N.: A novel

[143504]form of congenital dyserythropoietic anemia associated with deficiency

[143505]of erythroid CD44 and a unique blood group phenotype [In(a-b-), Co(a-b-)]. Blood 83:

[143506]860-868, 1994.

[143507]25698.

[143508]Pasquali, F.; Bernasconi, P.; Casalone, R.; Fraccaro, M.; Bernasconi,

[143509]C.; Lazzarino, M.; Morra, E.; Alessandrino, E. P.; Marchi, M. A.;

[143510]Sanger, R.: Pathogenic significance of 'pure' monosomy 7

in myeloproliferative

[143511]disorders: analysis of 14 cases.. Hum. Genet. 62: 40–51,
1982.

[143512]25699.

[143513]Preston, G. M.; Agre, P.: Isolation of the cDNA for erythro-
cyte

[143514]integral membrane protein of 28 kilodaltons: member of
an ancient

[143515]channel family. Proc. Nat. Acad. Sci. 88: 11110–11114,
1991.

[143516]25700.

[143517]Preston, G. M.; Smith, B. L.; Zeidel, M. L.; Moulds, J. J.;
Agre,

[143518]P.: Mutations in aquaporin-1 in phenotypically normal hu-
mans without

[143519]functional CHIP water channels. Science 265: 1585–1587,
1994.

[143520]25701.

[143521]Smith, B. L.; Baumgarten, R.; Nielsen, S.; Raben, D.; Zeidel,

[143522]M. L.; Agre, P.: Concurrent expression of erythroid and re-
nal aquaporin

[143523]CHIP and appearance of water channel activity in perinatal

rats. J.

[143524]Clin. Invest. 92: 2035–2041, 1993.

[143525]25702.

[143526]Smith, B. L.; Preston, G. M.; Spring, F.; Anstee, D. J.; Agre,

[143527]P.: Human red blood cell aquaporin CHIP: I. Molecular
characterization

[143528]of ABH and Colton blood group antigens. J. Clin. Invest.
94: 1043–1049,

[143529]1994.

[143530]25703.

[143531]Sui, H.; Han, B.–G.; Lee, J. K.; Walian, P.; Jap, B. K.: Struc-
tural

[143532]basis of water-specific transport through the AQP1 water
channel. Nature 414:

[143533]872–878, 2001.

[143534]25704.

[143535]Tang, W.; Cai, S.–P.; Eng, B.; Poon, M.–C.; Waye, J. S.; Il-
lum,

[143536]N.; Chui, D. H. K.: Expression of embryonic zeta-globin
and epsilon-globin

[143537]chains in a 10-year-old girl with congenital anemia. Blood

- 81: 1636–1640,
[143538]1993.
- [143539]25705.
- [143540]Wickramasinghe, S. N.; Illum, N.; Wimberley, P. D.: Con-
genital
[143541]dyserythropoietic anaemia with novel intra-erythroblastic
and intra-erythrocytic
[143542]inclusions. *Brit. J. Haemat.* 79: 322–330, 1991.
- [143543]25706.
- [143544]Yang, B.; Ma, T.; Verkman, A. S.: Erythrocyte water perme-
ability
[143545]and renal function in double knockout mice lacking aqua-
porin-1 and
[143546]aquaporin-3. *J. Biol. Chem.* 276: 624–628, 2001.
- [143547]25707.
- [143548]de The, H.; Chomienne, C.; Lanotte, M.; Degos, L.; Dejean,
A.:
[143549]The t(15;17) translocation of acute promyelocytic
leukaemia fuses
[143550]the retinoic acid receptor alpha gene to a novel tran-
scribed locus. *Nature* 347:

[143551]558–561, 1990.

[143552]25708.

[143553]de The, H.; Lavau, C.; Marchio, A.; Chomienne, C.; Degos, L.; Dejean,

[143554]A.: The PML–RAR–alpha fusion mRNA generated by the t(15;17) translocation

[143555]in acute promyelocytic leukemia encodes a functionally altered RAR. Cell 66:

[143556]675–684, 1991.

[143557]25709.

[143558]Di Croce, L.; Raker, V. A.; Corsaro, M.; Fazi, F.; Fanelli, M.;

[143559]Faretta, M.; Fuks, F.; Lo Coco, F.; Kouzarides, T.; Nervi, C.; Minucci,

[143560]S.; Pelicci, P. G.: Methyltransferase recruitment and DNA hypermethylation

[143561]of target promoters by an oncogenic transcription factor. Science 295:

[143562]1079–1082, 2002.

[143563]25710.

[143564]Dyck, J. A.; Maul, G. G.; Miller, W. H., Jr.; Chen, J. D.; Kak-izuka,

[143565]A.; Evans, R. M.: A novel macromolecular structure is a target of
[143566]the promyelocyte-retinoic acid receptor oncoprotein. Cell
76: 333-343,
[143567]1994.
[143568]25711.
[143569]Goddard, A. D.; Borrow, J.; Freemont, P. S.; Solomon, E.:
Characterization
[143570]of a zinc finger gene disrupted by the t(15;17) in acute
promyelocytic
[143571]leukemia. Science 254: 1371-1374, 1991.
[143572]25712.
[143573]Goddard, A. D.; Yuan, J. Q.; Fairbairn, L.; Dexter, M.; Bor-
row,
[143574]J.; Kozak, C.; Solomon, E.: Cloning of the murine homolog
of the
[143575]leukemia-associated PML gene. Mammalian Genome 6:
732-737, 1995.
[143576]25713.
[143577]Grignani, F.; Ferrucci, P. F.; Testa, U.; Talamo, G.; Fagioli,
[143578]M.; Alcalay, M.; Mencarelli, A.; Grignani, F.; Peschle, C.;

Nicoletti,

[143579]I.; Pelicci, P. G.: The acute promyelocytic leukemia-specific
PML-RAR-alpha

[143580]fusion protein inhibits differentiation and promotes survival of myeloid

[143581]precursor cells. Cell 74: 423-431, 1993.

[143582]25714.

[143583]Diverio, D.; Lo Coco, F.; D'Adamo, F.; Biondi, A.; Fagioli,
M.;

[143584]Grignani, F.; Rambaldi, A.; Rossi, V.; Avvisati, G.; Petti, M.
C.;

[143585]Testi, A. M.; Liso, V.; Specchia, G.; Fioritoni, G.; Recchia,
A.;

[143586]Frassoni, F.; Ciolli, S.; Pelicci, P. G.: Identification of DNA
rearrangements

[143587]at the retinoic acid receptor-alpha (RAR-alpha) locus in all
patients

[143588]with acute promyelocytic leukemia and mapping of APL
breakpoints within

[143589]the RAR-alpha second intron. Blood 79: 3331-3336,
1992.

[143590]25715.

[143591]Khan, M. M.; Nomura, T.; Kim, H.; Kaul, S. C.; Wadhwa, R.;
Shinagawa,

[143592]T.; Ichikawa-Iwata, E.; Zhong, S.; Pandolfi, P. P.; Ishii, S.:
Role

[143593]of PML and PML-RAR-alpha in Mad-mediated transcrip-
tional repression. Molec.

[143594]Cell 7: 1233-1243, 2001.

[143595]25716.

[143596]Lo Coco, F.; Diverio, D.; Falini, B.; Biondi, A.; Nervi, C.;
Pelicci,

[143597]P. G.: Genetic diagnosis and molecular monitoring in the
management

[143598]of acute promyelocytic leukemia. Blood 94: 12-22, 1999.

[143599]25717.

[143600]Pearson, M.; Carbone, R.; Sebastiani, C.; Cioce, M.; Fagioli,

[143601]M.; Saito, S.; Higashimoto, Y.; Appella, E.; Minucci, S.;
Pandolfi,

[143602]P. P.; Pelicci, P. G.: PML regulates p53 acetylation and pre-
mature

[143603]senescence induced by oncogenic Ras. Nature 406:
207-210, 2000.

[143604]25718.

[143605]Tong, J.-H.; Dong, S.; Geng, J.-P.; Huang, W.; Wang, Z.-Y.;
Sun,

[143606]G.-L.; Chen, S.-J.; Chen, Z.; Larsen, C.-J.; Berger, R.:
Molecular

[143607]rearrangements of the MYL gene in acute promyelocytic
leukemia (APL,

[143608]M3) define a breakpoint cluster region as well as some
molecular variants. Oncogene 7:

[143609]311-316, 1992.

[143610]25719.

[143611]Wang, Z. G.; Delva, L.; Gaboli, M.; Rivi, R.; Giorgio, M.;
Cordon-Cardo,

[143612]C.; Grosveld, F.; Pandolfi, P. P.: Role of PML in cell growth
and

[143613]the retinoic acid pathway. Science 279: 1547-1551, 1998.

[143614]25720.

[143615]Bunz, F.; Kobayashi, R.; Stillman, B.: cDNAs encoding the
large

[143616]subunit of human replication factor C. Proc. Nat. Acad.
Sci. 90:

[143617]11014-11018, 1993.

[143618]25721.

[143619]Lossie, A. C.; Haugen, B. R.; Wood, W. M.; Camper, S. A.;
Gordon,

[143620]D. F.: Chromosomal localization of the large subunit of
mouse replication

[143621]factor C in the mouse and human. Mammalian Genome 6:
58-59, 1995.

[143622]25722.

[143623]Luckow, B.; Bunz, F.; Stillman, B.; Lichter, P.; Schutz, G.:
Cloning,

[143624]expression, and chromosomal localization of the
140-kilodalton subunit

[143625]of replication factor C from mice and humans. Molec. Cell.
Biol. 14:

[143626]1626-1634, 1994.

[143627]25723.

[143628]Pennaneach, V.; Salles-Passador, I.; Munshi, A.; Brickner,
H.;

[143629]Regazzoni, K.; Dick, F.; Dyson, N.; Chen, T.-T.; Wang, J. Y.
J.; Fotedar,

[143630]R.; Fotedar, A.: The large subunit of replication factor C promotes

[143631]cell survival after DNA damage in an LxCxE motif- and Rb-dependent

[143632]manner. Molec. Cell 7: 715-727, 2001.

[143633]25724.

[143634]Uchiumi, F.; Ohta, T.; Tanuma, S.: Replication factor C recognizes

[143635]5-prime-phosphate ends of telomeres. Biochem. Biophys. Res. Commun. 229:

[143636]310-315, 1996.

[143637]25725.

[143638]Schmickel, R. D.: Contiguous gene syndromes: a component of recognizable

[143639]syndromes. J. Pediat. 109: 231-241, 1986.

[143640]25726.

[143641]Olson, E.; Srivastava, D.: Molecular pathways controlling heart

[143642]development. Science 272: 671-676, 1996.

[143643]25727.

[143644]Cunningham, J. M.; Vanin, E. F.; Tran, N.; Valentine, M.;
Jane,
[143645]S. M.: The human transcription factor CP2 (TFCP2), a com-
ponent of
[143646]the human gamma-globin stage selector protein, maps to
chromosome
[143647]region 12q13 and is within 250 kb of the NF-E2 gene. Ge-
nomics 30:
[143648]398-399, 1995.
[143649]25728.

[143650]Jane, S. M.; Nienhuis, A. W.; Cunningham, J. M.:
Hemoglobin switching
[143651]in man and chicken is mediated by a heteromeric complex
between the
[143652]ubiquitous transcription factor CP2 and a developmentally
specific
[143653]protein. EMBO J. 13: 197-105, 1995.
[143654]25729.

[143655]Lambert, J.-C.; Goumidi, L.; Wavrant-De Vrieze, F.;
Frigard, B.;
[143656]Harris, J. M.; Cummings, A.; Coates, J.; Pasquier, F.; Cotel,
D.;

- [143657] Gaillac, M.; St. Clair, D.; Mann, D. M. A.; Hardy, J.; London, C.
- [143658] L.; Amouyel, P.; Chartier-Harlin, M.-C.: The transcriptional factor
- [143659] LBP-1c/CP2/LSF gene on chromosome 12 is a genetic determinant of Alzheimer's
- [143660] disease. Hum. Molec. Genet. 9: 2275-2280, 2000.
- [143661] 25730.
- [143662] Swendeman, S. L.; Spielholz, C.; Jenkins, N. A.; Gilbert, D. J.;
- [143663] Copeland, N. G.; Sheffery, M.: Characterization of the genomic structure,
- [143664] chromosomal location, promoter, and developmental expression of the
- [143665] alpha-globin transcription factor CP2. J. Biol. Chem. 269: 11663-11671,
- [143666] 1994.
- [143667] 25731.
- [143668] Taylor, A. E.; Yip, A.; Brayne, C.; Easton, D.; Evans, J. G.; Xuereb,
- [143669] J.; Cairns, N.; Esiri, M. M.; Rubinsztein, D. C.: Genetic association

[143670] of an LBP-1c/CP2/LSF gene polymorphism with late onset
Alzheimer's

[143671] disease. J. Med. Genet. 38: 232-233, 2001.

[143672] 25732.

[143673] Bandara, L. R.; La Thangue, N. B.: Adenovirus E1a prevents
the

[143674] retinoblastoma gene product from complexing with a cel-
lular transcription

[143675] factor. Nature 351: 494-497, 1991.

[143676] 25733.

[143677] Chen, C.-R.; Kang, Y.; Siegel, P. M.; Massague, J.: E2F4/5
and

[143678] p107 as Smad cofactors linking the TGF-beta receptor to
c-myc repression. Cell 110:

[143679] 19-32, 2002.

[143680] 25734.

[143681] Girling, R.; Partridge, J. F.; Bandara, L. R.; Burden, N.;
Totty,

[143682] N. F.; Hsuan, J. J.; La Thangue, N. B.: A new component of
the transcription

[143683] factor DRTF1/E2F. Nature 362: 83-87, 1993. Note: Cor-

rection: Nature

[143684]365: 468 only, 1993.

[143685]25735.

[143686]Malas, S.; Duthie, S. M.; Mohri, F.; Lovell-Badge, R.;

Episkopou,

[143687]V.: Cloning and mapping of the human SOX1: a highly

conserved gene

[143688]expressed in the developing brain. Mammalian Genome 8:

866–868,

[143689]1997.

[143690]25736.

[143691]Zhang, Y.; Venkatraj, V. S.; Fischer, S. G.; Warburton, D.;

Chellappan,

[143692]S. P.: Genomic cloning and chromosomal assignment of

the E2F dimerization

[143693]partner TFDP gene family. Genomics 39: 95–98, 1997.

[143694]25737.

[143695] Moloney, D. J.; Panin, V. M.; Johnston, S. H.; Chen, J.; Shao,
[143696] L.; Wilson, R.; Wang, Y.; Stanley, P.; Irvine, K. D.; Halti-
wanger,
[143697] R. S.; Vogt, T. F.: Fringe is a glycosyltransferase that mod-
ifies
[143698] Notch. Nature 406: 369–375, 2000.
[143699] 25738.
[143700] Sestan, N.; Artavanis-Tsakonas, S.; Rakic, P.: Contact-
dependent
[143701] inhibition of cortical neurite growth mediated by Notch
signaling. Science 286:
[143702] 741–746, 1999.
[143703] 25739.
[143704] Tanigaki, K.; Nogaki, F.; Takahashi, J.; Tashiro, K.;
Kurooka,
[143705] H.; Honjo, T.: Notch1 and Notch3 instructively restrict
bFGF-responsive
[143706] multipotent neural progenitor cells to an astroglial fate.
Neuron 29:
[143707] 45–55, 2001.
[143708] 25740.

- [143709]Barton, D. E.; Foellmer, B. E.; Du, J.; Tamm, J.; Derynck, R.;
- [143710]Francke, U.: Chromosomal mapping of genes for trans-
forming growth
- [143711]factors beta-2 and beta-3 in man and mouse: dispersion
of TGF-beta
- [143712]gene family. *Oncogene Res.* 3: 323-331, 1988.
- [143713]25741.
- [143714]Dean, M.; Park, M.; Vande Woude, G. F.: Characterization
of the
- [143715]rearranged TPR-MET oncogene breakpoint. *Molec. Cell.*
Biol. 7: 921-924,
- [143716]1987.
- [143717]25742.
- [143718]Lemmink, H. H.; Mochizuki, T.; van den Heuvel, L. P. W. J.;
Schroder,
- [143719]C. H.; Barrientos, A.; Monnens, L. A. H.; van Oost, B. A.;
Brunner,
- [143720]H. G.; Reeders, S. T.; Smeets, H. J. M.: Mutations in the
type IV
- [143721]collagen alpha-3 (COL4A3) gene in autosomal recessive
Alport syndrome. *Hum.*

[143722]Molec. Genet. 3: 1269–1273, 1994.

[143723]25743.

[143724]Gibson, R. A.; Buchwald, M.; Roberts, R. G.; Mathew, C. G.:
Characterisation

[143725]of the exon structure of the Fanconi anaemia group C
gene by vectorette

[143726]PCR. Hum. Molec. Genet. 2: 35–38, 1993.

[143727]25744.

[143728]Gregersen, N.; Andresen, B. S.; Corydon, M. J.; Corydon, T.
J.;

[143729]Olsen, R. K. J.; Bolund, L.; Bross, P.: Mutation analysis in
mitochondrial

[143730]fatty acid oxidation defects: exemplified by acyl–CoA de-
hydrogenase

[143731]deficiencies, with special focus on genotype–phenotype
relationship. Hum.

[143732]Mutat. 18: 169–189, 2001.

[143733]25745.

[143734]Onkenhout, W.; Venizelos, V.; Scholte, H. R.; De Klerk, J. B.

[143735]C.; Poorthuis, B. J. H. M.: Intermediates of unsaturated
fatty acid

- [143736]oxidation are incorporated in triglycerides but not in phospholipids
- [143737]in tissues from patients with mitochondrial beta-oxidation defects. J.
- [143738]Inherit. Metab. Dis. 24: 337-344, 2001.
- [143739]25746.
- [143740]Lemmink, H. H.; Schroder, C. H.; Monnens, L. A. H.; Smeets, H.
- [143741]J. M.: The clinical spectrum of type IV collagen mutations. Hum.
- [143742]Mutat. 9: 477-499, 1997.
- [143743]25747.
- [143744]Goldmuntz, E.; Wang, Z.; Roe, B. A.; Budarf, M. L.: Cloning, genomic
- [143745]organization, and chromosomal localization of human citrate transport
- [143746]protein to the DiGeorge/velocardiofacial syndrome minimal critical
- [143747]region. Genomics 33: 271-276, 1996.
- [143748]25748.
- [143749]Heisterkamp, N.; Mulder, M. P.; Langeveld, A.; ten Hoeve,

J.; Wang,

[143750]Z.; Roe, B. A.; Groffen, J.: Localization of the human mitochondrial

[143751]citrate transporter protein gene to chromosome 22q11 in the DiGeorge

[143752]syndrome critical region. Genomics 29: 451–456, 1995.

[143753]25749.

[143754]Iacobazzi, V.; Lauria, G.; Palmieri, F.: Organization and sequence

[143755]of the human gene for the mitochondrial citrate transport protein. DNA

[143756]Seq. 7: 127–139, 1997.

[143757]25750.

[143758]Kaplan, R. S.; Mayor, J. A.; Wood, D. O.: The mitochondrial tricarboxylate

[143759]transport protein: cDNA cloning, primary structure, and comparison

[143760]with other mitochondrial transport proteins. J. Biol. Chem. 268:

[143761]13682–13690, 1993.

[143762]25751.

[143763]Stoffel, M.; Karayiorgou, M.; Espinosa, R., III; Le Beau, M.
M.

[143764]: The human mitochondrial citrate transporter gene
(SLC20A3) maps

[143765]to chromosome band 22q11 within a region implicated in
DiGeorge syndrome,

[143766]velo-cardio-facial syndrome and schizophrenia. Hum.
Genet. 98: 113-115,

[143767]1996.

[143768]25752.

[143769]Uberbacher, E. C.; Mural, R. J.: Locating protein-coding
regions

[143770]in human DNA sequences by a multiple sensor-neural
network approach. Proc.

[143771]Nat. Acad. Sci. 88: 11261-11265, 1991.

[143772]25753.

[143773]Gibson, R. A.; Hajianpour, A.; Buchwald, M.; Mathew, C.
G.: Screening

[143774]the FACC gene for mutations in patients with Fanconi
anaemia. (Abstract) Am.

[143775]J. Hum. Genet. 51 (suppl.): A216 only, 1992.

[143776]25754.

[143777]Vidal, R.; Frangione, B.; Rostagno, A.; Mead, S.; Revesz, T.;
Plant,

[143778]G.; Ghiso, J.: A stop-codon mutation in the BRI gene asso-
ciated with

[143779]familial British dementia. *Nature* 399: 776–781, 1999.

[143780]25755.

[143781]Joensuu, T.; Blanco, G.; Pakarinen, L.; Sistonen, P.; Kaari-
ainen,

[143782]H.; Brown, S.; de la Chapelle, A.; Sankila, E.-M.: Refined
mapping

[143783]of the Usher syndrome type III locus on chromosome 3,
exclusion of

[143784]candidate genes, and identification of the putative mouse
homologous

[143785]region. *Genomics* 38: 255–263, 1996.

[143786]25756.

[143787]Lamhonwah, A.-M.; Tein, I.: Carnitine uptake defect:
frameshift

[143788]mutations in the human plasmalemmal carnitine trans-
porter gene. *Biochem.*

[143789]Biophys. Res. Commun. 252: 396–401, 1998.

[143790]25757.

[143791]Matsuishi, T.; Hirata, K.; Terasawa, K.; Kato, H.; Yoshino, M.;

[143792]Ohtaki, E.; Hirose, F.; Nonaka, I.; Sugiyama, N.; Ohta, K.:
Successful

[143793]carnitine treatment in two siblings having lipid storage
myopathy

[143794]with hypertrophic cardiomyopathy. Neuropediatrics 16:
6–12, 1985.

[143795]25758.

[143796]Nezu, J.; Tamai, I.; Oku, A.; Ohashi, R.; Yabuuchi, H.;
Hashimoto,

[143797]N.; Nikaido, H.; Sai, Y.; Koizumi, A.; Shoji, Y.; Takada, G.;
Matsuishi,

[143798]T.; Yoshino, M.; Kato, H.; Ohura, T.; Tsujimoto, G.;
Hayakawa, J.;

[143799]Shimane, M.; Tsuji, A.: Primary systemic carnitine defi-
ciency is

[143800]caused by mutations in a gene encoding sodium ion-
dependent carnitine

[143801]transporter. Nature Genet. 21: 91–94, 1999.

[143802]25759.

[143803]Shoji, Y.; Koizumi, A.; Kayo, T.; Ohata, T.; Takahashi, T.;
Harada,

[143804]K.; Takada, G.: Evidence for linkage of human primary
systemic carnitine

[143805]deficiency with D5S436: a novel gene locus on chromo-
some 5q. Am.

[143806]J. Hum. Genet. 63: 101-108, 1998.

[143807]25760.

[143808]Tamai, I.; Ohashi, R.; Nezu, J.; Yabuuchi, H.; Oku, A.; Shi-
mane,

[143809]M.; Sai, Y.; Tsuji, A.: Molecular and functional identifica-
tion of

[143810]sodium ion-dependent, high affinity human carnitine
transporter OCTN2. J.

[143811]Biol. Chem. 273: 20378-20382, 1998.

[143812]25761.

[143813]Tein, I.; De Vivo, D. C.; Bierman, F.; Pulver, P.; De Meirleir,

[143814]L. J.; Cvitanovic-Sojat, L.; Pagon, R. A.; Bertini, E.; Dionisi-
Vici,

[143815]C.; Servidei, S.; Dimauro, S.: Impaired skin fibroblast car-

nitine

[143816]uptake in primary systemic carnitine deficiency manifested by childhood

[143817]carnitine-responsive cardiomyopathy. *Pediat. Res.* 28: 247–255, 1990.

[143818]25762.

[143819]Wang, Y.; Korman, S. H.; Ye, J.; Gargus, J. J.; Gutman, A.; Taroni,

[143820]F.; Garavaglia, B.; Longo, N.: Phenotype and genotype variation in

[143821]primary carnitine deficiency. *Genet. Med.* 3: 387–392, 2001.

[143822]25763.

[143823]Engelkamp, D.; Schafer, B. W.; Mattei, M. G.; Erne, P.; Heizmann,

[143824]C. W.: Six S100 genes are clustered on human chromosome 1q21: identification

[143825]of two genes coding for the two previously unreported calcium-binding

[143826]proteins S100D and S100E. *Proc. Nat. Acad. Sci.* 90: 6547–6551, 1993.

[143827]25764.

- [143828]Herrera, G. A.; Turbat-Herrera, E. A.; Lott, R. L.: S-100 protein
expression by primary and metastatic adenocarcinomas.
Am. J. Clin.
Path. 89: 168-176, 1988.
- [143831]25765.
- [143832]Isobe, T.; Tsugira, A.; Okuyama, T.: Amino acid sequence
of the
subunit structure of bovine brain S-100 protein (PAP 1-b).
J. Neurochem. 30:
921-923, 1978.
- [143835]25766.
- [143836]Moore, B. W.: A soluble protein characteristic of the ner-
vous
system. Biochem. Biophys. Res. Commun. 19: 739-744,
1965.
- [143838]25767.
- [143839]Morii, K.; Tanaka, R.; Takahashi, Y.; Minoshima, S.;
Fukuyama,
R.; Shimizu, N.; Kuwano, R.: Structure and chromosome
assignment

- [143841]of human S100 alpha and beta subunit genes. Biochem. Biophys. Res.
- [143842]Commun. 175: 185–191, 1991.
- [143843]25768.
- [143844]Most, P.; Bernotat, J.; Ehlermann, P.; Pleger, S. T.; Reppel, M.;
- [143845]Borries, M.; Niroomand, F.; Pieske, B.; Janssen, P. M. L.; Eschenhagen,
- [143846]T.; Karczewski, P.; Smith, G. L.; Koch, W. J.; Katus, H. A.; Remppis,
- [143847]A.: S100A1: a regulator of myocardial contractility. Proc. Nat.
- [143848]Acad. Sci. 98: 13889–13894, 2001.
- [143849]25769.
- [143850]Asada, Y.; Nadeau, J. H.: Fert is on mouse chromosome 11, not
- [143851]chromosome 17. Mammalian Genome 5: 830 only, 1994.
- [143852]25770.
- [143853]Hao, Q.–L.; Heisterkamp, N.; Groffen, J.: Isolation and sequence
- [143854]analysis of a novel human tyrosine kinase gene. Molec.

Cell. Biol. 9:

[143855]1587-1593, 1989.

[143856]25771.

[143857]Dryja, T. P.; Grondin, V. J.; Ringens, P.; Cotran, P.; Berson,

[143858]E. L.; Travis, G.: Isolation of human retinal cDNA fragments homologous

[143859]to the murine rds gene transcript. (Abstract) Invest. Ophthalmol. Vis.

[143860]Sci. 30 (suppl.): 43 only, 1989.

[143861]25772.

[143862]Farrar, G. J.; Jordan, S. A.; Kenna, P.; Humphries, M. M.; Kumar-Singh,

[143863]R.; McWilliam, P.; Allamand, V.; Sharp, E.; Humphries, P.: Autosomal

[143864]dominant retinitis pigmentosa: localization of a disease gene (RP6)

[143865]to the short arm of chromosome 6. Genomics 11: 870-874, 1991.

[143866]25773.

[143867]Farrar, G. J.; Kenna, P.; Jordan, S. A.; Kumar-Singh, R.; Humphries,

[143868]M. M.; Sharp, E. M.; Sheils, D. M.; Humphries, P.: A three-base-pair

[143869]deletion in the peripherin-RDS gene in one form of retinitis pigmentosa. Nature 354:

[143870]478-480, 1991.

[143871]25774.

[143872]Feist, R. M.; White, M. F., Jr.; Skalka, H.; Stone, E. M.: Choroidal

[143873]neovascularization in a patient with adult foveomacular dystrophy

[143874]and a mutation in the retinal degeneration slow gene (pro210-to-arg). Am.

[143875]J. Ophthalm. 118: 259-260, 1994.

[143876]25775.

[143877]Felbor, U.; Schilling, H.; Weber, B. H. F.: Adult vitelliform

[143878]macular dystrophy is frequently associated with mutations in the peripherin/RDS

[143879]gene. Hum. Mutat. 10: 301-309, 1997.

[143880]25776.

[143881]Gass, J. D. M.: A clinicopathologic study of a peculiar foveomacular

[143882]dystrophy. Trans. Am. Ophthal. Soc. 72: 139–155, 1974.

[143883]25777.

[143884]Jackson, K. E.; Mitchell, E. B.; Stone, E. M.; Ferrell, R. E.;

[143885]Gorin, M. B.: The identification of an exon–2 peripherin
mutation

[143886]in a family with heterogeneous manifestations of a butter-
fly pattern

[143887]macular dystrophy. (Abstract) Am. J. Hum. Genet. 53
(suppl.): 1177

[143888]only, 1993.

[143889]25778.

[143890]Jordan, S. A.; Farrar, G. J.; Kumar–Singh, R.; Kenna, P.;
Humphries,

[143891]M. M.; Allamand, V.; Sharp, E. M.; Humphries, P.: Autosomal dominant
mal dominant

[143892]retinitis pigmentosa (adRP; RP6): cosegregation of RP6
and the peripherin–RDS

[143893]locus in a late–onset family of Irish origin. Am. J. Hum.
Genet. 50:

[143894]634–639, 1992.

[143895]25779.

[143896]Billingsley, G. D.; Walter, M. A.; Hammond, G. L.; Cox, D.

W.:

[143897]Physical mapping of four serpin genes: alpha-

1-antitrypsin, alpha-1-antichymotrypsin,

[143898]corticosteroid-binding globulin, and protein C inhibitor,
within a

[143899]280-kb region on chromosome 14q32.1. Am. J. Hum.
Genet. 52: 343-353,

[143900]1993.

[143901]25780.

[143902]Clifton-Bligh, R. J.; Wentworth, J. M.; Heinz, P.; Crisp, M.
S.;

[143903]John, R.; Lazarus, J. H.; Ludgate, M.; Chatterjee, V. K.: Mu-
tation

[143904]of the gene encoding human TTF-2 associated with thy-
roid agenesis,

[143905]cleft palate and choanal atresia. Nature Genet. 19:
399-401, 1998.

[143906]25781.

[143907]Bird, T. D.; Koerker, R. M.; Leaird, B. J.; Vlcek, B. W.;
Thorning,

[143908]D. R.: Lipomembranous polycystic osteodysplasia (brain,

bone, and

[143909]fat disease): a genetic cause of presenile dementia. Neu-
rology 33:

[143910]81-86, 1983.

[143911]25782.

[143912]Paloneva, J.; Manninen, T.; Christman, G.; Hovanes, K.;
Mandelin,

[143913]J.; Adolfsson, R.; Bianchin, M.; Bird, T.; Miranda, R.;
Salmaggi,

[143914]A.; Tranebjaerg, L.; Konttinen, Y.; Peltonen, L.: Mutations
in two

[143915]genes encoding different subunits of a receptor signaling
complex

[143916]result in an identical disease phenotype. Am. J. Hum.
Genet. 71:

[143917]656-662, 2002.

[143918]25783.

[143919]Greco, A.; Ittmann, M.; Barletta, C.; Basilico, C.; Croce, C.
M.;

[143920]Cannizzaro, L. A.; Huebner, K.: Chromosomal localization
of human

[143921]genes required for G(1) progression in mammalian cells.

Genomics 4:

[143922]240–245, 1989.

[143923]25784.

[143924]Ittmann, M.; Greco, A.; Basilico, C.: Isolation of the human
gene

[143925]that complements a temperature-sensitive cell cycle mu-
tation in BHK

[143926]cells. Molec. Cell. Biol. 7: 3386–3393, 1987.

[143927]25785.

[143928]Bonyadi, M.; Cui, W.; Nagase, H.; Akhurst, R. J.: The TGF-
beta

[143929]type II receptor, Tgfbr2, maps to distal mouse chromo-
some 9. Genomics 33:

[143930]328–329, 1996.

[143931]25786.

[143932]Chen, R.-H.; Ebner, R.; Derynck, R.: Inactivation of the
type

[143933]II receptor reveals two receptor pathways for the diverse
TGF-beta

[143934]activities. Science 260: 1335–1338, 1993.

[143935]25787.

- [143936]Eshleman, J. R.; Markowitz, S. D.: Microsatellite instability
[143937]in inherited and sporadic neoplasms. *Curr. Opin. Oncol.* 7:
83–89,
[143938]1995.
[143939]25788.
- [143940]Hahm, K.-B.; Cho, K.; Lee, C.; Im, Y.-H.; Chang, J.; Choi,
S.-G.;
- [143941]Sorensen, P. H. B.; Thiele, C. J.; Kim, S.-J.: Repression of
the
[143942]gene encoding the TGF- β type II receptor is a major
target of the
[143943]EWS-FLI1 oncoprotein. *Nature Genet.* 23: 222–227, 1999.
[143944]25789.
- [143945]Lin, H. Y.; Wang, X.-F.; Ng-Eaton, E.; Weinberg, R. A.;
- Lodish,
[143946]H. F.: Expression cloning of the TGF- β type II receptor,
a functional
[143947]transmembrane serine/threonine kinase. *Cell* 68:
775–785, 1992.
[143948]25790.
- [143949]Lu, S.-L.; Kawabata, M.; Imamura, T.; Akiyama, Y.; No-

mizu, T.;

[143950]Miyazono, K.; Yuasa, Y.: HNPCC associated with germline mutation

[143951]in the TGF-beta type II receptor gene. (Letter) Nature Genet. 19:

[143952]17-18, 1998.

[143953]25791.

[143954]Markowitz, S.; Wang, J.; Myeroff, L.; Parsons, R.; Sun, L.; Lutterbaugh,

[143955]J.; Fan, R. S.; Zborowska, E.; Kinzler, K. W.; Vogelstein, B.; Brattain,

[143956]M.; Willson, J. K. V.: Inactivation of the type II TGF-beta receptor

[143957]in colon cancer cells with microsatellite instability. Science 268:

[143958]1336-1338, 1995.

[143959]25792.

[143960]Mathew, S.; Murty, V. V. V. S.; Cheifetz, S.; George, D.; Massague,

[143961]J.; Chaganti, R. S. K.: Transforming growth factor receptor gene

[143962]TGFBR2 maps to human chromosome band 3p22. Ge-

nomics 20: 114–115,

[143963]1994.

[143964]25793.

[143965]Myeroff, L. L.; Parsons, R.; Kim, S.-J.; Hedrick, L.; Cho, K.
R.;

[143966]Orth, K.; Mathis, M.; Kinzler, K. W.; Lutterbaugh, J.; Park,
K.; Bang,

[143967]Y.-J.; Lee, H. Y.; Park, J.-G.; Lynch, H. T.; Roberts, A. B.;
Vogelstein,

[143968]B.; Markowitz, S. D.: A transforming growth factor beta
receptor

[143969]type II gene mutation common in colon and gastric but
rare in endometrial

[143970]cancers with microsatellite instability. Cancer Res. 55:
5545–5547,

[143971]1995.

[143972]25794.

[143973]Parsons, R.; Myeroff, L. L.; Liu, B.; Willson, J. K. V.;
Markowitz,

[143974]S. D.; Kinzler, K. W.; Vogelstein, B.: Microsatellite instabil-
ity

[143975]and mutations of the transforming growth factor beta type

II receptor

[143976] gene in colorectal cancer. Cancer Res. 55: 5548–5550, 1995.

[143977] 25795.

[143978] Tanaka, S.; Mori, M.; Mafune, K.; Ohno, S.; Sugimachi, K.:

A

[143979] dominant negative mutation of transforming growth factor-beta receptor

[143980] type II gene in microsatellite stable oesophageal carcinoma. Brit.

[143981] J. Cancer 82: 1557–1560, 2000.

[143982] 25796.

[143983] Tannergard, P.; Liu, T.; Weger, A.; Nordenskjold, M.; Lindbloom,

[143984] A.: Tumorigenesis in colorectal tumors from patients with hereditary

[143985] non-polypoid colorectal cancer. Hum. Genet. 101: 51–55, 1997.

[143986] 25797.

[143987] Watanabe, T.; Wu, T.-T.; Catalano, P. J.; Ueki, T.; Satriano,

[143988] R.; Haller, D. G.; Benson, A. B., III; Hamilton, S. R.: Molecu-

lar

[143989]predictors of survival after adjuvant chemotherapy for
colon cancer. New

[143990]Eng. J. Med. 344: 1196–1206, 2001.

[143991]25798.

[143992]Bonnefont, J.-P.; Chretien, D.; Rustin, P.; Robinson, B.;
Vassault,

[143993]A.; Aupetit, J.; Charpentier, C.; Rabier, D.; Saudubray,
J.-M.; Munnich,

[143994]A.: Alpha-ketoglutarate dehydrogenase deficiency pre-
sented as congenital

[143995]lactic acidosis. J. Pediatr. 121: 255–258, 1992.

[143996]25799.

[143997]Guffon, N.; Lopez-Mediavilla, C.; Dumoulin, R.; Mousson,
B.; Godinot,

[143998]C.; Carrier, H.; Collombet, J. M.; Divry, P.; Mathieu, M.;
Guibaud,

[143999]P.: 2-Ketoglutarate dehydrogenase deficiency, a rare
cause of primary

[144000]hyperlactataemia: report of a new case. J. Inherit. Metab.
Dis. 16:

[144001]821–830, 1993.

[144002]25800.

[144003]Kohlschutter, A.; Behbehani, A.; Langenbeck, U.; Albani, M.; Heidemann,

[144004]P.; Hoffmann, G.; Kleineke, J.; Lehnert, W.; Wendel, U.: A familial

[144005]progressive neurodegenerative disease with 2-oxoglutaric aciduria. Europ.

[144006]J. Pediat. 138: 32-37, 1982.

[144007]25801.

[144008]Koike, K.: The gene encoding human 2-oxoglutarate dehydrogenase:

[144009]structural organization and mapping to chromosome 7p13-p14. Gene 159:

[144010]261-266, 1995.

[144011]25802.

[144012]Koike, K.; Urata, Y.; Goto, S.: Cloning and nucleotide sequence

[144013]of the cDNA encoding human 2-oxoglutarate dehydrogenase (lipoamide). Proc.

[144014]Nat. Acad. Sci. 89: 1963-1967, 1992.

[144015]25803.

[144016]Szabo, P.; Cai, X.; Ali, G.; Blass, J. P.: Localization of the
 [144017]gene (OGDH) coding for the E1 α component of the alpha-
 ketoglutarate
 [144018]dehydrogenase complex to chromosome 7p13-p11.2.
 Genomics 20: 324-326,
 [144019]1994.
 [144020]25804.

[144021]Kury, S.; Dreno, B.; Bezieau, S.; Giraudet, S.; Kharfi, M.;
 Kamoun,
 [144022]R.; Moisan, J.-P.: Identification of SLC39A4, a gene in-
 volved in
 [144023]acrodermatitis enteropathica. Nature Genet. 31: 239-240,
 2002.
 [144024]25805.

[144025]Rogers, E. E.; Eide, D. J.; Guerinot, M. L.: Altered selectivity
 [144026]in an Arabidopsis metal transporter. Proc. Nat. Acad. Sci.
 97: 12356-12360,
 [144027]2000.
 [144028]25806.

[144029]Wang, K.; Pugh, E. W.; Griffen, S.; Doheny, K. F.; Mostafa,
 W.

- [144030]Z.; al-Aboosi, M. M.; el-Shanti, H.; Gitschier, J.: Homozygosity
- [144031]mapping places the acrodermatitis enteropathica gene on chromosomal
- [144032]region 8q24.3. Am. J. Hum. Genet. 68: 1055–1060, 2001.
- [144033]25807.
- [144034]Haberle, J.; Pauli, S.; Linnebank, M.; Kleijer, W. J.; Bakker,
- [144035]H. D.; Wanders, R. J. A.; Harms, E.; Koch, H. G.: Structure of the
- [144036]human argininosuccinate synthetase gene and an improved system for
- [144037]molecular diagnostics in patients with classical and mild citrullinemia. Hum.
- [144038]Genet. 110: 327–333, 2002.
- [144039]25808.
- [144040]Weve, H.: Ablatio falciformis congenita (retinal fold). Brit.
- [144041]J. Ophthal. 22: 456–470, 1938.
- [144042]25809.
- [144043]Barbosa, J.; Rich, S.; Dunsworth, T.; Swanson, J.: Linkage disequilibrium

[144044]between insulin-dependent diabetes and the Kidd blood group Jk(b)

[144045]allele. J. Clin. Endocr. Metab. 55: 193-195, 1982.

[144046]25810.

[144047]Harper, P. A.; Healy, P. J.; Dennis, J. A.; O'Brien, J. J.; Rayward,

[144048]D. H.: Citrullinaemia as a cause of neurological disease in neonatal

[144049]Friesian calves. Aust. Vet. J. 63: 378-379, 1986.

[144050]25811.

[144051]Harper, P. A. W.; Healy, P. J.; Dennis, J. A.: Animal model of

[144052]human disease: citrullinemia (argininosuccinate synthetase deficiency). Am.

[144053]J. Path. 135: 1213-1215, 1989.

[144054]25812.

[144055]Jackson, M. J.; Surh, L. C.; O'Brien, W. E.; Beaudet, A. L.:

[144056]Assignment of the structural gene for argininosuccinate synthetase

[144057]to proximal mouse chromosome 2. Genomics 6: 545-547, 1990.

[144058]25813.

[144059]Kobayashi, K.; Ichiki, H.; Saheki, T.; Tatsuno, M.;
Uchiyama,

[144060]C.; Nukada, O.; Yoda, T.: Structure of an abnormal messenger RNA

[144061]for argininosuccinate synthetase in citrullinemia. Hum.
Genet. 76:

[144062]27-32, 1987.

[144063]25814.

[144064]Kobayashi, K.; Jackson, M. J.; Tick, D. B.; O'Brien, W. E.;
Beaudet,

[144065]A. L.: Characterization of nine mutant alleles causing citrullinemia.

[144066](Abstract) Am. J. Hum. Genet. 45 (suppl.): A201 only,
1989.

[144067]25815.

[144068]Kobayashi, K.; Jackson, M. J.; Tick, D. B.; O'Brien, W. E.;
Beaudet,

[144069]A. L.: Heterogeneity of mutations in argininosuccinate
synthetase

[144070]causing human citrullinemia. J. Biol. Chem. 265:
11361-11367, 1990.

[144071]25816.

[144072]Kobayashi, K.; Kakinoki, H.; Fukushige, T.; Shaheen, N.;
Terazono,

[144073]H.; Saheki, T.: Nature and frequency of mutations in the
argininosuccinate

[144074]synthetase gene that cause classical citrullinemia. Hum.
Genet. 96:

[144075]454–463, 1995.

[144076]25817.

[144077]Kobayashi, K.; Rosenbloom, C.; Beaudet, A. L.; O'Brien, W.
E.

[144078]: Additional mutations in argininosuccinate synthetase
causing citrullinemia. Molec.

[144079]Biol. Med. 8: 95–100, 1991.

[144080]25818.

[144081]Kobayashi, K.; Saheki, T.; Imamura, Y.; Noda, T.; Inoue, I.;
Matuo,

[144082]S.; Hagihara, S.; Nomiya, H.; Jinno, Y.; Shimada, K.:
Messenger

[144083]RNA coding for argininosuccinate synthetase in citrulline-
mia. Am.

- [144084]J. Hum. Genet. 38: 667–680, 1986.
- [144085]25819.
- [144086]Kobayashi, K.; Shaheen, N.; Terazono, H.; Saheki, T.: Mutations
in argininosuccinate synthetase mRNA of Japanese patients, causing
classical citrullinemia. Am. J. Hum. Genet. 55: 1103–1112, 1994.
- [144089]25820.
- [144090]Li, C.-M.; Chao, H.-K.; Liu, Y.-F.; Su, T.-S.: A nonsense mutation
is responsible for the RNA-negative phenotype in human citrullinaemia. Europ.
- [144092]J. Hum. Genet. 9: 685–689, 2001.
- [144093]25821.
- [144094]Northrup, H.; Lathrop, M.; Lu, S.-Y.; Daiger, S. P.; Beaudet, A. L.; O'Brien, W. E.: Multilocus linkage analysis with the
argininosuccinate synthetase gene. Genomics 5: 442–444, 1989.
- [144097]25822.

[144098] Sase, M.; Kobayashi, K.; Imamura, Y.; Saheki, T.; Nakano, K.;

[144099] Miura, S.; Mori, M.: Level of translatable messenger RNA coding for

[144100] argininosuccinate synthetase in the liver of the patients with quantitative-type

[144101] citrullinemia. Hum. Genet. 69: 130–134, 1985.

[144102] 25823.

[144103] Todd, S.; Naylor, S. L.: New chromosomal mapping assignments

[144104] for argininosuccinate synthetase pseudogene 1, interferon-beta-3 gene,

[144105] and the diazepam binding inhibitor gene. Somat. Cell Molec. Genet. 18:

[144106] 381–385, 1992.

[144107] 25824.

[144108] Hsieh, C.-L.; Swaroop, A.; Francke, U.: Chromosomal localization

[144109] and cDNA sequence of human RALB, a GTP binding protein. Somat. Cell

[144110] Molec. Genet. 16: 407–410, 1990.

[144111]25825.

[144112]Ahmad, F.; Goldstein, B. J.: Functional association between
the

[144113]insulin receptor and the transmembrane protein-tyrosine
phosphatase

[144114]LAR in intact cells. J. Biol. Chem. 272: 448-457, 1997.

[144115]25826.

[144116]Disteche, C. M.; Adler, D. A.; Tedder, T. F.; Saito, H.: Map-
ping

[144117]of the genes for LYAM1, a new lymphocyte adhesion
molecule, and for

[144118]LAR, a new receptor-linked protein tyrosine phosphatase,
to human

[144119]chromosome 1 (Abstract) Cytogenet. Cell Genet. 51: 990
only, 1989.

[144120]25827.

[144121]Harder, K. W.; Saw, J.; Miki, N.; Jirik, F.: Coexisting amplifi-
cations

[144122]of the chromosome 1p32 genes (PTPRF and MYCL1) en-
coding protein tyrosine

[144123]phosphatase LAR and L-myc in a small cell lung cancer
line. Genomics 27:

[144124]552-553, 1995.

[144125]25828.

[144126]Jirik, F. R.; Harder, K. W.; Melhado, I. G.; Anderson, L. L.;
Duncan,

[144127]A. M. V.: The gene for leukocyte antigen-related tyrosine
phosphatase

[144128](LAR) is localized to human chromosome 1p32, a region
frequently deleted

[144129]in tumors of neuroectodermal origin. Cytogenet. Cell
Genet. 61:

[144130]266-268, 1992.

[144131]25829.

[144132]Nam, H.-J.; Poy, F.; Krueger, N. X.; Saito, H.; Frederick, C.
A.

[144133]: Crystal structure of the tandem phosphatase domains of
RPTP LAR. Cell 97:

[144134]449-457, 1999.

[144135]25830.

[144136]O'Grady, P.; Krueger, N. X.; Streuli, M.; Saito, H.: Genomic
organization

[144137]of the human LAR protein tyrosine phosphatase gene and

alternative

[144138]splicing in the extracellular fibronectin type-III domains. J. Biol.

[144139]Chem. 269: 25193-25199, 1994.

[144140]25831.

[144141]Schaapveld, R. Q.; Schepens, J. T.; Robinson, G. W.; Attema, J.;

[144142]Oerlemans, F. T.; Fransen, J. A.; Streuli, M.; Wieringa, B.; Hennighausen,

[144143]L.; Hendriks, W. J.: Impaired mammary gland development and function

[144144]in mice lacking LAR receptor-like tyrosine phosphatase activity. Dev.

[144145]Biol. 188: 134-146, 1997.

[144146]25832.

[144147]Schaapveld, R. Q. J.; van den Maagdenberg, A. M. J. M.; Schepens,

[144148]J. T. G.; Olde Weghuis, D.; Geurts van Kessel, A.; Wieringa, B.; Hendriks,

[144149]W. J. A. J.: The mouse gene Ptp^{prf} encoding the leukocyte common antigen-related

[144150]molecule LAR: cloning, characterization, and chromoso-

mal localization. Genomics 27:

[144151]124–130, 1995.

[144152]25833.

[144153]Tsujioka, K.; Kawakami, N.; Uchino, Y.; Ichijo, T.; Furukawa,

[144154]T.; Saito, H.; Yamamoto, H.: Distinct functions of the two protein

[144155]tyrosine phosphatase domains of LAR (leukocyte common antigen-related)

[144156]on tyrosine dephosphorylation of insulin receptor. Molec. Endocr. 15:

[144157]271–280, 2001.

[144158]25834.

[144159]Ali, R. R.; Sarra, G.-M.; Stephens, C.; de Alwis, M.; Bainbridge,

[144160]J. W. B.; Munro, P. M.; Fauser, S.; Reichell, M. B.; Kinnon, C.; Hunt,

[144161]D. M.; Bhattacharya, S. S.; Thrasher, A. J.: Restoration of photoreceptor

[144162]ultrastructure and function in retinal degeneration slow mice by gene

[144163]therapy. Nature Genet. 25: 306–310, 2000.

[144164]25835.

[144165]Bascom, R. A.; Connell, G.; Garcia-Heras, J.; Collins, L.;
Ledbetter,

[144166]D.; Molday, R. S.; Kalnins, V.; McInnes, R. R.: Molecular
and ultrastructural

[144167]characterization of the products of the human retinopathy
candidate

[144168]genes ROM1 and RDS. (Abstract) Am. J. Hum. Genet. 47
(suppl.): A101

[144169]only, 1990.

[144170]25836.

[144171]Connell, G.; Bascom, R.; Molday, L.; Reid, D.; McInnes, R.
R.;

[144172]Molday, R. S.: Photoreceptor peripherin is the normal
product of

[144173]the gene responsible for retinal degeneration in the rds
mouse. Proc.

[144174]Nat. Acad. Sci. 88: 723-726, 1991.

[144175]25837.

[144176]Demant, P.; Ivanyi, D.; van Nie, R.: The map position of the
rds

[144177]gene on the 17th chromosome of the mouse. Tissue Anti-
gens 13: 53-55,

[144178]1979.

[144179]25838.

[144180]Inoue, C.; Shiga, K.; Takasawa, S.; Kitagawa, M.; Ya-
mamoto, H.;

[144181]Okamoto, H.: Evolutionary conservation of the insulinoma
gene rig

[144182]and its possible function. Proc. Nat. Acad. Sci. 84:
6659-6662,

[144183]1987.

[144184]25839.

[144185]Kitagawa, M.; Takasawa, S.; Kikuchi, N.; Itoh, T.; Teraoka,
H.;

[144186]Yamamoto, H.; Okamoto, H.: Rig encodes ribosomal pro-
tein S15: the

[144187]primary structure of mammalian ribosomal protein S15.
FEBS Lett. 283:

[144188]210-214, 1991.

[144189]25840.

[144190]Shiga, K.; Yamamoto, H.; Okamoto, H.: Isolation and char-

acterization

[144191] of the human homologue of rig and its pseudogenes: the functional

[144192] gene has features characteristic of housekeeping genes.

Proc. Nat.

[144193] Acad. Sci. 87: 3594–3598, 1990.

[144194] 25841.

[144195] Brody, L. C.; Abel, K. J.; Castilla, L. H.; Couch, F. J.; McKinley,

[144196] D. R.; Yin, G.-Y.; Ho, P. P.; Merajver, S.; Chandrasekharappa, S.

[144197] C.; Xu, J.; Cole, J. L.; Struwing, J. P.; Valdes, J. M.; Collins,

[144198] F. S.; Weber, B. L.: Construction of a transcription map surrounding

[144199] the BRCA1 locus of human chromosome 17. Genomics 25: 238–247, 1995.

[144200] 25842.

[144201] Hloch, P.; Schiedner, G.; Stahl, H.: Complete cDNA sequence of

[144202] the human p68 protein. Nucleic Acids Res. 18: 3045, 1990.

[144203] 25843.

[144204] Iggo, R.; Gough, A.; Xu, W.; Lane, D. P.; Spurr, N. K.: Chromosome

[144205] mapping of the human gene encoding the 68-kDa nuclear antigen (p68)

[144206] by using the polymerase chain reaction. Proc. Nat. Acad. Sci. 86:

[144207] 6211–6214, 1989.

[144208] 25844.

[144209] Iggo, R. D.; Xu, W.; Lane, D. P.; Gough, A.; Spurr, N. K.: Assignment

[144210] of P68 to chromosome 17 using PCR. (Abstract) Cytogenet. Cell Genet. 51:

[144211] 1017, 1989.

[144212] 25845.

[144213] Acker, J.; Mattei, M.-G.; Wintzerith, M.; Roeckel, N.; Depetris,

[144214] D.; Vigneron, M.; Keding, C.: Chromosomal localization of human

[144215] RNA polymerase II subunit genes. Genomics 20: 496–499, 1994.

[144216] 25846.

- [144217]Schulze, A.; Hansen, C.; Skakkebaek, N. E.; Brondum-Nielsen, K.;
- [144218]Ledbetter, D. H.; Tommerup, N.: Exclusion of SNRPN as a major determinant
- [144219]of Prader-Willi syndrome by a translocation breakpoint. Nature Genet. 12:
- [144220]452-454, 1996.
- [144221]25847.
- [144222]Schweizer, J.; Zynger, D.; Francke, U.: In vivo nuclease hypersensitivity
- [144223]studies reveal multiple sites of parental origin-dependent differential
- [144224]chromatin conformation in the 150 kb SNRPN transcription unit. Hum.
- [144225]Molec. Genet. 8: 555-566, 1999.
- [144226]25848.
- [144227]Shemer, R.; Hershko, A. Y.; Perk, J.; Mostoslavsky, R.; Tsuberi,
- [144228]B.; Cedar, H.; Buiting, K.; Razin, A.: The imprinting box of the
- [144229]Prader-Willi/Angelman syndrome domain. Nature Genet. 26: 440-443,

[144230]2000.

[144231]25849.

[144232]Sun, Y.; Nicholls, R. D.; Butler, M. G.; Saitoh, S.; Hainline,

[144233]B. E.; Palmer, C. G.: Breakage in the SNRPN locus in a balanced 46,XY,t(15;19)

[144234]Prader-Willi syndrome patient. Hum. Molec. Genet. 5: 517-524, 1996.

[144235]25850.

[144236]Sutcliffe, J. S.; Nakao, M.; Christian, S.; Orstavik, K. H.; Tommerup,

[144237]N.; Ledbetter, D. H.; Beaudet, A. L.: Deletions of a differentially

[144238]methylated CpG island at the SNRPN gene define a putative imprinting

[144239]control region. Nature Genet. 8: 52-58, 1994.

[144240]25851.

[144241]Wirth, J.; Back, E.; Huttenhofer, A.; Nothwang, H.-G.; Lich, C.;

[144242]Gross, S.; Menzel, C.; Schinzel, A.; Kioschis, P.; Tommerup, N.; Ropers,

[144243]H.-H.; Horsthemke, B.; Buiting, K.: A translocation break-

point cluster

[144244]disrupts the newly defined 3-prime end of the SNURF-SNRPN transcription

[144245]unit on chromosome 15. Hum. Molec. Genet. 10: 201-210, 2001.

[144246]25852.

[144247]Critcher, R.; Stitson, R. N. M.; Wade-Martins, R.; Easty, D. J.;

[144248]Farr, C. J.: Assignment of Sox4 to mouse chromosome 13 bands A3-A5

[144249]by fluorescence in situ hybridization; refinement of the human SOX4

[144250]location to 6p22.3 and of SOX20 to chromosome 17p12.3. Cytogenet.

[144251]Cell Genet. 81: 294-295, 1998.

[144252]25853.

[144253]Denny, P.; Swift, S.; Connor, F.; Ashworth, A.: An SRY-related

[144254]gene expressed during spermatogenesis in the mouse encodes a sequence-specific

[144255]DNA-binding protein. EMBO J. 11: 3705-3712, 1992.

[144256]25854.

- [144257]Farr, C. J.; Easty, D. J.; Ragoussis, J.; Collignon, J.; Lovell-Badge,
[144258]R.; Goodfellow, P. N.: Characterization and mapping of the
human
[144259]SOX4 gene. Mammalian Genome 4: 577–584, 1993.
[144260]25855.
- [144261]Hansen, G. M.; Skapura, D.; Justice, M. J.: Genetic profile
of
[144262]insertion mutations in mouse leukemias and lymphomas.
Genome Res. 10:
[144263]237–243, 2000.
[144264]25856.
- [144265]Li, J.; et al; et al: Leukaemia disease genes: large-scale
cloning
[144266]and pathway predictions. Nature Genet. 23: 348–353,
1999.
[144267]25857.
- [144268]Lund, A. H.; Turner, G.; Trubetskoy, A.; Verhoeven, E.;
Wientjens,
[144269]E.; Hulsman, D.; Russell, R.; DePinho, R. A.; Lenz, J.; van
Lohuizen,

- [144270]M.: Genome-wide retroviral insertional tagging of genes involved
- [144271]in cancer in Cdkn2a-deficient mice. Nature Genet. 32: 160–165, 2002.
- [144272]25858.
- [144273]Suzuki, T.; Shen, H.; Akagi, K.; Morse, H. C., III; Malley, J.
- [144274]D.; Naiman, D. Q.; Jenkins, N. A.; Copeland, N. G.: New genes involved
- [144275]in cancer identified by retroviral tagging. Nature Genet. 32: 166–174,
- [144276]2002. Note: Erratum: Nature Genet. 32: 331 only, 2002.
- [144277]25859.
- [144278]van de Wetering, M.; Oosterwegel, M.; van Norren, K.; Clevers,
- [144279]H.: Sox-4, an Sry-like HMG box protein, is a transcriptional activator
- [144280]in lymphocytes. EMBO J. 12: 3847–3854, 1993.
- [144281]25860.
- [144282]Shoyab, M.; McDonald, V. L.; Bradley, J. G.; Todaro, G. J.: Amphiregulin:
- [144283]a bifunctional growth-modulating glycoprotein produced

by the phorbol

[144284]12-myristate 13-acetate-treated human breast adenocarcinoma cell line

[144285]MCF-7. Proc. Nat. Acad. Sci. 85: 6528-6532, 1988.

[144286]25861.

[144287]Shoyab, M.; Plowman, G. D.; McDonald, V. L.; Bradley, J. G.; Todaro,

[144288]G. J.: Structure and function of human amphiregulin: a member of

[144289]the epidermal growth factor family. Science 243: 1074-1076, 1989.

[144290]25862.

[144291]Donahue, R. P.; Bias, W. B.; Renwick, J. H.; McKusick, V. A.:

[144292]Probable assignment of the Duffy blood group locus to chromosome 1

[144293]in man. Proc. Nat. Acad. Sci. 61: 949-955, 1968.

[144294]25863.

[144295]Jiang, Y.; Armstrong, D.; Albrecht, U.; Atkins, C. M.; Noebels,

[144296]J. L.; Eichele, G.; Sweatt, J. D.; Beaudet, A. L.: Mutation of the

[144297] Angelman ubiquitin ligase in mice causes increased cytoplasmic p53

[144298] and deficits of contextual learning and long-term potentiation. *Neuron* 21: 799–811, 1998.

[144300] 25864.

[144301] Kishino, T.; Lalande, M.; Wagstaff, J.: UBE3A/E6-AP mutations

[144302] cause Angelman syndrome. *Nature Genet.* 15: 70–73, 1997.

[144303] 25865.

[144304] Matsuura, T.; Sutcliffe, J. S.; Fang, P.; Galjaard, R.-J.; Jiang, Y.; Benton, C. S.; Rommens, J. M.; Beaudet, A. L.: De novo truncating

[144306] mutations in E6-AP ubiquitin-protein ligase gene (UBE3A) in Angelman

[144307] syndrome. *Nature Genet.* 15: 74–77, 1997.

[144308] 25866.

[144309] Ono, H.; Kuno, Y.; Tanaka, H.; Yamashina, M.; Tsuyoshi, T.; Kondo, N.; Orii, T.: A case of paroxysmal nocturnal hemoglobin-

uria without

[144311]deficiency of decay-accelerating factor on erythrocytes.

Blood 75:

[144312]1746–1747, 1990.

[144313]25867.

[144314]Petranka, J. G.; Fleenor, D. E.; Sykes, K.; Kaufman, R. E.;

Rosse,

[144315]W. F.: Structure of the CD59–encoding gene: further evidence of a

[144316]relationship to murine lymphocyte antigen Ly–6 protein.

Proc. Nat.

[144317]Acad. Sci. 89: 7876–7879, 1992.

[144318]25868.

[144319]Rosse, W. F.: Personal Communication. Durham, N. C.

6/3/1993.

[144320]25869.

[144321]Rosse, W. F.; Parker, C. J.: Paroxysmal nocturnal hemoglobinuria. Clin.

[144322]Haemat. 14: 105–125, 1985.

[144323]25870.

[144324]Rother, R. P.; Rollins, S. A.; Mennone, J.; Chodera, A.; Fi–

del,

[144325]S. A.; Bessler, M.; Hillmen, P.; Squinto, S. P.: Expression of recombinant

[144326]transmembrane CD59 in paroxysmal nocturnal hemoglobinuria B cells

[144327]confers resistance to human complement. Blood 84: 2604–2611, 1994.

[144328]25871.

[144329]Tone, M.; Walsh, L. A.; Waldmann, H.: Gene structure of human

[144330]CD59 and demonstration that discrete mRNAs are generated by alternative

[144331]polyadenylation. J. Molec. Biol. 227: 971–976, 1992.

[144332]25872.

[144333]Walsh, L. A.; Tone, M.; Thiru, S.; Waldmann, H.: The CD59 antigen--a

[144334]multifunctional molecule. Tissue Antigens 40: 213–220, 1992.

[144335]25873.

[144336]Yamashina, M.; Ueda, E.; Kinoshita, T.; Takami, T.; Ojima, A.;

[144337]Ono, H.; Tanaka, H.; Kondo, N.; Orii, T.; Okada, N.; Okada,

- H.; Inoue,
[144338]K.; Kitani, T.: Inherited complete deficiency of
20-kilodalton homologous
[144339]restriction factor (CD59) as a cause of paroxysmal noctur-
nal hemoglobinuria. New
[144340]Eng. J. Med. 323: 1184-1189, 1990.
[144341]25874.
- [144342]Milatovich, A.; Bolger, G.; Michaeli, T.; Francke, U.: Chro-
mosome
[144343]localizations of genes for five cAMP-specific phosphodi-
esterases in
[144344]man and mouse. Somat. Cell Molec. Genet. 20: 75-86,
1994.
[144345]25875.
- [144346]Loughney, K.; Martins, T. J.; Harris, E. A. S.; Sadhu, K.;
Hicks,
[144347]J. B.; Sonnenburg, W. K.; Beavo, J. A.; Ferguson, K.: Isola-
tion and
[144348]characterization of cDNAs corresponding to two human
calcium, calmodulin-regulated,
[144349]3-prime,5-prime-cyclic nucleotide phosphodiesterases. J.
Biol. Chem. 271:

[144350]796–806, 1996.

[144351]25876.

[144352]Wilson, D. E.; McKenna, L.: Assignment of the human gene
for phosphodiesterase

[144353]1A to chromosome 4. (Abstract) Am. J. Hum. Genet. 43:
A162 only,

[144354]1988.

[144355]25877.

[144356]Maruo, Y.; Nishizawa, K.; Sato, H.; Sawa, H.; Shimada, M.:
Prolonged

[144357]unconjugated hyperbilirubinemia associated with breast
milk and mutations

[144358]of the bilirubin uridine diphosphate–glu-
curonosyltransferase gene. Pediatrics 106:

[144359]e59, 2000. Note: Electronic Article.

[144360]25878.

[144361]Miles, J. S.; Moss, J. E.; Taylor, B. A.; Burchell, B.; Wolf,

[144362]C. R.: Mapping genes encoding drug–metabolizing en-
zymes in recombinant

[144363]inbred mice. Genomics 11: 309–316, 1991.

[144364]25879.

[144365]Moghrabi, N.; Clarke, D. J.; Boxer, M.; Burchell, B.: Identification

[144366]of an A-to-G missense mutation in exon 2 of the UGT1 gene complex

[144367]that causes Crigler-Najjar syndrome type 2. Genomics 18: 171-173,

[144368]1993.

[144369]25880.

[144370]Moghrabi, N.; Clarke, D. J.; Burchell, B.; Boxer, M.: Cosegregation

[144371]of intragenic markers with a novel mutation that causes Crigler-Najjar

[144372]syndrome type I: implication in carrier detection and prenatal diagnosis. Am.

[144373]J. Hum. Genet. 53: 722-729, 1993.

[144374]25881.

[144375]Moghrabi, N.; Sutherland, L.; Wooster, R.; Povey, S.; Boxer, M.;

[144376]Burchell, B.: Chromosomal assignment of human phenol and bilirubin

[144377]UDP-glucuronosyltransferase genes (UGT1A-subfamily).

Ann. Hum. Genet. 56:

[144378]81-91, 1992.

[144379]25882.

[144380]Mojarrabi, B.; Butler, R.; Mackenzie, P. I.: cDNA cloning
and

[144381]characterization of the human UDP glucuronosyltrans-
ferase, UGT1A3. Biochem.

[144382]Biophys. Res. Commun. 225: 785-790, 1996.

[144383]25883.

[144384]Monaghan, G.; McLellan, A.; McGeehan, A.; Li Volti, S.;
Mollica,

[144385]F.; Salemi, I.; Din, Z.; Cassidy, A.; Hume, R.; Burchell, B.:
Gilbert's

[144386]syndrome is a contributory factor in prolonged unconju-
gated hyperbilirubinemia

[144387]of the newborn. J. Pediat. 134: 441-446, 1999.

[144388]25884.

[144389]Nagai, F.; Homma, H.; Tanase, H.; Matsui, M.: Studies on
the

[144390]genetic linkage of bilirubin and androsterone UDP-

glucuronyltransferases

[144391]by cross-breeding of two mutant rat strains. *Biochem. J.*
252: 897-900,

[144392]1988.

[144393]25885.

[144394]Peters, W. H. M.; Jansen, P. L. M.; Nauta, H.: The molecular
[144395]weights of UDP-glucuronyltransferase determined with
radiation-inactivation

[144396]analysis: a molecular model of bilirubin UDP-
glucuronyltransferase. *J.*

[144397]*Biol. Chem.* 259: 11701-11705, 1984.

[144398]25886.

[144399]Ritter, J. K.; Chen, F.; Sheen, Y. Y.; Tran, H. M.; Kimura, S.;

[144400]Yeatman, M. T.; Owens, I. S.: A novel complex locus UGT1
encodes

[144401]human bilirubin, phenol, and other UDP-
glucuronosyltransferase isozymes

[144402]with identical carboxyl termini. *J. Biol. Chem.* 267:
3257-3261,

[144403]1992.

[144404]25887.

[144405]Ritter, J. K.; Crawford, J. M.; Owens, I. S.: Cloning of two

[144406]human liver bilirubin UDP-glucuronosyltransferase cDNAs
with expression

[144407]in COS-1 cells. J. Biol. Chem. 266: 1043-1047, 1991.

[144408]25888.

[144409]Ritter, J. K.; Yeatman, M. T.; Ferreira, P.; Owens, I. S.: Identification

[144410]of a genetic alteration in the code for bilirubin UDP-glucuronosyltransferase

[144411]in the UGT1 gene complex of a Crigler-Najjar type I patient. J. Clin.

[144412]Invest. 90: 150-155, 1992.

[144413]25889.

[144414]Ritter, J. K.; Yeatman, M. T.; Kaiser, C.; Gridelli, B.; Owens,

[144415]I. S.: A phenylalanine codon deletion at the UGT1 gene complex locus

[144416]of a Crigler-Najjar type I patient generates a pH-sensitive bilirubin

[144417]UDP-glucuronosyltransferase. J. Biol. Chem. 268: 23573-23579, 1993.

[144418]25890.

[144419]Rosatelli, M. C.; Meloni, A.; Faa, V.; Saba, L.; Crisponi, G.;
[144420]Clemente, M. G.; Meloni, G.; Piga, M. T.; Cao, A.: Molecular
analysis
[144421]of patients of Sardinian descent with Crigler–Najjar syn-
drome type
[144422]I. J. Med. Genet. 34: 122–125, 1997.
[144423]25891.
[144424]Sampietro, M.; Lupica, L.; Perrero, L.; Comino, A.; Martinez
di
[144425]Montemuros, F.; Cappellini, M. D.; Fiorelli, G.: The expres-
sion of
[144426]uridine diphosphate glucuronosyltransferase gene is a
major determinant
[144427]of bilirubin level in heterozygous beta–thalassaemia and
in glucose–6–phosphate
[144428]dehydrogenase deficiency. Brit. J. Haemat. 99: 437–439,
1997.
[144429]25892.
[144430]Sato, H.; Sakai, Y.; Koiwai, O.; Watanabe, T.: Mapping of
the
[144431]mouse bilirubin UDP–glucuronosyltransferase gene
(Gnt–1) to chromosome

[144432]1 by restriction fragment length variations. Biochem.

Genet. 30:

[144433]347–352, 1992.

[144434]25893.

[144435]Seppen, J.; Bosma, P. J.; Goldhoorn, B. G.; Bakker, C. T. M.;

[144436]Chowdhury, J. R.; Chowdhury, N. R.; Jansen, P. L. M.; Oude
Elferink,

[144437]R. P. J.: Discrimination between Crigler–Najjar type I and II
by

[144438]expression of mutant bilirubin uridine diphosphate–
glucuronosyltransferase. J.

[144439]Clin. Invest. 94: 2385–2391, 1994.

[144440]25894.

[144441]Strassburg, C. P.; Oldhafer, K.; Manns, M. P.; Tukey, R. H.:

[144442]Differential expression of the UGT1A locus in human liver,
biliary,

[144443]and gastric tissue: identification of UGT1A7 and UGT1A10
transcripts

[144444]in extrahepatic tissue. Molec. Pharm. 52: 212–220, 1997.

[144445]25895.

- [144446]Tukey, R. H.; Strassburg, C. P.: Human UDP-glucuronosyltransferases: metabolism, expression, and disease. Annu. Rev. Pharm. Toxicol. 40: 581-616, 2000.
- [144449]25896.
- [144450]Ueyama, H.; Koiwai, O.; Soeda, Y.; et al.: Analysis of the promoter of human bilirubin UDP-glucuronosyltransferase gene (UGT1*1) in relevance to Gilbert's syndrome. Hepatol. Res. 9: 152-163, 1997.
- [144453]25897.
- [144454]van Es, H. H.; Bout, A.; Liu, J.; Anderson, L.; Duncan, A. M.; Bosma, P.; Oude Elferink, R.; Jansen, P. L.; Roy Chowdhury, J.; Schurr, E.: Assignment of the human UDP glucuronosyltransferase gene (UGT1A1) to chromosome region 2q37. Cytogenet. Cell Genet. 63: 114-116, 1993.
- [144458]25898.
- [144459]Wooster, R.; Sutherland, L.; Ebner, T.; Clarke, D.; Da Cruz

e

[144460]Silva, O.; Burchell, B.: Cloning and stable expression of a
new member

[144461]of the human liver phenol/bilirubin:UDP-glucuronosyltransferase cDNA

[144462]family. *Biochem. J.* 278: 465-469, 1991.

[144463]25899.

[144464]Yamamoto, K.; Soeda, Y.; Kamisako, T.; Hosaka, H.;
Fukano, M.;

[144465]Sato, H.; Fujiyama, Y.; Adachi, Y.; Satoh, Y.; Bamba, T.:
Analysis

[144466]of bilirubin uridine 5-prime-diphosphate
(UDP)-glucuronosyltransferase

[144467]gene mutations in seven patients with Crigler-Najjar syndrome type

[144468]II. *J. Hum. Genet.* 43: 111-114, 1998.

[144469]25900.

[144470]James, P. F.; Grupp, I. L.; Grupp, G.; Woo, A. L.; Askew, G.
R.;

[144471]Croyle, M. L.; Walsh, R. A.; Lingrel, J. B.: Identification of a

[144472]specific role for the Na,K-ATPase alpha-2 isoform as a

regulator of

[144473]calcium in the heart. Molec. Cell 3: 555–563, 1999.

[144474]25901.

[144475]Kent, R. B.; Fallows, D. A.; Geissler, E.; Glaser, T.; Emanuel,

[144476]J. R.; Lalley, P. A.; Levenson, R.; Housman, D. E.: Genes

encoding

[144477]alpha and beta subunits of Na,K-ATPase are located on

three different

[144478]chromosomes in the mouse. Proc. Nat. Acad. Sci. 84:

5369–5373, 1987.

[144479]25902.

[144480]Yang-Feng, T. L.; Schneider, J. W.; Lindgren, V.; Shull, M.

M.;

[144481]Benz, E. J., Jr.; Lingrel, J. B.; Francke, U.: Chromosomal lo-

calization

[144482]of human Na⁺,K⁺-ATPase alpha- and beta-subunit

genes. Genomics 2:

[144483]128–138, 1988.

[144484]25903.

[144485]Kawakami, K.; Nojima, H.; Ohta, T.; Nagano, K.: Molecular

cloning

[144486]and sequence analysis of human Na,K-ATPase beta-subunit. Nucleic

[144487]Acids Res. 14: 2833-2844, 1986.

[144488]25904.

[144489]Lane, L. K.; Shull, M. M.; Whitmer, K. R.; Lingrel, J. B.:
Characterization

[144490]of two genes for the human Na,K-ATPase beta subunit.
Genomics 5:

[144491]445-453, 1989.

[144492]25905.

[144493]Gloor, S.; Antonicek, H.; Sweadner, K. J.; Pagliusi, S.;
Frank,

[144494]R.; Moos, M.; Schachner, M.: The adhesion molecule on
glia (AMOG)

[144495]is a homologue of the beta subunit of the Na,K-ATPase. J.
Cell Biol. 110:

[144496]165-174, 1990.

[144497]25906.

[144498]Hsieh, C.-L.; Cheng-Deutsch, A.; Gloor, S.; Schachner, M.;
Francke,

[144499]U.: Assignment of Amog (adhesion molecule on glia) gene

to mouse

[144500]chromosome 11 near Zfp-3 and Asgr-1,2 and to human chromosome 17. Somat.

[144501]Cell Molec. Genet. 16: 401-405, 1990.

[144502]25907.

[144503]Malo, D.; Schurr, E.; Levenson, R.; Gros, P.: Assignment of Na,K-ATPase

[144504]beta-(2)-subunit gene (Atpb-2) to mouse chromosome 11. Genomics 6:

[144505]697-699, 1990.

[144506]25908.

[144507]Martin-Vasallo, P.; Dackowski, P.; Emanuel, J. R.; Levenson, R.

[144508]: Identification of a putative isoform of the Na,K-ATPase beta subunit:

[144509]primary structure and tissue-specific expression. J. Biol. Chem. 164:

[144510]4613-4618, 1989.

[144511]25909.

[144512]Pagliusi, S.; Antonicek, H.; Gloor, S.; Frank, R.; Moos, M.; Schachner,

[144513]M.: Identification of a cDNA clone specific for the neural adhesion

[144514]molecule AMOG. J. Neurosci. Res. 22: 113–119, 1989.

[144515]25910.

[144516]Katzmarzyk, P. T.; Rankinen, T.; Perusse, L.; Deriaz, O.; Tremblay,

[144517]A.; Borecki, I.; Rao, D. C.; Bouchard, C.: Linkage and association

[144518]of the sodium potassium-adenosine triphosphatase alpha-2 and beta-1

[144519]genes with respiratory quotient and resting metabolic rate in the

[144520]Quebec Family Study. J. Clin. Endocr. Metab. 84: 2093–2097, 1999.

[144521]25911.

[144522]Rampazzo, C.; Gallinaro, L.; Milanesi, E.; Frigimelica, E.; Reichard,

[144523]P.; Bianchi, V.: A deoxyribonucleotidase in mitochondria: involvement

[144524]in regulation of dNTP pools and possible link to genetic disease. Proc.

[144525]Nat. Acad. Sci. 97: 8239–8244, 2000.

[144526]25912.

[144527]Akaba, K.; Kimura, T.; Sasaki, A.; Tanabe, S.; Wakabayashi, T.;

[144528]Hiroi, M.; Yasumura, S.; Maki, K.; Aikawa, S.; Hayasaka, K.:
Neonatal

[144529]hyperbilirubinemia and a common mutation of the bilirubin uridine

[144530]diphosphate-glucuronosyltransferase gene in Japanese. J. Hum. Genet. 44:

[144531]22-25, 1999.

[144532]25913.

[144533]Aono, S.; Yamada, Y.; Keino, H.; Sasaoka, Y.; Nakagawa, T.; Onishi,

[144534]S.; Mimura, S.; Koiwai, O.; Sato, H.: A new type of defect in the

[144535]gene for bilirubin uridine

5-prime-diphosphate-glucuronosyltransferase

[144536]in a patient with Crigler-Najjar syndrome type I. Pediat. Res. 35:

[144537]629-632, 1994.

[144538]25914.

- [144539]Beutler, E.; Gelbart, T.; Demina, A.: Racial variability in the
- [144540]UDP-glucuronosyltransferase 1 (UGT1A1) promoter: a
balanced polymorphism
- [144541]for regulation of bilirubin metabolism? Proc. Nat. Acad.
Sci. 95:
- [144542]8170–8174, 1998.
- [144543]25915.
- [144544]Bosma, P. J.; Goldhoorn, B.; Oude Elferink, R. P. J.; Sinaas-
appel,
- [144545]M.; Oostra, B. A.; Jansen, P. L. M.: A mutation in bilirubin
uridine
- [144546]5-prime-diphosphate-glucuronosyltransferase isoform 1
causing Crigler-Najjar
- [144547]syndrome type II. Gastroenterology 105: 216–220, 1993.
- [144548]25916.
- [144549]Akaba, K.; Kimura, T.; Sasaki, A.; Tanabe, S.; Ikegami, T.;
Hashimoto,
- [144550]M.; Umeda, H.; Yoshida, H.; Umetsu, K.; Chiba, H.; Yuasa,
I.; Hayasaka,
- [144551]K.: Neonatal hyperbilirubinemia and mutation of the
bilirubin uridine

[144552]diphosphate-glucuronosyltransferase gene: a common missense mutation

[144553]among Japanese, Koreans and Chinese. Biochem. Molec. Biol. Int. 46:

[144554]21-26, 1998.

[144555]25917.

[144556]Bosma, P. J.; Roy Chowdhury, J.; Bakker, C.; Gantla, S.; de Boer,

[144557]A.; Oostra, B. A.; Lindhout, D.; Tytgat, G. N. J.; Jansen, P. L. M.;

[144558]Oude Elferink, R. P. J.; Roy Chowdhury, N.: The genetic basis of

[144559]the reduced expression of bilirubin UDP-glucuronosyltransferase 1

[144560]in Gilbert's syndrome. New Eng. J. Med. 333: 1171-1175, 1995.

[144561]25918.

[144562]Bosma, P. J.; Roy Chowdhury, J.; Huang, T.-J.; Lahiri, P.; Oude

[144563]Elferink, R. P. J.; Van Es, H. H. G.; Lederstein, M.; Whittington,

[144564]P. F.; Jansen, P. L. M.; Roy Chowdhury, N.: Mechanisms of

inherited

[144565] deficiencies of multiple UDP-glucuronosyltransferase iso-
forms in two

[144566] patients with Crigler-Najjar syndrome, type I. FASEB J. 6:
2859-2863,

[144567] 1992.

[144568] 25919.

[144569] Thornton, M. A.; Poncz, M.; Korostishevsky, M.; Yakobson,
E.;

[144570] Usher, S.; Seligsohn, U.; Peretz, H.: The human platelet al-
pha-IIb

[144571] gene is not closely linked to its integrin partner beta-3.
Blood 94:

[144572] 2039-2047, 1999.

[144573] 25920.

[144574] von dem Borne, A. E. G.; Decary, F.: Nomenclature of
platelet

[144575] specific antigens. Brit. J. Haemat. 74: 239-240, 1990.

[144576] 25921.

[144577] Xiong, J.-P.; Stehle, T.; Diefenbach, B.; Zhang, R.; Dunker,
R.;

[144578]Scott, D. L.; Joachimiak, A.; Goodman, S. L.; Arnaout, M. A.: Crystal

[144579]structure of the extracellular segment of integrin alpha-V-beta-3. Science 294:

[144580]339-345, 2001.

[144581]25922.

[144582]Xiong, J.-P.; Stehle, T.; Zhang, R.; Joachimiak, A.; Frech, M.;

[144583]Goodman, S. L.; Arnaout, M. A.: Crystal structure of the extracellular

[144584]segment of integrin alpha-V-beta-3 in complex with an Arg-Gly-Asp

[144585]ligand. Science 296: 151-155, 2002.

[144586]25923.

[144587]Disteche, C. M.; Plowman, G. D.; Gronwald, R. G. K.; Kelly, J.;

[144588]Bowen-Pope, D.; Adler, D. A.; Murray, J. C.: Mapping of the amphiregulin

[144589]and the platelet-growth factor receptor alpha genes to the proximal

[144590]long arm of chromosome 4. (Abstract) Cytogenet. Cell Genet. 51:

[144591]990, 1989.

[144592]25924.

[144593]Gronwald, R. G. K.; Adler, D. A.; Kelly, J. D.; Disteché, C.
M.;

[144594]Bowen-Pope, D. F.: The human PDGF receptor alpha-
subunit gene maps

[144595]to chromosome 4 in close proximity to c-kit. Hum. Genet.
85: 383-385,

[144596]1990.

[144597]25925.

[144598]Hol, F. A.; Geurds, M. P. A.; Chatkupt, S.; Shugart, Y. Y.;
Balling,

[144599]R.; Schrandt-Stumpel, C. T. R. M.; Johnson, W. G.; Hamel,
B. C. J.;

[144600]Mariman, E. C. M.: PAX genes and human neural tube de-
fects: an amino

[144601]acid substitution in PAX1 in a patient with spina bifida. J.
Med.

[144602]Genet. 8: 655-660, 1996.

[144603]25926.

[144604]Hsieh, C.-L.; Navankasattusas, S.; Escobedo, J. A.;

Williams, L.

[144605]T.; Francke, U.: Chromosomal localization of the gene for
AA-type

[144606]platelet-derived growth factor receptor (PDGFRA) in hu-
mans and mice. Cytogenet.

[144607]Cell Genet. 56: 160-163, 1991.

[144608]25927.

[144609]Ikuno, Y.; Kazlauskas, A.: TGF-beta-1-dependent con-
traction of

[144610]fibroblasts is mediated by th PDGF-alpha receptor. Invest.
Ophthal.

[144611]Vis. Sci. 43: 41-46, 2002.

[144612]25928.

[144613]Joosten, P. H. L. J.; Hol, F. A.; van Beersum, S. E. C.; Peters,

[144614]H.; Hamel, B. C. J.; Afink, G. B.; van Zoelen, E. J. J.; Mari-
man,

[144615]E. C. M.: Altered regulation of platelet-derived growth fac-
tor receptor-alpha

[144616]gene-transcription in vitro by spina bifida-associated mu-
tant Pax1

[144617]proteins. Proc. Nat. Acad. Sci. 95: 14459-14463, 1998.

[144618]25929.

[144619]Sanke, T.; Bell, G. I.; Sample, C.; Rubenstein, A. H.; Steiner,

[144620]D. F.: An islet amyloid peptide is derived from an
89-amino acid

[144621]precursor by proteolytic processing. J. Biol. Chem. 263:
17243-17246,

[144622]1988.

[144623]25930.

[144624]Verchere, C. B.; D'Alessio, D. A.; Palmiter, R. D.; Weir, G.
C.;

[144625]Bonner-Weir, S.; Baskin, D. G.; Kahn, S. E.: Islet amyloid
formation

[144626]associated with hyperglycemia in transgenic mice with
pancreatic beta

[144627]cell expression of human islet amyloid polypeptide. Proc.
Nat. Acad.

[144628]Sci. 93: 3492-3496, 1996.

[144629]25931.

[144630]Westermarck, P.; Engstrom, U.; Johnson, K. H.; Westermarck,
G. T.;

[144631]Betsholtz, C.: Islet amyloid polypeptide: pinpointing amino

acid

[144632]residues linked to amyloid fibril formation. Proc. Nat.

Acad. Sci. 87:

[144633]5036–5040, 1990.

[144634]25932.

[144635]Hockaday, T. D. R.: Hypogonadism and life-long anosmia.

Postgrad.

[144636]Med. J. 42: 572–574, 1966.

[144637]25933.

[144638]Oliveira, L. M. B.; Seminara, S. B.; Beranova, M.; Hayes, F.

J.;

[144639]Valkenburgh, S. B.; Schipani, E.; Costa, E. M. F.; Latronico,

A. C.;

[144640]Crowley, W. F., Jr.; Vallejo, M.: The importance of autosomal genes

[144641]in Kallmann syndrome: genotype–phenotype correlations and neuroendocrine

[144642]characteristics. J. Clin. Endocr. Metab. 86: 1532–1538, 2001.

[144643]25934.

[144644]Bignon, C.; Roux–Dosseto, M.; Zeigler, M. E.; Mattei,

M.-G.; Lissitzky,

[144645]J.-C.; Wicha, M. S.; Martin, P.-M.: Genomic analysis of the
67-kDa

[144646]laminin receptor in normal and pathological tissues: cir-
cumstantial

[144647]evidence for retroposon features. Genomics 10: 481-485,
1991.

[144648]25935.

[144649]Gehlsen, K. R.; Dillner, L.; Engvall, E.; Ruoslahti, E.: The
human

[144650]laminin receptor is a member of the integrin family of cell
adhesion

[144651]receptors. Science 241: 1228-1229, 1988. Note: Erratum:
Science

[144652]245:342-343, 1989.

[144653]25936.

[144654]Jackers, P.; Minoletti, F.; Belotti, D.; Clausse, N.; Sozzi, G.;

[144655]Sobel, M. E.; Castronovo, V.: Isolation from a multigene
family of

[144656]the active human gene of the metastasis-associated mul-
tifunctional

[144657]protein 37LRP/p40 at chromosome 3p21.3. Oncogene 13:

495–503, 1996.

[144658]25937.

[144659]Lafreniere, R. G.; Brown, C. J.; Rider, S.; Chelly, J.; Taillon–
Miller,

[144660]P.; Chinault, A. C.; Monaco, A. P.; Willard, H. F.: 2.6 Mb
YAC contig

[144661]of the human X inactivation center region in Xq13: physical linkage

[144662]of the RPS4X, PHKA1, XIST and DXS128E genes. Hum.
Molec. Genet. 2:

[144663]1105–1115, 1993.

[144664]25938.

[144665]Montuori, N.; Muller, F.; De Riu, S.; Fenzi, G.; Sobel, M. E.;

[144666]Rossi, G.; Vitale, M.: Laminin receptors in differentiated
thyroid

[144667]tumors: restricted expression of the 67–kilodalton laminin
receptor

[144668]in follicular carcinoma cells. J. Clin. Endocr. Metab. 84:
2086–2092,

[144669]1999.

[144670]25939.

[144671]Sato, K.; Narumi, K.; Sakai, T.; Abe, T.; Kikuchi, T.; Matsushima,
[144672]K.; Sindoh, S.; Motomiya, M.: Cloning of 67-kDa laminin
receptor
[144673]cDNA and gene expression in normal and malignant cell
lines of the
[144674]human lung. Cancer Lett. 62: 199–203, 1992.
[144675]25940.
[144676]Tohgo, A.; Takasawa, S.; Munakata, H.; Yonekura, H.;
Hayashi, N.;
[144677]Okamoto, H.: Structural determination and characterization of a 40
[144678]kDa protein isolated from rat 40 S ribosomal subunit. FEBS
Lett. 340:
[144679]133–138, 1994.
[144680]25941.
[144681]Yow, H. K.; Wong, J. M.; Chen, H. S.; Lee, C. G.; Davis, S.;
Steele,
[144682]G. D., Jr.; Chen, L. B.: Increased mRNA expression of a
laminin-binding
[144683]protein in human colon carcinoma: complete sequence of
a full-length

- [144684]cDNA encoding the protein. Proc. Nat. Acad. Sci. 85:
6394–6398,
- [144685]1988. Note: Erratum: Proc. Nat. Acad. Sci. 86: 7032 only,
1989.
- [144686]25942.
- [144687]Oklu, R.; Hesketh, R.: The latent transforming growth fac-
tor beta
- [144688]binding protein (LTBP) family. Biochem. J. 352: 601–610,
2000.
- [144689]25943.
- [144690]Lloyd, D. J.; Trembath, R. C.; Shackleton, S.: A novel inter-
action
- [144691]between lamin A and SREBP1: implications for partial
lipodystrophy
- [144692]and other laminopathies. Hum. Molec. Genet. 11:
769–777, 2002.
- [144693]25944.
- [144694]Wydner, K. L.; McNeil, J. A.; Lin, F.; Worman, H. J.;
Lawrence,
- [144695]J. B.: Chromosomal assignment of human nuclear enve-
lope protein genes
- [144696]LMNA, LMNB1, and LBR by fluorescence in situ hybridiza-

tion. Genomics 32:

[144697]474–478, 1996.

[144698]25945.

[144699]Garbarz, M.; Devaux, I.; Bournier, O.; Grandchamp, B.;
Dhermy,

[144700]D.: Protein 4.1 Lille, a novel mutation in the downstream
initiation

[144701]codon of protein 4.1 gene associated with heterozygous
4,1(–) hereditary

[144702]elliptocytosis. Hum. Mutat. 5: 339–340, 1995.

[144703]25946.

[144704]Garbarz, M.; Dhermy, D.; Lecomte, M. C.; Feo, C.; Chave-
roche,

[144705]I.; Galand, C.; Bournier, O.; Bertrand, O.; Boivin, P.: A vari-
ant

[144706]of erythrocyte membrane skeletal protein band 4.1 asso-
ciated with

[144707]hereditary elliptocytosis. Blood 64: 1006–1015, 1984.

[144708]25947.

[144709]Geerdink, R. A.; Nijenhuis, L. E.; Huizinga, J.: Hereditary
elliptocytosis:

[144710]linkage data in man. Ann. Hum. Genet. 30: 363–378,
1967.

[144711]25948.

[144712]Jensson, O.; Jonasson, T.; Olafsson, O.: Hereditary ellipto-
cytosis

[144713]in Iceland. Brit. J. Haemat. 13: 844–854, 1967.

[144714]25949.

[144715]Kan, Y.-W.: Personal Communication. San Francisco, Calif.

[144716]2/28/1986.

[144717]25950.

[144718]Kuroda, S.; Takeuchi, T.; Nagamori, H.: Data on the linkage between

[144719]elliptocytosis and Rh blood type. Jpn. J. Hum. Genet. 5:
112-118,

[144720]1960.

[144721]25951.

[144722]Lambert, S.; Conboy, J.; Zail, S.: A molecular study of heterozygous

[144723]protein 4.1 deficiency in hereditary elliptocytosis. Blood
72: 1926-1929,

[144724]1988.

[144725]25952.

[144726]Lambert, S.; Zail, S.: Partial deficiency of protein 4.1 in
hereditary

[144727]elliptocytosis. Am. J. Hemat. 26: 263-272, 1987.

[144728]25953.

- [144729]Lipton, E. L.: Elliptocytosis with hemolytic anemia: the effects
[144730]of splenectomy. Pediatrics 15: 67–82, 1955.
[144731]25954.
- [144732]Lux, S. E.; Wolfe, L. C.: Inherited disorders of the red cell
[144733]membrane skeleton. Pediat. Clin. N. Am. 27: 463–486,
1980.
[144734]25955.
- [144735]Marchesi, S. L.; Conboy, J.; Agre, P.; Letsinger, J. T.;
Marchesi,
[144736]V. T.; Speicher, D. W.; Mohandas, N.: Molecular analysis of
insertion/deletion
[144737]mutations in protein 4.1 in elliptocytosis. I. Biochemical
identification
[144738]of rearrangements in the spectrin/actin binding domain
and functional
[144739]characterizations. J. Clin. Invest. 86: 516–523, 1990.
[144740]25956.
- [144741]McGuire, M.; Agre, P.: Three distinct variants of protein
4.1
[144742]in Caucasian hereditary elliptocytosis. (Abstract) Clin. Res.

35:

[144743]428A, 1987.

[144744]25957.

[144745]McGuire, M.; Smith, B. L.; Agre, P.: Distinct variants of erythrocyte

[144746]protein 4.1 inherited in linkage with elliptocytosis and Rh type in

[144747]three white families. Blood 72: 287–293, 1988.

[144748]25958.

[144749]Morle, L.; Garbarz, M.; Alloisio, N.; Girot, R.; Chaverroche, I.;

[144750]Boivin, P.; Delaunay, J.: The characterization of protein 4.1 Presles,

[144751]a shortened variant of RBC membrane protein 4.1. Blood 65: 1511–1517,

[144752]1985.

[144753]25959.

[144754]Morle, L.; Pothier, B.; Alloisio, N.; Ducluzeau, M.–T.; Marques,

[144755]S.; Olim, G.; Martins e Silva, J.; Feo, C.; Garbarz, M.; Chaverroche,

- [144756]I.; Boivin, P.; Delaunay, J.: Red cell membrane alteration involving
[144757]protein 4.1 and protein 3 in a case of recessively inherited
haemolytic
[144758]anemia. Europ. J. Haemat. 38: 447–455, 1987.
[144759]25960.
- [144760]Morton, N. E.: The detection and estimation of linkage between
[144761]the genes for elliptocytosis and the Rh blood type. Am. J.
Hum. Genet. 8:
[144762]80–96, 1956.
[144763]25961.
- [144764]Nielsen, J. A.; Strunk, K. W.: Homozygous hereditary elliptocytosis
[144765]as the cause of haemolytic anemia in infancy. Scand. J.
Haemat. 5:
[144766]486–496, 1968.
[144767]25962.
- [144768]Parra, M.; Gascard, P.; Walensky, L. D.; Snyder, S. H.; Mohandas,
[144769]N.; Conboy, J. G.: Cloning and characterization of 4.1G

(EPB41L2),

[144770]a new member of the skeletal protein 4.1 (EPB41) gene family. Genomics 49:

[144771]298–306, 1998.

[144772]25963.

[144773]Peters, J. C.; Rowland, M.; Israels, L. G.; Zipursky, A.: Erythrocyte

[144774]sodium transport in hereditary elliptocytosis. Canad. J. Physiol.

[144775]Pharm. 44: 817–827, 1966.

[144776]25964.

[144777]Roberts, J. A. F.: Genetic linkage in man, with particular reference

[144778]to the usefulness of very small bodies of data. Quart. J. Med. 14:

[144779]27–33, 1945.

[144780]25965.

[144781]Shi, Z.-T.; Afzal, V.; Coller, B.; Patel, D.; Chasis, J. A.; Parra,

[144782]M.; Lee, G.; Paszty, C.; Stevens, M.; Walensky, L.; Peters, L. L.;

- [144783]Mohandas, N.; Rubin, E.; Conboy, J. G.: Protein
4.1R-deficient mice
[144784]are viable but have erythroid membrane skeleton abnormalities. J.
[144785]Clin. Invest. 103: 331-340, 1999.
[144786]25966.
- [144787]Takakuwa, Y.; Tchernia, G.; Rossi, M.; Benabadji, M.; Mo-
handas,
[144788]N.: Restoration of normal membrane stability to unstable
protein
[144789]4.1-deficient erythrocyte membranes by incorporation of
purified protein
[144790]4.1. J. Clin. Invest. 78: 80-85, 1986.
[144791]25967.
- [144792]Tang, C.-J. C.; Tang, T. K.: Rapid localization of mem-
brane skeletal
[144793]protein 4.1 (EL1) to human chromosome 1p33-p34.2 by
nonradioactive
[144794]in situ hybridization. Cytogenet. Cell Genet. 57: 119,
1991.
[144795]25968.

[144796]Tang, T. K.; Leto, T. L.; Correas, I.; Alonso, M. A.; Marchesi,

[144797]V. T.; Benz, E. J., Jr.: Selective expression of an erythroid-specific

[144798]isoform of protein 4.1. Proc. Nat. Acad. Sci. 85: 3713–3717, 1988.

[144799]25969.

[144800]Tchernia, G.; Mohandas, N.; Shohet, S. B.: Deficiency of skeletal

[144801]membrane protein band 4.1 in homozygous hereditary elliptocytosis:

[144802]implications for erythrocyte membrane stability. J. Clin. Invest. 68:

[144803]454–460, 1981.

[144804]25970.

[144805]Tomaselli, M. B.; John, K. M.; Lux, S. E.: Elliptical erythrocyte

[144806]membrane skeletons and heat-sensitive spectrin in hereditary elliptocytosis. Proc.

[144807]Nat. Acad. Sci. 78: 1911–1915, 1981.

[144808]25971.

- [144809] Brands, J. H. G. M.; Maassen, J. A.; Van Hemert, F. J.;
Amons,
[144810] R.; Moller, W.: The primary structure of the alpha subunit
of human
[144811] elongation factor 1: structural aspects of guanine-nu-
cleotide-binding
[144812] sites. Europ. J. Biochem. 155: 167-171, 1986.
[144813] 25972.
[144814] Ditzel, H. J.; Masaki, Y.; Nielsen, H.; Farnaes, L.; Burton, D.
[144815] R.: Cloning and expression of a novel human antibody-
-antigen pair
[144816] associated with Felty's syndrome. Proc. Nat. Acad. Sci. 97:
9234-9239,
[144817] 2000.
[144818] 25973.
[144819] Lund, A.; Knudsen, S. M.; Vissing, H.; Clark, B.; Tom-
merup, N.
[144820]: Assignment of human elongation factor 1-alpha genes:
EEF1A maps
[144821] to chromosome 6q14 and EEF1A2 to 20q13.3. Genomics
36: 359-361,

[144822]1996.

[144823]25974.

[144824]Opdenakker, G.; Cabeza-Arvelaiz, Y.; Fiten, P.; Dijkmans, R.; Van

[144825]Damme, J.; Volckaert, G.; Billiau, A.; Van Elsen, A.; Van der Schueren,

[144826]B.; Van den Berghe, H.; Cassiman, J.-J.: Human elongation factor

[144827]1-alpha: a polymorphic and conserved multigene family with multiple

[144828]chromosomal localizations. Hum. Genet. 75: 339-344, 1987.

[144829]25975.

[144830]Cohen, A. J.; Li, F. P.; Berg, S.; Marchetto, D. J.; Tsai, S.;

[144831]Jacobs, S. C.; Brown, R. S.: Hereditary renal-cell carcinoma associated

[144832]with chromosomal translocation. New Eng. J. Med. 301: 592-595, 1979.

[144833]25976.

[144834]Gemmell, R. M.; West, J. D.; Boldog, F.; Tanaka, N.; Robinson,

[144835]L. J.; Smith, D. I.; Li, F.; Drabkin, H. A.: The hereditary renal

[144836]cell carcinoma 3;8 translocation fuses FHIT to a patched-related gene,

[144837]TRC8. Proc. Nat. Acad. Sci. 95: 9572–9577, 1998.

[144838]25977.

[144839]Nakano, A.; Pulkkinen, L.; Murrell, D.; Rico, J.; Lucky, A. W.;

[144840]Garzon, M.; Stevens, C. A.; Robertson, S.; Pfendner, E.; Uitto, J.

[144841]: Epidermolysis bullosa with congenital pyloric atresia: novel mutations

[144842]in the beta-4 integrin gene (ITGB4) and genotype/phenotype correlations. Pediat.

[144843]Res. 49: 618–626, 2001.

[144844]25978.

[144845]Pulkkinen, L.; Bruckner-Tuderman, L.; August, C.; Uitto, J.: Compound

[144846]heterozygosity for missense mutation (L156P) and non-sense (R554X)

[144847]mutations in the beta-4 integrin gene (ITGB4) underlies mild, nonlethal

[144848]phenotype of epidermolysis bullosa with pyloric atresia.

Am. J. Path. 152:

[144849]935–941, 1998.

[144850]25979.

[144851]Pulkkinen, L.; Rouan, F.; Bruckner–Tuderman, L.; Wallerstein, R.;

[144852]Garzon, M.; Brown, T.; Smith, L.; Carter, W.; Uitto, J.: Novel
ITGB4

[144853]mutations in lethal and nonlethal variants of epidermolysis bullosa

[144854]with pyloric atresia: missense versus nonsense. Am. J.
Hum. Genet. 63:

[144855]1376–1387, 1998.

[144856]25980.

[144857]Shaw, L. M.; Rabinovitz, I., Wang, H. H.–F.; Toker, A., Mercurio,

[144858]A. M.: Activation of phosphoinositide 3–OH kinase by the
alpha–6/beta–4

[144859]integrin promotes carcinoma invasion. Cell 91: 949–960,
1997.

[144860]25981.

[144861]Suzuki, S.; Naitoh, Y.: Amino acid sequence of a novel integrin

[144862]beta-4 subunit and primary expression of the mRNA in epithelial cells. EMBO

[144863]J. 9: 757-763, 1990.

[144864]25982.

[144865]Vidal, F.; Aberdam, D.; Miquel, C.; Christiano, A. M.; Pulkkinen,

[144866]L.; Uitto, J.; Ortonne, J.-P.; Meneguzzi, G.: Integrin beta-4 mutations

[144867]associated with junctional epidermolysis bullosa with pyloric atresia. Nature

[144868]Genet. 10: 229-234, 1995.

[144869]25983.

[144870]Chany, C.; Vignal, M.; Couillin, P.; Van Cong, N.; Boue, J.; Boue,

[144871]A.: Chromosomal localization of human genes governing the interferon-induced

[144872]antiviral state. Proc. Nat. Acad. Sci. 72: 3129-3133, 1975.

[144873]25984.

[144874]Aplin, H. M.; Hirst, K. L.; Crosby, A. H.; Dixon, M. J.: Map-

ping

[144875]of the human dentin matrix acidic phosphoprotein gene
(DMP1) to the

[144876]dentinogenesis imperfecta type II critical region at chro-
mosome 4q21. Genomics 30:

[144877]347-349, 1995.

[144878]25985.

[144879]Fisher, L. W.; McBride, O. W.; Termine, J. D.; Young, M. F.:
Human

[144880]bone sialoprotein: deduced protein sequence and chro-
mosomal localization.

[144881]J. Biol. Chem. 265: 2347-2351, 1990.

[144882]25986.

[144883]Goold, R. D.; diSibio, G. L.; Xu, H.; Lang, D. B.; Dadgar, J.;

[144884]Magrane, G. G.; Dugaiczky, A.; Smith, K. A.; Cox, D. R.;
Masters,

[144885]S. B.; Myers, R. M.: The development of sequence-tagged
sites for

[144886]human chromosome 4. Hum. Molec. Genet. 2:
1271-1288, 1993.

[144887]25987.

[144888]Kerr, J. M.; Fisher, L. W.; Termine, J. D.; Wang, M. G.;
McBride,

[144889]O. W.; Young, M. F.: The human bone sialoprotein gene
(IBSP): genomic

[144890]localization and characterization. Genomics 17: 408–415,
1993.

[144891]25988.

[144892]Kim, R. H.; Shapiro, H. S.; Li, J. J.; Wrana, J. L.; Sodek, J.

[144893]: Characterization of the human bone sialoprotein (BSP)
gene and its

[144894]promoter sequence. Matrix Biol. 14: 31–40, 1994.

[144895]25989.

[144896]Young, M. F.; Ibaraki, K.; Kerr, J. M.; Lyu, M. S.; Kozak, C.
A.

[144897]: Murine bone sialoprotein (BSP): cDNA cloning, mRNA ex-
pression, and

[144898]genetic mapping. Mammalian Genome 5: 108–111, 1994.

[144899]25990.

[144900]Ho, A. S. Y.; Liu, Y.; Khan, T. A.; Hsu, D.-H.; Bazan, J. F.;
Moore,

[144901]K. W.: A receptor for interleukin 10 is related to interferon

receptors. Proc.

[144902]Nat. Acad. Sci. 90: 11267–11271, 1993.

[144903]25991.

[144904]Liu, Y.; Wei, S. H.–Y.; Ho, A. S.–Y.; de Waal Malefyt, R.;

Moore,

[144905]K. W.: Expression cloning and characterization of a human

IL-10 receptor. J.

[144906]Immun. 152: 1821–1829, 1994.

[144907]25992.

[144908]Barker, D. F.; Hostikka, S. L.; Zhou, J.; Chow, L. T.;

Oliphant,

[144909]A. R.; Gerken, S. C.; Gregory, M. C.; Skolnick, M. H.; Atkin,

C. L.;

[144910]Tryggvason, K.: Identification of mutations in the COL4A5
collagen

[144911]gene in Alport syndrome. Science 248: 1224–1227, 1990.

[144912]25993.

[144913]Draptchinskaia, N.; Gustavsson, P.; Andersson, B.; Petters-
son,

[144914]M.; Willig, T.–N.; Dianzani, I.; Ball, S.; Tchernia, G.; Klar, J.;

[144915]Matsson, H.; Tentler, D.; Mohandas, N.; Carlsson, B.; Dahl,

N.: The

[144916] gene encoding ribosomal protein S19 is mutated in Diamond-Blackfan

[144917] anaemia. *Nature Genet.* 21: 169–175, 1999.

[144918] 25994.

[144919] Tanzi, R. E.; St. George-Hyslop, P. H.; Gusella, J. F.: Molecular

[144920] genetics of Alzheimer disease amyloid. *J. Biol. Chem.* 266: 20579–20582,

[144921] 1991.

[144922] 25995.

[144923] Thinakaran, G.; Borchelt, D. R.; Lee, M. K.; Slunt, H. H.; Spitzer,

[144924] L.; Kim, G.; Ratovitsky, T.; Davenport, F.; Nordstedt, C.; Seeger,

[144925] M.; Hardy, J.; Levey, A. I.; Gandy, S. E.; Jenkins, N. A.; Copeland,

[144926] N. G.; Price, D. L.; Sisodia, S. S.: Endoproteolysis of presenilin

[144927] 1 and accumulation of processed derivatives in vivo. *Neuron* 17:

[144928] 181–190, 1996.

[144929]25996.

[144930]Struhl, G.; Adachi, A.: Requirements for Presenilin-dependent

[144931]cleavage of Notch and other transmembrane proteins.
Molec. Cell 6:

[144932]625–636, 2000.

[144933]25997.

[144934]Struhl, G.; Greenwald, I.: Presenilin is required for activity

[144935]and nuclear access of Notch in Drosophila. Nature 398:
522–525,

[144936]1999.

[144937]25998.

[144938]Takahashi, Y.; Koizumi, K.; Takagi, A.; Kitajima, S.; Inoue,
T.;

[144939]Koseki, H.; Saga, Y.: Mesp2 initiates somite segmentation
through

[144940]the Notch signalling pathway. Nature Genet. 25: 390–396,
2000.

[144941]25999.

[144942]Takashima, A.; Sato, M.; Mercken, M.; Tanaka, S.; Kondo,
S.; Honda,

[144943]T.; Sato, K.; Murayama, M.; Noguchi, K.; Nakazato, Y.;
Takahashi,

[144944]H.: Localization of Alzheimer-associated presenilin 1 in
transfected

[144945]COS-7 cells. Biochem. Biophys. Res. Commun. 227:
423-426, 1996.

[144946]26000.

[144947]Trower, M. K.; Orton, S. M.; Purvis, I. J.; Sanseau, P.; Riley,

[144948]J.; Christodoulou, C.; Burt, D.; See, C. G.; Elgar, G.; Sher-
rington,

[144949]R.; Rogaev, E. I.; St. George-Hyslop, P.; Brenner, S.; Dykes,
C. W.

[144950]: Conservation of synteny between the genome of the
pufferfish (*Fugu*

[144951]*rubripes*) and the region on human chromosome 14
(14q24.3) associated

[144952]with familial Alzheimer disease (AD3 locus). Proc. Nat.
Acad. Sci. 93:

[144953]1366-1369, 1996.

[144954]26001.

[144955]Tysoe, C.; Whittaker, J.; Xuereb, J.; Cairns, N. J.; Cruts, M.;

- [144956] Van Broeckhoven, C.; Wilcock, G.; Rubinsztein, D. C.: A
presenilin-1
[144957] truncating mutation is present in two cases with autopsy-
confirmed
[144958] early-onset Alzheimer disease. Am. J. Hum. Genet. 62:
70-76, 1998.
[144959] 26002.
- [144960] Van Broeckhoven, C.; Backhovens, H.; Cruts, M.; De Win-
ter, G.;
[144961] Bruylant, M.; Cras, P.; Martin, J.-J.: Mapping of a gene
predisposing
[144962] to early-onset Alzheimer's disease to chromosome
14q24.3. Nature
[144963] Genet. 2: 335-339, 1992.
[144964] 26003.
- [144965] Verkkoniemi, A.; Somer, M.; Rinne, J. O.; Myllykangas, L.;
Crook,
[144966] R.; Hardy, J.; Viitanen, M.; Kalimo, H.; Haltia, M.: Variant
Alzheimer's
[144967] disease with spastic paraparesis: clinical characterization.
Neurology 54:
[144968] 1103-1109, 2000.

[144969]26004.

[144970]Wasco, W.; Pettingell, W. P.; Jondro, P. D.; Schmidt, S. D.;
Gurubhagaratula,

[144971]S.; Rodes, L.; DiBlasi, T.; Romano, D. M.; Guenette, S. Y.;
Kovacs,

[144972]D. M.; Growdon, J. H.; Tanzi, R. E.: Familial Alzheimer's
chromosome

[144973]14 mutations. (Letter) Nature Med. 1: 848, 1995.

[144974]26005.

[144975]Wilson, C. A.; Doms, R. W.; Zheng, H.; Lee, V. M.-Y.: Pre-
senilins

[144976]are not required for A-beta-42 production in the early se-
cretory pathway. Nature

[144977]Neurosci. 5: 849-855, 2002.

[144978]26006.

[144979]Wittenburg, N.; Eimer, S.; Lakowski, B.; Rohrig, S.;
Rudolph,

[144980]C.; Baumeister, R.: Presenilin is required for proper mor-
phology

[144981]and function of neurons in C. elegans. Nature 406:
306-309, 2000.

[144982]26007.

[144983]Wolfe, M. S.; Xia, W.; Ostaszewski, B. L.; Diehl, T. S.; Kimberly,

[144984]W. T.; Selkoe, D. J.: Two transmembrane aspartates in presenilin-1

[144985]required for presenilin endoproteolysis and gamma-secretase activity. Nature 398:

[144986]513-517, 1999.

[144987]26008.

[144988]Xia, X.; Qian, S.; Soriano, S.; Wu, Y.; Fletcher, A. M.; Wang,

[144989]X.-J.; Koo, E. H.; Wu, X.; Zheng, H.: Loss of presenilin 1 is associated

[144990]with enhanced beta-catenin signaling and skin tumorigenesis. Proc.

[144991]Nat. Acad. Sci. 98: 10863-10868, 2001.

[144992]26009.

[144993]Ye, Y.; Lukinova, N.; Fortini, M. E.: Neurogenic phenotypes and

[144994]altered Notch processing in Drosophila presenilin mutants. Nature 398:

[144995]525-529, 1999.

[144996]26010.

[144997]Yu, H.; Saura, C. A.; Choi, S.-Y.; Sun, L. D.; Yang, X.; Handler,

[144998]M.; Kawarabayashi, T.; Younkin, L.; Fedeles, B.; Wilson, M. A.; Younkin,

[144999]S.; Kandel, E. R.; Kirkwood, A.; Shen, J.: APP processing and synaptic

[145000]plasticity in presenilin-1 conditional knockout mice. *Neuron* 31:

[145001]713-726, 2001.

[145002]26011.

[145003]Zhang, Z.; Hartmann, H.; Minh Do, V.; Abramowski, D.; Sturchler-Pierrat,

[145004]C.; Staufenbiel, M.; Sommer, B.; van de Weterling, M.; Clevers, H.;

[145005]Saftig, P.; De Strooper, B.; He, X.; Yanker, B. A.: Destabilization

[145006]of beta-catenin by mutations in presenilin-1 potentiates neuronal

[145007]apoptosis. *Nature* 395: 698-702, 1998.

[145008]26012.

[145009]Karrman, C.; Backman, B.; Dixon, M.; Holmgren, G.; Forsman, K.

[145010] Mapping of the locus for autosomal dominant amelogenesis imperfecta

[145011](AIH2) to a 4-Mb YAC contig on chromosome 4q11-q21. Genomics 39:

[145012]164-170, 1997.

[145013]26013.

[145014]Rajpar, M. H.; Harley, K.; Laing, C.; Davies, R. M.; Dixon, M.

[145015]J.: Mutation of the gene encoding the enamel-specific protein, enamelin,

[145016]causes autosomal-dominant amelogenesis imperfecta. Hum. Molec. Genet. 10:

[145017]1673-1677, 2001.

[145018]26014.

[145019]Bleyl, S.; Nelson, L.; Odelberg, S. J.; Ruttenberg, H. D.; Otterud,

[145020]B.; Leppert, M.; Ward, K.: A gene for familial total anomalous pulmonary

[145021]venous return maps to chromosome 4p13-q12. Am. J. Hum. Genet. 56:

[145022]408–415, 1995.

[145023]26015.

[145024]Blixt, A.; Mahlapuu, M.; Aitola, M.; Pelto-Huikko, M.; Enerback,

[145025]S.; Carlsson, P.: A forkhead gene, FoxE3, is essential for lens epithelial

[145026]proliferation and closure of the lens vesicle. Genes Dev. 14: 245–254,

[145027]2000.

[145028]26016.

[145029]Semina, E. V.; Brownell, I.; Mintz-Hittner, H. A.; Murray, J. C.;

[145030]Jamrich, M.: Mutations in the human forkhead transcription factor

[145031]FOXE3 associated with anterior segment ocular dysgenesis and cataracts. Hum.

[145032]Molec. Genet. 10: 231–236, 2001.

[145033]26017.

[145034]Siegelmann-Danieli, N.; Buetow, K. H.: Constitutional genetic

[145035]variation at the human aromatase gene (Cyp19) and

breast cancer risk. Brit.

[145036]J. Cancer 79: 456–463, 1999.

[145037]26018.

[145038]Simpson, E. R.; Michael, M. D.; Agarwal, V. R.; Hinshelwood, M.

[145039]M.; Bulun, S. E.; Zhao, Y.: Expression of the CYP19 (aromatase) gene:

[145040]an unusual case of alternative promoter usage. FASEB J. 11: 29–36,

[145041]1997.

[145042]26019.

[145043]Sparkes, R. S.; Mohandas, T.; Chen, S.; Besman, M. J.; Zollman,

[145044]S.; Shively, J. E.: Assignment of the aromatase gene to human chromosome

[145045]15q21. (Abstract) Cytogenet. Cell Genet. 46: 696–697, 1987.

[145046]26020.

[145047]Toda, K.; Merashima, M.; Kawamoto, T.; Sumimoto, H.; Yokoyama,

[145048]Y.; Kuribayashi, I.; Mitsuuchi, Y.; Maeda, T.; Yamamoto, Y.;

Sagara,

[145049]Y.; Ikeda, H.; Shizuta, Y.: Structural and functional characterization

[145050]of human aromatase P-450 gene. *Europ. J. Biochem.* 193: 559-565,

[145051]1990.

[145052]26021.

[145053]Wang, Z. J.; Jeffs, B.; Ito, M.; Achermann, J. C.; Yu, R. N.;

[145054]Hales, D. B.; Jameson, J. L.: Aromatase (Cyp19) expression is up-regulated

[145055]by targeted disruption of Dax1. *Proc. Nat. Acad. Sci.* 98: 7988-7993,

[145056]2001.

[145057]26022.

[145058]Whitlock, J. P., Jr.: The regulation of cytochrome P-450 gene

[145059]expression. *Annu. Rev. Pharm. Toxicol.* 26: 333-369, 1986.

[145060]26023.

[145061]Zhou, D.; Pompon, D.; Chen, S.: Structure-function studies of

[145062]human aromatase by site-directed mutagenesis: kinetic properties of

[145063]mutants pro308-to-phe, tyr361-to-phe, tyr361-to-leu, and phe406-to-arg. Proc.

[145064]Nat. Acad. Sci. 88: 410-414, 1991.

[145065]26024.

[145066]Luttrell, L. M.; Ferguson, S. S. G.; Daaka, Y.; Miller, W. E.;

[145067]Maudsley, S.; Della Rocca, G. J.; Lin, F.-T.; Kawakatsu, H.; Owada,

[145068]K.; Luttrell, D. K.; Caron, M. G.; Lefkowitz, R. J.: Beta-arrestin-dependent

[145069]formation of beta-2 adrenergic receptor-Src protein kinase complexes. Science 283:

[145070]655-661, 1999.

[145071]26025.

[145072]McDonald, P. H.; Chow, C.-W.; Miller, W. E.; Laporte, S. A.; Field,

[145073]M. E.; Lin, F.-T.; Davis, R. J.; Lefkowitz, R. J.: Beta-arrestin

[145074]2: a receptor-regulated MAPK scaffold for the activation of JNK3. Science 290:

[145075]1574-1577, 2000.

[145076]26026.

[145077]Ferns, G. A. A.; Stocks, J.; Ritchie, C.; Galton, D. J.: Genetic

[145078]polymorphisms of apolipoprotein C-III and insulin in survivors of

[145079]myocardial infarction. Lancet II: 300-303, 1985.

[145080]26027.

[145081]Holt, M.; Hogan, P. F.; Nurse, G. T.: The ovalocytosis polymorphism

[145082]on the western border of Papua New Guinea. Hum. Biol. 53: 23-34,

[145083]1981.

[145084]26028.

[145085]Hsu, L.; Morrison, M.: A new variant of the anion transport protein

[145086]in human erythrocytes. Biochemistry 24: 3086-3090, 1985.

[145087]26029.

[145088]Ideguchi, H.; Okubo, K.; Ishikawa, A.; Futata, Y.; Hamasaki, N.

[145089]: Band 3-Memphis is associated with a lower transport rate of phosphoenolpyruvate. Brit.

[145090]J. Haemat. 82: 122–125, 1992.

[145091]26030.

[145092]Iwase, S.; Ideguchi, H.; Takao, M.; Horiguchi–Yamada, J.;
Iwasaki,

[145093]M.; Takahara, S.; Sekikawa, T.; Mochizuki, S.; Yamada, H.:
Band 3

[145094]Tokyo: thr837–ala837 substitution in erythrocyte band 3
protein associated

[145095]with spherocytic hemolysis. Acta Haemat. 100: 200–203,
1998.

[145096]26031.

[145097]Jarolim, P.; Palek, J.; Rubin, H. L.; Prchal, J. T.; Korsgren,

[145098]C.; Cohen, C. M.: Band 3 Tuscaloosa: pro327–to–arg327
substitution

[145099]in the cytoplasmic domain of erythrocyte band 3 protein
associated

[145100]with spherocytic hemolytic anemia and partial deficiency
of protein

[145101]4.2. (Abstract) Blood 78 (suppl.): 252a, 1991.

[145102]26032.

[145103]Jarolim, P.; Rubin, H. L.; Brabec, V.; Chrobak, L.; Zolotarev,

[145104]A. S.; Alper, S. L.; Brugnara, C.; Wichterle, H.; Palek, J.:

Mutations

[145105]of conserved arginines in the membrane domain of erythroid band 3

[145106]lead to a decrease in membrane-associated band 3 and to the phenotype

[145107]of hereditary spherocytosis. Blood 85: 634–640, 1995.

[145108]26033.

[145109]Jarolim, P.; Rubin, H. L.; Liu, S.-C.; Cho, M. R.; Brabec, V.;

[145110]Derick, L. H.; Yi, S. J.; Saad, S. T. O.; Alper, S.; Brugnara, C.;

[145111]Golan, D. E.; Palek, J.: Duplication of 10 nucleotides in the erythroid

[145112]band 3 (AE1) gene in a kindred with hereditary spherocytosis and band

[145113]3 protein deficiency (band 3–Prague). J. Clin. Invest. 93: 121–130,

[145114]1994.

[145115]26034.

[145116]Inaba, M.; Yawata, A.; Koshino, I.; Sato, K.; Takeuchi, M.; Takakuwa,

[145117]Y.; Manno, S.; Yawata, Y.; Kanzaki, A.; Sakai, J.; Ban, A.;
Ono, K.;

[145118]Maede, Y.: Defective anion transport and marked spherocytosis with

[145119]membrane instability caused by hereditary total deficiency
of red

[145120]cell band 3 in cattle due to a nonsense mutation. J. Clin.
Invest. 97:

[145121]1804–1817, 1996.

[145122]26035.

[145123]Inoue, T.; Kanzaki, A.; Kaku, M.; Yawata, A.; Takezono, M.;
Okamoto,

[145124]N.; Wada, H.; Sugihara, T.; Yamada, O.; Katayama, Y.; Nagata, N.;

[145125]Yawata, Y.: Homozygous missense mutation (band 3
Fukuoka: G130R):

[145126]a mild form of hereditary spherocytosis with near-normal
band 3 content

[145127]and minimal changes of membrane ultrastructure despite
moderate protein

[145128]4.2 deficiency. Brit. J. Haemat. 102: 932–939, 1998.

[145129]26036.

[145130]Jarolim, P.; Rubin, H. L.; Zakova, D.; Storry, J.; Reid, M. E.

[145131]: Characterization of seven low incidence blood group antigens carried

[145132]by erythrocyte band 3 protein. Blood 92: 4836–4843, 1998.

[145133]26037.

[145134]Jarolim, P.; Rubin, H. L.; Zhai, S.; Sahr, K. E.; Liu, S.-C.;

[145135]Mueller, T. J.; Palek, J.: Band 3 Memphis: a widespread polymorphism

[145136]with abnormal electrophoretic mobility of erythrocyte band 3 protein

[145137]caused by substitution AAG-to-GAG (lys-to-glu) in codon 56. Blood 80:

[145138]1592–1598, 1992.

[145139]26038.

[145140]Jay, D. G.: Role of band 3 in homeostasis and cell shape. Cell 86:

[145141]853–854, 1996.

[145142]26039.

[145143]Jenkins, P. B.; Abou-Alfa, G. K.; Dhermy, D.; Bursaux, E.; Feo,

[145144]C.; Scarpa, A. L.; Lux, S. E.; Garbarz, M.; Forget, B. G.; Gallagher,

[145145]P. G.: A nonsense mutation in the erythrocyte band 3 gene associated

[145146]with decreased mRNA accumulation in a kindred with dominant hereditary

[145147]spherocytosis. J. Clin. Invest. 97: 373–380, 1996.

[145148]26040.

[145149]Jones, G. L.; Edmundson, H. M.; Wesche, D.; Saul, A.: Human erythrocyte

[145150]band–3 has an altered N terminus in malaria-resistant Melanesian ovalocytosis. Biochim.

[145151]Biophys. Acta 1096: 33–40, 1990.

[145152]26041.

[145153]Kay, M. M. B.: Band 3 in aging and neurological disease. Ann.

[145154]N.Y. Acad. Sci. 621: 179–204, 1991.

[145155]26042.

[145156]Kay, M. M. B.; Bosman, G. J. C. G. M.; Lawrence, C.: Functional

[145157]topography of band 3: specific structural alteration linked

to functional

[145158]aberrations in human erythrocytes. Proc. Nat. Acad. Sci.

85: 492-496,

[145159]1988.

[145160]26043.

[145161]Kay, M. M. B.; Flowers, N.; Goodman, J.; Bosman, G.: Alter-
ation

[145162]in membrane protein band 3 associated with accelerated
erythrocyte

[145163]aging. Proc. Nat. Acad. Sci. 86: 5834-5838, 1989.

[145164]26044.

[145165]Kay, M. M. B.; Lawrence, C.; Bosman, G. J. C. G. M.: Molec-
ular

[145166]anatomy of an anemia. (Abstract) Clin. Res. 35: 599A,
1987.

[145167]26045.

[145168]Kay, M. M. B.; Marchalonis, J. J.; Hughes, J.; Watanabe, K.;
Schluter,

[145169]S. F.: Definition of a physiologic aging autoantigen by us-
ing synthetic

[145170]peptides of membrane protein band 3: localization of the

active antigenic

[145171]sites. Proc. Nat. Acad. Sci. 87: 5734–5738, 1990.

[145172]26046.

[145173]Kidson, C.; Lamont, G.; Saul, A.; Nurse, G. T.: Ovalocytic erythrocytes

[145174]from Melanesians are resistant to invasion by malaria parasites in

[145175]culture. Proc. Nat. Acad. Sci. 78: 5829–5832, 1981.

[145176]26047.

[145177]Langdon, R. G.; Holman, V. P.: Immunological evidence that band

[145178]3 is the major glucose transporter of the human erythrocyte membrane. Biochim.

[145179]Biophys. Acta 945: 23–32, 1988.

[145180]26048.

[145181]Lie-Injo, L. E.: Hereditary ovalocytosis and haemoglobin E-ovalocytosis

[145182]in Malayan aborigines. Nature 208: 1329, 1965.

[145183]26049.

[145184]Lie-Injo, L. E.; Fix, A.; Bolton, J. M.; Gilman, R. H.:

Haemoglobin

- [145185]E-hereditary elliptocytosis in Malayan aborigines. *Acta Haemat.* 47: 210–216, 1972.
- [145187]26050.
- [145188]Liu, S.-C.; Jarolim, P.; Rubin, H. L.; Palek, J.; Amato, D.; Hassan, K.; Zaik, M.; Sapak, P.: The homozygous state for the band 3 protein mutation in Southeast Asian ovalocytosis may be lethal. (Letter) *Blood* 84: 3590–3591, 1994.
- [145192]26051.
- [145193]Liu, S.-C.; Zhai, S.; Palek, J.; Golan, D. E.; Amato, D.; Hassan, K.; Nurse, G. T.; Babona, D.; Coetzer, T.; Jarolim, P.; Zaik, M.; Borwein, S.: Molecular defect of the band 3 protein in Southeast Asian ovalocytosis. *New Eng. J. Med.* 323: 1530–1538, 1990.
- [145197]26052.

[145198]Lux, S.; Bedrosian, C.; Shalev, O.; Morris, M.; Chasis, J.;
Davies,

[145199]K.; Savvides, P.; Telen, M.: Deficiency of band 3 in dominant hereditary

[145200]spherocytosis with normal spectrin content. (Abstract)
Clin. Res. 38:

[145201]300A, 1990.

[145202]26053.

[145203]Lux, S. E.; John, K. M.; Kopito, R. R.; Lodish, H. F.: Cloning

[145204]and characterization of band 3, the human erythrocyte
anion-exchange

[145205]protein (AE1). Proc. Nat. Acad. Sci. 86: 9089-9093, 1989.

[145206]26054.

[145207]Maillet, P.; Vallier, A.; Reinhart, W. H.; Wyss, E. J.; Ott, P.;

[145208]Texier, P.; Baklouti, F.; Tanner, M. J. A.; Delaunay, J.; Aloisio,

[145209]N.: Band 3 Chur: a variant associated with band
3-deficient hereditary

[145210]spherocytosis and substitution in a highly conserved position of transmembrane

[145211]segment 11. Brit. J. Haemat. 91: 804-810, 1995.

[145212]26055.

[145213]Carrington, M.; Kissner, T.; Gerrard, B.; Ivanov, S.; O'Brien,

[145214]S. J.; Dean, M.: Novel alleles of the chemokine–receptor
gene CCR5. Am.

[145215]J. Hum. Genet. 61: 1261–1267, 1997.

[145216]26056.

[145217]Chaganti, R. S. K.: Personal Communication. New York, N.
Y.

[145218]10/22/1993.

[145219]26057.

[145220]Chaudhuri, A.; Polyakova, J.; Zbrzezna, V .; Pogo, A. O.:
The

[145221]coding sequence of Duffy blood group gene in humans
and simians: restriction

[145222]fragment length polymorphism, antibody and malarial
parasite specificities,

[145223]and expression in nonerythroid tissues in Duffy–negative
individuals. Blood 85:

[145224]615–621, 1995.

[145225]26058.

[145226]Chaudhuri, A.; Polyakova, J.; Zbrzezna, V.; Williams, K.;
Gulati,

[145227]S.; Pogo, A. O.: Cloning of glycoprotein D cDNA, which
encodes the

[145228]major subunit of the Duffy blood group system and the
receptor for

[145229]the Plasmodium vivax malaria parasite. Proc. Nat. Acad.
Sci. 90:

[145230]10793–10797, 1993.

[145231]26059.

[145232]Cook, P. J. L.; Page, B. M.; Johnston, A. W.; Stanford, W. K.;

[145233]Gavin, J.: Four further families informative for 1q and the
Duffy

[145234]blood group. Cytogenet. Cell Genet. 22: 378–380, 1978.

[145235]26060.

[145236]Cook, P. J. L.; Robson, E. B.; Buckton, K. E.; Jacobs, P. A.;
Polani,

[145237]P. E.: Segregation of genetic markers in families with
chromosome

[145238]polymorphisms and structural rearrangements involving
chromosome no.

[145239]1. Ann. Hum. Genet. 37: 261-274, 1974.

[145240]26061.

[145241]Crawford, M. N.; Punnett, H. H.; Carpenter, G. G.: Deletion
of

[145242]the long arm of chromosome 16 and an unexpected Duffy
blood group

[145243]phenotype reveal a possible autosomal linkage. Nature
215: 1075-1076,

[145244]1967.

[145245]26062.

[145246]Gelpi, A. P.; King, M. C.: Association of Duffy blood groups

[145247]with the sickle cell trait. Hum. Genet. 32: 65-68, 1976.

[145248]26063.

[145249]Hadley, T. J.; David, P. H.; McGinniss, M. H.; Miller, L. H.:

[145250]Identification of an erythrocyte component carrying the
Duffy blood

[145251]group Fy-a antigen. Science 223: 597-599, 1984.

[145252]26064.

[145253]Hadley, T. J.; Peiper, S. C.: From malaria to chemokine re-

ceptor:

[145254]the emerging physiologic role of the Duffy blood group
antigen. Blood 89:

[145255]3077–3091, 1997.

[145256]26065.

[145257]Hamblin, M. T.; Di Rienzo, A.: Detection of the signature
of

[145258]natural selection in humans: evidence from the Duffy
blood group locus. Am.

[145259]J. Hum. Genet. 66: 1669–1679, 2000.

[145260]26066.

[145261]Hamblin, M. T.; Thompson, E. E.; Di Rienzo, A.: Complex
signatures

[145262]of natural selection at the Duffy blood group locus. Am. J.
Hum.

[145263]Genet. 70: 369–383, 2002.

[145264]26067.

[145265]Hessner, M. J.; Pircon, R. A.; Johnson, S. T.; Luhm, R. A.:
Prenatal

[145266]genotyping of the Duffy blood group system by allele-
specific polymerase

[145267]chain reaction. Prenatal Diag. 19: 41–45, 1999.

[145268]26068.

[145269]Horuk, R.; Chitnis, C. E.; Darbonne, W. C.; Colby, T. J.; Ry-
bicki,

[145270]A.; Hadley, T. J.; Miller, L. H.: A receptor for the malarial
parasite

[145271]Plasmodium vivax: the erythrocyte chemokine receptor.
Science 261:

[145272]1182–1184, 1993.

[145273]26069.

[145274]Howard, P. N.; Stoddard, G. R.; Goddard, M. W.; Seely, J.
R.:

[145275]Giemsa banding of chromosome 1qh+ and linkage analy-
sis. J. Med. Genet. 12:

[145276]44–48, 1975.

[145277]26070.

[145278]Iwamoto, S.; Li, J.; Sugimoto, N.; Okuda, H.; Kajii, E.: Char-
acterization

[145279]of the Duffy gene promoter: evidence for tissue-specific
abolishment

[145280]of expression in Fy(a–b–) of black individuals. Biochem.

Biophys.

[145281]Res. Commun. 222: 852–859, 1996.

[145282]26071.

[145283]Lautenberger, J. A.; Stephens, J. C.; O'Brien, S. J.; Smith, M.

[145284]W.: Significant admixture linkage disequilibrium across 30
cM around

[145285]the FY locus in African Americans. Am. J. Hum. Genet. 66:
969–978,

[145286]2000.

[145287]26072.

[145288]Lee, C. S. N.; Ying, K. L.; Bowen, P.: Position of the Duffy

[145289]locus on chromosome 1 in relation to breakpoints for
structural rearrangements. Am.

[145290]J. Hum. Genet. 26: 93–102, 1974.

[145291]26073.

[145292]Li, J.; Iwamoto, S.; Sugimoto, N.; Okuda, H.; Kajii, E.: Dinu-
cleotide

[145293]repeat in the 3-prime flanking region provides a clue to
the molecular

[145294]evolution of the Duffy gene. Hum. Genet. 99: 573–577,

1997.

[145295]26074.

[145296]Braun, A.; Kammerer, S.; Bohme, E.; Muller, B.; Roscher, A.
A.

[145297]: Identification of polymorphic sites of the human
bradykinin B(2)

[145298]receptor gene. Biochem. Biophys. Res. Commun. 211:
234-240, 1995.

[145299]26075.

[145300]Erdmann, J.; Hegemann, N.; Weidemann, A.; Kallisch, H.;
Hummel,

[145301]M.; Hetzer, R.; Fleck, E.; Regitz-Zagrosek, V.: Screening
the human

[145302]bradykinin B2 receptor gene in patients with cardiovascular
diseases:

[145303]identification of a functional mutation in the promoter and
a new

[145304]coding variant (T21M). Am. J. Med. Genet. 80: 521-525,
1998.

[145305]26076.

[145306]Hess, J. F.; Borkowski, J. A.; Young, G. S.; Strader, C. D.;
Ransom,

- [145307]R. W.: Cloning and pharmacological characterization of a human bradykinin
- [145308](BK-2) receptor. Biochem. Biophys. Res. Commun. 184: 260-268, 1992.
- [145309]26077.
- [145310]Kammerer, S.; Braun, A.; Arnold, N.; Roscher, A. A.: The human
- [145311]bradykinin B(2) receptor gene: full length cDNA, genomic organization
- [145312]and identification of the regulatory region. Biochem. Biophys. Res.
- [145313]Commun. 211: 226-233, 1995.
- [145314]26078.
- [145315]Ma, J.; Wang, D.; Ward, D. C.; Chen, L.; Dessai, T.; Chao, J.;
- [145316]Chao, L.: Structure and chromosomal localization of the gene (BDKRB2)
- [145317]encoding human bradykinin B2 receptor. Genomics 23: 362-369, 1994.
- [145318]26079.
- [145319]Powell, S. J.; Slynn, G.; Thomas, C.; Hopkins, B.; Briggs, I.;
- [145320]Graham, A.: Human bradykinin B2 receptor: nucleotide

sequence analysis

[145321]and assignment to chromosome 14. Genomics 15:
435–438, 1993.

[145322]26080.

[145323]Taketo, M.; Yokoyama, S.; Rochelle, J.; Kimura, S.; Higashida,

[145324]H.; Taketo, M.; Seldin, M. F.: Mouse B2 bradykinin receptor
gene

[145325]maps to distal chromosome 12. Genomics 27: 222–223,
1995.

[145326]26081.

[145327]Lu, Q.; Sun, E. E.; Klein, R. S.; Flanagan, J. G.: Ephrin-B re-
verse

[145328]signaling is mediated by a novel PDZ–RGS protein and se-
lectively inhibits

[145329]G protein–coupled chemoattraction. Cell 105: 69–79,
2001.

[145330]26082.

[145331]Amara, S. G.; Arriza, J. L.; Leff, S. E.; Swanson, L. W.;
Evans,

[145332]R. M.; Rosenfeld, M. G.: Expression in brain of a messen-
ger RNA encoding

[145333]a novel neuropeptide homologous to calcitonin gene-related peptide.

[145334]Science 229: 1094–1097, 1985.

[145335]26083.

[145336]Hoppener, J. W. M.; Steenbergh, P. H.; Zandberg, J.; Geurts van

[145337]Kessel, A. H. M.; Baylin, S. B.; Nelkin, B. D.; Jansz, H. S.; Lips,

[145338]C. J. M.: The second human calcitonin/CGRP gene is located on chromosome

[145339]11. Hum. Genet. 70: 259–263, 1985.

[145340]26084.

[145341]Steenbergh, P. H.; Hoppener, J. W. M.; Zandberg, J.; Lips, C. J.

[145342]M.; Jansz, H. S.: A second human calcitonin/CGRP gene. FEBS Lett. 183:

[145343]403–407, 1985.

[145344]26085.

[145345]Arthur, J. S. C.; Elce, J. S.; Hegadorn, C.; Williams, K.; Greer,

[145346]P. A.: Disruption of the murine calpain small subunit gene,

Capn4:

[145347]calpain is essential for embryonic development but not for
cell growth

[145348]and division. Molec. Cell. Biol. 20: 4474–4481, 2000.

[145349]26086.

[145350]Ohno, S.; Emori, Y.; Suzuki, K.: Nucleotide sequence of a
cDNA

[145351]coding for the small subunit of human calcium-dependent
protease. Nucleic

[145352]Acids Res. 14: 5559 only, 1986.

[145353]26087.

[145354]Ohno, S.; Minoshima, S.; Kudoh, J.; Fukuyama, R.; Ohmi-
Imajoh,

[145355]S.; Suzuki, K.; Shimizu, Y.; Shimizu, N.: Four genes for the
calpain

[145356]family locate on four distinct human chromosomes.
(Abstract) Cytogenet.

[145357]Cell Genet. 51: 1054–1055, 1989.

[145358]26088.

[145359]Ohno, S.; Minoshima, S.; Kudoh, J.; Fukuyama, R.; Shimizu,
Y.;

[145360] Ohmi-Imajoh, S.; Shimizu, N.; Suzuki, K.: Four genes for the calpain

[145361] family locate on four distinct human chromosomes. Cytogenet. Cell

[145362] Genet. 53: 225-229, 1990.

[145363] 26089.

[145364] Sakihama, T.; Kakidani, H.; Zenita, K.; Yumoto, N.; Kikuchi, T.;

[145365] Sasaki, T.; Kannagi, R.; Nakanishi, S.; Ohmori, M.; Takio, K.; Titani,

[145366] K.; Murachi, T.: A putative Ca(2+)-binding protein: structure of

[145367] the light subunit of porcine calpain elucidated by molecular cloning

[145368] and protein sequence analysis. Proc. Nat. Acad. Sci. 82: 6075-6079,

[145369] 1985.

[145370] 26090.

[145371] Berchtold, M. W.; Egli, R.; Rhyner, J. A.; Hameister, H.; Strehler,

[145372] E. E.: Localization of the human bona fide calmodulin genes CALM1,

[145373]CALM2, and CALM3 to chromosomes 14q24–q31,
2p21.1–p21.3, and 19q13.2–q13.3. Genomics 16:

[145374]461–465, 1993.

[145375]26091.

[145376]Chin, D.; Winkler, K. E.; Means, A. R.: Characterization of
substrate

[145377]phosphorylation and use of calmodulin mutants to ad-
dress implications

[145378]from the enzyme crystal structure of calmodulin–depen-
dent protein

[145379]kinase I. J. Biol. Chem. 272: 31235–31240, 1997.

[145380]26092.

[145381]Drum, C. L.; Yan, S.–Z.; Bard, J.; Shen, Y.–Q.; Lu, D.; Soe-
laiman,

[145382]S.; Grabarek, Z.; Bohm, A.; Tang, W.–J.: Structural basis for
the

[145383]activation of anthrax adenylyl cyclase exotoxin by
calmodulin. Nature 415:

[145384]396–402, 2002.

[145385]26093.

[145386]Koller, M.; Schnyder, B.; Strehler, E. E.: Structural organi-

zation

[145387]of the human CaMIII calmodulin gene. Biochim. Biophys.

Acta 1087:

[145388]180–189, 1990.

[145389]26094.

[145390]Kretsinger, R. H.; Rudnick, S. E.; Weissman, L. J.: Crystal
structure

[145391]of calmodulin. J. Inorganic Biochem. 28: 289–302, 1986.

[145392]26095.

[145393]McPherson, J. D.; Hickie, R. A.; Wasmuth, J. J.; Meyskens,
F. L.;

[145394]Perham, R. N.; Strehler, E. E.; Graham, M. T.: Chromosomal
localization

[145395]of multiple genes encoding calmodulin. (Abstract) Cyto-
genet. Cell

[145396]Genet. 58: 1951 only, 1991.

[145397]26096.

[145398]Pegues, J. C.; Friedberg, F.: Multiple mRNAs encoding hu-
man calmodulin. Biochem.

[145399]Biophys. Res. Commun. 172: 1145–1149, 1990.

[145400]26097.

- [145401]Rhyner, J. A.; Ottiger, M.; Wicki, R.; Greenwood, T. M.;
Strehler,
[145402]E. E.: Structure of the human CALM1 calmodulin gene and
identification
[145403]of two CALM1-related pseudogenes CALM1P1 and
CALM1P2. *Europ. J. Biochem.* 225:
[145404]71–82, 1994.
[145405]26098.
[145406]Scambler, P. J.; McPherson, M. A.; Bates, G.; Bradbury, N.
A.;
[145407]Dormer, R. L.; Williamson, R.: Biochemical and genetic ex-
clusion
[145408]of calmodulin as the site of the basic defect in cystic fi-
brosis. *Hum.*
[145409]*Genet.* 76: 278–282, 1987.
[145410]26099.
[145411]Schumacher, M. A.; Rivard, A. F.; Bachinger, H. P.; Adel-
man, J.
[145412]P.: Structure of the gating domain of a $\text{Ca}(2+)$ -activated
 K^+ channel
[145413]complexed with $\text{Ca}(2+)/$ calmodulin. *Nature* 410:
1120–1124, 2001.

[145414]26100.

[145415]Sen Gupta, B.; Detera-Wadleigh, S. D.; McBride, O. W.;
Friedberg,

[145416]F.: A calmodulin pseudogene on human chromosome 17.
Nucleic Acids

[145417]Res. 17: 2868 only, 1989.

[145418]26101.

[145419]Elbaz, A.; Vale-Santos, J.; Jurkat-Rott, K.; Lapie, P.;
Ophoff,

[145420]R. A.; Bady, B.; Links, T. P.; Piussan, C.; Vila, A.; Monnier,
N.;

[145421]Padberg, G. W.; Abe, K.; Feingold, N.; Guimaraes, J.;
Wintzen, A.

[145422]R.; van der Hoeven, J. H.; Saudubray, J. M.; Grunfeld, J. P.;
Lenoir,

[145423]G.; Nivet, H.; Echenne, B.; Frants, R. R.; Fardeau, M.;
Lehmann-Horn,

[145424]F.; Fontaine, B.: Hypokalemic periodic paralysis and the
dihydropyridine

[145425]receptor (CACNL1A3): genotype/phenotype correlations
for two predominant

[145426]mutations and evidence for the absence of a founder ef-

fect in 16 Caucasian

[145427]families. Am. J. Hum. Genet. 56: 374–380, 1995.

[145428]26102.

[145429]Gluecksohn-Waelsch, S.: Lethal genes and analysis of differentiation. Science 142:

[145430]1269–1276, 1963.

[145431]26103.

[145432]Gregg, R. G.; Couch, F.; Hogan, K.; Powers, P. A.: Assignment

[145433]of the human gene for the alpha-1 subunit of the skeletal muscle DHP-sensitive

[145434]Ca(2+) channel (CACNL1A3) to chromosome 1q31–q32.

Genomics 15: 107–112,

[145435]1993.

[145436]26104.

[145437]Hogan, K.; Gregg, R. G.; Powers, P. A.: The structure of the

[145438]gene encoding the human skeletal muscle alpha-1 subunit of the dihydropyridine-sensitive

[145439]L-type calcium channel (CACNL1A3). Genomics 31: 392–394, 1996.

[145440]26105.

[145441]lles, D. E.; Segers, B.; Weghuis, D. O.; Suijkerbuijk, R.;
Mikala,
[145442]G.; Schwartz, A.; Wieringa, B.: Refined localization of the
alpha-1-subunit
[145443]of the skeletal muscle L-type voltage-dependent calcium
channel (CACNL1A3)
[145444]to human chromosome 1q32 by in situ hybridization. Ge-
nomics 19:
[145445]561-563, 1994.
[145446]26106.
[145447]Jurkat-Rott, K.; Lehmann-Horn, F.; Elbaz, A.; Heine, R.;
Gregg,
[145448]R. G.; Hogan, K.; Powers, P. A.; Lapie, P.; Vale-Santos, J. E.;
Weissenbach,
[145449]J.; Fontaine, B.: A calcium channel mutation causing hy-
pokalemic
[145450]periodic paralysis. Hum. Molec. Genet. 3: 1415-1419,
1994.
[145451]26107.
[145452]Pai, A. C.: Developmental genetics of a lethal mutation,
muscular
[145453]dysgenesis (mdg), in the mouse. I. Genetic analysis and

gross morphology. Dev.

[145454]Biol. 11: 82–92, 1965.

[145455]26108.

[145456]Sillen, A.; Sorensen, T.; Kantola, I.; Friis, M. L.; Gustavson,

[145457]K.–H.; Wadelius, C.: Identification of mutations in the

CACNL1A3

[145458]gene in 13 families of Scandinavian origin having hypokalemic periodic

[145459]paralysis and evidence of a founder effect in Danish families. Am.

[145460]J. Med. Genet. 69: 102–106, 1997.

[145461]26109.

[145462]Stewart, S. L.; Hogan, K.; Rosenberg, H.; Fletcher, J. E.: Identification

[145463]of the arg1086his mutation in the alpha subunit of the voltage-dependent

[145464]calcium channel (CACNA1S) in a North American family with malignant

[145465]hyperthermia. Clin. Genet. 59: 178–184, 2001.

[145466]26110.

[145467]Tanabe, T.; Beam, K. G.; Powell, J. A.; Numa, S.: Restora–

tion

[145468]of excitation-contraction coupling and slow calcium current in dysgenic

[145469]muscle by dihydropyridine receptor complementary DNA.

Nature 336:

[145470]134-139, 1988.

[145471]26111.

[145472]Tanabe, T.; Takeshima, H.; Mikami, A.; Flockerzi, V.;

Takahashi,

[145473]H.; Kangawa, K.; Kojima, M.; Matsuo, H.; Hirose, T.; Numa,

S.: Primary

[145474]structure of the receptor for calcium channel blockers

from skeletal

[145475]muscle. Nature 328: 313-318, 1987.

[145476]26112.

[145477]Ambartsumian, N.; Tarabykina, S.; Grigorian, M.; Tulchin-

sky, E.;

[145478]Hulgaard, E.; Georgiev, G.; Lukanidin, E.: Characterization

of two

[145479]splice variants of metastasis-associated human mts1

gene. Gene 159:

[145480]125-130, 1995.

[145481]26113.

[145482]Jackson-Grusby, L. L.; Swiergiel, J.; Linzer, D. I.: A growth-related

[145483]mRNA in cultured mouse cells encodes a placental calcium binding protein.

[145484]Nucleic Acids Res. 15: 6677-6690, 1987.

[145485]26114.

[145486]Saris, C. J.; Kristensen, T.; D'Eustachio, P.; Hicks, L. J.; Noonan,

[145487]D. J.; Hunter, T.; Tack, B. F.: cDNA sequence and tissue distribution

[145488]of the mRNA for bovine and murine p11, the S100-related light chain

[145489]of the protein-tyrosine kinase substrate p36 (calpactin I). J. Biol.

[145490]Chem. 262: 10663-10671, 1987.

[145491]26115.

[145492]Lefort, A.; Lecocq, R.; Libert, F.; Lamy, F.; Swillens, S.; Vassart,

[145493]G.; Dumont, J. E.: Cloning and sequencing of a calcium-binding protein

[145494]regulated by cyclic AMP in the thyroid. EMBO J. 8:
111–116, 1989.

[145495]26116.

[145496]Loughlin, J.; Irven, C.; Sykes, B.: Exclusion of the cartilage
[145497]link protein and the cartilage matrix protein genes as the
mutant

[145498]loci in several heritable chondrodysplasias. Hum. Genet.
94: 698–700,

[145499]1994.

[145500]26117.

[145501]Deak, F.; Piecha, D.; Bachrati, C.; Paulsson, M.; Kiss, I.: Pri-
mary

[145502]structure and expression of matrilin-2, the closest relative
of cartilage

[145503]matrix protein within the von Willebrand factor type A-like
module

[145504]superfamily. J. Biol. Chem. 272: 9268–9274, 1997.

[145505]26118.

[145506]Deak, F.; Wagener, R.; Kiss, I.; Paulsson, M.: The matrilins:

[145507]a novel family of oligomeric extracellular matrix proteins.
Matrix

[145508]Biol. 18: 55-64, 1999.

[145509]26119.

[145510]Hansson, A.-S.; Heinegard, D.; Holmdahl, R.: A new animal model

[145511]for relapsing polychondritis, induced by cartilage matrix protein

[145512](matrilin-1). J. Clin. Invest. 104: 589-598, 1999.

[145513]26120.

[145514]Jenkins, R. N.; Osborne-Lawrence, S. L.; Sinclair, A. K.; Eddy,

[145515]R. L., Jr.; Byers, M. G.; Shows, T. B.; Duby, A. D.: Structure and

[145516]chromosomal location of the human gene encoding cartilage matrix protein. J.

[145517]Biol. Chem. 265: 19624-19631, 1990.

[145518]26121.

[145519]Lang, B.; Rothenfusser, A.; Lanchbury, J. S.; Rauh, G.; Breedveld,

[145520]F. C.; Urlacher, A.; Albert, E. D.; Peter, H.-H.; Melchers, I.: Susceptibility

[145521]to relapsing polychondritis is associated with HLA-DR4.

Arthritis

[145522]Rheum. 36: 660–664, 1993.

[145523]26122.

[145524]Muratoglu, S.; Krysan, K.; Balazs, M.; Sheng, H.; Zakany, R.; Modis,

[145525]L.; Kiss, I.; Deak, F.: Primary structure of human matrilin-2, chromosome

[145526]location of the MATN2 gene and conservation of an AT-AC intron in

[145527]matrilin genes. Cytogenet. Cell Genet. 90: 323–327, 2000.

[145528]26123.

[145529]Wagener, R.; Kobbe, B.; Paulsson, M.: Primary structure of matrilin-3,

[145530]a new member of a family of extracellular matrix proteins related

[145531]to cartilage matrix protein (matrilin-1) and von Willebrand factor. FEBS

[145532]Lett. 413: 129–134, 1997.

[145533]26124.

[145534]Zeuner, M.; Straub, R. H.; Rauh, G.; Albert, E. D.; Scholmerich,

[145535]J.; Lang, B.: Relapsing polychondritis: clinical and immunogenetic

[145536]analysis of 62 patients. J. Rheum. 24: 96–101, 1997.

[145537]26125.

[145538]Boldyreff, B.; Klett, C.; Gottert, E.; Geurts van Kessel, A.; Hameister,

[145539]H.; Issinger, O.–G.: Assignment of casein kinase 2 alpha sequences

[145540]to two different human chromosomes. Hum. Genet. 89: 79–82, 1992.

[145541]26126.

[145542]Doray, B.; Ghosh, P.; Griffith, J.; Geuze, H. J.; Kornfeld, S.

[145543]: Cooperation of GGAs and AP–1 in packaging MPRs at the trans–Golgi

[145544]network. Science 297: 1700–1703, 2002.

[145545]26127.

[145546]Meisner, H.; Heller–Harrison, R.; Buxton, J.; Czech, M. P.: Molecular

[145547]cloning of the human casein kinase II alpha subunit. Biochemistry 28:

[145548]4072–4076, 1989.

[145549]26128.

[145550]Wirkner, U.; Voss, H.; Lichter, P.; Ansorge, W.; Pyerin, W.:
The

[145551]human gene (CSNK2A1) coding for the casein kinase II
subunit alpha

[145552]is located on chromosome 20 and contains tandemly ar-
ranged Alu repeats. Genomics 19:

[145553]257-265, 1994.

[145554]26129.

[145555]Wirkner, U.; Voss, H.; Lichter, P.; Weitz, S.; Ansorge, W.;
Pyerin,

[145556]W.: Human casein kinase II subunit alpha: sequence of a
processed

[145557](pseudo)gene and its localization on chromosome 11.
Biochim. Biophys.

[145558]Acta 1131: 220-222, 1992.

[145559]26130.

[145560]Yang-Feng, T. L.; Zheng, K.; Kopatz, I.; Naiman, T.;
Canaani, D.

[145561]: Mapping of the human casein kinase II catalytic subunit
genes: two

[145562]loci carrying the homologous sequences for the alpha subunit. Nucleic

[145563]Acids Res. 19: 7125–7129, 1991.

[145564]26131.

[145565]Lozeman, F. J.; Litchfield, D. W.; Piening, C.; Takio, K.; Walsh,

[145566]K. A.; Krebs, E. G.: Isolation and characterization of human cDNA

[145567]clones encoding the alpha and the alpha-prime subunits of casein kinase

[145568]II. Biochemistry 29: 8436–8447, 1990.

[145569]26132.

[145570]Xu, X.; Toselli, P. A.; Russell, L. D.; Seldin, D. C.: Globozoospermia

[145571]in mice lacking the casein kinase II alpha-prime catalytic subunit. Nature

[145572]Genet. 23: 118–121, 1999.

[145573]26133.

[145574]Wickenheisser, J. K.; Quinn, P. G.; Nelson, V. L.; Legro, R. S.;

[145575]Strauss, J. F., III; McAllister, J. M.: Differential activity of

[145576]the cytochrome P450 17-alpha-hydroxylase and steroido-
genic acute regulatory

[145577]protein gene promoters in normal and polycystic ovary
syndrome theca

[145578]cells. J. Clin. Endocr. Metab. 85: 2304-2311, 2000.

[145579]26134.

[145580]Geissler, E. N.; Liao, M.; Brook, J. D.; Martin, F. H.; Zsebo,

[145581]K. M.; Housman, D. E.; Galli, S. J.: Stem cell factor (SCF), a
novel

[145582]hematopoietic growth factor and ligand for c-kit tyrosine
kinase receptor,

[145583]maps on human chromosome 12 between 12q14.3 and
12qter. Somat. Cell

[145584]Molec. Genet. 17: 207-214, 1991.

[145585]26135.

[145586]Heissig, B.; Hattori, K.; Dias, S.; Friedrich, M.; Ferris, B.;

[145587]Hackett, N. R.; Crystal, R. G.; Besmer, P.; Lyden, D.; Moore,
M. A.

[145588]S.; Werb, Z.; Rafii, S.: Recruitment of stem and progenitor
cells

[145589]from the bone marrow niche requires MMP-9 mediated

release of Kit-ligand. Cell 109:

[145590]625-637, 2002.

[145591]26136.

[145592]Martin, F. H.; Suggs, S. V.; Langley, K. E.; Lu, H. S.; Ting, J.;

[145593]Okino, K. H.; Morris, C. F.; McNiece, I. K.; Jacobsen, F. W.;

Mendiaz,

[145594]E. A.; Birkett, N. C.; Smith, K. A.; and 15 others: Primary
structure

[145595]and functional expression of rat and human stem cell fac-
tor DNAs. Cell 63:

[145596]203-211, 1990.

[145597]26137.

[145598]Mathew, S.; Murty, V. V. V. S.; Hunziker, W.; Chaganti, R. S.
K.

[145599]: Subregional mapping of 13 single-copy genes on the
long arm of chromosome

[145600]12 by fluorescence in situ hybridization. Genomics 14:
775-779,

[145601]1992.

[145602]26138.

[145603]Abbott, C.; Piaggio, G.; Ammendola, R.; Solomon, E.;

Povey, S.;

[145604]Gounari, F.; De Simone, V.; Cortese, R.: Mapping of the
gene TCF2

[145605]for the transcription factor LFB3 to human chromosome
17 by polymerase

[145606]chain reaction. Genomics 8: 165–167, 1990.

[145607]26139.

[145608]Bach, I.; Mattei, M.–G.; Cereghini, S.; Yaniv, M.: Two mem-
bers

[145609]of an HNF1 homeoprotein family are expressed in human
liver. Nucleic

[145610]Acids Res. 19: 3553–3559, 1991.

[145611]26140.

[145612]Bingham, C.; Ellard, S.; Allen, L.; Bulman, M.; Shepherd,
M.; Frayling,

[145613]T.; Berry, P. J.; Clark, P. M.; Lindner, T.; Bell, G. I.; Ryffel,

[145614]G. U.; Nicholls, A. J.; Hattersley, A. T.: Abnormal nephron
development

[145615]associated with a frameshift mutation in the transcription
factor

[145616]hepatocyte nuclear factor–1–beta. Kidney Int. 57:

898–907, 2000.

[145617]26141.

[145618]Horikawa, Y.; Iwasaki, N.; Hara, M.; Furuta, H.; Hinokio, Y.;
Cockburn,

[145619]B. N.; Lindner, T.; Yamagata, K.; Ogata, M.; Tomonaga, O.;
Kuroki,

[145620]H.; Kasahara, T.; Iwamoto, Y.; Bell, G. I.: Mutation in hepa-
tocyte

[145621]nuclear factor-1-beta gene (TCF2) associated with MODY.
(Letter) Nature

[145622]Genet. 17: 384–385, 1997.

[145623]26142.

[145624]Menzel, R.; Kaisaki, P. J.; Rjasanowski, I.; Heinke, P.;
Kerner,

[145625]W.; Menzel, S.: A low renal threshold for glucose in dia-
betic patients

[145626]with a mutation in the hepatocyte nuclear factor-1-alpha
(HNF-1-alpha)

[145627]gene. Diabet. Med. 15: 816–820, 1998.

[145628]26143.

[145629]Nishigori, H.; Yamada, S.; Kohama, T.; Tomura, H.; Sho,

K.; Horikawa,
[145630]Y.; Bell, G. I.; Takeuchi, T.; Takeda, J.: Frameshift mutation, A263fsinsGG,
[145631]in the hepatocyte nuclear factor-1-beta gene associated
with diabetes
[145632]and renal dysfunction. Diabetes 47: 1354-1355, 1998.
[145633]26144.

[145634]Wild, W.; Pogge von Strandmann, E.; Nastos, A.; Senkel, S.;
Lingott-Frieg,
[145635]A.; Bulman, M.; Bingham, C.; Ellard, S.; Hattersley, A. T.;
Ryffel,
[145636]G. U.: The mutated human gene encoding hepatocyte nuclear factor
[145637]1-beta inhibits kidney formation in developing *Xenopus*
embryos. Proc.
[145638]Nat. Acad. Sci. 97: 4695-4700, 2000.
[145639]26145.

[145640]Kingsmore, S. F.; Watson, M. L.; Seldin, M. F.: Genetic
mapping
[145641]of the T lymphocyte-specific transcription factor 7 gene
on mouse
[145642]chromosome 11. Mammalian Genome 6: 378 only, 1995.

[145643]26146.

[145644]Roose, J.; Huls, G.; van Beest, M.; Moerer, P.; van der Horn, K.;

[145645]Goldschmeding, R.; Logtenberg, T.; Clevers, H.: Synergy between tumor

[145646]suppressor APC and the beta-catenin-Tcf4 target Tcf1. Science 285:

[145647]1923–1926, 1999.

[145648]26147.

[145649]van de Wetering, M.; Oosterwegel, M.; Holstege, F.; Dooyes, D.;

[145650]Suijkerbuijk, R.; Geurts van Kessel, A.; Clevers, H.: The human T

[145651]cell transcription factor-1 gene: structure, localization, and promoter

[145652]characterization. J. Biol. Chem. 267: 8530–8536, 1992.

[145653]26148.

[145654]van de Wetering, M.; Suijkerbuijk, R.; Geurts van Kessel, A.; Clevers,

[145655]H.: Assignment of the human T lymphocyte-specific transcription factor

[145656]TCF-1 to chromosome 5, band q31.1. (Abstract) Cytogenet. Cell Genet. 58:

[145657]1906 only, 1991.

[145658]26149.

[145659]Verbeek, S.; Izon, D.; Hofhuis, F.; Robanus-Maandag, E.; te Riele,

[145660]H.; van de Wetering, M.; Oosterwegel, M.; Wilson, A.; MacDonald, H.

[145661]R.; Clevers, H.: An HMG-box-containing T-cell factor required for

[145662]thymocyte differentiation. Nature 374: 70-74, 1995.

[145663]26150.

[145664]Aplan, P. D.; Jones, C. A.; Chervinsky, D. S.; Zhao, X.; Ellsworth,

[145665]M.; Wu, C.; McGuire, E. A.; Gross, K. W.: An scl gene product lacking

[145666]the transactivation domain induces bony abnormalities and cooperates

[145667]with LMO1 to generate T-cell malignancies in transgenic mice. EMBO

[145668]J. 16: 2408-2419, 1997.

[145669]26151.

[145670]Aplan, P. D.; Lombardi, D. P.; Ginsberg, A. M.; Cossman, J.;
Bertness,

[145671]V. L.; Kirsch, I. L.: Disruption of the human SCL locus by
'illegitimate'

[145672]V-(D)-J recombinase activity. Science 250: 1426-1429,
1990.

[145673]26152.

[145674]Aplan, P. D.; Lombardi, D. P.; Reaman, G. H.; Sather, H. N.;
Hammond,

[145675]G. D.; Kirsch, I. R.: Involvement of the putative
hematopoietic transcription

[145676]factor SCL in T-cell acute lymphoblastic leukemia. Blood
79: 1327-1333,

[145677]1992.

[145678]26153.

[145679]Brown, L.; Cheng, J.-T.; Chen, Q.; Siciliano, M. J.; Crist, W.;

[145680]Buchanan, G.; Baer, R.: Site-specific recombination of the
tal-1

[145681]gene is a common occurrence in human T-cell leukemia.
EMBO J. 9:

[145682]3343–3351, 1990.

[145683]26154.

[145684]Collazo–Garcia, N.; Scherer, P.; Aplan, P. D.: Cloning and characterization

[145685]of a murine SIL gene. *Genomics* 30: 506–513, 1995.

[145686]26155.

[145687]Izraeli, S.; Lowe, L. A.; Bertness, V. L.; Good, D. J.; Dorward,

[145688]D. W.; Kirsch, I. R.; Kuehn, M. R.: The SIL gene is required for

[145689]mouse embryonic axial development and left–right specification. *Nature* 399:

[145690]691–694, 1999.

[145691]26156.

[145692]Shiratori, A.; Okumura, K.; Nogami, M.; Taguchi, H.; Onozaki, T.;

[145693]Inoue, T.; Ando, T.; Shibata, T.; Izumi, M.; Miyazawa, H.; Hanaoka,

[145694]F.; Murakami, Y.; Eki, T.: Assignment of the 49–kDa (PRIM1) and 58–kDa

[145695](PRIM2A and PRIM2B) subunit genes of the human DNA

primase to chromosome

[145696]bands 1q44 and 6p11.1-p12. Genomics 28: 350-353,
1995. Note: Erratum:

[145697]Genomics 44: 251 only, 1997.

[145698]26157.

[145699]Roy, A. L.; Du, H.; Gregor, P. D.; Novina, C. D.; Martinez,
E.;

[145700]Roeder, R. G.: Cloning of an Inr- and E-box binding pro-
tein, TFII-I,

[145701]that interacts physically and functionally with USF1. EMBO
J. 16:

[145702]7091-7104, 1997.

[145703]26158.

[145704]Steingrimsson, E.; Sawadogo, M.; Gilbert, D. J.; Zervos, A.
S.;

[145705]Brent, R.; Blonar, M. A.; Fisher, D. E.; Copeland, N. G.;
Jenkins,

[145706]N. A.: Murine chromosomal location of five bHLH-Zip
transcription

[145707]factor genes. Genomics 28: 179-183, 1995.

[145708]26159.

[145709]Zhong, G.; Fan, P.; Ji, H.; Dong, F.; Huang, Y.: Identification
of a chlamydial protease-like activity factor responsible
for the
degradation of host transcription factors. J. Exp. Med.
193: 935-942,
2001.
26160.
Aasland, R.; Olsen, L. C.; Spurr, N. K.; Krokan, H. E.; Hel-
land,
D. E.: Chromosomal assignment of human uracil-DNA
glycosylase to
chromosome 12. Genomics 7: 139-141, 1990.
26161.
Dinner, A. R.; Blackburn, G. M.; Karplus, M.: Uracil-DNA
glycosylase
acts by substrate autocatalysis. Nature 413: 752-755,
2001.
26162.
Haug, T.; Skorpen, F.; Kvaloy, K.; Eftedal, I.; Lund, H.;
Krokan,

[145722]H. E.: Human uracil–DNA glycosylase gene: sequence organization,

[145723]methylation pattern, and mapping to chromosome 12q23–q24.1. Genomics 36:

[145724]408–416, 1996.

[145725]26163.

[145726]Haug, T.; Skorpen, F.; Lund, H.; Krokan, H. E.: Structure of the

[145727]gene for human uracil–DNA glycosylase and analysis of the promoter

[145728]function. FEBS Lett. 353: 180–184, 1994.

[145729]26164.

[145730]Meyer–Sieglar, K.; Mauro, D. J.; Seal, G.; Wurzer, J.; deRiel,

[145731]J. K.; Sirover, M. A.: A human nuclear uracil DNA glycosylase is

[145732]the 37–kDa subunit of glyceraldehyde–3–phosphate dehydrogenase. Proc.

[145733]Nat. Acad. Sci. 88: 8460–8464, 1991.

[145734]26165.

[145735]Nilsen, H.; Rosewell, I.; Robins, P.; Skjelbred, C. F.; Andersen,

[145736]S.; Slupphaug, G.; Daly, G.; Krokan, H. E.; Lindahl, T.; Barnes, D.

[145737]E.: Uracil–DNA glycosylase (UNG)–deficient mice reveal a primary

[145738]role of the enzyme during DNA replication. Molec. Cell 5: 1059–1065,

[145739]2000.

[145740]26166.

[145741]Olsen, L. C.; Aasland, R.; Wittwer, C. U.; Krokan, H. E.; Helland,

[145742]D. E.: Molecular cloning of human uracil–DNA glycosylase, a highly

[145743]conserved DNA repair enzyme. EMBO J. 8: 3121–3125, 1989.

[145744]26167.

[145745]Vollberg, T. M.; Siegler, K. M.; Cool, B. L.; Sirover, M. A.:

[145746]Isolation and characterization of the human uracil DNA

glycosylase

[145747]gene. Proc. Nat. Acad. Sci. 86: 8693–8697, 1989.

[145748]26168.

[145749]Hooper, M.; Hardy, K.; Handyside, A.; Hunter, S.; Monk,
M.: HPRT-deficient

[145750](Lesch–Nyhan) mouse embryos derived from germline col-
onization by

[145751]cultured cells. Nature 326: 292–295, 1987.

[145752]26169.

[145753]Kunicki, T. J.; Pidard, D.; Cazenave, J.–P.; Nurden, A. T.;
Caen,

[145754]J. P.: Inheritance of the human platelet alloantigen, PI(A1),
in

[145755]type I Glanzmann's thrombasthenia. J. Clin. Invest. 67:
717–724,

[145756]1981.

[145757]26170.

[145758]Letellier, S. J.; Hunter, J. B.; Aster, R. H.: Probable genetic

[145759]linkage between genes coding for platelet-specific anti-
gens of the

[145760]PI(A) and Bak systems. Am. J. Hemat. 29: 139–143, 1988.

[145761]26171.

[145762]Newman, P. J.; Seligsohn, U.; Lyman, S.; Collier, B. S.: The
molecular

[145763]genetic basis of Glanzmann thrombasthenia in the Iraqi-
Jewish and

[145764]Arab populations in Israel. Proc. Nat. Acad. Sci. 88:
3160-3164,

[145765]1991.

[145766]26172.

[145767]Newman, P. J.; Seligsohn, U.; Lyman, S.; Poncz, M.; Collier,
B.

[145768]S.: The molecular genetic basis of Glanzmann thrombas-
thenia in the

[145769]Iraqi-Jewish and Arab populations in Israel. (Abstract)
Clin. Res. 38:

[145770]467A only, 1990.

[145771]26173.

[145772]Haan, E. A.; Hull, Y. J.; White, S.; Cockington, R.; Charlton,

[145773]P.; Callen, D. F.: Tricho-rhino-phalangeal and branchio-
oto syndromes

[145774]in a family with an inherited rearrangement of chromo-

some 8q. Am.

[145775]J. Med. Genet. 32: 490–494, 1989.

[145776]26174.

[145777]Momeni, P.; Glockner, G.; Schmidt, O.; von Holtum, D.; Albrecht,

[145778]B.; Gillissen-Kaesbach, G.; Hennekam, R.; Meinecke, P.; Zabel, B.;

[145779]Rosenthal, A.; Horsthemke, B.; Ludecke, H.-J.: Mutations in a new

[145780]gene, encoding a zinc-finger protein, cause trichorhino-phalangeal

[145781]syndrome type I. Nature Genet. 24: 71–74, 2000.

[145782]26175.

[145783]Andersson, S.; Russell, D. W.: Structural and biochemical properties

[145784]of cloned and expressed human and rat steroid 5-alpha-reductases. Proc.

[145785]Nat. Acad. Sci. 87: 3640–3644, 1990.

[145786]26176.

[145787]Ellis, J. A.; Stebbing, M.; Harrap, S. B.: Genetic analysis of

[145788]male pattern baldness and the 5-alpha-reductase genes.

J. Invest.

[145789]Derm. 110: 849–853, 1998.

[145790]26177.

[145791]Harris, G.; Azzolina, B.; Baginsky, W.; Cimis, G.; Rasmusson, G.

[145792]H.; Tolman, R. L.; Raetz, C. R. H.; Ellsworth, K.: Identification

[145793]and selective inhibition of an isozyme of steroid
5-alpha-reductase

[145794]in human scalp. Proc. Nat. Acad. Sci. 89: 10787–10791,
1992.

[145795]26178.

[145796]Hsieh, C.-L.; Milatovich, A.; Russell, D.; Francke, U.: Chromosomal

[145797]mapping of human steroid 5 alpha-reductase gene
(SRD5A1) and pseudogene

[145798](SRD5AP1) in human and mouse. (Abstract) Cytogenet.
Cell Genet. 58:

[145799]1897 only, 1991.

[145800]26179.

[145801]Jenkins, E. P.; Andersson, S.; Imperato-McGinley, J.; Wil-

son, J.

[145802]D.; Russell, D. W.: Genetic and pharmacological evidence
for more

[145803]than one human steroid 5- α -reductase. J. Clin. Invest.
89: 293-300,

[145804]1992.

[145805]26180.

[145806]Jenkins, E. P.; Hsieh, C.-L.; Milatovich, A.; Normington, K.;
Berman,

[145807]D. M.; Francke, U.; Russell, D. W.: Characterization and
chromosomal

[145808]mapping of a human steroid 5- α -reductase gene and
pseudogene and

[145809]mapping of the mouse homologue. Genomics 11:
1102-1112, 1991.

[145810]26181.

[145811]Thigpen, A. E.; Silver, R. I.; Guileyardo, J. M.; Casey, M. L.;

[145812]McConnell, J. D.; Russell, D. W.: Tissue distribution and
ontogeny

[145813]of steroid 5- α -reductase isozyme expression. J. Clin.
Invest. 92:

[145814]903-910, 1993.

[145815]26182.

[145816]DeBose-Boyd, R. A.; Brown, M. S.; Li, W.-P.; Nohturfft, A.;
Goldstein,

[145817]J. L.; Espenshade, P. J.: Transport-dependent proteolysis
of SREBP:

[145818]relocation of Site-1 protease from Golgi to ER obviates the
need for

[145819]SREBP transport to Golgi. Cell 99: 703-712, 1999.

[145820]26183.

[145821]Hua, X.; Wu, J.; Goldstein, J. L.; Brown, M. S.; Hobbs, H. H.:

[145822]Structure of the human gene encoding sterol regulatory
element binding

[145823]protein-1 (SREBF1) and localization of SREBF1 and SREBF2
to chromosomes

[145824]17p11.2 and 22q13. Genomics 25: 667-673, 1995.

[145825]26184.

[145826]Osborne, T. F.: CREating a SCAP-less liver keeps SREBPs
pinned

[145827]in the ER membrane and prevents increased lipid synthe-
sis in response

[145828]to low cholesterol and high insulin. Genes Dev. 15:

1873–1878, 2001.

[145829]26185.

[145830]Shimano, H.; Shimomura, I.; Hammer, R. E.; Herz, J.; Goldstein,

[145831]J. L.; Brown, M. S.; Horton, J. D.: Elevated levels of SREBP-2
and

[145832]cholesterol synthesis in livers of mice homozygous for a
targeted

[145833]disruption of the SREBP-1 gene. J. Clin. Invest. 100:
2115–2124,

[145834]1997.

[145835]26186.

[145836]Shimomura, I.; Hammer, R. E.; Richardson, J. A.; Ikemoto,
S.; Bashmakov,

[145837]Y.; Goldstein, J. L.; Brown, M. S.: Insulin resistance and di-
abetes

[145838]mellitus in transgenic mice expressing nuclear SREBP-1c
in adipose

[145839]tissue: model for congenital generalized lipodystrophy.
Genes Dev. 12:

[145840]3182–3194, 1998.

[145841]26187.

[145842]Tobe, K.; Suzuki, R.; Aoyama, M.; Yamauchi, T.; Kamon, J.;
Kubota,
[145843]N.; Terauchi, Y.; Matsui, J.; Akanuma, Y.; Kimura, S.;
Tanaka, J.;
[145844]Abe, M.; Ohsumi, J.; Nagai, R.; Kadowaki, T.: Increased ex-
pression
[145845]of the sterol regulatory element-binding protein-1 gene
in insulin
[145846]receptor substrate-2 -/- mouse liver. J. Biol. Chem. 276:
38337-38340,
[145847]2001.
[145848]26188.
[145849]Wang, X.; Sato, R.; Brown, M. S.; Hua, X.; Goldstein, J. L.:
SREBP-1,
[145850]a membrane-bound transcription factor released by
sterol-regulated
[145851]proteolysis. Cell 77: 53-62, 1994.
[145852]26189.
[145853]Yokoyama, C.; Wang, X.; Briggs, M. R.; Admon, A.; Wu, J.;
Hua,
[145854]X.; Goldstein, J. L.; Brown, M. S.: SREBP-1, a basic-he-
lix-loop-helix-leucine

[145855]zipper protein that controls transcription of the low density lipoprotein

[145856]receptor gene. Cell 75: 187–197, 1993.

[145857]26190.

[145858]Achermann, J. C.; Ito, M.; Ito, M.; Hindmarsh, P. C.; Jameson,

[145859]J. L.: A mutation in the gene encoding steroidogenic factor-1 causes

[145860]XY sex reversal and adrenal failure in humans. (Letter) Nature Genet. 22:

[145861]125–126, 1999.

[145862]26191.

[145863]Biaison-Lauber, A.; Schoenle, E. J.: Apparently normal ovarian

[145864]differentiation in a prepubertal girl with transcriptionally inactive

[145865]steroidogenic factor 1 (NR5A1/SF-1) and adrenocortical insufficiency. Am.

[145866]J. Hum. Genet. 67: 1563–1568, 2000.

[145867]26192.

[145868]Bland, M. L.; Jamieson, C. A. M.; Akana, S. F.; Bornstein, S.

R.;

[145869]Eisenhofer, G.; Dallman, M. F.; Ingraham, H. A.: Haploinsufficiency
of steroidogenic factor-1 in mice disrupts adrenal development leading
to an impaired stress response. Proc. Nat. Acad. Sci. 97:
14488-14493,
[145872]2000.

[145873]26193.

[145874]Joosten, P. H. L. J.; Toepoel, M.; Mariman, E. C. M.; Van
Zoelen,

[145875]E. J. J.: Promoter haplotype combinations of the platelet-derived

[145876]growth factor alpha-receptor gene predispose to human
neural tube

[145877]defects. Nature Genet. 27: 215-217, 2001.

[145878]26194.

[145879]Mackenzie, P. I.; Owens, I. S.; Burchell, B.; Bock, K. W.;
Bairoch,

[145880]A.; Belanger, A.; Fournel-Gigleux, S.; Green, M.; Hum, D.
W.; Iyanagi,

[145881]T.; Lancet, D.; Louisot, P.; Magalou, J.; Roy Chowdhury, J.;

Ritter,

[145882]J. K.; Schachter, H.; Tephly, T. R.; Tipton, K. E.; Nebert, D. W.

[145883]: The UDP glycosyltransferase gene superfamily: recommended nomenclature

[145884]update based on evolutionary divergence. Pharmacogenetics 7: 255–269,

[145885]1997.

[145886]26195.

[145887]Corydon, M. J.; Vockley, J.; Rinaldo, P.; Rhead, W. J.; Kjeldsen,

[145888]M.; Winter, V.; Riggs, C.; Babovic-Vuksanovic, D.; Smeitink, J.; De

[145889]Jong, J.; Levy, H.; Sewell, A. C.; Roe, C.; Matern, D.; Dsouki, M.;

[145890]Gregersen, N.: Role of common gene variations in the molecular pathogenesis

[145891]of short-chain acyl-CoA dehydrogenase deficiency. Pediat. Res. 49:

[145892]18–23, 2001.

[145893]26196.

[145894]Colombo, R.: Age and origin of the PRNP E200K mutation

causing

[145895]familial Creutzfeldt–Jacob disease in Libyan Jews. (Letter)
Am. J.

[145896]Hum. Genet. 67: 528–531, 2000.

[145897]26197.

[145898]Cousens, J.; Smith, P. G.; Ward, H.; Everington, D.; Knight,
R.

[145899]S. G.; Zeidler, M.; Stewart, G.; Smith–Bathgate, E. A. B.;
Macleod,

[145900]M.–A.; Mackenzie, J.; Will, R. G.: Geographical distribution
of variant

[145901]Creutzfeldt–Jakob disease in Great Britain, 1994–2000.
Lancet 357:

[145902]1002–1007, 2001.

[145903]26198.

[145904]de Silva, R.; Ironside, J. W.; McCardle, L.; Esmonde, T.; Bell,

[145905]J.; Will, R.; Windl, O.; Dempster, M.; Estibeiro, P.; Lathe, R.:

[145906]Neuropathological phenotype and 'prion protein' genotype
correlation

[145907]in sporadic Creutzfeldt–Jakob disease. Neurosci. Lett. 179:
50–52,

[145908]1994.

[145909]26199.

[145910]Deslys, J.-P.; Jaegly, A.; d'Aignaux, J. H.; Mouthon, F.; de
Villemeur,

[145911]T. B.; Dormont, D.: Genotype at codon 129 and susceptibility to Creutzfeldt-Jakob

[145912]disease. Lancet 351: 1251 only, 1998.

[145913]26200.

[145914]Dlouhy, S. R.; Hsiao, K.; Farlow, M. F.; Foroud, T.; Con-
neally,

[145915]P. M.; Johnson, P.; Prusiner, S. B.; Hodes, M. E.; Ghetti, B.:
Linkage

[145916]of the Indiana kindred of Gerstmann-Straussler-Scheinker
disease to

[145917]the prion protein gene. Nature Genet. 1: 64-67, 1992.

[145918]26201.

[145919]Doh-ura, K.; Kitamoto, T.; Sakaki, Y.; Tateishi, J.: CJD dis-
crepancy.

[145920](Letter) Nature 353: 801-802, 1991.

[145921]26202.

[145922]Doh-ura, K.; Tateishi, J.; Sasaki, H.; Kitamoto, T.; Sakaki, Y.

[145923]Pro-to-leu change at position 102 of prion protein is the most common

[145924]but not the sole mutation related to Gerstmann-Straussler syndrome. Biochem.

[145925]Biophys. Res. Commun. 163: 974-979, 1989.

[145926]26203.

[145927]Duchen, L. W.; Poulter, M.; Harding, A. E.: Dementia associated

[145928]with a 216 base pair insertion in the prion protein gene: clinical

[145929]and neuropathological features. Brain 116: 555-567, 1993.

[145930]26204.

[145931]Erginel-Unaltuna, N.; Peoc'h, K.; Komurcu, E.; Acuner, T. T.;

[145932]Issever, H.; Laplanche, J.-L.: Distribution of the M129V polymorphism

[145933]of the prion protein gene in a Turkish population suggests a high

[145934]risk for Creutzfeldt-Jacob disease. Europ. J. Hum. Genet.

9: 965–968,

[145935]2001.

[145936]26205.

[145937]Farlow, M. R.; Yee, R. D.; Dlouhy, S. R.; Conneally, P. M.;
Azzarelli,

[145938]B.; Ghetti, B.: Gerstmann–Straussler–Scheinker disease. I.
Extending

[145939]the clinical spectrum. *Neurology* 39: 1446–1452, 1989.

[145940]26206.

[145941]Flechsig, E.; Shmerling, D.; Hegyi, I.; Raeber, A. J.; Fischer,

[145942]M.; Cozzio, A.; von Mering, C.; Aguzzi, A.; Weissmann, C.:
Prion

[145943]protein devoid of the octapeptide repeat region restores
susceptibility

[145944]to scrapie in PrP knockout mice. *Neuron* 27: 399–408,
2000.

[145945]26207.

[145946]Forloni, G.; Angeretti, N.; Chiesa, R.; Monzani, E.;
Salmona,

[145947]M.; Bugiani, O.: Neurotoxicity of a prion protein fragment.
(Letter) *Nature* 362:

[145948]543–546, 1993.

[145949]26208.

[145950]Gabizon, R.; Rosenmann, H.; Meiner, Z.; Kahana, I.; Kahana, E.;

[145951]Shugart, Y.; Ott, J.; Prusiner, S. B.: Mutation and polymorphism

[145952]of the prion protein gene in Libyan Jews with Creutzfeldt–Jakob disease

[145953](CJD). Am. J. Hum. Genet. 53: 828–835, 1993.

[145954]26209.

[145955]Gajdusek, D. C.: The transmissible amyloidoses: genetical control

[145956]of spontaneous generation of infectious amyloid proteins by nucleation

[145957]of configurational change in host precursors: kuru–CJD–GSS–scrapie–BSE. Europ.

[145958]J. Epidemiol. 7: 567–577, 1991.

[145959]26210.

[145960]Gambetti, P.; Petersen, R.; Monari, L.; Tabaton, M.; Autilio–Gambetti,

[145961]L.: Fatal familial insomnia and the widening spectrum of

prion diseases. Brit.

[145962]Med. Bull. 49: 980–994, 1993.

[145963]26211.

[145964]Ghetti, B.; Tagliavini, F.; Masters, C. L.; Beyreuther, K.; Gi-
accione,

[145965]G.; Verga, L.; Farlow, M. R.; Conneally, P. M.; Dlouhy, S. R.;
Azzarelli,

[145966]B.; Bugiani, O.: Gerstmann–Straussler–Scheinker disease.
II. Neurofibrillary

[145967]tangles and plaques with PrP-amyloid coexist in an af-
fected family. Neurology 39:

[145968]1453–1461, 1989.

[145969]26212.

[145970]Giaccone, G.; Verga, L.; Bugiani, O.; Frangione, B.; Serban,
D.;

[145971]Prusiner, S. B.; Farlow, M. R.; Ghetti, B.; Tagliavini, F.: Prion

[145972]protein preamyloid and amyloid deposits in Gerstmann–
Straussler–Scheinker

[145973]disease, Indiana kindred. Proc. Nat. Acad. Sci. 89:
9349–9353, 1992.

[145974]26213.

[145975]Goldfarb, L. G.; Brown, P.; Cervenakova, L.; Gajdusek, D. C.:

[145976]Molecular genetic studies of Creutzfeldt-Jakob disease.
Molec. Neurobiol. 8:

[145977]89-97, 1994.

[145978]26214.

[145979]Goldfarb, L. G.; Brown, P.; Haltia, M.; Cathala, F.; McCombie,

[145980]W. R.; Kovanen, J.; Cervenakova, L.; Goldin, L.; Nieto, A.;
Godec,

[145981]M. S.; Asher, D. M.; Gajdusek, D. C.: Creutzfeldt-Jakob
disease cosegregates

[145982]with the codon 178-asn PRNP mutation in families of European origin. Ann.

[145983]Neurol. 31: 274-281, 1992.

[145984]26215.

[145985]Goldfarb, L. G.; Brown, P.; McCombie, W. R.; Goldgaber,
D.; Swergold,

[145986]G. D.; Wills, P. R.; Cervenakova, L.; Baron, H.; Gibbs, C. J.,
Jr.;

[145987]Gajdusek, D. C.: Transmissible familial Creutzfeldt-Jakob
disease

- [145988]associated with five, seven, and eight extra octapeptide coding repeats
- [145989]in the PRNP gene. Proc. Nat. Acad. Sci. 88: 10926–10930, 1991.
- [145990]26216.
- [145991]Goldfarb, L. G.; Brown, P.; Mitrova, E.; Cervenakova, L.; Goldin,
- [145992]L.; Korczyn, A. D.; Chapman, J.; Galvez, S.; Cartier, L.; Rubenstein,
- [145993]R.; Gajdusek, D. C.: Creutzfeldt–Jacob disease associated with the
- [145994]PRNP codon 200–lys mutation: an analysis of 45 families. Europ. J.
- [145995]Epidemiol. 7: 477–486, 1991.
- [145996]26217.
- [145997]Goldfarb, L. G.; Haltia, M.; Brown, P.; Nieto, A.; Kovanen, J.;
- [145998]McCombie, W. R.; Trapp, S.; Gajdusek, D. C.: New mutation in scrapie
- [145999]amyloid precursor gene (at codon 178) in Finnish Creutzfeldt–Jakob
- [146000]kindred. (Letter) Lancet 337: 425, 1991.

[146001]26218.

[146002]Goldfarb, L. G.; Mitrova, E.; Brown, P.; Toh, B. H.; Gaj-
dusek,

[146003]D. C.: Mutation in codon 200 of scrapie amyloid protein
gene in two

[146004]clusters of Creutzfeldt-Jakob disease in Slovakia. (Letter)
Lancet 336:

[146005]514-515, 1990.

[146006]26219.

[146007]Travis, G. H.; Sutcliffe, J. G.; Bok, D.: The retinal degenera-
tion

[146008]slow (rds) gene product is a photoreceptor disc mem-
brane-associated

[146009]glycoprotein. Neuron 6: 61-70, 1991.

[146010]26220.

[146011]Morris, C.; Heisterkamp, N.; Hao, Q. L.; Testa, J. R.; Grof-
fen,

[146012]J.: The human tyrosine kinase gene (FER) maps to chro-
mosome 5 and

[146013]is deleted in myeloid leukemias with a del(5q). Cytogenet.
Cell Genet. 53:

[146014]196–200, 1990.

[146015]26221.

[146016]Anderson, J.; Burns, H. D.; Enriquez–Harris, P.; Wilkie, A.
O.

[146017]M.; Heath, J. K.: Apert syndrome mutations in fibroblast
growth factor

[146018]receptor 2 exhibit increased affinity for FGF ligand. Hum.
Molec.

[146019]Genet. 7: 1475–1483, 1998.

[146020]26222.

[146021]Arman, E.; Haffner–Krausz, R.; Chen, Y.; Heath, J. K.;
Lonai, P.

[146022]: Targeted disruption of fibroblast growth factor (FGF) re-
ceptor 2

[146023]suggests a role for FGF signaling in pregastrulation mam-
malian development. Proc.

[146024]Nat. Acad. Sci. 95: 5082–5087, 1998.

[146025]26223.

[146026]Arman, E.; Haffner–Krausz, R.; Gorivodsky, M.; Lonai, P.:
Fgfr2

[146027]is required for limb outgrowth and lung–branching mor-

phogenesis. Proc.

[146028]Nat. Acad. Sci. 96: 11895–11899, 1999.

[146029]26224.

[146030]Chitayat, D.; Chun, K.: Reply to the letter to the editor by
Gripp

[146031]et al.--'not Antley–Bixler syndrome'. (Letter) Am. J. Med.
Genet. 83:

[146032]67–68, 1999.

[146033]26225.

[146034]Chun, K.; Siegel–Bartelt, J.; Chitayat, D.; Phillips, J.; Ray,

[146035]P. N.: FGFR2 mutation associated with clinical manifesta-
tions consistent

[146036]with Antley–Bixler syndrome. Am. J. Med. Genet. 77:
219–224, 1998.

[146037]26226.

[146038]Cohen, M. M., Jr.: Pfeiffer syndrome update, clinical sub-
types,

[146039]and guidelines for differential diagnosis. Am. J. Med.
Genet. 45:

[146040]300–307, 1993.

[146041]26227.

[146042]Cornejo-Roldan, L. R.; Roessler, E.; Muenke, M.: Analysis of the

[146043]mutational spectrum of the FGFR2 gene in Pfeiffer syndrome. Hum.

[146044]Genet. 104: 425-431, 1999.

[146045]26228.

[146046]den Dunnen, J. T.; Antonarakis, E.: Nomenclature for the description

[146047]of human sequence variations. Hum. Genet. 109: 121-124, 2001.

[146048]26229.

[146049]Deng, C.; Bedford, M.; Li, C.; Xu, X.; Yang, X.; Dunmore, J.;

[146050]Leder, P.: Fibroblast growth factor receptor-1 (FGFR-1) is essential

[146051]for normal neural tube and limb development. Dev. Biol. 185: 42-54,

[146052]1997.

[146053]26230.

[146054]Dionne, C. A.; Crumley, G.; Bellot, F.; Kaplow, J. M.; Searfoss,

[146055]G.; Ruta, M.; Burgess, W. H.; Jaye, M.; Schlessinger, J.:

Cloning

- [146056]and expression of two distinct high-affinity receptors cross-reacting
- [146057]with acidic and basic fibroblast growth factors. EMBO J. 9: 2685-2692,
- [146058]1990.
- [146059]26231.
- [146060]Dionne, C. A.; Modi, W. S.; Crumley, G.; O'Brien, S. J.; Schlessinger,
- [146061]J.; Jaye, M.: BEK, a receptor for multiple members of the fibroblast
- [146062]growth factor (FGF) family, maps to human chromosome 10q25.3-q26. Cytogenet.
- [146063]Cell Genet. 60: 34-36, 1992.
- [146064]26232.
- [146065]Gilbert, E.; Del Gatto, F.; Champion-Arnaud, P.; Gesnel, M.-C.;
- [146066]Breathnach, R.: Control of BEK and K-SAM splice sites in alternative
- [146067]splicing of the fibroblast growth factor receptor 2 pre-mRNA. Molec.
- [146068]Cell. Biol. 13: 5461-5468, 1993.

[146069]26233.

[146070]Gorlin, R. J.: Patient described by Chun et al. may not
present

[146071]Antley-Bixler syndrome. (Letter) Am. J. Med. Genet. 83: 64
only,

[146072]1999.

[146073]26234.

[146074]Gorry, M. C.; Preston, R. A.; White, G. J.; Zhang, Y.; Sing-
hal,

[146075]V. K.; Losken, H. W.; Parker, M. G.; Nwokoro, N. A.; Post, J.
C.;

[146076]Ehrlich, G. D.: Crouzon syndrome: mutations in two
spliceoforms of

[146077]FGFR2 and a common point mutation shared with Jack-
son-Weiss syndrome. Hum.

[146078]Molec. Genet. 4: 1387-1390, 1995.

[146079]26235.

[146080]Gripp, K. W.; Stolle, C. A.; McDonald-McGinn, D. M.;
Markowitz,

[146081]R. I.; Bartlett, S. P.; Katowitz, J. A.; Muenke, M.; Zackai, E.
H.

- [146082]: Phenotype of the fibroblast growth factor receptor 2
ser351-to-cys
- [146083]mutation: Pfeiffer syndrome type III. Am. J. Med. Genet.
78: 356-360,
- [146084]1998.
- [146085]26236.
- [146086]Gripp, K. W.; Zackai, E. H.; Cohen, M. M., Jr.: Not Antley-
Bixler
- [146087]syndrome. (Letter) Am. J. Med. Genet. 83: 65-66, 1999.
- [146088]26237.
- [146089]Hajihosseini, M. K.; Wilson, S.; De Moerlooze, L.; Dickson,
C.
- [146090]: A splicing switch and gain-of-function mutation in
Fgfr2-IIIc hemizygotes
- [146091]causes Apert/Pfeiffer-syndrome-like phenotypes. Proc.
Nat. Acad.
- [146092]Sci. 98: 3855-3860, 2001.
- [146093]26238.
- [146094]Hollway, G. E.; Suthers, G. K.; Haan, E. A.; Thompson, E.;
David,
- [146095]D. J.; Gecz, J.; Mulley, J. C.: Mutation detection in FGFR2

craniosynostosis

[146096]syndromes. Hum. Genet. 99: 251–255, 1997.

[146097]26239.

[146098]Houssaint, E.; Blanquet, P. R.; Champion–Arnaud, P.; Gesnel, M.

[146099]C.; Torriglia, A.; Courtois, Y.; Breathnach, R.: Related fibroblast

[146100]growth factor receptor genes exist in the human genome. Proc. Nat.

[146101]Acad. Sci. 87: 8180–8184, 1990.

[146102]26240.

[146103]Ibrahimi, O. A.; Eliseenkova, A. V.; Plotnikov, A. N.; Yu, K.;

[146104]Ornitz, D. M.; Mohammadi, M.: Structural basis for fibroblast growth

[146105]factor receptor 2 activation in Apert syndrome. Proc. Nat. Acad.

[146106]Sci. 98: 7182–7187, 2001.

[146107]26241.

[146108]Jackson, R.: The lines of Blaschko: a review and reconsideration. Brit.

[146109]J. Derm. 95: 349–360, 1976.

[146110]26242.

[146111]Johnson, D.; Wall, S. A.; Mann, S.; Wilkie, A. O. M.: A novel
[146112]mutation, ala315 to ser, in FGFR2: a gene-environment
interaction
[146113]leading to craniosynostosis? Europ. J. Hum. Genet. 8:
571-577, 2000.

[146114]26243.

[146115]Williams, T. M.; Montoya, G.; Wu, Y.; Eddy, R. L.; Byers, M.
G.;

[146116]Shows, T. B.: The TCF8 gene encoding a zinc finger pro-
tein (Nil-2-a)
[146117]resides on human chromosome 10p11.2. Genomics 14:
194-196, 1992.

[146118]26244.

[146119]van Nie, R.; Ivanyi, D.; Demant, P.: A new H-2-linked mu-
tation,

[146120]rds, causing retinal degeneration in the mouse. Tissue
Antigens 12:

[146121]106-108, 1978.

[146122]26245.

[146123]Vine, A. K.; Schatz, H.: Adult-onset foveomacular pigment

epithelial

[146124] dystrophy. Am. J. Ophthal. 89: 680–691, 1980.

[146125] 26246.

[146126] Weleber, R. G.; Carr, R. E.; Murphey, W. H.; Sheffield, V. C.;

[146127] Stone, E. M.: Phenotypic variation including retinitis pigmentosa,

[146128] pattern dystrophy, and fundus flavimaculatus in a single family with

[146129] a deletion of codon 153 or 154 of the peripherin/RDS gene. Arch.

[146130] Ophthal. 111: 1531–1542, 1993.

[146131] 26247.

[146132] Wells, J.; Wroblewski, J.; Keen, J.; Inglehearn, C.; Jubb, C.;

[146133] Eckstein, A.; Jay, M.; Arden, G.; Bhattacharya, S.; Fitzke, F.; Bird,

[146134] A.: Mutations in the human retinal degeneration slow (RDS) gene can

[146135] cause either retinitis pigmentosa or macular dystrophy.

Nature Genet. 3:

[146136] 213–218, 1993.

[146137] 26248.

- [146138]Wroblewski, J. J.; Wells, J. A., III; Eckstein, A.; Fitzke, F.;
- [146139]Jubb, C.; Keen, J.; Inglehearn, C.; Bhattacharya, S.; Arden,
G. B.;
- [146140]Jay, M.; Bird, A. C.: Macular dystrophy associated with mu-
tations
at codon 172 in the human retinal degeneration slow
gene. *Ophthalmology* 101:
12–22, 1994.
- [146142]26249.
- [146143]26249.
- [146144]Eph Nomenclature Committee: Unified nomenclature for
the Eph
family receptors and their ligands, the ephrins. *Cell* 90:
403–404,
1997.
- [146146]26250.
- [146147]26250.
- [146148]Hattori, M.; Osterfield, M.; Flanagan, J. G.: Regulated
cleavage
of a contact-mediated axon repellent. *Science* 289:
1360–1365, 2000.
- [146150]26251.
- [146151]Maru, Y.; Hirai, H.; Yoshida, M. C.; Takaku, F.: Evolution,

expression,

[146152]and chromosomal location of a novel receptor tyrosine kinase gene,

[146153]eph. Molec. Cell. Biol. 8: 3770–3776, 1988.

[146154]26252.

[146155]Yoshida, M. C.; Maru, H.; Hirai, H.; Takau, F.: Chromosomal location

[146156]of a novel receptor tyrosine kinase gene, EPH, on chromosome 7. (Abstract) Cytogenet.

[146157]Cell Genet. 51: 1113 only, 1989.

[146158]26253.

[146159]Boyd, A. W.; Ward, L. D.; Wicks, I. P.; Simpson, R. J.; Salvaris,

[146160]E.; Wilks, A.; Welch, K.; Loudovaris, M.; Rockman, S.; Busmanis, I.

[146161]: Isolation and characterization of a novel receptor-type protein

[146162]tyrosine kinase (hek) from a human pre-B cell line. J. Biol. Chem. 267:

[146163]3262–3267, 1992.

[146164]26254.

[146165]Fox, G. M.; Holst, P. L.; Chute, H. T.; Lindberg, R. A.; Janssen, A. M.; Basu, R.; Welcher, A. A.: cDNA cloning and tissue distribution of five human EPH-like receptor protein-tyrosine kinases. *Oncogene* 10: 897-905, 1995.

[146168]26255.

[146169]26255.

[146170]Wicks, I. P.; Lapsys, N. M.; Baker, E.; Campbell, L. J.; Boyd, A. W.; Sutherland, G. R.: Localization of a human receptor tyrosine kinase (ETK1) to chromosome region 3p11.2. *Genomics* 19: 38-41, 1994.

[146172]26256.

[146173]26256.

[146174]Wicks, I. P.; Wilkinson, D.; Salvaris, E.; Boyd, A. W.: Molecular cloning of HEK, the gene encoding a receptor tyrosine kinase expressed by human lymphoid tumor cell lines. *Proc. Nat. Acad. Sci.* 89: 1611-1615, 1992.

[146177]1992.

[146178]26257.

[146179]Agrawal, A.; Eastman, Q. M.; Schatz, D. G.: Transposition mediated

[146180]by RAG1 and RAG2 and its implications for the evolution of the immune

[146181]system. Nature 394: 744–751, 1998.

[146182]26258.

[146183]Blanquet, V.; Turleau, C.; Goossens, M.; Besmond, C.: Assignment

[146184]of the human recombination activating gene 1 (RAG1) to the 14q21.3–q22.2

[146185]region. Genomics 13: 488–489, 1992.

[146186]26259.

[146187]Fulop, G. M.; Phillips, R. A.: The scid mutation in mice causes

[146188]a general defect in DNA repair. Nature 347: 479–482, 1990.

[146189]26260.

[146190]Hikida, M.; Mori, M.; Takai, T.; Tomochika, K.; Hamatani, K.; Ohmori,

[146191]H.: Reexpression of RAG–1 and RAG–2 genes in activated

mature mouse

[146192]B cells. Science 274: 2091–2093, 1996.

[146193]26261.

[146194]Hiom, K.; Melek, M.; Gellert, M.: DNA transposition by the
RAG1

[146195]and RAG2 proteins: a possible source of oncogenic
translocations. Cell 94:

[146196]463–470, 1998.

[146197]26262.

[146198]Huye, L. E.; Purugganan, M. M.; Jiang, M.-M.; Roth, D. B.:
Mutational

[146199]analysis of all conserved basic amino acids in RAG–1 re-
veals catalytic,

[146200]step arrest, and joining-deficient mutants in the V(D)J re-
combinase. Molec.

[146201]Cell. Biol. 22: 3460–3473, 2002.

[146202]26263.

[146203]Ichihara, Y.; Hirai, M.; Kurosawa, Y.: Sequence and chro-
mosome

[146204]assignment to 11p13–p12 of human RAG genes. Immun.
Lett. 33: 277–284,

[146205]1992.

[146206]26264.

[146207]Janeway, C. A., Jr.: How the immune system works to protect the

[146208]host from infection: a personal view. Proc. Nat. Acad. Sci. 98:

[146209]7461–7468, 2001.

[146210]26265.

[146211]Kim, D. R.; Dai, Y.; Mundy, C. L.; Yang, W.; Oettinger, M. A.:

[146212]Mutations of acidic residues in RAG1 define the active site of the

[146213]V(D)J recombinase. Genes Dev. 13: 3070–3080, 1999.

[146214]26266.

[146215]Landree, M. A.; Wibbenmeyer, J. A.; Roth, D. B.: Mutational analysis

[146216]of RAG1 and RAG2 identifies three catalytic amino acids in RAG1 critical

[146217]for both cleavage steps of V(D)J recombination. Genes Dev. 13: 3059–3069,

[146218]1999.

[146219]26267.

[146220]McBlane, J. F.; van Gent, D. C.; Ramsden, D. A.; Romeo, C.;
Cuomo,

[146221]C. A.; Gellert, M.; Oettinger, M. A.: Cleavage at a V(D)J re-
combination

[146222]signal requires only RAG1 and RAG2 proteins and occurs
in two steps. Cell 83:

[146223]387–395, 1995.

[146224]26268.

[146225]Melek, M.; Gellert, M.: RAG1/2-mediated resolution of
transposition

[146226]intermediates: two pathways and possible consequences.
Cell 101:

[146227]625–633, 2000.

[146228]26269.

[146229]Mombaerts, P.; Iacomini, J.; Johnson, R. S.; Herrup, K.;
Tonegawa,

[146230]S.; Papaioannou, V. E.: RAG-1-deficient mice have no ma-
ture B and

[146231]T lymphocytes. Cell 68: 869–877, 1992.

[146232]26270.

[146233]Oettinger, M. A.: Personal Communication. Boston, Mass.
1/15/1993.

[146234]26271.

[146235]Sillekens, P. T. G.; Habets, W. J.; Beijer, R. P.; van Venrooij,

[146236]W. J.: cDNA cloning of the human U1 snRNA-associated A
protein: extensive

[146237]homology between U1 and U2 snRNP-specific proteins.
EMBO J. 6: 3841-3848,

[146238]1987.

[146239]26272.

[146240]Koyama, K.; Fukushima, Y.; Inazawa, J.; Tomotsune, D.;
Takahashi,

[146241]N.; Nakamura, Y.: The human homologue of the murine
Llg1h gene (LLGL)

[146242]maps within the Smith-Magenis syndrome region in
17p11.2. Cytogenet.

[146243]Cell Genet. 72: 78-82, 1996.

[146244]26273.

[146245]Forsberg, E.; Hirsch, E.; Frohlich, L.; Meyer, M.; Ekblom, P.;

[146246]Aszodi, A.; Werner, S.; Fassler, R.: Skin wounds and sev-
ered nerves

[146247] heal normally in mice lacking tenascin-C. Proc. Nat. Acad. Sci. 93:

[146248] 6594-6599, 1996.

[146249] 26274.

[146250] Gulcher, J. R.; Alexakos, M. J.; Le Beau, M. M.; Lemons, R. S.;

[146251] Stefansson, K.: Chromosomal localization of the human hexabrachion

[146252] (tenascin) gene and evidence for recent reduplication within the gene. Genomics 6:

[146253] 616-622, 1990.

[146254] 26275.

[146255] Gulcher, J. R.; Nies, D. E.; Alexakos, M. J.; Ravikant, N. A.;

[146256] Sturgill, M. E.; Marton, L. S.; Stefansson, K.: Structure of the

[146257] human hexabrachion (tenascin) gene. Proc. Nat. Acad. Sci. 88: 9438-9442,

[146258] 1991.

[146259] 26276.

[146260] Mitrovic, N.; Schachner, M.: Detection of tenascin-C in the nervous

[146261]system of the tenascin-C mutant mouse. J. Neurosci. Res.
42: 710-717,

[146262]1995.

[146263]26277.

[146264]Nies, D. E.; Hemesath, T. J.; Kim, J.-H.; Gulcher, J. R.; Ste-
fansson,

[146265]K.: The complete cDNA sequence of human hexabrachion
(tenascin):

[146266]a multidomain protein containing unique epidermal
growth factor repeats. J.

[146267]Biol. Chem. 266: 2818-2823, 1991.

[146268]26278.

[146269]Williams, T. M.; Moolten, D.; Burlein, J.; Romano, J.; Bhaer-
man,

[146270]R.; Godillot, A.; Mellon, M.; Rauscher, F. J., III; Kant, J. A.:

[146271]Identification of a zinc finger protein that inhibits IL-2
gene expression. Science 254:

[146272]1791-1794, 1991.

[146273]26279.

[146274]Ludecke, H.-J.; Schaper, J.; Meinecke, P.; Momeni, P.;
Gross, S.;

- [146275] von Holtum, D.; Hirche, H.; Abramowicz, M. J.; Albrecht, B.; Apacik, C.; Christen, H.-J.; Claussen, U.; and 28 others: Genotypic and phenotypic spectrum in tricho-rhino-phalangeal syndrome types I and III. *Am. J. Hum. Genet.* 68: 81-91, 2001.
- [146279] 26280.
- [146280] Arya, R.; Lalloz, M. R. A.; Bellingham, A. J.; Layton, D. M.: Evidence for founder effect of the glu104-to-asp substitution and identification of new mutations in triosephosphate isomerase deficiency. *Hum. Mutat.* 10: 290-294, 1997.
- [146284] 26281.
- [146285] Asakawa, J.; Iida, S.: Origin of human triosephosphate isomerase isozymes: further evidence for the single structural locus hypothesis with Japanese variants. *Hum. Genet.* 71: 22-26, 1985.
- [146288] 26282.

- [146289] McDonald, H.; Smailus, D.; Jenkins, H.; Adams, K.; Simpson, N.
- [146290] E.; Goodfellow, P. J.: Identification and characterization of a gene
- [146291] at D10S94 in the MEN2A region. Genomics 13: 344–348, 1992.
- [146292] 26283.
- [146293] Miwa, S.; Sato, T.; Murao, H.; Kozuru, M.; Ibayashi, H.: A new
- [146294] type of phosphofructokinase deficiency: hereditary non-spherocytic
- [146295] hemolytic anemia. Acta Haemat. Jpn. 35: 113–118, 1972.
- [146296] 26284.
- [146297] Vora, S.; DiMauro, S.; Spear, D.; Harker, D.; Danon, M. J.: Characterization
- [146298] of the enzymatic defect in late-onset muscle phosphofructokinase deficiency:
- [146299] new subtype of glycogen storage disease type VII. J. Clin. Invest. 80:
- [146300] 1479–1485, 1987.
- [146301] 26285.

- [146302]Bernardi, G.: Personal Communication. Paris, France
5/13/1992.
- [146303]26286.
- [146304]Boulard, M. R.; Bois, M.; Reviron, M.; Najean, Y.: Red-cell
phosphofructokinase
[146305]deficiency. New Eng. J. Med. 291: 978-979, 1974.
- [146306]26287.
- [146307]Chadefaux, B.; Rethore, M. O.; Allard, D.: Regional map-
ping of
[146308]liver type 6-phosphofructokinase isoenzyme on chromo-
some 21. Hum.
[146309]Genet. 68: 136-137, 1984.
- [146310]26288.
- [146311]Cox, D. R.; Kawashima, H.; Vora, S.; Epstein, C. J.: Regional
[146312]mapping of SOD-1, PRGS, and PFK-L on human chromo-
some 21. (Abstract) Cytogenet.
- [146313]Cell Genet. 37: 441-442, 1984.
- [146314]26289.
- [146315]Elson, A.; Levanon, D.; Brandeis, M.; Dafni, N.; Bernstein,
Y.;

- [146316]Danciger, E.; Groner, Y.: The structure of the human liver-type phosphofructokinase
- [146317]gene. Genomics 7: 47–56, 1990.
- [146318]26290.
- [146319]Etiemble, J.; Kahn, A.; Boivin, P.; Bernard, J. F.; Goude-
mand,
- [146320]M.: Hereditary hemolytic anemia with erythrocyte phospho-
phofructokinase
- [146321]deficiency. Hum. Genet. 31: 83–91, 1976.
- [146322]26291.
- [146323]Etiemble, J.; Picat, C.; Simeon, J.; Blatrix, C.; Boivin, P.:
- [146324]Inherited erythrocyte phosphofructokinase deficiency:
molecular mechanism. Hum.
- [146325]Genet. 55: 383–390, 1980.
- [146326]26292.
- [146327]Gehrich, S. C.; Gekakis, N.; Sul, H. S.: Liver (B-type)
phosphofructokinase
- [146328]mRNA: cloning, structure, and expression. J. Biol. Chem.
263: 11755–11759,
- [146329]1988.

[146330]26293.

[146331]Kahn, A.; Etiemble, J.; Meienhofer, M. C.; Boivin, P.: Erythrocyte

[146332]phosphofructokinase deficiency associated with an unstable variant

[146333]of muscle phosphofructokinase. Clin. Chim. Acta 61: 415-419, 1975.

[146334]26294.

[146335]Levanon, D.; Danciger, E.; Dafni, N.; Groner, Y.: Genomic clones

[146336]of the human liver-type phosphofructokinase. Biochem. Biophys. Res.

[146337]Commun. 141: 374-380, 1986.

[146338]26295.

[146339]Pantelakis, S. N.; Karaklis, A. G.; Alexiou, D.; Vardas, E.; Valaes,

[146340]T.: Red cell enzymes in trisomy 21. Am. J. Hum. Genet. 22: 184-193,

[146341]1970.

[146342]26296.

[146343]Van Keuren, M.; Drabkin, H.; Hart, I.; Harker, D.; Patter-

son,

[146344]D.; Vora, S.: Regional assignment of human liver-type
6-phosphofructokinase

[146345]to chromosome 21q22.3 by using somatic cell hybrids
and a monoclonal

[146346]anti-L antibody. Hum. Genet. 74: 34-40, 1986.

[146347]26297.

[146348]Vora, S.; Davidson, M.; Seaman, C.; Miranda, A. F.; Noble,
N.

[146349]A.; Tanaka, K. R.; Frenkel, E. P.; DiMauro, S.: Heterogeneity
of

[146350]the molecular lesions in inherited phosphofructokinase
deficiency. J.

[146351]Clin. Invest. 72: 1995-2006, 1983.

[146352]26298.

[146353]Vora, S.; Durham, S.; de Martinville, B.; Francke, U.: As-
signment

[146354]of the genes for liver type phosphofructokinase (PFKL) to
chromosome

[146355]21 and for muscle type (PFKM) to region p32-q32 of
chromosome 1. (Abstract) Cytogenet.

[146356]Cell Genet. 32: 324 only, 1982.

[146357]26299.

[146358]Vora, S.; Francke, U.: Assignment of the human gene for
liver

[146359]type phosphofructokinase isozyme to chromosome 21 us-
ing somatic cell

[146360]hybrids. (Abstract) Pediat. Res. 15: 570 only, 1981.

[146361]26300.

[146362]Vora, S.; Francke, U.: Assignment of the human gene for
liver-type

[146363]6-phosphofructokinase isozyme (PFKL) to chromosome 21
by using somatic

[146364]cell hybrids and monoclonal anti-L antibody. Proc. Nat.
Acad. Sci. 78:

[146365]3738-3742, 1981.

[146366]26301.

[146367]Wang, D.; Fang, H.; Cantor, C. R.; Smith, C. L.: A contigu-
ous

[146368]NotI restriction map of band q22.3 of human chromosome
21. Proc.

[146369]Nat. Acad. Sci. 89: 3222-3226, 1992.

[146370]26302.

[146371]Vidal, R.; Revesz, T.; Rostagno, A.; Kim, E.; Holton, J. L.;
Bek,

[146372]T.; Bojsen-Moller, M.; Braendgaard, H.; Plant, G.; Ghiso, J.;
Frangione,

[146373]B.: A decamer duplication in the 3-prime region of the BRI
gene originates

[146374]an amyloid peptide that is associated with dementia in a
Danish kindred. Proc.

[146375]Nat. Acad. Sci. 97: 4920-4925, 2000.

[146376]26303.

[146377]Fathallah-Shaykh, H.; Wolf, S.; Wong, E.; Posner, J. B.:
Cloning

[146378]of a leucine-zipper protein recognized by the sera of pa-
tients with

[146379]antibody-associated paraneoplastic cerebellar degenera-
tion. Proc.

[146380]Nat. Acad. Sci. 88: 3451-3454, 1991.

[146381]26304.

[146382]Gress, T.; Baldini, A.; Rocchi, M.; Furneaux, H.; Posner, J.
B.;

[146383]Siniscalco, M.: In situ mapping of the gene coding for a
leucine

[146384] zipper DNA binding protein (CDR 2) to the region between
two rare

[146385] fragile sites of autosome 16 (16p12–p13.1). (Abstract)
Cytogenet.

[146386] Cell Genet. 58: 1999–2000, 1991.

[146387] 26305.

[146388] Gress, T.; Baldini, A.; Rocchi, M.; Furneaux, H.; Posner, J.
B.;

[146389] Siniscalco, M.: In situ mapping of the gene coding for a
leucine

[146390] zipper DNA binding protein (CDR62) to 16p12–16p13.1.
Genomics 13:

[146391] 1340–1342, 1992.

[146392] 26306.

[146393] Raimondi, E.; Rubboli, F.; Moralli, D.; Chini, B.; Fornasari,
D.;

[146394] Tarroni, P.; De Carli, L.; Clementi, F.: Chromosomal local-
ization

[146395] and physical linkage of the genes encoding the human al-
pha-3, alpha-5,

[146396] and beta-4 neuronal nicotinic receptor subunits. Ge-
nomics 12: 849–850,

[146397]1992.

[146398]26307.

[146399]Anand, R.; Lindstrom, J.: Nucleotide sequence of the human nicotinic

[146400]acetylcholine receptor beta-2 subunit gene. Nucleic Acids Res. 18:

[146401]4272, 1990.

[146402]26308.

[146403]De Fusco, M.; Becchetti, A.; Patrignani, A.; Annesi, G.; Gambardella,

[146404]A.; Quattrone, A.; Ballabio, A.; Wanke, E.; Casari, G.: The nicotinic

[146405]receptor beta-2 subunit is mutant in nocturnal frontal lobe epilepsy. Nature

[146406]Genet. 26: 275-276, 2000.

[146407]26309.

[146408]Lueders, K. K.; Elliott, R. W.; Marenholz, I.; Mischke, D.; DuPree,

[146409]M.; Hamer, D.: Genomic organization and mapping of the human and

[146410]mouse neuronal beta-1-nicotinic acetylcholine receptor

genes. Mammalian

[146411]Genome 10: 900–905, 1999.

[146412]26310.

[146413]Phillips, H. A.; Favre, I.; Kirkpatrick, M.; Zuberi, S. M.;
Goudie,

[146414]D.; Heron, S. E.; Scheffer, I. E.; Sutherland, G. R.; Berkovic,
S.

[146415]F.; Bertrand, D.; Mulley, J. C.: CHRN2 is the second
acetylcholine

[146416]receptor subunit associated with autosomal dominant
nocturnal frontal

[146417]lobe epilepsy. Am. J. Hum. Genet. 68: 225–231, 2001.

[146418]26311.

[146419]Picciotto, M. R.; Zoli, M.; Lena, C.; Bessis, A.; Lallemant,
Y.;

[146420]LeNovere, N.; Vincent, P.; Pich, E. M.; Brulet, P.; Changeux,
J.–P.

[146421]: Abnormal avoidance learning in mice lacking functional
high-affinity

[146422]nicotine receptor in the brain. Nature 374: 65–67, 1995.

[146423]26312.

[146424]Zoli, M.; Picciotto, M. R.; Ferrari, R.; Cocchi, D.; Changeux,
[146425]J.-P.: Increased neurodegeneration during ageing in mice
lacking
[146426]high-affinity nicotine receptors. EMBO J. 18: 1235-1244,
1999.
[146427]26313.

[146428]Tarroni, P.; Rubboli, F.; Chini, B.; Zwart, R.; Oortgiesen, M.;
[146429]Sher, E.; Clementi, F.: Neuronal-type nicotinic receptors in
human
[146430]neuroblastoma and small-cell lung carcinoma cell lines.
FEBS Lett. 312:
[146431]66-70, 1992.
[146432]26314.

[146433]Bonner, T. I.; Buckley, N. J.; Young, A. C.; Brann, M. R.:
Identification
[146434]of a family of muscarinic acetylcholine receptor genes.
Science 237:
[146435]527-532, 1987.
[146436]26315.

[146437]Eng, C. E. L.; Strom, C. M.: Analysis of three restriction
fragment

[146438]length polymorphisms in the human type II procollagen gene. Am. J.

[146439]Hum. Genet. 37: 719–732, 1985.

[146440]26316.

[146441]Eyre, D. R.; Weis, M. A.; Moskowitz, R. W.: Cartilage expression

[146442]of a type II collagen mutation in an inherited form of osteoarthritis

[146443]associated with a mild chondrodysplasia. J. Clin. Invest. 87: 357–361,

[146444]1991.

[146445]26317.

[146446]Francomano, C. A.; Liberfarb, R. M.; Hirose, T.; Maumenee, I.

[146447]H.; Streeten, E. A.; Meyers, D. A.; Pyeritz, R. E.: The Stickler

[146448]syndrome: evidence for close linkage to the structural gene for type

[146449]II collagen. Genomics 1: 293–296, 1987.

[146450]26318.

[146451]Francomano, C. A.; Maumenee, I.; Liberfarb, R.; Pyeritz, R. E.

[146452]: Cosegregation of Stickler syndrome and type II collagen gene alleles.

[146453](Abstract) Cytogenet. Cell Genet. 46: 615 only, 1987.

[146454]26319.

[146455]Garofalo, S.; Vuorio, E.; Metsaranta, M.; Rosati, R.; Toman, D.;

[146456]Vaughan, J.; Lozano, G.; Mayne, R.; Ellard, J.; Horton, W.; de Crombrughe,

[146457]B.: Reduced amounts of cartilage collagen fibrils and growth plate

[146458]anomalies in transgenic mice harboring a glycine-to-cysteine mutation

[146459]in the mouse type II procollagen alpha-1-chain gene. Proc. Nat. Acad.

[146460]Sci. 88: 9648-9652, 1991.

[146461]26320.

[146462]Gupta, S. K.; Leonard, B. C.; Damji, K. F.; Bulman, D. E.: A

[146463]frame shift mutation in a tissue-specific alternatively spliced exon

[146464]of collagen 2A1 in Wagner's vitreoretinal degeneration. Am. J. Ophthal. 133:

[146465]203–210, 2002.

[146466]26321.

[146467]Holderbaum, D.; Malemud, C. J.; Moskowitz, R. W.; Haqqi, T. M.

[146468]: Human cartilage from late stage familial osteoarthritis transcribes

[146469]type II collagen mRNA encoding a cysteine in position 519. *Biochem.*

[146470]*Biophys. Res. Commun.* 192: 1169–1174, 1993.

[146471]26322.

[146472]Horton, W. A.: Personal Communication. Houston, Texas 1987.

[146473]26323.

[146474]Horton, W. A.; Machado, M. A.; Ellard, J.; Campbell, D.; Bartley,

[146475]J.; Ramirez, F.; Vitale, E.; Lee, B.: Characterization of a type

[146476]II collagen gene (COL2A1) mutation identified in cultured chondrocytes

[146477]from human hypochondrogenesis. *Proc. Nat. Acad. Sci.* 89: 4583–4587,

[146478]1992.

[146479]26324.

[146480]Huerre-Jeanpierre, C.; Mattei, M.-G.; Weil, D.; Grzeschik, K.

[146481]H.; Chu, M.-L.; Sangiorgi, F. O.; Sobel, M. E.; Ramirez, F.; Junien,

[146482]C.: Further evidence for the dispersion of the human fibrillar collagen

[146483]genes. Am. J. Hum. Genet. 38: 26-37, 1986.

[146484]26325.

[146485]Huerre-Jeanpierre, C.; Mattei, M. G.; Weil, D.; Grzeschik, K.

[146486]H.; Sangiorgi, F. O.; Ramirez, F.; Junien, C.: The gene for human

[146487]alpha-1 type II collagen (COL2A1) maps to band 12q131. (Abstract) Cytogenet.

[146488]Cell Genet. 40: 657-658, 1985.

[146489]26326.

[146490]Kaitila, I.; Korokko, J.; Marttinen, E.; Ala-Kokko, L.: Phenotypic

[146491]expressions of a gly154-to-arg mutation in type II collagen in two

[146492]unrelated patients with spondyloepimetaphyseal dysplasia

(SEMD). Am.

[146493]J. Med. Genet. 63: 111-122, 1996.

[146494]26327.

[146495]Kaprio, J.; Kujala, U. M.; Peltonen, L.; Koskenvuo, M.: Genetic

[146496]liability to osteoarthritis may be greater in women than in men. (Letter) Brit.

[146497]Med. J. 313: 232 only, 1996.

[146498]26328.

[146499]Kelly, T. E.; Lichtenstein, J. R.; Dorst, J. P.: An unusual familial

[146500]spondyloepiphyseal dysplasia: 'spondyloperipheral dysplasia.'. Birth

[146501]Defects Orig. Art. Ser. XIII(3B): 149-165, 1977.

[146502]26329.

[146503]Knowlton, R. G.; Katzenstein, P.; Moskowitz, R. W.; Weaver, E.

[146504]J.; Jimenez, S. A.; Pathria, M. N.; Malesud, C. J.; Prockop, D. J.

[146505]: Genetic linkage of the type II procollagen gene to primary generalized

[146506]osteoarthritis with chondrodysplasia. (Abstract) Cyto-
genet. Cell

[146507]Genet. 51: 1024 only, 1989.

[146508]26330.

[146509]Knowlton, R. G.; Katzenstein, P. L.; Moskowitz, R. W.;
Weaver,

[146510]E. J.; Malemud, C. J.; Pathria, M. N.; Jimenez, S. A.;
Prockop, D.

[146511]J.: Genetic linkage of a polymorphism in the type II pro-
collagen

[146512]gene (COL2A1) to primary osteoarthritis associated with
mild chondrodysplasia. New

[146513]Eng. J. Med. 322: 526-530, 1990.

[146514]26331.

[146515]Korkko, J.; Cohn, D. H.; Ala-Kokko, L.; Krakow, D.;
Prockop, D.

[146516]J.: Widely distributed mutations in the COL2A1 gene pro-
duce achondrogenesis

[146517]type II/hypochondrogenesis. Am. J. Med. Genet. 92:
95-100, 2000.

[146518]26332.

[146519]Law, M. L.; Tung, L.; Morse, H. G.; Berger, R.; Jones, C.;
Cheah,

[146520]K. S. E.; Solomon, E.: The human type II collagen gene
(COL2A1) assigned

[146521]to 12q14.3. Ann. Hum. Genet. 50: 131–137, 1986.

[146522]26333.

[146523]Law, M. L.; Tung, L.; Morse, H. G.; Berger, R.; Jones, C.;
Cheah,

[146524]K. S. E.; Stoker, N. G.; Solomon, E.: Regional assignment of
the

[146525]human type II collagen gene (COL2A1) on chromosome
12. (Abstract) Cytogenet.

[146526]Cell Genet. 40: 678 only, 1985.

[146527]26334.

[146528]Learmonth, I. D.; Christy, G.; Beighton, P.: Namaqualand
hip

[146529]dysplasia: orthopaedic implications. Clin. Orthop. 218:
142–147,

[146530]1987.

[146531]26335.

[146532]Li, S. W.; Prockop, D. J.; Helminen, H.; Fassler, R.; Lapvete–

lainen,

[146533]T.; Kiraly, K.; Peltarri, A.; Arokoski, J.; Lui, H.; Arita, M.;

Khillan,

[146534]J. S.: Transgenic mice with targeted inactivation of the

Col2a1 gene

[146535]for collagen II develop a skeleton with membranous and

periosteal

[146536]bone but no endochondral bone. Genes Dev. 9:

2821–2830, 1995.

[146537]26336.

[146538]Lovell–Badge, R. H.; Bygrave, A.; Bradley, A.; Robertson, E.;

[146539]Tilly, R.; Cheah, K. S. E.: Tissue–specific expression of the

human

[146540]type II collagen gene in mice. Proc. Nat. Acad. Sci. 84:

2803–2807,

[146541]1987.

[146542]26337.

[146543]Schwenk, W. F.; Rizza, R. A.; Mandarino, L. J.; Gerich, J. E.;

[146544]Hayles, A. B.; Haymond, M. W.: Familial insulin resistance

and acanthosis

[146545]nigricans: presence of a postbinding defect. Diabetes 35:

33–37,

[146546]1986.

[146547]26338.

[146548]Harris, E. L.: Personal Communication. Dunedin, New Zealand

[146549]11/27/2001.

[146550]26339.

[146551]Rossaak, J. I.; van Rij, A. M.; Jones, G. T.; Harris, E. L.:

[146552]Association of the 4G/5G polymorphism in the promoter region of plasminogen

[146553]activator inhibitor–1 with abdominal aortic aneurysms. J. Vasc. Surg. 31:

[146554]1026–1032, 2000.

[146555]26340.

[146556]Steinberger, D.; Mulliken, J. B.; Muller, U.: Crouzon syndrome:

[146557]previously unrecognized deletion, duplication, and point mutation

[146558]within FGFR2 gene. Hum. Mutat. 8: 386–390, 1996.

[146559]26341.

[146560]Oettinger, M. A.; Schatz, D. G.; Gorka, C.; Baltimore, D.:
RAG-1

[146561]and RAG-2, adjacent genes that synergistically activate
V(D)J recombination. Science 248:

[146562]1517-1523, 1990.

[146563]26342.

[146564]Steinberger, D.; Reinhartz, T.; Unsold, R.; Muller, U.:
FGFR2

[146565]mutation in clinically nonclassifiable autosomal dominant
craniosynostosis

[146566]with pronounced phenotypic variation. Am. J. Med. Genet.
66: 81-86,

[146567]1996.

[146568]26343.

[146569]Steinberger, D.; Vriend, G.; Mulliken, J. B.; Muller, U.: The

[146570]mutations in FGFR2-associated craniosynostoses are clustered in five

[146571]structural elements of immunoglobulin-like domain III of
the receptor. Hum.

[146572]Genet. 102: 145-150, 1998. Note: Erratum: Hum. Genet.
102: 495 only,

[146573]1998.

[146574]26344.

[146575]Tartaglia, M.; Di Rocco, C.; Lajeunie, E.; Valeri, S.; Velardi,

[146576]F.; Battaglia, P. A.: Jackson–Weiss syndrome: identification
of two

[146577]novel FGFR2 missense mutations shared with Crouzon and
Pfeiffer craniosynostotic

[146578]disorders. Hum. Genet. 101: 47–50, 1997.

[146579]26345.

[146580]Tartaglia, M.; Valeri, S.; Velardi, F.; Di Rocco, C.; Battaglia,

[146581]P. A.: Trp290Cys mutation in exon IIIa of the fibroblast
growth factor

[146582]receptor 2 (FGFR2) gene is associated with Pfeiffer syn-
drome. Hum.

[146583]Genet. 99: 602–606, 1997.

[146584]26346.

[146585]Teebi, A. S.; Kennedy, S.; Chun, K.; Ray, P. N.: Severe and
mild

[146586]phenotypes in Pfeiffer syndrome with splice acceptor mu-
tations in

[146587]lexon IIIc of FGFR2. Am. J. Med. Genet. 107: 43–47, 2002.

[146588]26347.

[146589]Tsukuno, M.; Suzuki, H.; Eto, T.: Pfeiffer syndrome caused
by

[146590]haploinsufficient mutation of FGFR2. J. Craniofac. Genet.
Dev. Biol. 19:

[146591]183–188, 1999.

[146592]26348.

[146593]Van Dissel–Emiliani, F. M.; De Boer–Brouwer, M.; De Rooij,
D.

[146594]G.: Effect of fibroblast growth factor–2 on Sertoli cells and
gonocytes

[146595]in coculture during the perinatal period. Endocrinology
137: 647–654,

[146596]1996.

[146597]26349.

[146598]Vogt, A.: Dyskephalie (dysostosis craniofacialis, maladie
De

[146599]Crouzon 1912) und eine neuartige Kombination dieser
Krankheit mit

[146600]Syndaktylie der 4 Extremitaeten (Dyskephalodaktylie).

Klin. Mbl.

[146601]Augenheilk. 90: 441–454, 1933.

[146602]26350.

[146603]Wang, T.-J.; Huang, C.-B.; Tsai, F.-J.; Wu, J.-Y.; Lai, R.-B.;

[146604]Hsiao, M.: Mutation in the FGFR2 gene in a Taiwanese patient with

[146605]Beare–Stevenson cutis gyrata syndrome. Clin. Genet. 61: 218–221,

[146606]2002.

[146607]26351.

[146608]Wilkie, A. O. M.: Abnormal spliceform expression associated with

[146609]splice acceptor mutations in exon IIIc of FGFR2. (Letter) Am. J.

[146610]Med. Genet. 111: 105 only, 2002.

[146611]26352.

[146612]Wilkie, A. O. M.: Personal Communication. Oxford, U.K. 2/24/1996.

[146613]26353.

[146614]Wilkie, A. O. M.; Slaney, S. F.; Oldridge, M.; Poole, M. D.;

Ashworth,

[146615]G. J.; Hockley, A. D.; Hayward, R. D.; David, D. J.; Pulleyn, L. J.;

[146616]Rutland, P.; Malcolm, S.; Winter, R. M.; Reardon, W.: Apert syndrome

[146617]results from localized mutations of FGFR2 and is allelic with Crouzon

[146618]syndrome. *Nature Genet.* 9: 165–172, 1995.

[146619]26354.

[146620]Wong, L.-J. C.; Chen, T.-J.; Dai, P.; Bird, L.; Muenke, M.: Novel

[146621]SNP at the common primer site of exon IIIa of FGFR2 gene causes error

[146622]in molecular diagnosis of craniosynostosis syndrome. *Am. J. Med.*

[146623]*Genet.* 102: 282–285, 2001.

[146624]26355.

[146625]Xu, X.; Weinstein, M.; Li, C.; Naski, M.; Cohen, R. I.; Ornitz,

[146626]D. M.; Leder, P.; Deng, C.: Fibroblast growth factor receptor 2 (FGFR2)–mediated

[146627]reciprocal regulation loop between FGF8 and FGF10 is es–

essential for

[146628]limb induction. Development 125: 753–765, 1998.

[146629]26356.

[146630]Yu, K.; Herr, A. B.; Waksman, G.; Ornitz, D. M.: Loss of fibroblast

[146631]growth factor receptor 2 ligand-binding specificity in Apert syndrome. Proc.

[146632]Nat. Acad. Sci. 97: 14536–14541, 2000.

[146633]26357.

[146634]Chan, J.; Watt, V. M.: Eek and erk, new members of the eph subclass

[146635]of receptor protein-tyrosine kinases. Oncogene 6: 1057–1061, 1991.

[146636]26358.

[146637]Park, S.; Frisen, J.; Barbacid, M.: Aberrant axonal projections

[146638]in mice lacking EphA8 (Eek) tyrosine protein kinase receptors. EMBO

[146639]J. 16: 3106–3114, 1997.

[146640]26359.

[146641]Ganju, P.; Shigemoto, K.; Brennan, J.; Entwistle, A.; Reith,

A.

- [146642]D.: The Eck receptor tyrosine kinase is implicated in pattern formation during gastrulation, hindbrain segmentation and limb development. *Oncogene* 9: 1613–1624, 1994.
- [146645]26360.
- [146646]Lindberg, R. A.; Hunter, T.: cDNA cloning and characterization of eck, an epithelial cell receptor–tyrosine kinase in the eph/elk family of protein kinases. *Molec. Cell. Biol.* 10: 6316–6324, 1990.
- [146649]26361.
- [146650]Sulman, E. P.; Tang, X. X.; Allen, C.; Biegel, J. A.; Pleasure, D. E.; Brodeur, G. M.; Ikegaki, N.: ECK, a human EPH–related gene, maps to 1p36.1, a common region of alteration in human cancers. *Genomics* 40: 371–374, 1997.
- [146654]26362.

- [146655]Arpaia, E.; Shahar, M.; Dadi, H.; Cohen, A.; Roifman, C. M.:
Defective
- [146656]T cell receptor signaling and CD8(+) thymic selection in
humans lacking
- [146657]Zap-70 kinase. Cell 76: 947-958, 1994.
- [146658]26363.
- [146659]Chan, A. C.; Iwashima, M.; Turck, C. W.; Weiss, A.: ZAP-
70: a
- [146660]70 kd protein-tyrosine kinase that associates with the
TCR zeta chain. Cell 71:
- [146661]649-662, 1992.
- [146662]26364.
- [146663]Chan, A. C.; Kadlecsek, T. A.; Elder, M. E.; Filipovich, A. H.;
- [146664]Kuo, W.-L.; Iwashima, M.; Parslow, T. G.; Weiss, A.: ZAP-
70 deficiency
- [146665]in an autosomal recessive form of severe combined im-
munodeficiency. Science 264:
- [146666]1559-1601, 1994.
- [146667]26365.
- [146668]Colucci, F.; Schweighoffer, E.; Tomasello, E.; Turner, M.;
Ortaldo,

[146669]J. R.; Vivier, E.; Tybulewicz, V. L. J.; Di Santo, J. P.: Natural
 [146670]cytotoxicity uncoupled from the Syk and ZAP-70 intracel-
 lular kinases. Nature
 [146671]Immun. 3: 288-294, 2002.
 [146672]26366.
 [146673]Elder, M. E.; Lin, D.; Clever, J.; Chan, A. C.; Hope, T. J.;
 Weiss,
 [146674]A.; Parslow, T. G.: Human severe combined immunodefi-
 ciency due to
 [146675]a defect in ZAP-70, a T cell tyrosine kinase. Science 264:
 1596-1599,
 [146676]1994.
 [146677]26367.
 [146678]Ku, G.; Malissen, B.; Mattei, M.-G.: Chromosomal location
 of the
 [146679]Syk and ZAP-70 tyrosine kinase genes in mice and hu-
 mans. Immunogenetics 40:
 [146680]300-302, 1994.
 [146681]26368.
 [146682]Roifman, C. M.; Hummel, D.; Martinez-Valdez, H.;
 Thorner, P.; Doherty,

- [146683]P. J.; Pan, S.; Cohen, F.; Cohen, A.: Depletion of CD8(+) cells in
[146684]human thymic medulla results in selective immune deficiency. J. Exp.
[146685]Med. 170: 2177–2182, 1989.
[146686]26369.
- [146687]Saito, T.; Matsuda, Y.; Ito, H.; Fusaki, N.; Hori, T.; Yamamoto,
[146688]T.: Localization of Zap70, the gene for a T cell-specific protein
[146689]tyrosine kinase, to mouse and rat chromosomes by fluorescence in situ
[146690]hybridization and molecular genetic linkage analyses. Mammalian Genome 8:
[146691]45–46, 1997.
[146692]26370.
- [146693]Toyabe, S.-I.; Watanabe, A.; Harada, W.; Karasawa, T.; Uchiyama,
[146694]M.: Specific immunoglobulin E responses in ZAP-70-deficient patients
[146695]are mediated by Syk-dependent T-cell receptor signalling. Immunology 103:

[146696]164–171, 2001.

[146697]26371.

[146698]Oettinger, M. A.; Stanger, B.; Schatz, D. G.; Glaser, T.; Call,

[146699]K.; Housman, D.; Baltimore, D.: The recombination activating genes,

[146700]RAG1 and RAG2, are on chromosome 11p in humans and chromosome 2p in

[146701]mice. Immunogenetics 35: 97–101, 1992.

[146702]26372.

[146703]Roman, C. A. J.; Baltimore, D.: Genetic evidence that the RAG1

[146704]protein directly participates in V(D)J recombination through substrate

[146705]recognition. Proc. Nat. Acad. Sci. 93: 2333–2338, 1996.

[146706]26373.

[146707]Santagata, S.; Gomez, C. A.; Sobacchi, C.; Bozzi, F.; Abinun,

[146708]M.; Pasic, S.; Cortes, P.; Vezzoni, P.; Villa, A.: N-terminal RAG1

[146709]frameshift mutations in Omenn's syndrome: internal methionine usage

[146710]leads to partial V(D)J recombination activity and reveals a
fundamental

[146711]role in vivo for the N-terminal domains. Proc. Nat. Acad.
Sci. 97:

[146712]14572–14577, 2000.

[146713]26374.

[146714]Schatz, D. G.; Oettinger, M. A.; Baltimore, D.: The V(D)J re-
combination

[146715]activating gene, RAG-1. Cell 59: 1035–1048, 1989.

[146716]26375.

[146717]Schultz, H. Y.; Landree, M. A.; Qiu, J.; Kale, S. B.; Roth, D.

[146718]B.: Joining-deficient RAG1 mutants block V(D)J recombina-
tion in vivo

[146719]and hairpin opening in vitro. Molec. Cell 7: 65–75, 2001.

[146720]26376.

[146721]Schwarz, K.; Gauss, G. H.; Ludwig, L.; Pannicke, U.; Li, Z.;
Linder,

[146722]D.; Friedrich, W.; Seger, R. A.; Hansen-Hagge, T. E.;
Desiderio, S.;

[146723]Lieber, M. R.; Bartram, C. R.: RAG mutations in human B
cell-negative

[146724]SCID. Science 274: 97–99, 1996.

[146725]26377.

[146726]Sherrington, P. D.; Forster, A.; Seawright, A.; van Heyning,
gen,

[146727]V.; Rabbitts, T. H.: Human RAG2, like RAG1, is on chromosome 11 band

[146728]p13 and therefore not linked to ataxia telangiectasia complementation

[146729]groups. Genes Chromosomes Cancer 5: 404–406, 1992.

[146730]26378.

[146731]Villa, A.; Santagata, S.; Bozzi, F.; Giliani, S.; Frattini, A.;

[146732]Imberti, L.; Gatta, L. B.; Ochs, H. D.; Schwarz, K.; Notarangelo,

[146733]L. D.; Vezzoni, P.; Spanopoulou, E.: Partial V(D)J recombination

[146734]activity leads to Omenn syndrome. Cell 93: 885–896, 1998.

[146735]26379.

[146736]Wienholds, E.; Schulte–Merker, S.; Walderich, B.; Plasterk, R.

[146737]H. A.: Target–selected inactivation of the zebrafish rag1

gene. Science 297:

[146738]99–102, 2002.

[146739]26380.

[146740]Yu, W.; Misulovin, Z.; Suh, H.; Hardy, R. R.; Jankovic, M.;
Yannoutsos,

[146741]N.; Nussenzweig, M. C.: Coordinate regulation of RAG1
and RAG2 by

[146742]cell type–specific DNA elements 5–prime of RAG2. Science
285: 1080–1084,

[146743]1999.

[146744]26381.

[146745]Richard, G.; Brown, N.; Smith, L. E.; Terrinoni, A.; Melino,
G.;

[146746]MacKie, R. M.; Bale, S. J.; Uitto, J.: The spectrum of muta-
tions

[146747]in erythrokeratodermias--novel and de novo mutations in
GJB3. Hum.

[146748]Genet. 106: 321–329, 2000.

[146749]26382.

[146750]Levitt, R. C.; Olckers, A.; Meyers, S.; Fletcher, J. E.; Rosen-
berg,

[146751]H.; Isaacs, H.; Meyers, D. A.: Evidence for the localization
of a

[146752]malignant hyperthermia susceptibility locus (MHS2) to human chromosome

[146753]17q. Genomics 14: 562–566, 1992.

[146754]26383.

[146755]Olickers, A.; Meyers, D. A.; Meyers, S.; Taylor, E. W.; Fletcher,

[146756]J. E.; Rosenberg, H.; Isaacs, H.; Levitt, R. C.: Adult muscle sodium

[146757]channel alpha-subunit is a gene candidate for malignant hyperthermia

[146758]susceptibility. Genomics 14: 829–831, 1992.

[146759]26384.

[146760]Naim, H. Y.; Sterchi, E. E.; Lentze, M. J.: Structure, biosynthesis,

[146761]and glycosylation of human small intestinal maltase-glucoamylase. J.

[146762]Biol. Chem. 263: 19709–19717, 1988.

[146763]26385.

[146764]Nichols, B. L.; Eldering, J.; Avery, S.; Hahn, D.; Quaroni, A.;

[146765]Sterchi, E.: Human small intestinal maltase-glucoamylase

cDNA cloning:

[146766]homology to sucrase-isomaltase. J. Biol. Chem. 273:
3076-3081, 1998.

[146767]26386.

[146768]Schuler, G. D.; Boguski, M. S.; Stewart, E. A.; Stein, L. D.;
Gyapay,

[146769]G.; Rice, K.; White, R. E.; Rodriguez-Tome, P.; Aggarwal,
A.; Bajorek,

[146770]E.; Bentolila, S.; Birren, B. B.; Butler, A.; Castle, A. B.; Chi-
annilkulchai,

[146771]N.; Chu, A.; Clee, C.; Cowles, S.; Day, P. J. R.; Dibling, T.;
and

[146772]84 others: A gene map of the human genome. Science
274: 540-546,

[146773]1996.

[146774]26387.

[146775]Levine, A.; Cantoni, G. L.; Razin, A.: Inhibition of promoter

[146776]activity by methylation: possible involvement of protein
mediators. Proc.

[146777]Nat. Acad. Sci. 88: 6515-6518, 1991.

[146778]26388.

[146779]Nan, X.; Meehan, R. R.; Bird, A.: Dissection of the methyl-CpG

[146780]binding domain from the chromosomal protein MeCP2.
Nucleic Acids

[146781]Res. 21: 4886-4892, 1993.

[146782]26389.

[146783]Ohki, I.; Shimotake, N.; Fujita, N.; Jee, J.-G.; Ikegami, T.;
Nakao,

[146784]M.; Shirakawa, M.: Solution structure of the methyl-CpG
binding domain

[146785]of human MBD1 in complex with methylated DNA. Cell
105: 487-497,

[146786]2001.

[146787]26390.

[146788]Carrera, C. J.; Eddy, R. L.; Shows, T. B.; Carson, D. A.: As-
signment

[146789]of the gene for methylthioadenosine phosphorylase to hu-
man chromosome

[146790]9 by mouse-human somatic cell hybridization. Proc. Nat.
Acad. Sci. 81:

[146791]2665-2668, 1984.

[146792]26391.

[146793]Chilcote, R. R.; Brown, E.; Rowley, J. D.: Lymphoblastic leukemia

[146794]with lymphomatous features associated with abnormalities of the short

[146795]arm of chromosome 9. New Eng. J. Med. 313: 286-291, 1985.

[146796]26392.

[146797]Nobori, T.; Takabayashi, K.; Tran, P.; Orvis, L.; Batova, A.; Yu,

[146798]A. L.; Carson, D. A.: Genomic cloning of methylthioadenosine phosphorylase:

[146799]a purine metabolic enzyme deficient in multiple different cancers. Proc.

[146800]Nat. Acad. Sci. 93: 6203-6208, 1996.

[146801]26393.

[146802]Olopade, O. I.; Jenkins, R. B.; Ransom, D. T.; Malik, K.; Pomykala,

[146803]H.; Nobori, T.; Cowan, J. M.; Rowley, J. D.; Diaz, M. O.: Molecular

[146804]analysis of deletions of the short arm of chromosome 9 in human gliomas. Cancer

[146805]Res. 52: 2523–2529, 1992.

[146806]26394.

[146807]Olopade, O. I.; Pomykala, H. M.; Hagos, F.; Sveen, L. W.;
Espinosa,

[146808]R., III; Dreyling, M. H.; Gursky, S.; Stadler, W. M.; Le Beau,
M.

[146809]M.; Bohlander, S. K.: Construction of a 2.8–megabase
yeast artificial

[146810]chromosome contig and cloning of the human methylthioadenosine phosphorylase

[146811]gene from the tumor suppressor region on 9p21. Proc.
Nat. Acad. Sci. 92:

[146812]6489–6493, 1995.

[146813]26395.

[146814]Ragione, F. D.; Takabayashi, K.; Mastropietro, S.; Mercurio,
C.;

[146815]Oliva, A.; Russo, G. L.; Pietra, V. D.; Borriello, A.; Nobori,
T.;

[146816]Carson, D. A.; Zappia, V.: Purification and characterization
of recombinant

[146817]human 5–prime–methylthioadenosine phosphorylase:
definite identification

[146818]of coding cDNA. Biochem. Biophys. Res. Commun. 223:
514–519, 1996.

[146819]26396.

[146820]Williams–Ashman, H. G.; Seidenfeld, J.; Galletti, P.: Trends
in

[146821]the biochemical pharmacology of

5–prime–deoxy–5–prime–methylthioadenosine. Biochem.

[146822]Pharm. 31: 277–288, 1982.

[146823]26397.

[146824]Kniest, W.: Zur Abgrenzung der Dysostosis enchondralis
von der

[146825]Chondrodystrophie. Z. Kinderheilk. 43: 633–640, 1952.

[146826]26398.

[146827]Mauensee, I. H.; Traboulsi, E. I.: The ocular findings in
Kniest

[146828]dysplasia. Am. J. Ophthal. 100: 155–160, 1985.

[146829]26399.

[146830]Siggers, D. C.; Rimoin, D. L.; Dorst, J. P.; Doty, S. B.;
Williams,

[146831]B. R.; Hollister, D. W.; Silberberg, R.; Granley, R. E.; Kauf-
man,

[146832]R. L.; McKusick, V. A.: The Kniest syndrome. Birth Defects
Orig.

[146833]Art. Ser. 10(9): 193–208, 1974.

[146834]26400.

[146835]Spranger, J.; Winterpacht, A.; Zabel, B.: Kniest dysplasia:
Dr.

[146836]W. Kniest, his patient, the molecular defect. Am. J. Med.
Genet. 69:

[146837]79–84, 1997.

[146838]26401.

[146839]Wilkin, D. J.; Artz, A. S.; South, S.; Lachman, R. S.; Rimoin,

[146840]D. L.; Wilcox, W. R.; McKusick, V. A.; Stratakis, C. A.; Fran-
comano,

[146841]C. A.; Cohn, D. H.: Small deletions in the type II collagen
triple

[146842]helix produce Kniest dysplasia. Am. J. Med. Genet. 85:
105–112,

[146843]1999.

[146844]26402.

[146845]Cirullo, R. E.; Wasmuth, J. J.: Assignment of the human
gene encoding

- [146846]methionyl-tRNA synthetase to chromosome 12. (Abstract)
Cytogenet.
- [146847]Cell Genet. 37: 437 only, 1984.
- [146848]26403.
- [146849]Cirullo, R. E.; Wasmuth, J. J.: Assignment of the human
MARS gene,
[146850]encoding methionyl-tRNA synthetase, to chromosome
12 using human
[146851]x Chinese hamster cell hybrids. Somat. Cell Molec. Genet.
10: 225-234,
[146852]1984.
- [146853]26404.
- [146854]Grewal, P. K.; Holzfeind, P. J.; Bittner, R. E.; Hewitt, J. E.
[146855]; Mutant glycosyltransferase and altered glycosylation of
alpha-dystroglycan
[146856]in the myodystrophy mouse. Nature Genet. 28: 151-154,
2001.
- [146857]26405.
- [146858]Evans, D. G. R.; Huson, S. M.; Donnai, D.; Neary, W.; Blair,
V.;
- [146859]Newton, V.; Harris, R.: A clinical study of type 2 neurofi-

bromatosis. Quart.

[146860]J. Med. 84: 603–618, 1992.

[146861]26406.

[146862]Honda, M.; Arai, E.; Sawada, S.; Ohta, A.; Niimura, M.:

Neurofibromatosis

[146863]2 and neurilemmomatosis gene are identical. J. Invest.

Derm. 104:

[146864]74–77, 1995.

[146865]26407.

[146866]Parry, D. M.; Eldridge, R.; Kaiser–Kupfer, M. I.; Bouzas, E.

A.;

[146867]Pikus, A.; Patronas, N.: Neurofibromatosis 2 (NF2): clinical

characteristics

[146868]of 63 affected individuals and clinical evidence for hetero–

geneity. Am.

[146869]J. Med. Genet. 52: 450–461, 1994.

[146870]26408.

[146871]Eddy, R. L.; Kretschmer, P. J.; Fairhurst, J. L.; Shows, T. B.;

[146872]Bohlen, P.; O'Hara, B.; Kovesdi, I.: A human gene family of

neurite

[146873]outgrowth–promoting proteins: the gene for a heparin

binding neurite

[146874]outgrowth-promoting factor maps to 7q22-qter.

(Abstract) Cytogenet.

[146875]Cell Genet. 58: 1920 only, 1991.

[146876]26409.

[146877]Lai, S.; Czubayko, F.; Riegel, A. T.; Wellstein, A.: Structure

[146878]of the human heparin-binding growth factor gene

pleiotrophin. Biochem.

[146879]Biophys. Res. Commun. 187: 1113-1122, 1992.

[146880]26410.

[146881]Li, Y.-S.; Hoffman, R. M.; Le Beau, M. M.; Espinosa, R., III;

Jenkins,

[146882]N. A.; Gilbert, D. J.; Copeland, N. G.; Deuel, T. F.: Charac-

terization

[146883]of the human pleiotrophin gene: promoter region and

chromosomal localization. J.

[146884]Biol. Chem. 267: 26011-26016, 1992.

[146885]26411.

[146886]Li, Y.-S.; Milner, P. G.; Chauhan, A. K.; Watson, M. A.;

Hoffman,

[146887]R. M.; Kodner, C. M.; Milbrandt, J.; Deuel, T. F.: Cloning

and expression

[146888]of a developmentally regulated protein that induces mitogenic and

[146889]neurite outgrowth activity. Science 250: 1690–1694, 1990.

[146890]26412.

[146891]Milner, P. G.; Shah, D.; Veile, R.; Donis-Keller, H.; Kumar, B.

[146892]V.: Cloning, nucleotide sequence, and chromosome localization of

[146893]the human pleiotrophin gene. Biochemistry 31: 12023–12028, 1992.

[146894]26413.

[146895]Brightbill, H. D.; Libraty, D. H.; Krutzik, S. R.; Yang, R.-B.;

[146896]Bellsie, J. T.; Bieharski, J. R.; Maitland, M.; Norgard, M. V.; Plevy,

[146897]S. E.; Smale, S. T.; Brennan, P. J.; Bloom, B. R.; Godowski, P. J.;

[146898]Modlin, R. L.: Host defense mechanisms triggered by microbial lipoproteins

[146899]through toll-like receptors. Science 285: 732–736, 1999.

[146900]26414.

- [146901]Jankovic, D.; Kullberg, M. C.; Hieny, S.; Caspar, P.; Collazo,
[146902]C. M.; Sher, A.: In the absence of IL-12, CD4+ T cell re-
sponses to
[146903]intracellular pathogens fail to default to a Th2 pattern and
are host
[146904]protective in an IL-10-/- setting. Immunity 16: 429-439,
2002.
[146905]26415.
- [146906]Jouanguy, E.; Lamhamedi-Cherradi, S.; Altare, F.; Fon-
daneche,
[146907]M.-C.; Tuerlinckx, D.; Blanche, S.; Emile, J.-F.; Gaillard,
J.-L.;
[146908]Schreiber, R.; Levin, M.; Fischer, A.; Hivroz, C.; Casanova,
J.-L.
- [146909]: Partial interferon-gamma receptor 1 deficiency in a child
with tuberculoid
[146910]bacillus Calmette-Guerin infection and a sibling with clinical tuberculosis. J.
[146911]Clin. Invest. 100: 2658-2664, 1997.
[146912]26416.
- [146913]Warrington, J. A.; Bengtsson, U.: High-resolution physical
mapping

[146914] of human 5q31-q33 using three methods: radiation hybrid mapping, interphase

[146915] fluorescence in situ hybridization, and pulsed-field gel electrophoresis. Genomics 24:

[146916] 395-398, 1994.

[146917] 26417.

[146918] Anderson, S. K.; Gallinger, S.; Roder, J.; Frey, J.; Young, H.

[146919] A.; Ortaldo, J. R.: A cyclophilin-related protein involved in the

[146920] function of natural killer cells. Proc. Nat. Acad. Sci. 90: 542-546,

[146921] 1993.

[146922] 26418.

[146923] Frey, J. L.; Bino, T.; Kantor, R. R. S.; Segal, D. M.; Giardina,

[146924] S. L.; Roder, J.; Anderson, S.; Ortaldo, J. R.: Mechanism of target

[146925] cell recognition by natural killer cells: characterization of a novel

[146926] triggering molecule restricted to CD3- large granular lymphocytes. J.

[146927] Exp. Med. 174: 1527-1536, 1991.

[146928]26419.

[146929]Rinfret, A.; Anderson, S. K.: IL-2 regulates the expression
of

[146930]the NK-TR gene via an alternate RNA splicing mechanism.
Molec. Immun. 30:

[146931]1307-1313, 1993.

[146932]26420.

[146933]Simons-Evelyn, M.; Young, H. A.; Anderson, S. K.: Charac-
terization

[146934]of the mouse Nktr gene and promoter. Genomics 40:
94-100, 1997.

[146935]26421.

[146936]Young, H. A.; Jenkins, N. A.; Copeland, N. G.; Simek, S.;
Lerman,

[146937]M. I.; Zbar, B.; Glenn, G.; Ortaldo, J. R.; Anderson, S. K.:
Localization

[146938]of a novel natural killer triggering receptor locus to hu-
man chromosome

[146939]3p23-p21 and mouse chromosome 9. Genomics 16:
548-549, 1993.

[146940]26422.

[146941] Scott, I. C.; Halila, R.; Jenkins, J. M.; Mehan, S.; Apostolou,
[146942] S.; Wingvist, R.; Callen, D. F.; Prockop, D. J.; Peltonen, L.;
Kadler,
[146943] K. E.: Molecular cloning, expression and chromosomal lo-
calization
[146944] of a human gene encoding a 33 kDa putative metallopep-
tidase (PRSM1). Gene 174:
[146945] 135–143, 1996.
[146946] 26423.
[146947] Baldwin, A. S., Jr.: The NF- κ B and I- κ B pro-
teins: new
[146948] discoveries and insights. Annu. Rev. Immun. 14: 649–683,
1996.
[146949] 26424.
[146950] Chen, F.; Castranova, V.; Shi, X.; Demers, L. M.: New in-
sights
[146951] into the role of nuclear factor- κ B, a ubiquitous tran-
scription
[146952] factor in the initiation of diseases. Clin. Chem. 45: 7–17,
1999.
[146953] 26425.

[146954]Liptay, S.; Schmid, R. M.; Perkins, N. D.; Meltzer, P.; Al-
therr,
[146955]M. R.; McPherson, J. D.; Wasmuth, J. J.; Nabel, G. J.: Related
subunits
[146956]of NF-kappa-B map to two distinct loci associated with
translocations
[146957]in leukemia, NFKB1 and NFKB2. Genomics 13: 287-292,
1992.
[146958]26426.

[146959]Mathew, S.; Murty, V. V. V. S.; Dalla-Favera, R.; Chaganti,
R.
[146960]S. K.: Chromosomal localization of genes encoding the
transcription
[146961]factors, c-rel, NF-kappa-Bp50, NF-kappa-Bp65, and lyt-
10 by fluorescence
[146962]in situ hybridization. Oncogene 8: 191-193, 1993.
[146963]26427.

[146964]Ozes, O. N.; Mayo, L. D.; Gustin, J. A.; Pfeffer, S. R.; Pfe-
fer,
[146965]L. M.; Donner, D. B.: NF-kappa-B activation by tumour
necrosis factor
[146966]requires the Akt serine-threonine kinase. Nature 401:

82–85, 1999.

[146967]26428.

[146968]Romashkova, J. A.; Makarov, S. S.: NF-kappa-B is a target
of

[146969]AKT in anti-apoptotic PDGF signalling. Nature 401:
86–90, 1999.

[146970]26429.

[146971]Zhong, H.; May, M. J.; Jimi, E.; Ghosh, S.: The phosphory-
lation

[146972]status of nuclear NF-kappa-B determines its association
with CBP/p300

[146973]or HDAC-1. Molec. Cell 9: 625–636, 2002.

[146974]26430.

[146975]Claudio, E.; Brown, K.; Park, S.; Wang, H.; Siebenlist, U.:
BAFF-induced

[146976]NEMO-independent processing of NF-kappa-B2 in matur-
ing B cells. Nature

[146977]Immun. 3: 958–965, 2002.

[146978]26431.

[146979]Neri, A.; Chang, C.-C.; Lombardi, L.; Salina, M.; Corradini,
P.;

- [146980]Maiolo, A. T.; Chaganti, R. S. K.; Dalla-Favera, R.: B cell lymphoma-associated
- [146981]chromosomal translocation involves candidate oncogene
- [146982]lyt-10, homologous
- [146982]to NF-kappa-B p50. Cell 67: 1075-1087, 1991.
- [146983]26432.
- [146984]Adams, B. S.; Leung, K.; Meltzer, P. S.; Lewis, K. A.; Wagner-McPherson,
- [146985]C.; Evans, G. A.; Nabel, G. J.: Localization of the gene encoding
- [146986]R-kappa-B (NFRKB), a tissue-specific DNA binding protein, to chromosome
- [146987]11q24-q25. Genomics 14: 270-274, 1992.
- [146988]26433.
- [146989]Sanyal, S.; Sandstrom, D. J.; Hoeffler, C. A.; Ramaswami, M.:
- [146990]AP-1 function upstream of CREB to control synaptic plasticity in Drosophila. Nature 416:
- [146991]870-874, 2002.
- [146992]26434.
- [146993]Arheden, K.; Mandahl, N.; Strombeck, B.; Isaksson, M.;

Mitelman,

[146994]F.: Chromosome localization of the human oncogene INT1 to 12q13 by

[146995]in situ hybridization. Cytogenet. Cell Genet. 47: 86–87, 1988.

[146996]26435.

[146997]Gavin, B. J.; McMahon, J. A.; McMahon, A. P.: Expression of multiple

[146998]novel Wnt–1/int–1–related genes during fetal and adult mouse development. Genes

[146999]Dev. 4: 2319–2332, 1990.

[147000]26436.

[147001]Kirikoshi, H.; Sekihara, H.; Katoh, M.: WNT10A and WNT6, clustered

[147002]in human chromosome 2q35 region with head–to–tail manner, are strongly

[147003]coexpressed in SW480 cells. Biochem. Biophys. Res. Commun. 283:

[147004]798–805, 2001.

[147005]26437.

[147006]McMahon, A. P.: The Wnt family of developmental regula–

tors. Trends

[147007]Genet. 8: 236–242, 1992.

[147008]26438.

[147009]Nusse, R.; Brown, A.; Papkoff, J.; Scambler, P.; Shackleford, G.;

[147010]McMahon, A.; Moon, R.; Varmus, H.: A new nomenclature for int-1 and

[147011]related genes: the Wnt gene family. (Letter) Cell 64: 231–232, 1991.

[147012]26439.

[147013]Nusse, R.; van't Veer, L.; Geurts van Kessel, A.; van Agthoven,

[147014]A.; Bootsma, D.; Varmus, H.: Chromosomal localization of a human

[147015]homologue of a putative mammary tumor oncogene. (Abstract) Cytogenet.

[147016]Cell Genet. 37: 556–557, 1984.

[147017]26440.

[147018]Pellegrino, J. E.; Lensch, M. W.; Muenke, M.; Chance, P. F.: Clinical

[147019]and molecular analysis in Joubert syndrome. Am. J. Med.

Genet. 72:

[147020]59–62, 1997.

[147021]26441.

[147022]Thomas, K. R.; Capecchi, M. R.: Targeted disruption of the murine

[147023]int-1 proto-oncogene resulting in severe abnormalities in midbrain

[147024]and cerebellar development. Nature 346: 847–850, 1990.

[147025]26442.

[147026]Andrews, J.; Erdjument, H.; Nicholson, D. C.: Hereditary coproporphyria:

[147027]incidence in a large English family. J. Med. Genet. 21: 341–349,

[147028]1984.

[147029]26443.

[147030]Barnes, H. D.; Whittaker, N.: Hereditary coproporphyria with acute

[147031]intermittent manifestations. Brit. Med. J. 2: 1102–1104, 1965.

[147032]26444.

[147033]Barohn, R. J.; Sanchez, J. A.; Anderson, K. E.: Acute pe-

ripheral

[147034]neuropathy due to hereditary coproporphyria. Muscle
Nerve 17: 793-799,

[147035]1994.

[147036]26445.

[147037]Berger, H.; Goldberg, A.: Hereditary coproporphyria. Brit.
Med.

[147038]J. 2: 85-88, 1955.

[147039]26446.

[147040]Brodie, M. J.; Thompson, G. G.; Moore, M. R.; Beattie, A.
D.; Goldberg,

[147041]A.: Hereditary coproporphyria: demonstration of the ab-
normalities

[147042]in haem biosynthesis in peripheral blood. Quart. J. Med.
46: 229-241,

[147043]1977.

[147044]26447.

[147045]Cacheux, V.; Martasek, P.; Fougerousse, F.; Delfau, M. H.;
Druart,

[147046]L.; Tachdjian, G.; Grandchamp, B.: Localization of the hu-
man coproporphyrinogen

[147047]oxidase gene to chromosome band 3q12. Hum. Genet.
94: 557–559, 1994.

[147048]26448.

[147049]Connon, J. J.; Turkington, V.: Hereditary coproporphyria.
Lancet II:

[147050]263–264, 1968.

[147051]26449.

[147052]Cripps, D. J.; Peters, H. A.: Stool porphyrins in acute inter-
mittent

[147053]and hereditary coproporphyria: adverse effects of tran-
quilizers. Arch.

[147054]Neurol. 23: 80–84, 1970.

[147055]26450.

[147056]Delfau-Larue, M.-H.; Martasek, P.; Grandchamp, B.: Co-
protophyrinogene

[147057](sic) oxidase: gene organization and description of a mu-
tation leading

[147058]to exon 6 skipping. Hum. Molec. Genet. 3: 1325–1330,
1994.

[147059]26451.

[147060]Doss, M.; von Tiepermann, R.; Kopp, W.: Harderoporphyryn

coproporphyria.

[147061](Letter) Lancet I: 292 only, 1984.

[147062]26452.

[147063]Elder, G. H.; Evans, J. O.; Thomas, N.; Cox, R.; Brodie, M.
J.;

[147064]Moore, M. R.; Goldberg, A.; Nicholson, D. C.: The primary
enzyme

[147065]defect in hereditary coproporphyria. Lancet II: 1217-1219,
1976.

[147066]26453.

[147067]Goldberg, A.; Rimington, C.; Lochhead, A. C.: Hereditary
coproporphyria. Lancet I:

[147068]632-636, 1967.

[147069]26454.

[147070]Grandchamp, B.; Phung, N.; Nordmann, Y.: Homozygous
case of hereditary

[147071]coproporphyria. (Letter) Lancet II: 1348-1349, 1977.

[147072]26455.

[147073]Grandchamp, B.; Weil, D.; Nordmann, Y.; Van Cong, N.; de
Verneuil,

[147074]H.; Foubert, C.; Gross, M.-S.: Assignment of the human

coproporphyrinogen

[147075]oxidase to chromosome 9. Hum. Genet. 64: 180–183, 1983.

[147076]26456.

[147077]Haeger–Aronsen, B.; Stathers, G.; Swahn, G.: Hereditary coproporphyria:

[147078]study of a Swedish family. Ann. Intern. Med. 69: 221–227, 1968.

[147079]26457.

[147080]Hunter, J. A. A.; Khan, S. A.; Hope, E.; Beattie, A. D.; Beveridge,

[147081]G. W.; Smith, A. W. M.; Goldberg, A.: Hereditary coproporphyria.

[147082]Photosensitivity, jaundice and neuropsychiatric manifestations associated

[147083]with pregnancy. Brit. J. Derm. 84: 301–310, 1971.

[147084]26458.

[147085]Kohno, H.; Furukawa, T.; Yoshinaga, T.; Tokunaga, R.; Taketani,

[147086]S.: Coproporphyrinogen oxidase: purification, molecular cloning,

[147087]and induction of mRNA during erythroid differentiation. J.

Biol.

[147088]Chem. 268: 21359–21363, 1993.

[147089]26459.

[147090]Lamoril, J.; Deybach, J.-C.; Puy, H.; Grandchamp, B.; Nordmann,

[147091]Y.: Three novel mutations in the coproporphyrinogen oxidase gene. Hum.

[147092]Mutat. 9: 78–80, 1997.

[147093]26460.

[147094]Lamoril, J.; Martasek, P.; Deybach, J.-C.; Da Silva, V.; Grandchamp,

[147095]B.; Nordmann, Y.: A molecular defect in coproporphyrinogen oxidase

[147096]gene causing harderoporphyria, a variant form of hereditary coproporphyria. Hum.

[147097]Molec. Genet. 4: 275–278, 1995.

[147098]26461.

[147099]Lamoril, J.; Puy, H.; Gouya, L.; Rosipal, R.; Da Silva, V.; Grandchamp,

[147100]B.; Foint, T.; Bader-Meunier, B.; Dommergues, J. P.; Deybach, J. C.;

- [147101]Nordmann, Y.: Neonatal hemolytic anemia due to inherited harderoporphyria:
[147102]clinical characteristics and molecular basis. Blood 91:
1453–1457,
[147103]1998.
[147104]26462.
- [147105]Lamoril, J.; Puy, H.; Whatley, S. D.; Martin, C.; Woolf, J. R.;
[147106]Da Silva, V.; Deybach, J.-C.; Elder, G. H.: Characterization
of mutations
[147107]in the CPO gene in British patients demonstrates absence
of genotype–phenotype
[147108]correlation and identifies relationship between hereditary
coproporphyria
[147109]and harderoporphyria. Am. J. Hum. Genet. 68:
1130–1138, 2001.
[147110]26463.
- [147111]Lomholt, J. C.; With, T. K.: Hereditary coproporphyria: a
family
[147112]with unusually few and mild symptoms. Acta Med. Scand.
186: 83–85,
[147113]1969.

[147114]26464.

[147115]Martasek, P.; Camadro, J. M.; Delfau-Larue, M.-H.; Dumas, J.-B.;

[147116]Montagne, J. J.; De Verneuil, H.; Labbe, P.; Grandchamp, B.: Molecular

[147117]cloning, sequencing, and functional expression of a cDNA encoding

[147118]human coproporphyrinogen oxidase. Proc. Nat. Acad. Sci. 91: 3024-3028,

[147119]1994.

[147120]26465.

[147121]Martasek, P.; Nordmann, Y.; Grandchamp, B.: Homozygous hereditary

[147122]coproporphyrinemia caused by an arginine to tryptophane substitution in

[147123]coproporphyrinogen oxidase and common intragenic polymorphisms. Hum.

[147124]Molec. Genet. 3: 477-480, 1994.

[147125]26466.

[147126]McIntyre, N.; Pearson, A. J. G.; Allan, D. J.; Craske, S.; West,

[147127]G. M. L.; Moore, M. R.; Beattie, A. D.; Paxton, J.; Goldberg, A.:

[147128]Hepatic delta-aminolaevulinic acid synthetase in an attack of hereditary

[147129]coproporphyria and during remission. Lancet I: 560-564, 1971.

[147130]26467.

[147131]Nordmann, Y.; Grandchamp, B.; de Verneuil, H.; Phung, L.; Cartigny,

[147132]B.; Fontaine, G.: Harderoporphyria: a variant hereditary coproporphyria. J.

[147133]Clin. Invest. 72: 1139-1149, 1983.

[147134]26468.

[147135]Roberts, D. T.; Brodie, M. J.; Moore, M. R.; Thompson, G. G.;

[147136]Goldberg, A.; MacSween, R. N. M.: Hereditary coproporphyria presenting

[147137]with photosensitivity induced by the contraceptive pill. Brit. J.

[147138]Derm. 96: 549-554, 1977.

[147139]26469.

[147140] Dickinson, D. P.; Zhao, Y.; Thiesse, M.; Siciliano, M. J.: Direct

[147141] mapping of seven genes encoding human type 2 cystatins to a single

[147142] site located at 20p11.2. Genomics 24: 172–175, 1994.

[147143] 26470.

[147144] Gopal Rao, V. V.; Schnittger, S.; Abrahamson, M.; Hansmann, I.

[147145] Cystatin–C (CST3), the candidate gene for the hereditary cystatin–C

[147146] amyloid angiopathy (HCCAA) maps to or close to human chromosome 20p11.22.

[147147] (Abstract) Cytogenet. Cell Genet. 58: 2029 only, 1991.

[147148] 26471.

[147149] Saitoh, E.; Kim, H.–S.; Smithies, O.; Maeda, N.: Human cysteine–proteinase

[147150] inhibitors: nucleotide sequence analysis of three members of the cystatin

[147151] gene family. Gene 61: 329–338, 1987.

[147152] 26472.

[147153] Saitoh, E.; Sabatini, L. M.; Eddy, R. L.; Shows, T. B.; Azen,

E.

[147154]A.; Isemura, S.; Sanada, K.: The human cystatin C gene (CST3) is

[147155]a member of the cystatin gene family which is localized on chromosome

[147156]20. Biochem. Biophys. Res. Commun. 162: 1324–1331, 1989.

[147157]26473.

[147158]Cruzen, M. E.; Bengtsson, U.; McMahon, J.; Wasmuth, J. J.; Arfin,

[147159]S. M.: Assignment of the cysteinyl-tRNA synthetase gene (CARS) to

[147160]11p15.5. Genomics 15: 692–693, 1993.

[147161]26474.

[147162]Takahashi, E.; Hori, T.; O'Connell, P.; Leppert, M.; White, R.

[147163]; R-banding and nonisotopic in situ hybridization: precise localization

[147164]of the human type II collagen gene (COL2A1). Hum. Genet. 86: 14–16,

[147165]1990.

[147166]26475.

[147167]Nebert, D. W.; Gonzalez, F. J.: P450 genes: structure, evolution,
[147168]and regulation. Annu. Rev. Biochem. 56: 945–993, 1987.
[147169]26476.

[147170]Paolini, M.; Cantelli–Forti, G.; Perocco, P.; Pedulli, G. F.;
[147171]Abdel–Rahman, S. Z.; Legator, M. S.: Co–carcinogenic effect of beta–carotene.
[147172](Letter) Nature 398: 760–761, 1999.
[147173]26477.

[147174]Espinoza, H.; Cox, C. J.; Semina, E. V.; Amendt, B. A.: A
molecular
[147175]basis for differential developmental anomalies in Axen–
feld–Rieger
[147176]syndrome. Hum. Molec. Genet. 11: 743–753, 2002.
[147177]26478.

[147178]Letinic, K.; Zoncu, R.; Rakic, P.: Origin of GABAergic neurons
[147179]in the human neocortex. Nature 417: 645–649, 2002.
[147180]26479.

[147181]McGuinness, T.; Porteus, M. H.; Smiga, S.; Bulfone, A.;

Kingsley,

[147182]C.; Qiu, M.; Liu, J. K.; Long, J. E.; Xu, D.; Rubenstein, J. L. R.

[147183]: Sequence, organization, and transcription of the *Dlx-1*
and the *Dlx-2*

[147184]locus. *Genomics* 35: 473–485, 1996.

[147185]26480.

[147186]Ozcelik, T.; Porteus, M. H.; Rubenstein, J. L. R.; Francke, U.

[147187]: *DLX2* (*TES1*), a homeobox gene of the Distal-less family,
assigned

[147188]to conserved regions on human and mouse chromosomes
2. *Genomics* 13:

[147189]1157–1161, 1992.

[147190]26481.

[147191]Porteus, M. H.; Brice, A. E. J.; Bulfone, A.; Usdin, T. B.; Ciaranello,

[147192]R. D.; Rubenstein, J. L. R.: Isolation and characterization of
a

[147193]library of cDNA clones that are preferentially expressed in
the embryonic

[147194]telencephalon. *Molec. Brain Res.* 12: 7–22, 1992.

[147195]26482.

- [147196] Qiu, M.; Bulfone, A.; Martinez, S.; Meneses, J. J.; Shimamura,
[147197] K.; Pedersen, R. A.; Rubenstein, J. L.: Null mutation of Dlx-
2 results
[147198] in abnormal morphogenesis of proximal first and second
branchial arch
[147199] derivatives and abnormal differentiation in the forebrain.
Genes
[147200] Dev. 9: 2523–2538, 1995.
[147201] 26483.
[147202] Salvador, J. M.; Hollander, M. C.; Nguyen, A. T.; Kopp, J. B.;
[147203] Barisoni, L.; Moore, J. K.; Ashwell, J. D.; Fornace, A. J., Jr.:
[147204] Mice lacking the p53–effector gene Gadd45a develop a
lupus–like syndrome. Immunity 16:
[147205] 499–508, 2002.
[147206] 26484.
[147207] Takekawa, M.; Saito, H.: A family of stress–inducible
GADD45–like
[147208] proteins mediate activation of the stress–responsive
MTK1/MEKK4 MAPKKK. Cell 95:
[147209] 521–530, 1998.

[147210]26485.

[147211]Dhermy, D.; Garbarz, M.; Lecomte, M. C.; Feo, C.;
Bournier, O.;

[147212]Chaverroche, I.; Gautero, H.; Galand, C.; Boivin, P.: Heredi-
tary elliptocytosis:

[147213]clinical, morphological and biochemical studies of 38
cases. Nouv.

[147214]Rev. Franc. Hemat. 28: 129-140, 1986.

[147215]26486.

[147216]Fairbanks, G.; Steck, T. L.; Wallach, D. F. H.: Elec-
trophoretic

[147217]analysis of the major polypeptides of the human erythro-
cyte membrane. Biochemistry 10:

[147218]2606-2617, 1971.

[147219]26487.

[147220]Feo, C. J.; Fischer, S.; Piau, J. P.; Grange, M. J.; Tchernia,

[147221]G.: Premiere observation de l'absence d'une proteine de la
membrane

[147222]erythrocytaire (bande 4-1) dans un cas d'anemie ellipto-
cytaire familiale. Nouv.

[147223]Rev. Franc. Hemat. 22: 315-325, 1980.

[147224]26488.

[147225]Fishbein, W. N.; Armbrustmacher, V. W.; Griffin, J. L.;
Davis,

[147226]J. I.; Foster, W. D.: Levels of adenylate deaminase, adenylate kinase,

[147227]and creatine kinase in frozen human muscle biopsy specimens relative

[147228]to type1/type2 fiber distribution: evidence for a carrier state of

[147229]myoadenylate deaminase deficiency. Ann. Neurol. 15:
271-277, 1984.

[147230]26489.

[147231]Fishbein, W. N.; Griffin, J. L.; Nagarajan, K.; Winkert, J. W.;

[147232]Armbrustmacher, V. W.: Immunologic uniqueness of muscle adenylate

[147233]deaminase (mAD) and genetic transmission of the deficiency state.

[147234](Abstract) Clin. Res. 27: 274A only, 1979.

[147235]26490.

[147236]Genetta, T.; Morisaki, H.; Morisaki, T.; Holmes, E. W.: A novel

[147237]bipartite intronic splicing enhancer promotes the inclusion
of a mini-exon

[147238]in the AMP deaminase 1 gene. J. Biol. Chem. 276:
25589-25597, 2001.

[147239]26491.

[147240]Kar, N. C.; Pearson, C. M.: Muscle adenylate deaminase
deficiency:

[147241]report of six new cases. Arch. Neurol. 38: 279-281, 1981.

[147242]26492.

[147243]Kelemen, J.; Bradley, W. G.; DiMauro, S.: Reply to J. B. Shu-
mate.

[147244](Letter) Neurology 33: 1534 only, 1983.

[147245]26493.

[147246]Kelemen, J.; Rice, D. R.; Bradley, W. G.; Munsat, T. L.; Di-
Mauro,

[147247]S.; Hogan, E. L.: Familial myoadenylate deaminase defi-
ciency and

[147248]exertional myalgia. Neurology 32: 857-863, 1982.

[147249]26494.

[147250]Lecky, B. R. F.: Failure of D-ribose in myoadenylate deam-
inase

[147251]deficiency. (Letter) Lancet I: 193 only, 1983.

[147252]26495.

[147253]Loh, E.; Rebbeck, T. R.; Mahoney, P. D.; DeNofrio, D.;
Swain,

[147254]J. L.; Holmes, E. W.: Common variant in AMPD1 gene pre-
dicts improved

[147255]clinical outcome in patients with heart failure. Circulation
99:

[147256]1422–1425, 1999.

[147257]26496.

[147258]Morisaki, H.; Higuchi, I.; Abe, M.; Osame, M.; Morisaki, T.:

[147259]First missense mutations (R388W and R425H) of AMPD1
accompanied with

[147260]myopathy found in a Japanese patient. Hum. Mutat. 16:
467–472, 2000.

[147261]26497.

[147262]Morisaki, H.; Morisaki, T.; Newby, L. K.; Holmes, E. W.: Al-
ternative

[147263]splicing: a mechanism for phenotypic rescue of a common
inherited

[147264]defect. J. Clin. Invest. 91: 2275–2280, 1993.

[147265]26498.

[147266]Morisaki, T.; Gross, M.; Morisaki, H.; Pongratz, D.; Zollner,

[147267]N.; Holmes, E. W.: Molecular basis of AMP deaminase deficiency in

[147268]skeletal muscle. Proc. Nat. Acad. Sci. 89: 6457-6461,
1992.

[147269]26499.

[147270]Morisaki, T.; Sabina, R. L.; Holmes, E. W.: Adenylate deaminase:

[147271]a multigene family in humans and rats. J. Biol. Chem. 265:
11482-11486,

[147272]1990.

[147273]26500.

[147274]Morton, C. C.; Eddy, R. L.; Shows, T. B.; Clark, P. R. H.;
Sabina,

[147275]R. L.; Holmes, E. W.: Human AMP deaminase-1 gene
(AMPD1) is mapped

[147276]to chromosome 1. (Abstract) Cytogenet. Cell Genet. 51:
1048-1049,

[147277]1989.

[147278]26501.

[147279]Moseley, W. S.; Morisaki, T.; Sabina, R. L.; Holmes, E. W.;
Seldin,

[147280]M. F.: Ampd-2 maps to distal mouse chromosome 3 in
linkage with Ampd-1. Genomics 6:

[147281]572-574, 1990.

[147282]26502.

[147283]Sabina, R. L.; Morisaki, T.; Clarke, P.; Eddy, R.; Shows, T.
B.;

[147284]Morton, C. C.; Holmes, E. W.: Characterization of the hu-
man and rat

[147285]myoadenylate deaminase genes. J. Biol. Chem. 265:
9423-9433, 1990.

[147286]26503.

[147287]Sabina, R. L.; Ogasawara, N.; Holmes, E. W.: Expression of
three

[147288]stage-specific transcripts of AMP deaminase during myo-
genesis. Molec.

[147289]Cell. Biol. 9: 2244-2246, 1989.

[147290]26504.

[147291]Sabina, R. L.; Swain, J. L.; Olanow, C. W.; Bradley, W. G.;
Fishbein,

[147292]W. N.; DiMauro, S.; Holmes, E. W.: Myoadenylate deaminase deficiency:

[147293]functional and metabolic abnormalities associated with disruption

[147294]of the purine nucleotide cycle. J. Clin. Invest. 73: 720–730, 1984.

[147295]26505.

[147296]Sabina, R. L.; Swain, J. L.; Patten, B. M.; Ashizawa, T.; O'Brien,

[147297]W. E.; Holmes, E. W.: Disruption of the purine nucleotide cycle:

[147298]a potential explanation for muscle dysfunction in myoadenylate deaminase

[147299]deficiency. J. Clin. Invest. 66: 1419–1423, 1980.

[147300]26506.

[147301]Shumate, J. B.: Myoadenylate deaminase deficiency--a nonfamilial,

[147302]nondisease? (Letter) Neurology 33: 1533–1534, 1983.

[147303]26507.

[147304]Shumate, J. B.; Kaiser, K. K.; Carroll, J. E.; Brooke, M. H.:

[147305]Adenylate deaminase deficiency in a hypotonic infant. J.

Pediat. 96:

[147306]885–887, 1980.

[147307]26508.

[147308]Sinkeler, S. P. T.; Joosten, E. M. G.; Wevers, R. A.; Oei, T.

[147309]L.; Jacobs, A. E. M.; Veerkamp, J. H.; Hamel, B. C. J.:

Myoadenylate

[147310]deaminase deficiency: a clinical, genetic, and biochemical
study in

[147311]nine families. Muscle Nerve 11: 312–317, 1988.

[147312]26509.

[147313]Valen, P. A.; Nakayama, D. A.; Veum, J.; Sulaiman, A. R.;
Wortmann,

[147314]R. L.: Myoadenylate deaminase deficiency and forearm is-
chemic exercise

[147315]testing. Arthritis Rheum. 30: 661–668, 1987.

[147316]26510.

[147317]van den Berghe, G.; Hers, H. G.: Abnormal AMP deaminase
in primary

[147318]gout. (Letter) Lancet II: 1090 only, 1980.

[147319]26511.

[147320]Eddy, R. L.; Mahnke-Zizelman, D. K.; Bausch-Jurken, M.
T.; Sabina,

[147321]R. L.; Shows, T. B.: Distribution of the AMP deaminase
multigene

[147322]family within the human genome: assignment of the
AMPD2 to chromosome

[147323]1p21-p34 and AMPD3 to chromosome 11p13-pter.
(Abstract) Human Genome

[147324]Mapping Workshop 93 24 only, 1993.

[147325]26512.

[147326]Ogasawara, N.; Goto, H.; Yamada, Y.; Nishigaki, I.; Itoh, T.;
Hasegawa,

[147327]I.; Park, K. S.: Deficiency of AMP deaminase in erythro-
cytes. Hum.

[147328]Genet. 75: 15-18, 1987.

[147329]26513.

[147330]Sermusvitayawong, K.; Wang, X.; Nagabukuro, A.; Mat-
suda, Y.; Morisaki,

[147331]H.; Toyama, K.; Mukai, T.; Morisaki, T.: Genomic organi-
zation of

[147332]Ampd3, heart-type AMPD gene, located in mouse chro-
mosome 7. Mammalian

- [147333]Genome 8: 767–769, 1997.
- [147334]26514.
- [147335]Yamada, Y.; Goto, H.; Ogasawara, N.: Cloning and nucleotide sequence
[147336]of the cDNA encoding human erythrocyte-specific AMP
deaminase. Biochim.
[147337]Biophys. Acta 1171: 125–128, 1992.
- [147338]26515.
- [147339]Yamada, Y.; Goto, H.; Ogasawara, N.: A point mutation responsible
[147340]for human erythrocyte AMP deaminase deficiency. Hum.
Molec. Genet. 3:
[147341]331–334, 1994.
- [147342]26516.
- [147343]Allegrucci, C.; Liguori, L.; Mezzasoma, I.; Minelli, A.: A1
adenosine
[147344]receptor in human spermatozoa: its role in the fertilization process. Molec.
[147345]Genet. Metab. 71: 381–386, 2000.
- [147346]26517.

[147347]Bennhold, H.; Peters, H.; Roth, E.: Ueber einen Fall von
kompletter

[147348]Analbuminaemie ohne wesentliche klinische Krankheits-
zeichen. Verh.

[147349]Dtsch. Ges. Inn. Med. 60: 630–634, 1954.

[147350]26518.

[147351]Blumberg, B. S.; Martin, J. R.; Melartin, L.: Alloalbumine-
mia:

[147352]albumin Naskapi in Indians of the Ungava. J.A.M.A. 203:
180–185,

[147353]1968.

[147354]26519.

[147355]Boman, H.; Hermodson, M.; Hammond, C. A.; Motulsky, A.
G.: Analbuminemia

[147356]in an American Indian girl. Clin. Genet. 9: 513–526, 1976.

[147357]26520.

[147358]Brennan, S. O.: The molecular abnormality of albumin
Parklands:

[147359]365 asp-to-his. Biochim. Biophys. Acta 830: 320–324,
1985.

[147360]26521.

[147361] Brennan, S. O.; Arai, K.; Madison, J.; Laurell, C.-B.; Gal-
liano,

[147362] M.; Watkins, S.; Peach, R.; Myles, T.; George, P.; Putnam, F.
W.:

[147363] Hypermutable CpG dinucleotides in the propeptide-en-
coding sequence

[147364] of the human albumin gene. Proc. Nat. Acad. Sci. 87:
3909-3913,

[147365] 1990.

[147366] 26522.

[147367] Brennan, S. O.; Carrell, R. W.: A circulating variant of hu-
man

[147368] proalbumin. Nature 274: 908-909, 1978.

[147369] 26523.

[147370] Brennan, S. O.; Fellowes, A. P.: Albumin Hawkes Bay; a low
level

[147371] variant caused by loss of a sulphhydryl group at position
177. Biochim.

[147372] Biophys. Acta 1182: 46-50, 1993.

[147373] 26524.

[147374] Brennan, S. O.; Herbert, P.: Albumin Canterbury (313 lys-

to-asn):

[147375]a point mutation in the second domain of serum albumin.

Biochim.

[147376]Biophys. Acta 912: 191–197, 1987.

[147377]26525.

[147378]Brennan, S. O.; Myles, T.; Peach, R. J.; Donaldson, D.;

George,

[147379]P. M.: Albumin Redhill (–1 arg, ala320-to-thr): a glycoprotein variant

[147380]of human serum albumin whose precursor has an aberrant signal peptidase

[147381]cleavage site. Proc. Nat. Acad. Sci. 87: 26–30, 1990.

[147382]26526.

[147383]Campagnoli, M.; Rossi, A.; Palmqvist, L.; Flisberg, A.;

Niklasson,

[147384]A.; Minchotti, L.; Galliano, M.: A novel splicing mutation causes

[147385]an undescribed type of analbuminemia. Biochim. Biophys. Acta 1586:

[147386]43–49, 2002.

[147387]26527.

- [147388]Carlson, J.; Sakamoto, Y.; Laurell, C.-B.; Madison, J.;
Watkins,
[147389]S.; Putnam, F. W.: Alloalbuminemia in Sweden: structural
study and
[147390]phenotypic distribution of nine albumin variants. Proc.
Nat. Acad.
[147391]Sci. 89: 8225-8229, 1992.
[147392]26528.
- [147393]Cormode, E. J.; Lyster, D. M.; Israels, S.: Analbuminemia in
[147394]a neonate. J. Pediat. 86: 862-867, 1975.
[147395]26529.
- [147396]Dammacco, F.; Miglietta, A.; D'Addabbo, A.; Fratello, A.;
Moschetta,
[147397]R.; Bonomo, L.: Analbuminemia: report of a case and re-
view of the
[147398]literature. Vox Sang. 39: 153-161, 1980.
[147399]26530.
- [147400]Darlington, G.: Personal Communication. New Haven,
Conn. and
[147401]New York, N. Y. 9/17/1974.

[147402]26531.

[147403]Darlington, G. J.; Bernhard, H. P.; Ruddle, F. H.: Human
serum

[147404]albumin phenotype activation in mouse hepatoma-human
leukocyte cell

[147405]hybrids. Science 185: 859-862, 1974.

[147406]26532.

[147407]Dayhoff, M. O.: Serum albumin. Atlas of Protein Sequence
and

[147408]Structure. Washington, D. C.: National Biomedical Re-
search Foundation

[147409](pub.) 5: 1972. Pp. D316.

[147410]26533.

[147411]DeCosimo, D. R.; Fang, S.-L.; Braverman, L. E.: Prevalence
of

[147412]familial dysalbuminemic hyperthyroxinemia in Hispanics.
(Letter) Ann.

[147413]Intern. Med. 107: 780-781, 1987.

[147414]26534.

[147415]Dugaiczuk, A.: Personal Communication. Riverside, Calif.
4/29/1989.

[147416]26535.

[147417]Dugaiczky, A.; Law, S. W.; Dennison, O. E.: Nucleotide sequence

[147418]and the encoded amino acids of human serum albumin mRNA. Proc. Nat.

[147419]Acad. Sci. 79: 71–75, 1982.

[147420]26536.

[147421]Earle, D. P.; Hutt, M. P.; Schmid, K.; Gitlin, D.: Observations

[147422]on double albumin: a genetically transmitted serum protein anomaly. J.

[147423]Clin. Invest. 38: 1412–1420, 1959.

[147424]26537.

[147425]Efremov, G. D.; Braend, M.: Serum albumin: polymorphism in man. Science 146:

[147426]1679–1680, 1964.

[147427]26538.

[147428]Esumi, H.; Okui, M.; Sato, S.; Sugimura, T.; Nagase, S.: Absence

[147429]of albumin mRNA in the liver of analbuminemic rats. Proc. Nat. Acad.

[147430]Sci. 77: 3215–3219, 1980.

[147431]26539.

[147432]Esumi, H.; Takahashi, Y.; Sato, S.; Nagase, S.; Sugimura, T.:

[147433]A seven–base–pair deletion in an intron of the albumin gene of analbuminemic

[147434]rats. Proc. Nat. Acad. Sci. 80: 95–99, 1983.

[147435]26540.

[147436]Esumi, H.; Takahashi, Y.; Sekiya, T.; Sato, S.; Nagase, S.; Sugimura,

[147437]T.: Presence of albumin mRNA precursors in nuclei of analbuminemic

[147438]rat liver lacking cytoplasmic albumin mRNA. Proc. Nat. Acad. Sci. 79:

[147439]734–738, 1982.

[147440]26541.

[147441]Fine, J. M.; Abdo, Y.; Rochu, D.; Rousseaux, J.; Dautrevaux, M.

[147442]: Identification of the human albumin variant 'Gainesville' with proalbumin

[147443]'Christchurch'. Blood Transf. Immunohemat. 26: 341–346,

1983.

[147444]26542.

[147445]Fine, J. M.; Marneux, M.; Rochu, D.: Human albumin genetic variants:

[147446]an attempt at a classification of European allotypes. Am. J. Hum.

[147447]Genet. 40: 278–286, 1987.

[147448]26543.

[147449]Franklin, S. G.; Wolf, S. I.; Ozdemir, Y.; Yuregir, G. T.; Isbir,

[147450]T.; Blumberg, B. S.: Albumin Naskapi variant in North American Indians

[147451]and Eti Turks. Proc. Nat. Acad. Sci. 77: 5480–5482, 1980.

[147452]26544.

[147453]Franklin, S. G.; Wolf, S. I.; Zweidler, A.; Blumberg, B. S.:

[147454]Localization of the amino acid substitution site in a new variant

[147455]of human serum albumin, albumin Mexico–2. Proc. Nat. Acad. Sci. 77:

[147456]2505–2509, 1980.

[147457]26545.

[147458]Fraser, G. R.; Harris, H.; Robson, E. B.: A new genetically determined

[147459]plasma-protein in man. Lancet I: 1023-1024, 1959.

[147460]26546.

[147461]Galliano, M.; Minchiotti, L.; Ferri, G.; Iadarola, P.; Zapponi,

[147462]M. C.; Fine, J. M.: Structural characterization of the human albumin

[147463]variant 'Pollibauer'. Blood Transf. Immunohemat. 27: 597-602, 1984.

[147464]26547.

[147465]Galliano, M.; Minchiotti, L.; Iadarola, P.; Ferri, G.; Zapponi,

[147466]M. C.; Castellani, A. A.: The amino acid substitution in albumin

[147467]Roma: 321 glu-to-lys. FEBS Lett. 233: 100-104, 1988.

[147468]26548.

[147469]Galliano, M.; Minchiotti, L.; Iadarola, P.; Porta, F.; Stoppini,

[147470]M.; Zapponi, M. C.; Ferri, G.; Castellani, A. A.: Genetic variants

[147471]of human serum albumin. Prog. Med. Lab. 2: 475-477, 1988.

[147472]26549.

[147473]Galliano, M.; Minchiotti, L.; Iadarola, P.; Stoppini, M.; Giagnoni,

[147474]P.; Watkins, S.; Madison, J.; Putnam, F. W.: Protein and DNA sequence

[147475]analysis of a 'private' genetic variant: albumin Ortonovo (glu505-to-lys). Biochim.

[147476]Biophys. Acta 1225: 27-32, 1993.

[147477]26550.

[147478]Galliano, M.; Minchiotti, L.; Iadarola, P.; Zapponi, M. C.; Ferri,

[147479]G.; Castellani, A. A.: Structural characterization of a chain termination

[147480]mutant of human serum albumin. J. Biol. Chem. 261: 4283-4287, 1986.

[147481]26551.

[147482]Galliano, M.; Minchiotti, L.; Porta, F.; Rossi, A.; Ferri, G.;

[147483]Madison, J.; Watkins, S.; Putnam, F. W.: Mutations in genetic variants

[147484]of human serum albumin found in Italy. Proc. Nat. Acad. Sci. 87:

[147485]8721–8725, 1990.

[147486]26552.

[147487]Galliano, M.; Minchiotti, L.; Stoppini, M.; Tarnoky, A. L.: A

[147488]new proalbumin variant: albumin Jaffna (–1 arg-to-leu).

FEBS Lett. 255:

[147489]295–299, 1989.

[147490]